A standardized database of Marine Isotopic Stage 5e sea-level proxies on tropical Pacific islands

Nadine Hallmann¹, Gilbert Camoin¹, Jody M. Webster², Marc Humblet³

¹Aix-Marseille Université, CNRS, IRD, INRAE, Coll France, CEREGE, Europôle Méditerranéen de l'Arbois, BP80, 13545 Aix-en-Provence cedex 4, France

 ²Geocoastal Research Group, School of Geosciences, The University of Sydney, Sydney, New South Wales 2006, Australia
 ³Department of Earth and Planetary Sciences, Graduate School of Environmental Studies, Nagoya University,

Nagoya 464-8601, Japan

10 Correspondence to: Nadine Hallmann (hallmann@cerege.fr)

Abstract

5

Marine isotope stage 5 deposits have been reported on many tropical Pacific islands. This paper presents a database compiled through the review of MIS 5e (Last Interglacial – LIG) coral reef records from islands belonging to

- 15 French Polynesia (Anaa, Niau, Makatea, Moruroa, Takapoto, Bora Bora), the Hawaiian Islands (Oahu, Lana'i, Midway Atoll), Tuvalu, Kiribati (Christmas Island, Tarawa), the Cook Islands (Mangaia, Atiu, Mitiaro, Mauke, Pukapuka, Rakahanga, Rarotonga), Tonga, Samoa, the Federal States of Micronesia, the Mariana Islands, the Marshall Islands (Enewetak, Bikini), New Caledonia, Papua New Guinea, <u>the Solomon Islands</u>, Vanuatu, Fiji and Niue. Studies reporting other sea-level indicators dated to other Pleistocene interglacials and Holocene sea-level
- 20 indicators were not inserted in the database, but are included in this data description paper for completeness. Overall, about 300 studies concerning Pleistocene and Holocene sea-level indicators have been reviewed, and finally <u>163_{rage}</u> data points and 94 relative sea-level (RSL) data points from 38_r studies on the MIS 5e have been inserted in the database. <u>Additional 155 age data points have been reviewed, i.e., the tropical Pacific islands</u> database contains 318 age data points. The main sea-level indicators include emerged coral reef terraces, but also
- 25 reef units recovered in drill cores from a few islands, thus reflecting the diversity of tectonic settings and sampling approaches. Future research should be directed towards better constrained RSL reconstructions, including more precise chronological data, more accurate elevation measurements as well as a better refinement of the palaeo-water depth significance of coralgal assemblages. The database for tropical Pacific islands is available open-access at this link: http://doi.org/10.5281/zenodo.3991672 (Hallmann and Camoin, 2020).

hat gelöscht: I

hat gelöscht: 284 hat gelöscht: 5

hat gelöscht: relative sea-level (
hat gelöscht:)	
hat gelöscht: T	
hat gelöscht: I	

1 Introduction

In this paper, we describe in detail the background information used to build the standardized database of tropical Pacific islands. This database was compiled as contribution to the World Atlas of Last Interglacial Shorelines

- 40 (WALIS) ESSD Special Issue. The database was created using the WALIS interface, that is available at this link: <u>https://warmcoasts.eu/world-atlas.html</u>. The WALIS interface allows the standardized insertion into a mySQL database data on relative sea-level indicators and associated ages. The database presented in this study has been downloaded from the interface as .xls file. The database presented here is open access and is available under this link: <u>http://doi.org/10.5281/zenodo.3991672</u> (Hallmann and Camoin, 2020). Each field in our database is
- 45 described at the following link: https://doi.org/10.5281/zenodo.3961543 (Rovere et al., 2020). The open access will enable the rapid and wide dissemination of the data that form the basement of this work and which in turn may trigger new studies in this field. The database presented here will be useful for scientists studying relative sea-level changes, reef geology, tectonics and geodynamics of the Pacific region, as well as palaeoceanographers. More than 200 studies on Marine Isotopic Stage (MIS) 5 coral reef records in the tropical Pacific have been
- 50 reviewed. Of those, data from 38 studies have been included in the database. The geographical distribution of the studied islands from fourteen nations is shown in Figure 1. Overall, the database contains 315 U-series data points and three electron spin resonance data points. Evaluation of the quality of radiometric ages includes the mineralogical assessment, but also the closed- versus open-system behaviour of radioisotopes. Data points obtained outside closed-system conditions are highlighted in the database. The information on the acceptance of
- 55 radiometric ages has been often directly taken from the original publication. In addition, the quality of the ages that are reported in the WALIS database has been reviewed by Chutcharavan and Dutton who compiled a U-series database for MIS 5e corals (see this Special Issue). We inserted in the database 163 age data points and 94 relative sea-level (RSL) data points from 38 studies on the MIS 5e. Additional 155 age data points inserted by Chutcharavan and Dutton have been reviewed, i.e., the tropical Pacific islands database contains 318 age data
- 60 points.

In the following, we first give an overview of the published literature related to <u>Last Interglacial (LIG)</u> sea-level indicators in the tropical Pacific islands, in order to give the reader a sense of the historical background upon which our review was done. Then, we describe the types of sea-level indicators as well as the elevation measurement techniques and sea-level datums reported in the literature and the method used to reconstruct MIS 5e palaeo-RSL.

65 We then report a detailed description of the data points located in each administrative province/region within the area of interest, where sea-level data was reviewed (Section 5). Section 6.1 contains the description of a number of studies where the presence of different peaks in the LIG sea-level record was discussed in literature. While in

hat gelöscht:	nine
hat gelöscht:	281

hat g	gelöscht:	have	been	inserted	in	the	database

	hat gelöscht: Those 284 data points have been summarized in 84 RSL indicators.
(hat gelöscht: I
(hat gelöscht:

our database we only reviewed specifically <u>MIS</u> 5e data, we encountered a number of studies where other Pleistocene shorelines were dated or described <u>(Sections 6.2 and 6.3)</u>, as well as Holocene sea-level data (Section 6.4). Major controversies regarding the MIS 5e are highlighted in Section 6.5, while uncertainties associated with estimating palaeo-RSL from the data into the database are detailed in Section 6.6. Large RSL uncertainties and

80 limitations in the accuracy of RSL reconstructions can be considered for studies concerning MIS 5e coral reef records from the tropical Pacific islands. The data quality is mostly affected by the quality of RSL data and, to a lesser extent, by the quality of age information (see Section 6.6), We further present future research directions (Section 7) that may stem from our data compilation. Future work using more accurate dating methods and elevation measurements as well as a more detailed identification of the collected coral samples are essential to

85 <u>better constrain the MIS 5e RSL dataset concerning the tropical Pacific islands.</u>

2 Literature overview

The sea-level database described by this article focuses on tropical Pacific islands, which cover most of the

- 90 intertropical realm of the Pacific Ocean from the Hawaiian archipelago to the North to New Caledonia to the South, and from Mariana Islands and Papua New Guinea to the West to French Polynesia to the East. Last Interglacial (LIG) coral reef records of relative sea-level (RSL) change on tropical Pacific islands were mostly investigated on subaerially exposed reef terraces. Since the pioneer work of Veeh (1966), which focused on the dating of corals from different islands (Hawaiian Islands, French Polynesia, Cook Islands), extensive studies have
- 95 been conducted in many islands across this vast region. Some of these studies have used the LIG shorelines to reconstruct tectonic uplift, especially in Papua New Guinea and Vanuatu. Submerged coral reef deposits related to the LIG, forming locally well-identified terraces, have been investigated since the beginning of the 20th century and the first drilling in Funafuti (Tuvalu). Drill cores including reef units related to the LIG have been collected between the 1950s and the 1980s on Enewetak (Marshall Islands), Midway Atoll, Moruroa (French Polynesia) and
- 100 Tarawa (Kiribati), and more recently in New Caledonia and Bora Bora (French Polynesia).
 All reviewed studies concerning LIG reef records from tropical Pacific islands (Fig. 1) are summarized hereafter.

hat gelöscht: Marine Isotopic Stage

hat gelöscht: . These are listed in

hat gelöscht:

In the final two sections, we discuss further details on other metadata on palaeo-sea-level indicators that are not included in our database, but that might be useful as research on Quaternary shorelines will progress in the tropical Pacific regions.



Figure 1: Map showing the tropical Pacific islands considered in this article (blue dots).

- French Polynesia (Fig. 2): Since the early 1960s, the geology of Moruroa has been investigated through the
 drilling of a number of deep vertical and inclined boreholes carried out in the modern lagoon and on the reef rim.
 Lalou et al. (1966) and Labeyrie et al. (1969) studied reef units from the last 300,000 years. Trichet et al. (1984) described the general stratigraphy of Moruroa based on three cores. Camoin et al. (2001) dated four 300 m-long drill cores from the reef rim and reconstructed RSL changes over the past 300,000 years, including the identification of MIS 5e. The most recent study on a Moruroa drill core by Braithwaite and Camoin (2011) focused
- 120 on the study of carbonate sedimentology and diagenetic features of these cores.



Figure 2: Map showing islands in French Polynesia considered in this article (blue dots). Study sites are indicated by white circles. © Google Earth 2020 (Figs 2B-F).

The earliest study on emerged Pleistocene coral reef terraces has been performed by Veeh (1966) who dated *in situ* fossil corals on Anaa, Niau and Makatea, followed by Pirazzoli et al. (1988a) who dated LIG reefs on Anaa. Montaggioni (1985), Montaggioni et al. (1985) and Montaggioni and Camoin (1997) studied LIG reef terraces on Makatea. More recent studies by Montaggioni et al. (2018, 2019a, b) considered that some reef units have formed

130 during Pleistocene high RSL, including MIS 5e, on Takapoto, but did not provide dating results. In addition, Gischler et al. (2016, 2019) reported the occurrence of a Pleistocene reef unit attributed to MIS 5e in Bora Bora drill cores. Cook Islands (Figs 3 and 4): In 1910, Schofield reported on late Quaternary RSL changes on Rarotonga in the southern Cook Islands. The earliest study providing datings on LIG reef limestones in the southern Cook Islands (Mangaia) has been performed by Veeh (1966). Subsequent studies have been conducted by Stoddart et al. (1985), Spencer et al. (1987, 1988, 1989) and Stoddart et al. (1990). Woodroffe et al. (1991) worked on the stratigraphy and chronology of the southern Cook Islands and presented new U-series ages for Atiu, Mitiaro and Mauke. Gray et al. (1992) studied the geochronology and subsurface stratigraphy of Pukapuka and Rakahanga in the northern

140 Cook Islands based on drill cores from the lagoon.



Figure 3: Map showing the northern Cook Islands. Study site on Pukapuka is indicated by a white circle. © Google Earth 2020.



Figure 4: Map showing islands in the southern Cook Islands considered in this article (white circles). Study sites are indicated by white circles. © Google Earth 2020.

Samoa: The early study by Kear and Wood (1959) described LIG reefs in Samoa. In 1968, Stice and McCoy studied the geology of the Manu'a Islands and reported on the occurrence of carbonate deposits.

Niue (Fig. 5): Different terraces on Niue have been explored at the beginning of the 20th century (Agassiz, 1903; David, 1904). Schofield (1959) identified seven terraces. A LIG reef sequence has been reported by Paulay (1988), Paulay and Spencer (1992), Spencer and Paulay (1994) and Wheeler (2000). More recent studies further described

155 the different terraces on Niue (Terry and Nunn, 2003; Nunn and Britton, 2004; Kennedy et al., 2012).



Figure 5: Map showing the island of Niue. Study site on Niue is indicated by a white circle. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA,

160 METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp (Fig. 5A). © Google Earth 2020 (Fig. 5B).

Tonga (Fig. 6): Taylor (1978) and Yonekura (1983) have attributed reef deposits from the Tongatapu block to the LIG.



165

Figure 6: Map showing Tonga. Study site on Tongatapu is indicated by a blue circle. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp.

- 170 Fiji (Fig. 7): Despite pioneer geomorphological works by Dana (1872) and Moore (1889), and reports on late Quaternary RSL changes by Schofield (1910), LIG deposits from Fiji were reported for the first time in the late 1970s by Taylor (1978) who dated reefs on Yasawa Island. In the late 1990s, Nunn and Omura (1999) studied exposed reef terraces around Kadavu Island. The most recent studies by Nunn et al. (2002) and Nunn and Omura (2003) summarized the levels of emerged LIG shorelines on the northeastern Fiji Islands and compared them to
- 175 those from the western Fiji Islands.



Figure 7: Map showing islands in Fiji considered in this article (white circles). Study sites on Kaibu Island are indicated by white circles. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp (Fig. 7A). © Google
 Earth 2020 (Figs 7B-C).

Tuvalu: Data regarding LIG deposits from Tuvalu are restricted to the 340 m-long drill core that was carried out by the Coral Reef Expedition of The Royal Society in 1896–1898 on Funafuti Atoll. This drill core was used to interpret the origin and history of atoll formation (Bonney, 1904) and test the Darwin's subsidence theory (Darwin, 1842). Despite a detailed study of these reef cores (Bonney, 1904), the core was originally interpreted as a modern reef deposit overlying Pleistocene limestone (Hinde, 1904) and no chronological frame was established before Ohde et al. (2002) who considered that LIG reef deposits must be deeper than 24.2 m below the modern surface. Since then, the Funafuti cores have been mentioned in general reviews regarding the development of coral reef islands (e.g., McLean and Woodroffe 1994; Spencer et al., 2008).

190 Kiribati (Fig. 8): In 1981, 30 m-long drill cores reaching Pleistocene limestone have been collected on Tarawa during a hydrogeological investigation by an Australian government team. These cores have been dated later by Marshall and Jacobson (1985). Woodroffe and McLean (1998) reported U-series ages from an outcrop of LIG limestone on Christmas Island.



195 Figure 8: Map showing islands in Kiribati considered in this article (white circles). Study sites on Tarawa and Christmas Island are indicated by white circles. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp (Figs 8A-B). © Google Earth 2020 (Figs 8C-D).

Solomon Islands: Stearns (1945) studied elevated benches in the South-West Pacific and correlated the different levels between the Solomon Islands, Vanuatu, the Mariana and Hawaiian Islands. The geology of the Solomon Islands was then studied in the 1950s by Grover (1958), while their geomorphology has been investigated later by
 Stoddart (1969) who reported on elevated shorelines, terraces and tidal notches on different island groups.

Vanuatu (Fig. 9): Uplifted Quaternary reef terraces on Vanuatu islands, specifically Santo, Malakula and Efate, have been identified and partially described since the late 1960s and the 1970s (Mitchell, 1966, 1969, 1971; Robinson, 1969; Mallick and Greenbaum, 1977; Ash et al., 1978). After these first descriptions, a reconnaissance

- 210 study has been carried out by Neef and Veeh (1977) who published the first U-series ages on Malo and Efate, immediately followed by Bloom et al. (1978) who focused on the dating of LIG reefs from Efate. In the 1980s, the vertical movements in the New Hebrides (Vanuatu) Island Arc have been reconstructed based on the dating of the uplifted coral reef terraces and several investigations provided additional dating results on terraces from Santo, Malakula, Efate and Torres Islands (Jouannic et al. 1980, 1982; Lecolle and Bernat, 1985; Taylor et al., 1980,
- 215 1985). Taylor et al. (1987) compared contemporary coseismic and nonseismic with quaternary vertical movements based on the analysis of partially emerged corals and reef terraces in the central Vanuatu Arc. Edwards et al. (1987a, b) have provided precise chronological data on exposed reef terraces from Efate. Chronological data obtained on uplifted coral reef terraces have been used by Taylor et al. (2005) to reconstruct rapid vertical movements in the New Hebrides forearc. Since the beginning of the current century, additional studies have been
- 220 carried out on Malakula (Cabioch and Ayliffe, 2001) and Tanna (Neef et al., 2003) and significantly advanced the chronological frame of Pleistocene raised reefs. On Tanna, the studied reefs and raised lagoonal deposits range in age from the Holocene to MIS 7 (Neef et al., 2003).



Figure 9: Map showing islands in Vanuatu considered in this article (green and orange circles). Study sites are indicated 225 by white and green circles. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp.

New Caledonia (Fig. 10): Early studies on Pleistocene reef terraces, including the LIG, have been carried out in the 1970s and early 1980s by Launay and Récy (1972), Coudray (1976), Bernat et al. (1976), Marshall and Launay (1978) and Gaven and Bourrouilh-Le Jan (1981), and were focused on their dating and use for the reconstruction

- 230 of neotectonic movements in the Loyalty Islands (Maré, Lifou and Ouvéa). These results have been summarized by Maurizot and Lafoy (2003) in their geological map of Maré Island. The drilling of modern reefs from the western coast of New Caledonia in the 1970s (Coudray, 1971), late 1990s (Cabioch et al., 1996, 1999) and after 2000 (Frank et al., 2006; Cabioch et al., 2008; Montaggioni et al., 2011; Hongo and Wirrmann, 2015) aimed at reconstructing the development pattern of fringing and barrier reefs. Drill cores from the Chesterfield Islands that
- 235 are located about 500 km to the WNW of New Caledonia have been studied with the same objective by Degaugue-Michalski (1993).





Figure 10: Map showing islands in New Caledonia considered in this article (green and white circles). Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp.

Papua New Guinea (Fig. 11): Uplifted Quaternary reef terraces from Huon Peninsula (Papua New Guinea) represent a classical site of RSL change studies in support of the astronomical theory of climate change. Early studies and first attempts to establish an accurate chronological frame for these coral reef terraces were carried out
in the 1970s by Veeh and Chappell (1970), Chappell (1974) and Bloom et al. (1974). Since then, the uplifted terraces from Huon Peninsula have been regularly investigated with the same general objective in the 1980s by Aharon (1983), Chappell (1983) and Aharon and Chappell (1986), in the 1990s by Stein et al. (1993) and Esat et al. (1999) and, in the early 2000s, by Cutler et al. (2003).

hat gelöscht: with





Figure 11: Map showing study sites on the Huon Peninsula, Papua New Guinea, considered in this article (orange circle). Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp.

Federated States of Micronesia: Ayers and Vacher (1986), Anthony (1996) and Fletcher and Richmond (2010) mentioned Pleistocene limestones, including the LIG period, in this region.

- 260 Mariana Islands (Fig. 12): The geologic history of Guam was first studied in 1937 by Stearns who then reported in 1940 on the deposition of Pleistocene limestone on volcanic substrate. Stearns (1945), Tayama (1952) and Emery (1962) described 'shore benches' related to sea level. In 1954, Cloud correlated stratigraphic sections throughout the West Pacific, including Guam. In the early 1950s, a programme by the Corps of Engineers, U. S. Army and the U. S. Geological Survey has focused on geology, soils and water of Guam (Tracey et al., 1959,
- 265 1964), including a detailed description of submarine and emerged coral reef terraces. Lower terraces on Guam, less than 15 m above modern sea level, are comparable to limestones that were described at the same elevation on Saipan, Northern Mariana Islands, (Tracey et al., 1964) and assigned to the late Pleistocene (Cloud et al., 1956). Further studies on Guam reef terraces included their lithological description (Emery, 1963), the provision of U-series ages obtained on MIS 5e limestone and estimated tectonic movements (Randall and Siegrist, 1996) and the
- 270 reconstruction of RSL changes since MIS 5e (Miklavič et al., 2012). Following up the work by Cloud et al. (1956) on the geology of Saipan, Weary and Burton (2011) mapped the island and stated that the Tanapag Limestone is usually found at elevations of less than 30 m. The latest study by Muhs et al. (2020) provided U-series ages for the emergent reef of the Tanapag Limestone in Saipan, which formed during the LIG period.



Marshall Islands: Five drill cores extracted from Bikini atoll in 1947 were first studied by Ladd et al. (1948) who described subaerial horizons in those cores. In 1954, Emery et al. further described cores from Bikini. Drillings on Enewetak have been performed in the early 1950s and drill cores have been studied by Ladd et al. (1953), Potratz et al. (1955), Ladd and Schlanger (1960), Schlanger (1963) and Thurber et al. (1965). Schlanger (1963) has shown

- 285 similarities between cores from Enewetak and Bikini. Two shallow drilling programmes have been initiated in the early 1970s: PACE (1971-1972; Henry et al., 1974) and EXPOE with 46 holes down to 90 m on Enewetak (1973-1974, Couch et al., 1975). Couch et al. (1975) confirmed the occurrence of subaerial unconformities in many drill cores. Tracey and Ladd (1974) described unconformities in the Bikini drill core, which could be partly also recognized in Enewetak drill cores. Szabo et al. (1985) presented dating results from two drill cores extracted from
- 290 the Enewetak reef flat during the PACE Program. During the Pacific Enewetak Atoll Crater Exploration (PEACE) Program (1984-1985), the U.S. Geological Survey investigated two craters formed by nuclear detonations in 1958 (Henry et al., 1986; Henry and Wardlaw, 1990). Ludwig et al. (1988) studied a 350 m-long drill core of Neogene lagoonal, shallow-water carbonates from Enewetak. Quinn and Matthews (1990), Wardlaw and Quinn (1991) and Quinn (1991) studied the post-Miocene sea-level history of Enewetak.

295

Hawaii (Fig. 12): Stearns (1935a, b, c) and Wentworth and Hoffmeister (1939) first studied Pleistocene shorelines on the islands of Oahu and Maui. In 1960, drill cores have been collected on Midway Atoll and studied in the following decades (e.g., Ladd et al., 1967, 1970; Wells, 1982). Stearns and Chamberlain (1967) described deep cores extracted from Oahu to better understand the geologic history. Several studies provided ages and

- 300 demonstrated that the Waimanalo limestone on Oahu correspond to the LIG. Veeh (1966) and Ku et al. (1974) provided the first dating results of those emerged terraces. Further chronological studies concerning the Waimanalo deposits included those from Sherman et al. (1993), Muhs and Szabo (1994) and Szabo et al. (1994). In 2002, Muhs et al. performed additional datings with a higher precision at the Szabo et al. (1994) study sites. The latest ages of reef deposits from Oahu have been reported by <u>Hearty et al. (2007) and McMurtry et al. (2010)</u>.
- 305 Rubin et al. (1995, 2000) provided U-series data for the Hulopoe gravel on Lana'i containing corals corresponding to MIS 5e and MIS 7.

hat gelöscht: 2

hat gelöscht: MIS 5e and 7



hat gelöscht: 2

310 Figure 13: Map showing islands in Hawaii considered in this article (green and orange circles). Study sites on Lana'i and Oahu are indicated by blue and green circles. Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp.

315 3 Sea-level indicators

The main features associated with Last Interglacial (LIG) relative sea levels (RSL) on tropical Pacific islands are exposed coral reef terraces. In addition, submerged coral reef deposits related to the LIG, forming locally wellidentified terraces, have been drilled on land on several islands, such as Funafuti (Tuvalu), Enewetak (Marshall Islands), Midway Atoll, Moruroa and Bora Bora (French Polynesia), Tarawa (Kiribati) and New Caledonia.

320

Name of RSL indicator	Description of RSL indicator	Description of Reference Water Level	Description of Indicative Range	Indicator reference(s)
Coral reef terrace (general description)	Coral-built flat surface, corresponding to shallow- water reef terrace to reef crest. The definition of indicative meaning is derived from Rovere et al., 2016, and it represents the broadest possible indicative range, that can be refined with information on coral living ranges.	(Mean Lower Low Water + Breaking depth)/2	Mean Lower Low Water - Breaking depth	Rovere et al., 2016

Table 1: Different types of RSL indicators reviewed in this study.

In general, palaeo_RSL is determined from the average elevation of the terrace or, if present, from the elevation of the highest *in situ* corals which are usually found on the palaeo reef crest (Rovere et al., 2016). However, the identification of coral assemblages that characterize coral reef terraces and the definition of the relevant water-depth range estimates may provide a more accurate indication of the palaeo_RSL. Single corals and coral assemblages have been described in some studies related to LIG reefs on tropical Pacific islands. Since their palaeo-water depth significance has been barely reported in the relevant studies, we have compiled data published in two databases: OBIS (Ocean Biodiversity Information System; https://obis.org) and IUCN (International Union

330 for Conservation of Nature; https://www.iucnredlist.org), thus following a similar approach to the one used by Hibbert et al. (2016, 2018). The datasets used are listed on the OBIS and IUCN websites and summarized <u>in Table</u> 2, which presents the best estimates of palaeo-water depth intervals for corals that were quoted in the literature. hat gelöscht:

hat gelöscht:

hat gelöscht: on

OBIS is a more detailed database, which allows to define distribution curves and to predict the depth ranges at which the relevant species can be found with the highest probability. Data were extracted from the OBIS database, using the Mapper tool. Two filters were used on Mapper: (1) Time range: 1990-2020, and (2) Depth range: 0-150 (to eliminate potential aberrant depths). Records which had the following characteristics were excluded from the

340 analysis: (1) those for which the difference between Max Depth and Min Depth were > 0,2 m, (2) those for which the "basis of record" was not "human observation" or "machine observation" (i.e., collection specimens were excluded).

All available data were used to define palaeo-water depth intervals for each species (Table 2). However, the data that have been compiled from OBIS and other databases cover the whole Indo-Pacific region. The value that has

- 345 been calculated for average and median depths can therefore differ when considering specific islands or areas of the studied area. In most cases, it has been difficult to assign a palaeo-water depth based on a single species, as most corals have a wide depth range. Accurate palaeo-water depth intervals can be usually defined based on coral assemblages. We have primarily selected data for which a precise depth, not a range, was available in the OBIS database.
- 350 For some species, which were not identified with certainty, we used close species to define a depth interval (e.g., for *Acropora* cf *aspera*, we have compiled data concerning close species that belong to the '*aspera* group' of Wallace, 1999: *A. aspera, A. pulchra*, and *A. millepora*).

Porites is the most frequent coral reported in LIG reefs from tropical Pacific islands. However, it cannot be generally used to constrain the palaeo-water depth interval as it displays a wide depth range when it is not identified

22

355 at the species level. The morphology of *Porites* colonies may give nevertheless some indications on a gross palaeowater depth distribution.

360

hat gelöscht: . hat gelöscht: ¶ Concerning hat gelöscht: , hat gelöscht: data were extracted

	OBIS Database				IUCN Database	
	Median depth	Average depth	Min depth	Max depth	Min depth	Max depth
<u>Acropora*</u>	<u>4.0</u>	<u>5.7</u>	<u>0.0</u>	<u>91.0</u>		
Acropora aspera	3.0	3.0	0.0	11.0	0.0	5.0
Acropora millepora	3.0	3.9	0.0	12.0	2.0	12.0
Acropora pulchra	3.0	4.0	0.0	18.0	1.0	20.0
Cyphastrea serailia	4.5	5.2	0.0	24.9	0.0	50.0
Dipsastraea	6.0	11.1	0.0	60.0		
Dipsastraea laxa	7.0	5.9	0.0	11.0	1.0	25.0
Gardineroseris planulata	3.0	5.9	0.0	50.0	2.0	30.0
Goniastrea	<u>6.0</u>	15.3	0.0	92.0		
Goniastrea retiformis	8.0	17.9	0.0	53.0	0.0	20.0
Goniastrea stelligera	0.0	13.6	0.0	91.0	0.0	20.0
Leptastrea	7.0	16.8	0.0	86.0		
Leptoria	4.0	7.5	0.0	86.0		
Leptoria phrygia	4.0	5.0	0.0	47.0	0.0	30.0
Oulophyllia crispa	6.0	6.4	0.0	50.0	2.0	30.0
Platygyra	4.0	6.9	0.0	84.0		
Platygyra lamellina	4.0	4.4	0.0	27.4	1.0	30.0
Platygyra sinensis	5.0	5.0	0.0	17.0	0.0	30.0
Plesiastrea versipora	6.0	5.8	0.0	30.0	1.0	40.0
Pocillopora	0.0	15.6	0.0	142.0		
Pocillopora meandrina	0.0	15.8	0.0	<u>98.0</u>	1.0	27.0
Porites*	0.0	18.5	0.0	142.0		
Porites lobata	0.0	19.2	0.0	98.0	0.0	30.0
Porites lutea	0.0	2.7	0.0	84.0	0.0	30.0

* Data for Indo-Pacific region only

Table 2: Modern depth distribution (in meters below <u>mean sea level</u> - MSL) of corals identified in LIG records from the tropical Pacific islands. This table is based on two databases OBIS (Ocean Biodiversity Information System; https://obis.org) and IUCN (International Union for Conservation of Nature; https://www.iucnredlist.org).

1

4 Elevation measurements

Almost all studies used auto/hand level to measure elevations of <u>relative sea-level (RSL)</u> indicators (Table 3); fewer studies used theodolite and rod or meter tape and rod. The elevation measurement technique has not been reported in many of the reviewed studies, but was most probably hand level or metered tape (Table 3). The sea-

390 level datums reviewed in this study include Mean Sea Level (MSL), Mean High Tide (MHT), Mean Low Tide (MLT), Mean Low Water Springs (MLWS) and modern reef (Table 4). In several studies the sea-level datum has not been reported, and has been assumed to be MSL (Table 4).

Measurement technique	Description	Typical accuracy
Total station or Auto/hand level	Total stations or levels measure slope distances from the instrument to a particular point and triangulate relative to the XYZ coordinates of the base station. The accuracy of this process depends on how well defined the reference point and on the distance of the surveyed point from the base station. Thus, it is necessary to benchmark the reference station with a nearby tidal datum, or use a precisely (DGPS) known geodetic point. The accuracy of the elevation measurement is also inversely proportional to the distance between the instrument and the point being measured.	±0.1/±0.2m for total station ±0.2/±0.4 m for hand level
Theodolite and rod	Elevation derived from triangulation with a theodolite.	Usually very precise, centimetric accuracy, depending on distance.
Barometric altimeter	Difference in barometric pressure between a point of known elevation (often sea level) and a point of unknown elevation. Not accurate and used only rarely in sea-level studies.	Up to ±20% of elevation measurement
Cross-section from publication	The elevation was extracted from a published sketch/topographic section.	Variable, depending on the scale of the sketch or topographic section.

Metered tape or rod	The end of a tape or rod is placed at a known elevation point, and the elevation of the unknown point is calculated using the metered scale and, if necessary, clinometers to calculate angles.	Up to ±10% of elevation measurement <u>.</u>
Distance from top of drill core	Distance from the top of drill core.	Depending on coring technique and sampling procedures.
Differential GPS	GPS positions acquired in the field and corrected either in real time or during post-processing with respect to the known position of a base station or a geostationary satellite system (e.g., Omnistar). Accuracy depends on satellite signal strength, distance from base station, and number of static positions acquired at the same location.	<u>±0.02/±0.08 m,</u> <u>depending on survey</u> <u>conditions and</u> <u>instruments used (e.g.,</u> <u>single-band vs dual-band</u> <u>receivers).</u>
Not reported	The elevation measurement technique was not reported, most probably hand level or metered tape.	20% of the original elevation reported added in root mean square to the sea level datum error <u>.</u>

Table 3: Measurement techniques used to establish the elevation of MIS 5e shorelines in the tropical Pacific islands.

Formatiert: Standard, Abstand Vor: 0 Pt.

395

I

Datum name	Datum description	Datum uncertainty
Mean Sea Level / General definition	General definition of MSL, with no indications on the datum to which it is referred to.	A datum uncertainty can be established on a case- by-case basis.
Modern reef	Upper surface of modern reef	A datum uncertainty may be established on a case- by-case basis.
Mean High Tide (MHT)	From NOAA definitions: "It is obtained by subtracting the mean of all the high waters from the mean of the higher high waters. One-half the average difference between the	Depending on the quality of tidal data used.

	two low waters of each tidal day observed over the National Tidal Datum Epoch."	
Mean Low Tide (MLT)	From tideschart.com: "The Mean Low Tide (or Mean Low Water) is the average height of all low tides in a given place, deriving from a long series of observations (NTDE) of all levels of low tide in that spot."	Depending on the quality of tidal data used.
Mean Low Water Springs (MLWS)	From www.coastalwiki.org: "The height of mean low water springs is the average throughout a year of the heights of two successive low waters during those periods of 24 hours (approximately once a fortnight) when the range of the tide is greatest."	Depending on the quality of tidal data used.
Not reported	The sea level datum is not reported and impossible to derive from metadata.	<u>N/A</u>

Table 4: Sea-level datums reviewed in this study.

5 Relative sea-level indicators

405

Because they correspond to easily accessible coral reef archives of sea-level change on many tropical and 400 subtropical coasts, the <u>Last Interglacial (LIG)</u> terraces have received much attention since the seminal works from Mayer (1917) and Daly (1920) on Samoa and other Pacific islands (Camoin and Webster, 2015).

Most of the tropical Pacific islands are underlain by the large northwestward-moving Pacific plate that is generated at the East Pacific Rise and the Pacific-Antarctic Ridge (see Fig. 1 in Neall and Trewick, 2008). The archipelagos and islands from the western part of the studied area developed in a complex geodynamic setting resulting from the relative motions of the Pacific and the Australian and Eurasian plates.

The sixteen archipelagos and islands reported in the database illustrate the exceptional diversity of tropical Pacific islands, which is mainly related to their dynamic and complex geological history. They have originated as linear chains of volcanic islands on the above plates, atolls, uplifted coral reef islands, fragments of continental crust, obducted portions of adjoining lithospheric plates, and have also resulted from subduction along convergent plate

410 margins (Neall and Trewick, 2008). This complex history has generated a great diversity of tectonic movements (uplift vs subsidence), which have guided the initiation and the evolution of LIG reefs described below.



The palaeo-relative sea level (RSL) has been reconstructed based on reported elevations of coral assemblages,

Figure 14: Reconstruction of palaeo-relative sea level (RSL) based on reported palaeo-elevations of Last Interglacial (LIG) corals and estimated palaeo-water depth intervals of their modern counterparts. The range bars indicate maximum, median and minimum depths. Modern corals are shown in grey and LIG corals in white.

420 5.1 French Polynesia

LIG reef deposits have been reported from outcrops on several Polynesian islands and from drill cores collected on Moruroa and Bora Bora.

The first dates on LIG reef terraces from French Polynesia, which can be found at elevations of +4 to +5 m, have been obtained by Veeh (1966) who dated *in situ* fossil corals from emerged Pleistocene reef terraces on Anaa,

Niau and Makatea. *Leptoria* specimens at elevations of +2 to +4 ± 0.5 m MLWS on Anaa have been dated at 110 ± 20 ka, 140 ± 30 ka and 150 ± 40 ka (WALIS U-series IDs 1731-1733; ANAA5, ANAA15 and ANAA16). However, the use of the alpha-counting technique resulted in large age uncertainties.

Pirazzoli et al. (1988a) used electron spin resonance (ESR) to date the 4-5 m-high raised reefs from Anaa (see Fig. 2 in Pirazzoli et al., 1988a). A *Leptoria*? coral colony in growth position with an aragonite content of 97% yielded

430 an age of 109 + 8 / -4 ka (WALIS ESR ID 103; 3AN14). The authors reported an elevation of 3.85 ± 0.2 m above mean sea level (MSL) and a palaeo-MSL at an elevation of +4.15 ± 0.2 m. Modern Leptoria is living at a median and average depths of 4 and 7.5 m, respectively (see Table 2). Consequently, the palaeo-RSL based on those four *Leptoria* specimen from Anaa is assumed to have been at $+8.75 \pm 2.3$ m (WALIS RSL ID 548).

- An emerged reef terrace at the same elevation has been dated by Veeh (1966) on Niau and Makatea. Two *Leptoria*specimens at hand-leveled elevations of +3 to +4 ± 0.5 m MLWS on Niau have been dated at 160 ± 40 ka and 120 ± 20 ka (WALIS U-series IDs 1735 and 1736; NIAU4 and NIAU6). The palaeo-RSL on Niau is assumed to have been at +9.25 ± 2.0 m (WALIS RSL ID 549). *Favia* (now called *Dipsastraea*; Huang et al., 2014) and *Leptoria* on Makatea at an elevation of +3 ± 0.5 m MLWS have yielded ages of 140 ± 30 ka and 100 ± 20 ka (WALIS U-series IDs 1737-1738; MAKA2 and MAKA3). The palaeo-RSL on Makatea is assumed to have been at +8.75 ± 1.8 m
- 440 (WALIS RSL ID 550). In 1997, Montaggioni and Camoin identified two well defined Pleistocene terraces on Makatea, one of which is at an elevation of +7 m and can be related to the 125-ka RSL highstand. It can be assumed that Makatea reached relative vertical stability just prior to the 120-ka marine transgression (Montaggioni et al., 1985; Montaggioni and Camoin, 1997).

LIG reef deposits have been reported in drill cores collected in Moruroa (Camoin et al., 2001) and Bora Bora
(Gischler et al., 2016). The reef units related to the LIG in Moruroa drill cores (see Figure 15, and Figure 2 in Braithwaite and Camoin, 2011) are mainly composed of coralgal frameworks built by an *Acropora gr. danairobusta / Porolithon onkodes* assemblage, which typifies a very shallow depositional environment (i.e., shallower

- robusta / Porolithon onkodes assemblage, which typifies a very shallow depositional environment (i.e., shallower than 6 m). However, only one out of six datings has been accepted (Camoin et al., 2001) and concerns a reworked colony of *Acropora* specimen at 92.7 m below MSL yielding an age of 124.4 ± 2 ka (WALIS U-series IDs 1017-1022; WALIS RSL ID 551; FIL5/30.92.7m). The subsidence rate of Moruroa has been estimated at 0.007-0.008
- 450

mm/yr (Camoin et al., 2001), i.e., about 1 m since 125 ka.



Figure 15: Distribution of radiometric ages and reconstruction of successive reef units (MIS = Marine Isotopic Stages) in Moruroa cores FIL 8/30 and 8/40 (from Camoin et al., 2001 and Braithwaite and Camoin, 2011). hat gelöscht: 3

Gischler et al. (2016) dated a barrier reef drill core from Bora Bora. Core TEV1 has been drilled at MSL and a *Pocillopora* specimen at 30.6 ± 0.6 m below MSL has given an age of 116.9 ± 1.1 ka (WALIS U-series ID 1869; TEV1). This sample is 100% aragonitic and the dating is considered as reliable. The subsidence rate of Bora Bora

460 $\,$ has been estimated at 0.05 to 0.14 mm/yr.

In a recent study by Montaggioni et al. (2019a) a 'eustatic sea-level' of about 7 m above MSL during MIS 5e has been considered on Takapoto, but no U-series age has been reported.

5.2 Cook Islands

455

LIG reef deposits have been reported from outcrops on several southern Cook Islands and from drill cores collected on Pukapuka, northern Cook Islands.

The southern Cook Islands are located in a complex tectonic setting related to volcanic processes since 1.65 Ma, the mean age for main shield-building phase (Te Manga) on Rarotonga (Thompson et al. 1998; Neal and Trewick, 2008). The formation of this volcanic system led to the subsidence of some islands (Rarotonga, Aitutaki, and Manuae) and the subsequent uplift of other islands related to a flexural response to loading of the ocean floor

(McNutt and Menard, 1978). However, this flexural response has been the subject of many debates and controversies regarding its course, timing and amplitude (see review in Woodroffe et al., 1991). Gray et al. (1992) reported an average subsidence rate of 0.03 to 0.06 mm/yr for the Cook Islands, resulting in a

- subsidence of up to 7.5 m since 125 ka. In contrast, uplift rates have been estimated at approximately 0.044 mm/yr 475 on Mangaia, 0.011 mm/yr on Atiu, and 0.005-0.007 mm/yr on Mitiaro and Mauke (Spencer et al., 1988). Uplift
- rates over the last 120 ka have been estimated at 0.07-0.10 mm/yr, 0.05 mm/yr, 0.03 mm/yr and 0.03-0.05 mm/yr for Mangaia, Atiu, Mitiaro and Mauke, respectively, based on a 6-m sea-level highstand during the LIG (Woodroffe et al., 1991). These rates would imply an uplift of these islands of about 3.6 to 12 m since 120 ka. Conversely, a subsidence rate of 0.02 mm/yr has been considered for Rarotonga (Woodroffe et al., 1991).
- 480 However, it is uncertain if the assumed 6-m LIG sea-level highstand for the Cook Islands is appropriate. Gray et al. (1992) reported a subsidence rate of 0.045 ± 0.028 mm/yr for Pukapuka in the northern Cook Islands. Exposed reef terraces can be found at different elevations, up to 15 m above MSL, on different southern Cook Islands, including Mangaia, Rarotonga, Atiu, Mauke and Mitiaro. The first datings of LIG reef limestones in the southern the Cook Islands have been provided by Veeh (1966). He dated *in situ* fossil corals from emerged
- 485 Pleistocene reef terraces on Mangaia using the alpha-counting technique. Two corals, *Favia* (now called *Dipsastraea*; Huang et al., 2014) and *Leptoria*, at a hand-leveled elevation of +2 ± 0.5 m MLWS have been dated at 90 ± 20 ka and 110 ± 20 ka, respectively (WALIS U-series IDs 1744 and 1746; MANG1 and MANG2; WALIS RSL ID 585). Modern *Leptoria* are living at median and average depths of 4 and 7.5 m, respectively; however, their depth range is large, from the surface down to more than 80 m (see Table 2). Spencer et al. (1988) dated four
- 490 Porites samples from an encrusting groove wall on Mangaia using the alpha-counting technique. All samples contained less than 1% of calcite and all datings have been accepted. The four samples at elevations of +15, +11, +3 and +13 m MLT have been dated at 118 ± 12 ka, 107 ± 18 ka, 101 ± 12 ka and 135 ± 15 ka, respectively (WALIS U-series IDs 1848 and 1850-1852; M2, M3, M47 and M54; 1-sigma uncertainty; WALIS RSL ID 584; see Fig. 1Q. The tidal range is approximately 80 cm, i.e., the sample elevations are of +15.4, +11.4, +3.4 and
- 495 +13.4 MSL. Porites is living at median and average depths of 0 and 18.5 m, respectively in modern environments; however, this genus displays a wide depth range, from the surface down to more than 100 m (see Table 2).



500 Figure 16: Sea-level indicators on Mangaia, Cook Islands. Selected profiles and dated samples are shown (from Woodroffe et al., 1991).

In 1985, Stoddart et al. presumed the occurrence of LIG reefs on Rarotonga, southern Cook Islands, at a maximum elevation of 3.5 m above MSL. They assumed that these reefs have not been tectonically uplifted.

Spencer et al. (1987, 1989) identified deposits of probable Late Pleistocene age at +12.2 m on Atiu, +10.0 m on
Mauke and +9.8 m on Mitiaro in the southern Cook Islands. Stoddart et al. (1990) found evidence of late
Pleistocene RSL up to +12.2 m on Atiu, +10.0 m on Mauke, and +9.8 m on Mitiaro. They also reported
unconformities within the Pleistocene at +1.75 to +2.65 m on Atiu, +1.5 to +2.75 m on Mauke and +4.1 to +6.0 m
on Mitiaro. Woodroffe et al. (1991) identified MIS 5e reefs on Atiu, Mauke and Mitiaro using the alpha-counting
technique. Three samples from Atiu have been dated (WALIS U-series IDs 1812-1817; AT8, AT9 and AT11; see

510 Fig. 17: 1) an *in situ Porites* on an erosional bench at an elevation of less than +2.7 m MLT (interpreted elevation +1.5 ± 1.5 m MSL) with an age of 132 ± 10 and 111 ± 7 ka (WALIS U-series IDs 1812 and 1813; AT8; WALIS RSL ID 557); 2) a *Porites* colony from the encrusting upper unit at an elevation of more than +2.7 m MLT (interpreted elevation +6.5 ± 3.5 m MSL) with an age of 135 ± 10 and 151 ± 12 ka (WALIS U-series IDs 1814 and 1815; AT9; WALIS RSL ID 558); and 3) a massive *in situ* coral on the cliff top at an elevation of +10 to +12

515 m MLT (interpreted elevation +11 ± 1 m MSL) with an age of 111 ± 6 and 113 ± 9 ka (WALIS U-series IDs 1816 and 1817; AT11; WALIS RSL ID 559).

hat gelöscht: 5



Figure 17: Sea-level indicators on Atiu, Cook Islands. Selected profiles and dated samples are shown (from Woodroffe et al., 1991).

Woodroffe et al. (1991) also dated four samples from Mauke (see Fig. 13): 1) a *Porites* colony from a perched reef block at an elevation of +7 to +8 m MLT (interpreted elevation +7.5 ± 1 m MSL) with an age of 136 ± 10 and 138
± 10 ka (WALIS U-series IDs 1818 and 1819; MK9; WALIS RSL ID 560); 2) an encrusting coral from the upper

- unit at an elevation of more than +2 m MLT (interpreted elevation +3 \pm 1 m MSL) with an age of 123 \pm 8 and 138 \pm 12 ka (WALIS U-series IDs 1820 and 1821; MK15; WALIS RSL ID 561); 3) a *Porites* colony from the encrusting upper unit at an elevation of +5 m MLT (interpreted elevation +5 \pm 1 m MSL) with an age of 136 \pm 11 and 145 \pm 10 ka (WALIS U-series IDs 1822 and 1823; MK16; WALIS RSL ID 561); and 4) a *Porites* boulder in
- 530 the upper unit at an elevation of +6 m MLT (interpreted elevation +6 ± 1 m MSL) with an age of 119 ± 8 and 119 ± 7 ka (WALIS U-series IDs 1824 and 1825; MK18; WALIS RSL ID 561). *Porites* is living at median and average depths of 0 and 18.5 m, respectively in modern environments; however, this genus displays a wide depth range (see Table 2).

hat gelöscht: 5

hat gelöscht: 6 hat formatiert: Schriftart: Nicht Kursiv



Figure 18: Sea-level indicators on Mauke, Cook Islands. Selected profiles and dated samples are shown (from Woodroffe et al., 1991).

Five samples from Mitiaro (see Fig. 19) have been dated by Woodroffe et al. (1991): 1) two specimens of Faviidae
(now called Merulinidae; Huang et al., 2014) from Te Unu at an elevation of more than +4 m (interpreted elevation +6.5 ± 2.5 m MSL) have been dated at 128 ± 6 ka, 140 ± 10 ka, 98.1 ± 5.4 ka, 112 ± 7 ka and 132 ± 9 ka (WALIS U-series IDs 1826-1830; MT1 and MT2; WALIS RSL ID 580); Merulinidae mainly indicate a depth range of 0-30 m; 2) one *Plesiastrea* specimen from the north coast at an elevation of more than +4 m (interpreted elevation +6.5 ± 2.5 m MSL) has been dated at 120 ± 6 ka and 132 ± 9 ka (WALIS U-series IDs 1832 and 1834; MT9;

- 545 WALIS RSL ID 581); in modern reef environments, *Plesiastrea* is mainly living at depths ranging from the surface down to 10 m; 3) one *Porites* specimen from the village at an elevation of more than +2 m (interpreted elevation +5.5 ± 3.5 m MSL) has yielded ages of 116 ± 8 ka, 144 ± 9 ka and 106.3 ± 5.4 ka (WALIS U-series IDs 1836-1838; MT13; WALIS RSL ID 582); and 4) one *Porites* specimen from Vaikoua at an elevation of more than +3 m (interpreted elevation +4 ± 1 m MSL revealed an age of 118 ± 7 (WALIS U-series IDs 1841 and 1842; MT15;
- 550 WALIS RSL ID 583). Porites is living at median and average depths of 0 and 18.5 m, respectively in modern environments; however, this genus displays a wide depth range (see Table 2). All samples contained less than 1-2% of calcite and all datings have been accepted (1-sigma uncertainty). Elevations have been hand-leveled and the tidal range is approximately 80 cm.

hat gelöscht: 6



Figure 19: Sea-level indicators on Mitiaro, Cook Islands. Selected profiles and dated samples are shown (from hat gelöscht: 7 Woodroffe et al., 1991).

Gray et al. (1992) extracted drill cores and described widespread LIG reefs on Pukapuka (see Fig. 3 in Gray et al.,
1992), however, they only obtained reliable ages on two *Porites* specimens (100% aragonite; WALIS RSL ID 586). Electron spin resonance (ESR) dating provided two ages of 144 ± 22 and 136 ± 20 ka for samples at depths of 21.77 and 24.85 m MSL, respectively (WALIS ESR IDs 104 and 105; 991 and 951). The *Porites* specimen at a depth of 24.85 m MSL was also dated using the alpha-counting technique, which revealed an age of 130 ± 9 ka (WALIS U-series ID 1855, 1-sigma uncertainty; 951). *Porites* is living at median and average depths of 0 and 18.5
565 m, respectively in modern environments.

hat gelöscht: 18

hat gelöscht: 18

Most of the dated samples from the southern Cook Islands by Woodroffe et al. (1991) can be related to the LIG RSL highstand, but a MIS 7 reef, i.e., the Penultimate Interglacial, has been also observed (see Fig. 20). Woodroffe et al. (1991) did not observe two periods of high RSL separated by a period of lower RSL during the LIG. However, 570 they describe several stratigraphic features, such as erosional benches and notches, that might indicate RSL

fluctuations during MIS 5e.



Figure 20; Uranium-series ages (± 1 standard error) from corals of the southern Cook Islands, and comparison with the late Quaternary sea-level curve of Chappell and Shackleton (1986) (from Woodroffe et al., 1991).

575

Overall, the large variability in elevations and the lack of accurate ages from the Cook Islands does not allow any precise conclusion about the LIG RSL.

580 5.3 Samoa

Exposed Pleistocene reef terraces have been described on several Samoan Islands, but no dates have been reported in the reviewed studies. Kear and Wood (1959) reported terraces occurring at up to 15 m and 10 m higher than present RSL on Upolu and Upolu's offshore islands, respectively and that might correspond to the LIG period (125 ka). They also described benches near Fagaloa Bay at elevations of +39 to +60 m, which might relate to the LIG or Penultimate Interglacial.

Stice and McCoy (1968) observed cliffs with benches at elevations of 4.5 to 1.5 m above MSL, probably indicating former higher RSL on three Manu'a Islands. The +3.7 to +3.0 m terrace is composed of sand and coral shingle or entirely of sand.

5.4 Niue

- 590 LIG reef deposits have been dated from outcrops by Kennedy et al. (2012). The earliest studies by Agassiz (1903), David (1904) and Schofield (1959) described different raised reef terraces on Niue. The widest terrace (Alofi Terrace; see Figs 21, and 22) is the most continuous one around the island (Agassiz, 1903). David (1904) recognized three terraces at +6 m, +24 to +27 m and +60 m and mentioned also a terrace at an elevation of 40 m above MSL. In 1959, Schofield distinguished the Alofi Terrace at about +23 m and the Mutalau Platform, a +69
- 595 m-high terrace, and identified five more terraces, including two submerged ones at -37 to -33 m and -15 to -11 m, and three exposed ones at +12 to +14 m, +35 to +40 m and at +55 m. At least parts of the Alofi Terrace date to 230-260 ka (Wheeler, 2000). The Alofi Terrace is overlain by a 6 m-thick LIG reef sequence (Paulay, 1988; Paulay and Spencer, 1992; Spencer and Paulay, 1994; Wheeler, 2000). Beach deposits and erosional features indicate a 6 m-high RSL (Paulay and Spencer, 1992). More recent studies describe the different terraces on Niue from close
- to present MSL up to an elevation of +60 m (Terry and Nunn, 2003; Nunn and Britton, 2004; Kennedy et al., 2012). Nunn and Britton (2004) indicated the presence of at least seven terraces, at +58, +52, +43.5, +34, +28, +23 and +18 m with ages back to 700 ka. Kennedy et al. (2012) dated the Alofi Terrace (see Fig. 22) based on the dating of one *Porites* specimen at 133 ± 0.9 ka (WALIS U-series ID 1285; NT11; WALIS RSL ID 3981) at an elevation of 3.1 m above the height of living corals. This coral head consisted of more than 95% of aragonite and
- 605 the dating is considered as reliable. *Porites* is living at median and average depths of 0 and 18.5 m, respectively in modern environments; however, this genus displays a wide depth range (see Table 2). Niue is affected by uplift with estimated rates of 0.13–0.16 mm/yr, with uplift considered to be ongoing (Dickinson, 2001), thus indicating an uplift of 16 to 20 m since 125 ka.

hat gelöscht: 19 hat gelöscht: 0




Figure 22: A schematic cross-section showing the stratigraphic relationships between the Pleistocene and modern terrace on Niue. Reef framework is not ubiquitous along the Alofi Terrace edge (from Kennedy et al., 2012).

hat gelöscht: 0

5.5 Tonga

LIG reef terraces in Tonga have been reported from outcrops at elevations of about +5 to +7 m. Taylor (1978) and Yonekura (1983) dated emerged reef terraces on the Tongatapu block using the alpha-counting technique. Taylor (1978) described a group of emerged notches and caves at 6.6-7.2 m above present MSL and dated a single sample
at 133 ± 12 ka (WALIS U-series ID 1787). Yonekura (1983) described LIG reefs at +5.5 m MHT and provided ages of 135 ± 15 ka and 133 ± 12 ka (WALIS U-series IDs 1785 and 1786; WALIS RSL ID 554). A wide palaeowater depth range of 0-30 m has been assumed as no taxa identification has been reported. The inaccurate dating of LIG reefs in Tonga does not allow any precise conclusion regarding the LIG RSL. Uplift rate is negligible in

630

620

5.6 Fiji

the Tonga arc (Yonekura, 1983).

The Fiji group of islands has a total land area of 18,270 km². Its geological history is complex owing to its proximity to the Australian–Pacific plate boundary. Several Fiji Islands (e.g., Kadavu, Kaibu) exhibit a series of coral reef terraces arranged around a volcanic core, the lowest one being attributed to the MIS 5e (Nunn and

Omura, 1999; Nunn et al., 2002). LIG deposits from Fiji were first dated by Taylor (1978) who reported ages of 126 ± 7 ka and 136 ± 12 ka (WALIS U-series IDs 1866 and 1867; Yasawas1 and Yasawas2) on two unidentified corals from a terrace at +2 to +3 m on Yasawa Island (WALIS RSL ID 512). However, the lack of any coral identification does not allow to precisely consider the sea-level position at that time. Nunn and Omura (1999) have

640 reported a similar age of 128.7 ± 1.6 ka from a coral sampled from the same reef terrace (WALIS U-series ID 1868; Yasawas3). In addition, Nunn and Omura (1999) reported on Kadavu Island morphological features that they have attributed to two RSL stands during the LIG without providing robust chronological constraints (see Fig. 4 in Nunn and Omura, 1999). In addition, an erosional bench reported at +5 m has been interpreted as an early LIG sea-level maximum at about 136 ka, while a distinctive level at +2 m would correspond to a second LIG

645 maximum at about 120 ka. Nunn and Omura (1999) have assumed that the LIG reefs are located below present MSL in the western part of Kadavu Island.

Nunn et al. (2002) and Nunn and Omura (2003) described LIG reef terraces from Kaibu Island (see Fig. 5 in Nunn et al., 2002; Figs 2 and 3 in Nunn and Omura, 2003), and reported three U-series ages. A *Porites* sampled at 1.45 m above the modern reef (WALIS RSL ID 511) yielded an age of 126.8 ± 2.1 ka (WALIS U-series ID 1865;

- 650 sample AO455). Two *Platygyra*, including *Platygyra lamellina* sampled between 3.6 and 3.85 m above the modern reef (WALIS RSL IDs 509 and 510) gave ages of 132.8 ± 2.3 ka (WALIS U-series ID 1864; AO454), and 131.1 ± 2.4 ka (WALIS U-series ID 1863; AO443), respectively. In modern environments, *Platygyra lamellina* typifies median and average depths of 4 and 4.4 m, respectively (Table 2; WALIS RSL IDs 509 and 510). Nunn et al. (2002) interpreted the age distribution as reflecting a double sea-level maximum with peaks around 133-130 ka
- and 123-120 ka, following the earlier interpretations by Nunn and Omura (1999). The earlier maximum has been reported about 2 m lower than the later and marked by the growth of a surface reef, the later maximum appearing to have involved only cutting of erosional shorelines at the +5.0 to +5.5 m level (Nunn and Omura, 2003). Nunn et al. (2002) and Nunn and Omura (2003) compared levels of emerged shorelines on northeastern Fiji Islands and show that a LIG shoreline has emerged in northeast Fiji by around +5.0 to +5.2 m (range +4.8 to +5.5 m). In
- 660 addition, they compared these levels to those from the western Fiji Islands and concluded that LIG RSL in the Fijian region ranged from 4.6 to 7.1 m above its present mean level.

5.7 Tuvalu

The only available chronological information concerning the potential occurrence of LIG on Tuvalu is based on the dating of a drill core carried out on Funafuti Atoll by the Coral Reef Expedition of The Royal Society (1896– 1898). It was obtained on an unidentified coral collected at 27.4 m (sample 274) below the reef surface and yielding a Sr isotope age of 0.14 ± 0.15 Ma (WALIS U-series ID 1734; Ohde et al., 2002). The carbonate sequence attributed to the LIG is reported from at least 24.4 to 27 m core depth and is limited by unconformities (Ohde et al., 2002).

5.8 Kiribati

- 670 LIG reefs have been reported from outcrops on Christmas Island and from a drill core collected on Tarawa. Marshall and Jacobson (1985) dated a drill core from an islet on Tarawa using the alpha-counting technique. The coral at a depth of 17-17.8 m below MSL revealed an age of 125 ± 9 ka (WALIS U-series ID 1811; 81390010; 1sigma uncertainty). The sample was composed of 100% aragonite and the dating is considered as reliable. Woodroffe and McLean (1998) dated an emerged reef terrace on Christmas Island and obtained a more accurate
- 675 age using a mass spectrometer. The coral sample at an elevation of +1 m MSL has been dated at 130 ± 5 ka (WALIS U-series ID 1810; 16a). A water depth range of 0-30 m has been assumed as no taxa identification has been reported. Christmas Island has remained stable throughout the Late Quaternary (Woodroffe and McLean, 1998).

5.9 Solomon Islands

- 680 The geomorphology of the Solomon Islands has been studied by Stearns (1945), Grover (1958) and Stoddart (1969) describing elevated terraces in different island groups. Grover (1958) described notches on Mone Island at an elevation of +0.76, +3 and +5.5 m on an 8.5 m-high cliff and at +22 m on a 16.8 to 29.6 m-high cliff. Stoddart (1969) mentioned the occurrence of Pleistocene reef limestones and reported terraces of clastic debris with reef material at elevations of 145 to 135, 80, 60, 44 and 27.5 m above present MSL. He also reported tidal notches on
- 685 Banika Island with an elevation of 1.5 m to 5 m above present MSL. A terrace on Banika island at 9 m above present MSL was older than 39,700 years. However, it was impossible to relate the relevant benches to specific Pleistocene RSL changes. The Solomon Islands are affected by uplift. There are 16 arc segments in the Solomon Islands comprising three major tectonic regimes (Chen et al., 2011).

690

5.10 Vanuatu

Uplifted Quaternary reef terraces cover wide areas of Vanuatu islands, specifically Santo, Malakula and Efate at elevations ranging from 600 m on Efate and the North of Malakula, to 860 m in the South of Malakula and up to

- 695 more than 1000 m in the western part of Santo (see Fig. IV-2 in Jouannic et al., 1982). They form the external part of a succession of terraces overhanging a Holocene coastal reef platform. Since the first geological and morphological investigations in the late 1960s and the 1970s (Mitchell, 1966, 1969, 1971; Robinson, 1969; Mallick and Greenbaum, 1977; Ash et al., 1978), these uplifted terraces have been the topic of studies dealing both with their chronology and the reconstruction of vertical movements in the New Hebrides forearc.
- 700 The first chronological study has been carried out by Neef and Veeh (1977) who published the first four U-series ages obtained on corals from MIS 5 raised reefs from Efate and Malo, which occur at +100 m and +96 m, respectively. These raised reefs have been interpreted as coeval to RSL stands VIIa (140,000 yr) and VIIb (120,000 yr) from Huon Peninsula (Chappell, 1974; Bloom et al., 1974), respectively. The two ages obtained by Neef and Veeh (1977) were of 134 ± 10 ka (WALIS U-series ID 1847; Efate 330) and 134 ± 6 ka (WALIS U-series ID; Ma
- 1) were judged to be more reliable than those which yielded ages of about 120 ka (Efate X and Efate 235) on the basis of mineralogical criteria used in the evaluation of samples (Neef and Veeh, 1977). However, corals that have been dated were not identified, thus hampering the accurate reconstruction of palaeo-water depth during reef growth. Subsequent studies by Bloom et al. (1978), Jouannic et al. (1980, 1982) and Lecolle and Bernat (1985) have been focused on the uplifted terraces from Efate. Two alpha spectrometry ages of 141 ± 8 ka (WALIS U-
- 710 series IDs 1952 and 1953; E-L-3 and E-T-2) were obtained by Bloom et al. (1978) on unidentified corals from the coral terrace located at +110 to +130 m in Port Havannah area (WALIS RSL IDs 491 and 492). Additional U/Th ages of 124 ± 7 ka (WALIS U-series ID 1950; sample I-7-5), 131 ± 11 ka (WALIS U-series ID 1951; E-L-1), 121.6 ± 7.3 ka (WALIS U-series ID 1942; EKB4-2) and 125.3 ± 7.5 ka (WALIS U-series ID 1941; EKB4) were obtained by Jouannic et al. (1982) and Lecolle and Bernat (1985) on the same terrace (WALIS RSL IDs 491 and
- 715 492). This terrace has been therefore attributed to MIS 5e and its duplication in the area of Port Havannah, similar to the one observed by Bloom et al. (1974) on Huon Peninsula (VIIa and VIIb terraces; Bloom et al., 1974), has been interpreted as resulting from low amplitude sea-level fluctuations during MIS 5e (Jouannic et al., 1982). Two ages of 130 ± 7 ka (WALIS U-series ID 1948; sample E-X-2) and 114 ± 6 ka (WALIS U-series ID 1949; sample E-X-4) have been obtained on the coral reef terrace at +85 to +95 m in the same area (Jouannic et al., 1982). The
- 720 MIS 5e terrace has been dated by Lecolle and Bernat (1985) at different altitudes in several other areas on Efate Island (see Fig. 2 in Lecolle and Bernat, 1985). An age of 121.4 ± 7.3 ka (WALIS U-series ID 1937; sample ESA3)

was reported at +50 m in Siviri area (WALIS RSL IDs 706 through 708), while ages of 128.0 ± 7.7 ka (WALIS U-series ID 1938; ERB1) and 133.5 ± 8 ka (WALIS U-series ID 1939; sample EWA03) were obtained on the terrace occurring at +80 m in Malafao area (WALIS RSL ID 709), and ages of 117.8 ± 7.0 ka (WALIS U-series

- 725 ID 1945; ETD1) and 123.6 ± 7.4 ka (WALIS U-series ID 1946; ETA4) on the terraces at +80 m and +100 m, respectively, in Tukutuk area (WALIS RSL ID 713). However, the palaeo-MSL cannot be further constrained due to the lack of coral identification. Chronological data obtained on Efate Island, combined with U-series ages obtained on Malakula and Santo islands, have been used to reconstruct vertical tectonic movements in the New Hebrides Island Arc (Jouannic et al., 1982; Lecolle and Bernat, 1985). The chronological frame of the MIS 5e
- 730 terrace from Efate has been reassessed by Edwards et al. (1987a, b) who identified and dated the two corals previously reported by Bloom et al. (1978) at 129.9 ± 1.1 ka and 129.2 ± 1.1 ka for *Oulophyllia crispa* (WALIS U-series IDs 1435 and 1436; E-T-2-A and E-T-2) and 125.5 ± 1.3 ka for *Porites lutea* (WALIS U-series ID 1854; E-L-3). The association of those two coral species indicates probable palaeo-water depths of less than 10 m (WALIS RSL ID 492).
- 735 The reef terrace corresponding to MIS 5e (from 119 to 132 ka) was not clearly identified on Malakula Island (Jouannic et al., 1982; Cabioch and Ayliffe, 2001; see Fig. 4 in Taylor et al., 1980). Jouannic et al. (1982) also suggested that some of the discrepancies in terrace altitudes in previous work may have been due to problems associated with coral reworking and errors in altitude assignment. An age of 119.8 ± 8.0 ka (WALIS U-series ID 1947; NH14) has been obtained on the coral reef terrace located at +180 m in Npénanavet/Tenmaru area by
- 740 Jouannic et al. (1982). A similar age of 121.3 ± 1.2 ka (WALIS U-series ID 1846; Espiegle1-188m) has been obtained at +188 m by Cabioch and Ayliffe (2001) on an unidentified coral (WALIS RSL ID 490; see Fig. 23). However, Cabioch and Ayliffe (2001) have considered that the MIS 5e terrace should correspond to the wide plateau culminating at +230 m and that, based on field evidence, its composite reef structure may result from minor sea-level variations during the MIS 5e RSL stand.



Figure 23: Profile showing the distribution of reef terraces on Malakula. Transect 1: Npénanavet/Tenmaru transect; transect 2: Espiègle Bay/Malua Bay transect (from Cabioch and Ayliffe, 2001).

hat gelöscht: 1

Taylor et al. (1985) have investigated raised coral reef terraces from the Torres Islands to reconstruct rapid vertical movements in the New Hebrides forearc. They interpreted the very wide terrace at about +100 m on Toga (WALIS RSL ID 488; see Fig. 24) as formed during the MIS 5e and deduced uplift rates of about 1 mm/yr for at least the past 125,000 years, i.e., 125 m since 125 ka. Ages assigned to MIS 5e have been reported from the reef terrace located at +80 m on Loh Island (WALIS RSL ID 487; see Fig. 25) where a *Porites* and a *Leptoria phrygia* yielded
ages of 122 ± 7 ka (WALIS U-series ID 1839; LO-B-2) and 135 ± 11 ka (WALIS U-series ID 1840; LO-B-3), respectively. In modern reefs, *Leptoria phrygia* typifies shallow-water environments with median and average depths of 4 and 5 m, respectively; however, this species displays a rather large depth range, from the surface down to 30 m or even 47 m (Table 2).

hat gelöscht: 2

hat gelöscht: 3

43



On Tanna Island, raised lagoonal deposits currently at +18 m (WALIS RSL ID 506; see Fig. 2() have been attributed to MIS 5e based on an age of 126.7 ± 1.5 ka (WALIS U-series ID 1861; Ta9) obtained on an *Acropora*,



which does not give precise indications on the palaeo-water depth during reef growth.



hat gelöscht: 4

5.11 New Caledonia

New Caledonia comprises the main island (Grande Terre), the Isle of Pines to the south, the Loyalty Islands to the east (Maré, Lifou, Tiga and Ouvéa), and the Belep Archipelago to the northwest.

LIG reef deposits have been reported both from outcrops in Loyalty Islands (Maré, Lifou and Ouvéa) and from

- 780 drill cores that have been carried out throughout the modern barrier and fringing reefs from the western coast of Grande Terre, as well as in the Chesterfield islands that are located about 500 km to the WNW of New Caledonia. Pioneering studies on Pleistocene reef terraces from the Loyalty Islands, including the LIG, have been carried out in the 1970s by Launay and Récy (1972) and Coudray (1976). The first radiometric ages on these raised reefs have been obtained by Bernat et al. (1976), Marshall and Launay (1978) and Gaven and Bourrouilh-Le Jan (1981), and 785 were used to reconstruct tectonic movements of these islands since Quaternary times.
- 230 Th/ 238 U ages reported by Bernat et al. (1976) on raised reefs occurring at +11 to +12 m elevation on Lifou and Maré range from 122 to 120 ka and would therefore indicate that this terrace corresponds to the MIS 5e. A coral from an elevation of +20 m on the Isle of Pines, which is located immediately to the southeast of New Caledonia, was dated at 126 ± 5 ka. However, this suite of corals did not satisfy the closed-system criteria because they were
- 790 partly recrystallized (Marshall and Launay, 1978). Marshall and Launay (1978) could not obtain reliable ages on a terrace located at +3.5 m on Lifou, but they have reported an age of 117 ± 6 ka on a *Platygyra* colony (WALIS U-series ID 1791; Ouvea3) sampled in the coral reef terrace occurring at +7.5 m on Ouvéa (WALIS RSL ID 737) and correlated this terrace with the 120 ka Waimanalo shoreline from Hawaii (Ku et al., 1974). The distribution of modern *Platygyra* is characterized by median and average depths of 4 and 7 m, respectively; however, their
- 795 depth range is very large and extends from the surface down to 80 m (Table 2). Marshall and Launay (1978) have concluded that there is evidence of LIG reef terraces at +11 to +12 m on Lifou, at +20 ± 3 m on the Isle of Pines (Bernat et al., 1976), and at +7.5 m on Ouvea but, because of diagenetic alteration of the Lifou and Isle of Pines coral samples, the exact age of these terraces could not be defined. Gaven and Bourrouilh-Le Jan (1981) have reported ages ranging from 100 to 295 ka on pristine corals sampled in terraces located between 0 and +9 m on
- 800 Maré. They have noted that some of these terraces have been karstified and attributed notches at +9 and +12 m to 125 and 205 ka RSL.

The study of drill cores from modern reefs from several areas of the west coast of New Caledonia (Amédée, Kendec and Ténia islets) have provided the opportunity to document reef units attributed to MIS 5e that occur at various depths, thus demonstrating differential tectonic movements along the west coast of New Caledonia (see

805 Fig. 1 in Cabioch et al., 2008).

Below fringing and barrier reefs, the 125 ka-old reef unit is generally capped by calcretes resulting from a relatively long period of emergence and is locally affected by freshwater alteration, even if the barrier reef is frequently unaltered. Several alpha counting U/Th dates have been obtained on these drill cores: 125 ± 20 ka in the Ténia

- 810 drill cores (Coudray, 1976), 135 ± 12.8 ka in the Ilot Vert drill cores (Degaugue-Michalski, 1993) and 125 ± 1 ka and 131 ± 1 ka at Ilot Amédée (in cores Amédée-1 and -2, respectively; Cabioch et al., 1996), where a framework of corals and coralline algae with a few bioclastic beds, typifies reef flat facies. In Amédée-4 drill core, the 125 ka-old reef occurs between 14 to 37 m core depth (WALIS RSL ID 474), and is predominantly composed of biocalcarenites and rare coral colonies (Cabioch et al., 1999). U/Th dating by TIMS carried out on coral samples
- 815 extracted from Amédée site has demonstrated that the distribution of U and Th isotopes was affected by subaerial diagenesis during emergence of the reef units, while the Holocene coral sequence seems largely unaffected (Frank et al., 2006; Cabioch et al., 2008). Various correction models were tested to get reliable ages from the late Pleistocene sequences and yielded ages ranging from 102.6 ± 4.4 to 127.5 ± 5.4 ka, with a majority of values around 122 ka (Frank et al. 2006; Cabioch et al., 2008). The reef unit occurring between 37 and 14 m core depth
- 820 has been therefore attributed to MIS 5, especially to MIS 5e (WALIS U-series IDs 1266 through 1283; WALIS RSL ID 474; see Fig. 27); based on these results, the local subsidence rate has been estimated to 0.16 mm /yr (Frank et al., 2006), thus a total of 20 m since 125 ka.



825 Figure 27; Synthetic log of the four cores drilled at Amédée Islet. D1 is the Pleistocene/Holocene boundary unconformity; D2 is the bottom of the LIG terrace (from Frank et al., 2006).

Similar Pleistocene reef units have been identified in drill cores retrieved from the Kendec (Cabioch et al., 2008)
 and Vert (Cabioch, 1988; Degaugue-Michalski, 1993) islets. Coral identifications (Hongo and Wirrmann, 2015)
 have shown that the MIS 5 reef units are dominated by important accumulations of shallow-water assemblages including arborescent colonies of *Acropora* (Kendec drill core), corymbose and tabular colonies of *Acropora* (Kendec and Amédée-4 drill cores) and massive *Porites* colonies (Amédée-5 drill core) that characterize reef edge and upper reef slope environments exposed to strong wave action.

- Below the barrier reef that occurs at Ténia islet, the 125 ka-old unit consists of a framework of corals, bryozoans and coralline algae trapped in bioclastic sediments rich in bryozoans, typifying reef flat facies (Coudray, 1976). The reef unit occurring between 14 and 16 m core depth in drill core extracted more recently in this area (Ténia-3; Montaggioni et al., 2011) has yielded U-series ages of 127.9 ± 3.3 ka (WALIS U-series ID 1788; 14T3), 130.55 ± 3.75 ka (WALIS U-series ID 1789; 16T3) and 132.65 ± 4.7 ka (WALIS U-series ID 1790; 18T3), according to
- 840 the U/Th dating procedure previously described by Frank et al. (2006).

5.12 Papua New Guinea

Quaternary coral reef terraces from Huon Peninsula (Papua New Guinea) are prominent morphological features that extend more than 80 km along the coast and have been uplifted at a rate that varies from 0.5 mm/yr in the northwest to nearly 4 mm/yr to the southeast (Esat et al., 1999). They range in elevations from 0 to 200 m (see

- 845 Figs 2% and 29 and consist of an upward succession of fringing and barrier reef types. Seven coral reef terraces, I through VII from the coast landwards, are spaced at about 20-kyr interval (see Figs 2% and 29) and provide a rather complete record of RSL, oxygen isotopes and temperature of the surface tropical ocean over the duration of the high RSL events of the last glacial cycle (Aharon and Chappell, 1986). These uplifted coral reef terraces have been recognized and extensively studied since the pioneer works of Veeh and Chappell (1970). Huon Peninsula now
- 850 represents a classical site of RSL change studies in support of the astronomical theory of climate change (Chappell, 1973).

Reef tract VII is a large structure with a barrier reef, a lagoon, and an inner fringing reef occurring at the crest of the relief (see Figs <u>30</u> and <u>31</u>). The main body of reef tract VII overlies the upper Tertiary limestone of the north Huon Peninsula (the Cromwell Limestone). Terraces VIb, VIa (above VIb), and V (below VIb) are platforms of

855 younger raised fringing reefs that offlap the front of reef tract VII (see Figs 28 to 31). Uplift rates for the Huon Peninsula have been estimated from terrace VIIb, which has been dated at 118 to 120 ka with an assumed palaeo-RSL at 0 to +5 m (Chappell, 1974; Bloom et al., 1974; Aharon and Chappell, 1986; Chappell and Polach, 1991; Chappell and Shackleton, 1986).

hat gelöscht: 6	
hat gelöscht: 7	
hat gelöscht: 6	
hat gelöscht: 7	

hat gelöscht: 28	
hat gelöscht: 29	
hat gelöscht: 6	
hat gelöscht: 29	



Figure 28: Location and generalized physiography of the emerged reef terraces on Huon Peninsula (from Stein et al., 1993). Spot elevations in meters. Reef numbering (I to VIII) follows Chappell (1974). Inset map shows position of study

hat gelöscht: 6

870 area in Papua New Guinea.



The first radiometric ages concerning Huon Peninsula reef terraces were obtained by Veeh and Chappell (1970) on coral colonies and molluses (*Tridacna gigas*). Three U-series ages (alpha spectrometry), which could be assigned to MIS 5 were obtained on unidentified corals from what they called the 'Reef Complex V', reported as

'Reef Complex VII' by Bloom et al. (1974) and interpreted as being built during the 125-120 ka time window. The oldest ages of 140 ± 10 ka and 133 ± 10 ka were obtained on a sample collected from fringing-reef terrace VIIa, on the landward side of the lagoon floor of reef complex VII on the Sialum profile. The youngest ages reported by Veeh and Chappell (1970) were of 116 ± 7 ka and 119 ± 7 ka and concerned a sample collected from

880

885 near the crest of the barrier reef at Sialum (near the Kwambu profile) referred to as VIIb crest by Bloom et al. (1974). The large errors in all these analyses did not allow a definite conclusion about the precise chronology of Terrace VII.

Bloom et al. (1974) reported a single datable coral sample from the undercut base of a large cliff at the front of the barrier of reef complex VII, about 20 m below the ridge crest, on the Nama traverse. The reported age is of $142 \pm$

- 890 8 ka and concerns a *Porites lutea* colony, which most commonly occurs at water depths of less than 10 m, with median and average depths of 0 and 2.7 m respectively in modern environments (Table 2). However, Bloom et al. (1974) had insufficient samples to document the age of terrace VII and selected an arbitrary age of 125 ka for this terrace. Based on the chronology provided by Bloom et al. (1974), Chappell (1974) has made a parallel between the described the raised coral terraces and coastal terraces from other parts of the world, especially Barbados where
- 895transgression maxima were previously established around 6.4, 29, 35, 60, 74, 118, 137, 185, 218 and >250,000 ka.The combined use of previous chronology and oxygen isotope records from *Tridacna gigas* occurring in uplifted
reef terraces allowed Aharon (1983) and Aharon and Chappell (1986) to quantitatively evaluate ice volume and
temperature factors. They established that sea level rose rapidly by 5 to 6.5 m above present MSL during the LIG,
which on stratigraphic grounds and δ^{18} O compositions, can be separated into an early phase whose δ^{18} O is similar
- 900 to modern values (reef VIIa dated at 138 ka) and a late phase that is 0.6% heavier (reef VIIb dated at 118 ka). A comprehensive chronological study of the LIG coral reef terraces from Huon Peninsula has been carried out by Stein et al. (1993). Valid U-series ages reported from the Kwambu section (Sialum area) at elevations ranging from +197 to +229 m fall into two tight groups centered at 118 ka and 134 ka (WALIS RSL IDs 464 through 470) and replicate with considerably more precision the few previous alpha-counting results. This confirms the

905 chronological frame of reef complex VII (see Fig. <u>30</u>), with ages around 118 ka to the VIIb barrier and 130-140 ka to the VIIa fringing reef, respectively (Chappell, 1974).





The group centered at 118 ka includes ages ranging from 116.4 ± 1.8 ka (WALIS U-series ID 1325; KIL-5b) to 119.5 ± 1.2 ka (WALIS U-series ID 1327; KIL-5a-1) and concern colonies of *Porites lutea* that likely indicate a

- 915 depositional environment shallower than 10 m, and colonies of *Platygyra lamellina* and *Gardineroseris planulata*, which also likely indicate a reef setting shallower than 10 m (WALIS RSL IDs 466 through 469). The group centered at 134 ka includes ages ranging from 131.9 ± 1.2 ka (WALIS U-series ID 1322; HP-23b) to 136.7 ± 1.2 (WALIS U-series ID 1313) ka and mostly concern colonies of *Gardineroseris planulata* and *Gardineroseris* sp., as well as *Cyphastrea serailia* typifying superficial environments shallower than 6 m (see Table 2; WALIS RSL
- 920 IDs 464, 465 and 470). These results show that the barrier VIIb includes corals of the two age groups, and that corals of 118 ka appear in the outer margin of VIIa, 3 m below the crest of VIIb and on the seaward surface of the barrier (30 m below the crest). This configuration suggests that the younger corals (118 ka) were unconformably deposited on the relief of the VIIb barrier. The main structure of the coral reef terrace VII is therefore believed to have developed during a major and continuous sea-level rise that commenced well before 134 ka and continued
- 925 until 118 ka (Stein et al., 1993). Results obtained by Stein et al. (1993) have been reassessed by Esat et al. (1999) who noted that these two age clusters are separated vertically by 7 m, thus indicating that the RSL at 136 ka could have been 43 m lower than at 118 ka, given the 1.9 mm/yr uplift rate (Stein et al., 1993).

Esat et al. (1999) have investigated a large ancient sea cave, Aladdin's Cave, at the base of terrace VIb (MIS 5b-5c) in the Kwangam section (see Fig. 31). Most of the acceptable ages of corals from Aladdin's Cave are within

- 930 the range of 126 to 134 ka (WALIS U-series IDs 1296 through 1299 and WALIS U-series IDs 1302 through 1312), with the exception of a coral sample dated at 115±0.9 ka (WALIS U-series ID 1301; AC-U12). The corals exposed within Aladdin's Cave include abundant and large heads of *Porites* (up to 1 m in diameter) and various Faviidae (now called Merulinidae; Huang et al., 2014), indicating water depth shallower than 20 m during their growth (WALIS RSL IDs 459, 462 and 463). Ages obtained independently by TIMS and alpha spectrometry on a large
- 935 Porites colony at the cave entrance are in excellent agreement (WALIS U-series IDs 1307 through 1309; AC-U11a-top, AC-U11a-bottom, AC-U11-bottom) and average 127.6 ± 0.6 ka. The mean age of corals from the crest of reef VII is 134 ± 1 ka, and the mean age of the Aladdin's Cave corals of this time interval is 130 ± 2 ka. The corals at the cave appear to have grown;~4000 years later than those that are found 10 m below VIIb or ~80 m above the cave and appear to have grown in conditions 6°C cooler than present. The conflict between samples
- 940 from Aladdin's Cave (about +100 m at about 130 ka) and from the coral reef terrace VII (up to +220 m at about 134 ka) has been explained by a sea-level drop of 60 to 80 m after the development of the reef terrace VIIb (Esat et al., 1999; see Fig. 32). A new rise in sea level after 130 ka is deduced from the dataset and occurred in response

hat gelöscht: 29

945 to the major insolation maximum at 126 to 128 ka (Esat et al., 1999). The 118-ka corals of reefs VIIa and VIIb may have also been deposited during an episode of sea-level rise at the end of the LIG (Esat et al., 1999).



950

Figure <u>31</u>; A profile of coral terraces V to VII, including Aladdin's Cave at the base of reef VIb at west Kwangam and the equivalent reefs at Kwambu. The horizontal scale for both sections is 1:15,000, which was calculated from air photographs; the vertical scale for Kwambu was calculated from a theodolite survey, and the vertical scale for Kwangam was calculated from calibrated digital altimeter (from Esat et al., 1999).



hat gelöscht: 0

Figure 32: (A) The age-height relation for dated terraces at Kwambu that was constructed by assuming that the top of terrace VIIa at 118 to 120 ka was 0 to 5 m above the present sea level. The dashed vertical lines indicate the possible range of water depth (0 to 20 m) in which corals grew. (B) Eustatic sea-level curve for the penultimate deglacial period and the Last Interglacial. The smooth sea-level curve glosses over co-seismic uplift events at Huon Peninsula, which are
 discrete and meter scale and would not be discernible at this scale. The diagonal dashed lines represent the uplift with

time for the crest of reef VII and Aladdin's Cave (from Esat et al., 1999).

Cutler et al. (2003) have dated coral samples from reef terraces VI (Sialum area) and VIIb (Kilasairo area) by applying both ²³⁰Th and ²³¹Pa dating techniques as a test of age accuracy. The two *Porites* samples from the coral reef terrace VI occurring at elevations of +169 to +177 m gave ages of 119.34 ± 0.76 ka and 121.87 ± 0.78 ka (WALIS U-series ID 1290 and 1291, respectively). The identified corals indicate a rather large palaeo-water depth range, from 0 to 20 m (WALIS RSL IDs 457 and 458). Four samples of *Gardineroseris planulata* from coral reef terrace VII bat +195 m yielded ages ranging from 117.77 ± 0.69 to 113.9 ± 0.65 ka (WALIS U-series ID 1289 and 1742, respectively; KIL3a and KIL3b). This result apparently confirms the U-series age of 116.6 ± 1.15 ka

previously obtained by Stein et al. (1993; WALIS U-series ID 1292), but discordant Pa-Th/U-series ages are reported (Cutler et. al., 2003). The reported corals indicate a shallow-water depositional environment, shallower than 10 m (WALIS RSL IDs 455 and 456), with median and average depths of 3 and 5.9 m respectively in modern environments. Based on their database, Cutler et al. (2003) concluded that MIS 5e ended prior to 113.1 ± 0.7 kyr,

980 when RSL was at -19 m. In addition, uplift rates of 1.85 mm/yr for Sialum, Kwambu, and Kilasairo, 2.6 mm/yr for Kanomi, and 2.8 mm/yr for Kanzarua, were calculated based on the dating results obtained on Terrace VIIb (Cutler et al., 2003).

Recently, Ayling et al. (2017) have investigated the chemical and isotopic distribution of uranium in *Tridacna* gigas from Marine Isotope Stage (MIS) 5e (128–116 ka) and MIS 11 (424–374 ka) reefs and demonstrated that

985 they cannot provide independent, reliable geochronological controls on the timing of past reef growth at Huon Peninsula.

5.13 Federated States of Micronesia

Pleistocene limestones, including the LIG period, have been described in this region, however, no dates have been reported in the reviewed studies. The hydrogeologic study on Pingelap and Pohnpei by Ayers and Vacher (1986)
indicated that the Pleistocene aquifer is located at a depth of 20-30 m. Anthony (1996a, b, c) and Anthony and Spengler (1996) provided several hydrogeologic reports on islands in Pohnpei and indicated the presence of Pleistocene deposits with a Holocene-Pleistocene contact at depths of 15-25 m below present MSL. Fletcher and Richmond (2010) summarized in a report that the Pleistocene limestone beneath the islands in this region at depths from about 8 to 28 m below the modern reef platform was deposited during the LIG period.

995 5.14 Mariana Islands

Early studies in the 1940s to 1960s, including Stearns (1945), Tayama (1952), Cloud (1954), Cloud et al. (1956), Tracey et al. (1959, 1964) and Emery (1962, 1963), described Pleistocene reef terraces in the Mariana Islands. Benches have been described at different elevations on Saipan and Guam: at +2, +5 to +10 and +20 m (Tayama, 1952), +1.22 to +2.44 m (Emery, 1962) and +1.52, +7.62, +13.72, +21.34 and +30.48 m (Stearns, 1945). The Last

1000 Interglacial Tanapag Limestone is usually found at elevations of less than 30 m (Weary and Burton, 2011). Muhs et al. (2020) carried out U/Th dating by high-precision thermal ionization mass spectrometry (TIMS) on coral samples from the Tanapag Limestone along the eastern and southern coasts of Saipan at elevations of about 13 to 30 m. Twelve samples of *Porites, Goniastrea* and *Acropora* have been collected from four sites; of those nine in growth position (see Fig. 12B; Fig. 4 in Muhs et al., 2020). Ten samples of these corals at elevations of +9 to +14

1005 m MSL have been dated at 125.7 ± 1.2 to 136.4 ± 1.1 ka (WALIS U-series IDs 3140-3146, 3149-3152; WALIS RSL IDs 3963, 3964, 3966), and two *Goniastrea* (growth position uncertain) at an elevation of +23 ± 3 m MSL have been dated at 126.9 ± 0.9 and 128.7 ± 1 ka (WALIS U-series IDs 3147, 3148; WALIS RSL ID 3965). All samples, except one, contain less than 3% of calcite and the ages can be considered as reliable and have been accepted. *Porites, Goniastrea* and *Acropora* are living in modern environments at median and average depths of 0 and 18.5 m, 6 and 15.3 m and 4 and 5.7 m, respectively.

The Mariana Islands are affected by uplift, but the rates are negligible in the Mariana arc (Yonekura, 1983). Miklavič et al. (2012) concluded an average uplift rate of about 0.1 to 0.2 mm/yr and Muhs et al. (2020) reported low late Quaternary uplift rates of 0.002 to 0.19 mm/yr

5.15 Marshall Islands

- 1015 Data regarding the LIG from the Marshall Islands have been obtained from drill cores. Szabo et al. (1985) presented alpha-counting U-series datings from two drill cores on the reef flat of Enewetak, which have been collected during the PACE program. Merulinidae in a coral-algal boundstone in core C-3 at a depth of 9.8 m revealed ages of 131 ± 7 and 132 ± 7 ka (WALIS U-series IDs 1780 and 1781; C-3A and C-3B). Three Merulinidae samples from core C-4, a coral-foraminiferal-skeletal grainstone containing numerous corals and abundant
- Halimeda, at a depth of 14.9 m have been dated at 128 ± 7 ka, 136 ± 8 ka and 129 ± 8 ka (WALIS U-series IDs 1782-1784; C-4A, C-4B and C-4C). The dated samples contain less than 3% of calcite and the ages can be considered as reliable. The elevation measurement method has not been reported, but the cores have been probably hand-leveled and the elevations refer to MSL. Modern Merulinidae mainly live at water depths shallower than 30 m (WALIS RSL IDs 555 and 556). The subsidence rate of Enewetak has been estimated at 0.038-0.052 mm/yr
 1025 (Deplete and MEdward 1000).
- 1025 (Paulay and McEdward, 1990).

5.16 Hawaii

Emerged marine carbonate deposits are observed in several Hawaiian Islands and palaeo-shorelines may have been affected by tsunamis and storms. The LIG Waimanalo Limestone fringes most of Oahu and consists of corals in growth position, which are often overlain by marine coral-basalt conglomerates. Stearns (1935a, b, c) has been the

1030 first who studied Pleistocene shorelines on the islands of Oahu and Maui describing benches at elevations of +6 to +10.7 m and have been related to the Late Pleistocene (Stearns, 1941). Beach deposits on Ulupau Head, Oahu, occur up to 10.7 to 12.2 m above present sea level (Stearns, 1935c; Wentworth and Hoffmeister, 1939).

Several studies have provided ages for the deposits of the Waimanalo Limestone on Oahu and Rubin et al. (2000) 1035 provided ages for Hulopoe gravel coral clasts on Lana'i.

LIG reef terraces on Oahu reach an elevation of up to 12.5 m MSL and Veeh (1966) and Ku et al. (1974) have been the first who dated these emerged terraces. Early studies by Veeh (1966), Ku et al. (1974) and Sherman et al. (1993) used the alpha-counting technique and later studies by Muhs and Szabo (1994), Szabo et al. (1994) and McMurtry et al. (2010) used mass spectrometry. In 2002, Muhs et al. performed additional datings with a higher

1040 precision at the Szabo et al. (1994) study sites using high-precision thermal ionization mass spectrometry (TIMS). Elevations have been mostly hand-leveled and sometimes the elevation measurement method has not been reported, but the samples have been probably hand-leveled. Uplift of Oahu was first quantified by Ku et al. (1974). Waimanalo LIG deposits have been dated from several sites around Oahu (see Fig. 13; Fig. 1 in McMurtry et al., 2010 and Fig. 1 in Szabo et al., 1994). Based on the depth distribution of their modern counterparts, the coral taxa

1045 described below indicate a median and average depth of 0 to 15.6 m; however, *Pocillopora* displays a wide depth range (see Table 2). Cross-sections for some of these sites are displayed on Figure 33. Information on each site is provided below.

```
    Kahuku Point (WALIS RSL ID 542): Eight Porites and Pocillopora, of those five in situ, at an elevation of +1
    ± 1 m MSL have been dated at 113 ± 0.7 to 137 ± 11 (alpha counting, 1-sigma uncertainty) and more accurately at 122.1 ± 1.3 ka (2-sigma uncertainty; WALIS U-series IDs 1221-1224, 1170, 1171, 1771, 1778; Ku et al., 1974; Szabo et al., 1994; Muhs et al., 2002). All ages have been accepted.
```

2) Mokapu Point:

1055 A) WALIS RSL ID 543; Porites and Pocillopora, of those six in situ, at an elevation of +9.5 ± 2.5 m MSL have been dated at 123.2 ± 2.6 to 134 ± 4 ka (n = 15 accepted out of 20 ages; WALIS U-series IDs 1193, 1194, 1226-1239, 1761-1762, 1800-1801; Ku et al., 1974; Szabo et al., 1994; Muhs and Szabo, 1994; Muhs et al., 2002). Muhs and Szabo (1994) dated reworked Pocillopora from coral-bearing beach or sublittoral sands (WALIS U-series ID 1800) and marine conglomerates with coral cobbles and heads (WALIS U-series ID 1801).

1060 B) WALIS RSL IDs 3979 and 3980 (Hearty et al., 2007): One *Porites lobata* from the lower reef at an elevation of +4.9 ± 0.5 m MSL has been dated at 131.24 ± 0.41 ka (WALIS U-series ID 1169; not reported if *in situ*). Two *in situ Pocillopora meandrina* from the upper reef at an elevation of +8.5 ± 1.0 m MSL have been dated at 130.9 ± 0.49 ka (WALIS U-series IDs 1167, 1168). *Porites lobata* and *Pocillopora meandrina* live at median and average depths of 0 and 19.2 m, and 0 and 15.8 m, respectively. All ages have not been accepted.

hat gelöscht: 2

hat gelöscht: 1

hat gelöscht: (

1070 3) Alala Point (WALIS RSL ID 539): Porites and one Pocillopora, not in situ or not reported if in situ, at an elevation of +2.4 ± 2 m MSL have been dated at 119.4 ± 3 to 134.5 ± 1.4 ka (n = 6 accepted out of 8 ages; WALIS U-series IDs 1172-1177, 1207, 1208; Szabo et al., 1994; Muhs et al., 2002).

4) Maunalua Bay (WALIS RSL ID 546): *Porites* and *Pocillopora*, not reported if *in situ*, at an elevation of +7.1 ±
1075 1 m MSL have been dated at 118 ± 3 to 121 ± 7 ka (n = 4; 1-sigma uncertainty; WALIS U-series IDs 1756-1759; Ku et al., 1974). All ages have been accepted.

5) Black Point (WALIS RSL ID 545): *Porites*, not reported if *in situ*, at an elevation of +7 ± 1 m MSL have been dated at 112 ± 6 to 128 ± 8 ka (n = 8; 1-sigma uncertainty; WALIS U-series IDs 1748-1755; Ku et al., 1974). All ages have been accepted. *Porites* is living at median and average depths of 0 and 18.5 m, respectively in modern environments.

6) Diamond Head (WALIS RSL ID 544): *Porites* and *Pocillopora*, not *in situ* or not reported if *in situ*, at an elevation of +1.25 ± 1.25 m MSL have been dated at 114.6 ± 2.6 to 137 ± 11 ka (n = 7; 1-sigma uncertainty;
1085 WALIS U-series IDs 1209, 1210, 1773-1777; Ku et al., 1974; Szabo et al., 1994). All ages have been accepted.

7) Barbers Point (WALIS U-series ID 1779): Sherman et al. (1993) dated a single, *in situ Porites lobata* from a coral bafflestone at an elevation of $+3 \pm 3$ m MSL revealing an (accepted) age of 115 ± 10 ka.

1090 8) Kahe Point:

A), WALIS RSL ID 530; Reworked *Porites* and *Pocillopora*, and one *in situ Leptoseris* (WALIS U-series ID 1240) at an elevation of +9.75 ± 2.75 m MSL have been dated at 110.9 ± 3.5 to 134 ± 4 ka (n = 14 accepted out of 17 ages; WALIS U-series IDs 1188-1192, 1216-1220, 1240-1244, 1802, 1803; Szabo et al., 1994; Muhs and Szabo, 1994; Muhs et al., 2002; McMurtry et al., 2010). Two samples have been collected from coral-bearing conglomerates (WALIS U-series IDs 1802 and 1803; Muhs and Szabo, 1994).

B) WALIS RSL IDs 3976 and 3977 (Kahe Beach State Park; Hearty et al., 2007): Two Porites lobata from the lower reef at an elevation of +4.0 ± 0.5 m MSL have been dated at 115.35 ± 1.15 ka and 118.31 ± 0.89 ka (WALIS U-series IDs 1163, 1164; not reported if *in situ*). One *in situ Porites lobata* from the upper reef at an elevation of

 $\pm 9.0 \pm 0.5$ m MSL has been dated at $\pm 119.37 \pm 0.57$ ka (WALIS U-series ID ± 1165). *Porites lobata* lives at median and average depths of 0 and ± 19.2 m. All ages have not been accepted.

9) Kaena Point State Park:

- A), WALIS RSL ID 540; *Porites, Pocillopora* and one *Pavona*, not *in situ* or not reported if *in situ*, at an elevation of +2 ± 2 m MSL have been dated at 115 ± 6 to 127.5 ± 0.8 ka (n = 11 accepted out of 13 ages; WALIS U-series IDs 1178-1187, 1214, 1215, 1760; Ku et al., 1974; Szabo et al., 1994; Muhs et al., 2002).
 B) WALIS RSL ID 3978 (Makua Valley; Hearty et al., 2007): One *in situ Porites lobata* from the upper reef at an elevation of +8.0 ± 0.5 m MSL has been dated at 123.8 ± 1.3 ka (WALIS U-series ID 1166). *Porites lobata* lives at median and average depths of 0 and 19.2 m. The age has been accepted, however, no calcite content has been
- reported.

10) East of Kaena Point (WALIS RSL ID 541): *Porites* and *Pocillopora*, of those eight *in situ*, at an elevation of $\pm 1.5 \pm 1.5$ m MSL have been dated at 110.5 ± 3.8 to 138 ± 4 ka (n = 15 accepted out of 22 ages; WALIS U-series

1115 IDs 1195-1206, 1211-1213, 1793-1799; Szabo et al., 1994; Muhs and Szabo, 1994; Muhs et al., 2002). Three samples have been collected from coral cobble conglomerates (WALIS U-series IDs 1797-1799; Muhs and Szabo, 1994). hat gelöscht: (hat gelöscht:)



1120

Figure 3.; Cross-sections of outcrops of the Waimanalo Formation exposed on (a) Alala Point, (b) Kahuku Point, (c) the north shore of Oahu east of Kaena Point and (d) Kahe Point along with reliable U-series ages of corals (from Muhs et al., 2002).

- 1125 Veeh (1966) studied several sites on Oahu, which are detailed in Veeh (1965). *In situ Leptastrea* at an elevation of +2.25 ± 1.25 m MLWS have been dated at 110 ± 20 to 140 ± 30 ka (n = 4; WALIS RSL ID 547; WALIS Useries IDs 1739-1741, 1743; OAHU2, OAHU5, OAHU9 and OAHU22). However, the use of the alpha-counting technique resulted in large age uncertainties. *Leptastrea* lives at median and average depths of 7 and 17 m, respectively.
- 1130 Emerged reef terraces on Oahu described in all reviewed studies reach an elevation of up to +12.5 m MSL. The majority of ages was obtained from the first 4 m of the reef terraces and they range between 110.5 ± 3.8 and 138 ± 4 ka (n = 48 accepted ages). The terrace at about +7 m reveals ages of 112 ± 6 to 128 ± 8 ka (n = 12 accepted

ages). The +9 to +12 m terraces have been dated at 110.9 ± 3.5 to 134 ± 4 ka (n = 29 accepted ages). Sherman et 1135 al. (1993) deduced the occurrence of two highstands during the LIG. Szabo et al. (1994) reported on a LIG sea-

- level high stand on Oahu, which lasted about 17,000 years from 114 ka to 131 ka. Based on corals in growth position, Szabo et al. (1994) estimated that RSL was at least 7 m higher than present at 131 ka and remained higher until at least 115 ka. These authors mention a considerable 1-27 m depth range of living *Porites* and *Pocillopora*. Muhs et al. (2002) performed additional datings with a higher precision at the Szabo et al. (1994) study sites and
- 1140 confirmed the concept of a long LIG period on Oahu as proposed by Szabo et al. (1994). In 2002, Muhs et al. performed additional datings with a higher precision at the Szabo et al. (1994) study sites and confirmed the concept of a long LIG on Oahu as proposed by Szabo et al. (1994). However, they did not report any evidence for two separate sea-level stands as concluded by Sherman et al. (1993). McMurtry et al. (2010) did not consider the age data by Szabo et al. (1994) (white triangles in Fig. 34) as they seem to be biased toward lower elevation
- 1145 estimates. Furthermore, these authors did not consider the ages of elevated deposits from stages 5e and 9 (colored triangles) as they derived from energetic deposits related to storms or tsunamis rather than being elevated reefs with corals in growth position. Besides MIS 5, other interglacials from MIS 7 to 13 can be found in the coral reef record from Oahu. MIS 7 reefs have been documented around Oahu where they form a gently sloping nearshore submerged terrace (Sherman et al., 2014). The most reliable maximum elevation and the mean age for each
- 1150 highstand or terrace from Oahu is summarized in Figure 34. The MIS 5 terrace, the Waimanalo formation, is usually found at an elevation of +7.5 m. Based on this dataset, a linear uplift for Oahu has been estimated at 0.060 \pm 0.001 mm/yr over the past 500 ka (McMurtry et al., 2010), resulting in an uplift of about 7.5 m since 125 ka.

hat gelöscht: 2



1155

Figure 34: Mean or best ages versus maximum measured terrace elevation for known elevated reefs on Oahu (from McMurtry et al., 2010).

hat gelöscht: 2

Uncertainties of elevation measurements and datings are often large and do not allow an accurate conclusion about 1160 the timing and magnitude of the LIG RSL on Oahu. Furthermore, taxa have been often not determined, and therefore, an accurate palaeo-RSL could not be reconstructed. Determinations at the genus level often allow only rough estimates of palaeo-RSL.

Rubin et al. (2000) provided high-precision TIMS U-series ages for emerged coral reef terraces on Lana'i. These authors dated Hulopoe gravel coral clasts (no taxa determination, not *in situ*) at elevations of $\pm 19.5 \pm 0.5$ m MSL

and +21 ± 2 m MSL and obtained ages of 130.9 ± 1.5 ka and 134.8 ± 0.9 ka (WALIS RSL ID 552; WALIS U-series IDs 1806 and 1807; Lan1-2h-1 and Lan3-6-2) as well as 136.4 ± 0.9 ka and 136.7 ± 1.0 ka (WALIS RSL ID 553; WALIS U-series IDs 1808 and 1809; Lan3-5-3 and Lan2-4-3), respectively. The dated samples contained less than 2.3% of calcite and the ages can be considered as reliable. The elevation measurement method has not

1170 been reported, but the samples have been probably hand-leveled and the elevations refer to MSL. A water depth range of 0-30 m has been assumed as no taxa identification has been reported. The mean uplift rate of Lana'i has been estimated at 0.15 ± 0.22 mm/vr (Rubin et al., 2000).

6 Further details

1175 6.1 Last Interglacial sea-level fluctuations

Studies regarding tropical Pacific islands have contributed to the long-lasting and ongoing debate regarding the course of the Last Interglacial (LIG) relative sea-level (RSL) curve, including the timing, duration and amplitude of the MIS 5e highstand. This is of prime importance for many tropical Pacific islands, as the elevation of the LIG peak has been considered as a reference to assess rates of vertical displacement, uplift or subsidence.

- 1180 The occurrence of two sea-level peaks during the LIG highstand was inferred from the study of LIG reef terraces from various tropical Pacific islands: Papua New Guinea (Chappell and Veeh, 1978), Vanuatu (Jouannic et al., 1982), Fiji (Nunn and Omura, 1999; Nunn et al., 2002) and Hawaii (Sherman et al., 1993) and, since then, in a number of regions, including the Caribbean (Blanchon et al., 2009; Thompson et al., 2011), even if the timing of this event is not always substantiated by radiometric ages (see Camoin and Webster, 2015).
- In Papua New Guinea, ages obtained on the VIIa and VIIb terraces, that correspond to fringing and barrier reefs, respectively (Bloom et al., 1974; Chappell, 1974), fall into two tight groups centered at 118 ka and 134 ka. These two age clusters are separated vertically by 7 m, thus indicating that the RSL at 136 ka could have been 43 m lower than at 118 ka, given the 1.9 mm/yr uplift rate (Stein et al., 1993). The conflict between samples from Aladdin's Cave (about 100 m above present mean sea level (MSL) and about 130 ka) and from the coral reef terrace VII (up to 220 m above present MSL and about 134 ka) has been explained by a sea-level drop of 60 to 80 m after the development of the reef terrace VIIb (Esat et al., 1999). A new rise in sea level after 130 ka is deduced from the dataset and occurred in response to the major insolation maximum at 126 to 128 ka (Esat et al., 1999). The 118-ka corals of reefs VIIa and VIIb may have also been deposited during an episode of sea-level rise at the end of the LIG (Esat et al., 1999).
 - On Vanuatu, the duplication of the MIS 5e terrace in the Port Havannah area displays striking similarities with the distribution of VIIa and VIIb reef terraces from Papua New Guinea (Bloom et al., 1974; Chappell, 1974) and has been interpreted as resulting from low amplitude sea-level fluctuations during MIS 5e (Jouannic et al., 1982).

- On Fiji Islands, the age distribution is thought to reflect a double sea-level maximum with peaks around 133-130 ka and 123-120 ka (Nunn et al., 2002). The earlier maximum has been reported about 2 m lower than the later and marked by the growth of a surface reef, the later maximum appearing to have involved only cutting of erosional shorelines at the +5.0 to +5.5 m level (Nunn and Omura, 2003).
- The occurrence of two separate sea-level highstands on Oahu during the LIG has been considered by
 Sherman et al. (1993). In contrast, the concept of a long LIG period of about 17,000 years, from 114 ka to 131 ka, on Oahu has been proposed by Szabo et al. (1994) and later confirmed by Muhs et al. (2002). This concept involves an apparent local sea-level position 7 m higher than its current one by 131 ka, which remained higher than present until at least 115 ka (Szabo et al., 1994).
 - On the Cook Islands, Woodroffe et al. (1991) did not observe two periods of high RSL separated by a
 period of lower RSL during the LIG. However, they describe several stratigraphic features, such as
 erosional benches and notches, that might indicate RSL fluctuations during MIS 5e. However, these
 fluctuations have not been substantiated by chronological results.
 - On Saipan, Northern Mariana Islands, no evidence has been found for the occurrence of two sea-level peaks during the LIG (Muhs et al., 2020).

Formatiert: Einzug: Links: 1,27 cm, Keine Aufzählungen oder Nummerierungen

6.2 MIS 5a and MIS 5c

1210

1215

1220

Since our database is specifically focused on MIS 5e reef records, the table below lists data concerning MIS 5a and 5c, which were obtained on tropical Pacific islands but not inserted in our database. These data might be useful for future research on MIS 5 shorelines in tropical Pacific regions and are reported in this description paper for completeness.

Region	Sub-region / Site	Description
Vanuatu	Malakula	5c terrace at +204 m with an age of 107.6 ± 1.1 ka (Cabioch and Ayliffe, 2001) Terrace at +30 m with an age of 98 ± 11 ka. The terrace is assumed to correspond to the 103 ka palaeo- RSL.

		Terraces correspond to 82 and 103 ka terraces from Bloom et al., 1974 (Jouannic et al., 1980, 1982)	
	Efate	Terrace at +80 m dated at 106.4 ± 6.4 ka BP and at 103.5 ka (Tututuk area; Lecolle and Bernat, 1985) Terraces correspond to 82 and 103 ka terraces from Bloom et al., 1974 (Lecolle and Bernat, 1985; Jouannic et al., 1982) Terrace at + 70 m dated at 108 ka (Pangona area; Lecolle and Bernat, 1985) Terrace at +20 m dated at 107.9 ka (Siviri area; Lecolle and Bernat, 1985)	
	Santo	Terraces correspond to 82 and 103 ka terraces from Bloom et al., 1974 (Jouannic et al., 1980, 1982)	
	Malo	Terraces correspond to 82 and 103 ka terraces from Bloom et al., 1974 (Jouannic et al., 1982)	
	Hiw, Tegua, Loh, Toga	Terraces correspond to 82 and 103 ka terraces from Bloom et al., 1974 (Taylor et al., 1985). These reefs probably correspond, respectively, to the 80 and 103 ka reefs on New Guinea and Barbados.	
	Tanna	Terraces attributed to stages 5a, 5b and 5c (Neef et al., 2003)	
New Caledonia	Lifou	Terrace at +2.2 m dated at 90 \pm 4 ka (Bernat et al., 1976)	
	Maré	Terrace at +2 to +5 m gave ages of the order of 90-98 ka (Bernat et al., 1976).	

	Grande Terre (Amédée drill cores)	Unit 10 (37 to 14 mbsl) assigned to MIS 5 (100 to 130 ka; Cabioch et al., 2008)
Papua New Guinea		 Prominent reef terrace of reef complex V dated at 85 ka (Bloom et al., 1974) Reef terraces V, VI, VIIb taken as 82, 103, and 124 ka (Bloom et al., 1974) High RSL at 105 ka (Chappell, 1974) Reef terraces V and VI dated at 85 ± 1.4 ka and 107 ± 7.5 ka, respectively (Aharon and Chappell, 1986) Reef crest VIa assumed to have formed at 105 ka at 10 m below the present MSL (Esat et al., 1999) MIS 5b RSL was -57 m at 92.6 ± 0.5 ka; drop of about 40 m in approximately 10 kyr during MIS 5b-5a transition, Sea-level rise more than 40 m during the MIS 5b-5a transition, also in about 10 kyr. MIS 5a lasted until at least 76.2 ± 0.4 kyr, at a level of -24 m at that time (Cutler et al., 2003).
Hawaii	Oahu	Presence of MIS 5a and/or 5c on Oahu (Sherman, 1999) Presence of MIS 5a-d on Oahu (Sherman et al., 2014)

 Table 5: MIS 5a and MIS 5c reviewed in this study.

1230

6.3 Other interglacials

The table below includes data concerning the description and/or the dating of other Pleistocene interglacial periods on tropical Pacific islands from a number of studies. These data are not included in our database, which concerns 1235 only MIS 5e reef records, but are of pivotal importance for further projects dealing with the reconstruction of sea-

level changes and	shoreline evolution	n during the Ple	istocene in tropica	l Pacific islands.
0		0	1	

Region	Sub-region / Site	Description	
French Polynesia	Moruroa	MIS 7 and 9 reported (Camoin et al., 2001; Braithwaite and Camoin, 2011)	
1 orynosia	Takapoto	MIS 7 and 9 reported (Montaggioni et al., 2019a)	
Cook	Atiu, Mauke, Mitiaro, Mangaia	MIS 7 reported (Woodroffe et al., 1991)	
Pukapuka, Rakahanga	MIS 7, 9, 11, 13 and 15 indicated (Gray et al., 1992)		
Niue		MIS 7 reported (Kennedy et al., 2012)	
Tonga	Tongatapu	MIS 7 probably present (Taylor and Bloom, 1977)	hat gelöscht: ¶
Fiji	Kadavu, Kaibu	Reef terrace at +3.2 to +4.8 m dated from at 223 \pm 7.6 to 207.2 \pm 5.9 ka on Kadavu Island (Nagigia area; Nunn and Omura, 1999).	Formatiert: Einzug: Links: 0,63 cm

		These dates may indicate that the two reefs cannot be considered as separate entities and that the Nagigia reef is thus interpreted as having grown upwards to a RSL at least 7.1 m higher than present MSL, perhaps c. 208 ka. Nunn and Omura (2003) have considered that, in the absence of U-series ages, the two higher terraces on Kaibu Island, at 8.0-9.2 m and 12-14 m, could have formed during Stage 7 RSL maxima	hat gelöscht: 1
Tuvalu	Funafuti	Reef sequence between 27 and 58 m core depth in Funafuti drill cores attributed to stage 7. A Sr isotope age of 0.21 ± 0.21 Ma obtained on an unidentified coral collected at 36.6 m below the reef surface (Ohde et al., 2002).	
	Tanna	MIS 7 reported (Neef et al., 2003)	hat gelöscht: ¶
Vanuatu	Futuna	MIS 7, 9, 11 and 13 (Neef and McCulloch, 2001)	
	Loyalty Islands	Terrace at +2.5 m dated at ca 225 ka (Bourrouilh-Le Jan, 1985)	
New Caledonia	Grande Terre (Amédée drill cores)	MIS 7 and probably 9 or 11 (Frank et al., 2006) MIS 7 (221 to 248 ka) relate to Unit 9 (40–37 m core depth; Cabioch et al., 2008)	Formatiert: Einzug: Links: 0,63 cm
	Grande Terre (Ténia drill cores)	MIS 7 not identified; MIS 9 and probably MIS 11 and 13 reported (Montaggioni et al., 2011)	

Papua New Guinea		MIS 7 reported (Omura et al., 1994)	
Marshall Islands	Enewetak	MIS 7 or 9, MIS 9 or 11 or 13 probably found (Szabo et al., 1985),	hat gelöscht: ¶
Hawaii	Oahu	MIS 11, 13 or 15 probably present (Szabo et al., 1994) MIS 7 reported (Sherman et al., 1999; 2014) MIS 11 reported (Hearty, 2002) MIS 7, 9 and 13 reported; no evidence for MIS 11 (McMurtry et al., 2010)	hat gelöscht: ¶
	Lana'i, Moloka'i	MIS 7 reported (Rubin et al., 2000)	

hat gelöscht: 1

Table 6: Other interglacials reviewed in this study.

1245 6.4 Holocene sea-level indicators

The table below summarizes data concerning Holocene sea-level records from tropical Pacific islands. It especially includes results obtained on a mid-Holocene RSL highstand that appears as a significant feature in this region (Grossman et al., 1998; Dickinson, 2001, 2003, 2004; Woodroffe and Horton, 2005; Nunn, 2007; Hallmann et al., 2018). Studies of Holocene sea-level indicators for the reviewed islands are listed chronologically. These data are

1250 not inserted in our database, which specifically concerns MIS 5e reef records of RSL changes.

Region	References

French Polynesia	Lalou et al. (1966); Guilcher et al. (1969); Labeyrie et al. (1969); Montaggioni and Pirazzoli (1984); Trichet et al. (1984); Montaggioni (1985); Pirazzoli et al. (1985a, b); Pirazzoli and Montaggioni (1986); Pirazzoli (1987); Pirazzoli and Montaggioni (1987); Pirazzoli et al. (1987); Pirazzoli and Montaggioni (1988); Pirazzoli et al. (1988a, b); Camoin et al. (2001); Braithwaite and Camoin (2011); Rashid et al. (2014); Gischler et al. (2016); Hallmann et al. (2018); Gischler et al. (2019); Montaggioni et al. (2019b)	
Cook Islands	Schofield (1970); Stoddart (1972, 1975); Scoffin et al. (1985); Spencer et al. (1988); Yonekura et al. (1988); Woodroffe et al. (1990, 1991); Allen (1992); Gray et al. (1992); Ellison (1994); Yonekura (1994); Chikamori (1996); Allen (1998); Chikamori (1998, 2001); Moriwaki et al. (2006); Goodwin and Harvey (2008); Allen et al. (2016)	
Samoa	Mayor (1921, 1924); Stearns (1944); Kear and Wood (1959); Fairbridge (1961); Grant-Taylor and Rafter (1962); Shepard (1963); Stice and McCoy (1968); Matsushima et al. (1984); Rodda (1988); Sugimura et al. (1988a); Nunn (1991); Goodwin and Grossman (2003)	
Niue	Schofield (1959); Paulay and Spencer (1992); Nunn and Britton (2004); Kennedy et al. (2012)	
Tonga	Taylor and Bloom (1977); Kirch (1978); Taylor (1978); Ellison (1988); Nunn (1991); Dickinson et al. (1994), Nunn (1994, 1995a); Nunn and Finau (1995); Spenneman (1997)	
Fiji	Taylor (1978); Berryman (1979); Green (1979); Rodda (1986); Ash (1987); Miyata et al. (1988); Nunn (1988); Roy (1988); Sugimura et al. (1988b);	
	Miyata et al. (1990); Nunn (1990a, b); Shepard (1990); Nunn (1995b); Spriggs (1997); Nunn (2000); Nunn et al. (2000); Nunn and Peltier (2001); Thomas et al. (2004); Mörner and Matlack-Klein (2017)	
-----------------------------------	---	--
Tuvalu	David and Sweet (1904); Schofield (1977a); Kaplin (1981); McLean and Hosking (1991); Dickinson (1999); Ohde et al. (2002)	
Kiribati	Tracey (1972); Schoffeld (1977a, b); Marshall and Jacobson (1985); Woodroffe and McLean (1998); Woodroffe and Morrison (2001); Woodroffe et al. (2012); Yamano et al. (2017)	
Solomon Islands	Kirch and Yen (1982); Thirumalai et al. (2015)	
Vanuatu	Jouannic et al. (1980, 1982); Taylor et al. (1985); Cabioch and Ayliffe (2001); Neef et al. (2003)	
New Caledonia	Baltzer (1970); Coudray and Delibrias (1972); Bourrouilh-Le Jan (1985); Cabioch et al. (1989, 1995); Frank et al. (2006); Cabioch et al. (2008); Wirrmann et al. (2011)	
Papua New Guinea	Chappell and Polach (1972, 1991); Ota et al. (1993); Chappell (1994); Chappell et al. (1996); Riker-Coleman et al. (2006)	
Federated States of Micronesia	Wiens (1962); Shepard et al. (1967); Tracey (1968); Curray et al. (1970); Bloom (1970); Newell and Bloom (1970); Leach and Ward (1981); Matsumoto et al. (1986); Athens (1995); Kawana et al. (1995); Fujimoto et al. (1996)	

Mariana Islands	Kuenen (1933); Stearns (1941); Cloud et al. (1956); Tracey et al. (1964); Curray et al. (1970); <u>Easton et al. (1978); Randall and Siegrist (1988);</u> Kayanne et al. (1993)	
Marshall Islands	Wiens (1962); Thurber et al. (1965); Shepard et al. (1967); Curray et al. (1970); Newell and Bloom (1970); Tracey and Ladd (1974); Buddemeier et al. (1975); Dickinson (1999); Kayanne et al. (2011)	
Hawaii	Stearns (1941); Gross et al. (1969); Ladd et al. (1970); Stearns (1974); Easton and Olson (1976); Athens and Ward (1991); Fletcher and Jones (1996); Allen (1997); Athens (1997); Grigg (1998); Grossman and Fletcher (1998); Matsumoto et al. (1988); Jones (1992, 1998); Calhoun and Fletcher (1996); Rubin et al. (2000); Carson (2003, 2004); Nunn et al. (2007)	

Table 7: Holocene sea-level indicators reviewed in this study.

1265

6.5 Controversies

Besides past debates regarding the course of LIG RSL curve (see 6.1 'Last Interglacial sea-level fluctuations'), the study of MIS 5e reefs on tropical Pacific islands has generated several controversies mostly related to the interpretation of reef terraces. For example, McMurtry et al. (2010) did not consider the ages obtained by Szabo
1270 et al. (1994) on MIS 5e reef terrace on Oahu as they interpreted these deposits as being derived from energetic deposits related to storms or tsunamis rather than as corals in growth position. A similar controversial issue has

concerned the interpretation of LIG deposits on Lana'i seen either as highstand deposits, then mega-tsunami deposits, and eventually re-interpreted as highstand shoreline features deposited during the last two interglacials (see review in Webster et al., 2007).

1275

6.6 Uncertainties and data quality

Overall, the database contains 318 age, data points summarized in <u>94</u> RSL indicators and based on <u>38</u> studies on <u>MIS 5e</u> coral reef records in the tropical Pacific. One key aspect concerning the data quality is that 25 studies are older than 20 years and 14 studies are even older than 30 years. This impacts the quality of elevation measurements

(hat gelöscht: 284

hat gelöscht: 8 hat gelöscht: 5 hat gelöscht: indicators have been inserted in the database and the age information as methods improved considerably over the last decades. The data quality is mostly affected by the quality of the RSL data and, to a lesser extent, by the quality of age information. Tables 8 and 9 summarize the evaluation of the RSL data and the age information quality.

Description	Quality rating
Final RSL uncertainty is between one and two meters.	<u>4 (good)</u>
Final RSL uncertainty is between two and three meters.	<u>3 (average)</u>
Final RSL uncertainty is higher than three meters.	<u>2 (poor)</u>
Taxa determination only at the genus or family level with a large modern	<u>1 (very poor)</u>
No taxa determination. Final RSL uncertainty is very large, about 15	<u>1 (very poor)</u>

Description	Quality rating
Accurate age determination with a very narrow age range and uncertainties of up to ± 2 ka.	<u>5 (excellent)</u>
Narrow age range and uncertainties of \pm 2-4 ka allowing the attribution to a specific substage of MIS 5.	<u>4 (good)</u>
With uncertainties of \pm 4-15 ka the RSL data point can be attributed only to a generic interglacial (e.g., MIS 5).	<u>3 (average)</u>

hat gelöscht: ¶

Age determination using the alpha-counting technique results in large uncertainties with more than \pm 15 ka.

<u>2 (poor)</u>

Table 9: Evaluation of the age information quality.

No study related to LIG reef systems from the tropical Pacific islands includes a detailed description of *in situ* reef assemblages. Over the last two decades, *in situ* reef assemblages, including corals and associated biota (i.e., coralline algae, vermetid gastropods and encrusting foraminifera) rather than monospecific coral communities, have increasingly been used to more accurately constrain palaeo-water depths as well as other important palaeo-environmental parameters in the study of last deglacial and Holocene reef sequences (see review in Camoin and Webster, 2015). Large RSL uncertainties and limitations in the accuracy of RSL reconstructions can be considered for studies concerning MIS 5e coral reef records from the tropical Pacific islands. In addition, sedimentological and morphological data are barely described, thus hampering the accurate reconstruction of the relevant reef

1300 systems.

About $8\frac{3}{6}$ of the data points reveal a very poor RSL quality as the coral taxa have been determined only at the genus (about $67\frac{9}{6}$ of the data points) or family (about 10% of the data points) level. For about $1\frac{3}{6}$ of the data points no coral taxa has been reported at all. This introduces very large RSL uncertainties of ± 7 -15 m and excludes

- 1305 any constraint regarding the RSL position during MIS 5e. Some coral genera, e.g. *Porites* and *Pocillopora* that are the most commonly studied genera, display wide depth ranges in modern environments with median-average depths of 0-18.5 m and 0-15.6 m, respectively, thus precluding an accurate RSL reconstruction. Additional 10% of the data points reveal a poor RSL quality with final RSL uncertainties of more than three metres. Only less than 10% of the RSL data are of an average or good quality with final RSL uncertainties between one and three metres.
- 1310 These data are based on coral genera, e.g. *Leptoria*, which live at narrower depth ranges and can therefore be used to reconstruct RSL with uncertainties of about ± 2 m. Consequently, <u>about</u> 90% of the RSL information is not reliable and does not allow accurate RSL reconstructions during MIS 5e. Coral assemblages would need to be identified carefully and their palaeo-water depth significance determined accurately. A special attention should be given to RSL indicators typified by a narrow depth range to better constrain RSL reconstructions.
- 1315 Age information quality of the data points is significantly higher than their RSL quality. Age information of $\frac{39\%}{2000}$ of the data points is of excellent quality, i.e., ages have been accurately measured with very narrow age ranges and uncertainties of up to ± 2 ka. About 28% of the data points are still of good to average quality concerning the age information. 'Good quality' means that the ages of the coral samples could be determined with narrow age ranges

hat gelöscht: 0 hat gelöscht: 70 hat gelöscht: 5

hat gelöscht: one third

hat gelöscht: Another third

and uncertainties of \pm 2-4 ka allowing the attribution to a specific substage of MIS 5. 'Average quality' refers to

- 1325 ages with uncertainties of \pm 4-15 ka so that the RSL data point can be attributed only to a generic interglacial (e.g., MIS 5). The use of the alpha-counting technique resulted in large age uncertainties (\pm 4-20 ka). Based on this method, 32% of the ages reveal large uncertainties with more than \pm 15 ka. Ages have been accepted in 54% of the original studies, 46% of the ages have not been accepted and for the data points it is not known. A further problem is that the preservation of the samples is often unknown, implying that the reliability of the ages presented
- 1330 in the reviewed studies cannot be addressed. Only 41% of the studies report a calcite content of less than 3%, i.e., the samples are well preserved and the obtained ages can be considered as reliable. <u>About 57%</u> of the studies do not mention the preservation of the dated coral samples, so that the published age cannot be confirmed. Another source of uncertainties are inaccurate elevation measurements. Many studies used hand levelling and metered tape to determine the elevations of coral samples resulting in uncertainties of about ± 0.5-1 m. Some
- 1335 studies did even not report elevations and thus they had to be extracted from published sketches, which introduced large uncertainties. Overall, the elevation uncertainties are significant and might be up to 2-4 m, but they are still overprinted by the very large palaeo-water depth uncertainties introduced by the common lack of accurate studies of coral assemblages. Accurate GPS measurements would be needed to better constrain the elevation of RSL indicators.
- 1340 A further important aspect for sea-level studies is that the corals need to be in growth position and *in situ* to provide reliable information on past sea levels. However, for about <u>65% and 51%</u> of the data points the original studies do not mention if the dated corals have been collected in growth position or *in situ*, respectively. This introduces large uncertainties, even when these corals have been accurately dated, as they cannot be therefore considered as reliable sea-level indicators.
- 1345 The overall quality of this database is very poor mainly due to the poorly constrained palaeo-water depth range of coral assemblages, which introduces the largest uncertainty. The inaccuracy of elevation measurements, the lack of information on coral preservation, as well as the poor/missing information regarding the coral occurrence in LIG reefs introduce additional biases. Future work using more accurate dating methods and elevation measurements as well as a more detailed identification of the collected coral samples are essential to better
- 1350 constrain the MIS 5e RSL dataset concerning the tropical Pacific islands.

7 Future research directions

RSL change reconstructions require the combination of reliable radiometric ages and elevation measurements, as well as an accurate estimate of palaeo-water depths deduced from the modern distribution of relevant reef

-(hat gelöscht: another third
-(hat gelöscht: about
(hat gelöscht: 0
(hat gelöscht: about 20
-(hat gelöscht: 0
-(hat gelöscht: The other 60

-(hat formatiert: Schriftart: Kursiv)
(hat gelöscht: 8)
-(hat formatiert: Schriftart: Kursiv)

communities (see review in Camoin and Webster, 2015). Our newly compiled database (Hallmann and Camoin, 2020) demonstrates that most of the studies that have been carried out on tropical Pacific islands do not satisfy these requirements, thus hampering the accurate reconstruction of LIG RSL history. <u>This implies that these LIG</u>
 reef systems will have to be studied in more details with such a perspective.

Future research directions may therefore require to revisit LIG reef records from tropical Pacific islands, especially the key 'reference sites' (e.g., Papua New Guinea, Hawaii, Vanuatu), in order to collect the missing information that is crucially needed to reconstruct properly LIG RSL changes.

More accurate constraints on the timing, duration and amplitude of the highstand is pivotal to better constrain

- 1370 glacial isostatic adjustments during the LIG. In addition, despite the extensive literature on LIG RSL changes, only a few papers have tried to estimate suborbital-scale interglacial RSL variability during that time window (see reviews in Dutton and Lambeck, 2012; Woodroffe and Webster, 2014; Camoin and Webster, 2015). Higher resolution dating combined with accurate sedimentological studies of LIG reefs will be needed to decipher and reconstruct high-frequency RSL oscillations related to shorter climate excursions during the LIG period.
- 1375 Finally, the full understanding of the impact of RSL changes at various time scales on LIG reef growth will require the study of reef development patterns that have been so far barely investigated. <u>Facies sequences are also an</u> important component to reconstruct sea-level change and have been considered in parallel to the analysis of reef assemblages in many studies focused on last deglacial and Holocene reefs from the Indo-Pacific (see Camoin and Webster, 2015 and references therein).
- 1380 The standardized MIS 5e database provides an excellent tool for future LIG sea-level research as any missing information can be added to the WALIS database at a later stage for a more accurate reconstruction of MIS 5e <u>RSL changes.</u>

8 Data availability

- 1385 The database is available open access, and kept updated as necessary, at the following link: http://doi.org/10.5281/zenodo.3991672 (Hallmann and Camoin, 2020). The files at this link were exported from the WALIS database interface on 22/04/2021, and collated into a single .xls file from the different users who inserted them (G. Camoin and N. Hallmann). Description of each field in the database is contained at this link: https://doi.org/10.5281/zenodo.3961543 (Rovere et al., 2020), that is readily accessible and searchable here:
- 1390 <u>https://walis-help.readthedocs.io/en/latest/</u>. More information on the World Atlas of Last Interglacial Shorelines

hat formatiert: Englisch (USA)

 hat gelöscht: 19

 hat gelöscht: 8

 hat gelöscht: 0

can be found here: <u>https://warmcoasts.eu/world-atlas.html</u>. Users of our database are encouraged to cite the 1395 original sources alongside with our database and this article.

Author contribution

NH and GC wrote the manuscript and compiled the database; MH provided the palaeo-water depth significance of coral assemblages; MH and JW contributed to the manuscript during the final stage; NH produced the figures

Competing interests

The authors declare that they have no conflict of interest.

1405

Acknowledgments

The data used in this study were compiled in WALIS, a sea-level database interface developed by the ERC Starting Grant "WARMCOASTS" (ERC-StG-802414), in collaboration with PALSEA (PAGES / INQUA) working group.

1410 The database structure was designed by A. Rovere, D. Ryan, T. Lorscheid, A. Dutton, P. Chutcharavan, D. Brill, N. Jankowski, D. Mueller, M. Bartz, E. Gowan and K. Cohen. The data points used in this study were contributed to WALIS by G. Camoin and N. Hallmann (163 age data points and 94 RSL indicators), and by P. Chutcharavan (155 age data points). Basemap: "National Geographic Map", with data from National Geographic, Esri, Garmin, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, INCREMENT P Corp (Figs 5A,

1415 6, 7A, 8A-B, 9-1<u>3</u>). The authors gratefully acknowledge the guidance and support of the two Editors Alessio Rovere and Alexander Simms as well as the two referees Clark Sherman and Blake Dyer for providing constructive comments. hat gelöscht: 2

1420 References

Agassiz, A.: The coral reefs of the tropical Pacific, Bull. Mus. Comp. Zool., 28, 171–174, 1903. Aharon, P.: 140,000-yr isotope climatic record from raised coral reefs in New Guinea, Nature, 304, 720–723, 1983. Aharon, P. and Chappell, J.: Oxygen isotopes, sea level changes and the temperature history of a coral reef environment in New Guinea over the last 105 years, Palaeogeogr., Palaeoclimatol. Palaeoecol., 56, 337–379, 1986. Allen, M.S.: Dynamic Landscapes and Human Subsistence: Archaeological Investigations on Aitutaki Island, Southern Cook Islands, Ph.D. thesis, University of Washington, Seattle, USA, 557 pp., 1992.

Allen, J.: Pre-contact landscape transformation and cultural change in windward O'ahu, in: Historical Ecology in the Pacific Islands, edited by: Kirch, P. V. and Hunt, T. L., Yale University Press, New Haven, USA, 230–247, 1007

1430 1997.

Allen, M. S.: Holocene sea-level change on Aitutaki, Cook Islands: landscape change and human response, J. Coast. Res., 14, 10–22, 1998.

Allen, M. S., Morrison, A. E., Lorrey, A. M., Zhao, J.-X., and Jacobsen, G.E.: Timing, magnitude and effects of late Holocene sea level drawdown on island habitability, Aitutaki, Cook Islands, Archaeol. Ocean., 51, 108–121, 2016.

Anthony, S.: Hydrogeology and ground-water resources of Pingelap Island, Pingelap Atoll, State of Pohnpei, Federated States of Micronesia, U.S. Geological Survey, Water-Resources Investigations Report 92-4005, 40 pp., 1996a.

Anthony, S.: Hydrogeology and ground-water resources of Kahlap Island, Mwoakilloa Atoll, State of Pohnpei,

1440 Federated States of Micronesia, U.S. Geological Survey, Water-Resources Investigations Report 91-4184, 44 pp., 1996b.

Anthony, S.: Hydrogeology and ground-water resources of Ngatik Island, Sapwuahfik Atoll, State of Pohnpei, Federated States of Micronesia, U.S. Geological Survey, Water-Resources Investigations Report 93-4117, 44 pp., 1996c.

1445 Anthony, S. and Spengler, S.: Geology and ground-water resources reconnaissance of Lenger Island, State of Pohnpei, Federated States of Micronesia, 1991, U.S. Geological Survey, Water-Resources Investigations Report 93-4217, 13 pp., 1996.

Ash, J.: Holocene sea levels in northern Viti Levu, Fiji, New Zeal. J. Geol. Geop., 30, 431-435, 1987.

Ash, R. P., Carney, J. N., and Macfarlane, A.: Geology of Efate and Offshore Islands, New Hebrides Geological
 Survey Regional Report, University of California, USA, 49 pp., 1978.

Athens, J. S.: Landscape archaeology: prehistoric settlement, subsistence, and environment of Kosrae, Eastern Caroline Islands, Micronesia, report prepared for the Kosrae State Government, Federated States of Micronesia, International Archaeological Research Institute, Honolulu, Hawaii, USA, 474 pp., 1995.

Athens, J. S.: Hawaiian native lowland vegetation in prehistory, in: Historical ecology in the Pacific Islands, edited by: Kirch, P. and Hunt, T. L., Yale University Press, New Haven, USA, 248–270, 1997. Athens, J. S. and Ward, J. V.: Paleoenvironmental and archaeological investigations, Kawainui March flood control project, Oahu, U.S. Army Engineering Division, Pacific Ocean Environmental Section, Fort Shafter, Hawaii, International Archaeological Research Institute, Honolulu, Hawaii, USA, 129 pp., 1991.

Ayers, J. F. and Vacher, H. L.: Hydrogeology of an Atoll Island: A conceptual model from detailed study of a 1460 Micronesian example, Ground Water, 24, 185–198, 1986.

Ayling, B. F., Eggins, S., McCulloch, M. T., Chappell, J., Grün, R., and Mortimer G.: Uranium uptake history, open-system behaviour and uranium-series ages of fossil *Tridacna gigas* from Huon Peninsula, Papua New Guinea, Geochim. Cosmochim. Acta, 213, 475–501, 2017.

Baltzer, F.: Datation absolue de la transgression holocène sur la côte Ouest de Nouvelle-Calédonie sur des 1465 échantillons de tourbes à palétuviers. Interprétation néotectonique, C. R. Acad. Sci, D 271, 2251–2254, 1970.

Bernat, M., Launay, J., and Recy, J.: Datation à l'ionium de quelques formations coralliennes emergées de Nouvelle-Calédonie et des Îles Loyauté, C. R. Acad. Sci., D 282, 9–12, 1976.

Berryman, K.: Seismotectonic zoning study of the Fiji Islands, New Zealand Geological Survey Report 70, 41 pp., 1979.

1470 Blanchon, P., Eisenhauer, A., Fietzke, J., and Liebetrau, V.: Rapid sea-level rise and reef back-stepping at the close of the last interglacial highstand, Nature, 458, 881–884, 2009.

Bloom, A. L.: Paludal stratigraphy of Truk, Ponape, and Kusai, Eastern Caroline Islands, Geol. Soc. Am. Bull., 81, 1895–1904, 1970.

Bloom, A. L., Broecker, W. S., Chappell, J., Matthews, R. K., and Mesolella, K. J.: Quaternary sea level 1475 fluctuations on a tectonic coast; New ²³⁰Th/²³⁴U dates from the Huon Peninsula, New Guinea, Quat. Res., 4, 185–

205, 1974.

Bloom, A. L., Jouannic, C., and Taylor, F. W.: Preliminary radiometric ages from the uplifted quaternary coral reefs of Efate, New Hebrides, in: Geology of Efate and Offshore Islands, edited by: Ash, R. P., Carney, J. N., and Macfarlane, A., New Hebrides Geological Survey Regional Report, University of California, USA, 47–49, 1978.

- Bonney, T. G.: The atoll of Funafuti: Borings into a coral reef and the results, being the Report of the Coral Reef Committee of the Royal Society, The Royal Society of London, London, UK, 19 pp., 1904.
 Bourrouilh-Le Jan, F. G.: An attempt at datation by isotopic stratigraphy of Holocene and Pleistocene notches of some so-called uplifted atolls (high carbonate islands), Proc. 5th Int. Coral Reef Congr. 3, Tahiti, 27 May–1 June 1985, 125–130, 1985.
- 1485 Braithwaite, C. J. R. and Camoin, G. F.: Diagenesis and sea-level change: lessons from Moruroa, French Polynesia, Sedimentology, 58, 259–284, 2011.

Buddemeier, R. W., Smith, S. V., and Kinzie, R.A.: Holocene windward reef-flat history, Enewetok Atoll, Geol. Soc. Am. Bull., 86, 1581–1584, 1975.

Cabioch, G.: Récifs frangeants de Nouvelle-Calédonie (Pacifique sud-ouest). Structure interne et influences de

1490 l'eustatisme et de la néotectonique, Ph.D. thesis, Université de Provence, Aix-en-Provence, France, 291 pp., 1988. Cabioch, G. and Ayliffe, L. K.: Raised coral terraces at Malakula, Vanuatu, Southwest Pacific, indicate high sea level during marine isotope stage 3, Quat. Res., 56, 357–365, 2001.

Cabioch, G., Thomassin, B. A., and Lecolle, J.: Age d'émersion des récifs frangeants holocènes autour de la 'Grande Terre' de Nouvelle-Calédonie; nouvelle interprétation de la courbe des niveaux marins depuis 8000 ans
BP, C. R. Acad. Sci., II 308, 419–425, 1989.

Cabioch, G., Montaggioni, L. F., and Faure, G.: Holocene initiation and development of New Caledonian fringing reefs, SW Pacific, Coral Reefs, 14, 131–140, 1995.

Cabioch, G., Récy, J., Jouannic, C., and Turpin L.: Contrôle environnemental et néotectonique de l'édification récifale en Nouvelle-Calédonie au tours du Quaternaire terminal, Bull. Soc. Géol. Fr., 167, 729–742, 1996.

- Cabioch, G., Correge, T., Turpin, L., Castellaro, C., and Récy, J., 1999. Development patterns of fringing and barrier reefs in New Caledonia (southwest Pacific), Oceanol. Acta, 22, 567–578, 1999.
 Cabioch, G., Montaggioni, L., Thouveny, N., Frank, N., Sato, T., Chazottes, V., Dalamasso, H., Payri, C., Pichon, M., and Sémah, A.-M.: The chronology and structure of the western New Caledonian barrier reef tracts, Palaeogeogr., Palaeoclimatol. Palaeoceol., 268, 91–105, 2008.
- 1505 Calhoun, R. S. and Fletcher, C. H.: Late Holocene coastal plain stratigraphy and sea-level history at Hanalei, Kauai, Hawaiian Islands, Quat. Res., 45, 47–58, 1996.

Camoin, G. F. and Webster, J. M.: Coral reef response to Quaternary sea-level and environmental changes: State of the science, Sedimentology, 62, 401–428, 2015.

Camoin, G. F., Ebren, P., Eisenhauer, A., Bard, E., and Faure, G.: A 300,000-yr coral reef record of sea level

1510 changes, Mururoa atoll (Tuamotu archipelago, French Polynesia), Palaeogeogr. Palaeoclimatol. Palaeoecol., 175, 325–341, 2001.

Carson, M. T.: Integrating fragments of a settlement pattern and cultural sequence in Wainiha Valley, Kaua'i, Hawaiian Islands, People and Culture in Oceania, 19, 83–105, 2003.

Carson, M. T.: Resolving the enigma of early coastal settlement in the Hawaiian Islands: The stratigraphic sequence of the Wainiha Beach site in Kaua'i, Geoarchaeology, 19, 99–118, 2004.

Chappell, J.: Astronomical theory of climatic change: Status and problem, Quat. Res., 3, 221-236, 1973.

Chappell, J.: Geology of coral terraces, Huon Peninsula, New Guinea: A study of Quaternary tectonic movements and sea level changes, Geol. Soc. Am. Bull., 85, 553–570, 1974.

Chappell, J.: A revised sea level record for the last 300,000 yrs from Papua New Guinea, Search, 4, 99-101, 1983.

1520 Chappell, J.: Upper Quaternary sea levels, coral terraces, oxygen isotopes and deep-sea temperatures, J. Geogr., 103, 828–840, 1994.

Chappell, J. and Polach, H. A.: Some effects of partial recrystallization on ¹⁴C dating late Pleistocene corals and molluses, Quat. Res., 2, 244–252, 1972.

Chappell, J. and Polach, H. A.: Post-glacial sea level rise from a coral record at Huon Peninsula, Papua New 1525 Guinea, Nature, 349, 147–149, 1991.

Chappell, J. and Shackleton, N. J.: Oxygen isotopes and sea level, Nature, 324, 137-140, 1986.

Chappell, J. and Veeh, H. H.: Late Quaternary tectonic movements and sea-level changes at Timor and Atauro Island, Geol. Soc. Am. Bull., 89, 356–368, 1978.

Chappell, J., Ota, Y., and Berryman, K.: Late Quaternary coseismic uplift history of Huon Peninsula, Papua New 1530 Guinea, Quat. Sci. Rev., 15, 7–22, 1996.

Chen, M.-C., Frohlich, C., Taylor, F. W., Burr, G., and van Ufford, A. Q.: Arc segmentation and seismicity in the Solomon Islands arc, SW Pacific, Tectonophysics, 507, 47–69, 2011.

Chikamori, M.: Development of coral reefs and human settlement: Archaeological research in the northern Cook Islands and Rarotonga, Bull. Indo-Pacific Prehist. Assoc., 15, 45–52, 1996.

1535 Chikamori, M.: Archaeological Studies in the Cook Islands, Series 2, Occasional Papers of the Department of Archaeology and Ethnology, Keio University, Tokyo, 12, 53 pp., 1998.

Chikamori, M.: Landscape changes in the coastal plains of fringing reef islands: Rarotonga in the Cook Islands, Shigaku, 70, 331–350, 2001.

Chutcharavan, P. M. and Dutton, A.: A Global Compilation of U-series Dated Fossil Coral Sea-level Indicators

1540 for the Last Interglacial Period (MIS 5e), Earth Syst. Sci. Data Discuss. [preprint], https://doi.org/10.5194/essd-2020-381, in review, 2020.

Cloud, P. E., Jr.: Superficial aspects of modern organic reefs, The Scientific Monthly, 79, 195–208, 1954.
 Cloud, P. E., Jr., Schmidt, R. G., and Burke, H.W.: Geology of Saipan, Mariana Islands, Part 1, General Geology,
 U.S. Geological Survey Professional Paper 280-A, 126 pp., 1956.

1545 Couch, R. F., Jr., Fetzer, J. A., Goter, E. R., Ristvet, B. L., Tremba, E. L., Walter, D. R., and Wendland, V. P.: Drilling Operations on Eniwetok Atoll during Project EXPOE, Air Force Weapons Laboratory, AFWL-TR-75-216, 270 pp., 1975. Coudray, J.: Nouvelles données sur la nature et l'origine du complexe récifal côtier de la Nouvelle Calédonie, Quat. Res., 1, 236–246, 1971.

1550 Coudray, J.: Recherches sur le Néogène et le Quaternaire marins de la Nouvelle- Calédonie: contribution de l'étude sédimentologique à la connaissance de l'histoire géologique post-Eocène, Éditions de la Fondation Singer-Polignac, Paris, France, 276 pp., 1976.

Coudray, J. and Delibrias, G.: Variations du niveau marin au-dessus de l'actuel en Nouvelle-Calédonie depuis 6000 ans, C. R. Acad. Sci., D 375, 2623–2626, 1972.

- Curray, J. R., Shepard, F. P., and Veeh, H. H.: Late Quaternary sea-level studies in Micronesia: CARMARSEL expedition, Geol. Soc. Am. Bull., 81, 1865–1880, 1970.
 Cutler, K. B., Edwards, R. L., Taylor, F. W., Cheng, H., Adkins, J., Gallup, C. D., Cutler, P. M., Burr, G. S., and Bloom, A. L.: Rapid sea-level fall and deep-ocean temperature change since the last interglacial period, Earth Planet. Sci. Lett., 206, 253–271, 2003.
- Daly, R. A.: A general sinking of sea-level in recent time, Proc. Natl. Acad. Sci. USA, 6, 246–250, 1920.
 Dana, J. D.: Corals and Coral Islands, Dodd and Mead, New York, USA, 428 pp., 1872.
 Darwin, C.: The structure and distribution of coral reefs: Being the first part of the geology of the voyage of the Beagle, under the command of Capt. Fitzroy, R. N. during the years 1832 to 1836, Smith Elder and Co., London, UK, 1842.
- 1565 David, T. W. E.: Section IV. Narrative of the second and third expeditions, in: The atoll of Funafuti: borings into a coral reef and the results, being the report of the Coral Reef Committee of the Royal Society, edited by: Armstrong, H. E., Blanford W. T., Bonney, T. G., Crookes, W., Darwin, F., Evans, J., Geikie, A., Hinde, G. J., Judd, J. W., Lankester, E. R., Lapworth, C., Murray, J., Sollas, W. J., Sorby, H. C., Teall, J. J. H., Wharton, W. J. L., Wolfe, B., Field, A. M., and Watts, W. W., The Royal Society of London, London, UK, 40–60, 1904.
- 1570 David, T. W. E. and Sweet, G.: The geology of Funafuti, in: The atoll of Funafuti: borings into a coral reef and the results, being the report of the Coral Reef Committee of the Royal Society, edited by: Armstrong, H. E., Blanford W. T., Bonney, T. G., Crookes, W., Darwin, F., Evans, J., Geikie, A., Hinde, G. J., Judd, J. W., Lankester, E. R., Lapworth, C., Murray, J., Sollas, W. J., Sorby, H. C., Teall, J. J. H., Wharton, W. J. L., Wolfe, B., Field, A. M., and Watts, W. W., The Royal Society of London, London, UK, 61–124, 1904.
- 1575 Degaugue-Michalski, F.-M.: Croissance et évolution d'édifices récifaux du Pacifique occidental (Nouvelle-Calédonie, Chesterfield) à l'Holocène et au Pléstocène, Ph.D. thesis, Université de Provence Aix-Marseille-I, France, 227 pp., 1993.

Dickinson, W. R.: Holocene sea-level record on Funafuti and potential impact of global warming on Central Pacific Atolls, Quat. Res., 51, 124–132, 1999.

1580 Dickinson, W. R.: Paleoshoreline record of relative Holocene sea levels on Pacific Islands. Earth-Sci. Rev., 55, 191–234, 2001.

Dickinson, W. R.: Impact of Mid-Holocene hydro-isostatic highstand in regional sea level on habitability of islands in Pacific Oceania, J. Coast. Res., 19, 489–502, 2003.

Dickinson, W. R.: Impacts of eustasy and hydro-isostasy on the evolution and landforms of Pacific atolls, 1585 Palaeogeogr., Palaeoclimatol. Palaeoecol., 213, 251–269, 2004.

Dickinson, W. R., Burley, D. V., and Shutler, R., Jr.: Impact of hydroisostatic Holocene sea-level change on the geologic context of island archaeological sites, Northern Ha'apai Group, Kingdom of Tonga, Geoarchaeology, 9, 85–111, 1994.

Dutton, A. and Lambeck, K.: Ice volume and sea level during the Last Interglacial, Science, 337, 216–219, 2012.

1590 Easton, W. H. and Olson, E. A.: Radiocarbon profile of Hanauma Reef, Oahu, Hawaii, Geol. Soc. Am. Bull., 87, 711–719, 1976.

Easton, W. H., Ku, T. L., and Randall, R. H.: Recent reefs and shore lines of Guam, Micronesica, 14, 1–11, 1978.
 Edwards, R. L., Chen, J. H., and Wasserburg, G. J.: ²³⁸U-²³⁴U-²³⁰Th-²³²Th systematics and the precise measurement of time over the past 500,000 years, Earth Planet. Sci. Lett., 81, 175–192, 1987a.

1595 Edwards, R. L., Chen, J. H., Ku, T.-L., and Wasserburg, G. J.: Precise timing of the last interglacial period from mass spectrometric determination of thorium-230 in corals, Science, 236, 1547–1553, 1987b. Ellison, J. C.: Holocene sea level record of Tongatapu, Kingdom of Tonga, from pollen analysis of mangrove sediments, Master thesis, Simon Fraser University, Canada, 94 pp., 1988.

Ellison, J. C.: Paleo-lake and swamp stratigraphic records of Holocene vegetation and sea-level changes, Mangaia, 1600 Cook Islands, Pac. Sci., 48, 1–15, 1994.

Emery, K. O.: Marine geology of Guam, U.S. Geological Survey Professional Paper 403-B, 76 pp., 1962.

Emery, K. O.: Marine geology of Guam, U.S. Geological Survey Professional Paper 403-B, 76 pp., 1963.

Emery, K. O., Tracey, J. I., Jr., and Ladd, H. S.: Geology of Bikini and nearby atolls, U.S. Geological Survey Professional Paper 260-A, 265 pp., 1954.

Esat, T. M., McCulloch, M. T., Chappell, J., Pillans, B., and Omura, A.: Rapid fluctuations in sea level recorded at Huon Peninsula during the Penultimate Deglaciation, Science, 283, 197–201, 1999.
 Fairbridge, R. W.: Eustatic charges in sea level, in: Physics and chemistry of the Earth, edited by: Ahrens, L. H., Press, F., Rankama, K., and Runcorn, S. K., Pergamon Press, New York, USA, 99–185, 1961.

Fletcher, C. H. and Jones, A. T.: Sea-level highstand recorded in Holocene shoreline deposits on Oahu, Hawaii, J.Sediment. Res., 66, 632–641, 1996.

Fletcher, C. H. and Richmond, B. M.: Climate change in the Federated States of Micronesia: food and water security, climate risk management, and adaptive strategies, report of findings 2010, University of Hawai'i Sea Grant College Program, USA, 32 pp., 2010.

Frank, N., Turpin, L., Cabioch, G., Blamart, D., Tressens-Fedou, M., Colin, C., and Jean-Baptiste, P.: Open system

1615 U-series ages of corals from a subsiding reef in New Caledonia: Implications for sea level changes, and subsidence rate, Earth Planet. Sci. Lett., 249, 274–289, 2006.

Fujimoto, K., Miyagi, T., Kikuchi, T., and Kawana, T.: Mangrove habitat formation and response to Holocene sea-level changes on Kosrae Island, Micronesia, Mangroves and Salt Marshes, 1, 47–57, 1996.

Gaven, C. and Bourrouilh-Le Jan, F.: Géochronologie (²³⁰Th-²³⁴U-²³⁸U) et sédimentologie des faciés récifaux
 pléstocénes à Maré, Archipel des Loyautés, SW Pacifique, Océanis, 7, 347–366, 1981.

Gischler, E., Hudson, J. H., Humblet, M., Braga, J. C., Eisenhauer, A., Isaack, A., Anselmetti, F. S., and Camoin, G. F.: Late Quaternary barrier and fringing reef development of Bora Bora (Society Islands, south Pacific): First subsurface data from the Darwin-type barrier-reef system, Sedimentology, 63, 1522–1549, 2016.

Gischler, E., Hudson, J. H., Humblet, M., Braga, J. C., Schmitt, D., Isaack, A., Eisenhauer, A., and Camoin, G. F.:

Holocene and Pleistocene fringing reef growth and the role of accommodation space and exposure to waves and currents (Bora Bora, Society Islands, French Polynesia), Sedimentology, 66, 305–328, 2019.
 Goodwin, I. and Grossman, E. E.: Middle to late Holocene coastal evolution along the south coast of Upolu Island, Samoa, Mar. Geol., 202, 1–16, 2003.

Goodwin, I. D. and Harvey, N.: Subtropical sea-level history from coral microatolls in the Southern Cook Islands,
 since 300 AD, Mar. Geol., 253, 14–25, 2008.

Grant-Taylor, T. L. and Rafter, T. A.: New Zealand radiocarbon age measurements, New Zeal. J. Geol. Geop., 5, 331–359, 1962.

Gray, S. C., Hein, J. R., Hausmann, R., and Radtke, U.: Geochronology and subsurface stratigraphy of Pukapuka and Rakahanga atolls, Cook Islands: Late Quaternary reef growth and sea level history, Palaeogeogr.,
Palaeoclimatol. Palaeoceol., 91, 377–394, 1992.

Green, R. C.: Lapita, in: The prehistory of Polynesia, edited by: Jennings, J. D., Australian National University, Canberra, Australia, 27–60, 1979.

Grigg, R. W.: Holocene coral reef accretion in Hawaii: a function of wave exposure and sea level history, Coral Reefs, 17, 263–272, 1998.

1640	Gross, M. G., Milliman, J. D., Tracey, J. I., and Ladd, H. S.: Marine geology of Kure and Midway atolls, Hawaii:	
	a preliminary report, Pac. Sci., 23, 17–25, 1969.	
	Grossman, E. E. and Fletcher, C. H.: Sea level higher than present 3500 years ago on the northern main Hawaiian	
	Islands, Geology, 26, 363-366, 1998.	
	Grossman, E. E., Fletcher, C. H., and Richmond, B. M.: The Holocene sea-level highstand in the equatorial Pacific:	
1645	analysis of the insular paleosea-level database, Coral Reefs, 17, 309-327, 1998.	
	Grover, J. C.: The Solomon Islands: geological exploration and research, 1953-1956, Geological Survey of the	
	British Solomon Islands, Memoir 2, 150 pp., 1958.	
	Guilcher, A., Berthois, L., Doumenge, F., Michel, A., Saint-Requier, A., and Arnold, R.: Les récifs et lagons	
	coralliens de Mopelia et de Bora Bora (Îles de la Société), Mémoires ORSTOM, 38, 107 pp., 1969.	
1650	Hallmann, N., Camoin, G., Eisenhauer, A., Botella, A., Milne, G. A., Vella, C., Samankassou, E., Pothin, V.,	
	Dussouillez, P., Fleury, J., and Fietzke, J.: Ice volume and climate changes from a 6000 year sea-level record in	
	French Polynesia, Nat. Commun., 9, 285, <u>https://doi.org/10.1038/s41467-017-02695-7</u> , 2018.	
	Hallmann, N. and Camoin, G.: Database of Last Interglacial (MIS 5e) sea-level proxies on tropical Pacific islands	hat gelöscht: WALIS d
	[dataset], Zenodo, http://doi.org/10.5281/zenodo.3991672, 2020.	
1655	Hearty, P. J.: The Ka'ena Highstand of O'ahu, Hawai'i: further evidence of Antarctic ice collapse during the Middle	
1	Pleistocene, Pac. Sci., 56, 65-81, 2002.	
	Hearty, P. J., Hollin, J. T., Neumann, A. C., O'Leary, M. J., and McCulloch M.: Global sea-level fluctuations	
	during the Last Interglaciation (MIS 5e), Quat. Sci. Rev., 26, 2090-2112.	
1	Henry, T. W. and Wardlaw, B. R.: Introduction: Enewetak Atoll and the PEACE Program, geological and	
1660	geophysical investigations of Enewetak Atoll, Republic of the Marshall Islands, U.S. Geological Survey	
	Professional Paper 1513-A, 29 pp., 1990.	
	Henry, R. W., Mercer, J. W., and Zbur, R. T.: Near surface geologic investigations at Eniwetok Atoll, in:	
1	Proceedings of the 2nd International Symposium on Coral Reefs, Great Barrier Reef Committee, Brisbane,	hat formatiert: Hochgestellt
1	Australia, 615–626, 1974.	
1665	Henry, T. W., Wardlaw, B. R., Skipp, B., Major, R. P. and Tracey, J. I., Jr.: Pacific Enewetak Crater Exploration	
	(PEACE) Program, Enewetak Atoll, Republic of the Marshall Islands, Part 1: Drilling operations and descriptions	
	of bore holes in vicinity of KOA and OAK craters, U.S. Geological Survey Open File Report 86-419, 583 pp.,	
	https://doi.org/10.3133/ofr86419, 1986.	

1670 Hibbert, F. D., Rohling, E. J., Dutton, A., Williams, F. H., Chutcharavan, P. M., Zhao, C., and Tamisiea, M. E.: Coral indicators of past sea-level change: a global repository of U-series dated benchmarks, Quat. Sci. Rev., 145, 1–56.

Hibbert, F. D., Williams, F. H., Fallon, S. J., and Rohling, E. J.: A database of biological and geomorphological sea-level markers from the Last Glacial Maximum to present, Scientific Data, 5:180088, 1675 doi:10.1038/sdata.2018.88.

- Hinde, G. J.: Report on the materials from the borings at the Funafuti atoll, in: The atoll of Funafuti: borings into a coral reef and the results, being the report of the Coral Reef Committee of the Royal Society, edited by:
 Armstrong, H. E., Blanford W. T., Bonney, T. G., Crookes, W., Darwin, F., Evans, J., Geikie, A., Hinde, G. J., Judd, J. W., Lankester, E. R., Lapworth, C., Murray, J., Sollas, W. J., Sorby, H. C., Teall, J. J. H., Wharton, W. J.
- 1680 L., Wolfe, B., Field, A. M., and Watts, W. W., The Royal Society of London, London, UK, 186–361, 1904. Hongo, C. and Denis Wirrmann, D.: Preliminary identification of key coral species from New Caledonia (Southwest Pacific Ocean), their significance to reef formation, and responses to environmental change, Isl. Arc, 24, 31–46, 2015.

Huang, D., Benzoni, F., Fukami, H., Knowlton, N., Smith, N. D., and Budd, A. F.: Taxonomic classification of the reef coral families Merulinidae, Montastraeidae, and Diploastraeidae (Cnidaria: Anthozoa: Scleractinia), Zool.

J. Linn. Soc., 171, 277-355, 2014.

Jones, A. T.: Holocene coral reef on Kauai, Hawaii: Evidence for a sea-level highstand in the central Pacific, in: Quaternary coasts of the United States: marine and lacustrine systems, edited by: Fletcher III, C. H. and Wehmiller, J. F., SEPM Special Publication, 48, 267–271, 1992.

1690 Jones, A. T.: Elevated fossil coral deposits in the Hawaiian Islands: A measure of island uplift in the Quaternary, Ph.D. thesis, University of Hawaii, Honolulu, USA, 274 pp., 1993.

Jones, A. T.: Late Holocene shoreline development in the Hawaiian Islands, J. Coast. Res., 14, 3-9, 1998.

Jouannic, C., Taylor, F. W., Bloom, A. L., and Bernat, M.: Late Quaternary uplift history from emerged reef terraces of Santo and Malekula islands, central New Hebrides island arc, UN ESCAP, CCOP/SOPAC Tech. Bull., 3, 91–108, 1980.

Jouannic, C., Taylor, F. W., and Bloom, A. L.: Sur la surrection et la déformation d'un arc jeune: l'arc des Nouvelles Hébrides, ORSTOM, 147, 223–246, 1982.

Kaplin, P.A.: Relief, age and types of Oceanic islands, New Zeal. Geogr., 36, 3-12, 1981.

 Kawana, T., Miyagi, T., Fujimoto, K., and Kikuchi, T.: Late-Holocene sea-level changes and mangrove
 development in Kosrae Island, the Carolines, Micronesia, in: Rapid sea-level rise and mangrove habitat, edited by: Kikuchi, T., Gifu University Institute for Basin Ecosystems, Gifu, Japan, 7 pp., 1995.
 Kayanne, H., Ishii, T., Matsumoto, E., and Yonekura, N.: Late Holocene sea-level change on Rota and Guam,

Mariana Islands, and its constraint on geophysical predictions, Quat. Res., 40, 189–200, 1993.

Kayanne, H., Yasukochi, T., Yamaguchi, T., Yamano, H., and Yoneda, M.: Rapid settlement of Majuro Atoll,

1705 central Pacific, following its emergence at 2000 years CalBP, Geophys. Res. Lett., 38, https://doi.org/10.1029/2011GL049163, 2011.

Kear, D. and Wood, B. L.: The geology and hydrology of Western Samoa, New Zeal. Geol. Surv. Bull., 63, New Zealand Department of Scientific and Industrial Research, 95 pp., 1959.

Kennedy, D. M., Marsters, T. H., Woods, J. L. D. and Woodroffe, C. D.: Shore platform development on an uplifting limestone island over multiple sea-level cycles, Niue, South Pacific, Geomorphology, 141–142, 170– 182, 2012.

Kirch, P. V.: The Lapitoid period in West Polynesia: excavations and survey on Niuatoputapu, Tonga, J. Field Arch., 5, 1–13, 1978.

Kirch, P. V. and Yen, D. E.: Tikopia: The prehistory and ecology of a Polynesian outlier, Bernice P. BishopMuseum Bulletin, 238, Bishop Museum Press, Honolulu, USA, 396 pp., 1982.

Ku, T.-L., Kimmel, M. A., Easton, W. H., and O'Neil, T. J.: Eustatic sea level 120,000 years ago on Oahu, Hawaii, Science, 183, 959–962, 1974.

Kuenen, P. H.: Geology of coral reefs: the Snellius Expedition in the Eastern Part of the Netherlands East Indies 1929-1930, geological results, volume 5, Kemink en Zoon, Utrecht, The Netherlands, 125 pp., 1933.

Labeyrie, J., Lalou, C., and Delibrias, G.: Étude des transgressions marines sur l'atoll de Mururoa par la datation des différents niveaux de corail, Cahiers du Pacifique, 13, 59–68, Monographie Mururoa, 202–212, 1969.
Ladd, H. S. and Schlanger, S. O.: Drilling operations on Eniwetok atoll: Bikini and nearby atolls, Marshall Islands, U.S. Geological Survey Professional Paper 260-Y, 863–903, 1960.

Ladd, H. S., Tracey, J. I., Jr., and Lill, G. G.: Drilling on Bikini Atoll, Marshall Islands, Science, 107, 51-55, 1948.

 Ladd, H. S., Ingerson, E., Townsend, R. C., Russell, M., and Stephenson, K.: Drilling on Eniwetok Atoll, Marshall Islands, Bull. Amer. Assoc. Petrol. Geol., 37, 2257–2280, 1953.
 Ladd, H. S., Tracey, J. I., Jr., and Gross, M. G.: Drilling on Midway Atoll, Hawaii, Science, 156, 1088–1094,

1967.

89

Ladd, H. S., Tracey, J. I., Jr., and Gross, M. G.: Deep drilling on Midway Atoll, U.S. Geological Survey 1730 Professional Paper 680-A, 22 pp., 1970.

Lalou, C., Labeyrie, J., and Delibrias, G.: Datation des calcaires coralliens de l'atoll de Mururoa (archipel des Tuamotu) de l'époque actuelle jusqu'à -500 000 ans, C. R. Acad. Sci., D 263, 1946–1949, 1966.

Launay J. and Récy J.: Variations relatives du niveau de la mer et néotectonique en Nouvelle-Calédonie au Pléistocène supérieur et à l'Oligocène, Rev. Géogr. Phy. Géol. Dyn., 14, 47–66, 1972.

- 1735 Leach, F. and Ward, G. K.: Archaeology on Kapingamarangi Atoll, a Polynesian outlier in the Eastern Caroline Islands, Studies in Prehistoric Anthropology, 16, University of Otago, New Zealand, 150 pp., 1981. Lecolle, J. and Bernat, M.: Late Quaternary surrection history of Efate island, New Hebrides island arc (Vanuatu): Th/U dates from uplifted terraces = Histoire de la surrection de l'île d'Efate, arc insulaire des Nouvelles-Hébrides (Vanuatu) au cours du Quaternaire récent: datations de terrasses soulevées par la méthode U/Th, Proc. 5th Int.
- Coral Reef Congr. 3, Tahiti, 27 May–1 June 1985, 179–184, 1985.
 Ludwig, K. R., Halley, R. B., Simmons, K. R., and Peterman, Z. E.: Strontium-isotope stratigraphy of Enewetak Atoll, Geology, 16, 173–177, 1988.

Mallick, D. I. J. and Greenbaum, D.: Geology of southern Santo, New Hebrides Geological Survey Regional Report, 84 pp., 1977.

1745 Marshall, J. F. and Jacobson, G.: Holocene growth of a mid-Pacific atoll: Tarawa, Kiribati, Coral Reefs, 4, 11–17, 1985.

Marshall, J.F. and Launay, J.: Uplift rates of the Loyalty Islands as determined by ²³⁰Th/²³⁴U dating of raised coral terraces, Quat. Res., 9, 186–192, 1978.

Matsumoto, E., Matsushima, Y., and Miyata, T.: Holocene sea-level studies of swampy coastal plains in Truk and
 Ponape, Micronesia, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1984 and 1985, edited by: Sugimura, Y., Kobe University Press, Kobe, Japan, 95–110, 1986.

Matsumoto, E., Matsushima, Y., Miyata, T., and Maeda, Y.: Holocene high sea-level stand on Kauai, Hawaii, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1986 and 1987, edited by: Yonekura, N., University of Tokyo, Japan, 91–99, 1988.

1755 Matsushima, Y., Sugimura, A., Berryman, K., Ishii, T., Maeda, Y., Matsumoto, E., and Yonekura, N.: Research report of B-party: Holocene sea-level changes in Fiji and Western Samoa, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1981, 1982 and 1983, edited by: Sugimura, A., Kobe University Press, Kobe, Japan, 137–185, 1984. Maurizot, P. and Lafoy, Y.: Notice explicative, Carte géol. Nouvelle-Calédonie (1 / 50 000), feuille Maré, Îles
Loyauté, Nouméa : Service des Mines et de l'Energie, Bureau de Recherches Géologiques et Minières, carte

géologique par P. Maurizot et al. (2003), 138 pp., 2003.

Mayer, A. G.: Coral reefs of Tutuila, with reference to the Murray-Agassiz solution theory, Proc. Natl. Acad. Sci. USA, 3, 522–526, 1917.

Mayor, A. G.: Rose Atoll, American Samoa, Proc. Am. Phil. Soc., 60, 62-70, 1921.

1765 Mayor, A. G.: Rose Atoll, American Samoa, papers from the Department of Marine Biology, Carnegie Institute of Washington, USA, 19, 73–91, 1924.

McLean, R. F. and Hosking, P. L.: Geomorphology of reef islands and atoll motu in Tuvalu, S. Pac. J. Nat. Sci., 11, 167–189, 1991.

McLean, R. F. and Woodroffe, C. D.: Coral atolls, in: Coastal evolution: Late Quaternary shoreline 1770 morphodynamics, edited by: Carter, R. W. G. and Woodroffe, C. D., Cambridge University Press, Cambridge,

UK, 267–302, 1994.
McMurtry, G. M., Campbell, J. F., Fryer, G. J., and Fietzke, J.: Uplift of Oahu, Hawaii, during the past 500 k.y. as recorded by elevated reef deposits, Geology, 38, 27–30, 2010.

McNutt, M. and Menard, H. W.: Lithospheric flexure and uplifted atolls, J. Geophys. Res., 83, 1206-1212, 1978.

Miklavič, B., Mylroie, J. E., Jenson, J. W., Randall, R. H., Banner, J. L., and Partin, J. W.: NSF Workshop: sealevel changes into the MIS 5: from observation to prediction, Palma de Mallorca, Spain, 10-14 April 2012.
 Mitchell, A. H. G.: Geology of South Malekula: Port Vila, Vanuatu, New Hebrides Condominium Geological Survey Report, 3, 41 pp., 1966.

Mitchell, A. H. G.: Raised reef-capped terraces and Plio-Pleistocene sea level changes, North Malekula, New 1780 Hebrides, J. Geol., 77, 56–67, 1969.

Mitchell, A. H. G.: Geology of Northern Malekula, New Hebrides Geological Survey Regional Report, 56 pp., 1971.

Miyata, T., Maeda, Y., Matsumoto, E., Matsushima, Y., Rodda, P., and Sugimura, A.: Emerged notches and microatolls on Vanua Levu, Fiji, in: in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC
Project in 1986 and 1987, edited by: Yonekura, N., University of Tokyo, Japan, 67–76, 1988.

Miyata, T., Maeda, Y., Matsumoto, E., Matsushima, Y., Rodda, P., Sugimura, A., and Kayanne, H.: Evidence for a Holocene high sea-level stand, Vanua Levu, Fiji, Quat. Res., 33, 352–359, 1990.

Mörner, N.-A. and Matlack-Klein, P.: New records of sea level changes in the Fiji Islands, Oceanogr. Fish. Open Access J., 5, doi:10.19080/OFOAJ.2017.05.555666, 2017.

1790 Montaggioni, L. F.: Makatea Island, Tuamotu Archipelago, Proc. 5th Int. Coral Reef Congr. 1, Tahiti, 27 May–1 June 1985, 103–158, 1985.

Montaggioni, L. F. and Camoin, G. F.: Geology of Makatea Island, Tuamotu Archipelago, French Polynesia, in: Geology and hydrogeology of carbonate islands, edited by: Vacher, H. L. and Quinn, T., Elsevier, Amsterdam, The Netherlands, 453–474, 1997.

- 1795 Montaggioni, L. F. and Pirazzoli, P. A.: The significance of exposed coral conglomerates from French Polynesia (Pacific Ocean) as indicators of recent relative sea level changes, Coral Reefs, 3, 29–42, 1984. Montaggioni, L. F., Richard, G., Bourrouilh-Le Jan, F., Gabrié, C., Humbert, L., Monteforte, M., Naim, O., Payri, C., and Salvat, B.: Geology and marine biology of Makatea, an uplifted atoll, Tuamotu Archipelago, Central Pacific Ocean, J. Coast. Res., 1, 165–171, 1985.
- 1800 Montaggioni, L. F., Cabioch, G., Thouveny, N., Frank, N., Sato, T., and Sémah, A. M.: Revisiting the Quaternary development history of the western New Caledonian shelf system: from ramp to barrier reef, Mar. Geol., 280, 57– 75, 2011.

Montaggioni, L. F., Salvat, B., Aubanel, A., Eisenhauer, A., and Martin-Garina, B.: The mode and timing of windward reef-island accretion in relation with Holocene sea-level change: a case study from Takapoto Atoll,
 French Polynesia, Geomorphology, 318, 320–335, 2018.

- Montaggioni, L. F., Collin, A., James, D., Salvat, B., Martin-Garin, B., Siu, G., Taiarui, M., and Chancerelle, Y.: Morphology of fore-reef slopes and terraces, Takapoto Atoll (Tuamotu Archipelago, French Polynesia, central Pacific): The tectonic, sea-level and coral-growth control, Mar. Geol., 417, https://doi.org/10.1016/j.margeo.2019.106027, 2019a.
- 1810 Montaggioni, L. F., Salvat, B., Aubanel, A., Pons-Branchu, E., Martin-Garin, B., Dapoigny, A., Goeldner-Gianella, L.: New insights into the Holocene development history of a Pacific, low-lying coral reef island: Takapoto Atoll, French Polynesia, Quat. Sci. Rev., 223, <u>https://doi.org/10.1016/j.quascirev.2019.105947</u>, 2019b. Moore, W. U.: Coral reefs, Nature, 40, 203–204, 1889.

Moriwaki, H., Chikamori, M., Okuno, M., and Nakamura, T.: Holocene changes in sea level and coastal 1815 environments on Rarotonga, Cook Islands, South Pacific Ocean, The Holocene, 16, 839–848, 2006.

Muhs, D. R. and Szabo, B. J.: New uranium-series ages of the Waimanalo Limestone, Oahu, Hawaii: implications for sea level during the last interglacial period, Mar. Geol., 118, 315–326, 1994.

Muhs, D. R., Simmons, K. R., and Steinke, B.: Timing and warmth of the Last Interglacial period: new U-series evidence from Hawaii and Bermuda and a new fossil compilation for North America, Quat. Sci. Rev., 21, 1355–

1820 1383, 2002.

Muhs, D. R., Schweig, E. S., and Simmons, K. R.: Late Quaternary sea-level history of Saipan, Commonwealth of the Northern Mariana Islands, USA: A test of tectonic uplift and glacial isostatic adjustment models, Geol. Soc. Am. Bull., 132, 863–883, 2020.

Neall, V. E. and Trewick, S. A.: The age and origin of the Pacific islands: a geological overview, Phil. Trans. R. 1825 Soc. B, 363, 3293–3308, 2008.

Neef, G. and McCulloch, M. T.: Pliocene-Quaternary history of Futuna Island, south Vanuatu, southwest Pacific, Aust. J. Earth Sci., 48, 805–814, 2001.

Neef, G. and Veeh, H. H.: Uranium series ages and late Quaternary uplift in the New Hebrides, Nature, 269, 682–683, 1977.

1830 Neef, G., Zhao, J. X., Collerson, K. D., and Zhang, F. S.: Late Quaternary uplift and subsidence of the west coast of Tanna, south Vanuatu, southwest Pacific: U-Th ages of raised coral reefs in the Median Sedimentary Basin, Aust. J. Earth Sci., 50, 39–48, 2003.

Newell, N. D. and Bloom, A. L.: The reef flat and 'two-meter eustatic terrace' of some Pacific atolls, GSA Bull., 81, 1881–1894, 1970.

1835 Nunn, P. D.: Studies in the tectonics and structure of southern Fiji, Working Paper 10, School of Social and Economic Development, University of the South Pacific, Suva, Fiji, 54 pp., 1988.

Nunn, P. D.: Coastal processes and landforms of Fiji: their bearing on Holocene sea-level changes in the south and west Pacific, J. Coast. Res., 6, 279–310, 1990a.

Nunn, P. D.: Coastal geomorphology of Beca and Yanuca Islands, South Pacific Ocean, and its significance for 1840 the tectonic history of the Vatulele-Beqa Ridge, Pac. Sci., 44, 348–365, 1990b.

Nunn, P. D.: Sea-level changes during the last 6000 years in Fiji, Tonga and Western Samoa: implications for future coastline development, workshop on coastal processes in the South Pacific Island Nations, Lae, Papua New Guinea, 1-8 October 1987, SOPAC Tech. Bull., 7, 79–90, 1991.

Nunn, P. D.: Oceanic islands, Blackwell, Oxford, UK, 413 pp., 1994.

1845 Nunn, P. D.: Holocene sea-level changes in the South and West Pacific, J. Coast. Res. Spec. Issue, 17, 311–319, 1995a.

Nunn, P. D.: Holocene tectonic histories for five islands in the south-central Lau Group, South Pacific, The Holocene, 5, 160–171, 1995b.

Nunn, P. D.: Significance of emerged Holocene corals around Ovalau and Moturiki islands, Fiji, southwest Pacific,
 Mar. Geol., 163, 345–351, 2000.

Nunn, P. D.: Climate, environment and society in the Pacific during the last millennium, in: Developments in Earth and Environmental Sciences, 6, Elsevier, 302 pp., 2007.

Nunn, P. D. and Britton, J. M. R.: The long-term evolution of Niue Island, in: Niue Island: geographic perspectives on the rock of Polynesia, edited by: Terry, J. P. and Murray, W. E., Insula, 31–73, 2004.

1855 Nunn, P. D and Finau, F. T.: Holocene emergence history of Tongatapu island, South Pacific, Z. Geomorphol., 39, 69–95, 1995.

Nunn, P. D. and Omura, A.: Penultimate Interglacial emerged reef around Kadavu Island, Southwest Pacific: implications for late Quaternary island-arc tectonics and sea-level history, New Zeal. J. Geol. Geop., 42, 219–227, 1999.

- 1860 Nunn, P. D. and Omura, A.: Quaternary shorelines of Kaibu Island, southwest Pacific Ocean: implications for Last Interglacial sea-level history and uplift of the Lau-Colville Ridge, Quat. Australasia, 21, 33–38, 2003. Nunn, P. D. and Peltier, W. R.: Far-field test of the ICE- 4G model of global isostatic response to deglaciation using empirical and theoretical Holocene sea-level reconstructions for the Fiji Islands, southwestern Pacific, Quate. Res., 55, 203–214, 2001.
- 1865 Nunn, P. D., Matararaba, S., and Ramos, J.: Investigations of anthropogenic sediments in Qaranilaca, Vanuabalavu Island, Fiji, Archaeol. New Zeal., 43, 73–79, 2000.

Nunn, P. D., Ollier, C., Hope, G., Rodda, P., Omura, A., and Peltier, W. R.: Late Quaternary sea-level and tectonic changes in northeast Fiji, Mar. Geol., 187, 299–311, 2002.

Nunn, P. D., Hunter-Anderson, R., Carson, M. T., Thomas, F., Ulm, S., and Rowland, M. J.: Times of plenty,times of less: Last-millennium societal disruption in the Pacific Basin, Hum. Ecol., 35, 385–401, 2007.

- Ohde, S., Greaves, M., Masuzawa, T., Buckley, H. A., VanWoesik, R., Wilson, P. A., Pirazzoli, P. A., and Elderfield, H.: The chronology of Funafuti atoll: revisiting an old friend, Proc. R. Soc. A, 458, 2289–2306, 2002.
 Omura, A., Chappell, J., Bloom, A. L., Pillans, B. J., McCulloch, M., Esat, T., Sasaki, K., and Kawada, Y.: Alphaspectrometric ²³⁰Th/²³⁴U dating of Pleistocene corals, in: Study on Coral Reef Terraces of the Huon Peninsula,
- 1875 Papua New Guinea, edited by: Ota, Y., Department of Geography, Senshu University, Kawasaki, Japan, 95–110, 1994.

Ota, Y., Chappell, J., Kelley, R., Yonekura, N., Matsumoto, E., Nishimura, T., and Head, J.: Holocene coral reef terraces and coseismic uplift of Huon Peninsula, Papua New Guinea, Quat. Res., 40, 177–188, 1993.

Paulay, G.: Effects of glacio-eustatic sea level fluctuations and local tectonics on the marine faunas of oceanic islands, Ph.D. thesis, University of Washington, Seattle, USA, 287 pp., 1988. Paulay, G. and McEdward, L. R.: A simulation model of island reef morphology: the effects of sea level fluctuations, growth, subsidence and erosion, Coral Reefs, 9, 51–62, 1990.

Paulay, G. and Spencer T.: Niue Island: geologic and faunatic history of a Pliocene atoll, Pac. Sci. Assoc. Infor. Bull., 44, 21–23, 1992.

1885 Pirazzoli, P. A.: A reconnaissance and geomorphological survey of Temoe atoll, Gambier Islands (South Pacific), J. Coast. Res., 3, 307–323, 1987.

Pirazzoli, P. A. and Montaggioni, L. F.: Late Holocene sea level changes in the northwest Tuamotu Islands, French Polynesia, Quat. Res., 25, 350–368, 1986.

Pirazzoli, P. A. and Montaggioni, L. F.: Les Îles Gambier et l'atoll de Temoe (Polynésie française): anciennes
lignes de rivage et comportement geodynamique, Géodynamique, 2, 13–28, 1987.

Pirazzoli, P. A. and Montaggioni, L. F.: Holocene sea-level changes in French Polynesia, Palaeogeogr., Palaeocclimatol. Palaeoecol., 68, 153–175, 1988.

Pirazzoli, P. A., Montaggioni, L. F., Delibrias, G., Faure, G., and Salvat, B.: Late Holocene sea-level changes in the Society Islands and in the northwest Tuamotu atolls, Proc. 5th Int. Coral Reef Congr. 3, Tahiti, 27 May–1 June 1985, 131–136, 1985a.

- Pirazzoli, P. A., Brousse, R., Delibrias, G., Montaggioni, L. F., Sachet, M. H., Salvat, B., and Sinoto, Y. H.: Îles Sous le Vent: Maupiti, Tupai, Bora Bora, Huahine, Archipel de la Société, in: Les récifs coralliens de Polynésie Française: connaissance des récifs et guides d'excursion, Proc. 5th Int. Coral Reef Congr., Tahiti, 27 May–1 June 1985, 19–72, 1985b.
- 1900 Pirazzoli, P. A., Montaggioni, L. F., Vergnaud-Grazzini, C., and Saliege, J. F.: Late Holocene sea levels and coral reef development in Vahitahi atoll, eastern Tuamotu Islands, Pacific Ocean, Mar. Geol., 76, 105–115, 1987. Pirazzoli, P. A., Koba, M., Montaggioni, L. F., and Person, A.: Anaa (Tuamotu Islands, central Pacific): an incipient rising atoll?, Mar. Geol., 82, 261–269, 1988a.

Pirazzoli, P. A., Montaggioni, L. F., Salvat, B., and Faure, G.: Late Holocene sea level indicators from twelve atolls in the central and eastern Tuamotus (Pacific Ocean), Coral Reefs, 7, 57–68, 1988b.

Potratz, H. A., Barnes, J. W., and Lang, E. J.: A radiochemical procedure for thorium and its application to the determination of ionium in coral limestone. U.S. Atomic Energy Commission, Los Alamos Scientific Laboratory, University of California, USA, 13 pp., 1955.

Quinn, T. M.: The history of post-Miocene sea level change: inferences from stratigraphic modeling of Enewetak
 Atoll, J. Geophys. Res., 96, 6713–6725, 1991.

Quinn, T. M. and Matthews, R. K.: Post-Miocene diagenetic and eustatic history of Enewetak Atoll: model and data comparison, Geology, 18, 942–945, 1990.

Randall, R. H. and Siegrist, H. G.: Geomorphology of the fringing reefs of northern Guam in response to Holocene sea level changes, in: Proceedings of the 6th International Coral Reef Symposium: Townsville, Australia, edited

1915 by: Choat, J. H., Barnes, D., Borowitzka, M. A., Coll, J. C., Davies, P. J., Flood, P., Hatcher, B. G., Hopley, D., Hutchings, P. A., Kinsey, D., Orme, G. R., Pichon, M., Sale, P. F., Sammarco, P., Wallace, C. C., Wilkinson, C., Wolomski, E., and Bellwood, O., 3, 473–477, 1988.

Randall, R. H_vand Siegrist, H. G.: The legacy of Tarague Embayment and its inhabitants, Anderson AFB, Guam, volume III, Geology, beaches, and coral reefs, prepared for Andersen Air Force Base, International Archaeology, 1920 Inc., Honolulu, USA, 462 pp., 1996.

Rashid, R., Eisenhauer, A., Stocchi, P., Liebetrau, V., Fietzke, J., Rüggeberg, A., and Dullo, W.-C.: Constraining mid to late Holocene relative sea level change in the southern equatorial Pacific Ocean relative to the Society Islands, French Polynesia, Geochem. Geophys. Geosyst., 15, doi:10.1002/2014GC005272, 2014.

Riker-Coleman, K. E., Gallup, C. D., Wallace, L. M., Webster, J. M., Cheng, H., and Edwards, R. L.: Evidence of
 Holocene uplift in east New Britain, Papua New Guinea, Geophys. Res. Lett., 33, doi:10.1029/2006GL026596,

2006.

Robinson, G. P.: The geology of north Santo, New Hebrides Geological Survey Regional Report, 77 pp., 1969. Rodda, P.: Late Quaternary geology of Fiji, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1984 and 1985, edited by: Sugimura, Y., Kobe University Press, Kobe, Japan, 15–24, 1986.

1930 Rodda, P.: Visit to Western Samoa with the HIPAC Team, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1986 and 1987, edited by: Yonekura, N., University of Tokyo, Japan, 85– 90, 1988.

Rovere, A., Raymo, M. E., Vacchi, M., Lorscheid, T., Stocchi, P., Gomez-Pujol, L., Harris, D. L., Casella, E., O'Leary, M. J., and Hearty, P. J.: The analysis of Last Interglacial (MIS 5e) relative sea-level indicators:
1935 reconstructing sea-level in a warmer world, Earth-Sci. Rev., 159, 404–427, 2016.

Rovere, A., Deirdre, R., Murray-Wallace, C., Simms, A., Vacchi, M., Dutton, A., Lorscheid, T., Chutcharavan, P., Brill, D., Bartz, M., Jankowski, N., Mueller, D., Cohen, K., and Gowan, E.: Descriptions of database fields for the World Atlas of Last Interglacial Shorelines (WALIS), doi:10.5281/zenodo.3961544, 2020.

Roy, P.: Coastal and nearshore geology of Savusavu Bay, Vanua Levu, Fiji, CCOP/SOPAC, Suva, Fiji, Techn.1940 Rep., 60, 33 pp., 1988.

hat gelöscht:

Rubin, K. H., Sherman, C. E., and Fletcher III, C. H.: Ages of emerged coral deposits in Kapihaa Gulch, Lana'i, Hawaiian Islands and speculation about their environment of deposition, Trans. Am. Geophys. Union, 76, F307, 1995.

1945 Rubin, K. H., Fletcher III, C. H., and Sherman, C.: Fossiliferous Lana'i deposits formed by multiple events rather than a single giant tsunami, Nature, 408, 675–681, 2000.

Schlanger, S. O.: Subsurface Geology of Eniwetok Atoll, U.S. Geological Survey Professional Paper 260-BB, 991–1048, 1963.

Schofield, J. C.: Notes on late Quaternary sea levels, Fiji and Rarotonga, New Zeal. J. Geol. Geop., 13, 199–206, 1950 1910.

Schoffeld, J. C.: The geology and hydrology of Niue Island, South Pacific, New Zeal. Geol. Surv. Bull., 62, 27 pp., 1959.

Schofield, J. C.: Notes on late Quaternary sea levels, Fiji and Rarotonga, New Zeal. J. Geol. Geop., 14, 240–241, 1970.

1955 Schofield, J. C.: Late Holocene sea level, Gilbert and Ellice Islands, west central Pacific Ocean, New Zeal. J. Geol. Geop., 20, 503–529, 1977a.

Schofield, J. C.: Effect of Late Holocene sea-level fall on atoll development, New Zeal. J. Geol. Geop., 20, 531–536, 1977b.

Scoffin, T. P., Stoddart, D. R., Tudhope, A. W., and Woodroffe, C. D.: Exposed limestones of Suwarrow Atoll,
 Proc. 5th Int. Coral Reef Congr. 3, Tahiti, 27 May–1 June 1985, 137–140, 1985.

Shepard, F. P.: Thirty-five thousand years of sea level, in: Essays in marine geology in honor of K. O. Emery, edited by: Clements, T., University of Southern California Press, Los Angeles, USA, 1–10, 1963.
Shepard, F. P.: The evolution of a moderate energy coast in Holocene time, Pacific Harbour, Viti Levu, Fiji, New Zeal. J. Geol. Geop., 33, 547–556, 1990.

- Shepard, F. P., Curray, J. R., Newman, W. A., Bloom, A. L., Newell, N. D., Tracey, J. I., Jr., and Veeh, H. H.: Holocene changes in sea level: evidence in Micronesia, Science, 157, 542–544, 1967.
 Sherman, C. E., Glenn, C. R., Jones, A. T., Burnett, W. C., and Schwarcz, H. P.: New evidence for two highstands of the sea during the Last Interglacial, oxygen isotope stage 5e, Geology, 21, 1079–1082, 1993.
 Sherman, C. E., Fletcher, C. H., and Rubin, K. H.: Marine and meteoric diagenesis of Pleistocene carbonates from
- 1970 a nearshore submarine terrace, Oahu, Hawaii, J. Sediment. Res., 69, 1083-1097, 1999.

Sherman, C. E., Fletcher, C. H., Rubin, K. H., Simmons, K. R., and Adey, W. H.: Sea-level and reef accretion history of Marine Oxygen Isotope Stage 7 and late Stage 5 based on age and facies of submerged late Pleistocene reefs, Oahu, Hawaii, Quat. Res., 81, 138–150, 2014.

Spencer, T. and Paulay, G.: Geomorphology, uplift and sea level history of Niue Island, southwest Polynesia, Proc.7th Int. Coral Reef Symp. 2, Guam, 22-27 June 1992, 1227, 1994.

Spencer, T., Stoddart, D. R., and Woodroffe, C. D.: Island uplift and lithospheric flexure; observations and cautions from the South Pacific, Z. Geomorphol., 63, 87–102, 1987.

Spencer, T., Stoddart, D. R., Woodroffe, C. D., and Harmon, R. S.: Lithospheric flexure and raised reef limestones, S. Cooks-Austral Islands, Proc. 6th Int. Coral Reef Congr. 3, Townsville, 8–12 August 1988, 485–489, 1988.

- 1980 Spencer, T., Stoddart, D. R., McLean, R. F.: Coral reefs, in: The history of the study of landforms volume 4: Quaternary and recent processes and forms (1890-1965) and the mid-century revolutions, edited by: Burt, T. P., Chorley, R. J., Brunsden, D., Cox, N. J., and Goudie, A. S., The Geological Society, London, UK, 863–922, 2008. Spenneman, D. H. R.: A Holocene sea-level history for Tongatapu, Kingdom of Tonga, in: Coastal and environmental Geoscience studies of the Southwest Pacific Islands, Suva: South Pacific, edited by: Howorth, R.
- and Rodda, P., Appl. Geosc. Comm. Techn. Bull., 9, 115–149, 1997.
 Spriggs, M.: The Island Melanesians, Wiley-Blackwell, Oxford, UK, 320 pp., 1997.
 Stearns, H. T.: Pleistocene shorelines on the islands of Oahu and Maui, Hawaii, Geol. Soc. Am. Bull., 46, 1927–1956, 1935a.

Stearns, H. T.: Shore benches on the island of Oahu, Hawaii, Geol. Soc. Am. Bull., 46, 1467-1482, 1935b.

1990 Stearns, H. T.: Geology and ground-water resources of the island of Oahu, Hawaii, Hawaii Div. Hydrogr. Bull., 1, 479 pp., 1935c.

Stearns, H. T.: Geologic history of Guam, Geol. Soc. Am. Bull., 52, 194, 1940.

Stearns, H. T.: Shore benches on North Pacific Islands, Geol. Soc. Am. Bull., 52, 773-780, 1941.

Stearns, H. T.: Geology of the Samoan Islands, Geol. Soc. Am. Bull., 55, 1279-1332, 1944.

1995 Stearns, H. T.: Eustatic shorelines in the Pacific, Geol. Soc. Am. Bull., 56, 1071–1078, 1945.

Stearns, H. T.: Submerged shorelines and shelves in the Hawaiian Islands and a revision of some of the eustatic emerged shorelines, Geol. Soc. Am. Bull., 85, 795–804, 1974.

Stearns, H. T. and Chamberlain, T. K.: Deep cores of Oahu, Hawaii and their bearing on the geologic history of the Central Pacific Basin, Pac. Sci., 11, 153–165, 1967.

- 2000 Stein, M., Wasserburg, G. J., Aharon, P., Chen, J. H., Zhu, Z. R., Bloom, A., and Chappell, J.: TIMS U-series dating and stable isotopes of the last interglacial event in Papua New Guinea, Geochim. Cosmochim. Acta, 57, 2541–2554, 1993.
 - Stice, G. D. and McCoy, F. W.: The geology of the Manu'a Islands, Samoa, Pac. Sci., 22, 427-457, 1968.
- Stoddart, D. R.: Marine biology, geomorphology of the Solomon Islands coral reefs, Phil. Trans. R. Soc. B, 255, 2005 355–382, 1969.
 - Stoddart, D. R.: Reef islands of Rarotonga, Atoll Res. Bull., 160, 1-7, 1972.

Stoddart, D. R.: Almost-atoll of Aitutaki: geomorphology of reefs and islands, in: Almost-Atoll of Aitutaki: Reef Studies in the Cook Islands, South Pacific, edited by: Stoddart, D. R. and Gibbs, P. E., Smithsonian Institution, Washington, USA, Atoll Research Bulletin, 190, 31–58, 1975.

2010 Stoddart, D. R., Spencer, T., and Scoffin, T. P.: Reef growth and karst erosion on Mangaia, Cook Islands: a reinterpretation in geomorphology of changing coastlines, Z. Geomorphol., 57, 121–140, 1985. Stoddart, D. R., Woodroffe, C. D., and Spencer, T.: Mauke, Mitiaro and Atiu: geomorphology of Makatea Islands in the southern Cooks, Atoll Res. Bull., 341, 1–65, 1990.

Sugimura, A., Maeda, Y., Matsushima, Y., and Rodda, P.: Further report on sea-level investigation in Western

2015 Samoa, in: Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1986 and 1987, edited by: Yonekura, N., University of Tokyo, Japan, 77–84, 1988a. Sugimura, A., Maeda, Y., Matsushima, Y., Rodda, P., and Matsumoto, E.: Lobau Lowland, Viti Levu, Fiji, in:

Sea-level changes and tectonics in the middle Pacific, report of the HIPAC Project in 1986 and 1987, edited by: Yonekura, N., University of Tokyo, Japan, 59–65, 1988b.

2020 Szabo, B. J., Tracey, J., and Goter, E.: Ages of subsurface stratigraphic intervals in the Quaternary of Enewetak Atoll, Marshall Islands, Quat. Res., 23, 54–61, 1985.

Szabo, B. J., Ludwig, K. R., Muhs, D. R., and Simmons, K. R.: Thorium-230 ages of corals and the duration of the last interglacial sea-level highstand on Oʻahu, Hawaii, Science, 266, 93–96, 1994.

Tayama, R.: Coral reefs in the South Seas, Bull. Hydrogr., 11, 1–292, 1952.

2025 Taylor, F. W.: Quaternary tectonic and sea-level history, Tonga and Fiji, Southwest Pacific, Ph.D. thesis, Cornell University, Ithaca, New York, USA, 355 pp., 1978.

Taylor, F. W. and Bloom, A. L.: Coral Reefs in tectonic blocks, Tonga Island Arc, Proc. 3rd Int. Coral Reef Symp., Miami, 275–281, 1977.

Taylor, F. W., Isacks, B. L., Jouannic, C., Bloom, A. L., Dubois, J.: Coseismic and Quaternary vertical tectonic
 movements, Santo and Malekula Islands, New Hebrides Island Arc, J. Geophys. Res., 85, 5367–5381, 1980.

Taylor, F. W., Jouannic, C., Bloom, A. L.: Quaternary uplift of the Torres Islands, Northern New Hebrides frontal arc: comparison with Santo and Malekula Islands, central New Hebrides frontal arc, Geology, 93, 419–438, 1985.
 Taylor, F. W., Frohlich, C., Lecolle, J., and Strecker, M.: Analysis of partially emerged corals and reef terraces in the central Vanuatu arc: comparison of contemporary coseismic and nonseismic with Quaternary vertical movements, J. Geophys. Res., 92, 4905–4933, 1987.

Taylor, F. W., Mann, P., Bevis, M. G., Edwards, R. L., Cheng, H., Cutler, K. B., Gray, S. C., Burr, G. S., Beck, J.
W., Phillips, D. A., Cabioch, G., and Recy, J.: Rapid forearc uplift and subsidence caused by impinging bathymetric features: examples from the New Hebrides and Solomon arcs, Tectonics, 24, TC6005, doi:10.1029/2004TC001650, 2005.

2040 Terry, P. and Nunn, P. D.: Interpreting features of carbonate geomorphology on Niue Island, a raised coral atoll, Z. Geomorph., 131, 43–57, 2003.

Thirumalai, K., Taylor, F. W., Shen, C.-C., Lavier, L. L., Frohlich, C., Wallace, L. M., Wu, C.-C., Sun, H., and Papabatu, A. K.: Variable Holocene deformation above a shallow subduction zone extremely close to the trench, Nat. Commun., 6, 7607, <u>https://doi.org/10.1038/ncomms8607</u>, 2015.

2045 Thomas, F. R., Nunn, P. D., Osborne, T., Kumar, R., Areki, F., Matararaba, S., Steadman, D., and Hope, G.: Recent archaeological findings at Qaranilaca Cave, Vanuabalavu Island, Fiji, Archaeol. Oceania, 39, 42–49, 2004. Thompson, W. G., Curran, H. A., Wilson, M. A., and White, B.: Sea-level oscillations during the last interglacial highstand recorded by Bahamas corals, Nat. Geosci., 4, 684–687, 2011. Thurber, D. L., Broecker, W. S., Blanchard, R. L., and Potratz, H. A.: Uranium-series ages of Pacific atoll coral,

2050 Science, 149, 55–58, 1965.

Tracey, J. I.: Recent reef features in the Caroline and Marshall Islands, Geol. Soc. Am. Spec. Pap., 115, 223, 1968. Tracey, J. I.: Holocene emergent reefs in the central Pacific, 2nd Conf. Am. Quat. Assoc., 51–52, 1972.

Tracey, J. I. and Ladd, H. S.: Quaternary history of Eniwetok and Bikini atolls, Marshall Islands, Proc. 2nd Int. Coral Reef Symp. 2, Great Barrier Reef Province, 22 June–2 July 1973, 537–550, 1974.

2055 Tracey, J. I., Stensland, C. H., Doan, D. B., May, H. G., Schlanger, S. O., Stark, J. T., Blumenstock, D. I., Emery, K. O., Fosberg, F. R.: Military geology of Guam, Mariana Islands, part I, Description of terrain and environment, part II, Engineering aspects of geology and soils, U.S. Army, Chief of Engineers, Washington D.C., USA, 282 pp., 1959.

Tracey, J. I., Schlanger, S. O., Stark, J. T., Doan, D. B., and May, H. G.: General geology of Guam: geology andhydrology of Guam, Mariana Islands, U.S. Geological Survey Professional Paper 403-A, 104 pp., 1964.

Trichet, J., Repellin, P., and Oustriere, P.: Stratigraphy and subsidence of the Mururoa Atoll (French Polynesia), Mar. Geol., 56, 241–257, 1984.

Veeh, H. H.: Th²³⁰/U²³⁸ and U²³⁴/U²³⁸ ages of elevated Pleistocene coral reefs and their geological implications, Ph.D. thesis, University of California, San Diego, USA, 182 pp., 1965.

2065 Veeh, H. H.: Th²³⁰/U²³⁸ and U²³⁴/U²³⁸ ages of Pleistocene high sea level stand, J. Geophys. Res., 71, 3379–3386, 1966.

Veeh, H. H. and Chappell, J.: Astronomical theory of climate change: support from New Guinea, Science, 167, 862–865, 1970.

Wallace, C. C.: Staghorn corals of the world: a revision of the coral genus Acropora (Scleractinia; Astrocoeniina;

2070 Acroporidae) worldwide, with emphasis on morphology, phylogeny and biogeography, CSIRO Publishing, 18, 422 pp., 1999.

Wardlaw, B. R. and Quinn, T. M.: The record of Pliocene sea-level change at Enewetak Atoll, Quat. Sci. Rev., 10, 247–258, 1991.

Weary, D. J. and Burton, W. C.: Preliminary Geologic Map of the Island of Saipan, Commonwealth of the Northern
 Mariana Islands: U.S. Geological Survey Open-File Report 2011-1234, scale 1:25,000, 2011.

Webster, J. M., Clague, D. A., and Braga J. C.: Support for the giant wave hypothesis: evidence from submerged terraces off Lanai, Hawaii, Int. J. Earth Sci., 96, 517–524, 2007.

Wells, J. W.: Fossil corals from Midway Atoll, U.S. Geological Survey Professional Paper 680-G, 19 pp., 1982.

Wentworth, C. K. and Hoffmeister, J. E.: Geology of Ulupau Head, Oahu, Geol. Soc. Am. Bull., 50, 1553–1572, 2080 1939.

Wheeler, C. W.: Stratigraphy, petrology, and geochemistry of a mid-oceanic carbonate platform: Niue Island, South Pacific, Ph.D. thesis, Luisiana State University Historical Dissertations and Theses, USA, 7397, 461 pp., 2000.

Wiens, H. J.: Atoll development and ecology, Yale University Press, New Haven, USA, 532 pp., 1962.

2085 Wirrmann, D., Sémah, A.-M., Debenay, J., and Chacornac-Rault, M.: Mid- to Late Holocene environmental and climatic changes in New Caledonia, Southwest tropical Pacific, inferred from the littoral plain Gouro-Déva, Quat. Res., 76, 229–242, 2011.

Woodroffe, S. A. and Horton, B. P.: Holocene sea-level changes in the Indo-Pacific, J. Asian Earth Sci., 25, 29– 43, 2005.

2090 Woodroffe, C. D. and McLean, R. F.: Pleistocene morphology and Holocene emergence of Christmas (Kiritimati) Island, Pacific Ocean, Coral Reefs, 17, 235–248, 1998. Woodroffe, C. D. and Morrison, R. J.: Reef-island accretion and soil development on Makin, Kiribati, central Pacific, Catena, 44, 245–261, 2001.

Woodroffe, C. D., Stoddart, D. R., Spencer, T., Scoffin, T. P., and Tudhope, A. W.: Holocene emergence in the 2095 Cook Islands, South Pacific, Coral Reefs, 9, 31–39, 1990.

Woodroffe, C. D., Short, S. A., Stoddart, D. R., Spencer, T., Harmon, R. S.: Stratigraphy and chronology of late pleistocene reefs in the Southern Cook Islands, south Pacific, Quat. Res., 35, 246–263, 1991.

Woodroffe, C. D., McGregor, H. V., Lambeck, K., Smithers, S. G., and Fink, D.: Mid-Pacific microatolls record sea-level stability over the past 5000 yr, Geology, 40, 951–954, 2012.

2100 Yamano, H., Kayanne, H., Yamaguchi, T., Inoue, T., Mochida, Y., and Baba, S.: Revisiting late Holocene sealevel change from the Gilbert Islands, Kiribati, west-central Pacific Ocean, Quat. Res., 88, 400–408, 2017. Yonekura, N.: Late Quaternary vertical crustal movements in and around the Pacific as deduced from former shoreline data, in: Geodynamics of the western Pacific-Indonesian region, edited by: Hilde, T. W. and Uyeda, S., Geodynamics Series, American Geophysical Union, USA, 41–50, 1983.

2105 Yonekura, N.: Geomorphic development of modern coral reefs and Holocene sea-level changes in the middle Pacific, J. Geogr., 103, 841–852, 1994.

Yonekura, N., Ishii, T., Saito, Y., Maeda, Y., Matsushima, Y., Matsumoto, E., and Kayanne, H.: Holocene fringing reefs and sea-level change in Mangaia Island, Southern Cook Islands, Palaeogeogr., Palaeoclimatol. Palaeoecol., 68, 171–188, 1988.

2110