



# Historical *K* index data collection of Soviet magnetic observatories, 1957–1992

Natalia Sergeyeva<sup>1</sup>, Alexei Gvishiani<sup>1,2</sup>, Anatoly Soloviev<sup>1,2</sup>, Lyudmila Zabarinskaya<sup>1</sup>, Tamara Krylova<sup>1</sup>, Mikhail Nisilevich<sup>1</sup>, Roman Krasnoperov<sup>1</sup>

5 <sup>1</sup>Geophysical Center of the Russian Academy of Sciences, Moscow, 119296, Russia

<sup>2</sup>Schmidt Institute of Physics of the Earth of the Russian Academy of Sciences, Moscow, 123242, Russia

*Correspondence to:* Roman Krasnoperov (r.krasnoperov@gcras.ru)

**Abstract.** *K* index is one of the oldest universal indices of geomagnetic activity, introduced in 1938 by Julius Bartels, that is still being widely used. Up to the present day, long-term timeseries of homogeneous *K* index records have been accumulated 10 at data repositories all over the world. The multidecadal practice of its application makes it an indispensable source of information for retrospective analysis of solar-terrestrial interaction for nearly eight Solar cycles. Most significantly, while studying the historical geomagnetic data, *K* index datasheets are in certain cases far easier for automated analysis than the conventional analogue magnetograms. The presented collection includes the results of the *K* index determination at 41 geomagnetic observatories of the former USSR for the period from July 1957 to early 1990s. This unique collection was 15 formed at the World Data Center for Solar-Terrestrial Physics in Moscow. The historical data, which are offered to the international scientific community, cover the second half of the 20th century and can be used for retrospective analysis and study of geomagnetic events in the past as well as for data validation or forecasting (Sergeyeva et al., 2020). The dataset is available at: <https://doi.org/10.1594/PANGAEA.922233>, last access: 16 September 2020.

## 1 Introduction

20 Preservation and providing access to historical observational data is an important and relevant problem in the Earth sciences. The World Data Center for Solar-Terrestrial Physics at the Geophysical Center of the Russian Academy of Sciences (GC RAS) consistently works on the organization of online open access to historical geomagnetic data stored in its archive. This paper presents the collection of the geomagnetic activity parameter – *K* index – obtained within the network of magnetic observatories and stations of the former USSR.

25 The most valuable part in this collection is the data of 1957–1959 related to the observation program of the International Geophysical Year (IGY), which was organized from July 1957 to December 1958, and the International Geophysical Cooperation of 1959 – the direct continuation of the IGY (Lyubovtseva et al., 2020; Nicolet, 1984).

To provide collecting, archiving and disseminating the data of all types of observations obtained within the IGY program the universal World Data Centers (WDC) were established in the USA – WDC A, and the USSR – WDC B (consisted of two 30 centers, B1 and B2), which received materials from the stations all over the world. The results of geomagnetic observations,



studies of the ionosphere, solar observations, data on cosmic rays and auroras were collected at WDC B2. It was established in 1956 at the Research Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation of the USSR Ministry of Communications (NIZMIR), which in 1959 was reassigned to the USSR Academy of Sciences (AS USSR) and renamed as the Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation of the AS USSR, widely known as IZMIRAN.

35 Later, in 1971 WDC B2 was reassigned to the Interdepartmental Geophysical Committee of the AS USSR (currently GC RAS) and designated as the World Data Center for Solar-Terrestrial Physics (WDC for STP). Currently WDC for STP is the part of the World Data System (WDS) (Raspovov et al., 2007; Rodnikov et al., 2009).

In the Soviet Union, more than 30 observatories were involved in geomagnetic research within the IGY program. The WDC for STP preserves photocopies and digital images of magnetograms, hourly mean and minute values of the Earth's magnetic field vector components, values of indices of geomagnetic activity, data on magnetic storms and other information obtained at these observatories.

40 It is difficult to overestimate the importance of data from high latitude observatories established in the Arctic at various times. Among others worth to mention: Tikhaya Bay on Franz Josef Land (established in 1929 during the development of the Northern Sea Route); Dixon, Cape Wellen (Uelen) and Cape Chelyuskin (established in 1931–1933 within the Second International 45 Polar Year program); Heiss Island and Tixie Bay (established in 1957 within the IGY program). Magnetic observations were performed at Antarctic stations Mirny, Pionerskaya, Oasis and Vostok that were established during the First and Second Soviet Complex Antarctic Expeditions in 1956–1957. At the same period, magnetic observations were made at the North Pole (romanized ‘Severny Polus’) drifting stations SP-6, SP-7, and SP-8. This data was used for daily calculation of the C index (Lyubovtseva et al., 2020).

50 At that time, the Soviet geomagnetic observatories were equipped mainly with foreign-manufactured instruments for registration of geomagnetic variations. For example, in 1957–1959 the observatories Irkutsk, Kiev, Lvov, Tbilisi, Tashkent, Yakutsk, Vladivostok, and Mirny were equipped with the Eschenhagen variation stations and La Cour variometers. The Edelman magnetometers and Lloyd scales were installed at the Kazan, Sverdlovsk, and Odessa observatories. The Tepfer variometers were installed at the Leningrad and Vladivostok observatories. The Schultz variometer was used at the Yuzhno-

55 Sakhalinsk observatory. The Schmidt scales were used at the Srednikan observatory. Some observatories used domestically manufactured instruments: the Yanovsky variometer in Srednikan, the Brunelli magnetovariational station in Ashkhabad. The polar observatories in the Arctic and Antarctic (including the North Pole drifting stations) were equipped with Soviet-manufactured Brunelli magnetovariational stations and La Cour variometers. The accuracy of these instruments was low due to the influence of temperature, humidity, and other factors (Belov et al., 2006).

60 In the mid-1960s the re-equipment of observatories (starting with Moscow, Odessa, and Tbilisi observatories in 1964) with the new Bobrov quartz variometers began (Bobrov, 1961; Belov et al., 2006). Automatic quartz magnetovariational stations were developed in IZMIRAN for polar geomagnetic stations that conducted research in severe climatic conditions in the Arctic and Antarctic (Burtsev et al., 1977). By the mid-1980s, the network of geomagnetic observatories in the USSR was equipped with digital quartz magnetovariational stations (Belov et al., 2006).



65 New geomagnetic observatories that were deployed in the USSR in the 1960s–1970s participated in major international projects: the International Year of the Quiet Sun (1964–1965), the International Year of the Active Sun (1969–1971), the International Magnetospheric Study (1976–1979). The observatories continued transferring data to the WDC system (Raspopov et al., 2007).

With the collapse of the USSR in December 1991 operation of a number of observatories was terminated. Some of them  
70 continued to transmit data to the WDC for STP until 1992.

To estimate the geomagnetic disturbances caused by extraterrestrial sources quantitatively, the indices of geomagnetic activity are widely used by the geomagnetic community (Lincoln, 1967). The majority of the indices are derived from the measurements carried out at magnetic stations and observatories (Mayaud, 1980) (Fig. 1). The International Service of  
75 Geomagnetic Indices (<http://isgi.unistra.fr/>, last access: 30 September 2020) handles and disseminates indices that are officially accepted by the International Association of Geomagnetism and Aeronomy (IAGA). They are divided into regional, planetary and source driven indices and express responses from various magnetospheric-ionospheric current systems. The first and the simplest *C* index was introduced in 1906 to estimate magnetic field activity over a day on a three-point scale (0, 1 and 2). Since that time, the index evolved into *K* index (Bartels, 1938), which is still widely used nowadays (Gvishiani and Soloviev, 2020). During the period from 1957 to 1971, all observatories and drifting stations SP-6, SP-7 and SP-8 determined *C* index and  
80 transferred it to the WDCs. Since July 1957, data on geomagnetic *K* and *C* indices were transferred to WDC B2 by 22 observatories, and since January 1958 – by 26 observatories (Fig. 2).

Among them are the oldest observatories, established in the Russian Empire in the 19th century: Kazan, St. Petersburg (Leningrad – Voeikovo), Yekaterinburg (Sverdlovsk), Tiflis (Tbilisi), Irkutsk; and observatories established in the Soviet Union in the 20th century before the IGY (in 1930s–1940s): Yakutsk, Vladivostok, Kiev, Odessa, Lvov, Moscow, Tashkent,  
85 Srednikan, Tomsk, Yuzhno-Sakhalinsk.

The collection of *K* index values, stored in the WDC for STP repository, contains data from 41 observatories located on the territory of the former USSR and the Antarctic (Table 1). The collection is unique and representative because it includes data obtained over a large area covering both high, medium, and low latitudes, and consists of a long series of *K* index values.

## 2 Description of *K* index

90 Comprehensive studies of the Earth magnetic field require more frequently defined geomagnetic activity measure, devoid of any subjectivity. Therefore, in 1939 the International Association of Terrestrial Magnetism and Electricity (now International Association of Geomagnetism and Aeronomy – IAGA) introduced a three-hour index *K* with a ten-point scale into the practice of magnetic observatories that was also proposed by Julius Bartels in 1938 (Bartels, 1938; 1939; Bartels et al., 1939). The designation *K* derives from the German word ‘Kennziffer’ (Bartels, 1938), which literally means a ‘characteristic parameter’  
95 or ‘identification number’. The *K* index was designed as a measure of the range of irregular and rapid geomagnetic activity, including geomagnetic storms. It implies to be insensitive to the long-term components of magnetic variations (Love and



Remick, 2007). It is a numerical characteristic of the variability of geomagnetic activity at a given observatory over 3-hour intervals. In this particular case the variability is understood as the amplitude of the observed horizontal field relative to the quiet daily variation  $Sq$ . The data source for  $K$  index calculating is magnetograms. Initially, when only analogue magnetograms 100 were available, magnetologists determined the amplitudes of geomagnetic disturbances and, consequently, the  $K$  index values manually (Fig. 3).

At Soviet observatories, indices were calculated once per 10 days or month based on the results of analog magnetograms' processing, following the procedure described in (Bartels et al., 1939). According to the procedure, for  $K$  index calculation it is necessary to determine and remove the regular  $Sq$  part of the record. Quiet daily variation  $Sq$  was determined based on five 105 daily magnetograms with clear and distinguishable characteristics of quiet days. Then amplitudes of variations of geomagnetic vector horizontal components ( $H$  and  $D$ ) for each three-hour interval were defined. Then the regular part of the quiet daily variation  $Sq$  was subtracted. To facilitate this procedure, special transparent flexible gauges (transparent pallets on celluloid or other similar materials) were used. In each observatory, they were drawn using the quiet daily variation line and the observatory 110  $K$  scale from Table 2 (Nechaev, 2006; Zabolotnaya, 2007). Considering that the amplitude of the geomagnetic disturbance depends on the latitude of the observation site (the maximum amplitude is observed in the aurora zone), the value of the  $K$  scale division is selected for each observatory. The largest value of the two amplitudes obtained for each of the three-hour 115 intervals using the observatory  $K$  scale was converted to the  $K$  index value and adopted as the final one. Only the horizontal  $H$  and  $D$  components are used to determine the  $K$  index. Component  $Z$  for observatories located 1,000 or 2,000 km from the auroral zone is extremely sensitive to the auroral electrojet and more than  $H$  and  $D$  sensitive to local effects of induced underground currents (Mayaud, 1967).

The  $K$  index for a particular observatory takes values from 0 to 9 for each 3-hour interval, starting from 00:00 UT, where 9 corresponds to the strongest geomagnetic disturbances. The value  $K = 9$  corresponds to a disturbance exceeding 2,500 nT in the auroral zone and about 300 nT at low latitudes (excluding the equator). For each observatory, a correspondence between the value  $K = 9$  and the disturbance amplitude was obtained by considering the extremely strong geomagnetic disturbance 120 observed on 16 April 1938. It was agreed that on this day between 06:00 and 09:00 UT the  $K$  index at all observatories was equal to 9 points, and the maximum value of the disturbance amplitude over the considered interval was taken as the lower amplitude limit corresponding to  $K = 9$ . An event with  $K \leq 2$  is quiet,  $K = 2–3$  is slightly disturbed,  $K = 4$  is disturbed,  $K = 5–6$  is a magnetic storm, and  $K \geq 7$  is a large magnetic storm.

With the advent of magnetic variometers with digital registration, data processing and calculation of  $K$  index was carried out 125 using computers. It became necessary to develop algorithms for calculating  $K$  index in automatic mode. Computer programs for calculating  $K$  index developed in the 1980s differ mainly in the method for determination of the quiet daily variation  $Sq$  (Menvielle et al., 1995). Two main approaches are used: averaging the magnetograms of the nearest quiet days (Takahashi et al., 2001; Nechaev, 2006; Dmitriev and Filippov, 2010) and by smoothing the current magnetogram (Hopgood, 1986; Jankowski et al., 1988; Golovkov et al., 1989; Sucksdorff et al., 1991).



130 The ex-Soviet observatories, such as Irkutsk (IRT) or Borok (BOX), use algorithms that are as close as possible to the method of Julius Bartels that allows to preserve the continuity of the observatory *K* index timeseries for previous years (Anisimov et al., 2015; Dmitriev and Filippov, 2010; Nechaev, 2006).

But the *K* index is a local indicator, describing geomagnetic disturbances in the vicinity of a particular observatory. In 1949 Julius Bartels suggested a new planetary *Kp* index for estimating the geomagnetic activity on the planetary scale over 3-hour intervals (Bartels, 1949). The planetary *Kp* index is calculated as the average of the *K* indices from 13 selected observatories situated in the subauroral zones. The *K* index values are currently used for the derivation of the IAGA planetary geomagnetic indices *Kp* (*ap*), *am* (*Km*), *an* (*Kn*), *as* (*Ks*) and *aa* (Berthelier, 1993; Mayaud, 1980; Menvielle and Berthelier, 1991; Menvielle et al., 1995).

### 3 Compilation of the historical *K* index data collection

140 Determination of the *K* index is a standard procedure for magnetic observatory practice. The data obtained using the abovementioned procedure were prepared as monthly arrays in the form of standard datasheets. The tables for these datasheets were developed and adopted by the beginning of the IGY (Fig. 4a). The IAGA datasheet was introduced in June 1977 (Fig. 4b). In the 1980s, datasheets with addition of information on magnetic storms, recommended by IZMIRAN, was used to represent data from Soviet observatories (Fig. 5). Datasheets with *K* index definitions were sent to the WDC B2 by mail and 145 stored as paper documents. Later, in the 1990s, the data exchange via e-mail started. Each datasheet indicated the value of the lower limit for *K* = 9.

To guarantee the preservation of this data and provide convenient and continuous on-line access and more efficient use, the entire array of paper datasheets was converted into digital form. The conversion was performed by scanning paper documents with transferring in the PDF format. By means of ScanSoft PaperPort software, electronic documents were edited: image crop 150 and alignment, removal of stains and excessive inscriptions (elements of “noise”). Where it was possible the text quality was manually improved. The following stage was visual verification of compliance of the digital version of documents with paper originals and necessary manual correction. The final structured data array in digital form (the array of digital documents) was published on the WDC for STP website with open access ([http://www.wdcb.ru/stp/data/K\\_indices/](http://www.wdcb.ru/stp/data/K_indices/), last access: 1 September 2020). Then the entire data set was converted into ASCII encoded text files. Most of the data represented as handwritten paper 155 documents was digitized manually. A small part of data was digitized using formatted text recognition and structured data extraction software, particularly the technology implemented in ABBYY FineReader software. The data files were visually checked and validated by calculating the total values in table line and comparing them to tabular daily values in Microsoft Excel 2010 software suite. Finally, all data files were converted into the uniform standard format adopted in the WDS (Fig. 6).



160 A detailed catalogue of data availability with monthly resolution was compiled. In total the collection includes 1054 annual files. The list of all observatories and stations that provided data for the presented collection is given in Table 1. It also shows the value of the  $K = 9$  lower amplitude limits for each of the observatories.

165 Data for 15 observatories (ASH, CCS, CWE, DIK, HIS, IRT, KIV, KZN, LNN, LVV, MMK, MOS, TKT, YAK, YSS) represent a single long time series of observations over 34–36 years without interruption. The OAS, PIO observatories in Antarctica, and TKH in the Arctic transferred their  $K$  index data to WDC only for 2 years during the IGY period of 1957–1958. They were closed shortly after IGY ending. Unfortunately, some sets have omissions, or data was not transferred or simply lost.

#### 4 Applicability of FAIR principles to the historical $K$ index data set

170 In the last decade, the term “FAIR Data” has become popular in scientific literature, as well as in general mass media. With time it became clear that FAIR data can significantly strengthen the results of scientific research and lead to their substantial and, sometimes, unexpected extensions. Scientists working in geophysics and the Earth sciences in general often claim nowadays that their results are obtained using FAIR pieces of information.

175 Following (<https://www.go-fair.org/fair-principles/>, last access: 1 September 2020) we, hereafter, shortly present the FAIR data formalism simultaneously showing that the historical  $K$  index data collection presented in this article fully satisfies the FAIR conditions.

The abbreviation FAIR stands for ‘Findable’, ‘Accessible’, ‘Interoperable’ and ‘Reusable’ data sets. Hereafter are the most valuable clarifications of what each of these four features means (Wilkinson et al., 2016). Along with that we show how the requirements are satisfied for the data set of the  $K$  indices under consideration (*in italic*).

##### To be Findable:

180 F1. data are assigned a unique and eternally persistent identifier (*see the DOI of the K index collection at PANGAEA data repository*).  
F2. data are described with rich metadata (*provided in the PANGAEA system along with a comprehensive description given in this paper*).  
F3. data are registered and indexed in a searchable resource (*in the PANGAEA data repository itself*).

##### 185 To be Accessible:

A1. data are retrievable by their identifier using a standardized communications protocol (*implemented by DataCite system, which ensures adequate DOI retrievals*).

A2. the protocol is open, free, and universally implementable (*the DataCite system fully satisfies this requirement*).

190 A3. metadata are accessible, even when the data are no longer available (*this is provided by independent storage of the metadata in the DataCite system*).

##### To be Interoperable:



I1. data and metadata use a formal and broadly applicable means for knowledge representation (*this is ensured by using the English language and standard Excel tables for data storage*).

I2. data and metadata include qualified references from other resources (*information description in this paper includes sufficient references on the dataset*).

195

**To be Reusable:**

R1. data have accurate and relevant attributes (*in the PANGAEA system the data are attributed with abstract, keywords, geographic coverage, dataset size, etc.*).

R2. (meta)data are released with a clear and accessible data usage license (*information on data usage license is provided by 200 the PANGAEA system*).

R3. (meta)data are associated with their provenance (*information on the provenance of the K index collection is given in detail in this paper as well as in the dataset description in the PANGAEA system*).

R4. (meta)data meet domain-relevant community standards (*the presented K index collection fully meets the requirements of IAGA and WDS*).

205 Hence, the FAIR requirements for the considered historical *K* index data collection are fully satisfied.

## 5 Conclusion

Implementation of digital magnetometers and data loggers that provide near real-time transfer of observational information to 210 geophysical data repositories has become a pivotal moment in the observatory practice. But fast and convenient access to digital magnetograms from observatories all over the world did not completely substitute the traditional parameters of geomagnetic activity. *K* index – the classical commonly used parameter of geomagnetic activity – serves as the measure of 215 local magnetic field variations within three-hour intervals. The *K* index values defined at different observatories are used for calculating other geomagnetic activity indices such as *Kp* (*ap*), *am* (*Km*), *an* (*Kn*), *as* (*Ks*) and *aa*. The *K* index is used for studying of magnetic storms – the most vivid reflection of the Sun’s impact on the Earth’s magnetosphere. The study of magnetic storms and geomagnetic variations that they induce is important for analyzing their impact on meteorological 220 processes, biosphere, technological infrastructure, etc.

The long-time practice of the *K* index application makes it an important instrument for retrospective complex analysis of solar-terrestrial interaction. The beginning of coordinated large-scale geomagnetic observations in the USSR was initiated during 225 the International Geophysical Year of 1957–1958. Although some of the observatories were temporary and were discontinued by the end of 1958, the observational network of the former USSR remained relatively dense and provided the sufficient spatial coverage. The World Data Center (WDC) B2 which later was renamed as WDC for Solar-Terrestrial Physics (WDC for STP) in Moscow became the main repository of data on geomagnetism in the former USSR. The *K* index datasheets were transferred from observatories to the WDC for STP where they are stored as paper documents. In 2000s and 2010s WDC for STP initiated an ambitious project for digitizing this archive of paper documents. The presented unique collection of historical *K* index



values (Sergeyeva et al., 2020) is the result of this activity. It is a vast massif of homogeneous data that cover more than three  
225 decades of geomagnetic measurements at observatories of the former USSR. Considering the FAIR principles in regard to the  
presented data collection, in this article we come up with a new example of FAIR data.

## 6 Data availability

The data on the *K* index from paper catalogues was digitized at Geophysical Center of RAS by the team of the World Data  
Center for Solar-Terrestrial Physics. Digital data are available at: <https://doi.org/10.1594/PANGAEA.922233>, last access: 16  
230 September 2020 (Sergeyeva et al., 2020). Photocopies of the datasheets and analogue magnetograms from geomagnetic  
observatories of the former USSR since 1957 are accessible via the web-page of the World Data Center for Solar-Terrestrial  
Physics: <http://www.wdcb.ru/stp/data.html>, last access: 4 September 2020.

**Author contributions.** NS, AG, AS, and RK – preparation of the manuscript and figures. LZ, TK, and MN – digitalization of  
analogue data, data validation, and preparation of the *K* index data collection. MN – publication of data in the PANGAEA  
235 system.

**Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgements.** The authors wish to thank the team of the World Data Center for Solar-Terrestrial Physics in Moscow,  
Russia, for preservation and making publicly available historical geophysical data. This work employed data provided by the  
Shared Research Facility “Analytical Geomagnetic Data Center” of the Geophysical Center of RAS (<http://ckp.gcras.ru/>, last  
240 access: 1 September 2020).

**Financial Support.** This work was conducted in the framework of budgetary funding of the Geophysical Center of RAS,  
adopted by the Ministry of Science and Higher Education of the Russian Federation.

## References

Anisimov, C. V., Dmitriev, E. M., Guryev, A. V., and Filippov, V. A.: Calculation of K-indices of geomagnetic activity in real  
245 time (in Russian), *Heliogeophysical research*, 13, 62–72, 2015.

Bartels, J.: Potsdamer erdmagnetische Kennziffern. 1 Mitteilung (in German), *Zeitschrift für Geophysik*, 14, 68–78, 1938.

Bartels, J.: Potsdamer erdmagnetische Kennziffern. 4 Mitteilung (in German), *Zeitschrift für Geophysik*, 15, 214–221, 1939.

Bartels, J.: The standardized index *K<sub>s</sub>*, and the planetary index *K*, *IATME Bull.*, 12b, 97–120, 1949.

Bartels, J., Heck, N. H., and Johnston, H. F.: The three-hour-range index measuring geomagnetic activity, *Journal of  
250 Geophysical Research*, 44(4), 411–454, doi:10.1029/te044i004p00411, 1939.



Belov, B. A., Burtsev, Yu. A., Kiriakov, V. Kh., and Lyubimov, V. V.: Digital quartz magnetic variation stations (in Russian), in “170 years of observational observations in the Urals: history and current state” (Proceedings of the international seminar, Yekaterinburg, July 17–23, 2006, Yekaterinburg), 19–23, 2006.

Berthelier, A.: The K-derived planetary indices: derivation, meaning and uses in solar terrestrial physics, in STPW-IV 255 proceedings, 3, 3–20, eds. Hruska, J. et al., US Gov. Publications Office, Boulder CO, 1993.

Bobrov, V. N.: Universal highly stable sensitive element with zero temperature coefficient for magnetometers, variometers and microvariometers registering any component of the Earth's magnetic field (in Russian), Trudy IZMIRAN (Works of IZMIRAN), 18 (28), 54, 1961.

Burtsev, Yu. A., Mansurov, S. M., and Timofeev, G. A.: Autonomous variation station for geomagnetic research in Antarctica 260 (in Russian), in Geomagnetic instrument engineering (ed. Sh. Sh. Dolginov), Moscow, Nauka Press, 1977.

Dmitriev, E. M. and Filippov, V. A.: Algorithm for calculation of indices of geomagnetic activity (in Russian), in proceedings of the V International conference “Solar-terrestrial relations and physics of earthquake precursors” (Petropavlovsk-Kamchatsky, IKIR Feb RAS), 110–113, 2010.

Golovkov, V. P., Papitashvili, V. O., and Papitashvili, N. E.: Automated calculation of K-indices using the method of natural 265 orthogonal components, *Geomagnetism and Aeronomy*, 29(4), 514–517, 1989.

Gvishiani, A. and Soloviev, A.: EMF Observations and Data Processing, in *Observations, Modeling and Systems Analysis in Geomagnetic Data Interpretation*, pp. 29–96, Springer International Publishing., doi: 10.1007/978-3-030-58969-1\_2, 2020.

Hopgood, P. A.: On the computer generation of geomagnetic K-indices from digital data., *Journal of geomagnetism and 270 geoelectricity*, 38(9), 861–871, doi:10.5636/jgg.38.861, 1986.

Jankowski, J. A., Ernst, T., Sucksdorff, C., Pirjola, R., and Ryno, J.: Experiences of a filter method and a standard curve method for determining K-indices, *Annales Geophysicae*, 6, 589–594, 1988.

Lincoln, J. V.: Geomagnetic indices, in *Physics of Geomagnetic Phenomena*, Vol. 1, (eds. S. Matsushita and W. H. Campbell), Academic, New York, 67–100, 1967.

Love, J. J. and Remick, K. J.: Magnetic Indices, in *Encyclopedia of Geomagnetism and Paleomagnetism*, 509–512, Springer 275 Netherlands, doi: 10.1007/978-1-4020-4423-6\_178, 2007.

Lyubovtseva, Y. S., Gvishiani, A. D., Soloviev, A. A., Samokhina, O. O., and Krasnoperov, R. I.: Sixtieth anniversary of the International Geophysical Year (1957–2017) – contribution of the Soviet Union, *Hist. Geo Space. Sci.*, 11(2), 157–171, doi:10.5194/hgss-11-157-2020, 2020.

Mayaud, P. N.: Atlas of indices K. Part 1, *IAGA Bulletin*, 21, 1967.

280 Mayaud, P. N.: Derivation, meaning and use of geomagnetic indices, *Geophysical Monograph*, 22, AGU, Washington, DC, 1980.

Menvielle, M. and Berthelier, A.: The K-derived planetary indices: Description and availability, *Reviews of Geophysics*, 29(3), 415–432, doi:10.1029/91rg00994, 1991.



285 Menvielle, M., Papitashvili, N., Hakkinen, L., and Sucksdorff, C.: Computer production of K indices: review and comparison of methods, *Geophysical Journal International*, 123(3), 866–886, doi:10.1111/j.1365-246x.1995.tb06895.x, 1995.

Nechaev, S. A.: Guide for stationary geomagnetic observations (in Russian), Irkutsk: Institute of geography SB RAS, 140 p., ISBN 5-94797-093-7, 2006.

Nicolet, M.: Historical aspects of the IGY, *History of Geophysics*, Vol. 1, AGU, 44–46, doi:10.1029/HG001p0044, 1984.

Raspopov, O. M., Kuz'min, I. A., and Kharin, E. P.: The 50<sup>th</sup> anniversary of International Geophysical Year (1957–1958): 290 from the First International Polar Year (1882–1883) to the International Heliophysical Year (2007–2008) and International Polar Year (2007–2009), *Geomagnetism and Aeronomy*, 47(1), 1–7, doi:10.1134/s001679320701001x, 2007.

Rodnikov, A. G., Sergeyeva, N. A., and Zabarinskaya, L. P.: Research on the Earth's Interior Conducted by Russia after IGY: The Geotraverse Project and “INTERMARGINS”, *Data Science Journal*, 8, S24–S34, doi:10.2481/dsj.8.S24, 2009.

Sergeyeva, N., Gvishiani, A., Soloviev, A., Zabarinskaya, L., Krylova, T., Nisilevich, M., and Krasnoperov, R.: Historical K 295 index data collection of Soviet magnetic observatories, 1957–1992. PANGAEA, doi:10.1594/PANGAEA.922233, 2020.

Sucksdorff, C., Pirjola, R., and Häkkinen, L.: Computer production of K-indices based on linear elimination, *Geophysical Transactions*, 36(3–4), 333–345, 1991.

Takahashi, K., Toth, B. A., and Olson, J. V.: An automated procedure for near-real-time K<sub>p</sub> estimates, *Journal of Geophysical Research: Space Physics*, 106(A10), 21017–21032, doi:10.1029/2000ja000218, 2001.

300 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., et al.: The FAIR Guiding Principles for scientific data management and stewardship, *Sci Data*, 3(1), doi:10.1038/sdata.2016.18, 2016.

Zabolotnaya, N. A.: Indices of geomagnetic activity: reference guide. 2nd ed. (in Russian), Moscow, LKI Publishing house, 88 p., 2007.



305 **Tables**

**Table 1: Geomagnetic observatories of the former USSR. K index data availability**

No.	Code	Name of the observatory	Country	Geographical coordinates		Altitude, m	Lower limit for K = 9, nT	K index Availability of data in the form of digital files, years
				Latitude	Longitude			
1	AAA	Alma- Ata	Kazakhstan	43.250	76.920	1300	300	1964–1981, 1983–1992
2	ARK	Arkhangelsk	Russia	64.583	40.5000		300	1991–1992
3	ARS	Arti (Yekaterinburg)	Russia	56.433	58.567	290	550	1976–1992
4	ASH	Ashkhabad (Vannovskaya)	Turkmenistan	37.950	58.100	300	300	1958–1991
5	CCS	Cape Chelyuskin	Russia	77.717	104.283	10	2500	1957–1992
6	CWE	Cape Wellen (Uelen)	Russia	66.170	−169.830	10	1250	1957–1992
7	DIK	Dikson (Dixon)	Russia	73.543	80.562	20	1500	1957–1992
8	HIS	Heiss Island (Druzhnaya)	Russia	80.620	58.050	20	2000	1957–1992
9	IRT	Irkutsk (Patrony)	Russia	52.167	104.450	460	350	1957–1992
10	KGD	Karaganda (Bereznyaki)	Kazakhstan	49.820	73.080		350	1973–1981, 1983–1990
11	KIV	Kiev (Dymer)	Ukraine	50.720	30.300	100	350	1958–1991
12	KNG	Kaliningrad	Russia	54.600	20.200		600	1977–1981, 1983–1992
13	KZN	Kazan (Zaymishe)	Russia	55.830	48.850	80	550	1957–1992
14	LNN	Leningrad (Voyeykovo)	Russia	59.950	30.700	70	600	1957–1992
15	LVV	Lvov	Ukraine	49.900	23.750	400	550	1957–1991
16	MGD	Magadan (Stekol'nyj)	Russia	60.120	151.020		550	1965–1972, 1976–1994
17	MIR	Mirny	Antarctica (Russia)	−66.550	93.020	20	2000	1957–1981, 1983–1990
18	MMK	Loparskaya (Murmansk)	Russia	68.250	33.080	210	2500	1957–1993
19	MNK	Minsk (Pleshchenicy)	Belorussia	54.100	26.520		550	1962–1981, 1983–1991
20	MOL	Molodezhnaya	Antarctica (Russia)	−67.670	45.850	0	2000	1975–1981, 1983–1990
21	MOS	Moscow (Krasnaya Pakhra)	Russia	55.467	37.312	190	550	1949–1992
22	NKK	Novokazalinsk	Kazakhstan	45.770	62.120		350	1973–1981, 1983–1990
23	NVL	Novolazarevskaya	Antarctica (Russia)	−70.770	11.820	460	1500	1973–1981, 1983–1988
24	NVS	Novosibirsk	Russia	55.030	82.900		500	1969–1992
25	OAS	Oasis	Antarctica (Russia)	−66.300	100.720		2000	1957–1958
26	ODE	Odessa (Stepanovka)	Ukraine	46.780	30.880	140	350	1957–1981, 1983–1991
27	PET	Petropavlovsk (Paratunka)	Russia	53.100	158.630		450	1969–1971, 1973–1992
28	PIO	Pionerskaya	Antarctica (Russia)	−69.730	95.500		2000	1957–1958
29	POD	Podkamennaya Tunguska	Russia	61.400	90.000		650	1969–1971, 1974–1992
30	SRE	Srednikan	Russia	62.430	152.320		550	1957–1966



31	SVD	Sverdlovsk (Vysokaya Dubrava)	Russia	56.730	61.070	290	550	1957–1977
32	TFS	Tbilisi (Dusheti)	Georgia	42.080	44.700	982	350	1957–1991
33	TIK	Tixie Bay (Tiksy)	Russia	71.580	129.000	40	1000	1957–1967, 1969–1991
34	TKH	Tikhaya Bay	Russia	80.300	52.800		2000	1957–1958
35	TKT	Tashkent (Yangi Bazar)	Uzbekistan	41.333	69.617	500	300	1957–1991
36	TMK	Tomsk	Russia	56.470	84.930	200	350	1958–1970
37	UBA	Ulan–Bator	Mongolia	47.850	106.750		300	1973–1975, 1978–1980, 1983–1989
38	VLA	Vladivostok (Gornotayozhnaya)	Russia	43.697	132.160	178	300	1957–1981, 1983–1988, 1990
39	VOS	Vostok	Antarctica (Russia)	–78.450	106.867	3500	2000	1958–1981, 1983–1990
40	YAK	Yakutsk	Russia	62.020	129.720	100	550	1957–1991
41	YSS	Yuzhno Sakhalinsk	Russia	46.950	142.717	70	350	1957–1990

**Table 2: Correspondence of geomagnetic disturbance amplitudes in nT to K index values depending on the geomagnetic latitude of observatory.**

Scale	K index										Lat
	0	1	2	3	4	5	6	7	8	9	
1	0–25	25–50	50–100	100–200	200–350	350–600	600–1000	1000–1650	1650–2500	>2500	64–90
2	0–20	20–40	40–80	80–160	160–280	280–480	480–800	800–1300	1300–2000	>2000	65–80
3	0–18	18–36	36–72	72–144	144–252	252–432	432–720	720–1188	1188–1800	>1800	80
4	0–15	15–30	30–60	60–120	120–210	210–360	360–600	600–1000	1000–1500	>1500	60–83
5	0–12	12–25	25–50	50–100	100–175	175–300	300–500	500–825	825–1250	>1250	62
6	0–10	10–20	20–40	40–80	80–140	140–240	240–400	400–660	660–1000	>1000	60–83
7	0–8	8–15	15–30	30–60	60–105	105–180	180–300	300–500	500–750	>750	58–62
8	0–6	6–12	12–24	24–48	48–85	85–145	145–240	240–400	400–600	>600	55–58
9	0–5	5–10	10–20	20–40	40–70	70–120	120–200	200–330	330–550	>550	48–54
10	0–5	5–10	10–20	20–40	40–70	70–120	120–200	200–330	330–550	>500	36–57
11	0–4	4–8	8–16	16–30	30–50	50–85	85–140	140–230	230–350	>350	31–47
12	0–3	3–6	6–12	12–24	24–40	40–70	70–120	120–220	200–300	>300	0–40



## Figures






Figure 1: The map shows the observatories used for calculating the geomagnetic indices: *Kp* (green circles); *AE, AU, AL, AO* (orange circles) and *Dst* (red circles).

315



**Figure 2: Location of Soviet geomagnetic observatories that provided data for the presented collection: a) continental territory of the USSR; b) the Antarctic.**



320

Figure 3: Photocopies of analogue magnetograms: a) Petropavlovsk (Paratunka) observatory, 16—17.08.1970, Bobrov variometer; Irkutsk (Patrony), 14.07.1960, Yanovsky variometer; c) manual determination of  $K$  (purple bars) and  $K_p$  values (<http://www.wdcb.ru/stp/index.en.html>, last access: 1 September 2020).



**a)** IRKUTSK  
 Иркутск Суточные характеристики Июль 1957 г.

Дата	Амплитуды			Характер	K-индекс								Дата	
	$\Delta D$	$\Delta H$	$\Delta Z$		0-3	3-6	6-9	9-12	12-15	15-18	18-21	21-24	ΣK	
1				2	4	4	5	3	2	3	6	3	30	1
2				2	2	1	3	4	6	5	3	2	26	2
3				1	4	4	4	4	5	1	2	2	26	3
4				0	2	2	2	2	1	1	2	4	16	4
5				1	5	6	6	5	2	1	2	3	30	5
6				0	3	3	3	3	2	2	3	3	22	6
7				0	3	3	2	3	2	1	2	2	18	7
8				0	2	3	2	2	2	1	2	2	16	8
9				0	2	2	2	2	1	2	1	1	13	9
10				0	2	2	2	2	0	0	1	1	10	10
11				0	-	1	2	2	4	1	1	1	-	11
12				0	3	2	3	2	2	2	2	2	18	12
13				0	1	2	2	2	0	1	1	2	11	13
14				0	2	2	2	2	2	1	2	3	16	14
15				0	3	2	1	2	1	0	0	2	11	15
16				1	3	2	4	3	3	3	2	3	23	16
17				0	3	3	2	3	2	2	2	3	20	17
28				0	2	2	3	3	2	3	2	3	20	18
19				1	2	4	4	3	3	4	3	3	26	19
20				0	2	3	2	3	2	1	3	2	18	20
21				0	2	2	2	3	1	1	1	2	14	21
22				1	3	3	4	4	3	3	4	3	27	22
23				1	3	4	3	1	1	1	1	3	17	23
24				0	2	3	3	4	3	3	2	2	22	21
25				0	2	2	3	1	2	3	2	-	-	25
26				0	1	1	1	0	2	1	1	2	09	26
27				0	2	1	1	2	1	2	3	3	15	27
28				0	3	2	2	3	1	1	1	2	15	28
29				1	3	3	4	4	3	3	1	1	22	29
30				0	2	2	3	3	1	1	1	1	14	30
31				0	3	3	2	2	1	1	2	3	17	31
Среднее	$\bar{\Delta D} = 5.54$			$\bar{\Delta H} = 4.57$			$\bar{\Delta Z} = 3.28$			$\bar{Kg} = 350$				

Зак. № 1093 2000

**b)** INTERNATIONAL UNION OF GEODESY AND GEOPHYSICS  
 INTERNATIONAL ASSOCIATION OF GEOMAGNETISM AND AERONOMY

INDICES OF GEOMAGNETIC ACTIVITY  
 OBSERVATORY *Patrony (Irkutsk)* SCALE-VALUES OF VARIOMETERS  
 MONTH *July* 1977 IN Y/MM  
 RANGE FOR K=0...350 D..... H..... Z.....

GR	K- INDICES FOR THREE-HOUR INTERVAL								SUM
	00 <sup>h</sup> 03 <sup>h</sup> 06 <sup>h</sup>	09 <sup>h</sup> 09 <sup>h</sup> 12 <sup>h</sup>	12 <sup>h</sup> 15 <sup>h</sup> 18 <sup>h</sup>	15 <sup>h</sup> 18 <sup>h</sup> 21 <sup>h</sup>	18 <sup>h</sup> 21 <sup>h</sup> 24 <sup>h</sup>	21 <sup>h</sup>			
1	3	0	0	5			2	4	3 3 20
2	3	3	3	2			2	2	3 4 22
3	3	2	4	3			3	2	2 2 21
4	-	-	-	-			-	-	- - -
5	2	I	3	2			3	2	3 I 17
6	3	3	4	4			6	5	4 3 32
7	4	3	4	3			3	3	2 3 25
8	3	3	3	4			3	3	2 3 24
9	3	3	3	4			2	2	2 4 23
10	3	3	3	3			4	3	3 3 25
11	3	3	4	2			3	2	3 3 23
12	2	2	2	2			0	0	0 0 0 08
13	2	3	3	2			3	2	3 3 21
14	4	5	5	4			3	2	2 3 28
15	2	2	3	2			2	2	2 4 19
16	2	4	4	-			-	-	- - - I 0
17	2	3	4	3			2	3	3 4 24
18	2	I	2	3			3	2	2 4 I 9
19	2	3	5	4			4	4	3 4 29
20	3	4	4	4			5	3	3 4 30
21	3	2	2	2			2	3	4 I 8
22	2	2	3	3			-	-	- - - I 0
23	2	2	3	3			2	3	3 2 20
24	2	2	2	2			2	I	2 2 I 5
25	2	3	3	2			I	2	I 3 I 7
26	0	2	2	2			I	2	I 3 I 3
27	I	I	I	I			I	I	I 2 0 9
28	I	I	2	2			I	2	I 3 I 3
29	5	6	5	5			7	5	2 3 38
30	5	3	4	4			2	2	2 2 24
31	2	2	2	3			2	2	2 3 I 8

Form № 1082.

Figure 4: Paper datasheets for K index values for observatory Irkutsk (Patrony): a) datasheet form adopted for the IGY (July 1957); b) datasheet form adopted by IAGA in 1977 (July 1977) ([http://www.wdcb.ru/stp/data/K\\_indices/IRT\\_Irkutsk/](http://www.wdcb.ru/stp/data/K_indices/IRT_Irkutsk/), last access: 1 September 2020).



МЕСЯЧНЫЙ ОБЗОР СОСТОЯНИЯ МАГНИТНОГО ПОЛЯ

МЕСЯЦ ЯНВАРЬ 1990 г.  
 ВРЕМЯ МИРОВОГО УТ

ОБСЕРВАТОРИЯ Душети (Ингебиши)  
 ВЕДОМСТВО Инст. Геофизики  
АН Груз. ССР

ДН ИУ	УТ												Ю БУРЫ	Начало ДАТА УТ	Конец ДАТА УТ	ПРО ДОЛЖ	АМПЛИТУДА	АКТИВНЫЕ ПЕРИОДЫ			
	U-3	3-6	6-9	9-12	12-15	15-18	18-21	21-24	ΣH	I	H	Z						НАЧАЛО ДАТА УТ	КОНЕЦ ДАТА УТ		
1	3	3	3	3	3	4	5	4	28	30.1.10 <sup>00</sup>	31.1.03 <sup>00</sup>	20	92	87	44	М	30.1.10 <sup>00</sup>	30.1.09 <sup>00</sup>			
2	4	2	3	3	3	2	3	4	24								30.1.13 <sup>00</sup>	30.1.15 <sup>00</sup>			
3	4	3	2	3	3	4	3	3	25								30.1.14 <sup>00</sup>	30.1.23 <sup>00</sup>			
4	2	2	3	4	3	2	2	3	21								31.1.01 <sup>00</sup>	31.1.03 <sup>00</sup>			
5	3	2	3	3	3	4	4	4	26												
6	3	3	2	2	2	2	2	2	18												
7	2	2	2	2	2	1	1	2	14												
8	2	2	2	2	4	5	4	4	25												
9	4	3	3	3	3	2	2	4	24												
10	4	2	3	4	4	4	4	3	28												
11	3	2	3	3	3	4	4	3	25												
12	3	3	3	4	4	3	4	2	26												
13	3	2	2	3	2	4	2	2	20												
14	2	3	2	3	3	3	2	2	20												
15	2	3	2	3	3	4	2	2	21												
16	2	3	2	3	2	4	4	4	24	Максимум магнитных бурь 30 <sup>0</sup> -31 <sup>0</sup>											
17	3	2	2	2	3	2	1	2	17	шестое постепенное начало.											
18	3	2	3	3	3	3	2	3	22												
19	2	2	1	2	2	3	3	3	18												
20	3	3	3	3	3	5	4	3	27												
21	4	3	3	3	4	4	4	3	28												
22	4	3	3	4	3	4	3	3	27												
23	3	3	3	4	3	3	5	3	27												
24	4	4	4	4	3	4	5	5	33												
25	5	3	3	4	4	3	2	2	26												
26	2	3	2	4	4	3	2	2	22												
27	2	3	1	2	1	1	2	1	13												
28	1	2	2	3	3	2	4	3	20												
29	4	3	4	5	4	3	4	4	31												
30	4	4	3	3	4	4	5	3	30												
31	4	3	3	3	4	4	3	2	26												

Обзор составил \_\_\_\_\_

Figure 5: Monthly K index values datasheet for observatory Tbilisi (TFS), Georgia (January 1990), compiled according to the form, proposed by IZMIRAN ([http://www.wdcb.ru/stp/data/K\\_indices/TFS\\_Tbilisi/TFS\\_1990\\_1-7+st.pdf](http://www.wdcb.ru/stp/data/K_indices/TFS_Tbilisi/TFS_1990_1-7+st.pdf), last access: 1 September 2020).



Line 1: from the 9th position - Observatory name and field value for k = 9 in nT.

Line 2: 1 - 3 positions - Observatory code.

Line 3: 1 - 4 i4 Year.

6 - 7 i2 Month.

9 - 10 i2 Day.

13 - 15 i3 Day of the year.

17 - 32 8i2 Eight 3-hourly values of the index for the day.

35 - 36 i2 The sum of 8 values of the index per day.

Line 4 and all subsequent to the end of the month:

6 - 7 i2 Month.

9 - 10 i2 Day.

13 - 15 i3 Day of the year.

17 - 32 8i2 Eight 3-hourly values of the index for the day.

35 - 36 i2 The sum of 8 values of the index per day.

Then each new month begins with the 2nd line.

A missing value is identified by “-1”.

**Figure 6:** Format description for K index data ([http://www.wdcb.ru/stp/geomag/format\\_K\\_index.html](http://www.wdcb.ru/stp/geomag/format_K_index.html), last access: 1 September 2020).