

Interactive comment on “Synoptic Analysis of a Decade of Daily Measurements of SO₂ Emission in the Troposphere from Volcanoes of the Global Ground-Based Network for Observation of Volcanic and Atmospheric Change” by Santiago Arellano et al.

Santiago Arellano et al.

santiago.arellano@chalmers.se

Received and published: 16 December 2020

We thank Dr. Arlin Krueger for the thorough and thoughtful review of our manuscript and for his many valuable suggestions.

Dr. Krueger's report provides a balanced and complete description of the content of this lengthy manuscript. His overall highly positive remarks are encouraging. In response to the most critical comments (between quotation marks) in Dr. Krueger's report we

C1

present the following arguments:

“In Sec. 2, I am bothered by treating the word ‘data’ as singular, as in ‘Data is transferred via a serial port. . .’ (line 180, and again in line 189) in the scientific literature. I realize in non-scientific literature ‘data’ can describe a single collection of facts, etc., but I learned to use “datum” for singular measurements and ‘data’ for multiple points.”

We thank for pointing out this grammatical error, which is now amended throughout the manuscript.

“Figure 3, consisting of 32 plots of station data, is impossible to read and should be presented in two or more sub-figures. In Fig. 5 (mis-referenced as fig. 4) the series of annual emissions from 12 volcanoes is compared with OMI data. These 12 panels are just legible on a printed page.”

We agree that the figures need to be improved. We have revised all figures, increased their size, and split Figure 3 into four figures for better visualization. We have also corrected the mis-referencing of Figure 4. Thanks for pointing out this mistake.

“Steps to deal with the problem of calculating long-term emission budgets from irregular sets of data are discussed. The adopted policy of filling missing days by interpolation is understandable but problematic.”

We agree that the procedure to attribute periods of null detection of volcanic plumes by the instruments to actual low levels of volcanic outgassing may not be entirely satisfactory, and in this sense “problematic”, due to lack of detailed information for all conceivable cases. Although the comment does not elaborate on details, we would like to stress that our procedure is a proposal to deal with irregular time series allowing a more reasonable comparison with the satellite-based dataset, which is assumed to have a more complete coverage. This is the simplest unbiased alternative to performing an interpolation between observed data-points: when all conditions are met for detecting a volcanic plume, and yet no emission is observed, the most likely ex-

C2

planation is that the emission was in fact low. This is quite different from assigning an emission equal to the mean-point between the two nearest observed points. Beyond the goal of comparing long-term emission budgets between two instruments, the proposed method could be very useful for monitoring purposes, because contextual information on wind direction and cloud cover allows to differentiate between true low emission of the volcano and other non-volcanic causes for missing the plume (weather, plume direction, instrumental failures). This type of information is crucial to interpret the variations in volcanic activity. Furthermore, we present both the original and the re-constructed time series resulting from applying this method, as well as an indication of the density of data for NOVAC. The cautious reader can then judge the potential bias that this method may cause, for example if the number of days with observed plumes is too low to serve for a conclusive comparison. We think these are valid arguments for keeping both time-series (and tabulated data) in the manuscript and appendix.

“Even on a log scale the differences between NOVAC and OMI annual emissions are large and the explanations are not convincing.”

Since this is the more substantial criticism to the manuscript, in the following we attempt to respond each assertion separately.

- “I would think the failure of OMI in separating emissions from two nearby volcanoes would not be important as ground observers certainly will notice emissions from the other volcano, as noted.”

We agree and, as mentioned in the comment, already noted that ground-based observations could help to discriminate between nearby sources. This is the case, for example, at the Virunga volcanoes Nyamuragira and Nyiragongo or the Central American volcanoes San Cristóbal and Telica. Inherent to the method to achieve tropospheric sensitivity from OMI observations, is a loss in spatial resolution of sources located just a few tens of km apart. We do not argue that OMI ‘fails’ in estimating the aggregated emissions of one source region, but rather that by comparing the two methods, as in

C3

the case of Nyiragongo presented in Figure 5, we can now estimate the contribution of each individual source to the aggregated emission. Therefore, the large discrepancy observed after 2011 should not be interpreted as a failure of OMI to quantify emissions from Nyiragongo, but instead demonstrate the value of combining both methods to obtain the emissions from each of the two individual volcanoes. We think that the text explains this well:

[...in the case of two nearby volcanoes, such as Nyiragongo-Nyamuragira (with a footprint of 13x24 km²), OMI cannot separate completely the contributions of each source, so they are reported as a complex. In this respect, NOVAC can aid to discriminate between these sources, since the stations are deployed with a focus on Nyiragongo and the finer time resolution allows disentangling contributions, especially during periods of heightened activity at any of them.].

- “Limitations of NOVAC data to daytime hours cannot explain factors of 2 - 4 differences even given the high variability of emissions. In line 451ff ‘whereas OMI could in principle detect. . .’ OMI can in fact detect any emissions that have occurred day or night prior to the overpass. However, one has to account for chemical or physical losses in order to calculate instantaneous emission rates and totals.”

We partially disagree with these assertions. Firstly, differences within the same order of magnitude are expected when we consider the uncertainty distributions for both methods. As discussed in section 2.3, NOVAC has an asymmetric uncertainty distribution skewed towards values lower than the mean with a standard deviation which is typically in the order of 50%. By averaging data for one day, the random component of uncertainty is reduced but the reported mean daily flux still lies within about -30 and +10% of the most probable value (as explained in section 2.3 of the manuscript). To compare with OMI, we further calculate statistics over an entire year (with and without the ‘filling’ procedure) which results in an unknown, but presumably larger uncertainty for the annual emission estimate. This value is more likely an underestimate than an overestimate of the most probable flux, with an uncertainty of around 50%. If we as-

C4

sume, due to lack of better information, that the daily emission derived from averaging of OMI images is symmetric and within $\pm 50\%$, then it is perfectly reasonable that differences on the order of a factor of 2-4 are sometimes encountered between the annual emissions derived from the two methods. It is worth-mentioning that the differences are largely dependent on the amount of valid data and that in many cases the agreement between the two methods is remarkably good. Secondly, we used the qualifier “could in principle”, because detection of all emissions within the OMI overpass-interval would be feasible for complete coverage of the source and surroundings, for images uncontaminated by clouds or large column density pixels, and by accounting for ‘losses’ due to chemical reactions, deposition on ground, adsorption on ash or aerosol surfaces, or dilution in the atmosphere below the detection limit of the sensor. For OMI, not all volcanoes can be observed completely every day. This is first a consequence of the orbit of the Aura satellite that leaves narrow gaps (‘stripes’), particularly at low latitudes. Since 2007, OMI spectra have also been affected by the so-called ‘row-anomaly’ (The OMI Team, Ozone Monitoring Instrument Data User’s Manual) which changes with time and which can have a significant effect on the retrieval, and therefore data affected by this anomaly are removed resulting in a further degradation of daily coverage. Finally, to produce the OMI dataset that we used in our comparison, Carn et al. (2017) applied filters for cloud cover above 20%, solar zenith angles below 70 degree, large SO₂ column densities (5-15 DU), and other selection criteria. Together, all of these effects lead to a significant reduction in spatial coverage and thus, filtered in this way, the OMI dataset is far from complete on daily timescales.

- “As stated, an in-depth study of discrepancies is certainly needed. This section detracts from the otherwise excellent presentation of the research work. I suggest reworking or removing this section.”

We respectfully disagree with this assertion. The dataset available from OMI (Carn et al., 2017 and related work) is the most complete complementary record of observations which overlaps in time with the ground-based measurements presented in this

C5

study. It is therefore highly desirable to compare the two datasets. We acknowledge the difficulties and complexities in performing a high-quality comparison and validation study and agree with the referee that a more thorough examination is needed than what is possible within the scope of this manuscript, which first and foremost focuses on presenting the ground-based data. However, we feel it is necessary to make a first comparison here in order to point out the complexities, difficulties, and potential pitfalls of such an exercise, lest other researchers with less experience in this field attempt an oversimplified comparison and possibly draw incorrect conclusions from it. We present, as transparently as possible, all the steps from data acquisition to generation of daily emission statistics. We then produce annual averages for comparison by proposing a method to more properly fill data gaps. Still, we find that each volcano should be treated separately because of different conditions of measurement, level of activity or proximity to nearby sources. We hope that this first comparison will elicit a more thorough study, ideally involving both teams behind the production of the two datasets.

- “Perhaps treating the two datasets as complementary will help explain the differences, as alluded to in the conclusions. It could be that the real value of NOVAC may be in characterizing the continuing low-level background emissions of volcanoes. Any thoughts of validating older satellite data appear to be gone. However, if the suggestion that low-level background emissions dominate the global volcanic SO₂ budget is borne out, then this effort will have been successful.”

We agree with this comment, that both datasets are to a large degree complementary. The assertion that low-level background volcanic emissions of SO₂ are dominant is not originally ours but was instead determined by global satellite-based measurements. Since the first detection of SO₂ emitted during the 1982 El Chichón eruption by our referee and with increasing sensitivity over the years, satellite observations have demonstrated the predominance of passive over eruptive volcanic degassing. We believe NOVAC could play a crucial role in refining this picture. The reason behind validating OMI data only in this study is the availability of a very thorough and comprehensive

C6

dataset for this sensor; however, extending this comparison to other sensors such as GOME-2 or OMPS, which have been operational during the same period, would be very interesting. This would allow an assessment of the effects of different spatial resolutions and other instrumental factors in producing time-series of volcanic gas emissions from satellite records.

“Detailed comments” We thank the reviewer for spotting these errors, all of which are now amended.

Interactive comment on Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2020-295>, 2020.