

# Construction of homogenized daily surface air temperature for Tianjin city during 1887-2019

Peng Si<sup>1</sup>, Qingxiang Li<sup>2, 4</sup>, and Phil Jones<sup>3</sup>

<sup>1</sup> Tianjin Meteorological Information Center, Tianjin Meteorological Bureau, Tianjin, China

<sup>2</sup> School of Atmospheric Sciences, Sun Yat-sen University, and Key Laboratory of Tropical Atmosphere-Ocean System, Ministry of Education, Zhuhai, China

<sup>3</sup> Climatic Research Unit, School of Environmental Sciences, University of East Anglia, Norwich, UK

<sup>4</sup> Southern Laboratory of Ocean Science and Engineering (Guangdong Zhuhai), Zhuhai, China

Resubmitted to *Earth System Science Data*

March 2021

## \*Corresponding author:

Prof. Qingxiang Li  
School of Atmospheric Sciences  
Sun Yat-Sen University  
Tangjiawan, Zhuhai Campus of SYSU  
Zhuhai, China, 519082  
Tel/Fax: 86-756-3668352  
E-Mail: liqingx5@mail.sysu.edu.cn

**Abstract.** The century-long continuous daily observations from some stations are important for the study of long-term trends and extreme climate events in the past. In this paper, three daily data sources: (1) Department of Industry Agency of British Concession in Tianjin covering Sep 1 1890-Dec 31 1931 (2) Water Conservancy Commission of North China covering Jan 1 1932-Dec 31 1950 and (3) monthly journal sheets for Tianjin surface meteorological observation records covering Jan 1 1951-Dec 31 2019 have been collected from the Tianjin Meteorological Archive. The completed daily maximum and minimum temperature series for Tianjin from Jan 1 1887 (Sep 1 1890 for minimum) to Dec 31 2019 has been constructed and assessed for quality control and an early extension from 1890 to 1887. Several significant breakpoints are detected by the Penalized Maximal T-test (PMT) for the daily maximum and minimum time series using multiple reference series around Tianjin from monthly Berkeley Earth (BE), CRUTS4.03 and GHCNV3 data. Using neighboring daily series the record has been homogenized with Quantile Matching (QM) adjustments. Based on the homogenized dataset, the warming trend in annual mean temperature in Tianjin averaged from the newly constructed daily maximum and minimum temperature is evaluated as  $0.154 \pm 0.013^{\circ}\text{C decade}^{-1}$  during the last 130 years. Trends of temperature extremes in Tianjin are all significant at the 5% level, and have much more coincident change than those from the raw, with amplitudes of  $-1.454 \text{ d decade}^{-1}$ ,  $1.196 \text{ d decade}^{-1}$ ,  $-0.140 \text{ d decade}^{-1}$  and  $0.975 \text{ d decade}^{-1}$  for cold nights (TN10p), warm nights (TN90p), cold days (TX10p) and warm days (TX90p) at the annual scale. The adjusted daily maximum, minimum and mean surface air temperature dataset for Tianjin city presented here is publicly available at <https://doi.org/10.1594/PANGAEA.924561> (Si and Li, 2020).

## 51    **1    Introduction**

52    Instrumental observation records at meteorological stations are the most widely used first-hand  
53    information about weather and climate change and variability. They have the advantages of better  
54    representativeness as well as accuracy compared to other data (Leeper et al., 2015; Dienst et al., 2017; Xu  
55    et al., 2018). The most representative long-term observational temperature datasets in the world since  
56    IPCC AR5 (2013) includes: Global Historical Climatology Network(GHCN)-monthly(GHCNm) dataset  
57    (Lawrimore et al., 2011; Menne et al., 2018), Climatic Research Unit (CRU) datasets (Jones et al., 2012;  
58    Harris et al., 2020), Goddard Institute for Space Studies (GISTEMP) dataset (Hansen et al., 2010; Lenssen  
59    et al., 2019) and Berkeley Earth surface temperature (BE) (Rohde et al, 2013; Rohde and Hausfather,  
60    2020). Recently, in order to make up for the limited coverage and the potential regional variability of data  
61    quality of current global climate datasets, Xu et al. (2018) has developed a new dataset of integrated and  
62    homogenized global land surface air temperature-monthly (C-LSAT). This has been updated to C-  
63    LSAT2.0, with the data extended to the period 1850-2019 (Li et al., 2020a; 2021). These datasets were all  
64    developed at the monthly scale based upon meteorological station records from different continents over  
65    the world through the integration of different data sources, quality control of climate outliers, time and  
66    space consistency, and the analysis of data homogenization. The Global Historical Climatology Network-  
67    daily (GHCNd) dataset has also been developed, to meet the needs of climate analysis and monitoring  
68    research, but about two-thirds of the stations contributing to this dataset report precipitation only. In  
69    addition, GHCNd dataset has not been homogenized for artifacts due to changes in reporting practice at  
70    different times at particular stations (i.e., systematic biases), although the entire dataset has been quality  
71    controlled (Menne et al., 2012).

72        Chinese scholars, since the 1980s, have also carried out many studies on the establishment of long-  
73    term observational time series in China, but they often mainly used tree rings, ice cores, historical materials  
74    and other proxy data as part of the restoration of time series before the 1950 (Wang et al., 1998; 2000;  
75    Zheng et al., 2015; Yu et al., 2018). The results based on these data are of great significance as they reveal

76 the characteristics of climate periodicity and multi-scale changes over the past hundred years, but they are  
77 insufficient to meet the needs of quantitative monitoring and detection of long-term extreme climate events.  
78 In particular, there are many limitations when homogenizing the time series before the 1950s (including  
79 the establishment of reference series) such as the lack of continuous observational data, detailed and  
80 reliable metadata information, leading to the increase of uncertainties for regional and/or local climate  
81 analysis (Li et al., 2020b). As a result there still exist many uncertainties in the characteristics of climate  
82 change from the 19th century to the mid-20th century (Li et al., 2010; 2017; Sun et al., 2017; Cao et al.,  
83 2017).

84 Since daily time series generally contain many more observations than monthly or annual series,  
85 daily analyses potentially have greater precision. As a result they are more useful in climate trend and  
86 variability studies, especially for extreme events (Vincent et al., 2012; Xu et al., 2013; Trewin, 2013;  
87 Hewaarachchi et al., 2017). However, due to difficulties in collecting and/or receiving daily data all over  
88 the world as well as non-climatic effects such as changes in observation times, there are numerous issues.  
89 For example, observations from temperature sites at principal stations in Canada were changed to be read  
90 at 0000UTC to 0600UTC (Vincent et al., 2002), making it is very difficult to form a global daily data  
91 product at century-long scales. This makes it extremely difficult to study global and/or regional extreme  
92 events over the past hundred years, especially before 1950. For some regional areas, daily instrumental  
93 observations may be extended to the 19th century and hence they are more valuable. Png et al. (2020) has  
94 compiled 463,530 instrumental observations of daily temperature, precipitation and sunshine from 319  
95 stations distributed over China during 1912-1951 mostly from the source of monthly reports from the  
96 Institute of Meteorology, Nanjing; observatories over Japan-occupied Manchuria and the Japanese Army  
97 for North China. Since this is a daily data, it is immensely useful for the analysis of mesoscale and sub-  
98 seasonal climate variations. Although the earliest instrumental observations in China began in the 1840s,  
99 observations at some sites were interrupted during 1940s due to wars (e.g. the War of Resistance Against  
100 Japan and the War of Liberation) and hence many pieces of information have likely been lost. Studies of

101 the rescuing, processing and constructing complete and continuous daily site data over China are  
102 somewhat rare.

103 Due to the historic reasons of leased territory in China, some local single sites often have multiple  
104 observational sources before 1950. For example, for Qingdao, monthly surface air temperature series  
105 during 1899 - 2014 have been constructed based on newly digitized and homogenized observations from  
106 the German National Meteorological Service from 1899 to 1913 (Li et al., 2018). Tianjin meteorological  
107 station is one of the typical stations with more than one hundred years of observed climate data in China  
108 (Yan et al., 2001; Si et al., 2017). This station also has multi-source observations before 1950 as observed  
109 at some other meteorological stations in China having century-long datasets. Thus, considering Tianjin  
110 station as an example, this paper aims to construct a new daily instrumental maximum and minimum  
111 temperature series on the century scale in China, through integration, quality control, extension and  
112 homogenization of the multiple daily observations. The newly constructed daily temperatures in Tianjin  
113 provide relatively longer, more complete and reliable climate series for studies of climate and extreme  
114 climate change over century-long scales.


115 The remainder of this paper is arranged as follows: Section 2 describes the station histories from  
116 Tianjin. The basic data and reference data sources and their pre-processing are introduced in section 3.  
117 Section 4 introduces the procedures of constructing new daily maximum and minimum temperature series.  
118 Section 5 presents average and extreme temperature trend change based on newly constructed series. The  
119 availability of the resulting dataset (Si and Li, 2020) is reported in section 6, and summary of results and  
120 some discussion are given in section 7.

## 121 **2 Historical evolution of Tianjin meteorological observation station**


122 Wu (2007) showed that Tianjin meteorological observation station was under the control of the  
123 Department of Industry Agency of the British Concession in Tianjin covering September 1887 to  
124 December 1941. During the period from September 1904 to December 1949, it was co-ordinated by many

125 departments, such as Japan Central Meteorological Station, Central Meteorological Bureau of the  
126 Republic of China, Aviation Department of North China Military Region of the People's Liberation Army  
127 (PLA), Shunzhi Water Conservancy Commission, Water Conservancy Commission of North China and  
128 Water Conservancy Engineering Bureau of North China. However, only the records of daily maximum  
129 and minimum temperatures from Department of Industry Agency of the British Concession in Tianjin  
130 (Fig.1a-b) and Water Conservancy Commission of North China (Fig.1c) have been collected by the Tianjin  
131 Meteorological Archive. Each of these is continuous and complete, and most importantly they can be  
132 connected to each other on 1 Jan 1932 without overlapping, thereby forming a complete and continuous  
133 time series before 1950.

134



135



136 **Figure 1.** The handwritten observation records of Tianjin before 1950, (a) and (b) for records of maximum and minimum  
137 temperatures from Department of Industry Agency of British Concession in Tianjin, (c) for records from Water  
138 Conservancy Commission of North China, and Tianjin Observatory at No. 22 Ziyou Road (d). These photographs were  
139 provided by Tianjin Meteorological Archive, Tianjin Meteorological Bureau.

140       The daily temperature records at Tianjin meteorological observation station that we have accessed  
141 begin with Sep 1 1890 collected by Tianjin Meteorological Archive (Fig.1a-c). The metadata history of  
142 the Tianjin observation station during Sep 1 1890 - Dec 31 2019 is listed in Table 1. The history is sorted  
143 according to the Surface Meteorological Observation Specifications in China (versions of 1950, 1954,  
144 1964, 1979, and 2003), metadata of China surface meteorological station and journal sheets of Tianjin  
145 surface meteorological records. Changes to observational times have been marked on the original time  
146 series of maximum and minimum temperatures for Tianjin city (see Fig. 2). As shown in Table 1, Tianjin  
147 observation station has relocated four times since 1890, several times in 1921 (Fig. 1d), Jan 1 1955, Jan 1  
148 1992, and Jan 1 2010 without prominent changes in elevations. The environment surroundings of Tianjin  
149 station changed from urban to suburban in 1955, accompanied by a number of instrument changes. In this  
150 period, changes to the instrument manufacturer have happened four times for both maximum and  
151 minimum temperature series, as well as changes of automatic observation instead of manual observation  
152 in 2004 and a new generation replacement of last automatic instrument in 2014. In documented metadata  
153 (Table 1), there have been changes of observing time four times for both maximum and minimum  
154 temperatures since 1951, but they were always recorded over a 24-hour observational window at Beijing  
155 Time (BT) or similar to BT (as for the period of Jan 1 1954 - Dec 31 1960). Moreover, it is important to  
156 mention here that since 1951 there were two stations viz., Tianjin and Xiqing collocated in the Tianjin  
157 area. Due to rapid urbanization at the surrounding environmental Tianjin area, the old Tianjin site gradually  
158 becomes less representative as a climate observation station and therefore since Jan 1 1992 afterwards,  
159 observations at Xiqing station are used to replace the old Tianjin station. This can also be considered as

160 Tianjin station being relocated to Xiqing station since then.


161 **Table 1.** The history logs at Tianjin meteorological observation station during Sep 1 1890 - Dec 31 2019.


Periods	Latitude	Longitude	Altitude (m)	Address (Surrounding environment)	Relocation description	Instrument change	Observing time
1890.9.1— 1921	39°07’	117°12’	unspecified	unspecified	——	unspecified	unspecified
1921— 1950.12.31	39°08’	117°11’	6.0	No. 22 Ziyou Road, The Third Distribution, Tianjin (urban)	unspecified	unspecified	unspecified
1951.1.1— 1953.12.31	39°08’	117°11’	6.0	Same as above	——	——	Tmax 18:00 Tmin 09:00
1954.1.1— 1954.12.31	39°08’	117°11’	6.0	Same as above	——	Tmax 1954.1.1 Tmin 1954.1.1	unspecified
1955.1.1— 1960.12.31	39°06’	117°10’	3.3	Zunyi Road, Hexi Distribution, Tianjin (suburban)	5 km north of the original site	——	unspecified
1961.1.1— 1991.12.31	39°06’	117°10’	3.3	Qixiangtai Road, Hexi Distribution, Tianjin (suburban)	——	Tmax 1961.1.1; 1973.1.1; 1989.1.1 Tmin 1961.1.1; 1966.1.1; 1989.1.1	20:00



Xidawa, Xiqing							
1992.1.1— 2003.12.31	39°05′	117°04′	2.5	Distribution, Tianjin (suburban)	unspecified	——	20:00
2004.1.1— 2009.12.31	39°05′	117°04′	2.5	Same as above	——	automatic observation	pick up from timing minutes data
2010.1.1— 2013.12.31	39°05′	117°03′	3.5	Jingfu Road, Xiqing Distribution, Tianjin (suburban)	1.5km southwest of the site in 1992	automatic observation	pick up from timing minutes data
2014.1.1 to now	39°05′	117°03′	3.5	Same as above	——	new generation of automatic observation equipment	pick up from timing minutes data

162 The straight line (——) indicates no change. The observing time is at Beijing Time (BT). Tmax and Tmin indicate the  
163 maximum and minimum temperature, respectively.





**Figure 2.** Original series of daily maximum temperature (a) and minimum temperature (b) at Tianjin meteorological observation station covering Sep 1 1890 - Dec 31 2019. Black stars with vertical dashed lines on the axes mark metadata times (The relocation point in 1921 is marked by green star with vertical dashed line due to no specific date).

### 3 Data sources

#### 3.1 Original data and preliminary quality control

Based on the analyses of metadata history at Tianjin station discussed in section 2, this paper selects three observation temperature data sources collected by Tianjin Meteorological Archive as the basic data to construct the daily maximum and minimum temperature time series at Tianjin station from 1890 (Fig. 2). These are the daily observation records from (1) Department of Industry Agency of British Concession in Tianjin covering Sep 1 1890 - Dec 31 1931; (2) Water Conservancy Commission of North China during Jan 1 1932 - Dec 31 1950; and (3) monthly journal sheets of Tianjin surface meteorological observation records covering Jan 1 1951 - Dec 31 2019. Since there are no missing data and overlap for each of the three daily sources, the three daily data resources are directly spliced into a complete time series. However, in view of the regime changes of operation and different station numbers between Tianjin station and Xiqing station that happened at Jan 1 1992 (Table 1), the daily records from Jan 1 1951 to Dec 31 1991 observed in the Hexi area and those observed at Xiqing area from Jan 1 1992 to Dec 31 2019 are used to form the basic daily time series from Jan 1 1951 to Dec 31 2019 for the long series.

A preliminary quality control procedure consisting of multiple steps was carried out on the original integrated daily time series of maximum and minimum temperature from Sep 1 1890 to Dec 31 2019 at Tianjin to remove any errors caused by manual observations, instrument malfunctions and digital inputs. Firstly, the range of the daily maximum or minimum temperature was scrutinized for magnitude beyond the limit of 60 °C and -80 °C as errors. Fortunately, both the series have no such error. Secondly based on anomalies from the 1961-1990 as reference period, climatic outliers of maximum and minimum temperature are assessed considering a magnitude exceeding five standard deviations of their monthly anomalies as outliers during 1890-2019. There is no outlier found in our validation. Finally, internal consistency is investigated by checking if there is any minimum temperature data greater than or equal to the maximum at the same date and no such inconsistencies were found. It is important to mention that there is a sudden rise in annual minimum temperature series during the year 1927 even after these three checks. The offsets of the discontinuities in 1927 compared with averages for the two sections before and after it are 4.2 °C and 3.4 °C, respectively. We repeated the steps of outlier checks once by reviewing the earlier condition with three times the standard deviation of monthly anomalies. Results indicate that most of the daily minimum data for April to October 1927 exceed the current condition and finally the data during this period so were set to missing values. Even though, the quality of original daily maximum and minimum temperatures during 1890 - 2019 at Tianjin station is good, these checks provide a good foundation for the subsequent construction of a reliable homogenized daily series.

### **3.2 Reference data**

Wu (2007) documented that although the earliest surface observation records at Tianjin station start with the year 1887, the observed daily maximum and minimum temperatures collected by Tianjin Meteorological Archive began with September 1890 (Fig. 2). Therefore, some additional reference data



205 sources are selected to extend the daily temperature series from January 1887 to August 1890 and lengthen  
 206 the established daily temperature data to as early as possible. In addition, it is extremely important to  
 207 establish an objective as well as a reasonable reference series for data homogenization. But due to non-  
 208 availability of observation records and station metadata before 1950 especially for daily data, it is  
 209 impossible to find a complete and reliable observed temperature series as a reliable reference series for  
 210 Tianjin. Based on few recently reported studies (Li et al., 2020a; Lenssen et al., 2019; Xu et al., 2018; and  
 211 Menne et al., 2018), we employ the station series or the interpolated temperature series using neighboring  
 212 grid boxes from three global land surface temperature observation series (Table 2) as reference data  
 213 sources for extension and establishment of reference data series used in data homogenization at Tianjin  
 214 station. Plots of the ‘Tianjin’ station from all three series are shown in Figure 3. The three global Land  
 215 Surface Temperature (LSAT) are (1) Berkeley Earth land temperature (BE; Rohde and Hausfather, 2020;  
 216 <http://berkeleyearth.org/data/>); (2) Climatic Research Unit (CRU) Time-Series (TS) version 4.03  
 217 (CRUTS4.03; Harris et al., 2020; [http://data.ceda.ac.uk/badc/cru/data/cru\\_ts/cru\\_ts\\_4.03/data/](http://data.ceda.ac.uk/badc/cru/data/cru_ts/cru_ts_4.03/data/)) and (3)  
 218 Global Historical Climatology Network (GHCN) version3 (GHCNV3; Lawrimore et al., 2011;  
 219 <https://www.ncdc.noaa.gov/ghcnd-data-access>).

220 The selected three LSAT are not independent as they likely use common input observations. The  
 221 multiple datasets provides a variety of useful checks because they employ different ways of handling data  
 222 problems such as incomplete spatial and temporal coverage and non-climatic influences on meteorological  
 223 measurement (Hansen et al., 2010). As shown in Table 2, the three LSAT involve quality control and  
 224 homogenization but using different methods. The records of BE were usually split into portions occurring  
 225 before and after known and presumed discontinuities (e.g., from station relocation or instrument changes)  
 226 without adjustment. For CRUTS4.03, most of these data have been adjusted, because the ultimate sources



227 of most station records are from National Meteorological Services (NMSs), so China Meteorological  
 228 Administration (CMA) for Tianjin. GHCNV3 station data used in this paper are the quality controlled and  
 229 adjusted (QCA) data, which were produced by the developers of GHCN-Monthly. Two types of grid data,  
 230 CRUTS4.03 and BE are both interpolated to the site level using the bilinear method.







231 From Table 2, only BE-daily maximum/minimum temperatures are available. So the maximum  
 232 temperature BE-daily data corresponding to the site level Tianjin station is selected as the extension data  
 233 for the period of Jan 1 1887-Aug 31 1890 (The specific information about extending the processing for  
 234 this period is in supplemental material S1-S3), and the daily minimum series still begins with the date on  
 235 Sep 1 1890 due to scarcity of reference data sources.

236



237






**Figure 3.** The annual and seasonal average anomalies of maximum (a-e) and minimum (f-j) temperatures based on the interpolated series of BE and CRUTS4.03 and the station series of GHCNV3 for Tianjin station.

**Table 2.** Information of reference data sources.

Data sources	Monthly	Daily	Gridded	Station	Temporal resolution	Spatial resolution	Time periods	Units	Quality control	Adjustment
CRUTS4.03	√	×	√	√	monthly	0.5°×0.5°gridded	1901.1-2018.12	°C	√	√
BE-monthly	√	√	√	√	monthly	1°×1°gridded	1872.12-2019.12	°C		
BE-daily	√	√	√	×	daily	1°×1°gridded	Tmax1880.1-2018.12 / Tmin1903.1-2018.12	°C	√	×
GHCNV3	√	√	√	√	monthly	station data	1904.1-1990.12	°C	√	√

#### 4 Construction of daily maximum and minimum temperature series from 1887 to 2019

On the basis of the quality controlled and the extended series, the daily homogenized maximum and minimum observation temperature series in Tianjin were constructed by means of the flow chart illustrated in the dashed-box part of Fig. 4. Homogenization is an important step to eliminate the discontinuities in observation records induced by non-climatic influences such as station relocation, instrument change, observing time change and so on. Most importantly, the true characteristics of climate change are preserved in this process (Quayle et al., 1991; Della-Marta and Wanner, 2006; Haimberger et al., 2012; Rahimzadeh and Zavareh, 2014; Hewaarachchi et al., 2017).



253 **Figure 4.** The flow chart of construction for century-long daily maximum and minimum temperature series in Tianjin.

#### 254 **4.1 Establishment of the reference series**

255 In the process of homogenization, reasonable reference series plays an important role in the reliability of  
256 the detected breakpoints. So in this section, we will establish monthly and daily reference series for the  
257 maximum or minimum temperature series at Tianjin station and use them for breakpoint detection and  
258 adjustment, respectively. Both reference series are established using a weighted average method. For  
259 monthly reference series, we will establish two types in order to make the detected breakpoints more  
260 reasonable and reliable. First, reference data are based on the combination of the interpolated temperature  
261 series from BE-monthly and CRUTS4.03 and station series from GHCNV3 data for Tianjin site (the three  
262 global LSAT datasets) and secondly based on the interpolated temperature series from BE-monthly data  
263 only. From the three LSAT data, the weight coefficients are the square of the correlation coefficients  
264 between each LSAT and Tianjin's observed data. The daily reference series we use is based on the  
265 interpolated temperature series from BE-daily data only.

266 In the case of the interpolated temperature series from BE-monthly or daily data only, the site level  
267 data are derived from the station network across the Beijing-Tianjin-Hebei area in China (Fig. 5). These  
268 stations are selected as follows: Firstly, the potential stations less than 300km at horizontal distances  
269 around Tianjin station and with altitude differences within 200m are chosen; Secondly, we will select 10  
270 stations those are closest to Tianjin station using a spherical distance; Finally, 9 stations are confirmed  
271 which are consistent between step 1 and step 2. In Figure 5 (the right), these 9 are identified by green solid  
272 circles with black or red stars, some metadata are also provided in Table 3. Thus, the interpolated  
273 temperature series from BE reference series are generated using the weighted average of the 9 stations.  
274 These weights are calculated as the square of correlation coefficients between the interpolated temperature




series from BE-monthly or daily data for each 9 stations and Tianjin's observed data. Recall also that, the missing values in the original daily minimum temperature at Tianjin for April to October 1927 (checked in section 3.1) were replaced by the corresponding data from the weighted 9 interpolated temperature series from BE-daily data.

**Table 3.** Metadata information of the 9 stations used to build reference series.

Station number	Latitude	Longitude	Altitude (m)	Surroundings
54517	39°04'	117°12'	2.2	urban
54518	39°07'	116°23'	9.0	rural
54526	39°05'	117°20'	1.9	urban
54528	39°14'	117°08'	3.4	rural
54530	39°13'	117°46'	0.5	rural
54619	38°55'	116°55'	5.5	urban
54622	38°57'	117°25'	1.5	rural
54623	39°03'	117°43'	4.8	urban

280




**Figure 5.** Geographical distribution of the surface weather stations (black solid circles) at Beijing-Tianjin-Hebei area in China and the selected 9 stations (green solid circles with black or red stars).

## 283 4.2 Breakpoints detection and adjustment

284 The RHtestsV4 software package is used to homogenize the daily maximum and minimum temperature  
285 data at Tianjin station. The software consists of the Penalized Maximal T-test (PMT) (Wang et al., 2007)  
286 and Quantile Matching (QM) adjustment (Wang et al., 2010; Bai et al., 2020; Lv et al., 2020), both of  
287 which are used to detect and adjust the known or presumed discontinuities. As observed in earlier reported  
288 studies (Vincent, 2012; Trewin, 2013; Xu et al., 2013), homogenization at the daily timescales is much  
289 more challenging than that at monthly or annual scales. Thus, firstly we test Tianjin's monthly observed  
290 maximum and minimum temperature series averaged from the daily ones to find the significant  
291 breakpoints by means of PMT at the 5% significance level using two types of monthly reference series.  
292 We then adjust the daily series at Tianjin station by QM-adjustment with or without the daily reference  
293 series.


294 The breakpoints in the segment before 1921 are mainly determined by objective judgment from the  
295 same shifts at the two monthly timescales simultaneously due to the scarcity of station metadata. Those  
296 after 1921 are additionally assessed together with the station metadata and PMT detection at the 5%  
297 significance level. According to Table 1, we made a list containing some possibilities that could cause  
298 shifts in Tianjin's daily maximum and minimum temperature series (Fig. 2 vertical dashed lines). The date  
299 of Jan 1 1932 and Jan 1 1951 are connection points for different data sources, 1921, Jan 1 1955, Jan 1  
300 1992 and Jan 1 2010 indicate station relocations, and the others are the times of instrument and/or  
301 observing time change. However, due to statistical non-significance, those potential discontinuities are not  
302 considered as the final discontinuities (Fig. 6). So potential dates do not always cause artificial  
303 discontinuities at the joining of the three observation segments for daily maximum or minimum  
304 temperature series. Also all the instrument changes that happened for maximum and minimum series have

also not introduced any significant shifts. In this regard, they do not look like the changes that happened with other networks around the world, such as the U.S. Cooperative Observer Program (COOP) network (Leeper et al., 2015). Moreover, different observing times (including automated observation system) also do not introduce any significant biases to the temperature time series, since the daily maximum and minimum temperatures are always recorded over a 24-hour observational window. Additionally, various versions of the surface meteorological observation specifications in China (e.g. versions of 1950, 1954, 1964, 1979 and 2003) imply that the observation principles of the highest and lowest thermometers are consistent, although there were a number of alterations of observing times.



**Figure 6.** QM-adjusted and raw series (after quality control and extension) of daily maximum temperature (a) and minimum temperature (b) at Tianjin meteorological observation station covering Jan 1 1887 (Sep 1 1890 for minimum)

317 to Dec 31 2019 (Rohde and Hausfather, 2020). Vertical solid lines demarcate the discontinuities at times Mar 1 1897, Jun  
 318 1 1907, Jan 1 1955 and Jan 1 1992.



319  
 320 **Figure 7.** The amplitudes of QM adjustment applied to daily maximum (a) and minimum (b) temperature data at Tianjin  
 321 meteorological observation station.


322 **Table 4.** The average QM adjustments at the monthly timescales applied to daily maximum and minimum temperature  
 323 data at Tianjin meteorological observation station (Units: °C).

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Maximum temperature	1.136	0.246	-0.616	-0.687	-1.322	-2.484	-2.817	-2.285	-1.046	-0.590	-0.582	0.566
Minimum temperature	-0.105	-0.297	-0.634	-0.979	-1.084	-1.090	-1.207	-1.184	-1.050	-0.951	-0.624	-0.317

324 Breakpoints in Mar 1 1897 and Jun 1 1907 are without metadata support, but those at Jan 1 1955 and  
 325 Jan 1 1992 are confirmed by metadata of station relocation. In Jan 1 1955 Tianjin meteorological station  
 326 was relocated from No. 22 Ziyou Road to Zunyi Road, where is only 5km north of the original site, as  
 327 well as being more open and clear, and in Jan 1 1992 it was relocated from Qixiangtai Road, Hexi District  
 328 to Xidawa, Xiqing District (Table 1). The significant breakpoints are given in Fig. 6 as vertical solid lines.

329 The amplitudes of QM adjustment applied to each individual daily maximum and minimum  
 330 temperature data are  $[-4.606, 2.621^{\circ}\text{C}]$  (Fig. 7a) and  $[-5.972, 1.897^{\circ}\text{C}]$  (Fig. 7b). The medians of QM

adjustment are  $-0.764^{\circ}\text{C}$  and  $-0.506^{\circ}\text{C}$  respectively. As shown in Fig. 7, there are about 75% of adjustments are covering  $-2.5 \sim 0.8^{\circ}\text{C}$  in daily maximum series. For the minimum ones, there are about 85% of adjustments are covering  $-0.8 \sim 0.5^{\circ}\text{C}$ . Table 4 provides the average amplitudes of QM adjustment at the monthly timescales. It shows that for the maximum data, the larger positive adjustments are mainly applied to series in January and December, while the larger negative adjustments are mainly in June, July and August. For the minimum data, all the average amplitudes of QM adjustment at the monthly timescales are negative, what is the same characteristic with the maximum ones, temperature series in the warm months (e.g. June, July and August) are adjusted with larger negative amplitudes, but smaller negative amplitudes for the series in the colder months (e.g. January, February and December).




**Figure 8.** Annual average series of maximum (a) and minimum (b) temperatures in Tianjin from 1887 (1891 for minimum) to 2019 based on newly constructed and raw daily data (after quality control and extension), correspondingly with annual


344 averaged data based on the interpolated series from BE-monthly (1873-2019) and CRUTS4.03 (1901-2018) and station  
 345 series from GHCNV3 (1905-1990) for Tianjin station.

346 The average annual maximum and minimum temperature series based on the adjusted daily data (the  
 347 red lines) in Tianjin are given in Fig. 8. The raw time series for Tianjin (after quality control and extension),  
 348 the interpolated series from BE and CRUTS4.03 and station series from GHCNV3 for Tianjin station  
 349 averaged from their monthly data are also displayed in Fig. 8. This shows that the newly constructed time  
 350 series has removed the large shifts in maximum and minimum temperature series before 1955 and 1992  
 351 (Fig. 8 red lines) compared with the raw ones (Fig. 8 black lines). Especially for the minimum data, the  
 352 QM-adjustments have offset the shifts between 1896 and 1908 to the greatest extent. Meanwhile, the  
 353 newly constructed temperature data has similar inter-annual variability and trend changes compared to  
 354 those of BE, CRUTS4.03 and GHCNV3 during the overlapping period.

355 **Table 5.** Definition of temperature extremes (Zhang et al., 2011). Days with maximum or minimum temperature above  
 356 the 90th and below the 10th percentiles are relative to the reference period of 1961 - 1990.

Index	Description	Definition	Units
TN10p	Cold nights	Days when daily minimum temperature <10th percentile	days
TN90p	Warm nights	Days when daily minimum temperature >90th percentile	days
TX10p	Cold days	Days when daily maximum temperature <10th percentile	days
TX90p	Warm days	Days when daily maximum temperature >90th percentile	days





**Figure 9.** Annual and monthly temperature extremes of Cold nights (TN10p) (a), Cold days (TX10p) (b), Warm nights (TN90p) (c) and Warm days (TX90p) (d) for daily newly constructed and raw temperatures (after quality control and extension) in Tianjin.

Table 5 provides the definition of temperature extremes (Zhang et al., 2011). They are calculated based on the newly constructed and raw series (after quality control and extension) in Tianjin. As shown in Fig. 9, the number of TN10p (Fig. 9a) and TX10p (Fig. 9b) at the monthly timescales are increased by 0.3-2.0 days and 0.6-2.3 days based on the newly constructed series, especially in August they are increased by 2.0 and 2.3 days respectively. TX10p in cold months (January, February and December) from newly data are all less than those from the raw data, with the number of days decreased by 3.1-6.1. This is mainly due to large positive adjustments applied to daily maximum temperature in these months (Table 4). In the opposite sense, the number of TN90p (Fig. 9c) and TX90p (Fig. 9d) are decreased by 0.2-3.8 days and 0.9-14.8 days respectively based on the newly constructed series. The decreasing numbers of the two indices between May and September are prominent, especially for TX90p from June to August the number is decreased by 11.8-14.8 days. This is due to the large negative adjustments applied to daily maximum temperature in these months (Table 4). The number of TN10p (Fig. 9a) from newly constructed series at the annual timescales are increased by 1.1 days compared to the raw ones while for TX10p (Fig. 9b), TN90p (Fig. 9c) and TX90p (Fig. 9d) are decreased by 0.3, 2.1 and 5.7 days respectively.

376 **5 The temperature change trend in Tianjin based on newly constructed series**

377 **5.1 Mean temperature trend during the last 130 years**

378 **Table 6.** Comparisons between newly constructed surface air temperatures and previous assessments of the annual trend  
379 change at the century-long scale in Tianjin with uncertainties at 95% significance level (Units: °C decade<sup>-1</sup>).

	Newly constructed	BE	CRUTS4.03
	1887(1891)-2019	(1873-2019)	(1901-2018)
Maximum temperature	0.119±0.015	0.099±0.010	0.062±0.015
Minimum temperature	0.194±0.013	0.156±0.010	0.217±0.015
Mean temperature	0.154±0.013	0.128±0.009	0.140±0.013

380 Table 6 shows that the annual trends of newly constructed maximum (1887-2019) and minimum  
381 temperature (1891-2019) series in Tianjin are 0.119±0.015°C decade<sup>-1</sup> and 0.194 ±0.013°C decade<sup>-1</sup>. Trend  
382 changes based on the newly constructed series are nearly consistent with those in BE and CRUTS4.03 on  
383 the century-long scale and these are 0.099±0.010°C decade<sup>-1</sup> and 0.156±0.010°C decade<sup>-1</sup>, 0.062±0.015°C  
384 decade<sup>-1</sup> and 0.217±0.015°C decade<sup>-1</sup> respectively. The trend of the mean temperature for the newly  
385 constructed series (0.154±0.013°C decade<sup>-1</sup>) is slightly larger than those from the interpolated series from  
386 BE, CRUTS4.03, and Cao et al. (2013) (0.128±0.009 °C decade<sup>-1</sup>, 0.140±0.013 °C decade<sup>-1</sup>, and  
387 0.098±0.017 °C decade<sup>-1</sup>, respectively). Moreover, annual trend change in mean temperature based on the  
388 newly constructed series at Tianjin is also a little larger than that over the whole China (Li et al., 2020b),  
389 which are 0.130±0.009°C decade<sup>-1</sup>, 0.114±0.009°C decade<sup>-1</sup> and 0.121±0.009°C decade<sup>-1</sup> respectively  
390 from CRUTEM4, GHCNV3 and C-LSAT (during 1900 - 2017). It conforms to the underlying changes  
391 across China. Increasing trends in northern China are more prominent than those from other regions in  
392 mainland China (Li et al., 2004; Zhai et al., 2004).

393 **5.2 Extreme events change trend during the last 130 years**



394 **Table 7.** Trends in annual and seasonal temperature extremes of Cold nights (TN10p), Cold days (TX10p), Warm nights  
 395 (TN90p) and Warm days (TX90p) for daily newly constructed temperatures in Tianjin (Units: d decade<sup>-1</sup>).

	TN10p	TX10p	TN90p	TX90p
Annual	-1.454*	-0.140*	1.196*	0.975*
Spring	-1.861*	-0.508*	1.423*	0.959*
Summer	-1.483*	-0.213*	1.443*	1.474*
Autumn	-0.798*	-0.221*	0.724*	0.621*
Winter	-1.555*	0.421*	1.119*	0.850*

396 The asterisks indicate trend changes are significant at the 5% level. The annual time periods of TN10p and TN90p cover  
 397 the 1891-2019 period, TX10p and TX90p cover 1887-2019. The time periods of TN10p and TN90p in spring and summer  
 398 cover 1891-2019, those in autumn and winter 1890-2019 (winter ending in 2018), and TX10p and TX90p in four seasons  
 399 all cover 1887-2019 (winter ending in 2018).

400 Table 7 illustrates that trends of temperature extremes based on the newly constructed series are all  
 401 significant at the 5% level and have much more coincident changes. The cold extremes (TN10p and TX10p)  
 402 at annual and seasonal timescales express significantly decreasing trends (except for TX10p in winter),  
 403 while the warm extremes (TN90p and TX90p) show increasing trends. Trends of TN10p, TX10p, TN90p  
 404 and TX90p at the annual timescales are -1.454 d decade<sup>-1</sup>, -0.140 d decade<sup>-1</sup>, 1.196 d decade<sup>-1</sup> and 0.975  
 405 d decade<sup>-1</sup>, all pass the significance test at the 5% level. For the seasonal change, the negative trends of  
 406 TN10p and TX10p in spring are the largest, reaching up to -1.861 d decade<sup>-1</sup> and -0.508 d decade<sup>-1</sup> during  
 407 the past 130 years. For TN90p and TX90p is in summer, with the amplitude of 1.443 d decade<sup>-1</sup> and 1.474  
 408 d decade<sup>-1</sup>.

## 409 6 Data availability

410 The newly homogenized daily surface air temperature for Tianjin city over century-long scales are  
 411 published at PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.924561>, last access: 17 April 2021)  
 412 under the DOI <https://doi.org/10.1594/PANGAEA.924561> (Si and Li, 2020). The dataset contain the  
 413 maximum, minimum and mean temperature time series before and after adjustment as well as new

estimates of average and extreme temperature trend change in Tianjin for the period of 1887-2019.

## **7 Conclusions and discussion**

This paper documents the various procedures necessary to construct a homogenized daily maximum and minimum temperature series since 1887 for Tianjin. These similar procedures could and should be used for other sufficiently long and complete series across the world. The newly constructed data have reduced non-climatic errors at the daily timescale, improved the accuracy and enhanced the real climate representation of average and extreme temperature over century-long scales.

Three sources of surface observation daily data collected by the Tianjin Meteorological Archive have been integrated viz., Department of Industry Agency of British Concession in Tianjin covering Sep 1 1890-Dec 31 1931, Water Conservancy Commission of North China covering Jan 1 1932-Dec 31 1950, and monthly journal sheets of Tianjin surface meteorological observation records covering Jan 1 1951-Dec 31 2019. These three have provided a good foundation for the construction of reliable homogenized daily series by quality control of climatic range checks, climatic outlier checks and internal consistency checks. Data extension has been undertaken in the interest of extending the length of the series as far back as possible, but it is carried out only for the daily maximum series due to length limitation of reference daily data.

Using the integration, quality control and extension, we detected and adjusted the statistically significant breakpoints in the daily maximum and minimum temperature time series from an objective perspective based on multiple reference series and statistical characteristics from homogenization detection by means of PMT as well as sophisticated manual data processing. This temperature series provides a set of new baseline data for the field of extreme climate change over the century-long scale and a reference for construction of other long-term reliable daily time series in the region. The annual trends

of newly constructed maximum and minimum temperature in Tianjin are  $0.119 \pm 0.015^{\circ}\text{C decade}^{-1}$  and  $0.194 \pm 0.013^{\circ}\text{C decade}^{-1}$  over the last 130 years, which are similar to those from Berkeley and CRUTS4.03. The trend of mean temperature averaged from the new series is  $0.154 \pm 0.013^{\circ}\text{C decade}^{-1}$ , which is of the same order as those over the whole China (Li et al., 2020a; 2020b). The new daily data also show improvements over the archived datasets for trend analyses of extremes. The trends in TN10p, TX10p, TN90p and TX90p are all significant at the 5% level, and they give a much more consistent set of trends. To some extent, changes in climate extremes can be analyzed with higher confidence using the newly constructed daily data in this paper.

However, in the current study, there may be some systematic biases (possibly some potential breakpoints missed) still in the adjusted time series, since metadata of Tianjin station are not consistently available in the climatological archives over the whole century as well as not being documented during the period before 1921. Climate data homogenization does not always follow a consistent pattern (Si et al., 2018; 2019). It is necessary to constantly improve the existing methodology and explore new techniques in order to obtain reliable homogenized data products. Accordingly, future work should involve more detailed station metadata and more advanced data processing techniques to produce much better daily datasets over century scales.

**Author contributions.** QL designed and implemented the dataset construction. PS collected the basic and reference data sources, constructed the dataset and written the paper. QL, PS and PJ all contributed to data analysis, discussion and writing of the paper.

**Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgements.** We thank the many people and /or institutions who contributed to the raw data and

457 metadata collection, data pre-processing and construction of this dataset.

458 **Financial support.** This research has been supported by National Natural Fund projects No. 41905132  
459 and No. 41975105.

## 460 **References**

461 Bai, K. X., Li, K., Wu, C. B., Chang, N. B., and Guo, J. P.: A homogenized daily in situ PM2.5  
462 concentration dataset from the national air quality monitoring network in China, *Earth Syst. Sci. Data*,  
463 12(4): 3067-3080, <https://doi.org/10.5194/essd-12-3067-2020>, 2020.

464 Cao, L. J., Yan, Z. W., Zhao, P., Zhu, Y. N., Yu, Y., Tang, G. L., and Jones P.: Climatic warming in China  
465 during 1901-2015 based on an extended dataset of instrumental temperature records, *Environ. Res. Lett.*,  
466 12(6): 064005, <https://doi.org/10.1088/1748-9326/aa68e8>, 2017.

467 Della-Marta, P. M., and Wanner, H.: A method of homogenizing the extremes and mean of daily  
468 temperature measurements, *J. Climate*, 19(17): 4179-4197, <https://doi.org/10.1175/JCLI3855.1>, 2006.

469 Dienst, M., Lindén, J., Engström, E., and Esper, J.: Removing the relocation bias from the 155-year  
470 Haparanda temperature record in Northern Europe, *Int. J. Climatol.*, 37(11): 4015-4026, <https://doi.org/10.1002/joc.4981>, 2017.

472 Hansen, J., Ruedy, R., and Sato Makiko K. Lo.: Global surface temperature change, *Rev. Geophys.*, 48,  
473 RG4004, <https://doi.org/10.1029/2010RG000345>, 2010.

474 Haimberger, L., Tavalato, C., Sperka, S.: Homogenization of the global radiosonde temperature dataset  
475 through combined comparison with reanalysis background series and neighboring stations, *J. Climate*,  
476 25(23): 8108-8131, <https://doi.org/10.1175/JCLI-D-11-00668.1>, 2012.

477 Hewaarachchi, A. P., Li, Y. G., Lund, R., and Rennie, J.: Homogenization of Daily Temperature Data, *J.*  
478 *Climate*, 30, 985-999, <https://doi.org/10.1175/JCLI-D-16-0139.1>, 2017.

479 Huang, J. Y., Liu, X. N., Li, Q. X.: The experimental study of reconstruction for summer precipitation and  
 480 temperature in China, *Journal of Applied Meteorological Science*, 15(2): 200-206, 2004(in Chinese).

481 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth  
 482 Assessment Report of the Intergovernmental Panel on Climate Change, Stocker TF, Qin D, Plattner G  
 483 K, TignorM, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. (eds.): 1535 pp. Cambridge  
 484 University Press: Cambridge, UK 25 and New York, NY, 2013.

485 Jones, P. D., Lister, D., Osborn, T. J., Harpham, C., Salmon, M., and Morice C.: Hemispheric and large-  
 486 scale land-surface air temperature variations: an extensive revision and an update to 2010, *J. Geophys.*  
 487 *Res.*, 117, D05127, [https:// doi.org/10.1029/2011JD017139](https://doi.org/10.1029/2011JD017139), 2012.

488 Lawrimore, J. H., Menne, M. J., Gleason, B., Williams, C. N., Wuertz, D. B., Vose, R. S., and Rennie, J.:  
 489 An overview of the Global Historical Climatology Network monthly mean temperature data set, Version  
 490 3, *J. Geophys. Res.*, 116, D19, <https://doi.org/10.1029/2011JD016187>, 2011.

491 Leeper, R. D., Rennie, J., Palecki, M. A.: Observation perspectives from U.S. Climate Reference Network  
 492 (USCRN) and Cooperative Observer Program (COOP) Network: temperature and precipitation  
 493 comparison, *J. Atmos. Oceanic Technol.*, 32(4): 703-721, [https://doi.org/10.1175/JTECH-D-14-](https://doi.org/10.1175/JTECH-D-14-00172.1)  
 494 [00172.1](https://doi.org/10.1175/JTECH-D-14-00172.1), 2015.

495 Lenssen, N. J. L., Schmidt, G. A., Hansen, J., Menne, M. J., Persin, A., Ruedy, R., and Zyss, D.:  
 496 Improvements in the GISTEMP uncertainty model, *J. Geophys. Res.*, 124(12): 6307-6326,  
 497 <https://doi.org/10.1029/2018JD029522>, 2019.

498 Li, Q. X., Sun, W. B., Huang, B. Y., Dong, W. J., Wang, X. L., Zhai, P. M., and Jones, P.: Consistency of  
 499 global warming trends strengthened since 1880s, *Science Bulletin*,  
 500 <https://doi.org/10.1016/j.scib.2020.06.009>, 2020a.

Li, Q. X., Sun, W. B., Huang, B. Y., Dong, W. J., Wang, X. L., Zhai, P. M., and Jones, P.: An updated evaluation on the global Mean Surface Temperature trends since the start of 20th century, *Climate Dynamics*, 56:635-650, <https://doi.org/10.1007/s00382-020-05502-0>, 2021..

Li, Q. X., Dong, W. J., and Jones, P.: Continental Scale Surface Air Temperature Variations: Experience Derived from the Chinese Region, *Earth-Science Reviews*, 200, 102998, <https://doi.org/10.1016/j.earscirev.2019.102998>, 2020b.

Li, Q. X., Zhang, L., Xu, W. H., Zhou, T. J., Wang, J. F., Zhai, P. M., and Jones, P.: Comparisons of time series of annual mean surface air temperature for China since the 1900s: Observation, Model simulation and extended reanalysis, *Bull. Amer. Meteor. Soc.*, 98(4):699-711, <https://doi.org/10.1175/BAMS-D-16-0092.1>, 2017.

Li, Q. X., Dong, W. J., Li, W., Gao, X. R., Jones P., Parker D., and Kennedy J.: Assessment of the uncertainties in temperature change in China during the last century, *Chin. Sci. Bull.* 55(19): 1974-1982, <https://doi.org/10.1007/s11434-010-3209-1>, 2010.

Li., Q. X., Zhang., H. Z., Liu., X. N., and Huang., J. Y.: Urban heat island effect on annual mean temperature during the last 50 years in China, *Theor. Appl. Climatol.*, 79, 165-174, <https://doi.org/10.1007/s00704-004-0065-4>, 2004.

Li, Y., Tinz, B., Storch, H., Wang, Q. Y., Zhou, Q. L., and Zhu, Y. N.: Construction of a surface air temperature series for Qingdao in China for the period 1899 to 2014, *Earth Syst. Sci. Data*, 10(1): 643-652, <https://doi.org/10.5194/essd-10-643-2018>, 2018.

Lv, Y. M., Guo, J. P., Yim, S. H., Yun, Y. X., Yin, J. F., Liu, L., Zhang, Y., Yang, Y. J., Yan, Y., and Chen, D. D.: Towards understanding multi-model precipitation predictions from CMIP5 based on China hourly merged precipitation analysis data, *Atmos. Res.*, 231(1): 104671,

<https://doi.org/10.1016/j.atmosres.2019.104671>, 2020. Menne, M. J., Durre, I., Vose, R. S., Gleason, B.,  
 and Houston, T. G.: An overview of the global historical climatology network-daily database, *J. Atmos.*  
*Ocean. Technol.*, 29(7): 897-910, <https://doi.org/10.1175/JTECH-D-11-00103.1>, 2012.

Menne, M. J., Williams, C. N., Gleason B. E., Rennie J. J., and Lawrimore, J. H.: The global historical  
 climatology network monthly temperature dataset, Version4, *J. Climate*, 31(24): 9835-9854,  
<https://doi.org/10.1175/JCLI-D-18-0094.1>, 2018.

Png, I. P. L., Chen, Y., Chu, J. H., Feng Y. K., Lin, E. K. H., and Tseng W. L.: Temperature, precipitation  
 and sunshine across China, 1912-51: A new daily instrumental dataset, *Geosci. Data J.*, 1-13,  
<https://doi.org/10.1002/gdj3.91>, 2020. Quayle, R. G., Easterling, D. R., Karl T. R., Hughes, P. Y.: Effects  
 of recent thermometer changes in the cooperative station network, *Bull. Amer. Meteor. Soc.*,  
 72(11):1718-1723, [https://doi.org/10.1175/1520-0477\(1991\)0722.0.CO;2](https://doi.org/10.1175/1520-0477(1991)0722.0.CO;2), 1991.

Rahimzadeh, F., and Zavareh, M. N.: Effects of adjustment for non-climatic discontinuities on  
 determination of temperature trends and variability over Iran, *Int. J. Climatol.*, 34(6): 2079-2096,  
<https://doi.org/10.1002/joc.3823>, 2014.

Rohde, R., A., Muller, R. A., Jacobsen, R., Muller, E., Perlmuter, S., Rosenfeld, A., Wurtele J., Groom,  
 D., and Wickham, C.: A new estimate of the average earth surface land temperature spanning 1753 to  
 2011, *Geoinfor. Geostat: An overview*, 1: 1, <https://doi.org/10.4172/2327-4581.1000101>, 2013.

Rohde, R., A., and Hausfather, Z.: The Berkeley Earth Land/Ocean Temperature Record, *Earth Syst. Sci.*  
*Data*, 12, 3469-3479, <https://doi.org/10.5194/essd-12-3469-2020>, 2020.

Si, P., Zheng, Z. F., Ren, Y., Liang, D. P., Li, M. C., and Shu, W. J.: Effects of urbanization on daily  
 temperature extremes in North China, *J. Geogr. Sci.*, 24(2): 349-362, [https://doi.org/10.1007/s11442-](https://doi.org/10.1007/s11442-014-1092-4)  
 014-1092-4, 2014.

Si, P., Hao, L. S., Luo, C. J., Cao, X. C., and Liang, D. P.: The interpolation and homogenization of long-term temperature time series at Baoding observation station in Hebei Province, *Climate Change Research*, 13(1): 41-51, 2017 (in Chinese).

Si, P., Luo, C. J., and Liang, D. P.: Homogenization of Tianjin monthly near-surface wind speed using RHtestsV4 for 1951-2014, *Theor. Appl. Climatol.*, 132 (3-4): 1303-1320, <https://doi.org/10.1007/s00704-017-2140-7>, 2018.

Si, P., Luo, C. J., and Wang, M.: Homogenization of Surface Pressure Data in Tianjin, China, *J. Meteor. Res.*, 33(6): 1131-1142, <https://doi.org/10.1007/s13351-019-9043-8>, 2019.

Si, P., Wang, J., Li, H. J., and Nian, F. X.: Homogenization and application of meteorological observation data at provincial level. Beijing: China Meteorological Press, 76-91, 2020(in Chinese).

Si, P., and Li, Q. X.: Tianjin homogenized daily surface air temperature over century-long scale. PANGAEA, <https://doi.org/10.1594/PANGAEA.924561>, 2020.

Sun, X. B., Ren, G. Y., Xu, W. H., Li, Q. X., and Ren, Y. Y.: Global land-surface air temperature change based on the new CMA GLSAT data set, *Science Bulletin*, 62(4): 236-238, <http://doi.org/10.1016/j.scib.2017.01.017>, 2017.

Trewin, B.: A daily homogenized temperature data set for Australia, *Int. J. Climatol.*, 33(6): 1510-1529, <http://doi.org/10.1002/joc.3530>, 2013.

Harris, I. C., Jones, P. D. CRU TS4.03: Climatic Research Unit (CRU) Time-Series (TS) version 4.03 of high-resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2018). Centre for Environmental Data Analysis, 22, <http://doi.org/10.5285/10d3e3640f004c578403419aac167d82>, 2020.

Vincent, L. A., Wang, X. L., Milewska, E. J., Wan, H. Yang, F., and Swail, V. R.: A second generation of homogenized Canadian monthly surface air temperature for climate trend analysis, *J. Geophys. Res.*, 117, D18110, <http://doi.org/10.1029/2012JD017859>, 2012.



567 Vincent, L. A., Zhang, X., Bonsal, B. R., and Hogg, W. D.: Homogenization of daily temperature over  
568 Canada, *J. Climate*, 15:1322-1334, [http:// doi.org/10.1175/1520-0442\(2002\)0152.0.CO;2](http://doi.org/10.1175/1520-0442(2002)0152.0.CO;2), 2002.

569 Wang, S. W., Ye, J. L., Gong, D. Y., Zhu, J. H., and Yao, T. D.: Construction of mean annual temperature  
570 series for the LSAT one hundred years in China, *Quarterly Journal of Applied Meteorology*, 9(4): 392-  
571 401, 1998(in Chinese).

572 Wang, S. W., Gong, D. Y., Ye, J. L., and Chen, Z. H.: Seasonal precipitation series of Eastern China since  
573 1880 and the variability, *Acta Geographica Sinica*, 35(3): 281-293, 2000 (in Chinese).

574 Wang, X. L., Wen, Q. H., and Wu Y. H.: Penalized maximal t test for detecting undocumented mean change  
575 in climate data series, *J. Appl. Meteor. Climatol.*, 46, 916-931, [http:// doi.org/10.1175/JAM2504.1](http://doi.org/10.1175/JAM2504.1), 2007.

576 Wang, X. L., Chen, H. F., Wu, Y. H., Feng, Y., and Pu, Q.: New techniques for the detection and adjustment  
577 of shifts in daily precipitation data series, *J. Appl. Meteor. Climatol.*, 49, 2416-2436, [http://](http://doi.org/10.1175/2010jamc2376.1)  
578 [doi.org/10.1175/2010jamc2376.1](http://doi.org/10.1175/2010jamc2376.1), 2010.

579 Wu, Z. X.: China Modern Meteorological Station. Beijing: China Meteorological Press, 180-182, 2007  
580 (in Chinese).

581 Xu, C. D., Wang, J. F., and Li, Q. X.: A new method for temperature spatial interpolation based on sparse  
582 historical stations, *J. Climate*, 31(5): 1757-1770, [http:// doi.org/10.1175/JCLI-D-17-0150.1](http://doi.org/10.1175/JCLI-D-17-0150.1), 2018.

583 Xu, W. H., Li, Q. X., Jones, P., Wang, X. L., Trewin, B., Yang, S., Zhu, C., Zhai, P. M., Wang, J. F., Vincent,  
584 L. A., Dai, A. G., Gao, Y., and Ding, Y. H.: A new integrated and homogenized global monthly land  
585 surface air temperature dataset for the period since 1900, *Clim. Dynam.*, 50(15): 2513-2536, [http://](http://doi.org/10.1007/s00382-017-3755-1)  
586 [doi.org/10.1007/s00382-017-3755-1](http://doi.org/10.1007/s00382-017-3755-1), 2018.

587 Xu, W. Q., Li, Q. X., Wang, X. L., Yang, S., Cao, L. J., and Feng, Y.: Homogenization of Chinese daily  
588 surface air temperatures and analysis of trends in the extreme temperature indices, *J. Geophys. Res.*

589 Atmos., 118, [http:// doi.org/10.1002/jgrd.50791](http://doi.org/10.1002/jgrd.50791), 2013.

590 Yan, Z. W., Chi, Y., and Jones, P.: Influence of inhomogeneity on the estimation of mean and extreme  
591 temperature trends in Beijing and Shanghai, *Adv. Atmos. Sci.*, 18(3): 309-322, [http://](http://doi.org/10.1007/BF02919312)  
592 [doi.org/10.1007/BF02919312](http://doi.org/10.1007/BF02919312), 2001.

593 Yu, J., Li, Q. X., Zhang, T. W., Xu, W. H., Zhang, L., and Cui, Y.: The merging test using measurements,  
594 paleoclimate reconstruction and climate model data based on Bayesian model, *Acta Meteorologica*  
595 *Sinica*, 76(2): 304-314, 2018(in Chinese).

596 Zhang, X., Alexander, L., Hegerl, G.C., Jones, P.D., Klein Tank, A., Peterson, T.C., Trewin, B. and Zwiers,  
597 F.W.: Indices for monitoring changes in extremes based on daily temperature and precipitation data.  
598 *WIREs Climate Change*, 2, 851-870, [http:// doi.org/10.1002/wcc/147](http://doi.org/10.1002/wcc/147), 2011.

599 Zheng, J. Y., Liu, Y., Ge Q. S., Hao, Z. X.: Spring phenodate records derived from historical documents  
600 and reconstruction on temperature change in Central China during 1850-2008, *Acta Geographica Sinica*,  
601 70(5): 696-704, 2015(in Chinese).

602 Zhai., P. M., Chao., Q. C., and Zou., X. K.: Progress in China's climate change study in the 20th century,  
603 *J. Geograph. Sci.*, 14(1): 3-11, <https://doi.org/10.1007/BF02841101>, 2004.

604