

# Changes in global air pollutant emissions during the COVID-19 pandemic: a dataset for atmospheric modeling

5 Thierno Doumbia<sup>1</sup>, Claire Granier<sup>1,2</sup>, Nellie Elguindi<sup>1</sup>, Idir Bouarar<sup>3</sup>, Sabine Darras<sup>4</sup>, Guy Brasseur<sup>3,5</sup>, Benjamin Gaubert<sup>5</sup>, Yiming Liu<sup>6</sup>, Xiaoqin Shi<sup>3</sup>, Trissevgeni Stavrakou<sup>7</sup>, Simone Tilmes<sup>5</sup>, Forrest Lacey<sup>5</sup>, Adrien Deroubaix<sup>3</sup>, Tao Wang<sup>6</sup>

<sup>1</sup>Laboratoire d'Aérodologie, Université de Toulouse, CNRS/UPS, Toulouse, France

<sup>2</sup>NOAA Chemical Sciences Laboratory and CIRES/University of Colorado, Boulder, CO, USA

10 <sup>3</sup>Max-Planck Institute for Meteorology, Hamburg, Germany

<sup>4</sup>Observatoire Midi-Pyrénées, Toulouse, France

<sup>5</sup>Atmospheric Chemistry Observations and Modeling, National Center for Atmospheric Research, Boulder, CO, USA

15 <sup>6</sup>Department of Civil and Environmental Engineering, The Hong Kong Polytechnic University, Hong Kong, China

<sup>7</sup>Royal Belgian Royal Institute for Space Aeronomy, Brussels, Belgium

*Correspondence to:* Thierno Doumbia ([thierno.doumbia@aero.obs-mip.fr](mailto:thierno.doumbia@aero.obs-mip.fr))

Claire Granier ([claire.granier@aero.obs-mip.fr](mailto:claire.granier@aero.obs-mip.fr))

20

25

**Abstract.** In order to fight the spread of the global COVID-19 pandemic, most of the world countries  
30 have taken control measures such as lockdowns during a few weeks to a few months. These  
lockdowns had significant impacts on economic and personal activities in many countries. Several  
studies using satellite and surface observations have reported important changes in the spatial and  
temporal distributions of atmospheric pollutants and greenhouse gases. Global and regional  
chemistry-transport model studies are being performed in order to analyze the impact of these  
35 lockdowns on the distribution of atmospheric compounds. These modeling studies aim at evaluating  
the impact of the regional lockdowns at the global scale. In order to provide input for the global and  
regional model simulations, a dataset providing adjustment factors (AFs) that can easily be applied  
to current global and regional emission inventories has been developed. This dataset provides, for  
the January-August 2020 period, gridded AFs at a 0.1x0.1 latitude/longitude degree resolution, on a  
40 daily or monthly basis for the transportation (road, air and ship traffic), power generation, industry  
and residential sectors. The quantification of AFs is based on activity data collected from different  
databases and previously published studies. A range of AFs is provided at each grid point for model  
sensitivity studies. The emission AFs developed in this study are applied to the CAMS global  
inventory (CAMS-GLOB-ANT\_v4.2\_R1.1), and the changes in emissions of the main pollutants  
45 are discussed for different regions of the world and the first six months of 2020. Maximum decreases  
in the total emissions are found in February in Eastern China, with an average reduction of 20-30 %  
in NO<sub>x</sub>, NMVOCs and SO<sub>2</sub> relative to the reference emissions. In the other regions, the maximum  
changes occur in April, with average reductions of 20-30 % for NO<sub>x</sub>, NMVOCs and CO in Europe  
and North America and larger decreases (30-50 %) in South America. In India and African regions,  
50 NO<sub>x</sub> and NMVOCs emissions are reduced on average by 15-30 %. For the others species, the  
maximum reductions are generally less than 15 %, except in South America, where large decreases  
in CO and BC are estimated. As discussed in the paper, reductions vary highly across regions and  
sectors, due to the differences in the duration of the lockdowns before partial or complete recovery.  
The dataset providing a range of AFs (average and average  $\pm$  standard deviation) is called  
55 CONFORM (COvid-19 adjustmeNt Factors fOR eMissions) (<https://doi.org/10.25326/88>). It is  
distributed by the Emissions of atmospheric Compounds and Compilation of Ancillary Data  
(ECCAD) database (<https://eccad.aeris-data.fr/>).

## 1 Introduction

60 The COVID-19 pandemic has triggered different measures in most world countries to protect  
citizens from the spread of the Severe Acute Respiratory Syndrome-Coronavirus 2 (SARS-CoV-2).  
The first measures, which included strict lockdowns, started in China at the end of January 2020. As  
the pandemic was spreading all over the world, lockdowns or other measures were gradually  
implemented in Asia, Europe, Oceania, North and South America and Africa. The restriction during  
65 the different lockdowns have resulted in significant changes in the emissions of greenhouse gases

(Le Quéré et al., 2020; Liu et al., 2020; Han et al., 2021; Zheng et al., 2020) and reactive air pollutants (Forster et al., 2020; Venter et al., 2020). The impact of the reductions on the emissions of primary pollutants was assessed in different regions based on surface observations (e.g. Shi and Brasseur, 2020; Lee et al., 2020; Kim et al., 2020) and satellite retrievals (e.g. Bauwens et al., 2020; Zhang et al., 2020; Diamond et al., 2020; Biswal et al., 2020). Modeling studies have been initiated to investigate the changes in the global and regional distributions of tropospheric chemical compounds during the pandemics (Gaubert et al., 2021; Venter et al., 2020; Xing et al., 2020; Keller et al., 2020; Barré et al., 2020). These modeling studies are based on estimates of emissions for primary species, and on consistent changes of the emissions during the COVID-19 lockdown periods.

Here we present a global gridded dataset of emission adjustment factors (AFs) at a  $0.1^\circ \times 0.1^\circ$  resolution and on a daily or monthly basis based on available activity data. Recent studies (Le Quéré et al., 2020; Forster et al., 2020; Liu et al., 2020) have discussed the changes in emissions at the global scale for several chemical compounds. In this paper, we propose a dataset of AFs for the individual sectors related to transportation including road, air and ship traffic, industry, residential activities and power generation. The advantage of such a dataset, which provides AFs for assessments of the impact of COVID-19 restrictions on pollutants emitted into the atmosphere, is that it can be applied directly to most global and regional inventory used in chemistry-climate and transport models in a flexible way. The inventories commonly used in these models (EDGAR (Crippa et al., 2018), ECLIPSE (Klimont et al., 2017), CAMS (Granier et al., 2019), MEIC (Li et al., 2017), etc.) include the sectors already mentioned (industrial processes, road transportation, power generation, residential, aviation and shipping). The emissions of these sectors are developed on the basis of a combination of several sub-sectors that are not provided separately in the emission inventories.

Section 2 describes the general methodology used in the development of the AFs. The following sub-sections analyze the sectoral activity data used to determine these factors. The availability of activity data depends strongly on the regions under consideration, and this might often lead to large uncertainties. We are therefore providing for each location and sector an estimated average, low and high values of the AFs. In Section 3, we present and discuss the sectoral changes in emission AFs over selected regions as a function of time.

This paper discusses the dataset developed for the January-August 2020 period: the dataset is called CONFORM (Covid-19 adjustmeNt Factors fOR eMissions) and will be updated to account for the latest available activity data information. The impact of the AFs on the emissions developed as part of the CAMS global anthropogenic emissions (Granier et al., 2019; Elguindi et al., 2020) are discussed for the NMVOC, CO, NO<sub>x</sub>, BC, OC and SO<sub>2</sub> species in Section 3.5.

## 2 Methodology

105 To quantify the amount of the corresponding global gridded daily/monthly changes, the adjustment factors (AFs) are determined following the general methodology schematized in Figure 1. The four different steps taken in our study to estimate the AFs are the following: 1) A data survey is first carried out to collect activity data for the power generation, industrial processes, residential, road transportation, shipping and aviation sectors. These sectors correspond to those considered in many  
110 global inventories such as the CAMS global anthropogenic emission inventory (CAMS-GLOB-ANT\_v4.2\_R1.1, Granier et al., 2019). The activity data used to estimate the emission AFs are available from numerous sources for different time scales, depending on the geographical area. It should be noted that, in several regions of the world, accurate and/or up-to-date data are not publicly available, which could lead to a significant source of uncertainty in estimating reduction rates. Table  
115 1 indicates the availability of data for the different sectors. Some of these data are used to support our analysis and contribute to the estimation of uncertainties on the AFs; 2) The collected data are then analyzed and an intercomparison of the changes in activity data of datasets providing similar or equivalent parameters is performed. The dataset that provides the most detailed information and which meets better our needs is then chosen. The non-gridded AFs are calculated for each country or state/province according to the availability of activity data, as detailed in the following sections ;  
120 3) The gridded daily/monthly netcdf files developed for each sector are obtained by assigning, to each cell, the value of the AFs in the whole country or state/province level corresponding to this grid cell: this is based on the fact that the lockdowns and restrictions have been generally taken at national or state level. For several sectors (road transport, industry and residential) the AF at the country or state/province level represents an average value calculated from several individual cities or locations; and 4) Finally, a comparison of the AFs derived from this study with other published data is performed, together with an evaluation of their impact on emissions, using the CAMS-GLOB-ANT\_v4.2\_R1.1 inventory as a basis.

125

130 The dataset discussed in this paper covers the period from 1 January to 31 August 2020. It will be regularly updated to account for the latest available activity data information.

The AFs are calculated as the ratio between the activity data for a given sector and day or month, and the median value of the activity data over the five week-period starting on 1 January 2020. This  
135 corresponds to the baseline used in Google mobility trends. The median value is calculated from 1 January to 4 February for almost all regions, except for China where the first lockdown started on 23 January 2020 in the Hubei province. The reference values for China's activity data are taken as the median daily value of the first three weeks (i.e. from 1 to 21 January 2020). The same baseline is used for all sectors, except for those for which only monthly activity data are available. For those  
140 sectors, we consider the value of January 2020 as the baseline. We have chosen the reference year as 2020 (pre-COVID-19), because detailed activity data such as mobility data are only publicly available for the year 2020.

## 145 2.1 Road transportation

Several studies related to the COVID-19 pandemic have used the Google COVID-19 Community Mobility data (<https://www.google.com/covid19/mobility/>) (Forster et al., 2020; Guevara et al., 2020). This dataset provides the number of visitors to specific locations based on mobile phone  
150 locations (e.g. parks, grocery and pharmacy stores, workplaces, retail and recreation, train stations and residential) every day relative to a baseline value. This baseline is calculated as the median value over the five-week period from January 3<sup>rd</sup> to February 6<sup>th</sup> 2020. We use measurements up to 31 August 2020 for all countries for which data are available (about 133 countries). For the USA, the analysis was performed for each state. Google daily data are provided at country/state level and  
155 generally cover several local areas (i.e. subregion or city). For example, in the USA, Google spans the 50 states and the District of Columbia, as well as several cities in each state. The AFs for the road transportation sector are derived from the estimation of transit usage (i.e. public transportation including train stations, bus and subways) made by Google. In order to make the calculated AFs comparable with those derived using the other data sources considered in this study, the AFs for the  
160 Google's categories are scaled to 1 using the following formula  $AF = 1 + \text{Google}/100$ , so that their values are less than 1 for a reduction in activity and above 1 otherwise.

It should be noted that Google mobility data are not available for all countries (e.g. China and for nearly two thirds of African countries). Other datasets are providing mobility trends, such as the  
165 Apple mobility trend reports (<https://covid19.apple.com/mobility>) and the TomTom ([https://www.tomtom.com/en\\_gb/traffic-index/ranking/](https://www.tomtom.com/en_gb/traffic-index/ranking/)) traffic congestion index. Apple measures the number of requests for directions scaled relative to 13 January 2020, while TomTom provides the percentage of the differences in time spent on a trip compared to uncongested conditions. Google provides a better spatial coverage than the other two datasets. In Africa for example, Apple and  
170 TomTom data are available for 3 and 2 countries, respectively, while Google provides data for 26 African countries. In China, TomTom reports measurements only for some cities. Unlike Google and Apple, TomTom markets its data, which are therefore not in open access. In order to evaluate the Google dataset, a regional comparison with Apple mobility data was performed (Supplementary Information, Figures S1-2). First, we looked separately at one Apple category (driving  
175 measurements, generated by counting the number of requested directions in transit or public transport on Apple applications) and three Google categories (retail/pharmacy, workplaces and grocery shopping destinations) according to three regions (Europe, USA and the rest of the world (ROW)). The comparisons of temporal series show that, in general, Apple driving displays much larger variations than the Google mobility changes (Figure S1). However, the patterns of the average  
180 values of the four categories show a significant decrease in Europe of up to 57 % for Apple driving and of 57, 65 and 31 % for Google workplaces, retail and grocery, respectively. The corresponding values in the USA are 43, 16, 41 and 40 %, respectively, while those in the ROW are 61, 31, 60 and 48 %. A comparison between monthly Google's non-residential (i.e. combining grocery, workplaces, transit and retail) and Apple driving data, which can be considered as equivalent

185 categories, displays high correlation coefficients varying from 0.79 to 0.97 depending on the region  
(Figure S2). The data of these two categories agree relatively well, particularly during the first  
months of the COVID-19 pandemic (February to May) when lockdowns were strict in most regions.  
The largest discrepancies are observed from June to August, especially in Europe and the USA.  
These differences are likely due to the fact that the two datasets do not represent the same parameters:  
190 Apple bases their values on the volume of directions requests on phone applications while Google  
uses mobile phone locations. These differences can be attributed to a combination of several factors  
including the spatial coverage, the mode and category of transportation considered, the location of  
the measurements within the country or state/province. The calculation methods are also very  
different from one dataset to another.

195 Comparisons of the changes in the ground transportation sector for the different mobility datasets  
have been performed in recent studies (Forster et al., 2020; Le Quéré et al., 2020). The correlation  
coefficients between Google and Apple transit categories calculated in our study are in line with the  
value of 0.8 reported by Forster et al. (2020), for the February to June period. Liu et al. (2020) show  
200 trends from both Google and TomTom mobility datasets with the same order of magnitude during  
the first quarter of 2020. This analysis highlights the importance in the choice of the data to be  
considered in the estimation of changes based on activity data, especially for the more recent months,  
as well as for the future, when the dataset will be extended. Based on this analysis, we have used the  
Google Mobility data for estimating the AFs in regions where data are available.

205 In China, the AFs for road transportation were calculated based on Baidu Migration Scale index  
(<https://qianxi.baidu.com/>, available for China only). These indices are aggregated from migration  
flows within China, based on the positioning requests on Baidu Map Services. The index indicates  
the ratio between the number of people traveling in a city and the population of this city. The data  
210 are available only from 1 January to 2 May 2020 and cover about 343 Chinese cities in all provinces.  
Baidu officially stopped updating the dataset on its platform on 8 May 2020. We compared the  
changes in the Baidu Migration Scale Index with the relative difference of TomTom congestion  
levels for the Beijing, Tianjin, Chongqing and Shanghai Chinese areas. TomTom archived data are  
not freely available, but they can be retrieved from published graphs  
215 ([https://www.tomtom.com/en\\_gb/traffic-index/ranking](https://www.tomtom.com/en_gb/traffic-index/ranking)) providing the weekly changes in 2020  
relative to the same periods in 2019. The results show a strong similarity between changes given by  
these two datasets for the period covered by the Baidu dataset (January to April), with a correlation  
coefficient of 0.9 (Figure S3). For the period from May to August 2020, Liu et al. (2020) showed  
that the difference in estimated CO<sub>2</sub> emissions from road transport between 2020 and 2019 is on  
220 average about 4% for May-November, with larger values for some days. Based on these analyses,  
and in order to cover the whole period of our study (January to August), we assume that changes in  
road traffic in China after May 2020 are relatively low and close to those before the spread of the  
COVID-19 virus. These comparisons show that the proposed method for calculating the AFs (i.e.  
ratio between the activity data and the median value of activity data over a defined reference period)  
225 is consistent with changes in 2020 relative to the same period in 2019.

## 2.2 Industrial processes

230 This sector includes industrial production processes such as manufactured products from fossil fuel  
combustion, and represents a significant part of the emission sources of atmospheric pollutants. The  
2020 data concerning industrial production are, however, not publicly available for many  
countries/regions. We used the crude steel production from the world steel association (Table 1),  
which is provided on a monthly basis, to estimate the rate of change in the industrial sector. Due to  
235 the difficulties to access the daily data in this source category, we assumed that changes in Google's  
workplace measures, which represent the percentage of people travelling to/from their workplaces,  
are representative of changes in industrial activities during the lockdowns. To verify our hypothesis,  
we compared the calculated monthly average AFs based on Google's workplace data for selected  
countries with those derived from crude steel productions (Figure S4). The average change in the  
240 first eight months of 2020 relative to the same periods in 2019 in crude steel production for the 24  
countries shown in Figure S4 is 17 %, while the corresponding value using Google's workplace  
measures is 27 %. This indicates a fair agreement between the two datasets. However, there are large  
differences in some countries between these data. For example, in Europe the greatest change in  
crude steel production is 24 % in comparison to 59 % estimated using Google's workplace category,  
245 indicating a high level of uncertainty in the AFs for the industry sector.

## 2.3 Power generation

250 The power sector emission AFs were estimated by compiling several sources such as the total  
electricity load from the ENTSO-E (European Network of Transmission System Operators for  
Electricity) transparent platform for the European Union and the United Kingdom, the regional  
electricity demand from the EIA (Energy Information Administration) for the United States, the  
daily reports of the electricity generation from fossil fuel including coal, lignite and gas, Naptha and  
255 Diesel provided by the POSOCO (Power System Operation Corporation) for India, the thermal  
electricity production from the ONS (Operator of the National Electricity System) for Brazil and the  
daily power generation provided by the United Power system of Russia. We also used the local  
electricity demand data in Singapore. For Canada, we assumed that the data in Ontario are  
representative of power generation trends (Table 1). It should be noted that we did not apply any  
260 temperature correction to the electricity load. Liu et al. (2020) indicated that the COVID-19 and  
related restrictions explain about 85 % of the reduction in the power sector from January to March  
and only 15 % are attributed to the temperature effect. For the rest of the world where data on  
electricity demand or thermal production are not publicly available, we estimate the AFs based on  
the data published by Le Quéré et al. (2020).

265

## 2.4 Air transportation and shipping

270 Due to a lack of public daily data on air transportation to determine the AFs related to the large  
decline in passenger flights, we used the monthly data published by the Knowledge Center on  
Migration and Demography (KCMD) Dynamic Data Hub (Table 1). In addition to the observed  
passenger volumes from around the world, the KCMD provides air traffic scenarios covering the  
COVID-19 period. Five scenarios are provided (Iacus et al., 2020). We selected the scenario called  
EUROC-L (L-shaped version) based on the Eurocontrol air traffic data, which provides global up-  
275 to-date monthly average air volumes. To assess the representativeness of the KCMD estimates, we  
compared the EUROC-L scenario data with the number of international scheduled flights from 14  
countries in 2020 provided by the OAG (Official Aviation Guide)  
(<https://www.oag.com/coronavirus-airline-schedules-data>). The results indicated that, when  
280 considering the global average, trends from both datasets are relatively similar, with a significant  
decrease of up to 65 % for OAG and 80 % for KCMD in April, May and June compared to the  
reference value in January 2020. However, since July there has been a slow recovery towards the  
pre-lockdown values (Figure S5, Supplementary Information). EUROC-L seems to overestimate the  
changes during the period when restrictions were the most severe, especially in China, Japan and  
South Korea. In contrast, there is an underestimation of trends since July 2020 in most of the  
285 countries as shown in Figure S5. A lockdown for the whole of China was declared on February 10,  
2020 and the country had its air traffic heavily impacted on the following days. This disruption does  
not appear in the KCMD data. This analysis leads us to the conclusion that the KCMD data can only  
be used to complement the OAG data. It should be noted that KCMD data have the advantage of  
being available for more than 230 countries and are regularly updated.

290 Detailed data on 2020 international and national shipping are not publicly available yet. In this study,  
the changes in shipping activity are determined based on the weekly containership port calls during  
the 31 weeks of 2020 (covering the period from 1 January to 2 August) compared to the same period  
in 2019 and reported by the United Nations Conference on Trade and Development (UNCTAD)  
295 (<https://unctad.org/news/covid-19-shipping-data-hints-some-recovery-global-trade>). UNCTAD  
provides official statistics from marine traffic for different regions: North America (Canada, Mexico,  
United States), East Coast of South America (Argentina, Brazil, Uruguay), Northern Europe  
(Belgium, Germany, Netherlands), Southern Europe (France, Italy, Spain), Northern and Western  
Africa (Egypt, Morocco, Nigeria, Togo), Eastern and Southern Africa (Kenya, Tanzania, South  
300 Africa), South Asia (Bangladesh, India, Pakistan, Sri Lanka), South East Asia (Indonesia, Malaysia,  
Singapore, Thailand), China and Hong Kong. Weekly values are directly extracted from graphs and  
monthly AFs are derived.

## 2.5 Residential sector

305 In this study, the AF calculations for the residential sector are performed using measurements from  
Google's residential category, which covers most of the countries in the world. Google's residential



category represents the duration during which people are constrained to their home through lockdown. This duration represents the additional time that people spent in places of residence due to the restrictions. In the CAMS-GLOB-ANT inventory, the residential sector includes mainly emissions due to cooking, heating and auxiliary engines that primarily use biomass or fossil fuels in households. The emissions from this sector vary according to the geographical zone due to the difference in lifestyles. We consider that, when people stay at home, the impact on heating use is moderate: heating systems generally continue to operate during working hours, at a lower intensity. The extra time spent at home contributes significantly to other domestic activities, namely cooking, heating water and activities using fossil fuels. Based on this analysis, we expect an increase in residential combustion during the lockdown period. In China, due to the lack of data, we use residential emissions published by Le Quéré et al. (2020) which are based on electricity demand for the city of London for the first fourth months of 2020. For the rest of the study period, we assumed that there is no change in the AFs for the residential sector in China.

### 3 Results

#### 3.1 Adjustment factors for the transportation sector

This section focuses on the changes in the transportation sector which include road transportation, air traffic and shipping.

##### 3.1.1 Road transportation

The AFs related to road transportation provide significantly lower emission values, when compared to typical values before the restrictions, and present large regional variations. Figure 2 displays the averages AFs in Europe, USA, South America, China, Africa and the rest of the world since the beginning of 2020. The regional variations of AFs (light pink color) are determined as the standard deviation of individual values for all the countries in the region or from local measurements in the country or state/province. We also calculated the minimum and maximum values within specific regions from all the values calculated for each of the countries in this region. An average daily reduction of up to 60 % in mid-April is shown in most regions of the world. In China and Europe, the reductions reached a maximum in mid-February and late March, respectively. In China, the lockdowns started on 23 January and the period from 24 January to 2 February coincides with Chinese Spring Festival holidays during which business activities were reduced. It has been reported that short term emissions and atmospheric composition reductions associated with the Spring Festival in China were about 10 % (Lin et al., 2011; He et al., 2021; Sun et al., 2020; Zhang et al., 2020; Zhang et al., 2021). The 2020 reductions in emissions from road transport are significantly larger and peak during February 2020, with no rebound after the Chinese New Year holiday (e.g. Kraemer et al., 2020; Miyazaki et al., 2020). In the USA where the variability is largest, the average decrease reached 40 % in mid-April. In Africa, the largest decrease occurred at the same period as in Europe and the USA, but with values of about 50 %, while the average decrease was much larger

350 in South America, reaching 70 %. These results are in agreement with the changes during the first  
quarter of 2020 reported by Le Quéré et al. (2020) and based on Apple, TomTom mobility trends  
and local traffic data for the USA provided by MS2 (Modern Transportation Analytics) Corporation.  
Based on the standard deviations calculated from the data for each region and the evaluation results,  
an uncertainty of  $\pm 10$  to  $\pm 30$  % is associated with the estimated AFs, depending on the region. Figure  
2 also indicates that some regions recovered to the pre-COVID-19 situation more rapidly than others  
355 (e.g. the USA, the European countries and China), while most countries in South America, Africa  
and the rest of the world continue to be affected by lockdown measures at the end of the period  
discussed in this paper.

### 360 **3.1.2 Air traffic and shipping**

As discussed in Section 2, the changes in air traffic were calculated using the monthly global  
scheduled flights from OAG combined with the passenger volumes reported from KCMD. Figure  
3a displays the monthly air traffic AFs with an average value as low as to 0.2 (a decline of 80 %) in  
most of countries in the world. This large decrease spreads over several months from April to June,  
365 while the aviation activity started to rebound in July but the activity remained below the pre-  
pandemic level.

As a consequence of the global lockdown measures, activities in shipping also declined. Due to the  
lack of up-to-date data, we assume that changes in container ship port calls, published on the  
370 UNCTAD website, are representative of trends in shipping activities. The average global change as  
well as the associated standard deviation and upper and lower limits values are represented in Figure  
3b. The number of port calls by container ship globally decline from January to August to  $7 \pm 6$  %  
relative to 2019. However, changes vary across regions. For example, we found a monthly reduction  
in North America and Europe up to 18 % and 20 % in June, respectively. These reductions are in the  
375 lower limit values of 20-30 % reported from the literature and based on forecast and published  
reports (Le Quéré et al., 2020; Liu et al., 2020).

### **3.2 Adjustment factors for the industrial sector**

380 Our estimates of AFs for the industry sector are derived from the Google's workplace measures,  
except for China for which factors are calculated based on CO<sub>2</sub> emissions published by Liu et al.  
2020. Results show that the level of activity in the industrial sector over the 214 considered countries  
started, on average, falling down in mid-March when most of these countries began to take restrictive  
measures. The average AF reaches a maximum decrease (up to 40 %) in April relative to the  
385 reference period (i.e., pre-COVID-19) before increasing until May. It remained relatively stable  
(approximately 20 % reduction) from the beginning of June to August. These levels of change are  
in line with those reported in Forster et al. (2020) for the first half of 2020. As Figure 4 illustrates,  
the average AF displays rather similar patterns across the different regions considered, except in

China. Based on Google data, industrial operations were subject to an important decrease in almost all regions, with a maximum daily average AF value close to 0.4 (60 % reduction) in the European and South American countries and in many other countries, especially in late March and early April. However, contrarily to the others countries, European countries show a second maximum daily average reduction (AF = 0.75 or 25 %) in mid-August, with a magnitude lower than the first peak. The impact of the lockdowns on the industrial activities is somewhat smaller in the USA, Africa and in the rest of the world, with a maximum reduction of about 30-40 % relative to the pre-COVID-19 pandemic values. In China, AFs fell in mid-February to their minimum average values of 0.60 (40 % decrease), but rapidly increased to a complete recovery at the beginning of March and exceeded the pre-pandemic level by an average of 25 % from April onward. The uncertainty range associated with the emission AFs for the industrial sector is evaluated to  $\pm 20$  to  $\pm 30$  %, depending on the region. It is noteworthy that for almost all countries except China, the pre-pandemic level of industrial activity has not been reached, eight months after the beginning of the first lockdown announcements.

### 3.3 Adjustment factors for the residential sector

Contrarily to other sectors, the AFs for the residential sector estimated from Google's mobility data show an increase of 20 to 30 % at its maximum, during the peaks of the lockdowns, depending on the region (Figure 5). The average percentage increase is 30 % in South America, while the maximum average AF value in China is less than 1.10 (i.e. about 10 % increase). As for the others activity sectors, there is a variability of 10 % in all regions, as shown by the standard deviation. This increase in the adjustment factors is mostly due to the fact that, in most countries, schools were closed and teleworking was widespread. As a result, most people in countries affected by strict lockdowns had to stay home most of the time. The impact of lockdowns on residential emissions is quite uncertain, as all the data including fuel use in commercial and residential buildings necessary to quantify this impact are not yet available worldwide. Our study, based on Google's residential measures reflecting time spent by people at their home, leads to increased emissions from the residential sector. However, Liu et al. (2020) reported a global decrease of about -2 % during the first seven months with an uncertainty of about 40 % using fuel consumption data in residential buildings from 2019 that was scaled to 2020 based on the population-weighted heating degree days variation. In this study, the estimated uncertainty range in the emission AFs for residential sector reached  $\pm 20$  %. Our global estimate is consistent with an average increase of around 5 % in activity data from the residential sector, mainly during the strict lockdown period, estimated in Le Quéré et al. (2020).

### 3.4 Adjustment factors for the power generation sector

The COVID-19 pandemic has implications on the share of energy use in industry, commercial and domestic operations. In order to quantify the impact of the lockdowns in this sector, we used the data of electricity demand. The global demand experienced a maximum average decrease of 20 % (AF=0.8) between late March and early April when the restrictions were most stringent in Europe, USA,

430 Africa and many other countries (Figure 6). From mid-June, there is an increase in the demand for  
electricity consumption in the USA which reaches a maximum value of about 20 % at the end of  
July and beginning of August but seems to be followed by a slow decrease. This can possibly be  
explained by an important demand in the commercial and residential sectors. It should be noted that  
435 most African countries did not implement strict lockdowns: as discussed in Section 2, activity data  
are available mostly for North African countries and for South Africa, which experienced severe  
restrictions. The maximum decline ranged from 20 to 30 % in South American countries and in  
China. As for the other sectors, the peak of the reduction factor for energy in China and South  
American countries happened at the end of February and early April, respectively. Our results are in  
line with the average reduction of 20 % or more of electricity demand in mid-March in several  
440 countries, as reported in the 2020 IEA energy review report (IEA, 2020). We estimated an  
uncertainty of  $\pm 15$  % for the power sector, in agreement with the average values of the standard  
deviations calculated over all regions.

### 3.5 Impact on surface emissions

445 The estimations of emission AFs for six economic activity sectors (road transportation, industry,  
power generation, residential, shipping and aviation) discussed in the previous sections are provided  
by the CONFORM dataset on a daily basis, except for aviation and shipping calculated on a monthly  
basis, for the period from January to August 2020 at a  $0.1^\circ \times 0.1^\circ$  grid resolution. This dataset has  
been also used in the Gaubert et al. (2021) paper which provides an analysis of the changes in  
450 secondary atmospheric pollutants during the 2020 COVID-19 Pandemic. Another assessment of  
CONFORM dataset has been performed in Liu et al. (2021)'s study on the analysis of observed and  
modeled surface ozone across China. Results show how the reductions in anthropogenic emissions of  
ozone precursors (NO<sub>x</sub> and VOCs) contributed to the surface ozone changes in agreement with the  
observations.

455 The impact of the AFs changes on the total emissions (sum of emissions from transportation (road  
and non-road traffic), industry, residential, power and shipping) has been analyzed for different  
compounds and selected regions, using the anthropogenic CAMS-GLOB-ANT\_v4.2\_R.1.1  
emission inventory (Granier et al., 2019; Elguindi et al., 2020). This dataset provides daily emissions  
460 of the main atmospheric compounds, including speciated volatile organic compounds at a  $0.1^\circ \times 0.1^\circ$   
resolution, from 2000 to 2020. Version R.1 of the CAMS-GLOB-ANT\_v4.2 dataset incorporates  
the MEIC1.3 regional inventory for China described by Zheng et al. (2018).

The percentage of global change in emissions during the COVID-19 compared to the reference 2020  
emissions is shown in Figure 7 for the main pollutants (NO<sub>x</sub>, CO, SO<sub>2</sub>, BC, OC and NMVOCs)  
465 from January to June 2020. The vertical bars indicate the estimated lower and upper limits of the  
changes in emissions due to the restrictions. Figure 7 shows that the changes depend on the chemical  
species, with decreases in monthly global emissions of 22 % (11-26 %) for NMVOCs, 17 % (12-24  
%) for NO<sub>x</sub>, 14 % (7-18 %) for CO, 10 % (6-18 %) for SO<sub>2</sub> and 9 % (3-13 %) for BC in April 2020

470 compared to a non-COVID-19 scenario. The global changes in OC emissions are different from the  
other species with an increase of 3 % (0-7 %). These results reflect the differences in the contribution  
of each sector to the total emissions for the different species as suggested in Figure S6  
(Supplementary material), which shows the absolute change in emission per sector for NO<sub>x</sub> and OC.  
Figure S6 indicates that the reductions in NO<sub>x</sub> emissions are mainly driven by the changes in road  
transport and industry sectors while there is a large contribution of the residential sector in the OC  
475 emissions.

Figure 8, which displays regional monthly changes in Eastern China (20°N-45°N, 80°E-125°E),  
Europe (35°N-70°N, 20°W-20°E), North America (20°N-50°N, 135°W-35°W), South America  
(60°S-20°N, 90°S-35°S), India (05°N-30°N, 60°E-90°E) and Africa (40°S-30°N, 20°W-40°E), also  
indicates large differences in the changes in emissions among the regions. In agreement with the  
480 changes in the activity data previously shown, the reductions are highest in February in Eastern  
China region for all species, except OC for which a relatively small increase is observed during that  
month. The average monthly NO<sub>x</sub> emissions in Eastern China derived from this study decreased in  
February 2020 by 29 % (24-37 %) compared to the reference emission scenario (i.e. without  
COVID-19 effect). The NMVOCs emissions decrease significantly, by 22 % (15-29 %). The  
485 decreases in the amount of SO<sub>2</sub> emissions are of the same order of magnitude. BC and CO show  
maximum average reductions of 8 and 10 %, respectively in the Eastern China region, while OC  
shows a slight increase of 1.6 % for February.

In the rest of the world, NO<sub>x</sub> emissions exhibit large decreases (13-42 %) during the strictest  
490 shutdown period (i.e. in April) when almost all sectors of activity slowed down or stopped. In  
Europe, the average reduction in NO<sub>x</sub> emissions is 25 % (20-35 %) in April. These values are in the  
same order of magnitude as the mean change of -33 % reported in Guevara et al. (2020) for Europe  
for the period from 23 March to 26 April 2020. For SO<sub>2</sub>, the average reduction in Europe calculated  
by Guevara et al. (2020) is in the same order of magnitude as our low estimation range of 9-23 %  
495 (average value of 14 %) in April. However, the percentage decline in NMVOCs emissions (21-40  
% with an average of 34 % as for April) derived from our study is much higher than the value from  
Guevara et al. (2020). Solvents and industrial processes are the main sectors contributing to  
anthropogenic NMVOCs emissions: in our study, the changes in the solvents sectors, for which no  
data are yet available, are assumed similar to the changes related to the industrial sector. The way  
500 the decrease in solvent emissions are handled in the different studies could explain the large  
differences in the changes in NMVOCs emissions in different regions. European CO, SO<sub>2</sub> and BC  
emissions during the lockdowns decreased by an average of 8, 14 and 18 % in April, respectively,  
relative to the reference emissions.

505 The results show that the lockdowns led to an average reduction in NO<sub>x</sub> emissions in the USA of 21  
% (15-35 %) during April, while these values reach 43 % (30-53 %) in South American regions, 29  
% (22-42 %) in India and 13 % (8-19 %) in Africa. For NMVOCs, the maximum average reduction  
is 33 % (21-43 %) in the USA in April, while these values reach 55 % (27-57 %) in South America,  
14 % (2-19 %) in India and 19 % (1.4-21 %) in Africa. The largest variabilities in the different

510 countries in Africa and in India are due to the large uncertainties associated with the activity data in these countries. The magnitude of the changes in total emissions are not homogenous within the same region and can be very different from one location to another. Figure 9 displays the spatial distribution of the absolute difference between the COVID-19 and the reference scenarios as well as the associated percentage changes in NO<sub>x</sub> emissions for April 2020, the strictest lockdown period  
515 in most countries in the world. During that month, substantial declines in almost all geographical areas and for most species are seen, with the main decreases happening in urban areas strongly affected by human activities (Figure 9a and Figure S7 in Supplementary material). The percentage changes (Figure 9b) reflect the regional average decreases shown in Figure 8, with, for example, reductions over China ranging from 10 to 20 % in April 2020. It can be noticed that the emission  
520 changes in China result mostly from the significant decrease of emissions in the densely populated and heavily industrialized North China Plain and megacities. As already mentioned, the largest decreases in large cities of Europe, USA and India occurred in April, and in February in China (Figure S8, Supplementary material). The changes in total emissions of each species are driven by changes in the predominant sectors contributing to the emissions of each species. The observed  
525 decrease over the oceans from the different figures is related to the reduction in shipping activities in response to the slowdown of the economy.

These results provide a global overview of the effect of the lockdowns in the emissions of different atmospheric compounds in different regions of the world.

#### 530 **4 Conclusions**

The restrictions and lockdowns resulting from the COVID-19 pandemic since the end of January 2020 have had important social, economic and environmental consequences. In this study, we have provided an estimate of adjustment factors (AFs) to quantify the changes in global emissions of major atmospheric pollutants during the lockdown periods. This dataset can be easily applied to the  
535 emissions used in global and regional models to simulate the impacts of the reduced human activity on the atmospheric composition and climate. To this purpose, we analyzed activity data from various sectors representative of transportation (road, air and ship), industry, residential and power generation. The resulting dataset provides daily or monthly sectoral AFs on a 0.1°x0.1° resolution over the globe, and can be used to quantify regional patterns in the distributions of emissions during  
540 the COVID-19 pandemic. When applied to the CAMS-GLOB-ANT\_v4.2\_R1.1 emissions dataset, large changes are estimated, with maximum decreases in the total emissions in February in Eastern China, with an average reduction of 20-30 % in NO<sub>x</sub>, NMVOCs and SO<sub>2</sub> emissions. In other regions, the maximum changes occur in April, with average reductions of 20-30 % for NO<sub>x</sub>, NMVOCs and CO in Europe and North America and larger decreases (30-50 %) in South America. In both India  
545 and Africa, NO<sub>x</sub> emissions decline by 10 to 40 % while NMVOCs emissions decrease by 2 to 20 %, which larger uncertainties compared to other regions. For the others species, the maximum average reductions are generally less than 15 %, except in the South American countries where large decreases in CO and BC are estimated. These changes in the total emissions are related to the

550 different changes in the individual sectors. The patterns of the reductions also show large differences at the regional level.

We acknowledge that the lack of data for some activity sectors in several regions and the absence of accurate information on all sectors might cause significant uncertainties in the estimation of the AFs. For that reason, besides average values, the dataset also includes low and high estimation of the AFs for the period over January-August 2020. This should give an estimate of the uncertainties on the distribution of chemical species in models. The AFs obtained in this study will be extended at least until the end of the year 2020, and will be revised and updated as new or improved information on economic and mobility activity becomes available.

560

## Data availability

565 The average estimated daily/monthly gridded AFs for the sectors considered in this study, i.e. transportation including air traffic and shipping, industries, residential and power generation are available as NetCDF files for the global domain at a resolution of  $0.1^\circ \times 0.1^\circ$  resolution. A range of AF is provided at each grid point as the average  $\pm$  standard deviation. The acronym of the dataset is CONFORM (Covid-19 adjustmeNt Factors fOR eMissions), and it is available at <https://doi.org/10.25326/88>. The files can be openly accessed through the Emissions of atmospheric Compounds and Compilation of Ancillary Data (ECCAD) database with a login account (<https://eccad.aeris-data.fr/>). In the ECCAD database, the dataset can be directly accessed using the link: <https://eccad.aeris-data.fr/essd-conform/>.

575

## Author contributions.

**Conceptualization:** Thierno Doumbia, Claire Granier

**Production of emissions:** Thierno Doumbia, Sabine Darras, Nellie Elguindi, Claire Granier

580 **Collection of datasets:** Thierno Doumbia, Yiming Liu, Xiaoqin Shi, Tao Wang

**Analysis of the AFs results:** Thierno Doumbia, Idir Bouarar, Simone Tilmes, Benjamin Gaubert, Trissevgeni Stavrou

**Test of the AFs in atmospheric models:** Benjamin Gaubert, Adrien Deroubaix, Forrest Lacey

**Writing -original draft:** Thierno Doumbia, Claire Granier, Guy Brasseur

585 **Writing -review and editing:** All authors.

**Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgments.** We acknowledge the support of the AQ-WATCH European project, a  
590 HORIZON 2020 Research and Innovation Action (GA 870301), the ESA ICOVAC project, and the  
SEEDS EU project. The CAMS-GLOB-ANT dataset has been developed with the support of the  
CAMS (Copernicus Atmosphere Monitoring Service, <https://atmosphere.copernicus.eu/>), operated  
by the European Centre for Medium-Range Weather Forecasts on behalf of the European  
Commission as part of the Copernicus Programme. The ECCAD database ([eccad.aeris-data.fr](http://eccad.aeris-data.fr)) is  
595 supported by the French AERIS data infrastructure ([aeris-data.fr](http://aeris-data.fr)). This material is based upon work  
supported by the National Center for Atmospheric Research, which is a major facility sponsored by  
the US National Science Foundation under cooperative agreement no. 1852977. T.W. and Y.L.  
acknowledge support by the Hong Kong Research Grants Council (T24-504/17-N and A-  
PolyU502/16).

600

## References

- Barré, J., Petetin, H., Colette, A., Guevara, M., Peuch, V.-H., Rouil, L., Engelen, R., Inness, A.,  
Flemming, J., Pérez García-Pando, C., Bowdalo, D., Meleux, F., Geels, C., Christensen, J. H., Gauss,  
605 M., Benedictow, A., Tsyro, S., Friese, E., Struzewska, J., Kaminski, J. W., Douros, J., Timmermans, R.,  
Robertson, L., Adani, M., Jorba, O., Joly, M., and Kouznetsov, R.: Estimating lockdown induced  
European NO<sub>2</sub> changes, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-995>, in review,  
2020.
- 610 Bauwens, M. S. Compernelle, T. Stavrakou, J.-F. Müller, J. van Gent, H. Eskes, P. F. Levelt, R. van der  
A, J. P. Veefkind, J. Vlietinck, H. Yu, C. Zehner: Impact of coronavirus outbreak on NO<sub>2</sub> pollution  
assessed using TROPOMI and OMI observations, *Geophys. Res. Lett.*, 47,  
[doi:10.1029/2020GL087978](https://doi.org/10.1029/2020GL087978), 2020.
- 615 Biswal, A., V. Singh, S. Singh, A. P. Kesarkar, K. Ravindra, R. S. Sokhl, M. P. Chipperfield, S. S.  
Dhomse, R. J. Pope, T. Singh and S. Mor: COVID-19 lockdown induced changes in NO<sub>2</sub> levels across  
India observed by multi-satellite and surface observations, *Atmos. Chem. Phys.*,  
<https://doi.org/10.5194/acp-2020-1023>, 2020.
- 620 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S.,  
Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air  
pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987–2013,  
<https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- 625 Diamond, M. S. and R. Wood: Limited Regional Aerosol and Cloud Microphysical Changes Despite  
Unprecedented Decline in Nitrogen Oxide Pollution During the February 2020 COVID-19 Shutdown  
in China, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2020GL088913>, 2020.



- 630 Elguindi, N., Granier, C., Stavrakou, T., Darras, S., Bauwens, M., Cao, H., C. Chen, H.A.C. Denier van der Gon, O. Dubovik, T.M. Fu, D.K. Henze, Z. Jiang, S. Keita, J.J.P. Kuenen, J. Kurokawa, C. Lioussé, J.F. Müller, Z. Qu, F. Solmon, B. Zheng: Intercomparison of magnitudes and trends in anthropogenic surface emissions from bottom-up inventories, top-down estimates, and emission scenarios, *Earth's Future*, 8, e2020EF001520. doi:10.1029/2020EF001520, 2020.
- 635 Forster, P. M., H. I. Forster, M. J. Evans, M. J. Gidden, C. D. Jones, C. A. Keller, R. D. Lamboll, C. Le Quéré, J. Rogelj, D. Rosen, C.-F. Schleussner, T. B. Richardson, C. J. Smith and S. T. Turnock: Current and future global climate impacts resulting from COVID-19, *Nat. Clim. Chan.*, 10, 913-919, doi:10.1038/s41558-020-0883-0, 2020.
- 640 Gaubert, B., Bouarar, I., Doumbia, T., Liu, Y., Stavrakou, T., Deroubaix, A., Darras, S., Elguindi, N., Granier, C., Lacey, F., Müller, J.F., Shi, X., Tilmes, S., Wang, T. and Brasseur, G. P. (2021). Global changes in secondary atmospheric pollutants during the 2020 COVID-19 pandemic. *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034213. <https://doi.org/10.1029/2020JD034213>.
- 645 Granier, C., S. Darras, H. Denier van der Gon, J. Doubalova, N. Elguindi, B. Galle, M. Gauss, M. Guevara, J.-P. Jalkanen, J. Kuenen, C. Lioussé, B. Quack, D. Simpson, K. Sindelarova: The Copernicus Atmosphere Monitoring Service global and regional emissions, Copernicus Atmosphere Monitoring Service (CAMS) report, doi:10.24380/d0bn-kx16, 2019.
- 650 Guevara, M., O. Jorba, A. Soret, H. Petetin, D. Bowdalo, K. Serradell, C. Tena, H. Denier van der Gon, J. Kuenen, V.-H. Peuch, C. P. Garcia-Pando: Time-resolved emission reductions for atmospheric chemistry modelling in Europe during the COVID-19 lockdowns, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-2020-686>, 2020.
- 655 Han, P. F., Cai, Q. X., Oda, T., Zeng, N., Shan, Y. L., Lin, X. H., and Liu, D.: Assessing the recent impact of COVID-19 on carbon emissions from China using domestic economic data, *Sci. of the Tot. Env.*, 750, 2021.
- 660 He, Y., Pan, Y., Gu, M., Sun, Q., Zhang, Q., Zhang, R., and Wang, Y.: Changes of Ammonia Concentrations in Wintertime on the North China Plain from 2018 to 2020, *Atmos. Res.*, doi: <https://doi.org/10.1016/j.atmosres.2021.105490>, 2021. 105490, 2021.
- IEA (International Energy Agency), *Global Energy Review 2020*, IEA, Paris <https://www.iea.org/reports/global-energy-review-2020>, 2020.
- 665 Iacus, S.M., F. Natale, C. Santamaria, S. Spyrtatos, M. Vespe: Estimating and projecting air passenger traffic during the COVID-19 coronavirus outbreak and its socio-economic impact, *Saf. Sci.*, 129, Article 104791, 10.1016/j.ssci.2020.104791, 2020.

- 670 Keller, C. A., Evans, M. J., Knowland, K. E., Hasenkopf, C. A., Modekurty, S., Lucchesi, R. A., Oda, T., Franca, B. B., Mandarino, F. C., Díaz Suárez, M. V., Ryan, R. G., Fakes, L. H., and Pawson, S.: Global Impact of COVID-19 Restrictions on the Surface Concentrations of Nitrogen Dioxide and Ozone, *Atmos. Chem. Phys. Discuss.* <https://doi.org/10.5194/acp-2020-685>, in review, 2020.
- 675 Kim, H. C., Kim, S., Cohen, M., Bae, C., Lee, D., Saylor, R., Bae, M., Kim, E., Kim, B.-U., Yoon, J.-H., and Stein, A.: Quantitative assessment of changes in surface particulate matter concentrations over China during the COVID-19 pandemic and their implications for Chinese economic activity, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-821>, in review, 2020.
- 680 Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos. Chem. Phys.*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.
- 685 Kraemer, M.U.G., Yang, C.-H., Gutierrez, B., Wu, C.-H., Klein, B., Pigott, D. M., open COVID-19 data working group, L. du Plessis, Faria, N. R., Li, R., Hanage, W. P., Brownstein, J. S., Layan, M., Vespignani, A., Tian, H., Dye, C., Cauchemez, S., Pybus, O., Scarpino, S. V.: The effect of human mobility and control measures on the COVID-19 epidemic in China, <https://doi.org/10.1126/science.abb4218>, 2020.
- 690 Lee, J. D., W. S. Drysdale, D. P. Finch, S. E. Wilde and P. I. Palmer: UK surface NO<sub>2</sub> levels dropped by 42 % during the COVID-19 lockdown: impact on surface O<sub>3</sub>, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2020-838, 2020.
- 695 Le Quéré, C., R. B. Jackson, M. W. Jones, A. J. P. Smith, S. Abernethy, R. M. Andrew, A. J. De-Gol, D. R. Willis, Y. Shan, J. G. Canadell, P. Friedlingstein, F. Creutzig and G. P. Peters: Temporary reduction in daily global CO<sub>2</sub> emissions during the COVID-19 forced confinement, *Nat. Clim. Chang.* 10, 647–665, doi:10.1038/s41558-020-0797-x, 2020.
- 700 Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., Tong, D., Zheng, B., Cui, H., Man, H., Zhang, Q., and He, K.: Anthropogenic emission inventories in China: a review, *Natl. Sci. Rev.*, 4, 834-866, doi:10.1093/nsr/nwx150, 2017.
- 705 Lin, J.T.; McElroy, M.B.: Detection from space of a reduction in anthropogenic emissions of nitrogen oxides during the Chinese economic downturn, *Atmos. Chem. Phys.*, 11, 8171-8188, 2011.
- 710 Liu, Z., Ciais, P., Deng, Z., Lei, R., Davis, S. J., Feng, S., Zheng, B., Cui, D., Dou, X., Zhu, B., Guo, R., Ke, P., Sun, T., Lu, C., He, P., Wang, Y., Yue, X., Wang, Y., Lei, Y., Zhou, H., Cai, Z., Wu, Y., Guo, R., Han, T., Xue, J., Boucher, O., Boucher, E., Chevallier, F., Tanaka, K., Wei, Y., Zhong, H., Kang, C., Zhang, N., Chen, B., Xi, F., Liu, M., Bréon, F.-M., Lu, Y., Zhang, Q., Guan, D., Gong, P., Kammen, D. M., He, K., Schellnhuber, H. J.: Near-real-time monitoring of global CO<sub>2</sub> emissions


- reveals the effects of the COVID-19 pandemic, *Nat. Commun.* 11, 5172, <https://doi.org/10.1038/s41467-020-18922-7>, 2020.
- 715 Miyazaki, K., Bowman, K., Sekiya, T., Jiang, Z., Chen, X., Eskes, H., M. Ru, Y. Zhang, and D. Shindell: Air quality response in China linked to the 2019 novel coronavirus (COVID-19) lockdown, *Geophys. Res. Lett.*, 47, e2020GL089252. <https://doi.org/10.1029/2020GL089252>, 2020.
- 720 Shi, X. and G. P. Brasseur: The response in air quality to the reduction of Chinese Economic Activities during the COVID-19 Outbreak, *Geophys. Res. Lett.*, 47, 11, doi:10.1029/2020GL088070, 2020.
- 725 Sun, Y., Lei, L., Zhou, W., Chen, C., He, Y., Sun, J., Li, Z., Xu, W., Wang, Q., Ji, D., Fu, P., Wang, Z., and Worsnop, R. D.: A chemical cocktail during the COVID-19 outbreak in Beijing, China: Insights from six-year aerosol particle composition measurements during the Chinese New Year holiday, *Sci. of The Tot. Env.*, 742, 2020.
- 730 Venter, Z.S., K. Aunan, S. Chowdhury, and J. Lelieveld: COVID-19 lockdowns cause global air pollution declines, *Proc. Nat. Acad. of Sci.*, 117 (32) 18984-18990, doi:10.1073/pnas.2006853117, 2020.
- 735 Xing, J., Li, S., Jiang, Y., Wang, S., Ding, D., Dong, Z., Zhu, Y., and Hao, J.: Quantifying the emission changes and associated air quality impacts during the COVID-19 pandemic in North China Plain: a response modeling study, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-522>, in review, 2020
- 740 Zhang, R., Y. Zhang, H. Lin, X. Feng, T.M. Fu, and Y. Wang: NO<sub>x</sub> Emission Reduction and Recovery during COVID-19 in East China, *Atmosphere*, 11, 433, DOI:10.1038/s41558-020-0797-x, 2020.
- 745 Zhang, Q., Pan, Y., He, Y., Walters, W. W., Ni, Q., Liu, X., Xu, G., Shao, J., and Jiang, C.: Substantial nitrogen oxides emission reduction from China due to COVID-19 and its impact on surface ozone and aerosol pollution, *Sci. of The Tot. Env.*, 753, 142238, 2021.
- 750 Zheng, B., Geng, G. N., Ciais, P., Davis, S. J., Martin, R. V., Meng, J., Wu, N. N., Chevallier, F., Broquet, G., Boersma, F., van der Ronald, A., Lin, J. T., Guan, D. B., Lei, Y., He, K. B., and Zhang, Q.: Satellite-based estimates of decline and rebound in China's CO<sub>2</sub> emissions during COVID-19 pandemic, *Sci. Adv.*, 6, 2020.
- 755 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095–14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.

## Tables


Table 1: Data sources of activity data used to estimate the emission AFs. In this table, Rest Of World (ROW) refers to all world countries except China.

| Sectors                    | Country/<br>Region                | Data  | Data sources  |
|----------------------------|-----------------------------------|---|---|
| Road transport (TRO)       | China                             | Baidu Migration Scale Index (daily)<br>TomTom Congestion Index <sup>1</sup> (weekly)<br><small><sup>1</sup>TomTom congestion traffic index are used to evaluate Baidu migration scale index</small>               | China Data Lab, 2020, "Baidu Mobility Data", <a href="https://doi.org/10.7910/DVN/FAEZIO">https://doi.org/10.7910/DVN/FAEZIO</a><br><a href="https://www.tomtom.com/en_gb/traffic-index/ranking">https://www.tomtom.com/en_gb/traffic-index/ranking</a> |
|                            | Rest Of World (ROW)               | Google's transit stations (daily)<br>Apple mobility <sup>2</sup> (daily)<br><small><sup>2</sup>Apple mobility data are used to evaluate Google activity data</small>  | <a href="https://www.google.com/covid19/mobility/">https://www.google.com/covid19/mobility/</a><br><a href="https://covid19.apple.com/mobility">https://covid19.apple.com/mobility</a>  |
| Residential (RES)          | China                             | Emissions from residential sector (daily)   | Le Quéré et al. (2020)  |
|                            | ROW                               | Google's residential (daily)  | <a href="https://www.google.com/covid19/mobility/">https://www.google.com/covid19/mobility/</a>   |
| Industrial processes (IND) | China                             | Coal consumption from the six main coal producers (daily)   | Liu et al. (2020)   |
|                            | ROW                               | Google's workplaces (daily)   | <a href="https://www.google.com/covid19/mobility/">https://www.google.com/covid19/mobility/</a>   |
|                            | World (ROW + China)               | Crude steel production <sup>3</sup> (monthly)<br><small><sup>3</sup>Monthly used to help in the analysis of the adjustment factors in the industrial sector, estimated from Google's workplaces category.</small> | <a href="https://www.worldsteel.org/">https://www.worldsteel.org/</a>   |
| Power Generation (ENE)     | India                             | Production of Coal, Lignite, and Gas Naphtha and Diesel (daily)   | Power System Operation Corporation Limited ( <a href="https://posoco.in/reports/daily-reports/">https://posoco.in/reports/daily-reports/</a> )  |
|                            | USA (regional data)               | Regional electricity load (daily)   | Energy Information Administration (EIA) ( <a href="https://www.eia.gov/beta/states/states/ca/data/dashboard/electricity">https://www.eia.gov/beta/states/states/ca/data/dashboard/electricity</a> )   |
|                            | Europe                            | Total electricity load (daily)  | ENTSO-E Transparent platform ( <a href="https://transparency.entsoe.eu/dashboard/">https://transparency.entsoe.eu/dashboard/</a> )  |
|                            | Brazil                            | Thermal Production (daily)  | Operator of the National Electricity System ( <a href="http://www.ons.org.br/Paginas/">http://www.ons.org.br/Paginas/</a> )   |
|                            | Russia                            | Power Generation (daily)  | United Power System of Russia ( <a href="http://www.so-ups.ru/index.php">http://www.so-ups.ru/index.php</a> )   |
|                            | Singapore                         | Electricity demand (daily)  | <a href="https://www.ema.gov.sg/Statistics.aspx">https://www.ema.gov.sg/Statistics.aspx</a>   |
|                            | Canada                            | Electricity demand Ontario (daily)  | <a href="http://reports.ieso.ca/public/Demand/">http://reports.ieso.ca/public/Demand/</a>   |
|                            | Other regions (e.g. Asia, Africa) | Emissions from Power sector (daily)   | Forster et al. (2020)   |
| Air transportati on (AVI)  | China and ROW                     | Projection of air traffic volume (monthly)  | Knowledge Center on Migration and Demography (KCMD) Dynamic Data Hub <a href="https://bluehub.jrc.ec.europa.eu/migration/app/index.html#">https://bluehub.jrc.ec.europa.eu/migration/app/index.html#</a>  |
| Shipping (SHP)             | World                             | Container shipping (monthly)  | United Nation Conference on Trade and Development (UNCTAD) <a href="https://unctad.org/news/covid-19-shipping-data-hints-some-recovery-global-trade">https://unctad.org/news/covid-19-shipping-data-hints-some-recovery-global-trade</a>                |

## Figures




760 Figure 1: Schematic view of the different steps for estimating the emission adjustment factors (AFs).




765

Figure 2: Daily AFs for the road transportation sector from January to August 2020 over Europe, USA, South America, China, Africa and the rest of the world. These estimations are derived from Google's transit category. The standard deviation values (light pink) result from the AFs for the individual countries or states/provinces. The light blue color indicates the range of the minimum and maximum values.

770




775 Figure 3: AF time series for air traffic and shipping as a function of month from January to August 2020. The standard deviation values (light pink) are calculated, based on the AFs of all individual countries, while the light blue color indicates the range of the minimum and maximum values.



780 Figure 4: Daily AFs for the industrial sector from January to August 2020 over Europe, USA, South America, China, Africa and the rest of the world. These estimations are derived from Google's workplaces category. The standard deviation values (light pink) result from the AFs for the individual countries or states/provinces. The light blue color indicates the range of the minimum and maximum values.


785




790 Figure 5: Daily AFs for residential sector from January to August 2020 over Europe, USA, South America, China, Africa and the rest of the world. These estimations are derived from Google’s residential category. The standard deviation values (light pink) result from the AFs for the individual countries or states/provinces. The light blue color indicates the minimum and maximum values.

795






800 Figure 6: Daily AFs for power sector from January to August 2020 over Europe, USA, South America, China, Africa and the rest of the world. These estimations are derived from multiple data sources (Table 1). The standard deviation values (light pink) result from the AFs for the individual countries or states. The light blue color indicates the range of the minimum and maximum values.




805

Figure 7: Global percentage change in total emissions (combination of emissions from ground transportation, industry, power, residential and shipping) of the main atmospheric compounds from January to June 2020. The vertical lines show the uncertainties associated to the estimated regional AFs.

810



815 Figure 8: Percentage change in total emissions (combination of emissions from ground transportation, industry, power, residential and shipping), as a function of month for selected regions: a) Europe (35°N-70°N, 20°W-20°E), b) Eastern China (20°N-45°N, 80°E-125°E), c) South America (60°S-20°N, 90°S-35°S), d) North America (20°N-50°N, 135°W-35°W), e) Africa (40°S-30°N, 20°W-40°E) and f) India (05°N-30°N, 60°E-90°E). The vertical lines show the uncertainties associated to the estimated regional AFs.



820

Figure 9: Spatial distribution of a) the absolute change and b) the percentage change in total NO<sub>x</sub> emission (combination of emissions from ground transportation, industry, power, residential and shipping) for April 2020.

825