

Interactive comment on “A global monthly climatology of oceanic total dissolved inorganic carbon: a neural network approach” by Daniel Broullón et al.

Anonymous Referee #3

Received and published: 26 May 2020

The authors provide a nice description of global TCO₂ distribution based on monthly climatology from a neural network approach. The addition input variable “year” is reasonable and important for TCO₂. It is great to see that the neural network outputs match very well with the measured TCO₂ for the independent time series locations. This manuscript is well organized and easy to follow. I would like to see the publication of this work.

Below are specific comments.

Lines 71-74, include the influence of mixing on TCO₂ variability. For example, upwelling-induced increase in TCO₂, which is described in lines 83-86.

Printer-friendly version

Discussion paper



Lines 75-76, the temperature and salinity influence on TCO₂ through the modification of the solubility of CO₂ affects the seawater pCO₂ (which is almost instantaneous) and thus the air-sea CO₂ flux, which eventually drives the change in TCO₂ over time.

Line 131, delete de in “minimize de errors”

Lines 142-143, move the full list of the input variables to line 153 to make it ahead of the sentence, “In addition to the target variable. . .”

Line 147, specify which generalization method was used to prevent overfitting.

Line 258, provide how many hidden layers are used in the neural network

Line 263, the errors cannot be avoided in any case. It is likely that a much smaller weight is assigned to a variable that contains a large error.

Line 276, provide how depth-weighted RMSE is calculated

Lines 349-351, the error may come from the bias in total alkalinity. I agree that the error involved in the calculation of pCO₂ from AT and TCO₂ can be large, but the monthly averaged error is another case. In addition, the choice of K1 and K2 affects the calculated pCO₂. Lueker et al. (2000) is a better choice in the case of calculating pCO₂. Provide the bias information for total alkalinity in each region in Table 4.

Line 420-430, Fig. 9, the differences in the surface water at BATS in May, June, July, and August seem large as compared to the values in other months. Provide an explanation.

Line 445-448, the bias for pCO₂, considering it is the averaged value over space and time, seems large as compared to the bias in Landschutzer et al. (2017) as shown in Table 4. Therefore, I don't agree that climatologies of other seawater CO₂ system variables can be confidently computed. Furthermore, this statement compromises the value of this study if TCO₂ can be confidently computed from pCO₂ and AT.

Fig. 2, color-coded depth instead of the log₁₀ frequency would provide more informa-

[Printer-friendly version](#)[Discussion paper](#)

tion regarding the distribution of errors. Fig. S4, add another panel with a similar generated from measured bottle TCO₂ data. Furthermore, add one more panel to show the differences between the derived TCO₂ and the bottle TCO₂. The differences are partially shown in Table 2 with the average differences by region. An additional panel regarding the differences would help to determine the reliability of this new dataset on a smaller scale (e.g., 1 degree by 1 degree).

Interactive comment on Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2020-37>, 2020.

[Printer-friendly version](#)

[Discussion paper](#)

