



## A global monthly climatology of oceanic total dissolved inorganic carbon: a neural network approach

Daniel Broullón<sup>1</sup>, Fiz F. Pérez<sup>1</sup>, Antón Velo<sup>1</sup>, Mario Hoppema<sup>2</sup>, Are Olsen<sup>3</sup>, Taro Takahashi<sup>4,†</sup>, Robert M. Key<sup>5</sup>, Toste Tanhua<sup>6</sup>, J. Magdalena Santana-Casiano<sup>7</sup> and Alex  
5 Kozyr<sup>8</sup>

<sup>1</sup>Instituto de Investigaciones Marinas, CSIC, Eduardo Cabello 6, 36208 Vigo, Spain

<sup>2</sup>Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Postfach 120161, 27515 Bremerhaven, Germany

10 <sup>3</sup>Geophysical Institute, University of Bergen and Bjerknes Centre for Climate Research, Allégaten 70, 5007 Bergen, Norway

<sup>4</sup>Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY 10964, USA

<sup>5</sup>Atmospheric and Oceanic Sciences, Princeton University, 300 Forrestal Road, Sayre Hall, Princeton, NJ 08544, USA

<sup>6</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20D-24105 Kiel, Germany

15 <sup>7</sup>Instituto de Oceanografía y Cambio Global, IOCAG, Universidad de Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain

<sup>8</sup>NOAA National Centers for Environmental Information, 1315 East-West Hwy Silver Spring, MD 20910 USA

<sup>†</sup>Deceased

20

*Correspondence to:* Daniel Broullón (dbroullon@iim.csic.es)

### Abstract




Anthropogenic emissions of CO<sub>2</sub> to the atmosphere have modified the carbon cycle for more than two centuries. As the ocean stores most of the carbon on our planet, there is an important task in unraveling the  
25 natural and anthropogenic processes that drive the carbon cycle at different spatial and temporal scales. We contribute to this by designing a global monthly climatology of total dissolved inorganic carbon (TCO<sub>2</sub>) which offers a robust basis in carbon cycle modeling but also for other studies related to this cycle. A feedforward neural network (dubbed NNGv2LDEO) was configured to extract from the Global Ocean Data Analysis Project version 2.2019 (GLODAPv2.2019) and the Lamont-Doherty Earth Observatory (LDEO)  
30 datasets the relations between TCO<sub>2</sub> and a set of variables related to the former's variability. The global root-mean-squared error (RMSE) of mapping TCO<sub>2</sub> is relatively low for the two datasets (GLODAPv2.2019: 7.2 μmol kg<sup>-1</sup>; LDEO: 11.4 μmol kg<sup>-1</sup>) and also for independent data, suggesting that the network does not overfit possible errors in data. The ability of NNGv2LDEO in capturing the monthly



variability of TCO<sub>2</sub> was testified through the good reproduction of the seasonal cycle in ten time-series  
35 stations spread over different regions of the ocean (RMSE: 3.6 to 13.1 μmol kg<sup>-1</sup>). The climatology was  
obtained by passing through NNGv2LDEO the monthly climatological fields of temperature, salinity and  
oxygen from World Ocean Atlas 2013, and phosphate, nitrate and silicate computed from a neural network  
fed with the previous fields. The resolution is 1° x 1° in the horizontal, 102 depth levels (0-5500m) and  
40 monthly (0-1500 m) to annual (1550-5500 m), and it is centered in the year 1995. The uncertainty of the  
climatology is low when compared with climatological values derived from measured TCO<sub>2</sub> in the largest  
time-series stations. Furthermore, a computed climatology of partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>) from a  
previous climatology of total alkalinity and the present one of TCO<sub>2</sub> supports the robustness of this product  
through the good correlation with a widely used pCO<sub>2</sub> climatology (Landschützer et al., 2016). Our TCO<sub>2</sub>  
climatology is distributed through the data repository of the Spanish National Research Council (CSIC;  
45 <http://dx.doi.org/10.20350/digitalCSIC/10551>, Broullón et al., 2020).

## 1 Introduction

The ocean is the major carbon reservoir of the Earth. Most of this carbon occurs as dissolved inorganic  
carbon (TCO<sub>2</sub>, also known as DIC or C<sub>T</sub>) (Ciais et al., 2013; Tanhua et al., 2013). Three species make up  
TCO<sub>2</sub>: dissolved CO<sub>2</sub> (generally considered as the sum of the dissolved CO<sub>2</sub> itself (CO<sub>2</sub>(aq)) and carbonic  
50 acid (H<sub>2</sub>CO<sub>3</sub>)), bicarbonate ion (HCO<sub>3</sub><sup>-</sup>) and carbonate ion (CO<sub>3</sub><sup>2-</sup>). The relative concentrations of these  
species with respect to each other determine the seawater pH (Zeebe and Wolf-Gladrow, 2001). The  
seawater CO<sub>2</sub> chemistry system can be represented as a set of chemical equilibria reactions that describes  
the speciation of the various ions of TCO<sub>2</sub> as follows:



Since the Industrial Revolution, the concentration of TCO<sub>2</sub> in the global ocean has increased, generally to  
a certain depth level (depending on the particular processes in each ocean area) due to the entry of CO<sub>2</sub> into  
60 the seawater from the atmosphere (Sarmiento and Gruber, 2002; Doney et al., 2009; Vázquez-Rodríguez et  
al., 2009; Bates et al., 2012; Sallée et al., 2012; Khatiwala et al., 2013). The uptake is driven by the  
increasing partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>) in the atmosphere relative to the ocean, generated by the  
anthropogenic emissions of CO<sub>2</sub> that cause an annual net flux of this gas into the ocean (Le Quéré et al.,  
2018). Accompanying the change in TCO<sub>2</sub>, the pH and carbonate ion concentration have been declining  
65 because of the anthropogenic process previously mentioned, being these changes reflected in the  
proportions of the chemical species of TCO<sub>2</sub> (Kleypas and Langdon, 2000; Orr et al., 2005). These changes  
in seawater chemistry framed in the ocean acidification process can negatively influence various processes



involving marine organisms such as calcification, growth and survival (Orr et al., 2005; Fabry et al., 2008; Hendriks et al., 2010; Hoegh-Guldberg and Bruno, 2010; Kroeker et al., 2013).

70 In addition to the secular trends driven by the uptake of anthropogenic CO<sub>2</sub>, ocean TCO<sub>2</sub> varies both temporally and spatially as a consequence of several natural processes. This variability may reach values of 15% of the mean TCO<sub>2</sub> value in the ocean (Lee et al., 2000). The processes that increase TCO<sub>2</sub> are: net flux of CO<sub>2</sub> from the atmosphere to the ocean, organic matter remineralization and the dissolution of calcium carbonate (CaCO<sub>3</sub>). The processes that reduce TCO<sub>2</sub> are: net flux of CO<sub>2</sub> from the ocean to the atmosphere, primary production and calcification. In the surface ocean, the main variables influencing de  
75 TCO<sub>2</sub> variability are temperature and salinity (Weiss et al., 1982; Lee et al., 2000; Wu et al., 2019) through the modification of the solubility of CO<sub>2</sub>, and therefore, the seawater pCO<sub>2</sub>. Nutrients and oxygen can also reflect the processes that modify the concentration of TCO<sub>2</sub> through their consumption and release, like during the cycling of organic matter (Körtzinger et al., 2001; Bauer et al., 2013). From products generated with measured data (Key et al., 2004; Takahashi et al., 2014; Lauvset et al., 2016) and in modeling studies (e.g., Doi et al., 2015), it is known that the global surface distribution of TCO<sub>2</sub> follows a zonal gradient: there is a reduction of its concentration from the poles to the equator reflecting the processes that control its variability. Key et al. (2004) emphasize that this distribution is associated to the distribution pattern of nutrients. Recently, Wu et al. (2019) found that the distribution of surface salinity-normalized TCO<sub>2</sub> (nDIC)  
80 has two main drivers: temperature and upwelling. At depth, the variation shown in almost any measured profile of TCO<sub>2</sub> mainly reflects the remineralization of organic matter and, to a lesser extent, the dissolution of CaCO<sub>3</sub>, resulting in an increase in TCO<sub>2</sub> from the surface to the intermediate depths.

Understanding the distribution and variability of TCO<sub>2</sub> in the ocean and its secular trends driven by anthropogenic carbon uptake is needed to assess the magnitude and possible impacts of ocean acidification.  
90 It is also necessary for the evaluation of numerical models that include the carbon cycle and their estimates of past, current and future ocean carbon cycle behavior (e.g., Yool et al., 2013; Aumont et al., 2015; Butenschön et al., 2016; Le Quéré et al., 2016; Goris et al., 2018). Seasonality of TCO<sub>2</sub> and the horizontal and vertical variability underscore the necessity to design a climatology with both monthly and spatial resolutions according to the processes that influence this variable on a global scale. The existing climatologies of TCO<sub>2</sub> do not include all these characteristics collected together. Key et al. (2004) and Lauvset et al. (2016) built an annual climatology in 33 depth levels using interpolation techniques over data from Global Ocean Data Analysis Project version 1 (GLODAPv1) (Key et al., 2004) and GLODAPv2 (Key et al., 2015; Olsen et al., 2016), respectively. Takahashi et al. (2014) published a monthly climatology for the surface ocean computed from climatologies of pCO<sub>2</sub> and total alkalinity (A<sub>T</sub>). Other studies used the  
100 covariability between TCO<sub>2</sub> and other more commonly measured variables discussed above for mapping/gap-filling via empirical regressions and neural networks. Lee et al. (2000) used temperature and nitrate to compute surface nDIC with an area-weighted error of ±7 μmol kg<sup>-1</sup>. Sauzède et al. (2017) and Bittig et al. (2018) trained neural networks with GLODAPv2 data to compute TCO<sub>2</sub> over the depth range 0-8000 m with an accuracy of ±9 μmol kg<sup>-1</sup> and ±7.1 μmol kg<sup>-1</sup>, respectively. The predictor variables used  
105 in those studies were location, pressure, temperature, salinity and dissolved oxygen.



In the present study, we introduce the use of neural networks for going one step further in the design of a climatology. We have generated a climatology of  $\text{TCO}_2$  with a resolution consistent with that of the climatology of  $A_T$  of Broullón et al. (2019): horizontal resolution of  $1^\circ \times 1^\circ$ , 102 depth levels between 0 and 5500 m and a monthly (0-1500 m) and annual (1550-5500 m) temporal resolution. The availability of global databases containing variables of the seawater  $\text{CO}_2$  system with more and more data (e.g., GLODAPv2.2019, Lamont-Doherty Earth Observatory database (LDEO; Takahashi et al., 2017), Surface Ocean  $\text{CO}_2$  Atlas (SOCAT; Bakker et al., 2016)) and the great ability of the neural networks to interpolate as shown in other climatology studies about  $\text{CO}_2$  system variables (Landschützer et al., 2014; Broullón et al., 2019), show the appropriateness of this approach for generating a global monthly climatology covering more than the surface ocean.

## 2 Methodology

### 2.1 Neural network design

A feed-forward neural network was configured to compute  $\text{TCO}_2$  throughout the entire global ocean and to create a global climatology based on the good results previously obtained with this method in similar studies (e.g., Broullón et al., 2019). Briefly, a neural network of this type (Fig. S1a) is used to extract relationships between a set of input variables and a target one through a training process. At this stage, the inputs are passed through different parallel layers composed by a tunable number of neurons to reach values as closest as possible to the target ones (Fig. S1a). Initially, all inputs enter in each neuron of the first layer where they are being multiplied by different weights depending on the neuron they go. Inside the neurons (Fig. S1b), the results of the previous operation are summed and a bias is added. The obtained value inside each neuron is passed through an activation function which yields an output. The outputs of each neuron in each layer go to the following layer suffering the same process described to this point. In the last layer, which is composed by one neuron, a unique value for the target variable is calculated for each pair of inputs-target. This value is compared to the desired one and the difference between both values is backpropagated through the entire network in order to adjust the weights and biases, and to start again the processes and reach an accurate output value after multiple iterations. A complete description of the most common algorithms used to backpropagate and minimize de errors can be founded in Rumelhart et al. (1986), Levenberg (1944) and Marquardt (1963).

The method used here is equivalent to that fully described by Broullón et al. (2019) for  $A_T$ . In addition to the target variable ( $\text{TCO}_2$  instead of  $A_T$ ), the main changes in the present study compared to that of Broullón et al (2019) are the inclusion of the input variable “year”, accounting for the anthropogenic increase of the  $\text{TCO}_2$  pool, and the use of the  $p\text{CO}_2$  database from LDEO (Takahashi et al., 2017) in addition to the extended GLODAPv2.2019 (Olsen et al. 2019) to enable more robust  $\text{TCO}_2$  estimates in the surface ocean. Similar to Broullón et al. (2019), the neural networks were trained using the Levenberg-Marquardt method (Levenberg, 1944; Marquardt, 1963) through the *trainlm* function (detailed in Beale et al., 2018) in MATLAB. The splitting of the database used in this study (see Sect. 2.2) in the sets needed for training and testing the network is depicted in Fig. 1. The data were randomly associated to each dataset to capture (training) and evaluate (test) all possible variability. The input variables are temperature, salinity,



phosphate, nitrate, silicate, oxygen, sample position and year (Fig. S1a). The number of neurons tested to  
145 find the best neural network was 16, 32, 64, 128 and 256. Ten networks were trained for each number of  
neurons. The criteria to select the final number of neurons are based on a trade-off between the root-mean-  
squared error (RMSE — between the measured  $\text{TCO}_2$  and that estimated by the neural network) on the one  
hand, and the generalization of the network (to prevent overfitting) as in Broullón et al. (2019) on the other  
hand. Furthermore, an additional criterion based on the influence of each input variable on the  $\text{TCO}_2$   
150 extracted with the connection weight approach (Olden and Jackson, 2002) was followed to ensure that  
biogeochemical input variables have a larger influence on the  $\text{TCO}_2$  estimates than the input variables  
related to sample position for selecting a proper network.

## 2.2 Data

We included the LDEO database version 2016 (Takahashi et al., 2017;  
155 [https://www.nodc.noaa.gov/ocads/oceans/LDEO\\_Underway\\_Database](https://www.nodc.noaa.gov/ocads/oceans/LDEO_Underway_Database), last access: 13 November 2017),  
because it contains significantly more data in the surface layer than GLODAPv2.2019. Since the higher  
variability in the surface layer may lead to high errors in modeling variables of the seawater  $\text{CO}_2$  system  
(e.g., Carter et al., 2018; Bittig et al., 2018; Broullón et al., 2019), including the LDEO database should  
force the network to reach a more robust fit. The idea is that these additional data probably have more  
160 different relationships between input variables and  $\text{TCO}_2$  to help the neural network to adequately capture  
spatiotemporal variability. The  $\text{pCO}_2$ , temperature and salinity data from LDEO were monthly-averaged  
for each year in a  $1^\circ \times 1^\circ$  grid. The points where the standard deviation (std) of the averaged  $\text{pCO}_2$ ,  
temperature and salinity were greater than  $\pm 20 \mu\text{atm}$ ,  $1.5^\circ\text{C}$  and 0.5, respectively, were discarded, since the  
objective is to capture the monthly variability and therefore an extremely high sub-monthly variability could  
165 lead to errors. To obtain  $\text{TCO}_2$  values from the LDEO data, an additional variable of the  $\text{CO}_2$  system is  
necessary, for which we take  $A_T$  computed using the neural network NNGv2 of Broullón et al. (2019). The  
input variables required by NNGv2 were obtained from: 1) temperature and salinity from LDEO; 2) filtered  
oxygen from World Ocean Atlas version 2013 (WOA13; see Broullón et al., 2019); 3) phosphate, nitrate  
and silicate computed with CANYON-B (Bittig et al., 2018) using the previous variables as inputs. Finally,  
170  $\text{TCO}_2$  was calculated from this  $A_T$  and the averaged  $\text{pCO}_2$  using the MATLAB-version of the  $\text{CO}_2\text{SYS}$   
program (van Heuven et al., 2011); we used the dissociation constants of Mehrbach et al. (1973) (as refit  
by Dickson and Millero, 1987) and the borate dissociation constant of Dickson (1990). Note that we used  
this software and set of constants for all seawater  $\text{CO}_2$  chemistry calculations in the present study. The thus  
calculated  $\text{TCO}_2$  and the associated input variables were used as a part of the training and testing data for  
175 the neural networks created here. The final number of data points derived from LDEO was 54572.

To represent interior ocean conditions, the GLODAPv2.2019 database (Olsen et al., 2019) was added to  
the LDEO dataset for training and testing the neural network. Only samples which had data for all input  
variables and  $\text{TCO}_2$  were used. This database was included in two ways: 1) Only samples where all  
variables passed the quality control ( $n=287953$ ) (Olsen et al., 2016; Olsen et al., 2019; hereafter abbreviated  
180 Gv2QC) and 2) all samples ( $n=321647$ ) (hereafter abbreviated Gv2). Therefore, two neural networks  
options were trained and tested: NNGv2QLDEO and NNGv2LDEO, respectively.



### 2.3 Comparison of methods

We compared our method with CANYON-B of Bittig et al. (2018), where also  $\text{TCO}_2$  values were computed from multiple input variables. Both methods are based on neural networks but with certain differences as summarized in Table 1.

An error analysis was carried out in the same areas for which this was done by Broullón et al. (2019) for  $A_T$  and in several depth ranges (0-50 m, 50-200 m, 200-500m, 500-1000 m and 1000 m-bottom) for the two methods (our method and CANYON-B) and for the two datasets (Gv2QC and LDEO). The Gv2QC database was analyzed in this section instead of Gv2 because in the designing of CANYON-B only quality-controlled data were included. The analysis of CANYON-B using the LDEO dataset is useful to evaluate the validity of the approach followed by converting  $\text{pCO}_2$  to  $\text{TCO}_2$  since CANYON-B has not been trained with this dataset.

Computed  $\text{pCO}_2$  from  $A_T$  and  $\text{TCO}_2$  derived from neural networks was also evaluated in the LDEO dataset to assess the adequacy of including this dataset in our approach and to assess the ability of NNGv2 of Broullón et al. (2019) and the present  $\text{TCO}_2$  neural network to compute other variables of the seawater  $\text{CO}_2$  system. Furthermore, we compared the magnitude of the errors with the ones obtained by Landschützer et al. (2014), in which  $\text{pCO}_2$  is computed directly with a neural network, to evaluate the accuracy of our computed  $\text{pCO}_2$ .

### 2.4 Validation

In addition to the ability of computing  $\text{TCO}_2$  using the Gv2 and LDEO test sets, the neural network has been tested using independent data from ten ocean time series, located in different regions of the world ocean (data were obtained from [https://www.nodc.noaa.gov/ocads/oceans/time\\_series\\_moorings.html](https://www.nodc.noaa.gov/ocads/oceans/time_series_moorings.html), last access: 4 June 2019): Hawaii Ocean Time-series (HOT ALOHA and HOT ALOHA SURFACE; Dore et al., 2009), Bermuda Atlantic Time-series Study (BATS; Bates et al., 2012), European Station for Time-series in the Ocean at the Canary Islands (ESTOC; González-Dávila et al., 2010), Iceland Sea Time-series (ICELAND; Olafsson et al., 2010) Irminger Sea Time-series (IRMINGER; Olafsson et al., 2010), Kyodo North Pacific Ocean Time-series (KNOT; Wakita et al., 2010), K2 (Wakita et al., 2010), Ocean Weather Station Mike (OWS; Gislefoss et al., 1998) and Kerguelen Islands in the Indian sector of the Southern Ocean (KERFIX; Jeandel et al., 1998). The data from these time series allow us to test the ability of the neural network to reconstruct not only the seasonal variability of  $\text{TCO}_2$  at the various locations and depths sampled, but also its long-term trends. CANYON-B was also used to compute  $\text{TCO}_2$  in the time series to show the differences between that method and ours. The  $\text{TCO}_2$  values were obtained by feeding the neural networks with the measured values of the input variables at each time series. For the trend analyses, the measured and estimated  $\text{TCO}_2$  values were deseasonalized following Bates et al. (2014).

As an additional test,  $\text{pCO}_2$  calculated from measured  $\text{TCO}_2$  and  $A_T$  at the time series stations were compared with  $\text{pCO}_2$  calculated from the neural network generated values of  $A_T$  and  $\text{TCO}_2$ . This provides



insight on the combined performance of the two networks. Furthermore, we compared the magnitude of the errors to that obtained by Landschützer et al. (2014) for some of the time series.

## 2.5 Climatology of TCO<sub>2</sub>

220 We used the selected network, based on the results of the analyses described above, to construct a  
climatology of TCO<sub>2</sub>. Climatologies of the input variables were passed through the network to obtain the  
climatological fields of TCO<sub>2</sub>. The spatiotemporal resolution of the product is determined by that of the  
climatologies used as inputs: 1° x 1° horizontal resolution, 102 first depth levels of the WOA13 and monthly  
(for 0-1500 m depth) to annual (for 1550-5500 m depth) temporal resolution. Temperature and salinity  
225 climatologies were obtained from WOA13 objectively analyzed fields (Locarnini et al., 2013; Zweng et al.,  
2013; <https://www.nodc.noaa.gov/OC5/woa13/woa13data.html>, last access: 6 February 2017). Oxygen,  
phosphate, nitrate and silicate climatologies were taken from Broullón et al. (2019)  
(<http://dx.doi.org/10.20350/digitalCSIC/8644>, last access: 1 August 2019). These climatologies of nutrients  
were created using the objectively analyzed climatologies of temperature, salinity and oxygen (Garcia et  
230 al., 2014 filtered, see Broullón et al. (2019)) from WOA13 in CANYON-B (Bittig et al., 2018). As a year  
input is needed, we decided to center the TCO<sub>2</sub> climatology in 1995 based on the time distribution of the  
data used to create the WOA13 climatologies: World Ocean Database 2013 (Boyer et al., 2013).

The computed climatological values were compared with those from measured data to assess the  
uncertainty of the climatology, since WOA13 does not offer an uncertainty field with the objectively  
235 analyzed climatologies. Unfortunately, only two locations have enough measured data to calculate a pure  
climatological value of TCO<sub>2</sub> for each month: HOT ALOHA and BATS. The measured values were  
monthly averaged at several depth levels and the anthropogenic carbon as calculated by Lauvset et al.  
(2016) was added or subtracted to correct the data to the reference year of the climatology according to:

$$TCO_2^{year_2} = TCO_2^{year_1} - C_{ant_{2002}}[(1 + 0.0191)^{(year_1 - 2002)} - (1 + 0.0191)^{year_2 - 2002}] \quad \text{Eq. (1)}$$

240 where  $TCO_2^{year_2}$  is the TCO<sub>2</sub> corrected to  $year_2$ , which is the reference year of the climatology,  $TCO_2^{year_1}$   
is the TCO<sub>2</sub> measured in  $year_1$ ,  $C_{ant_{2002}}$  is the anthropogenic carbon for 2002 and 0.0191 is the annual  
increase rate derived from the scaling factor determined by Gruber et al. (2019) for the global ocean  
between 1994 and 2007.

We compared our climatology with previously published climatologies of TCO<sub>2</sub>. The surface monthly  
245 climatology created by Takahashi et al. (2014) was used to assess the spatiotemporal differences in the  
surface layer. The annual climatology of Lauvset et al. (2016) was used to evaluate the spatial differences  
in the deeper parts of the ocean. For the comparisons, the climatologies of Takahashi et al. (2014) and  
Lauvset et al. (2016) were adjusted to the year 1995 subtracting the anthropogenic carbon ( $C_{ant}$ ) of Lauvset  
et al. (2016) as in the Eq. (1).

250 Finally, a surface climatology of pCO<sub>2</sub> was computed from the TCO<sub>2</sub> climatology of the present study and  
the A<sub>T</sub> climatology of Broullón et al. (2019) to assess the potential of computing climatologies of other



variables of the seawater CO<sub>2</sub> system. For comparison, the updated monthly pCO<sub>2</sub> climatology from Landschützer et al. (2016) (Landschützer et al., 2017, last access: 30 July 2019) was used. The values between 1981 and 2015 were averaged to obtain the climatological year 1995. The variable selected from  
255 Landschützer et al. (2017) was that labeled as spco2\_raw.

### 3. Results

#### 3.1 Neural network analysis

Following the established criteria to obtain the optimal number of neurons, the configuration with 128  
neurons in the hidden layer was selected. From the ten networks trained with this number of neurons for  
260 each approach (NNGv2LDEO and NNGv2QCLDEO), the ones with the lowest influence of the position  
input variables were selected. These two networks present a similar RMSE in both training and test datasets,  
showing there is no overfitting. Because in Gv2QC both NNGv2LDEO and NNGv2QCLDEO produce the  
same global RMSE (6.1 μmol kg<sup>-1</sup>), it is likely that the Gv2 dataset contains high-quality measurements  
and the possible errors on the non-QC data of this dataset are clearly avoided by the network; otherwise  
265 NNGv2LDEO should have a higher RMSE in the test dataset than NNGv2QCLDEO because of an  
overfitting of the errors in the Gv2 dataset. The same holds for the LDEO dataset. The network properly  
fitted TCO<sub>2</sub> derived from LDEO, since it does not significantly increase the global RMSE relative to a  
network only trained with Gv2. Therefore, we decided to continue with NNGv2LDEO only since it has  
fitted more relationships between variables (e.g., Gv2 has more data points than Gv2QC in the  
270 Mediterranean Sea) providing a more robust fitting.

The global RMSE is quite low for the Gv2 dataset and for the LDEO dataset (Fig. 2). The measured and  
the computed data are highly correlated (Fig. 2) and the bias is negligible in both datasets. The higher  
RMSE in the LDEO dataset likely results from the higher variability of the TCO<sub>2</sub> in the surface layer and  
from uncertainties in its calculation from pCO<sub>2</sub>.

275 The RMSE by area and depth for NNGv2LDEO and CANYON-B is shown in Table 2. The highest errors  
for the two methods are in the 0-50 m layer for Gv2QC dataset and the LDEO dataset. These errors get  
smaller with increasing depth for all areas and the depth-weighted RMSE of the two methods is not  
significantly different. In the LDEO dataset, NNGv2LDEO produces a lower error than CANYON-B,  
except for two areas: East GIN (Greenland, Iceland and Norwegian) Seas and Main South Indian (Table  
280 2). Interestingly, CANYON-B is able to reproduce the TCO<sub>2</sub> data derived from the complete LDEO dataset  
with a lower error than the one it obtains for the complete Gv2QC dataset in the surface ocean (RMSE  
LDEO: 16.4 μmol kg<sup>-1</sup>; RMSE Gv2QC (0-5 m): 17.8 μmol kg<sup>-1</sup>), supporting the approach of computing  
reliable TCO<sub>2</sub> values from the pCO<sub>2</sub> of LDEO and the A<sub>T</sub> computed with NNGv2 (Broullón et al., 2019),  
since CANYON-B was not trained with the LDEO database. This result was also obtained for  
285 NNGv2LDEO but with a higher difference between the two errors (RMSE LDEO: 11.4 μmol kg<sup>-1</sup>; RMSE  
Gv2QC (0-5m): 17.1 μmol kg<sup>-1</sup>). Finally, the surface RMSE towards LDEO data of NNGv2LDEO is clearly  
lower than that of CANYON-B. This shows the value of including pCO<sub>2</sub> derived surface TCO<sub>2</sub> among the  
training data, through which there are more fitted relations in our new method.





In general, the highest differences between measured and estimated TCO<sub>2</sub> occurs in the high latitude surface  
290 oceans (Figs. 3 and 4). In Gv2, 40% of the samples with differences beyond  $\pm 3\text{RMSE}$  (3 times RMSE;  
threshold selected to show samples with large residuals) are in latitudes greater than 70° N. In the LDEO  
dataset, 39% of the data are from latitudes south of 70° S. These samples are 7.5% of the total north of 70°  
N in Gv2 and 42% of the total south of 70° S in LDEO. The samples with low salinities have the highest  
295 errors (Fig. 4). 41.5% of the samples in Gv2 and 43% in LDEO with differences beyond  $\pm 3\text{RMSE}$  have  
salinities below 33. Furthermore, in the LDEO dataset, the number of samples with residuals beyond  
 $\pm 3\text{RMSE}$  increases with increasing std of both pCO<sub>2</sub> and salinity obtained in the monthly averaging in each  
pixel in the LDEO subset (Fig. S2). This result shows the difficulty of modeling areas with a high sub-  
monthly variability in pCO<sub>2</sub> and salinity and supports the exclusion of the averaged LDEO data with a high  
std since it could cause the network to interpret the sub-monthly variability as monthly variability – note  
300 that the purpose of this study is to capture the monthly variability.

Like for modeling A<sub>T</sub> (Takahashi et al., 2014; Broullón et al., 2019), the Arctic Ocean is one of the regions  
with the highest RMSE of neural network estimated TCO<sub>2</sub>. The major Arctic rivers contribute with TCO<sub>2</sub>  
concentrations ranging between 400 and 3600  $\mu\text{mol kg}^{-1}$  (estimated by Tank et al., 2012), derived mainly  
from carbonated rocks in the watersheds. Other areas like the Okhotsk Sea also show a high RMSE (Table  
305 2 and Fig. 3), probably because of the high riverine input of TCO<sub>2</sub> (Watanabe et al., 2009). An input variable  
accounting for the contribution of the rivers to the TCO<sub>2</sub> pool would improve the neural network  
performance in areas like these, but is not available.

The errors of the pCO<sub>2</sub> computed in LDEO with TCO<sub>2</sub> from NNGv2LDEO and A<sub>T</sub> from NNGv2 (Broullón  
et al., 2019) are similar to the errors obtained by Landschützer et al. (2014) for the SOCAT database in  
310 some of the areas (10-16  $\mu\text{atm}$ , Table 2). This result shows the potential of computing pCO<sub>2</sub> values with  
neural networks trained for other variables of the seawater CO<sub>2</sub> system, at least in some ocean regions. The  
global error of the pCO<sub>2</sub> in the LDEO dataset is clearly higher than that obtained by Landschützer et al.  
(2014) for the SOCAT dataset (22 vs. 12  $\mu\text{atm}$ , respectively), although the critical areas are mainly the  
same (Fig. S3): equatorial Pacific upwelling system, Arctic and subarctic waters around the Alaska  
315 Peninsula, the Southern Ocean, the Gulf Stream and the North Atlantic Current. At this point, the following  
should be considered: 1) the pCO<sub>2</sub> computed in the present study derives from A<sub>T</sub> and TCO<sub>2</sub> and not from  
specific modeling for pCO<sub>2</sub>, and therefore, it contains errors associated to this computation (~6  $\mu\text{atm}$ ;  
Millero, 1995); 2) the present study includes the Arctic region where the highest errors occur (Table 2;  
Beaufort Sea and High Arctic areas); and 3) there is a longer temporal range in the present study (1973-  
320 2016). The analysis of Landschützer et al. (2014) in the LDEO dataset for data that differs from SOCAT  
shows a global error higher than the one obtained in the present study for all LDEO data between 1998 and  
2011 (25.9 vs. 21.3  $\mu\text{atm}$ , respectively). The error between 40° S-40° N is similar in the two studies  
(Landschützer et al. (2014): 16.5  $\mu\text{atm}$ ; NNGv2LDEO: 16.4  $\mu\text{atm}$ ). Although it is not the main objective  
of this work, these two last results show how NNGv2LDEO and NNGv2 (Broullón et al., 2019) have the  
325 potential to compute pCO<sub>2</sub> values between 40° S-40° N with similar errors as the method with the lower  
error in the pCO<sub>2</sub> modeling to obtain a climatology and with lower errors in high latitudes, for the LDEO  
dataset; even taking into account the inclusion of the critical area of the Arctic in the computation of the



error of the  $p\text{CO}_2$  from the present study (it is not included in Landschützer et al., 2014) and the higher amount of data from high latitudes in the present study (15479 vs. 3799).

### 330 3.2 Time-series validation

The good generalization of the network in the test dataset containing data from Gv2 and LDEO by the similar RMSE that the one reached in the training set is also evidenced through independent time-series data (Table 3). Except for KERFIX, where the number of data points is very low and Olsen et al. (2019) suggested an adjustment to the original data of  $-39 \mu\text{mol kg}^{-1}$ , NNGv2LDEO and CANYON-B computed  
335  $\text{TCO}_2$  at the time-series locations are characterized by low errors and biases (Table 3). NNGv2LDEO computes  $\text{TCO}_2$  with a lower RMSE and bias than CANYON-B for most of the time-series stations (Table 3). CANYON-B reaches a lower RMSE in HOT ALOHA SURFACE and ESTOC than NNGv2LDEO, but the bias is considerably higher in these time series for CANYON-B.

The seasonal variability is well captured by NNGv2LDEO showing its great potential to design a monthly  
340 climatology. In the surface layer, where the seasonal variability is the highest, the computed values are strongly correlated with the measured  $\text{TCO}_2$  in all the time series (Fig. 5). In addition, the high correlation holds for all depths. The location of the time series in different oceanographic regimes allows to complement the good  $\text{TCO}_2$  computation by NNGv2LDEO already shown in the previous independent sets in almost any region of the ocean.

345 Assessing the potential of neural networks to obtain values of other variables of the seawater  $\text{CO}_2$  system in the time series,  $p\text{CO}_2$  calculated with  $A_T$  from NNGv2 (Broullón et al., 2019) and  $\text{TCO}_2$  from NNGv2LDEO compared quite well with  $p\text{CO}_2$  as measured or calculated from  $A_T$  and  $\text{TCO}_2$  at the time-series stations (Table 4). Except for BATS, the  $p\text{CO}_2$  obtained in the present study has a lower error than that reported by Landschützer et al. (2014) (Table 4). In contrast, the bias in the present study is higher,  
350 except for ESTOC. Considering the error involved in the calculation of  $p\text{CO}_2$  from  $A_T$  and  $\text{TCO}_2$  ( $\sim 6 \mu\text{atm}$ ; Millero, 1995), our results demonstrate again the ability of NNGv2 and NNGv2LDEO to calculate other variables of the seawater  $\text{CO}_2$  system with a relatively low error.

Using NNGv2LDEO, it is also possible to reproduce the secular trends in  $\text{TCO}_2$ . Using seasonal detrending to enhance the multi-annual changes, similar trends in the longer time series are found for the measured  
355 and the neural network computed  $\text{TCO}_2$  (Table 5). The same holds for  $p\text{CO}_2$  (Table 5), although at IRMINGER site the trend obtained from the neural network generated data is significantly lower than that from measured data. The neural networks seem to capture the anthropogenic influence in the seawater  $\text{CO}_2$  system and therefore the ocean acidification process (Fig. 6). Furthermore, NNGv2LDEO increases the amount of  $\text{TCO}_2$  data where the various inputs were measured but not  $\text{TCO}_2$  itself. This allows for  
360 evaluation of high frequency changes (Fig. 6) and for calculation of interannual trends with a low error (as temporal sampling biases are reduced).

### 3.3 Climatology



Using NNGv2LDEO we have demonstrated its ability to compute TCO<sub>2</sub> values with low errors and, especially, to capture the monthly variability of this variable. In addition, the climatologies of the input variables used to create the climatology of TCO<sub>2</sub> have been satisfactorily evaluated previously for the construction of an A<sub>T</sub> climatology (Broullón et al., 2019). Considering these results, a monthly climatology of TCO<sub>2</sub> is obtained by passing the input climatologies through NNGv2LDEO.

The spatial distribution of the surface annual mean climatology of TCO<sub>2</sub> (Fig. 7a) is similar to two recent climatologies: those of Takahashi et al. (2014) and Lauvset et al. (2016). The largest surface TCO<sub>2</sub> concentrations occur in the Southern Ocean, Subpolar North Atlantic, Nordic Seas and Mediterranean Sea (note that the latter is not included in these previous climatologies). In general, surface TCO<sub>2</sub> decreases from high to low latitudes. The Indian and the Pacific oceans are characterized by lower concentrations of TCO<sub>2</sub> at higher latitudes than the Atlantic, the latter being the ocean with the highest surface TCO<sub>2</sub> by area. TCO<sub>2</sub> increases with depth in all oceans, in particular in the upwelling regions, where this increase is expanded eastwards with depth (Fig. 7b and video in <http://dx.doi.org/10.20350/digitalCSIC/10551>). Depending on the area, the values reach a maximum at certain intermediate depths and below it the concentration gradually decreases or remains almost constant (Fig. S4).

The largest seasonal variability occurs at the surface in high latitudes, in the Pacific upwelling region, the equatorial African coasts and in the area under influence of the Amazon River (Fig. 8a). At depth, the seasonal variability decreases, except for the Pacific upwelling region where it increases and moves progressively northward between 30 and 150 m (Fig 8b). This increase is correlated with the high seasonal variability of the climatologies of nutrients, oxygen and temperature at these depths. Czeschel et al. (2012) also showed an increase in the subsurface variability of oxygen from measured profiles. Similar increases also occur in the Indian Ocean north of 20° S between 50 and 100 m and in the equatorial Atlantic Ocean in the same depth range. At 1500 m level, the seasonal variability is below 10 μmol kg<sup>-1</sup> in most of the ocean (Fig. 8c). This last result shows that an annual climatology below 1500 m is sufficient.

Although the surface patterns of the annual mean of the TCO<sub>2</sub> climatology are very similar to those of the other recent climatologies (Takahashi et al., 2014; Lauvset et al., 2016), differences do occur. The annual mean of the climatology of the present study is closest to that of Takahashi et al. (2014) (Table 6). The largest differences between these two climatologies are located in the Arctic, North Pacific, Peru upwelling area, western South Pacific and the area of influence of the Antarctic Circumpolar Current (Fig. S5a). The Atlantic and the Indian oceans do not show significant differences. Our climatology shows more deviations to that of Lauvset et al. (2016), compared in the grid of Takahashi et al. (2014) (Table 6). The highest differences are found in the North Pacific, around Antarctica, Nordic Seas, South and North Atlantic and in several less localized areas around the oceans (Fig. S5b). When the climatology of Takahashi et al. (2014) is compared to that of Lauvset et al. (2016), the differences are even higher (Table 6) and the critical areas are the same of those of the previous comparison. Although it is clear that discrepancies between the three climatologies derive from the different methods used, the higher similarity between ours and the one of Takahashi et al. (2014) is probably due to the influence of the same source used to create them, the World Ocean Atlas.



The comparison of our climatology with that of Lauvset et al. (2016) at the 33 depth levels of Lauvset et al. (2016) shows a reduction of the RMSE with depth. Between 0 and 1000 m, the RMSE is reduced from ~32 to 7  $\mu\text{mol kg}^{-1}$  (Table S1) (note the higher RMSE at surface compared to one obtained for the grid of Takahashi et al. (2014) because of the inclusion of areas which are not included in the latter's grid, and the difficulty of modeling  $\text{TCO}_2$  in some areas, like the Arctic and the Mediterranean Sea). This reduction with  
405 depth is probably due to the reduction of the variability in most of the ocean below the surface. The surface values in Lauvset et al. are likely characteristic from months in which most of the sampling was carried out. Because of the lower variability of  $\text{TCO}_2$  in depth, the values are closer to the annual mean and therefore the two compared climatologies are more similar at depth than in surface depth levels. Below  
410 1000 m, the differences between the two climatologies are not significant, with a RMSE around 5  $\mu\text{mol kg}^{-1}$  and a bias around 0.5  $\mu\text{mol kg}^{-1}$  at each depth level (Table S1).

Our monthly climatology shows a high correspondence with that of Takahashi et al. (2014), although the RMSE values show that there are also large differences in certain areas (Table 7). These areas are mainly the same of those in the comparison of the annual mean climatologies, but some other small regions with  
415 high differences appear for each month all through the ocean (Fig. S6).

Unfortunately, the uncertainty of the  $\text{TCO}_2$  climatology cannot be assessed globally and robustly. As Broullón et al. (2019) stated, the unavailability of an uncertainty field associated to the WOA13 objectively analyzed climatologies does not allow to perform a proper global uncertainty assessment. Therefore, the analysis is relegated to the areas where repeated sampling of  $\text{TCO}_2$  has been carried out monthly over a  
420 long period, that is, the HOT ALOHA and BATS time-series stations. The climatology of  $\text{TCO}_2$  from NNGv2LDEO is consistent with the monthly climatological values at these two places (Fig. 9). In general, the profiles from the  $\text{TCO}_2$  climatology are within the variability range (shadow area in Fig. 9) of the monthly averaged measured data for each depth level. For HOT ALOHA, the RMSE of the profiles of the  $\text{TCO}_2$  climatology oscillates between 3.6 and 9.2  $\mu\text{mol kg}^{-1}$  with a mean value of 6.3  $\mu\text{mol kg}^{-1}$  (bias range:  
425 -3.8 to 1.2  $\mu\text{mol kg}^{-1}$ ; mean bias: -1.4  $\mu\text{mol kg}^{-1}$ ). At BATS, the RMSE is lower than for HOT ALOHA: 1.1 to 8.5  $\mu\text{mol kg}^{-1}$  with a mean value of 4.4  $\mu\text{mol kg}^{-1}$  (bias range: -1.4 to 7.6  $\mu\text{mol kg}^{-1}$ ; mean bias: -2.5  $\mu\text{mol kg}^{-1}$ ). Furthermore, the seasonal variability of the  $\text{TCO}_2$  climatology is quite similar to that of the measured data at BATS and HOT ALOHA (Fig. S7). Although in other time series there are not enough measured data to obtain climatological values, these pseudo climatological values also correlate very well  
430 with the  $\text{TCO}_2$  climatology (data not shown). These results suggest that the climatology is robust in different oceanographic regimes and adequately captures the seasonal cycle of  $\text{TCO}_2$ .

It has been demonstrated in this study that  $\text{pCO}_2$  and possibly other variables of the seawater  $\text{CO}_2$  system can be computed from  $A_T$  and  $\text{TCO}_2$  derived from neural networks with a relatively low error in different datasets (LDEO and time series). The  $\text{pCO}_2$  climatology computed from the  $\text{TCO}_2$  climatology of the  
435 present study and the  $A_T$  climatology of Broullón et al. (2019) is very similar to that of Landschützer et al. (2017). The differences between the annual mean climatology of the two studies are below 15  $\mu\text{atm}$  in most of the ocean (RMSE: 8.3  $\mu\text{atm}$ ; bias: 2.9  $\mu\text{atm}$ ;  $r^2$ : 0.82). The differences above this threshold are mainly located in the Pacific equatorial upwelling system, the east part of the South Pacific Gyre, Nordic Seas,



440 Labrador Sea, Atlantic section of the Southern Ocean, Bay of Bengal and the waters surrounding the east margin of Asia (Fig. S8). In most of these areas, both methods have the greatest errors (Figs. 2 and 4 in Landschützer et al., 2014 and Fig. S3 of the present study).

445 On a monthly basis, the RMSE between the two pCO<sub>2</sub> climatologies is between 13.6 and 15.6 μatm and the correlation is lower than for the annual mean comparison ( $r^2$ : 0.55-0.72 vs 0.72). The areas with the higher differences are the same as in the annual comparison but other small regions appear along the ocean month by month (Fig. S9). Furthermore, the seasonal variability in the two climatologies matches in a great extension of the ocean, although there are areas with notable differences (Fig. S10). In general, the pCO<sub>2</sub> climatology is quite similar to that of Landschützer et al. (2017) and this result contributes to show that both the TCO<sub>2</sub> climatology of the present study and the A<sub>T</sub> climatology of Broullón et al. (2019) are mostly robust and suggest that climatologies of other seawater CO<sub>2</sub> system variables can be confidently computed.

#### 450 **4 Data availability**

The climatologies of TCO<sub>2</sub> and pCO<sub>2</sub> and NNGv2LDEO designed in this study are available at the data repository of the Spanish National Research Council (CSIC; <http://dx.doi.org/10.20350/digitalCSIC/10551>, Broullón et al., 2020).

#### **5 Conclusions**

455 We presented a tool for computing TCO<sub>2</sub> in the global ocean. Compared to previous methods, the uncertainties in such computations have been reduced. Including two updated datasets containing thousands of measurements of inorganic carbon variables across the ocean in the training of the neural network, we were able to capture a wide range of variability of TCO<sub>2</sub>. The low errors obtained in independent subsets as in time-series stations, are further evidence of the potential of the network in computing TCO<sub>2</sub>.

460 Our global monthly climatology created with a neural network is the first that covers the oceans from the surface to the abyss at such temporal resolution. In addition to the accuracy of the network, the low uncertainty of the climatology in different regions and its usefulness in creating climatologies of other seawater CO<sub>2</sub> chemistry variables (i.e. pCO<sub>2</sub>) show its robustness. Therefore, we present the global climatology of TCO<sub>2</sub> to the scientific community to complement the recently designed climatology of A<sub>T</sub> by Broullón et al. (2019) for its use in the initialization and evaluation of models or any other analysis related to the carbon cycle.

#### **6 Author contributions**

DB, FFP and AV designed the study. The manuscript was written by DB and revised and discussed by all the authors. The dataset of the climatology and the neural network were created by DB.

#### 470 **7 Competing interests**

The authors declare that they have no conflict of interest.



## 8 Acknowledgements

The authors want to thank the comments of Siv K. Lauvset to improve the study. The paper is dedicated to Taro Takahashi, who immensely contributed to the ocean's role in the carbon cycle and passed away in  
475 December 2019.

## 9 Financial support

This research was supported by Ministerio de Educación, Cultura y Deporte (FPU grant FPU15/06026), Ministerio de Economía y Competitividad through the ARIOS (CTM2016-76146-C3-1-R) project co-funded by the Fondo Europeo de Desarrollo Regional 2014-2020 (FEDER) and EU Horizon 2020 through  
480 the AtlantOS project (grant agreement 633211). Are Olsen was supported by the Norwegian Research Council through ICOS (245927).

## 10 References

- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L. and Gehlen, M.: PISCES-v2: An ocean biogeochemical model for carbon and ecosystem studies, *Geosci. Model Dev.*, 8(8), 2465–2513, doi:10.5194/gmd-8-2465-2015, 2015.  
485
- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S. I., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W. J., Castle, R. D., Chen, L., Chierici, M.,  
490 Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T.,  
495 Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., Van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J. and Xu, S.: A multi-decade record of high-quality fCO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth Syst. Sci. Data*, 8(2), 383–413, doi:10.5194/essd-8-383-2016, 2016.
- 500 Bates, N. R., Best, M. H. P., Neely, K., Garley, R., Dickson, A. G. and Johnson, R. J.: Detecting anthropogenic carbon dioxide uptake and ocean acidification in the North Atlantic Ocean, *Biogeosciences*, 9(7), 2509–2522, doi:10.5194/bg-9-2509-2012, 2012.
- Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S. and Regnier, P. A. G.: The changing carbon cycle of the coastal ocean, *Nature*, 504(7478), 61–70, doi:10.1038/nature12857, 2013.



- 505 Beale, M. H., Hagan, T. M., and Demuth, H. B.: Deep Learning Toolbox™, User's Guide, Release 2018a, The MathWorks, Inc., Natick, Massachusetts, United States, available at: [https://es.mathworks.com/help/pdf\\_doc/deeplearning/nnet Ug.pdf](https://es.mathworks.com/help/pdf_doc/deeplearning/nnet Ug.pdf) (last access: 20 August 2018), 2018.
- Bittig, H. C., Steinhoff, T., Claustre, H., Fiedler, B., Williams, N. L., Sauzède, R., Körtzinger, A. and Gattuso, J.-P.: An Alternative to Static Climatologies: Robust Estimation of Open Ocean CO<sub>2</sub> Variables and Nutrient Concentrations From T, S, and O<sub>2</sub> Data Using Bayesian Neural Networks, *Front. Mar. Sci.*, 5,328, doi:10.3389/fmars.2018.00328, 2018.
- 510 Boyer, T.P., J. I. Antonov, O. K. Baranova, C. Coleman, H. E. Garcia, A. Grodsky, D. R. Johnson, R. A. Locarnini, A. V. Mishonov, T.D. O'Brien, C.R. Paver, J.R. Reagan, D. Seidov, I. V. Smolyar, and M. M. Zweng, 2013: World Ocean Database 2013, NOAA Atlas NESDIS 72, S. Levitus, Ed., A. Mishonov, Technical Ed.; Silver Spring, MD, 209 pp., <http://doi.org/10.7289/V5NZ85MT>, 2013.
- 515 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., González-Dávila, M., Jeansson, E., Kozyr, A. and van Heuven, S. M. A. C.: A global monthly climatology of total alkalinity: a neural network approach, *Earth Syst. Sci. Data*, 11(3), 1109–1127, doi:10.5194/essd-11-1109-2019, 2019.
- 520 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, M., Tanhua, T., Santana-Casiano, J.M. and Kozyr, A.: A global monthly climatology of oceanic total dissolved inorganic carbon: a neural network approach (Discussions version) [Dataset], <http://dx.doi.org/10.20350/digitalCSIC/10551>, 2020.
- 525 Butenschön, M., Clark, J., Aldridge, J. N., Icarus Allen, J., Artioli, Y., Blackford, J., Bruggeman, J., Cazenave, P., Ciavatta, S., Kay, S., Lessin, G., Van Leeuwen, S., Van Der Molen, J., De Mora, L., Polimene, L., Saille, S., Stephens, N. and Torres, R.: ERSEM 15.06: A generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels, *Geosci. Model Dev.*, 9(4), 1293–1339, doi:10.5194/gmd-9-1293-2016, 2016.
- 530 Carter, B. R., Feely, R. A., Williams, N. L., Dickson, A. G., Fong, M. B. and Takeshita, Y.: Updated methods for global locally interpolated estimation of alkalinity, pH, and nitrate, *Limnol. Oceanogr. Methods*, 16(2), 119–131, doi:10.1002/lom3.10232, 2018.
- 535 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Quéré, C. Le, Myneni, R., Piao, S. and Thornton, P.: Carbon and Other Biogeochemical Cycles, in *Climate Change 2013 - The Physical Science Basis*, edited by Intergovernmental Panel on Climate Change, pp. 465–570, Cambridge University Press, Cambridge., 2013.
- Czeschel, R., Stramma, L. and Johnson, G. C.: Oxygen decreases and variability in the eastern equatorial Pacific, *J. Geophys. Res. Ocean.*, 117(C11), C11019, doi:10.1029/2012JC008043, 2012.



- Dickson, A. G.: Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K, *Deep Sea Res. Part A.*, 37(5), 755–766, doi:10.1016/0198-0149(90)90004-F, 1990.
- 540 Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep Sea Res. Part A.*, 34(10), 1733–1743, doi:10.1016/0198-0149(87)90021-5, 1987.
- Doi, T., Osafune, S., Sugiura, N., Kouketsu, S., Murata, A., Masuda, S. and Toyoda, T.: Multidecadal change in the dissolved inorganic carbon in a long-term ocean state estimation, *J. Adv. Model. Earth Syst.*, 545 7(4), 1885–1900, doi:10.1002/2015MS000462, 2015.
- Doney, S. C., Fabry, V. J., Feely, R. A. and Kleypas, J. A.: Ocean Acidification: The Other CO<sub>2</sub> Problem, *Ann. Rev. Mar. Sci.*, 1(1), 169–192, doi:10.1146/annurev.marine.010908.163834, 2009.
- Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J. and Karl, D. M.: Physical and biogeochemical modulation of ocean acidification in the central North Pacific, *Proc. Natl. Acad. Sci. U. S. A.*, 106(30), 550 12235–12240, doi:10.1073/pnas.0906044106, 2009.
- Fabry, V. J., Seibel, B. A., Feely, R. A., Fabry, J. C. O. and Fabry, V. J.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES J. Mar. Sci.*, 65, 414–432, doi:10.1093/icesjms/fsn048, 2008.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., and Johnson, D. R.: *World Ocean Atlas 2013, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation*, edited by: Levitus, S. and Mishonov, A., Technical Ed., NOAA Atlas NESDIS 75, 555 27 pp., 2014.
- Gislefoss, J. S., Nydal, R., Slagstad, D., Sonninen, E. and Holmén, K.: Carbon time series in the Norwegian sea, *Deep. Res. Part I Oceanogr. Res. Pap.*, 45(2–3), 433–460, doi:10.1016/S0967-0637(97)00093-9, 1998.
- González-Dávila, M., Santana-Casiano, J. M., Rueda, M. J. and Llinás, O.: The water column distribution of carbonate system variables at the ESTOC site from 1995 to 2004, *Biogeosciences*, 7(10), 3067–3081, 560 doi:10.5194/bg-7-3067-2010, 2010.
- Goris, N., Tjiputra, J. F., Olsen, A., Schwinger, J., Lauvset, S. K. and Jeansson, E.: Constraining projection-based estimates of the future North Atlantic carbon uptake, *J. Clim.*, doi:10.1175/JCLI-D-17-0564.1, 2018.
- Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., 565 Kozyr, A., Lauvset, S. K., Monaco, C. Lo, Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T. and Wanninkhof, R.: The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007, *Science*, 363(6432), 1193–1199, doi:10.1126/science.aau5153, 2019.
- Hendriks, I. E., Duarte, C. M. and Álvarez, M.: Vulnerability of marine biodiversity to ocean acidification: A meta-analysis, *Estuar. Coast. Shelf Sci.*, 86(2), 157–164, doi:10.1016/J.ECSS.2009.11.022, 2010.





- 570 Hoegh-Guldberg, O. and Bruno, J. F.: The Impact of Climate Change on the World's Marine Ecosystems, *Science*, 328(5985), 1523–1528, doi:10.1126/science.1189930, 2010.
- Jeandel, C., Ruiz-Pino, D., Gjata, E., Poisson, A., Brunet, C., Charriaud, E., Dehairs, F., Delille, D., Fiala, M., Fravallo, C., Carlos Miquel, J., Park, Y. H., Pondaven, P., Quéguiner, B., Razouls, S., Shauer, B. and Tréguer, P.: KERFIX, a time-series station in the Southern Ocean: A presentation, in *Journal of Marine Systems*, 17, 555–569., 1998.
- 575 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C. and Peng, T. H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cycles*, 18(4), 1–23, doi:10.1029/2004GB002247, 2004.
- Key, R. M., Olsen, A., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishi, M., Perez, F. F. and Suzuki, T.: Global Ocean Data Analysis Project, Version 2 (GLODAPv2), ORNL/CDIAC-162, NDP-093, doi:10.3334/CDIAC/OTG.NDP093\_GLODAPv2, 2015.
- 580 Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F. and Sabine, C. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences*, 10(4), 2169–2191, doi:10.5194/bg-10-2169-2013, 2013.
- 585 Kleypas, J. and Langdon, C.: Overview of CO<sub>2</sub>-induced changes in seawater chemistry, *Proc. 9th Int. Coral Reef Symp. Bali.*, 2, 2000.
- Körtzinger, A., Hedges, J. I. and Quay, P. D.: Redfield ratios revisited: Removing the biasing effect of anthropogenic CO<sub>2</sub>, *Limnol. Oceanogr.*, doi:10.4319/lo.2001.46.4.0964, 2001.
- 590 Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M. and Gattuso, J.-P.: Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming, *Glob. Chang. Biol.*, 19(6), 1884–1896, doi:10.1111/gcb.12179, 2013.
- Landschützer, P., Gruber, N., Bakker, D. C. E. and Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochem. Cycles*, 28(9), 927–949, doi:10.1002/2014GB004853, 2014.
- 595 Landschützer, P., Gruber, N. and Bakker, D. C. E.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochem. Cycles*, 30(10), 1396–1417, doi:10.1002/2015GB005359, 2016.
- Landschützer, P., Gruber, N. and Bakker, D.C.E.: An updated observation-based global monthly gridded sea surface pCO<sub>2</sub> and air-sea CO<sub>2</sub> flux product from 1982 through 2015 and its monthly climatology (NCEI Accession 0160558). Version 2.2. NOAA National Centers for Environmental Information. Dataset. [2017-07-11], (last access: 15 July 2019), 2017.
- 600 Lauvset, S. K., Key, R. M., Olsen, A., Van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., Suzuki, T. and Watelet,



- S.: A new global interior ocean mapped climatology: The  $1^\circ \times 1^\circ$  GLODAP version 2, *Earth Syst. Sci. Data*, 8(2), 325–340, doi:10.5194/essd-8-325-2016, 2016.
- 605 Lee, K., Wanninkhof, R., Feely, R. A., Millero, F. J. and Peng, T.-H.: Global relationships of total inorganic carbon with temperature and nitrate in surface seawater, *Global Biogeochem. Cycles*, 14(3), 979–994, doi:10.1029/1998GB001087, 2000.
- Le Quéré, C., Buitenhuis, E. T., Moriarty, R., Alvain, S., Aumont, O., Bopp, L., Chollet, S., Enright, C., Franklin, D. J., Geider, R. J., Harrison, S. P., Hirst, A. G., Larsen, S., Legendre, L., Platt, T., Prentice, I. C., Rivkin, R. B., Sailley, S., Sathyendranath, S., Stephens, N., Vogt, M. and Vallina, S. M.: Role of zooplankton dynamics for Southern Ocean phytoplankton biomass and global biogeochemical cycles, *Biogeosciences*, 13(14), 4111–4133, doi:10.5194/bg-13-4111-2016, 2016.
- 615 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S. and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, 10(4), 2141–2194, doi:10.5194/essd-10-2141-2018, 2018.
- Levenberg, K.: A Method for the solution of certain non-linear problems in least squares., *Q. Appl. Math.*, 625 II(2), 164–168, 1944.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: *World Ocean Atlas 2013, Volume 1: Temperature*, edited by: Levitus, S. and Mishonov, A., NOAA Atlas NESDIS 73, 40 pp., 2013.
- Marquardt, D.: An Algorithm for Least-Squares Estimation of Nonlinear Parameters, *J. Soc. Ind. Appl. Math.*, 630 11(2), 1963.
- Mehrbach, C., Culberson, C. H., Hawley, J. E. and Pytkowicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18(6), 897–907, doi:10.4319/lo.1973.18.6.0897, 1973.
- Millero, F. J.: Thermodynamics of the carbon dioxide system in the oceans, *Geochim. Cosmochim. Acta*, 635 59(4), 661–677, doi:10.1016/0016-7037(94)00354-O, 1995.



- Olafsson, J., Olafsdottir, S. R., Benoit-Cattin, A. and Takahashi, T.: The Irminger Sea and the Iceland Sea time series measurements of sea water carbon and nutrient chemistry 1983–2008, *Earth Syst. Sci. Data*, 2(1), 99–104, doi:10.5194/essd-2-99-2010, 2010.
- 640 Olden, J. D. and Jackson, D. A.: Illuminating the “black box”: a randomization approach for understanding variable contributions in artificial neural networks, *Ecol. Modell.*, 154, 135–150, doi:10.1016/S0304-3800(02)00064-9, 2002.
- Olsen, A., Lange, N., Key, R. M., Tanhua, T., Álvarez, M., Becker, S., Bittig, H. C., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jones, S. D., Jutterström, S., Karlsen, M. K., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Murata, A., Pérez, F. F., Pfeil, B., Schirnick, C., Steinfeldt, R., Suzuki, T., Telszewski, M., Tilbrook, B., Velo, A. and Wanninkhof, R.: 645 GLODAPv2.2019 - an update of GLODAPv2, *Earth Syst. Sci. Data*, 11, 1437–1461, <https://doi.org/10.5194/essd-11-1437-2019>, 2019.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., 650 Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y. and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437(7059), 681–686, doi:10.1038/nature04095, 2005.
- Sallée, J.-B., Matear, R. J., Rintoul, S. R. and Lenton, A.: Localized subduction of anthropogenic carbon 655 dioxide in the Southern Hemisphere oceans, *Nat. Geosci.*, 5(8), 579–584, doi:10.1038/ngeo1523, 2012.
- Sarmiento, J. L. and Gruber, N.: Sinks for Anthropogenic Carbon, *Phys. Today*, 55(8), 30–36, doi:10.1063/1.1510279, 2002.
- Sauzède, R., Claustre, H., Pasqueron de Fommervault, O., Bittig, H. C., Gattuso, J.-P., Legendre, L. and Johnson, K. S.: Estimates of water-column nutrients concentration and carbonate system parameters in the 660 global ocean: A novel approach based on neural networks, *Front. Mar. Sci.*, 4, 128, doi:10.3389/fmars.2017.00128, 2017.
- Schlitzer, R., Ocean Data View, available at: <http://odv.awi.de> (last access: 21 May 2018), 2016.
- Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G. and Ho, C.: Climatological distributions of pH, pCO<sub>2</sub>, total CO<sub>2</sub>, alkalinity, and CaCO<sub>3</sub> saturation in the global surface ocean, and temporal changes 665 at selected locations, *Mar. Chem.*, 164, 95–125, doi:10.1016/j.marchem.2014.06.004, 2014.
- Takahashi, T., S.C. Sutherland and A. Kozyr.: Global Ocean Surface Water Partial Pressure of CO<sub>2</sub> Database: Measurements Performed During 1957-2016 (Version 2016). ORNL/CDIAC-161, NDP-088(V2015). Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, doi: 10.3334/CDIAC/OTG.NDP088(V2016), 2017



670 Tanhua, T., Bates, N. R. and Körtzinger, A.: The Marine Carbon Cycle and Ocean Carbon Inventories, in International Geophysics, vol. 103, edited by J. Siedler, G., Griffies, S., Gould, J. and Church, pp. 787–815, Academic Press., 2013.

Tank, S. E., Raymond, P. A., Striegl, R. G., McClelland, J. W., Holmes, R. M., Fiske, G. J. and Peterson, B. J.: A land-to-ocean perspective on the magnitude, source and implication of DIC flux from major Arctic rivers to the Arctic Ocean, *Global Biogeochem. Cycles*, 26(4), doi:10.1029/2011GB004192, 2012.

van Heuven, S., D. Pierrot, J.W.B. Rae, E. Lewis, and D.W.R. Wallace: MATLAB Program Developed for CO<sub>2</sub> System Calculations. ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. [https://doi.org/10.3334/CDIAC/otg.CO2SYS\\_MATLAB\\_v1.1](https://doi.org/10.3334/CDIAC/otg.CO2SYS_MATLAB_v1.1), 2011.

680 Vázquez-Rodríguez, M., Touratier, F., Lo Monaco, C., Waugh, D. W., Padin, X. A., Bellerby, R. G. J., Goyet, C., Metzl, N., Ríos, A. F. and Pérez, F. F.: Anthropogenic carbon distributions in the Atlantic Ocean: data-based estimates from the Arctic to the Antarctic, *Biogeosciences*, 6(3), 439–451, doi:10.5194/bg-6-439-2009, 2009.

685 Wakita, M., Watanabe, S., Murata, A., Tsurushima, N. and Honda, M.: Decadal change of dissolved inorganic carbon in the subarctic western North Pacific Ocean, *Tellus B Chem. Phys. Meteorol.*, 62(5), 608–620, doi:10.1111/j.1600-0889.2010.00476.x, 2010.

Weiss, R. F., Jahnke, R. A. and Keeling, C. D.: Seasonal effects of temperature and salinity on the partial pressure of CO<sub>2</sub> in seawater, *Nature*, 300(5892), 511–513, doi:10.1038/300511a0, 1982.

690 Wu, Y., Hain, M. P., Humphreys, M. P., Hartman, S. and Tyrrell, T.: What drives the latitudinal gradient in open-ocean surface dissolved inorganic carbon concentration?, *Biogeosciences*, 16(13), 2661–2681, doi:10.5194/bg-16-2661-2019, 2019.

Yool, A., Popova, E. E. and Anderson, T. R.: MEDUSA-2.0: An intermediate complexity biogeochemical model of the marine carbon cycle for climate change and ocean acidification studies, *Geosci. Model Dev.*, 6(5), 1767–1811, doi:10.5194/gmd-6-1767-2013, 2013.

695 Zeebe and Wolf-Gladrow: CO<sub>2</sub> in seawater: Equilibrium, Kinetics, Isotopes, in Elsevier oceanography series, edited by D. Halpem, Elsevier Oceanography Series., 2001.

Zweng, M.M, J.R. Reagan, J.I. Antonov, R.A. Locarnini, A.V. Mishonov, T.P. Boyer, H.E. Garcia, O.K. Baranova, D.R. Johnson, D. Seidov, M.M. Biddle.: World Ocean Atlas 2013, Volume 2: Salinity. S. Levitus, Ed., A. Mishonov Technical Ed.; NOAA Atlas NESDIS 74, 39 pp., 2013.

700



**Table 1. Differences between the methods used in the present study and in CANYON-B (Bittig et al., 2018).**

	Bittig et al. (2018)	This study
Training technique	Bayesian regularization	Levenberg-Marquardt
Predictor variables	Temperature, salinity, oxygen, position and time	Temperature, salinity, oxygen, phosphate, nitrate, silicate, position and time
Datasets	GLODAPv2 (Olsen et al., 2016)	GLODAPv2.2019 (Olsen et al., 2019) LDEOv2016 (Takahashi et al., 2017)

**Table 2. Error analysis by area and depth for TCO<sub>2</sub> and pCO<sub>2</sub> computed with CANYON-B and NNGv2LDEO in Gv2QC and LDEO datasets. Units are micromoles per kilogram ( $\mu\text{mol kg}^{-1}$ ) for TCO<sub>2</sub> and microatmospheres ( $\mu\text{atm}$ ) for pCO<sub>2</sub>.**

Area	TCO <sub>2</sub>										pCO <sub>2</sub>			
	0-50m				50-200m		200-500m		500-1000m		1000-bottom		0m	
	Gv2QC		LDEO (0m)		Gv2QC		Gv2QC		Gv2QC		Gv2QC		LDEO	
	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B	NNGv2LDEO	CANYON-B
West GIN Seas	14.2	15.3	16.8	21.2	6.4	6.7	5.5	6.8	4.1	5.0	4.0	4.1	28.6	34.9
East GIN Seas	10.2	11.4	11.1	9.2	5.9	6.2	4.6	5.2	3.5	4.0	3.9	3.7	17.3	15.0
High Arctic	20.2	24.1	13.3	32.4	11.4	12.3	6.5	6.6	6.8	7.6	6.1	6.2	43.0	79.1
Beaufort Sea	54	53.1	29.7	42.0	14.5	13.3	8.1	9.2	7.5	8.4	6.5	7.2	58.7	135.6
Labrador Sea	10.5	11.6	5.3	7.5	5.4	6.6	4.1	4.3	3.5	3.7	2.7	3.1	10.7	13.6
Subarctic Atlantic	9	11.1	11.7	14.6	4.5	5.4	4.3	4.8	4.2	4.6	4.0	4.3	21.9	26.7
North Atlantic Drift	13.3	14.5	11.1	12.9	9.9	10.4	7.8	7.8	4.3	4.5	3.5	3.6	21.0	24.7
Central Atlantic	15.8	14.9	7.9	9.6	6.6	6.5	5.2	5.1	4.6	4.5	4.3	4.4	13.5	17.4
South Atlantic Transition Zone	7.2	7.8	7.7	13.8	5.4	5.7	5.7	4.7	5.0	4.8	4.3	4.2	14.6	25.0



Antarctic (Atlantic)	8.6	10	11.8	19.2	4.6	5.4	3.5	3.8	3.1	3.1	3.1	3.1	25.6	41.5
Kuroshio-Alaska Gyre	8.5	12.2	10.9	12.3	6.4	8.1	5.0	5.2	4.5	4.3	3.7	3.9	20.7	23.7
North Central Pacific	9.6	15	26.3	34.5	6.8	8.3	4.2	4.7	4.0	4.1	3.4	3.8	46.7	56.6
Okhotsk Sea	23.1	16.4	-	-	11.3	6.8	6.3	5.1	5.2	3.4	4.1	3.5	-	-
Central Tropical North Pacific	7.9	8.8	8.0	9.7	7.2	7.2	4.9	5.0	4.3	4.5	3.6	3.8	14.2	17.2
Tropical East North Pacific	10.9	13.8	11.1	14.5	5.9	8.5	2.6	3.4	2.1	2.1	2.4	2.1	20.8	28.2
Central South Pacific	10.2	9.5	12.5	17.4	6.5	3.9	4.0	5.8	3.8	3.2	4.2	4.3	25.8	38.8
East Central South Pacific	10.3	10.9	10.1	12.9	8.9	9.4	4.4	4.5	3.8	3.8	3.3	3.5	18.6	24.3
Subpolar South Pacific	10.6	15.2	10.7	15.4	6.9	7.5	4.1	2.8	3.8	3.5	3.3	3.0	24.1	34.2
Antarctic (Pacific)	5.8	7.7	6.9	7.9	5.0	5.4	2.9	2.8	4.7	4.8	4.1	4.4	13.7	16.0
Main North Indian	8.3	7	19.5	29.3	3.6	4.1	2.7	3.3	2.6	2.9	2.6	2.1	34.1	53.8
Red Sea	10.5	12.9	10.8	13.0	8.1	7.8	3.2	3.3	2.4	2.6	3.1	3.7	19.8	23.5
Bengal Basin	12	16.8	18.3	20.9	9.4	8.7	7.6	7.9	7.4	5.7	3.3	7.2	28.0	32.3
Main South Indian	9.8	7.4	6.0	3.7	7.4	6.4	2.0	2.1	1.9	2.1	2.0	2.2	10.7	6.7
South Indian Transition	9.1	10	8.1	10.4	7.1	6.9	3.8	3.8	4.2	4.5	3.4	3.8	14.0	17.7
Antarctic (Indian)	5.3	5.3	8.0	9.4	4.1	4.4	4.0	3.7	3.9	3.4	3.4	3.7	16.0	18.8
Circumpolar Southern Ocean	6	6.8	9.7	11.8	4.1	4.7	3.0	3.3	2.5	2.6	2.7	2.6	23.2	28.7
Weighted RMSE	7.6	8.2	15.9	24.5	4.4	4.9	3.1	3.4	2.9	2.9	2.9	2.7	21.1	29.6

**Table 3. RMSE and bias between measured and computed TCO<sub>2</sub> concentrations in several time series. The comparison was done using only water samples where all the input variables for NNGv2LDEO and the TCO<sub>2</sub> were measured in the same water sample.**

Time series	Location	Time period	n	NNGv2LDEO		CANYON-B	
				RMSE (μmol kg <sup>-1</sup> )	bias (μmol kg <sup>-1</sup> )	RMSE (μmol kg <sup>-1</sup> )	bias (μmol kg <sup>-1</sup> )
BATS	31.7°N, 64.2°W	1988-2014	4121	8	0.1	8	-0.6
HOT ALOHA	22.8°N, 158°W	1988-2017	4054	5.4	-0.5	5.1	-2
HOT ALOHA SURF	22.8°N, 158°W	1988-2016	281	6.3	-1.6	5.8	-5.1
ESTOC	29.3°N, 15.5°W	1995-2008	1697	7.2	0.8	6.6	4.7
ICELAND	68°N, 12.7°W	1985-2013	1322	5.8	5.6	6.9	5.3
IRMINGER	64.3°N, 28°W	1991-2013	1086	5.5	3.3	7.6	6.6
K2	47°N, 160°E	1999-2008	615	3.6	1.3	6.4	2.4
KNOT	44°N, 155°E	1997-2008	1321	5.8	-0.8	7.2	-1.9
OWS	66°N, 2°E	2001-2007	803	7.3	-1	10.7	-4.7
KERFIX	50.4°S, 68.2°E	1992-1994	98	13.1	26.4	13	28.9

710

**Table 4. RMSE and bias between measured (and in some cases, computed from measured A<sub>T</sub> and TCO<sub>2</sub> in time series where pCO<sub>2</sub> was not measured) and computed pCO<sub>2</sub> with A<sub>T</sub> from NNGv2 (Broullón et al., 2019) and TCO<sub>2</sub> from NNGv2LDEO in several time series. The time period for pCO<sub>2</sub> from this study is the same as in Table 3. Consult Table 2 in Landschützer et al. (2014) for its time period. The depth range is 0-15 m. Only time series with more than 30 data points are included.**

715

Time series	NNGv2LDEO		Landschützer et al., 2014	
	RMSE (μatm)	bias (μatm)	RMSE (μatm)	bias (μatm)
BATS	17.2	9.7	15.6	0.4



HOT ALOHA SURF	10.3	-3.6	11.6	0.1
ESTOC	10.6	2.7	14.5	-7.1
ICELAND	16	14.8	-	-
IRMINGER	13.1	-1.8	22.6	-1.1
K2	18.1	-3.2	27.8	-0.2
KNOT	20.8	8.6	-	-

**Table 5. Long-term trends (seasonally detrended) of the measured and computed TCO<sub>2</sub> and pCO<sub>2</sub> from neural networks at time-series locations in the depth range 0-15 m.**

Time series	TCO <sub>2</sub> (μmol kg <sup>-1</sup> year <sup>-1</sup> )		pCO <sub>2</sub> (μatm year <sup>-1</sup> )	
	Measured	Computed	Measured*	Computed
BATS	1.2	1.1	1.8	1.7
HOT ALOHA SURF	1.7	1.3	1.8	1.4
ICELAND	0.9	0.9	1.5	1.6
IRMINGER	0.6	0.5	2.5	1.7

\*Computed from measured A<sub>T</sub> and TCO<sub>2</sub> in time series where pCO<sub>2</sub> was not measured.

720

**Table 6: Comparison of four annual mean surface climatologies of TCO<sub>2</sub>.**

RMSE (μmol kg <sup>-1</sup> ) r <sup>2</sup>	NNGv2	Lauvset et al. 2016*	Takahashi et al. 2014
NNGv2LDEO		0.93	0.97
Lauvset et al. 2016*	19.8		0.90
Takahashi et al. 2014	13.2	23.7	

\*The domain analyzed is the same as in Takahashi et al. (2014) for coherency reasons.


725

**Table 7. Comparison of the monthly TCO<sub>2</sub> climatology of Takahashi et al. (2014) and the one of the present study.**

Month	RMSE (μmol kg <sup>-1</sup> )	Bias (μmol kg <sup>-1</sup> )	r <sup>2</sup>
January	16	3.0	0.95
February	16.7	1.5	0.94
March	15.8	2.5	0.95
April	17	2.6	0.95
May	16.8	2.5	0.95
June	17.2	3.2	0.95
July	22.6	4.0	0.92
August	17.8	3.4	0.95
September	15.5	2.5	0.97




October	15.6	2.3	0.96
November	15.7	2.7	0.96
December	17.6	4.3	0.95



730


**Figure 1. Division of the complete database in the data sets needed to train the neural network. The percentages in each level are relative to the number of data in the previous one. Data in the data sets of the first level are always the same for each network. Data in the sets of the second level are randomly associated to each set for each network to find the best network weights, because of the different starting points in the error-weight space of the training process (see also Broullón et al., 2019).**




735

**Figure 2. Regression of TCO<sub>2</sub> computed using NNGv2LDEO and TCO<sub>2</sub> in Gv2 and LDEO. The graph is divided in pixels. The color of each pixel is determined by the number of points inside it. Note the logarithmic scale of the pixels accounting for the large amount of data.**





740 **Figure 3.** The absolute differences between (a) Gv2 TCO<sub>2</sub> and NNGv2LDEO TCO<sub>2</sub> (0-30 m) and (b) LDEO TCO<sub>2</sub> and NNGv2LDEO TCO<sub>2</sub> (0 m).



**Figure 4.** Histograms and box plots of differences between measured and neural network computed  $\text{TCO}_2$  in (a) Gv2 and (b) LDEO. \* $\text{TCO}_2$  computed from measured  $\text{pCO}_2$  and neural network derived  $A_T$ .









Figure 5. Measured (blue line) and computed (orange line) TCO<sub>2</sub> with NGv2LDEO for the depth range 0-15 m (0-30 m in (b)) for several time series. (a) BATS, (b) HOL ALOHA SURFACE, (c) ESTOC, (d) ICELAND, (e) IRMINGER, (f) KNOT, (g) K2 and (h) OWS.




750 Figure 6. Time series of TCO<sub>2</sub> using NNGv2LDEO at (a) BATS and (b) HOT ALOHA locations. The water column shows a higher concentration of TCO<sub>2</sub> year by year. This figure was made with Ocean Data View (Schlitzer, 2016).



755 Figure 7. Annual mean climatology of TCO<sub>2</sub> at (a) 0 m, (b) 100 m and (c) 1000 m. This figure was made with Ocean Data View (Schlitzer, 2016).



**Figure 8.** Seasonal amplitude of TCO<sub>2</sub> at (a) 0 m, (b) 100 m and (c) 1500 m. The contour lines of 25, 50, 75 and 100  $\mu\text{mol kg}^{-1}$  are shown. This figure was made with Ocean Data View (Schlitzer, 2016).



760 **Figure 9.** Comparison of the monthly climatological profiles of TCO<sub>2</sub> computed from measured data (red profile; shadow area is the standard deviation of the averaged values at each depth level) and those from the

<https://doi.org/10.5194/essd-2020-37>  
Preprint. Discussion started: 11 March 2020  
© Author(s) 2020. CC BY 4.0 License.



**TCO: climatology at (a) BATS and (b) HOT ALOHA locations. Units on the x axis are micromoles per kilogram ( $\mu\text{mol kg}^{-1}$ ).**