

CASCADE - The Circum-Arctic Sediment Carbon Database

Jannik Martens¹, Evgeny Romankevich², Igor Semiletov^{3,4,5}, Birgit Wild¹, Bart van Dongen^{1,6}, Jorien Vonk^{1,7}, Tommaso Tesi^{1,8}, Natalia Shakhova⁹, Oleg V. Dudarev³, Denis Kosmach³, Alexander Vetrov², Leopold Lobkovsky², Nikolay Belyaev², Robie Macdonald¹⁰, Anna J. Pieńkowski¹¹, Timothy I. Eglinton¹², Negar Haghipour¹², Salve Dahle¹³, Michael L. Carroll¹³, Emmelie K.L. Åström¹⁴, Jacqueline M. Grebmeier¹⁵, Lee W. Cooper¹⁵, Göran Possnert¹⁶, and Örjan Gustafsson¹

¹Department of Environmental Science and Bolin Centre for Climate Research, Stockholm University, Sweden

²Shirshov Institute of Oceanology, Moscow, Russia

³Il'ichov Pacific Oceanological Institute FEB RAS, Vladivostok, Russia

⁴Tomsk State University, Tomsk, Russia

⁵Tomsk Polytechnic University, Tomsk, Russia

⁶Department of Earth and Environmental Sciences and Williamson Research Centre for Molecular Environmental Science, University of Manchester, United Kingdom

⁷Department of Earth Sciences, Vrije Universiteit Amsterdam, The Netherlands

⁸Institute of Polar Sciences, National Research Council, Bologna, Italy

⁹Department of Chemistry, Moscow State University, Moscow, Russia

¹⁰Institute of Ocean Sciences, Department of Fisheries and Oceans, Sidney, Canada

¹¹Department of Arctic Geology, The University Centre in Svalbard (UNIS), Svalbard, Norway (current address: Norwegian Polar Institute, Longyearbyen, Svalbard, Norway)

¹²Laboratory of Ion Beam Physics and Geological Institute, ETH Zürich, Switzerland

¹³Akvaplan-niva, FRAM - High North Research Centre for Climate and the Environment, Tromsø, Norway

¹⁴Department of Arctic and Marine Biology, UiT-The Arctic University of Norway, Tromsø, Norway

¹⁵Chesapeake Biological Laboratory, University of Maryland Center for Environmental Science, Solomons, USA

¹⁶Department of Physics and Astronomy, Tandem Laboratory, Uppsala University, Sweden

Correspondence to: Örjan Gustafsson (orjan.gustafsson@aces.su.se)

29 Abstract

30 Biogeochemical cycling in the semi-enclosed Arctic Ocean is strongly influenced by land-ocean transport of carbon
31 and other elements, and is vulnerable to environmental and climate changes. Sediments of the Arctic Ocean are
32 important part of biogeochemical cycling in the Arctic, and provide the opportunity to study present and historical
33 input and fate of organic matter (e.g., through permafrost thawing).

34 Comprehensive sedimentary records are required to compare differences between the Arctic regions and to study
35 Arctic biogeochemical budgets. To this end, the Circum-Arctic Sediment Carbon Database (CASCADE) was
36 established to curate data primarily on concentrations of organic carbon (OC) and OC isotopes ($\delta^{13}\text{C}$, $\Delta^{14}\text{C}$), yet
37 also on total N (TN) as well as of terrigenous biomarkers and other sediment geochemical and physical properties.
38 This new database builds on the published literature and earlier unpublished records through an extensive
39 international community collaboration.

40 This paper describes the establishment, structure and current status of CASCADE. The first public version includes
41 OC concentrations in surface sediments at 4244 oceanographic stations including 2317 with TN concentrations,
42 1555 with $\delta^{13}\text{C}$ -OC values, 268 with $\Delta^{14}\text{C}$ -OC values and 653 records with quantified terrigenous biomarkers (high
43 molecular weight *n*-alkanes, *n*-alkanoic acids and lignin phenols). CASCADE also includes data from 326 sediment
44 cores, retrieved by shallow box- or multi-coring, deep gravity/piston coring, or sea-bottom drilling. The
45 comprehensive dataset reveals large-scale features of both OC content and OC sources between the shelf sea
46 recipients. This offers insight to release of pre-aged terrigenous OC to the East Siberian Arctic shelf and younger
47 terrigenous OC to the Kara Sea. Circum-Arctic sediments thereby reveal patterns of terrestrial OC remobilization
48 and provide clues about e.g. thawing of permafrost.

49 CASCADE enables synoptic analysis of OC in Arctic Ocean sediments and facilitates a wide array of future
50 empirical and modelling studies of the Arctic carbon cycle. The database is openly and freely available online
51 (<https://doi.org/10.17043/cascade>; Martens et al., 2020b), is provided in various machine-readable data formats
52 (data tables, GIS shapefile, GIS raster), and also provides ways for contributing data for future CASCADE versions.
53 We will continuously update CASCADE with newly published and contributed data over the foreseeable future as
54 part of the database management of the Bolin Centre for Climate Research at Stockholm University.

55 1 Introduction

56 The Arctic Ocean receives large input of terrestrial organic matter from rivers and coastal erosion, making it a
57 valuable receptor system for studying both large-scale terrestrial carbon remobilization and marine
58 biogeochemistry. Rising temperatures cause multiple changes to the Arctic, including reduced sea-ice cover,
59 accelerated erosion of ice-rich permafrost shorelines and enhanced river runoff, which changes the input of
60 terrestrial organic matter to the Arctic Ocean (AMAP, 2017). This affects nutrients and the detrital load, the ocean
61 optical field, marine primary productivity, ocean acidification and many other aspects of biogeochemical cycling
62 (Stein and Macdonald, 2004; Vonk and Gustafsson, 2013). On land, climate change causes warming and thaw of
63 terrestrial permafrost (Biskaborn et al., 2019), potentially remobilizing parts of its large dormant pool of OC (1300
64 Pg; Hugelius et al., 2014) into active carbon cycling. Rising temperatures may thus shift balances in the Arctic
65 carbon cycle by transformation and translocation of previously frozen organic matter, which leads to system
66 hysteresis effects and translocated carbon-climate feedback (e.g., Vonk and Gustafsson, 2013). Couplings between
67 the large permafrost-carbon pools and amplified climate warming in the Arctic represent a potential “tipping point”
68 in the climate system (Lenton, 2012). These perturbations may affect both OC sequestration in the biosphere and
69 release of climate-forcing greenhouse gases (e.g., AMAP, 2017; IPCC, 2019) as well as the coupling between
70 permafrost carbon remobilization and ocean acidification across the extensive shelf seas (Semiletov et al., 2016).

71
72 Continental shelves cover less than 10% of the global ocean area but account for the largest part of OC accumulation
73 in marine sediments and thereby provide an excellent archive for both terrestrial carbon input and marine
74 productivity (Hedges et al., 1997). The Arctic Ocean is semi-enclosed and dominated by its extensive shelves,
75 including the World’s largest continental shelf system, the East Siberian Arctic Shelf (ESAS; the Laptev, East
76 Siberian and Russian part of the Chukchi Sea). This further accentuates the particular importance of shelf sediments
77 for carbon cycling in the Arctic (Stein et al., 2004; Vetrov and Romankevich, 2004). Earlier landmark contributions
78 have provided comprehensive observational perspectives on the distribution of organic matter in marine sediments
79 at the global scale (e.g., Berner, 1982; Romankevich, 1984; Hedges and Keil, 1995). Focusing in greater detail on
80 carbon in the Arctic, the book by Vetrov and Romankevich (2004) “*Carbon Cycle in the Russian Arctic Seas*” and
81 the book edited by Stein and Macdonald (2004) “*The Organic Carbon Cycle in the Arctic Ocean*” provided the
82 first more comprehensive perspectives on the Arctic land-ocean carbon couplings across various regions. Therein,
83 the authors synthesized the collected knowledge of carbon sources, transformations and burial in Arctic marginal
84 seas and the central Arctic Ocean. These compilations demonstrated substantial regional variations in carbon

85 cycling between different Arctic shelf seas, while also acknowledging the near lack of observational data for key
86 parameters and regions. Substantial progress was made by individual and region-specific studies since then; with
87 key advances in isotope and organic geochemistry that expand the variety of biogeochemical proxies to trace both
88 sources and organic matter degradation. Stable carbon isotopes ($\delta^{13}\text{C}$ -OC) have been widely used to distinguish
89 between marine and terrigenous sources in Arctic Ocean sediments (e.g., Naidu et al., 1993; Mueller-Lupp et al.,
90 2000; Semiletov et al., 2005) and have since then been greatly supplemented by an expanded use of natural
91 abundance radiocarbon ($\Delta^{14}\text{C}$ -OC). This has improved source apportionment of OC in bulk sediments across Arctic
92 regions and time scales (e.g., Vonk et al., 2012; Goñi et al., 2013; Martens et al., 2020) and in sediment density
93 fractions (Tesi et al., 2016b), suspended particulate organic matter (e.g., Vonk et al., 2010, 2014; Karlsson et al.,
94 2016), and at the molecular level (e.g., Drenzek et al., 2007; Gustafsson et al., 2011; Feng et al., 2013). Extensive
95 studies of a wide set of molecular biomarkers (e.g., Fahl and Stein, 1997; Goñi et al., 2000; Belicka et al., 2004;
96 Yunker et al., 2005; van Dongen et al., 2008; Tesi et al., 2014; Sparkes et al., 2015; Bröder et al., 2016) have
97 provided growing insights in OC distribution and fate, particularly for terrigenous organic matter. Access to this
98 growing number of observational data in a readily-accessible interactive format would be greatly beneficial to
99 wider system assessments and interpretations of organic matter in the Arctic Ocean.

100

101 The overarching objective of this effort is to curate and harmonize all available data on OC in Arctic Ocean
102 sediments in an open and freely available database. The Circum-Arctic Sediment CARbon DatabasE (CASCADE)
103 builds on previously-published and unpublished collections holding information on OC and total N (TN)
104 concentrations, as well as OC isotopes ($\delta^{13}\text{C}$ -OC, $\Delta^{14}\text{C}$ -OC) in sediments of all continental shelves and the deep
105 central basins of the Arctic Ocean. Furthermore, CASCADE contains molecular data with an initial focus on
106 terrestrial biomarkers (i.e., high molecular weight - HMW *n*-alkanes, *n*-alkanoic acids, lignin phenols) to facilitate
107 studies of terrestrial OC remobilization. The backbone of CASCADE are large data collections, including i) OC
108 concentrations, $\delta^{13}\text{C}/\Delta^{14}\text{C}$ -isotope data and biomarkers from the informal two-decades long Swedish-Russian
109 collaboration network the International Siberian Shelf Study (ISSS; Semiletov and Gustafsson, 2009) (e.g., Guo et
110 al., 2004; Semiletov et al., 2005; van Dongen et al., 2008; Vonk et al., 2012; Tesi et al., 2016a; Bröder et al., 2018;
111 Martens et al., 2019, 2020; Muschitiello et al., 2020); ii) OC concentrations from the Arctic portion of the “Carbon
112 Database” of the Shirshov Institute of Oceanology, Russian Academy of Sciences (Romankevich, 1984; Vetrov
113 and Romankevich, 2004); iii) previously-published databases and online collections (e.g., Pangaea.de) with many
114 contributions from German-Russian partnerships and cruises involving the Alfred-Wegener-Institute, Germany

115 (e.g., Stein et al., 1994; Mueller-Lupp et al., 2000; Stein and Macdonald, 2004; Xiao et al., 2015); iv) US and
116 Canadian based research (e.g., Naidu et al., 1993, 2000; Goñi et al., 2000, 2013; Grebmeier et al., 2006); and v)
117 data from various other contributors that are acknowledged in the database. The initial version also includes
118 previously unpublished data, with some generated here in the upstart of CASCADE, to fill gaps for particularly
119 data-lean regions such as the Barents and Kara Seas, the Canadian Arctic Archipelago, and the Chukchi Sea.

120

121 The aim of the CASCADE effort is to provide a foundation for future studies. These may include large-scale
122 assessments of the carbon cycle, such as characteristics of OC input, and its distribution and fate in the Arctic
123 Ocean. This paper describes the creation and the structure of CASCADE, including a discussion of data availability
124 and quality.

125

126 **2 Data collection and methods**

127 **2.1 The physical compartments: Arctic shelf seas and interior Arctic Ocean basins**

128 The CASCADE includes OC data from the entire Arctic Ocean with special focus on the seven Arctic continental
129 shelf seas (Fig. 1; Table 1). Accordingly, a distinction is made among the central Arctic Ocean and the following
130 marginal seas: Beaufort Sea, Chukchi Sea, East Siberian Sea, Laptev Sea, Kara Sea, Barents Sea (incl. White Sea),
131 and the Canadian Arctic Archipelago (we exclude data from Baffin Bay, Foxe Basin and Hudson Bay, as they are
132 outside the Circum-Arctic scope of the database). For defining the limits of these Arctic shelf seas, Jakobsson
133 (2002) is followed, which distinguishes the Arctic Ocean constituent seas using hypsometric criteria. Therein, *shelf*
134 is defined as the seaward extension of the continental margin until the increase in steepness at the shelf break
135 (Jakobsson, 2002). Data for the central Arctic Ocean was treated as one individual unit that covers all area beyond
136 the shelf break and includes the continental slope, rise, deep basins and mid-ocean ridges.

137

138 **2.2 Georeferencing and sampling**

139 The coordinate system used for CASCADE is WGS1984 and coordinates are kept in machine-readable decimal
140 degrees (Latitude in °N, Longitudes in the -180° to 180° format) to harmonize the data across all GIS applications.
141 The collection of data from oceanographic stations is the main part of CASCADE and is organized in a table format

142 that contains columns for the station number ('STATION') and geographical coordinates ('LAT'; 'LON'). The
143 spatial references also include information about the sediment depth interval that reported data represent
144 ('UPPERDEPTH'; 'LOWERDEPTH'), where the upper depth is equal to 0 cm in the case of surface sediments. In
145 addition, the table contains a column for water depth ('WATERDEPTH') as reported by the data source. In cases
146 where the water depth was not reported, the water depth was estimated using the latest version (v4) of the
147 bathymetric map of IBCAO (Jakobsson et al., 2020) corresponding to the position of the oceanographic station and
148 reported in a separate column ('IBCAODEPTH'). Furthermore, the name of the expedition and/or ship
149 ('EXPEDITION') and the year when the sample was taken ('YEAR') are reported. For samples where the sampling
150 year was unknown, users may use the year of publication instead.

151

152 **2.3 Surface sediments and sediment cores**

153 The first stage of the CASCADE development focused on maximizing spatial coverage for surface sediments of
154 the seven Circum-Arctic shelf sea systems and the central Arctic Ocean. Here, surface sediments are defined as
155 those collected from the water-sediment interface to a depth of maximum 5 cm. Data for surface sediments are
156 provided in a table ('CASCADEsurfscd') as .txt and .xlsx files and in a ready-to-use GIS shapefile format. This
157 database also includes deeper sediments from sediment cores, which represent longer time-scales and add a third
158 dimension to the geographical referencing. Types of sediment cores are distinguished in CASCADE such that
159 different biogeochemical processes, acting on three depositional time scales, may be addressed. The three time
160 scales are:

161

- 162 1. Centennial scale cores (core scale 1) in upper sediments of the Arctic Ocean, e.g., multi corer, Gemini
163 corer, box corer, van Veen grab sampler, other short gravity corers up to 1 m length;
- 164 2. Millennial scale cores (core scale 2) of shelf sediments roughly covering the depositional time frame from
165 the late Holocene to the last glacial/interglacial transition, by e.g., piston corer, long gravity corer, kasten
166 corer;
- 167 3. Glacial cycles scale cores (core scale 3) from the continental slopes or the deeper Arctic Ocean basins
168 covering periods from earlier than the Last Glacial Maximum, including e.g. drill coring on the Circum-
169 Arctic shelves or deep-sea piston cores.

170

Downcore data are stored in three separate data tables ('CASCADEcorescale1'; 'CASCADEcorescale2'; 'CASCADEcorescale3') in addition to the surface sediment files, including a column for the sampling depth of core subsamples in cm below the sediment surface ('COREDEPTH').

2.4 Database parameters

CASCADE contains information about the concentration, isotopic and molecular composition of OC in marine Arctic sediments. In addition to i) OC concentrations (column 'OC'), the database includes ii) concentrations of TN ('TN') and iii) the gravimetric ratio of OC/TN ('OC/TN'), which may provide additional information about the organic matter source (e.g., Goñi et al., 2005; van Dongen et al., 2008). Furthermore, CASCADE contains data of iv) $\delta^{13}\text{C}$ -OC ('d13C') as a parameter to distinguish between marine and terrestrial sources (e.g., Fry and Sherr, 1989), and v) $\Delta^{14}\text{C}$ -OC ('D14C') to assess the presence of aged organic matter released from e.g., permafrost deposits (e.g., Gustafsson et al., 2011; Vonk et al., 2012) or from petrogenic sources such as sedimentary rocks (e.g., Yunker et al., 2005; Goñi et al., 2013) in marine sediments. More details about the CASCADE parameters and their units are provided in Table 2.

Data of terrigenous biomarkers may facilitate further investigations of terrigenous OC input (Table 2). The first version of CASCADE compiles total concentrations of *n*-alkanes with high molecular weight (HMW) and C_{21} - C_{31} carbon atoms ($\sum \text{C}_{21}\text{-C}_{31}$; column 'HMWALK'), as well as the often separately-reported more specific *n*-alkanes $\sum \text{C}_{27}+\text{C}_{29}+\text{C}_{31}$ ('HMWALK_SPEC'). CASCADE also contains the sum of the HMW *n*-alkanoic acids $\sum \text{C}_{20}\text{-C}_{30}$ ('HMWACID'). Both compound classes stem mostly from terrigenous compartments as they derive from epicuticular leaf waxes of land plants with a typical pattern of dominating odd-numbered homologues for HMW *n*-alkanes and even-numbered homologues for HMW *n*-alkanoic acids (Eglinton and Hamilton, 1967). Furthermore, the database holds concentrations of lignin phenols (\sum syringyl, vanillyl, cinnamyl; 'LIGNIN'), which are products from the break-up of the lignin biopolymer, a compound only produced by vascular plants (Hedges and Mann, 1979). These three compound classes are frequently used as tracers of the sources and fate of terrestrial organic matter sequestered in Arctic Ocean sediments (Fahl and Stein, 1997; Goñi et al., 2000; Tesi et al., 2014; Bröder et al., 2016). It is recognized that there are more parameters that could be included and CASCADE can add further extensions in future versions.

200 2.5 Reference to the original publication

201 Each data source added to CASCADE is fully cited (in the formatting style of Earth Systems Science Data; ESSD)
202 to maintain a high level of transparency. When applicable, citations also include a digital object identifier (doi) that
203 is linked to the reference in the primary literature next to each parameter column. Accordingly, the CASCADE
204 data sheet distinguishes between a common reference for OC, TN and OC/TN data ('CN_CITATION') as they are
205 often combined in one measurement, and separate references for OC isotopes ('d13C_CITATION';
206 'D14C_CITATION') and concentrations of biomarkers ('BM_CITATION'). This facilitates to register multiple
207 measurements based on the same or split sediment sample material for individual oceanographic stations. A full
208 list of references is separately provided on the CASCADE website and in the Supplementary Information of this
209 paper.

211 2.6 Data source and quality

212 A part of CASCADE builds on previous separate and partly inaccessible databases of OC parameters that key
213 partners of the CASCADE consortium and others have collected over the years. This includes data from the
214 informal Swedish-Russian led collaboration network called the International Siberian Shelf Study (ISSS; Semiletov
215 and Gustafsson, 2009) and the "Carbon" database of the Shirshov Institute of Oceanology. This basis for
216 CASCADE was strengthened by an extensive survey of the peer-reviewed literature and data mining in the grey
217 literature of scientific cruise reports. To facilitate quality assurance criteria by the end users the database also
218 records metadata (e.g., sampling technique in the field, sample storage) and quality data when available. The quality
219 assurance information for data in CASCADE are:

- 220 • Data need to be (geo-)referenceable and located in the target region (i.e. the Arctic Ocean).
- 221 • Information about the analysis method is provided by the data source.
- 222 • For OC concentrations, values were generated by elemental analyzer (EA) or Rock Eval pyrolysis and
223 reported as weight-% OC. Total N concentrations and OC/TN ratios are based on EA only.
- 224 • For $\delta^{13}\text{C}$ -OC data stored in CASCADE are based on isotope ratio mass spectrometry (IRMS), often coupled
225 to an EA and calibrated against the PDB/V-PDB analytical standards.
- 226 • For $\Delta^{14}\text{C}$ -OC the measurements of ^{14}C data are based on mass spectrometry with ^{14}C data reported as $\Delta^{14}\text{C}$,
227 fraction modern (F_m) or conventional ^{14}C age in the original publication. We also kept records of the
228 ^{14}C /AMS lab code of the sample if given.

- Terrigenous biomarker analysis was carried out by solvent extraction (for HMW *n*-alkanes and *n*-alkanoic acids) or by alkaline CuO oxidation of the lignin biopolymer (for lignin phenols) of the sediments, followed by wet chemistry purification and quantification using gas chromatography analysis with either flame ionization or mass spectrometry detection.

In addition to the above-mentioned information, the aim was to also include information about carbonate removal by acid treatment prior to the measurement of OC, $\delta^{13}\text{C}$ -OC and $\Delta^{14}\text{C}$ -OC. However, details about applied procedures were missing in most cases and it is therefore assumed that the carbonate fraction was removed from total carbon prior to OC, $\delta^{13}\text{C}$ -OC and $\Delta^{14}\text{C}$ -OC measurements. All meta information (sampling, storage, analysis) to each CASCADE entry is included in a respective column in the data spreadsheet (Table 2).

2.7 New gap filling analyses

2.7.1 Bulk OC and carbon isotopes

Gap filling was performed in surface sediments of regions with particularly poor data density. These efforts thus focused on areas north of western Siberia (Barents and Kara Sea region) and in the Canadian Arctic Archipelago, using archived sample material that was provided by CASCADE collaborators. For OC, TN and $\delta^{13}\text{C}$ -OC analysis, about 10 mg each of a total of 153 freeze-dried sediment samples were weighed in silver capsules and acidified drop-wise with 3 M HCl in order to remove carbonates. The measurement was carried out using a Carlo Erba NC2500 elemental analyzer coupled to an isotope-ratio mass spectrometer (Finnigan DeltaV Advantage) in the Department of Geological Sciences, Stockholm University, with $\pm 3\%$ precision for OC analysis and $\pm 0.15\%$ precision for $\delta^{13}\text{C}$ -OC isotopic measurements.

Furthermore, a subset of 95 samples was selected for gap-filling bulk-level $\Delta^{14}\text{C}$ -OC analysis at the Tandem Laboratory, Department of Physics, Uppsala University. A sample amount corresponding to 1 mg OC was weighed in tin capsules and acidified with 3 M HCl to remove carbonates. Samples with low OC concentrations ($< 0.5\%$) were placed in small beakers and exposed to acid fumes in a desiccator for 24 h to remove carbonates and combusted to CO_2 in evacuated quartz tubes prior to graphitization at the ^{14}C /AMS laboratory. An additional set of 30 gap-filling samples was analyzed for $\Delta^{14}\text{C}$ at the ^{14}C laboratory of ETH Zurich after acid fumigation. The measurements at Uppsala University had a precision of on average $\pm 1.9\%$ while the precision at ETH Zurich was on average $\pm 1.1\%$ (based on ^{14}C counting statistics).

258 In CASCADE, all new data points are labelled by citing the database ('Martens et al., 2021. CASCADE - The
 259 Circum-Arctic Sediment CARbon DatabasE. Bolin Centre for Climate Research, Stockholm University, Sweden.
 260 doi:10.17043/cascade, 2021.') in the respective reference columns.
 261

262 2.7.2 Analysis of lignin phenols

263 Gap-filling analysis was also performed for lignin phenols as molecular biomarkers for terrestrial organic matter
 264 using a set of 64 samples from data-lean regions. To extract lignin phenols from marine sediments we applied an
 265 alkaline CuO oxidation protocol using a microwave-based method as originally presented by Goñi and
 266 Montgomery (2000) and followed the same laboratory routine as described in greater detail elsewhere (Tesi et al.,
 267 2014; Martens et al., 2019).
 268

269 2.8 Data conversion and harmonization

270 Recalculations of literature data (e.g., for unit conversions) were in some cases necessary to harmonize the data to
 271 the standard units as defined in Table 2.

- 272 • In CASCADE the concentration of OC is reported in percent (%) of the dry weight; values previously
 273 published as mg OC per g dry weight were divided by a factor of 10.
- 274 • CASCADE uses $\Delta^{14}\text{C}$ with age correction (equation 1) to report the activity of radiocarbon according to
 275 convention (Stuiver and Polach, 1977; Stenström et al., 2011). For radiocarbon values that were reported
 276 as conventional ^{14}C ages we used equation 2 to calculate the age-corrected $\Delta^{14}\text{C}$.

$$277 \quad \Delta^{14}\text{C} = (F_m \cdot e^{\lambda_c(1950-Y_c)} - 1) \cdot 1000\text{‰} \quad (1)$$

$$278 \quad \Delta^{14}\text{C} = (e^{-\lambda_L \cdot T_{14\text{C-years}}} \cdot e^{\lambda_c(1950-Y_c)} - 1) \cdot 1000\text{‰} \quad (2)$$

279 Where F_m is the fraction modern, λ_c the decay constant of the Cambridge half-life of ^{14}C ($T_{1/2-C} = 5730$; λ_c
 280 $= 1/8267$), Y_c the year of sample collection, λ_L the decay constant of the Libby half-life of ^{14}C ($T_{1/2-L} =$
 281 5568 ; $\lambda_c = 1/8033$ and $T_{14\text{C-years}}$ the conventional ^{14}C age (Stuiver and Polach, 1977).

- 282 • All biomarker concentrations of HMW *n*-alkanes and *n*-alkanoic acids are reported as μg per g OC while
 283 lignin phenols are reported as mg per g OC. Biomarker concentrations that in the original publication were
 284 reported as normalized to dry sediment weight were for CASCADE normalized to the OC concentration
 285 of the sample.

287 **2.9 Data interpolation**

288 CASCADE provides interpolated files (GEOtiff, ASCII; coordinate system WGS 1984 Arctic Polar Stereographic)
 289 for OC content, $\delta^{13}\text{C}$ -OC and for $\Delta^{14}\text{C}$ -OC in surface sediments across the Arctic Ocean. OC data was mapped in
 290 ArcGIS 10.6 and interpolated to a resolution of 5x5 km per grid cell using the Empirical Bayesian Kriging function
 291 (EBK; Gribov and Krivoruchko, 2020) in the commercially-available ArcGIS 10.8 software package (ESRI).
 292 Kriging builds on the assumption that two points located in proximity are more similar than two points further
 293 distant and creates a gridded surface of predicted values using an empirical semivariogram model. As an
 294 advancement to Kriging, EBK repeatedly simulates semivariogram models in subsets of up to 200 data points and
 295 thus not only improves the prediction but also optimizes interpolation across areas with strongly varying data
 296 availability in the Arctic Ocean (e.g., shelf seas vs. central basins).

297 **3 Results and Discussion**

298 **3.1 Data set inventory**

299 Surface sediments show by far the largest data availability. The dataset of OC concentrations in CASCADE
 300 includes 4244 different locations across the Arctic Ocean (Fig. 2), while the concentration of TN and the OC/TN
 301 ratio are known for 2317 locations (Table 1). For carbon isotopes, the number of individual $\delta^{13}\text{C}$ -OC values is 1555
 302 and for $\Delta^{14}\text{C}$ -OC it is 268. CASCADE also holds concentrations of terrigenous biomarkers at 131-213 locations
 303 per compound group. Most of the biomarker data is for HMW *n*-alkanes, with either concentrations of HMW *n*-
 304 alkanes ($\Sigma\text{C}_{21}\text{-C}_{31}$; 213 stations) or chain lengths more specific for higher plants (ΣC_{27} , C_{29} , C_{31} ; 164). Fewer data
 305 are available for concentrations of HMW *n*-alkanoic acids ($\Sigma\text{C}_{20}\text{-C}_{30}$; 131) and the concentrations of lignin phenols
 306 (145).

307

308 In addition to surface sediments, a total number of 326 sediment cores (79 centennial, 229 millennial, and 18 glacial
 309 cycle scale cores) is included in the first version of CASCADE. Combined, these hold another 10552 observations
 310 of OC concentrations, 4769 concentrations of TN and 2122 $\delta^{13}\text{C}$ -OC ratios in core samples from across the Arctic
 311 Ocean.

312

313 3.2 Spatial distribution of data

314 The data coverage for surface sediments is highly variable among the shelf seas, yet improved by the extensive
315 gap-filling analysis (Table 1). The largest number of OC concentrations is in the Barents Sea (1092; Table 1).
316 Despite the large total number of available Arctic sediment OC concentrations, there are only 236 samples analyzed
317 for $\delta^{13}\text{C}$ -OC and 33 with $\Delta^{14}\text{C}$ -OC in the Barents Sea, and of these most are located in the Norwegian (western)
318 sector of the Barents Sea. For the eastern Siberian Arctic and the North American sector of the Arctic Ocean,
319 observations of OC concentrations are lower but the availability of $\delta^{13}\text{C}$ -OC data is higher (Table 1, Fig 2b, c).
320 Accordingly, the Kara, Laptev, East Siberian and Chukchi Seas each support more than 200 $\delta^{13}\text{C}$ -OC observations.
321 The number of $\Delta^{14}\text{C}$ -OC observations is generally lower but reveals highest coverage in near-coastal areas, with
322 28 values in the Kara Sea, 42 values in the Laptev and 71 values in the East Siberian Sea. Data availability in the
323 Chukchi Sea for $\Delta^{14}\text{C}$ -OC is lower ($n=12$), stressing the need for future analysis. The lowest availability of data is
324 in the Canadian Arctic Archipelago. Gap-filling analysis of OC here increased the number of OC concentrations
325 from 21 to 54, with a similar number for carbon isotopes (51 of $\delta^{13}\text{C}$ -OC; 22 of $\Delta^{14}\text{C}$ -OC) distributed over its vast
326 area of 1,171,000 km². The largest individual regime area is covered by the interior basins of the central Arctic
327 Ocean, which holds 529 observations of OC concentrations, 130 of $\delta^{13}\text{C}$ -OC and 27 of $\Delta^{14}\text{C}$ -OC values.

328

329 3.3 Assessment of data quality

330 Based on the quality assurance data available, CASCADE provides detailed information about the techniques
331 involved in analyzing OC concentrations, isotopes and biomarkers. The development of CASCADE included the
332 collection of meta information about sampling, storage and analysis, as described in section 2.6. This information
333 is included and detailed in CASCADE. The quality assurance information shows that 86% of the reported OC
334 concentrations were analyzed using EA and only a minority was analyzed by Rock Eval pyrolysis. For $\delta^{13}\text{C}$ -OC,
335 in 66% of the cases IRMS coupled to EA was reported as the method of analysis. Regarding sample storage,
336 information was given in about 59% of all data sources that the samples were kept frozen between sampling and
337 analysis, while for <1% of the cases it was documented that the samples were stored refrigerated; this means that
338 for 40% of the samples, there was no information provided about sample storage. For 78% of the $\Delta^{14}\text{C}$ -OC values,
339 the laboratory ^{14}C /AMS label was documented and thus also added to the CASCADE sheet.

340

341 3.4 Circum-Arctic carbon features

342 Visualization of CASCADE data directly reveals several large-scale features of OC in Arctic Ocean sediments.
343 These include clear differences in both OC concentration and source-diagnostic isotope composition among the
344 shelf seas. For instance, interpolated OC concentrations (Fig. 2) indicate that high sedimentary OC content is found
345 both in regions of high terrestrial input (e.g., Kara Sea, Laptev Sea, East Siberian Sea and Beaufort Sea) and in
346 regions of high nutrient availability and marine primary productivity (Barents Sea and Chukchi Sea). The
347 combination of $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ isotope values delineate large-scale differences in OC sources. Values of $\delta^{13}\text{C}$ -OC
348 close to marine OC (-21‰; Fry and Sherr, 1989) and $\Delta^{14}\text{C}$ reflecting contemporary carbon are consistent with high
349 marine primary productivity in the Barents Sea and Chukchi Sea. The Kara Sea receives input from major West
350 Siberian catchments (Ob and Yenisey rivers), with sediment OC that appears to reflect OC from contemporary
351 terrestrial sources (~-27‰; Fry and Sherr, 1989). By contrast, the terrigenous OC fraction in the Laptev and East
352 Siberian Seas is much older with a presumably substantial contribution from remobilization of thawing permafrost
353 or other old deposits via erosional or fluvial processes (Fig. 1; Fig. 2). These and other features can now be
354 investigated through CASCADE at greater quantitative detail over large intra- and inter-system scales.

355

356 4 Vision and future development

357 The CASCADE is the largest and most comprehensive open-access database of OC parameters for Arctic Ocean
358 sediments. It is a resource that can facilitate a wide range of investigations on OC cycling in the high northern
359 latitudes. For instance, CASCADE may help research on sources of organic matter, marine primary production,
360 OC degradation, OC transport both in the offshore direction and vertically from the sea surface to the sediment;
361 and all this both in the contemporary and the historical perspectives. CASCADE provides opportunities to expand
362 our still limited understanding of how sensitive terrestrial permafrost in different circum-Arctic regions is towards
363 remobilization in both, the current and over earlier periods of rapid climate change. Future versions of CASCADE
364 may also expand on parameters by adding more compound classes of terrestrial biomarkers, marine biomarkers,
365 environmental contaminants (e.g., Hg and organic legacy and emerging substances) and others to investigate
366 biogeochemical distribution and fate of these in the Arctic Ocean.

367

368 **5 Data availability**

369 CASCADE will be hosted and actively updated and extended by a database management at the Bolin Centre for
370 Climate Research at Stockholm University. CASCADE is accessible at the Bolin Centre data repository
371 (<https://doi.org/10.17043/cascade>; Martens et al., 2020b). When using the CASCADE, this paper and the database
372 should be cited. The website also includes contact details, which can be used to submit new data for incorporation
373 into future versions of CASCADE – a community effort and resource.

374 **Author contributions**

375 The CASCADE database was conceptualized and planned by a team led by ÖG, IS and ER. JM, NB, BW and ÖG
376 developed the technical framework of the CASCADE. JM executed the development of the CASCADE, populated
377 the database with published and unpublished data from the literature and internal records, coordinated gap-filling
378 analyses and created maps. JM drafted and coordinated the manuscript in close collaboration with ÖG and BW.
379 All authors contributed to the realization of the CASCADE database and participated in the editing of the
380 manuscript.

381 **Acknowledgements**

382 We thank collaborators throughout and beyond the International Siberian Shelf Study (ISSS) network, and all
383 participants of the Arctic Partner Forum 2018 for their advice in constructing the CASCADE database and for
384 pointing out data sources during the development of the database. We also thank the crew and the scientific party
385 of the ISSS-08 expedition onboard *r/v* Yacob Smirnitskyi, the SWERUS-C3 expedition onboard the *i/b* Oden in
386 2014 and various other field campaigns organized by the ISSS in 2004, 2005, 2007, 2008, 2011, 2016 and 2017.
387 Furthermore, Lisa Bröder and Rickard Wennström are thanked for their help with gap-filling analysis of lignin
388 phenols. We also thank August Andersson and Henry Holmstrand for their long-term assistance and advice during
389 various field campaigns, laboratory analyses and computer-based work that contributed to the realization of this
390 database.

391 **Funding**

392 Development of the CASCADE was supported by the European Research Council (ERC Advanced Grant CC-TOP
393 695331 to ÖG), the EU H2020-funded project Nunataryuk (Grant 773421) and the Swedish Research Council
394 (Grant 2017-01601). Field campaigns to obtain gap-filling samples were supported by the Knut and Alice
395 Wallenberg Foundation (KAW contract 2011.0027 to ÖG) as part of the SWERUS-C3 program, as well as by the
396 Russian Science Foundation (grant 21-77-30001 to IS). Furthermore, this study was supported by the assignment
397 of the Russian Academy of Sciences (grant 0128-2021-0005) and the Russian Science Foundation (grant 18-05-
398 60214) to the Shirshov Institute of Oceanology (ER, AV). The collection of sample material in the Barents Sea was
399 supported by the Research Council of Norway (grant 228107 to MLC; grant 223259) and VISTA (grant 6172 to
400 EKLÅ). Gap-filling samples from the Canadian Arctic were supported by the Research Council of Canada (NSERC

401 Discovery Grant RGPIN-2016-05457 to AJP). BvD was supported by NERC research grant (NE/I024798/1) and
402 JV was supported by the Dutch-NWO (Veni grant # 863.12.004).

403

404 **Competing interests**

405 The authors declare that they have no conflict of interest.

406 References

- 407 AMAP: Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017. Arctic Monitoring and Assessment
408 Programme (AMAP), Oslo, Norway., 2017.
- 409 Belicka, L. L., Macdonald, R. W., Yunker, M. B. and Harvey, H. R.: The role of depositional regime on carbon
410 transport and preservation in Arctic Ocean sediments, *Mar. Chem.*, 86(1–2), 65–88,
411 doi:10.1016/j.marchem.2003.12.006, 2004.
- 412 Berner, R. A.: Burial of organic carbon and pyrite sulfur in the modern ocean: Its geochemical and environmental
413 significance, *Am. J. Sci.*, 282(4), 451–473, doi:10.2475/ajs.282.4.451, 1982.
- 414 Biskaborn, B. K., Smith, S. L., Noetzi, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P., Romanovsky,
415 V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R.,
416 Diekmann, B., Drozdov, D., Etzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T., Isaksen, K.,
417 Ishikawa, M., Johansson, M., Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P., Kröger, T.,
418 Lambiel, C., Lanckman, J.-P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M., Phillips, M.,
419 Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q., Yoshikawa, K.,
420 Zheleznyak, M. and Lantuit, H.: Permafrost is warming at a global scale, *Nat. Commun.*, 10(1), 264,
421 doi:10.1038/s41467-018-08240-4, 2019.
- 422 Bröder, L., Tesi, T., Salvadó, J. A., Semiletov, I. P., Dudarev, O. V and Gustafsson, Ö.: Fate of terrigenous organic
423 matter across the Laptev Sea from the mouth of the Lena River to the deep sea of the Arctic interior,
424 *Biogeosciences*, 13(17), 5003–5019, doi:10.5194/bg-13-5003-2016, 2016.
- 425 Bröder, L., Tesi, T., Andersson, A., Semiletov, I. and Gustafsson, Ö.: Bounding the role of cross-shelf transport
426 and degradation in land-ocean carbon transfer, *Nat. Commun.*, 9(1), 806, doi:10.1038/s41467-018-03192-1, 2018.
- 427 van Dongen, B. E., Semiletov, I., Weijers, J. W. H. and Gustafsson, Ö.: Contrasting lipid biomarker composition
428 of terrestrial organic matter exported from across the Eurasian Arctic by the five great Russian Arctic rivers, *Global*
429 *Biogeochem. Cycles*, 22(1), doi:10.1029/2007GB002974, 2008.
- 430 Drenzek, N. J., Montluçon, D. B., Yunker, M. B., Macdonald, R. W. and Eglinton, T. I.: Constraints on the origin
431 of sedimentary organic carbon in the Beaufort Sea from coupled molecular ^{13}C and ^{14}C measurements, *Mar.*
432 *Chem.*, 103(1–2), 146–162, doi:10.1016/j.marchem.2006.06.017, 2007.
- 433 Eglinton, G. and Hamilton, R. J.: Leaf Epicuticular Waxes, *Science*, 156(3780), 1322–1335,
434 doi:10.1126/science.156.3780.1322, 1967.
- 435 Fahl, K. and Stein, R.: Modern organic carbon deposition in the Laptev Sea and the adjacent continental slope:
436 Surface water productivity vs. terrigenous input, *Org. Geochem.*, 26(5–6), 379–390, doi:10.1016/S0146-
437 6380(97)00007-7, 1997.
- 438 Feng, X., Vonk, J. E., van Dongen, B. E., Gustafsson, Ö., Semiletov, I. P., Dudarev, O. V, Wang, Z., Montluçon,
439 D. B., Wacker, L. and Eglinton, T. I.: Differential mobilization of terrestrial carbon pools in Eurasian Arctic river
440 basins., *Proc. Natl. Acad. Sci. U. S. A.*, 110(35), 14168–73, doi:10.1073/pnas.1307031110, 2013.

- 441 Fry, B. and Sherr, E. B.: $\delta^{13}\text{C}$ Measurements as Indicators of Carbon Flow in Marine and Freshwater Ecosystems,
442 in *Stable Isotopes in Ecological Research*. Ecological Studies (Analysis and Synthesis), pp. 196–229, Springer,
443 New York, NY., 1989.
- 444 Goñi, M. A. and Montgomery, S.: Alkaline CuO Oxidation with a Microwave Digestion System: Lignin Analyses
445 of Geochemical Samples, *Anal. Chem.*, 72(14), 3116–3121, doi:10.1021/ac991316w, 2000.
- 446 Goñi, M. A., Yunker, M. B., MacDonald, R. W. and Eglinton, T. I.: Distribution and sources of organic biomarkers
447 in arctic sediments from the Mackenzie River and Beaufort Shelf, *Mar. Chem.*, 71(1–2), 23–51,
448 doi:10.1016/S0304-4203(00)00037-2, 2000.
- 449 Goñi, M. A., Yunker, M. B., Macdonald, R. W. and Eglinton, T. I.: The supply and preservation of ancient and
450 modern components of organic carbon in the Canadian Beaufort Shelf of the Arctic Ocean, *Mar. Chem.*, 93(1), 53–
451 73, doi:10.1016/j.marchem.2004.08.001, 2005.
- 452 Goñi, M. A., O'Connor, A. E., Kuzyk, Z. Z., Yunker, M. B., Gobeil, C. and Macdonald, R. W.: Distribution and
453 sources of organic matter in surface marine sediments across the North American Arctic margin, *J. Geophys. Res.*
454 *Ocean.*, 118(9), 4017–4035, doi:10.1002/jgrc.20286, 2013.
- 455 Grebmeier, J. M., Cooper, L. W., Feder, H. M. and Sirenko, B. I.: Ecosystem dynamics of the Pacific-influenced
456 Northern Bering and Chukchi Seas in the Amerasian Arctic, *Prog. Oceanogr.*, 71(2–4), 331–361,
457 doi:10.1016/j.pocean.2006.10.001, 2006.
- 458 Gribov, A. and Krivoruchko, K.: Empirical Bayesian kriging implementation and usage, *Sci. Total Environ.*, 722,
459 137290, doi:10.1016/j.scitotenv.2020.137290, 2020.
- 460 Guo, L., Semiletov, I., Gustafsson, Ö., Ingri, J., Andersson, P., Dudarev, O. and White, D.: Characterization of
461 Siberian Arctic coastal sediments: Implications for terrestrial organic carbon export, *Global Biogeochem. Cycles*,
462 18(1), doi:10.1029/2003gb002087, 2004.
- 463 Gustafsson, Ö., van Dongen, B. E., Vonk, J. E., Dudarev, O. V and Semiletov, I. P.: Widespread release of old
464 carbon across the Siberian Arctic echoed by its large rivers, *Biogeosciences*, 8(6), 1737–1743, doi:10.5194/bg-8-
465 1737-2011, 2011.
- 466 Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation: an assessment and speculative synthesis,
467 *Mar. Chem.*, 49(2–3), 81–115, doi:10.1016/0304-4203(95)00008-F, 1995.
- 468 Hedges, J. I. and Mann, D. C.: The characterization of plant tissues by their lignin oxidation products, *Geochim.*
469 *Cosmochim. Acta*, 43(11), 1803–1807, doi:10.1016/0016-7037(79)90028-0, 1979.
- 470 Hedges, J. I., Keil, R. G. and Benner, R.: What happens to terrestrial organic matter in the ocean?, in *Organic*
471 *Geochemistry*, vol. 27, pp. 195–212, Pergamon., 1997.
- 472 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G.,
473 Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J. and
474 Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified

- 475 data gaps, *Biogeosciences*, 11(23), 6573–6593, doi:10.5194/bg-11-6573-2014, 2014.
- 476 Jakobsson, M.: Hypsometry and volume of the Arctic Ocean and its constituent seas, *Geochemistry, Geophys.*
 477 *Geosystems*, 3(5), 1–18, doi:10.1029/2001GC000302, 2002.
- 478 Jakobsson, M., Mayer, L. A., Bringsenparr, C., Castro, C. F., Mohammad, R., Johnson, P., Ketter, T., Accettella,
 479 D., Amblas, D., An, L., Arndt, J. E., Canals, M., Casamor, J. L., Chauché, N., Coakley, B., Danielson, S., Demarte,
 480 M., Dickson, M. L., Dorschel, B., Dowdeswell, J. A., Dreutter, S., Fremand, A. C., Gallant, D., Hall, J. K.,
 481 Hehemann, L., Hodnesdal, H., Hong, J., Ivaldi, R., Kane, E., Klaucke, I., Krawczyk, D. W., Kristoffersen, Y.,
 482 Kuipers, B. R., Millan, R., Masetti, G., Morlighem, M., Noormets, R., Prescott, M. M., Rebesco, M., Rignot, E.,
 483 Semiletov, I., Tate, A. J., Travaglini, P., Velicogna, I., Weatherall, P., Weinrebe, W., Willis, J. K., Wood, M.,
 484 Zarayskaya, Y., Zhang, T., Zimmermann, M. and Zinglensen, K. B.: The International Bathymetric Chart of the
 485 Arctic Ocean Version 4.0, *Sci. Data*, 7(176), doi:10.1038/s41597-020-0520-9, 2020.
- 486 Karlsson, E., Gelting, J., Tesi, T., van Dongen, B., Andersson, A., Semiletov, I., Charkin, A., Dudarev, O. and
 487 Gustafsson, Ö.: Different sources and degradation state of dissolved, particulate, and sedimentary organic matter
 488 along the Eurasian Arctic coastal margin, *Global Biogeochem. Cycles*, 30(6), 898–919 [online] Available from:
 489 <http://doi.wiley.com/10.1002/2015GB005307>, 2016.
- 490 Lantuit, H., Overduin, P. P., Couture, N., Wetterich, S., Aré, F., Atkinson, D., Brown, J., Cherkashov, G., Drozdov,
 491 D., Donald Forbes, L., Graves-Gaylord, A., Grigoriev, M., Hubberten, H. W., Jordan, J., Jorgenson, T., Ødegård,
 492 R. S., Ogorodov, S., Pollard, W. H., Rachold, V., Sedenko, S., Solomon, S., Steenhuisen, F., Streletskaia, I. and
 493 Vasiliev, A.: The Arctic Coastal Dynamics Database: A New Classification Scheme and Statistics on Arctic
 494 Permafrost Coastlines, *Estuaries and Coasts*, 35(2), 383–400, doi:10.1007/s12237-010-9362-6, 2012.
- 495 Lenton, T. M.: Arctic climate tipping points, *Ambio*, 41(1), 10–22, doi:10.1007/s13280-011-0221-x, 2012.
- 496 Martens, J., Wild, B., Pearce, C., Tesi, T., Andersson, A., Bröder, L., O'Regan, M., Jakobsson, M., Sköld, M.,
 497 Gemery, L., Cronin, T. M., Semiletov, I., Dudarev, O. V. and Gustafsson, Ö.: Remobilization of Old Permafrost
 498 Carbon to Chukchi Sea Sediments During the End of the Last Deglaciation, *Global Biogeochem. Cycles*, 33(1), 2–
 499 14, doi:10.1029/2018GB005969, 2019.
- 500 Martens, J., Wild, B., Muschitiello, F., O'Regan, M., Jakobsson, M., Semiletov, I., Dudarev, O. V. and Gustafsson,
 501 Ö.: Remobilization of dormant carbon from Siberian-Arctic permafrost during three past warming events, *Sci.*
 502 *Adv.*, 6(42), 6546–6562, doi:10.1126/sciadv.abb6546, 2020a.
- 503 Martens, J., Romankevich, E., Semiletov, I., Wild, B., Dongen, B. van, Vonk, J., Tesi, T., Shakhova, N., Dudarev,
 504 O. V., Kosmach, D., Vetrov, A., Lobkovsky, L., Belyaev, N., Macdonald, R., Pieńkowski, A. J., Eglinton, T. I.,
 505 Haghipour, N., Dahle, S., Carroll, M. L., Åström, E. K. L., Grebmeier, J. M., Cooper, L. W., Possnert, G. and
 506 Gustafsson, Ö.: The Circum-Arctic Sediment Carbon Database — CASCADE, version 1.0, Bolin Centre Database,
 507 , doi:10.17043/cascade, 2020b.
- 508 Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh,
 509 A., Melbourne-Thomas, J., Muelbert, M. M. C., Ottersen, G., Pritchard, H. and Schuur, E. A. G.: Polar Regions, in
 510 IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by H.-O. Pörtner, D. C. Roberts,

- 511 V. Masson-Delmotte, M. T. P. Zhai, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold,
512 B. Rama, and N. M. Weyer., 2019.
- 513 Mueller-Lupp, T., Bauch, H. A., Erlenkeuser, H., Hefter, J., Kassens, H. and Thiede, J.: Changes in the deposition
514 of terrestrial organic matter on the Laptev Sea shelf during the Holocene: evidence from stable carbon isotopes,
515 *Int. J. Earth Sci.*, 89(3), 563–568, doi:10.1007/s005310000128, 2000.
- 516 Muschitiello, F., O'Regan, M., Martens, J., West, G., Gustafsson, Ö. and Jakobsson, M.: A new 30 000-year
517 chronology for rapidly deposited sediments on the Lomonosov Ridge using bulk radiocarbon dating and
518 probabilistic stratigraphic alignment, *Geochronology*, 2(1), 81–91, doi:10.5194/gchron-2-81-2020, 2020.
- 519 Naidu, A. S., Scalan, R. S., Feder, H. M., Goering, J. J., Hameedi, M. J., Parker, P. L., Behrens, E. W., Caughey,
520 M. E. and Jewett, S. C.: Stable organic carbon isotopes in sediments of the north Bering-south Chukchi seas,
521 *Alaskan-Soviet Arctic Shelf, Cont. Shelf Res.*, 13(5–6), 669–691, doi:10.1016/0278-4343(93)90099-J, 1993.
- 522 Naidu, A. S., Cooper, L. W., Finney, B. P., Macdonald, R. W., Alexander, C. and Semiletov, I. P.: Organic carbon
523 isotope ratio ($\delta^{13}\text{C}$) of Arctic Amerasian Continental shelf sediments, *Int. J. Earth Sci.*, 89(3), 522–532,
524 doi:10.1007/s005310000121, 2000.
- 525 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling,
526 B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K.,
527 Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J. and Zou, D.: Northern Hemisphere
528 permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Science Rev.*, 193, 299–316,
529 doi:10.1016/j.earscirev.2019.04.023, 2019.
- 530 Romankevich, E. A.: *Geochemistry of Organic Matter in the Ocean*, Springer Berlin Heidelberg, Berlin,
531 Heidelberg, doi:10.1007/978-3-642-49964-7., 1984.
- 532 Semiletov, I. and Gustafsson, Ö.: East Siberian shelf study alleviates scarcity of observations, *Eos (Washington*
533 *DC)*, 90(17), 145–146, doi:10.1029/2009eo170001, 2009.
- 534 Semiletov, I., Dudarev, O., Luchin, V., Charkin, A., Shin, K. H. and Tanaka, N.: The East Siberian Sea as a
535 transition zone between Pacific-derived waters and Arctic shelf waters, *Geophys. Res. Lett.*, 32(10), 1–5,
536 doi:10.1029/2005GL022490, 2005.
- 537 Semiletov, I., Pipko, I., Gustafsson, Ö., Anderson, L. G., Sergienko, V., Pugach, S., Dudarev, O., Charkin, A.,
538 Gukov, A., Bröder, L., Andersson, A., Spivak, E. and Shakhova, N.: Acidification of East Siberian Arctic Shelf
539 waters through addition of freshwater and terrestrial carbon, *Nat. Geosci.*, 9(5), 361–365, doi:10.1038/NEGO2695,
540 2016.
- 541 Sparkes, R. B., Dořul Selver, A., Bischoff, J., Talbot, H. M., Gustafsson, Semiletov, I. P., Dudarev, O. V and Van
542 Dongen, B. E.: GDGT distributions on the East Siberian Arctic Shelf: Implications for organic carbon export, burial
543 and degradation, *Biogeosciences*, 12(12), 3753–3768, doi:10.5194/bg-12-3753-2015, 2015.
- 544 Stein, R. and Macdonald, R. W.: *The Organic Carbon Cycle in the Arctic Ocean*, edited by R. Stein and R. W.
545 MacDonald, Springer Berlin Heidelberg, Berlin, Heidelberg, doi:10.1007/978-3-642-18912-8., 2004.

- 546 Stein, R., Grobe, H. and Wahsner, M.: Organic carbon, carbonate, and clay mineral distributions in eastern central
547 Arctic Ocean surface sediments, *Mar. Geol.*, 119(3–4), 269–285, doi:10.1016/0025-3227(94)90185-6, 1994.
- 548 Stein, R., Macdonald, R. W., Naidu, A. S., Yunker, M. B., Gobeil, C., Cooper, L. W., Grebmeier, J. M., Whitley,
549 T. E., Hameedi, M. J., Petrova, V. I., Batova, G. I., Zinchenko, A. G., Kursheva, A. V., Narkevskiy, E. V., Fahl, K.,
550 Vetrov, A., Romankevich, E. A., Birgel, D., Schubert, C., Harvey, H. R. and Weiel, D.: Organic Carbon in Arctic
551 Ocean Sediments: Sources, Variability, Burial, and Paleoenvironmental Significance, in *The Organic Carbon Cycle*
552 *in the Arctic Ocean*, edited by R. Stein and R. W. MacDonald, pp. 169–314, Springer Berlin Heidelberg, Berlin,
553 Heidelberg., 2004.
- 554 Stenström, K. E., Skog, G., Georgiadou, E., Genberg, J. and Johansson, A.: A guide to radiocarbon units and
555 calculations, Lund University, Nuclear Physics., 2011.
- 556 Stuiver, M. and Polach, H. A.: Discussion Reporting of ^{14}C Data, *Radiocarbon*, 19(03), 355–363,
557 doi:10.1017/S0033822200003672, 1977.
- 558 Tesi, T., Semiletov, I., Hugelius, G., Dudarev, O., Kuhry, P. and Gustafsson, Ö.: Composition and fate of
559 terrigenous organic matter along the Arctic land–ocean continuum in East Siberia: Insights from biomarkers and
560 carbon isotopes, *Geochim. Cosmochim. Acta*, 133, 235–256, doi:10.1016/j.gca.2014.02.045, 2014.
- 561 Tesi, T., Muschitiello, F., Smittenberg, R. H., Jakobsson, M., Vonk, J. E., Hill, P., Andersson, A., Kirchner, N.,
562 Noormets, R., Dudarev, O. V., Semiletov, I. P. and Gustafsson, Ö.: Massive remobilization of permafrost carbon
563 during post-glacial warming, *Nat. Commun.*, 7, 13653, doi:10.1038/ncomms13653, 2016a.
- 564 Tesi, T., Semiletov, I., Dudarev, O., Andersson, A. and Gustafsson, Ö.: Matrix association effects on hydrodynamic
565 sorting and degradation of terrestrial organic matter during cross-shelf transport in the Laptev and East Siberian
566 shelf seas, *J. Geophys. Res. Biogeosciences*, 121(3), 731–752, doi:10.1002/2015JG003067, 2016b.
- 567 Vetrov, A. A. and Romankevich, E. A.: *Carbon Cycle in the Russian Arctic Seas*, Springer Berlin Heidelberg,
568 Berlin, Heidelberg., 2004.
- 569 Vonk, J., Sánchez-García, L., van Dongen, B. E., Alling, V., Kosmach, D., Charkin, A., Semiletov, I. P., Dudarev,
570 O. V., Shakhova, N., Roos, P., Eglinton, T. I., Andersson, A., Gustafsson, Ö. and Gustafsson, O.: Activation of old
571 carbon by erosion of coastal and subsea permafrost in Arctic Siberia, *Nature*, 489(7414), 137–140,
572 doi:10.1038/nature11392, 2012.
- 573 Vonk, J. E. and Gustafsson, Ö.: Permafrost-carbon complexities, *Nat. Geosci.*, 6(9), 675–676,
574 doi:10.1038/ngeo1937, 2013.
- 575 Vonk, J. E., Sánchez-García, L., Semiletov, I., Dudarev, O., Eglinton, T., Andersson, A. and Gustafsson, Ö.:
576 Molecular and radiocarbon constraints on sources and degradation of terrestrial organic carbon along the Kolyma
577 paleoriver transect, East Siberian Sea, *Biogeosciences*, 7(10), 3153–3166 [online] Available from:
578 <http://www.biogeosciences.net/7/3153/2010/>, 2010.
- 579 Vonk, J. E., Semiletov, I. P., Dudarev, O. V., Eglinton, T. I., Andersson, A., Shakhova, N., Charkin, A., Heim, B.
580 and Gustafsson, Ö.: Preferential burial of permafrost-derived organic carbon in Siberian-Arctic shelf waters, *J.*

581 Geophys. Res. Ocean., 119(12), 8410–8421 [online] Available from: <http://doi.wiley.com/10.1002/2014JC010261>,
582 2014.

583 Xiao, X., Fahl, K., Müller, J. and Stein, R.: Sea-ice distribution in the modern Arctic Ocean: Biomarker records
584 from trans-Arctic Ocean surface sediments, *Geochim. Cosmochim. Acta*, 155, 16–29,
585 doi:10.1016/j.gca.2015.01.029, 2015.

586 Yunker, M. B., Belicka, L. L., Harvey, H. R. and Macdonald, R. W.: Tracing the inputs and fate of marine and
587 terrigenous organic matter in Arctic Ocean sediments: A multivariate analysis of lipid biomarkers, *Deep. Res. Part*
588 *II Top. Stud. Oceanogr.*, 52(24–26), 3478–3508, doi:10.1016/j.dsr2.2005.09.008, 2005.

589
590





Figure 1: Overview map of the Arctic Ocean compartments defined and used in CASCADE, with the permafrost distribution based on numerical modelling (Obu et al., 2019), rates of coastal erosion (Lantuit et al., 2012); and the latest IBCAO v4 bathymetry (Jakobsson et al., 2020). Black lines delineate the extent of the Arctic Ocean shelf seas and each respective watershed on land.



596
 597 **Figure 2:** CASCADe data location for OC concentrations (panel A), carbon isotopes $\delta^{13}\text{C}$ (panel B) and $\Delta^{14}\text{C}$
 598 (panel C) marked as red dots, with interpolated fields as indicated by the inserted color scale and as described in
 599 the main text.

600 **Table 1: CASCADE data availability per circum-Arctic shelf sea and for the interior basin**

Shelf area		Area	OC	TN	$\delta^{13}\text{C}$	New	$\Delta^{14}\text{C}$	New	Alk1 ¹	Alk2 ²	Acid ³	Lignin ⁴	New
		10 ³ km ²	<i>n</i>		<i>n</i>	$\delta^{13}\text{C}$	<i>n</i>	$\Delta^{14}\text{C}$	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	lignin
					<i>n</i>		<i>n</i>						<i>n</i>
1.	Barents Sea ⁵	1626	1092	353	236	48	33	33	0	13	0	0	0
2.	Kara Sea	942	637	201	262	22	29	22	2	90	2	0	0
3.	Laptev Sea	505	312	110	214	8	42	14	33	46	31	36	19
4.	East Siberian Sea	1000	259	217	187	17	71	16	28	13	10	68	40
5.	Chukchi Sea	639	1084	950	256	9	12	10	67	14	58	3	0
6.	Beaufort Sea	183	247	122	219	5	32	3	5	1	2	11	0
7.	Canadian Arctic Archipelago ⁶	1171	92	87	55	29	22	19	0	0	0	9	0
8.	Central Arctic Ocean ⁷	4500	529	282	130	15	27	10	29	36	28	18	5
Total		10566	4252	2322	1559	153	268	127	164	213	131	145	64

601 ^{1.} Alk1: HMW *n*-alkanes $\Sigma\text{C}_{21}\text{-C}_{31}$
602 ^{2.} Alk2: HMW *n*-alkanes $\text{C}_{27}+\text{C}_{29}+\text{C}_{31}$
603 ^{3.} Acid: HMW *n*-alkanoic acids $\Sigma\text{C}_{20}\text{-C}_{30}$
604 ^{4.} Lignin: lignin phenols syringyl, vanillyl and cinnamyl
605 ^{5.} incl. White sea and shelf northwest of Svalbard
606 ^{6.} incl. shelf northeast of Greenland
607 ^{7.} incl. continental slope, rise and abyssal plain

608 **Table 2: Parameter description and name of the respective columns in the CASCADE data sheet**

Parameters	Description	Column name
CASCADE entry ID	Serial number	ID
<i>Georeference and sampling information</i>		
Sample code	Expedition station ID	STATION
Latitude	Decimal latitude according to WGS1984	LAT
Longitude	Decimal longitude according to WGS1984	LON
Upper sample depth (cm)	Sample depth (for surface sediments only)	UPPERDEPTH
Lower sample depth (cm)	Sample depth (for surface sediments only)	LOWERDEPTH
Median sample depth (cm)	Median sample depth (for core samples only)	COREDEPTH
Water depth (m b.s.l.)	Water depth of sampling according to shipboard measurement	WATERDEPTH
Water depth based on IBCAO (m b.s.l.)	Water depth according to IBCAOv4	IBCAODEPTH
Expedition or vessel name	Vessel name, expedition name, cruise number	EXPEDITION
Sampling year	Year when the sample was taken as reported in literature	YEAR
<i>Carbon and Nitrogen (CN) data</i>		
OC (%)	Total OC concentration of the bulk sediment; carbonate removal assumed	OC
TN (%)	Total N concentration of the bulk sediment	TN
OC/TN	OC/TN ratio (gravimetric); published values or calculated	OC_TN
<i>Carbon isotopes</i>		
$\delta^{13}\text{C}$ (‰ VPDB)	$\delta^{13}\text{C}$ -OC; carbonate removal assumed	d13C
$\Delta^{14}\text{C}$ (‰)	$\Delta^{14}\text{C}$ -OC corrected for age; carbonate removal assumed	D14C
<i>Biomarkers</i>		
<i>n</i> -alkanes C ₂₁₋₃₁ (μg g ⁻¹ OC)	OC-normalized concentration of HMW <i>n</i> -alkanes	HMWALK
<i>n</i> -alkanes C _{27,29,31} (μg g ⁻¹ OC)	OC-normalized concentration of specific HMW <i>n</i> -alkanes	HMWALK_SPEC
<i>n</i> -alkanoic acids C ₂₀₋₃₀ (μg g ⁻¹ OC)	OC-normalized concentration of HMW <i>n</i> -alkanoic acids	HMWACID
lignin phenols (mg g ⁻¹ OC)	OC-normalized concentration of syringyl, vanillyl, cinnamyl	LIGNIN
<i>Quality parameter and meta information</i>		
Sediment sampler	Method of sediment sampling	SAMPLER
Sample storage	0 unknown, 1 frozen, 2 refrigerated, 3 dried onboard	STORAGE
CN measurement	Description of the method of analysis of the OC and TN data	CN_METHOD
$\delta^{13}\text{C}$ measurement	Description of the method of analysis of $\delta^{13}\text{C}$ -OC	d13C_METHOD

AMS/ ¹⁴ C label	Laboratory number of the $\Delta^{14}\text{C}$ measurement	D14C_LABEL
<i>Citation of the data source</i>		
Citation of CN data	Full citation in ESSD style incl. info about publication format	CN_CITATION
Citation of $\delta^{13}\text{C}$ data	Authors, title, journal, volume, pages, doi, year	d13C_CITATION
Citation of $\Delta^{14}\text{C}$ data	Full citation in ESSD style incl. info about publication format	D14C_CITATION
Citation of biomarker data	Full citation in ESSD style incl. info about publication format	BM_CITATION