

Greenland liquid water discharge from 1958 through 2019

Kenneth D. Mankoff¹, Brice Noël⁵, Xavier Fettweis², Andreas P. Ahlstrøm¹, William Colgan¹, Ken Kondo³, Kirsty Langley⁴, Shin Sugiyama³, Dirk van As¹, and Robert S. Fausto¹

¹Department of Glaciology and Climate, Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

²Department of Geography, University of Liège, Belgium

³Institute of Low Temperature Science, Hokkaido University, Japan

⁴Asiaq-Greenland Survey, Nuuk, Greenland

⁵Institute for Marine and Atmospheric Research, Utrecht University, The Netherlands

Correspondence: Ken Mankoff (kdm@geus.dk)

Abstract. Greenland runoff, from ice mass loss and increasing rainfall, is increasing. That runoff then discharges and impacts the physical, chemical, and biological properties of the adjacent fjords. However, where and when the discharge occurs is not readily available in an open database. Here we provide datasets of high-resolution Greenland hydrologic outlets, basins, and streams, and a daily 1958 through 2019 time series of Greenland liquid water discharge for each outlet. The data include 24507 ice marginal outlets and upstream basins, and 29635 land coast outlets and upstream basins, derived from the 100 m ArcticDEM and 150 m BedMachine. At each outlet there are daily discharge data for 22644 days - ice sheet runoff routed subglacially to ice margin outlets, and land runoff routed to coast outlets - from two regional climate models (RCMs; MAR and RACMO). Our sensitivity study of how outlet location changes for every inland cell based on subglacial routing assumptions, shows that most inland cells where runoff occurs are not highly sensitive to those routing assumptions, and outflow location does not move far. We compare RCM results with 10 streams instrumented with gauges spanning four orders of magnitude of daily discharge. Results show that for daily discharge at individual basin scale the 95 % prediction interval generally falls within plus-or-minus a factor of five (half an order of magnitude, or +500%/-80%). Results from this study are available at doi:10.22008/promice/freshwater (Mankoff, 2020a) and code is available at <http://github.com/mankoff/freshwater> (Mankoff, 2020b).

1 Introduction

Over the past decades, liquid runoff from Greenland has increased (Mernild and Liston, 2012; Bamber et al., 2018; Trusel et al., 2018; Perner et al., 2019) contributing to mass decrease (Sasgen et al., 2020). When that runoff leaves the ice sheet and discharges into fjords and coastal seas, it influences a wide range of physical (Straneo et al., 2011; An et al., 2012; Mortensen et al., 2013; Bendtsen et al., 2015; Cowton et al., 2015; Mankoff et al., 2016; Fried et al., 2019; Cowton et al., 2019; Beckmann et al., 2019), chemical (Kanna et al., 2018; Balmonte et al., 2019), and biological (Kamenos et al., 2012; Kanna et al., 2018; Balmonte et al., 2019) systems (Catania et al., 2019). The scales of the impacts range from instantaneous at the ice-ocean boundary to decadal in the distal ocean (Gillard et al., 2016). The influence of freshwater on multiple domains and disciplines (Catania et al., 2019) is the reason several past studies have estimated runoff and discharge at various temporal and spatial

scales (e.g. Mernild et al. (2008, 2009, 2010a); Langen et al. (2015); Ahlstrøm et al. (2017); Citterio et al. (2017); van As et al. (2018); Bamber et al. (2018); Perner et al. (2019); Slater et al. (2019)).

To date no product provides discharge estimates at stream spatial resolution (~100 m), daily temporal resolution, for all of Greenland, covering a broad time span (1958 through 2019), from multiple regional climate models (RCMs), and with a simple database access software to support downstream users. Here we present these data. In the following description and methods, we document the inputs, assumptions, methodologies, and results we use to estimate Greenland discharge from 1958 through 2019. This product is available at doi:10.22008/promice/freshwater (Mankoff, 2020a).

Freshwater discharge from Greenland primarily takes three forms: solid ice from calving at marine terminating glaciers, submarine meltwater from ice-ocean boundary melting at marine terminating glaciers, and liquid runoff from melted inland surface ice, rain, and condensation. A recent paper by Mankoff et al. (2020) targets the solid ice discharge plus submarine melt budget by estimating the ice flow rate across gates 5 km upstream from all fast-flowing marine terminating glaciers in Greenland. Complementing that paper, this paper targets Greenland's point-source liquid water discharge budget by partitioning RCM runoff estimates to all ice margin and coastal outlets. The sum of these data and Mankoff et al. (2020) is an estimate of the majority of freshwater (in both liquid and solid form) volume flow rates into Greenland fjords. Those two terms comprise the bulk but not all freshwater - they exclude precipitation directly onto the fjord or ocean surface, and relatively minor contributions from evaporation and condensation, sea ice formation and melt, or subglacial basal melting.

2 Input and validation data

2.1 Static data

The static products (streams, outlets, and basins (Fig. 1)) are derived from an ice-sheet surface digital elevation model (DEM), an ice-sheet mask, the land surface DEM, and an ocean mask. For the surface DEM, we use ArcticDEM v7 100 m (Porter et al., 2018). Subglacial routing uses ArcticDEM and ice thickness from BedMachine v3 (Morlighem et al., 2017a, b). For the ice mask we use the Programme for Monitoring of the Greenland Ice Sheet (PROMICE) Ice Extent (Citterio and Ahlstrøm, 2013). For the ocean mask we use the Making Earth System Data Records for Use in Research Environments (MEaSUREs) Greenland Ice Mapping Project (GIMP) Land Ice and Ocean Classification Mask, Version 1 (Howat, 2017b; Howat et al., 2014).

2.2 RCM time series

The time series product (daily discharge) is derived from daily runoff estimates from RCM calculations over the land and ice areas of Greenland. We use the Modèle Atmosphérique Régional (MAR; Fettweis et al. (2017)) and the Regional Atmospheric Climate Model (RACMO; Noël et al. (2019)). Runoff, R , is defined by

$$R = ME + RA - RT - RF. \quad (1)$$

In Eq. 1, *ME* is melt, *RA* is rainfall, *RT* is retention, and *RF* is refreezing. In RACMO, retention occurs only when firn is present (not with bare ice). MAR does have a delay for bare ice runoff. Neither have a delay for land runoff. Both RCM results were provided regridded to the same 1 km grid using an offline statistical down-scaling technique based on local vertical runoff gradient applied to the sub-grid topography (Noël et al., 2016; Fettweis et al., 2020). MAR (v 3.11; Delhasse et al. (2019)) ran with 7.5 km resolution and ERA 6-hour forcing. RACMO (v 2.3p2; Noël et al. (2018)) ran with 5.5 km resolution and ERA-Interim 6-hour forcing. Runoff is assigned an uncertainty of $\pm 15\%$.

2.3 River discharge observations

We use 10 river discharge daily time series to validate the results of this work. The name, location, time coverage, and relevant data and scientific publications associated with each of these observational data are listed in Table 1.

Table 1. Table of observation locations, time spans, and associated references. Coordinates are decimal degree W and N.

Location	Lon	Lat	Time	Data	Publication	Fig(s).
Kiattuut Sermiat	45.33	61.21	2013	Hawkings et al. (2016a)	Hawkings et al. (2016b)	1 4 5 6 10
Kingigtorsuaq (Nuuk)	51.5801	64.1387	2008-2018	Langley (2020)		1 4 5 11
Kobbefjord (Nuuk)	51.3810	64.1336	2006-2017	Langley (2020)		1 4 5 14
Leverett Glacier	50.17	67.06	2009-2012	Tedstone et al. (2017)	Hawkings et al. (2015)	1 4 5 6 9
Oriartorfik (Nuuk)	51.4066	64.1707	2007-2018	Langley (2020)		1 4 5 12
Qaanaaq	69.3030	77.4753	2017-2018	Kondo and Sugiyama (2020)	Sugiyama et al. (2014)	1 4 5 6 17
Røde Elv (Disko)	53.4989	69.2534	2017	Langley (2020)		1 6 4 5 6 15
Teqinngalip (Nuuk)	51.5484	64.1586	2007-2018	Langley (2020)		1 4 5 13
Watson River	50.68	67.01	2006-2019	van As et al. (2018)	van As et al. (2018)	1 4 5 6 7 8
Zackenbergl	20.5628	74.4722	1996-2018	Langley (2020)		1 4 5 6 16

3 Methods

3.1 Terminology

We use the following terminology throughout the document:

- Runoff refers to the unmodified RCM data products - melted ice, rain, condensation, and other RCM outputs that are inputs to this work.
- Discharge refers to the runoff after it has been processed by this work - routed to and aggregated at the outlets. Depending on context, discharge may also refer to the observed stream discharge (Table 1).
- Basins refer to the 100 m x 100 m resolution basins derived from a combination of the ArcticDEM product and the mask.

- 70 – Mask refers to the surface classification at that 100 m x 100 m resolution and is one of ice, land, or ocean (also called fjord or water). When referring to the surface classification in the RCM, we explicitly state "RCM mask".
- MAR and RACMO refer to the RCMs, but when comparing discharge estimates between them or to observations, we use MAR and RACMO to refer to our discharge product derived from the MAR and RACMO RCM runoff variables, rather than repeatedly explicitly stating "discharged derived from [MAR|RACMO] runoff". The use should be clear from
- 75 context.
- The $\pm 95\%$ quantile range refers to data $< 95\%$, or $> 5\%$, and is therefore only 90% of the data.

3.2 Streams, outlets, and basins

Streams are calculated from the hydraulic head h which is the DEM surface for land surface routing, or the subglacial pressure head elevation for subglacial routing. h is defined as

$$80 \quad h = z_b + k \frac{\rho_i}{\rho_w} (z_s - z_b), \quad (2)$$

with z_b the ice-free land surface and basal topography, k the flotation fraction, ρ_i the density of ice (917 kg m^{-3}), ρ_w the density of water (1000 kg m^{-3}), and z_s the land surface for both ice free and ice covered surfaces.

Eq. 2 comes from Shreve (1972) where the hydropotential has units Pa, but here is divided by gravitational acceleration g times the density of water ρ_w to convert the units from Pa to m. We compute h and from that streams, outlets, basins, and

85 runoff for a range of subglacial pressures, implemented as a range of k values: 0.8, 0.9, and 1.0. We use these three scenarios to estimate sensitivity of the outlet location for all upstream cells, but otherwise only use results from the $k = 1.0$ scenario. Eq. 2 makes the assumption that when ice is present all water routes subglacially, meaning that water flows from the surface to the bed in the grid cell where it is generated. In reality, internal catchments and moulins likely drain waters to the bed within a few km of their source (Yang and Smith, 2016). The difference between some supraglacial flow and immediate subglacial flow is

90 not likely to impact results because discharge is reported only at the outlet locations.

We use the GRASS GIS software (Neteler et al., 2012; GRASS Development Team, 2018) and the `r.stream.extract` command configured for single-flow direction from eight neighbors (SFD-8) to calculate streams and outlets at the ice edge and coast. Streams are defined only if their upstream contributing area is above a threshold, so small basins may have outlets but no streams. The software fills all sinks so that all water flows to the domain edge. We then use the `r.stream.basins`

95 tool (Jasiewicz and Metz, 2011) to calculate basins upstream from each outlet. Basins $< 1 \text{ km}^2$ are absorbed into their largest neighbor and the associated outlets are dropped.

3.2.1 Outlet sensitivity

The three choices of k generate three scenarios of basins and outlets, and we use this to show sensitivity of every ice grid cell to these choices. After three k -scenarios, each cell has three possible outlets, where each outlet is an (x,y) coordinate. To show

100 results in a map view, we reduced these six properties (three 2D coordinates) to a single property. For every grid cell in the ice domain we compute the distance between each outlet and the other two (six becomes three), and then select the maximum (three becomes one). Fig. 2 displays the maximum distance - a worst-case scenario - of how far the outlet of every inland ice cell may move due to basal routing assumptions.

3.3 Discharge and RCM coverage

105 RCM runoff is summed over each basin for each day of RCM data, and assigned to each outlet for that day. This assumes routing between the runoff and the outlet is instantaneous, so all analyses done here include a seven-day smooth applied as in van As et al. (2017). The released data do not include any smoothing.

The alignment of the RCM and the basins do not always agree. Each 100 m x 100 m ArcticDEM pixel is classified as ice (Citterio and Ahlström, 2013), ocean (Howat, 2017b), or land (defined as neither ice nor ocean). However, the classification of
110 the mask cells and the 1 km² RCM domains do not always agree - for example, when a mask cell is classified as ice but the matching RCM cell is land. This disagreement occurs almost everywhere along the ice margin because the 1 km resolution RCM boundary and the 100 m mask boundary rarely perfectly align. The ice margin is where most runoff occurs due to the highest temperatures at the lowest ice elevations, so small changes in masks in these locations can introduce large changes in RCM outputs.

115 We adjust for this imprecise overlap and scale the RCM results to the basin area. Where the mask reports ice and a RCM reports land, the RCM land runoff fraction is discarded, and the RCM ice runoff fraction over this basin is adjusted for the uncovered basin cells (and vice versa for basin land and RCM ice). Small basins with no RCM coverage of the same type have no runoff.

Runoff adjustments using this method are underestimated for large basins with large inland high elevation regions with
120 low runoff, because this method fills in misaligned cells with each days average discharge, but the misalignment (missing runoff) occurs at the ice sheet edge where maximum runoff occurs. However, given that the basin is large, misalignment is proportionally small, and therefore errors are proportionally small. When misalignment is proportionally large (e.g. a basin is only 1 % covered by the same RCM classification), that implies a small basin with the covered region near the uncovered region, and the filling method therefore uses nearby data, and not distal high-elevation data, and there should not be a large
125 underestimate.

RCM inputs are also scaled by the projection area error between the EPSG:3413 map projection of the RCM and an approximation of the true earth spheroid. This error is up to 8 % for some grid cells, but ranges from - 6 % to + 8 % over Greenland and the cumulative error for the entire ice sheet is < 8 %.

3.4 Validation

130 We validate the modeled outlet discharge against the observations first in bulk and then individually. Bulk comparisons are done with scatter plots (Figs. 3 & 4), and modified Tukey plots comparing observations vs. differences (Fig. 5, based on Tukey mean-difference plots, also known as Bland-Altman plots (Altman and Bland, 1983; Martin Bland and Altman, 1986)). When

comparing modeled with observed discharge, we drop any days where observed discharge is zero or modeled discharge is less than $1 \text{ m}^3 \text{ day}^{-1}$.

135 We introduce the graphics here as part of the methods to reduce replication in figure captions - we show 10 nearly identical graphics (Figs. 7 and 9 through 17) for 10 different observation locations, and each graphic uses the same template of six panels.

For each figure (Figs. 7, 9 to 17), the top panel (a) shows a satellite basemap with the land portion of the basin of interest (if it exists) outlined in dark green, the streams within that basin in light green, the basin outlet as an orange filled diamond, and the stream gauge location as an orange unfilled diamond. Ice basin(s) that drain to the outlet are outlined in thick dark blue if they exist, and all other ice basins in thin dark blue. The RCM ice domain is in light blue, and RCM land domain not shown, but is outside the light blue ice domain (not including the water). The scale of each map varies, but the basins lines (green and dark blue) are discretized at 100 m resolution, and the RCM grid cells (light blue) are at 1 km resolution.

Panel b shows an example time series - whatever data are available for the last calendar year of the observations.

145 Panel c shows a scatter plot of observations vs. RCM-derived discharge. This is the same data shown in Fig. 3, but subset to just the basin of interest. Color encodes day-of-year, and a kernel density estimation (KDE) of the discharge values highlights where most points occur - not necessarily visible without the KDE because the points overlap (total number of plotted points is printed on the graphic near "n:"). The r^2 correlation coefficient for each RCM-derived discharge is displayed. The gray band shows the 95 % prediction interval, and the three solid lines mark the 1:1, 1:5, and 5:1 ratios.

150 Panel d shows observations vs. difference. This is the same data shown in Fig. 5, but subset to just the basin of interest. Color denotes sample density (similar to the KDE in panel c). The horizontal lines mark the mean, 0.05, and 0.95 quantile of the scale difference between the RCM and the observations. Scale difference means that a value of 1 (or 10^0) is agreement between observations and the RCM, and a value of 2 or 0.5 is a factor of 2 or a +100/-50 % disagreement. The horizontal split marks the bottom 1/3rd and top 2/3rds quantiles of discharge.

155 4 Product evaluation and assessment

Results of this work include 1) ice-margin terminating streams, outlets, and basins, 2) coast-terminating streams, outlets, and basins, 3) discharge at the ice-marginal outlets from ice runoff and 4) discharge at the coastal outlets from land runoff. Discharge products are provided from both the MAR and RACMO RCMs. We note that our subglacial streams represent where the model routes the water, and does not indicate actual streams, unlike the land streams that do appear near actual streams when compared to satellite imagery. Even so, these streams routed using simple subglacial theory show remarkable alignment with ice surface streams and lakes visible in satellite imagery. This may support the theory that basal topography exerts a strong control on supraglacial hydrology (Lampkin and VanderBerg, 2011; Sergienko, 2013; Crozier et al., 2018), or may indicate a poorly represented and smooth bed in BedMachine, and therefore Eq. 2 is effectively applying surface routing in these locations.

165 Of the 361,950 km² of basin land cells, the RCMs cover 339,749 km² (~94 %) with their land grid cells, and 22,201 km² (~6
%) of basin grid cells are filled in with our coverage algorithm (Sect. 3.3; the RCMs have these as ice or ocean). Alternatively,
51,532 km² of RCM land are discarded because the basins classify part or all of these cells as ice or ocean. Of the 1,781,816
km² of basin ice cells, the RCMs cover 1,760,912 km² (~99 %) with their ice cells, and 20,904 km² (~1 %) of basin grid cells
170 are filled in (the RCMs have these as land or ocean). Alternatively, 21,793 km² of RCM ice are discarded, because the basins
classify part or all of these cells as land or ice (Table and data available in Supplemental Online Material).

Our coverage correction (Sect. 3.3) adjusts RCM ice runoff values by ~3 %. Discarding RCM ice runoff that does not match
the underlying mask ice cells results in a 5 % reduction in discharge. However, applying our coverage algorithm to adjust RCM
inputs for regions where basins have ice but the RCMs do not results in an 8 % increase from the reduced discharge (net gain
of ~3 %). A similar adjustment occurs for RCM land runoff.

175 **4.1 Comparison with previous similar work**

Our static products - streams, outlets, and basins - have been previously estimated. Lewis and Smith (2009) identified 293
distinct hydrologic ice basins and provided a data set of ice basins and ice margin outlets. Our work, a decade later, has ~2
orders of magnitude more basins and outlets because of the higher resolution of the input data, and includes additional data. We
provide ice basins, ice margin outlets, ice streams with metadata, land basins, coastal outlets, and land streams with metadata.
180 Lewis and Smith (2009) generated basins from a 5 km DEM, compared to the 100 m DEM used here. Routing with a 5 km
DEM that does not capture small-scale topography is likely to cause some basins and outlets to drain into an incorrect fjord -
we find that some land basins delineated with even the 150 m BedMachine land surface may drain into the incorrect fjord, but
we did not find similar errors with the 100 m ArcticDEM product used in this work.

Our time-series product - discharge, also has existing similar products. The most recent of these is from Bamber et al. (2018)
185 who provide a data product at lower spatial resolution (5 km), lower temporal resolution (monthly), and only coastal discharge,
not coastal basins, ice basins, or ice margin outlets and discharge. However, Bamber et al. (2018) surpasses our product in
that spatial coverage includes a larger portion of the Arctic including Iceland, Svalbard, and Arctic Canada. Furthermore, by
providing data at 5 km spatial and monthly temporal resolution, Bamber et al. (2018) implements the main strategy suggested
here to increase the signal-to-noise ratio of the data - averaging discharge in space or time (see Sect. 4.3.5).

190 We show both the geospatial and temporal differences between this product and the Bamber et al. (2018) for an example
location - Disko Island (Fig. 6). Spatially our product allows assessment of discharge at interior locations, necessary when
comparing with observations that are not at the coast (for example, the Leverett Glacier observations (Fig. 9)). Temporally, the
MAR and RACMO runoff summed over all of Disko Island and to monthly resolution is similar to the monthly Disko discharge
of Bamber et al. (2018), but the daily resolution shows increased variability and individual discharge events (from warm days
195 or rain) not seen in the monthly view.

A similar GIS workflow was presented by Pitcher et al. (2016) only focusing on the discharge uncertainty from basal routing
assumptions (the k parameter in Eq 2). We find these differences to be smaller than the differences between RCMs or between
RCM and observations (see Sect. 4.3).

4.2 Validation against observations

200 Here we compare our results to all publicly accessible observations we could find, or willing to become open and publicly accessible as part of this work (Table 1).

This validation compares discharge derived from RCM runoff estimated far inland on the ice sheet. That runoff is both spatially and temporally disconnected from the stream discharge observations used here. Disagreement is expected and does not indicate any specific issues in the RCMs, but are instead likely due to our routing algorithm (Sect. 3.3).

205 Below we discuss first the validation for all points, then the individual outlets. For the individual outlets we begin by focusing on the "problematic" results in order of severity: Watson River (Figs. 7 & 8), Leverett Glacier (Fig. 9), and Kiattuut Sermiat (Fig. 10), and show that for two of these three, simple solutions are available, although manual intervention is needed to detect the issue and then adjust results.

4.2.1 Bulk validation

210 A comparison of every day of observational data with discharge > 0 (15,778 days) and the two RCMs (Fig. 3) shows good agreement with r^2 of 0.45 and 0.88 for discharge derived from MAR and RACMO runoff respectively (hereafter "MAR" and "RACMO"). For RACMO this is within a factor of five spanning four orders of magnitude, although both RCMs report only ~50 % of the observed discharge for the largest volumes at the Watson River outlet (Fig. 7). The reason for the disagreement at the Watson River outlet is discussed in detail in Sect. 4.2.2.

215 The four near-Nuuk GEM basins (Table 1, Sect. 4.2.5) have ice basins but either no or limited coverage in the RCMs. When excluding these basins from the comparison the r^2 agreement changes to 0.59 and 0.78 for MAR and RACMO respectively and the 95 % prediction interval is significantly smaller for MAR (red band in Fig. 3). The largest disagreements throughout this work comes from these small basins with no RCM coverage. These disagreements are therefore indicative of differences between the land/ice classification mask used by the RCMs compared with the basin masks used here, not necessarily the
220 ability of the models to simulate melting ice or local weather.

Fig. 4 shows a similar view as Fig. 3, but here each observational data set and associated daily discharge is summed by year for all and only the days in that year that observations exist (hence units m^3 and not $\text{m}^3 \text{ yr}^{-1}$; for example the single "Ks" and "R" means is only one calendar year with some observations at the Kiattuut Sermiat and Røde Elv outlets, respectively). Here it is more clear that the Watson River outlet (Sect. 4.2.2) reports ~50 % of the observed discharge, the Kiattuut Sermiat
225 outlet (Sect. 4.2.4) over-estimates discharge, and the remainder fall within the factor-of-two lines, except for low discharge at Kingigtorsuaq in the MAR RCM where the RCMs do not cover that small glacier (Sect. 4.2.5).

Because discharge spans a wide range (~4 orders of magnitude, Fig. 3), a high correlation (r^2 of 0.88, Fig. 3) may be due primarily to the range which is larger than the error (Altman and Bland, 1983; Martin Bland and Altman, 1986). Fig. 5 compensates for this and more clearly shows bias and the range of errors. This graphic again excludes the four near-Nuuk GEM
230 basins. From Fig. 5, the top 2/3rds of observed discharge has modeled discharge under-estimated by a scale of 0.78 (MAR) and 0.73 (RACMO), and ± 95 % quantile of 0.30 to 2.06. The top 2/3rds of discharge spans ~2 orders of magnitude (width

of horizontal line, from $\sim 10^1$ to $\sim 10^3$ $\text{m}^3 \text{s}^{-1}$), and has a ± 95 % quantile uncertainty of $\sim \pm 0.5$ order of magnitude, or half of the range of the data. Put differently, days with high observed discharge may have modeled discharge within ± 0.5 order of magnitude, or plus-or-minus a factor of five, or $+500/-80$ %. The modeled discharge is not likely to move farther than this from the observations, and high discharge remains high.

The bottom third of discharge is where the largest disagreement occurs. The mean model values are near the observed - discharge is scaled by 0.69 for MAR (~ 31 % low) and 1.08 for RACMO (~ 8 % high), but the ± 95 % quantile range is large. Although large uncertainties for low discharge may not seem to matter for some uses (e.g. estimates of total discharge from Greenland, which is dominated by the largest quantities of discharge), it may matter for other uses. The bottom 1/3 quantile of observed discharge spans 3 orders of magnitude (10^{-2} to $\sim 10^1$) but the uncertainty spans ~ 4 and ~ 2 orders of magnitude for MAR and RACMO respectively ($\sim 10^{-3}$ to $\sim 2.2 \times 10^1$ MAR; $\sim 10^{-1}$ to 2.2×10^1 RACMO).

4.2.2 Watson River

The Watson River discharge basin area is 1882 km^2 , of which 521 km^2 (28 %) are land and 1361 km^2 (72 %) are ice (Fig 7a). The partial (last calendar year) discharge time series shows MAR and RACMO agree well with each other, but have a maximum of $500 \text{ m}^3 \text{ s}^{-1}$ while observations are up to 4x more (Fig. 7b). Low discharge (both early and late season) is over-estimated and high discharge is under-estimated, approximately equal for both RCMs (Fig. 7c). The low discharge over-estimate ranges from a mean multiple of 1.64 (MAR) and 1.55 (RACMO) to a $+95$ % quantile scale ~ 70 (MAR) and ~ 50 (RACMO). The high-discharge under-estimate has a mean multiple of 0.5 for both MAR and RACMO, and a ± 95 quantile range of between 0.23 to 1.06.

The Watson River discharge presented here is approximately half of the van As et al. (2018) discharge for high discharge. The large underestimate for high discharge may be due to either errors in the basin delineation used in this study, errors in the stage-discharge relationship used by van As et al. (2018), errors in the RCM runoff estimates, or a combination of the above three. All three of these error sources increase with high discharge (and associated melt): Basin delineation becomes less certain with inland distance from the ice sheet margin. The river stage-discharge conversion becomes less certain at high stage levels. Runoff calculations become less certain from a snow surface than an ice surface, because of e.g. snow density, subsurface refreezing, and surface darkening.

The complexity of estimating the area of the Watson River catchment is described by Monteban et al. (2020), who note that previous studies have used values ranging from 6131 km^2 (Mernild et al., 2010b) to 12547 km^2 van As et al. (2012). Our basin is smaller than the basin used in van As et al. (2018) and similar to Mernild et al. (2018) who attributed the difference between their modeled outflow and observations from van As et al. (2017) to their decision to use surface rather than subglacial routing, and applied a correction term. We find that our basin does not include a separate basin here to the south that is part of the Watson River ice basin in van As et al. (2018) (from Lindbäck et al. (2015) and Lindbäck et al. (2014)). We are able to recreate the van As et al. (2018) basin but only when using the Lindbäck et al. (2014) bed and the Bamber et al. (2013) surface. When using one or zero of those and any combination of BedMachine v2 (Morlighem et al., 2014), BedMachine v3, or ArcticDEM surface elevations and BedMachine v2 or v3 bed elevations, and any range of k values, we are unable to match

the Lindbäck et al. (2015) basin. Instead all our basins resemble the basin shown in Fig 7. To solve this, we manually select two large ice basins to the south of the Watson River ice basin. Modeled and observed discharge agree after including these two basins (Fig. 8), suggesting basin delineation, not stage-discharge or RCM runoff is the primary cause for this disagreement. Furthermore, it is the additional width at lower elevation from the larger basin, not the increased inland high-elevation area, that likely contributes the runoff needed to match the observations, because 85 % of all surface runoff occurs below 1350 m, and almost all below 1850 van As et al. (2017).

There is no way to predict the disagreement between our and observed discharge. The observations are needed to highlight the disagreement. It is also not clear what to do to reduce the disagreement, without the previous efforts by Lindbäck et al. (2015) and Lindbäck et al. (2014). Basin delineation is discussed in more detail in the Uncertainty section (Sect. 4.3.2). The other two "problematic" areas can be detected and improved without observational support.

4.2.3 Leverett Glacier

The Leverett Glacier basin area is 1361 km² and 100 % ice (Fig 9a). The partial (last calendar year) discharge time series shows MAR and RACMO agree well with each other and with the observations (Fig. 9b), with no seasonal dependence (Fig 9c). The 95 % prediction interval for MAR is generally within the 1:5 and 5:1 bands, with a larger spread for RACMO (Fig 9c). High model discharge is 3 % higher than observed in MAR and 25 % higher than observed in RACMO, and the ± 95 quantile range is between 0.74 and 1.62 (MAR) and 0.82 and 2.02 (RACMO). Low model discharge is also centered near the observations, but as always larger errors exist for low discharge (Fig 9d).

This basin is problematic because the basin feeding the outlet is small (< 5 km²), but even without the observational record satellite imagery shows a large river discharging from the ice sheet here. Meanwhile, a large (100s of km²) ice basin does discharge just a few 100 m away, but not upstream of this gauge location. We therefore adjust the gauge location onto the ice so that our database access software selects what appears to be the correct basin given the size of the stream in the satellite imagery (Fig. 9).

The plots shown here use the adjusted gauge location and modeled discharge appears to match the observed discharge. When plotting (not shown) the modeled discharge for the outlet just upstream of the true gauge location, results are clearly incorrect. This issue - small basins at the margin and incorrect outlet location - is persistent throughout this product and discussed in more detail in Sect. 4.3.2.

The Leverett Glacier basin is a subset of the Watson River outlet basin (Sect. 4.2.2). The strong agreement here supports our claim that the Watson River disagreement is not from the RCM runoff or the stage-discharge relationship, but more likely due to basin area. The correct Watson River basin should include some basins outside of the Leverett Glacier basin that still drain to the Watson River outlet gauge location.

4.2.4 Kiattuut Sermiat

The Kiattuut Sermiat discharge basin area is 693 km², of which 391 km² (56 %) are land and 302 km² (44 %) are ice. The basin area is incorrectly large because the land basin reported and shown includes the entire basin that contains the discharge point,

of which some is downstream (Fig 10a). However, only ~25 % of runoff comes from the land, and only a small portion of the
300 land basin is downstream of the gauge location, so this is not enough to explain the discharge vs. observation disagreement.
The partial (last calendar year) discharge time series shows MAR and RACMO agree well with each other, but are significantly
higher than the observations (Fig. 10b). Both low and high discharge are over-estimated, but the 95% quantile range are within
a factor of five (Fig 10c), with a mean scale factor between 1.71 (RACMO bottom 1/3rd of discharge) to 2.47 (MAR high
2/3rds discharge)

305 The Kiattuut Sermiat gauge is in a problematic location in terms of determining the actual (non-theoretical) upstream con-
tributing area. Similar to the Leverett Glacier gauge location, the issues here can be estimated independent of observational
data. Specifically, it is not clear if this stream includes water from the larger glacier to the east and ENE that feeds this glacier
(Fig. 10a) - in our delineation it does not. Furthermore, several glaciers to the NNE and detached from the glacier near the
stream gauge appear to drain into a lake that then drains under the glacier with the stream gauge. This latter issue is observ-
310 able in any cloud-free satellite imagery (for example Google Earth) and does not need the basin delineations provided here
to highlight the complexities of this field site. Nonetheless, RCM discharge estimates are only slightly more than double the
observations.

The Kiattuut Sermiat gauge location may have been selected in part due to its accessibility - it is walking distance from the
Narsarsuaq airport. The data may also suit their intended purpose well and there are likely many results that can be derived
315 independent of the area or location of the upstream source water. However, if the location or area of the upstream contributions
are important, then gauge location should balance ease of access and maintenance with the ease with which the data can be
interpreted in the broader environment.

4.2.5 GEM observations near Nuuk

Four Greenland Ecosystem Monitoring Programme (GEM) stream gauges are located near Nuuk with similar basin properties.
320 All are small (7.56 to 37.52 km²), and 10 % to 25 % ice in the basin mask, but two of the four (Kingigtorsuaq (Fig. 11) and
Oriartorfik (Fig. 12)) contain small glaciers contributing to observed discharge but no RCM ice cells cover those glaciers, and
the remaining two (Teqinngalip (Fig. 13) and Kobbefjord (Fig. 14)) have several small glaciers, but only one per basin has
RCM ice coverage.

All four of these basins show some weak agreement. The maximum r^2 is 0.47 (Fig. 13c) and the minimum is 0.11 (Fig
325 11c), but we note that the worst agreement comes from a basin with no glaciers in the RCM domain, and that in all cases the
mean high discharge agrees well, suggesting high discharge in these small basins with few small glaciers may be due to rain
(captured in the RCMs) rather than warm days and melted ice. These agreements exist even though our modeled discharge
comes from the RCMs that are focused on and validated against the large ice Greenland ice sheet.

4.2.6 Remaining observations

330 Three additional stream gauges remain: Røde Elv, Zackenberg, and Qaanaaq.

The Røde Elv basin is situated at the southern edge of Disko Island (Fig. 6). It has an area of 100 km², of which 72 km² are land and 28 km² are ice (Fig 15a). The partial (last calendar year) discharge time series shows MAR, RACMO, and the observations all in approximately the same range but with high variability (Fig. 15b). Of the few samples here (n = 98), most are within the factor-of-five bands for MAR and a few more are outside the bands for RACMO (Fig. 15c). Mean discharge offset ranges from a scale factor of 0.86 (RACMO low) to 1.93 (MAR low), with high discharge estimates slightly closer to observations - a 48 % and 77 % overestimate for MAR and RACMO respectively (Fig. 15d).

The Zackenberg basin in NE Greenland has an area of 487 km², of which 378 km² (78 %) are land and 109 km² (22 %) are ice (Fig. 16a). The partial (last calendar year) discharge time series shows disagreement between MAR and RACMO that generally bound the observations (Fig. 16b). RACMO-derived discharge is consistently high for low discharge early in the year, but both discharge products fall mostly within the factor-of-five bands (Fig. 16c). For high discharge, mean modeled discharge is 9 % high (MAR) and 24 % low (RACMO), and has worst-case ± 95 quantile range low by a factor of 0.29 (Fig. 16d).

The Qaanaaq basin in NW Greenland has an area of 13.2 km², of which 2.2 km² (17 %) are land and 11 km² (83 %) are ice (Fig. 17a). The partial (last calendar year) discharge time series shows disagreement between MAR and RACMO that generally bound the observations (Fig 17b). Of the few samples (n = 82), MAR preferentially over-estimates and RACMO under-estimates discharge, but both generally within a factor of 5 (Fig 17c). Mean high discharge offset scale is 1.14 (MAR) and 0.36 (RACMO) from Fig. 17d.

4.3 Uncertainty

The volume of data generated here is such that manually examining all of it or editing it to remove artifacts or improve the data would be time and cost prohibitive. A similar warning is provided with the ArcticDEM data used here. However, any ArcticDEM issues interior to a basin do not impact results here that are aggregated by basin. Any ArcticDEM issues that cross a basin boundary should impact only the part of the basins it intersects.

Uncertainty from RCM inputs and observations are considered external to this work, although they are still discussed (Sects. 4.3.3 and 4.3.4). In this work, we introduce one new source of uncertainty - the routing model, which generates both temporal (runoff delay) and spatial (basin delineation) uncertainty.

4.3.1 Temporal uncertainty

The RCMs include a time lag between when water melts in the model and when it leaves a grid cell. RACMO retention occurs only when there is firn cover (no retention when bare ice melts); MAR includes a time delay of up to 10 days that is primarily a function of surface slope (Zuo and Oerlemans, 1996; Yang et al., 2019). However, neither model includes a subglacial system. Properly addressing time delays with runoff requires addressing storage and release of water across a variety of timescales in a variety of media: firn (e.g. Munneke et al. (2014); Vandecrux et al. (2019)), supraglacial streams and lakes (e.g. Zuo and Oerlemans (1996); Smith et al. (2015); Yang et al. (2019)), the subglacial system (e.g. Rennermalm et al. (2013)), possibly terrestrial streams and lakes (e.g. van As et al. (2018)) and a variety of other physical processes that are not within the scope of surface mass balance (SMB) modeling. Runoff delay can be implemented outside the RCMs (e.g. Liston and Mernild (2012);

Mernild et al. (2018)), but for this version of the product we assume that once an RCM classifies meltwater as "runoff", it is
365 instantly transported to the outlet. Actual lags between melt and discharge range from hours to years (Colgan et al., 2011; van
As et al., 2017; Rennermalm et al., 2013; Livingston et al., 2013).

Data released here includes no additional lag beyond the RCM lag, although a 7-day running mean (van As et al., 2017) is
included in all of the results presented here except Fig. 6 which shows monthly summed data, and Fig. 4 which shows yearly
summed data. When increasing the signal to noise by summing by year (Fig. 4 vs. Fig. 3), model results more closely match
370 observations.

4.3.2 Basin uncertainty

Basin uncertainty is a function of the subglacial routing assumptions (the k parameter in Eq. 2, which in reality varies in both
space and time). However, basin uncertainty does not necessary translate to discharge uncertainty. For example, two large
almost-overlapping ice basins may change their outlet location by one or a few grid cells between two k values, with a new
375 micro-basin occupying the same outlet as one of the old basin outlets. Large variation in discharge between one of these
theoretical large basins and its "replacement" at the same outlet for a different k is not an appropriate estimate of uncertainty
- rather the two large almost entirely overlapping basins, but with different outlets, should be compared. This fluidity of
basins and outlets between k -scenarios makes it almost impossible to define, identify, and compare basins between scenarios,
unless working manually with individual basins (as we did, for example, at the Leverett Glacier observation location, modeled
380 upstream basin, and adjusted upstream basin (see Sect. 4.2.3)).

Another example is that for two different k values, the same ice outlet may theoretically have two different upstream basins
that only overlap at the single grid cell containing the outlet, but otherwise have no overlap, yet these two basins (possibly of
different size) may have the same discharge values. Put differently, although inland grid cells may change their outlet location
by large distances under different routing assumptions (Fig. 2), that does not imply upstream basin area changes under different
385 routing assumptions. Large changes in upstream catchment area are possible (Chu et al., 2016), but we note Chu et al. (2016)
highlight changes at only a few outlets and under the extreme scenario of $k = 1.11$ describing an over-pressured system.
Because $\rho_w/\rho_i = 1.09$, setting $k = 1.09$ reduces Eq. 2 to $h = z_s$, and is equivalent to an over-pressured system with surface
routing of the water. In a limited examination comparing our results with $k \in [0.8, 0.9, 1.0]$, we did not detect basins with large
changes in upstream area. In addition all time series graphics show the mean RCM discharge for $k = 1.0$, but the uncertainty
390 among all three k values (not shown) is small enough it is usually difficult to distinguish the three separate uncertainty bands -
the difference between RCMs or between RCMs and observations is much larger than uncertainty from the k parameter.

The above issues are specific to ice basins. Land basin outlets do not change location, and the range of upstream runoff to a
land outlet provides one metric of uncertainty introduced by the k parameter. This uncertainty among all three k values is small
at ice margin outlets. It is even smaller at land outlets which act as spatial aggregators and increase the signal-to-noise ratio.

395 Below, we discuss the known uncertainties, ranging from least to most uncertain.

The basins presented here are static approximations based on 100 m resolution surface DEM of a dynamic system. Land
basin boundaries are likely to be more precise and accurate than ice basins, because the land surface is better resolved, has

larger surface slopes, has negligible sub-surface flow, and is less dynamic than the ice surface. Even if basins and outlets seem visually correct from the 100 m product, the basin outline still has uncertainty on the order of hundreds of meters and will therefore include many minor errors and non-physical properties, such as drainage basin boundaries bisecting lakes. However, all artefacts we did find are significantly smaller than the resolution of the 1 km² RCM inputs. We do not show but note that when doing the same work with the 150 m BedMachine land surface DEM, some basins change their outlet locations significantly - draining on the opposite side of a spit or isthmus and into a different fjord than the streams do when observed in satellite imagery. We have not observed these errors in streams and basins derived from the 100 m ArcticDEM in a visual comparison with Google Earth, although they may still exist.

Moving from land basins to subglacial ice basins, the uncertainty increases because subglacial routing is highly dynamic on timescales from minutes to seasons (e.g. Werder et al. (2013)). This dynamic system may introduce large spatial changes in outflow location (water or basin "piracy", Ahlstrøm et al. (2002); Lindbäck et al. (2015); Chu et al. (2016)), but Stevens et al. (2018) suggests basins switching outlet locations may not be as common as earlier work suggests, and our sensitivity analysis suggests that near the margin where the majority of runoff occurs, outlet location often changes by less than 10 km under different routing assumptions (Fig. 2). The largest (> 100 km) changes in outlet location in Fig. 2 occur when the continental or ice flow divides move, and one or two of the *k* scenario(s) drain cells to an entirely different coast or sector of the ice sheet.

The regions near the domain edges - both the land coast and the ice margin - are covered by many small basins, and in this work basins < 1 km² are absorbed into their largest neighbor (see Methods section). By definition these basins are now hydraulically incorrect. An example can be seen in the Zackenberg basin (Fig. 16a, southwest corner of the basin), where one small basin on the southern side of a hydraulic divide was absorbed into the large Zackenberg basin that should be defined by and limited to the northern side of the mountain range.

Near the ice margin quality issues exist. At the margin, many of the small basins (absorbed or not) may be incorrect because the relative uncertainty between the bed to the surface increases. Minor mask mis-alignments may cause hydraulic jumps (waterfalls) at the margin, or sinks that then need to be filled by the algorithm, and may overflow away from the real stream. The solution for individual outlets is to visually examine modeled outlet location, nearby streams in satellite imagery, and the area of upstream catchments, as we did for the Leverett Glacier outlet (Sect 4.2.3). Alternatively, selecting several outlets in an area will likely include the nearby "correct" outlet. This can be automated and an effective method to aggregate all the micro-ice basins that occur at the domain edge is to select the downstream land basin associated with one ice outlet, and then all upstream ice outlets for that land basin.

4.3.3 RCM uncertainty

In addition to the basin delineation issues discussed above, the runoff product from the RCMs also introduces uncertainty into the product generated here. The RCM input products do not provide formal time- or space-varying error estimates, but of course do contain errors because they represent a simplified and discretised reality. RCM uncertainty is shown here with a value of $\pm 15\%$. The MAR uncertainty comes from an evaluation by the Greenland SMB Model Intercomparison Project (GrSMBMIP; Fettweis et al. (2020)) that examined the uncertainty of modelled SMB for 95% of the 10767 in-situ measurements over the

main ice sheet. The mean bias between the model and the measurements was 15% with a maximum of 1000 mmWE yr⁻¹. GrSMBMIP uses integrated values over several months of SMB, suggesting larger uncertainty of modeled runoff at the daily time scale. The RACMO uncertainty comes from an estimated average 5% runoff bias in RACMO2.3p2 compared to annual cumulative discharge from the Watson River (Noël et al., 2019). The bias increases to a maximum of 20% for extreme runoff years (e.g. 2010 and 2012), so here we select 15 %, a value between the reported 5 % and the maximum 20 % that matches the MAR uncertainty. We display ±15 % uncertainty in the graphics here and suggest this is a minimum value for daily runoff data.

The 15 % RCM uncertainty is represented graphically in the time series plots when comparing to each of the observations. It is not shown in the scatter plots because the log-log scaling and many points makes it difficult to display. In the time series plots, we show the mean value from the $k = 1.0$ scenario, and note that discharge from the other two k scenarios covered approximately the same range.

4.3.4 Observational uncertainty

When comparing against observations, additional uncertainty is introduced because the stage-discharge relationship is neither completely precise or accurate. We use published observation uncertainty when it exists. Only two observational data sets come with uncertainty: Watson River and Qaanaaq. Similar to the RCM uncertainty, they are displayed in the time series but not the scatter plot for each observation graphic.

4.3.5 Mitigating uncertainties

Traditional uncertainty propagation is further complicated because it is not clear to what extent the three uncertainties (observational, RCM, and routing model) should be treated as independent from each other - all three uncertainties are likely to show some correlation with elevation, slope, air temperature, or other shared physical processes.

Many of the uncertainties discussed here can be mitigated by increasing the signal to noise ratio of the product. Because we provide a high spatial and temporal resolution product, this is equivalent many signals, each of which has some uncertainty (noise). Averaging results spatially or temporally, if possible for a downstream use of this product, will increase the signal to noise ratio and reduce uncertainty.

For example, because we provide basins for the entire ice sheet, total discharge is not subject to basin uncertainty. Any error in the delineation of one basin must necessarily be corrected by the inclusion (if underestimate) or exclusion (if overestimate) of a neighboring basin, although neighboring basins may introduce their own errors. Therefore, summing basins reduces the error introduced by basin outline uncertainty, and should be done if a downstream product does not need an estimate of discharge from a single outlet. This feature is built-in to coastal outlet discharge which is not as sensitive to our routing algorithm as ice margin outlet discharge because most coast outlets include a range of upstream ice margin outlets (e.g. Fig. 7 v. 9). Conversely, at the ice margin, outlet location and discharge volume is more uncertain than at the land coast. However, most runoff is generated near the ice margin and as runoff approaches the margin, there are fewer opportunities to change outlet location (Fig. 2).

465 Our coverage algorithm only fills in glaciated regions that have at least some RCM coverage. When working with basins that have glaciated areas and no RCM coverage as in the case for all four of the GEM outlets near Nuuk, discharge can be approximated by estimating discharge from the nearest covered glaciated area with a similar climatic environment.

Temporally, errors introduced by this study's assumption of instantaneous discharge can be reduced by summing or averaging discharge over larger time periods, or applying a lag function to the time series as done here and in van As et al. (2017).
470 Although a given volume of water may remain in storage long term, if one assumes that storage is roughly steady state, then long-term storage shown by, for example, dye trace studies, can be ignored - the volume with the dye may be stored, but a similar volume should be discharged in its place.

4.3.6 Quality control

The scale of the data are such that manual editing to remove artifacts is time and cost prohibitive. Here we provide one example
475 of incorrect metadata. The elevation of each outlet is included as metadata by looking up the bed elevation in the BedMachine data set at the location of each outlet. Errors in BedMachine or in the outlet location (defined by the GIMP ocean mask) introduce errors in outlet elevation.

A large basin in NW Greenland has metadata outlet elevation > 0 (gray in Fig. 1) but appears to be marine terminating when viewed in satellite imagery. Elsewhere the land vs. marine terminating color coding in Fig. 1 appears to be mostly correct, but
480 this view only provides information about the sign of the elevation, not the magnitude (i.e. if the reported depth is correct). Ice outlets can occur above, at, or below 0 m. It is easier to validate the land terminating basins, which should in theory all have an outlet elevation of 0 m. That is not the case (Fig. 18). It is possible for land outlets to be correctly assigned an elevation > 0 m, if a land basin outlet occurs at a waterfall off a cliff (as might occur the edges of Petermann fjord) or due to DEM discretization of steep cells. However, most of the land outlets at elevations other than 0 are likely due to mask misalignment pushing the
485 coast into fjords (negative land elevation) or inland (positive land elevation). The bulk of land discharge does occur within the 10 m bin at 0 m elevation. More than 75 % of land outlets occur within ± 10 m, and 90 % within 30 m (Fig. 18).

4.4 Other sources of freshwater

The liquid water discharge product provided here is only one source of freshwater that leaves the ice sheet and affects fjords and coastal seas. The other primary freshwater source is iceberg calving and submarine melt at the ice/ocean boundary of
490 marine terminating glaciers. A companion to the liquid water discharge product introduced here is provided by Mankoff et al. (2019, 2020), which estimates solid ice volume flow rates across gates near marine terminating glaciers. That downstream ice enters fjords as either calving icebergs or liquid water from submarine melting.

Both this product and Mankoff et al. (2020) provide liquid or solid freshwater volume flow rates at outlets (this product) or grounding lines (Mankoff et al., 2020), but actual freshwater discharge into a fjord occurs at a more complicated range of
495 locations. Solid ice melts throughout the fjord and beyond (e.g. Enderlin et al. (2016); Moon et al. (2017)), and the freshwater discharge presented here may enter at the reported depth (Sect. 4.3.6), but rapidly rises up the ice front and eventually flows into

the fjord at some isopycnal (see Mankoff et al. (2016)). The eventual downstream location of the fresh water is not addressed in this work.

Freshwater inputs directly to the water surface are also not included in this product. The flux (per square meter) to the water surface should be similar to the flux to the non-ice-covered land surface - assuming the orographic effects on precipitation produce similar fluxes to the near-land water surface.

Finally, basal melt from 1) geothermal heating (e.g. Fahnestock et al. (2001)) 2) frictional heating (e.g. Echelmeyer and Harrison (1990)) and 3) viscous heat dissipation from all previous freshwater sources (c.f. Mankoff and Tulaczyk (2017)) contributes additional discharge (see for example Jóhannesson et al. (2020)) to the surface melt. Geothermal and frictional heating are approximately in steady state and contribute freshwater throughout the winter months.

4.5 Summary

Of the 20 comparisons between the two RCMs and the 10 observations,

- In general this product shows good agreement between observations and the modeled discharge from the RCM runoff routed to the outlets, when comparing across multiple basins, especially when ignoring small basins with small glaciers that are not included in the RCMs (Fig. 3). The agreement is not as good when estimating the discharge variability within individual basins. From this, the product is more appropriately used to estimate the magnitude of the discharge from any individual basin, and perhaps provide some idea of the statistical variability, but not necessarily the precise amount of discharge for any specific day.
- The majority of the 20 comparisons have the 95 % prediction interval between scales of 1:5 and 5:1. From this, the model results match observations within plus-or-minus a factor of five, or half an order-of-magnitude. Put differently, the daily RCM values for single or few basins have an uncertainty of +500 % or -80 %.
- The uncertainty of +500%/-80% is for "raw" data: daily discharge for one or few basins with a simple temporal lag. When averaging spatially or temporally over larger areas or longer times, uncertainty decreases (Sect. 4.3). For example, when moving from daily data (Fig. 3) to annual sum (Fig. 4), the uncertainty is reduced to +100%/-50%.
- The two RCMs agree best with each other for the three observations dominated by large ice domains (Watson River (Sect. 4.2.2 & Fig. 7), Leverett Glacier (Sect. 4.2.3 & Fig. 9) which is a subset of the Watson River basin, and Kiattuut Sermiat (Sect. 4.2.4 & Fig. 10)). RCMs agree best with observations for ice-dominated basins with well-resolved bed topography in BedMachine (i.e. correct basins modeled in this work) - here only Leverett Glacier (Sect. 4.2.3 & Fig. 9) meets this criterion.
- Runoff errors increase with low discharge (panels 'd' in Figs. 7, 9 to 17).
- For land basins, errors are dominated by RCM runoff uncertainty, and may be systematic (bias).

530

- For ice basins, errors are dominated by basin uncertainty. Errors between similar-sized and neighboring basins are likely to offset and may even cancel each other. Even so, a conservative treatment might consider errors between basins as random and reduce by the sum of the squares when summing discharge from multiple similar-sized and neighboring basins.

5 Product description

These data contain a static map of Greenland’s hydrological outlets, basins, and streams and a times-series of discharge from each outlet.

The output data are provided in the following formats: The stream data are provided as a GeoPackage standard GIS product and a metadata CSV that includes the stream type (start or intermediate segment), network, stream along-flow length, stream straight length, sinuosity, source elevation, outlet elevation, and a variety of stream indices such as the Strahler, Horton, Shreve, Hack, and other parameters (Jasiewicz and Metz, 2011). We note that the subglacial streams are unvalidated with respect to actual subglacial conduits, and they should be used with caution. The outlet data are also provided as a GeoPackage and CSV, each of which include the outlet ID (linked to the basin ID), the longitude, latitude, EPSG:3413 x and y, and the outlet elevation. The outlet elevation is the BedMachine bed elevation at the outlet location, and users should be aware of quality issues identified in Sect. 4.3.6. The ice outlet metadata includes the ID, lon, lat, x, and y of the downstream land outlet, if one exists. The basin product GeoPackage includes the geospatial region that defines the basin. The metadata CSV includes the basin ID (linked to the outlet ID), and the area of each basin. The time-series discharge product is provided as four NetCDF files per year, one for each domain (ice margin, land coast) and one for each RCM (MAR and RACMO). The NetCDF files contain an unlimited time dimension, usually containing 365 or 366 days, much of the same metadata as the outlets CSV file, including the outlet (a.k.a station) ID, the latitude, longitude, and altitude of the outlet, and a runoff variable with dimensions (station, time) and units $\text{m}^3 \text{s}^{-1}$.

5.1 Database access software

The data can be accessed with custom code from the raw data files. However, to support downstream users we provide a tool to access the outlets, basins, and discharge for any region of interest (ROI). The ROI can be a point, a list describing a polygon, or a file, with units in longitude,latitude (EPSG:4326) or meters (EPSG:3413). If the ROI includes any land basins, an option can be set to include all upstream ice basins and outlets, if they exist. The script can be called from the command line (CLI) and returns CSV formatted tables, or within Python and returns standard Python data structures (from the `GeoPandas` or `xarray` package).

For example, to query for discharge at one point (50.5 °W, 67.2 °N), the following command is issued:

```
python ./discharge.py --base ./freshwater --roi=-50.5,67.2 --discharge,
```

where `discharge.py` is the provided script, `./freshwater` is the folder containing the downloaded data, and `--discharge` tells the program to return RCM discharge (as opposed to `--outlets` which would return basin and outlet information). The program documentation and usage examples are available at <http://github.com/mankoff/freshwater> (Mankoff, 2020b).

560 Because the `--upstream` option is not set, the `--discharge` option is set, and the point is over land, the results of this command are a time series for the MAR and RACMO land discharge for the basin containing this point. A small subset (the first 10 days of June 2012) are shown as an example:

time	MAR _{land}	RACMO _{land}
2012-06-01	0.043025	0.382903
2012-06-02	5.5e-05	0.095672
2012-06-03	5e-05	0.009784
2012-06-04	9e-06	-0.007501
2012-06-05	0.008212	0.007498
2012-06-06	28.601947	0.607345
2012-06-07	0.333926	0.05691
2012-06-08	0.489437	0.204384
2012-06-09	0.038816	0.167325
2012-06-10	5.1e-05	0.011415

565 If the `upstream` option is set, two additional columns are added: One for each of the two RCM ice domains. A maximum of six columns may be returned: 2 RCM times (1 land + 1 ice + 1 upstream ice domain), because results are summed across all outlets within each domain when the script is called from the command line (summing is not done when the script is accessed from within Python).

If the `--outlets` option is set instead of the `--RCM` option, then results are a table of outlets. For example, moving 10 ° east and over the ice,

570 `python ./discharge.py --base ./freshwater --roi=-40.5,67.2 --outlets`
 results in

index	id	lon	lat	x	y	elev	domain	upstream	coast _{id}	...
0	118180	-38.071	66.33	313650	-2580750	-78	land	False	-1	...
1	67133	-38.11	66.333	311850	-2580650	-58	ice	False	118180	...

575 If the script is accessed from within Python, then the `RCM` option returns an `xarray Dataset` of discharge, without aggregating by outlet, and the `outlets` option returns a `GeoPandas GeoDataFrame`, and includes the geospatial location of all outlets and outline of all basins, and can be saved to GIS-standard file formats for further analysis.

6 Conclusions

We provide a 100 m spatial resolution data set of streams, outlets, and basins, and a 1 day temporal resolution data set of discharge through those outlets for the entire ice-sheet area from 1958 through 2019. Access to this database is made simple for non-specialists with a Python script. Comparing the two RCM-derived discharge products to 10 gauged streams shows the
580 uncertainty is approximately plus-or-minus a factor of five, or half an order-of-magnitude, or +500%/-80%, when comparing daily discharge for single or few basins.

Because of the high spatial (individual basins) and temporal (daily) resolution, larger uncertainty exists than when working over larger areas or time-steps. These larger areas and times can be achieved through spatial or temporal averaging (or implementing a lag function) of this product.

585 This liquid freshwater volumetric flow rate product is complemented by a solid ice discharge product (Mankoff et al., 2020). Combined, these provide an estimate of the majority of freshwater (total solid ice and liquid) flow rates from the Greenland ice sheet into fjords and coastal seas, at high temporal resolution and process-level spatial resolution (i.e. glacier terminus for solid ice discharge, stream for liquid discharge).

This estimate of freshwater volume flow rate into Greenland fjords aims to support further studies of the impact of freshwater
590 on ocean physical, chemical, and biological properties; fjord nutrient, sediment, and ecosystems; and larger societal impacts of freshwater on the fjord and surrounding environments.

7 Code and data availability

The data from this work is available at doi:10.22008/promice/freshwater (Mankoff, 2020a).

The code and a website for post-publication updates is available at <https://github.com/mankoff/freshwater> (Mankoff, 2020b)
595 where we document ongoing changes to this work and use the GitHub Issues feature to collect suggested improvements, document those improvements as they are implemented, document problems that made it through review, and mention related works not yet published. This version of the document is generated with git commit version 834cd97 .

8 Figures

8.1 Overview




Figure 1. Map of Greenland showing all basins and the location of 10 gauged streams used for comparison. Land basins shown in green. Ice basins in blue when outlet elevation < 0 , and gray when outlet elevation ≥ 0 (outlet error elevation is discussed in Sect. 4.3.6). Black boxes and labels mark location of stream gauge observation locations (see Table 1).




Figure 2. Map of Greenland showing maximum possible distance between outlet locations for all cells, based on three effective basal pressure regimes ($k \in [0.8, 0.9, 1.0]$, Eq 2). Contour line in Greenland shows 1500 m elevation contour - most runoff occurs below this elevation.

8.3 Bulk observation v. RCM scatter plots




Figure 3. Daily runoff vs. observations for 10 outlets and a total of 17,370 days. Solid lines show 1:1 (center), 1:5 (upper), and 5:1 (lower). Grey band shows 95 % prediction interval. Red band shows 95 % prediction interval when removing the GEM stations near Nuuk (Table 1) that have small glaciers not included in the RCMs




Figure 4. Similar to Figure 3, except here showing annual sum of observed runoff - all days within each year when observations exist are summed. Days without observation are excluded from this comparison. Solid lines show 1:1 (center), 1:2 (upper), and 2:1 (lower). Grey band shows 95 % prediction interval.

8.4 Modified Tukey plot for all observations




Figure 5. Observation vs. "RCM minus observations" for MAR (left) and RACMO (right), discussed in Sect. 4.2.1. Number of samples at a location is represented by color. Horizontal solid line shows mean, dashed lines $\pm 95\%$ quantile range, and horizontal split denotes the bottom 1/3 and top 2/3rds quantiles of observed discharge. The four near-Nuuk GEM basins which have glaciers not included in the RCM domain are excluded.

8.5 Bamber 2018




Figure 6. Disko Island comparison between this product and Bamber et al. (2018). Light green are land basins with dark green outlet dots. Light blue are ice basins with dark blue outlet dots. Brown and hatched blue 5 km² cells are the land and ice runoff locations, respectively, from Bamber et al. (2018). Bottom graphs show ice (upper) and land (lower) runoff for the 2012 runoff calendar year.

8.6 Watson River




Figure 7. Graphical summary of Watson River outlet, basin, and discharge (W in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.2 for discussion of the Watson River basin. Basemap from Howat et al. (2014); Howat (2017a).




Figure 8. Watson River and manually adjusted basin area. Top panel: map view showing land and ice basin from this work (green and orange, respectively, same as region shown in 7, and two additional basins to the south in blue. Vertical dashed lines denote approximate location of 1500 m and 1850 m elevation. Bottom panel: Kernel density estimate (concentration of points) comparing observed vs. average of RACMO and MAR RCM runoff for the default land and ice basin (orange; filled) and with the additional southern basins (blue; lines). Solid and dashed lines are 1:1 and 2:1 (respectively) observed-to-RCM ratios.

8.8 Leverett Glacier




Figure 9. Graphical summary of Leverett Glacier outlet, basin, and discharge (L in Fig. 1). Red X in panel A marks actual observation location, but adjusted here to orange diamond within the ice basin. See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.3 for discussion of the Leverett Glacier basin. Basemap from Howat et al. (2014); Howat (2017a).

8.9 Kiattuut Sermiat




Figure 10. Graphical summary of Kiattuut Sermiat outlet, basin, and discharge (Ks in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.4 for discussion of the Kiattuut Sermiat basin. Basemap from Howat et al. (2014); Howat (2017a).

8.10 Kingigtorsuaq




Figure 11. Graphical summary of Kingigtorsuaq outlet, basin, and discharge (K in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.5 for discussion of the Kingigtorsuaq basin. Basemap from Howat et al. (2014); Howat (2017a).

8.11 Oriartorfik




Figure 12. Graphical summary of Oriartorfik outlet, basin, and discharge (O in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.5 for discussion of the Oriartorfik basin. Basemap from Howat et al. (2014); Howat (2017a).




Figure 13. Graphical summary of Teqinngalip outlet, basin, and discharge (T in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.5 for discussion of the Teqinngalip basin. Basemap from Howat et al. (2014); Howat (2017a).

8.13 Kobbefjord




Figure 14. Graphical summary of Kobbefjord outlet, basin, and discharge (Kb in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.5 for discussion of the Kobbefjord basin. Basemap from Howat et al. (2014); Howat (2017a).

8.14 Røde Elv




Figure 15. Graphical summary of Røde Elv outlet, basin, and discharge (R in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.6 for discussion of the Røde Elv basin. Basemap from Howat et al. (2014); Howat (2017a).

8.15 Zackenberg




Figure 16. Graphical summary of Zackenberg outlet, basin, and discharge (Z in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.6 for discussion of the Zackenberg basin. Basemap from Howat et al. (2014); Howat (2017a).

8.16 Qaanaaq




Figure 17. Graphical summary of Qaanaaq outlet, basin, and discharge (Q in Fig. 1). See Sect. 3.4 for general overview of graphical elements, and Sect. 4.2.6 for discussion of the Qaanaaq basin. Basemap from Howat et al. (2014); Howat (2017a).




Figure 18. Top: Histogram of outlet elevations. Bottom: Cumulative distribution of absolute land outlet elevation. More than 75 % of land outlets occur within ± 10 m, and 90 % within 30 m.

Appendix A: Software

This work was performed using only open-source software, primarily GRASS GIS (Neteler et al., 2012), CDO (Schulzweida, 2019), NCO (Zender, 2008), GDAL (GDAL/OGR contributors, 2020), and Python (Van Rossum and Drake Jr, 1995), in particular the Jupyter (Kluyver et al., 2016), dask (Dask Development Team, 2016; Rocklin, 2015), pandas (McKinney, 2010), geopandas (Jordahl et al., 2020), numpy (Oliphant, 2006), x-array (Hoyer and Hamman, 2017), and Matplotlib (Hunter, 2007) packages. The entire work was performed in Emacs (Stallman, 1981) using Org Mode (Schulte et al., 2012) on GNU/Linux and using many GNU utilities (See Supplemental Material). The parallel (Tange, 2011) tool was used to speed up processing. We used proj4 (PROJ contributors, 2018) to compute the errors in the EPSG 3413 projection. The color map for Fig. 2 comes from Brewer (2020).

All code used in this work is available in the Supplemental Online Material.

Author contributions. KDM produced this work - wrote the code and the text. APA and DVA helped with discussions of methods and quality control. WC, RSF, and DVA helped with writing. KK and SS supplied Qaanaaq data. XF and BN supplied RCM inputs. KL provided GEM data.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Funding was provided by the Programme for Monitoring of the Greenland Ice Sheet (PROMICE). Parts of this work were funded by the INTAROS project under the European Union's Horizon 2020 research and innovation program under grant agreement No. 727890. BN was funded by NWO VENI grant VI.Veni.192.019. DEMs were provided by the Polar Geospatial Center under NSF-OPP awards 1043681, 1559691, and 1542736. Data from the Greenland Ecosystem Monitoring Programme (GEM) were provided by Asiaq – Greenland Survey, Nuuk, Greenland. We thank Dorthe Petersen (ASIAQ) for help with basin quality control.

635 **References**

- Ahlstrøm, A. P., Petersen, D., Langen, P. L., Citterio, M., and Box, J. E.: Abrupt shift in the observed runoff from the southwestern Greenland ice sheet, *Science Advances*, 3, e1701169, <https://doi.org/10.1126/sciadv.1701169>, <http://dx.doi.org/10.1126/sciadv.1701169>, 2017.
- Ahlstrøm, A. P., Bøggild, C. E., Mohr, J. J., Reeh, N., Christensen, E. L., Olesen, O. B., and Keller, K.: Mapping of a hydrological ice-sheet drainage basin on the West Greenland ice-sheet margin from ERS-1/-2 SAR interferometry, ice-radar measurement and modelling, *Annals of Glaciology*, 34, 309–314, <https://doi.org/10.3189/172756402781817860>, <http://dx.doi.org/10.3189/172756402781817860>, 2002.
- Altman, D. G. and Bland, J. M.: Measurement in Medicine: The Analysis of Method Comparison Studies, *The Statistician*, 32, 307, <https://doi.org/10.2307/2987937>, <http://dx.doi.org/10.2307/2987937>, 1983.
- An, S.-I., Kim, H., and Kim, B.-M.: Impact of freshwater discharge from the Greenland ice sheet on North Atlantic climate variability, *Theoretical and Applied Climatology*, 112, 29–43, <https://doi.org/10.1007/s00704-012-0699-6>, 2012.
- 645 Balmonte, J. P., Hasler-Sheetal, H., Glud, R. N., Andersen, T. J., Sej, M. K., Middelboe, M., Teske, A., and Arnosti, C.: Sharp contrasts between freshwater and marine microbial enzymatic capabilities, community composition, and DOM pools in a NE Greenland fjord, *Limnology and Oceanography*, <https://doi.org/10.1002/lno.11253>, <http://dx.doi.org/10.1002/lno.11253>, 2019.
- Bamber, J. L., Griggs, J. A., Hurkmans, R. T. W. L., Dowdeswell, J. A., Gogineni, S. P., Howat, I., Mouginot, J., Paden, J., Palmer, S., Rignot, E., and Steinhage, D.: A new bed elevation dataset for Greenland, *The Cryosphere*, 7, 499–510, <https://doi.org/10.5194/tc-7-499-2013>, 650 2013.
- Bamber, J. L., Tedstone, A. J., King, M. D., Howat, I. M., Enderlin, E. M., van den Broeke, M. R., and Noel, B.: Land Ice Freshwater Budget of the Arctic and North Atlantic Oceans: 1. Data, Methods, and Results, *Journal of Geophysical Research: Oceans*, 123, 1827–1837, <https://doi.org/10.1002/2017jc013605>, 2018.
- Beckmann, J., Perrette, M., Beyler, S., Calov, R., Willeit, M., and Ganopolski, A.: Modeling the response of Greenland outlet glaciers to global warming using a coupled flow line–plume model, *The Cryosphere*, 13, 2281–2301, <https://doi.org/10.5194/tc-13-2281-2019>, 655 <http://dx.doi.org/10.5194/tc-13-2281-2019>, 2019.
- Bendtsen, J., Mortensen, J., Lennert, K., and Rysgaard, S.: Heat sources for glacial ice melt in a West Greenland tidewater outlet glacier fjord: The role of subglacial freshwater discharge, *Geophysical Research Letters*, <https://doi.org/10.1002/2015GL063846>, 2015.
- Brewer, C. A.: ColorBrewer.org Website, <http://www.ColorBrewer.org>, accessed 2020-06-02, 2020.
- 660 Catania, G., Stearns, L., Moon, T., Enderlin, E., and Jackson, R.: Future Evolution of Greenland’s Marine-Terminating Outlet Glaciers, *Journal of Geophysical Research: Earth Surface*, <https://doi.org/10.1029/2018jg004873>, <http://dx.doi.org/10.1029/2018JF004873>, 2019.
- Chu, W., Creyts, T. T., and Bell, R. E.: Rerouting of subglacial water flow between neighboring glaciers in West Greenland, *Journal of Geophysical Research: Earth Surface*, 121, 925–938, <https://doi.org/10.1002/2015JF003705>, 2016.
- Citterio, M. and Ahlstrøm, A. P.: Brief communication “The aerophotogrammetric map of Greenland ice masses”, *The Cryosphere*, 7, 445–665 449, <https://doi.org/10.5194/tc-7-445-2013>, <http://dx.doi.org/10.5194/tc-7-445-2013>, 2013.
- Citterio, M., Sej, M. K., Langen, P. L., Mottram, R. H., Abermann, J., Hillerup Larsen, S., Skov, K., and Lund, M.: Towards quantifying the glacial runoff signal in the freshwater input to Tyrolerfjord–Young Sound, NE Greenland, *Ambio*, 46, 146–159, <https://doi.org/10.1007/s13280-016-0876-4>, <http://dx.doi.org/10.1007/s13280-016-0876-4>, 2017.
- Colgan, W., Rajaram, H., Anderson, R., Steffen, K., Phillips, T., Joughin, I. R., Zwally, H. J., and Abdalati, W.: The annual glaciohydrology cycle in the ablation zone of the Greenland ice sheet: Part 1. Hydrology model, *Journal of Glaciology*, 57, 697–709, 2011.
- 670

- Cowton, T. R., Slater, D., Sole, A., Goldberg, D., and Nienow, P.: Modeling the impact of glacial runoff on fjord circulation and submarine melt rate using a new subgrid-scale parameterization for glacial plumes, *Journal of Geophysical Research: Oceans*, 120, 796–812, <https://doi.org/10.1002/2014JC010324>, 2015.
- Cowton, T. R., Todd, J. A., and Benn, D. I.: Sensitivity of tidewater glaciers to submarine melting governed by plume locations, *Geophysical Research Letters*, <https://doi.org/10.1029/2019gl084215>, <http://dx.doi.org/10.1029/2019GL084215>, 2019.
- Crozier, J., Karlstrom, L., and Yang, K.: Basal control of supraglacial meltwater catchments on the Greenland Ice Sheet, *The Cryosphere*, 12, 3383–3407, <https://doi.org/10.5194/tc-12-3383-2018>, <http://dx.doi.org/10.5194/tc-12-3383-2018>, 2018.
- Dask Development Team: Dask: Library for dynamic task scheduling, <https://dask.org>, 2016.
- Delhasse, A., Kittel, C., Amory, C., Hofer, S., and Fettweis, X.: Brief communication: Interest of a regional climate model against ERA5 to simulate the near-surface climate of the Greenland ice sheet, *The Cryosphere Discussions*, <https://doi.org/10.5194/tc-2019-96>, <https://doi.org/10.5194/tc-2019-96>, 2019.
- Echelmeyer, K. and Harrison, W. D.: Jakobshavns Isbræ, West Greenland: Seasonal Variations in Velocity - or Lack Thereof, *Journal of Glaciology*, 36, 82–88, <https://doi.org/10.3189/s0022143000005591>, 1990.
- Enderlin, E. M., Hamilton, G. S., Straneo, F., and Sutherland, D. A.: Iceberg meltwater fluxes dominate the freshwater budget in Greenland's iceberg-congested glacial fjords, *Geophysical Research Letters*, <https://doi.org/10.1002/2016gl070718>, 2016.
- Fahnestock, M., Abdalati, W., Joughin, I. R., Brozena, J., and Gogineni, P. S.: High geothermal heat flow, basal melt, and the origin of rapid ice flow in central Greenland, *Science*, 294, 2338–2342, <https://doi.org/10.1126/science.1065370>, 2001.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., van As, D., Machguth, H., and Gallée, H.: Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1033, <https://doi.org/10.5194/tc-11-1015-2017>, <http://dx.doi.org/10.5194/tc-11-1015-2017>, 2017.
- Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., and et al.: GrSMBMIP: Intercomparison of the modelled 1980–2012 surface mass balance over the Greenland ice sheet, *The Cryosphere Discussions*, <https://doi.org/10.5194/tc-2019-321>, <http://dx.doi.org/10.5194/tc-2019-321>, 2020.
- Fried, M., Carroll, D., Catania, G., Sutherland, D., Stearns, L., Shroyer, E., and Nash, J.: Distinct frontal ablation processes drive heterogeneous submarine terminus morphology, *Geophysical Research Letters*, <https://doi.org/10.1029/2019gl083980>, <http://dx.doi.org/10.1029/2019GL083980>, 2019.
- GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, Open Source Geospatial Foundation, <https://gdal.org>, 2020.
- Gillard, L. C., Hu, X., Myers, P. G., and Bamber, J. L.: Meltwater pathways from marine terminating glaciers of the Greenland ice sheet, *Geophysical Research Letters*, 43, 10,873–10,882, <https://doi.org/10.1002/2016gl070969>, <http://dx.doi.org/10.1002/2016GL070969>, 2016.
- GRASS Development Team: Geographic Resources Analysis Support System (GRASS GIS) Software, Open Source Geospatial Foundation, USA, <https://grass.osgeo.org>, 2018.
- Hawkings, J., Wadham, J., Tranter, M., Lawson, E., Sole, A., Cowton, T., Tedstone, A., Bartholomew, I., Nienow, P., Chandler, D., and et al.: The effect of warming climate on nutrient and solute export from the Greenland Ice Sheet, *Geochemical Perspectives Letters*, p. 94–104, <https://doi.org/10.7185/geochemlet.1510>, <http://dx.doi.org/10.7185/geochemlet.1510>, 2015.
- Hawkings, J., Wadham, J., Telling, J., Bagshaw, E., Beaton, A., Chandler, D., and Dubnick, A.: Proglacial discharge measurements, Kiattuut Sermiat glacier, south Greenland (near Narsarsuaq), <https://doi.org/10.5281/zenodo.3685976>, 2016a.

- Hawkings, J., Wadham, J., Tranter, M., Telling, J., Bagshaw, E., Beaton, A., Simmons, S.-L., Chandler, D., Tedstone, A., and Nienow, P.: The Greenland Ice Sheet as a hot spot of phosphorus weathering and export in the Arctic, *Global Biogeochemical Cycles*, 30, 191–210, <https://doi.org/10.1002/2015gb005237>, 2016b.
- 710 Howat, I.: MEaSURES Greenland Ice Mapping Project (GIMP) 2000 Image Mosaic, Version 1, <https://doi.org/10.5067/4RNRTRRE4JCYD>, updated 2018. Used all subsets. Accessed 2020-02-04. NASA National Snow and Ice Data Center Distributed Active Archive Center, 2017a.
- Howat, I.: MEaSURES Greenland Ice Mapping Project (GIMP) Land Ice and Ocean Classification Mask, Version 1, <https://doi.org/10.5067/B8X58MQBFUPA>, used all subsets; Accessed 2019-03-21. NASA National Snow and Ice Data Center Distributed Active Archive Center, 2017b.
- 715 Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8, 1509–1518, <https://doi.org/10.5194/tc-8-1509-2014>, <http://dx.doi.org/10.5194/tc-8-1509-2014>, 2014.
- Hoyer, S. and Hamman, J. J.: xarray: N-D labeled Arrays and Datasets in Python, *Journal of Open Research Software*, 5, <https://doi.org/10.5334/jors.148>, <http://dx.doi.org/10.5334/jors.148>, 2017.
- 720 Hunter, J. D.: Matplotlib: A 2D graphics environment, *Computing In Science & Engineering*, 9, 90–95, 2007.
- Jasiewicz, J. and Metz, M.: A new GRASS GIS toolkit for Hortonian analysis of drainage networks, *Computers & Geosciences*, 37, 1162–1173, <https://doi.org/10.1016/j.cageo.2011.03.003>, <http://dx.doi.org/10.1016/j.cageo.2011.03.003>, 2011.
- Jordahl, K., den Bossche, J. V., Wasserman, J., McBride, J., Fleischmann, M., Gerard, J., Tratner, J., Perry, M., Farmer, C., Hjelle, G. A., Gillies, S., Cochran, M., Bartos, M., Culbertson, L., Eubank, N., Bilogur, A., and maxalbert: *geopandas/geopandas: v0.7.0*, <https://doi.org/10.5281/zenodo.3669853>, <https://doi.org/10.5281/zenodo.3669853>, 2020.
- 725 Jóhannesson, T., Pálmason, B., Hjartarson, J., Jarosch, A. H., Magnússon, E., Belart, J. M. C., and Gudmundsson, M. T.: Non-surface mass balance of glaciers in Iceland, *Journal of Glaciology*, p. 1–13, <https://doi.org/10.1017/jog.2020.37>, 2020.
- Kamenos, N. A., Hoey, T. B., Nienow, P. W., Fallick, A. E., and Claverle, T.: Reconstructing Greenland ice sheet runoff using coralline algae, *Geology*, 40, 1095–1098, <https://doi.org/10.1130/G33405.1>, 2012.
- 730 Kanna, N., Sugiyama, S., Ohashi, Y., Sakakibara, D., Fukamachi, Y., and Nomura, D.: Upwelling of Macronutrients and Dissolved Inorganic Carbon by a Subglacial Freshwater Driven Plume in Bowdoin Fjord, Northwestern Greenland, *Journal of Geophysical Research: Biogeosciences*, 123, 1666–1682, <https://doi.org/10.1029/2017jg004248>, <http://dx.doi.org/10.1029/2017JG004248>, 2018.
- Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B., Bussonnier, M., Frederic, J., Kelley, K., Hamrick, J., Grout, J., Corlay, S., Ivanov, P., Avila, D., Abdalla, S., and Willing, C.: *Jupyter Notebooks – a publishing format for reproducible computational workflows*, 2016.
- 735 Kondo, K. and Sugiyama, S.: Discharge measurement at the outlet stream of Qaanaaq Glacier in the summer 2017-2019, https://doi.org/10.22008/hokkaido/data/meltwater_discharge/qaanaaq, GEUS Data Center, 2020.
- Lampkin, D. J. and VanderBerg, J.: A preliminary investigation of the influence of basal and surface topography on supraglacial lake distribution near Jakobshavn Isbrae, western Greenland, *Hydrological Processes*, 25, 3347–3355, <https://doi.org/10.1002/hyp.8170>, <http://dx.doi.org/10.1002/hyp.8170>, 2011.
- 740 Langen, P. L., Mottram, R. H., Christensen, J. H., Boberg, F., Rodehacke, C. B., Stendel, M., van As, D., Ahlstrøm, A. P., Mortensen, J., Rysgaard, S., Petersen, D., Svendsen, K. H., Aħalgeirsdóttir, G., and Cappelen, J.: Quantifying energy and mass fluxes controlling Godthåbsfjord freshwater input in a 5 km simulation (1991-2012), *Journal of Climate*, 28, 3694–3713, <https://doi.org/10.1175/JCLI-D-14-00271.1>, 2015.
- 745 Langley, K.: GEM river discharge measurements, https://doi.org/10.22008/asiaq/data/meltwater_discharge/gem, GEUS Data Center, 2020.

- Lewis, S. M. and Smith, L. C.: Hydrologic drainage of the Greenland ice sheet, *Hydrological Processes*, 23, 2004–2011, <https://doi.org/10.1002/hyp.7343>, 2009.
- Lindbäck, K., Pettersson, R., Doyle, S. H., Helanow, C., Jansson, P., Kristensen, S. S., Stenseng, L., Forsberg, R., and Hubbard, A. L.: High-resolution ice thickness and bed topography of a land-terminating section of the Greenland Ice Sheet, *Earth System Science Data*, 6, 331–338, <https://doi.org/10.5194/essd-6-331-2014>, <http://dx.doi.org/10.5194/essd-6-331-2014>, 2014.
- Lindbäck, K., Pettersson, R., Hubbard, A. L., Doyle, S. H., van As, D., Mikkelsen, A. B., and Fitzpatrick, A. A.: Subglacial water drainage, storage, and piracy beneath the Greenland Ice Sheet, *Geophysical Research Letters*, <https://doi.org/10.1002/2015GL065393>, 2015.
- Liston, G. E. and Mernild, S. H.: Greenland freshwater runoff. Part I: A runoff routing model for glaciated and nonglaciated landscapes (HydroFlow), *Journal of Climate*, 25, 5997–6014, <https://doi.org/10.1175/JCLI-D-11-00591.1>, 2012.
- Livingston, S. J., Clark, C. D., Woodward, J., and Kingslake, J.: Potential subglacial lake locations and meltwater drainage pathways beneath the Antarctic and Greenland ice sheets, *The Cryosphere*, 7, 1721–1740, <https://doi.org/10.5194/tc-7-1721-2013>, 2013.
- Mankoff, K. D.: Freshwater runoff, <https://doi.org/10.22008/promice/freshwater>, last access: 10 August 2020, 2020a.
- Mankoff, K. D.: Freshwater, GitHub <https://github.com/mankoff/freshwater>, last access: 10 August 2020, 2020b.
- Mankoff, K. D. and Tulaczyk, S. M.: The past, present, and future viscous heat dissipation available for Greenland subglacial conduit formation, *The Cryosphere*, 11, 303–317, <https://doi.org/10.5194/tc-11-303-2017>, 2017.
- Mankoff, K. D., Straneo, F., Cenedese, C., Das, S. B., Richards, C. G., and Singh, H.: Structure and dynamics of a subglacial discharge plume in a Greenlandic fjord, *Journal of Geophysical Research: Oceans*, 121, 8670–8688, <https://doi.org/10.1002/2016JC011764>, 2016.
- Mankoff, K. D., Colgan, W., Solgaard, A., Karlsson, N. B., Ahlstrøm, A. P., van As, D., Box, J. E., Khan, S. A., Kjeldsen, K. K., Mouginot, J., and Fausto, R. S.: Greenland Ice Sheet solid ice discharge from 1986 through 2017, *Earth System Science Data*, 11, 769 – 786, <https://doi.org/10.5194/essd-11-769-2019>, <https://doi.org/10.5194/essd-11-769-2019>, 2019.
- Mankoff, K. D., Solgaard, A., Colgan, W., Ahlstrøm, A. P., Khan, S. A., and Fausto, R. S.: Greenland Ice Sheet solid ice discharge from 1986 through March 2020, *Earth System Science Data*, 12, 1367–1383, <https://doi.org/10.5194/essd-12-1367-2020>, <http://dx.doi.org/10.5194/essd-12-1367-2020>, 2020.
- Martin Bland, J. and Altman, D. G.: Statistical methods for assessing agreement between two methods of clinical measurement, *The Lancet*, 327, 307–310, [https://doi.org/10.1016/s0140-6736\(86\)90837-8](https://doi.org/10.1016/s0140-6736(86)90837-8), [http://dx.doi.org/10.1016/S0140-6736\(86\)90837-8](http://dx.doi.org/10.1016/S0140-6736(86)90837-8), 1986.
- McKinney, W.: Data Structures for Statistical Computing in Python, in: *Proceedings of the 9th Python in Science Conference*, edited by van der Walt, S. and Millman, J., pp. 51 – 56, 2010.
- Mernild, S., Howat, I., Ahn, Y., Liston, G., Steffen, K., Jakobsen, B., Hasholt, B., Fog, B., and van As, D.: Freshwater flux to Sermilik Fjord, SE Greenland, *The Cryosphere*, 4, 453–465, <https://doi.org/10.5194/tc-4-453-2010>, 2010a.
- Mernild, S. H. and Liston, G. E.: Greenland Freshwater Runoff. Part II: Distribution and Trends, 1960–2010, *Journal of Climate*, 25, 6015–6035, <https://doi.org/10.1175/JCLI-D-11-00592.1>, 2012.
- Mernild, S. H., Liston, G. E., Hiemstra, C. A., and Steffen, K.: Surface melt area and water balance modeling on the Greenland ice sheet 1995–2005, *Journal of Hydrometeorology*, 9, 1191–1211, <https://doi.org/10.1175/2008JHM957.1>, 2008.
- Mernild, S. H., Liston, G. E., Hiemstra, C. A., Steffen, K., Hanna, E., and Christensen, J. H.: Greenland Ice Sheet surface mass-balance modelling and freshwater flux for 2007, and in a 1995–2007 perspective, *Hydrological Processes*, 23, 2470–2484, <https://doi.org/10.1002/hyp.7354>, 2009.

- Mernild, S. H., Liston, G. E., Steffen, K., van den Broeke, M., and Hasholt, B.: Runoff and mass-balance simulations from the Greenland Ice Sheet at Kangerlussuaq (Søndre Strømfjord) in a 30-year perspective, 1979–2008, *The Cryosphere*, 4, 231–242, <https://doi.org/10.5194/tc-4-231-2010>, <http://dx.doi.org/10.5194/tc-4-231-2010>, 2010b.
- 785 Mernild, S. H., Liston, G. E., van As, D., Hasholt, B., and Yde, J. C.: High-resolution ice sheet surface mass-balance and spatiotemporal runoff simulations: Kangerlussuaq, west Greenland, Arctic, Antarctic, and Alpine Research, 50, S100008, <https://doi.org/10.1080/15230430.2017.1415856>, <http://dx.doi.org/10.1080/15230430.2017.1415856>, 2018.
- Monteban, D., Pedersen, J. O. P., and Nielsen, M. H.: Physical oceanographic conditions and a sensitivity study on meltwater runoff in a West Greenland fjord: Kangerlussuaq, *Oceanologia*, <https://doi.org/10.1016/j.oceano.2020.06.001>, <http://dx.doi.org/10.1016/j.oceano.2020.06.001>, 2020.
- 790 Moon, T., Sutherland, D. A., Carroll, D., Felikson, D., Kehrl, L., and Straneo, F.: Subsurface iceberg melt key to Greenland fjord freshwater budget, *Nature Geoscience*, <https://doi.org/10.1038/s41561-017-0018-z>, <http://dx.doi.org/10.1038/s41561-017-0018-z>, 2017.
- Morlighem, M., Rignot, E. J., Mouginot, J., Seroussi, H., and Larour, E.: Deeply incised submarine glacial valleys beneath the Greenland ice sheet, *Nature Geoscience*, 7, 418–422, <https://doi.org/10.1038/NGEO2167>, 2014.
- 795 Morlighem, M., Williams, C., Rignot, E., An, L., Arndt, J. E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B., O’Cofaigh, C., Palmer, S. J., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K.: IceBridge BedMachine Greenland, Version 3, <https://doi.org/10.5067/2CIX82HUV88Y>, used all subsets; Accessed 2018-10-28, 2017a.
- 800 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I. M., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., Cofaigh, C. Ó., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo sounding combined with mass conservation, *Geophysical Research Letters*, <https://doi.org/10.1002/2017gl074954>, 2017b.
- 805 Mortensen, J., Bendtsen, J., Motyka, R. J., Lennert, K., Truffer, M., Fahnestock, M., and Rysgaard, S.: On the seasonal freshwater stratification in the proximity of fast-flowing tidewater outlet glaciers in a sub-Arctic sill fjord, *Journal of Geophysical Research: Oceans*, 118, 1382–1395, <https://doi.org/10.1002/jgrc.20134>, 2013.
- Munneke, P. K., Ligtenberg, S. R. M., van den Broeke, M. R., van Angelen, J. H., and Forster, R. R.: Explaining the presence of perennial liquid water bodies in the firn of the Greenland Ice Sheet, *Geophysical Research Letters*, 41, 476–483, <https://doi.org/10.1002/2013GL058389>, 2014.
- 810 Neteler, M., Bowman, M. H., Landa, M., and Metz, M.: GRASS GIS: a multi-purpose Open Source GIS, *Environmental Modelling & Software*, 31, 124–130, <https://doi.org/10.1016/j.envsoft.2011.11.014>, 2012.
- Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., and van den Broeke, M. R.: A daily, 1 km resolution data set of downscaled Greenland ice sheet surface mass balance (1958–2015), *The Cryosphere*, 10, 2361–2377, <https://doi.org/10.5194/tc-10-2361-2016>, <http://dx.doi.org/10.5194/tc-10-2361-2016>, 2016.
- 815 Noël, B., van de Berg, W. J., van Wessem, J. M., van Meijgaard, E., van As, D., Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., van Ulft, L. H., and et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, <http://dx.doi.org/10.5194/tc-12-811-2018>, 2018.

- 820 Noël, B., van de Berg, W. J., Lhermitte, S., and van den Broeke, M. R.: Rapid ablation zone expansion amplifies north Greenland mass loss, *Science Advances*, 5, eaaw0123, <https://doi.org/10.1126/sciadv.aaw0123>, <https://doi.org/10.1126/sciadv.aaw0123>, 2019.
- Oliphant, T. E.: *A guide to NumPy*, vol. 1, Trelgol Publishing USA, 2006.
- Perner, K., Moros, M., Otterå, O. H., Blanz, T., Schneider, R. R., and Jansen, E.: An oceanic perspective on Greenland's recent freshwater discharge since 1850, *Scientific Reports*, 9, <https://doi.org/10.1038/s41598-019-53723-z>, <http://dx.doi.org/10.1038/s41598-019-53723-z>,
825 2019.
- Pitcher, L. H., Smith, L. C., Gleason, C. J., and Yang, K.: CryoSheds: a GIS modeling framework for delineating land-ice watersheds for the Greenland Ice Sheet, *GIScience & Remote Sensing*, 53, 707–722, <https://doi.org/10.1080/15481603.2016.1230084>, <http://dx.doi.org/10.1080/15481603.2016.1230084>, 2016.
- Porter, C., Morin, P., Howat, I., Noh, M.-J., Bates, B., Peterman, K., Keeseey, S., Schlenk, M., Gardiner, J., Tomko, K., Willis, M.,
830 Kelleher, C., Cloutier, M., Husby, E., Foga, S., Nakamura, H., Platson, M., Wethington, Michael, J., Williamson, C., Bauer, G., Enos, J., Arnold, G., Kramer, W., Becker, P., Doshi, A., D'Souza, C., Cummins, P., Laurier, F., and Bojesen, M.: ArcticDEM, <https://doi.org/10.7910/DVN/OHHUKH>, date accessed: 2019-11-14, 2018.
- PROJ contributors: PROJ coordinate transformation software library, Open Source Geospatial Foundation, <https://proj4.org/>, 2018.
- Rennermalm, A. K., Smith, L. C., Chu, V. W., Box, J. E., Forester, R. R., van den Broeke, M. R., van As, D., and Moustafa, S. E.: Evidence
835 of meltwater retention within the Greenland ice sheet, *The Cryosphere*, 7, 1433–1445, <https://doi.org/10.5194/tc-7-1433-2013>, 2013.
- Rocklin, M.: Dask: Parallel Computation with Blocked algorithms and Task Scheduling, in: *Proceedings of the 14th Python in Science Conference*, edited by Huff, K. and Bergstra, J., pp. 130 – 136, 2015.
- Sasgen, I., Wouters, B., Gardner, A. S., King, M. D., Tedesco, M., Landerer, F. W., Dahle, C., Save, H., and Fettweis, X.: Return to rapid ice loss in Greenland and record loss in 2019 detected by the GRACE-FO satellites, *Communications Earth Environment*, 1,
840 <https://doi.org/10.1038/s43247-020-0010-1>, <http://dx.doi.org/10.1038/s43247-020-0010-1>, 2020.
- Schulte, E., Davison, D., Dye, T., and Dominik, C.: A multi-language computing environment for literate programming and reproducible research, *Journal of Statistical Software*, 46, 1–24, 2012.
- Schulzweida, U.: CDO User Guide, <https://doi.org/10.5281/zenodo.3539275>, <https://doi.org/10.5281/zenodo.3539275>, 2019.
- Sergienko, O. V.: Glaciological twins: basally controlled subglacial and supraglacial lakes, *Journal of Glaciology*, 59, 3–8,
845 <https://doi.org/10.3189/2013JoG12J040>, 2013.
- Shreve, R. L.: Movement of water in glaciers, *Journal of Glaciology*, 11, 205–214, 1972.
- Slater, D. A., Straneo, F., Felikson, D., Little, C. M., Goelzer, H., Fettweis, X., and Holte, J.: Estimating Greenland tidewater glacier retreat driven by submarine melting, *The Cryosphere*, 13, 2489–2509, <https://doi.org/10.5194/tc-13-2489-2019>, <http://dx.doi.org/10.5194/tc-13-2489-2019>, 2019.
- 850 Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y., and Balog, J.: Efficient meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *Proceedings of the National Academy of Sciences*, 112, 1001–1006, <https://doi.org/10.1073/pnas.1413024112>, 2015.
- Stallman, R. M.: EMACS the extensible, customizable self-documenting display editor, *Proceedings of the ACM SIGPLAN SIGOA symposium on Text manipulation -*, <https://doi.org/10.1145/800209.806466>, 1981.
855

- Stevens, L. A., Hewitt, I. J., Das, S. B., and Behn, M. D.: Relationship Between Greenland Ice Sheet Surface Speed and Modeled Effective Pressure, *Journal of Geophysical Research: Earth Surface*, 123, 2258–2278, <https://doi.org/10.1029/2017jf004581>, <http://dx.doi.org/10.1029/2017JF004581>, 2018.
- 860 Straneo, F., Curry, R. G., Sutherland, D. A., Hamilton, G. S., Cenedese, C., Våge, K., and Stearns, L. A.: Impact of fjord dynamics and glacial runoff on the circulation near Helheim Glacier, *Nature Geoscience*, 4, 322–327, <https://doi.org/10.1038/NGEO1109>, 2011.
- Sugiyama, S., Sakakibara, D., Matsuno, S., Yamaguchi, S., Matoba, S., and Aoki, T.: Initial field observations on Qaanaaq ice cap, northwestern Greenland, *Annals of Glaciology*, 55, 25–33, <https://doi.org/10.3189/2014aog66a102>, <http://dx.doi.org/10.3189/2014AoG66A102>, 2014.
- Tange, O.: GNU Parallel - The Command-Line Power Tool, ;login: The USENIX Magazine, 36, 42–47, 865 <https://doi.org/10.5281/zenodo.16303>, <http://www.gnu.org/s/parallel>, 2011.
- Tedstone, A., Bartholomew, I., Chandler, D., Cowton, C., Mair, D., Sole, A., Wadham, J., , and Nienow, P.: Proglacial discharge measurements, Leverett Glacier, south-west Greenland (2009–2012), <https://doi.org/10.5285/17c400f1-ed6d-4d5a-a51f-aad9ee61ce3d>, accessed 2020-02-21, 2017.
- Trusel, L. D., Das, S. B., Osman, M. B., Evans, M. J., Smith, B. E., Fettweis, X., McConnell, J. R., Noël, B. P. Y., and van den Broeke, M. R.: 870 Nonlinear rise in Greenland runoff in response to post-industrial Arctic warming, *Nature*, 564, 104–108, <https://doi.org/10.1038/s41586-018-0752-4>, <http://dx.doi.org/10.1038/s41586-018-0752-4>, 2018.
- van As, D., Hubbard, A., Hasholt, B., Mikkelsen, A. P. B., van den Broeke, M. R., and Fausto, R.: Large surface meltwater discharge from the Kangerlussuaq sector of the Greenland ice sheet during the record-warm year 2010 explained by detailed energy balance observations, *The Cryosphere*, 6, 199–209, <https://doi.org/10.5194/tc-6-199-2012>, 2012.
- 875 van As, D., Bech Mikkelsen, A., Holtegaard Nielsen, M., Box, J. E., Claesson Liljedahl, L., Lindbäck, K., Pitcher, L., and Hasholt, B.: Hypsometric amplification and routing moderation of Greenland ice sheet meltwater release, *The Cryosphere*, 11, 1371–1386, <https://doi.org/10.5194/tc-11-1371-2017>, <http://dx.doi.org/10.5194/tc-11-1371-2017>, 2017.
- van As, D., Hasholt, B., Ahlstrøm, A. P., Box, J. E., Cappelen, J., Colgan, W., Fausto, R. S., Mernild, S. H., Mikkelsen, A. B., Noël, B. P. Y., Petersen, D., and van den Broeke, M. R.: Reconstructing Greenland Ice Sheet meltwater discharge through the Watson River (1949–2017), 880 *Arctic, Antarctic, and Alpine Research*, <https://doi.org/10.1080/15230430.2018.1433799>, 2018.
- Van Rossum, G. and Drake Jr, F. L.: Python reference manual, Centrum voor Wiskunde en Informatica Amsterdam, 1995.
- Vandecrux, B., MacFerrin, M., Machguth, H., Colgan, W. T., van As, D., Heilig, A., Stevens, C. M., Charalampidis, C., Fausto, R. S., Morris, E. M., and et al.: Firn data compilation reveals widespread decrease of firn air content in western Greenland, *The Cryosphere*, 13, 845–859, <https://doi.org/10.5194/tc-13-845-2019>, <http://dx.doi.org/10.5194/tc-13-845-2019>, 2019.
- 885 Werder, M. A., Hewitt, I. J., Schoof, C. G., and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, *Journal of Geophysical Research*, 118, 2140–2158, <https://doi.org/10.1002/jgrf.20146>, 2013.
- Yang, K. and Smith, L. C.: Internally drained catchments dominate supraglacial hydrology of the southwest Greenland Ice Sheet, *Journal of Geophysical Research: Earth Surface*, 121, 1891–1910, <https://doi.org/10.1002/2016jf003927>, <http://dx.doi.org/10.1002/2016JF003927>, 2016.
- 890 Yang, K., Smith, L. C., Fettweis, X., Gleason, C. J., Lu, Y., and Li, M.: Surface meltwater runoff on the Greenland ice sheet estimated from remotely sensed supraglacial lake infilling rate, *Remote Sensing of Environment*, 234, 111 459, <https://doi.org/10.1016/j.rse.2019.111459>, <http://dx.doi.org/10.1016/j.rse.2019.111459>, 2019.

- Zender, C. S.: Analysis of self-describing gridded geoscience data with netCDF Operators (NCO), *Environmental Modelling & Software*, 23, 1338–1342, <https://doi.org/10.1016/j.envsoft.2008.03.004>, <https://doi.org/10.1016/j.envsoft.2008.03.004>, 2008.
- 895 Zuo, Z. and Oerlemans, J.: Modelling albedo and specific balance of the Greenland ice sheet: calculations for the Søndre Strømfjord transect, *Journal of Glaciology*, 42, 305–317, <https://doi.org/10.3189/s0022143000004160>, <http://dx.doi.org/10.3189/S0022143000004160>, 1996.