



Reconstruction of daily snowfall accumulation at 5.5km resolution over Dronning Maud Land, Antarctica, from 1850 to 2014 using an analog-based downscaling technique

Nicolas Ghilain¹, Stéphane Vannitsem¹, Quentin Dalaiden², Hugues Goosse², Lesley De Cruz¹, and Wenguang Wei³

¹Royal Meteorological Institute of Belgium, Brussels, Belgium

²Georges Lemaître Centre for Earth and Climate Research, Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium

³Key Laboratory of Regional Climate–Environment for Temperate East Asia, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

Correspondence: Stéphane Vannitsem (stephane.vannitsem@meteo.be)

Abstract. The surface mass balance (SMB) over the Antarctic Ice Sheet displays large temporal and spatial variations. Due to the complex Antarctic topography, modelling the climate at high resolution is crucial to accurately represent the dynamics of SMB. While ice core records provide a means to infer the SMB over centuries, the view is very spatially constrained. General circulation models (GCMs) estimate its spatial distribution over centuries, but with a resolution that is too coarse to capture the large variations due to local orographic effects. We have therefore explored a methodology to statistically downscale snowfall accumulation, the primary driver of SMB, from climate model historical simulations (1850–present day) over the coastal region of Dronning Maud Land. An analog method is set up over a period of 30 years with the ERA-Interim and ERA5 reanalyses (1979–2010 AD) and associated with snowfall daily accumulation forecasts from the Regional Atmospheric Climate Model (RACMO2.3) at 5.5 km spatial resolution over Dronning Maud in East Antarctica. The same method is then applied to the period from 1850 to present day using an ensemble of ten members from the CESM2 model. This method enables to derive a spatial distribution of the accumulation of snowfall, the principal driver of the SMB variability over the region. A new dataset of daily and yearly snowfall accumulation based on this methodology is presented in this paper (MASS2ANT dataset, <http://doi.org:10.5281/zenodo.4287517>, Ghilain et al. (2021)), along with comparisons with ice core data and available spatial reconstructions. It offers a more detailed spatio-temporal view of the changes over the past 150 years compared to other available datasets, allowing a possible connection with the ice core records, and provides information that may be useful in identifying the large-scale patterns associated to the local precipitation conditions and their changes over the past century.

1 Introduction

In the context of the global climate warming, polar ice sheets have increasingly gained attention, due to the threat of a massive sea level rise at the global scale (Garbe et al. , 2020). While the Greenland Ice sheet is eroding at an increasing speed both from the base and the surface (Lenaerts et al. , 2019), the Antarctic Ice Sheet is sometimes viewed as subject to a mitigation



mechanism to the observed melting of the ice shelf through an increased coastal precipitation due to a higher atmospheric humidity (Shepherd et al. , 2018; Krinner et al. , 2014; Agosta et al. , 2013; Medley and Thomas , 2019). This effect has been documented regionally and has been evidenced to be large over the Antarctic Peninsula (Thomas et al. , 2008) and over some coastal areas (Frezzoti et al. , 2013), but is less clear for the rest of the Antarctic Ice Sheet (Monaghan et al. , 2006; van den Berg et al. , 2005; Lenaerts et al. , 2012). However, the study of other coastal regions of Antarctica has revealed methodological limitations in estimating the surface mass balance, especially because the tools and observations available have a too low spatial resolution in regions where snowfall is orography induced and spatial variations are high (Eisen et al. , 2008). Dronning Maud Land (DML, 20°W – 45°E, map on Figure 2) is one of the sectors for which such uncertainty exists. Like any other coastal sector of Antarctica, the gradient of snowfall accumulation follows the large scale topography, with maxima over the coast and minima in the interior (Rotschky et al. , 2007), but it is highly variable at the scale of a few kilometers because of the ice rises and rumples that punctuate the coast (Lenaerts et al. , 2014). The available reconstitutions of surface mass balance distribution over the last 200 years are deduced from the interpolation of the quality controlled SMB estimations from ice cores drilled at several locations (Favier et al. , 2013), assuming an averaged (Rotschky et al. , 2007) or a time-dependent spatial distribution (Medley and Thomas , 2019).

General circulation models provide another source of information for the SMB over the last millenium, but, as the local topography is not well resolved by these models, the estimation of snowfall accumulation over the coast is inaccurate (Tetzner et al. , 2019). On another side, regional climate models (RCMs) have been adapted over the polar regions (van Wessem et al. , 2016; Agosta et al. , 2019) allowing the modelling of more detailed processes determining SMB at a scale of a few kilometers (Agosta et al. , 2019; Mottram et al. , 2020), but with the limitation of the period considered, generally not before the satellite era (past 40 years). The time frame of 40 years, though of relevance for climate analysis, seems too short to determine trends in snowfall accumulation potentially related to climate warming due to the uncertainty on the process at play, the large interannual variability in the region and the unknown history of the local SMB.

Indications of a recent increase of SMB in Dronning Maud Land have been found at some locations (Lenaerts et al. , 2013; Schlosser et al. , 2016; Medley et al. , 2018; Philippe et al. , 2016), but a stationary or decreasing trend has been found elsewhere (Thomas et al. , 2017; Vega et al. , 2016; Altnau et al. , 2015; Schlosser et al. , 2014). Most of the coastal ice core drilling sites are situated on top of an ice rise, a choice that may be due to the technical difficulties posed by the lateral flow of the icefields and the low volume of snow accumulated, both preventing to accurately date the strata (Matsuoka et al. , 2015; Goel et al. , 2020). But the drilling sites may not be representative of larger areas (Cavitte et al. , 2020; Kausch et al. , 2020). Extrapolating the findings from ice core sites to large areas may thus be difficult.

Statistical methods can be set up to emulate the precipitation field relating the spatial pattern of precipitation given by a RCM to large scale dynamics of the global climate model. We have explored such a methodology to statistically downscale the dominant SMB component over the area , the accumulated precipitation, from the climate model historical simulations (1850-present day). A method based on the search for analogs has been set up over a period of 30 years with the ERA-Interim reanalysis (1979-2010 AD) and associated with snowfall from the Regional Atmospheric Climate Model (RACMO2.3) at 5.5 km spatial resolution over Dronning Maud in East Antarctica. The method is then applied to the period from 1850 to present



day using an ensemble of 10 simulations from the CESM2 model. This method enables to derive 10 time-dependent spatial distribution of the accumulated snowfall over Dronning Maud Land at 5.5 km resolution from CESM2 ensemble members. In addition, we provide the time series of the 10 first principal components of each of the 4 variables used that can be used to characterize the regional synoptic situations. We present first the method and the data used, then the structure of the database, and finally the validation with ice core records and a discussion of the uncertainties.

2 Method and input data

Statistical downscaling follows a general scheme composed of three to four steps, involving several sources of information. In our case, at least three sources are required: 1) a GCM with coarse resolution but spanning a long time period, 2) a global reanalysis with coarse resolution spanning a shorter (recent) time frame, but of high quality, and finally 3) a regionally optimized RCM, with a finer resolution but spanning a short recent period. The objective is to evaluate the relationship (called here Perfect Prog - PP -) between the reanalysis and the RCM fields and then to apply it to the GCM in order to downscale its results to the scale of the RCM. Applying the PP directly to the GCM turned out to be unsuccessful due to discrepancies between the GCM climatology and the reanalyses. Therefore, three essential steps (Figure 1) are envisaged: the setting of a PP between the reanalysis and the RCM, the definition of a correction scheme of the GCM with the reanalysis over a common time frame to produce an “emulated” GCM, and finally the use of the PP on the “emulated” GCM for the whole period of the GCM integration.

2.1 Input data

Four datasets are used in the development of the downscaled GCM: the reanalyses ERA-Interim and ERA5, the GCM runs composed of are the 10 -members- CESM2 simulations, and the optimized reanalysis-driven RCM integration RACMO2.3.

The reanalyses ERA-Interim (Dee et al. , 2011) reanalysis has been largely validated all over the Earth surface, and also over Antarctica, where the surface mass balance deduced from it was found to be consistent with satellites observations of CloudSat (Palermo et al. , 2017), and exhibits a significant correlation with the observed interannual variability (Wang et al. , 2016). Because of its relatively coarse spatial resolution (0.75° in latitude-longitude) which cannot resolve the atmospheric circulation over sharp local orographic changes, it has been however observed that the local scale snowfall over DML coast, which can be heavy due to orographic uplift, is not properly estimated by ERA-Interim (Palermo et al. , 2017). Heavy snowfall days from ERA-Interim correspond to less than 20% of the days identified by an automatic weather station near Kohlen (Welker et al. , 2014). By contrast, large-scale fields over the continent and Southern Ocean have been assumed to be more accurate. Therefore, ERA-Interim fields of relative humidity at 850 hPa, geopotential height at 500 hPa, mean sea level pressure, sea ice extent, sea surface temperature and air temperature at 850 hPa were extracted from ECMWF archives. Snowfall was extracted the same way, for comparison. More recently, the ERA5 reanalysis (Hersbach et al. , 2020) has been released with a higher spatio-temporal resolution in comparison with ERA-Interim (0.25° and 1 hour). The same fields have been extracted from the Copernicus Climate Change Service Data Store (C3S CDS). The comparison of ERA5 snow accumulation with meteorological




Figure 1. The statistical downscaling can be generally decomposed into 3 successive steps, including 1) the association of the reanalysis with the regional climate model by a Perfect Prog (PP, Maraun and Widmann (2018)), 2) a correction of the historical GCM to make it unbiased compared to the reanalysis using Model Output Statistics, and 3) the successive correction of the GCM followed by the application of the Perfect Prog., giving way to an emulated RCM estimation using the corrected GCM.

observations over the Antarctic Peninsula indicates an increased ability to identify strong precipitation events, which indicates a strong consistency between the synoptic weather patterns and the observed precipitation (Tetzner et al. , 2019).

90 **The Regional Climate Model** The Regional Atmospheric Climate Model version 2.3 (RACMO2.3) was forced at its boundaries by the ERA Interim reanalysis, including an upper-air relaxation. This provides a simulation of the atmospheric variables and precipitations over the Antarctic Ice Sheet from 1979 to 2016 (Lenaerts et al. , 2013; van den Berg and Medley , 2016; van Wessem et al. , 2016). A simulation at 5.5 km resolution over specific regions of Antarctica, including Dronning Maud Land, allows to better emphasize the orographic effects of precipitation and the potential wind redistribution of snow.

95 The latter configuration at higher resolution allows studying in more detail a region with complex surface topography and SMB records (Lenaerts et al. , 2018, 2014). RACMO2.3 has been recognized to have the best fit to recent AIS SMB observations compared to other atmospheric and reanalysis models (Wang et al. , 2016; Rignot et al. , 2019).

The General Circulation Model The Community Earth System Model version 2 (Danabasoglu et al. , 2020) provides a model framework that allows the reconstruction of the evolution of atmospheric variables at 1 to 2 degree resolution over the globe. A set of ten historical CESM2 runs using forcing from CMIP6 have been made available for the industrial period (1850 - 100 2014, Historical CESM2 CAM6). A different initialization has been generated for each member. The CESM2 CAM6 provides new historical runs issued from the latest development of the CESM model, and is shown to be an improvement over CESM1 (Danabasoglu et al. , 2020).



2.2 The Method

105 The comparison of the meteorological and snowfall fields estimated from the high resolution (5.5 km) RCM RACMO2.3
and from the low resolution (regridDED to 1°latitude-longitude) reanalysis ERA-Interim has revealed large local differences
in the snowfall amounts and spatial distribution over the Antarctic coast in Dronning Maud Land. Two methods have first
been unsuccessfully tried. First, the analysis of back-trajectories (HYSPLIT model on NCEP/NCAR reanalysis (Stein et al. ,
2015)) of strong snowfall events has not lead to a strong relation between the location of the origin of those air masses and
110 the intensity of associated high precipitation amounts. This result contrasts with the success obtained with ERA-Interim and
ERA5 in identifying the atmospheric rivers responsible of such events (Gorodetskaya et al. , 2020). Secondly, another approach
based on the Random Forest technique was tested, but failed probably due to unclear and non-systematic relations between
large scale atmospheric patterns and snowfall intensity. In addition, since causal links can be model-dependent (Vannitsem et
al. , 2019), a method that does not over-exploit those causal links is required to guarantee the transferability to other models in
115 presence of large model uncertainties.

A downscaling method for snowfall based on the search for analogs has then been set up (Maraun and Widmann , 2018).
The search for analogs by passes the problem of non-systematic relations, as long as similar synoptic situations provide on
average similar precipitation amounts. The choice of a daily time step is in accordance with the duration of high precipitation
events over Antarctica's coast, which account for about 60% of the annual snowfall (Schlosser et al. , 2010; Reijmer and
120 van den Broeke , 2003; Turner et al. , 2019), last between 12 hours to 7 days long, and therefore can most of the time be
associated to synoptic atmospheric situations (Reijmer and van den Broeke , 2003; Schlosser et al. , 2010; Welker et al. ,
2014; Gorodetskaya et al. , 2020). The meteorological fields identified to be the most explanatory (and possibly replicable by
climate models) for precipitation rates over the Antarctic continent have been decomposed in Empirical Orthogonal Functions
(EOFs). The principal components weights (PCs) of the dominant EOFs (the first ten) are used for the selection of the analogs
125 (Sneyers and Gossens , 1988; Hannachi , 2004). To select the large-scale meteorological fields of interest for the downscaling
method, we have built for a set of 50 points from the RACMO2.3 domain an analog database from the association between the
principal components weights (PCs) of different fields from ERA-Interim and the RACMO2.3 precipitation over a 11 years
period (1979-1990). These points correspond to the highest annual snowfall accumulation over the area. The fields tested were:
geopotential height at 700 and 500 hPa, air temperature at 500 and 850 hPa, relative humidity at 700 and 500 hPa, surface
130 atmospheric pressure, sea-ice cover, and total precipitation (liquid and solid). One to four fields were used in the different
tests. The statistical evaluation of the performance was based on correlation and root-mean square difference at both daily and
annual scales after a bias correction (quantile mapping). The analysis over the validation period (1991-2000) indicates that the
optimal choice is reached with four fields: 1) geopotential height at 500 hPa, 2) surface pressure, 3) relative humidity at 700
hPa and 4) total precipitation. The same evaluation methodology was used to select 1) the optimum number of closest analogs
135 used for the estimation, namely 20 analogs, 2) the way to select the best estimation from the distribution of analogs: the mean
of the ensemble seems the most appropriate, 3) the optimum minimum geographical spanning of the reanalysis fields: latitude
40°S-90°S, longitude 10°W-60°E, and 4) the length of the training period: 3 is a minimum, but we use 11 years without a




Figure 2. The map shows the Princess Ragnhild Coast, which is part of Dronning Maud Land (red box), and the whole Antarctic continent. Verification of the performance of the downscaling scheme trained on 11 years (1979-1990) of ERA-Interim data and RACMO2.3 daily snowfall and applied on ERA-Interim for a 10 years period (1991-2000). The monthly time series comparison over a location of maximum of annual snowfall accumulation shows a high degree of accuracy, while the spatial comparison of the accumulated snowfall (in mm) over 1996 on Princess Ragnhild Coast in presence of ice rises (eg Derwael Ice Rise – DIR -, Lokeryggen Ice Rise – LIR and Hammaryggen Ice Rise - HIR) illustrates the high degree of fidelity of the analog method (top right) in reproducing the RCM (bottom right) accumulation patterns, especially in the West-East difference of accumulation around ice rises (Kausch et al. , 2020).

large statistical gain. As expected, the spatial and temporal variations of RACMO2.3 have been preserved (Figure 2 shows an example of time series comparison and a comparison over Princess Ragnhild Coast characterized by the presence of ice rises near the coast). The root mean square difference on a daily basis over the validation period ranges from 10% in large accumulation areas near the coast to 15% in the inner regions. The advantage of the analog method is that it allows one to identify the major types of weathers delivering precipitation and their occurrence over time in the form of PCs. This could be of interest for the analysis of a frequency change in the precipitation over long time periods.

The downscaling method was then repeated over the ensemble of the ten climate runs from the CESM2 model over the period 1850-2014. Before the downscaling could be applied, we first needed to make the PCs compatible with the reanalyses. A simple bias correction based on the linear regression of EOFs (Feudale and Tompkins , 2011; Yu et al, 2018), followed by a quantile mapping, transforms the CESM2 original PCs into “Reanalysis-like” PCs. The operation is done after the verification




Figure 3. The analogs method follows 3 steps: 1) the building of the analogs database using the association of the reanalyses PCs to the snowfall accumulation from RACMO2.3, 2) the transformation of CESM2 into “reanalysis like” fields, 3) the search in the analogs database. The second step ensures the compatibility of the principal components between the database and the GCM.

of the similarities of the EOFs among the members. The principal components can then be compared to the analog database in search for the closest events. As during the training process, quantile mapping defined over a 10 years period with the reanalysis
150 is applied to the complete time series to obtain the final downscaled estimations (Figure 3). No significantly high differences are expected from any one member versus another, as the initializations have been carefully controlled and the spectral analysis of the time series of PCs give very similar results (Danabasoglu et al. , 2020).

3 The dataset

3.1 Structure of the dataset

155 The dataset is composed of 46 files (netcdf4 format) in total, which accounts for about 500GB, when uncompressed. The yearly accumulated snowfall maps are provided in polar stereographic projection at 5.5 km resolution, conforming to the RACMO2.3p5.5 run set-up: 1 file per downscaled CESM2 run, for each re-analysis used for training (20 files). The daily dataset is provided in the form of time series (20 files): 10 files (netcdf4 format), each containing the time series of daily snowfall accumulation over each grid point of the RACMO2.3 domain on the ice sheet (DML coastal region) for the period
160 1850-2014, 365 days per year. Each file corresponds to a downscaling of one of the 10 runs from CESM2, based on the training on ERA-Interim and RACMO2.3. Latitude, longitude, corresponding line and column in the RACMO2.3 domain are stored for each grid point. Extraction of time series for a specific location is therefore straightforward, and, thanks to the column and line, it is possible to recompose daily maps. Due to the size of the daily files, only a set of 2 files are stored along with the annual data files on Zenodo, and the whole set is available on request. Another set of 10 files with the same structure is stored with the
165 results from the application of the same method trained on ERA5. In addition the time series of the 40 principal components used are stored in separate files, one for ERA-Interim trained, and one for ERA5 trained simulations (Figure 4), all provided with embedded metadata. PCs time series for the 40 EOFs for all the CESM2 members are stored in 2 additional files. The




Figure 5. Recomposed maps of downscaled daily snowfall (in mm) over DML on the 1st day of 1850 for members 1 to 8. A saturation at 10 mm per day has been defined for visualization.




Figure 6. Map of downscaled snowfall accumulated over a year (CESM2 member 1).

While a large disparity is observed on daily maps, the effect is much smaller on the annual accumulation (Figure 6). We illustrate this by a daily map (members 1 to 8) and an annual accumulation over the entire domain of the downscaled CESM2 member 1, trained with ERA5 (Figure 5).




Figure 7. The configurations of the 9 first EOFs for surface atmospheric pressure with their relative occurrence are ordered by importance in the decomposition of the variability of the field (represented here unitless, covering the total area 40°S - 90°S , 10°W - 60°E). The time series of the associated PCs for one year show the different amplitudes.

For each date, the PCs are provided, and associated to the EOFs for each meteorological field. An analysis of the association between snowfall regimes with the synoptic patterns is then possible. In Figure 7, we show an illustration of the first 9 EOFs for Surface atmospheric pressure over the GCM domain considered (40°S - 90°S , 10°W - 60°E), as well as the time series of the associated PCs, showing their various amplitudes.

180 3.3 Comparison with ice core measurements

The annual variation of the surface mass balance is estimated from ice cores retrieved from a set of locations over the coastal DML. The length of the records differs between the different sites and can cover more than 150 years. The annual accumulation computed from the downscaled daily snowfall is compared to SMB from the different ice core records from Thomas et al.



Table 1. Ice cores drilled over Dronning Maud Land used for comparison, extracted from Thomas et al. (2017)

Ice Core	Latitude	Longitude	Start date	End date	Reference
S100	-70.2439	4.8	1850	1999	Karczmaraska et al. (2004)
S20	-70.2472	4.8183	1956	1996	Isaksson et al. (1999)
H72	-69.2	41.08	1850	1999	Nishio et al. (2002)
B04	-70.6167	-8.3667	1892	1981	Schlosser (1999)
DIR	-70.25	26.34	1850	2011	Philippe et al. (2016)
E91	-73.6	-12.4333	1932	1991	Isaksson et al. (1996)
DML16 FB9813	-75.1669	5.0006	1850	1997	Oerter et al. (2000)

(2017), after a conversion of ice core estimates to water equivalent height (Figure 1, Figure 8). A good overall match between
185 the downscaled snowfall and the ice cores is found, as well as a large bias reduction compared to the use of CESM2 without
downscaling.

3.4 Comparison with other sources and other studies

Almost all the available datasets of spatial estimates of snowfall accumulation or SMB over Dronning Maud Land are 1) at high
resolution (20-30 km (Lenaerts et al. , 2018; Agosta et al. , 2019)) but spanning the last 30 years, 2) spanning a longer period
190 but at low resolution (0.75° , Medley and Thomas (2019)), or 3) at high resolution but averaged in time (Rotschky et al. , 2007).
Over other regions, downscaling using a physical orographic model has been used to obtain maps at high resolution from a
RCM over a longer period of time (Agosta et al. , 2013). In this study, we compare our dataset to two other sources. The first
reconstruction covers Western Dronning Maud Land (Rotschky et al. , 2007), which offers a static view of the accumulation
patterns based on the spatial interpolation of ice core data (data available from the PANGAEA website [http://doi.pangaea.de/
10.1594/PANGAEA.472297](http://doi.pangaea.de/10.1594/PANGAEA.472297)). A visual comparison with the CESM2 member 1 downscaled (trained with ERA5) shows an
195 enhanced variability along sharp elevation changes (Figure 9). The second reconstruction over the entire domain is based on
ice core datasets and climate patterns from GCM over the last 150 years on a yearly basis (Medley and Thomas , 2019), of
which we only consider the reconstruction using ERA-Interim (data available at [https://earth.gsfc.nasa.gov/cryo/data/antarctic-
accumulation-reconstructions](https://earth.gsfc.nasa.gov/cryo/data/antarctic-accumulation-reconstructions)). The downscaled CESM2 member 1 averaged over the total period seems in agreement with the
200 averaged reconstruction based on the ice core records, especially to reconstruct the gradient from the coast to inner land, but
depicts much more details related to topography (Figure 10).

4 Uncertainties

The uncertainties in the daily snowfall estimates compiled into this dataset can arise from 1) the choice and accuracy of
the model sources (regional climate model, re-analysis, “historical” climate model), and 2) the accuracy of the downscaling




Figure 8. The time series of yearly accumulated snowfall extracted from the database (CESM2 member 1, training with ERA-Interim, and all members in shaded gray) is compared to the SMB estimated from ice cores available from (Thomas et al. , 2017) (Ice core data available on the webpage of PAGES2k/Antarctica2k and in <https://data.bas.ac.uk/full-record.php?id=GB/NERC/BAS/PDC/00940>). RACMO2.3 time series has been superimposed (cyan) and the 10 CESM2 simulations (shaded blue), showing the large bias reduction thanks to the use of RACMO2.3 for the training. A 4-year moving average has been applied on the time series for the visualization.

205 method (choice of parameters, choice of predictive variables). In this section, we report a (non-exhaustive) quantification of the uncertainty level related to both the choice of model sources and of the parameters of the method.

4.1 Sampling in choice of analogs

A simple way to assess the effect of the sampling of the analogs on the estimation uncertainty is to use the bootstrap techniques. We have created a distribution of 100 ensembles of analogs, with the 20 analogs randomly drawn with repetition at the ice core sites (Figure 11). The downscaled CESM2 member 1 with 100 different analogs draws represented at the ice core sites. The uncertainty is between 5 and 10 % of the mean value.

210




Figure 9. The visual comparison of the average annual snowfall accumulation over Western Dronning Maud Land between the database from (Rotschky et al. , 2007) and the downscaled CESM2 member 1 (trained with ERA5) shows more pronounced asymmetrical patterns near sharp surface elevation changes. The difference in magnitude should be taken with caution, as only the snowfall component is provided by the MASS2ANT database, while (Rotschky et al. , 2007) is the integration of the total surface mass balance.




Figure 10. The coast-to-inland gradient of the downscaled CESM2 member 1 averaged over the total period seems in agreement with the averaged reconstruction based on the ice core records of (Medley and Thomas , 2019), but depicts much more details that could be useful in detailed analysis of ice core records representativity (same color scale).

4.2 Choice of reanalysis used for training

The choice of reanalysis used for training the analogs database has an effect on the downscaling obtained from CESM2. The choice of reanalysis influences the result, at equivalent training scores (Figure 12). This is one of the reasons for extending the database to ERA5 as training reanalysis instead of ERA-Interim. More independent reanalyses could be used to further assess the uncertainty. The difference CESM2 downscaling using ERA-Interim or ERA5 in the training reveals an uncertainty of 25% in average at the ice core sites.

4.3 Choice of “historical” GCM runs

CESM2 simulations consist of 10 members, corresponding to different initializations (Danabasoglu et al. , 2020). The inherent model uncertainty is therefore impacting the results and is quantified here (Figure 13). The scatter of the 10 members of



Table 2. Summary of the uncertainty levels associated to the choices of input model or sampling method. The uncertainties (mean uncertainties) are estimated on yearly accumulation of snowfall, relative to the basic methodology followed to derive the database, for at least one time series over the domain.

Snowfall Uncertainty (% mean yearly accumulation)	
Sampling in choice of analogs ⁽¹⁾	5 to 10 %
Choice of reanalysis for training ⁽²⁾	25%
Choice of “historical” GCM runs ⁽³⁾	17-22 % (std) to max 30-38%




Figure 11. The downscaled CESM2 member 1 with 100 different analogs draws represented at the ice core sites. The uncertainty is between 5 and 10 % of the mean value.

CESM2 downscaling using ERA-Interim in the training reveals an uncertainty of 20% (standard deviation) and up to 40% at some ice core sites, maybe a direct consequence of the CESM2 members not being temporally correlated.

4.4 Choice of RCM model

At last, the choice of the RCM critically drives the dynamical link between large-scale patterns and local snowfall, and the average and maximum amplitude of snowfall accumulation. An assessment of the uncertainty caused by the choice of RCM could be done using another RCM, with equally proven quality. For example, the Modele Atmospherique Regional (MAR) is

225




Figure 12. The difference CESM2 downscaling using ERA-Interim or ERA5 in the training reveals an uncertainty of 25% on average at the ice core sites.

another recognized RCM used for polar climates at high resolution. The newest simulations over Antarctica at 30 km resolution (Agosta et al. , 2019) have shown similarities with RACMO2 at 27 km resolution, but revealed localized differences over the ice sheet related to a more realistic sublimation of falling snowfall in comparison with RACMO2 (Gallée et al. , 2013). A simulation at the same resolution with another RCM optimized for polar regions, like MAR, could be of great interest to better frame the uncertainty linked to RCM physics. The uncertainty linked to the choice of RCM has not been evaluated in this study and could be envisaged to extend the database if new RCM simulations at such resolution are available in the future.

5 Conclusions

We propose here a reconstruction of snowfall evolution over Dronning Maud Land, Antarctica, at 5.5 km resolution using an analog-based downscaling technique. This technique has allowed us to exploit the detailed spatio-temporal estimation of snowfall from 30-year RACMO2.3 simulations in combination with synoptic patterns from recent reanalyses to statistically downscale the historical runs from CMIP6 (CESM2 model). The resulting database stores the ensembles of daily accumulated snowfall from 1850 to 2014, the pertinent information for synoptic patterns analysis (the principal components weights and empirical orthogonal functions from four large-scale meteorological fields), and the annual evolution of accumulated snowfall over Dronning Maud Land. The database can be used to analyze the detailed contribution of snowfall to the surface mass




Figure 13. The scatter of the 10 members of CESM2 downscaling using ERA-Interim in the training reveals an uncertainty of 20% (standard deviation) and up to 40% at some ice core sites.

balance over the region, its evolution and its association to synoptic weather conditions. The method can be easily replicated with new RCM and GCM simulations.

6 Data availability

The files of the dataset (the annual snowfall, the PCs and the EOFs) are available on Zenodo platform (<http://doi.org/10.5281/zenodo.4287517>). However, due to size limitations, only 2 daily snowfall files out of 20 have been stored there, the whole set is available on request to the contact author.

Author contributions. Project proposal and funding: SV & HG; Scientific supervision: SV; Design of the method: LDC, HG, SV & NG; Climatological Analysis NG, WW & SV; Implementation, tests, method tuning, verifications, database creation, validation: NG & SV; Analysis of the results: all authors; Preparation data material: QD & NG. Manuscript drafting: NG; Manuscript revisions: all authors.

250 *Competing interests.* The authors declare that they have no conflict of interest.



Acknowledgements. The authors thank the Santander Meteorology group for making available the MeteoLab Toolbox Matlab software (Cofino et al. , 2013), which was re-used and partly re-coded in fortran90 using multi-core parallel computing for the processing of the database. We thank Chad Greene for providing the Climate Toolbox for Matlab, which was used to verify intermediate steps. We thank Dr Jan Lenaerts for providing the RACMO2.3 fields over DML Dr Marie Cavitte for suggestions on the use of ice core datasets, and Dr
255 Jean-Louis Tison, Dr Stef Lhermite and Sarah Wauthy for fruitful discussions. The study was funded in the framework of the MASS2ANT project (<https://www.elic.ucl.ac.be/users/klein/Mass2Ant/index.html>, contract No BR/165/A2) by the Belgian Science Policy.



References

- Adusumilli, S., Fricker, H.A., Medley, B., Padman, L., and Siegfried, M.R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nature Geoscience*, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- 260 Agosta, C., Favier, V., Krinner, G., Gallée, H., Fettweis, X., and Genthon, C.: High-resolution modelling of the Antarctic surface mass balance, application for the twentieth, twenty first and twenty second centuries, *Clim Dyn*, <https://doi.org/10.1007/s00382-013-1903-9>, 2013.
- Agosta, C., Amory, C., Kittel, C., Orsi, A., Favier, V., Gallée, H., van den Broeke, M. R., Lenaerts, J. T. M., van Wessem, J. M., van de Berg, W. J., and Fettweis, X.: Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979–2015) and
265 identification of dominant processes, *The Cryosphere*, 13, 281–296, <https://doi.org/10.5194/tc-13-281-2019>, 2019.
- Altnau, S., Schlosser, E., Isaksson, E., and Divine, D.: Climatic signals from 76 shallow firn cores in Dronning Maud Land, East Antarctica, *The Cryosphere*, 9, 925–944, <https://doi.org/10.5194/tc-9-925-2015>, 2015.
- Bardossy, A., and Pegram, G.: Downscaling precipitation using regional climate models and circulation patterns toward hydrology, *Water Resour Res*, 47, W04505, <https://doi.org/10.1029/2010WR009689>, 2011.
- 270 Beersma, J.J., and Adri Buishand, T.: Multi-site simulation of daily precipitation and temperature conditional on the atmospheric circulation, *Clim Res*, 25, 121–133, 2003.
- Biau, G., Zorita, E., von Storch, H., and Wackernagel, H.: Estimation of precipitation by kriging in the EOF space of the sea level pressure field, *J Clim*, 12, 1070–1085, 1998.
- Cavitte, M. G. P., Dalaiden, Q., Goosse, H., Lenaerts, J. T. M., and Thomas, E.R.: Reconciling the surface temperature–surface mass balance
275 relationship in models and ice cores in Antarctica over the last 2 centuries, *The Cryosphere*, 14, 114083–4102, [10.5194/tc-14-4083-2020](https://doi.org/10.5194/tc-14-4083-2020), 2020.
- Cofino, A.S., Ancell, R., San-Martin, D., Herrera, S., Gutierrez, J.M., and Manzananas, R.: An open-source Matlab toolbox for Meteorology & Climate, <https://www.meteo.unican.es/en/software/meteolab>, 2013.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D.A., DuVivier, A.K., Edwards, J., Emmons, L.K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M.M., Large, W.G., Lauritzen, P.H., Lawrence, D.M., Lenaerts, J.T.M., Lindsay, K., Lipscomb, W.H.,
280 Mills, M.J., Neale, R., Oleson, K.W., Otto-Bliesner, B., Phillips, A.S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J.E., Kinnison, D., Kushner, P.J., Larson, V.E., Long, M.C., Mickelson, S., Moore, J.K., Nienhouse, E., Polvani, L., Rasch, P.J., and Strand, W.G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 285 Dalaiden, Q., Goosse, H., Klein, F., Lenaerts, J. T. M., Holloway, M., Sime, L., and Thomas, E. R.: How useful is snow accumulation in reconstructing surface air temperature in Antarctica? A study combining ice core records and climate models, *The Cryosphere*, 14, 1187–1207, <https://doi.org/10.5194/tc-14-1187-2020>, 2020.
- Dee, D.P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.:
290 The ERA-Interim Reanalysis: Configuration and Performance of the Data Assimilation system, *Quarterly Journal of Royal Meteorological Society*, Vol. 137, No. 656, 2011, pp. 553–597. doi:10.1002/qj.828, 2011.
- Deppeler, S.L., and Davidson, A.T.: Southern ocean phytoplankton in a changing climate, *Front. Mar. Sci.*, 4, 40, <https://doi.org/10.3389/fmars.2017.00040>, 2017.



- 295 Donat-Magnin, M., Jourdain, N.C., Gallée, H., Amory, C., Kittel, C., Fettweis, X., Wille, J.D., Favier, V., Drira, A., and Agosta, C.:
Interannual variability of summer surface mass balance and surface melting in the Amundsen sector, West Antarctica, Cryosphere,
<https://doi.org/10.5194/tc-2019-109>, 2019.
- Drews, R., Matsuoka, K., Martin, C., Callens, D., Bergeot, N., and Pattyn, F.: Evolution of Derwael Ice Rise in Dronning Maud Land,
Antarctica, over the last millenia, *J. Geophys. Res. Earth Surf.*, 120, <https://doi.org/10.1002/2014JF003246>, 2015.
- Eisen, O., Frezzotti, M., Genthon, C., Isaksson, E., Magand, O., van den Broeke, M.R., Dixon, D.A., Ekaykin, A., Holmlund, H. P., Takao
300 Kameda, T., Karlöf, L., Kaspari, S., Lipenkov, V.Y., Oerter, H., Takahashi, S., and Vaughan, D.G.: Ground-based measurements of spatial
and temporal variability of snow accumulation in East Antarctica, *Rev. Geophys.*, 46, RG2001, <https://doi.org/10.1029/2006RG000218>,
2008.
- Favier, V., Agosta, C., Parouty, S., Durand, G., Delaygue, G., Gallée, H., Drouet, A.-S., Trouvilliez, A., and Krinner, G.: An updated and
quality controlled surface mass balance dataset for Antarctica, *The Cryosphere*, 7, 583–597, <https://doi.org/10.5194/tc-7-583-2013>, 2013.
- 305 Feudale, L., and Tompkins, A.M.: A simple bias correction technique for modeled monsoon precipitation applied to West Africa, *Geophysical
Research Letters*, 38, L03803, <https://doi.org/10.1029/2010GL045909>, 2011.
- Frezzotti, M., Scarchilli, C., Becagli, S., Proposito, M., and Urbini, S.: A synthesis of the Antarctic surface mass balance during the last 800
yr, *The Cryosphere*, 7, 303-319, <https://doi.org/10.5194/tc-7-303-2013>, 2013.
- Gallée, H., Agosta, C., Genial, L., Favier, V., and Krinner, G.: A downscaling approach toward high-resolution surface mass balance over
310 Antarctica, *Surv. Geophys*, 32, 507-518, 2011.
- Gallée, H., Trouvilliez, A., Agosta, C., Genthon, C., Favier, V., and Naaim-Bouvet, F.: Transport of snow by the wind: a comparison between
observations in Adelie Land, Antarctica, and simulations made with the regional climate model MAR, *Bound.-Lay. Meteorol.*, 146, 133-
147, <https://doi.org/10.1007/s10546-012-9764z>, 2013.
- Garbe J., Albrecht T., Levermann A., et al, (2020), The hysteresis of the Antarctic Ice Sheet, *Nature*, 85, 538 – 544,
315 <https://doi.org/10.1038/s41586-020-2727-5>
- Ghilain, N., Vannitsem, S., Dalaiden, Q., Goosse, H., and De Cruz, L.: MASS2ANT Snowfall Dataset (Downscaling @5.5km over Dronning
Maud Land, Antarctica, 1850 - 2014), doi:10.5281/zenodo.4287517, 2021.
- Goel, V., Matsuoka, K., Berger, C., Lee, I., Dall, J., and Forsberg, R.: Characteristics of ice rises and ice rumples in Dronning Maud Land
and Enderby Land, Antarctica, *Journal of Glaciology*, 66(260), 1064-1078, <https://doi.org/10.1017/jog.2020.77>, 2020.
- 320 Gorodetskaya, I.V., Silva, T., Schmithüsen, H., and Hirasawa, H.: Atmospheric River Signatures in Radiosonde Profiles and Reanalyses at
the Dronning Maud Land Coast, East Antarctica, *Adv. Atmos. Sci.*, 37, 455–476, <https://doi.org/10.1007/s00376-020-9221-8>, 2020.
- Greene, C.A., Gwyther, D.E., and Blankenship, D.D.: Antarctic mapping tools for matlab, *Computers & Geosciences*, 104, 151-157,
<https://doi.org/10.1016/j.cageo.2016.08.003>, 2017.
- He, X., Chaney, N.W., Schleiss, M., and Sheffield, J.: Spatial downscaling of precipitation using adaptable random forests, *Water Resour.
325 Res.*, 52, 8217-8237, <https://doi.org/10.1002/2016WR019034>, 2016.
- Hannachi, A.: A primer for EOF analysis of Climate Data, 33p., <http://www.o3d.org/eas-6490/lectures/EOFs/eofprimer.pdf>, 2004.
- Hersbach, H., Bell, B., Berrisford, P., et al.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999 - 2049,
<https://doi.org/10.1002/qj.3803>, 2020.
- Hochman, A., Kunin, P., Alpert, P., Harpaz, T., Saaroni, H., and Rotskier-Edelstein, D.: Weather regimes and analogues of seasonal precipi-
330 tation for the 21st century: A case study over Israel, *Int. J. Clim.*, <https://doi.org/10.1002/joc.6318>, 2019.



- Horton, P., Obled, Ch., Jaboyedoff, M.: The analogue method for precipitation prediction: finding better analogue situations at sub-daily time step, *Hydrol. Earth Syst. Sci.*, 21, 3307-3323, 2017.
- Isaksson, E., Karlen W., Gundestrup, N., Mayewski, P., Whitlowand, S., and Twickler, M.; A century of accumulation and temperature changes in Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 101(D3), 7085–7094, 1996.
- 335 Isaksson, E., van den Broeke, M.R., Winther, J.-G., Karlof, L., Pinglot, J. F., and Gundestrup, N.: Accumulation and proxy-temperature variability in Dronning Maud Land, Antarctica, determined from shallow firn cores, *Ann. Glaciol.*, 29, 17–22, 1999.
- Kaczmarek, M., Isaksson, E., Karlof, L., Winther, J.-G., Kohler, J., Godtlielsen, F., Olsen, L., Hofstede, C., van den Broeke, M., Wal, R.S.W., and Gundestrup, N.: Accumulation variability derived from an ice core from coastal Dronning Maud Land, Antarctica, *Annals of Glaciology*, 39, <https://doi.org/10.3189/172756404781814186>, 2004.
- 340 Kausch, T., Lhermitte, S., Lenaerts, J. T. M., Wever, N., Inoue, M., Pattyn, F., Sun, S., Wauthy, S., Tison, J.-L., and van de Berg, W. J.: Impact of coastal East Antarctic ice rises on surface mass balance: insights from observations and modeling, *The Cryosphere*, <https://doi.org/10.5194/tc-2020-66>, 2020.
- Krinner, G., Lhermitte, S., Ménégoz, M., Agosta, C., and Brutel-Vuilmet, C.: Oceanic Forcing of Antarctic Climate Change: A Study Using a Stretched-Grid Atmospheric General Circulation Model, *Journal of Climate*, 27(15), 5786-5800, 2014.
- 345 Kurita, N., Hirasawa, N., Koga, S., Matsuhita, J., Steen-Larsen, H. C., Masson-Delmotte, V., and Fujiyoshi, Y.: Identification of air masses responsible for warm events on the East Antarctic Coast, *SOLA*, 12, 307-313, <https://doi.org/10.2151/sola.2016-060>, 2016.
- Lenaerts, J.T.M., van den Broeke, M.R., van de Berg, W.J., van Meijgaard, E., and Kuipers Munneke, P.: A new high-resolution surface mass balance map of Antarctica (1979–2010) based on regional atmospheric climate modeling, *Geophysical Research Letters*, 39, L04501/1 - L04501/5, 2012.
- 350 Lenaerts, J. T. M., van Meijgaard, E., van den Broeke, M. R., Ligtenberg, S. R. M., Horwath, M., and Isaksson, E.: Recent snowfall anomalies in Dronning Maud Land, East Antarctica, in a historical and future climate perspective, *Geophys. Res. Lett.*, 40, 2684 - 2688, doi:10.1002/grl.50559, 2013.
- Lenaerts, J.T.M., Brown, J., van den Broeke, M.R., Matsuoka, K., Drews, R., Callens, D., Philippe, M., Gorodetskaya, I.V., van Meijgaard, E., Reijmer, C.H., Pattyn, F., and van Lipzig, N.P.M.: High variability of climate and surface mass balance induced by Antarctic ice rises, *J Glaciology*, 60, 224, 1101-1110, <https://doi.org/10.3189/2014JG14J040>, 2014.
- 355 Lenaerts, J., Ligtenberg, S., Medley, B., van de Berg, W., Konrad, H., Nicolas, J., Van Wessem, J. M., Trusel, L.D., Mulvaney, R., Tuckwell, R. J., Hogg, A.E., and Thomas, E.: Climate and surface mass balance of coastal West Antarctica resolved by regional climate modelling. *Annals of Glaciology*, 59, 29-41, doi:10.1017/aog.2017.42, 2018.
- Lenaerts, J.T., Medley, B., van den Broeke, M.R., Wouters, B.: Observing and modeling ice sheet surface mass balance, *Reviews of Geophysics*, 57, <https://doi.org/10.1029/2018RG000622>, 2019.
- 360 Liang, Y.-C., Mazloff, M.R., Rosso, I., Fang, S.-W., and Yu, J.-Y.: A multivariate empirical orthogonal function method to construct nitrate maps in the Southern Ocean, *J. Atmos. Ocea. Tech.*, 35, 1505-1519, <https://doi.org/10.1175/JTECH-D-18-0018.1>, 2018.
- Maraun D. & Widmann M. (2018), *Statistical Downscaling and Bias Correction for Climate Research*, Cambridge University Press, doi:10.1017/9781107588783
- 365 Massom, R.A., Pook, M.J., Comiso, J.C., Adams, N., Turner, J., Lachlan-Cope, T., and Gibson, T.T.: Precipitation over the interior East Antarctic Ice sheet related to midlatitude blocking-high activity, *J. Clim.*, 17, 1914-1928, 2003.
- Matsuoka, K., Hindmarsh, R.C.A., Moholdt, G., Bentley, M.J., Pritchard, H.D., Brown, J., Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts, J.T.M., Martín, C., Mulvaney, R., Nicholls, K.W., Pattyn, F., Ross, N., Scambos, T. and



- Whitehouse, P.L.: Antarctic ice rises and rumples: Their properties and significance for ice-sheet dynamics and evolution, *Earth-Science Reviews*, 150, 724 - 745, <https://doi.org/10.1016/j.earscirev.2015.09.004>, 2015.
- 370 Medley, B., McConnell, J.R., Neumann, T.A., Reijmer, C.H., Chellman, N., Sigl, M., and Kipfstuhl, S.: Temperature and snowfall in Western Queen Maud Land increasing faster than climate model projections, *Geophysical Research Letters* 45, 1472-1480, 2018.
- Medley, B., and Thomas, E.R.: Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise, *Nature Climate Change*, 9, 34-39, <https://doi.org/10.1038/s41558-018-0356-x>, 2019.
- 375 Monaghan, A.J., Bromwich, D., Wang, S.-H.: Recent trends in Antarctic snow accumulation from Polar MM5 simulations, *Phil. Trans. R. Soc.*, 364, 1683-1708, <https://doi.org/10.1098/rsta.2006.1795>, 2006.
- Mottram, R., Hansen, N., Kittel, C., van Wessem, M., Agosta, C., Amory, C., Boberg, F., van de Berg, W. J., Fettweis, X., Gossart, A., van Lipzig, N. P. M., van Meijgaard, E., Orr, A., Phillips, T., Webster, S., Simonsen, S. B., and Souverijns, N.: What is the Surface Mass Balance of Antarctica? An Intercomparison of Regional Climate Model Estimates, *The Cryosphere Discuss.*, [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-2019-333)
380 2019-333, 2020.
- Nishio, F., Furukawa, T., Hashida, G., Igarashi, M., Kameda, T., Kohno, M., Motoyama, H., Naoki, K., Satow, K., Suzuki, K., Morimasa, T., Toyama, Y., Yamada, T., and Watanabe, O.: Annual-layer determinations and 167 year records of past climate of H72 ice core in east Dronning Maud Land, Antarctica, *Ann. Glaciol.*, 35, 471-479, 2002.
- Noone, D., Turner, J., and Mulvaney, R.: Atmospheric signals and characteristics of accumulation in Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 104, D16, 19191-19211, 1999.
- 385 Oerter, H., Wilhelms, F., Jung-Rothenhäusler, F., Göktas, F., Miller, H., Graf, W., and Sommer, S.: Accumulation rates in Dronning Maud Land, Antarctica, as revealed by dielectric-profiling measurements of shallow firn cores, *Annals of Glaciology*, 30, 27-34, <https://doi.org/10.3189/172756400781820705>, 2000.
- Palermo, C., Claud, C., Dufour, A., Genthon, C., Wood, N.B., and L'Ecuyer, T.: Evaluation of Antarctic snowfall in global meteorological reanalyses, *Atmos. Res.*, 190, 104-112, <https://doi.org/10.1016/j.atmosres.2017.02.015>, 2017.
- 390 Paolo, F.S., Fricker, H.A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 6232, 327-331, <https://doi.org/10.1126/science.aaa0940>, 2015.
- Philippe, M., Tison, J.-L., Fjosne, K., Hubbard, B., Kjaer, H.A., Lenaerts, J.T.M., Drews, R., Sheldon, S.G., De Bondt, K., Claeys, P., and Pattyn, F.: Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica, *The Cryosphere*, 10, 2501-2516, <https://doi.org/10.5194/tc-10-2501-2016>, 2016.
- 395 Reijmer, C.H., and van den Broeke, M.R.: Temporal and spatial variability of the surface mass balance in Dronning Maud Land, Antarctica, *J. Glaciol.*, 49(167), 512-520, doi:10.3189/172756503781830494, 2003.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M.J., Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979-2017, *PNAS*, 116, 4, 1095-1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- 400 Rodwell, M.J., Ferranti, L., Haiden, T., Magnusson, L., Bidlot, J., Bormann, N., Dahoui, M., De Chiara, G., Duffy, S., Forbes, R., Holm, E., Ingleby, B., Janousek, M., Lang, S.T.K., Morgensen, K., Prates, F., Rabier, F., Richardson, D.S., Tsonevsky, I., Vitart, F., and Yamaguchi M.: New developments in the diagnosis and verification of high-impact weather forecasts, *ECMWF Technical Memorandum 759*, November 2015, 1-44, 2015.
- Rotschky, G., Holmlund, P., Isaksson, E., Mulvaney, R., Oerter, H., van den Broeke, M.R., and Winther, J.-G.: A new surface accumulation map for western Dronning Maud Land, Antarctica, from interpolation of point measurements, *Journal of Glaciology*, 53, 182, 385-398, <https://doi.org/10.3189/002214307783258459>, 2007.
- 405



- Russel, A., McGregor, G.R., and Marshall, G.J.: 340 years of atmospheric circulation characteristics reconstructed from an eastern Antarctic Peninsula ice core, *Geophys. Res. Lett.*, 33, L08702, <https://doi.org/10.1029/2006GL025899>, 2006.
- Russel, A., McGregor, G.R., and Marshall, G.J.: Eastern Antarctic peninsula precipitation delivery mechanisms: process studies and back trajectory evaluation, *Atmos. Sci. Lett.*, 9, 214-221, <https://doi.org/10.1002/asl.190>, 2008.
- 410 Scarchilli, C., Frezzotti, M., and Ruti, P.M.: Snow precipitation at four ice core sites in East Antarctica: provenance, seasonality and blocking factors, *Clim. Dyn.*, 37, 2107-2125, <https://doi.org/10.1007/s00382-010-0946-4>, 2011.
- Schlosser, E.: Effects of seasonal variability of accumulation on yearly mean $\delta^{18}\text{O}$ values in Antarctic snow, *J. Glaciol.*, 45, 151, 463-468, 1999.
- 415 Schlosser, E., Manning, K.W., Powers, J.G., Duda, M.G., Birnbaum, G., and Fujita, K.: Characteristics of high-precipitation events in Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 115, D14107, <https://doi.org/10.1029/2009JD013410>, 2010.
- Schlosser, E., Anshütz, H., Divine, D., Martma, T., Sinisalo, A., Altnau, S., and Isaksson, E.: Recent climate tendencies on an East Antarctic ice shelf inferred from a shallow firn core network, *J. Geophysic. Res.*, 119, 11, 6549-6562, doi:10.1002/2013JD020818, 2014.
- Schlosser, E., Stenni, B., Valt, M., Cagnati, A., Powers, J.G., Manning, K.W., Raphael, M., and Duda, M.G.: Precipitation and synoptic regime in two extreme years 2009 and 2010 at Dome C, Antarctica – implications for ice core interpretation, *Atmos. Chem. Phys.*, 16, 4757-4770, 2016.
- 420 Sellevold, R., van Kampenhout, L., Lenaerts, J.T.M., Noël, B., Lipscomb, W.H., and Vizcaino, M.: Surface mass balance downscaling through elevation classes in an Earth system model: analysis, evaluation and impacts on the simulated climate, *The Cryosphere, Discussion*, <https://doi.org/10.5194/tc-2019-122>, 2019.
- 425 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A, G., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V., Blazquez, A., Bonin, J., Csatho, B., Cullather, R., Felikson, D., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert, L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mouginit, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noel, B., Ootaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg-Sorensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W.J., van der Wal, W., van Wessel, M., Vishwakarma, B.D., Wiese, D., Wouters, B.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 430 Sneyers, R., and Goossens, Chr.: L'analyse par la methode des composantes principales: application à la climatologie et à la météorologie, Annexe au rapport du rapporteur pour les méthodes statistiques présenté à la 9ème session de la Commission de climatologie, Genève, décembre 1985, 1-59, 1988.
- Souvereinjs, N., Gossart, A., Gorodetskaya, I.V., Lhermitte, S., Mangold, A., Laffineur, Q., Delcloc, A., and van Lipzig, N.P.M.: How does the ice sheet surface mass balance relate to snowfall? Insights from a ground-based precipitation radar in East Antarctica, *The Cryosphere*, 12, 1987-2003, <https://doi.org/10.5194/tc-12-1987-2018>, 2018.
- 440 Sparnocchia, S., Pinardi, N., and Demirov, E.: Multivariate empirical orthogonal function analysis of the upper thermocline structure of the Mediterranean sea from observations and model simulations, *Annales Geophysicae*, 21, 167-187, 2003..
- Stein, A.F., Draxler, R.R., Rolph, G.D., Stunder, B.J.B., Cohen, M.D., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *BAMS*, 2059-2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.



- 445 Tetzner, D., Thomas, L., and Allen, C.: A validation of ERA5 reanalysis data in the southern Antarctic Peninsula – Ellsworth Land region, and its implications for ice core studies, *Geosciences*, 9, <https://doi.org/10.3390/geosciences9070289>, 2019.
- Thomas, E. R., Marshall, G. J., and McConnell, J. R.: A doubling in accumulation in the western Antarctic Peninsula since 1850, *Geophys. Res. Lett.*, 35, L01706, doi:10.1029/2007GL032529, 2008.
- Thomas, E.R., van Wessem, J.M., Roberts, J., Isaksson, E., Schlosser, E., Fudge, T.J., Vallelonga, P., Medley, B., Lenaerts, J., Bertler, N., van den Broeke, M.R., Dixon, D.A., Frezzotti, M., Stenni, B., Curran, M., and Ekaykin, A.A.: Regional Antarctic snow accumulation over the
450 past 1000 years, *Clim. Past*, 13, 1491-1513, <https://doi.org/10.5194/cp-13-1491-2017>, 2017.
- Turner, J., Phillips, T., Thamban, M., Rahaman, W., Marshall, G.J., Wille, J.D., Favier, V., Holly, V., Winton, L., Thomas, E., Wang, Z., van den Broeke, M., Scott Hosking, J., and Lachlan-Cope, T.: The dominant role of extreme precipitation events in Antarctic Snowfall variability, *Geophys Res. Letters*, 46, 3502-3511, <https://doi.org/10.1029/2018GL081517>, 2019.
- Van De Berg, W., Van Den Broeke, M., Reijmer, C., and Van Meijgaard, E.: Characteristics of the Antarctic surface mass balance, 1958–2002,
455 using a regional atmospheric climate model, *Annals of Glaciology*, 41, 97-104, doi:10.3189/172756405781813302, 2005.
- van de Berg, W. J., and Medley, B.: Brief Communication: Upper-air relaxation in RACMO2 significantly improves modelled interannual surface mass balance variability in Antarctica, *The Cryosphere*, 10, 459–463, <https://doi.org/10.5194/tc-10-459-2016>, 2016.
- van Wessem, J. M., Ligtenberg, S. R. M., Reijmer, C. H., van de Berg, W. J., van den Broeke, M. R., Barrand, N. E., Thomas, E. R., Turner, J., Wuite, J., Scambos, T. A., and van Meijgaard, E.: The modelled surface mass balance of the Antarctic Peninsula at 5.5 km horizontal
460 resolution, *The Cryosphere*, 10, 271–285, <https://doi.org/10.5194/tc-10-271-2016>, 2016.
- Vannitsem, S., Dalaiden, Q., and Goosse, H.: Testing for dynamical dependence: Application to the surface mass balance over Antarctica, *Geophysical Research Letters*, 46, 12,125-12,135, <https://doi.org/10.1029/2019GL084329>, 2019.
- Vega, C.P., Schlosser, E., Divine, D.V., Kohler, J., Martma, T., Eichler, A., Schwikowski, M., and Isaksson, E.: Surface mass balance and water stable isotopes derived from firn cores on three ice rises, Fimbul Ice Shelf, Antarctica, *The Cryosphere*, 10, 2763-2777,
465 <https://doi.org/10.5194/tc-10-2763-2016>, 2016.
- Wang, Y., Ding, M., van Wessem, J. M., Schlosser, E., Altnau, S., van den Broeke, M. R., Lenaerts, J. T. M., Thomas, E. R., Isaksson, E., Wang, J., and Sun, W.: A Comparison of Antarctic Ice Sheet Surface Mass Balance from Atmospheric Climate Models and In Situ Observations, *Journal of Climate*, 29(14), 5317-5337, 2016.
- Weare, B.C., and Nasstrom, J.S.: Examples of extended empirical orthogonal function analysis, *Monthly Weather Review*, 110, 481-485, 1982.
- 470 Welker, C., Martius, O., Froidevaux, P., Reijmer, C. H., and Fischer, H.: A climatological analysis of high-precipitation events in Dronning Maud Land, Antarctica, and associated large-scale atmospheric conditions, *J. Geophys. Res. Atmos.*, 119, 11,932 – 11,954, <https://doi.org/10.1002/2014JD022259>, 2014.
- Yu, Y., Lin, Z.-H., and Qin, Z.-K.: Improved EOF-based bias correction method for seasonal forecasts and its application in IAP AGCM4.1, *Atmospheric and Oceanic Science Letters*, 11:6, 499-508, <https://doi.org/10.1080/16742834.2018.1529532>, 2018.