



Arctic sea surface height maps from multi-altimeter combination

Pierre Prandi¹, Jean-Christophe Poisson², Yannice Faugère¹, Amandine Guillot³, and Gerald Dibarboure³

¹CLS, Ramonville, France

²VorteX-Io, Toulouse France, previously at CLS

³CNES, Toulouse France

Correspondence: Pierre Prandi (pprandi@groupcls.com)

Abstract. We present a new Arctic sea level anomaly dataset, based on the combination of three altimeter missions using an optimal interpolation scheme. Measurements from SARAL/AltiKa, CryoSat-2 and Sentinel-3A are blended together providing an unprecedented resolution for this type of products. The final gridded fields cover all latitudes north of 50°N, on a 25 km EASE2 grid, with one grid every three days over three years from July 2016 to April 2019. We use the Adaptive retracker to process both open ocean and lead echoes on SARAL/AltiKa thus removing the need to estimate a bias between open ocean an
5 ice covered areas. SARAL/AltiKa also provides the baseline for the cross-calibration of CryoSat-2 and Sentinel-3A data. When compared to independent data, the combined product exhibits a much better performance than previously available datasets based on the analysis of a single mission.

10 1 Introduction

All components of the Arctic are undergoing large climate changes (Meredith et al., 2019). Among them the Arctic sea ice is certainly the most striking, with dramatic extent (Stroeve and Notz, 2018), thickness and volume losses (e.g. Kwok, 2018). Physical characteristics of the Arctic Ocean are also changing. Ocean temperature is increasing both in the mixed layer (Timmermans et al., 2017) and at depths (e.g. Polyakov et al., 2017). Regarding salinity, the Beaufort Gyre region is freshening
15 (Proshutinsky et al., 2015) while freshwater content decline is reported in other parts of the basin (Armitage et al., 2016). Changes in the Arctic Ocean circulation are also documented, with a strengthening of surface geostrophic currents (Armitage et al., 2017) or intensification of eddy activity (Zhao et al., 2016) in some parts of the basin.

Despite those pressing matters and due to harsh conditions the Arctic Ocean remains poorly observed (Smith et al., 2019). In this context remote sensing, and satellite altimetry in particular is of great interest. While satellite altimetry was designed to measure the global ocean circulation (Stammer and Cazenave, 2018), it is also used in the Arctic Ocean to retrieve sea level
20 (SL) and sea ice freeboard (Quartly et al., 2019). First estimates of Arctic SL variability from satellite altimetry were made by Peacock and Laxon (2004). The same methodology was successfully used by Giles et al. (2012) to estimate freshwater



variations in the Beaufort Gyre. More recently a state of the art Arctic SL dataset (Armitage et al., 2016) has been used to estimate the shape and extent of the Beaufort Gyre (Regan et al., 2019).

25 Current state-of-the-art Arctic SL datasets (Armitage et al., 2016; Rose et al., 2019) rely on the measurements of one altimeter at a time to produce SL maps with monthly temporal and 1000km spatial resolutions, which are not enough to characterize the mesoscale activity (Regan et al., 2020). The combination of several altimeters can improve the resolution of global SL maps (Ducet et al., 2000; Pascual et al., 2006). This multi-mission combination is operational in the DUACS system (Pujol et al., 2016; Taburet et al., 2019). In this paper we present a new Arctic SL dataset based on the combination of
30 three satellite altimetry missions: SARAL/AltiKa (SRL), Sentinel-3A (S3A) and CryoSat-2(C2). Section 2.2 details the along-track data processing scheme as well as the standards used. The multi-mission cross-calibration and combination methods are described in section 2.3. Results of the data quality assessment are presented in section 4.

2 Data and methods

SL estimation in the ice-covered Arctic Ocean relies on the identification of radar waveforms originating from leads (cracks) in the ice pack, where the ocean surfaces (Quartly et al., 2019). All groups (Armitage et al., 2016; Rose et al., 2019) follow the same general workflow, but processing details can be different. In this section we describe our regional Arctic sea level processing: data sources (2.1), echo classification (2.2.1) and retracking (2.2.2), lead selection (2.2.3), geophysical corrections (2.2.4), data editing (2.2.5) and mapping (2.3).

2.1 Satellite radar altimetry missions

40 We processed measurements from three satellite altimetry missions: SARAL/AltiKa, Sentinel-3A and CryoSat-2. Some relevant characteristics of these missions are summarized in table 1. A brief description of each mission is given below.

2.1.1 SARAL/AltiKa

SARAL/AltiKa (herein abbreviated SRL) is a joint French (CNES) and Indian (ISRO) satellite radar altimetry mission (Verron et al., 2015). Its main instrument is a Ka-band pulse limited radar altimeter. The higher frequency compared to Ku-band
45 missions, combined with a higher pulse repetition frequency permits a higher along-track sampling. A better range resolution is also achieved thanks to a larger bandwidth (Steunou et al., 2015). SRL was launched in February 2013 and is still operational today. Initially launched on the Envisat orbit it provides measurements up to 81.5° latitude. SRL data used in this study are taken from the CNES PEACHI project dataset (Valladeau et al., 2015).

2.1.2 CryoSat-2

50 CryoSat-2 (herein abbreviated C2) was launched in April 2010 and is an ESA satellite radar altimetry mission designed to monitor the Earth's cryosphere (Wingham et al., 2006). Polar regions are observed up to 88° thanks to its 92° orbit inclination. The SIRAL radar onboard CryoSat-2 can operate in low resolution mode (LRM), synthetic aperture mode (SAR) or synthetic



aperture interferometric mode (SARIn). The switch from one mode to another is based on a geographical mode mask. SAR mode is generally used over sea-ice areas and provides an unprecedented along-track resolution of 300m. In this study, only SAR mode data from C2 is used and the area covered therefore varies over time with the geographical mode mask. C2 data used in this study are L1b data taken from ESA PDGS Ice Baseline C processor. The PDGS Ice processor includes 0 padding and Hamming windowing which reduces the impact of antenna side lobes and increases the range resolution over peaky echoes (Smith and Scharroo, 2015).

2.1.3 Sentinel-3A

Sentinel-3A (herein abbreviated S3A) is a Copernicus mission providing sea surface topography measurements (among other variables) thanks to its SRAL SAR mode altimeter (Donlon et al., 2012). S3A provides measurements up to 81.5° latitudes. Compared to C2, S3A always operates in SAR mode, whether over open ocean or ice covered areas. S3A was launched in February 2016 with an expected lifetime of 7 years. For S3A no operational ground segments implements 0 padding and Hamming windowing (Lawrence et al., 2019). We therefore rely on data from the CNES S3PP processing prototype which does include these algorithms.

2.2 Along-track data processing

2.2.1 Waveform classification

Waveform classification aims at separating radar altimetry waveforms based on their shape to identify echoes from leads, floes and open ocean. In Arctic SL studies, classification generally relies on the pulse peakiness (Peacock and Laxon, 2004, e.g.): peaky echoes are associated to leads. Here we use the neural network based classification method proposed by Poisson et al. (2018). A dedicated classifier is used for each mission. This classification methods provides a wealth of information with 16 output classes. Classification outputs were validated against coincident SAR images from Sentinel-1 by Longépé et al. (2019), especially for sea-ice leads detection. For the purpose of this study we only select echoes labeled as class 1 (brownian echoes, associated to open ocean) and class 2 (peaky echoes, associated to sea ice leads). For C2, the geographical SAR mode mask varies over time to match sea ice extent. And since we only process C2 SAR mode data, only a very small fraction of the open ocean is observed. These data (Brownian C2 echoes) are discarded from the analysis.

2.2.2 Retracking

Retracking designates the process of extracting geophysical parameters from the radar waveforms. There are a variety a re-tracking algorithms, from purely empirical algorithms to physical ones which require a waveform model. In this study several retracking algorithms are used, depending on the measurements mode (LRM or SAR) and echo type (brownian or peaky). On SRL, all echoes are retracked by the Adaptive algorithm (Poisson et al., 2018). This physical retracker provides a processing continuity between open ocean and ice covered areas thus removing the need to estimate a bias between the two surfaces. This represents an important difference with respect to other Arctic SL datasets (Armitage et al., 2016) where different retracking



algorithms are used to process open ocean and leads. S3A operates in SAR mode and there is currently no equivalent of the
85 Adaptive retracking algorithm available. We therefore fallback to using different retracking algorithms over sea ice and open
ocean areas: peaky echoes are retracked by the TFMRA algorithm (50% threshold, Helm et al. (2014)) while brownian echoes
are retracked using an ocean MLE3 algorithm. On C2 only peaky echoes are kept in the analysis, they are retracked using the
TFMRA algorithm (50% threshold).

2.2.3 Ocean/Lead selection

90 After waveform classification and retracking, the ocean/lead selection algorithm is applied. This algorithm is based on three pa-
rameters: sea ice concentration (SIC) taken from OSI-450 product (Lavergne et al., 2019), waveform class and radar backscatter
coefficient. All class 1 (brownian) echoes in areas where SIC is lower than 30% are considered to represent the open ocean. All
class 2 echoes with enough backscattered power over areas where SCI is greater than 30% are considered to be lead echoes.
Backscattering distributions differ for each mission and therefore backscattering thresholds are different for SRL, S3A and C2.
95 Thresholds used in this study are given in table 2.

Leads act as bright targets in the radar footprint and their contribution tend to be dominant, even when they represent a
small fraction of the radar footprint. This can lead to the retracker following off-nadir targets and biasing range estimates.
This effect is called "snagging" (Peacock and Laxon, 2004) or "hooking" (Boergens et al., 2016). SRL, as an LRM altimeter,
is more prone to this phenomenon than S3A and C2. Here we use the method proposed by Poisson et al. (2018) to remove
100 measurements identified as leads that may be affected by hooking. SAR altimetry is less sensitive to such effects due to the
much smaller along-track resolution and no hooking flag is applied on S3A and C2. Note that this does not account for cross-
track hooking errors which were investigated by Armitage and Davidson (2014) resulting in an estimated -1 to -4 cm bias
on ocean topography itself and are neglected in the present study.

2.2.4 Sea level anomaly estimation

105 Once relevant radar echoes are selected and retracked to estimate the radar range, one can estimate sea levels. Classically sea
level is estimated following equation 1 (e.g. Chelton et al., 2001).

$$SLA = Orbit - Range - \sum Corrections - MSS \quad (1)$$

Where the corrections account for a range of geophysical and instrumental effects. The standards and models used in this
study are given in table 3. The corrections used are not uncommon and are mainly derived from DUACS processing (Taburet
110 et al., 2019), with three notable exceptions:

- we use the DTU15 mean sea surface (and, 2016), which thanks to its use of C2 is defined over the whole Arctic domain,
- over leads, the sea state bias correction is set to zero, we consider that over these small open water stretches, waves are small,



– the radiometer wet tropospheric correction is undefined over sea ice, and we therefore use a modeled wet tropospheric
115 correction over the whole product domain.

2.2.5 Data editing

The data editing is a crucial step for final product quality and is always the result of a balance between data quality and
coverage. The most fundamental editing is the ocean/lead selection algorithm described above, yet this may retain erroneous
measurements. Here we tried to remove obvious outliers while retaining as many measurements as possible to build the final
120 SL product. First we apply a basic thresholding to remove any SL anomalies greater than two meters (absolute deviation).
Over open ocean areas, an iterative editing process is applied, which is designed to remove outliers with respect to the local
along-track SL anomaly distribution. For this iterative editing to work continuous segments are required. This requirement is
not met over ice covered areas, and the iterative editing is not applied to lead measurements. Following Rose et al. (2019)
we also apply a statistical editing based on local SL anomaly variance levels derived from a coarse gridding (200km) of each
125 mission data. Measurements that are locally further than 2.5σ from a 3 month running mean are discarded. This methods takes
into account local SL variance levels, and the running mean prevents from systematically removing measurements at highs and
lows of the seasonal cycle.

2.2.6 Ocean/lead bias correction

S3A data are retracked by different algorithms over open ocean and leads. This introduces a discontinuity that must be corrected
130 empirically (Giles et al., 2012). Armitage et al. (2016) faced a similar issue and proposed a correction method comparing sea
ice an open ocean echoes near the sea ice edge. Using a similar methodology leads to an average lead/ocean bias of around 16
cm on S3A, with a large uncertainty. Using the Adaptive retracker on SRL means that a baseline is available to estimate the
lead/ocean bias on S3A. Comparing SRL and S3A over leads and ocean leads to a S3A lead/ocean bias estimate of 11 cm. This
emphasizes the importance of processing continuity for at least one mission, which can be used as a reference, to ensure final
135 product accuracy. As we only process leads echoes on C2, no lead/ocean bias estimation is required.

2.3 Multi-mission combination

After the operations described in section 2.2 we are left with an ensemble of valid SL anomaly measurements along the track
of three different satellite radar altimeters. While this is already enough to estimate mono-mission products (see section 3), our
goal here is to combine all three missions together to increase SL maps resolution. The methods used here are derived from the
140 DUACS processing (e.g. Le Traon and Dibarboure, 1999; Pascual et al., 2006) with adaptations to fit the Arctic Ocean.

2.3.1 Cross-calibration

Cross-calibration is designed to remove long-wavelength errors in along-track altimetry prior to the optimal interpolation. In
a typical global ocean processing inter-mission cross-calibration would be performed through empirical orbit error estimation



(Le Traon and Ogor, 1998). This requires the estimation of crossovers over the whole globe which is not possible here, as
145 we processed only areas north of 50°N. Moreover empirical orbit errors are not well constrained over the Arctic which is
surrounded by large continental areas where no crossovers are available. As a result a much simpler cross-calibration technique
is used here. Time dependent regional SL differences between C2 and SRL and S3A and SRL are estimated through box
averages. Time series of these regional differences are shown on figure 1. In both cases differences exhibit an annual pattern
with amplitude around one centimeter. We fit and remove a 1 year period sine wave to correct for the time-dependent part of
150 the intermission bias. Geographically dependent biases were also investigated: maps of average biases are shown on figure 2
for C2 and S3A with respect to SRL. Differences remain small (below 2.5 cm) except along the coasts, and at the sea ice edge
for C2. While some geographical patterns are observed with higher SLA values on C2 and S3A in the multi-year ice region,
they are not easily modeled and are left uncorrected.

2.3.2 Along-track filtering and subsampling

155 Before the optimal interpolation open-ocean along track SL anomaly measurements are filtered and sub-sampled to reach a
5Hz resolution. Using 5Hz measurements rather than the full altimeter resolution significantly reduces the processing time for
the optimal interpolation, with almost no impact on the estimated fields. Measurements over the ice covered Arctic Ocean are
left at the full resolution of the altimeter.

2.3.3 Optimal interpolation

160 We use an optimal interpolation (OI) scheme to map SL anomaly fields on a regular grid. OI is based on an inverse formulation,
first introduced in oceanography by Bretherton et al. (1976). The methodology used here is derived from the DUACS global
processing (Le Traon and Dibarboure, 1999; Ducet et al., 2000) with some adaptations to the Arctic Ocean which are described
below. The quality of interpolated fields depends on the accuracy of several prior fields such as signal variance, covariance
scales and error levels.

165 Signal variance and error levels were adapted from currently used global values (Taburet et al., 2019) to fit the Arctic. One
important adaptation is an updated prior for signal variance. The signal variance map currently used in the DUACS global
processing is shown on figure 4 (left) and shows large drop at latitudes inside the Arctic Ocean. An analysis of the CPOM
dataset (Armitage et al., 2016), or the CCI dataset (Rose et al., 2019) doesn't show the same pattern. Unrealistically low
variance levels will cause the interpolation to dampen signals during the mapping, which is unwanted. We estimate an updated
170 signal variance map by locally taking the maximum variance level among DUACS, CCI and CPOM data for latitudes greater
than 60°N. The resulting variance map is shown on figure 4 (right).

Two error terms control the error level on the input data: noise and long wavelength errors (LWE). Accurate noise levels
will prevent error artifacts to be interpreted as real signals during the interpolation. Both were tuned to account for regional
characteristics.

175 DUACS standard processing uses different noise levels for different missions, reflecting the actual level of noise of each
mission, plus the unobservable part of the ocean dynamics. While SRL and S3A show very low noise levels in the Arctic



Ocean, which are unrealistically optimistic, C2 defaults to a very high noise level in the Arctic Ocean, especially above 82°N, due to high MSS errors. While this is not a problem for DUACS, as these areas are rarely mapped due to sea-ice cover, this is not something we want to convey in our analysis. Here we construct two noise level files, based on the existing DUACS noise levels, depending on whether measurements are over open ocean or over sea-ice. The assumptions used build these estimates are:

- Over ocean, our measurements should be slightly noisier than the standard processing due to the modeled wet tropospheric correction and a less accurate mean sea surface model,
- Over sea-ice, noise levels should be even higher due to increased errors in geophysical corrections and in range retrieval from peaky waveforms,
- Noise levels should be scaled to account for the fact that we are using 5Hz or 20(40)Hz measurements depending on the surface.

For the open ocean, the noise level is derived from the SRL file (to avoid unrealistic values inside the basin) augmented to match C2 noise levels in open ocean. The ice covered noise level is derived from open ocean noise by adding 5 cm^2 everywhere and accounting for the use of high resolution measurements. Open and ice-covered ocean noise levels are shown on figure 5.

LWEs are designed to absorb correlated errors along-track, coming from errors in geophysical models such as the tide and DAC corrections. For this regional analysis, they should also be able to absorb residual orbit errors, which are not corrected due to the lack of a proper cross-calibration. Again we derive the LWE priors from existing DUACS files. We start from C2 LWE error, which exhibits the most variance at high latitudes and set a minimum LWE variance of 10 cm^2 for all latitudes greater than 68°N. The resulting LWE variance distribution, which is used for all three missions considered here is shown on figure 6.

Correlation scales are kept unchanged from the DUACS global processing and SL anomaly fields are interpolated onto a 25 km EASE2 grid (Brodzik et al., 2012, 2014) every 3 days.

3 Product description

The combined product is distributed as a single NetCDF file containing maps of SL anomalies and absolute dynamic topography (ADT). ADT fields are obtained by adding the DTU15 MDT (and, 2016) to SL anomaly fields.

We also provide three mono-mission products for SRL, C2 and S3A which are available over longer periods (see table 4) depending on input data availability at product generation time. These are the result of simple box-averages of along-track measurements, with no cross-calibration. Due to their mono-mission nature, they are only available at a lower resolution (75km, one month).



4 Data quality assessment

In this section we present some product validation results. Assessing the product accuracy is difficult in the Arctic Ocean as few independent validation data are available (Smith et al., 2019). Here rely on comparisons between mono-missions products (??), analysis of regional statistics (4.2), comparisons to the DUACS global product (4.3) and comparisons to tide gauges data
210 available in the basin (4.4) .

4.1 Mono-mission products comparisons

Time series of the regional average sea level in the Arctic Ocean from SRL, S3A and C2 are shown on figure 1 while maps of the average SL anomalies are shown on figure 2. All three missions exhibit very consistent behaviors considering SL variations over time, as well as geographical patterns, even before any type of cross-calibration is applied. This is already a
215 good indication that observed variations are not artifacts arising from errors in the data. Differences between missions remain small when considering regional average SL (see figure 1, right) with differences below 2 cm Differences in geographical patterns (see figure 3) are slightly larger with differences (up to 5 cm) found at the coast, near the sea ice edge, and to smaller extent in the thicker ice zones north of Greenland and the Canadian Arctic Archipelago.

4.2 Regional statistics

220 Maps of the mean and variance of SL in the Arctic Ocean, derived from the combined product are shown on figure 7. SL variance levels are consistent throughout the Arctic basin. This suggests a good product accuracy as high variance levels in the interior of the Arctic Ocean would be an indication of errors. The variance distribution is consistent with previous datasets (Armitage et al., 2016; Rose et al., 2019) with high variance levels along the Russian Arctic coasts. Such variance levels could result from continental shelf waves propagation (Danielson et al., 2020). Another prominent feature is the SL variance slight
225 drop above 81.5°N. This is expected, as in this area C2 is the only radar altimeter mission, and the product is therefore unable to reach the same resolution than at lower latitudes.

Figure 7 also shows the mean Arctic Ocean SL from the combined products. First, there is no large bias between the open ocean and the seasonally ice covered areas, suggesting that our lead detection and retracking algorithms perform well. A negative geometric patch is visible north of the Canadian Arctic Archipelago in the the so-called "Wingham box" where C2
230 has been operating in SARIn mode from April 2011 to July 2014 (geographical mode mask versions 3.2, 3.3 and 3.4). There are no SARIn data over the period considered here and this bias is certainly coming from the MSS. Another bias is clearly visible between the Labrador sea and Hudson Bay, and extends up north into the western parts of Baffin Bay. The origin of this pattern remains to be investigated: it could again be an error in the MSS, or an issue in our current classification and retracking methods in this area.



235 4.3 Comparisons to the DUACS global product

The Arctic regional product presented here focuses on SL estimation in ice-covered areas, and its accuracy over open ocean might be hindered by some processing choices such as using the modelled wet tropospheric correction, or the lack of a proper cross-calibration prior to the optimal interpolation. Comparing the Arctic Ocean product with the CMEMS global dataset (Taburet et al., 2019) is a way to assess that despite these processing choices, we still have an acceptable performance over
240 open ocean surfaces. To perform this comparison, CMEMS grids are bilinearly interpolated onto the 25 km EASE2 grid used for the Arctic Ocean product. The mean and variance of SLA differences are shown on figure 8. Largest differences are found in the interior of the Arctic Ocean, as expected. In this area, the CMEMS global product is largely inaccurate (all measurements affected by sea-ice are removed by the editing). In the permanently open ocean areas (North Atlantic Ocean, Pacific Ocean) the regional product appears biased with respect to DUACS, but with no geographically varying pattern. In these areas the variance
245 of SLA differences is generally low indicating a good agreement between both products. There is a strong bias gradient in the Atlantic Ocean, between Iceland and Norway, likely related to the different mean sea surfaces used in both products (DTU15 rather CNES/CLS15).

4.4 Comparisons to tide gauges

Tide gauges provide an independent measurements of SL variability, and are very valuable for the validation of satellite
250 altimetry products (e.g Cipollini et al., 2016). They are relatively scarce in the Arctic Ocean. Here we use data from the GLOSS/CLIVAR tide gauges dataset from the UHSLC (Caldwell et al., 2010). 27 stations are left after removal of records with obvious issues or large data gaps. Table 5 summarizes the correlations and RMSd of altimetry and in-situ SL differences for the combined product and for the three mono-mission products. For the combined product, the mean correlation across all stations is 0.78 and the mean rmsd is 5.3 *cm*, indicating an excellent agreement between tide gauges and co-located altimeter
255 data. The combined product performs better than any mono-mission product, indicating that the product accuracy benefits from the multi-mission combination.

To illustrate the level of agreement between altimetry and tide gauges, the Prudhoe Bay station records are shown on figure ???. Prudhoe Bay is one of the few stations providing high quality tide gauge data in an area that is seasonally ice covered. At this station, where the ocean is seasonally ice covered there is a good agreement between the tide gauge and the all satellite
260 radar altimeter products (mono-mission and combined). However, the combined product is able to capture the high frequency SL variability observed by the tide gauge much better than any single altimeter product.

We also use tide gauges records to estimate the effective temporal resolution of the altimeter product following Ballarotta et al. (2019). Let the product error be the altimeter minus tide gauges SL differences while the true signal is given by the tide gauge record. Then the product resolution corresponds to the frequency at which the error spectrum becomes greater than half
265 the signal spectrum. Results are summarized on figure 10 The spread of the spectrum ratio remains large, as the time span available is limited to three years. However the improvement in resolution from a single altimeter product to a combined one



is large. At best the resolution of the S3A-only product is around 3 months. For the combined product resolution can be as low as 1.5 months.

5 Conclusions

270 In this paper, we document a new Arctic Ocean sea level dataset based on the combination of measurements from 3 satellite radar altimetry missions: SARAL/AltiKa, CryoSat-2 and Sentinel-3A. The processing applied to those three missions is described, as well as the optimal interpolation scheme used to create sea level anomaly fields.

Combined sea level anomaly and absolute dynamic topography fields are available over three years (from mid 2016 to mid 2019) on a 25km grid, every three days, for all latitudes greater than 50°N. Comparisons to the DUACS global product suggest
275 that despite focusing on high latitudes and ice-covered areas, the product performs well in permanently open ocean areas at lower latitudes. Comparisons with tide gauges available in the Arctic Ocean show that the combined product is able to capture some of the high frequency sea level variability observed by tide gauges and generally performs better than any single-altimeter analysis. This unprecedented resolution may be useful for the characterization of small scale Arctic Ocean circulation features.

6 Data availability

280 The different gridded datasets described in this manuscript are freely available, after registration, on the AVISO website (<https://www.aviso.altimetry.fr/>) at <https://doi.org/10.24400/527896/a01-2020.001> (Prandi, 2020).

Author contributions. PP generated the dataset, JCP provided algorithms and advice for waveform processing. GD and YF helped with multi-mission combination. AG led the project for CNES. All authors contributed to the manuscript.

Competing interests. The authors declare no competing interests.

285 *Acknowledgements.* The work presented here was performed within the frame of the CNES Alti-Doppler Glaciologie contract. Radar altimeter data were provided by CNES (SARAL/AltiKA, Sentinel-3A) and ESA (CryoSat-2).



References

- 290 The DTU15 MSS (Mean Sea Surface) and DTU15LAT (Lowest Astronomical Tide) reference surface, ESA Living planet symposium 2016, Prague, Czech Republic, 2016.
- Armitage, T. W. K. and Davidson, M. W. J.: Using the Interferometric Capabilities of the ESA CryoSat-2 Mission to Improve the Accuracy of Sea Ice Freeboard Retrievals, *IEEE Transactions on Geoscience and Remote Sensing*, 52, 529–536, 2014.
- Armitage, T. W. K., Bacon, S., Ridout, A. L., Thomas, S. F., Aksenov, Y., and Wingham, D. J.: Arctic sea surface height variability and change from satellite radar altimetry and GRACE, 2003–2014, *Journal of Geophysical Research: Oceans*, 121, 4303–4322, 295 <https://doi.org/10.1002/2015JC011579>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JC011579>, 2016.
- Armitage, T. W. K., Bacon, S., Ridout, A. L., Petty, A. A., Wolbach, S., and Tsamados, M.: Arctic Ocean surface geostrophic circulation 2003–2014, *The Cryosphere*, 11, 1767–1780, <https://doi.org/10.5194/tc-11-1767-2017>, <https://tc.copernicus.org/articles/11/1767/2017/>, 2017.
- Ballarotta, M., Ubelmann, C., Pujol, M.-I., Taburet, G., Fournier, F., Legeais, J.-F., Faugère, Y., Delepouille, A., Chelton, D., Dibarboure, 300 G., and Picot, N.: On the resolutions of ocean altimetry maps, *Ocean Science*, 15, 1091–1109, <https://doi.org/10.5194/os-15-1091-2019>, <https://doi.org/10.5194/os-15-1091-2019>, 2019.
- Boergens, E., Dettmering, D., Schwatke, C., and Seitz, F.: Treating the Hooking Effect in Satellite Altimetry Data: A Case Study along the Mekong River and Its Tributaries, *Remote Sensing*, 8, <https://doi.org/10.3390/rs8020091>, <https://www.mdpi.com/2072-4292/8/2/91>, 2016.
- 305 Bretherton, F. P., Davis, R. E., and Fandry, C.: A technique for objective analysis and design of oceanographic experiments applied to MODE-73, *Deep Sea Research and Oceanographic Abstracts*, 23, 559–582, [https://doi.org/10.1016/0011-7471\(76\)90001-2](https://doi.org/10.1016/0011-7471(76)90001-2), [https://doi.org/10.1016/0011-7471\(76\)90001-2](https://doi.org/10.1016/0011-7471(76)90001-2), 1976.
- Brodzik, M., Billingsley, B., Haran, T., Raup, B., and Savoie, M.: Correction: Brodzik, M.J., et al. EASE-Grid 2.0: Incremental but Significant Improvements for Earth-Gridded Data Sets. *ISPRS International Journal of Geo-Information* 2012, 1, 32–45, *ISPRS International* 310 *Journal of Geo-Information*, 3, 1154–1156, <https://doi.org/10.3390/ijgi3031154>, <https://doi.org/10.3390/ijgi3031154>, 2014.
- Brodzik, M. J., Billingsley, B., Haran, T., Raup, B., and Savoie, M. H.: EASE-Grid 2.0: Incremental but Significant Improvements for Earth-Gridded Data Sets, *ISPRS International Journal of Geo-Information*, 1, 32–45, <https://doi.org/10.3390/ijgi1010032>, <https://doi.org/10.3390/ijgi1010032>, 2012.
- Caldwell, P. C., US DOC/NOAA/NESDIS > National Centers For Environmental Information, Merrifield, M. A., and Thompson, 315 P. R.: Sea level measured by tide gauges from global oceans as part of the Joint Archive for Sea Level (JASL) since 1846, <https://doi.org/10.7289/V5V40S7W>, <https://accession.nodc.noaa.gov/JIMAR-JASL>, 2010.
- Carrère, L. and Lyard, F.: Modeling the barotropic response of the global ocean to atmospheric wind and pressure forcing - comparisons with observations, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2002gl016473>, <https://doi.org/10.1029/2002gl016473>, 2003.
- 320 Carrère, L., Lyard, F., Cancet, M., Guillot, A., and Picot, N.: FES 2014, a new tidal model - Validation results and perspectives for improvements, 2016.
- Cartwright, D. E. and Edden, A. C.: Corrected Tables of Tidal Harmonics, *Geophysical Journal International*, 33, 253–264, <https://doi.org/10.1111/j.1365-246x.1973.tb03420.x>, <https://doi.org/10.1111/j.1365-246x.1973.tb03420.x>, 1973.




- 325 Chelton, D. B., Ries, J. C., Haines, B. J., Fu, L.-L., and Callahan, P. S.: Chapter 1 Satellite Altimetry, in: *International Geophysics*, pp. 1–ii, Elsevier, [https://doi.org/10.1016/s0074-6142\(01\)80146-7](https://doi.org/10.1016/s0074-6142(01)80146-7), [https://doi.org/10.1016/s0074-6142\(01\)80146-7](https://doi.org/10.1016/s0074-6142(01)80146-7), 2001.
- Cipollini, P., Calafat, F. M., Jevrejeva, S., Melet, A., and Prandi, P.: Monitoring Sea Level in the Coastal Zone with Satellite Altimetry and Tide Gauges, *Surveys in Geophysics*, 38, 33–57, <https://doi.org/10.1007/s10712-016-9392-0>, <https://doi.org/10.1007/s10712-016-9392-0>, 2016.
- 330 Danielson, S. L., Hennon, T. D., Hedstrom, K. S., Pnyushkov, A. V., Polyakov, I. V., Carmack, E., Filchuk, K., Janout, M., Makhotin, M., Williams, W. J., and Padman, L.: Oceanic Routing of Wind-Sourced Energy Along the Arctic Continental Shelves, *Frontiers in Marine Science*, 7, <https://doi.org/10.3389/fmars.2020.00509>, <https://doi.org/10.3389/fmars.2020.00509>, 2020.
- Desai, S., Wahr, J., and Beckley, B.: Revisiting the pole tide for and from satellite altimetry, *Journal of Geodesy*, 89, 1233–1243, <https://doi.org/10.1007/s00190-015-0848-7>, <https://doi.org/10.1007/s00190-015-0848-7>, 2015.
- 335 Donlon, C., Berruti, B., Buongiorno, A., Ferreira, M.-H., FÃ©mÃ©nias, P., Frerick, J., Goryl, P., Klein, U., Laur, H., Mavrocordatos, C., Nieke, J., Rebhan, H., Seitz, B., Stroede, J., and Sciarra, R.: The Global Monitoring for Environment and Security (GMES) Sentinel-3 mission, *Remote Sensing of Environment*, 120, 37 – 57, <https://doi.org/https://doi.org/10.1016/j.rse.2011.07.024>, <http://www.sciencedirect.com/science/article/pii/S0034425712000685>, the Sentinel Missions - New Opportunities for Science, 2012.
- Ducet, N., Le Traon, P. Y., and Reverdin, G.: Global high-resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and -2, *Journal of Geophysical Research: Oceans*, 105, 19477–19498, <https://doi.org/10.1029/2000jc900063>, <https://doi.org/10.1029/2000jc900063>, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2000JC900063><https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JC900063><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2000JC900063>, 2000.
- 340 Giles, K. a., Laxon, S. W., Ridout, A. L., Wingham, D. J., and Bacon, S.: Western Arctic Ocean freshwater storage increased by wind-driven spin-up of the Beaufort Gyre, *Nature Geoscience*, 5, 194–197, <https://doi.org/10.1038/ngeo1379>, <http://www.nature.com/doi/10.1038/ngeo1379>, 2012.
- 345 Helm, V., Humbert, A., and Miller, H.: Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8, 1539–1559, <https://doi.org/10.5194/tc-8-1539-2014>, <https://tc.copernicus.org/articles/8/1539/2014/>, 2014.
- Iijima, B., Harris, I., Ho, C., Lindqwister, U., Mannucci, A., Pi, X., Reyes, M., Sparks, L., and Wilson, B.: Automated daily process for global ionospheric total electron content maps and satellite ocean altimeter ionospheric calibration based on Global Positioning System data, *Journal of Atmospheric and Solar-Terrestrial Physics*, 61, 1205–1218, [https://doi.org/10.1016/s1364-6826\(99\)00067-x](https://doi.org/10.1016/s1364-6826(99)00067-x), [https://doi.org/10.1016/s1364-6826\(99\)00067-x](https://doi.org/10.1016/s1364-6826(99)00067-x), 1999.
- 350 Kwok, R.: Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018), *Environmental Research Letters*, 13, 105 005, <https://doi.org/10.1088/1748-9326/aae3ec>, <https://doi.org/10.1088/1748-9326/aae3ec>, 2018.
- Lavergne, T., Sørensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, G., Eastwood, S., Gabarro, C., Heygster, G., Killie, M. A., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S., and Pedersen, L. T.: Version 2 of the EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records, *The Cryosphere*, 13, 49–78, <https://doi.org/10.5194/tc-13-49-2019>, <https://tc.copernicus.org/articles/13/49/2019/>, 2019.
- Lawrence, I. R., Armitage, T. W., Tsamados, M. C., Stroeve, J. C., Dinardo, S., Ridout, A. L., Muir, A., Tilling, R. L., and Shepherd, A.: Extending the Arctic sea ice freeboard and sea level record with the Sentinel-3 radar altimeters, *Advances in Space Research*, <https://doi.org/10.1016/j.asr.2019.10.011>, <https://doi.org/10.1016/j.asr.2019.10.011>, 2019.




- 360 Le Traon, P. Y. and Dibarboure, G.: Mesoscale Mapping Capabilities of Multiple-Satellite Altimeter Missions, *Journal of Atmospheric and Oceanic Technology*, 16, 1208–1223, [https://doi.org/10.1175/1520-0426\(1999\)016<1208:mmcoms>2.0.co;2](https://doi.org/10.1175/1520-0426(1999)016<1208:mmcoms>2.0.co;2), [https://doi.org/10.1175/1520-0426\(1999\)016<1208:mmcoms>2.0.co;2](https://doi.org/10.1175/1520-0426(1999)016<1208:mmcoms>2.0.co;2), 1999.
- Le Traon, P.-Y. and Ogor, F.: ERS-1/2 orbit improvement using TOPEX/POSEIDON: The 2 cm challenge, *Journal of Geophysical Research: Oceans*, 103, 8045–8057, <https://doi.org/10.1029/97jc01917>, <https://doi.org/10.1029/97jc01917>, 1998.
- 365 Long  p  , N., Thibaut, P., Vadaine, R., Poisson, J., Guillot, A., Boy, F., Picot, N., and Borde, F.: Comparative Evaluation of Sea Ice Lead Detection Based on SAR Imagery and Altimeter Data, *IEEE Transactions on Geoscience and Remote Sensing*, 57, 4050–4061, <https://doi.org/10.1109/TGRS.2018.2889519>, 2019.
- Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh, A., Melbourne-Thomas, J., Muelbert, M., Ottersen, G., Pritchard, H., and Schuur, E.: *Polar Regions*, 2019.
- 370 Pascual, A., Faug  re, Y., Larnicol, G., and Le Traon, P. Y.: Improved description of the ocean mesoscale variability by combining four satellite altimeters, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL024633>, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2005GL024633><https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005GL024633><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2005GL024633>, 2006.
- Peacock, N. R. and Laxon, S.: Sea surface height determination in the Arctic Ocean from ERS altimetry, *Journal of Geophysical Research*, 375, C07 001, <https://doi.org/10.1029/2001JC001026>, <http://doi.wiley.com/10.1029/2001JC001026>, 2004.
- Poisson, J., Quartly, G. D., Kurekin, A. A., Thibaut, P., Hoang, D., and Nencioli, F.: Development of an ENVISAT Altimetry Processor Providing Sea Level Continuity Between Open Ocean and Arctic Leads, *IEEE Transactions on Geoscience and Remote Sensing*, 56, 5299–5319, 2018.
- 380 Polyakov, I. V., Pnyushkov, A. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Carmack, E. C., Goszczko, I., Guthrie, J., Ivanov, V. V., Kanzow, T., Krishfield, R., Kwok, R., Sundfjord, A., Morison, J., Rember, R., and Yulin, A.: Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356, 285–291, <https://doi.org/10.1126/science.aai8204>, <https://science.sciencemag.org/content/356/6335/285>, 2017.
- Prandi, P.: Gridded Sea Level Heights - Arctic Ocean, <https://doi.org/10.24400/527896/a01-2020.001>, 2020.
- Proshutinsky, A., Dukhovskoy, D., Timmermans, M.-L., Krishfield, R., and Bamber, J. L.: Arctic circulation regimes, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373, 20140 160, <https://doi.org/10.1098/rsta.2014.0160>, <https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2014.0160>, 2015.
- Pujol, M.-I., Faug  re, Y., Taburet, G., Dupuy, S., Pelloquin, C., Ablain, M., and Picot, N.: DUACS DT2014: the new multi-mission altimeter data set reprocessed over 20 years, *Ocean Science*, 12, 1067–1090, <https://doi.org/10.5194/os-12-1067-2016>, <https://os.copernicus.org/articles/12/1067/2016/>, 2016.
- 390 Quartly, G., Rinne, E., Passaro, M., Andersen, O., Dinardo, S., Fleury, S., Guillot, A., Hendricks, S., Kurekin, A., M  ller, F., Ricker, R., Skourup, H., and Tsamados, M.: Retrieving Sea Level and Freeboard in the Arctic: A Review of Current Radar Altimetry Methodologies and Future Perspectives, *Remote Sens.*, 11, 881, 2019.
- Regan, H., Lique, C., Talandier, C., and Meneghello, G.: Response of total and eddy kinetic energy to the recent spinup of the Beaufort Gyre, *Journal of Physical Oceanography*, 50, 575–594, <https://doi.org/10.1175/JPO-D-19-0234.1>, 2020.
- 395 Regan, H. C., Lique, C., and Armitage, T. W. K.: The Beaufort Gyre Extent, Shape, and Location Between 2003 and 2014 From Satellite Observations, *Journal of Geophysical Research: Oceans*, 124, 844–862, <https://doi.org/10.1029/2018JC014379>, <https://onlinelibrary.wiley.com/doi/abs/10.1029/2018JC014379>, 2019.



- Rose, S. K., Andersen, O. B., Passaro, M., Ludwigsen, C. A., and Schwatke, C.: Arctic Ocean Sea Level Record from the Complete Radar Altimetry Era: 1991–2018, *Remote Sensing*, 11, 1672, <https://doi.org/10.3390/rs11141672>, <https://www.mdpi.com/2072-4292/11/14/1672>, 2019.
- Smith, G. C., Allard, R., Babin, M., Bertino, L., Chevallier, M., Corlett, G., Crout, J., Davidson, F., Delille, B., Gille, S. T., Hebert, D., Hyder, P., Intrieri, J., Lagunas, J., Larnicol, G., Kaminski, T., Kater, B., Kauker, F., Marec, C., Mazloff, M., Metzger, E. J., Mordy, C., OCarroll, A., Olsen, S. M., Phelps, M., Posey, P., Prandi, P., Rehm, E., Reid, P., Rigor, I., Sandven, S., Shupe, M., Swart, S., Smedstad, O. M., Solomon, A., Storto, A., Thibaut, P., Toole, J., Wood, K., Xie, J., Yang, Q., and , t. W. P. S. G.: Polar Ocean Observations: A Critical Gap in the Observing System and Its Effect on Environmental Predictions From Hours to a Season, *Frontiers in Marine Science*, 6, 429, <https://doi.org/10.3389/fmars.2019.00429>, <https://www.frontiersin.org/article/10.3389/fmars.2019.00429>, 2019.
- Smith, W. H. F. and Scharroo, R.: Waveform Aliasing in Satellite Radar Altimetry, *IEEE Transactions on Geoscience and Remote Sensing*, 53, 1671–1682, <https://doi.org/10.1109/TGRS.2014.2331193>, 2015.
- Stammer, D. and Cazenave, A., eds.: *Satellite Altimetry Over Oceans and Land Surfaces*, CRC Press, <https://doi.org/https://doi.org/10.1201/9781315151779>, 2018.
- Steunou, N., Desjonquères, J. D., Picot, N., Sengenès, P., Noubel, J., and Poisson, J. C.: AltiKa Altimeter: Instrument Description and In Flight Performance, *Marine Geodesy*, 38, 22–42, <https://doi.org/10.1080/01490419.2014.988835>, <https://doi.org/10.1080/01490419.2014.988835>, 2015.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environmental Research Letters*, 13, 103 001, <https://doi.org/10.1088/1748-9326/aade56>, <https://doi.org/10.1088%2F1748-9326%2Faade56>, 2018.
- Taburet, G., Sanchez-Roman, A., Ballarotta, M., Pujol, M.-I., Legeais, J.-F., Fournier, F., Faugere, Y., and Dibarboue, G.: DUACS DT2018: 25 years of reprocessed sea level altimetry products, *Ocean Science*, 15, 1207–1224, <https://doi.org/10.5194/os-15-1207-2019>, <https://os.copernicus.org/articles/15/1207/2019/>, 2019.
- Timmermans, M.-L., C., L., and Wood, K.: Sea surface temperature. Arctic Report Card, <https://arctic.noaa.gov/Report-Card/Report-Card-2017/ArtMID/7798/ArticleID/698/Sea-Surface-Temperature>, 2017.
- Valladeau, G., Thibaut, P., Picard, B., Poisson, J. C., Tran, N., Picot, N., and Guillot, A.: Using SARAL/AltiKa to Improve Ka-band Altimeter Measurements for Coastal Zones, Hydrology and Ice: The PEACHI Prototype, *Marine Geodesy*, 38, 124–142, <https://doi.org/10.1080/01490419.2015.1020176>, <https://doi.org/10.1080/01490419.2015.1020176>, 2015.
- Verron, J., Sengenès, P., Lambin, J., Noubel, J., Steunou, N., Guillot, A., Picot, N., Coutin-Faye, S., Sharma, R., Gairola, R. M., Murthy, D. V. A. R., Richman, J. G., Griffin, D., Pascual, A., Rémy, F., and Gupta, P. K.: The SARAL/AltiKa Altimetry Satellite Mission, *Marine Geodesy*, 38, 2–21, <https://doi.org/10.1080/01490419.2014.1000471>, <https://doi.org/10.1080/01490419.2014.1000471>, 2015.
- Wingham, D., Francis, C., Baker, S., Bouzinac, C., Brockley, D., Cullen, R., de Chateau-Thierry, P., Laxon, S., Mallow, U., Mavrocardatos, C., Phalippou, L., Ratier, G., Rey, L., Rostan, F., Viau, P., and Wallis, D.: CryoSat: A mission to determine the fluctuations in Earth’s land and marine ice fields, *Advances in Space Research*, 37, 841–871, <https://doi.org/10.1016/j.asr.2005.07.027>, <https://doi.org/10.1016/j.asr.2005.07.027>, 2006.
- Zhao, M., Timmermans, M.-L., Cole, S., Krishfield, R., and Toole, J.: Evolution of the eddy field in the Arctic Ocean’s Canada Basin, 2005–2015, *Geophysical Research Letters*, 43, 8106–8114, <https://doi.org/10.1002/2016GL069671>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL069671>, 2016.



(a)



(b)

Figure 1. time series of (a) Arctic regional sea level for SRL, S3A and C2, and (b) regional sea level differences between C2 and SRL and S3A and SRL






Figure 2. mean Arctic SLA maps for SRL (a), S3A (b) and C2 (c)



(a) S3A-SRL



(b) C2-SRL

Figure 3. maps of mean SL differences between (a) S3A and SRL and (b) C2 and SRL




Figure 4. a priori signal variance maps used in (a) DUACS global processing and (b) in the Arctic regional processing




Figure 5. noise levels over open ocean (left) and ice covered (right) areas used in the regional analysis




Figure 6. variance of LWE errors used for the Arctic Ocean optimal interpolation






Figure 7. maps of (a) mean and (b) variance of SL in the Arctic Ocean from the combined product



(a) mean



(b) variance

Figure 8. differences with between arctic product and CMEMS global products (a) mean difference and (b) variance of the differences expressed as fraction of signal variance




Figure 9. Sea level anomaly time series at station Prudhoe Bay from the tide gauge station (dark gray), the combined altimetry product (red) and mono-mission sea levels for SRL (blue) S3A (green) and C2 (purple)




Figure 10. Ratio of the error power spectrum to the signal power spectrum at tide gauges (gray lines), the mean (blue) and median (red) ratio are shown. The resolution threshold is displayed as a dashed black line. Resolution values are displayed (in cyc/day) on the bottom panel. Left is for a single altimeter (S3A) product, right for the combined product.



Table 1. some characteristics of altimeters used

mission name	cycle duration (days)	inclination	mode	input product	sampling freq
SARAL/AltiKa	35	98.538°	LRM	SGDR-T Patch 2	40 Hz
CryoSat-2	369 (30 days pseudo-cycle)	92°	SAR	PDGS Ice Baseline C	20 Hz
Sentinel-3A	27	98.65°	SAR	CNES S3PP with 0pad/Hamming	20 Hz



Table 2. backscattering thresholds

mission name	backscatter (dB)
SARAL/AltiKa	20
CryoSat-2	23
Sentinel-3A	13



Table 3. Arctic SL altimeter standards

	SARAL/AltiKa	Sentinel-3A	CryoSat-2
orbit	POE-D	from SRAL product	POE-E
ionospheric correction		GIM (Iijima et al., 1999)	
dry tropospheric correction		ECMWF model	
wet tropospheric correction		ECMWF model	
DAC		MOG2D model (Carrère and Lyard, 2003)	
pole tide		Desai et al. (2015)	
ocean tide		FES2014 (Carrère et al., 2016)	
solid earth tide	elastic response to tidal potential (Cartwright and Edden, 1973)		
sea state bias		non parametric (ocean only)	
mean sea surface		DTU15 (and, 2016)	



Table 4. product characteristics

product name	grid resolution (km)	time step (days)	first date	last date
SARAL/AltiKa monomission	75	10	2013/03/28	2019/08/24
CryoSat-2 monomission	75	10	2010/07/30	2019/04/24
Sentinel-3A monomission	75	10	2016/07/01	2019/05/01
combined	25	3	2016/07/01	2019/04/29



Table 5. comparisons to tide gauges, *RMSd* values are given in m

name	station		SARAL/AltiKa		Sentinel-3A		CryoSat-2		combined	
	latitude	longitude	<i>R</i>	<i>RMSd</i>	<i>R</i>	<i>RMSd</i>	<i>R</i>	<i>RMSd</i>	<i>R</i>	<i>RMSd</i>
Castletownsend	51.5	-9.2	0.72	0.06	0.55	0.05	–	–	0.69	0.05
Adak	51.9	-176.6	0.54	0.03	0.51	0.03	–	–	0.62	0.03
Unalaska	53.9	-166.5	0.76	0.03	0.68	0.04	–	–	0.76	0.03
Prince Rupert	54.3	-130.3	0.81	0.05	0.83	0.05	–	–	0.63	0.07
Ketchikan	55.3	-131.6	0.66	0.07	0.83	0.05	–	–	0.73	0.06
Sand Point	55.3	-160.5	0.87	0.06	0.88	0.05	–	–	0.92	0.04
Malin Head	55.4	-7.3	0.63	0.07	0.78	0.04	–	–	0.80	0.04
Nain	56.5	-61.7	0.61	0.06	0.72	0.06	0.50	0.07	0.62	0.06
Sitka	57.1	-135.3	0.69	0.06	0.74	0.05	–	–	0.85	0.04
Goteborg	57.7	11.8	0.65	0.09	0.62	0.09	–	–	0.74	0.07
Kodiak Island	57.7	-152.5	0.80	0.04	0.85	0.03	–	–	0.81	0.03
Tregde	58.0	7.5	0.68	0.06	0.70	0.06	–	–	0.64	0.06
Smogen	58.3	11.2	0.65	0.08	0.65	0.08	–	–	0.64	0.08
Stockholm	59.3	18.1	0.87	0.08	0.88	0.07	–	–	0.98	0.04
Yakutat	59.5	-139.7	0.77	0.07	0.82	0.06	–	–	0.90	0.04
Seward	60.1	-149.4	0.81	0.06	0.83	0.06	–	–	0.87	0.05
Qaqortoq	60.7	-46.0	0.37	0.12	0.57	0.08	–	–	0.62	0.07
Maloy	61.2	5.1	0.76	0.07	0.85	0.06	–	–	0.88	0.05
Reykjavik	64.2	-21.9	0.82	0.04	0.77	0.03	–	–	0.81	0.03
Godthaab	64.2	-51.0	0.58	0.09	0.46	0.10	–	–	0.56	0.09
Nome	64.5	-165.4	0.49	0.18	0.61	0.18	0.42	0.21	0.90	0.10
Rorvik	64.8	11.3	0.83	0.07	0.88	0.06	–	–	0.90	0.05
Andenes	69.3	16.1	0.78	0.07	0.85	0.06	–	–	0.92	0.04
Vardo	70.3	31.1	0.82	0.06	0.82	0.06	–	–	0.90	0.04
Prudhoe Bay	70.4	-148.5	0.64	0.13	0.62	0.12	0.35	0.15	0.85	0.08
Honningsvag	70.9	26.0	0.72	0.06	0.84	0.05	–	–	0.93	0.03
Thule	76.0	-68.0	0.55	0.06	0.48	0.06	0.51	0.08	0.50	0.06
mean			0.71	0.07	0.72	0.06	0.45	0.13	0.78	0.05