

Greenland ice sheet mass balance from 1840 through next week

Kenneth D. Mankoff¹, Xavier Fettweis², Peter L. Langen³, Martin Stendel⁴, Kristian K. Kjeldsen¹, Nanna B. Karlsson¹, Brice Noël⁵, Michiel R. van den Broeke⁵, Anne Solgaard¹, William Colgan¹, Jason E. Box¹, Sebastian B. Simonsen⁶, Michalea D. King⁷, Andreas P. Ahlstrøm¹, Signe Bech Andersen¹, and Robert S. Fausto¹

¹Department of Glaciology and Climate, Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

²SPHERES research unit, Department of Geography, University of Liège, Liège, Belgium

³Department of Environmental Science, iClimate, Aarhus University, Roskilde, Denmark

⁴Danish Meteorological Institute (DMI), Copenhagen, Denmark

⁵Institute for Marine and Atmospheric Research, Utrecht University, The Netherlands

⁶Geodesy and Earth Observation, DTU Space, Technical University of Denmark, Lyngby, Denmark

⁷Polar Science Center, University of Washington, Seattle, WA, United States

Correspondence: Ken Mankoff (kdm@geus.dk)

Abstract. The mass of the Greenland ice sheet is declining as mass gain from snow accumulation is exceeded by mass loss from surface meltwater runoff, marine-terminating glacier calving and submarine melting, and basal melting. Here we use the input/output (IO) method to estimate mass change from 1840 through next week. Surface mass balance (SMB) gains and losses come from a semi-empirical SMB model from 1840 through 1985, and three regional climate models (RCMs; 5 HIRHAM/HARMONIE, MAR, and RACMO) from 1986 through next week. Additional non-SMB losses come from a marine terminating glacier ice discharge product and a basal mass balance model. From these products we provide an annual estimate of Greenland ice sheet mass balance from 1840 through 1985 and a daily estimate at sector and region scale from 1986 through next week. This product updates daily and is the first IO product to include the basal mass balance which is a source of an additional ~ 24 Gt yr⁻¹ of mass loss. Our results demonstrate an accelerating ice-sheet-scale mass loss and general agreement 10 (coefficient of determination, r^2 , ranges from 0.62 to 0.94) among six other products, including gravitational, volume, and other IO mass balance estimates. Results from this study are available at <https://doi.org/10.22008/FK2/OHI23Z> (Mankoff et al., 2021).

1 Introduction

Over the past several decades, mass loss from the Greenland ice sheet has increased (Khan et al., 2015; The IMBIE Team, 15 2019). Different processes dominate the regional mass loss of the ice sheet, and their relative contribution has fluctuated in time (Mouginot and Rignot, 2019). For example, in the 1970s nearly all sectors gained mass due to positive SMB, except the northwest sector where discharge losses dominated. More recently in the 2010s, all sectors lost mass, with some sectors losing mass almost entirely via negative SMB, and others primarily due to discharge (Fig. 1).

There are three common methods for estimating mass balance – changes in gravity (Barletta et al., 2013; Groh et al., 2019; 20 The IMBIE Team, 2019; Velicogna et al., 2020), changes in volume (Simonsen et al., 2021a; Sørensen et al., 2011; Zwally and

Giovinetto, 2011; Sasgen et al., 2012; Smith et al., 2020), and the input/output (IO) method (Colgan et al., 2019; Mougnot et al., 2019; Rignot et al., 2019; King et al., 2020). Each provides some estimate of where, when, and how the mass is lost or gained, and each method has some limitations. The gravity mass balance (GMB) estimate has low ~100 km spatial resolution (where), monthly temporal resolution (when), and little information on the processes contributing to changes in mass balance components (how). The volume change (VC) mass balance estimate has ~1 km spatial resolution (where), often provided on annual or multi-year temporal resolution (when), and with little information on the driving processes (how).

The IO method has a complex spatial resolution (where). The inputs typically come from regional climate models (RCMs) which can reach a spatial resolution of up to 1 km. However, that spatial resolution is generally reduced in the final output to sector or region scale – typically higher than GMB but now lower resolution than VC. The IO temporal resolution (when) is limited by ice velocity data updates, which for the past several years occur every 12 days year-round after the launch of the Sentinel missions (Solgaard et al., 2021). The primary issue with the IO method is unknown ice thickness in some locations (e.g. Mankoff et al. (2020b)). Finally, the IO method can provide insight into the processes (how) by distinguishing between changes caused by SMB (which may be due to changes in positive and/or negative SMB components) vs. changes in other mass loss terms (e.g., calving). Our IO method is also the first IO product to include the basal mass balance (Karlsson et al., 2021) – a term implicitly included in the GMB and VC methods but neglected by all previous IO estimates.

In this work we introduce the new PROMICE Greenland ice sheet mass balance dataset based on the IO method, updating the previous product from Colgan et al. (2019). We use the SMB field from one empirical model from 1840 through 1985, and three RCMs from 1986 onward. The combined SMB field used here is comprised of positive SMB terms (precipitation in the form of snowfall, rainfall, and condensation/riming) and negative SMB terms (surface melt, evaporation, sublimation, and snow drift erosion/deposition). We also use the basal mass balance, and an estimate of dynamic ice discharge. Spatial resolution is effectively per sector (Zwally et al., 2012) or region (Mougnot and Rignot, 2019). Temporal resolution is annual from 1840 through 1985, and effectively daily since 1986 - the RCM fields are updated daily and forecasted through next week, and the marine mass balance (i.e., discharge at marine terminating glaciers) is updated every 12 days with ~12 day resolution, interpolated to daily, and forecasted using historical and seasonal trends through next week. Thus, this study provides a daily-updating estimate of Greenland mass changes from 1840 through next week.

2 Terminology

We use the following terminology throughout the document:

- This Study refers to the new results presented in this study.
- Recent refers to the new 1986 through next week daily temporal resolution data at region and sector scale
- Reconstructed refers to the adjusted Kjeldsen et al. (2015) annual temporal resolution data at ice sheet scale used to extend this product from 1986 back through 1840. The 1986 through 2012 portion of the Kjeldsen et al. (2015) data set is used only to adjust the reconstructed data, then discarded.

- ROI (region of interest) refers to one or more of the ice sheet sectors or regions (Fig. 1).
- Sector refers to one of the Zwally et al. (2012) sectors (Fig. 1), expanded here to cover the RCM ice domains which exist slightly outside these sectors in some locations.
- Region refers to the Mouginot and Rignot (2019) regions (Fig. 1), expanded here to cover the RCM ice domains.
- SMB is the surface mass balance from an RCM.
- MMB is the marine mass balance, which in other works is often referred to as solid ice discharge or D. It includes both calving and submarine melting.
- BMB is the basal mass balance. It comes from geothermal flux (BMB_{GF}), frictional heating from ice velocity ($BMB_{friction}$), and viscous heat dissipation (BMB_{VHD}).
- MB is the total mass balance including the BMB term (Eq. 3).
- MB* is the mass balance not including the BMB term (Eq. 4).
- HIRHAM/HARMONIE, MAR, and RACMO refer to the RCMs, which only provide SMB, and runoff in the case of MAR. However, when referencing the different MB products, we use, for example, "MAR MB" rather than repeatedly explicitly stating "MB derived from MAR SMB minus BMB and MMB". The use should be clear from context.

3 Product description

The output of this work is two NetCDF files and one CSV file containing a time series of mass balance and the components used to calculate mass balance. The only difference between the two NetCDF files is the ROI – one for Zwally et al. (2012) sectors and one for Mouginot and Rignot (2019) regions. Each NetCDF file includes the ice sheet mass balance (MB), MB per region of interest (ROI; sector or region), MB per ROI per RCM, ice sheet surface mass balance (SMB), SMB per ROI, ice sheet marine mass balance (MMB), MMB per ROI, ice sheet basal mass balance (BMB), and BMB per ROI. The CSV file contains a copy of the ice-sheet summed data.

An example of the output is shown in Fig. 2, where the top panel shows mass balance for the entire Greenland ice sheet, in addition to SMB, and MMB, at annual resolution. The lower panel shows an example two years at daily temporal resolution. The ice-sheet-wide product includes data from 1840 through next week, but the sector and region-scale products only includes data from 1986 through next week, because the 1840 through 1985 reconstructed only exists at ice-sheet scale (Fig. 1).

4 Data sources

This section introduces data products that exist prior to and are external to this work (Table 1). In the following Methods section we introduce both the intermediate products we generate using these data sources, and the final product that is the output of This Study.




Figure 1. Annual mass balance (black lines), surface mass balance (blue lines) and marine plus basal mass balance (dashed grey) in Gt yr^{-1} for each of the seven Mouginot and Rignot (2019) regions. The map shows both the named regions (Mouginot and Rignot, 2019) and the numbered sectors (Zwally et al., 2012). Marine mass balance gates are marked in black. Only recent (post-1986) data are shown because reconstructed data are not separated into regions or sectors. Next week is defined as 2021-08-20 based on the date this document was compiled.

The inputs to this work are the recent SMB fields from the three RCMs, the recent marine mass balance data from Mankoff et al. (2020b) (data: Mankoff and Solgaard (2020)), and the recent basal mass balance fields, of which BMB_{GF} and $\text{BMB}_{\text{friction}}$




Figure 2. Mass balance and its major components. Top panel: Annual average surface mass balance (blue line), marine mass balance (gray dashed), and their mass balance sum (black line). Here the marine and basal mass balance (MMB + BMB) are shown with sign inverted (e.g. $-1 \times (\text{MMB} + \text{BMB})$) Lower panel: Same data at daily resolution and limited to 2019 and 2020.

are direct outputs from Karlsson et al. (2021) (data: Karlsson (2021)), but the BMB_{VHD} calculations are redone here (see
 85 Methods Sect. 5.3) using the MAR runoff field. The reconstructed data (pre 1986) are surface and marine mass balance from
 Kjeldsen et al. (2015) (data: Box et al. (2021)), but adjusted here using the overlapping period (see Methods Sect. 5.4), and
 runoff from Kjeldsen et al. (2015) (data: Box et al. (2021)) as a proxy and scaled for BMB_{VHD} (see Methods Sect. 5.3).

4.1 Surface mass balance

We use one reconstructed SMB from 1840 through 1985, and three recent SMB from 1986 through last month (HIRHAM/HARMONIE,
 90 MAR, and RACMO), two through yesterday (HIRHAM/HARMONIE and MAR) and one through next week (MAR).

4.1.1 HIRHAM/HARMONIE

The HIRHAM/HARMONIE product from the Danmarks Meteorologiske Institut (Danish Meteorological Institute; DMI) is
 based on an offline subsurface firn/SMB model (Langen et al., 2017), which is forced with surface fluxes of energy (turbu-

Table 1. Summary of data products used as inputs to This Study.

Product	Period	Reference	Data / Notes
Reconstructed SMB	1840 through 1985	Kjeldsen et al. (2015)	Box et al. (2021)
Reconstructed MMB	1840 through 1985	Kjeldsen et al. (2015)	Box et al. (2021)
HIRHAM/HARMONIE SMB	1986 through yesterday	Langen et al. (2017)	
MAR SMB	1986 through next week	Fettweis et al. (2020)	
RACMO SMB	1986 through last month	Noël et al. (2019)	
MMB	1986 through last month	Mankoff et al. (2020b)	Mankoff and Solgaard (2020)
BMB_{GF} ; $BMB_{friction}$	1840 through next week	Karlsson et al. (2021)	Karlsson (2021)
BMB_{VHD}	1840 through 1985	Kjeldsen et al. (2015)	Box et al. (2021) reconstructed runoff
BMB_{VHD}	1986 through next week	Fettweis et al. (2020)	MAR runoff

lent and downward radiative) and mass (snow, rain, evaporation, and sublimation). These surface fluxes are derived from the
95 HIRHAM5 regional climate model for the reconstructed part of the simulation and from DMI's operational numerical weather
forecast model HARMONIE (Iceland-Greenland domain "B", which covers Iceland, Greenland, and the adjacent seas) for the
real-time part. HIRHAM5 is used until 2017-08-31 after which HARMONIE is used.

The HIRHAM5 regional climate model (Christensen et al., 2007) combines the dynamical core of the HIRLAM7 numerical
weather forecasting model (Eerola, 2006) with physics schemes from the ECHAM5 general circulation model (Roeckner et al.,
100 2003). In the Greenland setup employed here (Lucas-Picher et al., 2012), it has a horizontal resolution of $0.05^\circ \times 0.05^\circ$ on a
rotated pole grid (corresponding to 5.5 km resolution), and 31 atmospheric levels in the vertical. It is forced at 6 hr intervals
on the lateral boundaries with horizontal wind vectors, temperature, and specific humidity from the ERA-Interim reanalysis
(Dee et al., 2011). ERA-Interim sea surface temperatures and sea ice concentration are prescribed in ocean grid points. Surface
fluxes from HIRHAM5 are passed to the offline subsurface model.

105 The offline subsurface model was developed to improve firn details for the HIRHAM5 experiments (Langen et al., 2017).
The subsurface consists of 32 layers with time-varying fractions of snow, ice and liquid water. Layer thicknesses increase with
depth from 6.5 cm water equivalent (w.e.) at the top to 9.2 m w.e. at the bottom giving a full model depth of 60 m w.e. The
processes governing the firn evolution include snow densification, varying hydraulic conductivity, irreducible water saturation
and other effects on snow liquid water percolation, and retention. Runoff is calculated from liquid water in excess of the
110 irreducible saturation with a characteristic local timescale that depends on surface slope (Zuo and Oerlemans, 1996; Lefebvre,
2003). The offline subsurface model is run on the HIRHAM5 5.5 km grid.

For the real-time data we use DMI's operational numerical weather forecast model HARMONIE (Bengtsson et al., 2017),
a nonhydrostatic model in terrain-following sigma coordinates based on the fully compressible Euler equations (Simmons and
Burridge, 1981; Laprise, 1992). HARMONIE is run at 2.5 km horizontal resolution and with 65 vertical levels. Compared to
115 previous model versions, upper air 3D variational data assimilation of satellite wind and radiance data, radio occultation data,
radiosonde, aircraft, and surface observations are incorporated. This greatly improves the number of observations in the model,

as in situ observations from ground stations and radiosondes only make up approximately 20 % of observations in Greenland (Wang and Randriamampianina, 2021; Yang et al., 2018). The model is driven at the boundaries with European Centre for Medium-Range Weather Forecasts (ECMWF) high-resolution data at 9 km resolution. The 2.5 km HARMONIE output is regridded to the 5.5 km HIRHAM grid before input to the offline subsurface model. The HIRHAM5 and the offline model both employ the Citterio and Ahlstrøm (2013) ice mask interpolated to the 5.5 km grid.

4.1.2 MAR

The Modèle Atmosphérique Régional (MAR) RCM has been developed by the University of Liège (Belgium) with a focus on the polar regions (Fettweis et al., 2020). The MAR atmosphere module (Gallée and Schayes, 1994) is fully coupled with the soil-ice-snow energy balance vegetation model SISVAT (Gallée et al., 2001) simulating the evolution of the 30 first meters of snow/ice over the ice sheet with the help of 30 snow layers (with time varying thickness) or the 10 first meter of soil over the tundra area. At its lateral boundary, MAR is 6 hourly forced by the reanalysis ERA5 and runs at a resolution of 20 km. The snow pack has been initialised in 1950 from a former MARv3.11 based simulation. Its snow model is based on a former version of the CROCUS snow model (Vionnet et al., 2012) dealing with all the snowpack processes including the meltwater retention, transformation of melting snow and grain size, compaction of snow, formation of ice lenses impacting meltwater penetration, warming of the snowpack from rainfall, and complex snow/bare ice albedo. MAR uses the Greenland Ice Mapping Project (GIMP) ice sheet mask and ice sheet topography (Howat et al., 2014).

The new version, MAR 3.12, is used here. With respect to version 3.9 intensively validated over Greenland (Fettweis et al., 2020) or the 20 km based MARv3.10 set-up used in Tedesco and Fettweis (2020), MARv3.12 now uses the common polar stereographic projection EPSG 3413. With respect to MARv3.11 fully described in Amory et al. (2021), MARv3.12 assures now the full conservation of water mass into both soil and snowpack at each time step, takes into account of the geographical projection deformations in its advection scheme, better deals with the snow/rain temperature limit with a continuous temperature threshold between 0 and -2°C, increases the evaporation above snow thanks to a saturated humidity computation in SISVAT adapted to freezing temperatures, disallows melt below the 30 m of the resolved snowpack, includes small improvements and bug fixes with the aim of improving the evaluation of MAR (with both in situ and satellite products) as presented in Fettweis et al. (2020) in addition to small computer time improvements in the parallelisation of its code.

In addition to providing SMB, MAR also provides daily runoff over both permanent ice and tundra area – this is used for the daily BMB_{VHD} estimate (Section 5.3).

As the recent SMB decrease (successfully evaluated with GRACE based estimates in Fettweis et al. (2020)) has been fully driven by the increase in runoff (Sasgen et al., 2020), we assume the same degree of accuracy between SMB simulated by MAR (evaluated with the PROMICE SMB database (Fettweis et al., 2020)) and the runoff simulated by MAR.

Weather-forecasted SMB: To provide a real-time state of the Greenland ice sheet, MAR is forced automatically everyday by the run of 00 h UTC from the Global Forecast System (GFS) model providing weather forecasting initialised by the snowpack behaviours of the MAR run from the previous day. This continuous GFS forced time series (without any reinitialisation of MAR) provides SMB and runoff estimates between the period covered by ERA5 and the next 7 days. At the end of each

day, ERA5 is used to update the GFS forced MAR time series until about 5 days before the current date and to provide an homogeneous ERA5 forced MAR times series from 1950 to a few days before the current date. We use both the forecasted SMB and forecasted runoff (for BMB_{VHD}) fields

4.1.3 RACMO

155 The Regional Atmospheric Climate MOdel (RACMO) v2.3p2 has been developed at the Koninklijk Nederlands Meteorologisch Instituut (Royal Netherlands Meteorological Institute; KNMI). It incorporates the dynamical core of the High-Resolution Limited Area Model (HIRLAM) and the physics parameterizations of the ECMWF Integrated Forecast System cycle CY33r1. A polar version (p) of RACMO has been developed at the Institute for Marine and Atmospheric research of Utrecht University (UU-IMAU), to assess the surface mass balance of glaciated surfaces. The current version RACMO2.3p2 has been described
160 in detail in Noël et al. (2018), and here we repeat the main characteristics.

The ice sheet has an extensive dry interior snow zone, a relatively narrow runoff zone along the low-lying margins, and a percolation zone of varying width in between. To capture these processes in first order, the original single-layer snow model in RACMO has been replaced by a 40-layer snow scheme that includes expressions for dry snow densification and a simple tipping bucket scheme to simulate meltwater percolation, retention, refreezing, and runoff (Ettema et al., 2010). The snow layers are
165 initialized in September 1957 using temperature and density from a previous run with the offline IMAU Firn Densification Model (Ligtenberg et al., 2018). To simulate drifting snow transport and sublimation, Lenaerts et al. (2012) implemented a drifting snow scheme. Snow albedo depends on snow grain size, cloud optical thickness, solar zenith angle, and impurity content (van Angelen et al., 2012). Bare ice albedo is assumed constant and estimated as the fifth percentile value of albedo time series (2000-2015) from the 500 m resolution MODIS 16-day albedo product (MCD43A3). Minimum/maximum values
170 of 0.30/0.55 are applied to the bare ice albedo, representing ice with high/low impurity content (cryoconite, algae).

To simulate as accurately as possible the contemporary climate and surface mass balance of the ice sheet, the following boundary conditions have been applied. The glacier ice mask and surface topography have been down-sampled from the 90 m resolution Greenland Ice Mapping Project (GIMP) digital elevation model (Howat et al., 2014). At the lateral boundaries, model temperature, specific humidity, pressure, and horizontal wind components at the 40 vertical model levels are relaxed
175 towards 6-hourly ECMWF reanalysis (ERA) data. For this we use ERA-40 between 1958 and 1978 (Uppala et al., 2005), ERA-Interim between 1979 and 1989 (Dee et al., 2011), and ERA-5 between 1990 and 2020 (Hersbach et al., 2020). The relaxation zone is 24 grid cells (~130 km) wide to ensure a smooth transition to the domain interior. This run has active upper atmosphere relaxation (van de Berg and Medley, 2016). Over glaciated grid points, surface aerodynamic roughness is assumed constant for snow (1 mm) and ice (5 mm). In this run, RACMO2.3p2 has 5.5 km horizontal resolution over Greenland and the
180 adjacent oceans and land masses, but it was found previously that this is insufficient to resolve the many narrow outlet glaciers. The 5.5 km product is therefore statistically downscaled onto a 1 km grid sampled from the GIMP DEM (Noël et al., 2019), employing corrections for biases in elevation and bare ice albedo using a MODIS albedo product at 1 km resolution (Noël et al., 2016).

4.1.4 Reconstructed

185 The Kjeldsen et al. (2015) 173-year (1840 through 2012) mass balance reconstruction is based on the Box (2013) 171-year (1840 through 2010) statistical reconstruction. Kjeldsen et al. (2015) add a more sophisticated meltwater retention scheme (Pfeffer et al., 1991); weighting of in situ records in their contribution to the estimated value; dispersal of annual accumulation to monthly; and extend the reconstruction in time through 2012.

The Box (2013) 171-year (1840-2010) reconstruction is developed from linear regression parameters that describe the least squares regression between a) spatially discontinuous in situ monthly air temperature records (Cappelen et al., 2011; Cappelen, 2001; Cappelen et al., 2006; Vinther et al., 2006)) or firn/ice cores (Box et al., 2013) and b) spatially continuous outputs from regional climate model RACMO version 2.1 (Ettema et al., 2010). A 43-year overlap period (1960 through 2012) with the RACMO data are used to determine regression parameters (slope, intercept) on a 5 km grid cell basis. Temperature data define melting degree days, which have a different coefficient for bare ice than snow cover, determined from hydrological-
195 year cumulative SMB. A fundamental assumption is that the calibration factors, regression slope, and offset for the calibration period 1960 through 2012 are stationary over time for which there is some evidence of in Fettweis et al. (2017). Box et al. (2013) describes the methods in more detail.

The reconstructed surface mass balance is adjusted as described in the Methods Sect. 5.4 (Fig. 3).

4.2 Marine mass balance

200 The recent marine mass balance data are the discharge (D) product from Mankoff et al. (2020b) (data: Mankoff and Solgaard (2020)). This product covers all fast-flowing (> 100 m yr⁻¹) marine-terminating glaciers. The marine mass balance in Mankoff et al. (2020b) is computed at flux gates ~5 km upstream from glacier termini (Mankoff, 2020), using a wide range of velocity products, and ice thickness from BedMachine v4. Discharge across flux gates is derived with a 200 m spatial resolution grid, but then summed and provided at glacier resolution. Temporal coverage begins in 1986 with a few velocity estimates, and is
205 updated each time a new velocity product is released, which is every ~12 days with a ~30 day lag (Solgaard et al., 2021) (data: Solgaard and Kusk (2021)).

Some changes have been implemented since the last publication describing the marine mass balance product (i.e., Mankoff et al. (2020b)). These are minor and include updating the Khan et al. (2016) (data: Khan (2017)) surface elevation change product from 2015 through 2019, updating various MEaSURES velocity products to their latest version, updating the PROMICE
210 Sentinel ice velocity product from Edition 1 (doi:10.22008/promice/data/sentinel1icevelocity/greenlandicesheet/v1.0.0) to Edition 2 (Solgaard et al. (2021); Solgaard and Kusk (2021)), and updating from BedMachine v3 (supplemented in the SE with Millan et al. (2018)) to use only BedMachine v4 (Morlighem et al., 2021).

The reconstructed marine mass balance data (Kjeldsen et al., 2015) are estimated via a linear fit between unsmoothed annual marine mass balance spanning 2000 to 2012 (Enderlin et al., 2014) and runoff data from (Kjeldsen et al., 2015) using a 6-year
215 trailing average. The method for scaling discharge from runoff was introduced by (Rignot et al., 2008), who scaled the SMB anomaly with discharge. Sensitivity analyses conducted by Box and Colgan (2013) showed runoff to be the more effective

discharge predictor, and include a discussion of the physical basis. Although the fitting period of the present dataset includes an anomalous period of discharge (2000 through 2005; e.g., Boers and Rypdal (2021)), the discharge data used by Rignot et al. (2008) and Box and Colgan (2013) also includes years 1958 and 1964 that lie near the regression line (See Box and Colgan (2013) Fig. 4 and related section 4. Physical basis). Further, while 2000 through 2005 cover a changing period in Greenlandic discharge (Mankoff et al., 2020b; King et al., 2020), there were likely other anomalous periods in the past, when glaciers in Greenland experienced considerable increases in discharge as inferred by geological and geodetic investigations (Andresen et al., 2012; Bjørk et al., 2012; Khan et al., 2015, 2020).

The reconstructed marine mass balance is adjusted as described in the Methods Sect. 5.4.

225 4.3 Basal mass balance

The basal mass balance (BMB; Karlsson et al. (2021)) comes from mass lost at the bed from geothermal flux (BMB_{GF}), frictional heating ($BMB_{friction}$) from the basal shear velocity, and viscous heat dissipation (BMB_{VHD}) from surface runoff routed to the bed (i.e. the volume of the subglacial conduits formed from surface runoff; Mankoff and Tulaczyk (2017)).

230 These fields (data: Karlsson (2021)) are provided as steady state annual estimates. We use the BMB_{GF} and $BMB_{friction}$ products and apply 1/365th to each day, each year. Because BMB_{VHD} is proportional to runoff, an annual estimate is not appropriate for this work with daily resolution. We therefore re-calculate the BMB_{VHD} -induced basal melt as described in Methods Sect. 5.3.

4.3.1 Geothermal Flux

235 Due to a lack of direct observations, the geothermal flux is poorly constrained under most of the Greenland ice sheet. Different approaches have been employed to infer the value of the BMB_{GF} often with diverging results (see e.g., Rogozhina et al. (2012); Rezvanbehbahani et al. (2019)). Lacking substantial validation that favours one BMB_{GF} map over the others, Karlsson et al. (2021) instead use the average of three widely used BMB_{GF} estimates: Fox Maule et al. (2009); Shapiro and Ritzwoller (2004), and Martos et al. (2018). The BMB_{GF} melt rate is calculated as

$$\dot{b}_m = E_{GF} \rho_i^{-1} L^{-1}, \quad (1)$$

240 where E_{GF} is available energy at the bed, here the geothermal flux in units $W m^{-2}$, ρ_i is the density of ice (917 kg m^{-3}), and L is the latent heat of fusion (335 kJ kg^{-1} ; Cuffey and Paterson (2010)). BMB_{GF} melting is only calculated where the bed is not frozen. We use the MacGregor et al. (2016) estimate of temperate bed extent and scale Eq. 1 by 0, 0.5, or 1 where the bed is frozen ($\sim 25\%$ of the ice sheet area), uncertain ($\sim 33\%$), or thawed ($\sim 42\%$), respectively.

4.3.2 Friction

245 This heat term stems from the friction produced as ice slides over the bedrock. The term has only been measured in a handful of places (e.g., Ryser et al. (2014); Maier et al. (2019)) and it is unclear how representative those measurements are at ice-sheet

scales. Karlsson et al. (2021) therefore estimate the frictional heating using the Full Stokes Elmer/Ice model that resolves all stresses while relating basal sliding and shear stress using a linear friction law (Gillet-Chaulet et al., 2012; Maier et al., 2021). The model is tuned to match a multi-decadal surface velocity map (Joughin et al., 2018) covering 1995-2015 and it returns an
 250 estimated basal friction heat that is used to calculate the basal melt due to friction, similar to Eq. 1:

$$\dot{b}_m = E_f \rho_i^{-1} L^{-1}, \quad (2)$$

where E_f is energy due to friction. We also apply the 0, 0.5, and 1 scale as used for the BMB_{GF} term (MacGregor et al., 2016) in order to mask out areas that are likely frozen.

4.4 Other

255 ROI regions come from Mouginit and Rignot (2019) and ROI sectors come from Zwally et al. (2012).

4.5 Products used for validation

We validate This Study against five other data products (See Table 2 and Sect. 6). These products are the most recent IO product (Mouginit et al., 2019), the previous PROMICE mass balance product (Colgan et al. (2019); data: Colgan (2021)), the two mostly-independent methods of estimating ice sheet mass change: GMB (Barletta et al. (2013); data: Barletta et al. (2020)) and
 260 VC (Simonsen et al. (2021a); data: Simonsen et al. (2021b)), and the IMBIE2 data (The IMBIE Team, 2019). In addition to this we evaluate the reconstructed Kjeldsen et al. (2015) (data: Box et al. (2021)) and This Study data during the overlapping period 1986 through 2012.

5 Methods

The total mass balance for all of Greenland and all the different ROIs involves summing each field (SMB, MMB, BMB) by
 265 each ROI, then subtracting the MMB and BMB from the SMB fields, or,

$$MB = SMB - MMB - BMB. \quad (3)$$

Products that do not include the BMB term (i.e., Mouginit et al. (2019); Colgan et al. (2019), and Kjeldsen et al. (2015)) have total mass balance defined as

$$MB^* = SMB - MMB, \quad (4)$$

270 and when comparing This Study to those products, we compare like terms, never comparing our MB to a different product MB^* , except in Fig. 4 where all products are shown together.

Prior to calculating the mass balance, we perform the following steps.

5.1 Surface mass balance

In This Study we generate an output based on each of the three RCMs (HIRHAM/HARMONIE, MAR, and RACMO), however, 275 in addition to these we generate a final and 4th SMB field defined as a combination of 1) the adjusted reconstructed SMB from 1840 through 1985 (Sect. 5.4), and 2) the average of HIRHAM/HARMONIE, MAR, and RACMO from 1986 through a few months ago when RACMO drops are available, the average of HIRHAM/HARMONIE and MAR from a few months ago through yesterday, and MAR from yesterday through next week. See the Appendix A for differences among This Study MB and MB derived using each of the RCM SMBs. There is no obvious change or step function at the 1985 to 1986 reconstructed- 280 to-recent change, nor as the RACMO and then HIRHAM/HARMONIE RCMs become unavailable a few months ago and yesterday, respectively.

5.2 Projected marine mass balance

We project the marine mass balance from the last observed point from Mankoff et al. (2020b) (generally between 2 weeks and 1 month old) to seven days into the future at each glacier. We define the long-term trend as the linear least squares fit to the 285 last three years of observed data. We define the seasonal signal as the mean of the last three years of observed data during the temporal window of interest that spans from the most recently available observation through to present plus one week. We then assign the long term trend plus the seasonal signal value to next week's MMB (i.e., now + 7 days), and finally linearly interpolate between the last observed value and the forecasted next week's value, to provide the recent past projected and future forecasted MMB.

290 Marine mass balance does not change sign and changes magnitude by approximately 6 % annually over the entire ice sheet (King et al., 2018), but surface mass balance changes sign and has much larger and short-term variability. From this, the statistical forecast for marine mass balance described above does not impact results as much as the physically-based model forecast for surface mass balance.

5.3 Basal mass balance

295 Because Karlsson et al. (2021) provide a steady-state annual-average estimate of the BMB_{GF} and BMB_{friction} fields by 365 to estimate daily average. This is a reasonable treatment of the BMB_{GF} field, which does not have an annual cycle. The BMB_{friction} field does have a small annual cycle that matches the annual velocity cycle. However, when averaged over all of Greenland, this is only a ~6 % variation (King et al., 2018), and Karlsson et al. (2021) found that basal melt rates are 5 % higher for summer maps. Thus, the intra-annual changes are less than the uncertainty. The BMB_{VHD} field 300 varies significantly throughout the year, because it is proportional to surface runoff. We therefore generate our own BMB_{VHD} for this study.

To estimate recent BMB_{VHD} we use daily MAR runoff (see Mankoff et al. (2020a)) and BedMachine v4 (Morlighem et al., 2017, 2021) to derive subglacial routing pathways, similar to Mankoff and Tulaczyk (2017). We assume that all runoff travels to the bed within the grid cell where it is generated, the bed is pressurized by the load of the overhead ice, and the runoff

305 discharges on the day it is generated. We calculate subglacial routing from the gradient of the subglacial pressure head surface, h , defined as

$$h = z_b + k \frac{\rho_i}{\rho_w} (z_s - z_b), \quad (5)$$

with z_b the basal topography, k the flotation fraction (1), ρ_i the density of ice (917 kg m^{-3}), ρ_w the density of water (1000 kg m^{-3}), and z_s the ice surface. Eq. 5 comes from Shreve (1972), where the hydropotential has units of pascals (Pa), but here it is
310 divided by gravitational acceleration g times the density of water ρ_w to convert the units from pascals to meters (Pa to m).

We compute h and from h streams and outlets, and both the pressure and elevation difference between the source and outlet. The energy available for basal melting is the elevation difference (gravitational potential energy) and two-thirds of the pressure difference, with the remaining one third consumed to warm the water to match the changing phase transition temperature (Liestøl, 1956; Mankoff and Tulaczyk, 2017). We assume all energy, E_{VHD} (in Joules), is used to melt ice with

$$315 \quad b_m = E_{\text{VHD}} \rho_i^{-1} L^{-1}. \quad (6)$$

Because results are presented per ROI and to reduce the computational load of this daily estimate, we only calculate the integrated energy released between the RCM runoff source cell and the outlet cell, and then assign that to the ROI containing the runoff source cell.

To estimate reconstructed basal mass balance, we treat BMB_{GF} and $\text{BMB}_{\text{friction}}$ as steady state as described at the start of
320 this section. For BMB_{VHD} we use the fact that VHD comes from runoff by definition, and from this, reconstructed BMB_{VHD} is calculated using scaled runoff as a proxy. VHD theory suggests that a unit volume of runoff that experiences a 1000 m elevation drop will release enough heat to melt an additional 3 % (Liestøl, 1956). To estimate the scale factor we use the 1986 through 2012 overlap between Kjeldsen et al. (2015) runoff and This Study recent BMB_{VHD} from MAR runoff described above. The correlation between the two has an r^2 value of 0.75, slope of 0.03, and an intercept of -3 Gt yr^{-1} (Appendix D). From this, we
325 scale the Kjeldsen et al. (2015) reconstructed runoff by 3 % (from the 0.03 slope, unrelated to the theoretical 1000 m drop described earlier) to estimate reconstructed BMB_{VHD} .

5.4 Reconstructed adjustment

We use the reconstructed and recent surface (SMB) and marine (MMB) mass balance overlap from 1986 through 2012 to adjust the reconstructed data. This Study vs reconstructed SMB has a slope of 0.6 and an intercept of 165 Gt yr^{-1} (Fig. 3 SMB), and
330 This Study vs reconstructed MMB has a slope of 1.1 and an intercept of -6 Gt yr^{-1} (Fig. 3 MMB). The unadjusted reconstructed data slightly underestimates years with high SMB and overestimates years with low SMB (see 1986, 2010, 2011, and 2012 in Fig. 3 SMB). The unadjusted reconstructed data slightly overestimates years with low MMB and overestimates years with high MMB.

We adjust the reconstructed data until the reconstructed vs. recent slope is 1 and intercept is 0 Gt yr⁻¹ for each of the surface
335 and marine mass balance comparisons (Fig. 3). We then derive the BMB_{VHD} term for reconstructed basal mass balance (Sect.
5.3 and Appendix D), bring in the other BMB terms (Sect. 5.3), and use Eq. 3 to compute the adjusted reconstructed mass
balance.

For reconstructed SMB and MMB, the mean of the recent uncertainty is added to the reconstructed uncertainty during the
adjustment. Reconstructed MB uncertainty is then re-calculated as the square root of the sum of the squares of the reconstructed
340 SMB and MMB uncertainty.

For surface mass balance, the adjustment is effectively a rotation around the mean values, with years with low SMB de-
creasing and years with high SMB increasing after the adjustment. For marine mass balance, years with low MMB are slightly
reduced, and years with high MMB have a higher reduction to better match the overlapping estimates.

The adjustment described above treats all biases in the reconstructed data. The primary assumption of our adjustment is that
345 the bias contributions do not change in proportion to each other over time. We attribute the disagreement and need for the
adjustment to the demonstrated too-high biases in accumulation and ablation estimates in the 1840-2012 reconstructed SMB
field (Fettweis et al., 2020), an offset resulting from differences in ice masks (Kjeldsen et al., 2015), the inclusion of peripheral
glaciers (Kjeldsen et al., 2015), other accumulation rate inaccuracies (Lewis et al., 2017, 2019), and other unknowns.

5.5 Domains, boundaries, and regions of interest

350 Few of the ice masks used here are spatially aligned. The Zwally et al. (2012) sectors and the Mouginot and Rignot (2019)
regions are often smaller than the RCM ice domains. For example, the RACMO ice domain is 1,718,959 km², of which
1,696,419 km² (99 %) are covered by the Mouginot and Rignot (2019) regions, and 22,540 km² (1 %) are not, or 1,678,864
km² (98 %) are covered by the Zwally et al. (2012) and 40,095 km² (2 %) are not.

Cropping the RCM domain edges would remove the edge cells where the largest SMB losses occur. This effect is minor
355 when SMB is high (years with low runoff, assuming SMB magnitude is dominated by the runoff term). This effect is large
when SMB is low (years with high runoff). As an example for the 2010 decade, RACMO SMB has a mean of 251 Gt yr⁻¹ for
the decade, with a low of 45 Gt in 2019, a high of 420 Gt in 2018. For these same extreme years RACMO cropped to Mouginot
and Rignot (2019) has a low of 76 Gt (68 % high) and a high of 429 Gt (2 % high). RACMO cropped to Zwally et al. (2012)
has a low of 84 Gt (85 % high) and a high of 429 Gt (2 % high).

360 We therefore grow the ROIs to cover the RCM domains. ROIs are grown by expanding them outward, assigning the new cells
the value (ROI classification, that is sector number or region name, see Fig. 1) of the nearest non-null cell, and then clipping
to the RCM ice domain. This is done for each ROI and RCM. Appendix E provides a graphical display of the HIRHAM RCM
domain, the Mouginot and Rignot (2019) domain, and our expanded Mouginot and Rignot (2019) domain.

BMB_{VHD} comes from the MAR ice domain runoff, but is generated on the BedMachine ice thickness grid, which is smaller
365 than the ice domain in some places. Therefore, the largest runoff volumes per unit area (from the low-elevation edge of the ice
sheet) are discarded in these locations.

6 Product evaluation and assessment

We compare to six related data sets (see Table 2 and Sect. 4.5): The most similar and recent IO product (Mouginot et al., 2019), the previous PROMICE assessment (Colgan et al., 2019), the two mostly independent methods (GMB (Barletta et al., 2013) and VC (Simonsen et al., 2021a)), IMBIE2 (The IMBIE Team, 2019), and the unadjusted reconstructed/recent overlap (Kjeldsen et al., 2015).

Our initial comparison (Fig. 4) shows all seven products overlaid in a time series accumulating at the product resolution (daily to annual) from the beginning of the first overlap (1972, Mouginot et al. (2019)) until seven days from now (now defined as 2021-08-13). Each data set is manually aligned vertically so that the last timestamps appear to overlap, allowing disagreements to grow back in time. We also assume errors are smallest at present and allow errors to grow back in time. The errors for this product are described in the Uncertainty section.

In the sections below, we compare This Study to each of the validation data in more detail. The Mouginot et al. (2019) and Colgan et al. (2019) products allow term-level (SMB, MMB, and MB^{*}) comparison, and the GMB, VC, and IMBIE2 only MB-level comparison. The MB or MB^{*} comparison for each product is summarized in Table 2. All have different masks. Bias [Gt yr⁻¹] is defined as $\frac{1}{n} \sum_{i=1}^n (x_i - y_i)$. RMSE [Gt yr⁻¹] is defined as $\sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - y_i)^2}$. Sums are computed using ice-sheet wide annual values, where x is This Study, y is the other product, and a positive bias means that This Study has a larger value.

Table 2. Summary of correlation, bias, and RMSE between different products during their overlap periods with This Study. Basal mass balance not included in This Study when comparing against Mouginot and Rignot (2019), Colgan et al. (2019), or Kjeldsen et al. (2015). Peripheral ice masses never included in This Study.

Other product	r ²	bias	RMSE	Fig.	Overlap	Notes
Mouginot et al. (2019)	0.94	9	36	5	1986 – 2018	No basal mass balance
Colgan et al. (2019)	0.87	-35	61	6	1995 – 2015	No basal mass balance
GMB	0.86	30	62	7	2002 – 2020	Includes peripheral masses
VC	0.62	-14	85	7	1992 – 2019	Multi-year smooth
IMBIE2	0.89	-10	44	7	1992 – 2018	No BMB when using IO; BMB when using GMB or VC
Kjeldsen et al. (2015)	0.80	3	59	3	1986 – 2012	No basal mass balance; Includes peripheral masses

6.1 Mouginot (2019)

The Mouginot et al. (2019) product spans the 1972 through 2018 period. The SMB comes from RACMO v2.3p2 downscaled at 1 km, and agrees very well with SMB from This Study (r² 0.94, bias 9, RMSE 36, slope 1.1). The minor SMB differences are likely due to mask differences, or our use of a three-RCM average SMB estimate.

Mouginot et al. (2019) discharge and our MMB from Mankoff et al. (2020b) have a -33 Gt yr⁻¹ bias. This difference can mainly be attributed to different discharge estimates in the Southeast and Central east sector (Appendix: Mouginot regions). When we include BMB in This Study (diamonds in middle panel shifting values to the right), it adds ~25 Gt yr⁻¹ to This Study.

Because MB^* is a linear combination of SMB and MMB terms (Eq 4), the MB^* differences between this product and Mouginito et al. (2019) are dominated by the MMB term, although it is not apparent because interannual variability is dominated by SMB.

6.2 Colgan (2019)

The Colgan et al. (2019) product spans 1995 through 2015. The SMB term is broadly similar to the RCM-averaged SMB term in This Study, although Colgan et al. (2019) use only an older version of MAR (Fig. 6 top panel). The Colgan et al. (2019) SMB is spatially interpolated over the PROMICE ice-sheet ice mask (Citterio and Ahlstrøm, 2013), which contains more detail on the ice sheet periphery, and therefore a larger ablation area than the native coarser MAR ice mask. This Study does not interpolate the SMB field and instead works on the SMB ice domain and.

The most critical difference between This Study and Colgan et al. (2019) is that the latter estimate grounding line ice discharge based on corrections to ice volume flow rate measured across the approximately 1700 m elevation contour. This ice volume flow rate measurement is far inland relative to the grounding line flux gates used in This Study (from Mankoff (2020)). This introduces uncertainty in the Colgan et al. (2019) MMB term induced by the SMB corrections between the 1700 m elevation contour and the terminus (see large disagreement in Fig. 6 mid panel). This disagreement increases when BMB is included in the results of This Study (shown by dots shifting the annual values to the right).

The MMB disagreement is represented differently across sectors (Appendix: Colgan 2019), where sectors 1, 2, 5, and 6 all have correlation coefficients less than ~ 0.1 , while the remaining sectors 3, 4, 7, and 8 all have correlation coefficients greater than 0.5.

This Study assesses greater MMB bias (43 Gt yr^{-1}) than Colgan et al. (2019). While Colgan et al. (2019) did not assess BMB, the majority of this discrepancy likely results from Colgan et al. (2019) aliasing the aforementioned downstream correction terms. For example, while This Study shows very little interannual variability in ice discharge in the predominantly land-terminating SW region, Colgan et al. (2019) infer large interannual variability in ice discharge based on large interannual variability in SMB and changes in ablation area ice volume in their Sector 6. The discrepancy between This Study and Colgan et al. (2019) MMB [+BMB] is largest during the earliest part of the record (i.e. 1995-2000), decreasing towards present-day, which may suggest that Colgan et al. (2019) particularly overestimated the response in ice discharge to 1990s climate variability.

Similar to the comparison with Mouginito et al. (2019), the disagreement between This Study and Colgan et al. (2019) is dominated by MMB disagreement, although it is again not apparent because interannual variability is dominated by SMB.

6.3 Gravimetric Mass Balance (GMB)

Unlike this Study, the GMB method includes mass losses and gains on peripheral ice masses which should introduce a bias of ~ 10 to 15% (Colgan et al., 2015; Bolch et al., 2013). The inclusion of peripheral ice in the GMB product is because the spatial resolution is so low that it cannot distinguish between them and the main ice sheet. There is also signal leakage from other glaciated areas, e.g., the Canadian Arctic. This can have an effect on the estimated signal, especially in sectors 1 and 8

or regions NW and NO. There is also leakage between basins, which becomes a larger issue for smaller basins or where major outlet glaciers are near basin boundaries. GMB may also have an amplified seasonal signal due to changing snow loading in the surrounding land areas that may be mapped as ice sheet mass change variability. This would enhance the seasonal amplitude but not have an impact on the interannual mass change rates. Additionally, different glacial isostatic adjustment (GIA) corrections applied to the gravimetric signal may also lead to differences in GMB estimates on ice sheet scale, but also on sector scale (e.g. Sutterley et al. (2014); Khan et al. (2016)).

GMB and the IO method (This Study) both report changes in ice sheet mass, but they are measuring two fundamentally different things. The IO method tracks volume flow rate across the ice sheet boundaries. Typically this is meltwater across the ice sheet surface and solid ice across flux gates near the calving edge of the ice sheet, and in This Study also meltwater across the ice sheet basal boundary. That volume is then converted to mass. We consider that mass is ‘lost’ as soon as it crosses the boundary (i.e. the ice melts or ice crosses the flux gate). The GMB method tracks the regional mass changes. Melting ice has no impact on this, until the meltwater enters the ocean and a similar mass leaves the far-field GMB footprint. From these differences, the GMB method may be a better estimate of sea level rise, while the IO method may be a better representation of the state of the Greenland ice sheet.

6.4 Volume Change (VC)

When deriving surface elevation change from satellite altimetry, data from multiple years are needed to give a stable ice sheet-wide prediction. Hence, the altimetric mass balance estimates are often reported as averages of single satellite missions.

Although This Study has a small (-14 Gt yr^{-1}) bias in comparison to Simonsen et al. (2021a) VC, there is a relatively high RMSE of 85 Gt yr^{-1} and a mid-range correlation ($r^2 = 0.62$). This suggests that while both This Study and VC agree on the total mass loss of the ice sheet, they disagree on the precise temporal distribution of this mass loss. It is possible the outlying 1992 and 2019 years are influenced by the edge of the time series record if not fully sampled, but other outliers exist - the 1992 extreme low melt year and the 2019 extreme melt year, as well as the 1995 through 1998 period, stand out as years with poor agreement.

We suggest that this is due to climate influences on the effective radar horizon across the ice sheet during these years. Weather-driven changes in the effective scatter horizon, mapped by Ku-band in the upper snow layer of ice sheets hampers the conversion of radar-derived elevation change into mass change (Nilsson et al., 2015). Simonsen et al. (2021a) used a machine learning approach to derive a temporal calibration field for converting the radar elevation change estimates into mass change. This approach relied on precise mass balance estimates from ICESat to train the model and thereby was able to remove the effects of the changing scattering horizon in the radar data. This VC mass balance is given for monthly time steps (Simonsen et al., 2021a), however the running-mean applied to derive radar elevation change will dampen the interannual variability of the mass balance estimate from VC. This is especially true prior to 2010, after which the novel radar altimeter onboard CryoSat-2 allowed for a shortening of the data windowing from 5 to 3 years. This smoothing of the interannual variability is also seen in the intercomparison between This Study and the VC MB, where in addition to the two end members of the time series (1992

455 and 2019) the years 1995, 1996, and 1998 seem to be outliers (Fig. 7). These years are notable for high MB which seems to be captured less precisely by the older radar altimeters due to the longer temporal averaging.

6.5 IMBIE

The most widely cited estimate of Greenland mass balance today is the Ice-Sheet Mass Balance Inter-Comparison Exercise 2 (IMBIE2, The IMBIE Team (2019)). IMBIE2 seeks to provide a consensus estimate of monthly Greenland mass balance
460 between 1992 and 2018 that is derived from altimetry, gravimetry, and input-output ensemble members. There are two critical methodological differences between This Study and IMBIE2. Firstly, the gravimetry members of IMBIE2 assess mass balance of all Greenlandic land ice, including peripheral ice masses, while This Study only assesses mass balance of the ice sheet proper. Secondly, the input-output members of IMBIE2 do not assess BMB, while This Study does.

The IMBIE2 composite record of ice-sheet mass balance equally weights three methods of assessing ice-sheet mass balance:
465 input-output, altimetry and gravimetry. Prior to c. 2003, however, IMBIE2 is derived solely from IO studies that explicitly exclude BMB (MB is actually MB^{*}). After c. 2003, by comparison, IMBIE2 includes both satellite altimetry and gravimetry records implicitly sample BMB. The representation of BMB in the composite IMBIE2 mass balance record therefore shifts before and after c. 2003.

In comparison to mass balance assessed by IMBIE2, This Study has a small bias of $\sim -10 \text{ Gt yr}^{-1}$ over the 26 calendar year
470 comparison period. This apparent agreement may be attributed to the compensating effects of IMBIE2 effectively sampling peripheral ice masses and ignoring BMB, while This Study does the opposite and ignores peripheral ice masses but samples BMB, equal to $\sim 25 \text{ Gt yr}^{-1}$. Over the entire 26-year comparison period, the RMSE with IMBIE2 is 44 Gt yr^{-1} and the correlation is 0.89 . This relatively high correlation highlights good agreement in interannual variability between studies, and the RMSE suggests that formal stated uncertainties of each study (c. ± 30 to $\pm 63 \text{ Gt yr}^{-1}$ for IMBIE2 and mean of 86 Gt yr^{-1}
475 for This Study) are indeed good estimates of the true uncertainty, as assessed by inter-study discrepancies.

7 Uncertainty

We treat the three inputs to the total mass balance (surface, marine, and basal mass balance, or SMB, MMB, and BMB) as independent when calculating the total error. This is a simplification – the RCM SMB and the BMB_{VHD} from RCM runoff are related, and MMB ice thickness and BMB_{VHD} pressure gradients are related, and other terms may have dependencies. However,
480 the two dominant IO terms, SMB inputs and MMB outputs, are independent on annual time scales, and for simplification we treat all terms as independent. We use Eq 3 and standard error propagation for SMB, MMB, and BMB terms (i.e., the square root of the sum of the squares of the SMB plus MMB plus BMB error terms). For the MMB, extra work is done to calculate uncertainty between the last Mankoff 2020 MMB data (up to 30 days old, with error of $\sim 9\%$ or $\sim 45 \text{ Gt yr}^{-1}$) and the forecasted now-plus-7-day MMB (see Sect. 7.1). Table 3 provides a summary of the uncertainty for each input.

485 The final This Study MB uncertainty value shown in Table 3 comes from mean of the annual sum of the MB error term.

Term	Uncertainty [\pm]	Notes
HIRHAM / HARMONIE SMB	15 %	Langen et al. (2017). The mean accumulation bias (-5%) and ablation bias (-7%) tend to cancel out, but this cannot be expected to be the case on single-basin, short-term scales where uncertainty is estimated to be larger.
MAR SMB	15 %	Fettweis et al. (2020). The mean bias between the model and the measurements was 15 % with a maximum of 1000 mmWE yr ⁻¹ . GrSMBMIP uses integrated values over several months of SMB, suggesting larger uncertainty of modeled runoff at the daily timescale.
RACMO SMB	15 %	Noël et al. (2019). Average 5% runoff bias compared to annual cumulative discharge from the Watson River. Increases to a maximum of 20 % for extreme runoff years.
This Study SMB	9 %	Average of 15 % SMB uncertainties above, assuming uncorrelated.
Reconstructed SMB	~ 20 %	From Kjeldsen et al. (2015) Table 1.
Recent MMB	~ 45 Gt yr ⁻¹	~ 9 %. Mankoff et al. (2020b) updated (Mankoff, 2021).
Reconstructed MMB	~ 10 %	From Kjeldsen et al. (2015) Table 1.
BMB _{GF}	50 %	5.3 +4/-1.4 Gt yr ⁻¹ from Karlsson et al. (2021) Table 1, using the average of the three available methods.
BMB _{friction}	20 %	11.8 \pm 3.4 Gt yr ⁻¹ from Karlsson et al. (2021) Table 1.
BMB _{VHD}	15 %	MAR runoff uncertainty.
This Study MB	~ 86 Gt yr ⁻¹	Eq 3, assuming all uncertainty is uncorrelated.

Table 3. Summary of uncertainty estimates for products used in This Study. This is an approximate and simplified representation – RCM uncertainties are calculated separately for gain and loss terms, because SMB near 0 does not mean uncertainty is near 0. This is also why the final This Study uncertainty is presented with units [Gt yr⁻¹].

7.1 Marine mass balance

The MMB uncertainty is discussed in detail in Mankoff et al. (2020b), but the main uncertainties come from unknown ice thickness, the assumption of no vertical shear at fast-flowing marine-terminating outlet glaciers, and ice density of 917 kg m⁻³. Regional ice density can be significantly reduced by crevasses. For example, Mankoff et al. (2020c) identified a snow-covered
490 crevasse field with 20 % crevasse density, meaning at that location regional firn density should be reduced by 20 %.

Temporally, MMB at daily resolution comes from ~12 day observations up-sampled to daily, and those ~12 day resolution observations come from longer time period observations (Solgaard et al., 2021). Because the velocity method using feature tracking, it is correct on average but misses variability within each sample period (e.g., Greene et al. (2020)).

Spatially, MMB discharge is estimated ~5 km upstream from the grounding lines for ice velocities as low as 100 m yr⁻¹. That
495 ice accelerates toward the margin, but even ice flowing steady at 1 km yr⁻¹ would take 5 years before that mass is lost. However, at any given point in time, ice that had previously crossed the flux gate is calving or melting into the fjord. The discrepancy here between the flux gate estimated mass loss and the actual mass lost at the downstream terminus is only significant for

glaciers that have had large velocity changes at some point in the recent past, large changes in ice thickness, or large changes in the location (retreat or advance) of the terminus. We do not consider SMB changes downstream of the flux gate, because the gates are temporally near the terminus for most of the ice that is fast-flowing, and the largest SMB uncertainty is at the ice sheet margin where there are both mask issues and high topographic variability.

The forecasted MMB uncertainty is estimated from the same forecasted calendar days of the last three years of observed MMB at each glacier. MMB uncertainty for the first forecasted day is the baseline MMB uncertainty plus the largest daily change observed during the forecast period from the last three years. The MMB uncertainty for the second forecasted day is the baseline uncertainty plus the uncertainty from the first forecasted day plus the second largest daily change observed during the forecast period from the last three years. We repeat this for the approximately 30 days of forecasted MMB. This implementation takes into account the larger variability (uncertainty) during the seasonal transition between the lower winter and higher summer discharge, or the smaller variability during the winter period.

7.2 Regions of interest (ROI)

We work on the three different domains of the three RCMs, and expand the ROIs to match the RCMs (see Appendix E). However, some alignment issues cannot be solved. For example, we use BedMachine ice thickness to estimate BMB_{VHD} . Often, the largest BMB_{VHD} occurs near the ice margin under ice with the steepest surface slopes. This is also where the largest runoff often occurs, because the ice margin, at the lowest elevations, is exposed to the warmest air. If these RCM ice grid cells with high runoff are anywhere inside the BedMachine ice domain, that runoff is still included in our BMB_{VHD} estimates because it flows outward and passes through the BedMachine near-ice-edge grid cells with the large pressure gradients. However, if these RCM ice grid cells with high runoff are outside the BedMachine ice domain (ice thickness is 0), there is no reasonable way to include that runoff in our BMB_{VHD} budget, and these grid cells are ignored.

The MAR ice domain is 1,828,800 km² of which 1,711,200 km² (94 %) are covered by the BedMachine ice mask, and 26,000 km² (6 %) are not. This 6 % area contributes ~15 % of the runoff (excluded from the VHD calculations) and likely a higher percent of the VHD, because the border region of the ice sheet has the steepest gradients and the largest volume of subglacial flow.

We encourage RCM developers, BedMachine, and others to use a common and up-to-date mask (see Kjeldsen et al. (2020)).

7.3 Accumulating uncertainties

When accumulating errors as in Fig. 4, we use only the MMB and BMB_{GF} uncertainty. The MMB uncertainty is primarily due to unknown ice thickness and is invariant in time, and the geothermal heat flux is steady state. SMB uncertainty is assumed to have errors randomly distributed in time (for the purposes of Fig. 4). There may be time-invariant biases in the $BMB_{friction}$ and SMB fields, but treating all uncertainties as biases is incorrect - evidence for that comes from the six other MB estimates. This distinction between bias and random uncertainty is only done for Fig. 4 where errors accumulate in time. The provided data product contains one uncertainty field and does not distinguish between systematic and random uncertainty.

530 The shaded region in Fig. 4 representing the uncertainty for This Study is computed as a 365 day rolling smooth from 1840 through 1999 of the above-described uncertainty, 1/365th of the annual error at now + 7 days, and a linear blend, from 2000 to now + 7 days, between the smoothed reconstructed uncertainty and the present and future more variable uncertainty.

The Mouginot et al. (2019), Colgan et al. (2019), and Kjeldsen et al. (2015) products all provide an error estimate, but do not distinguish between temporally fixed errors (biases; should accumulate in time) vs. temporally random errors.

535 We treat the Mouginot et al. (2019) data the same as This Study. Marine mass balance uncertainty is treated as a bias and accumulates, and surface mass balance uncertainty is treated as random and does not accumulate.

The Colgan et al. (2019) vs. this study bias and RMSE are -35 and 61 Gt yr^{-1} respectively. This suggests that in any given year, there could be up to -35 ± 61 or $+26/-96 \text{ Gt yr}^{-1}$ departure from This Study. From this, we assign a 35 Gt yr^{-1} bias (36 %; accumulates in time) and a 61 Gt yr^{-1} RMSE (64 %; random in time).

540 The adjusted Kjeldsen et al. (2015) data have 0 surface and marine mass balance bias by definition (Sect. 5.4), but Fig. 4 displays the unadjusted data, and we apply a 36 Gt yr^{-1} accumulating uncertainty from the unadjusted MMB bias (Fig. 3).

7.4 Peripheral ice masses

Greenland's peripheral glaciers and ice caps are not included in this product. Nonetheless, we briefly summarize recent mass balance estimates of these areas. Greenlandic peripheral ice contributes more runoff per unit area than the main ice sheet –
545 they are $< 5 \%$ of the total ice area but contribute ~ 15 to 20% of the whole island mass loss (Bolch et al., 2013). From 2003 to 2009 and using the VC method (altimetry), Gardner et al. (2013) estimate $-38 \pm 7 \text{ Gt yr}^{-1}$ peripheral mass balance. From 2006 to 2016 and using the VC method (DEM differencing), Zemp et al. (2020) estimate $-51 \pm 17 \text{ Gt yr}^{-1}$ peripheral mass balance, using Rastner et al. (2012) delineations.

8 Results

550 From the 181 years of data, the mean mass balance is $-80 \pm 125 \text{ Gt yr}^{-1}$, with a minimum of $-429 \pm 109 \text{ Gt}$ in 2012 (SMB of $86 \pm 8 \text{ Gt}$, MMB of $485 \pm 46 \text{ Gt}$, BMB of $29 \pm 6 \text{ Gt}$) and a maximum of $137 \pm 83 \text{ Gt yr}^{-1}$ in 1996 (SMB of $578 \pm 52 \text{ Gt}$, MMB of $420 \pm 39 \text{ Gt}$, BMB of $21 \pm 5 \text{ Gt}$).

At the decadal average, the following trends are apparent. Surface mass balance has decreased from a high of $\sim 450 \text{ Gt yr}^{-1}$ in the 1860s to low of $\sim 260 \text{ Gt yr}^{-1}$ in the 2010s. SMB variability has also increased during this time. Marine mass balance has
555 increased slightly from a low of $\sim 375 \text{ Gt yr}^{-1}$ in the 1860s to a high of $\sim 490 \text{ Gt yr}^{-1}$ in the 2010s. Basal mass balance, from runoff as a proxy, had a high of $26 \pm 16 \text{ Gt yr}^{-1}$ in the 1930s and a low of $22 \pm 5 \text{ Gt yr}^{-1}$ in the 1990s, but as with runoff, is increasing in recent decades.

The total mass balance decadal trend from the 1840s through the 2010s is one of general mass decrease and increased intra-decadal variability. The record begins in the 1840s with $\sim -10 \text{ Gt yr}^{-1}$, has only one (of 19) decades with a mass gain ($\sim 50 \text{ Gt}$
560 yr^{-1} in the 1860s), and a record low of $\sim -250 \text{ Gt yr}^{-1}$ in the 2010s.

9 Data availability

The RCM surface mass balance, and the VHD basal mass balance components are updated daily, the marine mass balance approximately every 12 days, and all are used to produce the final daily-updating product. The data area available at <https://doi.org/10.22008/FK2/YG3IWC> (Mankoff et al., 2021), with all historical (daily updated) versions archived.

565 As part of our commitment to make continual and improving updates to the data product, we introduce a GitHub database (https://github.com/GEUS-Glaciology-and-Climate/mass_balance/; last visited August 13, 2021) where users can track progress, make suggestions, discuss, report and respond to issues that arise during use of this product.

10 Conclusions

This study is the first to provide a dataset containing more than a century and real time estimates detailing the state of Greenland
570 ice sheet mass balance, with regional or sector spatial and daily temporal resolution products of surface mass balance, marine mass balance, basal mass balance, and the total mass balance.

IMBIE2 highlights that during the GRACE satellite gravimetry era (2003 through 2017), there are usually more than twenty independent estimates of annual Greenland ice sheet mass balance. Just two independent estimates, however, are available prior to 2003. This study will therefore provide additional insight on ice sheet mass balance during the late 1980s and 1990s.
575 IMBIE2 also highlights how the availability of mass balance estimates declines in the year prior to IMBIE2 publication. This reflects a lag period during which mass balance assessments from non-operational products are undergoing peer-review. The operational nature of this product supports the timely inclusion of annual MB estimates in community consensus reports such as those from IMBIE and the IPCC.

As such, the data products provided in this study present the first operational monitoring of the Greenland ice sheet total mass
580 balance and its components. One property of the input-output approach used in This Study is the explanatory capabilities of the data products, allowing scrutiny of the physical origins of recorded mass changes. By excluding peripheral ice masses, this study allows and invites anyone to keep an eye on the current evolution of the Greenland ice sheet proper. However, as the spatial resolution of RCMs increase and estimates of peripheral ice thickness become available, our setup allows inclusion of these ice masses to generate a full Greenland-wide product. Moreover, as the determination of each of the individual components of the
585 ice sheet mass balance is expected to improve over time through international research efforts, the total mass balance product presented will also be able to improve, as it is sustained by the Danish-Greenlandic governmental long-term monitoring effort – the Programme for Monitoring of the Greenland ice sheet (PROMICE).




Figure 3. Comparison between This Study and the reconstructed (Kjeldsen et al., 2015). All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the years for the unadjusted data sets. The matching colored diamonds show the adjusted data. MB^* shown here does not include BMB for either the reconstructed or This Study data. Arrows show statistical properties before and after the adjustment. No adjustment is made to MB^* , but it is computed from Eq. 4 both before (numbered) and after (diamonds) the surface and marine mass balance adjustments.




Figure 4. Comparison between This Study and other mass balance time series. Note that various products do or do not include basal mass balance or peripheral ice masses (see Table 2). This Study annual-resolution data prior to 1986 is the Kjeldsen et al. (2015) data adjusted as described in Sect. 5.4. Sea level rise calculated as $-Gt/361.8$. Inset highlights changes since 2010.




Figure 5. Comparison of This Study vs Mouginot et al. (2019). All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the year. Matching colored diamonds show the data when BMB is added to This Study. Printed numbers (r^2 , bias, RMSE, slope) compare values without BMB.




Figure 6. Comparison of This Study vs Colgan et al. (2019). All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the year. Matching colored diamonds show the data when BMB is added to This Study. Printed numbers (r^2 , bias, RMSE, slope) compare values without BMB.




Figure 7. This Study total mass balance (MB) vs. the gravimetric method (GMB), volume change method (VC) and IMBIE2 estimates of MB. All three include BMB. All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the year. GRACE and IMBIE2 include peripheral ice masses.

Appendix A: RCM differences




Figure A1. Comparison of This Study combined RCM product and the HIRHAM/HARMONIE, MAR, and RACMO RCMs. Results shown here are MB, not SMB, but the same MMB and BMB have been subtracted from each SMB product. Top panel: annual MB for entire time series. Middle panel: Example two years (2019 and 2020) at daily resolution. Bottom panel: Difference between the three RCM MB products and This Study RCM-averaged product, for the same data shown in the middle panel.

Appendix B: Mougnot 2019 by region




Figure B1. Comparison between This Study (excluding BMB) and Mouginot et al. (2019). Same data and display as Fig. 5 except here displayed by Mouginot and Rignot (2019) region. Numbers in each graph show r^2 , bias, and RMSE, from top to bottom, respectively. All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the year.




Figure C1. Comparison between This Study (excluding BMB) and Colgan et al. (2019). Same data and display as Fig. 6 except here displayed by Zwally et al. (2012) sector. Numbers in each graph show r^2 , bias, and RMSE, from top to bottom, respectively. All axes units are Gt yr^{-1} . Plotted numbers represent the last two digits of the year.

Appendix D: Reconstructed runoff




Figure D1. Comparison between MAR runoff and basal viscous heat dissipation derived from that runoff. The slope is used to estimate the reconstructed BMB_{VHD} from reconstructed runoff (see Sect. 5.3). Axes units are Gt yr⁻¹. Plotted numbers represent the last two digits of the year.

Appendix E: RCM coverage




Figure E1. HIRHAM RCM coverage by Mougino and Rignot (2019). Coverage of HIRHAM by Zwally et al. (2012), and MAR and RACMO by Mougino and Rignot (2019) and Zwally et al. (2012) is similar to graphic shown here (See section 5.5 for discussion of RACMO coverage issues). HIRHAM latitude and longitude covers the equator because we work on the native HIRHAM rotated pole coordinate system.

Appendix F: Software

This work was performed using only open-source software, primarily GRASS GIS (Neteler et al., 2012), CDO (Schulzweida, 2019), NCO (Zender, 2008), GDAL (GDAL/OGR contributors, 2020), and Python (Van Rossum and Drake Jr, 1995), in particular the Jupyter (Kluyver et al., 2016), dask (Dask Development Team, 2016; Rocklin, 2015), pandas (McKinney, 2010), geopandas (Jordahl et al., 2020), numpy (Oliphant, 2006), x-array (Hoyer and Hamman, 2017), and Matplotlib (Hunter, 2007) packages. The entire work was performed in Emacs (Stallman, 1981) using Org Mode (Schulte et al., 2012) on GNU/Linux and using many GNU utilities. The parallel (Tange, 2011) tool was used to speed up processing.

Appendix G: CRediT




Figure G1. Author contributions following the CRediT system (Allen et al., 2014; Brand et al., 2015; Allen et al., 2019)

Author contributions. {

Author contribution is captured following the CRediT system (Allen et al., 2014; Brand et al., 2015; Allen et al., 2019) and shown graphically in Figure G1. The following authors contributed in the following ways. Conceptualization: KDM, APA, and RSF. Curation: KDM, XF, PL, MS, KK, NBK, BN, MvdB, AS, and JB. Implementation: KDM, XF, PL, KK, and MKD. Funding: AS, APA, SBA, and RSF. SMB methods: XF, PL, BN, and MvdB. MMB methods: KDM, WC, AS, MKD, APA, and RSF. BMB methods: NBK and KDM. Validation (general): KDM. Validation GRACE: WC. Validation VC: WC and SS. Reconstruction methods: KK, JB, and KDM. Project admin: KDM, APA, SBA, and RSF. Resources: KDM, XF, PL, MS, KK, NBK, BN, MvdB, AS, and SBA. Software: KDM, XF, PL, AS, and MKD. Visualization: KDM. Writing: KDM, XF, PL, MS, KK, NBK, BN, MvdB, WC, JB, SS, APA, and RSF. }

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The editor, one anonymous reviewer (Anonymous, 2021), and Andrew Shepherd (Shepherd, 2021) provided feedback and helped improve this paper.

Financial Support: Funding was provided by the Programme for Monitoring of the Greenland ice sheet (PROMICE). Parts of this work
615 were funded by the INTAROS project under the European Union's Horizon 2020 research and innovation program under grant agreement no.
727890. B. Noël was funded by the NWO VENI grant VI.Veni.192.019. MvdB acknowledges support from the Netherlands Earth System
Science Centre (NESSC).

References

- Allen, L., Scott, J., Brand, A., Hlava, M., and Altman, M.: Publishing: Credit where credit is due, *Nature*, 508, 312–313, 620 <https://doi.org/10.1038/508312a>, 2014.
- Allen, L., O’Connell, A., and Kiermer, V.: How can we ensure visibility and diversity in research contributions? How the Contributor Role Taxonomy (CRediT) is helping the shift from authorship to contributorship, *Learned Publishing*, 32, 71–74, <https://doi.org/10.1002/leap.1210>, 2019.
- Amory, C., Kittel, C., Le Toumelin, L., Agosta, C., Delhasse, A., Favier, V., and Fettweis, X.: Performance of MAR (v3.11) in simulating 625 the drifting-snow climate and surface mass balance of Adélie Land, East Antarctica, *Geoscientific Model Development*, 14, 3487–3510, <https://doi.org/10.5194/gmd-14-3487-2021>, 2021.
- Andresen, C. S., Straneo, F., Ribergaard, M. H., Bjørk, A. A., Andersen, T. J., Kuijpers, A., Nørgaard-Pedersen, N., Kjær, K. H., Schjøth, F., Weckström, K., and Ahlstrøm, A. P.: Rapid response of Helheim Glacier in Greenland to climate variability over the past century, *Nature Geoscience*, 5, 37 – 41, <https://doi.org/10.1038/NGEO1349>, 2012.
- 630 Anonymous: RC1: Comment on essd-2021-131, <https://doi.org/10.5194/essd-2021-131-rc1>, 2021.
- Barletta, V. R., Sørensen, L. S., and Forsberg, R.: Scatter of mass changes estimates at basin scale for Greenland and Antarctica, *The Cryosphere*, 7, 1411–1432, <https://doi.org/10.5194/tc-7-1411-2013>, 2013.
- Barletta, V. R., Sørensen, L. S., Simonsen, S. B., and Forsberg, R.: Gravitational mass balance of Greenland 2003 to present v2.1, <https://doi.org/10.11583/DTU.12866579.v1>, 2020.
- 635 Bengtsson, L., Andrae, U., Aspelien, T., Batrak, Y., Calvo, J., de Rooy, W., Gleeson, E., Hansen-Sass, B., Homleid, M., Hortal, M., Ivarsson, K.-I., Lenderink, G., Niemelä, S., Nielsen, K. P., Onvlee, J., Rontu, L., Samuelsson, P., Muñoz, D. S., Subias, A., Tijm, S., Toll, V., Yang, X., and Kjøltzow, M. : The HARMONIE–AROME Model Configuration in the ALADIN–HIRLAM NWP System, *Monthly Weather Review*, 145, 1919–1935, <https://doi.org/10.1175/mwr-d-16-0417.1>, 2017.
- Bjørk, A. A., Kjær, K. H., Korsgaard, N. J., Khan, S. A., Kjeldsen, K. K., Andresen, C. S., Box, J. E., Larsen, N. K., and Funder, S.: An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland, *Nature Geoscience*, 5, 427 – 432, 640 <https://doi.org/10.1038/NGEO1481>, 2012.
- Boers, N. and Rypdal, M.: Critical slowing down suggests that the western Greenland Ice Sheet is close to a tipping point, *Proceedings of the National Academy of Sciences*, 118, e2024192 118, <https://doi.org/10.1073/pnas.2024192118>, 2021.
- Bolch, T., Sørensen, L. S., Simonsen, S. B., Mölg, N., Machguth, H., Rastner, P., and Paul, F.: Mass loss of Greenland’s glaciers and ice caps 645 2003-2008 revealed from ICESat data, *Geophysical Research Letters*, 40, 875–881, <https://doi.org/10.1002/grl.50270>, 2013.
- Box, J., Colgan, W. T., and Kjeldsen, K.: Greenland SMB, D and TMB annual time series 1840-2012, <https://doi.org/10.22008/FK2/L1QOZ5>, 2021.
- Box, J. E.: Greenland ice sheet mass balance reconstruction. Part II: surface mass balance (1840-2010), *Journal of Climate*, 26, 6974 – 6989, <https://doi.org/10.1175/JCLI-D-12-00518.1>, 2013.
- 650 Box, J. E. and Colgan, W.: Greenland ice sheet mass balance reconstruction. Part III: marine ice loss and total mass balance (1840-2010), *Journal of Climate*, 26, 6990 – 7002, <https://doi.org/10.1175/JCLI-D-12-00546.1>, 2013.
- Box, J. E., Cressie, N., Bromwich, D. H., Jung, J.-H., van den Broeke, M. R., van Angelen, J. H., Forster, R. R., Miège, C., Mosley-Thompson, E., Vinther, B., and McConnell, J. R.: Greenland ice sheet mass balance reconstruction. Part I: Net snow accumulation (1600–2009), *Journal of Climate*, 26, 3919–3934, <https://doi.org/10.1175/JCLI-D-12-00373.1>, 2013.

- 655 Brand, A., Allen, L., Altman, M., Hlava, M., and Scott, J.: Beyond authorship: attribution, contribution, collaboration, and credit, *Learned Publishing*, 28, 151–155, <https://doi.org/10.1087/20150211>, 2015.
- Cappelen, J.: The observed climate of Greenland, 1958-99-with climatological standard normals, 1961-90, Danish Meteorological Institute, 2001.
- Cappelen, J., Laursen, E. V., Jørgensen, P. V., and Kern-Hansen, C.: DMI Monthly Climate Data Collection 1768-2005: Denmark, The Faroe
660 Islands and Greenland, DMI, 2006.
- Cappelen, J., Laursen, E., Jørgensen, P., and Kern-Hansen, C.: DMI monthly climate data collection 1768-2010, Denmark, the Faroe Islands and Greenland, Tech. rep., DMI Technical Report 11-05. Copenhagen, 2011.
- Christensen, O. B., Drews, M., Hesselbjerg Christensen, J., Dethloff, K., Ketelsen, K., Hebestadt, I., and Rinke, A.: The HIRHAM Regional Climate Model. Version 5 (beta), no. 06-17 in Denmark. Danish Meteorological Institute. Technical Report, Danish Climate Centre,
665 Danish Meteorological Institute, 2007.
- Citterio, M. and Ahlstrøm, A. P.: Brief communication “The aerophotogrammetric map of Greenland ice masses”, *The Cryosphere*, 7, 445–449, <https://doi.org/10.5194/tc-7-445-2013>, 2013.
- Colgan, W.: Greenland ice sheet mass balance assessment (1995-2019), <https://doi.org/10.22008/FK2/XOTO3K>, 2021.
- Colgan, W., Abdalati, W., Citterio, M., Csatho, B., Fettweis, X., Luthcke, S., Moholdt, G., Simonsen, S. B., and Stober, M.: Hybrid glacier
670 Inventory, Gravimetry and Altimetry (HIGA) mass balance product for Greenland and the Canadian Arctic, *Remote Sensing of Environment*, 168, 24–39, <https://doi.org/10.1016/j.rse.2015.06.016>, 2015.
- Colgan, W., Mankoff, K. D., Kjeldsen, K. K., Bjørk, A. A., Box, J. E., Simonsen, S. B., Sørensen, L. S., Khan, S. A., Solgaard, A. M., Forsberg, R., Skourup, H., Stenseng, L., Kristensen, S. S., Hvidegaard, S. M., Citterio, M., Karlsson, N., Fettweis, X., Ahlstrøm, A. P., Andersen, S. B., van As, D., and Fausto, R. S.: Greenland ice sheet mass balance assessed by PROMICE (1995 – 2015), *Geological Survey
675 of Denmark and Greenland Bulletin*, 43, e2019430 201, <https://doi.org/10.34194/GEUSB-201943-02-01>, 2019.
- Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, fourth edn., 2010.
- Dask Development Team: Dask: Library for dynamic task scheduling, <https://dask.org>, 2016.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, I., Biblot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Greer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Mong-Sanz, B. M., Morcette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., , and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Eerola, K.: About the performance of HIRLAM version 7.0, *Hirlam Newsletter*, 51, 93–102, 2006.
- 685 Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M.-J., van Angelen, J. H., and van den Broeke, M. R.: An improved mass budget for the Greenland Ice Sheet, *Geophysical Research Letters*, 41, 866–872, <https://doi.org/10.1002/2013GL059010>, 2014.
- Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Box, J. E., and Steffen, K.: Climate of the Greenland ice sheet using a high-resolution climate model – Part 1: Evaluation, *The Cryosphere*, 4, 511–527, <https://doi.org/10.5194/tc-4-511-2010>, 2010.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., Van As, D., Machguth, H., and Gallée, H.: Reconstructions of
690 the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1033, <https://doi.org/10.5194/tc-11-1015-2017>, 2017.

- Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhout, L., Wilton, D., Wouters, B., Ziemen, F., and Zolles, T.: GrSMBMIP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet, *The Cryosphere*, 14, 3935–3958, <https://doi.org/10.5194/tc-14-3935-2020>, 2020.
- Fox Maule, C., Purucker, M. E., and Olsen, N.: Inferring magnetic crustal thickness and geothermal heat flux from crustal magnetic field models, 2009.
- 700 Gallée, H. and Schayes, G.: Development of a Three-Dimensional Meso- γ Primitive Equation Model: Katabatic Winds Simulation in the Area of Terra Nova Bay, Antarctica, *Monthly Weather Review*, 122, 671–685, [https://doi.org/10.1175/1520-0493\(1994\)122<0671:doatdm>2.0.co;2](https://doi.org/10.1175/1520-0493(1994)122<0671:doatdm>2.0.co;2), 1994.
- Gallée, H., Guyomarc'h, G., and Brun, E.: Impact Of Snow Drift On The Antarctic Ice Sheet Surface Mass Balance: Possible Sensitivity To Snow-Surface Properties, *Boundary-Layer Meteorology*, 99, 1–19, <https://doi.org/10.1023/a:1018776422809>, 2001.
- 705 Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., Van Den Broeke, M. R., and Paul, F.: A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*, 340, 852–857, <https://doi.org/10.1126/science.1234532>, 2013.
- GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, Open Source Geospatial Foundation, <https://gdal.org>, 2020.
- 710 Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.
- Greene, C. A., Gardner, A. S., and Andrews, L. C.: Detecting seasonal ice dynamics in satellite images, *The Cryosphere*, 14, 4365–4378, <https://doi.org/10.5194/tc-14-4365-2020>, 2020.
- 715 Groh, A., Horwath, M., Horvath, A., Meister, R., Sørensen, L. S., Barletta, V. R., Forsberg, R., Wouters, B., Ditmar, P., Ran, J., Klees, R., Su, X., Shang, K., Guo, J., Shum, C. K., Schrama, E., and Shepherd, A.: Evaluating GRACE Mass Change Time Series for the Antarctic and Greenland Ice Sheet – Methods and Results, *Geosciences*, 9, 415, <https://doi.org/10.3390/geosciences9100415>, 2019.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G. D., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 720 Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8, 1509–1518, <https://doi.org/10.5194/tc-8-1509-2014>, 2014.
- Hoyer, S. and Hamman, J. J.: xarray: N-D labeled Arrays and Datasets in Python, *Journal of Open Research Software*, 5, <https://doi.org/10.5334/jors.148>, 2017.
- Hunter, J. D.: Matplotlib: A 2D graphics environment, *Computing In Science Engineering*, 9, 90–95, 2007.

- 730 Jordahl, K., den Bossche, J. V., Wasserman, J., McBride, J., Fleischmann, M., Gerard, J., Tratner, J., Perry, M., Farmer, C., Hjelle, G. A., Gillies, S., Cochran, M., Bartos, M., Culbertson, L., Eubank, N., Bilogur, A., and maxalbert: *geopandas/geopandas: v0.7.0*, <https://doi.org/10.5281/zenodo.3669853>, 2020.
- Joughin, I., Smith, B. E., and Howat, I.: Greenland Ice Mapping Project: ice flow velocity variation at sub-monthly to decadal timescales, *The Cryosphere*, 12, 2211–2227, <https://doi.org/10.5194/tc-12-2211-2018>, 2018.
- 735 Karlsson, N. B.: Greenland Ice Sheet Basal Melt, <https://doi.org/10.22008/FK2/PLNUEO>, 2021.
- Karlsson, N. B., Solgaard, A. M., Mankoff, K. D., Gillet-Chaulet, F., MacGregor, J. A., Box, J. E., Citterio, M., Colgan, W. T., Larsen, S. H., Kjeldsen, K. K., Korsgaard, N. J., Benn, D. I., Hewitt, I., and Fausto, R. S.: A first constraint on basal melt-water production of the Greenland ice sheet, *Nature Communications*, 12, <https://doi.org/10.1038/s41467-021-23739-z>, 2021.
- Khan, S. A.: Greenland Ice Sheet Surface Elevation Change, https://doi.org/10.22008/promice/data/DTU/surface_elevation_change/v1.0.0,
740 GEUS Data Center, 2017.
- Khan, S. A., Aschwanden, A., Bjørk, A. A., Wahr, J., Kjeldsen, K. K., and Kjær, K. H.: Greenland ice sheet mass balance: a review, *Reports on Progress in Physics*, 78, 046 801, <https://doi.org/10.1088/0034-4885/78/4/046801>, 2015.
- Khan, S. A., Sasgen, I., Bevis, M., van Dam, T., Bamber, J. L., Wahr, J., Willis, M., Kjær, K. H., Wouters, B., Helm, V., Csatho, B., Fleming, K., Bjørk, A. A., Aschwanden, A., Knudsen, P., and Munneke, P. K.: Geodetic measurements reveal similarities between
745 post-Last Glacial Maximum and present-day mass loss from the Greenland ice sheet, *Science Advances*, 2, e1600 931–e1600 931, <https://doi.org/10.1126/sciadv.1600931>, 2016.
- Khan, S. A., Bjørk, A. A., Bamber, J. L., Morlighem, M., Bevis, M., Kjær, K. H., Mouginot, J., Løkkegaard, A., Holland, D. M., Aschwanden, A., Zhang, B., Helm, V., Korsgaard, N. J., Colgan, W., Larsen, N. K., Lui, L., Hansen, K., Barletta, V., Dahl-Jensen, T. S., Søndergaard, A. S., Csatho, B. M., Sasgen, I., Box, J., and Schenk, T.: Centennial response of Greenland’s three largest outlet glaciers,
750 *Nature Communications*, 11, <https://doi.org/10.1038/s41467-020-19580-5>, 2020.
- King, M. D., Howat, I. M., Jeong, S., Noh, M. J., Wouters, B., Noël, B., and van den Broeke, M. R.: Seasonal to decadal variability in ice discharge from the Greenland Ice Sheet, *The Cryosphere*, 12, 3813–3825, <https://doi.org/10.5194/tc-12-3813-2018>, 2018.
- King, M. D., Howat, I. M., Candela, S. G., Noh, M. J., Jeong, S., Noël, B. P. Y., van den Broeke, M. R., Wouters, B., and Negrete, A.: Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat, *Communications Earth Environment*, 1,
755 <https://doi.org/10.1038/s43247-020-0001-2>, 2020.
- Kjeldsen, K. K., Korsgaard, N. J., Bjørk, A. A., Khan, S. A., Box, J. E., Funder, S., Larsen, N. K., Bamber, J. L., Colgan, W., Broeke, M. v. d., Siggaard-Andersen, M.-L., Nuth, C., Schomacker, A., Andresen, C. S., Willerslev, E., and Kjær, K. H.: Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900, *Nature*, 528, 396–400, <https://doi.org/10.1038/nature16183>, 2015.
- Kjeldsen, K. K., Khan, S. A., Colgan, W. T., MacGregor, J. A., and Fausto, R. S.: Time-Varying Ice Sheet Mask: Implications on Ice-Sheet
760 Mass Balance and Crustal Uplift, *Journal of Geophysical Research: Earth Surface*, 125, <https://doi.org/10.1029/2020jf005775>, 2020.
- Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B., Bussonnier, M., Frederic, J., Kelley, K., Hamrick, J., Grout, J., Corlay, S., Ivanov, P., Avila, D., Abdalla, S., and Willing, C.: Jupyter Notebooks – a publishing format for reproducible computational workflows, <https://doi.org/10.3233/978-1-61499-649-1-87>, 2016.
- Langen, P. L., Fausto, R. S., Vandecrux, B., Mottram, R. H., and Box, J. E.: Liquid Water Flow and Retention on the Greenland Ice Sheet in the Regional Climate Model HIRHAM5: Local and Large-Scale Impacts, *Frontiers in Earth Science*, 4,
765 <https://doi.org/10.3389/feart.2016.00110>, 2017.

- Laprise, R.: The Euler Equations of Motion with Hydrostatic Pressure as an Independent Variable, *Monthly Weather Review*, 120, 197–207, [https://doi.org/10.1175/1520-0493\(1992\)120<0197:teecomw>2.0.co;2](https://doi.org/10.1175/1520-0493(1992)120<0197:teecomw>2.0.co;2), 1992.
- Lefebvre, F.: Modeling of snow and ice melt at ETH Camp (West Greenland): A study of surface albedo, *Journal of Geophysical Research*, 770 108, <https://doi.org/10.1029/2001jd001160>, 2003.
- Lenaerts, J. T. M., van den Broeke, M. R., van Angelen, J. H., van Meijgaard, E., and Déry, S. J.: Drifting snow climate of the Greenland ice sheet: a study with a regional climate model, *The Cryosphere*, 6, 891–899, <https://doi.org/10.5194/tc-6-891-2012>, 2012.
- Lewis, G., Osterberg, E., Hawley, R., Whitmore, B., Marshall, H. P., and Box, J.: Regional Greenland accumulation variability from Operation IceBridge airborne accumulation radar, *The Cryosphere*, 11, 773–788, <https://doi.org/10.5194/tc-11-773-2017>, 2017.
- 775 Lewis, G., Osterberg, E., Hawley, R., Marshall, H. P., Meehan, T., Graeter, K., McCarthy, F., Overly, T., Thundercloud, Z., and Ferris, D.: Recent precipitation decrease across the western Greenland ice sheet percolation zone, *The Cryosphere*, 13, 2797–2815, <https://doi.org/10.5194/tc-13-2797-2019>, 2019.
- Liestøl, O.: *Glacier Dammed Lakes in Norway*, vol. 81, Fabritius Sønners Forlag, Oslo, 1956.
- 780 Ligtenberg, S. R. M., Kuipers Munneke, P., Noël, B. P. Y., and van den Broeke, M. R.: Brief communication: Improved simulation of the present-day Greenland firn layer (1960–2016), *The Cryosphere*, 12, 1643–1649, <https://doi.org/10.5194/tc-12-1643-2018>, 2018.
- Lucas-Picher, P., Wulff-Nielsen, M., Christensen, J. H., Aðalgeirsdóttir, G., Mottram, R., and Simonsen, S. B.: Very high resolution regional climate model simulations over Greenland: Identifying added value, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011jd016267>, 2012.
- 785 MacGregor, J. A., Fahnestock, M. A., Catania, G. A., Aschwanden, A., Clow, G. D., Colgan, W. T., Gogineni, P. S., Morlighem, M., Nowicki, S. M. J., Paden, J. D., Price, S. F., and Seroussi, H.: A synthesis of the basal thermal state of the Greenland Ice Sheet, *Journal of Geophysical Research: Earth Surface*, 121, 1328–1350, <https://doi.org/10.1002/2015JF003803>, 2016.
- Maier, N., Humphrey, N., Harper, J., and Meierbachtol, T.: Sliding dominates slow-flowing margin regions, *Greenland Ice Sheet, Science Advances*, 5, eaaw5406, <https://doi.org/10.1126/sciadv.aaw5406>, 2019.
- 790 Maier, N., Gimbert, F., Gillet-Chaulet, F., and Gilbert, A.: Basal traction mainly dictated by hard-bed physics over grounded regions of Greenland, *The Cryosphere*, 15, 1435–1451, <https://doi.org/10.5194/tc-15-1435-2021>, 2021.
- Mankoff, K.: Greenland Ice Sheet solid ice discharge from 1986 through last month: Gates, https://doi.org/10.22008/promice/data/ice_discharge/gates/v02, 2020.
- Mankoff, K. and Solgaard, A.: Ice Discharge, https://dataverse01.geus.dk/dataverse/ice_discharge, 2020.
- 795 Mankoff, K., Fettweis, X., Solgaard, A., Langen, P., Stendel, M., Noël, B., van den Broeke, M. R., Karlsson, N., Box, J. E., and Kjeldsen, K.: Greenland ice sheet mass balance from 1840 through next week, <https://doi.org/10.22008/FK2/YG3IWC>, 2021.
- Mankoff, K. D.: Total Mass Balance, GitHub https://github.com/GEUS-Glaciology-and-Climatology/ice_discharge/issues, last access: August 13, 2021, 2021.
- Mankoff, K. D. and Tulaczyk, S. M.: The past, present, and future viscous heat dissipation available for Greenland subglacial conduit formation, *The Cryosphere*, 11, 303–317, <https://doi.org/10.5194/tc-11-303-2017>, 2017.
- 800 Mankoff, K. D., Noël, B., Fettweis, X., Ahlstrøm, A. P., Colgan, W., Kondo, K., Langley, K., Sugiyama, S., van As, D., and Fausto, R. S.: Greenland liquid water discharge from 1958 through 2019, *Earth System Science Data*, 12, 2811–2841, <https://doi.org/10.5194/essd-12-2811-2020>, 2020a.

- Mankoff, K. D., Solgaard, A., Colgan, W., Ahlstrøm, A. P., Khan, S. A., and Fausto, R. S.: Greenland Ice Sheet solid ice discharge from 1986 through March 2020, *Earth System Science Data*, 12, 1367–1383, <https://doi.org/10.5194/essd-12-1367-2020>, 2020b.
- Mankoff, K. D., van As, D., Lines, A., Bording, T., Elliott, J., Kraghede, R., Cantalloube, H., Oriot, H., Dubois-Fernandez, P., Ruault du Plessis, O., Vest Christiansen, A., Auken, E., Hansen, K., Colgan, W., and Karlsson, N. B.: Search and recovery of aircraft parts in ice-sheet crevasse fields using airborne and in situ geophysical sensors, *Journal of Glaciology*, 66, 496–508, <https://doi.org/10.1017/jog.2020.26>, 2020c.
- Martos, Y. M., Jordan, T. A., Catalán, M., Jordan, T. M., Bamber, J. L., and Vaughan, D. G.: Geothermal Heat Flux Reveals the Iceland Hotspot Track Underneath Greenland, *Geophysical Research Letters*, <https://doi.org/10.1029/2018gl078289>, 2018.
- McKinney, W.: Data Structures for Statistical Computing in Python, in: *Proceedings of the 9th Python in Science Conference*, edited by van der Walt, S. and Millman, J., pp. 51 – 56, <https://doi.org/10.25080/Majora-92bf1922-00a>, 2010.
- Millan, R., Rignot, E., Mouginot, J., Wood, M., Bjørk, A. A., and Morlighem, M.: Vulnerability of Southeast Greenland glaciers to warm Atlantic Water from Operation IceBridge and Ocean Melting Greenland data., *Geophysical Research Letters*, 45, 2688–2696, <https://doi.org/10.1002/2017gl076561>, 2018.
- Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I. M., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., Cofaigh, C. Ó., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo sounding combined with mass conservation, *Geophysical Research Letters*, 44, <https://doi.org/10.1002/2017gl074954>, 2017.
- Morlighem, M., Williams, C., Rignot, E., An, L., Arndt, J. E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B., O’Cofaigh, C., Palmer, S. J., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K.: IceBridge BedMachine Greenland, Version 4, <https://doi.org/10.5067/VLJ5YXKCNXGO>, used all subsets; Accessed 2021-06-30, 2021.
- Mouginot, J. and Rignot, E.: Glacier catchments/basins for the Greenland Ice Sheet, <https://doi.org/10.7280/d1wt11>, 2019.
- Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, p. 201904242, <https://doi.org/10.1073/pnas.1904242116>, 2019.
- Neteler, M., Bowman, M. H., Landa, M., and Metz, M.: GRASS GIS: a multi-purpose Open Source GIS, *Environmental Modelling Software*, 31, 124–130, <https://doi.org/10.1016/j.envsoft.2011.11.014>, 2012.
- Nilsson, J., Vallelonga, P., Simonsen, S. B., Sørensen, L. S., Forsberg, R., Dahl-Jensen, D., Hirabayashi, M., Goto-Azuma, K., Hvidberg, C. S., Kjær, H. A., et al.: Greenland 2012 melt event effects on CryoSat-2 radar altimetry, *Geophysical Research Letters*, 42, 3919–3926, 2015.
- Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., and van den Broeke, M. R.: A daily, 1 km resolution data set of downscaled Greenland ice sheet surface mass balance (1958–2015), *The Cryosphere*, 10, 2361–2377, <https://doi.org/10.5194/tc-10-2361-2016>, 2016.
- Noël, B., van de Berg, W. J., van Wessem, J. M., van Meijgaard, E., van As, D., Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., van Ulft, L. H., van de Wal, R. S. W., and van den Broeke, M. R.: Modelling the climate and surface mass balance of

- polar ice sheets using RACMO2 – Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.
- Noël, B., van de Berg, W. J., Lhermitte, S., and van den Broeke, M. R.: Rapid ablation zone expansion amplifies north Greenland mass loss, *Science Advances*, 5, eaaw0123, <https://doi.org/10.1126/sciadv.aaw0123>, 2019.
- 845 Oliphant, T. E.: *A guide to NumPy*, vol. 1, Trelgol Publishing USA, 2006.
- Pfeffer, W. T., Meier, M. F., and Illangasekare, T. H.: Retention of Greenland runoff by refreezing: Implications for projected future sea level change, *Journal of Geophysical Research*, 96, 22 117, <https://doi.org/10.1029/91jc02502>, 1991.
- Rastner, P., Bolch, T., Mölg, N., Machguth, H., Le Bris, R., and Paul, F.: The first complete inventory of the local glaciers and ice caps on
850 Greenland, *The Cryosphere*, 6, 1483–1495, <https://doi.org/10.5194/tc-6-1483-2012>, 2012.
- Rezvanbehbahani, S., Stearns, L. A., van der Veen, C. J., Oswald, G. K. A., and Greve, R.: Constraining the geothermal heat flux in Greenland at regions of radar-detected basal water, *Journal of Glaciology*, 65, 1023–1034, <https://doi.org/10.1017/jog.2019.79>, 2019.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessel, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>,
855 2019.
- Rignot, E. J., Box, J. E., Burgess, E. W., and Hanna, E.: Mass balance of the Greenland ice sheet from 1958 to 2007, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL035417>, 2008.
- Rocklin, M.: Dask: Parallel Computation with Blocked algorithms and Task Scheduling, in: *Proceedings of the 14th Python in Science Conference*, edited by Huff, K. and Bergstra, J., pp. 130 – 136, 2015.
- 860 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornbluh, L., Manzini, E., et al.: The atmospheric general circulation model ECHAM 5. Part I: Model description, 2003.
- Rogozhina, I., Hagedoorn, J. M., Martinec, Z., Fleming, K., Soucek, O., Greve, R., and Thomas, M.: Effects of uncertainties in the geothermal heat flux distribution on the Greenland Ice Sheet: An assessment of existing heat flow models, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002098>, 2012.
- 865 Ryser, C., Lüthi, M. P., Andrews, L. C., Hoffman, M. J., Catania, G. A., Hawley, R. L., Neumann, T. A., and Kristensen, S. S.: Sustained high basal motion of the Greenland ice sheet revealed by borehole deformation, *Journal of Glaciology*, 60, 647 – 660, <https://doi.org/10.3189/2014JoG13J196>, 2014.
- Sasgen, I., Van Den Broeke, M., Bamber, J. L., Rignot, E., Sørensen, L. S., Wouters, B., Martinec, Z., Velicogna, I., and Simonsen, S. B.: Timing and origin of recent regional ice-mass loss in Greenland, *Earth and Planetary Science Letters*, 333-334, 293–303,
870 <https://doi.org/10.1016/j.epsl.2012.03.033>, 2012.
- Sasgen, I., Wouters, B., Gardner, A. S., King, M. D., Tedesco, M., Landerer, F. W., Dahle, C., Save, H., and Fettweis, X.: Return to rapid ice loss in Greenland and record loss in 2019 detected by the GRACE-FO satellites, *Communications Earth Environment*, 1, <https://doi.org/10.1038/s43247-020-0010-1>, 2020.
- Schulte, E., Davison, D., Dye, T., and Dominik, C.: A multi-language computing environment for literate programming and reproducible
875 research, *Journal of Statistical Software*, 46, 1–24, 2012.
- Schulzweida, U.: CDO User Guide, <https://doi.org/10.5281/zenodo.3539275>, 2019.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring Surface Heat Flux Distributions Guided by a Global Seismic Model: Particular Application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>, 2004.
- Shepherd, A.: RC2: Comment on *essd-2021-131*, <https://doi.org/10.5194/essd-2021-131-rc2>, 2021.

- 880 Shreve, R. L.: Movement of water in glaciers, *Journal of Glaciology*, 11, 205–214, 1972.
- Simmons, A. J. and Burridge, D. M.: An Energy and Angular-Momentum Conserving Vertical Finite-Difference Scheme and Hybrid Vertical Coordinates, *Monthly Weather Review*, 109, 758–766, [https://doi.org/10.1175/1520-0493\(1981\)109<0758:aeaamc>2.0.co;2](https://doi.org/10.1175/1520-0493(1981)109<0758:aeaamc>2.0.co;2), 1981.
- Simonsen, S. B., Barletta, V. R., Colgan, W., and Sørensen, L. S.: Greenland Ice Sheet mass balance (1992–2020) from calibrated radar altimetry, *Geophysical Research Letters*, <https://doi.org/10.1029/2020gl091216>, see <http://doi.org/10.11583/DTU.13353062>. See citet:simonsen2021a, 2021a.
- Simonsen, S. B., Barletta, V. R., Colgan, W., and Sørensen, L. S.: Greenland Ice Sheet mass balance (1992–2020) from calibrated radar altimetry, <https://doi.org/10.11583/DTU.13353062.v1>, https://data.dtu.dk/articles/dataset/Greenland_Ice_Sheet_mass_balance_1992-2020_from_calibrated_radar_altimetry/13353062, see <http://doi.org/10.1029/2020GL091216>. See citet:simonsen2021, 2021b.
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., and et al.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, p. eaaz5845, <https://doi.org/10.1126/science.aaz5845>,
885 2020.
- Solgaard, A. and Kusk, A.: Greenland Ice Velocity from Sentinel-1 Edition 2, <https://doi.org/10.22008/promice/data/sentinel1icevelocity/greenlandicesheet>,
2021.
- Solgaard, A., Kusk, A., Boncori, J. P. M., Dall, J., Mankoff, K. D., Ahlstrøm, A. P., Andersen, S. B., Citterio, M., Karlsson, N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., and Fausto, R. S.: Greenland ice velocity maps from the PROMICE project, *Earth System Science*
890 Data, 13, 3491–3512, <https://doi.org/10.5194/essd-13-3491-2021>, 2021.
- Stallman, R. M.: EMACS the Extensible, Customizable Self-Documenting Display Editor, in: Proceedings of the ACM SIGPLAN SIGOA Symposium on Text Manipulation, p. 147–156, Association for Computing Machinery, New York, NY, USA, <https://doi.org/10.1145/800209.806466>, 1981.
- Sutterley, T. C., Velicogna, I., Rignot, E. J., Mouginot, J., Flament, T., van den Broeke, M. R., van Wessem, J. M., and Reijmer, C. H.: Mass
895 loss of the Amundsen Sea Embayment of West Antarctica from four independent techniques, *Geophysical Research Letters*, 41, 8421–8428, <https://doi.org/10.1002/2014GL061940>, 2014.
- Sørensen, L. S., Simonsen, S. B., Nielsen, K., Lucas-Picher, P., Spada, G., Adalgeirsdottir, G., Forsberg, R., and Hvidberg, C. S.: Mass balance of the Greenland ice sheet (2003–2008) from ICESat data – the impact of interpolation, sampling and firn density, *The Cryosphere*, 5, 173–186, <https://doi.org/10.5194/tc-5-173-2011>, 2011.
- 900Tange, O.: GNU Parallel - The Command-Line Power Tool, ;login: The USENIX Magazine, 36, 42–47, <https://doi.org/10.5281/zenodo.16303>,
<http://www.gnu.org/s/parallel>, 2011.
- Tedesco, M. and Fettweis, X.: Unprecedented atmospheric conditions (1948–2019) drive the 2019 exceptional melting season over the Greenland ice sheet, *The Cryosphere*, 14, 1209–1223, <https://doi.org/10.5194/tc-14-1209-2020>, 2020.
- The IMBIE Team: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, [https://doi.org/10.1038/s41586-019-1855-2](https://doi.org/10.1038/s41586-019-
905 1855-2), 2019.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Da Costa Bechtold, V., Fiorino, M., Gibson, J., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Anderson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, I., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P.,
910 Sterl, A., Trenbreth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., , and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, <https://doi.org/10.1256/qj.04.176>, 2005.

- van Angelen, J. H., Lenaerts, J. T. M., Lhermitte, S., Fettweis, X., Kuipers Munneke, P., van den Broeke, M. R., van Meijgaard, E., and Smeets, C. J. P. P.: Sensitivity of Greenland Ice Sheet surface mass balance to surface albedo parameterization: a study with a regional climate model, *The Cryosphere*, 6, 1175–1186, <https://doi.org/10.5194/tc-6-1175-2012>, 2012.
- 915 van de Berg, W. J. and Medley, B.: Brief Communication: Upper-air relaxation in RACMO2 significantly improves modelled interannual surface mass balance variability in Antarctica, *The Cryosphere*, 10, 459–463, <https://doi.org/10.5194/tc-10-459-2016>, 2016.
- Van Rossum, G. and Drake Jr, F. L.: Python reference manual, Centrum voor Wiskunde en Informatica Amsterdam, 1995.
- Velicogna, I., Mohajerani, Y., A. G., Landerer, F., Mouginot, J., Noel, B., Rignot, E., Sutterley, T., Broeke, M., Wessem, M., and et al.: Continuity of Ice Sheet Mass Loss in Greenland and Antarctica From the GRACE and GRACE Follow-On Missions, *Geophysical Research Letters*, 47, 920 <https://doi.org/10.1029/2020gl087291>, 2020.
- Vinther, B. M., Andersen, K. K., Jones, P. D., Briffa, K. R., and Cappelen, J.: Extending Greenland temperature records into the late eighteenth century, *Journal of Geophysical Research*, 111, <https://doi.org/10.1029/2005jd006810>, 2006.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- 925 Wang, Z. Q. and Randriamampianina, R.: The Impact of Assimilating Satellite Radiance Observations in the Copernicus European Regional Reanalysis (CERRA), *Remote Sensing*, 13, 426, <https://doi.org/10.3390/rs13030426>, 2021.
- Yang, X., Palmason, B., Sattler, K., Thorsteinsson, S., Amstrup, B., Dahlbom, M., Hansen Sass, B., Pagh Nielsen, K., and Petersen, G. N.: IGB, the Upgrade to the Joint Operational HARMONIE by DMI and IMO in 2018, *ALADIN-HIRLAM Newsletter*, pp. 93–96, 2018.
- Zemp, M., Huss, M., Eckert, N., Thibert, E., Paul, F., Nussbaumer, S. U., and Gärtner-Roer, I.: Brief communication: Ad hoc estimation of 930 glacier contributions to sea-level rise from the latest glaciological observations, *The Cryosphere*, 14, 1043–1050, <https://doi.org/10.5194/tc-14-1043-2020>, 2020.
- Zender, C. S.: Analysis of self-describing gridded geoscience data with netCDF Operators (NCO), *Environmental Modelling & Software*, 23, 1338–1342, <https://doi.org/10.1016/j.envsoft.2008.03.004>, 2008.
- Zuo, Z. and Oerlemans, J.: Modelling albedo and specific balance of the Greenland ice sheet: calculations for the Søndre Strømfjord transect, 935 *Journal of Glaciology*, 42, 305–317, <https://doi.org/10.3189/s0022143000004160>, 1996.
- Zwally, H. J. and Giovinetto, M. B.: Overview and Assessment of Antarctic Ice-Sheet Mass Balance Estimates: 1992–2009, *Surveys in Geophysics*, 32, 1–26, <https://doi.org/10.1007/s10712-011-9123-5>, 2011.
- Zwally, H. J., Giovinetto, M. B., Beckley, M. A., and Saba, J. L.: Antarctic and Greenland Drainage Systems, http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php, GSFC Cryospheric Sciences Laboratory, 2012.