

The Southern Ocean RADiolarian (SO-RAD) dataset: a new compilation of modern radiolarian census data

Kelly-Anne Lawler¹, Giuseppe Cortese², Matthieu Civel-Mazens^{3,4}, Helen Bostock^{5,6}, Xavier Crosta⁷,
5 Amy Leventer⁸, Vikki Lowe⁵, John Rogers¹, Leanne K. Armand¹

¹Research School of Earth Sciences, The Australian National University, Canberra, ACT, 2601, Australia

²GNS Science, 1 Fairway Drive, Avalon, Lower Hutt 5010, New Zealand

10 ³Graduate School of Integrated Arts and Sciences, Kochi University, Japan

⁴Geological Survey of Japan, AIST, Tsukuba, Japan

⁵School of Earth and Environmental Sciences, University of Queensland, Australia

⁶National Institute of Water and Atmospheric Research (NIWA), Wellington, New Zealand

⁷Université de Bordeaux, CNRS, EPHE, UMR 5805 EPOC, 33615 Pessac, France

15 ⁸Geology Department, Colgate University, Hamilton NY, USA

Correspondence to: Kelly-Anne Lawler (kelly-anne.lawler@anu.edu.au)

Abstract

20 Radiolarians (holoplanktonic Protozoa) preserved in marine sediments are commonly used as palaeoclimate proxies for reconstructing past Southern Ocean environments. Generating reconstructions of past climate based on microfossil abundances, such as radiolarians, requires a spatially and environmentally comprehensive reference dataset of modern census counts. The Southern Ocean RADiolarian (SO-RAD) dataset includes census counts for 238 radiolarian taxa from 228 surface sediment samples located in the Atlantic, Indian and southwest Pacific sectors of the Southern Ocean. This compilation is the
25 largest radiolarian census dataset derived from surface sediment samples in the Southern Ocean. The SO-RAD dataset may be used as a reference dataset for palaeoceanographic reconstructions, or for studying modern radiolarian biogeography and species diversity. As well as describing the data collection and collation, we include recommendations and guidelines for cleaning and subsetting the data for users unfamiliar with the procedures typically used by the radiolarian community. The SO-RAD dataset is available to download from doi.pangaea.de/10.1594/PANGAEA.929903 (Lawler et al., 2021).

30 1 Introduction

The Southern Ocean is an important part of the global climate system as a major hub of oceanic circulation and nutrient redistribution. Future changes to its physical, chemical, and biological properties will therefore have global climate implications. Palaeoceanographic reconstructions of the Southern Ocean reveal its response to past climate and how it might respond to future climate change. Marine microfossils are commonly used as climate proxies to obtain such reconstructions
35 (Yasuhara et al., 2020). Plankton distribution and abundance are related to the oceanic conditions prevalent in the water masses where these organisms live. After death, their skeletons settle to the ocean floor, becoming part of the sedimentary record. In the Southern Ocean, two main siliceous microfossil groups, diatoms and radiolarians, have been used to determine past environmental conditions such as ocean temperature (Cortese and Abelmann, 2002; Esper and Gersonde, 2014b; Panitz et al., 2015) and sea-ice coverage (Bianchi and Gersonde, 2004; Crosta et al., 2004; Gersonde et al., 2005; Esper and Gersonde,
40 2014a; Ferry et al., 2015).

Radiolarians are unicellular, eukaryotic marine microzooplankton, generally ranging in size from 30 – 300 μm , that inhabit all depths throughout the world's oceans (Lazarus, 2005). Belonging to the supergroup Rhizaria, Radiolaria consist of 600-800 extant species in five orders: Acantharia, Taxopodia, and the polycystine orders Collodaria, Nassellaria and Spumellaria (Suzuki and Not, 2015; Boltovskoy et al., 2017). Radiolarians belonging to the polycystine orders are the most likely to be
45 found in marine sediments and are therefore the focus of most micropaleontological studies (Lazarus, 2005; Suzuki & Not, 2015). Radiolaria feed on diatoms, other small algae, and bacteria, and some also have photosynthetic algal symbionts (Lazarus, 2005; Boltovskoy et al., 2017). Little is known about the radiolarian life cycle as they are notoriously difficult to culture (Lazarus, 2005; Suzuki and Not, 2015). In the Southern Ocean, radiolarian productivity is highest (up to 90% of annual

flux) during the Austral summer (Abelmann and Gersonde, 1991; Abelmann, 1992). Unlike diatoms, radiolarians do not reach
50 maximum biomass in Southern Ocean surface waters, but rather peak between 100-400 m water depth south of 45°S
(Abelmann and Gowing, 1997; Boltovskoy, 2017). Lazarus et al., (2021) provide an extensive and up-to-date review on
radiolarian biology as well as their distribution both in the water column and sediments.

Siliceous microfossils, particularly diatoms, are useful for paleoclimate reconstructions at high latitudes and in deep-sea
regions where carbonate microfossils are generally not well preserved (Crosta et al., 2005; Gersonde et al., 2005). After
55 diatoms, radiolarians are the second largest source of biogenic silica in marine sediments (Anderson, 2019). Fossil radiolarians
have long been used as a palaeo-proxy for Southern Ocean sea-surface temperature (Lozano and Hays, 1976; Cortese and
Abelmann, 2002; Cortese et al., 2007; Rogers and De Deckker, 2011; Panitz et al., 2015) and, more recently, have been used
for reconstructing subsurface ocean temperature (Hernández-Almeida et al., 2020; Civel-Mazens et al., 2021a, 2021b). Such
palaeoceanographic reconstructions rely on a spatially and environmentally comprehensive radiolarian reference dataset.

60 In this paper we present, for the first time, the Southern Ocean RADiolarian (SO-RAD) dataset consisting of census data for
238 radiolarian taxa from 228 surface sediment samples. We illustrate the spatial and environmental coverage of the dataset,
briefly describe the distribution and abundance of taxa, and discuss how the dataset might be cleaned and/or split into subsets
prior to analysis. The SO-RAD dataset can be used as a reference dataset for palaeoceanographic reconstructions and to explore
new parameters for estimation, as standalone data to assist studies in the biogeography and diversity of modern Southern Ocean
65 radiolarian species, or for the development of new statistical methods in microfossil analysis.

2 Methods

2.1 Southern Ocean Radiolarian Dataset

The Southern Ocean Radiolarian (SO-RAD) dataset includes census counts for 238 taxa/taxonomic groups from 228 surface
sediment samples located in Atlantic, Indian, and southwest Pacific sectors of the Southern Ocean (Figure 1). This compilation
70 is the largest dataset available for radiolarian-based palaeoenvironmental reconstructions constrained to the Southern Ocean.
Sector-specific datasets have been previously published as follows: Atlantic sector (Abelmann et al., 1999; Cortese and
Abelmann, 2002), southwest Pacific sector (Cortese and Prebble, 2015), and Indian sector (Rogers and De Deckker, 2007).
These existing data, after a recount of the Indian sector sites to ensure taxonomic consistency, have been compiled as one
dataset, which additionally includes new census data from 26 previously unpublished sites in the Ross Sea, and the Indian
75 sector of the Southern Ocean. Sediment coring in the Southern Ocean, particularly in areas close to the ice edge where sea-ice
coverage varies from season to season, is expensive and difficult to plan. The SO-RAD dataset is, thus, the result of cooperative
international partnerships based on decades of oceanographic voyages undertaken by scientists from many institutions and





Figure 1: Surface sediment sites included in the SO-RAD database shown with a) Southern Ocean frontal boundaries (Orsi et al., 1995), and ocean basin boundaries (WOCE Data Products Committee, 2002), and b) the 1981 - 2010 median sea-ice extent for February (broken line) and September (solid line) (Fetterer et al., 2017).

nations. In a compilation such as the SO-RAD dataset, sample locations are not chosen according to uniform spatial coverage
85 criteria, but rather dictated by the availability of, and access to, radiolarian-bearing surface sediment samples.

2.1.1 Census sources and handling protocols

In bringing together census data for this global dataset from various sources, the following data limitations and handling
90 procedures are detailed.

Census data from the Atlantic sector include radiolarian counts from 64 sites. Data from 44 sites were published by Abelman
et al. (1999) and 20 sites were subsequently added and published in Cortese and Abelman (2002). Sites from Abelman et al.
(1999) and Cortese and Abelman (2002) do not include the full counts of all taxa observed at each site, rather they include
95 only a selection of the taxa observed. Taxa were excluded from these sites prior to their original publication based on criteria
outlined in Abelman et al. (1999) such as their relative abundances (i.e., excluding rare taxa, with the threshold set at 2%),
depth preferences, and clarity of identification. These sites have been included in the dataset, despite containing counts for
only a percentage of the taxa observed, because they contain valuable information about species useful for palaeo-
reconstructions and for single-species studies.

Radiolarian census data from 87 sites located in the southwest Pacific sector published by Cortese and Prebble (2015) contained
100 full census counts and required no further treatment.

The Indian Ocean sector includes census data from 41 sites first published by Rogers and De Deckker (2007). The slides from
these sites were recounted to ensure consistency with the taxonomy of the SO-RAD compilation presented here (described in
further detail in section 2.1.3). Also located in the Indian Ocean sector are census data from four sites from Abelman et al.
(1999) and six from Cortese and Abelman (2002).

105 We also include radiolarian census data from 26 previously unpublished sites. These include counts from 14 archive samples
from sites in the Ross Sea, retrieved during multiple expeditions by the National Institute of Water and Atmospheric Research
(NIWA) and its predecessor New Zealand Oceanographic Institute; five sites near the Sabrina Coast, East Antarctica (three
retrieved during the 2017 RV *Investigator* voyage IN2017-V01 (Armand et al., 2018), two during the 2014 RVIB *Nathaniel*
B. Palmer voyage NBP-1402 (Leventer and Science Party, 2014); one previously unpublished site in the southwest Pacific
110 sector; and a total of six samples retrieved during the MD185 INDIEN-SUD1 (Mazaud and Michel, 2011), MD189 INDIEN-
SUD2 (Mazaud and Michel, 2012) and MD218 CROTALE (Crosta, 2019) voyages.

2.1.2 Slide preparation and microscopy

In general, radiolarian slides were prepared using a similar method as described in Cortese and Prebble (2015). Sediment
115 samples were dried, weighed, placed in beakers with 100 mL of distilled water, and heated to 60-70 °C on a hotplate. Organic
matter was removed by adding a mixture of diluted hydrogen peroxide (ca. 20 % concentration) and sodium
hexametaphosphate (Calgon™) and left to react for around one hour. To remove carbonate, 10 % hydrochloric acid was added
to the beaker. On completion of the reaction, usually within ca. 30 minutes, the content of the beaker was washed through a
sieve. The residue was placed in a beaker or storage bottle with distilled water, stirred into suspension, and a pipette used to
120 place a few drops of it on a coverslip. After the cover slip had dried on a hotplate, it was mounted on a slide by using Canada
balsam or Norland™ Optical Adhesive (NOA61) as a mounting medium. Finally, the mounted slide was either placed on a
hotplate again for ca. 10-20 minutes to allow the Canada balsam to harden, or under ultraviolet light for the NOA61 to cure.
Note that the specific chemical separation steps may differ between authors, as may the mesh size used. A mesh size of 40-45
µm has been used with the exception of Rogers & DeDecker (2007) who used a 63 µm mesh size. For detailed methods on
125 slide preparation for previously published data, refer to the references listed in section 2.1.1.

Radiolarian census counts were determined using transmitted-light microscopy at 100-400x magnification. The number of
individuals counted per site ranges from 44 to 706, with a median of 381.

2.1.3 Taxonomy

Species identification is mainly based on the taxonomic concepts of Petrushevskaya (1967, 1971), Nigrini and Moore (1979)
130 and Boltovskoy (1998). This was supplemented by consulting Lazarus et al. (2015), the online dataset at www.radiolaria.org,
(Radiolaria.org, n.d.) with a final check on currently valid taxonomic concepts according to the World Register of Marine
Species (WoRMS Editorial Board, 2020). Some naming adjustments were made to taxa in the previously published datasets
where more recent, accepted names were available (Supplement 1). If individuals could not be identified at species level, they
were assigned to a higher rank taxonomic group e.g., from genus level identifications to those identified only as belonging to
135 the Nassellaria or Spumellaria orders.

A total of 238 radiolarian taxa have been recognised. Radiolarian taxa identified in SO-RAD are from the super-class
Polycystinea, which is divided into three orders, Nassellaria, Spumellaria and Collodaria. When present, phaeodarian species
were identified, counted, and recorded. The binomial names for each of the taxa included in the SO-RAD dataset are listed in
Supplement 2.

140 2.2 Complementary Data

To spatially categorise the dataset, each site was assigned to one of four zones: the Antarctic Zone (AZ; south of the Antarctic
Polar Front), Polar Frontal Zone (PFZ; between the Polar Front and the Subantarctic Front), Subantarctic Zone (SAZ; between
the Subantarctic Front and the Subtropical Front), or Subtropical Zone (STZ; north of the Subtropical Front), based on frontal

boundaries delineated by Orsi et al., (1995). Sites were also assigned to one of three sectors, either the Atlantic, Indian, or southwest Pacific sector, based on the longitudinal boundaries used by the World Ocean Circulation Experiment (WOCE Data Products Committee, 2002).

To illustrate the environmental coverage of the dataset, Austral summer (January - March) observations of temperature (°C), salinity (psu), and dissolved oxygen (µmol/kg), silicate (µmol/kg), nitrate (µmol/kg) and phosphate (µmol/kg), were extracted from the World Ocean Atlas 2018 (Locarnini et al., 2018; Garcia et al., 2019a, 2019b; Zweng et al., 2019) using Ocean Data View software (Schlitzer, 2020). These data represent a collation of 1°x1° gridded data from 1955 – 2017 and, given that radiolarian abundances are highest below the surface, we elected to use data reported at 100 m depth. Future users of the dataset should select the environmental variables, water depths and data source that best suits their needs. The median sea-ice extent boundaries from 1981 – 2010 representing minimum (February) and maximum (September) sea-ice extents were retrieved from the National Snow and Ice Data Center repository (Fetterer et al., 2017). These monthly sea-ice extent boundaries are based on >15% sea-ice concentration.

2.3 Structure of the SO-RAD dataset

The SO-RAD dataset consists of 228 rows (plus header rows) and 251 columns. Each row contains data for one site from which a surface sediment sample was obtained. Columns 1-13 contain information relating to the site or sample, while columns 14 - 251 are the radiolarian taxa observed (one taxon per column). The SO-RAD dataset is available to download from doi.pangaea.de/10.1594/PANGAEA.929903 (Lawler et al., 2021).

Table 1: Description of variables in the SO-RAD dataset

Variable name	Description
Event label	The sampling site/station at which the data was obtained.
Site	The sampling site/station at which the data was obtained. Site names act as unique identifier for each record (row).
Latitude	All latitudes are negative as all sites are in the southern hemisphere. Units are decimal degrees.
Longitude	Western hemisphere longitudes are negative (e.g., 155°00'W = -155.00) Units are decimal degrees.
Zone	The location of the site according to the boundaries described in section 2.2. AZ = Antarctic Zone PFZ = Polar Frontal Zone SAZ = Subantarctic Zone STZ = Subtropical Zone

Sector	The location of the site according to the boundaries described in section 2.2. ATL = Atlantic sector IND = Indian sector SWP = Southwest Pacific sector
Depth, bathymetric	Bathymetric depth i.e., depth below sea level. Units are metres.
Sample method	The method used to extract the sediment core/sample from the seafloor.
Depth, top/min	The upper depth of the sample. Units are metres.
Depth bottom/max	The lower depth of the sample. Units are metres.
Depth, sediment/rock	The average of the upper and lower depth of the sample. Units are metres.
Radiolarians	The number of individual radiolarians observed in each sample. This data can be used to convert count data to relative abundances.
Reference	Reference for the original source of radiolarian census data.
Radiolarian taxa	Columns 14 - 251 contain census data for each taxonomic grouping in the dataset. Units are the number of individuals observed in a sample.

2.4 Software

All figures and summary statistics in this paper were generated using R statistical software (R core team, 2020) and the packages *mapproj* (McIlroy, 2020), *maps* (Becker, 2018), *orsifronts* (Sumner, 2015), *rgdal* (Bivand et al., 2018) and *tidyverse* (Wickham et al., 2019).

3 Results


3.1 Spatial distribution of data

The SO-RAD dataset is the most comprehensive Southern Ocean radiolarian surface sediment dataset available. However, the spatial coverage is still uneven with, for example, more Antarctic Zone sites in the Atlantic sector than in the Indian and southwest Pacific sectors, and more Subtropical Zone sites in the southwest Pacific sector (Table 2). The latter sector has better overall coverage than the Atlantic and Indian sectors, nevertheless there is a paucity of sites from the Pacific Ocean east of the Ross Sea. Sites represent locations covering a large water depth gradient, from the Antarctic continental shelf (235 metres below sea level) to the abyssal plain (5883 metres below sea level).

175

3.2 Environmental distribution/coverage of data


Sites included in the SO-RAD dataset are representative of a broad range of environmental conditions. Temperature values range from $-1.8\text{ }^{\circ}\text{C}$ to $26.4\text{ }^{\circ}\text{C}$, salinity from 33.8 to 35.7 psu, and dissolved oxygen from 180.3 to 337.9 $\mu\text{mol/kg}$. The SO-RAD dataset also includes sites representing a large range of nutrient concentrations: nitrate (0.1 - 31.4 $\mu\text{mol/kg}$), phosphate (0.1 - 2.2 $\mu\text{mol/kg}$) and silicate (0.6 - 89.3 $\mu\text{mol/kg}$) (Figures 2 and 3). Forty-three sites lie on, or within, the median September sea-ice extent boundary, however no sites lie within the median February sea-ice extent boundary. Some sites that are slightly north of the median September sea-ice extent boundary may be affected by sea ice in certain years.



185 **Figure 2:** Locations of sites (black dots) and the six environmental variables at 100 m water depth used to demonstrate the environmental coverage of the SO-RAD dataset (Data source: Locarnini et al., 2018; Garcia et al., 2019a, 2019b; Zweng et al., 2019).

Table 2: Number of sites located in each zone and sector of the Southern Ocean.

	Atlantic Sector	Indian Sector	Southwest Pacific Sector	Total sites per zone
Subtropical/Tropical Zone	14	13	58	85
Subantarctic Zone	17	26	21	64
Polar Frontal Zone	7	9	5	21
Antarctic Zone	26	14	18	58
Total sites per sector	64	62	102	228



195 **Figure 3:** Distribution of sites across observed environmental variables at 100 m water depth (Locarnini et al., 2018; Garcia et al., 2019a, 2019b; Zweng et al., 2019) identified by sector – Atlantic Ocean sector (ATL; green), Indian Ocean sector (IND; orange), southwest Pacific sector (SWP; purple).

3.3 Radiolarian distribution and abundance

200 The SO-RAD dataset documents 238 identified taxa characterised by subspecies/species level identifications through to individuals identified only as belonging to the Nassellaria or Spumellaria orders.

Radiolarian species (i.e., not including taxa of genus level and above, or taxa that had not been formally described) with the highest average relative abundances were *Antarctissa denticulata* and *Antarctissa strelkovi*. These were the only species with average relative abundances >10%. The *Antarctissa* genus (consisting of *A. denticulata*, *A. strelkovi* and *Antarctissa* spp.) is the most abundant radiolarian genus in the Southern Ocean and was observed at 146 sites. It has a relative abundance of > 40% at several sites in the SO-RAD dataset (n=29), with some sites in the Ross Sea consisting of > 80% *Antarctissa* species.



205

Other well-distributed and abundant species were *Tetrapyle octacantha/stenozona* (average relative abundance = 4.3%, observed at 161 sites), *Cenosphaera cristata* (3.75%, 123 sites), *Cycladophora bicornis* (3.72%, 165 sites), *Lithomelissa setosa* (3.52%, 77 sites), *Lithomelissa* sp. A (3.4%, 95 sites), *Phorticium clevei* (3.24%, 129 sites) and *Lithelius minor* (3.19%, 170 sites) (Figure 4). Only one other species, *Sphaerozoum punctatum* (3.28%), had an average relative abundance > 3%, however it was observed at only 9 sites.



Thirty-eight taxa were found at more than 100 sites. *Spongopyle osculosa* was the most widely distributed species having been observed at 200 sites.

There were 71 taxa (30% of the total number of taxa) in the SO-RAD dataset considered to be 'rare' i.e., they had a maximum relative abundance <1% and/or were observed at only one site. If the maximum relative abundance threshold for a rare categorisation was increased to <2%, then 123 taxa (52%) would be considered rare.



Antarctissa denticulata





Antarctissa strelkovi





Cenosphaera cristata





Cycladophora bicornis





Lithelius minor





Lithomelissa setosa





Lithomelissa Sp. A



Phorticium clevei



Tetrapyle octacantha/stenozona



Sites
Subtropical Front
Subantarctic Front
Polar Front

Figure 4: Distribution maps for the nine most abundant and widely distributed radiolarian taxa. Ocean front boundaries from Orsi et al. (1995).

4 Discussion

While the SO-RAD dataset is predominantly a Southern Ocean dataset, there are several sites located considerably north of the Subtropical Front (the northern boundary of the Southern Ocean). One of the anticipated uses of the SO-RAD dataset is as a reference dataset for radiolarian-based palaeoenvironmental reconstructions from sediment cores retrieved from the Southern Ocean and Subtropical southern hemisphere. One of the key requirements of a reference dataset is that it should cover a wide range of environmental conditions. It has been demonstrated elsewhere that radiolarian assemblages from temperate latitudes, such as offshore New Zealand, show large and distinct changes over glacial and interglacial periods (e.g., Lüer et al., 2009). The broad coverage of the SO-RAD modern reference dataset facilitates identifying future faunal shifts, and imparts robustness to the environmental variable estimates derived from them. In a broader sense, the large spatial and environmental coverage of this dataset can aid in the refining of palaeoenvironmental reconstructions using radiolarian assemblages from periods that were substantially cooler or warmer than the present. The SO-RAD database is a valuable resource for exploring how physical, biological, and chemical characteristics of the Southern Ocean influence modern radiolarian biogeography, ecology, and species diversity. This ecological information and species diversity may provide key insights in the past palaeceanography. The SO-RAD dataset also holds potential for the exploration of sea-ice coverage/concentration as a causal mechanism underlying the abundance patterns for some of the species occurring in the seasonally ice-covered zone. It may be possible to establish surface-dwelling radiolarian species as sea-ice proxies, as routinely done for diatom species, such as *Fragilariopsis curta/cylindrus* and *F. obliquecostata* (Gersonde and Zielinski, 2000; Armand et al., 2005), and in diatom-based transfer functions (Crosta et al., 1998; Esper and Gersonde, 2014a; Ferry et al., 2015). Radiolarian species inhabiting sub-surface and intermediate depths may also be affected by sea-ice related changes deep in the water column, such as salinity and stratification changes due to brine rejection during sea-ice formation, and act as a palaeo-indicator of past sea-ice presence. This should be explored in the future.

4.1 General limitations of microfossil reference datasets

When constructing any reference dataset based on surface sediment samples there are possible sources of error and assumptions that apply. Reproducibility of species abundance data can be affected by differing taxonomic concepts held by the analysts. We have minimized this potential source of error by including counts by radiolarists who regularly collaborate on taxonomic concepts. Furthermore, the vast majority of the samples included here have been counted by only one of us (Cortese). As more samples are added to the SO-RAD dataset, only samples for which quality control of the census data is possible, will be included in the database in an effort to minimise this source of error.

The species-level taxonomy of all microfossil groups has been based on phenotype/morphology, and not on molecular biology. Identification can be difficult when there are only minor differences in morphology between species, or where there are fragments of an individual, or additionally, where a juvenile specimen has been encountered. In such instances individual radiolarians have been assigned to higher rank taxonomic categories i.e., genus or family rather than species or subspecies. When using surface sediment samples as a modern analogue record it is assumed that the fossil assemblages found within those sediments are representative of modern assemblages that live in the water column above. This is problematic for two reasons. The first is that the coring device can affect the quality/age of the surface sediment sample. For example, using a multi-corer allows for easier observation, and intact retrieval of, the surface sediment/bottom water interface (Barnett et al., 1986). With other coring devices it is not possible to observe if the uppermost sediment layer is the true surface sediment sample, or if the top of the coring device was buried beneath the surface. This means the ‘core top’ sample may be from several centimetres below the surface and is not the most recently deposited sediment. Some of these sediment samples have been radiocarbon dated using planktic foraminifera in the same samples (Cortese and Prebble, 2015), but this is not possible with all samples, particularly those that have no carbonate present. Secondly, even when the sediment sample is truly from the surface, the transportation or reworking of sediment, or preferential dissolution of species, mean that the assemblages observed in the sediment may not be truly representative of the assemblages that lived in the water column above.

4.2 Cleaning and subsetting the SO-RAD dataset based on sites

Where possible, we have presented census counts in their complete form to cater for a variety of analyses as well as the development of novel techniques for radiolarian data analysis. We have opted not to present a ‘reconstruction ready’ dataset as we believe the full dataset is useful for more than just palaeoenvironmental reconstructions. However, for most analyses the dataset would not be used in its raw form. Here we offer practical suggestions for preparing and producing a subset from this dataset based on practices utilised by radiolarists in the past.

4.2.1 Quality control of sites

Users of the dataset may wish to extract a subset of sites from the SO-RAD dataset for quality control reasons. For example, placing a lower bound on the total number of individuals counted per site may depend on which species are of interest to the user. In general, lower total counts are sufficient when only the dominant species will be used in the analysis, while higher total counts are needed when a user is interested in species comprising as little as 1% of the population. For example, there is a 30% probability of failing to observe a species that comprises 1% of the population when counting 100 individuals but, by increasing the count to 300 individuals, this probability is reduced to less than 1% (Fatela and Taborda, 2002).

Sites in the Atlantic sector, as well as some in the Indian sector, did not have full census counts available and therefore a percentage of the assemblage is recorded as ‘Other radiolarians’. Rules for the exclusion of species at these sites can be found in Abelmann et al. (1999).

280 The coring apparatus used to retrieve a core can affect the quality of the sample collected, as well as its age. In most studies, the latter is usually assumed to be modern, but this can only be confirmed by radiocarbon dating of each surface sediment sample. Of the sites included in the SO-RAD dataset, the only radiocarbon dated samples are those reported in Cortese & Prebble (2015). Where known, the apparatus used to retrieve the core has been listed in the ‘Sample Method’ column to facilitate the extraction of a data subset based on this variable.

285 **4.2.2 Ensuring spatial/environmental coverage is suitable**

Palaeoenvironmental reconstructions require a reference dataset that provides appropriate spatial and environmental coverage. Reference sites may be included based on regional needs e.g., by sub-setting the data based on latitudinal and/or longitudinal boundaries, based on sector or oceanographic zone, or based on certain values of the environmental variable of interest.

4.3 Cleaning and sub-setting the SO-RAD dataset based on taxa

290 Here we outline some practices to consider when refining the SO-RAD dataset for use as a reference dataset in palaeoceanographic reconstructions. These include selecting only taxa that can be clearly identified and with clear taxonomic descriptions, selecting appropriate taxa for the depth level of the reconstruction, grouping species/subspecies with similar environmental affinities into higher-order groupings, and removing rare species.

4.3.1 Removing questionable taxonomic groupings

295 If the focus of a study is on environmental reconstructions, the signal-to-noise ratio in radiolarian census data can be increased by removing broad, often higher order, counting categories having no clear taxonomic boundaries/definitions, and hence questionable environmental significance. The inclusion of unidentifiable individuals in higher-order taxonomic groupings in the SO-RAD dataset ensures the correct calculation of species relative abundances, however these taxa are not useful for analyses that rely on genus- or species-level data and might be removed.

300 Broad, higher-order categories that can be taken into consideration for removal from the SO-RAD dataset prior to analysis, include the indeterminate Nassellaria and Spumellaria taxonomic groups, along with taxa including, but not necessarily limited to, Litheliidae/Pyloniidae, *Spongotrochus/Spongopyle* sp., Trissocyclidae, and *Zygocircus productus* group. Some Phaeodarea taxa have been included in the census counts. Studies focusing only on polycystine radiolarians should remove the phaeodarians *Lirella melo*, and all species belonging to the *Euphysetta* and *Protocystis* genera.

305 **4.3.2 Grouping taxa**

Radiolarian reference datasets often contain taxonomic groupings made up of species/subspecies grouped at genus level, or of unrelated species with the same environmental affinities that may be routinely confused for one another during the counting procedure (e.g., Cortese and Prebble, 2015; Matsuzaki and Itaki, 2017). Grouping species is especially relevant for less abundant taxa, and for those with a particularly complex taxonomy. Grouping species and/or subspecies that have similar

310 distributions according to water column and modern sediment sampling can increase the statistical weight (and thus their chances of inclusion in a working reference dataset) of rare species that are known to carry a strong environmental signal. For example, Collosphaeridae is a taxonomic group that can be composed of several to tens of species that are most abundant at low latitudes (Cortese & Prebble, 2015; Biard et al., 2016). At high latitudes, their presence alone, even in very small numbers, may be indicative of warmer than usual conditions. It is, therefore, useful to group these rare species so that their signal is not
315 lost when removing rare species prior to analysis. Grouping species, while obviously leading to the loss of species-level information, also increases inter-comparability of results between different researchers/teams, as higher-level categories are generally more accommodating and less prone to identification mistakes based on fine skeletal differences. Recommended groupings are listed in Supplement 3 and are adapted from Cortese and Prebble (2015).

4.3.3 Removing rare taxa

320 Previous radiolarian-based palaeoenvironmental reconstructions and biogeographical studies have removed rare taxa using one, or both, of the following methods 1) removing species observed at fewer than a certain number of sites (e.g., Abelman et al., 1999; Boltovskoy & Correa, 2016) and/or 2) removing species based on their low abundances (e.g. <1%). In the latter situation, species were removed from previous datasets when their maximum relative abundance did not exceed 1% (Boltovskoy, 1987; Cortese & Prebble, 2015), 2% (Lozano & Hays, 1976; Abelman et al., 1999), or had fewer than a certain
325 number of individuals (Rogers & De Deckker, 2007). Conversely, some studies (e.g. Hernández-Almeida et al., 2020) elect not to remove rare species.

4.3.4 Removing taxa that do not inhabit the depth being reconstructed

As radiolarians are present throughout the water column, a surface sediment sample contains a collection of taxa that, while living, do not inhabit the same depth, but eventually end up in the same surface sediment sample. It is possible to limit the
330 reference dataset to taxa that are known to live at the depth for which a specific environmental variable is being reconstructed. Examples of studies where the authors have used a subset of taxa based on their vertical distributions include Abelman et al. (1999), Cortese and Abelman (2002), and Matsuzaki and Itaki (2017).

4.4 Update strategy and future plans for the SO-RAD dataset

There are noticeable gaps in the spatial coverage of the SO-RAD dataset, in particular the area between the Ross Sea and the
335 Antarctic Peninsula, comprising a vast part of the high-latitude south Pacific Ocean, and between Australia and the Antarctic continent in the Indian Ocean. The intention of the authors is that the SO-RAD dataset will constantly evolve as more surface sediment samples, collected during upcoming oceanographic expeditions, become available for census counts.

5 **Data Availability**

The SO-RAD dataset is available to download from doi.pangaea.de/10.1594/PANGAEA.929903 (Lawler et al., 2021).

340 6 **Conclusion**

The SO-RAD dataset contains census data for 238 radiolarian taxa from 228 Southern Ocean sites. The dataset is the most comprehensive Southern Ocean radiolarian census dataset to date. We have provided a detailed description of the methods used to build the dataset, along with an overview of the spatial and taxonomic coverage of the data. The SO-RAD dataset may be used as a reference dataset for palaeoenvironmental reconstructions, or in studies of modern radiolarian biogeography, ecology, and species diversity.

7 **Appendices**

- 1: Key to taxonomic homogenisation.
- 2: List of radiolarian taxa included in the SO-RAD dataset.
- 3: Recommended taxonomic groupings to simplify/harmonise the dataset.

350 8 **Author Contribution**

K-AL was responsible for the data curation (along with GC), formal analysis, investigation (along with GC and MC-M), data visualisation and preparation of the original draft. GC conceptualised and acquired funding for this study. Resources (access to samples) was provided by GC, HB, XC, AL, JR and LKA. GC and LKA supervised this project. All authors contributed to the review and editing of the manuscript.

355 9 **Competing Interests**

The authors declare that they have no conflict of interest.

10 **Acknowledgements**

This research was supported by the New Zealand Ministry of Business, Innovation and Employment through the Antarctic Science Platform (ANTA1801) and the Global Change through Time Programme (GCT SSIF, contract C05X1702), and through an IdEx (Initiative d'Excellence de l'Université de Bordeaux) Visiting Professorship grant (GC). K-AL is supported by an Australian Research Training Program (RTP) scholarship. A series of national and international core repositories

provided the surface sample material used in this study. These include the core repository at the National Institute of Water and Atmospheric Research (NIWA, New Zealand); three of IODP's (International Ocean Discovery Program) core repository centres: Kochi Core Center (jointly managed by Kochi University and JAMSTEC, the Japan Agency for Marine-Earth Science and Technology), the IODP West Coast repository at Scripps Institution of Oceanography, and the IODP Gulf Coast Repository at Texas A&M University; the Oregon State University Marine Geology Repository located at the College of Earth, Ocean, and Atmospheric Sciences; the Wood Hole Oceanographic Institution core repository; the Alfred Wegener Institute for Polar and Marine Research (AWI) core facility in Bremerhaven; and the IfM/GEOMAR core repository in Kiel. The help of these repositories, and the funding agencies providing support to them, is hereby greatly appreciated. The authors wish to thank the crews and scientists that collected the sediment core material used in this project, as well as Sonja Penafiel-Bermudez (GNS, Lower Hutt), Ute Bock (AWI, Bremerhaven) and Natalie Kozlowski (Colgate University) for preparing the microscopic slides.

Samples not previously published were collected through the listed missions and supported by the following research support bodies.

Sabrina Coast material: We thank the Marine National Facility (MNF) IN2017-V01 scientific party-led by the Chief Scientists LA and P. O'Brien, CSIRO MNF support staff and ASP crew members led by Capt. M. Watson on board the RV *Investigator*. Support was provided through the Australian Government's Australian Antarctic Science Grant Program (AAS #4333) and Australian Research Council (DP170100557).

Ross Sea material was supplied from the NIWA surface sediment and core repository and collected on numerous research voyages by NIWA and its predecessor NZOI since the 1950's. The samples have been collected by a range of different researchers on research voyages on a number of different vessels, funded by the New Zealand government through various funding programs over the last 70 years.

We thank Dr. Alain Mazaud and Dr. Elisabeth Michel for providing access to cores MD11-3353 and MD12-3396CQ. We are also grateful to the crews and scientists on-board of the R/V Marion Dufresne II during MD185 and MD189 INDIEN-SUD 1 and 2, and MD218 CROTALE expeditions. We thank the Institut Polaire Français, the Flotte Océanographique Française and the EquipEx CLIMCOR (ANR-11-EQPX- 0009- CLIMCOR) for their logistical and technical support.

We thank the scientific party, ASC support staff and ECO crew on RV/IB *Nathaniel B. Palmer* cruise NBP1402, supported by NSF Antarctic Integrated Systems Science Program grants #1143834, 1143836, 1143837, 1143843 and 1313826.

390 **References**

- Abelmann, A.: Radiolarian flux in Antarctic waters (Drake Passage, Powell Basin, Bransfield Strait), *Polar Biol.*, 12(3), 357–372, <https://doi.org/10.1007/BF00243107>, 1992.
- Abelmann, A. and Gersonde, R.: Biosiliceous particle flux in the Southern Ocean, *Mar. Chem.*, 35(1–4), 503–536, [https://doi.org/10.1016/S0304-4203\(09\)90040-8](https://doi.org/10.1016/S0304-4203(09)90040-8), 1991.
- 395 Abelmann, A. and Gowing, M. M.: Spatial distribution pattern of living polycystine radiolarian taxa - baseline study for paleoenvironmental reconstructions in the Southern Ocean (Atlantic sector), *Mar. Micropaleontol.*, 30, 3–28, 1997.
- Abelmann, A., Brathauer, U., Gersonde, R., Sieger, R. and Zielinski, U.: Radiolarian-based transfer function for the estimation of sea surface temperatures in the Southern Ocean (Atlantic Sector), *Paleoceanography*, 14(3), 410–421, <https://doi.org/10.1029/1998PA900024>, 1999.
- 400 Anderson, O. R.: Protozoa, Radiolarians, in *Encyclopedia of Ocean Sciences: A Derivative of the Encyclopedia of Ocean Sciences*, vol. 4, pp. 651–655, Elsevier Science & Technology, 2019.
- Armand, L., O’Brien, P. and Science Party.: Post-survey report: Interactions of the Totten Glacier with the Southern Ocean through multiple glacial cycles (IN2017-V01). <https://doi.org/10.4225/13/5acea64c48693>, 2018.
- Armand, L. K., Crosta, X., Romero, O. and Pichon, J.-J.: The biogeography of major diatom taxa in Southern Ocean sediments
- 405 1. Sea ice related species, *Palaeogeogr. Palaeoclimatol.*, 223(1–2), 93–126, <https://doi.org/10.1016/j.palaeo.2005.02.015>, 2005.
- Barnett, P. R. O., Watson, J. and Connelly, D.: A multiple corer for taking virtually undisturbed samples from shelf, bathyal and abyssal sediments, *Proc. R. Soc. Edinb. Sect. B Biol. Sci.*, 88, 304–305, <https://doi.org/10.1017/S0269727000014846>, 1986.
- Becker, R. A., Wilks, A. R., Brownrigg, R., Minka, T. P. and Deckmyn, A.: maps: Draw Geographical Maps. <https://CRAN.R-project.org/package=maps>, 2018.
- 410 Bianchi, C. and Gersonde, R.: Climate evolution at the last deglaciation: the role of the Southern Ocean, *Earth Planet. Sc. Lett.*, 228, 407–424, <https://doi.org/10.1016/j.epsl.2004.10.003>, 2004.
- Biard, T., Stemmann, L., Picheral, M., Mayot, N., Vandromme, P., Hauss, H., Gorsky, G., Guidi, L., Kiko, R. and Not, F.: In situ imaging reveals the biomass of giant protists in the global ocean, *Nature*, 532(7600), 504–507,
- 415 <https://doi.org/10.1038/nature17652>, 2016.
- Bivand, R., Keitt, T. and Rowlingson, B.: rgdal: Bindings for the “Geospatial” Data Abstraction Library. <https://CRAN.R-project.org/package=rgdal>, 2020.
- Boltovskoy, D.: Sedimentary Record of Radiolarian Biogeography in the Equatorial to Antarctic Western Pacific Ocean, *Micropaleontology*, 33(3), 267–281, <https://doi.org/10.2307/1485643>, 1987.
- 420 Boltovskoy, D.: Classification and distribution of South Atlantic Recent polycystine Radiolaria, *Palaeontol. Electron.*, <https://doi.org/10.26879/98006>, 1998.

- Boltovskoy, D.: Vertical distribution patterns of Radiolaria Polycystina (Protista) in the World Ocean: living ranges, isothermal submersion and settling shells, *J. Plankton Res.*, 39(2), 330–349, <https://doi.org/10.1093/plankt/fbx003>, 2017.
- 425 Boltovskoy, D. and Correa, N.: Biogeography of Radiolaria Polycystina (Protista) in the World Ocean, *Prog. Oceanogr.*, 149, 82–105, <https://doi.org/10.1016/j.pocean.2016.09.006>, 2016.
- Boltovskoy, D., Anderson, O. R. and Correa, N. M.: Radiolaria and Phaeodaria, in *Handbook of the Protists*, edited by J. M. Archibald, A. G. B. Simpson, C. H. Slamovits, L. Margulis, M. Melkonian, D. J. Chapman, and J. O. Corliss, pp. 1–33, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-319-32669-6_19-2, 2017.
- 430 Civel-Mazens, M., Crosta, X., Cortese, G., Michel, E., Mazaud, A., Ther, O., Ikehara, M., and Itaki, T.: Antarctic Polar Front migrations in the Kerguelen Plateau region, Southern Ocean, over the past 360 kyrs, *Global Planet. Change*, 202, 103526, <https://doi.org/10.1016/j.gloplacha.2021.103526>, 2021a.
- Civel-Mazens, M., Crosta, X., Cortese, G., Michel, E., Mazaud, A., Ther, O., Ikehara, M., and Itaki, T.: Impact of the Agulhas Return Current on the oceanography of the Kerguelen Plateau region, Southern Ocean, over the last 40 kyrs, *Quaternary Sci. Rev.*, 251, 106711, <https://doi.org/10.1016/j.quascirev.2020.106711>, 2021b.
- 435 Cortese, G. and Abelmann, A.: Radiolarian-based paleotemperatures during the last 160 kyr at ODP Site 1089 (Southern Ocean, Atlantic Sector), *Palaeogeogr. Palaeoclimatol.*, 182(3), 259–286, [https://doi.org/10.1016/S0031-0182\(01\)00499-0](https://doi.org/10.1016/S0031-0182(01)00499-0), 2002.
- Cortese, G. and Prebble, J.: A radiolarian-based modern analogue dataset for palaeoenvironmental reconstructions in the southwest Pacific, *Mar. Micropaleontol.*, 118, 34–49, <https://doi.org/10.1016/j.marmicro.2015.05.002>, 2015.
- Cortese, G., Abelmann, A. and Gersonde, R.: The last five glacial-interglacial transitions: A high-resolution 450,000-year record from the subantarctic Atlantic, *Paleoceanography*, 22(4), n/a-n/a, <https://doi.org/10.1029/2007PA001457>, 2007.
- 440 Crosta, X.: MD 218 / CROTALE cruise, RV Marion Dufresne. <https://doi.org/10.17600/18000886>, 2019.
- Crosta, X., Pichon, J.-J. and Burckle, L. H.: Application of modern analog technique to marine Antarctic diatoms: Reconstruction of maximum sea-ice extent at the last glacial maximum, *Paleoceanography*, 13(3), 284–297, <https://doi.org/10.1029/98PA00339>, 1998.
- 445 Crosta, X., Sturm, A., Armand, L. and Pichon, J.-J.: Late Quaternary sea ice history in the Indian sector of the Southern Ocean as recorded by diatom assemblages, *Mar. Micropaleontol.*, 50(3–4), 209–223, [https://doi.org/10.1016/S0377-8398\(03\)00072-0](https://doi.org/10.1016/S0377-8398(03)00072-0), 2004.
- Crosta, X., Romero, O., Armand, L. K., and Pichon, J.-J.: The biogeography of major diatom taxa in Southern Ocean sediments: 2. Open ocean related species, *Palaeogeogr. Palaeoclimatol.*, 223, 66–92, <https://doi.org/10.1016/j.palaeo.2005.03.028>, 450 2005.
- Esper, O. and Gersonde, R.: New tools for the reconstruction of Pleistocene Antarctic sea ice, *Palaeogeogr. Palaeoclimatol.* 399, 260–283, <https://doi.org/10.1016/j.palaeo.2014.01.019>, 2014a.
- Esper, O. and Gersonde, R.: Quaternary surface water temperature estimations: New diatom transfer functions for the Southern Ocean, *Palaeogeogr. Palaeoclimatol.*, 414, 1–19, <https://doi.org/10.1016/j.palaeo.2014.08.008>, 2014b.

- 455 Fatela, F. and Taborda, R.: Confidence limits of species proportions in microfossil assemblages, *Mar. Micropaleontol.*, 45(2), 169–174, [https://doi.org/10.1016/S0377-8398\(02\)00021-X](https://doi.org/10.1016/S0377-8398(02)00021-X), 2002.
- Ferry, A. J., Crosta, X., Quilty, P. G., Fink, D., Howard, W. and Armand, L. K.: First records of winter sea ice concentration in the southwest Pacific sector of the Southern Ocean, *Paleoceanography*, 30(11), 1525–1539, <https://doi.org/10.1002/2014PA002764>, 2015.
- 460 Fetterer, F., Knowles, K., Meier, W. N., Savoie, M. and Windnagel, A. K.: Sea Ice Index, Version 3, Boulder, Colorado USA. doi: <https://doi.org/10.7265/N5K072F8>, 2017.
- Garcia, H., Weathers, K., Paver, C., Smolyar, I., Boyer, T., Locarnini, M., Zweng, M., Mishonov, A., Baranova, O. and Seidov, D.: World Ocean Atlas 2018. Vol. 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+ nitrite, silicate), 2019a.
- Garcia, H., Weathers, K., Paver, C., Smolyar, I., Boyer, T., Locarnini, M., Zweng, M., Mishonov, A., Baranova, O. and Seidov, D.: World Ocean Atlas 2018, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Dissolved Oxygen Saturation., 2019b.
- Gersonde, R. and Zielinski, U.: The reconstruction of late Quaternary Antarctic sea-ice distribution—the use of diatoms as a proxy for sea-ice, *Palaeogeogr. Palaeoclimatol.*, 162(3), 263–286, [https://doi.org/10.1016/S0031-0182\(00\)00131-0](https://doi.org/10.1016/S0031-0182(00)00131-0), 2000.
- Gersonde, R., Crosta, X., Abelmann, A., and Armand, L.: Sea-surface temperature and sea ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum—a circum-Antarctic view based on siliceous microfossil records, 24, 869–896, <https://doi.org/10.1016/j.quascirev.2004.07.015>, 2005.
- 470 Hernández-Almeida, I., Boltovskoy, D., Kruglikova, S. B. and Cortese, G.: A new radiolarian transfer function for the Pacific Ocean and application to fossil records: Assessing potential and limitations for the last glacial-interglacial cycle, *Glob. Planet. Change*, 190, 103186, <https://doi.org/10.1016/j.gloplacha.2020.103186>, 2020.
- 475 Lawler, K.-A., Cortese, G., Civel-Mazens, M., Bostock, H., Crosta, X., Leventer, A., Lowe, V., Rogers, J., and Armand, L. K.: Southern Ocean RADiolarian dataset (SO-RAD). PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.929903>, 2021.
- Lazarus, D.: A brief review of radiolarian research, *Paläontol. Z.*, 79(1), 183–200, <https://doi.org/10.1007/BF03021761>, 2005.
- Lazarus, D., Suzuki, N., Caulet, J.-P., Nigrini, C., Goll, I., Goll, R., Dolven, J. K., Diver, P. and Sanfilippo, A.: An evaluated list of Cenozoic-Recent radiolarian species names (Polycystinea), based on those used in the DSDP, ODP and IODP deep-sea drilling programs, *Zootaxa*, 3999(3), 301, <https://doi.org/10.11646/zootaxa.3999.3.1>, 2015.
- 480 Lazarus, D., Suzuki, N., Ishitani, Y. and Takahashi, K.: *Paleobiology of the Polycystine Radiolaria*, Wiley Blackwell, 2021.
- Leventer, A. and Science Party: Sabrina Coast: Marine Record of Cryosphere - Ocean Dynamics., 2014.
- Locarnini, M., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Seidov, D., Weathers, K., Paver, C. and Smolyar, I.: World ocean atlas 2018, volume 1: Temperature, 2018.
- 485 Lozano, J. A. and Hays, J. D.: Relationship of Radiolarian Assemblages to Sediment Types and Physical Oceanography in the Atlantic and Western Indian Ocean Sectors of the Antarctic Ocean, in *Geol. Soc. Am. Mem.*, vol. 145, pp. 303–336, Geological Society of America, <https://doi.org/10.1130/MEM145-p303>, 1976.

- Lüer, V., Cortese, G., Neil, H. L., Hollis, C. J. and Willems, H.: Radiolarian-based sea surface temperatures and paleoceanographic changes during the Late Pleistocene–Holocene in the subantarctic southwest Pacific, *Mar. Micropaleontol.*, 490 70(3–4), 151–165, <https://doi.org/10.1016/j.marmicro.2008.12.002>, 2009.
- Matsuzaki, K. M. and Itaki, T.: New northwest Pacific radiolarian data as a tool to estimate past sea surface and intermediate water temperatures, *Paleoceanography*, 32(3), 218–245, <https://doi.org/10.1002/2017PA003087>, 2017.
- Mazaud, A. and Michel, E.: MD 185 / Indien Sud-2 Cruise Report, RV Marion Dufresne. <https://doi.org/10.17600/11200030>, 2011.
- 495 Mazaud, A. and Michel, E.: MD 189 / Indien Sud-2 Cruise Report, RV Marion Dufresne. <https://doi.org/10.17600/12200010>, 2012.
- McIlroy, D.: mapproj: Map Projections. <https://CRAN.R-project.org/package=mapproj>, 2020.
- Nigrini, C. and Moore, T. C. J.: A guide to modern Radiolaria, Cushman Foundation for Foraminiferal Research, Washington, D.C., 1979.
- 500 Orsi, A. H., Whitworth, T. and Nowlin, W. D.: On the meridional extent and fronts of the Antarctic Circumpolar Current, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 42(5), 641–673, [https://doi.org/10.1016/0967-0637\(95\)00021-W](https://doi.org/10.1016/0967-0637(95)00021-W), 1995.
- Panitz, S., Cortese, G., Neil, H. L. and Diekmann, B.: A radiolarian-based palaeoclimate history of Core Y9 (Northeast of Campbell Plateau, New Zealand) for the last 160 kyr, *Mar. Micropaleontol.*, 116, 1–14, <https://doi.org/10.1016/j.marmicro.2014.12.003>, 2015.
- 505 Petrushevskaya, M. G.: Radiolarians of orders Spumellaria and Nassellaria of the Antarctic region, in *Biological Reports of the Soviet Antarctic Expedition (1955 - 1958)*, edited by A. P. Andriyashev and P. V. Ushakov, pp. 2–186, 1967.
- Petrushevskaya, M. G.: On the natural system of polycystine Radiolaria (Class Sarcodina), in *Proceedings Second Planktonic Conference*. Tecnoscienza, Rome, edited by A. Farinacci, pp. 981–992, 1971.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, 510 Austria. <https://www.R-project.org/>, 2020.
- Radiolaria.org, <http://www.radiolaria.org/>, last access: 27 September 2019, n.d.
- Rogers, J. and De Deckker, P.: Radiolaria as a reflection of environmental conditions in the eastern and southern sectors of the Indian Ocean: A new statistical approach, *Mar. Micropaleontol.*, 65(3), 137–162, <https://doi.org/10.1016/j.marmicro.2007.07.001>, 2007.
- 515 Rogers, J. and De Deckker, P.: Environmental reconstructions of the upper 500 m of the southern Indian Ocean over the last 40 ka using Radiolarian (Protista) proxies, *Quaternary Sci. Rev.*, 30, 876–886, <https://doi.org/10.1016/j.quascirev.2011.01.006>, 2011.
- Schlitzer, R.: Ocean Data View. <https://odv.awi.de>, 2020.
- Sumner, M.D.: orsifronts: Southern Ocean Frontal Distributions (Orsi), 2015

- 520 Suzuki, N. and Not, F.: Biology and Ecology of Radiolaria, in *Marine Protists: Diversity and Dynamics*, edited by S. Ohtsuka, T. Suzuki, T. Horiguchi, N. Suzuki, and F. Not, pp. 179–222, Springer Japan, Tokyo, https://doi.org/10.1007/978-4-431-55130-0_8, last access: 27 September 2019, 2015.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemond, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K.,
- 525 Vaughan, D., Wilke, C., Woo, K. and Yutani, H.: Welcome to the Tidyverse, *Journal of Open Source Software*, 4(43), 1686, <https://doi.org/10.21105/joss.01686>, 2019.
- WOCE Data Products Committee: WOCE Data Products Committee, 2002, WOCE International Project Office, Southampton, UK., 2002.
- WoRMS Editorial Board: WoRMS - World Register of Marine Species, <http://www.marinespecies.org/> at VLIZ, last access:
- 530 27 September 2019, 2020.
- Yasuhara, M., Huang, H.-H., Hull, P., Rillo, M., Condamine, F., Tittensor, D., Kučera, M., Costello, M., Finnegan, S., O’Dea, A., Hong, Y., Bonebrake, T., McKenzie, R., Doi, H., Wei, C.-L., Kubota, Y. and Saupe, E.: Time Machine Biology: Cross-Timescale Integration of Ecology, Evolution, and Oceanography, *Oceanography*, 33(2), <https://doi.org/10.5670/oceanog.2020.225>, 2020.
- 535 Zweng, M., Seidov, D., Boyer, T., Locarnini, M., Garcia, H., Mishonov, A., Baranova, O., Weathers, K., Paver, C. and Smolyar, I.: *World ocean atlas 2018, volume 2: Salinity*, 2019.