

The Plankton Lifeform Extraction Tool: A digital tool to increase the discoverability and usability of plankton time-series data

Clare Ostle^{1*}, Kevin Paxman¹, Carolyn A. Graves², Mathew Arnold¹, Felipe Artigas³, Angus Atkinson⁴, Anaïs Aubert⁵, Malcolm Baptie⁶, Beth Bear⁷, Jacob Bedford⁸, Michael Best⁹, Eileen Bresnan¹⁰, Rachel Brittain¹, Derek Broughton¹, Alexandre Budria^{5,11}, Kathryn Cook¹², Michelle Devlin⁷, George Graham¹, Nick Halliday¹, Pierre Hélaouët¹, Marie Johansen¹³, David G. Johns¹, Dan Lear¹, Margarita Machairopoulou¹⁰, April McKinney¹⁴, Adam Mellor¹⁴, Alex Milligan⁷, Sophie Pitois⁷, Isabelle Rombouts⁵, Cordula Scherer¹⁵, Paul Tett¹⁶, Claire Widdicombe⁴, and Abigail McQuatters-Gollop⁸

1¹The Marine Biological Association (MBA), The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK.

2²Centre for Environment Fisheries and Aquaculture Science (Cefas), Weymouth, UK.

3³Université du Littoral Côte d'Opale, Université de Lille, CNRS UMR 8187 LOG, Laboratoire d'Océanologie et de Géosciences, Wimereux, France.

4⁴Plymouth Marine Laboratory, Prospect Place, Plymouth, PL1 3DH, UK.

15 5⁵Muséum National d'Histoire Naturelle (MNHN), CRESCO, 38 UMS Patrinat, Dinard, France.

6⁶Scottish Environment Protection Agency, Angus Smith Building, Maxim 6, Parklands Avenue, Eurocentral, Holytown, North Lanarkshire ML1 4WQ, UK.

7⁷Centre for Environment Fisheries and Aquaculture Science (Cefas), Lowestoft, UK.

8⁸Marine Conservation Research Group, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, UK.

20 9⁹The Environment Agency, Kingfisher House, Goldhay Way, Peterborough, PE4 6HL, UK.

10¹⁰Marine Scotland Science, Marine Laboratory, 375 Victoria Road, Aberdeen, AB11 9DB, UK.

11¹¹Department of Coastal Systems, NIOZ Royal Netherlands Institute for Sea Research, Den Burg, Texel, The Netherlands.

12¹²National Oceanography Centre, European Way, Southampton, S014 3ZH, UK.

13¹³Swedish meteorological and hydrological institute, Sven Kallfelts gata 15, 426 71 Vastra Frolunda. Sweden.

25 14¹⁴Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute, 18a Newforge Lane, Belfast BT9 5PX.

15¹⁵Trinity Centre for Environmental Humanities, Department of History, School of Histories and Humanities, Trinity College, University of Dublin, Ireland.

16¹⁶Scottish Association for Marine Science, Scottish Marine Institute, Oban, PA37 1QA, UK.

Correspondence: *Clare Ostle (claost@mba.ac.uk)

1 Abstract

Plankton form the base of the marine food web and are sensitive indicators of environmental change. Plankton time-series are therefore an essential part of monitoring progress towards global biodiversity goals, such as the Convention on Biological Diversity Aichi Targets, and for informing ecosystem-based policy, such as the EU Marine Strategy Framework Directive.

35 Multiple plankton monitoring programmes exist in Europe, but differences in sampling and analysis methods prevent the integration of their data, constraining their utility over large spatio-temporal scales. The Plankton Lifeform Extraction Tool brings together disparate European plankton datasets into a central database from which it extracts abundance time-series of plankton functional groups, called ‘lifeforms’, according to shared biological traits. This tool has been designed to make complex plankton datasets accessible and meaningful for policy, public interest, and scientific discovery. It allows examination

40 of large-scale shifts in lifeform abundance or distribution (for example, holoplankton being partially replaced by meroplankton), providing clues to how the marine environment is changing. The lifeform method enables datasets with different plankton sampling and taxonomic analysis methodologies to be used together to provide insights into the response to multiple stressors and robust policy evidence for decision making. Lifeform time-series generated with the Plankton Lifeform Extraction Tool currently inform plankton and food web indicators for the UK’s Marine Strategy, the EU’s Marine Strategy

45 Framework Directive, and for the Convention for the Protection of the Marine Environment of the North- East Atlantic (OSPAR) biodiversity assessments. The Plankton Lifeform Extraction Tool currently integrates 155,000 samples, containing over 44 million plankton records, from 9 different plankton datasets within UK and European Seas, collected between 1924 and 2017. Additional datasets can be added, and time-series updated. The Plankton Lifeform Extraction Tool is hosted by The Archive for Marine Species and Habitats Data (DASSH) at <https://www.dassh.ac.uk/lifeforms/>. The lifeform outputs are linked

50 to specific, doi-ed, versions of the Plankton Lifeform Traits Master List and each underlying dataset.

2 Introduction

Plankton form the foundation of the marine food web, help to regulate ocean chemistry, and provide approximately half of the world's oxygen (Capuzzo et al., 2018; Falkowski, 2012). Globally, plankton communities are undergoing significant changes in distribution (Reid et al., 2016), community composition (Beaugrand et al., 2002), phenology (Edwards and Richardson, 2004), and productivity (Kulk et al., 2020). These changes vary in space and time, reflecting both direct and locally acute anthropogenic pressure on the marine environment, such as nutrient loading, and wider-scale climate-driven changes in ocean chemistry and temperature (Beaugrand et al., 2010; Bedford et al., 2020a).

Plankton have short life cycles, drift freely in the ocean and have wide distributions. For these reasons they are considered to be particularly sensitive indicators to climate change (Richardson, 2008). Changes in the composition and abundance of plankton can have negative impacts on industries such as fisheries and aquaculture (Richardson et al., 2009; Schmidt et al., 2020). As the base of the food web, they are a key element of the ecosystem approach to marine management (Morishita, 2008). Monitoring plankton communities over wide spatial and long temporal scales can help tease apart the prevailing footprint of climate change on marine ecosystems from other, more localised pressures, for example, pollution, nutrient loading and fishing (Bedford et al., 2020b). Consequently, plankton time-series play an increasingly important role in decision-making and provision of advice. Plankton indicators contribute to the delivery of global, regional and national policy drivers such as the Convention on Biological Diversity's Aichi targets (Chiba et al., 2018), the regional Convention for the Protection of the Marine Environment of the North-East Atlantic (OSPAR) (OSPAR, 2017), and biodiversity state in the European Union's Marine Strategy Framework Directive (MSFD) and the UK Marine Strategy (McQuatters-Gollop et al., 2019).

To map changes in ocean colour, Earth Observation (EO) satellite tools provide unparalleled spatial coverage, and now offer the prospect of 20 years of ocean colour data, with increasingly resolved information, for example on trends of specific size-fractions of Chlorophyll (Schmidt et al., 2020). However, the EO techniques are still not yet sufficiently developed to obtain information on changes in abundance of the key component planktonic functional groups, particularly for the zooplankton.

80 Additionally, some taxonomic datasets now have up to 90 years of data which provide a critical perspective in assessing long-term change and which is unparalleled by satellites. We therefore need to maintain direct monitoring approaches for a holistic view of the plankton, and northwest European waters are particularly well-blessed with these time-series.


85 Although there are a number of programmes that monitor plankton in Northwest European waters, they operate at different spatial scales, from fixed-point sampling stations to long-distance continuously sampled ship transects (O'Brien et al., 2017). Furthermore, European plankton surveys employ different sampling methods, enumerate specimens at a variety of taxonomic levels and employ different counting regimes (Raybaud et al., 2011). These methodological differences and the lack of direct comparability between datasets has meant that the tools to use all available datasets together to produce a comprehensive assessment have only recently been developed (Bedford et al., 2020a; McQuatters-Gollop et al., 2019).
90 While most datasets are regularly submitted to appropriate data repositories (e.g.: the Ocean Biodiversity Information System: OBIS; the British Oceanographic Data Centre: BODC, or the PANGAEA data publisher for earth and environmental science) and some are available through institutional websites or data centres, the aggregation of plankton data into functional groups (or 'lifeforms' e.g.: diatoms, dinoflagellates, holoplankton, meroplankton) has not yet been linked to traceable dataset versions or been
95 possible to apply in an accessible, transparent and centralised way. Accordingly, understanding of plankton change across multiple spatial and temporal scales has been limited. The International Group of Marine Ecological Time Series IGMETS (<https://igmets.net>; O'Brien et al., 2017), represents valuable progress towards this goal: it provides a global-scale compilation of pelagic time series, with a tool to summarise visualisations of trends across a variety of temporal and spatial scales. However, this initiative
100 summarises time trends of highly aggregated variables (e.g.: total zooplankton) for multiple sites. Trajectories of the key component plankton functional groups are not described, and the underlying data products are not made available to users for further analysis. Aggregating these disparate plankton datasets increases the spatial-temporal scope of analysis, increases their robustness and provides decision makers with more scientifically robust evidence.

105 Building on previous work (Gowen et al., 2011; Scherer et al., 2014; Tett et al., 2008, 2013) an indicator of shifts in plankton structure based on time-series of broad plankton functional groups, called 'lifeforms',

has been developed for use in policy assessments (McQuatters-Gollop *et al.*, 2019). The term ‘lifeform’ is derived from work carried out by Margalef (1978), to distinguish between diatoms and dinoflagellates based on traits related to survival in specific hydrodynamic conditions. Lifeforms differ slightly from the term ‘Plankton Functional Type’ (PFT), in that PFTs are often used to describe plankton based on their ecosystem function and not on their traits. This indicator enables plankton datasets with different sample collection and analysis routines to be used congruently to investigate changes in pelagic habitat functioning. By using these pre-defined lifeforms to group plankton taxa, the new Plankton Lifeform Extraction Tool (PLET), hosted by the Archive for Marine Species and Habitats Data (DASSH, <https://www.dassh.ac.uk>), brings together disparate plankton datasets, increasing their accessibility and promoting compliance with the FAIR data principles (Wilkinson *et al.*, 2016). The PLET enables the user to investigate multiple datasets to assess changes in plankton ecology at multiple spatio-temporal scales using a consistent plankton indicator for the first time. As time-series grow in length and/or spatial distribution, and new plankton time-series are established, additional plankton taxa and datasets can be added to the PLET in order to improve future biodiversity assessments. The tool is a key step towards transparent and standardised assessment allowing the integration of information from multiple datasets at multiple spatial and temporal scales.

3 Plankton datasets

In its current form, the PLET integrates 155 thousand samples containing over 44 million plankton records from 9 different data providers around the UK and European Seas, collected between 1924 and 2017 (**Table 1**). There are a number of plankton trait datasets and plankton compilation efforts that are complementary to the PLET with the potential to feed into future versions of the tool, such as the nutrient utilisation traits dataset (Edwards et al., 2015), the Baltic Sea phytoplankton traits dataset (Klais et al., 2017), and the French lakes phytoplankton traits database (Laplace-Treyture et al., 2021). While these are highly valuable resources, the authors are not aware of a platform to bring such information together and disseminate it in a consistent format. The design of PLET allows for this lifeform extraction and dissemination, with the aim to incorporate further plankton trait datasets in future versions. Flexibility of the PLET design allows existing time-series to be updated and new time-series to be added, continuing the expansion of integrated data sets beyond the UK, where policy reporting motivated its initial development. Plankton time-series have been collected both along transects and at fixed-point stations (**Figure 1**). These datasets, which underpin the PLET lifeform outputs, enumerate plankton in taxon groupings (see the Plankton Lifeform Traits Master List (UK Pelagic Habitats Expert Group, 2020); <https://www.dassh.ac.uk/doitool/data/1709>).



140 **Figure 1. Spatial coverage of plankton measurements** currently integrated with the PLET tool for lifeform extraction. See data provider and station information in **Table 1**, individual station names are given next to the symbols, while symbols designate data providers as shown in the legend. The sampling transects for the CPR are coloured by date sampled, with pre-1980 in light grey, 1980-1999 in darker grey, and the most recent 2000-onwards in black. Coloured regions indicate how data are summarised for presentation of Lifeform outputs in **Section 6**: blue: Celtic Seas; red: northern North Sea; green: southern North Sea; light yellow: Channel.

145 **Table 1. Plankton data currently held in PLET** and used to produce the aggregated lifeform outputs. For un-aggregated plankton data, contact information and institute-specific data holdings (where available) are given for each data institute. Most of these time series are ongoing and many sample at higher temporal resolution than the monthly average data held in PLET. Prospective users for these higher resolution versions of the respective time series are encouraged to consult with the contact people listed below.

Institute, dataset name, <i>primary contact</i> ; data web address, [PLET doi]	Region or Station Name	Sampling Period	
		Phytoplankton	Zooplankton
The Marine Biological Association (MBA) , Continuous Plankton Recorder Survey, <i>David Johns</i> (djoh@mba.ac.uk); https://data.cprsurvey.org/datacatalog/ https://doi.org/10.17031/1629 (CPR and Johns, 2019)	UK and European Seas	1958-2017	1958-2017
The Marine Biological Association (MBA) , Station sampling, <i>Rachel Brittain</i> (racbri@mba.ac.uk) https://doi.org/10.17031/1636 (MBA, 2019)	L5	not determined	1924 - 1940, 1945 - 1987, 2001 - 2013
	E1		
Plymouth Marine Laboratory (PML) <i>Angus Atkinson</i> (aat@pml.ac.uk); http://www.westernchannelobservatory.org.uk/ https://doi.org/10.17031/1632 (PML, 2019)	L4	1992-2015	1988-2017
Centre for Environment Fisheries and Aquaculture Science (Cefas) , Smartbuoys, <i>Michelle Devlin</i> (michelle.devlin@cefas.ac.uk); https://www.cefas.co.uk/cefas-data-hub/smartbuoys/ https://doi.org/10.17031/1634 (CEFAS, 2019)	Dowsing	2000-2017	not determined
	Gabbard and West Gabbard	2001-2017	
	Liverpool Bay	2002-2017	
	Warp	2001-2012	
Environment Agency (EA) <i>Mike Best</i> (mike.best@environment-agency.gov.uk) https://doi.org/10.17031/1635 (EA, 2019)	UK coastal and transitional waters	2010-2017	not determined
Marine Scotland Science (MSS) <i>Eileen Bresnan & Margarita Machairopoulou</i> (Eileen.Bresnan@gov.scot ; Margarita.Machairopoulou@gov.scot) https://data.marine.gov.scot/search/type/dataset https://doi.org/10.17031/1637 (MSS, 2019)	Stonehaven	1997-2017	1999-2017
	Loch Ewe	2002-2017	2002-2017
	Scapa (Orkney Islands)	2001-2017	not determined
	Scalloway (Shetland Islands)	2001-2017	
Swedish Meteorological and Hydrological Institute (SMHI) <i>Marie Johansen</i> (marie.johansen@smhi.se) https://sharkweb.smhi.se/ https://doi.org/10.17031/1633 (SMHI, 2019)	Swedish west coast	1986-2015	1998-2015
Scottish Environment Protection Agency (SEPA) <i>Malcolm Baptie</i> (Malcolm.Baptie@sepa.org.uk) https://doi.org/10.17031/b84a-7951 (SEPA, 2020)	Forth	2007-2017	2014 - 2017
	Clyde		
Scottish Association for Marine Science (SAMS) <i>Paul Tett</i> (Paul.Tett@sams.ac.uk) https://doi.org/10.17031/nz24-br35 (SAMS, 2020)	Lorne Pelagic Observatory	1970-2015	not determined

3.1 Plankton sampling and analysis methodology

All individual datasets that have been added to the PLET have been pre-processed to ensure suitability for extraction of monthly-aggregated lifeform data products. Pre-processing was the responsibility of the individual data providers. Examples of pre-processing required are (i) the exclusion of instances of
155 ‘double counting’ where, for example, a taxon is included in both higher and lower taxonomic groupings within the same dataset and (ii) the removal of taxa that have not been looked for (recorded) over the entire time-period to avoid apparent changes in lifeform abundance due to methodological changes. When new datasets are submitted to the PLET the data providers supply aphia IDs of all of the taxa within their dataset. Following the pre-processing of the data by the data providers, the data manager of PLET and
160 the manager of the Plankton Lifeform Traits Master List does a check of the submitted aphia IDs to highlight any missing taxa. Any taxa that are not included within the Plankton Lifeform Traits Master List are checked for compatibility with the lifeforms and their traits are added in discussion with an expert group and the data providers.

Existing datasets were gathered through a data call issued by OSPAR in 2016. The purpose of the data
165 call was to gather plankton datasets to use for assessment and reporting for the European Union’s and individual Member States’ Marine Strategy Framework Directive initial biodiversity assessment in 2017 (<https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/>). A simple data submission template was developed as part of this process and is now available on the PLET website for wider use.
170 To make data submission as simple and easy as possible, the template allows data-holders to submit the datasets in either list (long) or matrix (wide) formats. A data archiving and access permission agreement form is also available from the PLET website, which allows data-holders to specify their preferred level of data access, such as full access to raw data or access to lifeform data products only.

All plankton records currently included have been identified using light microscopy. For simplicity we
175 use the term “phytoplankton” to mean protist cells, mindful that these include a spectrum of auto-, mixo- and heterotrophic forms (Flynn et al., 2013). This terminology is used to differentiate from “zooplankton” which are the metazoans usually counted from net haul samples. For quality assurance, analysts

participate in NMBAQC (the Northeast Atlantic Marine Biological Analytical Quality Control Scheme) and the International Phytoplankton Intercomparison external identification ring trials, although these do not cover the full length of some of the historical data sets. Field abundance, in individuals per unit volume, is calculated as sample abundance multiplied by subsample factor, divided by the sampled water volume. Concentrations of phytoplankton identified by light microscopy are typically expressed as numbers (cells) per mL, and those of zooplankton are typically expressed as numbers (individuals) per m^{-3} .

185 3.1.1 *Continuous Plankton Recorder Survey (Marine Biological Association)*

The Continuous Plankton Recorder (CPR) is a marine sampler that is towed behind volunteer ships of opportunity at speeds of up to ~ 20 knots and samples at a depth of ~ 7 m below the surface. Plankton have been sampled on routes crossing the North Atlantic and NW European shelf seas using a consistent methodology since 1958.

190 The CPR unit is a metal casing in the shape of a ~ 1 m torpedo that houses a roll of silk which automatically rotates using a geared propeller system. The seawater enters the front aperture where plankton and small particles are captured onto the rotating silk, which has a mesh size of $270 \mu\text{m}$. This silk is stored in 4 % buffered formalin to preserve the sample until microscopic analysis at the laboratory in Plymouth. The silk is cut into pre-defined sections that represent one sample and equate to 10 nautical
195 miles of tow. Phytoplankton and zooplankton are identified and counted at different stages of the microscopic analysis: semi-quantitative count of phytoplankton across 20 fields of view per sample, quantitative count of all zooplankton ≥ 2 mm (these are picked off the silk for identification), and semi-quantitative traverse count of all zooplankton < 2 mm.

For a more in-depth description of the sampling methodology please refer to Richardson et al. (2006).
200 CPR monthly abundance counts from 1958 to 2017 are available from the following open access data portal: <https://data.cprsurvey.org/datacatalog/>.

3.1.2 *Western Channel Observatory (Marine Biological Association and Plymouth Marine Laboratory)*

The Marine Biological Association (MBA) and Plymouth Marine Laboratory (PML) jointly sample at three offshore stations in the western English Channel as part of their Western Channel Observatory

205 (<https://westernchannelobservatory.org.uk>). These stations are termed: L4 (50.25°N, 4.3°W; approx. 55 m water depth) 13 km south-west of Plymouth, which can be regarded as a coastal station, albeit in transitionally stratified water; L5 (50.18°N, 4.3°W; approx. 58 m depth) is positioned between coastal and offshore waters, and E1 (50.03°N, 4.37°W; approx. 70 m depth) is 40 km offshore in seasonally stratified water. Sampling at these historical sites began in 1924 with interruptions between 1940-45 and 1987-
210 2001. Sampling frequency has varied between weekly and fortnightly; current sampling is weekly at station L4 and, weather-permitting, fortnightly at L5 and E1.

The phytoplankton and zooplankton time-series at L4 are provided by PML. Sampling for phytoplankton began in 1992 and for mesozooplankton in 1988. Detailed phytoplankton taxonomic microscope counts are from water samples collected at 10 m depth. These samples are preserved in 2% acid Lugol's iodine
215 solution and enumerated for all taxa larger than approximately 2 μm using the Utermöhl (1958) technique, usually settling 50 ml (Widdicombe et al., 2010). Mesozooplankton are collected each week in two replicate 0-50 m vertical hauls with a WP2 net (0.57 m diameter, 200 μm mesh-size). Each of these are analysed in two aliquots, the first being a stempel pipette – derived small subsample for enumeration of the more numerous taxa and the second larger fraction, often one-half to one-eighth, analysed for the
220 larger or rarer taxa.

Weekly densities are calculated as the average of the two separate net hauls. Environmental conditions and the mesozooplankton sampling and analysis methods are described in detail in Atkinson et al. (2015). Macroplankton and larval fish sampling at the WCO sites is carried out by the MBA. Although net design and methods of deployment have changed on several occasions, care has been taken to ensure that
225 sampling characteristics have not altered appreciably. The 1 m^2 Young Fish Trawl (YFT), fitted with a 700 μm knitted mesh is hauled for 20 minutes in an oblique profile to an ideal depth \sim 5 m above the seabed. Depth and temperature profiles are occasionally recorded and the volume of water filtered calculated using flow data recorded by a flowmeter fitted across the net mouth. The samples are preserved in 4 % buffered formalin and analysed as soon as possible after collection using a WILD M5 binocular
230 microscope. Results are standardized to number of individuals per 4000 m^3 in order to mitigate historical changes in sampling gear and deployment.

A comprehensive summary of these macroplankton sampling methods and analysis is given in Southward et al. (2004) and references therein.

3.1.3 *Smartbuoys (Centre for Environment Fisheries and Aquaculture Science)*

235 Water samples for phytoplankton analysis are collected from several of the Centre of Environment Fisheries and Aquaculture Science (Cefas) ‘Smartbuoy’ moorings using automated water samplers mounted at 1 m below the surface. Time-series at approximately monthly resolution from four buoy stations are available: Dowsing off the Humber estuary (51.53°N, 1.05°E, sampled 2000-present), Gabbard/West Gabbard off the Thames estuary (51.95°N, 2.11°E, sampled 2001-present), Warp in the
240 outer Thames estuary (51.52°N, 1.028°E, sampled 2001-2012), and Liverpool Bay (53.53 °N, 3.35°W, sampled 2002-present).

Water samplers are pre-programmed to collect 150 mL samples on an approximately weekly cycle into sample bags pre-spiked with acidified Lugol’s iodine solution. Phytoplankton samples are returned for analysis at Cefas every 1–3 months, where they are decanted into 175 mL glass jars and topped up with
245 acidified Lugol’s iodine. A minimum of one sample per month is selected for analysis from each deployment location where sample availability allows. Samples are analysed at Cefas using the Utermöhl (1958) technique under inverted Olympus microscopes within 1 year of collection. Species are identified and enumerated to the lowest possible taxonomic level and counts recorded in cells per litre.

More detailed methodology is available in Weston et al. (2008) and Greenwood (2019). Plankton and
250 environmental parameters from the Smartbuoy monitoring programme are available from the Cefas Data Hub: <https://www.cefas.co.uk/data-and-publications/smartbuoys/>.

3.1.4 *England’s Estuarine and Coastal Waters (Environment Agency)*

The Environment Agency (EA) and its predecessors have been collecting phytoplankton on targeted campaigns since the 1990’s, however from the inception of EU Water Framework Directive (WFD; EU,
255 2000) monitoring in 2006, Environment Agency routine phytoplankton samples have been collected from sites in near-shore WFD waterbodies from boats, or occasionally, jetties or bridges in estuaries Devlin et al. (2012).

Sampling in WFD transitional and coastal waters typically consists of one sample per calendar month from 3 to 5 sites per water body. Ideally, samples should be 28-31 days apart throughout the year. There must be at least a 14-day interval between sampling occasions at each site. Phytoplankton samples are taken in the mixed surface layer usually between 1-2 m below the water surface using a standard NIO/Niskin-style water sampler, avoiding the surface film and without disturbing bottom sediments. In coastal or non-turbid waters >5m depth, the diurnal vertical migration of phytoplankton with light availability is accommodated by collection during daylight hours. However, for some samples, the use of integrated depth sampling using a Lund-type tube system negated the need to constrain the sampling window to daylight hours. Samples are collected in 250ml clear PET bottles filled to approximately 90%, leaving sufficient headspace to allow for preservation and homogenisation. Samples are preserved with acidified Lugol's iodine, and stored in the dark, ideally at a temperature of $3^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for no longer than 6 months. Samples are analysed using the Utermöhl (1958) method under inverted microscopes. Analysis was conducted at Cefas until 2013, then at both Cefas and an external laboratory from 2013 onwards. Some samples are analysed by multiple analysts to check for comparability of results.

3.1.5 The Scottish Coastal Observatory (Marine Scotland Science)

Marine Scotland Science (MSS) routinely samples the plankton in Scottish waters as part of the Scottish Coastal Observatory. Weekly phytoplankton samples have been collected from Stonehaven (56.96°N , 2.11°W) since 1997, Scapa (Orkney Islands; 58.74°N , 2.97°W) since 2001, Loch Ewe (57.84°N , 5.65°W) since 2023 and Scalloway (Shetlands, 60.18°N , 1.28°W) since 2001. Meso-zooplankton have been sampled, also weekly at Stonehaven since 1999 and Loch Ewe since 2002.

Phytoplankton samples are collected using a 10 m integrated tube sampler. A 1 L subsample is preserved with 0.5% acidic Lugol's iodine and returned to the MSS Marine Laboratory. Phytoplankton samples are analysed using a modified Utermöhl (1958) technique. Phytoplankton samples are analysed using an inverted Zeiss Axiovert microscope. The presence/absence of all cells in the chamber are recorded and fields of view across a transect are counted at X200 magnification. Phytoplankton are identified to the lowest taxonomic level possible, however due to the limitations of light microscopy and the Lugol's fixative in some instances a genus level identification or 'unidentified' category is assigned.

285 Zooplankton samples are collected using 40 cm diameter bongo nets fitted with 200 μm mesh and filtering
cod ends. The nets are hauled vertically from near bottom (45 m at Stonehaven and 35 m at Loch Ewe)
to surface at a speed of 1 m s^{-1} . The samples are immediately fixed in 4 % borax buffered formaldehyde
for later analysis in the laboratory. Zooplankton samples are analysed in the laboratory using a Zeiss
Stemi SV-11 stereomicroscope. Larger zooplankton categories (such as *Calanus* spp., chaetognaths,
290 jellyfish, euphausiids etc.) are identified and enumerated from the whole sample. The remaining
zooplankton categories are identified and enumerated from a series of subsamples (of variable volumes
depending on concentration of animals but a minimum 2.5 % of the whole sample) so that at least 100
animals of the most common taxa are recorded. Most taxa are identified to the lowest taxonomic level
possible, whilst other animals are recorded at Class or Phylum level.

295 More detailed methodology is available in Bresnan et al. (2016). Phytoplankton monthly densities from
Stonehaven, Loch Ewe, Scapa and Scalloway, and zooplankton weekly densities from Stonehaven and
Loch Ewe are available from: <https://data.marine.gov.scot/search/type/dataset>.

3.1.6 Scotland Coastal Stations (Scottish Environment Protection Agency)

The Scottish Environment Protection Agency (SEPA) collects plankton samples at two near-shore
300 stations (Forth: 56.03°N, 3.18°W; Clyde: 55.95°N, 4.89°W). Monthly samples for phytoplankton have
been collected since 2007 and zooplankton since 2014.

Phytoplankton samples are collected using an integrated tube column water sampler with foot valve and
closure tap, which is lowered open to 10 m depth. The closure tap is then moved to the closed position
and the sampler retrieved. The foot valve is opened and the contents of the sampler are emptied into a
305 rinsed bucket. A 250 ml sample bottle prefilled with 2.5 ml of 5% w/v Lugol's iodine solution is gently
submerged to fill with water from the bucket. The sample bottle is gently inverted to mix preservative
and stored in the dark in a refrigerator at 4 °C. Samples of phytoplankton are removed from cold storage
and left to acclimatise at room temperature for 24 hours after which they are gently inverted 100 times to
re-suspend settled cells and a volume of sample, typically 50 ml, poured into a sample tube and left to
310 settle for 24 hours. After this time, 40ml of supernatant is drawn off slowly and discarded. The remaining
10 ml of sample is then gently inverted 100 times before being carefully poured into a 10 ml Utermöhl

(1958) counting chamber. This is then left to settle for a further 24 hours before being analysed on an inverted microscope (Leica DM IRB or Leica DMI4000B – Wetzlar, Germany or Zeiss Axiovert S100 – Jena, Germany). The chamber plate is scanned to assess rough composition of the sample and to determine
315 if settled cells are randomly distributed. Depending on the cell type, size and density, cell counts are made of the whole counting chamber, a number of transects of the widest point, or a number of random fields of view. At least 400 cells are counted when employing transect or field of view counting strategies. Field abundance in cells L⁻¹ is calculated by multiplying sample count by microscope subsample factor and 1000 divided by settled volume.

320 Zooplankton samples are taken with a 27 cm diameter net fitted with a 200 µm mesh with a non-filtering 1 litre cod end. A Hydro-bios (Kiel-Altenholz, Germany) digital flow meter with a back run stop is fitted to the mouth of the net in order to determine the volume hauled, and therefore abundance in individuals m³. The net is deployed vertically from near-bottom to the surface at approximately 0.5 m s⁻¹. Upon recovery, the net is rinsed with seawater and the contents of the cod ends are transferred into a sample
325 bottle and preserved in 4 % borax buffered formaldehyde. These samples are gently rinsed through a 63 µm wire mesh sieve for microscopic analysis using Leica (Wetzlar, Germany) M165C microscopes. Abundance is determined by counting in the full sample any zooplankton larger than stage IV *Calanus* (including and from copepodite stage V) and enumerating all other zooplankton in a subsample taken using a Folsom or Motoda splitter or a plunge sampler as appropriate to achieve an acceptable density of
330 zooplankton, being no less than 100 of the most abundant taxa.

3.1.7 Lorne Pelagic Observatory (Scottish Association of Marine Science)

Phytoplankton samples have been collected weekly at the Lorne Pelagic Observatory (56.48 °N, 5.5°W) since 1970 by the Scottish Association of Marine Science (SAMS). Water samples for microplankton (i.e.: phytoplankton and pelagic micro-heterotrophs) are taken with water bottles and in some cases with
335 a 10 m integrating hose. They are preserved with 0.5% acidic Lugol's iodine and volumes of 10 to 50 mL sedimented for counting using the Utermöhl (1958) technique and Wild and Zeiss inverted microscopes equipped with phase contrast. Depending on abundance and organism size, a variety of counting patterns are used, ranging from examination of the whole base of the sedimentation chamber at low power to narrow transects or a few fields at high power.


340 3.1.8 *Swedish West Coast (Swedish Meteorological and Hydrological Institute)*

The Swedish Meteorological and Hydrological Institute (SMHI) samples both phytoplankton and zooplankton at four stations on the Swedish west coast. Phytoplankton are sampled as an integrated sample using a hose (0-10 m) and preserved in acidic Lugol's; alkaline Lugol's is used for counts of coccolithophores. Twenty-five ml of each sample is analysed using the Utermöhl (1958) method. The
345 samples are stored in the dark and at room temperature prior to analyses. Zooplankton are sampled with a WP2 net (100 µm mesh size), and an integrated sample is taken from 0-25 m. Samples are preserved in formalin and stored in the dark prior to analyses. The subsample volume used when counting depends on the concentration of copepods in the sample to enable statistically sound data. Data available from:
[https://www.smhi.se/en/services/open-data/national-archive-for-oceanographic-data/download-data-
350 1.153150.](https://www.smhi.se/en/services/open-data/national-archive-for-oceanographic-data/download-data-1.153150)

3.2 *Spatio-temporal data distribution*

355 The plankton datasets currently available for lifeform extraction by the PLET have variable spatial and temporal extents, summarised herein into the four regions shown in **Figure 1**. Within each region, the availability of plankton data over time differs between datasets (**Figure 2**). Due to their high spatial coverage the CPR and EA datasets contain the largest numbers of samples available for any month, within each region. The number of samples at fixed-point sampling stations shows variations in sampling
360 frequency; in some cases this has changed over the course of the time-series (for example, in the Celtic Seas both Cefas Smartbuoy and the SAMS dataset).

While sampling is typically carried out weekly or monthly in order to capture the seasonal cycle of the plankton community structure and rapid changes associated with plankton bloom events, several datasets include sampling gaps and changes in sampling intensity for a variety of reasons (**Figure 3**). For example,
365 the EA dataset sampling frequency (and spatial distribution of samples) increased alongside the implementation of the Water Framework Directive (EU, 2000) in 2007, while the SAMS time-series stopped between 1982 and 1999. Missing months in the Cefas Smartbuoy time-series largely indicate failures of the automated sampling system, or sample loss related to logistical delays in buoy servicing. Ongoing sampling of all time-series are at risk of additional reductions in sampling frequency and quality
370 related to funding (McQuatters-Gollop et al., 2017; Zingone et al., 2015).



75

Figure 2. Sampling effort of each dataset within each region: number of sampling timepoints collected per month for each dataset within each of the regions defined in **Figure 1**, except for MSS stations in the northern North Sea where only Stonehaven is shown as an example of the three MSS stations in that region. Note that axis limits are not fixed between panels. Bar colour indicates spatial region (see **Figure 1**); blue: Celtic Seas; red: northern North Sea; green: southern North Sea; light yellow: Channel.




Figure 3. Number of months sampled per year for each dataset within each region: number of months sampled within each year, for each dataset within each of the regions defined in **Figure 1**, except for MSS stations in the northern North Sea where only Stonehaven is shown as an example of the three MSS stations in that region. Widths of the bars indicate the total time-series length. Bar colour indicates spatial region (see **Figure 1**); blue: Celtic Seas; red: northern North Sea; green: southern North Sea; light yellow: Channel.

4 Plankton lifeforms

The PLET uses a trait look-up table to aggregate plankton taxa into lifeforms. The lifeforms have been pre-defined using biological traits to represent groups of plankton which perform similar ecological functional roles (McQuatters-Gollop et al., 2019; Scherer et al., 2014). Details of each lifeform, and lifeform pairings with ecological relevance for assessment, are given in **Table 2**. It should be noted that these traits have been developed for marine taxa only (see list of included taxon groups in UK Pelagic Habitats Expert Group, (2020); <https://www.dassh.ac.uk/doitool/data/1709>), and traits in **Table A1**), with the goal of simplifying plankton datasets for use in assessments; they are not intended as a fully comprehensive list of plankton traits.

The trait look-up table (Plankton Lifeform Traits Master List (UK Pelagic Habitats Expert Group, 2020) was developed by using a combination of extensive literature synthesis and expert opinion. The World Register of Marine Species (WoRMS Editorial Board, 2020) Aphia IDs are used to link the taxa to their associated traits. Confidence in the lifeforms extracted is assigned based on a combination of the ability to identify each of the taxa it comprises reliably by light microscopy and the ability to assign traits to each taxon (**Table 3**). The confidence associated with each lifeform is described in **Table 4**. Only lifeforms with a ‘high’ confidence rating are provided in the PLET outputs. In some cases, confidence assignment reflects the limitations of identification by light microscopy by nature of the datasets around which the table was developed. Similarly, the size-based lifeforms (as currently defined and used) reflect the size limitations of sampling and identification of the currently included datasets. For example, the lifeform ‘small phytoplankton’, defined as phytoplankton with size < 20 µm, is more appropriately termed ‘small micro-phytoplankton’ because while the lower size limit of identification by routine light microscopy will include some large nanophytoplankton, it excludes pico-plankton and the smaller nano- phytoplankton.

410

Table 2. List of lifeforms and their ecological importance. The definitions of the lifeforms (see also McQuatters-Gollop et al. (2019) based on the trait lookup table are given in **Table B1**.

Lifeform	Definition ¹	Ecological Importance
(micro)Phytoplankton (size range determined by possible enumeration by light microscopy)	Protista taxa that contribute to primary production	Encompasses key primary producers, with notable exclusion of pico-, small nano- and microphytoplankton. Important for food web support, dynamics and biogeochemical cycling.
Large microphytoplankton	≥ 20 µm individual cell diameter	Changes in relative abundance provides a size-based indicator of the efficiency of energy flow to higher trophic levels. (Schmidt <i>et al.</i> , 2020).
Small microphytoplankton	< 20 µm individual cell diameter, with lower size limit determined by current enumeration by light microscopy.	
Diatoms	Taxa of the class Bacillariophyceae	Key groups of primary producers. Changes in abundance, and relative abundance in particular, are used to monitor changes in ecosystem functioning (Hinder et al., 2012; Wasmund et al., 2017). Dominance of dinoflagellates over diatoms may be an indicator of eutrophication or of change in water column stability, indicating changes in nutrient concentration or stratification (Devlin et al., 2009; Wasmund et al., 2017). When dinoflagellates are mainly heterotrophic, then they can account for a significant part of diatom grazing (as in the Eastern English Channel; Grattepanche et al. (2011)).
Dinoflagellates	Taxa of the phylum Dinoflagellata	
Autotrophic and mixotrophic dinoflagellates	Autotrophic: nutrition by photosynthesis; Mixotrophic: capable of obtaining nourishment via photo(auto)trophy and phago(hetero)trophy, as well as via osmo(hetero)trophy (see Flynn <i>et al.</i> , 2019)	Shift in primary producers may indicate eutrophication (Gowen et al., 2012).
Pelagic diatoms	Diatoms living in the water column.	Changes in relative abundance provides an indicator of benthic disturbance and frequency of resuspension events (Cibic et al., 2012).
Tychopelagic diatoms	Benthic diatoms which can become mixed into the water column.	
Potentially toxic or nuisance diatoms	Diatoms and dinoflagellates which are either ‘toxic’, defined as <i>capable of producing toxins which can cause illness or death in humans, animals and/or fish</i> , or ‘nuisance’ defined as taxa producing effects which are detrimental to aquaculture and benthos via physical harm or causing anoxia or produce water discolorations, scums or foams that can be aesthetically, socially, or economically negative.	These groups include species which have the potential for negative impacts on human health and provision of ecosystem services for people as well as other higher trophic levels of the system (Hallegraeff et al., 2021; Wells et al., 2020).
Potentially toxic or nuisance dinoflagellates		
Ciliates	Protozoans characterized by the presence of cilia.	Increases in abundance could indicate a shift from primarily autotrophic to a more heterotrophic system (Scherer, 2012).
Holoplankton	Zooplankton taxa which spend their entire lifecycle in the plankton.	Changes in relative abundance provides an indicator of change in pelagic-benthic food web structure (Bedford et al., 2020a; Kirby et al., 2008).

Lifeform	Definition¹	Ecological Importance
Meroplankton	Taxa which spend part of their lifecycle as zooplankton.	
Gelatinous zooplankton	Taxa of the phyla Cnidaria and Ctenophora only.	Changes in relative abundance indicate potential alternative energy flows through the food web of varying importance to fisheries, aquaculture, tourism etc. (Richardson <i>et al.</i> , 2009).
Fish larvae and fish eggs	Includes fish eggs as well as larvae.	
Carnivorous zooplankton	Taxa which prey mainly on other zooplankton.	Non-carnivorous functionally refers to zooplankton that could be grazers on phytoplankton, at some point in their lifecycle. Changes in relative abundance of carnivorous and non-carnivorous zooplankton indicates a shift in energy flow and balance between primary consumers and secondary consumers.
Non-carnivorous zooplankton	Zooplankton with less-carnivorous feeding mechanisms, i.e.: predominately suspension or filter feeders, omnivores which can use both carnivorous and herbivorous feeding, or ambiguous (diet uncertain).	
Crustaceans	Taxa of the Subphylum Crustacea	Crustaceans are important for commercial fisheries, either directly or in food chains that fuel them. Changes in crustacean zooplankton abundance can reflect both bottom-up and top-down controls and may indicate changes in food availability for exploited fish stocks (Capuzzo <i>et al.</i> , 2018).
Large copepod species (≥ 2 mm)	≥ 2 mm adult total body length	Changes in relative abundance provide a size-based indicator of food web structure and energy flow (Daufresne <i>et al.</i> , 2009).
Small copepod species (< 2 mm)	< 2 mm adult total body length	

¹ Modified from McQuatters-Gollop *et al.* (2019)

Table 3. Lifeform confidence assignment matrix, where ‘High’, ‘Medium’, or ‘Low’ are based on the ability to identify and assign traits for the constituent taxa groups of a lifeform.

	Can assign traits to constituent taxa	Cannot assign traits to constituent taxa
Can identify constituent taxa	high	medium
Cannot identify constituent taxa	medium	low

Table 4. Lifeform confidences based on ability to identify and assign traits, applying rationale in **Table 3**. Only lifeforms with a ‘high’ confidence rating are provided in the PLET outputs.

Lifeform	Confidence	Reason for confidence (where not ‘high’)
(micro)Phytoplankton	High	
Large microphytoplankton	Medium	Can reliably identify individual plankton species size class but cannot always reliably assign the size trait if the group counted spans taxa that are both larger and smaller than 20 microns.
Small microphytoplankton	Medium	
Diatoms	High	
Dinoflagellates	High	
Autotrophic and mixotrophic dinoflagellates	Medium	Can identify taxa, but assigning feeding mechanism trait is not always clear (see discussion in Flynn et al. (2019))
Pelagic diatoms	High	
Tycho pelagic diatoms	High	
Potentially toxic and nuisance diatoms	Low	Designation of some algal blooms as “harmful” (i.e.: Harmful Algal Blooms, ‘HABS’), relates more to societal assessment than plankton traits, these ‘lifeforms’ are therefore not currently recommended for use though they are defined in the traits list and will be the focus of future development work. Specific issues include: <ul style="list-style-type: none"> The toxin producing diatom genus <i>Pseudo-nitzschia</i> contains both amnesic shellfish toxin-producers which can render shellfish unfit for human consumption and potentially negatively impact the health of marine mammals, and non toxin-producing species/individuals. It is not possible to identify these cells to species level using routine light microscopy; some toxin and non-toxin producing species are morphologically identical.
Potentially toxic and nuisance dinoflagellates	Low	<ul style="list-style-type: none"> The genus <i>Alexandrium</i> contains both paralytic shellfish toxin- and non-toxin-producing species/strains and it is not possible to distinguish these using routine light microscopy; some toxin and non-toxin producing species are morphologically identical. Currently it is unknown if the negative impact from <i>Karenia mikimotoi</i> in European waters is via toxin production or anoxia arising from high biomass blooms. Not all datasets included in PLET reliably record key species (e.g.: CPR does not record <i>Alexandrium</i>)
Ciliates	Low	<ul style="list-style-type: none"> Ecological function can be duplicated by heterotrophic and mixotrophic dinoflagellates. Ciliates do not preserve well in the standard 0.5% Lugol’s iodine preservative used to preserve phytoplankton samples and some (but not all) are too small and too fragile to be well sampled by many of the datasets currently in PLET.
Holoplankton	Medium	<ul style="list-style-type: none"> May not identify taxa specifically enough to determine traits. Some of the rarer species are resuspended from the seabed and definition of their holo- or meroplanktonic status is difficult
Meroplankton	Medium	
Gelatinous zooplankton	High	
Fish larvae	High	
Carnivorous zooplankton	Medium	Can identify taxa, but assigning diet trait is unclear, especially at different life stages.
Non-carnivorous zooplankton	Medium	
Crustaceans	High	
Small copepods	High	
Large copepods	High	

5 Plankton lifeform extraction tool functionality

The PLET is accessed through a web-based user-interface (see **Figure 4**). To generate custom plankton lifeform outputs, users select: time-series start and end dates, a spatial area, and a data set. Because of methodological differences in sampling and analysis it is not appropriate to produce average lifeforms across the multiple datasets, as such all sample locations within the selected spatial area for any data set are aggregated into a single lifeform data product but stations from different datasets are never aggregated. The resulting data product, monthly averaged aggregated lifeform abundance, is generated either within the web-browser, or for download in *.csv* or *.json* format. The output data includes the number of individual samples from which each monthly average was derived, as well as a list of component taxon groupings. Blank output component taxon groupings indicate that the originally submitted sample data did not include information in the (optional) 'Taxon Name' field.

The PLET also has a simple API (Application Programming Interface), which provides the option of bypassing the webpage interface and sending queries to the tool using the URL only. The base URL is 'https://www.dassh.ac.uk/lifeforms/cgi-bin/get_form.py'. The parameters are: *startdate* (YYYY-MM-DD), *enddate* (YYYY-MM-DD), *north* (northern edge of bounding box, in decimal degrees), *south* (southern edge of bounding box, in decimal degrees), *east* (eastern edge of bounding box, in decimal degrees), *west* (western edge of bounding box, in decimal degrees), *dataset* (currently: CPR, L4_phyto, L4_zoo, SMHI, CEFAS_SmartBuoy, EA_PHYTO_2000-2017, MBA_E1_L5, MSS_phyto, MSS_zoo, SEPA_Zooplankton, SEPA_Phytoplankton or SAMS-LPO), *format* (*csv*, *json* or *pretty*). For example, to retrieve results from the CPR dataset for May 1975 between 50 and 60 degrees of Latitude and -5 and 5 degrees of Longitude, and return in CSV format, the URL request is:

```
https://www.dassh.ac.uk/lifeforms/cgi-bin/get_form.py?startdate=1975-05-01&enddate=1975-05-30&north=60&south=50&east=5&west=-5&dataset=CPR&format=CSV. Sending such URL commands via Curl or similar tools allows the PLET to be used programmatically if desired.
```

There are a number of options for defining the spatial domain of the lifeform data product. A rectangular extent can be manually defined by the northern, southern, western, and eastern edges of a

rectangular bounding box, by simply drawing a rectangle on the interactive map display which shows sample locations for each data set. Similarly, a polygon shaped extent can be manually defined in ‘well-known text’ (WKT) format or through the interactive map. A query specifying spatial extent by polygon
450 instead of bounding box can be constructed for the API by designating the parameter *wkt* instead of *north*, *south*, *east* and *west*. A more complex area, for example a formal assessment region, can be used by uploading a shapefile to the tool.

All integrated datasets are listed within the web interface, with full metadata. The trait look-up table can also be accessed and downloaded. To facilitate submission of new and updated plankton data, templates
455 for data submission are also provided.

DASSH Home Search data Submit data **Plankton Lifeform Extraction Tool**

Plankton Lifeform Extraction Tool

Fill out the form below to generate monthly lifeform abundances from the selected dataset.

Click here to download the current master species list, a data submission template, or the DASSH data agreement:

Start Date
End Date

Select all dates

Area of Interest

Enter the North, South, West and East edges of a rectangular bounding box (values in decimal degrees)

OR upload a **zipped shapefile with your bounding polygon**

No file chosen

OR draw a box or polygon on the map below (using the square or pentagon icons, respectively).

OR manually define a polygon in WKT format

Select all latitudes and longitudes

Dataset

Output Format

Figure 4. The Plankton Lifeform Extraction Tool. Screenshot of the Plankton Lifeform Extraction Tool (v1).

6 Lifeform outputs

The spatial and temporal patterns of plankton lifeforms, based on the data currently held in PLET, are summarised to highlight seasonal patterns in **Figure 5** (phytoplankton) and **Figure 6** (zooplankton), and inter-annual patterns in **Figure 7** and **Figure 8**. In order to facilitate visualisation across the different lifeforms and datasets, where absolute lifeform abundances are extremely variable, within-lifeform and dataset changes are shown as standardised z-scores that indicate the difference from the overall time-series mean values (Glover et al., 2005).


Plankton abundance peaks in spring and summer are associated with nearly all plankton lifeforms, across all datasets. However, the timing, duration, and intensity of these peaks differ between lifeforms and datasets (see **Figure 5** and **Figure 6**), in some cases partly because of spatial aggregation of the data. In the CPR dataset, which samples furthest offshore, seasonal zooplankton lifeform abundance peaks last longer than those of the phytoplankton lifeforms. The EA datasets, which represent estuarine and coastal waters, include much shorter seasonal phytoplankton lifeform abundance peaks than the CPR for the corresponding regions, and differences in bloom timing are also evident, highlighting the small-scale spatial variability in plankton abundance. This is further evidenced in the comparison between seasonal patterns in PML's L4 station in the Channel and both the EA and CPR data aggregation for the same larger region. This heterogeneity demonstrates the added value of integrating datasets to achieve a representative description of plankton community seasonal succession within even a relatively localised sea area, particularly where different programmes sample different subsets of the plankton community (in this case the fragile dinoflagellates being less well preserved by CPR compared to PML and EA sampling). Comparing across larger spatial scales, differences in seasonal patterns between English and Scottish waters are likely influenced by the latitudinal gradients (Fanjul et al., 2017, 2019; Uriarte et al., 2021) as well as local hydrographic conditions (e.g.: Atlantic inflow). The Swedish stations, located in the Kattegat, show the most divergent lifeform seasonality compared to the other datasets, notably in the timing of abundance peaks (e.g.: the absence of April plankton blooms) which likely reflect their distinct oceanographic setting and the influence of Baltic Sea outflow waters.

Interannual trends in lifeform abundance can be related to changes in pressures within the marine system
485 (Bedford et al. (2020a) and McQuatters-Gollop et al. (2019)). Given the strong seasonal variability,
summarising plankton abundance to compare across years is non-trivial. Representative data coverage,
typically at least monthly is needed to ensure that inter-annual differences are not due to missing samples.
For example, the WFD eutrophication assessment procedure (Greenwood (2019)) requires phytoplankton
data for at least 9 months of every year assessed. The Plankton Lifeform Index (Tett et al., 2007, 2008),
490 by looking at changes from a reference envelope defined by 3-5 years of adequate data (i.e.: at least
monthly sampling) is robust against missing samples (months) so long as these are not biased to particular
times of the year. The Pelagic Habitat Expert Group has recommended at least monthly sampling to
adequately take account of seasonal changes in the balance of plankton lifeforms, while noting that higher
temporal resolution would provide greater confidence that all transient bloom events (which may last less
495 than a month) were observed. Given the tool's robustness against data loss, annual assessments can be
made reliably when one to three months have been lost, so long as there is no persistent bias in lost months
over several years.


Despite missing months being an important consideration for annual aggregation (in some datasets in
particular, see (**Figure 4**) the interannual trends in phytoplankton (**Figure 7**) and zooplankton (**Figure 8**)
500 lifeforms show considerable changes in lifeform abundance among years in all datasets and regions. The
longest time-series (MBA L5&E1 since 1924, CPR since 1960 and SAMS phytoplankton since 1970)
capture decreases over several years followed by subsequent increases, which caution against over-
interpretation of the shorter time-series. For example, there have been decreases in all zooplankton
lifeforms at the MSS Celtic Seas station since 2013 that cannot be seen in the nearby SEPA Celtic Seas
505 station which only has observations from 2014 onwards; while both in the Celtic Seas area, these two
sites are characterised by very different hydrographical settings. The importance of considering both
short- and long-term changes in plankton lifeforms is discussed in detail in (Bedford et al., 2020b).

Bringing diverse datasets together to extend both spatial and temporal coverage is a key tool for
distinguishing small-scale, short-term fluctuations from larger-scale longer-term changes. For example,
510 Bedford *et al.*, 2020a identified regional-scale trends in lifeform changes (increasing diatom abundance
in the northern North Sea and increasing mesozooplankton abundance across almost the whole North-


West European shelf) using timeseries data from 5 different UK plankton surveys and linked some of these changes to changing sea surface temperature. Assessment of the status of the marine pelagic habitat (McQuatters-Gollop et al., 2019) requires linking changes to pressures (Scherer et al., 2016), which relies
515 on high temporal and spatial resolution good quality observations, such as climate (Bedford *et al.*, 2020a) and eutrophication (Gowen et al., 2015; Greenwood, 2019).



520 **Figure 5. Phytoplankton lifeform monthly means, by data provider and region.** Colour indicates lifeform abundance relative to the long-term mean of each lifeform within each region and dataset as standardised z-score (Glover et al., 2005); Scores of zero are equal to the long-term mean, positive scores (in green/yellow) signify values above the long-term mean and negative scores values below the long-term mean (in blue). Only those lifeforms that have been assigned a confidence level of ‘high’ are shown (see Table 2 and Table 3). Regions are defined in Figure 1.



525 **Figure 6. Zooplankton lifeform monthly means, by data provider and region.** Colour indicates lifeform abundance relative to the long-term mean of each lifeform within each region and dataset as standardised z-score (Glover et al., 2005): Scores of zero are equal to the long-term mean, positive scores (in green/yellow) signify values above the long-term mean and negative scores values below the long-term mean (in blue). Only those lifeforms that have been assigned a confidence level of ‘high’ are shown (see **Table 2** and **Table 3**). Regions are defined in **Figure 1**.



530

Figure 7. Phytoplankton lifeform annual means, by data provider and region. Colour indicates lifeform abundance relative to the long-term mean of each lifeform within each region and dataset as standardised z-score (Glover et al., 2005): Scores of zero are equal to the long-term mean, positive scores (in green/yellow) signify values above the long-term mean and negative scores values below the long-term mean (in blue). Only those lifeforms that have been assigned a confidence level of ‘high’ are shown (see **Table 2** and **Table 3**). Regions are defined in **Figure 1**.

535




Figure 8. Zooplankton lifeform annual means, by data provider and region. Colour indicates lifeform abundance relative to the long-term mean of each lifeform within each region and dataset as standardised z-score (Glover et al., 2005): Scores of zero are equal to the long-term mean, positive scores (in green/yellow) signify values above the long-term mean and negative scores values below the long-term mean (in blue). Only those lifeforms that have been assigned a confidence level of ‘high’ are shown (see **Table 2** and **Table 3**). Regions are defined in **Figure 1**.

540

7 Discussion

Large-scale trends in the abundance of individual species are challenging to compare across multiple
545 time-series due to difficulties in sampling and in counting at the species level, particularly at the limits of
the geographic range of a species. Conversely, trends in bulk indices such as total zooplankton biomass
and abundance or total chlorophyll *a* concentration can miss important underlying details. Our method to
aggregate at the level of functional groups (lifeforms) provides a tractable approach to reveal meaningful
information at an intermediate level of organisation that is still ecologically relevant.

550 While the main available UK plankton time series are included here as well as Swedish data from the
south eastern North Sea, further extending the geographical and temporal extent of plankton time-series
held in the PLET will improve robustness of evidence underpinning pelagic habitat assessment. As a
transparent and accessible source of diverse plankton datasets, the PLET also facilitates exploration of
associated research questions in an integrated way. Observations of changes in size-based plankton
555 lifeform abundances (Greenwood *et al.*, 2019; McQuatters-Gollop *et al.*, 2019, Bedford *et al.*, 2020),
alongside methodological developments in measuring the complete plankton size spectrum (Atkinson *et al.*,
2021), provide an improved ability to understand what is driving changes in plankton size and species
composition across the full spatial-temporal scales of the component datasets (Schmidt *et al.*, 2020). Data
from different data sources are not aggregated within PLET in order to maintain the scientific robustness
560 of the outputs, and incorporate a range of methodologies (units, scale, fixed-point, transect etc.). The data
is outputted in a unified way from PLET to encourage comparison and interpretation of the changes in
the plankton, if the user wishes to combine the outputs within a specified area and time period a
normalisation technique can be applied, however care needs to be taken to ensure compatibility and
coverage does not bias the combined results. For example, due to the differing taxonomic resolution and
565 sampling methods across the various component time series, we do not recommend simple comparisons
of indices of species richness or diversity. There is also the flexibility to improve confidence in low
confidence groupings, and to potentially incorporate new types of plankton data into the tool in the future,
such as the use of flow cytometry data.

Time-series datasets are critical for identifying and assessing changes in the marine environment. Given
570 the expense and effort which goes into producing and maintaining these invaluable datasets, tools that
make them more widely available, transparent, and accessible to the broader user community are needed.
The PLET provides a centralised, easily accessible, source for version-controlled time-series data and
metadata and is an essential component of a robust assessment process as well as a tool to support the
research, which is needed to underpin assessment. This includes exploring new ecologically relevant
575 lifeform groupings and improving the understanding of lifeforms currently designated as medium and
low confidence, and which will in turn feed back into the process of assessing the health of the marine
environment.

The PLET is not a static resource; it is designed to readily accept additional datasets and be updated to
support future assessments as the assessment procedure continues to evolve. This is a critical step towards
580 using multiple datasets collected with diverse methods to populate and assess a common indicator,
allowing the assessment of pelagic ecosystems at regional scales. As the tool is expanded with additional
datasets, its ability to detect change in plankton communities will increase, and the policy evidence it
provides will continue to become more robust, providing decision makers with critical information to
inform management measures. As pelagic habitat assessments continue to improve and adapt to the
585 changing policy landscape (Boyes and Elliott, 2016) and to evolving plankton data availability, the
PLET's flexibility will allow it to continue to underpin assessment.

8 Data Availability and Citation

The Plankton Lifeform Extraction Tool is hosted by the Archive for Marine Species and Habitats Data
(DASSH), which is accredited as the UK Node of the Ocean Biodiversity Information System (OBIS)
590 and through the Marine Environmental Data and Information Network (MEDIN), the UK partnership of
organisations committed to improving access to UK marine data, and core-funded by the Department for
the Environment, Food and Rural Affairs (Defra) and the Scottish Government. Lifeform data products
can be generated at: <https://www.dassh.ac.uk/lifeforms/>.

The PLET's lifeform data products are generated by applying the Plankton Lifeform Traits Master List
595 trait look-up table (UK Pelagic Habitats Expert Group, 2020; <https://www.dassh.ac.uk/doitool/data/1709>) to
standardised-format versions of the integrated plankton datasets (see **Table 1** for details (CPR and Johns,
2019; MBA, 2019; PML, 2019; CEFAS, 2019; EA, 2019; MSS, 2019; SMHI, 2019; SEPA, 2020; SAMS,
2020)). These time-series may be updated in the future to include ongoing plankton monitoring, and more
datasets may be added. Versions of several of these datasets are also available through other data
600 repositories (e.g.: institute-specific websites provided in **Table 1**, or the Ocean Biodiversity Information
System: OBIS). However, PLET provides the first centralised database for all time-series feeding into
UK Marine Strategy and OSPAR biodiversity lifeform-based assessments, and importantly: a format and
structure compatible with extraction of lifeform time-series. By attaching doi's to the underlying dataset
versions held within PLET and the traits list, the tool provides full transparency and reproducibility for
605 the generated lifeform outputs. Users of these datasets are encouraged to appropriately cite the data
sources by means of their doi's as well as this data paper, so that usage can be more easily traced. Doing
so provides evidence of data uptake, enhancing the possibility of continued funding for valuable time
series and for plankton indicator development.

610

9 Acknowledgements

We are greatly indebted to the crews and scientists for collecting the component samples over the last 100 years, to the large number of taxonomic analysts for processing the samples, and to those individuals who strived to keep plankton time-series going under difficult circumstances. Funding that supports this work and the data collected has come from the European Union (EU) Grant No. 11.0661/2015/712630/SUB/ENVC.2 OSPAR, UK Natural Environment Research Council, Grant/Award Number: NE/R002738/1 and NE/M007855/1; EMFF; Climate Linked Atlantic Sector Science, Grant/Award Number: NE/ R015953/1, DEFRA UK ME-5308 and ME-414135, NSF USA OCE-1657887, DFO CA F5955- 150026/001/HAL, NERC UK NC-R8/H12/100, Horizon 2020: 862428 Atlantic Mission, iCPR SBFF-2019-36526, IMR Norway, DTU Aqua Denmark and the French Ministry of Environment, Energy, and the Sea (MEEM). Recent funding for the development of PLET and the Pelagic Habitats Indicator has been provided by: HBDSEG/Defra, and MMO/EMFF. The MSS Scottish Coastal Observatory data and analyses are funded and maintained by the Scottish Government Schedules of Service ST05a and ST02H, MSS Stonehaven Samplers, North Atlantic Fisheries College, Shetland, Orkney Islands Harbour Council, and Isle Ewe Shellfish.

10 Author contribution

AMG co-ordinated the project. All authors developed the Plankton lifeform traits master list and provided input on the lifeform traits and groupings. CO and GG designed the Plankton Lifeform Extraction Tool (PLET) and wrote the initial code. KP developed the PLET and performed the maintenance of the site, with support from DL and DB. DGJ, CO, AA, CW, RB, MB, MD, CG, BB, EB, KC, MM, MJ, and PT provided datasets. CO, CG and AMG prepared the manuscript with contributions from all co-authors.

11 Competing interests

The authors declare that they have no conflict of interest.

635

12 References

- Atkinson, A., Harmer, R. A., Widdicombe, C. E., McEvoy, A. J., Smyth, T. J., Cummings, D. G., Somerfield, P. J., Maud, J. L. and McConville, K.: Questioning the role of phenology shifts and trophic mismatching in a planktonic food web, *Prog. Oceanogr.*, (MAY), doi:10.1016/j.pocean.2015.04.023, 2015.
- 640 Atkinson, A., Lilley, M. K. S., Hirst, A. G., McEvoy, A. J., Tarran, G. A., Widdicombe, C., Fileman, E. S., Woodward, E. M. S., Schmidt, K., Smyth, T. J. and Somerfield, P. J.: Increasing nutrient stress reduces the efficiency of energy transfer through planktonic size spectra, *Limnol. Oceanogr.*, 66(2), 422–437, doi:10.1002/lno.11613, 2021.
- 645 Beaugrand, G., Reid, P. C., Ibañez, F., Lindley, J. A. and Edwards, M.: Reorganization of North Atlantic marine copepod biodiversity and climate. *Supplementary Material, Science*, 296(5573), 1692–4, doi:10.1126/science.1071329, 2002.
- Beaugrand, G., Edwards, M. and Legendre, L.: Marine biodiversity, ecosystem functioning, and carbon cycles, *Proc. Natl. Acad. Sci. U. S. A.*, 107(22), 10120–10124, doi:10.1073/pnas.0913855107, 2010.
- 650 Bedford, J., Ostle, C., Johns, D. G., Atkinson, A., Best, M., Bresnan, E., Machairopoulou, M., Graves, C. A., Devlin, M., Milligan, A., Pitois, S., Mellor, A., Tett, P. and Mcquatters-gollop, A.: Lifeform indicators reveal large-scale shifts in plankton across the North-West European shelf, *Glob. Chang. Biol.*, (December 2019), 1–16, doi:10.1111/gcb.15066, 2020a.
- Bedford, J., Ostle, C., Johns, D. G., Budria, A. and Mcquatters-gollop, A.: The influence of temporal scale selection on pelagic habitat biodiversity indicators, *Ecol. Indic.*, 114(March), 106311, doi:10.1016/j.ecolind.2020.106311, 2020b.
- 655 Boyes, S. J. and Elliott, M.: Brexit: The marine governance horrendogram just got more horrendous!, *Mar. Pollut. Bull.*, 111(1–2), 41–44, doi:10.1016/j.marpolbul.2016.08.020, 2016.
- Bresnan, E., Cook, K., Hindson, J., Hughes, S., Lacaze, J.-P., Walsham, P., Webster, L. and Turrell W., R.: The Scottish Coastal Observatory 1997-2013, Part 2: Description of Scotland’s Coastal Waters, *Scottish Mar. Freshw. Sci. Rep.*, Vol 7(No 26), 278 pp., 2016.
- 660 Capuzzo, E., Lynam, C. P., Barry, J., Stephens, D., Forster, R. M., Greenwood, N., McQuatters-Gollop, A., Silva, T., van Leeuwen, S. M. and Engelhard, G. H.: A decline in primary production in the North Sea over 25 years, associated with reductions in zooplankton abundance and fish stock recruitment, *Glob. Chang. Biol.*, 24(1), e352–e364, doi:10.1111/gcb.13916, 2018.
- 665 CEFAS: 2000 to date North, Irish & Celtic Seas Centre for Environment, Fisheries and Aquaculture Science Smartbuoy Marine Monitoring. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., doi:https://doi.org/10.17031/1634, 2019.
- Chiba, S., Batten, S., Martin, C. S., Ivory, S., Miloslavich, P. and Weatherdon, L. V: Zooplankton monitoring to contribute towards addressing global biodiversity conservation challenges, *J. Plankton Res.*, 00(00), 1–10, doi:10.1093/plankt/fby030, 2018.
- 670 Cibic, T., Comici, C., Bussani, A. and Del Negro, P.: Benthic diatom response to changing environmental conditions, *Estuar. Coast. Shelf Sci.*, 115, 158–169, doi:10.1016/j.ecss.2012.03.033, 2012.
- CPR and Johns, D. G.: Continuous Plankton Recorder Survey (CPR Survey). The Marine Biological Association of the UK, CPR Survey. (Dataset)., doi:https://doi.org/10.17031/1629, 2019.
- 675 Daufresne, M., Lengfellner, K. and Sommer, U.: Global warming benefits the small in aquatic

- ecosystems, *Proc. Natl. Acad. Sci. U. S. A.*, 106(31), 12788–12793, doi:10.1073/pnas.0902080106, 2009.
- 680 Devlin, M., Barry, J., Painting, S. and Best, M.: Extending the phytoplankton tool kit for the UK Water Framework Directive: Indicators of phytoplankton community structure, *Hydrobiologia*, 633(1), 151–168, doi:10.1007/s10750-009-9879-5, 2009.
- Devlin, M., Best, M., Bresnan, E., Scanlan, C. and Baptie, M.: *Water Framework Directive: The development and status of phytoplankton tools for ecological assessment of coastal and transitional waters.*, 2012.
- EA: *Phytoplankton Dataset provided by the Environment Agency (EA) 2000-2017 . The Archive for Marine Species and Habitats Data (DASSH).* (Dataset)., , doi:https://doi.org/10.17031/1635, 2019.
- 685 Edwards, K. F., Klausmeier, C. A. and Litchman, E.: Nutrient utilization traits of phytoplankton, *Ecology*, 96(8), 2311–2311, doi:10.1890/14-2252.1, 2015.
- Edwards, M. and Richardson, A. J. A. J.: Impact of climate change on marine pelagic phenology and trophic mismatch, *Nature*, 430, 881–884, doi:10.1038/nature02808, 2004.
- 690 EU: Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy., *Off. J. Eur. Commun.*, 327, 1–72, 2000.
- Falkowski, P.: Ocean Science: The power of plankton, *Nature*, 483(7387), S17–S20, doi:10.1038/483S17a, 2012.
- Fanjul, A., Villate, F., Uriarte, I., Iriarte, A., Atkinson, A. and Cook, K.: Zooplankton variability at four 695 monitoring sites of the Northeast Atlantic Shelves differing in latitude and trophic status, *J. Plankton Res.*, 39(6), 891–909, doi:10.1093/plankt/fbx054, 2017.
- Fanjul, A., Iriarte, A., Villate, F., Uriarte, I., Artiach, M., Atkinson, A. and Cook, K.: Latitude, distance offshore and local environmental features as modulators of zooplankton assemblages across the NE Atlantic Shelves Province, *J. Plankton Res.*, 41(3), 293–308, doi:10.1093/plankt/fbz015, 2019.
- 700 Flynn, K. J., Stoecker, D. K., Mitra, A., Raven, J. A., Glibert, P. M., Hansen, P. J., Granéli, E. and Burkholder, J. M.: Misuse of the phytoplankton-zooplankton dichotomy: The need to assign organisms as mixotrophs within plankton functional types, *J. Plankton Res.*, 35(1), 3–11, doi:10.1093/plankt/fbs062, 2013.
- Flynn, K. J., Mitra, A., Anestis, K., Anschütz, A. A., Calbet, A., Ferreira, G. D., Gypens, N., Hansen, P. 705 J., John, U., Martin, J. L., Mansour, J. S., Maselli, M., Medić, N., Norlin, A., Not, F., Pitta, P., Romano, F., Saiz, E., Schneider, L. K., Stolte, W. and Traboni, C.: Mixotrophic protists and a new paradigm for marine ecology: Where does plankton research go now?, *J. Plankton Res.*, 41(4), 375–391, doi:10.1093/plankt/fbz026, 2019.
- Glover, D., Jenkins, W. and Doney, S.: *Modeling methods for marine science*, Cambridge University 710 Press., 2005.
- Gowen, R. J., Mcquatters-gollop, A., Tett, P., Best, M., Bresnan, E., Castellani, C., Cook, K., Forster, R. M., Scherer, C. and Mckinney, A.: The Development of UK Pelagic (Plankton) Indicators and Targets for the MSFD, *Ecosystems*, (June 2011), 1–41, doi:10.13140/RG.2.1.5181.1920, 2011.
- Gowen, R. J., Tett, P. and Smayda, T. J.: Phytoplankton and the balance of nature: An opinion, *Estuar. 715 Coast. Shelf Sci.*, 113, 317–323, doi:10.1016/j.ecss.2012.08.009, 2012.
- Gowen, R. J., Collos, Y., Tett, P., Scherer, C., Bec, B., Abadie, E., Allen, M. and O’Brien, T.: Response of diatom and dinoflagellate lifeforms to reduced phosphorus loading: A case study in the Thau lagoon, France, *Estuar. Coast. Shelf Sci.*, 162, 45–52, doi:10.1016/j.ecss.2015.03.033, 2015.

- Grattepanche, J. D., Breton, E., Brylinski, J. M., Lecuyer, E. and Christaki, U.: Succession of primary producers and micrograzers in a coastal ecosystem dominated by *Phaeocystis globosa* blooms, *J. Plankton Res.*, 33(1), 37–50, doi:10.1093/plankt/fbq097, 2011.
- Greenwood, N.: Utilizing eutrophication assessment directives from transitional to marine systems in the Thames estuary and Liverpool Bay, UK, *Front. Mar. Sci.*, doi:10.3389/fmars.2019.00116, 2019.
- Hallegraeff, G., Enevoldsen, H. and Zingone, A.: Global harmful algal bloom status reporting, *Harmful Algae*, 102, doi:10.1016/j.hal.2021.101992, 2021.
- Hinder, S. L., Hays, G. C., Edwards, M., Roberts, E. C., Walne, A. W. and Gravenor, M. B.: Changes in marine dinoflagellate and diatom abundance under climate change, *Nat. Clim. Chang.*, 2(4), 271–275, doi:10.1038/nclimate1388, 2012.
- Kirby, R. R., Beaugrand, G. and Lindley, J. A.: Climate-induced effects on the meroplankton and the benthic-pelagic ecology of the North Sea, *Limnol. Oceanogr.*, 53(5), 1805–1815, doi:10.4319/lo.2008.53.5.1805, 2008.
- Klais, R., Norros, V., Lehtinen, S., Tamminen, T. and Olli, K.: Community assembly and drivers of phytoplankton functional structure, *Funct. Ecol.*, 31(3), 760–767, doi:10.1111/1365-2435.12784, 2017.
- Kulk, G., Platt, T., Dingle, J., Jackson, T., Jönsson, B. F., Bouman, H. A., Babin, M., Brewin, R. J. W., Doblin, M., Estrada, M., Figueiras, F. G., Furuya, K., González-Benítez, N., Gudfinnsson, H. G., Gudmundsson, K., Huang, B., Isada, T., Kovač, Ž., Lutz, V. A., Marañón, E., Raman, M., Richardson, K., Rozema, P. D., van de Poll, W. H., Segura, V., Tilstone, G. H., Uitz, J., van Dongen-Vogels, V., Yoshikawa, T. and Sathyendranath, S.: Primary production, an index of climate change in the ocean: Satellite-based estimates over two decades, *Remote Sens.*, 12(5), doi:10.3390/rs12050826, 2020.
- Laplace-Treyture, C., Derot, J., Prévost, E., Le Mat, A. and Jamoneau, A.: Phytoplankton morpho-functional trait dataset from French water-bodies, *Sci. Data*, 8(1), 1–9, doi:10.1038/s41597-021-00814-0, 2021.
- Margalef, R.: Life-forms of phytoplankton as survival alternatives in an unstable environment, *Oceanol. Acta*, 1, 493–509, 1978.
- MBA: 1924-2013 MBA L4 and E1 Young Fish Survey. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., , doi:https://doi.org/10.17031/1636, 2019.
- McQuatters-Gollop, A., Johns, D. G., Bresnan, E., Skinner, J., Rombouts, I., Stern, R., Aubert, A., Johansen, M., Bedford, J. and Knights, A.: From microscope to management: The critical value of plankton taxonomy to marine policy and biodiversity conservation, *Mar. Policy*, 83, 1–10, doi:10.1016/j.marpol.2017.05.022, 2017.
- McQuatters-Gollop, A., Atkinson, A., Aubert, A., Bedford, J., Best, M., Bresnan, E., Cook, K., Devlin, M., Gowen, R. and Johns, D. G.: Plankton lifeforms as a biodiversity indicator for regional-scale assessment of pelagic habitats for policy, *Ecol. Indic.*, 101, 913–925, 2019.
- Morishita, J.: What is the ecosystem approach for fisheries management?, *Mar. Policy*, 32(1), 19–26, doi:10.1016/j.marpol.2007.04.004, 2008.
- MSS: Phyto and zooplankton records provided by Marine Science Scotland (MSS) from Loch Ewe, Scalloway, Scapa and Stonehaven. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., , doi:https://doi.org/10.17031/1637, 2019.
- O’Brien, T. D., Lorenzoni, L., Isensee, K. and Valdés, L.: What are Marine Ecological Time Series telling us about the ocean? A status report. IOC-UNESCO, IOC, OC-UNESCO, IOC, (July), 297 [online]

- Available from: <http://igmets.net/report>, 2017.
- OSPAR: Changes in Phytoplankton and Zooplankton Communities., Intermediate Assessment 2017. [online] Available from: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/changes-phytoplankton-and-zooplankton-communities/>, 2017.
- 765 PML: 1992 - Ongoing Plymouth Marine Laboratory (PML) Zooplankton and Phytoplankton data collected at L4. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., , doi:<https://doi.org/10.17031/1632>, 2019.
- Raybaud, V., Heroin, D., Raud, T., J-m, B., Stemmann, L. and Sautour, B.: Census and analysis of zooplankton metadata of the French coasts since 1955, *J. Oceanogr. Res. Data*, 4, 11–37, 2011.
- 770 Reid, P. C., Hari, R. E., Beaugrand, G., Livingstone, D. M., Marty, C., Straile, D., Barichivich, J., Goberville, E., Adrian, R., Aono, Y., Brown, R., Foster, J., Groisman, P., Hélaouët, P., Hsu, H. H., Kirby, R., Knight, J., Kraberg, A., Li, J., Lo, T. T., Myneni, R. B., North, R. P., Pounds, J. A., Sparks, T., Stübi, R., Tian, Y., Wiltshire, K. H., Xiao, D. and Zhu, Z.: Global impacts of the 1980s regime shift, *Glob. Chang. Biol.*, 22(2), 682–703, doi:10.1111/gcb.13106, 2016.
- 775 Richardson, A. J.: In hot water: zooplankton and climate change, *Ices J. Mar. Sci.*, 65, 279–295, 2008.
- Richardson, A. J., Walne, A. W. and John, A. W. G.: Using continuous plankton recorder data, *Prog. Oceanogr.*, 68(1), 27–74, doi:10.1016/j.pocean.2005.09.011, 2006.
- Richardson, A. J., Bakun, A., Hays, G. C. and Gibbons, M. J.: The jellyfish joyride: causes, consequences and management responses to a more gelatinous future, *Trends Ecol. Evol.*, 24(6), 312–322, doi:10.1016/j.tree.2009.01.010, 2009.
- 780 SAMS: Plankton dataset provided by Scottish Association for Marine Science (SAMS) from the Lorn Pelagic Observatory (LPO) . The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., , doi:<https://doi.org/10.17031/nz24-br35>, 2020.
- Scherer, C.: Developing and testing an index of change in microplankton community structure in temperate shelf seas, Edinburgh Napier University. ISNI: 0000 0004 2721 3014., 2012.
- 785 Scherer, C., Gowen, R. J., Tett, P., Atkinson, A., Baptie, M., Best, M., Bresnan, E., Cook, K., Forster, R., Keeble, S. and Mcquatters-gollop, A.: Development of a UK Integrated Plankton Monitoring Programme - A final report of the Lifeform and State Space project Prepared for The Department of Environment , Food and Rural Affairs December., 2014.
- 790 Scherer, C., Gowen, R. J. and Tett, P.: Assessing the State of the Pelagic Habitat: A Case Study of Plankton and Its Environment in the Western Irish Sea, *Front. Mar. Sci.*, 3(November), doi:10.3389/fmars.2016.00236, 2016.
- Schmidt, K., Birchill, A. J., Atkinson, A., Brewin, R. J. W., Clark, J. R., Hickman, A. E., Johns, D. G., Lohan, M. C., Milne, A., Pardo, S., Polimene, L., Smyth, T. J., Tarran, G. A., Widdicombe, C. E., 795 Woodward, E. M. S. and Ussher, S. J.: Increasing picocyanobacteria success in shelf waters contributes to long-term food web degradation, *Glob. Chang. Biol.*, 26(10), 5574–5587, doi:10.1111/gcb.15161, 2020.
- SEPA: Photo- and Zooplankton dataset provided by Scottish Environment Protection Agency (SEPA) collected from the Forth and Clyde. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., , doi:<https://doi.org/10.17031/b84a-7951>, 2020.
- 800 SMHI: 1987-2015 Swedish Meteorological and Hydrological Institute (SMHI) Plankton Swedish West Coast. The Archive for Marine Species and Habitats Data (DASSH). (Dataset)., ,

doi:<https://doi.org/10.17031/1633>, 2019.

- Southward, A. J., Langmead, O., Hardman-Mountford, N. J., Aiken, J., Boalch, G. T., Dando, P. R.,
805 Genner, M. J., Joint, I., Kendall, M. A., Halliday, N. C., Harris, R. P., Leaper, R., Mieszkowska, N.,
Pingree, R. D., Richardson, A. J., Sims, D. W., Smith, T., Walne, A. W. and Hawkins, S. J.: Long-term
oceanographic and ecological research in the western English Channel, *Adv. Mar. Biol.*, 47, 1–105,
doi:10.1016/S0065-2881(04)47001-1, 2004.
- Tett, P., Gowen, R., Mills, D., Fernandes, T., Gilpin, L., Huxham, M., Kennington, K., Read, P., Service,
810 M., Wilkinson, M. and Malcolm, S.: Defining and detecting undesirable disturbance in the context of
marine eutrophication, *Mar. Pollut. Bull.*, 55(1–6), 282–297, doi:10.1016/j.marpolbul.2006.08.028, 2007.
- Tett, P., Carreira, C., Mills, D. K., Van Leeuwen, S., Foden, J., Bresnan, E. and Gowen, R. J.: Use of a
Phytoplankton Community Index to assess the health of coastal waters, *ICES J. Mar. Sci.*, 65(8), 1475–
1482, doi:10.1093/icesjms/fsn161, 2008.
- 815 Tett, P., Gowen, R., Painting, S., Elliott, M., Foster, R., Mills, D., Bresnan, E., Capuzzo, E., Fernandes,
T., Foden, J., Geider, R., Gilpin, L., Huxham, M., McQuatters-Gollop, A., Malcolm, S., Saux-Picart, S.,
Platt, T., Racault, M.-F. and Sathyendranath, M.: A framework for understanding marine ecosystem
health, *Mar. Ecol. Prog. Ser.*, 494, 1–27, doi:10.3354, 2013.
- UK Pelagic Habitats Expert Group: Plankton lifeform traits master list (v4). The Archive for Marine
820 Species and Habitats Data (DASSH). (Dataset). <https://doi.org/10.17031/1709>, [online] Available from:
<https://doi.org/10.17031/1709>, 2020.
- Uriarte, I., Villate, F., Iriarte, A., Fanjul, Á., Atkinson, A. and Cook, K.: Opposite phenological responses
of zooplankton to climate along a latitudinal gradient through the European Shelf, *ICES J. Mar. Sci.*,
doi:10.1093/icesjms/fsab008, 2021.
- 825 Utermöhl, H.: Zur vervollkommnung der quantitativen Phytoplankton Methodik., *Mittl. Int Verein Theor
Angew Limnol*, 9, 1–38, 1958.
- Wasmund, N., Kownacka, J., Göbel, J., Jaanus, A., Johansen, M., Jurgensone, I., Lehtinen, S. and
Powilleit, M.: The diatom/dinoflagellate index as an indicator of ecosystem changes in the Baltic Sea 1.
principle and handling instruction, *Front. Mar. Sci.*, 4(FEB), doi:10.3389/FMARS.2017.00022, 2017.
- 830 Wells, M. L., Karlson, B., Wulff, A., Kudela, R., Trick, C., Asnaghi, V., Berdalet, E., Cochlan, W.,
Davidson, K., De Rijcke, M., Dutkiewicz, S., Hallegraef, G., Flynn, K. J., Legrand, C., Paerl, H., Silke,
J., Suikkanen, S., Thompson, P. and Trainer, V. L.: Future HAB science: Directions and challenges in a
changing climate, *Harmful Algae*, 91, doi:10.1016/j.hal.2019.101632, 2020.
- Weston, K., Greenwood, N., Fernand, L., Pearce, D. J. and Sivy, D. B.: Environmental controls on
835 phytoplankton community composition in the Thames plume, U.K., *J. Sea Res.*, 60, 246–254,
doi:10.1016/j.seares.2008.09.003, 2008.
- Widdicombe, C. E., Eloire, D., Harbour, D., R.P., H. and Somerfield, P. J.: Long-term phytoplankton
community dynamics in the Western English Channel, *J. Plankton Res.*, 32, 643–655, 2010.
- Wilkinson, M., Dumontier, M. and Aalbersberg, I.: The FAIR Guiding Principles for scientific data
840 management and stewardship. *Sci. Data* 3 (2016), *Sci. data*, 3, 1–9, 2016.
- WoRMS Editorial Board: World Register of Marine Species. Available from
<http://www.marinespecies.org> at VLIZ. Accessed 2020-10-05. doi:10.14284/170, 2020.
- Zingone, A., Harrison, P. J., Kraberg, A., Lehtinen, S., McQuatters-Gollop, A., O'Brien, T., Sun, J. and
Jakobsen, H. H.: Increasing the quality, comparability and accessibility of phytoplankton species

13 Appendices

850 **Table A1:** Relevant traits in lifeform traits list. The trait list reflects the lifeforms and types of plankton data in the datasets used to date in lifeform-based assessment. The list is a living document, whose status reflects ongoing efforts to refine and improve the lifeform approach as well as a necessary compromise between focusing on traits which inform the lifeforms currently used for assessments and including additional information and traits where these are known and readily available. Trait groups for the plankton type “Protozoa” have recently been added to the trait list but are not shown here as they are not finalised and not used in any of the current lifeforms. The inclusion of ‘*protozoa*’ and ‘*ciliates*’ (which are protozoa) designation under ‘*phytoplankton type*’ is to ensure these key taxa are captured into lifeforms. One important example is the abundant group ‘*Flagellates*’ which has ‘*Plankton Type = phytoplankton*’ to allow assignment of the phytoplankton trait groups, despite including both phototrophic and heterotrophic taxa and thus *Phytoplankton Type = protozoa*’.

860 In the traits list: spaces between words are omitted, e.g. ‘*Plankton Type*’ is written as ‘*PlanktonType*’, and Zooplankton is often abbreviated as ‘*Zoo*’. For all trait categories, ‘*Y*’: yes (trait applies); ‘*N*’: no (trait does not apply); The following definitions apply to all columns (trait categories) of the list, with additional details given in the table where relevant:

1. ‘*Ambiguous*’: taxa cannot be reliably assigned to any one category for this trait, mostly because taxa within this group can fall under more than one trait category (e.g.: taxa categories which include individuals of both ‘large’ and ‘small’ size classes);
2. ‘*[blank]*’: trait is not use for this Plankton Type (e.g.: “PhytoHabitat” is blank for all zooplankton taxa groups);
- 865 3. ‘*NA*’: (Not applicable) trait is used for this Plankton Type but not relevant for this taxa (e.g.: the line “fish larvae” has a “ZooHabitat” of *NA* because they are neither meroplankton nor holoplankton, the line is intended to only contribute to the “fish larvae” lifeform);
4. ‘*NYA*’: (Not Yet Assigned) trait is used for this Plankton Type but is not yet assigned (e.g. phytoplankton which have not been assigned any of “tychopelagic”, “pelagic”, or “ambiguous” under “PhytoDepth”).

Trait Category	Trait Assigned	Description/notes (see also main text, Table 2)
Plankton Type	Phytoplankton	The ‘type’ defines which of the following groups of traits applies to the taxa group in question.
	Protozoa	
	Zooplankton	
Phytoplankton Type	(various)	Types included are: Cercozoa; Charophyte; Chlorophyte; Chrysophyte; Ciliate*; Cryptophyte; Cyanobacteria; Diatom; Dictyochophyte; Dinoflagellate; Euryarchaeote; Eustigmatophyceae; Haptophyte; Protozoa*; Raphidophyte; Silicoflagellate; Xanthophyceae. *Note that while the trait groups for the plankton type “Protozoa” have recently been added to the master taxa list, “protozoa” and “ciliates” (which are protozoa) remain designated under “phytoplankton type” to ensure these key taxa are captured in the lifeforms currently used in assessments. This reflects the ‘living’ nature of the Master Taxa List.
	Plankton Size	Sm

	Size Class	Lg	Large (> 20 µm individual cell diameter)
		1	Used to differentiate between taxa/groups that are of ambiguous size but have size information recorded in the raw datasets. 1 is large 'Plankton Size'; 2 is small 'Plankton Size'
		2	
		[blank]	Additional Size Class not required.
	Phytoplankton Depth	Pelagic	Living in the water column.
		Tychopelagic	Benthic taxa, which can become mixed into the water column.
	Phytoplankton Feeding Mechanism ('PhytoFeedingMech')	Auto	Autotrophic: nutrition by photosynthesis.
		Auto/Mixo	Auto- and mixotrophic
		Hetro	Heterotrophic: non-photosynthesising.
		Mixo	Mixotrophic: capable of obtaining nourishment via photo(auto)trophy and phago(hetero)trophy, as well as via osmo(hetero)trophy.
	Phytoplankton Habitat	Freshwater	
		Marine	
	Potentially Toxic or Nuisance	Ambiguous	Cannot assign trait: Some taxa in the group may be toxic. Taxa in this group cannot be identified to the taxonomic level required to confirm if they are a toxin producing strain or species using routine monitoring techniques. Toxin production in the relevant species may be strain dependent requiring confirmation using molecular methods. The mechanism of harm for this species may not be confirmed (e.g.: fish mortalities caused by either anoxia or ichthyotoxins).
Nuisance		Taxa produce effects which are detrimental to aquaculture and benthos via physical harm or causing anoxia or produce water discolourations, scums or foams that can be aesthetically, socially, or economically negative	
Non-Toxic		Taxa do not produce toxins which pose a risk to marine biota or human health, and do not produce nuisance effects, and do not produce nuisance effects.	
Zooplankton Traits	Zooplankton Type	(various)	Types included are: Bryozoa; Cephalochordate; Cephalopod; Chaetognath; Cladoceran; Crustacean; Echinoderm; Fish; Gastropod; Gelatinous; Hemichordate; Mollusc; Nematode; Nemertea; Phoronid; Polychaete; Rotifer; Sipuncula; Tunicate.
	Habitat	Holoplankton	Zooplankton taxa which spend their entire lifecycle in the plankton.
		Meroplankton	Taxa which spend part of their lifecycle as zooplankton.
	Diet	Carnivore	Taxa which prey mainly on other zooplankton.
		Herbivore	Taxa which are predominately suspension or filter feeders.
		Omnivore	Can use both carnivorous and herbivorous feeding.
		Ambiguous	Cannot assign trait: feeding mechanism variable or none of carnivore, herbivore or omnivore.
		Parasite	Feeds attached to food source either internally or externally.
	Crustacean	Y	Taxa of the Subphylum Crustacea
		N	
	Copepod	Y	Taxa of the Subclass Copepoda.
		N	
	Gelatinous	Y	Taxa of the phyla Cnidaria and Ctenophora only.
N			
Zooplankton Size	Sm	< 2 mm adult total body length.	
	Lg	≥ 2 mm adult total body length.	
	Ambiguous	Cannot assign trait: taxa in group includes those both under and over 2 mm.	

Table B1: Definitions of lifeforms by trait as defined in the traits list. See Table 2 for descriptions of traits used.

Lifeform	Definition (Trait(s))
(micro)Phytoplankton	PlanktonType = <i>Phytoplankton</i>
Large (micro)phytoplankton (>= 20 μm)	PlanktonType = <i>Phytoplankton</i> AND PhytoplanktonSize = <i>Lg</i>
Small (micro)phytoplankton (< 20 μm)	PlanktonType = <i>Phytoplankton</i> AND PhytoplanktonSize = <i>Sm</i>
Diatoms	PhytoplanktonType = <i>Diatom</i>
Pelagic diatoms	PhytoplanktonType = <i>Diatom</i> AND PhytoDepth = <i>Pelagic</i>
Tychopelagic diatoms	PhytoplanktonType = <i>Diatom</i> AND PhytoDepth = <i>Tychopelagic</i>
Potentially toxic or nuisance diatoms	PhytoplanktonType = <i>Diatom</i> AND Toxic_Nuisance = (<i>Toxic</i> OR <i>Nuisance</i>)
Dinoflagellates	PhytoplanktonType = <i>Dinoflagellate</i>
Autotrophic and mixotrophic dinoflagellates	PhytoplanktonType = <i>Dinoflagellate</i> AND PhytoFeedingMech = (<i>Auto</i> OR <i>Auto/Mixo</i>)
Potentially toxic or nuisance dinoflagellates	PhytoplanktonType = <i>Dinoflagellate</i> AND Toxic_Nuisance = (<i>Toxic</i> OR <i>Nuisance</i>)
Ciliates	PhytoplanktonType = <i>Ciliate</i>
Holoplankton	ZooHabitat = <i>Holoplankton</i>
Meroplankton	ZooHabitat = <i>Meroplankton</i>
Gelatinous zooplankton	PlanktonType = <i>Zooplankton</i> AND Gelatinous = <i>Y</i>
Carnivorous zooplankton	PlanktonType = <i>Zooplankton</i> AND ZooDiet = <i>Carnivore</i>
Non-carnivorous zooplankton	PlanktonType = <i>Zooplankton</i> AND ZooDiet = (<i>Herbivore</i> OR <i>Omnivore</i> OR <i>Ambiguous</i>)
Crustaceans	Crustacean = <i>Y</i>
Large copepod species(>=2 mm)	Copepod = <i>Y</i> AND ZooSize = <i>Lg</i>
Small copepod species (< 2 mm)	Copepod = <i>Y</i> AND ZooSize = <i>Sm</i>
Fish larvae	ZooType = <i>Fish</i>