

1 The ABoVE L-band and P-band Airborne SAR Surveys

2 Charles E. Miller¹, Peter C. Griffith², Elizabeth Hoy³, Naiara S. Pinto¹, Yunling Lou¹, Scott Hensley¹,
3 Bruce D. Chapman¹, Jennifer Baltzer⁴, Kazem Bakian-Dogaheh⁵, W. Robert Bolton⁶, Laura Bourgeau-
4 Chavez⁷, Richard H. Chen¹, Byung-Hun Choe⁸, Leah Clayton⁹, Thomas A. Douglas¹⁰, Nancy French⁷,
5 Jean E. Holloway¹¹, Gang Hong⁸, Lingcao Huang¹², Go Iwahana⁶, Liza Jenkins⁷, John S. Kimball¹³,
6 Tatiana Loboda¹⁴, Michelle Mack¹⁵, Philip Marsh¹⁶, Roger J. Michaelides¹⁷, Mahta Moghaddam⁵,
7 Andrew Parsekian¹⁸, Kevin Schaefer¹², Paul R. Siqueira¹⁹, Debjani Singh²⁰, Alireza Tabatabaenejad,⁵
8 Merritt Turetsky²¹, Ridha Touzi,⁸ Elizabeth Wig²², Cathy J. Wilson²³, Paul Wilson⁸, Stan D.
9 Wullschleger²⁰, Yonghong Yi^{1,24}, Howard A. Zebker²², Yu Zhang⁸, Yuhuan Zhao⁵, Scott J. Goetz²⁵

10

11 ¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, 91109, USA

12 ²NASA Goddard Space Flight Center / SSAI, Greenbelt, Maryland, USA

13 ³NASA Goddard Space Flight Center / GST, Inc., Greenbelt, Maryland, USA

14 ⁴Department of Biology, Wilfrid Laurier University, Waterloo, Ontario, Canada

15 ⁵Department of Electrical Engineering, University of Southern California, Los Angeles, CA, USA

16 ⁶International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, AK 99775-7340 USA

17 ⁷Michigan Tech Research Institute, Michigan Technological University, Ann Arbor, MI 48105 USA

18 ⁸Canada Centre for Remote Sensing, Ottawa, Ontario K1A0E4, Canada

19 ⁹Department of Earth & Planetary Sciences, Yale University, New Haven, CT, USA

20 ¹⁰U.S. Army Cold Regions Research and Engineering Laboratory, Fort Wainwright, Alaska 99709 USA

21 ¹¹Department of Geography, University of Ottawa, Ottawa ON K1N 6N5, Canada

22 ¹²National Snow and Ice Data Center (NSIDC), University of Colorado, Boulder, Colorado 80309-0449 USA

23 ¹³NTSG, WA Franke College of Forestry & Conservation, The University of Montana, Missoula, Montana USA

24 ¹⁴Department of Geographical Sciences, University of Maryland, College Park, Maryland 20742, USA

25 ¹⁵Center for Ecosystem Science and Society and Department of Biological Sciences, Northern Arizona University, Flagstaff,
26 AZ 86011 USA

27 ¹⁶Department of Geography and Cold Regions Research Centre, Wilfrid Laurier University, Waterloo, Ontario, Canada

28 ¹⁷Department of Geophysics, Colorado School of Mines, Golden, CO, USA

29 ¹⁸Department of Geology & Geophysics, University of Wyoming, 1000 E University Ave., Laramie, WY, USA

30 ¹⁹Department of Electrical and Computer Engineering, University of Massachusetts, Amherst, MA 01003-9284 USA

31 ²⁰Biological and Environmental Systems Science Directorate, Oak Ridge National Laboratory, Oak Ridge, Tennessee USA

32 ²¹Ecology and Evolutionary Biology Department, University of Colorado Boulder

33 ²²Department of Geophysics, Stanford University, Stanford, CA, USA

34 ²³Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, New Mexico USA

35 ²⁴Joint Institute for Regional Earth System Science & Engineering, The University of California, Los Angeles, CA 90095-
36 7228 USA

37 ²⁵School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, AZ 86011 USA

38

39

40 *Correspondence to:* Charles E. Miller (charles.e.miller@jpl.nasa.gov)

41

42 **Abstract**

43 Permafrost-affected ecosystems of the Arctic-boreal zone in northwestern North America are
44 undergoing profound transformation due to rapid climate change. NASA’s Arctic Boreal Vulnerability
45 Experiment (ABoVE) is investigating characteristics that make these ecosystems vulnerable or resilient
46 to this change. ABoVE employs airborne synthetic aperture radar (SAR) as a powerful tool to
47 characterize tundra, taiga, peatlands, and fens. Here, we present an annotated guide to the L-band and P-
48 band airborne SAR data acquired during the 2017, 2018, 2019, and 2022 ABoVE airborne campaigns.
49 We summarize the ~80 SAR flight lines and how they fit into the ABoVE experimental design. The
50 Supplemental Information provides hyperlinks to extensive maps, tables, and every flight plan as well
51 as individual flight lines. We illustrate the interdisciplinary nature of airborne SAR data with examples
52 of preliminary results from ABoVE studies including: boreal forest canopy structure from tomoSAR
53 data over Delta Junction, AK and the Boreal Ecosystem Research and Monitoring Sites (BERMS) site
54 in northern Saskatchewan and active layer thickness and soil moisture data product validation. This
55 paper is presented as a guide to enable interested readers to fully explore the ABoVE L- and P-band
56 airborne SAR data.

57

58

59 **Short Summary**

60 NASA’s Arctic Boreal Vulnerability Experiment (ABoVE) conducted airborne synthetic aperture radar
61 (SAR) surveys of over 120,000 km² in Alaska and northwestern Canada during 2017, 2018, 2019, and
62 2022. This paper summarizes those results and provides links to details on ~80 individual flight lines.
63 This paper is presented as a guide to enable interested readers to fully explore the ABoVE L- and P-
64 band SAR data.

65

66

67 **Keywords:** Airborne Synthetic Aperture Radar (SAR), Interferometric SAR (InSAR), Polarimetric
68 SAR (PolSAR), Tomographic SAR (tomoSAR), Arctic, tundra, taiga, boreal forest, permafrost,
69 wildfire, soil moisture, active layer thickness (ALT), Arctic Boreal Vulnerability Experiment (ABoVE)

70

71

72 **Copyright Statement:**

73 © California Institute of Technology. Government funding acknowledged.


74

75

76

77 **1 Introduction**

78 The Arctic region contains a remarkable diversity of cold-adapted biota, habitats, and permafrost-
79 affected ecosystems [McGuire et al. 2009; Vincent et al. 2011]. As with other components of the Arctic
80 system, Arctic ecosystems are strongly interdependent and the rapid degradation of the Arctic
81 cryosphere is altering their physical, biogeochemical, and biological linkages in ways that may be
82 irreversible [Vincent et al. 2011; Hinzman et al. 2013]. Understanding characteristics that make Arctic
83 ecosystems vulnerable or resilient to this change is the overarching objective of NASA's Arctic Boreal
84 Vulnerability Experiment (ABoVE, <https://above.nasa.gov/>). Miller et al. [2019] describes how airborne
85 campaigns fit into the broader ABoVE research strategy and how the foundational synthetic aperture
86 radar (SAR) measurements formed the framework around which all other airborne data acquisitions
87 were planned.
88
89



90
91 **Figure 1. Flight lines for the L-band and P-band PollnSAR measurements capture critical bioclimatic, permafrost, and geographic**
92 **gradients as well as key field sites and long-term measurement records across the 4 Mkm² ABoVE domain. The flight lines**
93 **are collected into 10 composites which roughly correspond to the Alaskan (A1-A4) and Canadian (C1-C6) regions sampled on**
94 **individual flight days. © Google Maps**

95 ABoVE SAR flight lines (Figure 1) were planned to leverage legacy L- and P-band SAR transects
96 acquired during the pre-ABoVE period; remotely-sensed permafrost active layer thickness time series
97 derived from satellite interferometric SAR observations (ReSALT) [Schaefer et al. 2015]; SAR data
98 from PALSAR, PALSAR-2, RadarSat, RadarSat-2, and Sentinel-1; historic or planned airborne LiDAR
99 acquisitions; and data from existing field sites [Hoy et al. 2018]. Legacy airborne SAR flight lines
00 include the L-band grid acquired over the Boreal Ecosystem Research and Monitoring Sites (BERMS)
01 area near Prince Albert, SK during SMAP CanEx 2010 [Magagi et al. 2012], the P-band lines over the
02 BERMS area acquired from 2012-2015 during the Airborne Microwave Observatory of Subcanopy and
03 Subsurface (AirMOSS) Earth Ventures Sub-orbital (EV-S1) investigation [Allen et al. 2010;
04 Moghaddam et al. 2016], and a collection of 10 L- and P-band flight lines acquired over the Seward
05 Peninsula, Northwestern Interior, and North Slope of Alaska during 2014 and 2015 [Chen et al. 2019a,
06 2019b]. The BERMS area observations, in particular, link ABoVE to the Boreal Ecosystem–
07 Atmosphere Study (BOREAS) studies of the 1990s [Sellers et al. 1995; 1997].
08

09 Hoy et al. [2018] compiled information on more than 6,700 field sites and previous remote sensing data
10 sets to help plan the SAR flight lines and the ABoVE Airborne Campaigns [Miller et al. 2019]. This
11 compilation is intended to help investigators understand flight line choices and identify ground locations
12 used to anchor individual flight lines. SAR data users may also search for the underlying data available
13 within each flight line. Key anchor points for the SAR flight lines include: Active layer thickness
14 measurements from the Circumpolar Active Layer Monitoring network (CALM); Permafrost
15 temperatures and annual thaw depths from the Global Terrestrial Network for Permafrost (GTN-P)
16 database; Soil moisture and permafrost state data from the Department of Energy’s Next Generation
17 Ecological Experiment-Arctic (NGEE-Arctic) field sites on the Seward Peninsula and near Utqiagvik
18 (formerly Barrow), AK; Extensive in situ terrestrial and aquatic ecosystem data as well as airborne
19 LiDAR and spectral imagery from NSF’s National Ecological Observatory Network (NEON) D18
20 tundra field sites near Utqiagvik (Barrow), AK and Toolik Lake, AK, and from the D19 taiga field sites
21 near Caribou/Poker Creek, AK, Delta Junction, AK, and Healy, AK; Detailed ecological and physical
22 climate time series from NSF’s Long Term Ecological Research (LTER) Arctic (Toolik Lake) and
23 Boreal Forest (Bonanza Creek) sites; Long-term boreal forest inventory data from the Canadian
24 Forestry Service’s (CFS) Climate Impacts on Productivity and Health of Aspen (CIPHA) and High
25 Elevation & Latitude Climate Change Impacts & Adaptation (HELICIA) plots; and Long term
26 permafrost, hydrology and ecology time series records from the Canadian Changing Cold Regions
27 Network (CCRN) sites at Trail Valley Creek, NWT, Havikpak Creek, NWT, Scotty Creek, NWT,
28 Baker Creek, NWT, Wolf Creek Research Basin, YT, and the BERMS site at White Gull Creek, SK.
29

30 Airborne SAR data enable numerous ecosystem and ecosystem change research investigations [NRC
31 2014]. ABoVE researchers are using the airborne L- and P-band data to: Quantify permafrost active
32 layer thickness and soil moisture content [Bakian-Dogahneh et al. 2020]; Complement AirSWOT Ka-
33 band acquisitions to determine water surface elevations in Arctic lakes, wetlands, and rivers [Pitcher et
34 al. 2019a,b]; Investigate boreal forest and tundra fire scars, especially in conjunction with fire
35 disturbance plots [Tank et al. 2018; Walker et al. 2018 a,b; 2019a,b; French et al. 2020; Holloway et al.
36 2020; Loboda et al. 2021]; Map tree density and distribution across the Tundra-Taiga ecotone; Provide

37 control point data for the ArcticDEM [Porter et al. 2018; Meddens et al. 2018]; Investigate lidar-radar
38 fusion remote sensing for boreal forest characterization as a precursor to NISAR/IceSAT-2
39 investigations [Silva et al. 2021]; Quantify expansion and sediment mass flow from massive
40 retrogressive thaw slumps – so-called megaslumps – on the Peel Plateau along the Dempster Hwy west
41 of Fort McPherson [Kokelj et al. 2013; 2015]; Classify Arctic wetlands and habitats [French 2020]; and
42 Support algorithm development for NISAR (L-band) and BIOMASS (P-band) estimates of boreal forest
43 structure and above ground biomass [Quegan et al. 2019; Saatchi et al. 2019]. Goetz et al. [2021]
44 summarizes how the ABoVE airborne SAR data are helping advance Arctic-boreal understanding and
45 the remaining knowledge gaps still to be addressed.

46
47 This paper presents an annotated guide to enable interested readers to fully explore the ABoVE L- and
48 P-band SAR data acquired during the 2017, 2018, 2019, and 2022 ABoVE airborne campaigns. Section
49 2 provides details on the L- and P-band SAR instruments and the flight line catalog. Section 3
50 summarizes the daily sorties from each airborne campaign. Section 4 briefly describes the tomographic
51 SAR (tomoSAR) experiments flown over Delta Junction, AK and the BERMS site near Prince Albert,
52 SK. Section 5 describes some of the ABoVE SAR data products and their validation. Section 6
53 highlights the synergies between the L- and P-band airborne SAR data and other airborne sensors.
54 Section 7 summarizes access to the data products. Section 8 discusses potential future acquisitions and
55 outlooks for exploiting these data. Additionally, we include an Appendix which describes the ~80
56 ABoVE SAR flight lines and how each line fits into the ABoVE experimental design. The Appendix
57 also provides extensive maps and tables for every flight plan and individual flight lines as well as a list
58 of the acronyms and abbreviations. The Supplemental Information includes hyperlinked versions of the
59 tables for direct access to flight lines and flight plans.

60 **2 The L-Band and P-band Airborne SAR Instruments and Data Acquisition**

61 Both the L- and P-band airborne SARs are sensitive to geometrical and material properties of
62 vegetation, soil surface, and subsurface profiles [Saatchi and Moghaddam 2000; Tabatabaenejad et al.
63 2011; 2015]. The joint use of both L- and P-band gives enhanced sensitivity to near-surface (< 5 cm, L-
64 band) and root zone (10-40 cm, P-band) portions of the subsurface profile compared to use of either
65 wavelength alone [Du et al. 2015]. Airborne acquisitions with both SARs provide 6-10 m spatial
66 resolution, ~15 km swaths and transect lengths of 100 – 200 km, making them ideal for surveying
67 above-ground biomass and vegetation canopy structure [Hensley et al. 2014; 2016] as well as the
68 tundra-taiga ecotone [Montesano et al. 2016]. Special tomoSAR data were acquired over the well
69 characterized BERMS site in northern Saskatchewan and the NEON site in Delta Junction, AK to
70 quantify the performance of both SARs in reproducing the structure and biomass of boreal forests.
71

72 **2.1 The L-band SAR Instrument**

73 NASA's airborne L-band SAR (initially named the Uninhabited Aerial Vehicle Synthetic Aperture
74 Radar (UAVSAR) system) is a compact pod-mounted polarimetric instrument for interferometric

75 repeat-track observations that was developed at the NASA Jet Propulsion Laboratory (JPL) and Dryden
76 Flight Research Center (DFRC) in Edwards, CA. It operates at a center frequency of 1.2575 GHz
77 (wavelength = 23.8 cm) with 80 MHz bandwidth. It is deployed on a Gulfstream III aircraft and images
78 the Earth surface from a nominal 12.5 km altitude. The image swath is collected off-nadir in a ~22 km
79 wide with incidence angles ranging from ~22°–67°. The instrument spatial resolution is 0.8m (along
80 flight-line) by 1.7m (slant range, along line-of-sight (LOS) from the antenna to the ground).
81 Topographic information is derived from phase measurements that, in turn, are obtained from two or
82 more passes over a given target region. Its 1.26 GHz frequency results in radar images that are well-
83 correlated from pass to pass. Polarization agility facilitates terrain and land-use classification.

84
85 All L-band SAR data are publicly available at <http://uavsar.jpl.nasa.gov/> as individual InSAR products
86 or as a single look complex (SLC) stack product of coregistered images for individual flight lines.
87 Products are also available from the UAVSAR data portal at the Alaska SAR Facility Distributed
88 Active Archive Center (<https://asf.alaska.edu/data-sets/sar-data-sets/uavsar/>). The L-band SAR provides
89 key pre-launch algorithm development and validation data sets [Saatchi et al. 2019] for the NASA-
90 ISRO SAR (NISAR) mission [Rosen et al. 2017].
91

92 **2.2 The P-band SAR Instrument**

93 The P-band SAR was developed circa 2012 for the Earth Ventures Sub-orbital (EV-S1) Airborne
94 Microwave Observatory of Subcanopy and Subsurface (AirMOSS) investigation [Allen et al. 2010;
95 Moghaddam et al. 2016]. The radar is based on JPL's L-band UAVSAR system. The P-band SAR
96 inherits UAVSAR's existing L-band RF and digital electronics subsystems. New up- and down-
97 converters convert the L-band signals to UHF frequencies (280-440 MHz). The passive antenna is based
98 on the legacy GeoSAR design [Chapin et al. 2012].
99

00 The P-band SAR was flown more than 1200 h from 2012 to 2015, covering regions of 2500 km² spread
01 over nine major biomes in North America during the AirMOSS EV-S1 investigation [Tabatabaenejad
02 et al. 2020]. Legacy acquisitions in Alaska [Chen 2019a, b] and over the BERMS site in northern
03 Saskatchewan [Chapin et al. 2012, 2018] provide an opportunity for extended time series analysis. All
04 P-band SAR data are publicly available at <http://uavsar.jpl.nasa.gov/>. Additionally, ABoVE P-band
05 SAR data will provide valuable insights into the characterization of boreal forest and tundra ecosystems
06 by the upcoming BIOMASS mission [Le Toan et al. 2011; Quegan et al. 2019].
07

08 **2.3 The Platform Precision Autopilot (PPA) System**

09 To support cm-precision interferometric land surface characterization, repeat pass measurements
10 acquired by the SARs need to be taken from flight paths that are nearly identical. Both the L- and P-
11 band SARs utilize real-time GPS that interfaces with the platform flight management system (FMS) to
12 confine the repeat flight path to within a 10 m tube over a 200 km course in conditions of calm to light
13 turbulence. The FMS is also referred to as the Platform Precision Autopilot (PPA). Additionally, the

14 radar vector from the aircraft to the ground target area must be similar from pass to pass. This is
15 accomplished with an actively scanned antenna designed to support electronic steering of the antenna
16 beam with a minimum of 1° increments over a range to exceed ±15° in the flight direction.

17
18 ABoVE SAR measurements were typically acquired with platform RMS deviations less than ±3 m. Any
19 platform deviations larger than ±10 m from the programmed flight path resulted in the acquisition being
20 terminated and a real time decision made to reacquire the line from the beginning or to continue with
21 the flight plan and proceed to the next line. This decision balanced the science priority of the flight line,
22 fuel consumption and remaining endurance, the number of flight lines yet to be acquired in the day's
23 flight plan, distance from our base of operations, and whether there would be future opportunities to
24 collect a given line by adding it to an upcoming flight plan. The flight team was extremely efficient in
25 executing these decisions, resulting in 95% flight line acquisition success across the 2017-2019 period.
26

27 **2.4 Airborne SAR Flight Line and Flight Plan Designations**

28 The JPL SAR team devised a convenient and powerful way to identify airborne SAR data acquisitions
29 for the Facility and PI instruments under their charge. Each L-band or P-band SAR flight line receives a
30 unique 5-digit identifier consisting of the three-digit GPS compass heading followed by a two-digit
31 index. A 6-character text string is also associated with each line for ease of identification. The text
32 string proceeds the numerical ID and usually provides abbreviated geographic or infrastructure
33 information that characterizes the line. For example, L-band flight line Teller_04901 identifies the flight
34 line on the Seward Peninsula that overflies the NGEE-Arctic Teller watershed. The flight line identifier
35 is a constant and, once assigned, is used whenever a line is reflown. In some cases, there are
36 overlapping or nearly identical flight lines which differ slightly in their ID number. L-band and P-band
37 flight lines use the same flight line identification system, allowing rapid identification of overlapping L-
38 and P-band data acquisitions.

39
40 Flight Plans are assembled from the composite flight lines for a given sortie. Each flight plan also
41 receives a unique 5-digit identifier based on the year flown (digits 1 and 2) and the flight number for
42 that year (digits 3-5). For example, L-band flight plan 17093 was flown in 2017 and was the 93rd sortie
43 flown that year. Note that there may be more than one sortie flown on a given day, in which case each
44 would have a unique flight plan identifier even though they were flown on the same calendar day and
45 may include some or all of the same flight lines.

46
47 In the Supplemental Information we provide hyperlinks to the JPL UAVSAR data portal
48 (<https://uavsar.jpl.nasa.gov/cgi-bin/data.pl>). This provides links to the individual flight line data, maps,
49 and related flight plans that acquired data over one or more of the individual flight lines. We hope this
50 enables interested readers to explore the ABoVE L- and P-band SAR data more fully. These data and all
51 other airborne data from the ABoVE campaigns may be explored on NASA's EarthData ABoVE Portal
52 (<https://search.earthdata.nasa.gov/portal/above/search>). Ground sites used to design the orientation and
53 locations of the flight lines are archived at the ORNL DAAC [Hoy 2018].

55 **3 The ABoVE Airborne SAR Campaigns**

56 The L-band (Figure 2) and P-band (Figure 3) SARs were considered foundational measurements in the
57 ABoVE airborne campaign strategy [Miller et al. 2019]. The ~80 flight lines described in the Appendix
58 formed the framework for the remainder of the airborne remote sensing acquisitions. The ABoVE SAR
59 strategy was to execute same day acquisitions of both L- and P-band flight lines (Figure 1) for a given
60 sortie during 2017 to optimize dual frequency retrievals; however, technical issues forced us to fly the
61 instruments sequentially. The baseline L-band campaigns were flown in June (DOY 164-173) and
62 September (DOY 251-263) of 2017 to characterize the land surface during periods of minimum and
63 maximum active layer thickness, respectively. Subsequent L-band campaigns in 2018 (DOY 231-241),
64 2019 (DOY 247-260) and 2022 (DOY 226-237) provide a time series synched to maximum annual
65 active layer thickness. P-band campaigns were conducted in May-June (DOY 142-157) and August
66 (DOY 219-227) of 2017. There was a 2-day P-band mini-campaign in October 2017 to extend the
67 legacy time series of early cold season acquisitions over the Seward Peninsula, NW Alaska and North
68 Slope Alaska (DOY 280-283).

69



70
71
72
73
74

Figure 2. Sahtu students Mandy Bayha (front left) and Joanne Speakman (front center) pose with their mentor Cindy Gilday (front right) and NASA flight crew in Yellowknife, NT after completing a L-band SAR survey flight around the Great Slave Lake Region on 22 August 2018 (Flight Plan 18048). This experience gave these Northerners a new appreciation for how NASA was helping understand, preserve, and protect their lands. Photo Credit: Stephen M. Fochuk, Government of Northwest Territories.

75
76
77



78

79
80

Figure 3. The P-band SAR team with the NASA JSC G-III (N992NA) on the tarmac in Fairbanks, AK on 18 August 2017 after completing a survey of the Upper Mackenzie Valley (Flight plan 17083). Photo Credit: M. Moghaddam.

81

82 3.1 Alaskan Flight Lines

83 The Alaskan SAR flight lines are broken into four main regional collections: A1) North Slope Alaska,
84 A2) Seward Peninsula and Northwest Alaska, A3) Eastern Interior, and A4) Southwest Alaska and the
85 Yukon-Kuskokwim Delta (**Figure 1**). Individual flight
86 lines were planned based on long-term ground
87 monitoring sites [Hoy et al. 2018], existing or planned
88 field research, recent disturbances, important geographic
89 or ecological gradients, complementary remote sensing
90 data, and consultation with indigenous peoples and
91 governments [Miller et al. 2019]. Legacy L- and P-band
92 flight lines from the AirMOSS EV-S1 investigation
93 [Allen et al. 2010; Moghaddam et al. 2016] in the
94 Seward Peninsula, NW Alaska, and the North Slope
95 were adapted for ABoVE use. Acquisition of P-band
96 flight lines in the central Interior was not possible due to
97 a military radar keep-out zone centered near Clear, AK.
98 The keep-out zone is shown in all P-band flight plan
99 maps (Ex. **Figure 4**).




Figure 4. The military radar at Clear, AK creates a large P-band operations keep-out zone in the central Interior (red areas). The aircraft symbol marks our Fairbanks International Airport (PAFA) base of operations. Data acquisitions (blue bars) are from Flight Plan 17054. © Google Maps

01 3.2 Canadian Flight Lines

02 The Canadian SAR flight lines are broken into six regional collections: C1) Lower Mackenzie Valley
03 and Northern Yukon Territory, C2) Southern Yukon Territory, C3) Upper Mackenzie Valley, C4) Great
04 Slave Lake Region, C5) Transboundary Watershed, and C6) Southern Boreal Forest/BERMS.
05 Individual lines were planned based on long-term ground monitoring sites [Hoy et al. 2018], existing or
06 planned field research, recent disturbances, important geographic or ecological gradients,
07 complementary remote sensing data, and consultation with local inhabitants and governments [Miller et
08 al. 2019]. Legacy L- and P-band flight lines in the BERMS area from the CANEX 2010 campaign
09 [Magagi et al. 2012] and the AirMOSS EV-S1 investigation [Chapin et al. 2012, 2018] provide the
10 potential to establish longer time series.

11
12 Flight planning for the Canadian transects benefited tremendously from consultations with our
13 Canadian colleagues and interested parties in Yellowknife, NT and Whitehorse, YT in 2015 and 2016.
14 Extensive discussions with the Government of the Northwest Territories (GNWT), the Government of
15 the Yukon Territory, First Nations representatives, and scientists from Polar Knowledge Canada
16 (POLAR), the NWT Center for Geomatics, and the Canadian Forestry Service (CFS) Northern Forestry
17 Centre (NoFC) were critical to designing a strategy that captured many of their observing priorities.
18 Subsequent discussions in Yellowknife during 2017 and 2018 enabled us to disseminate preliminary
19 results and coordinate the flights with same-day field data acquisitions.

20


21 4 TomoSAR Measurements of Boreal Forest Structure

22 SAR tomographic methods have proven extremely adept at measuring vegetation vertical structure at a
23 variety of wavelengths including L- and P-bands. The three-dimensional vegetation structure and its
24 changes resulting from either natural or anthropogenic causes are key ecosystem monitoring parameters.
25 ABoVE collected tomographic L- and P-band SAR data over the boreal forest near Delta Junction, AK
26 in September of 2017. UAVSAR (L-band) and the German Space Agency's F-SAR (L- and S-bands)
27 acquired coordinated tomographic SAR data at the BERMS site near Saskatoon, SK in August 2018.
28 Ground truth data sets and LiDAR data from the NASA LVIS system were also acquired at BERMS in
29 2017 [Blair et al. 2018]. We compared L- and P-band tomography at Delta Junction and L-band and S-
30 band tomography from the two systems, to each other, and to the LiDAR data sets at BERMS. Here we
31 provide a preliminary analysis of the data acquired at BERMS.

32
33 BERMS is a southern boreal forest site with gentle topography dominated by Jack Pine and Aspen
34 stands. There is active logging in the area and the site contains clear cut areas and new growth stands in
35 various maturity states. The tomography data acquisition at BERMS was planned jointly in cooperation
36 with the German Space Agency (DLR) who flew the F-SAR radar and acquired data at L-band and S-
37 band. The UAVSAR and F-SAR flight lines were designed to overlap each other and LVIS data
38 acquired at the site in 2017. LVIS reacquired BERMS area data again in 2019 with the LVIS-F and
39 LVIS-C instruments [<https://lvis.gsfc.nasa.gov/Data/Maps/ABOVE2019Map.html>]. Figure 5 (left)
40 shows swaths for the UAVSAR and F-SAR radars along with the LVIS data. UAVSAR acquired L-
41 band tomography data on a racetrack pattern to get multiple incidence angle data for most points in the
42 swath. Because UAVSAR and F-SAR fly at 12500 mAGL and 4200 mAGL, respectively, it is not
43 possible to acquire data with the same incidence angles across the swath. Thus, we configured the flight
44 lines to overlap so that the 40° incidence angle points would coincide. Figure 5 (right) shows photos
45 collected at four of our seventeen ground truth sites during the tomoSAR acquisitions.


46
47 LVIS full waveform LiDAR provides surface elevations and tree height estimates as well as LiDAR
48 echo strength throughout the canopy and thereby information on the canopy internal structure. From
49 LVIS waveforms many products are possible including surface elevation, tree height, moments of the
50 returned waveform distribution and cumulative percentile elevations. We compared these waveforms to
51 radar tomographic profiles for the different radar wavelengths.

52
53 BERMS field measurements consist of soil moisture measurements at the 17 sites using the average of
54 15 measurements distributed over 60 m × 60 m plots on the day of the UAVSAR radar observations. At
55 BERMS the soil was very dry, roughly 10% volumetric soil moisture or less, during the radar
56 observations. During the summer of 2020 diameter at breast height (DBH) measurements, used to
57 estimate biomass, for a subset of our selected sites was planned but postponed due to COVID-19.
58



59
 60 **Figure 5.** LEFT: Experimental design for the BERMS area TomoSAR flights in August 2018. The large White and yellow
 61 boxes show the ~18 km-wide UAVSAR L-band swaths. The offset is due to the off-nadir viewing angle of the L-band SAR –
 62 it is pointed to the south when flying the 270° swath and pointed north when flying the 90° swath. The small white box near
 63 the center of the image marks the ~3 km-wide F-SAR swath. LVIS LiDAR data are the ~1.5 km-wide colored swaths across
 64 the image; the color scaling reflects the canopy height. RIGHT: Photos from four of the 17 plots used for in situ ground truth
 65 measurement at BERMS. Vegetation at BERMS was mostly Jack Pine and Aspen with many areas having dense understory
 66 vegetation. Most areas have substantial detritus and ground litter left over from previous logging operations. © Google Earth

67 **Figure 6** compares UAVSAR L-band tomography with F-SAR L-band and S-band tomography and
 68 LVIS LiDAR data along a transect shown as a yellow line on the right in the figure. Tree height along
 69 the transect varied from 10-20 m. Middle of Figure 2 is the UAVSAR L-band transect. Top of the figure
 70 shows the LVIS RH25m RH50, RH75 and RH95 profiles overlaid on the UAVSAR tomogram and
 71 below are the radar and LiDAR vertical profiles. On the left of Figure 2 are the F-SAR L-band and S-
 72 band tomographic profiles along with the LVIS RH100 data. The L-band radar profiles exhibit power
 73 concentrated at the base of the canopy whereas the LVIS LiDAR data show more return from the
 74 middle portion of the canopy. S-band obtains greater returns in the upper canopy compared to L-band
 75 and show more uniform scattering within the canopy.



77
 78 **Figure 6.** On the left shows location of transect as yellow line overlaid on UAVSAR imagery in grayscale and LVIS RH95 data is color
 79 where UAVSAR L-band, DLR F-SAR L and S-band and LVIS LiDAR profiles are compared. Center figure shows UAVSAR L-band
 80 tomographic profiles along the transect along with the corresponding LVIS LiDAR profiles. On the right are the corresponding F-SAR L
 81 and S-band tomographic profiles. The L-band radar profiles exhibit power concentrated at the base of the canopy whereas the LVIS LiDAR
 82 data show more return from the middle portion of the canopy. S-band obtains greater returns in the upper canopy compared to L-band and
 83 show more uniform scattering within the canopy.

84 5 ABoVE SAR Data Products

85 Here we highlight some data sets enabling or derived from the ABoVE L- and P-band airborne SAR
86 acquisitions. They represent the current state of the art for the study of permafrost-affected ecosystems
87 using SAR. The ABoVE science team continues to develop additional products and the insights from
88 these studies will be published separately. Links to the repositories for each of these data sets are
89 provided below.

90

91 5.1 Active Layer Thickness (ALT)

92 The Permafrost Dynamics Observatory (PDO) data product estimates seasonal subsidence, active layer
93 thickness (ALT), soil Volumetric Water Content (VWC), and uncertainties at 30-m resolution for 66
94 flight lines across Alaska and Northwest Canada [Michaelides et al. 2021; Chen et al. 2021a,b]. The
95 PDO retrieval uses L-band Synthetic Aperture Radar (SAR) data acquired by the Uninhabited Aerial
96 Vehicle Synthetic Aperture Radar (UAVSAR) instrument and P-band data acquired by the Airborne
97 Microwave Observatory of Subcanopy and Subsurface (AirMOSS) instrument. The PDO results for
98 each flight line appear in separate netcdf files. Each line has a spatial resolution of 30 meters on the
99 ABoVE common grid with a width of 22 km based on the swath width of the AirMOSS instrument. The
00 flight lines as a whole cover many ecosystem types and provide north-south and east-west gradients in
01 ALT and soil moisture across the ABoVE domain ([https://daac.ornl.gov/cgi-
02 bin/dsviewer.pl?ds_id=1796](https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1796)).

03 **Table 1** defines all variables in PDO data files. The first eight variables represent the four primary
04 outputs of the PDO algorithm and associated uncertainties. Subsidence and ALT represent one-time
05 measurements for the 2017 thaw season [Schaefer et al. 2021]. VWC, defined as the ratio of water
06 volume to total soil volume, represents soil moisture at maximum thaw for 2017. We assume a vertical
07 profile of VWC and estimate Sw0 and wtd, the parameters that define the exact shape of the assumed
08 profile. The product includes a Python script that will create a map of VWC averaged over any user
09 specified depth range. We included maps of VWC averaged over depth ranges of hand-held soil
10 moisture probes commonly used in ABoVE fieldwork.

11

12 **Table 1. Variables in the Permafrost Dynamics Observatory (PDO) data files**

Variable	Full Name	Units	Description
alt	Active Layer Thickness	M	Maximum thaw depth at the end of summer
sub	Subsidence	M	Surface subsidence from start of thaw after snow melt to maximum thaw depth in August or September
Sw0	Surface Saturation Fraction	m ³ /m ³	The ratio of water volume to pore space volume at the surface or zero meters depth
wtd	Water Table Depth	M	The depth from the surface to the level where the soil is 100% saturated

alt_unc	Uncertainty ALT	M	Uncertainty of estimated ALT
sub_unc	Uncertainty Subsidence	M	Uncertainty of estimated seasonal subsidence
Sw0_unc	Uncertainty Surface Saturation Fraction	m3/m3	Uncertainty of estimated surface water saturation fraction
wtd_unc	Uncertainty Water Table Depth	M	Uncertainty of estimated water table depth
mv_6cm	VWC from 0 to 6 cm	m3/m3	The ratio of water volume to soil volume averaged over zero to 6 cm depths
mv_12cm	VWC from 0 to 12 cm	m3/m3	The ratio of water volume to soil volume averaged over zero to 12 cm depths
mv_20cm	VWC from 0 to 20 cm	m3/m3	The ratio of water volume to soil volume averaged over zero to 20 cm depths
mv_alt	VWC from 0 to ALT	m3/m3	The ratio of water volume to soil volume averaged over the entire active layer, from zero to ALT

13 5.2 Alaska Active Layer and Soil Moisture Properties from Airborne P-band SAR

14 Chen et al. [2019b] synthesized the P-band polarimetric synthetic aperture radar (PolSAR) data
15 collected in August and October of 2014 and 2015 during the AirMOSS EV-S1 investigation with the
16 ABoVE P-band measurements collected in August and October of 2017 to estimate soil geophysical
17 properties over 12 study sites in Northern Alaska (see Figure S2). Soil properties reported include the
18 ALT, soil dielectric constant, soil moisture profile, surface roughness, and their respective uncertainty
19 estimates at 30-m spatial resolution (<https://doi.org/10.3334/ORNLDAAAC/1657>).

20
21 Most of the study sites are located within the continuous permafrost zone and where the aboveground
22 vegetation consisting mainly of dwarf shrub and tussock/sedge/moss tundra has a minimal impact on P-
23 band radar backscatter. These data were used as inputs to the L-band ReSALT data described in Section
24 5.1.

25 5.3 In Situ Soil Moisture and Thaw Depth Measurements

26 In situ measurements of soil moisture, thaw depth, and other quantities are essential to calibrate and
27 validate ABoVE SAR retrievals. The ABoVE team established a set of standardized measurement
28 protocols for field plots to ensure uniform data products and quality in measurements collected by
29 different groups across the ABoVE domain and across multiple years. Numerous teams collected in situ
30 data during the initial 2017 Airborne Campaign, with more targeted field acquisitions conducted in
31 2018 and 2019 [Bourgeau-Chavez et al. 2019a,b, 2021; Bakian-Dogaheh et al. 2020; Loboda et al.
32 2021].

33
34 Bourgeau-Chavez and coworkers [2019a,b; 2021] collected soil moisture at 6, 12, 20, and 50 cm depths,
35 ALT, soil profiles and biophysical measurements of aboveground canopy and ground layers in the
36 Greater Slave Lake Region (C4). These data provide vegetation community characteristics and

37 biophysical data collected in 2018 from areas that were burned by wildfire in 2014 and 2015, and from
38 nine unburned validation sites. Vegetation data include vegetation inventories, ground cover, regrowth,
39 tree diameter and height, and woody seedling/sprouting data at burned sites, and similar vegetation
40 community characterization at unburned validation sites. Additional measurements included soil
41 moisture, collected for validation of the UAVSAR airborne collection, and depth to frozen ground at the
42 nine unburned sites. This 2018 fieldwork completes four years of field sampling at the wildfire areas.

43
44 Bakian-Dogaheh et al. [2020] measurements included soil dielectric properties, temperature, and
45 moisture profiles, active layer thickness (ALT), and measurements of soil organic matter, bulk density,
46 porosity, texture, and coarse root biomass from the surface to permafrost table in soil pits at selected
47 sites along the Dalton Highway in Northern Alaska (A1). Their investigation sites included Franklin
48 Bluffs, Sagwon, Happy Valley, Ice Cut, and Innavait Creek. Measurements collected at Franklin Bluffs
49 were concurrent with an August 2018 ABoVE L-band flight.
50 (<https://doi.org/10.3334/ORNLDAAAC/1759>).

51
52 Loboda et al [2022] collected field measurements from unburned sites and single and repeated burns in
53 the Noatak River valley and the Seward Peninsula regions of the Alaska tundra in July-August in the
54 years 2016-2018. The data include ocular assessment of vegetation cover, soil moisture at 6 and 12 cm,
55 soil temperature at 10 cm, organic soil thickness, thaw depth, and weather measurements.
56 (<https://doi.org/10.3334/ORNLDAAAC/1919>)

57
58 The strong partnership between the ABoVE and NGEE-Arctic projects also resulted in coordinated
59 same-day acquisition of airborne L- and P-band SAR data with in situ soil moisture and thaw depth
60 measurements over the NGEE-Arctic study site at Barrow (Utqiagvik), AK and the Seward Peninsula
61 watersheds near Teller, AK, Council, AK, and Kougarak, AK [Wilson et al. 2018]. These data provide
62 critical calibration for the ABoVE SAR retrievals under continuous (Utqiagvik) and discontinuous
63 (Seward Peninsula) permafrost conditions. Version 2 (V2) of the in situ soil moisture and thaw depth
64 measurements covering years 2017-2019 was released in November 2020.
65 (<https://doi.org/10.5440/1423892>)

66 **6 Synergy with Other Airborne Sensors**

67 Miller et al. [2019] described the overall ABoVE Airborne Campaign design strategy and anticipated
68 airborne sensor synergies. Here, we highlight three SARs and a LiDAR – AirSWOT (NASA), F-SAR
69 (DLR), LS-ASAR (ISRO) and LVIS (NASA) – whose acquisitions in the ABoVE domain were
70 specifically designed to complement and leverage the ABoVE L- and/or P-band SAR acquisitions.
71 Many other airborne sensor synergies are being exploited by the ABoVE science team and are reported
72 separately.

73 **6.1 AirSWOT**

74 NASA's AirSWOT airborne instrument suite has been developed to support the Surface Water and
75 Ocean Topography (SWOT) mission. The heart of AirSWOT is the Ka-band SWOT Phenomenology
76 Airborne Radar (KaSPAR). KaSPAR collects two swaths of across-track interferometry data: one swath
77 from nadir to 1 km and a second swath that extends from 1 km to 5 km off-nadir. AirSWOT flight lines
78 for ABoVE were designed to center the AirSWOT swath on the center of the P-band swath for
79 maximum overlap. KaSPAR is complemented by a high-resolution color-infrared (CIR) Digital Camera
80 System [Kyzivat et al. 2019a,b] and a Precision Inertial Measurement Unit (IMU) for accurate attitude
81 and positioning information. In 2015 AirSWOT made pre-ABoVE deployments to the Tanana River
82 Valley [Altenau et al. 2017] and the Yukon Flats [Pitcher et al. 2019a, 2019b] in Region A3.

83
84 In 2017, AirSWOT deployed to acquire
85 early season (May-June) and late season
86 (August) WSEs across the ABoVE
87 domain. **Figure 7** shows the concentration
88 of AirSWOT lines in wetlands complexes
89 in the boreal forest, across the Canadian
90 Shield, along the Mackenzie River Valley,
91 and into the Arctic tundra. All of these
92 regions contain overlapping L- and P-
93 band acquisitions. Of special interest are
94 the lines in the Peace-Athabasca Delta
95 (36000: PADeE and 18035: PADeW)
96 and the Yukon Flats (21508: YFlatW,
97 21609: YflatE, and 04707: FtYuko) and
98 Trail Valley Creek, NT (01703: TukHwy)
99 where extensive on-water measurements
00 were made [Pitcher et al. 2020]. Future
01 joint analyses of the Ka- and L-band data
02 will highlight the advances possible in
03 pan-Arctic hydrology from the upcoming NISAR and SWOT missions.




Figure 7. AirSWOT flight lines acquired during the 2017 ABoVE airborne campaign sampled wetlands ranging from the Arctic Ocean coast to the southern boreal forest. AirSWOT's Ka-band acquisitions were designed to overlap with the L- and P-band SAR near-field acquisitions (See Fig. 1). © Google Earth

14
15
16

17 6.2 F-SAR

18 The German Space Agency (DLR) developed
19 the F-SAR instrument as an advanced
20 airborne SAR testbed for technology and
21 remote sensing applications [Reigber et al.
22 2013]. F-SAR operates fully polarimetric at
23 X-, C-, S-, L- and P-bands and features
24 single-pass polarimetric interferometric SAR
25 (PolInSAR) capabilities in X- and S-bands
26 [Reigber et al. 2013]. The radar covers an
27 off-nadir angle range of 25 to 60 degrees and
28 provides sub-meter scale spatial resolution
29 from flight altitudes up to 6000 mAGL.

30
31 During August 2018 and April 2019, F-SAR
32 was deployed to northern Canada as part of
33 DLR's permafrost airborne SAR experiment
34 (PerMASAR). It was configured in X-, C-, S-
35 and L-band mode and flew onboard a Dornier
36 Do 228-212 research aircraft. Measurements
37 were acquired from ~4500 mAGL.

38 Coordinated tomoSAR transects were flown
39 over the BERMS site in the southern boreal
40 forest on 18 August (UAVSAR) and 23
41 August (F-SAR). Preliminary results [Hensley
42 et al. 2020] are summarized in Section 6. F-
43 SAR also acquired data over the Scotty Creek
44 watershed, flux towers, and AOIs (Figure 8),
45 the Smith Creek flux tower (Wrigley, NT),
46 Baker Lake, Havipak Creek, Trail Valley Creek,
47 and Herschel Island, providing extensive opportunities to cross-
48 compare F-SAR and the ABoVE SAR
49 acquisitions.

50




Figure 8. Overlap of the F-SAR acquisition at Scotty Creek, NT (yellow box) and the ABoVE L-band SAR line 16713 (red box & polarized SAR false color). The ABoVE line also captures Scotty Creek AOIs 2 – 6 (see Region C3 details, Sec. 5.3). The SAR data will complement and benefit from the extensive ground-based data acquired in this area [Quinton et al. 2019]. © Google Earth

51 6.3 LS-ASAR (ISRO)

52 The Indian Space Research Organisation (ISRO)
53 and NASA are jointly developing the NASA-
54 ISRO Synthetic Aperture Radar (NISAR), which
55 will map Earth's surface in L-band and S-band
56 every 12 days [Rosen et al. 2017]. As a precursor
57 to the NISAR mission, ISRO has developed a L-
58 and S-Band-Airborne SAR (LS-ASAR) to
59 prepare the community to maximize the scientific
60 and societal benefits of NISAR data [Ramanujam
61 et al. 2016; 2019; Mehra et al. 2019]. LS-ASAR
62 operates in Dual, Quad, and Hybrid Polarization
63 modes in both L- and S-bands. It covers
64 incidence angles from 24°-77° with swaths
65 ranging from 5.5 km to 15 km.

66
67 In December 2019 LS-ASAR flew a series of
68 Arctic sea ice sorties from Fairbanks, AK. During
69 this deployment, LS-ASAR also acquired data
70 over a number of the ABoVE flight lines in
71 Regions A1 (North Slope) and A3 (Eastern
72 Interior) as well as over a number of glacier sites
73 in the Alaska Range. The acquisitions are
74 available via the NASA & ISRO ASAR
75 Campaign page ([https://uavsar.jpl.nasa.gov/cgi-
76 bin/deployment.pl?id=L20191101](https://uavsar.jpl.nasa.gov/cgi-bin/deployment.pl?id=L20191101)) and are summarized in Figure 9. These data provide snow-on
77 coverage that was a known deficiency of previous ABoVE airborne campaigns. Additionally, the LS-
78 ASAR data extend coverage of these regions to S-band.

80 6.4 LVIS

81 The Land, Vegetation, and Ice Sensor (LVIS) is an airborne, full waveform scanning laser altimeter
82 which produces topographic maps with decimeter accuracy as well as vegetation vertical height and
83 structure measurements [Blair et al. 1999a,b]. Flight lines for LVIS (~1.4 km swath) were slaved to the
84 centerline of the P-band swath during ABoVE, except where deviations were required to capture critical
85 ground sites. LVIS-C (classic configuration) was deployed in 2017 aboard a B-200 and achieved limited
86 coverage (Figure 10, left panel). During 2019, the new LVIS Facility instrument (LVIS-F) as well as
87 LVIS-C were deployed on the NASA Gulfstream-V and achieved coverage of all legacy SAR lines
88 (Figure 10, right panel).





Figure 9. L- and S-band SAR lines acquired over Alaska during the December 2019 ASAR Campaign. ASAR flight lines on the North Slope, in the Yukon Flats, and in the Western Interior exactly overlap ABoVE flight lines. These early winter acquisitions provide a preliminary look at cold season SAR data that will be explored in greater detail in the planned ABoVE-SnowEx campaign. © Google Earth

90 LVIS' unique capability for measuring the sub-meter topography beneath boreal forest canopies
91 complemented the tomoSAR acquisitions over Delta Junction, AK and the BERMS site in northern
92 Saskatchewan [Hensley et al. 2020; Section 6]. LVIS altimetry will also prove valuable in analyses of
93 such variables as permafrost degradation, active layer thickness, and water surface elevation; however,
94 LVIS' significantly narrower swath limits the spatial extent over which these analyses may be
95 performed.

96
97 In June-July 2017, the NASA LVIS Facility was deployed to sites in northern Canada and Alaska as
98 part of NASA's Arctic-Boreal Vulnerability Experiment (ABoVE) 2017 airborne campaign. During the
99 4-week deployment of LVIS-F, a total of 15 flights were flown over diverse science targets based out of
00 multiple airports in Canada and Alaska. Data are available in both Level1B and Level2 formats (**Table**
01 **2**). The Level1b data files contain the geolocated laser waveform data for each laser footprint. The
02 Level2 data files contain canopy top and ground elevations and relative heights derived from the
03 Level1b data. ABoVE LVIS L1B Geolocated Return Energy Waveforms, Version 1 [Blair and Hofton,
04 2018a] and L2 Geolocated Surface Elevation Product, Version 1 [Blair and Hofton, 2018b] may be
05 obtained from the National Snow and Ice Data Center via <https://doi.org/10.5067/UMRAWS57QAFU>
06 and <https://doi.org/10.5067/IA5WAX7K3YGY>, respectively.

07



08
09 **Figure 10. Flight lines for the LVIS 2017 flights (left) and 2019 flights (right) were designed to overlap with the near-filed portions**
10 **of the L-band and P-band SAR swaths to maximize opportunities for synergistic science. Aircraft and weather limited coverage**
11 **during the 2017 campaign, but complete coverage of the SAR flight lines was achieved in 2019. These data will enable studies of**
12 **SAR/LiDAR fusion over the Arctic-boreal regions as a precursor to NISAR/IceSat-2 studies. © National Geographic Society**

13

14 **Table 2. 2017 LVIS Data Products**

LVIS Data Products	Format
Flight Trajectories	KMZ
Camera Trajectories	KMZ
LVIS L1A Camera Imagery	JPG*
LVIS L1B Geolocated Waveforms	HDF, LDS 2.0.2
LVIS L2 Elevation and Height Products	ASCII TXT, LDS 2.0.2a

15

16 In July-August 2019, the NASA LVIS Facility and LVIS Classic were deployed to sites in northern
 17 Canada and Alaska as part of NASA's ABoVE 2019 airborne campaign. The increased range and
 18 endurance of the Gulfstream-V platform enabled extensive sampling, including: all L-band SAR lines,
 19 multiple IceSAT-2 underflights, and numerous ABoVE field sites. The available data products are given
 20 in **Table 3**.

21

22 **Table 3. 2019 LVIS Data Products**

LVIS Data Products	Format
Flight Trajectories	KMZ
Coverage Maps	KMZ
LVIS Classic L1B Geolocated Waveforms	HDF, LDS 2.0.3
LVIS Classic L2 Elevation and Height Products	ASCII TXT, LDS 2.0.3
LVIS Facility L1B Geolocated Waveforms	HDF, LDS 2.0.3
LVIS Facility L2 Elevation and Height Products	ASCII TXT, LDS 2.0.3

23 **6.5 G-LiHT**

24 Zhao et al. [2022] used ABoVE airborne L- and P-band SAR to map boreal forest species and canopy
 25 height in the Tanana Valley State Forest (TVSF) near Delta Junction, AK. They employed machine
 26 learning (random forests) to train separate regression models for canopy height mapping and a
 27 classification model for forest species mapping. Data derived from NASA's Goddard LiDAR,
 28 Hyperspectral, and Thermal Imager (G-LiHT) system [Cook et al. 2013] were treated as ground truth
 29 for the canopy height model (CHM). Forest species prediction were referenced against (TVSF) Timber
 30 Inventory and Forest Inventory and Analysis (FIA) data. The experimental results show the proposed
 31 method yields a root-mean-square error of 1.90 m for forest height estimation and overall accuracy of
 32 79.5% for forest species classification. A significant finding was that PolSAR decomposition
 33 parameters, such as volume scattering and entropy, strongly influenced the canopy height estimates.
 34 Interestingly, topography played a crucial role in the species classification.

35

36

37

38 7 Data Availability

39

40 Links to the ABoVE L- and P-band SAR products, supporting data, derived products, and ancillary
41 measurements are provided in the Appendix. Formal citations to all DOIs are provided in the
42 References. The L- and P-band SAR data may be found at the JPL UAVSAR data portal,
43 <https://uavsar.jpl.nasa.gov/cgi-bin/data.pl>. L-band data may also be accessed via the UAVSAR portal at
44 the Alaska SAR Facility (ASF) DAAC (<https://asf.alaska.edu/data-sets/sar-data-sets/uavsar/>) while
45 AirMOSS P-band data may be accessed via the ORNL DAAC, https://daac.ornl.gov/get_data/#projects,
46 select “AirMOSS”.

47

48 Miller et al. [2023; <https://doi.org/10.3334/ORNLDAAC/2150>] provides a detailed description of all 80
49 SAR flight lines and how each fits into the ABoVE experimental design. Extensive maps, tables, and
50 hyperlinks give direct access to every flight plan as well as individual flight lines. It is a guide to enable
51 interested readers to fully explore the ABoVE L- and P-band SAR data.

52

53

54 8 Summary

55 The ABoVE project conducted airborne L-band PolInSAR surveys in 2017, 2018, 2019 and 2022 across
56 Alaska and northwestern Canada. These were complemented by a P-band PolInSAR survey in 2017
57 along the same transects. This time series provides a powerful data set with which to evaluate the state
58 of permafrost, active layer thickness, soil moisture, boreal forest structure, above ground biomass, and
59 water surface elevation. Additional studies leverage the PolInSAR data to address fire disturbance and
60 recovery, thermokarst feature development, and retrogressive permafrost thaw megaslumps. Many of
61 these analyses are in progress and will be published separately.

62

63 Miller et al. [2023] provides extensive, fully hyperlinked notes on the airborne SAR data. Researchers
64 may discover these data via daily sorties and/or individual flight lines. Alternatively, they may be
65 explored via the interactive map at the JPL UAVSAR data portal, ([https://uavsar.jpl.nasa.gov/cgi-
66 bin/data.pl](https://uavsar.jpl.nasa.gov/cgi-bin/data.pl)) which provides links to the individual flight line data, maps, and related flight plans that
67 acquired data over one or more of the individual flight lines. We have also identified the ground-based
68 anchor points for each flight line to facilitate comparisons with those data. Calibration and validation
69 data sets as well as many derived products produced by the ABoVE Science Team may be found at the
70 Arctic Boreal Vulnerability Experiment (ABoVE) landing page at the ORNL DAAC
71 (https://daac.ornl.gov/cgi-bin/dataset_lister.pl?p=34).

72

73 The example studies (Sections 4 and 5) and multi-instrument synergies (Section 6) described here are
74 only a small portion of the studies currently being undertaken by the ABoVE Science Team and the
75 SAR Working Group. We anticipate many new and innovative uses of the L-band and P-band SAR data
76 as the ABoVE team expands its range of synthesis activities in Phase 3.

77

78 The ABoVE L-band SAR flights planned for 2020 and 2021 were postponed due to the global COVID-
79 19 pandemic and safety considerations; however, flights were resumed in 2022 and we anticipate at
80 least one more thaw season campaign in 2024. Finally, we note that the data and analyses discussed
81 here set the stage for the upcoming NISAR mission (expected launch in 2023). NISAR will deliver
82 global L- and S-band imagery with a 12-day revisit. Its emphasis on snow- and ice-covered surfaces has
83 obvious applications in the ABoVE domain, and its global coverage will allow researchers to test the
84 methods developed for the ABoVE domain across the pan-Arctic. Beyond NISAR, NASA is studying
85 architectures for the Surface Deformation and Change (SDC) Earth System Observatory Mission and
86 ESA are developing the Rose-L Copernicus expansion mission. SDC and Rose-L are also envisioned as
87 an L-band sensors. NISAR, SDC, and Rose-L will all benefit from the ABoVE SAR studies,
88 algorithmic advances, and lessons learned.

89 **9 Supplemental Information**

90 The Supplemental Information (SI) contains detailed descriptions of all L-band flight lines plus tables
91 with hyperlinks to all L-band lines and sorties. Additionally, the SI includes tables and links to the P-
92 band flight lines acquired during the ABoVE campaigns and to all Legacy L-band and P-band flight
93 lines. The SI is identical to the file <Summary ABoVE L- & P-Band SAR Surveys - hyperlinked.pdf>
94 that may be found in the /data folder of the uncompressed data download from Miller et al. [2023;
95 <https://doi.org/10.3334/ORNLDAAC/2150>].
96

97 **10 Author Contributions**

98 SJG, CEM, PCG and the ABoVE Science Definition Team developed the preliminary ABoVE
99 Implementation Plan; this was updated by SJG, CEM, PCG and the ABoVE Science Team. CEM, PCG,
00 and EH developed the initial flight lines based on the ABoVE Implementation Plan and consultations
01 with the ABoVE Science and partners. NSP translated the notional flight lines into the UAVSAR
02 planning system. NSP, YL, MM, PCG, , ELH, and CEM served as Scientist on Board during the data
03 acquisition flights. YL, SH, BC, NP and the JPL Sub-orbital Radar Science and Engineering Team
04 (334F) processed the L- and P-band SAR data. CW, KS, MT, JB, LBC, RHC, MM, SW, MM, RJM,
05 TL, LJ, PS AT, and RD collected cal/val field data. CW and SW coordinated ground cal/val data
06 acquisitions at the NGEE-Arctic sites. SH, PS, NSP performed the tomoSAR analyses and SH
07 contributed the text and images for Section 6. KS, MM and the ABoVE SAR Working Group calibrated
08 the L-band SAR data. MM, AT, RHC processed the P-band data. NSP and YL coordinated the LS-
09 ASAR flights and the joint BERMS area tomoSAR flights with F-SAR. DS coordinated all data product
10 submissions to the ORNL DAAC. CEM wrote the initial manuscript and all co-authors contributed to
11 the final version.

12 11 Competing Interests

13 The authors declare that they have no conflict of interest

14 12 Acknowledgments

15 The L- and P-band SAR data acquisitions would not have been possible without the indefatigable
16 support of our NASA pilots and flight crews. We thank John McGrath and the NASA AFRC C-20
17 (N30502) team as well as Derek Rutovic and the NASA JSC G-III (N995NA) team. We also thank the
18 instrument scientists, operators, and data processing team from the JPL Suborbital Radar Science and
19 Engineering Team (334F) who were essential to the successful execution of these experiments and rapid
20 processing of the resulting data products. The field work supporting the SAR campaigns was made
21 possible by the excellent support from Dan Hodkinson, Sarah Sackett, and the ABoVE Logistics Office.
22 Finally, we thank the data curation team at the Oak Ridge Distributed Active Archive Center for their
23 support and expert advice.

24

25 This work was supported by the NASA Terrestrial Ecology Program's Arctic-Boreal Vulnerability
26 Experiment (ABoVE). A portion of this work was performed at the Jet Propulsion Laboratory,
27 California Institute of Technology, under contract with National Aeronautics and Space Administration
28 (80NM0018D0004). Government funding acknowledged.

29

30 References

- 31 Abe, T., Iwahana, G., Efremov, P. V., Desyatkin, A. R., Kawamura, T., Fedorov, A., Zhegusov, Y., Yanagiya,
32 K., and Tadono, T.: Surface displacement revealed by L-band InSAR analysis in the Mayya area, Central
33 Yakutia, underlain by continuous permafrost, *Earth Planets Space*, 72, 138, <https://doi.org/10.1186/s40623-020-01266-3>, 2020.
- 34 Airborne Imaging: Final LiDAR processing & vertical accuracy report: Prepared for the U.S. Fish and Wildlife
35 Service, in: *LiDAR Imagery & DEM Model for Yukon Delta National Wildlife Refuge—Near Angyaravak
36 Bay, Alaska*, Airborne Imaging, Calgary, AB, Canada, 28, 2011.
- 37 Allen, B. D., Braun, S. A., Crawford, J. H., Jensen, E. J., Miller, C. E., Moghaddam, M., and Maring, H.:
38 Proposed investigations from NASA's Earth Venture-1 (EV-1) airborne science selections, in: 2010 IEEE
39 International Geoscience and Remote Sensing Symposium, 2010 IEEE International Geoscience and Remote
40 Sensing Symposium, 2575–2578, <https://doi.org/10.1109/IGARSS.2010.5651920>, 2010.
- 41 Alonzo, M., Dial, R. J., Schulz, B. K., Andersen, H.-E., Lewis-Clark, E., Cook, B. D., and Morton, D. C.:
42 Mapping tall shrub biomass in Alaska at landscape scale using structure-from-motion photogrammetry and
43 lidar, *Remote Sens. Environ.*, 245, 111841, <https://doi.org/10.1016/j.rse.2020.111841>, 2020.
- 44 Altenau, E. H., Pavelsky, T. M., Moller, D., Lion, C., Pitcher, L. H., Allen, G. H., Bates, P. D., Calmant, S.,
45 Durand, M., and Smith, L. C.: AirSWOT measurements of river water surface elevation and slope: Tanana
46 River, AK, *Geophys. Res. Lett.*, 44, 181–189, <https://doi.org/10.1002/2016GL071577>, 2017.
- 47 Kenaston / Brightwater Creek Mesonet Site, SK - CCRNetwork:
48 <http://www.ccrnetwork.ca/science/WECC/prairie/brightwater-creek.php>, last access: 30 April 2021.
- 49

- 50 Bakian-Dogaheh, K., Chen, R. H., Moghaddam, M., and Tabatabaenejad, A.: Electromagnetic scattering
51 behavior of a new organic soil dielectric model for long-wavelength radar retrieval of permafrost active layer
52 soil properties, 2020 IEEE International Geoscience and Remote Sensing Symposium, 2020.
- 53 Bakian-Dogaheh, K., R.H. Chen, M. Moghaddam, Y. Yi, and A. Tabatabaenejad. 2020. ABoVE:
54 Active Layer Soil Characterization of Permafrost Sites, Northern Alaska, 2018. ORNL DAAC, Oak
55 Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1759>
- 56 Banks, S. N., Ullmann, T., Duffe, J., Roth, A., King, D., Demers, A.-M., Hogg, A., Schmitt, A., Baumhauer, R.,
57 and Dech, S.: Multi-frequency analysis of high resolution quad-pol Radarsat-2 and dual-pol TerraSAR-X
58 data for land cover classification in Arctic Coastal Ecosystems, Mackenzie Delta, beaufort sea, 3548–3551,
59 <https://doi.org/10.1109/IGARSS.2012.6350653>, 2012.
- 60 Banks, S. N., King, D. J., Merzouki, A., and Duffe, J.: Characterizing Scattering Behaviour and Assessing
61 Potential for Classification of Arctic Shore and Near-Shore Land Covers with Fine Quad-Pol RADARSAT-2
62 Data, *Can. J. Remote Sens.*, 40, 291–314, <https://doi.org/10.1080/07038992.2014.979487>, 2014.
- 63 Barr, A. G., Morgenstern, K., Black, T. A., McCaughey, J. H., and Nestic, Z.: Surface energy balance closure by
64 the eddy-covariance method above three boreal forest stands and implications for the measurement of the
65 CO₂ flux, *Agric. For. Meteorol.*, 140, 322–337, <https://doi.org/10.1016/j.agrformet.2006.08.007>, 2006.
- 66 Bennett, A. J., Thompson, W. L., and Mortenson, D. C.: Vital Signs Monitoring Plan Southwest Alaska Network:
67 Inventory and Monitoring Program, *Natl. Park Serv.*, 2006.
- 68 Blair, J. B and Hofton, M.: ABoVE LVIS L2 Geolocated Surface Elevation Product, Version 1,
69 <https://doi.org/10.5067/IA5WAX7K3YGY>, 2018.
- 70 Blomberg, E., Ulander, L. M. H., Tebaldini, S., and Ferro-Famil, L.: Evaluating P-Band TomoSAR for Biomass
71 Retrieval in Boreal Forest, *IEEE Trans. Geosci. Remote Sens.*, 59, 3793–3804,
72 <https://doi.org/10.1109/TGRS.2020.3020775>, 2021.
- 73 Bourgeau-Chavez, L. L., French, N. H. F., Grelick, S. E., Jenkins, L., Battaglia, M., Serocki, E., and Billmire, M.:
74 ABoVE: Burn Severity, Fire Progression, Landcover and Field Data, NWT, Canada, 2014, ORNL DAAC,
75 <https://doi.org/10.3334/ORNLDAAAC/1307>, 2016.
- 76 Bourgeau-Chavez, L. L., Grelick, S. E., Jenkins, L., Battaglia, M., Serocki, E., and Billmire, M.: ABoVE: Burn
77 Severity, Fire Progression, and Field Data, NWT, Canada, 2015-2016, ORNL DAAC,
78 <https://doi.org/10.3334/ORNLDAAAC/1548>, 2017.
- 79 Bourgeau-Chavez, L. L., Graham, J. A., Grelick, S. E., French, N. H. F., Battaglia, M., Hansen, D., and Tanzer,
80 D.: ABoVE: Ecosystem Map, Great Slave Lake Area, Northwest Territories, Canada, 1997-2011, ORNL
81 DAAC, <https://doi.org/10.3334/ORNLDAAAC/1695>, 2019a.
- 82 Bourgeau-Chavez, L. L., Battaglia, M., Kane, E. S., Cohen, L. M., and Tanzer, D.: ABoVE: Post-Fire and
83 Unburned Vegetation Community and Field Data, NWT, Canada, 2018, ORNL DAAC,
84 <https://doi.org/10.3334/ORNLDAAAC/1703>, 2019b.
- 85 Burns, L. E., Fugro Airborne Survey Corp., and Stevens Exploration Management Corp.: Final processed
86 database for the airborne geophysical surveys of the Alaska Highway corridor, east-central Alaska: Alaska
87 Division of Geological & Geophysical Surveys Geophysical Report 2006-8, 1 DVD.,
88 <https://doi.org/10.14509/17761>, 2008.
- 89 Campbell, E. M., Antos, J. A., and vanAkker, L.: Resilience of southern Yukon boreal forests to spruce beetle
90 outbreaks, *For. Ecol. Manag.*, 433, 52–63, <https://doi.org/10.1016/j.foreco.2018.10.037>, 2019.
- 91 Chapin, E., Chau, A., Chen, J., Heavey, B., Hensley, S., Lou, Y., Machuzak, R., and Moghaddam, M.: AirMOSS:
92 An Airborne P-band SAR to measure root-zone soil moisture, in: 2012 IEEE Radar Conference, 2012 IEEE
93 Radar Conference, 0693–0698, <https://doi.org/10.1109/RADAR.2012.6212227>, 2012.

- 94 Chapin, E., Flores, S., Harcke, L., Hawkins, B. P., Hensley, S., Michel, T. R., Muellerschoen, R. J., Shimada, J.
95 G., Tung, W. W., and Veeramachaneni, C.: AirMOSS: L1 S-0 Polarimetric Data from AirMOSS P-band
96 SAR, BERMS, Canada, 2012–2015, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1406>, 2018.
- 97 Chen, A. C., Parsekian, A. D., Schaefer, K., Jafarov, E. E., Panda, S. K., Liu, L., Zhang, T., and Zebker, H. A.:
98 Pre-ABoVE: Ground-penetrating Radar Measurements of ALT on the Alaska North Slope, ORNL DAAC,
99 <https://doi.org/10.3334/ORNLDAAC/1265>, 2015.
- 00 Chen, R. H., Tabatabaenejad, A., and Moghaddam, M.: A time-series active layer thickness retrieval algorithm
01 using P- and L-band SAR observations, 2016 IEEE International Geoscience and Remote Sensing
02 Symposium (IGARSS), 3672–3675, <https://doi.org/10.1109/IGARSS.2016.7729951>, 2016.
- 03 Chen, R. H., Tabatabaenejad, A., and Moghaddam, M.: Retrieval of permafrost active layer properties using P-
04 band airmoss and L-band UAVSAR data, 2017 IEEE International Geoscience and Remote Sensing
05 Symposium (IGARSS), 1415–1418, <https://doi.org/10.1109/IGARSS.2017.8127230>, 2017.
- 06 Chen, R. H., Tabatabaenejad, A., and Moghaddam, M.: P-Band Radar Retrieval of Permafrost Active Layer
07 Properties: Time-Series Approach and Validation with In-Situ Observations, IGARSS 2018 - 2018 IEEE
08 International Geoscience and Remote Sensing Symposium, 6777–6779,
09 <https://doi.org/10.1109/IGARSS.2018.8518179>, 2018.
- 10 Chen, R. H., Tabatabaenejad, A., and Moghaddam, M.: ABoVE: Active Layer and Soil Moisture Properties
11 from AirMOSS P-band SAR in Alaska, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1657>, 2019a.
- 12 Chen, R. H., Bakian-Dogaheh, K., Tabatabaenejad, A., and Moghaddam, M.: Modeling and Retrieving Soil
13 Moisture and Organic Matter Profiles in the Active Layer of Permafrost Soils From P-Band Radar
14 Observations, IGARSS 2019 - 2019 IEEE International Geoscience and Remote Sensing Symposium,
15 10095–10098, <https://doi.org/10.1109/IGARSS.2019.8899802>, 2019b.
- 16 Chen, R. H., Tabatabaenejad, A., and Moghaddam, M.: Retrieval of Permafrost Active Layer Properties Using
17 Time-Series P-Band Radar Observations, *IEEE Trans. Geosci. Remote Sens.*, *57*, 6037–6054,
18 <https://doi.org/10.1109/TGRS.2019.2903935>, 2019c.
- 19 Chen, R. H., Michaelides, R. J., Sullivan, T. D., Parsekian, A. D., Zebker, H. A., Moghaddam, M., and Schaefer,
20 K. M.: Joint Retrieval of Soil Moisture and Permafrost Active Layer Thickness Using L-Band Insar and P-
21 Band Polsar, IGARSS 2020 - 2020 IEEE International Geoscience and Remote Sensing Symposium, 4606–
22 4609, <https://doi.org/10.1109/IGARSS39084.2020.9324660>, 2020a.
- 23 Chen, R. H., Pinto, N., Duan, X., Tabatabaenejad, A., and Moghaddam, M.: Mapping Tree Canopy Cover and
24 Canopy Height with L-Band SAR Using LiDAR Data and Random Forests, IGARSS 2020 - 2020 IEEE
25 International Geoscience and Remote Sensing Symposium, 4136–4139,
26 <https://doi.org/10.1109/IGARSS39084.2020.9323738>, 2020b.
- 27 Chen, R. H., Michaelides, R. J., Sullivan, T. D., Parsekian, A. D., Zebker, H. A., Moghaddam, M., & Schaefer,
28 K. M. (2021a). Joint retrieval of soil moisture and activere layer thickness in the northern circumpolar
29 permafrost region using l-band InSAR and p-band porsar. *Earth and Space Science*.
- 30 Chen, Richard H., Roger J. Michaelides, Yuhuan Zhao, Lingcao Huang, Elizabeth Wig, Taylor D. Sullivan,
31 Andrew D. Parsekian, Howard A. Zebker, Mahta Moghaddam, and Kevin M. Schaefer. "Permafrost
32 Dynamics Observatory: Retrieval of Active Layer Thickness and Soil Moisture from Airborne Insar and
33 Polsar Data." In *2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS*, pp. 1444-
34 1447. IEEE, 2021b. DOI: 10.1109/IGARSS47720.2021.9554288
- 35 Colliander, A., Cosh, M. H., Misra, S., Jackson, T. J., Crow, W. T., Powers, J., McNairn, H., Bullock, P., Berg,
36 A., Magagi, R., Gao, Y., Bindlish, R., Williamson, R., Ramos, I., Latham, B., O'Neill, P., and Yueh, S.:
37 Comparison of high-resolution airborne soil moisture retrievals to SMAP soil moisture during the SMAP
38 validation experiment 2016 (SMAPVEX16), *Remote Sens. Environ.*, *227*, 137–150,
39 <https://doi.org/10.1016/j.rse.2019.04.004>, 2019.

- 40 Cook, B. D., L. W. Corp, R. F. Nelson, E. M. Middleton, D. C. Morton, J. T. McCorkel, J. G. Masek, K. J.
41 Ranson, V. Ly, and P. M. Montesano (2013): NASA Goddard's Lidar, Hyperspectral and Thermal (G-LiHT)
42 airborne imager. *Remote Sensing* 5:4045-4066, doi:10.3390/rs5084045.
- 43 Cooley, S. W., Smith, L. C., Stepan, L., and Mascaro, J.: Tracking Dynamic Northern Surface Water Changes
44 with High-Frequency Planet CubeSat Imagery, *Remote Sens.*, 9, 1306, <https://doi.org/10.3390/rs9121306>,
45 2017.
- 46 Crasto, N., Hopkinson, C., Forbes, D. L., Lesack, L., Marsh, P., Spooner, I., and van der Sanden, J. J.: A LiDAR-
47 based decision-tree classification of open water surfaces in an Arctic delta, *Remote Sens. Environ.*, 164, 90-
48 102, <https://doi.org/10.1016/j.rse.2015.04.011>, 2015.
- 49 Danby, R.: A Muniscale Study of Tree-Line Dynamics in Southwestern Yukon, Arctic, 56, 427-429, 2003.
- 50 Danby, R. K., Williams, A., and Hik, D. S.: Fifty Years of Science at the Kluane Lake Research Station, Arctic,
51 67, iii-viii, <https://doi.org/10.14430/arctic4398>, 2014.
- 52 Davey, C. A., Redmond, K. T., and Simeral, D. B.: Weather and Climate Inventory National Park Service
53 Southwest Alaska Network, National Park Service, Fort Collins, Colorado, 2007.
- 54 Davidson, S. J., Santos, M. J., Sloan, V. L., Watts, J. D., Phoenix, G. K., Oechel, W. C., and Zona, D.: Mapping
55 Arctic Tundra Vegetation Communities Using Field Spectroscopy and Multispectral Satellite Data in North
56 Alaska, USA, *Remote Sens.*, 8, 978, <https://doi.org/10.3390/rs8120978>, 2016.
- 57 Davidson, S. J., Santos, M. J., Sloan, V. L., Reuss-Schmidt, K., Phoenix, G. K., Oechel, W. C., and Zona, D.:
58 Upscaling CH₄ Fluxes Using High-Resolution Imagery in Arctic Tundra Ecosystems, *Remote Sens.*, 9, 1227,
59 <https://doi.org/10.3390/rs9121227>, 2017.
- 60 Dengel, S. and Torn, M.: NGEA Arctic CO₂, CH₄ and Energy Eddy-Covariance (EC) Flux Tower Auxiliary
61 Measurements, Council Road Mile Marker 71, Seward Peninsula, Alaska, 2017 - Ongoing,
62 <https://doi.org/10.5440/1526749>, 2020.
- 63 Douglas, T. A.: ABoVE: Soil Active Layer Thaw Depths at CRREL sites near Fairbanks, Alaska, 2014-2018,
64 ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1701>, 2019.
- 65 Douglas, T. A., Hiemstra, C. A., and Barker, A. J.: ABoVE: End of Season Snow Depth at CRREL sites near
66 Fairbanks, Alaska, 2014-2019, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1702>, 2019.
- 67 Douglas, T. A., Turetsky, M. R., and Koven, C. D.: Increased rainfall stimulates permafrost thaw across a variety
68 of Interior Alaskan boreal ecosystems, *Npj Clim. Atmospheric Sci.*, 3, 1-7, [https://doi.org/10.1038/s41612-
69 020-0130-4](https://doi.org/10.1038/s41612-020-0130-4), 2020.
- 70 Douglas, T. A., Hiemstra, C. A., Anderson, J. E., Barbato, R. A., Bjella, K. L., Deeb, E. J., Gelvin, A. B., Nelsen,
71 P. E., Newman, S. D., Saari, S. P., and Wagner, A. M.: Recent degradation of Interior Alaska permafrost
72 mapped with ground surveys, geophysics, deep drilling, and repeat airborne LiDAR, *Cryosphere Discuss.*, 1-
73 39, <https://doi.org/10.5194/tc-2021-47>, 2021.
- 74 Du, J., Kimball, J. S., and Moghaddam, M.: Theoretical Modeling and Analysis of L- and P-band Radar
75 Backscatter Sensitivity to Soil Active Layer Dielectric Variations, *Remote Sens.*, 7, 9450-9472,
76 <https://doi.org/10.3390/rs70709450>, 2015.
- 77 Engstrom, R., Hope, A., Kwon, H., Stow, D., and Zamolodchikov, D.: Spatial distribution of near surface soil
78 moisture and its relationship to microtopography in the Alaskan Arctic coastal plain, *Hydrol. Res.*, 36, 219-
79 234, <https://doi.org/10.2166/nh.2005.0016>, 2005.
- 80 Euskirchen, E. S. and Edgar, C.: FLUXNET-CH₄ US-BZB Bonanza Creek Thermokarst Bog, ,
81 <https://doi.org/10.18140/FLX/1669668>, 2014a.
- 82 Euskirchen, E. S. and Edgar, C.: FLUXNET-CH₄ US-BZF Bonanza Creek Rich Fen, ,
83 <https://doi.org/10.18140/FLX/1669669>, 2014b.
- 84 Euskirchen, E. S. and Edgar, C.: FLUXNET-CH₄ US-BZS Bonanza Creek Black Spruce, ,
85 <https://doi.org/10.18140/FLX/1669670>, 2015.

- 86 Euskirchen, E. S., Bret-Harte, M. S., Scott, G. J., Edgar, C., and Shaver, G. R.: Seasonal patterns of carbon
87 dioxide and water fluxes in three representative tundra ecosystems in northern Alaska, *Ecosphere*, 3, art4,
88 <https://doi.org/10.1890/ES11-00202.1>, 2012.
- 89 Fayne, J. V., Smith, L. C., Pitcher, L. H., Kyzivat, E. D., Cooley, S. W., Cooper, M. G., Denbina, M. W., Chen,
90 A. C., Chen, C. W., and Pavelsky, T. M.: Airborne observations of arctic-boreal water surface elevations
91 from AirSWOT Ka-Band InSAR and LVIS LiDAR, *Environ. Res. Lett.*, 15, 105005,
92 <https://doi.org/10.1088/1748-9326/abadcc>, 2020.
- 93 French, N. H. F., Bourgeau-Chavez, L. L., and Chapman, B.: Wetland monitoring in high northern latitudes for
94 carbon and habitat assessment using synthetic aperture radar, in: AGU Fall Meeting Abstracts, H31N-1951,
95 2019.
- 96 French, N. H. F., Graham, J. A., Bourgeau-Chavez, L. L., and Whitman, E.: ABoVE: Burn Severity of Soil
97 Organic Matter, Northwest Territories, Canada, 2014-2015, ORNL DAAC,
98 <https://doi.org/10.3334/ORNLDAAC/1694>, 2020a.
- 99 French, N. H. F., Graham, J., Whitman, E., Bourgeau-Chavez, L. L., French, N. H. F., Graham, J., Whitman, E.,
00 and Bourgeau-Chavez, L. L.: Quantifying surface severity of the 2014 and 2015 fires in the Great Slave Lake
01 area of Canada, *Int. J. Wildland Fire*, 29, 892–906, <https://doi.org/10.1071/WF20008>, 2020b.
- 02 Frost, G. V., Loehman, R. A., Nelson, P. R., and Paradis, D. P.: ABoVE: Vegetation Composition across Fire
03 History Gradients on the Y-K Delta, Alaska, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1772>,
04 2020.
- 05 Gamon, J. A., Huemmrich, K. F., Peddle, D. R., Chen, J., Fuentes, D., Hall, F. G., Kimball, J. S., Goetz, S., Gu,
06 J., McDonald, K. C., Miller, J. R., Moghaddam, M., Rahman, A. F., Roujean, J.-L., Smith, E. A., Walthall, C.
07 L., Zarco-Tejada, P., Hu, B., Fernandes, R., and Cihlar, J.: Remote sensing in BOREAS: Lessons learned,
08 *Remote Sens. Environ.*, 89, 139–162, <https://doi.org/10.1016/j.rse.2003.08.017>, 2004.
- 09 Goetz, SJ, Miller, CE, Griffith, PC, et al., An overview of NASA’s Arctic Boreal Vulnerability Experiment
10 (ABoVE): An integrated research campaign to assess ecosystem vulnerability and its implications within the
11 Arctic and boreal domain, *Environmental Research Letters* (ABoVE Special Collection), Manuscript ERL-
12 112259, in revision, 2022
- 13 Greaves, H. E., Vierling, L., Eitel, J., Boelman, N., Magney, T., Prager, C., and Griffin, K.: High-Resolution
14 Shrub Biomass and Uncertainty Maps, Toolik Lake Area, Alaska, 2013, ORNL DAAC,
15 <https://doi.org/10.3334/ORNLDAAC/1573>, 2018.
- 16 Gusmeroli, A., Liu, L., Schaefer, K., Zhang, T., Schaefer, T., and Grosse, G.: Active Layer Stratigraphy and
17 Organic Layer Thickness at a Thermokarst Site in Arctic Alaska Identified Using Ground Penetrating Radar,
18 *Arct. Antarct. Alp. Res.*, 47, 195–202, <https://doi.org/10.1657/AAAR00C-13-301>, 2015.
- 19 He, J., Loboda, T. V., Jenkins, L., and Chen, D.: Mapping fractional cover of major fuel type components across
20 Alaskan tundra, *Remote Sens. Environ.*, 232, 111324, <https://doi.org/10.1016/j.rse.2019.111324>, 2019.
- 21 He, J., Loboda, T. V., Jenkins, L., and Chen, D.: ABoVE: Distribution Maps of Wildland Fire Fuel Components
22 across Alaskan Tundra, 2015, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1761>, 2020.
- 23 He, J., Chen, D., Jenkins, L., and Loboda, T. V.: Impacts of wildfire and landscape factors on organic soil
24 properties in Arctic tussock tundra. *Environmental Research Letters.*, *Environ. Res. Lett.*, in review, n.d.
- 25 Hensley, S., Oveisgharan, S., Saatchi, S., Simard, M., Ahmed, R. and Haddad, Z., 2014. An error model
26 for biomass estimates derived from polarimetric radar backscatter. *IEEE Transactions on*
27 *Geoscience and Remote Sensing*, 52(7), pp.4065-4082.
- 28 Hensley, S., Lou, Y., Michel, T., Muellerschoen, R., Hawkins, B., Lavalley, M., Pinto, N., Reigber, A.
29 and Pardini, M., 2016, July. UAVSAR PolInSAR and tomographic experiments in Germany. In

- 30 *Geoscience and Remote Sensing Symposium (IGARSS), 2016 IEEE International* (pp. 7517-7520).
31 IEEE.
- 32 Hensley, S., Chapman, B., Lavalley, M., Hawkins, B., Riel, B., Michel, T., Muellerschoen, R., Lou, Y., and
33 Simard, M.: UAVSAR L-Band and P-Band Tomographic Experiments in Boreal Forests, IGARSS 2018 -
34 2018 IEEE International Geoscience and Remote Sensing Symposium, 8679–8682,
35 <https://doi.org/10.1109/IGARSS.2018.8518784>, 2018.
- 36 Hensley, S., Ahmed, R., Chapman, B., Hawkins, B., Lavalley, M., Pinto, N., Pardini, M., Papathanassiou, K.,
37 Siqueria, P., and Treuhaft, R.: Boreal Forest Radar Tomography at P, L and S-Bands at Berms and Delta
38 Junction, IGARSS 2020 - 2020 IEEE International Geoscience and Remote Sensing Symposium, 96–99,
39 <https://doi.org/10.1109/IGARSS39084.2020.9323337>, 2020.
- 40 Higuera, P. E., Barnes, J., Chipman, M. L., Urban, M., and Hu, F. S.: The Burning Tundra: A Look Back at the
41 Last 6,000 Years of Fire in the Noatak National Preserve, Northwestern Alaska (U.S. National Park Service),
42 *Alsk. Park Sci.*, 10, 36–41, 2011.
- 43 Hinzman, L. D., Deal, C. J., McGuire, A. D., Mernild, S. H., Polyakov, I. V., and Walsh, J. E.: Trajectory of the
44 Arctic as an integrated system, *Ecol. Appl.*, 23, 1837–1868, <https://doi.org/10.1890/11-1498.1>, 2013.
- 45 Hogg, E. H., Michaelian, M., Hook, T. I., and Undershultz, M. E.: Recent climatic drying leads to age-
46 independent growth reductions of white spruce stands in western Canada, *Glob. Change Biol.*, 23, 5297–
47 5308, <https://doi.org/10.1111/gcb.13795>, 2017.
- 48 Hogg, E. H. (Ted) H. H., Brandt, J. P. B. P., and Michaelian, M. M.: Impacts of a regional drought on the
49 productivity, dieback, and biomass of western Canadian aspen forests, *Can. J. For. Res.*,
50 <https://doi.org/10.1139/X08-001>, 2008.
- 51 Holloway, J. E., Lewkowicz, A. G., Douglas, T. A., Li, X., Turetsky, M. R., Baltzer, J. L., and Jin, H.: Impact of
52 wildfire on permafrost landscapes: A review of recent advances and future prospects, *Permafr. Periglac.*
53 *Process.*, 31, 371–382, <https://doi.org/10.1002/ppp.2048>, 2020.
- 54 Hopkinson, C., Crasto, N., Marsh, P., Forbes, D., and Lesack, L.: Investigating the spatial distribution of water
55 levels in the Mackenzie Delta using airborne LiDAR, *Hydrol. Process.*, 25, 2995–3011,
56 <https://doi.org/10.1002/hyp.8167>, 2011.
- 57 Horn, R., Jaeger, M., Keller, M., Limbach, M., Nottensteiner, A., Pardini, M., Reigber, A., and Scheiber, R.: F-
58 SAR - recent upgrades and campaign activities, in: 2017 18th International Radar Symposium (IRS), 2017
59 18th International Radar Symposium (IRS), 1–10, <https://doi.org/10.23919/IRS.2017.8008092>, 2017.
- 60 Hoy, E. E., Griffith, P., Miller, C. E., and Team, A. S.: ABoVE: Directory of Field Sites Associated with 2017
61 ABoVE Airborne Campaign, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1582>, 2018.
- 62 Hu, F. S., Higuera, P. E., Walsh, J. E., Chapman, W. L., Duffy, P. A., Brubaker, L. B., and Chipman, M. L.:
63 Tundra burning in Alaska: Linkages to climatic change and sea ice retreat, *J. Geophys. Res. Biogeosciences*,
64 115, <https://doi.org/10.1029/2009JG001270>, 2010.
- 65 Iwahana, G., Harada, K., Uchida, M., Tsuyuzaki, S., Saito, K., Narita, K., Kushida, K., and Hinzman, L. D.:
66 Geomorphological and geochemistry changes in permafrost after the 2002 tundra wildfire in Kougarak,
67 Seward Peninsula, Alaska, *J. Geophys. Res. Earth Surf.*, 121, 1697–1715,
68 <https://doi.org/10.1002/2016JF003921>, 2016a.
- 69 Iwahana, G., Uchida, M., Liu, L., Gong, W., Meyer, F. J., Guritz, R., Yamanokuchi, T., and Hinzman, L.: InSAR
70 Detection and Field Evidence for Thermokarst after a Tundra Wildfire, Using ALOS-PALSAR, *Remote*
71 *Sens.*, 8, 218, <https://doi.org/10.3390/rs8030218>, 2016b.
- 72 Jafarov, E. E., Parsekian, A. D., Schaefer, K., Liu, L., Chen, A. C., Panda, S. K., and Zhang, T.: Estimating active
73 layer thickness and volumetric water content from ground penetrating radar measurements in Barrow,
74 Alaska, *Geosci. Data J.*, 4, 72–79, <https://doi.org/10.1002/gdj3.49>, 2017.

- 75 Jenkins, L. K., Bourgeau-Chavez, L. L., French, N. H. F., Loboda, T. V., and Thelen, B. J.: Development of
76 Methods for Detection and Monitoring of Fire Disturbance in the Alaskan Tundra Using a Two-Decade Long
77 Record of Synthetic Aperture Radar Satellite Images, *Remote Sens.*, 6, 6347–6364,
78 <https://doi.org/10.3390/rs6076347>, 2014.
- 79 Johnston, C. E., Ewing, S. A., Harden, J. W., Varner, R. K., Wickland, K. P., Koch, J. C., Fuller, C. C., Manies,
80 K., and Jorgenson, M. T.: Effect of permafrost thaw on CO₂ and CH₄ exchange in a western Alaska peatland
81 chronosequence, *Environ. Res. Lett.*, 9, 085004, <https://doi.org/10.1088/1748-9326/9/8/085004>, 2014.
- 82 Jones, B. M., Kolden, C. A., Jandt, R., Abatzoglou, J. T., Urban, F., and Arp, C. D.: Fire Behavior, Weather, and
83 Burn Severity of the 2007 Anaktuvuk River Tundra Fire, North Slope, Alaska, *Arct. Antarct. Alp. Res.*, 41,
84 309–316, <https://doi.org/10.1657/1938-4246-41.3.309>, 2009.
- 85 Jones, B. M., Grosse, G., Arp, C. D., Miller, E., Liu, L., Hayes, D. J., and Larsen, C. F.: Recent Arctic tundra fire
86 initiates widespread thermokarst development, *Sci. Rep.*, 5, 15865, <https://doi.org/10.1038/srep15865>, 2015.
- 87 Jones, M. C., Harden, J., O'Donnell, J., Manies, K., Jorgenson, T., Treat, C., and Ewing, S.: Rapid carbon loss
88 and slow recovery following permafrost thaw in boreal peatlands, *Glob. Change Biol.*, 23, 1109–1127,
89 <https://doi.org/10.1111/gcb.13403>, 2017.
- 90 Jorgenson, M. T., Harden, J., Kanevskiy, M., O'Donnell, J., Wickland, K., Ewing, S., Manies, K., Zhuang, Q.,
91 Shur, Y., Striegl, R., and Koch, J.: Reorganization of vegetation, hydrology and soil carbon after permafrost
92 degradation across heterogeneous boreal landscapes, *Environ. Res. Lett.*, 8, 035017,
93 <https://doi.org/10.1088/1748-9326/8/3/035017>, 2013.
- 94 Karion, A., Sweeney, C., Miller, J. B., Andrews, A. E., Commane, R., Dinardo, S., Henderson, J. M., Lindaas, J.,
95 Lin, J. C., Luus, K. A., Newberger, T., Tans, P., Wofsy, S. C., Wolter, S., and Miller, C. E.: Investigating
96 Alaskan methane and carbon dioxide fluxes using measurements from the CARVE tower, *Atmospheric*
97 *Chem. Phys.*, 16, 5383–5398, <https://doi.org/10.5194/acp-16-5383-2016>, 2016.
- 98 Kobayashi, H., Hiroki, I., and Suzuki, R.: AmeriFlux US-Prr Poker Flat Research Range Black Spruce Forest,
99 Ver. 3-5, <https://doi.org/10.17190/AMF/1246153>, 2019.
- 00 Kokelj, S. V., Lacelle, D., Lantz, T. C., Tunnicliffe, J., Malone, L., Clark, I. D., and Chin, K. S.: Thawing of
01 massive ground ice in mega slumps drives increases in stream sediment and solute flux across a range of
02 watershed scales, *J. Geophys. Res. Earth Surf.*, 118, 681–692, <https://doi.org/10.1002/jgrf.20063>, 2013.
- 03 Kokelj, S. V., Tunnicliffe, J., Lacelle, D., Lantz, T. C., Chin, K. S., and Fraser, R.: Increased precipitation drives
04 mega slump development and destabilization of ice-rich permafrost terrain, northwestern Canada, *Glob.*
05 *Planet. Change*, 129, 56–68, <https://doi.org/10.1016/j.gloplacha.2015.02.008>, 2015.
- 06 Kokelj, S. V., Lantz, T. C., Tunnicliffe, J., Segal, R., and Lacelle, D.: Climate-driven thaw of permafrost
07 preserved glacial landscapes, northwestern Canada, *Geology*, 45, 371–374, <https://doi.org/10.1130/G38626.1>,
08 2017.
- 09 Kyzivat, E. D., Smith, L. C., Pitcher, L. H., Fayne, J. V., Cooley, S. W., Cooper, M. G., Topp, S. N., Langhorst,
10 T., Harlan, M. E., Horvat, C., Gleason, C. J., and Pavelsky, T. M.: A High-Resolution Airborne Color-
11 Infrared Camera Water Mask for the NASA ABoVE Campaign, *Remote Sens.*, 11, 2163,
12 <https://doi.org/10.3390/rs11182163>, 2019.
- 13 Kyzivat, E. D., Smith, L. C., Pitcher, L. H., Fayne, J. V., Cooley, S. W., Cooper, M. G., Topp, S., Langhorst, T.,
14 Harlan, M. E., Gleason, C. J., and Pavelsky, T. M.: ABoVE: AirSWOT Water Masks from Color-Infrared
15 Imagery over Alaska and Canada, 2017, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1707>, 2020.
- 16 Lafleur, P. M. and Humphreys, E. R.: Tundra shrub effects on growing season energy and carbon dioxide
17 exchange, *Environ. Res. Lett.*, 13, 055001, <https://doi.org/10.1088/1748-9326/aab863>, 2018.
- 18 Lantz, T. C. and Turner, K. W.: Changes in lake area in response to thermokarst processes and climate in Old
19 Crow Flats, Yukon, *J. Geophys. Res. Biogeosciences*, 120, 513–524, <https://doi.org/10.1002/2014JG002744>,
20 2015.

- 21 Lavalle, M., Hawkins, B., and Hensley, S.: Tomographic imaging with UAVSAR: Current status and new results
22 from the 2016 AfriSAR campaign, 2017 IEEE International Geoscience and Remote Sensing Symposium
23 (IGARSS), 2485–2488, <https://doi.org/10.1109/IGARSS.2017.8127498>, 2017.
- 24 Le Toan, T., Quegan, S., Davidson, M. W. J., Balzter, H., Paillou, P., Papathanassiou, K., Plummer, S., Rocca, F.,
25 Saatchi, S., Shugart, H., and Ulander, L.: The BIOMASS mission: Mapping global forest biomass to better
26 understand the terrestrial carbon cycle, *Remote Sens. Environ.*, 115, 2850–2860,
27 <https://doi.org/10.1016/j.rse.2011.03.020>, 2011.
- 28 Liljedahl, A., Hinzman, L., Busey, R., and Yoshikawa, K.: Physical short-term changes after a tussock tundra
29 fire, Seward Peninsula, Alaska, *J. Geophys. Res. Earth Surf.*, 112, <https://doi.org/10.1029/2006JF000554>,
30 2007.
- 31 Lindsay, C., Zhu, J., Miller, A. E., Kirchner, P., and Wilson, T. L.: Deriving Snow Cover Metrics for Alaska
32 from MODIS, *Remote Sens.*, 7, 12961–12985, <https://doi.org/10.3390/rs71012961>, 2015.
- 33 Lipovsky, P. S.: Summary of Yukon Geological Survey permafrost monitoring network results, 2008-2013, in:
34 Yukon Exploration and Geology 2014, edited by: MacFarlane, K. E., Nordling, M. G., and Sack, P. J., Yukon
35 Geological Survey, 113–122, 2015.
- 36 Liu, L., Schaefer, K., Zhang, T., and Wahr, J.: Estimating 1992–2000 average active layer thickness on the
37 Alaskan North Slope from remotely sensed surface subsidence, *J. Geophys. Res. Earth Surf.*, 117,
38 <https://doi.org/10.1029/2011JF002041>, 2012.
- 39 Liu, L., Jafarov, E. E., Schaefer, K. M., Jones, B. M., Zebker, H. A., Williams, C. A., Rogan, J., and Zhang, T.:
40 InSAR detects increase in surface subsidence caused by an Arctic tundra fire, *Geophys. Res. Lett.*, 41, 3906–
41 3913, <https://doi.org/10.1002/2014GL060533>, 2014.
- 42 Liu, L., Schaefer, K., Chen, A. C., Gusmeroli, A., Jafarov, E. E., Panda, S. K., Parsekian, A. D., Schaefer, T.,
43 Zebker, H. A., and Zhang, T.: Pre-ABoVE: Remotely Sensed Active Layer Thickness, Barrow, Alaska, 2006-
44 2011, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1266>, 2015a.
- 45 Liu, L., Schaefer, K., Chen, A. C., Gusmeroli, A., Jafarov, E. E., Panda, S. K., Parsekian, A. D., Schaefer, T.,
46 Zebker, H. A., and Zhang, T.: Pre-ABoVE: Remotely Sensed Active Layer Thickness, Prudhoe Bay, Alaska,
47 1992-2000, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1267>, 2015b.
- 48 Loboda, T. V., French, N. H. F., Hight-Harf, C., Jenkins, L., and Miller, M. E.: Mapping fire extent and burn
49 severity in Alaskan tussock tundra: An analysis of the spectral response of tundra vegetation to wildland fire,
50 *Remote Sens. Environ.*, 134, 194–209, <https://doi.org/10.1016/j.rse.2013.03.003>, 2013.
- 51 Loboda, T.V., L.K. Jenkins, D. Chen, J. He, and A. Baer. 2022. Burned and Unburned Field Site Data, Noatak,
52 Seward, and North Slope, AK, 2016-2018. ORNL DAAC, Oak Ridge, Tennessee,
53 USA. <https://doi.org/10.3334/ORNLDAAC/1919>
- 54 Lopez-Sanchez, J. M., Ballester-Berman, J. D., Vicente-Guijalba, F., Cloude, S. R., McNairn, H., Shang, J.,
55 Skriver, H., Jagdhuber, T., Hajnsek, I., Pottier, E., Marechal, C., Hubert-Moy, L., Corgne, S., Wdowinski, S.,
56 Touzi, R., Gosselin, G., Brooks, R., Yamaguchi, Y., and Singh, G.: Agriculture and Wetland Applications,
57 in: *Polarimetric Synthetic Aperture Radar: Principles and Application*, vol. 25, edited by: Hajnsek, I. and
58 Desnos, Y.-L., Springer International Publishing, Cham, 119–178, [https://doi.org/10.1007/978-3-030-56504-
59 6_3](https://doi.org/10.1007/978-3-030-56504-6_3), 2021.
- 60 Lou, Y., Shimada, J. G., Michel, T. R., Muellerschoen, R. J., Zheng, Y., and Moghaddam, M.: Pre-ABoVE: L1
61 S-0 Polarimetric Data from AirMOSS P-band SAR, Alaska, 2014-2015, ORNL DAAC,
62 <https://doi.org/10.3334/ORNLDAAC/1678>, 2019.
- 63 Mack, M. C., Bret-Harte, M. S., Hollingsworth, T. N., Jandt, R. R., Schuur, E. A. G., Shaver, G. R., and Verbyla,
64 D. L.: Carbon loss from an unprecedented Arctic tundra wildfire, *Nature*, 475, 489–492,
65 <https://doi.org/10.1038/nature10283>, 2011.

- 66 Magagi, R., Berg, A. A., Goita, K., Belair, S., Jackson, T. J., Toth, B., Walker, A., McNairn, H., O'Neill, P. E.,
67 Moghaddam, M., Gherboudj, I., Colliander, A., Cosh, M. H., Burgin, M., Fisher, J. B., Kim, S.-B.,
68 Mladenova, I., Djamai, N., Rousseau, L.-P. B., Belanger, J., Shang, J., and Merzouki, A.: Canadian
69 Experiment for Soil Moisture in 2010 (CanEx-SM10): Overview and Preliminary Results, *IEEE Trans.*
70 *Geosci. Remote Sens.*, 51, 347–363, <https://doi.org/10.1109/TGRS.2012.2198920>, 2013.
- 71 Malone, T., Liang, J., and Packee, E. C.: Cooperative Alaska Forest Inventory, Gen Tech Rep PNW-GTR-785
72 Portland US Dep. Agric. For. Serv. Pac. Northwest Res. Stn. 42 P, 785, [https://doi.org/10.2737/PNW-GTR-](https://doi.org/10.2737/PNW-GTR-785)
73 785, 2009.
- 74 Marsh, P., Russell, M., Pohl, S., Haywood, H., and Onclin, C.: Changes in thaw lake drainage in the Western
75 Canadian Arctic from 1950 to 2000, *Hydrol. Process.*, 23, 145–158, <https://doi.org/10.1002/hyp.7179>, 2009a.
- 76 Marsh, P., Lesack, L., Hicks, F., Roberts, A., Hopkinson, C., Solomon, S., Forbes, D., Russell, M., and Haywood,
77 H.: Hydrology of the Mackenzie Delta: off-channel water storage and delta interaction with the Beaufort Sea,
78 2009b.
- 79 Marsh, P., Bartlett, P., MacKay, M., Pohl, S., and Lantz, T.: Snowmelt energetics at a shrub tundra site in the
80 western Canadian Arctic, *Hydrol. Process.*, 24, 3603–3620, <https://doi.org/10.1002/hyp.7786>, 2010.
- 81 McCune, B., Arup, U., Breuss, O., Di Meglio, J., Esslinger, T. L., Magain, N., Miadlikowska, J., Miller, A. E.,
82 Muggia, L., Nelson, P. R., Rosentreter, R., Schultz, M., Sheard, J. W., Tønsberg, T., and Walton, J.:
83 Biodiversity and ecology of lichens of Katmai and Lake Clark National Parks and Preserves, Alaska,
84 *Mycosphere*, 9, 859–930, 2018.
- 85 McGuire, A. D., Anderson, L. G., Christensen, T. R., Dallimore, S., Guo, L., Hayes, D. J., Heimann, M.,
86 Lorenson, T. D., Macdonald, R. W., and Roulet, N.: Sensitivity of the carbon cycle in the Arctic to climate
87 change, *Ecol. Monogr.*, 79, 523–555, <https://doi.org/10.1890/08-2025.1>, 2009.
- 88 Meddens, A. J. H., Vierling, L. A., Eitel, J. U. H., Jennewein, J. S., White, J. C., and Wulder, M. A.: Developing
89 5 m resolution canopy height and digital terrain models from WorldView and ArcticDEM data, *Remote Sens.*
90 *Environ.*, 218, 174–188, <https://doi.org/10.1016/j.rse.2018.09.010>, 2018.
- 91 Mehra, R. and Ramanujam, V. M.: L S Band Airborne SAR Data Products Calibration, in: 2019 URSI Asia-
92 Pacific Radio Science Conference (AP-RASC), 2019 URSI Asia-Pacific Radio Science Conference (AP-
93 RASC), 1–1, <https://doi.org/10.23919/URSIAP-RASC.2019.8738681>, 2019.
- 94 Meyer, G., Humphreys, E. R., Melton, J. R., Cannon, A. J., and Lafleur, P. M.: Simulating shrubs and their
95 energy and carbon dioxide fluxes in Canada's Low Arctic with the Canadian Land Surface Scheme Including
96 biogeochemical Cycles (CLASSIC), *Biogeosciences Discuss.*, 1–34, <https://doi.org/10.5194/bg-2020-458>,
97 2020.
- 98 Michaelian, M., Hogg, E. H., Hall, R. J., and Arsenault, E.: Massive mortality of aspen following severe drought
99 along the southern edge of the Canadian boreal forest, *Glob. Change Biol.*, 17, 2084–2094,
00 <https://doi.org/10.1111/j.1365-2486.2010.02357.x>, 2011.
- 01 Michaelides, R. J., Schaefer, K., Zebker, H. A., Parsekian, A., Liu, L., Chen, J., Natali, S., Ludwig, S., and
02 Schaefer, S. R.: Inference of the impact of wildfire on permafrost and active layer thickness in a
03 discontinuous permafrost region using the remotely sensed active layer thickness (ReSALT) algorithm,
04 *Environ. Res. Lett.*, 14, 035007, <https://doi.org/10.1088/1748-9326/aaf932>, 2019.
- 05 Michaelides, R. J., Chen, R. H., Zhao, Y., Schaefer, K., Parsekian, A. D., Sullivan, T., et al. (2021).
06 Permafrost Dynamics Observatory—part I: Postprocessing and calibration methods of UAVSAR L-
07 band InSAR data for seasonal subsidence estimation. *Earth and Space Science*, 8, e2020EA001630.
08 <https://doi.org/10.1029/2020EA001630>
- 09 Miller, C. E., Griffith, P. C., Goetz, S. J., Hoy, E. E., Pinto, N., McCubbin, I. B., Thorpe, A. K., Hofton, M.,
10 Hodkinson, D., Hansen, C., Woods, J., Larson, E., Kasischke, E. S., and Margolis, H. A.: An overview of

- 11 ABoVE airborne campaign data acquisitions and science opportunities, *Environ. Res. Lett.*, 14, 080201,
12 <https://doi.org/10.1088/1748-9326/ab0d44>, 2019.
- 13 Miller, C.E., P. Griffith, E.E. Hoy, N. Pinto, Y. Lou, S. Hensley, B. Chapman, J.L. Baltzer, K. Bakian-
14 Dogaheh, W.R. Bolton, L.L. Bourgeau-Chavez, R.H. Chen, B-H. Choe, L.K. Clayton, T.A.
15 Douglas, N.H.F. French, J.E. Holloway, G. Hong, L. Huang, G. Iwahana, L.K. Jenkins, J.S.
16 Kimball, T.V. Loboda, M.C. Mack, P. Marsh, R.J. Michaelides, M. Moghaddam, A.D. Parsekian, K.
17 Schaefer, P. Siqueira, D. Singh, A. Tabatabaenejad, M.R. Turetsky, R. Touzi, E. Wig, P. Wilson,
18 C.J. Wilson, S.D. Wulschleger, Y. Yi, H.A. Zebker, Y. Zhang, Y. Zhao, and S.J. Goetz. 2023.
19 Summary of the ABoVE L-band and P-band Airborne SAR Surveys. ORNL DAAC, Oak Ridge,
20 Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/2150>
- 21 Minions, C., Natali, S., Watts, J. D., Ludwig, S., and Risk, D.: ABoVE: Year-Round Soil CO₂ Efflux in Alaskan
22 Ecosystems, Version 2, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1762>, 2020.
- 23 Minsley, B. J., Abraham, J. D., Smith, B. D., Cannia, J. C., Voss, C. I., Jorgenson, M. T., Walvoord, M. A.,
24 Wylie, B. K., Anderson, L., Ball, L. B., Deszcz-Pan, M., Wellman, T. P., and Ager, T. A.: Airborne
25 electromagnetic imaging of discontinuous permafrost, *Geophys. Res. Lett.*, 39,
26 <https://doi.org/10.1029/2011GL050079>, 2012.
- 27 Moghaddam, M., Tabatabaenejad, A., Chen, R. H., Saatchi, S. S., Jaruwatanadilok, S., Burgin, M., Duan, X.,
28 and Truong-Loi, M. L.: AirMOSS: L2/3 Volumetric Soil Moisture Profiles Derived From Radar, 2012-2015,
29 ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1418>, 2016.
- 30 Montesano, P. M., Sun, G., Dubayah, R. O., and Ranson, K. J.: Spaceborne potential for examining taiga-tundra
31 ecotone form and vulnerability, *Biogeosciences*, 13, 3847–3861, <https://doi.org/10.5194/bg-13-3847-2016>,
32 2016.
- 33 Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K. D., Macias-Fauria,
34 M., Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P., Hermanutz, L., Trant, A., Collier, L. S.,
35 Weijers, S., Rozema, J., Rayback, S. A., Schmidt, N. M., Schaepman-Strub, G., Wipf, S., Rixen, C., Ménard,
36 C. B., Venn, S., Goetz, S., Andreu-Hayles, L., Elmendorf, S., Ravolainen, V., Welker, J., Grogan, P., Epstein,
37 H. E., and Hik, D. S.: Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities,
38 *Environ. Res. Lett.*, 6, 045509, <https://doi.org/10.1088/1748-9326/6/4/045509>, 2011.
- 39 Natali, S. M., Schuur, E. A. G., Webb, E. E., Pries, C. E. H., and Crummer, K. G.: Permafrost degradation
40 stimulates carbon loss from experimentally warmed tundra, *Ecology*, 95, 602–608,
41 <https://doi.org/10.1890/13-0602.1>, 2014.
- 42 National Research Council: Opportunities to Use Remote Sensing in Understanding Permafrost and Related
43 Ecological Characteristics: Report of a Workshop, The National Academies Press, Washington, DC, 2014.
- 44 Oechel, W. C., Vourlitis, G. L., Hastings, S. J., Zulueta, R. C., Hinzman, L., and Kane, D.: Acclimation of
45 ecosystem CO₂ exchange in the Alaskan Arctic in response to decadal climate warming, *Nature*, 406, 978–
46 981, <https://doi.org/10.1038/35023137>, 2000.
- 47 Pan, C. G., Kirchner, P. B., Kimball, J. S., and Du, J.: A Long-Term Passive Microwave Snowoff Record for the
48 Alaska Region 1988–2016, *Remote Sens.*, 12, 153, <https://doi.org/10.3390/rs12010153>, 2020.
- 49 Pavelsky, T. M. and Smith, L. C.: Remote sensing of hydrologic recharge in the Peace-Athabasca Delta, Canada,
50 *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL033268>, 2008.
- 51 Pietroniro, A., Peters, D. L., Yang, D., Fiset, J.-M., Saint-Jean, R., Fortin, V., Leconte, R., Bergeron, J., Siles, G.
52 L., Trudel, M., Garnaud, C., Matte, P., Smith, L. C., Gleason, C. J., and Pavelsky, T. M.: Canada's
53 Contributions to the SWOT Mission – Terrestrial Hydrology (SWOT-C TH), *Can. J. Remote Sens.*, 45, 116–
54 138, <https://doi.org/10.1080/07038992.2019.1581056>, 2019.

- 55 Pitcher, L. H., Smith, L. C., Pavelsky, T. M., Fayne, J. V., Cooley, S. W., Altenau, E. H., Moller, D. K., and
56 Arvesen, J.: ABoVE: AirSWOT Radar, Orthomosaic, and Water Masks, Yukon Flats Basin, Alaska, 2015,
57 ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1655>, 2019a.
- 58 Pitcher, L. H., Pavelsky, T. M., Smith, L. C., Moller, D. K., Altenau, E. H., Allen, G. H., Lion, C., Butman, D.,
59 Cooley, S. W., Fayne, J. V., and Bertram, M.: AirSWOT InSAR Mapping of Surface Water Elevations and
60 Hydraulic Gradients Across the Yukon Flats Basin, Alaska, *Water Resour. Res.*, *55*, 937–953,
61 <https://doi.org/10.1029/2018WR023274>, 2019b.
- 62 Pitcher, L. H., Smith, L. C., Cooley, S. W., Zaino, A., Carlson, R., Pettit, J., Gleason, C. J., Minear, J. T., Fayne,
63 J. V., Willis, M. J., Hansen, J. S., Easterday, K. J., Harlan, M. E., Langhorst, T., Topp, S. N., Dolan, W.,
64 Kyzivat, E. D., Pietroniro, A., Marsh, P., Yang, D., Carter, T., Onclin, C., Hosseini, N., Wilcox, E., Moreira,
65 D., Berge-Nguyen, M., Cretaux, J.-F., and Pavelsky, T. M.: Advancing Field-Based GNSS Surveying for
66 Validation of Remotely Sensed Water Surface Elevation Products, *Front. Earth Sci.*, *8*,
67 <https://doi.org/10.3389/feart.2020.00278>, 2020.
- 68 Plaza, C., Pegoraro, E., Bracho, R., Celis, G., Crummer, K. G., Hutchings, J. A., Hicks Pries, C. E., Mauritz, M.,
69 Natali, S. M., Salmon, V. G., Schädel, C., Webb, E. E., and Schuur, E. A. G.: Direct observation of
70 permafrost degradation and rapid soil carbon loss in tundra, *Nat. Geosci.*, *12*, 627–631,
71 <https://doi.org/10.1038/s41561-019-0387-6>, 2019.
- 72 Porter, C., Morin, P., Howat, I., Noh, M.-J., Bates, B., Peterman, K., Keesey, S., Schlenk, M., Gardiner, J.,
73 Tomko, K., Willis, M., Kelleher, C., Cloutier, M., Husby, E., Foga, S., Nakamura, H., Platson, M.,
74 Wethington, M., Williamson, C., Bauer, G., Enos, J., Arnold, G., Kramer, W., Becker, P., Doshi, A.,
75 D’Souza, C., Cummins, P., Laurier, F., and Bojesen, M.: ArcticDEM,
76 <https://doi.org/10.7910/DVN/OHHUKH>, 2018.
- 77 Potter, C.: Ecosystem carbon emissions from 2015 forest fires in interior Alaska, *Carbon Balance Manag.*, *13*, 2,
78 <https://doi.org/10.1186/s13021-017-0090-0>, 2018.
- 79 Quegan, S., Le Toan, T., Chave, J., Dall, J., Exbrayat, J.-F., Minh, D. H. T., Lomas, M., D’Alessandro, M. M.,
80 Paillou, P., Papathanassiou, K., Rocca, F., Saatchi, S., Scipal, K., Shugart, H., Smallman, T. L., Soja, M. J.,
81 Tebaldini, S., Ulander, L., Villard, L., and Williams, M.: The European Space Agency BIOMASS mission:
82 Measuring forest above-ground biomass from space, *Remote Sens. Environ.*, *227*, 44–60,
83 <https://doi.org/10.1016/j.rse.2019.03.032>, 2019.
- 84 Quinton, W., Berg, A., Braverman, M., Carpino, O., Chasmer, L., Connon, R., Craig, J., Devoie, É., Hayashi, M.,
85 Haynes, K., Olefeldt, D., Pietroniro, A., Rezanezhad, F., Schincariol, R., and Sonnentag, O.: A synthesis of
86 three decades of hydrological research at Scotty Creek, NWT, Canada, *Hydrol. Earth Syst. Sci.*, *23*, 2015–
87 2039, <https://doi.org/10.5194/hess-23-2015-2019>, 2019.
- 88 Quinton, W. L., Adams, J. R., Baltzer, J. L., Berg, A. A., Craig, J. R., and Johnson, E.: Permafrost Ecosystems in
89 Transition: Understanding and Predicting Hydrological and Ecological Change in the Southern Taiga Plains,
90 Northeastern British Columbia and Southwestern Northwest Territories, *6*, 2015.
- 91 Ramanujam, V. M. and Mehra, R.: L S Band SAR Data Processing and Products, in: 2019 URSI Asia-Pacific
92 Radio Science Conference (AP-RASC), 2019 URSI Asia-Pacific Radio Science Conference (AP-RASC), 1–
93 1, <https://doi.org/10.23919/URSIAP-RASC.2019.8738235>, 2019.
- 94 Ramanujam, V. M., Suneela, T. J. V. D., and Bhan, R.: ISRO’s dual frequency airborne SAR pre-cursor to
95 NISAR, in: *Earth Observing Missions and Sensors: Development, Implementation, and Characterization IV*,
96 *Earth Observing Missions and Sensors: Development, Implementation, and Characterization IV*, 98810A,
97 <https://doi.org/10.1117/12.2228086>, 2016.
- 98 Rocha, A. V. and Shaver, G. R.: Burn severity influences postfire CO₂ exchange in arctic tundra, *Ecol. Appl.*, *21*,
99 477–489, <https://doi.org/10.1890/10-0255.1>, 2011a.

- 00 Rocha, A. V. and Shaver, G. R.: Postfire energy exchange in arctic tundra: the importance and climatic
01 implications of burn severity, *Glob. Change Biol.*, 17, 2831–2841, <https://doi.org/10.1111/j.1365->
02 2486.2011.02441.x, 2011b.
- 03 Rosen, P.A., Kim, Y., Kumar, R., Misra, T., Bhan, R. and Sagi, V.R., 2017, May. Global persistent SAR
04 sampling with the NASA-ISRO SAR (NISAR) mission. In *Radar Conference (RadarConf), 2017 IEEE* (pp.
05 0410-0414). IEEE.
- 06 Roy, A., Toose, P., Mavrovic, A., Pappas, C., Royer, A., Derksen, C., Berg, A., Rowlandson, T., El-Amine, M.,
07 Barr, A., Black, A., Langlois, A., and Sonnentag, O.: L-Band response to freeze/thaw in a boreal forest stand
08 from ground- and tower-based radiometer observations, *Remote Sens. Environ.*, 237, 111542,
09 <https://doi.org/10.1016/j.rse.2019.111542>, 2020.
- 10 Saatchi, S. S. and Moghaddam, M.: Estimation of crown and stem water content and biomass of boreal forest
11 using polarimetric SAR imagery, *IEEE Trans. Geosci. Remote Sens.*, 38, 697–709,
12 <https://doi.org/10.1109/36.841999>, 2000.
- 13 Saatchi, S., Xu, L., Yang, Y., and Yu, Y.: Evaluation of NISAR Biomass Algorithm in Temperate and Boreal
14 Forests, *IGARSS 2019 - 2019 IEEE International Geoscience and Remote Sensing Symposium*, 7363–7366,
15 <https://doi.org/10.1109/IGARSS.2019.8898657>, 2019.
- 16 Sadeghi, M., Tabatabaenejad, A., Tuller, M., Moghaddam, M., and Jones, S. B.: Advancing NASA’s AirMOSS
17 P-Band Radar Root Zone Soil Moisture Retrieval Algorithm via Incorporation of Richards’ Equation,
18 *Remote Sens.*, 9, 17, <https://doi.org/10.3390/rs9010017>, 2017.
- 19 Schaefer, K., Chen, A. C., Chen, J., Chen, R. H., Dogaheh, K., Jafarov, E., Liu, L., Michaelides, R. J.,
20 Moghaddam, M., Parsekian, A. D., Sullivan, T. D., Tabatabaenejad, A., Thompson, J., and Zebker, H.: The
21 Permafrost Dynamics Observatory, 2018a.
- 22 Schaefer, K., Chen, A. C., Chen, J., Chen, R. H., Dogaheh, K., Jafarov, E., Liu, L., Michaelides, R. J.,
23 Moghaddam, M., Parsekian, A. D., Sullivan, T. D., Tabatabaenejad, A., Thompson, J., and Zebker, H.: The
24 Permafrost Dynamics Observatory (PDO), 2018b.
- 25 Schaefer, K., Michaelides, R. J., Chen, R. H., Sullivan, T. D., Parsekian, A. D., Bakian-Dogaheh, K.,
26 Tabatabaenejad, A., Moghaddam, M., Chen, J., Chen, A. C., Liu, L., and Zebker, H. A.: ABoVE: Active
27 Layer Thickness Derived from Airborne L- and P-band SAR, Alaska, 2017, ORNL DAAC,
28 <https://doi.org/10.3334/ORN LDAAC/1676>, 2019.
- 29 Schaefer, K., R.J. Michaelides, R.H. Chen, T.D. Sullivan, A.D. Parsekian, Y. Zhao, K. Bakian-Dogaheh, A.
30 Tabatabaenejad, M. Moghaddam, J. Chen, A.C. Chen, L. Liu, and H.A. Zebker. 2021a. ABoVE: Active
31 Layer Thickness Derived from Airborne L- and P-band SAR, Alaska, 2017. ORNL DAAC, Oak Ridge,
32 Tennessee, USA. <https://doi.org/10.3334/ORN LDAAC/1796>
- 33 Schaefer, K., L.K. Clayton, M.J. Battaglia, L.L. Bourgeau-Chavez, R.H. Chen, A.C. Chen, J. Chen, K. Bakian-
34 Dogaheh, T.A. Douglas, S.E. Grelick, G. Iwahana, E. Jafarov, L. Liu, S. Ludwig, R.J. Michaelides, M.
35 Moghaddam, S. Natali, S.K. Panda, A.D. Parsekian, A.V. Rocha, S.R. Schaefer, T.D. Sullivan, A.
36 Tabatabaenejad, K. Wang, C.J. Wilson, H.A. Zebker, T. Zhang, and Y. Zhao. 2021b. ABoVE: Soil Moisture
37 and Active Layer Thickness in Alaska and NWT, Canada, 2008-2020. ORNL DAAC, Oak Ridge, Tennessee,
38 USA. <https://doi.org/10.3334/ORN LDAAC/1903>
- 39 Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of
40 permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559,
41 <https://doi.org/10.1038/nature08031>, 2009.
- 42 Schuur, T.: AmeriFlux US-EML Eight Mile Lake Permafrost thaw gradient, Healy Alaska (3.5),
43 <https://doi.org/10.17190/AMF/1418678>, 2019.
- 44 Sellers, P., Hall, F., Margolis, H., Kelly, B., Baldocchi, D., Hartog, G. den, Cihlar, J., Ryan, M. G., Goodison, B.,
45 Crill, P., Ranson, K. J., Lettenmaier, D., and Wickland, D. E.: The Boreal Ecosystem–Atmosphere Study

46 (BOREAS): An Overview and Early Results from the 1994 Field Year, *Bull. Am. Meteorol. Soc.*, 76, 1549–
 47 1577, [https://doi.org/10.1175/1520-0477\(1995\)076<1549:TBESAO>2.0.CO;2](https://doi.org/10.1175/1520-0477(1995)076<1549:TBESAO>2.0.CO;2), 1995.

48 Sellers, P. J., Hall, F. G., Kelly, R. D., Black, A., Baldocchi, D., Berry, J., Ryan, M., Ranson, K. J., Crill, P. M.,
 49 Lettenmaier, D. P., Margolis, H., Cihlar, J., Newcomer, J., Fitzjarrald, D., Jarvis, P. G., Gower, S. T.,
 50 Halliwell, D., Williams, D., Goodison, B., Wickland, D. E., and Guertin, F. E.: BOREAS in 1997:
 51 Experiment overview, scientific results, and future directions, *J. Geophys. Res. Atmospheres*, 102, 28731–
 52 28769, <https://doi.org/10.1029/97JD03300>, 1997.

53 Serrouya, R., Dickie, M., Lamb, C., van Oort, H., Kelly, A. P., DeMars, C., McLoughlin, P. D., Larter, N. C.,
 54 Hervieux, D., Ford, A. T., and Boutin, S.: Trophic consequences of terrestrial eutrophication for a threatened
 55 ungulate, *Proc. R. Soc. B Biol. Sci.*, 288, 20202811, <https://doi.org/10.1098/rspb.2020.2811>, 2021.

56 Sherriff, R. L., Miller, A. E., Muth, K., Schriver, M., and Batzel, R.: Spruce growth responses to warming vary
 57 by ecoregion and ecosystem type near the forest-tundra boundary in south-west Alaska, *J. Biogeogr.*, 44,
 58 1457–1468, <https://doi.org/10.1111/jbi.12968>, 2017.

59 Shugar, D. H., Clague, J. J., Best, J. L., Schoof, C., Willis, M. J., Copland, L., and Roe, G. H.: River piracy and
 60 drainage basin reorganization led by climate-driven glacier retreat, *Nat. Geosci.*, 10, 370–375,
 61 <https://doi.org/10.1038/ngeo2932>, 2017.

62 Silva, C. A., Duncanson, L., Hancock, S., Neuenschwander, A., Thomas, N., Hofton, M., Fatoyinbo, L., Simard,
 63 M., Marshak, C. Z., Armston, J., Lutchke, S., and Dubayah, R.: Fusing simulated GEDI, ICESat-2 and
 64 NISAR data for regional aboveground biomass mapping, *Remote Sens. Environ.*, 253, 112234,
 65 <https://doi.org/10.1016/j.rse.2020.112234>, 2021.

66 Smith, L. C., Pavelsky, T., Lettenmaier, D. P., Gleason, C. J., Pietroniro, A., Applejohn, A., Arvesen, J. C.,
 67 Bjella, K., Carter, T., Chao, R., Cooley, S. W., Cooper, M. G., Cretaux, J. F., Douglass, T., Faria, D., Fayne,
 68 J., Fiset, J. M., Goodman, S., Hanna, B., Harlan, M., Langhorst, T., Marsh, P., Moreira, D. M., Minear, J. T.,
 69 Onclin, C., Overstreet, B. T., Peters, D., Pettit, J., Pitcher, L. H., Russell, M., Spence, C., Topp, S., Turner, K.
 70 W., Vimal, S., Wilcox, E., Woodward, J., Yang, D., and Zaino, A.: AirSWOT flights and field campaigns for
 71 the 2017 Arctic-Boreal Vulnerability Experiment (ABoVE), in: AGU Fall Meeting Abstracts, C21F-1176,
 72 2017a.

73 Smith, S. L., Roy, L.-P., Lewkowicz, A. G., and Chartrand, J.: Ground thermal data collection along the Alaska
 74 Highway corridor (KP1559-1895), Yukon, summer 2016, Geological Survey of Canada, 2017b.

75 Spence, C. and Hedstrom, N.: Hydrometeorological data from Baker Creek Research Watershed, Northwest
 76 Territories, Canada, *Earth Syst. Sci. Data*, 10, 1753–1767, <https://doi.org/10.5194/essd-10-1753-2018>, 2018.

77 Sun, G., K. J. Ranson, D. S. Kimes, J. B. Blair, and K. Kovacs (2008), Forest vertical structure from GLAS: An
 78 evaluation using LVIS and SRTM data, *Remote Sens. Environ.*, **112**, 107–117, doi:[10.1016/j.rse.2006.09.036](https://doi.org/10.1016/j.rse.2006.09.036).

79 Tabatabaenejad, A., and M. Moghaddam, 2011. Retrieval of surface and deep soil moisture and effect
 80 of moisture profile on inversion accuracy. *IEEE Geosci. Remote Sensing Lett.*, vol. 8, no. 3, pp.
 81 477- 481, May 2011.

82 Tabatabaenejad, A., Burgin, M., Duan, X., and Moghaddam, M.: P-Band Radar Retrieval of Subsurface Soil
 83 Moisture Profile as a Second-Order Polynomial: First AirMOSS Results, *IEEE Trans. Geosci. Remote Sens.*,
 84 53, 645–658, <https://doi.org/10.1109/TGRS.2014.2326839>, 2015.

85 Tabatabaenejad, A., Chen, R. H., Burgin, M. S., Duan, X., Cuenca, R. H., Cosh, M. H., Scott, R. L., and
 86 Moghaddam, M.: Assessment and Validation of AirMOSS P-Band Root-Zone Soil Moisture Products, *IEEE*
 87 *Trans. Geosci. Remote Sens.*, 58, 6181–6196, <https://doi.org/10.1109/TGRS.2020.2974976>, 2020.

88 Tank, S., Olefeldt, D., Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada,
 89 Quinton, W., Centre for Cold Regions and Water Science, Wilfred Laurier University, Waterloo, Ontario,
 90 Canada, Spence, C., Environment and Climate Change Canada, Saskatoon, Saskatchewan, Canada, Dion, N.,
 91 Water Resources Department, Government of Northwest Territories, Yellowknife, Northwest Territories,

- 92 Canada, Ackley, C., Centre for Cold Regions and Water Science, Wilfred Laurier University, Waterloo,
93 Ontario, Canada, Burd, K., Department of Renewable Resources, University of Alberta, Edmonton, Alberta,
94 Canada, Hutchins, R., Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada,
95 Mengistu, S., and Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada:
96 Fire in the Arctic: The effect of wildfire across diverse aquatic ecosystems of the Northwest Territories, *Polar*
97 *Knowl. Aquat. Rep.*, 1, 31–38, <https://doi.org/10.35298/pkc.2018.04>, 2019.
- 98 Touzi, R., Deschamps, A., and Rother, G.: Phase of Target Scattering for Wetland Characterization Using
99 Polarimetric C-Band SAR, *IEEE Trans. Geosci. Remote Sens.*, 47, 3241–3261,
00 <https://doi.org/10.1109/TGRS.2009.2018626>, 2009.
- 01 Touzi, R., Omari, K., Sleep, B., and Jiao, X.: Scattered and Received Wave Polarization Optimization for
02 Enhanced Peatland Classification and Fire Damage Assessment Using Polarimetric PALSAR, *IEEE J. Sel.*
03 *Top. Appl. Earth Obs. Remote Sens.*, PP, 1–26, <https://doi.org/10.1109/JSTARS.2018.2873740>, 2018.
- 04 Touzi, R., Hong, G., Motohka, T., Shinichi, S., and De Lisle, D.: Investigation of Compact SAR L and C band
05 Complementarity for Permafrost Characterization In Arctic Regions, *IGARSS 2019 - 2019 IEEE*
06 *International Geoscience and Remote Sensing Symposium*, 4665–4667,
07 <https://doi.org/10.1109/IGARSS.2019.8898510>, 2019a.
- 08 Touzi, R., Pawley, S., Hosseini, M., and Jiao, X.: Polarimetric L-band PALSAR2 for Discontinuous Permafrost
09 Mapping In Peatland Regions, in: *IGARSS 2019 - 2019 IEEE International Geoscience and Remote Sensing*
10 *Symposium, IGARSS 2019 - 2019 IEEE International Geoscience and Remote Sensing Symposium*,
11 Yokohama, Japan, xvii–ccxxii, <https://doi.org/10.1109/IGARSS.2019.8900243>, 2019b.
- 12 Tsuyuzaki, S., Iwahana, G., and Saito, K.: Tundra fire alters vegetation patterns more than the resultant
13 thermokarst, *Polar Biol.*, 41, 753–761, <https://doi.org/10.1007/s00300-017-2236-7>, 2018.
- 14 Ullmann, T., Banks, S. N., Schmitt, A., and Jagdhuber, T.: Scattering Characteristics of X-, C- and L-Band
15 PolSAR Data Examined for the Tundra Environment of the Tuktoyaktuk Peninsula, Canada, *Appl. Sci.*, 7,
16 595, <https://doi.org/10.3390/app7060595>, 2017.
- 17 Vincent, W. F., Callaghan, T. V., Dahl-Jensen, D., Johansson, M., Kovacs, K. M., Michel, C., Prowse, T., Reist,
18 J. D., and Sharp, M.: Ecological Implications of Changes in the Arctic Cryosphere, *AMBIO*, 40, 87–99,
19 <https://doi.org/10.1007/s13280-011-0218-5>, 2011.
- 20 Walker, B., Wilcox, E. J., and Marsh, P.: Accuracy assessment of late winter snow depth mapping for tundra
21 environments using Structure-from-Motion photogrammetry1, *Arct. Sci.*, [https://doi.org/10.1139/as-2020-](https://doi.org/10.1139/as-2020-0006)
22 0006, 2020.
- 23 Walker, X. J., Rogers, B. M., Baltzer, J. L., Cummings, S. R., Day, N. J., Goetz, S. J., Johnstone, J. F., Turetsky,
24 M. R., and Mack, M. C.: ABoVE: Wildfire Carbon Emissions and Burned Plot Characteristics, NWT, CA,
25 2014–2016, ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1561>, 2018a.
- 26 Walker, X. J., Rogers, B. M., Baltzer, J. L., Cumming, S. G., Day, N. J., Goetz, S. J., Johnstone, J. F., Schuur, E.
27 A. G., Turetsky, M. R., and Mack, M. C.: Cross-scale controls on carbon emissions from boreal forest
28 megafires, *Glob. Change Biol.*, 24, 4251–4265, <https://doi.org/10.1111/gcb.14287>, 2018b.
- 29 Walker, X. J., Baltzer, J. L., Laurier, W., Cumming, S. G., Day, N. J., Goetz, S. J., Johnstone, J. F., Potter, S.,
30 Rogers, B. M., Schuur, E. a. G., Turetsky, M. R., and Mack, M. C.: ABoVE: Characterization of Carbon
31 Dynamics in Burned Forest Plots, NWT, Canada, 2014, ORNL DAAC,
32 <https://doi.org/10.3334/ORNLDAAC/1664>, 2019a.
- 33 Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., Johnstone, J. F., Potter, S., Rogers,
34 B. M., Schuur, E. A. G., Turetsky, M. R., and Mack, M. C.: Increasing wildfires threaten historic carbon sink
35 of boreal forest soils, *Nature*, 572, 520–523, <https://doi.org/10.1038/s41586-019-1474-y>, 2019b.
- 36 Whalen, D., Forbes, D. L., Hopkinson, C., Lavergne, J. C., Manson, G. K., Marsh, P., and Solomon, S. M.:
37 Topographic LiDAR-Providing a new perspective in the Mackenzie Delta, 2009.

- 38 Whitley, M., Frost, G. V., Jorgenson, M. T., Macander, M., Maio, C. V., and Winder, S. G.: ABoVE: Permafrost
39 Measurements and Distribution Across the Y-K Delta, Alaska, 2016, ORNL DAAC,
40 <https://doi.org/10.3334/ORNLDAAC/1598>, 2018a.
- 41 Whitley, M. A., Frost, G. V., Jorgenson, M. T., Macander, M. J., Maio, C. V., and Winder, S. G.: Assessment of
42 LiDAR and Spectral Techniques for High-Resolution Mapping of Sporadic Permafrost on the Yukon-
43 Kuskokwim Delta, Alaska, *Remote Sens.*, 10, 258, <https://doi.org/10.3390/rs10020258>, 2018b.
- 44 Wilcox, E. J., Keim, D., Jong, T. de, Walker, B., Sonnentag, O., Sniderhan, A. E., Mann, P., and Marsh, P.:
45 Tundra shrub expansion may amplify permafrost thaw by advancing snowmelt timing, *Arct. Sci.*,
46 <https://doi.org/10.1139/as-2018-0028>, 2019.
- 47 Wilson, C., Dann, J., Bolton, R., Charsley-Groffman, L., Jafarov, E., Musa, D., and Wullschleger, S.: In Situ Soil
48 Moisture and Thaw Depth Measurements Coincident with Airborne SAR Data Collections, Barrow and
49 Seward Peninsulas, Alaska, 2017, <https://doi.org/10.5440/1423892>, 2021.
- 50 Woodward, A. and Beever, E. A.: Conceptual ecological models to support detection of ecological change on
51 Alaska National Wildlife Refuges, 2011.
- 52 Yi, Y., Kimball, J. S., Chen, R. H., Moghaddam, M., Reichle, R. H., Mishra, U., Zona, D., and Oechel, W. C.:
53 Characterizing permafrost active layer dynamics and sensitivity to landscape spatial heterogeneity in Alaska,
54 *The Cryosphere*, 12, 145–161, <https://doi.org/10.5194/tc-12-145-2018>, 2018.
- 55 Zhang, Y., Touzi, R., Feng, W., Hong, G., Lantz, T. C., and Kokelj, S. V.: Landscape-scale variations in near-
56 surface soil temperature and active-layer thickness: Implications for high-resolution permafrost mapping,
57 *Permafr. Periglac. Process. J.*, 2021. <https://doi.org/10.1002/ppp.2104>
- 58 Zhao, Y., Chen, R.H., Bakina-Dogaheh, K., Whitcomb, J., Yi, Y., Kimball, J. S., Moghaddam, M.: Mapping
59 Boreal Forest Species and Canopy Height using Airborne SAR and Lidar Data in Interior Alaska, IGARSS
60 2022 - 2022 IEEE International Geoscience and Remote Sensing Symposium, Kuala Lumpur, Malaysia,
61 2022, pp. 4955-4958, doi: 10.1109/IGARSS46834.2022.9883311.

62
63
64