

PAPILA dataset: a regional emission inventory of reactive gases for South America based on the combination of local and global information

Paula Castesana^{1,2,3,*}, Melisa Diaz Resquin^{2,4,5,*}, Nicolás Huneeus^{5,6}, Enrique Puliafito^{1,7}, Sabine Darras⁸, Darío Gómez^{2,4}, Claire Granier^{8,9}, Mauricio Osse Alvarado¹⁰, Néstor Rojas¹¹, and Laura Dawidowski^{2,3}

¹Consejo Nacional de Investigaciones Científicas y Técnicas, Buenos Aires, Argentina.

²Comisión Nacional de Energía Atómica, Gerencia Química, Buenos Aires, Argentina.

³Instituto de Investigación e Ingeniería Ambiental, Universidad Nacional de San Martín, Buenos Aires, Argentina.

⁴Facultad de Ingeniería, Universidad de Buenos Aires, Buenos Aires, Argentina.

⁵Center for Climate and Resilience Research (CR)², Santiago, Chile.

⁶Departamento de Geofísica, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Santiago, Chile

⁷Mendoza Regional Faculty – National Technological University (FRM-UTN), Mendoza, Argentina

⁸Laboratoire d’Aérologie, Université de Toulouse, CNRS, UPS, France

⁹CIRES, University of Colorado and NOAA Chemical Sciences Laboratory, Boulder, United States

¹⁰Departamento Ingeniería Mecánica, Universidad Técnica Federico Santa María (UTFSM), Santiago, Chile

¹¹Air Quality Research Group, Universidad Nacional de Colombia, Bogotá, Colombia

*These authors contributed equally to this work.

Correspondence: Paula Castesana (pcastesana@unsam.edu.ar), Melisa Diaz Resquin (mdiazresquin@fi.uba.ar)

Abstract. The multidisciplinary project Prediction of Air Pollution in Latin America and the Caribbean (PAPILA) is dedicated to the development and implementation of an air quality analysis and forecasting system to assess pollution impacts on human health and economy. In this context, a comprehensive emission inventory for South America was developed on the basis of the existing data on the global dataset ~~CAMS-GLOB-ANT~~ [CAMS-GLOB-ANT](#) v4.1 (developed by joining CEDS trends

5 and ~~EDGARv4~~ [EDGAR v4](#).3.2 historical data), enriching it with ~~derived data~~ [data derived](#) from locally available emission inventories for Argentina, Chile and Colombia. This work presents the results of the first joint effort of South American

researchers and European colleagues to generate regional maps of emissions, together with a methodological approach to continue incorporating information into future versions of the dataset. This version of the PAPILA dataset includes CO, NO_x, NMVOCs, NH₃ and SO₂ annual emissions from anthropogenic sources for the period ~~2014–2016~~ [2014–2016](#), with a spatial

10 resolution of 0.1° x 0.1° over a domain that covers ~~32°–120°W and 34°N–58°S~~ [32°–120°W and 34° N–58° S](#). PAPILA

dataset is presented as netCDF4 files and is available in an open access data repository under a CC-BY 4 license: <http://dx.doi.org/10.17632/btf2mz4fhf.3>. A comparative assessment of ~~PAPILA–CAMS~~ [PAPILA–CAMS](#) datasets was carried out for (i) the South American region, (ii) the countries with local data (Argentina, Colombia and Chile), and (iii) downscaled

15 emission maps for urban domains with different environmental and anthropogenic factors. Relevant differences were ~~obtained~~ [found](#) both at country and urban ~~level~~ [levels](#) for all the compounds ~~analysed~~ [analyzed](#). Among them, we found that when comparing ~~total emissions of PAPILA~~ [PAPILA total emissions](#) versus CAMS datasets at the national level, higher levels of

NO_x and considerably lower levels of the other species were obtained for Argentina, higher levels of SO₂ and lower levels of CO and NO_x for Colombia, and considerably higher levels of CO, NMVOCs and SO₂ for Chile. These discrepancies are mainly related to the representativeness of the local practices in the local emissions emission estimates, to the improvements made in the spatial distribution of the locally estimated emissions, or both. Both datasets were evaluated relative to against surface concentrations of CO and NO_x by using them as input data to the WRF-Chem WRF-Chem model for one of the analysed analyzed domains, the Metropolitan Area of Buenos Aires, for summer and winter of 2015. For winter, PAPILA-based results had lower PAPILA-based modelling results had a smaller bias for CO and NO_x concentrations, for which in winter while CAMS-based results tended to be underestimated for the same period tended to deliver an underestimation of these concentrations. Both inventories exhibited similar performances for CO in summer, while PAPILA simulation outperformed CAMS for NO_x concentrations. These results highlight the importance of refining global inventories with local data to obtain accurate results with high-resolution air quality models.

1 Introduction

South America (SA) is a region of complex political and social contrasts, fluctuating economies and the highest inequality levels worldwide (The World Bank, 2019). Demographically, SA is a region with a growing population and an increasing trend towards urban agglomeration and the demand for goods and services (Huneeus et al., 2020a). From the energy standpoint Regarding energy use, SA has significantly low coal consumption levels and a higher share of hydroelectricity in comparison with other world regions (IEA, 2020). The use of alternative fuels, such as biomass or waste, is usually not well covered by national statistics and therefore global information does not accurately represent the sectoral mix of fuels consumed in the different countries. Road transport in the region is characterized by a fleet older and in poorer operating and maintenance conditions than that circulating in developed countries. Moreover, the use of motorcycles has increased in the region, being of particular concern in some cities such as Lima and Bogotá (Romero et al., 2020; Ortegon-Sanchez and Oviedo Hernandez, 2016). In addition to diesel oil and gasoline, different fuels are consumed for road transport in the region: compressed natural gas (CNG) covers a significant fraction of fuel use by passenger vehicles in Argentina, a high share of liquefied petroleum gas (LPG) is used in Peru, while pure ethanol and gasoline-ethanol gasoline-ethanol blends are broadly used by flex fuel vehicles in Brazil (Belincanta et al., 2016). Adding to this diversity, legislation on sulfur content in fuels is very restrictive in some countries such as Chile and Colombia and much more flexible, particularly concerning diesel oil used by trucks and off-road off-road vehicles, in others (Forquera, 2002) (Huneeus et al., 2020a). With respect to land use, SA is one of the least densely populated places in the world although it is highly urbanized (United Nations, 2015). This often implies poor or lacking information on the level and spatial distribution of some anthropogenic activities. This is the case, for example of the extended use in SA of wood and wood-waste for cooking and for heating in colder zones of the region, e.g., southern Chile (Villalobos et al., 2017). Another relevant land use characteristic is the sustained trend of larger increasing harvested land areas largely due to conversion from forests to agriculture agricultural lands (The World Bank, 2020), and partly as a consequence of rising temperatures and changes in rainfall patterns, resulting in a shift of the agricultural border like as occurred in Argentina (Bar-

50 ros and Camilloni, 2016). Lastly, a unique feature of the region concerns hotspots of sulfur dioxide identified by the Ozone Monitoring Instrument (OMI) satellite sensor. For SA they are attributable mainly to volcanoes and the smelting of sulfides of copper and other metal ores in Chile and Peru, differing remarkably from other regions worldwide where these hotspots are mainly ~~the responsibility of emitted by~~ thermal power plants and oil and gas activities (Fioletov et al., 2016).

55 These regional particularities have direct consequences not only on the level and chemical profiles of the pollutants discharged to the atmosphere, but also on the specific locations where these emissions occur and on the population exposed to their environmental and health effects. Assessing the impact of atmospheric emissions as well as designing mitigation strategies, require reliable atmospheric emissions inventories (AEIs), which include spatially disaggregated emissions covering the entire region of interest in a transparent and consistent way in terms of emission sources and estimation methodologies (Kuenen et al., 2014). There is a wide range of global AEIs covering SA for different species and periods that meet the
60 mentioned requirements. Some of the AEIs worth mentioning include: the Emissions Database for Global Atmospheric Research (EDGAR) (Janssens-Maenhout et al., 2019; EDGAR, 2021), the Evaluating the Climate and Air Quality Impacts of ~~Short-Lived~~ ~~Short-Lived~~ Pollutants (ECLIPSE) (Stohl et al., 2015), the Community Emissions Data System (CEDS) (Hoesly et al., 2018), the integrated assessment model Greenhouse gas - Air pollution Interactions and Synergies (GAINS) (Klimont et al., 2017), or the Copernicus Atmosphere Monitoring Service datasets (CAMS) (Granier et al., 2019).

65 Across the region, government efforts on AEIs are mainly focused on GHGs in line with the international commitments under the United Nations Convention on Climate Change (UNFCCC). The regional community of GHG inventory compilers has grown remarkably in the last two decades and in many cases has helped to improve the collection of activity data including specific areas of national statistics systems. In parallel, research groups in SA have built inventories of ozone precursors and particles to be used as input data to air quality models. Links between several of these groups have recently been strengthened
70 by the creation of a regional initiative focused on the construction of inventories of species not covered by governments in their reports to the UNFCCC (Huneeus et al., 2017, 2020a).

75 Completeness in terms of species, represented sectors and time series is a strength of global AEIs while locally developed inventories seldom fully cover all three aspects. On the other hand, although the most current versions of global AEIs accurately reflect the emissions from sectors for which regional information is well documented in global statistics, they may miss some specificity and accuracy associated with local practices and technologies that is often better represented in local AEIs (Huneeus et al., 2020a). From this, it is plausible to assume that better emission estimates would be obtained by enriching the comprehensive global AEIs with locally generated information. This mosaic approach is an idea that has been successfully applied in the framework of the Task Force on Hemispheric Transport of Air Pollution (HTAP), an international cooperative effort to improve the understanding of the intercontinental transport of air pollution across the Northern Hemisphere. In this context, the HTAP_v2.2 air pollutant grid maps were developed combining available regional information within a complete global dataset (Janssens-Maenhout et al., 2015), and have been widely used even outside of HTAP.

80 This work presents what to our knowledge constitutes the first AEIs from anthropogenic sources covering the ~~entire~~ continental SA region, which combines ~~in a proper and rigorous way~~ local available information with a global database in a proper and rigorous way. For this purpose, the dataset ~~CAMS-GLOB-ANT~~ CAMS-GLOB-ANT v4.1 (Granier et al., 2019), developed

85 by joining CEDS trends and EDGAR v4.3.2 historical data, was used as a basis (hereinafter CAMS dataset), enriching it with
locally developed inventories available in the ~~bibliography~~ literature until 2019, and selecting those with national coverage and
with availability of data for the period and species of interest. The dataset presented in this work, hereinafter called PAPILA,
focuses on the group of species known as reactive gases, given their relevance in atmospheric chemistry as precursors of O₃ and
PM_{2.5}: carbon monoxide (CO), nitrogen oxides (NO_x), ~~non-methane~~ volatile organic compounds (NMVOCs),
90 ammonia (NH₃) and sulfur dioxide (SO₂) (Sharma et al., 2017). Due to the availability of data in the local AEIs and the
completeness of the sectors represented, the ~~2014–2016~~ period was selected for this first version of the PAPILA
dataset, including local information from ~~the continental areas of~~ Argentina (Puliafito et al., 2017; Castesana et al., 2018), Chile
(Mazzeo et al., 2018; Gallardo et al., 2018) and Colombia (IDEAM, 2017). In addition, a comparison of the performance of
95 both AEIs (PAPILA and CAMS) is presented using ~~near-surface~~ CO and NO_x mixing ~~ratio~~ simulated by the
Weather Research and ~~Forecasting–Chemistry (WRF–Chem)~~ (Grell et al., 2005) at a high
spatial resolution (3 km) against in situ observations made in Buenos Aires during ~~February–March and August–September~~
~~February–March and August–September~~ 2015.

This work was carried out within the framework of the Prediction of Air Pollution in Latin America and the Caribbean
(PAPILA, 2020) and Emission Inventories in South America (EMISA, 2020) projects. PAPILA combines for the first time
100 an ensemble of ~~state-of-the-art models, high-resolution state-of-the-art models, high-resolution~~ emission inventories, space
observations and surface measurements to provide real time forecasts and analysis of regional air pollution in the ~~LAC~~ Latin
America and the Caribbean region. Thus, an important aspect of the project is the development of appropriate and consistent
surface emission inventories as input data for air quality models. EMISA initiative was created to lay the foundations for
constructing robust and transparent inventories of the same set of species that have been consistently estimated across South
105 American countries using the same methodological approach. Local information on emissions was gathered from the countries
that participate in the EMISA project: Argentina, Brazil, Chile, Colombia and Peru. Relevant research groups in Brazil and
Peru have developed emission inventories for different cities (Policarpo et al., 2018; Dos Santos Lucon and Moutinho Dos
Santos, 2005; Vivanco and Andrade, 2006; Romero et al., 2020; Dawidowski et al., 2014); however as far as we know they
have ~~have~~ not developed inventories covering the entire countries for the species included in this study. ~~For this reason, the~~
110 ~~information~~ Since national territories are the common administrative entities that can be exchanged with the global inventory,
the local information on emissions from these countries was not included in this first version of the combined dataset. However,
this work is expected to be the starting point for the preparation of comprehensive emission inventories in South America
enriched with local information. For this purpose, we include a flow chart with the general methodology that we have applied
in combining local information with a global dataset(see Figure 2).

115 The paper is organized as follows. Section 2 describes, for each country, the approach and sources of information used to
develop the PAPILA dataset and also discusses the application of this inventory in an air quality model. Section 3 provides the
main differences of PAPILA and CAMS datasets for SA and other smaller domains and the results of the air quality simulations.
Section 4 provides a description of the data availability, and finally Section 5 presents the main conclusions of this work.

2 Methods

120 2.1 PAPILA dataset overview

The PAPILA dataset is a collection of CO, NO_x, NMVOCs, NH₃ and SO₂ inventories of annual emissions from anthropogenic sources in South America for the period 2014–2016. The inventories are presented as netCDF4 files, one for each species gridded with a spatial resolution of 0.1° x 0.1° covering the domain 32° W–120° W and 34° N–58° S. Each file contains 12 variables corresponding to the emissions in Tg y⁻¹ from the following categories, which are organized and denominated using the nomenclature given by CAMS: thermal power plants (ENE); residential and commercial combustion (RES); road transportation (TRO); ~~non-road~~ ~~non-road~~ transportation (TNR); fugitive emissions (FEF); industries, including fuel consumption in manufacturing industries and construction, refineries, industrial processes and solvent and other products use, (IND); agricultural soils (AGS); ~~agriculture~~ ~~agricultural~~ livestock (AGL); ~~inland navigation, which includes domestic, coastal, deep-sea and inland waterborne navigation, domestic and international navigation~~ (SHP); ~~international navigation (SHP-INT)~~; waste (including solid waste, wastewater and incineration) (SWD); and the sum of all categories (SUM). This grouping of ~~activities and sectors categories~~ was carried out following the CAMS sectoral disaggregation, except for the use of solvents, reported under IND in the PAPILA dataset. ~~International navigation emissions were taken entirely from the CAMS database. To be consistent with the base inventory (CAMS-GLOB-ANT v4.1) used for our mosaic inventory, aviation emissions were not included in this first version of the PAPILA dataset.~~ Agricultural fires were removed to allow the use of the inventories together with fire products such as GFEDv3 (van der Werf et al., 2010), FINNv1 (Wiedinmyer et al., 2011) and GFASv1 FINN v1 (Wiedinmyer et al., 2011) and GFAS v1 (Kaiser et al., 2012), avoiding double counting of these fires. It is worth mentioning that by "sum of all categories" we refer to all ~~the sectors~~ ~~those~~ included in PAPILA, both for the presentation of our results and for comparative purposes with CAMS. A broader description of the activities contemplated under each category is presented in the Table A1 in the Annex A, together with the equivalences with the IPCC 1996 reporting code.

The PAPILA dataset (Castesana et al., 2021) combines surface emissions from the comprehensive CAMS dataset with local information of those countries that, at the time of development of this emission inventory, had emission estimates of the mentioned species and covering the entire national territory: Argentina, Chile and Colombia. That information was collected and assessed in terms of species, emission categories and spatial coverage, selecting the most appropriate and representative data for each country, as described in the following subsections and summarized in Figure 1.

2.2 Designing and building the PAPILA dataset

Figure 2 presents a flow chart of the general methodology applied in combining local information with CAMS. Data from Argentina, Chile and Colombia ~~was~~ ~~were~~ assessed in terms of species availability. In addition, and as described in the next subsections, the transparency on the methodology applied in emission estimates, and the representativeness and completeness of emission ~~sectors categories~~ were revised in line with the CAMS emission reporting system. For those species and/or categories with absence of local data, CAMS inventory was used to fill the gaps.




Figure 1. Local and global information on emissions and their spatial distribution disaggregation included in the PAPILA dataset, by species, categories and countries. ENE: thermal power plants; RES: residential, commercial and other combustion; TR: road transportation; TRN: non-road non-road transportation; FEF: fugitive emissions; IND: industrial process; AGS: agricultural soils; AGL: agriculture livestock; SHP: inland navigation; SHP-INT- domestic and international navigation; SWD: wastes. CAMS refers to the global dataset CAMS-GLOB-ANT CAMS-GLOB-ANT v4.1, and "Rest of SA" to the rest of the countries of the South American region. NO: Not occurring.

One of the challenges of combining different local inventories to a common regional database is bringing them to a single, uniform and homogeneous grid. For this purpose, it was necessary to resolve all conflicts arising from cells shared by more




Figure 2. Flow Chart illustrating the general methodology applied to combine local information with the global dataset **CAMS-GLOB-ANT** **CAMS-GLOB-ANT** v4.1.

than one country or coastal cells. ~~This To be consistent with the base inventory used in this work, this~~ problem was solved 155 using the country and continent masks ~~applied by CAMS (CIESIN and CIAT, 2005), which are~~ created at ~~0.1° resolution (CIESIN and CIAT, 2005) that assign a unique ° resolution assigning a unique country~~ value for each cell.

2.2.1 Argentina

Spatially disaggregated emission inventories for all the species included in this work are available for Argentina. They cover all categories except SWD. Emissions from the following categories ENE, RES, TRO, TNR, FEF, IND and SHP were taken 160 from the GEAA inventory (Puliafito et al., 2017) which consists of a ~~high-resolution high-resolution~~ ~~(0.025° × 0.025°)~~ in-

ventory of 2014 annual emissions. For each category, GEAA covers: (i) for energy industries, the precise location of power plants, plus fuel consumption by technology and by fuel of each utility; (ii) for residential and commercial sources, spatially distributed fuel consumption estimated using energy use by province and census based population maps; (iii) for road transportation, fleet composition, fuel consumption by refueling stations, geographically distributed considering road maps by type and distance to the refueling stations; (iv) for ~~off-road~~ off-road transportation, emissions from railways, with fuel consumption data, geographically distributed with rail maps; (v) for fugitive emissions, including those from refining, storage, venting and flaring, and those from distribution of oil products and natural gas, annual data from national statistics, spatially distributed with the exact location of the facilities; (vi) for inland navigation ~~, fuel consumption, (namely, domestic plus international navigation on the continental area of Argentina), fuel consumption~~ spatially distributed with the geographical identification of the berths~~routs~~ routes and ports boundaries. The GEAA inventory has been updated for this work including emissions from IND, which were not covered in the published version (Puliafito et al., 2017). ~~With these changes for manufacturing industries, the database considers fuel consumption by fuels, petroleum refining and emissions from the~~ These emissions include (i) those from fuel consumption and from production process itself for the main industries, disaggregated by fuel and spatially distributed with the ~~location of the main industries and distributing the rest as area sources in the whole territory. In all these categories the~~ combustion of fossil and biomass fuels was considered. A precise location of each facility, and (ii) those from fuel consumption of small industries, whose consumption is known by activity and by district, and whose spatial disaggregation of emissions was carried out using the population density of each district as a proxy. We noted that a different allocation of fugitive emissions from the distribution of oil products and natural gas (mainly consisting of NMVOCs) ~~between the~~ exists between CAMS and the Argentinean inventory ~~is worth mentioning.:~~ CAMS includes these emissions under the IND ~~sector category~~ (see Table A1, Annex A) while ~~in the Argentinean inventory~~ they are reported under FEF in the Argentinean inventory. This does not imply omission or double counting of emissions.

To construct the PAPILA inventory, this new version of GEAA was updated to 2015 and 2016 by applying CEDS trends by emission categories ~~for Argentina and Chile and country-specific trends for Colombia~~. Final emissions were adapted to a homogeneous grid of $0.1^{\circ} \times 0.1^{\circ}$, and combined with agricultural local inventories described below and with the CAMS information on emissions from SWD and from SHP outwards from the Argentine coast.

Ammonia emissions from agricultural activities were taken from the ~~2000–2012~~ 2000–2012 estimates by Castesana et al. (2018). The time series was updated to the period ~~2014–2016~~ 2014–2016 by applying the methodology and local activity data sources detailed in the cited work. The rest of the studied species emitted from activities under AGL and AGS (*i.e.*, NO_x and NMVOCs) were estimated according to the 2016 methodology of the European Monitoring and Evaluation Program (EMEP, 2017), following the general expression:

$$E_i = AD \cdot EF_i \quad (1)$$

where E_i is the emission amount of the species i , AD is the activity data, and EF_i represents the emission factor of the species i related to that activity. Both the activity data and their spatial distribution were based on the previous work by Castesana et al. (2018, 2020), while the emission factors were those suggested by the EMEP according to the level of detail described

Table 1. Level of detail of the EMEP 2016 methodology applied in estimating NO_x and NMVOCs emissions from agricultural activities in Argentina.

Category	Description of sources	EMEP 2016 approach	
		NO _x	NMVOCs
AGL	<u>Manure management</u> (Dairy cattle, <u>non-dairy cattle, non-dairy cattle</u> , and other livestock)	Tier 2	
AGS	<u>Inorganic N-fertilisers</u>	<u>Inorganic N-fertilisers</u> Tier 1	NO
	Manure in pasture (all livestock)	Tier 1*	Tier 2
	Crops	NO	Tier 1

* There is no EMEP Tier 2 method.

Emissions from grazing (deposited on pasture) are reported as agricultural soils. NO: Not occurring. AGS: Agricultural soils.

AGL: Agricultural livestock.

195 for each activity in Table 1. These emissions have been estimated ad hoc to be included as part of the PAPILA dataset. Since
they have not been published thus far, interested readers may find a more complete description of the results in ~~the~~ Annex A,
including resulting emissions from fertilizers, crop production and animal excreta (dairy and beef cattle, poultry, swine, sheep,
goats, horses and other livestock). Consistent with the referenced studies cited above, emissions from managed excreta are
200 reported as AGL, and those deposited in pasture during animal grazing are reported under AGS. Resulting inventories of
annual emissions from agricultural activities spatially disaggregated at district level were ~~taken converted~~ to grids with a 0.1°
x 0.1°
resolution.

2.2.2 Chile

Chilean annual emissions were taken from the CR2-MMA-CR2-MMA dataset (CR2-MMA, 2018), based on the works of
Gallardo et al. (2018) and Mazzeo et al. (2018). ~~Such~~ This dataset is presented with a spatial resolution of 0.01°
205 x 0.01°, and includes 2014 emissions of reactive gases, GHG and particles, reported under the following aggregation: industries (which
includes emissions from energy), urban and non-urban road transportation (which only includes CO and NO_x for
reactive gases), residential consumption, and agricultural and forest fires.

Emission from industrial sources corresponds to the compilation of self-declared estimates by each facility
to the Chilean Repository of Emissions and Pollutants Transport. Neither the methodology nor the emission factors used to
210 estimate these emissions could be traced. ~~However, given that~~ For the particular case of SO₂, the local methodology for ~~the~~
emission estimates is based on sulfur content in fuels and in mass-flow balances in copper production processes,
which constitute the main SO₂ emitter activity in Chile (González-Rojas et al., 2021), ~~we have considered~~. For this reason,

215 and assuming that the information on sulfur content that is handled locally is reliable, and we have included the spatially distributed emissions as estimated in Chile in our dataset. For the rest of the species, we decided to exclusively adopt the spatial distribution and the share of the locally reported emissions and distributed the CAMS estimates by weighing them on the ~~CR2-MMA~~ ~~CR2-MMA~~ spatial distribution as follows:

$$E_{cell}(i, j, k) = E_{local}(i, j, k) \frac{\sum_{k=1}^N E_{CAMS}(i, j, k)}{\sum_{k=1}^N E_{local}(i, j, k)} \quad (2)$$

220 where $E_{cell}(i, j, k)$ are the emissions of species i and ~~sector category~~ j assigned to the cell grid k , N is the total number of grid cells covering the country, E_{local} represents the emissions locally estimated and E_{CAMS} the corresponding estimate from the global database.

225 Emission estimates from residential sources in the ~~CR2-MMA~~ ~~CR2-MMA~~ dataset cover only firewood combustion for all species considered in our work (Mazzeo et al., 2018). According to local experts, one of the most relevant aspects of air quality in the coldest regions in southern Chile is the presence of CO, NMVOCs and particles from burning of firewood in households. Although this practice is included under the residential category of global inventories, the corresponding estimated emissions 230 do not seem to be consistent with the magnitude of the air pollution situation observed at the local level (Huneeus et al., 2020b). In addition, by downscaling global inventories in the Metropolitan Region of Santiago (~~MRS~~, Chile's central region), Huneeus et al. (2020a) found out that residential emissions were strongly overestimated in global databases and attributed this inaccuracy to the use of population density as a proxy for the spatial distribution. Residential emissions from firewood burning are less relevant in the more temperate northern areas where air pollution is mostly linked to emissions of SO₂ and particles from the metal industry (Huneeus et al., 2020b). From this and assuming that: (i) residential firewood burning is a predominant source in the southern region, (ii) in central and northern regions this work improves the representation of the diversity of sources and local practices for the other fuel combustion categories, we have decided to replace the residential emissions of ~~the~~ CAMS with those of the ~~CR2-MMA~~ ~~CR2-MMA~~, at the risk of underestimating residential emissions in the central and northern regions by omitting those from fuels other than firewood (see (section 3.1)).

235 ~~Emissions~~ Local estimates of CO and NO_x ~~emissions~~ from urban and ~~non-urban~~ ~~non-urban~~ road transportation were ~~added~~ aggregated and reported in PAPILA dataset under the TRO category. Given that the ~~local inventory reports~~ ~~magnitudes and the spatial distribution of emissions from~~ ENE and IND (including use of solvent) ~~emissions together and that insufficient information for spatial disaggregation was available, we to report ENE + IND are reported in an aggregate way in the Chilean inventory, we decided to report them~~ under the IND ~~sector for the ease of Chile~~ category. Emissions taken from ~~CR2-MMA~~ ~~CR2-MMA~~ were extrapolated to 2015 and 2016 by applying CEDS trends (Hoesly et al., 2018) and projected to a 0.1° x 0.1° grid. Emissions from categories and species not estimated by the local inventory were taken from CAMS inventory.

2.2.3 Colombia

For the purpose of the PAPILA dataset, the only available information for Colombia was the emission estimates of CO, NO_x and SO₂ at the national level from all the categories of interest, except agricultural soils. These estimates were developed for 245 the Third National Communication of Colombia to the UNFCCC, covering the period 2010–2014 2010–2014 (IDEAM, 2017). Annual emissions from Colombia were extrapolated to 2015 and 2016 by applying linear regression forecast using the local time series and disaggregated using the spatial distribution of sources of the ~~base~~ CAMS inventory as follows:

$$E_{cell}(i, j, k) = E_{CAMS}(i, j, k) \frac{\sum_{k=1}^N E_{local}(i, j, k)}{\sum_{k=1}^N E_{CAMS}(i, j, k)} \quad (3)$$

where variables and indexes are those described in Eq. 2.

250 Although in this context the country ~~estimates~~ ~~reports~~ CO, NO_x and SO₂ emissions from ~~solid waste, wastewater and waste incineration, SWD~~ ~~emissions were taken from CAMS-SWD~~, CAMS reports them as zero. The latter precluded the spatial assignment of the locally estimated emissions, and for this reason it was decided to take the SWD category from CAMS. The reason for this decision was that although the magnitude of the emissions was available, there was no information on their spatial distribution and it was not possible to apply the methodology described above, since CAMS considers zero SWD 255 emissions for these species in Colombia.

2.3 Comparison of local and global datasets

A spatial analysis was performed following a similar approach to ~~that by~~ Trombetti et al. (2018) in their work on spatial ~~inter-comparison of top-down~~ ~~inter-comparison of top-down~~ emission inventories in European urban areas, in which the analysis was made in terms of normalized emission values by group of categories and for different urban domains in order to 260 become independent ~~from~~ ~~of~~ emission levels and to show the relative contribution of a certain group of emission activities in different areas. ~~However, we only need to compare two inventories and are also interested in observing~~ ~~Since in our work we are interested in comparing only two inventories without losing sight of~~ the differences in terms of magnitude, we ~~therefore propose a comparison of~~ ~~have adapted this approach by comparing~~ normalized emissions by category and urban domain normalizing them with respect to those from the CAMS dataset, ~~such~~ as shown in Eq. 4. In this way, we were able to compare both datasets 265 in relative terms and without losing information on the shares of each group of categories and the differences in the emission levels of each dataset.

$$\forall i, \forall J : E_{i,J}^*(d, area) = \frac{\sum_{k=1}^N E_{i,J}(d, k)}{\sum_{k=1}^N E_{i,J}(CAMS, k)} \quad (4)$$

where $E_{i,J}^*(d, area)$ and $E_{i,J}(d, k)$ are the normalized emissions and the emission levels, respectively, of the species i and group of categories J corresponding to the dataset d and the area (region, country or urban domain) covered by the total number N of cell grids k .

For this analysis, we grouped categories as ENE + IND, RES, TRO, and “[Other categories](#) [Others](#)”, and applied the analysis to (i) SA region, (ii) countries with local data (Argentina, Chile and Colombia), and (iii) urban domains from those countries [with local information on the spatial disaggregation that have implemented their own methodologies for the spatial distribution](#) of emissions. Urban domains were selected seeking to represent a wide variety of environmental and anthropogenic factors. In Chile, we have chosen three regions with different air quality concerns: Antofagasta (Northern region) with a strong presence of mining activity, Osorno (Southern region) a cold region where firewood burning dominates residential emissions, and the [MRS](#) [Metropolitan Region of Santiago](#) (Central region, [hereinafter Santiago](#)) with a mix of emission sources and a strong presence of road transport. In Argentina, we have chosen three urban domains where relevant research groups are located, hoping that this analysis would contribute to these activities. Those sites are: the Metropolitan Area of Buenos Aires ([MABA](#) [to facilitate reading, hereinafter Buenos Aires](#)) which is a coastal city and one of the main megacities in South America, Bahía Blanca (B. Blanca) which is a port city with an important industrial park, and Mendoza, one of the most important cities in the country that borders the Andes mountain range. A broader description of the studied areas is included in Table A3 and Figure A1 of the Annex A.

2.4 [Dataset performance evaluation](#) [WRF–Chem simulations](#): case study in [MABA](#) [Buenos Aires](#)

The performance of the PAPILA dataset in comparison with [that of CAMS](#) [CAMS](#), can be assessed using both inventories as input data [to air quality models was assessed of a regional model, implemented in the whole domain where local data have been integrated into the global dataset. This vast region, that includes the tropical Andes in Colombia, the dry Andes in Southern Chile and the Argentinean plateau towards the Atlantic coast, is characterized by diverse topographic features and vegetation patterns. In order to capture the differences in boundary layer process and surface energy budget in the whole area, a high-resolution model is needed, setup in each area where the main PAPILA/CAMS datasets changes have been made.](#) As a first step of this verification exercise, we present here a study focused on Buenos Aires, using the Weather Research and Forecasting Chemistry ([WRF–Chem](#) regional model version 4.1.2 ([WRF–Chem](#) v4.1.2) [regional model. The site chosen for this case study was the MABA, a megacity. This megacity is](#) strongly influenced not only by mobile and residential sources, but also by the presence of four big thermal power plants, an important industrial park and an international port. Simulations were conducted using the model over three nested domains with a highest horizontal resolution of 3 km centered in [MABA](#) [Buenos Aires](#). Two time periods were selected to cover summer (from 7 February to 5 March 2015) and winter (from 26 August to 17 September 2015) to assess the role of the emission estimates from the two inventories in the simulated air pollutant concentrations in the different seasons.

2.4.1 Model Description and Simulation Configuration

300 ~~WRF-Chem is a fully-coupled~~ WRF-Chem is a fully-coupled online chemistry transport model that simultaneously predicts weather and atmospheric composition (Grell et al., 2005). The simulations were done over 3 nested domains. The lowest resolution domain (d01) has a grid size of 18 km x 18 km (~~51° W-78° W, 15° S-57° S~~ 51° W-78° W, 15° S-57° S), and the highest resolution domain (d03) has a 3 km x 3 km grid covering ~~MABA~~ the metropolitan region of Buenos Aires and the surroundings. The coverage of the domains can be seen in Figure A2. All the simulations conducted in this study were 305 performed using a spin-up time of two weeks.

~~Lambert-conformal~~ Lambert-conformal projections were used. The physical parameterizations adopted for the three domains were: a) ~~The~~ Thompson scheme (Thompson et al., 2008) for Microphysics, b) ~~The~~ Grell 3D scheme for cumulus parameterization, c) The Yonsei University scheme for boundary layer processes, d) ~~The~~ MM5 similarity scheme for surface processes, e) ~~The~~ RRTMG scheme to compute long and shortwave radiation. The chemistry in these simulations was modelled 310 using the GOCART bulk aerosol scheme for aerosol phase chemistry along with RADM2 for ~~gas-phase chemistry~~for aerosol phase chemistry~~gas-phase chemistry~~. The initial and boundary conditions were taken from the NCEP Final Operational Global Analysis data (FNL), available at a resolution of 1 degree every 6 hours (NOAA, 2000).

~~The~~ FINN Fire Database was used for fire emissions (Wiedinmyer et al., 2011), ~~the~~ MEGAN Biogenic database for biogenic emissions (Guenther et al., 2006) and sea salt emissions from GOCART were also included.

315 Reported annual emissions in 2015 from the two inventories were processed to produce ~~hourly-resolved~~ hourly-resolved emissions at resolution of each of the domains. ~~For reactive gases we used the emission inventory presented in this article. The impact of aerosols emissions was also included in the analysis, enriching PAPILA database with EDGARv4. Since PAPILA dataset only includes reactive gases, for aerosol emissions, the ones of the CAMS simulation were used: EDGAR v4.3.2 Global emission database for PM_{2.5} and PM₁₀ and CAMSv4 CAMS v4.1 for OC, BC and VOCs. Temporal emission patterns were computed by an iterative method, focusing on capturing daily and seasonal variations of CO, and PM surface mixing ratios observed. However, the results presented in this article will only cover the species included in the PAPILA inventory.~~

320 For ENE, TRO and RES monthly emission patterns were defined to breakdown total annual emissions into monthly fluxes (see Fig. A2). Emissions from other categories were evenly distributed throughout the year. RES monthly cycle was established from the reports on natural gas consumption reported in national statistics (ENARGAS, 2021) for residential and commercial activities in the ~~MABA~~ activities in Buenos Aires. This profile shows a maximum during winter linked to the increase in residential heating. Similarly, TRO monthly cycle was defined from the total fossil fuel consumption from the road transport reported in the statistics for the entire country (Secretaría de Energía, 2021). For ENE, the same source of data was used to obtain monthly fossil fuel sales for thermal power plants in the Buenos Aires Province. Weekly cycles taken from ~~PREP-CHEM~~ PREP-CHEM (Freitas et al., 2011) were applied to the resulting total monthly emission fluxes. The ~~hourly emission patterns were computed with the model by an iterative method in preliminary simulations, taken as seed the diurnal variations for transportation emissions~~ diurnal cycles were adapted from those reported by Wang et al. (2010) ~~and focusing to capture the observed daily cycles and seasonal variation of CO, and PM surface concentrations measured, focusing on reproducing Buenos Aires traffic patterns observed~~ in the two monitoring stations, ~~Parque Centenario and Córdoba~~ Parque Centenario and Córdoba. With this approach, the best configuration obtained with the simulations includes three hourly emission patterns: one related to diesel vehicle emis-

335 sions, defined using PM observations, and the other associated with gasoline car emissions in winter and in summer, defined with the measured concentrations of CO and NO_x.

2.4.2 Model evaluation

Highest resolution model outputs using these two emission inventories were evaluated against CO and NO_x ~~ground-based~~ ground-based observations from the available monitoring stations in ~~MABA~~ Buenos Aires (See locations of the sites in Figure A2). Air pollutant data from the Environmental Protection Agency of Buenos Aires (APRA) include hourly measurements of NO_x and CO in two sites, Cordoba (34.60° S, 58.39° W) and Parque Centenario (34.61° S, 58.44° W). Cordoba's site is located in a commercial area with high vehicular flow and very low incidence of stationary sources while Parque Centenario is located in a residential area next to an arboreal space with medium vehicular flow and also very low incidence of stationary sources. As the air quality database is at hourly resolution, the model was also sampled at every hour.

345 The model evaluation was mainly focused on the effects of enriching the CAMS inventory with local inventories on the simulated air pollutant concentrations. For this purpose, median and percentiles for the entire period were evaluated. Also, mean daily concentrations were calculated to inspect whether model performance of both inventories was consistent and satisfactory. ~~Well-accepted~~ Well-accepted statistical measures such as normalized mean bias (NMB), normalized mean gross error (NMGE) and the fraction of predictions within a factor of two (FAC2) were used (Wang et al., 2021). These statistical 350 metrics were calculated using the following expressions:

$$NMB = \frac{1}{N} \frac{\sum_{k=1}^N S_k - O_k}{\sum_{k=1}^N O_k} \quad (5)$$

$$NMGE = \frac{1}{N} \frac{\sum_{k=1}^N |S_k - O_k|}{\sum_{k=1}^N O_k} \quad (6)$$

$$FAC2 : \frac{N(0.5 < \frac{S_k}{O_k} < 2)}{N} \quad (7)$$

355 where S_k and O_k are the simulated and observed hourly average concentrations respectively, and N is the total number of observations. The model was sampled at each measuring location using grid interpolation and compared with the ~~ground-based~~ ground-based observations for the calculation of statistical performance metrics.

3 RESULTS AND DISCUSSION

Table 2 reports the resulting 2015 annual emissions of CO, NO_x, NMVOCs, NH₃ and SO₂ corresponding to the sum of all categories in the PAPILA dataset, for different domains: South America, Argentina, ~~Chile~~ Buenos Aires, Bahía Blanca, ~~MABA~~,

360 ~~Mendoza, Antofagasta~~^{Mendoza, MRS and Osorno} Chile, Santiago, Antofagasta, Osorno and Colombia, in comparison with the corresponding emissions levels from CAMS. In addition, Figure 3 depicts the shares of each group of categories (ENE + IND, RES, TRO, and Others) to the total emissions of each species and for all the aforementioned domains. They are expressed in a normalized way with respect to the sum of all categories here ~~analysed~~^{analyzed} in CAMS for each corresponding species and domain. In this way, the emissions of each species in CAMS correspond to the sum of the share of each group of categories, 365 adding up to a total of 1.00. On the other hand, the shares of each ~~sector~~^{category} in PAPILA can be compared with those in CAMS and add up to a total greater or less than 1.00 according to the differences in the sum of all the categories indicated in Table 2.

370 The spatial distribution of 2015 annual emissions of CO, NO_x, NMVOCs, NH₃ and SO₂ is shown in Figure 4 for the sum of all categories. In addition, this figure includes maps with the differences between PAPILA and CAMS datasets for each species, depicting the differences in terms of intensity and location of emission sources. For comparative purposes, emissions from agricultural fires have been subtracted from the sum of all categories in CAMS database.

375 In what follows results are presented firstly by species, highlighting the most relevant aspects of the 2015 emission (section 3.1). Then, surface concentrations of CO and NO_x obtained from the use of PAPILA dataset as input information in a chemical transport model in ~~MABA~~^{Buenos Aires}, compared to those obtained using CAMS are presented (section 3.2). The section ends with an analysis of the local aspects that may have generated the difference between both emission databases (section 3.3).

3.1 Local-global comparison by species

3.1.1 Carbon monoxide

380 Local estimates of CO for Argentina and Colombia presented lower CO annual emissions than those in CAMS, the largest differences occurred under the TRO category. Although the local estimates for TRO in Chile also showed significant smaller levels this difference was masked in the total CO national estimates by the larger emissions from the residential category, even after having omitted CO emissions from fuel combustion other than firewood. Lower CO PAPILA emissions in Argentina (-39%) and Colombia (-56-54%) were compensated by larger PAPILA emissions in Chile (58%)~~resulting in~~^{According to} CAMS dataset these three countries are only responsible for 25% of the total SA emissions and hence the impact of the changes 385 introduced in this work on total SA is very limited. This same situation occurs with the other species, for which it is found that these three countries are only responsible for 20-25% in the case of NO_x, NMVOCs and NH₃ and 31% for SO₂ (Table 2). However, when analyzing the impact on these three countries together, and even when these countries compensate for each other, a difference with CAMS for South America of only -5% CO CAMS emissions of -19% is observed.

390 At the urban level, in ~~the~~^{MABA} Buenos Aires domain PAPILA emission estimates resulted 12% higher than those from CAMS, this difference was mainly associated with higher local emissions from TRO and ENE + IND even ~~when lower emissions resulted with lower emissions~~. In the same way, Mendoza and B. Blanca exhibited lower CO total levels, mainly associated with differences in TRO and in less extent in RES. In B. Blanca, this difference masked the larger emissions

Table 2. Summary of annual emissions by domain for 2015 (Gg y⁻¹)

		CO		NO _x		NMVOCs		NH ₃		SO ₂	
2015 (Gg y ⁻¹)		PAPILA	CAMS	PAPILA	CAMS	PAPILA	CAMS	PAPILA	CAMS	PAPILA	CAMS
South America	South America	31225.2	32780.5	5791.0	5637.1	12186.0	11297.0	4885.5	5113.3	3248.4	3158.0
		31263	32780	5803	5637	12186	11297	4886	5113	3275	3158
Argentina	Argentina	2289.37	3740.3	901.2	660.2	548.7	1064.8	323.4	536.1	101.5	252.2
		2289	3740	901	660	549	1065	323	536	102	252
Colombia	Buenos Aires	1040.1	2343.4	328.3	367.7	798.0	798.0	395.1	395.1	177.5	146.3
		336	300	118							
Chile	Chile	3279.6	2080.5	307.3	355.0	1938.5	533.5	213.8	228.8	782.5	572.6
		335.7	299.9	118.4		80.9	63.8	3.5	5.9	16.3	49.1
							210				
	Bahia Blanca	9.6	21.5	16.8	7.3	2.6	7.3	0.3	0.4	3.7	16.1
	Mendoza	39.9	49.1	16.8	10.5	8.0	25.6	1.0	0.6	1.2	5.5
MRS	Chile	112.2	345.4	307	355	1939	533	214	229	782	573
		3280	2081								
	Santiago			35.0	31.8	30.7		1.8	9.1	14.7	38.9
		112	345				136.6				
							137				
	Antofagasta	24.5	24.5	2.0	2.0	1.5	7.6	0.1	0.6	7.9	2.9
	Osorno		19.5	4.4	1.6	88.7	3.6	1.5	1.4	0.4	0.3
		136.7									
		137									
Colombia		1078	2343	341	368	798	798	395	395	204	146

by a factor of five in the local estimates of ENE + IND with respect to the global dataset. By downscaling B. Blanca urban domain we identified the absence of emissions from shipping activities (**inland: SHP, and international: SHP-INT**) in the global

395 inventory. While emissions from SHP **within the continental area** were estimated locally, **estimates for SHP-INT were not available and therefore they were taken from global estimates** offshore emissions were taken from CAMS, which reports zero




Figure 3. Normalized ~~sectoral~~ breakdown of PAPILA emissions compared with CAMS inventory by domain [and category group](#) for 2015. Total CAMS emissions for each domain equal to 1. ENE + IND: energy and industries; RES: residential and commercial combustion; TRO: road transportation; Others: non-road transportation, fugitive emissions, agricultural soils, agriculture livestock, navigation and waste.




Figure 4. (left) Spatial distribution of 2015 PAPILA emissions (Gg y^{-1}) and (right) Difference between PAPILA and CAMS inventories by species in 2015.

400 emissions for this region. In this domain, international navigation is emissions from navigation activities are a concern since its port activity is almost as relevant as that of the international port of the Buenos Aires city (Ports, 2021). Although the absence of this source was not reflected in this comparative analysis, it is relevant to point it out that it could lead to underestimation of surface concentrations when modelling air quality in the region.

405 In the MRS Santiago, local estimates of total emissions were almost 70% lower than global ones, this difference is attributable to the two locally estimated categories that were included in this work (TRO, RES) but also to ENE + IND, categories for which we used a combination of the CAMS emission estimates with national information of location and emission shares. In Antofagasta, although total emissions levels from both datasets were similar, there were substantial differences in the contributions by category: emissions from ENE + IND are almost seven times larger in PAPILA than in CAMS, and while RES and TRO emissions are negligible in the local estimates, according to global estimates they contribute to almost 90% of the domain's emissions. On the contrary, in Osorno local estimates for the sum of all categories were seven times larger than those in CAMS, emissions coming almost entirely from residential firewood burning.

3.1.2 Nitrogen oxides

410 Local estimates of national NO_x emissions for Chile and Colombia were lower by 13% and 117%, respectively, than those in CAMS. In both countries, the main responsible for these differences was the TRO category and to a lesser extent the lower emissions of RES, which in the case of Chile were due in part to the omission of the burning of fossil fuels in this category. In Colombia, local SHP estimates contributed to this difference, being the difference was partially offset by considerably higher emissions from TNR. Local estimates for Argentina resulted in higher total NO_x emissions (37%) with very different sectoral
415 sector contributions to this difference. The contributions by category (from highest to lowest) were TRO, ENE, AGS and RES, partially offset by lower emissions from IND. All in all, NO_x estimated emissions with local data for the whole SA region
were 3three countries together were 12% higher than those reported by CAMS.

420 As seen in Figure 3, all urban domains showed higher local emissions, except Antofagasta with a barely noticeable difference. In B. Blanca, the most relevant differences were the larger emissions in ENE + IND and SHP, category for which CAMS reported zero emissions while according to the local data it represented 12% of the domain's emissions, despite the omission of the international port as a source of emission. Relevant larger emissions existed for TRO, ENE and RES in the MABA
425 Buenos Aires and Mendoza, together with significantly smaller emissions from IND. The larger emissions from other categories in the MABABuenos Aires and B. Blanca are mainly attributable to SHP, and although the impact of Other sectorsother categories on the total budget in these domains was negligible, local estimates showed considerably higher levels than CAMS for emissions from agricultural activities and FEF, category for which the global database attributed zero emissions in the three urban domains.

430 In the MRS Santiago, local emission estimates were slightly higher than those from CAMS, a difference mainly attributable to larger emissions from TRO, partially offset by smaller emissions in ENE + IND. In Antofagasta, by contrast, the larger emissions from ENE + IND sectorscategories were almost completely masked by smaller TRO emissions. Local emissions from RES were strongly underestimated in these two domains where according to CAMS data, RES is a minorityminor

emission source. In Osorno, local estimates of total emissions exceeded by a factor of almost three those of CAMS, being ENE + IND the ~~sectors~~categories with the greatest contribution to this difference and to a lesser extent TRO and RES. It is worth mentioning that the differences observed in ENE + IND in Chilean urban domains are exclusively associated with the local information on the spatial distribution and shares of NO_x emissions, and not with a local estimate of the magnitudes.

435 3.1.3 ~~Non-methane~~Non-methane volatile organic compounds

Local estimates of NMVOCs emissions for Argentina were 48% lower, this difference is mainly attributable to IND (which in this work includes solvent production and use). CAMS did not report NMVOCs emissions from agricultural activities (neither livestock nor soils) for any country in South America, while the local estimates for Argentina showed that 11% of the NMVOCs emissions came from these activities. PAPILA estimates of RES emissions in Chile (the only category locally 440 estimated) exceeded those of CAMS by more than an order of magnitude, which was reflected in a total emission level three times larger for the country. ~~The contribution of local Argentinean information, and that from RES in Chile, Although Argentina partially compensates for the difference introduced by Chile's local data, considering both countries jointly, the changes made~~ resulted in larger ~~regional emissions by 8% emissions by 56%~~.

Local estimates showed important differences in total emissions for the three Argentinean domains (around ~~60-70~~60-70% lower than CAMS), being IND the main contributor. Smaller emissions from FEF were observed in B. Blanca and Mendoza; while TRO contributed to these differences in the first domain and counteracted them in the second. In ~~the MABA~~Buenos Aires emissions from FEF and TRO were considerably larger than those in CAMS. Even when estimates from RES in PAPILA were around 80% lower than those of CAMS they exhibited lesser impact on the differences between the two datasets and on the total emissions in each domain.

450 Local estimates for ~~the MRS~~Santiago and Antofagasta were significantly smaller (around 80%) than the global ones, the difference is mainly attributable to the adopted local information on locations and emission shares for ENE + IND. On the contrary, local estimates for Osorno showed emissions more than 24 times larger than those of CAMS, almost exclusively attributed to the incorporation of local information on firewood consumption in the RES category in cold areas of the country.

3.1.4 Ammonia

455 Similarly to NMVOCs, the only two countries with local data on NH₃ are Chile and Argentina. At ~~the~~ national level, the inclusion of local information is reflected in differences of -7% of NH₃ emissions in Chile (only attributable to RES) and -40% in Argentina, where smaller emissions from AGS were the main responsible for that difference partially offset by larger emissions from AGL. These two categories represent the main sources of NH₃ emissions in the country with a contribution of 72% from soils and 24% from livestock, according to the local estimates. Smaller emissions in local estimates of Argentina 460 and Chile were reflected in a difference with CAMS of ~~only 4% of total emissions in South America~~30% of the emissions of both countries together.

Although the impact of emissions from urban domains on the total levels of each country was negligible (around 1%), big differences were found at the category level between the two datasets. In Mendoza, local estimates resulted in larger

emissions by around 70% in total levels, mainly attributable to larger emissions from agricultural activities partly countered
465 by substantially lower emissions from IND. In B. Blanca, the difference in total levels was around -30% mainly attributable to ENE and AGS, while in ~~the MABA the Buenos Aires the~~ difference was around -40% where the main contribution to this difference was IND, partially offset by larger emissions from TRO, RES and other categories as SHP and AGL. Locally estimated emissions from RES were larger in the three urban domains.

Both Antofagasta and ~~the MRS Santiago~~ showed smaller total emissions by around 80% as a result of relocating ENE + IND
470 emission sources, and replacing emissions from RES by the local inventory. In Osorno, slightly larger local estimates from RES and IND were observed. However, as in the case of urban domains in Argentina, the contribution of each domain to the total emissions in Chile was negligible.

3.1.5 Sulfur dioxide

Local estimates of SO₂ emissions in Argentina were 60% lower than those by CAMS for the sum of all categories, being
475 IND, ENE and RES the main contributors to that difference (and SHP in a lesser extent), while TRO emissions were considerable higher (around eight times) in local estimates. For this country, these larger CAMS emissions were associated with the sulfur content adopted, mainly from solid fuels, since the national mineral coal has lower sulfur content (370 kg SO₂ Tj⁻¹) than those imported (1100 kg SO₂ Tj⁻¹), and because the national/imported ratio presented high variability between 2011 and 2015 (TCN, 2015). Colombia showed larger emissions from ENE, IND and TRO, partially offset by lower emissions from
480 RES ~~and SHP~~ in the local estimates, and although negligible at the national level the emissions from FEF were significantly higher than in CAMS. As in Argentina, the sulfur content in the coal used was highly variable, due to the different sulfur levels that the country's coal fields present. In the same way as Colombia, Chile showed larger emissions from the sum of all categories as a consequence of the inclusion of local data in ENE + IND, differences mainly related with sulfur emissions from the relevant copper mining activities that take place along the country, which were ~~non-not~~ fully covered by CAMS. The
485 lower TRO emissions reported by CAMS for Argentina and Colombia seem to be related in part to the methodology used for projections, that assumes a sustained reduction in sulfur content from 2012 to 2015. Nevertheless, this reduction did not occur in any of the countries: while Colombia introduced prior to 2012 strong restrictions to fuel quality, in Argentina these restrictions for the fuels used by heavy duty trucks (the main emitters) did not take place. ~~All this together results-Although the differences introduced by the local data for Chile and Colombia are partially offset by Argentina, all these together result~~
490 in larger emissions by ~~only 3% in the entire region~~^{12%}.

Local estimates in ~~the MABA Buenos Aires~~ showed smaller emissions by 67%, mainly associated with lower emissions from IND and RES (80-90%), the latter with less impact on the totals. This situation may be related to the fact that the proportion of ~~S-emitting industries in the MABA sulfur emitting industries in Buenos Aires~~ is lower than in the rest of the country. Also with little impact, and offsetting these aforementioned differences, increases were observed in estimates from
495 thermal power plants, inland navigation and transportation (TRO and TRN). Both B. Blanca and Mendoza showed smaller emissions by 77%, mainly attributable to ENE in the first case and IND and RES in the second, where at the same time an

increment in emissions from ENE was observed. Although the contribution of the TRO to the total emissions was minor in the urban domains of the country, the larger emissions estimated locally with respect to ~~the~~-CAMS is particularly noticeable.

500 ~~The MRS-Santiago~~ showed a difference in local estimates of around -62% mainly attributable to ENE + IND, while the result of having included local estimates of emissions from these ~~sectors~~ ~~categories~~ in Antofagasta was reflected in total levels of SO₂ almost three times larger than in CAMS. In these two domains, local estimates attributed to ENE + IND a contribution of more than 99% of total emissions. In Osorno, although local emission estimates for RES were lower than CAMS, total emissions in the domain resulted 16% larger than in global estimates as a consequence of the difference in emissions from ENE + IND.

3.2 ~~Model~~ Case study: model evaluation and results

505 Table 3 summarizes the overall model performance of PAPILA and ~~CAMS-based~~ ~~CAMS-based~~ results for hourly CO and NO_x concentrations ~~for Buenos Aires case study~~. For the winter period, ~~PAPILA-based~~ ~~PAPILA-based~~ results had lower normalized mean error than ~~CAMS-based~~ ~~CAMS-based~~ results; the negative bias was larger for the CAMS emission run, exceeding in all cases more than 12% for both CO and NO_x except for NO_x in Parque Centenario. FAC2 was also better in PAPILA simulation. Differences in the concentrations resulting from both runs were consistent with those exhibited between 510 the inventories. In terms of CO emissions, PAPILA dataset emissions were 12% higher than CAMS being road transportation the main responsible followed by industry and residential sources. On the other hand, NO_x emissions were 46% higher in PAPILA, with significant discrepancies in emissions mainly from TRO followed by ENE and IND. Emissions in ~~the MABA~~ ~~Buenos Aires~~ are typically lower in summer because of decreasing traffic levels, ~~null~~ ~~no~~ heating requirements and lesser use 515 of liquid fuels by thermal power plants, which burn almost exclusively natural gas during the warmer period. Lower emissions levels coupled with favorable meteorological conditions for air pollutant dispersion conduct to lower concentrations levels in summer. Thus, the ~~goodness of the PAPILA-based results exhibited for winter were not that apparent for summer~~ results for the summer simulations were not as conclusive as for winter simulations. NMB was still negative with CAMS emissions, consistent with winter period results and better FAC2 for PAPILA's run. Therefore, this highlights the importance of having accurate inventories especially for winter when the highest emissions and worst dispersion conditions occur.

520 Figure 5 shows median and percentiles for ~~CO and~~ NO_x hourly concentrations. The site Parque Centenario (residential) was better reproduced by the model with both inventories compared to Córdoba (commercial area with high traffic flow), especially in winter where the traffic flow in the city has a greater influence on the dynamics of these pollutants. However, we must also highlight that Cordoba monitoring station is located in a corner with high traffic flow and the sampling is at 2 m high. Therefore, the measurements may be capturing an overestimation of the real average activity in the area.

525 Scatter plots of daily mean concentrations using ~~the~~ PAPILA inventory (Figure 6) depicted a good agreement between observations and model ~~results~~, in winter more than in summer for CO and the other way around for NO_x. Concentrations estimated with ~~the~~ CAMS inventory tend to be underestimated in all cases.

530 All in all these results show a better agreement between observations and simulations using PAPILA ~~dataset~~ than CAMS, to represent surface concentrations of CO and NO_x in ~~the MABA~~ ~~Buenos Aires~~. However, these emission improvements do not fully explain the underestimations of the model especially for CO concentrations with respect to ~~the~~ measured data.

Table 3. Summary of statistical metrics used for evaluating model performance to simulate surface CO and NO_x concentrations in PAPILA and CAMS experiments for both sites. In Bold is highlighted the inventory that has the best performance for that metric in each period.

		Córdoba				Parque Centenario			
		Winter		Summer		Winter		Summer	
		PAPILA	CAMS	PAPILA	CAMS	PAPILA	CAMS	PAPILA	CAMS
CO	NMB	-0.18	-0.41	-0.15	-0.39	0.02	-0.39	-2E-3	-0.28
	NMGE	0.43	0.51	0.58	0.56	0.39	0.50	0.60	0.52
	FAC2	0.74	0.53	0.59	0.51	0.83	0.54	0.62	0.62
NO _x	NMB	0.07	-0.32	-0.08	-0.41	-0.07	-0.28	0.13	-0.12
	NMGE	0.58	0.63	0.52	0.69	0.62	0.63	0.61	0.63
	FAC2	0.66	0.46	0.66	0.37	0.55	0.45	0.64	0.52

NMB: Normalised mean bias, NMGE: normalised mean gross error, FAC2: fraction of predictions within a factor of two.

3.3 ~~PAPILA-CAMS~~^{PAPILA-CAMS} main differences

Our results show relevant differences between the PAPILA and CAMS datasets, both in terms of emission levels and their spatial distribution. They also served to exemplify the ~~goodness of the PAPILA-based quality of the PAPILA-based~~ simulated surface mixing ratios in the city where concentrations were ~~analysed~~^{analyzed}.

535 The reasons behind the observed differences are diverse and are mainly linked to the activity data and to the methodologies applied to estimate and spatially distribute the emissions. Given the limited availability of local data, the emission factors for the compounds covered in this work were mostly based on default values in both the global and local datasets considered. In general, it is observed that the ~~CAMS-GLOB-ANTv4~~^{CAMS-GLOB-ANT} v4.1 applies linear trends on the emissions of the aggregated categories, based on the ~~global~~ estimates of the ~~EDGARv4~~^{EDGAR} v4.3.2 of the year 2012. ~~This is different from what has been done in this work~~, ~~since for the three countries~~ In contrast, in our work three different situations can be identified: (1) for Colombia, locally estimated trends were applied based on 2014 ~~was taken as the base year, and while the local emission estimates~~, (2) for some categories in Argentina annual estimates for the entire period were applied, and (3) for the remaining categories in Argentina and for the Chilean inventory the same methodology than ~~that the~~ used by CAMS was applied ~~to extrapolate to 2015 and 2016 for Chile and some categories for Argentina, locally estimated trends were applied for Colombia~~, ~~but based on local estimates for 2014~~. In this way, local inventories are not only based on ~~region-specific~~ ^{region-specific} information, but also the extrapolation of a shorter time period reduces the uncertainties associated with the activity data strongly linked to ~~short-term~~ ^{short-term} variations and technological changes.

540 For those activities related to fuel consumption, local inventories used the information reported in the national energy balance and other national energy statistics while global inventories are based on the information reported by the countries to the International Energy Agency ([IEA](#)), which is consistent but not exactly the same ~~than as~~ that reported in the national statistics since the IEA processes the information received (IEA, 2020). Moreover, although these statistics adequately represent




Figure 5. Box-Plots of hourly CO and NO_x in both sites for summer and winter simulations. The dots represent the average value, the thick lines represent the medians for each period. The vertical hinges represent data points between the 25th and 75th percentiles. The whiskers represent data points between the 5th and 95th percentiles. CO concentrations are in ppm and NO_x in ppb.

the national energy balances, it is worth pointing out the lack of specificity in terms of spatial disaggregation. A relevant aspect of the fuel consumption patterns for power generation in the three countries analysed, is their inter-annual and inter-regional variation, which in turn are strongly correlated with hydraulic availability, the

555 water availability for hydropower generation, not captured by the extrapolations. In addition, in order that electricity supply matches demand, some short-term technological changes are often resorted to. For example, the incorporation of diesel-based motor diesel-fuelled generators located in different urban centers in Argentina such as the MABA-Buenos Aires in 2014 (CAMESA, 2021). Although the diesel consumed in these generators was reflected in international statistics, they did not distinguish between the gas oil used for this purpose and that used by combined gas cycles and could not reflect 560 their location and operating regimes.

Another relevant aspect of national and therefore international statistics is the lack of reliable information on firewood consumption, widely used in rural and urban areas of SA and even in some urban areas, such as the cold regions of Chile. This fact also impacts on the correct representation of the replacement of firewood by LPG or Natural Gas that took




Figure 6. Scatter plots for observed and modeled daily concentrations for both sites. Values within the dotted lines represent the fraction of values that are within a factor of 2 of the observations (FAC2). CO concentrations are in ppm and NO_x in ppb.

place in Argentina in the last decade, due to higher production of ~~non-conventional~~ non-conventional shale gas in Vaca Muerta 565 basin (El País, 2015) and the resulting reduction in fossil fuel prices. ~~In addition, the use~~ Additional differences between local and global datasets are related with different resolutions of the population ~~as a driver~~ distribution maps used as proxies for the spatial distribution of the emissions from some categories, ~~such as RES, implies another difficulty that is linked to the resolution used. Generally, international databases use country and subnational borders as proxy population data for the spatial disaggregation (Janssens-Maenhout et al., 2019), and this low-resolution population density, in comparison with that~~ 570 from locally developed inventories, may also be a source of spatial uncertainties. The latter ~~Local inventories use population density information based on higher resolution maps than those used by the global ones. This~~ is clearly noticeable not only in the ~~local-global-local-global~~ differences found by downscaling urban domains, but also in the spatial coverage of RES emissions in global inventories, where emissions are assigned even to large ~~non-populated~~ non-populated areas, such as the Amazon rainforest or some desert areas of the region (Figure 4). ~~It is also worth mentioning that, unlike local inventories,~~ 575 ~~CAMS treats countries uniformly without correcting for climatic zones, which vary widely within many of the SA countries.~~ Broader discussions on emissions from RES are given by Puliafito et al. (2021) and Álamos et al. (this issue) for Argentina and Chile, respectively.

For ~~sources other than fuel combustion for energy purposes~~ non-combustion sources, as many industrial processes, ~~population estimates are used as drivers for the~~ CAMS projections (based on CEDS trends) ~~use population as driver~~ (Hoesly et al., 2018). 580 This approach may not be the most appropriate for many countries in the region, where changes in economic policies and even

the occurrence of economic crises are frequent, affecting not only consumption patterns but also the relocation of activities dependent on regional economies. Added to this, substantial differences have been observed in terms of the location of the IND sources in both inventories, probably attributable to the drivers used by the global databases for this purpose, with a very low presence of sources throughout the Argentine territory in CAMS, and a striking abundance of sources distributed over the 585 central and northern region in Chile, which however do not reflect the heterogeneous share of emissions shown by the local distribution.

~~With the advancement of satellite tools, there are many aspects related to the spatial disaggregation of emissions that are improving. Although the representativeness of agricultural emissions in global databases continues to improve over time, as the allocation of agricultural emissions. However, there are many aspects that the global inventory development global methodologies have so far failed to replicate, whereas. In this sense, the incorporation of information generated from a local perspective has provided local inventories with greater precision for estimating emissions. In addition to local information has given the inventories a greater capacity to reflect the specific agricultural practices of the region, such as the predominant use of grazing for cattle farming or, the use of large proportions of urea for the fertilization of crops, during the last decades as well as economic, natural and technological changes have occurred that have impacted on the sector's emissions and which are not reflected in global inventories that have occurred during the last decades~~ (Castesana et al., 2018, 2020).

Lastly, and with the aim of contributing some aspects that may improve both global inventories and the dataset presented here, we list some information gaps identified when using the version ~~CAMS-GLOB-ANT~~ v4.1 as a base inventory for the region: (i) emissions from domestic and international civil aviation are not included; (ii) ~~domestic and international waterborne navigation~~ are not reflected by downscaling the port city B. Blanca in Argentina; 600 (iii) CAMS assigns zero FEF emissions for species other than NMVOCs in all the downscaled urban domains; (iv) in the whole region CAMS assigns zero emissions of NMVOCs from agricultural activities, both for those from animal excreta (manure ~~managed-AGL~~ and deposited in ~~pasture-AGS~~) and those from crop production (AGS); and (v) CAMS assigns zero emissions of CO, NO_x and SO₂ from SWD ~~in Chile and Colombia even when the latter reports the occurrence of emissions in its local estimates for all SA, except for Brazil and the Guianas~~. Item (iii) may be attributed to the 605 spatial distribution of FEF emissions in the global inventory, but given that this pattern is repeated in the six ~~analysed-analyzed~~ domains and since in most of the urban domains considered there exist industrial facilities that may use venting and flaring to dispose their waste gases, the assumption of zero emissions for species other than NMVOCs seems to be more related to the omission of venting and flaring as emission sources under FEF (Granier et al., 2019). For its part, the item (v) brings to light the problems around waste management in the region, ~~where and consequently around the representativeness of the associated emissions. In SA, waste is often disposed of in open dumpsites and, particularly regarding CO, NO_x and SO₂ emissions, uncontrolled open burning occurs, and these practices are not reflected in local or global emission maps (UNEP, 2018). All these gaps in the base inventory were replicated in addressed in the~~ PAPILA dataset, except ~~for the case of~~ FEF emissions and NMVOCs emissions from agricultural activities in Argentina.

4 Data availability

615 Gridded maps with all inventories per species and year are available as netCDF4 (Network Common Data Format) files for
the regional domain (32° W– 120° W, 34° N– 58° S) at a resolution of $0.1^{\circ} \times 0.1^{\circ}$ (PAPILA dataset) and can be accessed through
the open access data repository <http://dx.doi.org/10.17632/btf2mz4fhf.3>, under a CC-BY 4 license (Castesana et al., 2021). The
dataset includes all necessary attributes to be easily used in air quality simulations, such as molecular weight, projection and
units. In addition, temporal profiles applied in the inventory evaluation for the Metropolitan Area of Buenos Aires, Argentina,
620 are available in Appendix A of this work.

5 Conclusions

This work presents the results of the first joint effort of South American countries to generate regional maps of emissions inventories. ~~Although it includes the emissions estimates made for a limited number of countries, it shows the enormous potential for the improvement of air quality studies in the region, by including local data into global databases and building together an inventory of the reactive gases CO.~~ With a mosaic approach, we developed a new anthropogenic emission inventory 625 for South America, named PAPILA, harmonizing inventories of Argentina, Chile and Colombia in a uniform format. To such end, locally developed emission inventories of these countries have been supplemented with the existing data of the global dataset CAMS-GLOB-ANT v4.1 to fill the gaps. PAPILA dataset consists in annual emission gridded maps of reactive gases, namely CO, NO_x, ~~NMVOCS~~, NMVOCs, NH₃ and SO₂, for the period 2014–2016 with a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$ 630 over a domain defined by 32° – 120° W and 34° N– 58° S.

~~The results obtained through modelling using the inventory presented here as input are promising. This study introduces significant improvements to global inventories mainly observed in the spatial distribution of emissions, in a better representation of the interannual variability of emissions strongly related with short-term variations and technological changes, and in an improvement of the representation of emissions from informal activities, such as the use of wood for heating and cooking.~~ 635 Relevant differences were found between both inventories at country and urban levels for all the compounds analyzed. At the national level, PAPILA presents lower CO emissions (-39% and -54%, respectively) for Argentina and Colombia while the opposite is observed in Chile (58%). For NO_x, PAPILA emissions for Chile and Colombia are lower than CAMS by 13% and 7%, respectively whereas Argentina presents larger emissions (37%). For NMVOCs and NH₃, the only two countries with local data are Argentina and Chile. While for NMVOCs PAPILA estimates lower emissions for Argentina (-48%) and around 640 three times larger for Chile, for NH₃ PAPILA estimates lower emissions in both countries, namely -7% for Chile and -40% for Argentina. Larger SO₂ PAPILA emissions in Chile and Colombia are offset by lower emissions in Argentina (-60%) and for the three countries together result in larger PAPILA SO₂ emission by 12%. However, based on CAMS emissions in these three countries represent only 31% of the total SA emissions for SO₂ and between 20 and 25% for the rest of the species. Therefore, 645 further studies are necessary to complement this work with local data from other countries with significant emission levels, such as Brazil, Peru and Venezuela.

650 A complete analysis of the performance of the inventory developed in this work could be provided by comparing model simulations with surface and satellite observations in the whole domain where local data have been integrated into the global dataset. Since South America is a wide region with significant differences in terms of latitudinal extension, topography and vegetation patterns, a high-resolution model would be needed to capture the boundary layer process and surface energy budget of each area where the main changes have been introduced. As a first step of this verification exercise, we have presented a case study focused on the Metropolitan Area of Buenos Aires, using WRF-chem regional model. Although there is room for improvements that may be linked to both emissions and other processes, the CO and NO_x ~~results in the Metropolitan Area of Buenos Aires were able to reasonably~~ modeling results showed a better performance with PAPILA dataset over CAMS to reproduce surface observations, ~~improving those obtained from the use of global inventories~~ in this big urban area.

655 Nevertheless, further modeling studies with PAPILA and CAMS datasets are needed over other key areas of Argentina, Chile and Colombia.

660 This work ~~not only~~ highlights the strengths and weaknesses of ~~not only global inventories but also~~ global inventories, but also ~~those of~~ local ones. Although the latter improve the representativeness of the estimates, the groups that generate information on emissions in the region do not necessarily have the same objectives (some are mainly oriented to the generation of input information for models, others towards mitigation measures that respond to air pollution concerns of their region) ~~which is reflected in the uneven levels of development of local inventories~~. In this sense, it is worth mentioning that although the resources in the region often limit their growth, the capacities of the groups are growing, which is partly reflected in the development of local databases published in this same special issue (Puliafito et al., 2021; Álamos et al., this issue; Osse et al., this issue).

665 In addition to individual advances, we want to emphasize the role of the PAPILA project and the EMISA initiative, which promote collaboration between groups in the region, enhancing efforts aimed at the development of appropriate and consistent surface emission inventories. In this ~~auspicious~~ context, we trust that this work will be a starting point for the development of comprehensive emission inventories ~~in for~~ South America enriched with local information. To this end, the first step will be to join the efforts of other countries in this endeavor, encouraging those with inventory capabilities to broaden their focus beyond 670 cities by building national emission maps. ~~In addition to the aspects that can be improved both in local and global inventories, we found it relevant to point out the key role of transparency in the allocation of sources in the categories for the development of coherent databases.~~

675 *Author contributions.* Paula Castesana: Conceptualization, Formal analysis, Visualization, Writing - original draft, review & editing. Melisa Diaz Resquin: Conceptualization, Formal analysis, Visualization, Writing - original draft, review & editing. Nicolas Huneeus: Formal analysis, Writing - original draft, review & editing. Enrique Puliafito: Formal analysis, Writing - review & editing. Sabine Darras: Data processing. Darío Gomez: Formal analysis, Writing - review & editing. Claire Granier: Conceptualization. Mauricio Osse Alvarado: Writing - review & editing. Nestor Rojas: Writing - review & editing. Laura Dawidowski: Conceptualization, Formal analysis, Writing - original draft, review & editing.




Figure A1. Location of the small domains ~~analysed~~analyzed.

Competing interests. The authors declare that they have no known competing financial interests or personal relationships that could have

680 appeared to influence the work reported in this paper.

Acknowledgements. This work was conducted within the framework of the Prediction of Air Pollution in Latin America and the Caribbean (PAPILA) project. The research leading to these results has received funding from the European Union H2020 program PAPILA (GA 777544). This work was carried out in part with the aid of grants [PICT-O 2016-4802](#) [PICT-O 2016-4802](#) (Agencia Nacional de Promoción Científica y Tecnológica, Argentina) and [PICT 2016-3590](#), [PICT 2016-3590](#) (Fondo para la Investigación Científica y Tecnológica) and

685 [ANID PIA/APOYO AFB180002](#) ([Centro Científico y Tecnológico de Valparaíso](#)). We greatly acknowledge ECCAD (<https://eccad3.sedoo.fr/>) for the archiving and distribution of the datasets [CNEA-3iA-GEIA](#) [CNEA-3iA-GEIA](#) (from Argentina), [CR2-MMA](#) [CR2-MMA](#) (from Chile) and [CAMS-GLOB-ANT](#) [CAMS-GLOB-ANT](#) v4.1 used in this work. The authors also wish to thank the Environmental Protection Agency of Buenos Aires (APRA) for sharing the air quality data for this study.

Appendix A: Supporting Material

Table A1. Description of [sectors](#) [categories](#) and codes considered in PAPILA inventory and their equivalences with [those of](#) CAMS [sectors](#).

PAPILA description	PAPILA code	IPCC 1996 code	IPCC 1996 description	CAMS 4.1 v4.1 code
Thermal power plants	ENE	1A1	Energy industries	ENE
Residential, commercial and other combustion	RES	1A4	Residential, commercial and other combustion	RES
Road transportation	TRO	1A3b	Road transportation	TRO
Non-road Non-road transportation	TNR	1A3c, 1A3e	Rail and other transportation	TNR
Fugitive emissions	FEF	1B1a, 1B2a1, 1B2a2, 1B2a3, 1B2a4, 1B2c, 7A	Coal Mining, Exploration, Production, Transport, Refining/Storage, Venting and Flaring	FEF
Industries (fuel combustion + refining + industrial processes + product use)	IND	1A1b, 1A1c, 1A2, 1A5, 1B1b, 1B2a5, 1B2a6, 1B2b5, 2, 3	Petroleum refining, Manufacture of solid fuels and other energy industries, Other fuel combustion activities, Fuel combustion from manufacturing industries and construction, Fugitive emissions from solid fuel transformation, Fugitive emissions from distribution of oil products and gas natural, Industrial processes, Solvent and other product use.	IND + SLV
Agricultural soils	AGS	4D	Agricultural soils: Synthetic fertilisers, Manure in pasture, Crops	AGS
Agricultural livestock	AGL	4B	Manure Management	AGL
International ships Navigation	SHP-INT 1A3d International navigation SHP Ships	SHP-1A3d1 , 1A3d2	Inland navigation Domestic and International Water-borne Navigation	SHP
Waste	SWD	6	Waste	SWD

Table A2. Summary of NO_x and NMVOCs total agricultural emissions by ~~sector and categories~~ category for years ~~2014–2016~~ 2014–2016.

		NO _x (Gg y ⁻¹)			NMVOCs (Gg y ⁻¹)		
		2014	2015	2016	2014	2015	2016
Manure management							
AGL	Dairy cattle	3.2E-03	3.2E-03	3.2E-03	0.62	0.62	0.63
	Non-dairy cattle	0.29	0.29	0.30	1.32 <ins>1.27</ins>	1.32 <ins>1.30</ins>	1.32
	Other livestock	0.95	0.96	0.94	11.04	11.08	10.92
	Total from AGL	1.24	1.25	1.24	12.98 <ins>12.93</ins>	13.03 <ins>13.00</ins>	12.86
Manure in pasture							
AGS	Dairy cattle	7.80	7.73	7.38	1.25	1.25	1.25
	Non-dairy cattle	58.01	58.50	58.87	13.01 <ins>12.82</ins>	13.01 <ins>12.93</ins>	13.01
	Other livestock	12.88	13.55	13.64	0.10	0.11	0.11
	Total from Manure in pasture	78.70	79.78	79.90	14.36 <ins>14.17</ins>	14.37 <ins>14.29</ins>	14.37
Fertilizers							
Crops		31.54	24.12	36.54	NO	NO	NO
Total from AGS		NO	NO	NO	29.88	34.23	33.73
		110.24	103.90	116.44	44.24 <ins>44.05</ins>	48.60 <ins>48.52</ins>	48.10

Table A3. Description of the selected domains.

Urban domain	Population (10 ⁶ inhab.)	Mean (daily max - daily min) [^o ^o C]	Temp.	Characteristics	Coverage (km ²)	Central coordinates
Antofagasta	0.404	16.4° C (20.1 - 13.7)		PM and SO ₂ Emissions from the copper extraction, smelting operations, heavy machinery operation, surface detonations and the transport of minerals in different degrees of fragmentation. ~11k ^{~11 thousand} industries located in the area.	680 (20x34)	23.62° S; 70.3° W, 2260 m.a.s.l.
Metropolitan Region of San- tiago de Chile (MRS Santiago)	6.257	14.4° C (22.5 - 8.3)		It is a relatively large city located between two mountain ranges, the Andes on the east and the Coastal Range on the west. Pollution levels are very high in winter because of low wind speeds and strong temperature inversions (Gramsch et al., 2014).	990 (30x33)	33.74° S, 70.49° W, 567 m.a.s.l.
Osorno	0.161	10.5° C (16.5 - 5.4)		Economy based on agriculture and livestock. ~5k ^{~5 thousand} industries settled in the area. Cold region, residential/urban area.	378 (18x21)	40.53° S; 73.18° W, 32 m.a.s.l.
Metropolitan Area of Buenos Aires (MABA Buenos Aires)	13.3	17.9° C (9.5 - 23.4)		It is a Metropolitan area that concentrates 32% of the population of the entire country. It has several industrial poles and thermoelectric plants.	1258 (37x34)	34.57° S; 58.44° W, 14 m.a.s.l.
Bahía Blanca	0.291	15° C (8 - 21.8 - 21)		City located near a Petrochemical Pole and a Port Area.	572 (26x22)	38.70° S; 62.22° W, 99 m.a.s.l.
Mendoza	1.1	17.7° C (11.4 - 25 - 25)		It is one of the main cities of the country. It has a main industrial pole linked to the wine industry.	924 (28x33)	32.9° S; 68.7° W, 680 m.a.s.l.




Figure A2. (left) Location of [WRF-Chem](#) Domains. Blue dots are the [monitoring](#) sites location. (right) Hourly, weekly and monthly variations used for the temporal disaggregation of emissions within this work.

690 **References**

Álamos, N., Huneeus, N., Opazo, M., Osses, M., Puja, S., Pantoja, N., Denier van der Gon, H., Schueftan, A., Reyes, R., and Calvo, R.: High resolution inventory of atmospheric emissions from transport, industrial, energy, mining and residential sectors of Chile., Submitted to Earth Syst. Sci. Data ESSD2021-208, Special Issue Surface emissions for atmospheric chemistry and air quality modelling, this issue.

Barros, V. and Camilloni, I.: La Argentina y el cambio climático. De la física a la política, EUDEBA, 2016.

695 Belincanta, J., Alchorne, J. A., and Teixeira Da Silva, M.: The Brazilian experience with ethanol fuel: Aspects of production, use, quality and distribution logistics, *Braz. J. Chem. Eng.*, 33, 1091–1102, <https://doi.org/10.1590/0104-6632.20160334s20150088>, 2016.

CAMESA: <https://portalweb.cammesa.com/default.aspx>, accesed 2021-06-27, 2021.

Castesana, P. S., Dawidowski, L. E., Finster, L., Gómez, D. R., and Taboada, M. A.: Ammonia emissions from the agriculture sector in Argentina; 2000–2012, *Atmos. Env.*, 178, 293–304, <https://doi.org/10.1016/j.atmosenv.2018.02.003>, 2018.

700 Castesana, P. S., Vázquez-Amábile, G., Dawidowski, L. H., and Gómez, D. R.: Temporal and spatial variability of nitrous oxide emissions from agriculture in Argentina, *Carbon Manag.*, 11, 251–263, <https://doi.org/10.1080/17583004.2020.1750229>, 2020.

Castesana, P. S., Diaz Resquin, M. C., Huneeus, N., Puliafito, E., Darras, S., Gómez, D., Granier, C., Osses Alvarado, M., Rojas, N., and Dawidowski, L.: Data for: PAPILA dataset: a regional emission inventory of reactive gases for South America based on the combination of local and global information, v3, Mendeley Data [data set], <https://doi.org/10.17632/btf2mz4fhf.3>, 2021.

705 CIESIN and CIAT: Gridded Population of the World, Version 3 (GPWv3): National Identifier Grid, Center for International Earth Science Information Network - CIESIN - Columbia University, and Centro Internacional de Agricultura Tropical - CIAT. NASA Socioeconomic Data and Applications Center (SEDAC), [data set], <https://doi.org/10.7927/H49W0CDN>, 2005.

CR2-MMA: CR2 Emission inventory for Chile. Emissions of atmospheric Compounds and Compilation of Ancillary Data (ECCAD), [data set], <https://permalink.aeris-data.fr/CR2-MMA>, 2018.

710 Dawidowski, L., Sánchez-Ccoyollo, O., and Alarcón, N.: Estimación de emisiones vehiculares en Lima Metropolitana, Informe final. Lima: SENAMHI/SAEMC, 2014.

Dos Santos Lucon, O. and Moutinho Dos Santos, E.: The HORUS model - Inventory of atmospheric pollutant emissions from industrial combustion in Sao Paulo, Brazil, *Environ. Impact Assess. Rev.*, 25, 197–214, <https://doi.org/10.1016/j.eiar.2004.06.010>, 2005.

EDGAR: EDGAR - Emissions Database for Global Atmospheric Research, https://edgar.jrc.ec.europa.eu/emissions_data_and_maps, 2021.

715 El Pais: https://elpais.com/internacional/2015/06/14/actualidad/1434286413_160142.html, accesed 2021-06-27, 2015.

EMEP: EMEP/EEA air pollutant emission inventory guidebook 2016., European Environment Agency, Copenhagen, Denmark., 2017.

EMISA: Emission Inventories in South America, <https://igacproject.org/activities/emisa>, accessed: 2021-06-07, 2020.

ENARGAS: Ente Nacional Regulador del Gas. Datos Abiertos: Datos operativos de Transporte y Distribución de Gas, <https://www.enargas.gob.ar/secciones/datos-abiertos/datos-abiertos.php>, 2021.

720 Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S., and Moran, M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 16, 11 497–11 519, 2016.

Freitas, S., Longo, K.M. and Alonso, M., Pirre, M., Marecal, V., Grell, G., Stockler, R., Mello, R., and Gacita, M.: PREP-CHEM-SRC-1.0: a preprocessor of trace gas and aerosol emission fields for regional and global atmospheric chemistry models, *Geosci. Model Dev.*, 4, 419 – 433, 2011.

725 Gallardo, L., Barraza, F., Ceballos, A., Galleguillos, M., Huneeus, N., Lambert, F., Ibarra, C., Munizaga, M., O’Ryan, R., Osses, M., Tolvett, S., Urquiza, A., and Véliz, K. D.: Evolution of air quality in Santiago: The role of mobility and lessons from the science-policy interface, *Elementa: Science of the Anthropocene*, 6, <https://doi.org/10.1525/elementa.293>, 2018.

González-Rojas, C. H., Leiva-Guzmán, M., Manzano, C. A., Morales S., R. G., and Araya, R. T.: Short-term air pollution events in the Atacama desert, Chile, *J. S. Am. Earth Sci.*, 105, 103 010, <https://doi.org/https://doi.org/10.1016/j.jsames.2020.103010>, 2021.

730 Gramsch, E., Cáceres, D., Oyola, P., Reyes, F., Vásquez, Y., Rubio, M., and Sánchez, G.: Influence of surface and subsidence thermal inversion on PM2.5 and black carbon concentration, *Atmos. Env.*, 98, 290–298, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2014.08.066>, 2014.

735 Granier, C., Darras, S., Denier Van Der Gon, H., Jana, D., Elguindi, N., Bo, G., Michael, G., Marc, G., Jalkanen, J.-P., and Kuenen, J.: The Copernicus Atmosphere Monitoring Service global and regional emissions (April 2019 version), Copernicus Atmosphere Monitoring Service, pp. 1–55, <https://doi.org/10.24380/d0bn-kx16>, 2019.

Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Env.*, 39, 6957–6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.

740 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., et al.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, 2018.

Huneeus, N., Granier, C., Dawidowski, L., van der Gon, H. D., Alonso, M., Castesana, P., Diaz, M., Frost, G. J., Gallardo, L., Gomez, D., et al.: Anthropogenic emissions in South America for air quality and climate modelling, in: 2017 International Emission Inventory Conference “Applying Science and Streamlining Processes to Improve Inventories”, 2017.

745 Huneeus, N., Denier van der Gon, H., Castesana, P., Menares, C., Granier, C., Granier, L., Alonso, M., de Fatima Andrade, M., Dawidowski, L., Gallardo, L., Gomez, D., Klimont, Z., Janssens-Maenhout, G., Osses, M., Puliafito, S. E., Rojas, N., Sánchez-Ccoylo, O., Tolvett, S., and Ynoue, R. Y.: Evaluation of anthropogenic air pollutant emission inventories for South America at national and city scale, *Atmos. Env.*, 235, 117 606, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2020.117606>, 2020a.

750 Huneeus, N., A., U., Gayó, E., Osses, M., Arriagada, R., Valdés, M., Álamos, N., Amigo, C., Arrieta, D., Basoa, K., Billi, M., Blanco, G., Boisier, J., Calvo, R., Casielles, I., Castro, M., Chahuán, J., Christie, D., Cordero, L., Correa, V., Cortés, J., Fleming, Z., Gajardo, N., Gallardo, L., Gómez, L., Insunza, X., Iriarte, P., Labran˜a, J., Lambert, F., Mun˜oz, A., Opazo, M., O’Ryan, R., Osses, A., Plass, M., Rivas, M., Salinas, S., Santander, S., Seguel, R., Smith, P., and Tolvett, S.: Informe a las Naciones. El aire que respiramos: pasado, presente, futuro. Contaminación atmosférica por MP2,5 en el centro y sur de Chile, available in www.cr2.cl/contaminacion/, 2020b.

IDEAM: Prediction of Air Pollution in Latin America and the Caribbean. Institute of Hydrology, Meteorology and Environmental Studies (IDEAM), <http://www.cambioclimatico.gov.co/3ra-comunicacion-cambio-climatico>, accessed: 2021-06-07, 2017.

IEA: Consistent, accurate and timely energy data and statistics, International Energy Agency, <https://www.iea.org/data-and-statistics>, accessed: 2021-06-22, 2020.

760 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., et al.: HTAP_v2. 2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmospheric Chemistry and Physics*, 15, 11 411–11 432, 2015.

765 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P., Pagliari, V., Olivier, J. G. J.,
Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu, A. M. R., Solazzo, E., and Oreggioni, G. D.: EDGAR
v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012, *Earth Syst. Sci. Data*, 11, 959–1002,
<https://doi.org/10.5194/essd-11-959-2019>, 2019.

770 Jorquer, H.: Air quality at Santiago, Chile: a box modeling approach—I. Carbon monoxide, nitrogen oxides and sulfur dioxide, *Atmos. Env.*, 36, 315–330, [https://doi.org/10.1016/S1352-2310\(01\)00417-4](https://doi.org/10.1016/S1352-2310(01)00417-4), 2002.

Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M.,
775 and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative
power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.

Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions
of particulate matter including black carbon, *Atmos. Chem. Phys.*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.

Kuenen, J., Visschedijk, A., Jozwicka, M., and Denier Van Der Gon, H.: TNO-MACC_II emission inventory; a multi-year (2003–2009)
780 consistent high-resolution European emission inventory for air quality modelling, *Atmos. Chem. Phys.*, 14, 10 963–10 976, 2014.

Mazzeo, A., Huneeus, N., Ordoñez, C., Orfanoz-Chequela, A., Menut, L., Mailler, S., Valari, M., van der Gon, H. D., Gallardo, L., Muñoz,
R., et al.: Impact of residential combustion and transport emissions on air pollution in Santiago during winter, *Atmos. Env.*, 190, 195–208,
2018.

785 NOAA: NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July 1999, Research Data Archive at the National
Center for Atmospheric Research, Computational and Information Systems Laboratory. National Centers for Environmental Prediction,
National Weather Service, U.S. Department of Commerce, <https://doi.org/10.5065/D6M043C6>, 2000.

Ortegon-Sanchez, A. and Oviedo Hernandez, D.: Assessment of the potential for modal shift to non-motorised transport in a developing
context: Case of Lima, Peru, *Res. Transp. Econ.*, 60, 3–13, <https://doi.org/10.1016/j.retrec.2016.05.010>, transportation and
Land Development: A Global View, 2016.

790 795 Osses, M., Rojas, N., Ibarra, C., Valdevenito, V., Laengle, I., Pantoja, N., Osses, D., Basoa, K., Tolvett, S., Hunneus, N., and Gallardo, L.:
High-definition spatial distribution maps of on-road transport exhaust emissions in Chile, 1990–2020, Submitted to *Earth Syst. Sci. Data*
ESSD2021-208, Special Issue Surface emissions for atmospheric chemistry and air quality modelling, this issue.

PAPILA: Prediction of Air Pollution in Latin America and the Caribbean, <http://papila-h2020.eu/papila>, accessed: 2021-06-07, 2020.

Policarpo, N. A., Silva, C., Lopes, T. F. A., Araújo, R. d. S., Cavalcante, F. S. Á., Pitombo, C. S., and de Oliveira, M. L. M.: Road ve-
hicle emission inventory of a Brazilian metropolitan area and insights for other emerging economies, *Transp. Res. D*, 58, 172–185,
<https://doi.org/10.1016/j.trd.2017.12.004>, 2018.

Ports: Estadísticas de carga. Puertos, Vías Navegables y Marina Mercante. Ministerio de Transporte, <https://www.argentina.gob.ar/puertos-vias-navegables-y-marina-mercante/estadisticas-de-carga>, 2021.

Puliafito, S. E., Allende, D. G., Castesana, P. S., and Ruggeri, M. F.: High-resolution atmospheric emission in-
ventory of the argentine energy sector. Comparison with edgar global emission database, *Helion*, 3, e00489,
<https://doi.org/10.1016/j.heliyon.2017.e00489>, 2017.

Puliafito, S. E., Bolaño Ortiz, T. R., Fernandez, R. P., Berná, L. L., Pascual-Flores, R. M., Urquiza, J., López-Noreña, A. I., and Tames,
M. F.: High resolution seasonal and decadal inventory of anthropic gas-phase and particle emissions for Argentina, *Earth Syst. Sci. Data*
Discuss., 2021, 1–56, <https://doi.org/10.5194/essd-2021-81>, 2021.

800 Romero, Y., Chicchon, N., Duarte, F., Noel, J., Ratti, C., and Nyhan, M.: Quantifying and spatial disaggregation of air pollution emissions from ground transportation in a developing country context: Case study for the Lima Metropolitan Area in Peru, *Sci. Total Environ.*, 698, <https://doi.org/10.1016/j.scitotenv.2019.134313>, 2020.

Secretaría de Energía: Datos Energía, Secretaría de Energía, Ministerio de Economía de la Nación, [http://datos\[minem\].gob.ar/](http://datos[minem].gob.ar/), 2021.

805 Sharma, S., Sharma, P., and Khare, M.: Photo-chemical transport modelling of tropospheric ozone: A review, *Atmos. Env.*, 159, 34–54, <https://doi.org/10.1016/j.atmosenv.2017.03.047>, 2017.

810 Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestvedt, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivé, D., Quaas, J., Quennehen, B., Raut, J.-C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E., and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15, 10 529–10 566, <https://doi.org/10.5194/acp-15-10529-2015>, 2015.

TCN: Tercera Comunicación Nacional de la República Argentina a la Convención Marco de la Naciones Unidas para el cambio Climático, Tech. rep., Secretaría de Ambiente y Desarrollo Sustentable de la Nación, <https://www.argentina.gob.ar/ambiente/cambio-climatico/tercera-comunicacion>, 2015.

815 The World Bank: Gini Index (World Bank estimate), <https://data.worldbank.org/indicator/SI.POV.GINI>, accessed: 2021-06-22, 2019.

The World Bank: Agricultural land (% of land area), <https://data.worldbank.org/indicator/AG.LND.AGRI.ZS>, accessed: 2021-06-22, 2020.

Thompson, G., Field, P. R., Rasmussen, R. M., and Hall, W. D.: Explicit Forecasts of Winter Precipitation Using an Improved Bulk Microphysics Scheme. Part II: Implementation of a New Snow Parameterization, *Mon. Weather Rev.*, 136, 5095 – 5115, <https://doi.org/10.1175/2008MWR2387.1>, 2008.

820 Trombetti, M., Thunis, P., Bessagnet, B., Clappier, A., Couvidat, F., Guevara, M., Kuenen, J., and López-Aparicio, S.: Spatial inter-comparison of Top-down emission inventories in European urban areas, *Atmos. Env.*, 173, 142–156, <https://doi.org/10.1016/j.atmosenv.2017.10.032>, 2018.

UNEP: Waste Management Outlook for Latin America and the Caribbean., Programa de las Naciones Unidas para el Medio Ambiente, Oficina para América Latina y el Caribe. Ciudad de Panamá, <https://wedocs.unep.org/20.500.11822/26448>, 2018.

825 United Nations: World Urbanization Prospects The 2014 Revision, United Nations, Department of Economic and Social Affairs, Population Division, (ST/ESA/SER.A/366), 2015.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11 707–11 735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.

830 Villalobos, A. M., Barraza, F., Jorquera, H., and Schauer, J. J.: Wood burning pollution in southern Chile: PM2.5 source apportionment using CMB and molecular markers, *Environ. Pollut.*, 225, 514–523, <https://doi.org/10.1016/j.envpol.2017.02.069>, 2017.

Vivanco, M. G. and Andrade, M. D. F.: Validation of the emission inventory in the São Paulo Metropolitan Area of Brazil, based on ambient concentrations ratios of CO, NMOC and NO_x and on a photochemical model, *Atmos. Env.*, 40, 1189–1198, <https://doi.org/10.1016/j.atmosenv.2005.10.041>, 2006.

835 Wang, K., Zhang, Y., and Yahya, K.: Decadal application of WRF/Chem over the continental U.S.: Simulation design, sensitivity simulations, and climatological model evaluation, *Atmos. Env.*, 253, 118 331, <https://doi.org/10.1016/j.atmosenv.2021.118331>, 2021.

Wang, X., Liang, X. Z., Jiang, W., Tao, Z., Wang, J. X., Liu, H., Han, Z., Liu, S., Zhang, Y., Grell, G. A., and Peckham, S. E.: WRF-Chem simulation of East Asian air quality: Sensitivity to temporal and vertical emissions distributions, *Atmos. Env.*, 44, 660–669, <https://doi.org/10.1016/j.atmosenv.2009.11.011>, 2010.

840 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.