High temporal resolution hydrometeorological data collected in the tropical Cordillera Blanca, Peru (2004-2020)

3 4 5	Emilio I. Mateo ¹ , Bryan G. Mark ¹ , Robert Å. Hellström ² , Michel Baraer ³ , Jeffrey M. McKenzie ⁴ , Thomas Condom ⁵ , Alejo Cochachín Rapre ⁶ , Gilber Gonzales ⁶ , Joe Quijano Gómez ⁶ , Rolando Cesai Crúz Encarnación ⁶				
6 7	¹ Department of Geography, Byrd Polar and Climate Research Center, The Ohio State Universit Columbus, OH, USA				
8	² Department of Geography, Bridgewater State University, Bridgewater, MA, USA				
9 10	³ Département de génie de la construction, École de technologie supérieure, Montreal, QC, Canada				
11	⁴ Department of Earth and Planetary Sciences, McGill University, Montreal, QC, Canada				
12 13	⁵ Université Grenoble Alpes, CNRS, IRD, Grenoble-INP, Institut des Géosciences de l'Environnement (IGE, UMR 5001), Grenoble, France				
14	⁶ Peruvian National Water Authority, Division of Glaciers and Water Resources, Huaraz, Peru				
15	Correspondence to: Emilio Mateo (mateo.9@osu.edu)				
16					
17					
18					
19					
20					
21					
22					
23					
24					
25					

26	Abstract. This article presents a comprehensive hydrometeorological dataset collected over the
27	past two decades throughout the Cordillera Blanca, Peru. The data recording sites, located in the
28	upper portion of the Rio Santa valley, also known as the Callejon de Huaylas, span an elevation
29	range of 3738 - 4750 m a.s.l. As many historical hydrological stations measuring daily discharge
30	across the region became defunct after their installation in the 1950s, there was a need for new
31	stations to be installed and an opportunity to increase the temporal resolution of the streamflow
32	observations. Through inter-institutional collaboration the hydrometeorological network
33	described in this paper was deployed with goals to evaluate how progressive glacier mass loss
34	was impacting stream hydrology, and to better understand the local manifestation of climate
35	change over diurnal to seasonal and interannual time scales. The four automatic weather stations
36	supply detailed meteorological observations, and are situated in a variety of mountain
37	landscapes, with one on a high-mountain pass, another next to a glacial lake, and two in glacially
38	carved valleys. Four additional temperature and relative humidity loggers complement the
39	weather stations within the Llanganuco valley by providing these data across an elevation
40	gradient. The six streamflow gauges are located in tributaries to the Rio Santa and collect high
41	temporal resolution runoff data. The datasets presented here are available freely from
42	https://doi.org/10.4211/hs.059794371790407abd749576df8fd121 (Mateo et al., 2021).
43	Combined, the hydrological and meteorological data collected throughout the Cordillera Blanca
44	enable detailed research of atmospheric and hydrological processes in tropical high-mountain
45	terrain.
46	
40	
47	
48	
40	
49	
50	
51	
50	
52	
53	
54	
55	
55	
56	

57 1 Introduction

58 Glaciers and water resources in the Cordillera Blanca, Peru, have been under close 59 observation for nearly a century. In the 1930s, an Austrian geographer from Universitat 60 Innsbruck, Hans Kinzl, laid the groundwork for glaciological research in the region by surveying 61 and mapping the glaciers and identifying other natural features in the mountainous landscape 62 (Kaser and Osmaston, 2002). While the first systematic Peruvian effort to observe glacier tongue 63 variations in the Cordillera Blanca was initiated in 1944 by Broggi (Petersen et al., 1969), 64 glaciers as a source of security and water resources became the object of study in the subsequent decades. The Unidad de Glaciología e Hidrología (later Unidad de Glaciología y Recursos 65 66 Hidricos (UGRH) of Electroperú S.A.) was initiated by the Corporación Peruana del Santa, a 67 government company for energy development, and took on glacier and lake monitoring after many glacial lake outburst flooding (GLOF) events damaged communities downstream (Ames, 68 69 1998). GLOFs in 1941, 1945, and 1950 killed over 6000 people and destroyed a third of the Ancash district capital of Huaraz (Carey, 2010; Carey et al., 2012). Other glacier hazards have 70 71 had detrimental impacts on the communities as well, including the 1970 avalanche, triggered by 72 a massive earthquake, which killed approximately 6000 people and along with the debris flow it 73 produced, covered the city of Yungay (Carey, 2010). Lliboutry, Morales and Schneider (1977) 74 investigated two glaciers in the mountain range in relation to the danger they presented for 75 flooding a downstream power plant. From the 1970s to the 1990s, a local Peruvian from Huaraz, 76 Alcides Ames, provided many important contributions to the present knowledge of glaciers in 77 the Cordillera Blanca while employed at UGRH, and after his retirement continued to dedicate 78 much of his time to studying the glaciers and sharing his knowledge with other researchers 79 (Ames et al., 1989; Hastenrath and Ames, 1995; Ames and Hastenrath, 1996; Ames, 1998; Kaser 80 and Osmaston, 2002). In the 1980s, Georg Kaser, an Austrian geographer, began studying the 81 Cordillera Blanca with a focus on the extent, causes, and possible consequences of the observed glacier retreat (Kaser and Osmaston, 2002). For many decades, much of the research in the 82 83 region focused on developing better maps of the mountains and observing the marked glacier retreat in the tropics. Although the meteorological and hydrological stations installed by UGRH 84 had been in place for nearly 40 years, it was not until the mid-1990s when studies began to 85 86 assess water resources in the Cordillera Blanca, but with minimal focus on flooding and GLOFs 87 (Kaser and Georges 1997; Mark and Seltzer, 2003).

The UGRH spent many years collecting an abundance of climatological, hydrological, 88 and glaciological data, which was useful for future researchers, hydroelectric and mining 89 90 companies, and other stakeholders throughout the Cordillera Blanca and its glaciers. Since the 91 mid-1990s, most of the daily-resolution discharge data from the Rio Santa watershed has been 92 cataloged and collected at a much reduced number of sites by private energy companies who 93 took over after Electroperu was privatized. The rapid turnover of the energy companies holding 94 the discharge observations makes it difficult to track down all available historical data. Also, 95 many of the hydrological stations have become defunct since they were installed in the 1950s,

96 creating a need for new stations to be installed for continuous monitoring of the region. A further

97 benefit of installing a new hydrological monitoring network in the 2000s was to significantly98 increase the temporal resolution of the discharge observations.

99 Collaborating researchers from The Ohio State University, Bridgewater State University 100 and McGill University initiated a network of embedded environmental sensors in 2005. The 101 main goals of the instrument arrays were to evaluate how progressive glacier mass loss was 102 impacting stream hydrology, and to better understand the local manifestation of climate change 103 on the variability and controls of local weather phenomena over diurnal to seasonal and 104 interannual time scales. In order to create a sustainable network for continuous, long term 105 observation, this project has been maintained in close collaboration with Peruvian researchers 106 and government agencies, as well as with other international scientists to leverage resources in 107 maintaining instruments, in exchange for openly sharing data.

108 The instrumentation for the data collection we present in this work was installed and maintained under a collaborative work agreement ("convenio") formalized with the Peruvian 109 110 government agency overseeing the office in Huaraz (Peruvian Institute of Natural Resources 111 (INRENA) and the Autoridad Nacional del Agua (ANA)). This work agreement involved a 112 secondary collaboration with other international researchers who shared in installing and 113 calibrating the instrumentation. Specifically, Dr. Thomas Condom of the French Institut de 114 Recherche pour le Developpement (IRD) joined the agreement to install and maintain a series of 115 stream gauges logging water stage at 15 minute intervals. We also collaborated with the Austrian 116 research team of Dr. Irme Juen and Professor Georg Kaser from the Universitat Innsbruck, 117 Austria, who co-located a precipitation gauge with our weather station in Llanganuco. Further 118 details about the instrumentation are provided below.

119 Inter-institutional collaboration in this fashion has provided an effective partnership to 120 aid in maintaining the instrumentation over time, but also introduces challenges of logistical coordination and data continuity. Visitation as non-resident international scientists to the field 121 122 sites has been feasible only one or two times annually. Thus, the Peruvians in our collaboration 123 have incorporated our instruments within their routine monitoring network. This has permitted 124 regular observations of stations and instruments to download data loggers, perform stream 125 discharge measurement to build a rating curve, and undertake a limited range of repair work. 126 However, limitations in local resources and manpower in Peru have often prevented recordings 127 of stage observation and discharge measurement to constrain rating curves. Likewise, having 128 multiple operators also increases some risk for data recovery errors. For instance, loggers that are 129 improperly relaunched after data downloads can jeopardize subsequent acquisitions. Having 130 more frequent site visits can allow for interventions, but also incurs increased risk of operator 131 error. Furthermore, Peruvian domestic political changes have disrupted the operations by 132 introducing different leadership, with altered operational priorities and resources, which directly 133 or indirectly interrupt the continuity of trained personnel responsible for data recovery and

134 preservation. Overcoming the many challenges of maintaining the instrumentation and

- 135 constructing rating curves has required regular cross-cultural communication, multiple and
- annual visitation to the region. Likewise, the rating curves continue to be refined as additional
- 137 measurements are made.

138 Our collaborative observations have provided important new insights into how the 139 hydroclimate of the region is changing on different scales. Updated discharge and climate observations in specific glacier catchments documented important shifts in seasonal supply of 140 141 glacier storage to the Yanamarey catchment (Bury et al., 2011) as well as suggesting regional 142 thresholds in glacier melt provision (Mark et al., 2010). The discharge constraints also provided 143 important validation for a novel hydrochemical basin characterization method to quantify 144 proportionate glacier melt and groundwater contributions to streamflow in tributaries of varying 145 glacierized coverage (Baraer et al., 2009, 2015). Regionally, the gauge network provided the key 146 constraint for a model of time-progressive hydrograph evaluation that verified significantly that 147 the main catchment had already passed "peak water" in the wake of strong glacier recession 148 underway for multiple decades (Baraer et al., 2012). Our embedded temperature and humidity 149 loggers distributed over elevation and linked to weather stations in the Llanganuco valley have 150 revealed novel diurnal to seasonal variations in lapse rates linked to catchment-specific valley 151 wind dynamics validated with downscaled climate models (Hellstrom et al., 2017). These studies 152 demonstrate the importance of collecting in situ hydrometeorological data and indicate the need

- 153 for continued data collection in the high Andes (Condom et al., 2020).
- In this paper, we document available data from the Cordillera Blanca area collected over
 the past 15 years. It is separated into (i) meteorological data recorded by permanently installed
 automatic temperature and relative humidity loggers (Lascars), or automatic weather stations,
 and (ii) hydrological data consisting of stage and discharge data from multiple sub-catchments.
- The data are stored in .csv extension format on Consortium of Universities for the Advancementof Hydrologic Science, Inc. (CUAHSI) Hydroshare:
- 160 https://doi.org/10.4211/hs.059794371790407abd749576df8fd121 (Mateo et al., 2021). The data
- 161 collection consists of multiple time series of point observations from both meteorological and
- 162 hydrological measurement sites. These high-temporal resolution data provide detailed insight
- 163 into the complex hydro-meteorological system of the tropical Cordillera Blanca.
- 164 2 Study Area Cordillera Blanca
- 165 The Rio Santa (Santa River) captures runoff from the western side of the glacierized 166 Cordillera Blanca and the eastern side of the non-glacierized Cordillera Negra, encompassing a 167 drainage area of 11636 km² at the outlet to the Pacific Ocean. While the headwaters of the Rio 168 Santa are found at Laguna Conococha at 4100 m above sea level (a.s.l.), the highest point in the 169 basin (and in Peru) is the summit of Huascaran at 6768 m a.s.l.. The average slope of the entire 170 drainage basin is 20.6° and its average elevation is 3374 m a.s.l. calculated from a 3 m resolution

- 171 DEM of the region. The Rio Santa flows over 300 km northwest from its origin at Laguna
- 172 Conococha, an alpine lake at 4000 m a.s.l., to its outlet into the Pacific Ocean, near Chimbote.
- 173 The Callejon de Huaylas refers to the upper section of the Rio Santa, comprising approximately
- 174 4773 km² located above the Cañon del Pato 50 MW hydroelectric generation plant in Huallanca
- 175 (Figure 1). The Rio Santa basin is home to three other hydroelectric plants and provides water to
- the expansive Chavimochic irrigation district near the coast (Mark and McKenzie, 2007). The
- 177 discharge on the Rio Santa has been carefully monitored since the Cañon del Pato dam began
- 178 operating in the 1950s, however, only one station near the dam remains active which is situated
- 179 slightly upstream at La Balsa (labeled as LBQ in Figure 1).
- 180 Figure 1.
- 181 Map of the Callejon de Huaylas showing the locations of the hydrological and meteorological
- 182 stations. (Base layers of the map originated from: © Esri, © NASA, © NGA, © USGS, © FAO,
- 183 © NOAA; all other layers were created and edited by authors of this article.)







Rio Santa discharge experiences a strong seasonal contrast with the lowest discharge occurring between July and September, while peak discharge, nearly 20 times greater, typically occurs in March. Calculated from daily streamflow observations between 1954 and 2015, mean annual streamflow at La Balsa station on the Rio Santa, is 87 m³s⁻¹, while the average annual

- 190 minimum discharge is $25 \text{ m}^3\text{s}^{-1}$, and the average annual maximum discharge is $445 \text{ m}^3\text{s}^{-1}$. The
- 191 Callejon de Huaylas is approximately 10% covered in ice (RGI Consortium, 2017), has an
- 192 average slope of 19° and an average elevation of 4055 m a.s.l.. Glacial melt in the Rio Santa at
- La Balsa provides 10-20% of the total annual discharge and may exceed 40% in the dry season
- 194 (Mark et al., 2005; Condom et al., 2012).
- The climate of the Callejon de Huaylas is semi-arid and displays distinct precipitation
 seasonality, with the wet season between October and May being responsible for 80% of the
 800-1200 mm/year of precipitation (Baraer et al., 2009), and the dry season lasting from June to
 September (Figure 2). As a localized example, the Llanganuco valley displays an average total of
- 199 8 mm during the dry season, and an average of total of 258 mm during the wet season months of
- 200 December, January, and February, based on monthly totals from 1953 to 2010 (Hellstrom et al.,
- 201 2017). Variations in river discharge are largely driven by seasonality of precipitation.
- 202 Temperature remains nearly constant in the outer tropics, with the annual variation in
- 203 temperature being smaller than diurnal variation (Kaser et al., 1990).
- 204 Figure 2.
- A climograph <u>from LlanWX ofin</u> the Cordillera Blanca displaying the strong contrast in
- precipitation between wet and dry seasons, while maintaining a steady temperature throughoutthe entire year.



The geology in the Rio Santa basin is dictated by numerous tectonic and erosional processes because the basin is situated along the active detachment fault of the Cordillera Blanca (McNulty et al., 1998; Garver et al., 2005; Eddy et al., 2017). The highest peaks of the Cordillera Blanca are composed largely of a granodiorite batholith intruded into a metamorphic unit that includes hornfels, gneiss, and sulfide-rich lithologies, such as pyrite schist, phyllite, and pyritebearing quartzite (Giovanni et al., 2010; Eddy et al., 2017). Granodiorite is found predominantly in the northern portion of the range, while the southern portion of the range is made up of a

- 216 variety of metasediments, including quartzites and carbonates (Garver et al., 2005). The main
- 217 valley of the Rio Santa watershed is covered by recent sediment deposits, including alluvium,
- 218 landslide deposits, and glacial-fluvial fill. The impact of past and present glacier extent in the
- 219 topography is visible in geomorphic features throughout the range, including steep walled U-
- shaped valleys, moraines, and proglacial lakes (Eddy et al., 2017).

221 3 The Data

In the following section, the data collected from the Cordillera Blanca are presented in two collections. First, we provide a description of the meteorological stations and embedded sensor network of Lascar data loggers, and then we present the time series data from the stations. Second, we provide details of the setup and context of the discharge gauging station network, and conclude by presenting the time series of discharge data and general statistics from them.

227 3.1 Meteorological Data

228 The Servicio Nacional de Meteorología e Hidrología del Perú (SENAMHI) has operated 229 a relatively dense national network of meteorological stations since 1964. Air temperature is 230 provided as daily T_{max} and T_{min} (additionally measured at 07:00, 13:00 and 19:00 local time), and 231 precipitation is measured once a day. Some SENAMHI stations still provide daily temperature 232 and precipitation data through their web portal, providing data back to 1980 233 (https://www.senamhi.gob.pe/?&p=descarga-datos-hidrometeorologicos), but very few stations 234 within the Rio Santa Valley or Cordillera Blanca, Peru, remain active. Note that additional data 235 from SENAMHI are available upon request from the Swiss MeteoDat GmbH team through a 236 data portal (Schwarb et al. 2011, http://www.meteodat.ch/). In addition, the Universidad 237 Nacional Santiago Antúnez de Mayolo (UNASAM) has maintained a network of what was 238 originally 16 meteorological stations located at different elevations in the Cordillera Blanca 239 (Ancash district) since 2012. Monthly totals of precipitation data for the Rio Santa watershed 240 have been collected by Electroperú South America since 1953 although measurements were 241 interrupted in the mid-1990's4 with the privatization of the respective institutions when most of 242 the stations were abandoned and quality control was an issue (Kaser et al., 2003). More recent 243 installations of weather stations in the Cordillera Blanca have been mostly associated with short-244 lived research projects typically lasting less than three years (Hofer et al., 2010; Georges and 245 Kaser, 2002). The hourly meteorological observations we describe below were collected 246 primarily from instrumentation we installed in two sub-catchments of the Rio Santa drainage 247 basin, Llanganuco, and Quilcayhuanca.

The Llanganuco valley is situated on the western side of the Cordillera Blanca across a southwest (~240°) to northeast (~60°) axis with elevations ranging from 3400 m a.s.l. to 6746 m a.s.l. Llanganuco is one of most glacierized valleys in the mountain range at 30% glacier coverage, making it one of the most glacierized tropical valleys in the world. In the Llanganuco catchment, our group has installed multiple weather stations dating back to 2007, although only

- 253 one remains active. This currently active station, labeled LlanWX, is located at 3835 m.a.s.l. near
- the lower of the two largest valley lakes (Table 1).

255 Table 1.

258

- 256 This table provides general information about each meteorological station (WX) and lascar data
- 257 logger (LAS) location within the embedded sensor network.

Station	Elevation (m a.s.l.)	Period of Operation	Lascar Error Adjustment	Slope Angle (°)	Slope Aspect (°)	Temporal Resolution
Cuchillacocha - CuchWX	4642	2013-Present		0	NA	30-min
Casa de Agua - CDAWX	3924	2013-Present		0	NA	30-min
Llanganuco - LlanWX	3835	2007-Present	Yes	0	NA	60-min
Portachuelo - PortWX	4750	2006-2015		0	NA	60-min
LlanUp-1/1A LAS	3955	2006-2015; 2015-Present	No	14	301	60-min
LlanUp-2/2A LAS	4122	2006-2014; 2015-Present	No	33	259	60-min
LlanUp-3/3A LAS	4355	2006-2015; 2015-Present	No	31	236	60-min
LlanUp-4/4A LAS	4561	2006-2015; 2018-Present	Yes	33	225	60-min
Portachuelo LAS	4767	2007-2015	Yes	9	180	60-min

259 The embedded sensor network (ESN), as described by Hellström et al. (2010) and 260 Hellström and Mark (2006), provides the in-situ meteorological data provided in the paper. In 261 July 20074 the first automatic weather station (AWS), LlanWX, was installed near the lower lake in the Llanganuco valley to collect a continuous record of air temperature, wind speed, wind 262 direction, relative humidity, solar irradiance, and soil temperature/moisture (Table 2). The AWS 263 264 shown in Figure 3A is located in an open area on the valley floor and is surrounded by *Polylepis* 265 trees. A lascar data logger was also hung at this site for static calibration. The site is protected in Huascaran National Park, and the location was previously used by the University of Innsbruck 266 for precipitation measurements. The most significant wind obstructions are the steep bedrock 267 268 walls of the valley toward the northwest and southeast which exceed 1000 meters above the 269 valley floor. Northerly and southerly winds are occasionally recorded by LlanWX, and are likely caused by turbulence or lateral winds from the uneven heating of the valley walls. Winds flow 270 271 parallel to the axis of the valley during approximately 92% of the recorded time and are not greatly obstructed by surface vegetation. The sensors for LlanWX were originally sourced from 272 273 the Onset Computer Corporation and logged with an Onset HOBO® data logger 274 (http://www.onsetcomp.com) until 2014 when these loggers were replaced with an Iridium® 275 satellite Data Garrison logger (http://www.upwardinnovations.com). A 102-mm diameter 276 radiation shield was used to reduce air temperature error caused by the sun except during the 277 following years: 2007, 2011-, 2012, 2013, and 2014.; During these years when a separate data logger was used as the primary source for air temperature and relative humidity observations. In 278 279 2013, our team upgraded the station by replacing the wind sensors with two new units from Onset and a new pyranometer from Apogee (http://www.apogeeinstruments.com/pyranometer) 280 which exceeded the 1277 Wm⁻² maximum reading of the previous Onset sensors (Covert, 2016). 281 The observations from LlanWX are largely continuous since 20075, with data gaps occurring 282 283 occasionally between 2010 and 2014.

284 Table 2.

285 This table details the variables, sensors, and their accuracy which are collected at the

286 meteorological stations throughout the region.

Variable	Sensor	Accuracy	Unit
Air temperature	Onset HOBO S- THB- M002 Temperature RH Smart Sensor	± 0.2 °C	°C
Precipitation	Onset HOBO S- RGB-M002 0.2 mm Rainfall tipping bucket Smart Sensor	± 0.2 mm	mm
Relative humidity	Onset HOBO Temperature RH Smart Sensor: S- THB-M002	± 2.5 %	%
Wind speed	Onset HOBO S- WSB-M003 Wind Speed Smart Sensor	± 1.1 m/s	m/s
Wind direction	Onset HOBO S- WDA-M003 Wind Direction Smart Sensor	±5°	o
Incoming solar radiation	Onset HOBO S- LIN-M003 Solar Radiation Smart Sensor	± 10 W/m²	W/m²
Atmospheric pressure	Onset HOBO S- BPB-CM50 Smart Barometric Pressure Sensor	± 3.0 mb	mb
Soil temperature	Onset HOBO S- TMB-M002 12-bit Temperature Smart Sensor	± 0.2 °C	°C

287

A second AWS, situated at Portachuelo (referred to as PortWX in Figure 1), was installed in July 2006 on a high pass at the top of the Llanganuco valley (4742 m a.s.l.). The station was situated on a steep, rocky ridge between the Llanganuco valley and the Vaqueria valley. There were only wind obstructions to the north due to a steep rock wall within 10 m of the station location. The AWS at Portachuelo had the same sensors as its counterpart, LlanWX, however was lacking an air temperature and relative humidity sensor with a radiation shield until it was installed in July 2015. Prior to this upgrade, these variables were recorded each hour by a Lascar data logger part of the ESN discussed in the following section. The Portachuelo station was
stolen in 2015 after recording data for nine years; a replacement station installed in 2016 was
also stolen so the site was abandoned.

298 Figure 3.

299 A) The typical setup for the HOBO automatic weather station, shown here is the LlanWX station

300 near the lower lake in Llanganuco. Measurements include: air temperature, humidity, wind speed

301 and direction, incoming solar radiation, rainfall, and soil temperature and soil moisture at -10cm.

Note also the <u>two Lascars</u> with different radiationin shields hanging from crossbar. B) Example

303 of a Lascar data logger location shaded from sunlight and hidden from view in a polylepis tree.

304 Photographs taken by R. Hellström (Covert, 2016).





305

306 Smaller loggers measuring air temperature were installed in June 2005 to begin collecting 307 near surface temperatures to calculate lapse rates (Table 1). These data loggers were a series of 8 308 nickel sized iButton Thermochron® temperature loggers (Figure 3^B). Hellström et al. (2010) 309 demonstrated the effectiveness of using the Thermochron loggers for purposes of observing near 310 surface lapse rates within the Llanganuco valley between the elevations 3470 and 4740 m a.s.l. 311 (Covert, 2016). The ESN of iButtons was replaced in July 2006 with a more robust network of 312 Lascar El-USB2 data loggers (www.lascarelectronics.com) which were setup to measure air 313 temperature and relative humidity at one--hour intervals. Each Lascar logger is attached to a 314 custom-designed and locally crafted radiation shield made of a thin tin cone and two Styrofoam 315 pieces in order to reduce error caused by direct sunlight, as shown in Figure 3B. Figure 4 (from: Covert, 2016) provides photos for a visual landscape context for each Lascar logger. While using 316 317 the weather observations provided, note that Because the LlanWX, LlanUp-4, and PortWX 318 Lascar loggers are nearly entirely exposed to sunlight, the recorded which led to higher than 319 expected air temperature values in the datasets are higher than expected and contain greater error 320 thanfrom actual temperatures during the day (see section 4.1 for details). Each Lascar logger is 321 attached to a custom-designed and locally crafted radiation shield made of a thin tin cone and two Styrofoam pieces in order to reduce error caused by direct sunlight, as shown in Figure 3B. 322 323 This Lascar network is still recording hourly data.

- 324 Figure 4.
- Photos A-F show the locations of all Lascar dataloggers <u>labeled by names matching Table</u>
- <u>1</u>which gathered data provided in this paper. Photographs taken by B. Mark and R. Hellström
 (Covert, 2016).



Quilcayhuanca contains two weather stations (Table 1) similar to the LlanWX station in Llanganuco valley, the lower is located at 3924 m a.s.l. near an old river diversion station called Casa de Agua (<u>station</u> referred to as CDAWX in Figure 1), and the upper is located at 4642 m a.s.l. slightly above an alpine lake, Cuchillacocha (<u>station</u> referred to as CuchWX in Figure 1). The AWS at Casa de Agua is still active since its installation in July 2013 and mostly continuous ³³⁴ over this duration. CDAWX station collects all of the same variables as LlanWX, which are

- described in (Table 2), is still active since its installation in July 2013 and data are mostly
- 336 <u>continuous over this duration</u>. The Cuchillacocha AWS CuchWX was also installed in July 2013,
- and is still presently active, and only has one data gap lasting longer than a month (from
- **338** November 2016 June 2017).

339 <u>3.1.1 Relative Humidity Error Correction for LlanWX</u>

340 <u>Data inspection revealed two systematic errors that were corrected including: supersaturation of</u>

341 <u>the humidity sensor during extended wet periods, and radiation error of temperature during the</u>
 342 late afternoon with direct solar radiation heating the sides of the Lascar data loggers.

343 In 2006 and 2007 saturation of the humidity sensor at LLanWX resulted in relative 344 humidity values greater than 100% (as high as 110%) resulting from dust and condensation on 345 the humidity sensor and persisting during the wet season for periods of up to about 48 hours, 346 with periods of reasonable humidity values during driver periods. In light of data preservation, 347 and simplicity, these erroneously supersaturated values were simply replaced with 100% relative 348 humidity;- this correction is referred to as "Lascar Error Adjustment" in Table 1. Note that the 349 dew point was equal to the air temperature for values of relative humidity 100% or greater, so no 350 adjustment was made for dew point. The humidity sensor was replaced in June 2007 and the correction was no longer needed. You will find bBoth the original and corrected values are 351

retained in the LlanWX data file. There was no similar humidity error observed for PortWX or
 any of the LAS sensors.

354 <u>3.1.2 Radiation Error Correction for Lascar Temperature Loggers</u>

A year-long comparison between the Lascar setup-(Figure 5) and LlanWX AWS 355 temperature sensors indicated an expected error in the Lascar thatwhich was largely dependent 356 357 on the time of day. Variability in temperature bias due to solar radiative heating is likely caused by changes in solar angle and cloud cover patterns, particularly during the dry season. It is well 358 documented that daytime temperature error rises on sunny days with calm wind and is reduced 359 on windy and cloudy days regardless of the season, but is far more prominent during the dry 360 season in the tropical Andes (Georges and Kaser, 2002). It is important to note that snow cover, 361 which can create larger biases in temperature by reflected solar radiation, is minimal throughout 362 the year at the locations of the Lascars in this study. Over an entire month, nightly error was 363 within the 0.5 °C resolution of the Lascar sensor. The occasional occurrence of greater nighttime 364 error could be caused by the sensor capturing longwave radiation emitted by the surface. Unlike 365 366 the AWS radiation shield in which the temperature sensor is fully enclosed, the Lascar is exposed from below meaning it is able to capture longwave radiation at night. Further analysis 367 will need to be done to verify this as the cause of the nocturnal error, though these errors are 368 369 considered minimal for the uses of these datasets. The primary source of consistent 370 instrumentation error for air temperature measurements is the heating of radiation shields from 371 those locations exposed to direct sunlight, which required bias correction according to Figure 5.

372 We bias-corrected for radiation error using comparison with the LlanWX shielded temperature and Lascar logger hanging next to it under the same exposure, including the LlanWX, LlanUp-4 373 374 LAS, and PortWX station locations. We conducted T this same comparison was conducted when 375 replacing Lascar sensors in 2015, and found no notable significant differences between old and 376 new sensors. Because solar radiation is the most significant predominant source of error, 377 corrections were only applied to daylight hours between 07:00 to 19:00 (Covert, 2016). During 378 nighttime hours the error was within the sensor's output resolution and no corrections were 379 applied. 380 381 Figure 5. 382 Image A displays a Lascar data logger connected to its cone radiation shield. Two white 383 Styrofoam plates beneath the cone insulate the sensor from radiative heating. Image B shows the radiation shields used on the AWS in comparison to the Lascar setup in image A. Image C shows 384 385 polylepis trees which are used to hide and shade Lascars in order to obtain accurate temperature 386 and relative humidity values. A comparison test between the Lascar setup and the AWS radiation 387 shield with no shading yielded the expected errors shown in graph D. Positive values indicate 388 that the Lascar reported a temperature higher than the AWS sensor. Photographs taken by R. 389 Hellström (Covert, 2016). 390



391

392

393 394 3

3.1.3 Data Gaps and Other Quality Control Checks

The most common reasons for-larger gaps in data was either loss of sensors due to theft or inability to access sensors because of logistical reasons, such as poor weather conditions or human error during downloading or redeploying of sensors. In addition to corrections for solar radiation heating and the relative humidity correction, humidity and temperature records were assessed for accuracy and outliers removed if there were unrealistic deviations from the previous 400 24-hour trend of conditions within the valley. Most of these deviations occurred during time 401 intervals coincident with sensor deployment, battery replacement or data downloadwere due to 402 an hour where sensors were downloaded and held by hand or in a pocket, although data (less 403 than 1%) were also removed if there was a concern with the integrity of the sensor. Due to theft 404 or replacement of radiation shields, tThe four Lascar loggers were replaced or had new radiation 405 shields installed in June 2015, and likewise renamed in the dataset asrenamed to LlanUp-1A, 2A, 406 3A, and 4A. in June 2015 and this is reflected in the database. The most common reason for 407 larger gaps in data was either loss of sensors due to theft or inability to access sensors because of logistical reasons, such as poor weather conditions or human error during downloading or 408

409 redeploying of sensors.

410 3.2 Hydrological Data

411 The La Balsa gauge (8.87° S, 77.82° W at 1880 m a.s.l.) (Duke Energy, Orazul Energy, 412 Inkia Energy) near the diversion canal for the Cañon del Pato damhydroelectric plant has a nearly complete record since 1954, acting as a long-term reference point for discharge on the Rio 413 414 Santa at the base of the Callejon de Huaylas basin. Many other stream gauges and precipitation 415 stations were put in place by the hydroelectric companies, however, due to the lack of 416 maintenance these stations became unusable after the mid-1990's. In 2008, The Ohio State University, McGill University, IRD, and the Peruvian glaciology unit of ANA commenced a 417 joint project to reinstitute a stream gauging network throughout the Cordillera Blanca (Baraer et 418 al., 2012). Many of these redeployed stream gauges in sub-catchments of the Rio Santa are still 419 420 in working condition and are continuing to be monitored.

421 The stream gauges comprise custom-designed and locally crafted steel stilling wells 422 containing two Solinst leveloggers (Figure 65), with one measuring total pressure (water plus 423 atmospheric), and the other measuring barometric pressure. The Model 3001 Solinst levelogger 424 Edge has an accuracy of ± 0.05 kPa and a resolution of 0.002% Full Scale (FS) for pressure and a temperature accuracy of $\pm 0.05^{\circ}$ C, and a temperature resolution of 0.003° C 425 426 (www.solinst.com). Water level is continuously monitored at these gauges at a temporal 427 resolution of 15-minutes by subtracting atmospheric pressure from the total pressure. The 428 adjusted water level variable in the discharge datasets indicates the adjustment based on where 429 the water level logger was located in the water column. This measurement is conducted in the 430 field using a built-in meter stick at each gauging station. Rating curves at all stations were 431 established and verified by conducting discharge measurements using the velocity area method. 432 Unavailable data is denoted by "NA" in the dataset. Linear interpolation of missing data was 433 only calculated for missing periods of less than 1 hour, coinciding with the time the leveloggers 434 were being downloaded in the field. Other than these short periods of interpolated data, the 435 datasets provided consist of raw data.

436 Figure <u>6</u>5.

Figure 65 illustrates the design of the stilling wells used to collect total pressure (water +

barometric) and barometric pressure. <u>The photograph shows the stilling well and the insert tube</u>containing the pressure data loggers.



440

441 The stream gauges that were installed in 2008 and 2009 and which are continuing to 442 collect discharge are Casa de Agua, Cuchillacocha, Pachacoto, Llanganuco, and Ouerococha 443 (Figure 1). The Pumapampa gauge went out of commission in 2016 but is also included in this 444 dataset (Table 3). These stream gauges have variable lengths of record due to periods of missing 445 data, but still provide valuable streamflow information in a region where stream records are 446 limited. As shown in the map (Figure 1), two pairs of these stream gauges are located at different 447 points along the same Rio Santa tributary in their respective sub-catchments: Pumapampa and 448 Pachacoto in the Pachocoto valley; and Cuchillacocha and Casa de Agua in the Quilcayhuanca 449 valley. All discharge measurements are calculated from rating curves containing a minimum of 450 eight point measurements throughout all times of the year (as mentioned above, rating curves) 451 will be provided upon request). Rating curves are not directly provided in this data paper because 452 they are variable in nature as we collect new discharge measurements in the field. Rating curves 453 are available upon request. The following two subsections provide a brief overview and 454 summary for each gauging station, organized by general location in the Cordillera Blanca, 455 moving from the southern end of the mountain range to the northern. Note that all glacier 456 coverage is calculated from RGI 6.0 (RGI Consortium, 2017).

457 Table 3.

458 This table provides geographical information about each of the hydrological gauging stations.

459 Latitude and longitude are not provided for the security of the leveloggers. Note that all glacier

460 coverage is calculated from RGI 6.0 (RGI Consortium, 2017).

Station	Latitude	Longitude	Elevation (m a.s.l.)	Period of Operation	Contributing Area (km ²)	Percent basin covered in ice (%)
Cuchillacocha - CuchQ	-9.41	-77.35	4631	2008-Present	4.1	60.49
Casa de Agua - CDAQ	-9.46	-77.37	3948	2009-Present	66.9	24.78
Pumapampa - PumQ	-9.88	-77.24	4287	2008-2016	58.1	8.33
Pachacoto - PachQ	-9.85	-77.4	3738	2008-Present	202.3	4.95
Llanganuco - LlanQ	-9.07	-77.65	3850	2008-Present	86.9	30.32
Querococha - QuerQ	-9.72	-77.33	4005	2008-Present	62.9	1.53

462 **3.2.1** The southern Cordillera Blanca

461

The Pumapampa catchment encompasses 58 km² with about 8% being covered by 463 464 glaciers (Table 3). Pumapampa gauge (referred to as PumQ in Figure 1) is located at 4287 m 465 a.s.l., below Nevado Pastoruri at the southern end of the Cordillera Blanca. This stream gauge is 466 situated in a channel through a low-lying meadow, where wet-season flow exceeds bankfull 467 stage, water overflowed its banks during high volume flows causing inaccuracies in high discharge measurements. Most notably, a series of values in late-February 2014 indicated a peak 468 469 discharge of 25 m³s⁻¹, well above extremes measured from the rating curve. It was determined that during this period the leveloggers were overrun by water and measurements became 470 471 unreliable solely during this flooding event. Outside of this event, discharge at Pumapampa is 472 consistently within the measured values on the rating curve. Mean annual streamflow (averaged 473 from three years of data missing less than one-month during the entire year) at this stream gauge was 2.3 m³s⁻¹ and a specific discharge of 1248 mm a⁻¹. Mean streamflow during the dry season 474 (May through September) in Pumapampa was 1 m³s⁻¹, while the mean streamflow during the wet 475 season (October through April) was 2.9 m³s⁻¹. Highest mean monthly streamflow occurred in 476 March 2014 with an average of 5.3 m^3s^{-1} (Figure 76). 477

478 The Pachacoto gauge (referred to as PachQ in Figure 1) measures discharge 19 km 479 downstream from Pumapampa, at 3738 m a.s.l., just above its confluence with the Rio Santa. 480 Streamflow at Pachacoto is significantly greater than Pumapampa as it is the drainage point for a 481 larger area of 202 km², 5% of which is covered by glaciers. The mean annual streamflow at 482 Pachacoto was 3.2 m³s⁻¹, averaged from the seven near-complete years of data, with a calculated specific discharge of 499 mm a⁻¹. Dry season runoff accounted for approximately 20% of the 483 484 annual streamflow by volume, while the majority of the remaining wet season runoff occurred 485 during February and March.

486 Streamflow at another sub-tributary catchment in the southern Cordillera Blanca is 487 recorded at the Querococha gauge (referred to as QuerQ in Figure 1) at 4005 m a.s.l., located 100 488 m downstream of a largeQuerococha lake and defining a drainage area of 63 km², including 489 1.5% of which is glacierized. This drainage collects from two sub-catchmentswatersheds, one of 490 which is entirely glacier-freedeglaciated, while the otherand one which has a contains rapidly receding ice masses including the Yanamarey glaciermass of ice. Using five years of complete 491 492 data, the mean annual streamflow at this gauging station was calculated to be 1.4 m³s⁻¹ and the 493 specific discharge was 702 mm a⁻¹. The streamflow from this gauge is somewhat regulated by Querococha, the second largestlarge lake in the Cordillera Blancaresiding 100 m upstream. This 494

- 495 basin <u>catchment has featured</u> is home to many previous <u>research</u> studies from our research group,
- and many other instruments throughout the basin are still consistently monitored by ANA.
- 497 Figure <u>7</u>6.
- Figure $\frac{76}{6}$ shows the monthly discharge, in m^3s^{-1} , at each of the gauging stations described in
- this study calculated from <u>all</u> complete months of data for each station. The strong seasonal
- 500 pattern is clearly visible at most of the stream gauges.



502 3.2.2 The central and northern Cordillera Blanca

Another catchment that contains two streamflow gauges is Quilcayhuanca, a valley directly above the region's most populous city, Huaraz. The higher elevation stream gauge is Cuchillacocha (referred to as CuchQ in Figure 1) which is situated at 4631 m a.s.l. and measures discharge below a high-alpine lake and two cirque glaciers encompassing a drainage area of 4 km² (with 61% of the basin covered in ice). This station displays a noticeably different discharge pattern throughout the year than the lower gauging stations which collect greater runoff. Cuchillacocha discharge is not defined by a strong wet-dry season fluctuation, instead displaying
 more variability each month and rising to peak values much earlier in the wet season than other

- 511 stream gauges. The average streamflow values at this location are also an order of magnitude
- bla lower than other gauging stations due to the small drainage area it collects from (Figure $\frac{67}{2}$).

513 The stream gauge located at a lower elevation in the Ouilcayhuanca valley is Casa de 514 Agua (referred to as CDAQ in Figure 1) at 3948 m a.s.l. This gauging station is found near a 515 channel cut in the stream, in a large meadow, where water was previously rerouted for agricultural purposes. This gauge collects drainage from an area of 67 km² below the confluence 516 517 of two upper catchments, Cuchillacocha and Cayesh, well above the city of Huaraz (3100 m 518 a.s.l.). From this drainage point in the watershed, the basin is approximately 25% glacierized. 519 The mean annual streamflow, calculated from six complete years of data at Casa de Agua, was 2 520 m^3s^{-1} , with a specific discharge of 943 mm a^{-1} .

521 The last streamflow gauge still recording data, Llanganuco (referred to as LlanQ in 522 Figure 1), is named after the valley it is located in and is positioned at 3850 m a.s.l., below two 523 valley lakes and a catchment area of 87 km^2 , 30% of which is currently glacierized. The mean 524 annual streamflow, calculated from three complete years of data, was 2.3 m³s⁻¹, with a specific 525 discharge of 835 mm a⁻¹. A stream gauge has been in commission at Llanganuco on and off for 526 68 years, beginning as a gauge for UGRH and hydroelectric companies and later, after the 527 original station became defunct, a site for our new network of gauges.

528 **4 Data availability**

- 529 The datasets presented here are available freely from
- 530 https://doi.org/10.4211/hs.059794371790407abd749576df8fd121 (Mateo et al., 2021). The
- 531 hydrological and meteorological data from the Cordillera Blanca have been used in short
- 532 segments in previous studies (Baraer et al., 2012; Baraer et al., 2015). Data availability varies
- 533 from station to station depending on location of loggers and instrumental errors which caused
- periods of time to lapse without data being recorded. These datasets represent a majority of the
- 535 streamflow and meteorological data collected by our research group over the past two decades
- collected at a high temporal resolution of 15-30 minutes. These include point dischargemeasurements and short-time periods of 1-minute temporal resolution meteorological
- measurements and short-time periods of 1-minute temporal resolution meteorological
 measurements. The data provided here will be added to as future field seasons occur and the
- effort to provide hydrometeorological data to the scientific community will continue at all of the
- 540 involved universities and institutions.
- 541 Figure 7.
- 542 Image A displays a Lascar data logger connected to its cone radiation shield. Two white
- 543 Styrofoam plates beneath the cone insulate the sensor from radiative heating. Image B shows the
- 544 radiation shields used on the AWS in comparison to the Lascar setup in image A. Image C shows
- 545 polylepis trees which are used to hide and shade Lascars in order to obtain accurate temperature

- 546 and relative humidity values. A comparison test between the Lascar setup and the AWS radiation
- 547 shield with no shading yielded the expected errors shown in graph D. Positive values indicate
- 548 that the Lascar reported a temperature higher than the AWS sensor. Photographs taken by R.
- 549 Hellström (Covert, 2016).
- 550



552 4.1 Radiation Error for Lascar Temperature Loggers

A year-long comparison between the Lascar setup (Figure 7) and LlanWX AWS 553 indicated an expected error which was largely dependent on the time of day. Variability in 554 555 temperature bias due to solar radiative heating is likely caused by changes in solar angle and 556 cloud cover patterns, particularly during the dry season. It is well documented that temperature 557 error rises on sunny days with calm wind and is reduced on windy and cloudy days regardless of 558 the season, but is far more prominent during the dry season in the tropical Andes (Georges and 559 Kaser, 2002). It is important to note that snow cover, which can create larger biases in 560 temperature by reflected solar radiation, is minimal throughout the year at the locations of the Lascars in this study. Over an entire month, nightly error was within the 0.5 °C resolution of the 561 Lascar sensor. The occasional occurrence of greater nighttime error could be caused by the 562 563 sensor capturing longwave radiation emitted by the surface. Unlike the AWS radiation shield in 564 which the temperature sensor is fully enclosed, the Lascar is exposed from below meaning it is 565 able to capture longwave radiation at night. Further analysis will need to be done to verify this as 566 the cause of the nocturnal error, though these errors are considered minimal for the uses of these 567 datasets. The primary source of consistent instrumentation error for air temperature 568 measurements is the heating of radiation shields from those locations exposed to direct sunlight, 569 which required bias correction according to Figure 7. We bias corrected for radiation error using 570 comparison with the LlanWX shielded temperature and Lascar logger hanging next to it under 571 the same exposure, including the LlanWX, LlanUp-4 LAS and PortWX station locations. 572 Because solar radiation is the most significant source of error, corrections were only applied to

573 daylight hours between 07:00 to 19:00 (Covert, 2016). During nighttime hours the error was 574 within the sensor's output resolution and no corrections were applied. In addition to corrections 575 for solar radiation heating, humidity and temperature records were assessed for accuracy and 576 removed if there were unrealistic deviations from the previous 24-hour trend of conditions within 577 the valley. Most of these deviations were due to an hour where sensors were downloaded and 578 held by hand or in a pocket, although data (less than 1%) were also removed if there was a 579 concern with the integrity of the sensor. Due to theft or replacement of radiation shields, the four 580 Lascar loggers were renamed to LanUp-1A, 2A, 3A, and 4A in June 2015 and this is reflected in 581 the database. The most common reason for larger gaps in data was either loss of sensors due to theft or inability to access sensors because of logistical reasons, such as poor weather windows 582 583 or human error in downloading or redeploying sensors.

584 5 Conclusions

585 The Cordillera Blanca in the tropical Andes of Peru is a unique, high mountain region 586 with a wealth of where high resolution meteorological and hydrological time series observations 587 collected from a network of instruments over the past two decades have been compiled in a new 588 dataset. The region has been the focus of glacier monitoring efforts for nearly a century. While 589 daily-to-monthly time series of meteorological and hydrological observations have been recorded 590 discontinuouslyoff and on for nearly 70 years, there was no generalized sub-daily data monitoring until the early 2000s. Maintaining this network of instruments recording high-591 592 temporal resolution data has involveds traveling to remote areas of the Cordillera Blanca, 593 protecting the instruments from the harsh weather conditions at high elevations, and concealing 594 instruments to prevent theft. There is also difficulty in developing long-term strategies because 595 most funding agencies are focused on short-term based projects and do not fund continuous 596 monitoring and maintenance of data collection networks. In the context of ongoing globalWith 597 climate change with variable localized extremes, as an ongoing global situation, and its harsh impacts at local levels, long-term monitoring of hydrometeorological variables is will become of 598 599 utmost importance to document and understand the changes that are occurring.

600 The datasets collected and described in this paper support investigations into climate 601 evolution, water resource availability, and hydrological changes across the Cordillera Blanca in 602 the past twenty-years. These datasets will also provide valuable, easy-to-access observations for 603 local water resource managers. This paper provides an overview of the variables which have 604 been measured and what is being made available as of 2021. Future measurements recorded in 605 the field will be made available as they are collected, to further build on the available hydro-606 meteorological database in the Cordillera Blanca.

607 Author Contribution

608 EM prepared the manuscript with contributions from co-authors. EM, RH and MB curated the 609 data in preparation for this article. BGM initiated research investigation and acquired <u>the</u> initial

- 610 funding for this international research collaboration and instrumental data collection network. All
- authors <u>collaborated in fieldwork</u> assisted in gathering data and maintaining the
- 612 <u>instrumentsnetwork of and dataloggers collecting stations</u>.

Competing Interests

614 The authors declare that they have no conflict of interest.

615 Acknowledgements

- 616 Funding for instrumentation, installation and field maintenance was provided by various
- 617 agencies and institutions associated with overlapping research projects and educational initiatives
- 618 through respective institutions. BGM acknowledges Ohio State University, National Science
- 619 Foundation (NSF #0752175 with REU supplement; NSF #1010384; NSF #1316432), National
- 620 Geographic Society, NASA (New Investigator Program), and US Fulbright (Award #4506).

- 637
- 638
- 639
- 640
- 641 References
- Ames, A., Munoz, G., Verastegui, J., Vigil, R., Zamora, M., and Zapata, M.: Glacier inventory of
 Peru, Huaraz, Peru, 1989.
- 644 Ames, A. and Hastenrath S.: Mass balance and iceflow of the Uruashraju Glacier, Cordillera
- Blanca. Peru, Zeitschrift für Gletscherkunde und Glazialgeologie, 32 (2), 83-89, 1996.
- Ames, A.: A documentation of glacier tongue variations and lake development in the Cordillera
 Blanca, Peru, Zeitschrift für Gletscherkunde und Glazialgeologie, 34 (1), 1-36, 1998.
- 648 Baraër, M., McKenzie, J. M., Mark, B. G., Bury, J. and Knox, S.: Characterizing contributions of
- 649 glacier melt and groundwater during the dry season in a poorly gauged catchment of the
- 650 Cordillera Blanca (Peru), Advances in Geosciences, 22, 41-49, 2009.
- Baraër, M., Mark, B. G., McKenzie, J. M., Condom, T., Bury, J., Huh, K., Portocarrero, C.,
- 652 Gomez, J., and Rathay, S.: Glacier recession and water resources in Peru's Cordillera Blanca,
- 653 Journal of Glaciology, 58, 134-150, doi: 10.3189/2012JoG11J186, 2012.
- 654 Baraër, M., McKenzie, J., Mark, B. G., Gordon, R., Bury, J., Condom, T., Gomez, J., Knox, S.,
- and Fortner, S. K.: Contribution of groundwater to the outflow from ungauged glacierized
- 656 catchments: a multi-site study in the tropical Cordillera Blanca, Peru, Hydrological Processes, 29
- 657 (11), 2561–2581, https://doi.org/10.1002/hyp.10386, 2015.
- Bury, J. T., Mark, B. G., McKenzie, J. M., French, A., Baraër, M., Huh, K. I., and Gómez López,
- R. J.: Glacier recession and human vulnerability in the Yanamarey watershed of the Cordillera
 Blanca, Peru, Climate Change, 105 (12), 179-206, 2011.
- 661 Carey M.: In the Shadow of Melting Glaciers: Climate Change and Andean Society, Oxford662 University Press, New York, 2010.
- 663 Carey, M., Huggel., C., Bury, J., Portocarrero, C., and Haeberli, W.: An integrated socio-
- environmental framework for glacier hazard management and climate change adaptation: lessons
 from Lake 513, Cordillera Blanca, Peru, Climate Change, 112, 733-767, 2012.

- 666 Condom, T., Escobar, M., Purkey, D., Pouget, J.C., Suarez, W., Ramos, C., Apaestegui, J., Tacsi,
- A., and Gomez, J.: Simulating the implications of glaciers' retreat for water management: a case
 study in the Rio Santa basin, Peru, Water International, 37, 442–459, 2012.
- 669 Condom, T., Martínez, R., Pabón, J.D., Costa, F., Pineda, L., Nieto, J.J., López, F., and Villacis,
- 670 M.: Climatological and Hydrological Observations for the South American Andes: In situ
- 671 Stations, Satellite, and Reanalysis Data Sets, Frontiers in Earth Science, 8, 92, 2020.
- 672 Covert, J. M.: Observational Analysis of Inter-annual Boundary Layer Processes within the
- 673 Glaciated Llanganuco Valley, Peru. In BSU Honors Program Theses and Projects. Item 167.
- 674 Available at: http://vc.bridgew.edu/honors_proj/167 Copyright © 2016 Jason Covert, 2016.
- 675 Eddy, A. M., Mark, B. G., Baraër, M., McKenzie, J. M., Fernández, A., Welch, S., and Fortner,
- 676 S.: Exploring patterns and controls on the hydrochemistry of proglacial streams in the Upper
- 677 Santa River, Peru, Revista de Glaciares y Ecosistemas de Montaña, 3, 41-57, 2017.
- 678 Garver, J. I., Reiners, P. W., Walker, L. J., Ramage, J. M., and Perry, S. E.: Implications for
- 679 timing of Andean uplift from thermal resetting of radiation damaged zircon in the Cordillera
- Huayhuash, northern Peru, Journal of Geology, 113 (2), 117-138, 2015.
- 681 Georges, C. and Kaser, G.: Ventilated and unventilated air temperature measurements for
- glacier-climate studies on a tropical high mountain site, Journal of Geophysical Research, 107
 (D24), 4775, doi:10.1029/2002JD002503, 2002.
- Giovanni, M. K., Horton, B. K., Garzione, C. N., McNulty, B., and Grove, M.: Extensional basin
 evolution in the Cordillera Blanca, Peru: Stratigraphic and isotopic records of detachment
 faulting and orogenic collapse in the Andean hinterland, Tectonics, 29 (6), TC6007,
- 687 doi:10.1029/2010TC002666, 2010.
- Hastenrath, S. and Ames, A.: Recession of Yanamarey Glacier in Cordillera Blanca, Peru, during
 the 20th century, J. Glaciology, 41, 191–196, https://doi.org/10.1029/94JD03108, 1995.
- 690 Hellström, R. Å., and Mark, B. G.: An embedded sensor network for measuring
- 691 hydrometeorological variability within a tropical alpine valley, Proceedings of the 63rd Eastern
- 692 Snow Conference, U. Delaware, Newark, DE, USA, 2006.
- 693 Hellström, R. Å., Higgins, A., Ferris, D., Mark, B. G., and Levia, D. F.: Impacts of complex
- 694 terrain on evapotranspiration within a tropical alpine valley in the Peruvian Andes, Proceedings
- 695 of the 67th Eastern Snow Conference, Jiminy Peak Mountain Resort, Hancock, MA, USA, 2010.
- 696 Hellström, R. Å., Fernandez, A., Mark, B. G., Covert, J. M., Cochachin, A., Gomez, J.:
- 697 Incorporating autonomous sensors and climate modeling to gain insight into seasonal
- 698 hydrometeorological processes within a tropical glacierized valley, Annals of the American

- Association of Geographers, 107 (2), 260-273, https://doi.org/10.1080/24694452.2016.1232615,
 2017.
- 701 Hofer, M., Mölg, T., Marzeion, B., and Kaser, G.: Empirical-statistical downscaling of reanalysis
- 702 data to high-resolution air temperature and specific humidity above a glacier surface (Cordillera
- 703 Blanca, Peru), J. Geophysical Research, 115, D12120, doi:10.1029/2009JD012556, 2010.
- Juen, I., Kaser, G., and Georges, C.: Modelling observed and future runoff from a glacierized
- tropical catchment (Cordillera Blanca, Perú), Global and Planetary Change, 59, 1-4, 37-48, 2007.
- Kaser, G., Ames, A., and Zamora, M.: Glacier fluctuations and climate in the Cordillera Blanca,
 Peru, Annals of Glaciology, 14, 136–140, 1990.
- 708 Kaser, G. and Georges, C.: Changes of the equilibrium-line altitude in the tropical Cordillera
- Blanca, Peru, 1930-50, and their spatial variations, Annals of Glaciology, 24, 344-349, 1997.
- Kaser, G. and Osmaston, H. A.: Tropical glaciers, Cambridge University Press, Cambridge,United Kingdom, 2002.
- 712 Lliboutry, L., Morales, B., Schneider, B.: Glaciological problems set by the control of dangerous
- 713 Lakes in Cordillera Blanca, Peru. III. Study of Moraines and mass balances at Safuna, Journal of
- 714 Glaciology, 18 (79), 275–290, 1977.
- Mark, B. G. and Seltzer, G. O.: Tropical glacier meltwater contribution to stream discharge: A
 case study in the Cordillera Blanca, Peru, Journal of Glaciology, 49 (165), 271-281, 2003.
- Mark, B. G. and McKenzie, J. M.: Tracing increasing tropical Andean glacier melt with stable
 isotopes in water, Environmental Science and Technology, 41 (20), 6955-6960, 2007.
- 719 Mark, B. G., Bury, J., McKenzie, J. M., French, A., and Baraer, M.: Climate change and tropical
- Andean glacier recession: evaluating hydrologic changes and livelihood vulnerability in the
- 721 Cordillera Blanca, Peru, Annals of the Association of American Geographers 100, 794-805,
- 722 2010.
- Mark, B. G., McKenzie, J. M., and Gomez, J.: Hydrochemical evaluation of changing glacier
 meltwater contribution to stream discharge: Callejon de Huaylas, Peru, Hydrological Sciences
 Journal, 50, 975-987, 2005.
- 726 Mateo, E. I., Mark, B. G., Hellström, R. Å., Baraer, M., McKenzie, J. M., Condom, T., Rapre, A.
- 727 C., Gonzales, G., Gómez, J. Q., Encarnación, R. C. C.: High temporal resolution
- hydrometeorological data collected in the tropical Cordillera Blanca, Peru (2004-2020),
- 729 HydroShare, https://doi.org/10.4211/hs.059794371790407abd749576df8fd121, 2021.

- 730 McNulty, B. A., Farber, D. L., Wallace, G. S., Lopez, R., and Palacios, O.: Role of plate
- kinematics and plate-slip-vector partitioning in continental magmatic arcs: Evidence from the
 Cordillera Blanca, Peru, Geology, 26 (9), 827–830, 1998.
- Petersen, U., Sassarini, L., Plenge, R.,: Glaciar Yanasinga (Central Peru): 24 years of
 measurements, Journal of Glaciology, 8 (54), 487–489, 1969.
- 735 RGI Consortium: Randolph Glacier Inventory A Dataset of Global Glacier Outlines: Version
- 6.0: Technical Report, Global Land Ice Measurements from Space, Colorado, USA, Digital
- 737 Media, DOI: https://doi.org/10.7265/N5-RGI-60, 2017.
- 738 Schauwecker, S., Rohrer, M., Acuna, D., Cochachin, A., Davila, L., Frey, H., Giraldez, C.,
- Gomez, J., Huggel, C., Jacques-Coper, M., Loarte, E., Salzmann, N., and Vuille, M.: Climate
- trends and glacier retreat in the Cordillera Blanca, Peru, revisited, Global and Planetary Change,
- 741 119, 85-97, doi:10.1016/j.gloplacha.2014.05.005, 2014.
- Schwarb, M., Acuña, D., Konzelmann, T., Rohrer, M., Salzmann, N., Serpa Lopez, B., Silvestre,
 <u>E.: A data portal for regional climatic trend analysis in a Peruvian High Andes region, Advances</u>
 in Science and Research, 6, 219–226, 2011.
- 745 Vuille, M., Carey, M., Huggel, C., Buytaert, W., Rabatel, A., Jacobsen, D., Soruco, A., Villacis,
- 746 M., Yarleque, C., Elison Timm, O., Condom, T., Salzmann, N., and Sicart, J.-E.: Rapid decline
- of snow and ice in the tropical Andes impacts, uncertainties and challenges ahead, Earth-
- 748 Science Reviews, 176, 195–213, https://doi.org/10.1016/j.earscirev.2017.09.019, 2018.