

# An eleven year record of XCO<sub>2</sub> estimates derived from GOSAT measurements using the NASA ACOS version 9 retrieval algorithm

Thomas E. Taylor<sup>1</sup>, Christopher W. O'Dell<sup>1</sup>, David Crisp<sup>2</sup>, Akhiko Kuze<sup>3</sup>, Hannakaisa Lindqvist<sup>4</sup>, Paul O. Wennberg<sup>5</sup>, Abhishek Chatterjee<sup>6,7,\*</sup>, Michael Gunson<sup>2</sup>, Annmarie Eldering<sup>2</sup>, Brendan Fisher<sup>2</sup>, Matthäus Kiel<sup>2</sup>, Robert R. Nelson<sup>2</sup>, Aronne Merrelli<sup>8</sup>, Greg Osterman<sup>2</sup>, Frédéric Chevallier<sup>9</sup>, Paul I. Palmer<sup>10</sup>, Liang Feng<sup>10</sup>, Nicholas M. Deutscher<sup>11</sup>, Manvendra K. Dubey<sup>12</sup>, Dietrich G. Feist<sup>13,14,15</sup>, Omaira E. García<sup>16</sup>, David W. T. Griffith<sup>11</sup>, Frank Hase<sup>17</sup>, Laura T. Iraci<sup>18</sup>, Rigel Kivi<sup>19</sup>, Cheng Liu<sup>20</sup>, Martine De Mazière<sup>21</sup>, Isamu Morino<sup>22</sup>, Justus Notholt<sup>23</sup>, Young-Suk Oh<sup>24</sup>, Hirofumi Ohyama<sup>22</sup>, David F. Pollard<sup>25</sup>, Markus Rettinger<sup>26</sup>, Coleen M. Roehl<sup>27</sup>, Matthias Schneider<sup>26</sup>, Mahesh Kumar Sha<sup>21</sup>, Kei Shiomi<sup>3</sup>, Kimberly Strong<sup>28</sup>, Ralf Sussmann<sup>26</sup>, Yao Té<sup>29</sup>, Voltaire A. Velazco<sup>11,30</sup>, Mihalís Vrekoussis<sup>31,23</sup>, Thorsten Warneke<sup>23</sup>, and Debra Wunch<sup>28</sup>

<sup>1</sup>Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO, USA.

<sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA.

<sup>3</sup>Japan Aerospace Exploration Agency, Tsukuba-City, Ibaraki, Japan

<sup>4</sup>Finnish Meteorological Institute, Helsinki, Finland

<sup>5</sup>California Institute of Technology, Pasadena, CA, USA.

<sup>6</sup>Universities Space Research Association, Columbia, MD, USA

<sup>7</sup>Goddard Space Flight Center, Greenbelt, MD, USA

\* now at Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA.

<sup>8</sup>Space Science and Engineering Center, University of Wisconsin - Madison, Madison WI, 53706, USA.

<sup>9</sup>Laboratoire des Sciences du Climat et de l'Environnement/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91198 Gif-sur-Yvette, France

<sup>10</sup>National Centre for Earth Observation, University of Edinburgh, Edinburgh, UK

<sup>11</sup>Centre for Atmospheric Chemistry, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, Australia

<sup>12</sup>Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>13</sup>Max Planck Institute for Biogeochemistry, Jena, Germany

<sup>14</sup>Ludwig-Maximilians-Universität München, Lehrstuhl für Physik der Atmosphäre, Munich, Germany

<sup>15</sup>Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

<sup>16</sup>Izaña Atmospheric Research Centre (IARC), State Meteorological Agency of Spain (AEMET), Santa Cruz de Tenerife, Spain

<sup>17</sup>Karlsruhe Institute of Technology, IMK-ASF, Karlsruhe, Germany

<sup>18</sup>NASA Ames Research Center, Moffett Field, CA, USA

<sup>19</sup>Finnish Meteorological Institute, FMI, Sodankylä, Finland

<sup>20</sup>Department of Precision Machinery and Precision Instrumentation, University of Science and Technology of China, Hefei 230026, China

<sup>21</sup>Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

<sup>22</sup>National Institute for Environmental Studies (NIES), Tsukuba, Japan

<sup>23</sup>Institute of Environmental Physics, University of Bremen, Bremen, Germany

<sup>24</sup>Global Atmosphere Watch Team, Innovative Meteorological Research Department, National Institute of Meteorological Sciences, Jeju-do, Republic of Korea

<sup>25</sup>National Institute of Water and Atmospheric Research Ltd (NIWA), Lauder, New Zealand

<sup>26</sup>Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

<sup>27</sup>Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

<sup>28</sup>Department of Physics, University of Toronto, Toronto, Ontario, Canada

<sup>29</sup>Laboratoire d'Etudes du Rayonnement et de la Matière en Astrophysique et Atmosphères (LERMA-IPSL), Sorbonne Université, CNRS, Observatoire de Paris, PSL Université, Paris, France

<sup>30</sup>Deutscher Wetterdienst, Meteorological Observatory Hohenpeissenberg, 82383 Germany

<sup>31</sup>Climate and Atmosphere Research Center (CARE-C), The Cyprus Institute, Nicosia, Cyprus

**Correspondence:** Tommy Taylor (tommy.taylor@colostate.edu)

**Abstract.** The Thermal And Near infrared Sensor for carbon Observation - Fourier Transform Spectrometer (TANSO-FTS) on the Japanese Greenhouse gases Observing SATellite (GOSAT) has been returning data since April 2009. The version 9 (v9) Atmospheric Carbon Observations from Space (ACOS) Level 2 Full Physics (L2FP) retrieval algorithm (Kiel et al., 2019) was used to derive estimates of carbon dioxide (CO<sub>2</sub>) dry air mole fraction (XCO<sub>2</sub>) from the TANSO-FTS measurements collected over its first eleven years of operation. The bias correction and quality filtering of the L2FP XCO<sub>2</sub> product were evaluated using estimates derived from the Total Carbon Column Observing Network (TCCON) as well as values simulated from a suite of global atmospheric inversion systems (models) which do not assimilate satellite derived CO<sub>2</sub>. In addition, the v9 ACOS GOSAT XCO<sub>2</sub> results were compared with collocated XCO<sub>2</sub> estimates derived from NASA's Orbiting Carbon Observatory-2 (OCO-2), using the version 10 (v10) ACOS L2FP algorithm.

These tests indicate that the v9 ACOS GOSAT XCO<sub>2</sub> product has improved throughput, scatter, and bias, when compared to the earlier v7.3 ACOS GOSAT product, which extended through mid 2016. Of the 37 million (M) soundings collected by GOSAT through June 2020, approximately 20% were selected for processing by the v9 L2FP algorithm after screening for clouds and other artifacts. After post-processing, 5.4% of the soundings (2 M out of 37 M) were assigned a “good” XCO<sub>2</sub> quality flag, as compared to 3.9% in v7.3 (<1 M out of 24 M). After quality filtering and bias correction, the differences in XCO<sub>2</sub> between ACOS GOSAT v9 and both TCCON and models have a scatter (one sigma) of approximately 1 ppm for ocean-glint observations and 1 to 1.5 ppm for land observations. Global mean biases against TCCON and models are less than approximately 0.2 ppm. Seasonal mean biases relative to the v10 OCO-2 XCO<sub>2</sub> product are of order 0.1 ppm for observations over land. However, for ocean-glint observations, seasonal mean biases relative to OCO-2 range from 0.2 to 0.6 ppm, with substantial variation in time and latitude.

The ACOS GOSAT v9 XCO<sub>2</sub> data are available on the NASA Goddard Earth Science Data and Information Services Center (GES-DISC) ([OCO-2 Science Team et al., 2019a, b](#)), in both the per-orbit full format (OCO-2 Science Team et al., 2019b, <https://doi.org/10.5067/VWSABTO7ZI4>) and in the per-day lite format (OCO-2 Science Team et al., 2019a, <https://doi.org/10.5067/VWSABTO7ZI4>). In addition, a new set of monthly super-lite files, containing only the most essential variables for each satellite observation, has been generated to provide entry level users with a light weight satellite product for initial exploration (CaltechDATA, <https://orcid.org/0000-0003-1080-9921>). The v9 ACOS Data User's Guide (DUG) describes best-use practices for the GOSAT data (O'Dell et al., 2020). This dataset The GOSAT v9 data set should be especially useful for studies of carbon cycle phenomena that span a full decade or more, and may serve as a useful complement to the shorter OCO-2 v10 dataset data set, which begins in September 2014.



## 1 Introduction

A new era of dedicated satellite observations of greenhouse gases began in 2009, with the successful launch of GOSAT (Kuze et al., 2009). Each day, GOSAT's Thermal And Near infrared Sensor for carbon Observation - Fourier Transform Spectrometer (TANSO-FTS) acquires approximately ten thousand high spectral resolution measurements of reflected sunlight ( $\simeq 36.5$  M in  
 35 ten years). Soundings that are determined to be sufficiently clear of clouds and aerosols are processed by retrieval algorithms to produce estimates of  $\text{XCO}_2$ . Both the quality of the GOSAT TANSO-FTS spectra and the derived  $\text{XCO}_2$  estimates have been continually refined over the past twelve years. While the official GOSAT L2 products are available from the National Institute for Environmental Studies (NIES; [http://www.gosat.nies.go.jp/en/about\\_5\\_products.html](http://www.gosat.nies.go.jp/en/about_5_products.html); Yoshida et al., 2013) a number of independent research institutes have developed their own products e.g., Butz et al., 2011; Crisp et al., 2012; Cogan et al., 2012;  
 40 Heymann et al., 2015.

One of these groups, the Atmospheric  $\text{CO}_2$  Observations from Space (ACOS) team, used a Level 2 Full Physics (L2FP) retrieval algorithm developed for the NASA Orbiting Carbon Observatory (OCO) to derive estimates of  $\text{XCO}_2$  from the GOSAT data (O'Dell et al., 2012; Crisp et al., 2012). Early  $\text{XCO}_2$  estimates from these efforts had large biases and random errors when  
 45 compared to  $\text{XCO}_2$  estimates from the Total Carbon Column Observing Network (TCCON) and other standards. For example, the v2.8 ACOS GOSAT L2FP product had biases of 7 to 8 ppm relative to TCCON (Crisp et al., 2012). These biases were reduced to 1-2 ppm in the v2.9 product. The next major release was v3.5 in 2014, which spanned approximately 4 years. This data product showed additional reductions in bias and scatter against TCCON, as well as reasonable agreement in seasonal cycle phase and amplitude (Lindqvist et al., 2015; Kulawik et al., 2016).

50

These early space-based  $\text{XCO}_2$  products were rapidly adopted by the carbon cycle science community. Early studies based on GOSAT ACOS retrievals included Basu et al. (2013), Deng et al. (2014), Chevallier et al. (2014), and Feng et al. (2016). These studies provided the first comprehensive insights into regional flux estimates from space-based observations of carbon dioxide. Houweling et al. (2015) conducted an extensive inter-comparison of the early GOSAT-based atmospheric inversion system  
 55 studies and reported a reduction in the global land sink for  $\text{CO}_2$  and a shift in the terrestrial net uptake of carbon from the tropics to the extratropics. However, these studies also highlighted the role of spatiotemporal systematic errors in the satellite retrievals and the negative impact they can have on estimation of  $\text{CO}_2$  sources and sinks using atmospheric inversion systems.

Motivated by these early studies, as well as the launch of the OCO-2 sensor in July 2014, the ACOS team continued to refine the  
 60 L2FP retrieval algorithm. In 2016, the ACOS GOSAT v7.3 product was distributed. No formal results of the  $\text{XCO}_2$  estimates were published by the algorithm team, although internal analysis showed small improvement over v3.5, as well as an extension of the record to 7 years. A number of atmospheric inversion studies were published using the v7.3 product. For example, Chat-

terjee et al. (2017) and Liu et al. (2017) used v7.3 to define the climatological background in their studies of the impact of the 2015-2016 El Niño on the tropical carbon cycle. Palmer et al. (2019) used this data product in a global study, concluding that the tropical land regions were a net annual source of CO<sub>2</sub> emissions, including unexpectedly large net emissions from northern tropical Africa. Wang et al. (2019) found that the ACOS GOSAT v7.3 XCO<sub>2</sub> yielded a stronger carbon land sink than the v7 OCO-2 product. Byrne et al. (2020) used the ACOS GOSAT 7.3 product to study interannual variability in the carbon cycle across North America, and Jiang et al. (2021) investigated interannual variability of the carbon cycle across the globe with v7.3.

Most recently, the v9 ACOS L2FP retrieval algorithm, first applied to OCO-2 (Kiel et al., 2019), was used to generate estimates of XCO<sub>2</sub> from an eleven year record of GOSAT measurements, spanning April 2009 through June 2020. This both extends the time record over v7.3, and produces an ACOS GOSAT product that is more directly comparable to the newest OCO-2 product, which is now using version 10.

The paper is organized as follows; Section 2 discusses the GOSAT TANSO-FTS instrument and measurements as related to the ACOS XCO<sub>2</sub> estimates. In Section 3, updates to the ACOS v9 L2FP algorithm are detailed, and an assessment is given of the v9 XCO<sub>2</sub> data product volume. The XCO<sub>2</sub> quality filtering and bias correction procedures, specific to ACOS GOSAT v9, are also discussed. Section 4 provides an evaluation of the v9 XCO<sub>2</sub> product using estimates of XCO<sub>2</sub> from TCCON and from a suite of 4 atmospheric inversion systems (models). In addition, a comparison to collocated XCO<sub>2</sub> estimates derived from NASA’s OCO-2 sensor is presented. A summary of the results is provided in Section 5.

## 2 The GOSAT instrument and measurements

The GOSAT mission is a joint project between the Japan Aerospace Exploration Agency (JAXA), the National Institute for Environmental Studies (NIES), and the Ministry of the Environment (MOE) (Kuze et al., 2009). GOSAT was launched on 23 January 2009 into a sun-synchronous orbit with a local overpass time of approximately 12:49PM and a 3 day ground repeat cycle. Its TANSO-FTS collects high resolution spectra of reflected sunlight that can be analyzed to yield estimates of carbon dioxide (CO<sub>2</sub>) (Yoshida et al., 2011, 2013).

### 2.1 GOSAT TANSO-FTS instrument

TANSO-FTS collects high resolution spectra of reflected sunlight in the near infrared (NIR) and short-wave infrared (SWIR) spectral ranges that include the the oxygen A-band near 0.76  $\mu\text{m}$  (ABO2 band) at approximately  $0.36\text{ cm}^{-1}$  spectral resolution, and weak and strong CO<sub>2</sub> absorption features near 1.6  $\mu\text{m}$  (WCO2 band) and 2.0  $\mu\text{m}$  (SCO2 band), respectively, at  $0.27\text{ cm}^{-1}$  spectral resolution. All three channels simultaneously measure two orthogonal components of polarization approximately ev-

ery 4.6 seconds.

95

Each GOSAT sounding has a circular ground footprint with a diameter of approximately 10.5 km when viewing the local nadir. An agile, two-axis pointing system allows cross-track and along-track motions of  $\pm 35^\circ$  and  $\pm 20^\circ$ , respectively. Before August 2010, a 5-point cross-track scan was used, yielding footprints that were separated by approximately 150 km in both the down-track and along-track dimensions. Since that time, a 3-point cross-track scan has been used, yielding footprint separation  
100 of approximately 260 km (Kuze et al., 2016).

Over water, the TANSO-FTS scan mechanism targets the field of view to collect observations in the direction of the local glint spot, where sunlight is specularly reflected from the surface. Early in the mission, glint observations were collected only within  $\pm 20^\circ$  of the sub-solar latitude. In May 2013, to increase the latitudinal extent of the GOSAT Ocean-Glint measurements, the  
105 scanning strategy was improved to better track the actual specular glint spot, which varies by latitude and season. The latitude range for glint observation was further extended three times in increments of  $3^\circ$  in September 2014, June 2015, and January 2016, by not only tracking the exact specular point but also tracking along the principal plane of the specular reflection when the glint spot was out of range of the scan mechanism. In addition, more observations over fossil fuel emission target sites such as mega-cities and power plants have been made in recent years, allowing for detailed emission source studies (e.g., Kuze  
110 et al., 2020). Daily observation patterns can be found at [https://www.eorc.jaxa.jp/GOSAT/currentStatus\\_10.html](https://www.eorc.jaxa.jp/GOSAT/currentStatus_10.html).

The TANSO-FTS detectors can be read out using independent medium-gain and high-gain signal chains. Most measurements over land use the instrument's high-gain signal chain (Land H-gain), while brighter land surfaces are measured using the medium-gain signal chain (Land M-gain) to avoid saturating the detectors. Over oceans, which appear dark in the SWIR spec-  
115 tral bands, measurements are collected using the high-gain signal chain (Ocean H-gain or Ocean-Glint) to maximize the signal.

During the first 7 years of GOSAT operations (2009-2015), data acquisition was temporarily suspended due to one spacecraft and two instrument anomalies, as highlighted in Kuze et al. (2016). A rotation failure of a solar paddle in 2014 resulted in a data loss of 6 days. A switch from the primary to secondary pointing mirror in January 2015 resulted in a data loss of approximately  
120 6 weeks, while a temporary shutdown of the cryocooler in August 2015 resulted in a data loss of 13 days.

Since 2015, three additional anomalies interrupted data acquisition. An unexpected shutdown of the instrument occurred in May 2018, resulting in the loss of a week of data. A failure of the second solar panel caused a significant loss of data spanning more than a month in November and December of 2018, and an anomaly of the FTS alignment laser, caused a loss of a week of  
125 data in June of 2020. In all these cases, the system was able to recover full functionality either through utilization of on-board back-up systems, or through mitigation strategies, and as of the summer of 2021, TANSO-FTS continues to collect science data.

## 2.2 ACOS GOSAT v9 L1b measurements

The JAXA L1b algorithm, which has been updated more than 10 times over the eleven year data record, produces an internally  
130 consistent set of geometrically, radiometrically, and spectrally calibrated TANSO-FTS radiances. The raw spectral measure-  
ments are interferograms, which are calibrated and Fourier transformed to yield spectra. The version 205/210 Level 1b (L1b)  
geolocated and calibrated radiances provided by JAXA have been used for the ACOS v9 reprocessing. A list of L1b updates  
for v205/210 can be found in Table 3 of the ACOS v9 Data Users Guide (DUG) (O'Dell et al., 2020). Note that while the  
current L1b version is now 230, the only differences between this version and 205/210 are in the thermal infrared band (5.6 -  
135 14.3  $\mu\text{m}$ ), which is not used in the ACOS XCO<sub>2</sub> retrieval.

After obtaining the calibrated L1b product from JAXA, the ACOS team converts the files to the format needed as input to the  
ACOS L2 algorithms. The L2FP algorithm uses a simple average of the S and P linear polarizations to produce an approxima-  
tion of the total measured intensity. Due to cooperation agreements between JAXA and the California Institute of Technology,  
140 the distribution of the ACOS GOSAT L1b product is restricted and therefore not publicly available on the NASA DISC. How-  
ever, the data may be procured by submitting a request to the GOSAT project.

## 3 The ACOS v9 L2FP XCO<sub>2</sub> retrieval algorithm

The ACOS Level 2 full physics (L2FP) retrieval algorithm is well documented, most recently in O'Dell et al. (2018) for v8  
145 and in Kiel et al. (2019) for v9. A Bayesian optimal estimation framework is used to derive estimates of XCO<sub>2</sub> from spectral  
measurements of reflected solar radiation. A post-processing step assigns a simple good/bad quality flag (QF) to each XCO<sub>2</sub>  
value based on successful L2FP algorithm convergence and a series of empirically derived filters. An empirical bias correction  
(BC) to the estimated XCO<sub>2</sub> values, derived from comparisons with TCCON derived XCO<sub>2</sub> and CO<sub>2</sub> fields from a suite of  
atmospheric inversion systems, is included in the Lite File product. Here we provide a summary of the recent evolution of the  
150 ACOS algorithm and discuss retrieval parameters and setup specific to GOSAT.

### 3.1 ACOS L2FP algorithm updates

Table 1 summarizes the evolution of the ACOS L2FP retrieval algorithm from v7 to v10. A similar table, complete through v8,  
can be found in O'Dell et al. (2018). The trace gas absorption coefficient tables (ABSCO) were updated from v4.2 (Thompson  
155 et al., 2012) in ACOS v7 to ABSCO v5.0 (Oyafuso et al., 2017) in ACOS v8/9. The ACOS v9 L2FP algorithm is unmodi-  
fied relative to v8 (Kiel et al., 2019). However, changes were made in v9 regarding the sampling of the meteorological prior,  
which does affect ACOS GOSAT estimates of XCO<sub>2</sub>. The source of the prior meteorology was switched from the European  
Center for Medium-range Weather Forecast (ECMWF) in ACOS v7, to the NASA Goddard Modeling and Assimilation Of-

**Table 1.** Updates to recent versions of the ACOS L2FP retrieval algorithm. N/C stands for No Change.

		ACOS v7	ACOS v8/v9	ACOS v10
1	Spectroscopy	ABSCO v4.2	ABSCO v5.0	ABSCO v5.1
2	Meteorology prior source	ECMWF	GEOS5 FP-IT	N/C
3	Aerosol prior source	MERRA monthly climatology	N/C	GEOS5 FP-IT with tightened prior uncertainty
4	Retrieved aerosol types	water + ice + 2 MERRA types	+ stratospheric aerosol	N/C N/C
5	AOD prior value (per type)	0.0375	0.0125	N/C
6	CO <sub>2</sub> prior source	TCCON ggg2014	N/C	TCCON ggg2020
7	Land surface model	Lambertian	BRDF	N/C

160 fice (GMAO) Goddard Earth Observing System (GEOS) Forward Processing - Instrument Team (FP-IT) product for ACOS v8/9. Both v7 and v8/9 used aerosol priors based on a simple monthly 1° latitude by 1° longitude climatology constructed from the output aerosol fields of the GMAO Modern-Era Retrospective analysis for Research and Applications (MERRA) product (Rienecker et al., 2011). However, between v7 and v8/9, an additional stratospheric aerosol layer was introduced, as described in Section 3.1.1 of O’Dell et al. (2018). In addition, the prior value of the aerosol optical depth (AOD) for each retrieved aerosol type was lowered from 0.0375 in ACOS v7 to 0.0125 in ACOS v8/9 based on extensive testing. There was  
165 no change in the source of the CO<sub>2</sub> prior from ACOS v7 to v8/9; both versions adopted the prior developed by the TCCON team for use in the ggg2014 algorithm (Wunch et al., 2015). An additional change from ACOS v7 to v8/9 was a switch from a purely Lambertian land surface model, to a more sophisticated bi-directional reflectance distribution function (BRDF) model.

Several important components of the v9 ACOS L2FP retrieval configured for GOSAT have not changed from v7.3; (i) the sur-  
170 face pressure prior constraint remains set at  $\pm 2$  hPa, (ii) three Empirical Orthogonal Functions (EOFs) are fit in each spectral band (see Section 3.3 in O’Dell et al. (2018) for a full discussion of ACOS EOFs), and (iii) a zero level offset (ZLO) is fit in the state vector to account for non-linearity in the ABO2 signal chain on GOSAT TANSO-FTS (Crisp et al., 2012).

To support comparisons of the ACOS GOSAT v9 XCO<sub>2</sub> product with the OCO-2 v10 product, Table 1 includes the most recent  
175 updates to the ACOS v10 L2FP algorithm. For v10, the ABSCO tables were again updated from v5.0 to v5.1 (Payne et al., 2020). The aerosol prior was updated from the MERRA monthly climatology to daily GEOS-FT-IT values, with a tightened prior uncertainty (Nelson and O’Dell, 2019). Finally, the CO<sub>2</sub> priors developed by the TCCON team for use in ggg2014 were updated to a revised set of priors developed for use in ggg2020.

**Table 2.** Accounting of the soundings in the eleven year long GOSAT ACOS v9 dataset at each stage of the data processing chain. The final line summarizes the number of good quality XCO<sub>2</sub> soundings used in the evaluation section of this work.


	Reference	Number of soundings (N)	Total (%)	Fraction of Selected (%)    Valid (%)	
Total in ACOS GOSAT v9 record	Figure 1A	37.4 M	100.	–	–
Selected for L2FP	Figure 1B	7.0 M	18.8	100.	–
Non-convergence (terminated at unphysical state)	–	0.3 M	0.7	3.9	–
Non-convergence (exceeded iteration limit)	–	0.2 M	0.6	3.2	–
Non-convergence (exceeded diverging steps limit)	–	0.4 M	1.1	5.9	–
Valid, converged L2FP XCO <sub>2</sub> result	–	6.1 M	16.4	87.0	100.
Lite file aggregator IDP filtering	–	0.3 M	0.7	3.9	4.5
Bad L2Lite quality flag	–	3.9 M	10.4	55.3	63.6
Good L2Lite quality flag	Figure 1C&D	2.0 M	5.4	28.6	32.9

180    **3.2    ACOS GOSAT v9 L2FP sounding selection and convergence**

GOSAT data from April 20, 2009 through June 30, 2020 were passed through the ACOS L2FP algorithm pipeline, which includes a series of stages where soundings can be rejected or selected for further processing. The throughput of each of these stages for ACOS GOSAT v9 is summarized in Table 2 and Figure 1. The pipeline begins with a series of preprocessing steps, which reject corrupted spectra and screen the remainder to eliminate those with optically-thick clouds and/or aerosols (Taylor et al., 2016). From the full set of measurements (Panel A of Figure 1), the remaining soundings are accepted by the L2FP algorithm (18.8% of the 37.4 M measured soundings contained in the ACOS GOSAT v9 record) (Panel B of Figure 1) and a retrieval of XCO<sub>2</sub> is attempted. The majority of the selected soundings successfully converge to a valid solution; 87% for ACOS GOSAT v9 (16.4% of the total measured soundings). Soundings can fail to converge for a variety of reasons, including (i) producing non physical values, such as negative gas mixing ratios or surface pressures (3.9% of the selected), (ii) converging too slowly and exceeding a predefined number of iterations (3.2% of the selected), or (iii) having more diverging steps than the predefined maximum (5.9% of the selected). The 6.1 M valid soundings were then run through the quality filtering and bias correction procedure discussed in the next section.

195    **3.3    ACOS GOSAT v9 XCO<sub>2</sub> quality filtering and bias correction**

All GOSAT soundings that converged to a valid XCO<sub>2</sub> value within the L2FP retrieval were input to the quality filtering and bias correction procedure. A modest fraction (4.5% of the valid soundings) were removed from the final L2Lite product based on screening via the IMAP-DOAS Preprocessor (IDP) CO<sub>2</sub> ratio, which indicated the presence of clouds or aerosols. Based on



**Figure 1.** The total measured sounding density per  $2.5^\circ$  by  $5^\circ$  latitude/longitude grid cell in the eleven year (April 2009 - June 2020) ACOS GOSAT v9 data record (A). The fraction of the total soundings selected to run through the L2FP algorithm (B). The fraction of the total soundings that converged in the L2FP and were assigned a good L2FP QF (C). The sounding density of the good QF data per  $2.5^\circ$  by  $5^\circ$  latitude/longitude grid cell (D).

a series of screening criteria derived from comparisons with TCCON and modeled  $\text{CO}_2$  fields, each sounding that converged within the L2FP is assigned either a “good” (=0) or “bad” (=1)  $\text{XCO}_2$  quality flag. Generally, for global or regional studies, it is recommended that users retain only the “good” quality soundings, as the soundings flagged as “bad” quality are likely to include biases that compromise their utility for some applications. A global map of the ACOS GOSAT v9 “good”  $\text{XCO}_2$  sounding density is provided in panels C and D of Figure 1. A subset of data variables from the per-orbit L2Std files (OCO-2 Science Team et al., 2019b), along with the quality filter flag and bias corrected  $\text{XCO}_2$ , are repackaged into the daily aggregated L2Lite NetCDF files (OCO-2 Science Team et al., 2019a).

205

A fundamental aspect of the quality filtering and bias correction procedures (QF/BC) is the need for  $\text{XCO}_2$  truth metrics with which to compare the satellite derived estimates (O’Dell et al., 2018). The development of ACOS GOSAT v9 used  $\text{XCO}_2$  truth metrics derived from both TCCON measurements, and the median  $\text{CO}_2$  distributions determined from a suite of four atmospheric inversion systems, which do not assimilate satellite  $\text{CO}_2$  measurements.

210

TCCON is a well established validation transfer standard for space-based estimates of XCO<sub>2</sub> (Wunch et al., 2011a, 2017b). For the ACOS GOSAT v9 QF/BC, estimates of XCO<sub>2</sub> derived from TCCON measurements using the ggg2014 retrieval algorithm were used (Wunch et al., 2015). Individual GOSAT soundings were compared to TCCON daily mean XCO<sub>2</sub> values. TCCON data were included if: (i) they were flagged good (flag = 0), (ii) they fell within 3 standard deviations of a daily quadratic fit  
215 against time (to remove outliers, e.g. due to unscreened cloud), (iii) they covered at least 15 minutes within a given day, (iv) there were at least 3 good soundings within the day, and (v) the standard deviation of the good soundings for the day was less than 3 ppm. In the GOSAT-TCCON comparisons described here, an averaging kernel correction was applied to each TCCON XCO<sub>2</sub> estimate following Nguyen et al. (2014), prior to calculating the daily mean value.

220 Following the criteria defined in Wunch et al. (2017b), the spatial collocation criteria for GOSAT soundings were those falling within  $\pm 2.5^\circ$  latitude and  $\pm 5^\circ$  longitude of a TCCON station for most sites. For southern hemisphere (SH) sites poleward of  $25^\circ$  S latitude, where the variation of CO<sub>2</sub> is low, the spatial box was increased to  $\pm 10^\circ$  latitude by  $20^\circ$  longitude to increase the number of collocations. For the Edwards TCCON station, which lies in an arid region just north of the polluted Los Angeles metropolitan area, a very specific collocation box of [34.68, 37.46] latitude and [-127.88, -112.88] longitude was used to avoid  
225 contamination from the city. Similarly, for the Caltech site, located in Pasadena, California, a latitude box of [33.38, 34.27] and longitude box of [-118.49, -117.55] was used. This avoids collocating GOSAT soundings measured over ocean, the San Gabriel Mountains, and regions too far outside of the Los Angeles basin with the Caltech TCCON data. Finally, only GOSAT soundings acquired within  $\pm 2$  hours of the mean TCCON measurement time were considered. For the quality filtering and bias correction procedure, single sounding level collocations are used to maximize the number of fit points.

230 Estimates of CO<sub>2</sub> from atmospheric inversion systems, or models, provide a useful metric for evaluating satellite based estimates of XCO<sub>2</sub> (O'Dell et al., 2018). In this work, a suite of four models (CarbonTracker, CAMS, CarboScope, and Univ. of Edinburgh) were sampled at the GOSAT sounding times and locations. Brief descriptions of each, along with references, are provided in Table 3. The models use a variety of land biosphere prior fluxes, inverse solvers and transport models, and assimilate CO<sub>2</sub> data only from flasks and continuous analyzers on a wide variety of platforms, e.g., observatories, towers, aircraft,  
235 and ships. Specifically, no data from GOSAT, OCO-2, or TCCON are assimilated. The CO<sub>2</sub> concentration fields of the models capture the known features of the global atmospheric CO<sub>2</sub> distribution, including seasonality, time trends and inter-annual variability (IAV) due to ENSO. For each GOSAT sounding, the vertical profiles of CO<sub>2</sub> from the corresponding grid box of each of the four models are spatiotemporally interpolated (linear in latitude, longitude, and time) to the GOSAT observation  
240 point, and the GOSAT averaging kernel is applied to each vertical profile to produce a modeled XCO<sub>2</sub> as if viewed from the satellite.

For each GOSAT sounding, a multi-model median (MMM) XCO<sub>2</sub> was calculated from the models having a valid XCO<sub>2</sub> estimate for that location and time. Unless otherwise noted, the model XCO<sub>2</sub> is taken to be that which a perfect OCO-2 would



**Table 3.** Carbon inversion systems used for ACOS GOSAT v9.

Model Name	Institute	Land Biosphere Prior	Transport Model	Inverse Method	Citations
CarbonTracker	NOAA Global Monitoring Laboratory	CASA	TM5	EnKF	Peters et al. (2007) CarbonTracker (2021)
CarboScope	Max Planck Institute for Biogeochemistry	Zero	TM3	4D-Var	Rödenbeck (2005); Rödenbeck et al. (2018) CarboScope (2021)
CAMS	Copernicus Atmosphere Monitoring Service	ORCHIDEE	LMDZ	4D-Var	Chevallier et al. (2010) CAMS (2021)
UoE	University of Edinburgh Atmospheric Composition Modelling Group	CASA	GEOS-Chem	EnKF	Feng et al. (2009) UoE (2021)

245 have observed,  $XCO_{2,ak}$ ; that is, an averaging kernel correction is applied to account for differences between the model profile of  $CO_2$  and the ACOS prior in the unmeasured part of the profile:

$$XCO_{2,ak} = \sum_{i=1}^{20} h_i \{a_i u_{m,i} + (1 - a_i) u_{a,i}\}, \tag{1}$$

where  $h_i$  is the pressure weighting function on the  $i = 1 \dots 20$  ACOS model levels,  $a$  is the normalized ACOS averaging kernel for  $CO_2$ ,  $u_m$  is the model profile of  $CO_2$ , and  $u_a$  is the ACOS prior profile of  $CO_2$ .

250

To exclude outliers, models with  $XCO_2$  that deviated more than  $\pm 1.5$  ppm from the initial MMM for that sounding were not included. The sounding was then rejected if more than one of the four models had been excluded, or if the standard deviation amongst the valid models was  $> 1$  ppm. Approximately 85% of the GOSAT v9 soundings with a good L2FP quality flag had a valid MMM  $XCO_2$  value for analysis. The regions with the highest fraction of rejections occur along the southern ocean (latitude  $-60^\circ$ ), the Amazon and Congo rain forests, and a broad region across northern Asia. Table 4 lists the model version numbers used for the QF/BC procedure, as well as that used in the evaluation of the final good quality  $XCO_2$  product that will be presented later.

255

Table 5 lists the quality filtering variables used for ACOS GOSAT v9 and their corresponding thresholds. Many of the same variables (18 out of 31) were also used in the OCO-2 v9 quality filtering, as seen in Table 5 of Kiel et al. (2019). This includes the IDP  $CO_2$  and  $H_2O$  ratios (Frankenberg et al., 2005), and the A-Band Preprocessor  $dP$ , i.e., the difference between the retrieved and prior surface pressure from the Oxygen-A band (Taylor et al., 2016). Another common variable used for quality

260

**Table 4.** Carbon inversion system data sets used for the QF/BC and XCO<sub>2</sub> evaluation of ACOS GOSAT v9.

Model	QF/BC	Evaluation
CarbonTracker	CT2017 (through 20170429) CT-NRT.v2019-2 (through 20190330)	CT2019 (through 20181231)
CarboScope	Jena_s04c_v4.3	Jena_s10oc-v2020
CAMS	v18r2 (second release of CAMS data that extends through 2018)	v20r1 (first release of CAMS data that extends through 2020)
UoE	v4.0 (used in Palmer et al. (2019))	v4.0a (an extension of v4.0, using near real-time in situ data for 2019.

filtering is the perturbation in the L2FP CO<sub>2</sub> vertical profile relative to the prior, a quantity called “CO<sub>2</sub> grad del” ( $\delta \nabla_{\text{CO}_2}$ ), as defined in equation 5 of O’Dell et al. (2018). A number of aerosol related retrieval parameters are also used, similar to OCO-2  
265 v9. Section 2.5 of the ACOS GOSAT v9 DUG provides additional details on the quality filtering (O’Dell et al., 2020).

Spurious correlations in the estimates of XCO<sub>2</sub> with other retrieval variables due to inadequacies in the modeled physics motivate the application of a bias correction (Wunch et al., 2011b; O’Dell et al., 2018). Generally such spurious correlations are found with state vector elements such as retrieved surface pressure, various aerosol parameters, and  $\delta \nabla_{\text{CO}_2}$ . For each  
270 sensor there are also typically offsets by viewing mode, e.g., land versus ocean-glint, which are accounted for via the bias correction. A general discussion of the ACOS XCO<sub>2</sub> bias correction methodology is provided in Section 4 of O’Dell et al. (2018), where the fundamental equation is defined as:

$$\text{XCO}_{2,\text{bc}} = \frac{\text{XCO}_{2,\text{raw}} - \text{C}_\text{P}(\text{mode}) - \text{C}_\text{F}(j)}{\text{C}_0(\text{mode})}, \tag{2}$$

where  $\text{C}_\text{P}$  is the mode-dependent parametric bias,  $\text{C}_\text{F}$  is a footprint-dependent bias for footprints 1...8, and  $\text{C}_0$  represents a  
275 mode-dependent global scaling factor. Note that for GOSAT there is no footprint-dependent bias correction term, as is necessary for OCO due to low level calibration errors across the detector frame. Further, to be consistent with previous ACOS GOSAT data versions, the global divisor is replaced by an additive offset, which is effectively the same because the range of XCO<sub>2</sub> variability ( $\sim 20$  ppm) is small relative to the mean XCO<sub>2</sub> ( $\sim 400$  ppm).

280 The explicit formula for application of the ACOS GOSAT v9 correction is provided in Section 2.5.6 of the DUG (O’Dell et al., 2020). For both land H-gain and M-gain, a set of five BC variables are used, while Ocean-Glint uses only 3 variables. The difference between the H- and M-gain bias correction over land is minor. New for ACOS GOSAT v9 is the use of a correction against time, which is made possible with an eleven year data record; the corrections are 0.05 ppm/yr over land and 0.10 ppm/yr over water. The source of this spurious drift in the bias-corrected XCO<sub>2</sub> is currently unclear and is the subject of on going study.  
285 Although there is some commonality in the quality filtering and bias correction variables used for ACOS GOSAT v9 (compare

**Table 5.** ACOS GOSAT v9 L2FP quality filtering variables and thresholds. Descriptions of the variables can be found in the DUG (O’Dell et al., 2020). Soundings falling outside of the data ranges are assigned a bad XCO<sub>2</sub> quality flag. The second column identifies variables that were also used for OCO-2 v9 quality filtering, as taken from Table 5 of Kiel et al. (2019).

Variable	Used for OCO-2 v9	Ocean-Glint	Land H-gain	Land M-gain
Geo altitude $\sigma$	Y	NA	< 250.0	< 250.0
Geo airmass	N	< 3.0	–	–
L1b SCO2/WCO2 signal ratio	N	> 0.58	–	–
IDP CO <sub>2</sub> ratio	Y	[0.989, 1.02]	[0.95, 1.02]	[0.989, 1.012]
IDP H <sub>2</sub> O ratio	Y	–	[0.80, 1.04]	[0.88, 1.05]
ABP $dP$ (retrieved - prior $p_{\text{surf}}$ )	Y	[-25.0, 14.0]	[-7.0, 7.0]	[-10.0, 7.0]
L2FP outcome flag	Y	1 or 2	1 or 2	1 or 2
L2FP Total AOD	Y	< 0.5	[0.02, 0.3]	< 0.4
L2FP AOD ice cloud	Y	< 0.07	< 0.06	[0.002, 0.05]
L2FP AOD sulfate aerosol	N (used organic carbon and sea salt, independently)	–	< 0.20	–
L2FP AOD stratospheric aerosol	Y	–	–	[0.0008, 0.015]
L2FP AOD DWS (dust + sea salt + water cloud)	Y	–	–	[0.0001, 0.35]
L2FP AOD fine (organic carbon + sulfate aerosol)	N	< 0.18	–	< 0.04
L2FP ice cloud pressure height	Y	[-0.50, 0.40]	[-0.12, 0.40]	[-0.12, 0.30]
L2FP dust aerosol pressure height	N	–	[0.75, 1.4]	[0.80, 1.4]
L2FP XCO <sub>2</sub> uncertainty	Y	–	< 2.0	< 1.5
L2FP CO <sub>2</sub> grad del ( $\delta \nabla_{\text{CO}_2}$ )	Y	[-19.0, 10.0]	[-40.0, 100.0]	[-10.0, 100.0]
L2FP $dP$ (retrieved - prior $p_{\text{surf}}$ )	Y	[-0.75, 5.5]	[-2.0, 10.0]	[-6.0, 5.0]
L2FP WCO2 albedo	N	[0.017, 0.030]	–	–
L2FP SCO2 albedo	Y	–	[0.04, 1.0]	[0.10, 1.0]
L2FP ABO2 albedo slope	N	–	[-4E-5, 2E5]	–
L2FP WCO2 albedo slope	Y	[3E-5, 2.7E-5]	–	–
L2FP SCO2 albedo slope	Y	[0.0, 5E-5]	[-1E-4, 2.5E-4]	–
L2FP ABO2 $\chi^2$	N	–	< 1.2	< 1.25
L2FP WCO2 $\chi^2$	Y	< 1.4	–	< 1.4
L2FP SCO2 $\chi^2$	N	< 1.35	–	< 1.9
L2FP ABO2 offset	N	–	–	[-1.5, 0.1]
L2FP temperature offset	N	–	[-1.0, 10.0]	[-0.7, 1.2]
L2FP ABO2 EOF 3 scaling	N	[-0.05, 0.04]	–	–
L2FP SCO2 EOF 2 scaling	N (used EOF 3)	[-0.15, 0.35]	–	–
L2FP wind speed	Y	[2.0, 24.0]	–	–

Tables 5 and 6), they do differ somewhat, as is typically the case with each sensor and data version.

Table 6 compares the bias correction variables used for ACOS GOSAT v9 with the variables used in the previous ACOS GOSAT v7.3, as well as with OCO-2 v9 and v10. The same few variables have appeared in all recent versions, including L2FP  $\delta \nabla_{\text{CO}_2}$ , L2FP  $dP$ , and L2FP DWS for land soundings. For ocean soundings the bias correction variables have evolved, with the only common one being  $\delta \nabla_{\text{CO}_2}$ .

**Table 6.** ACOS L2FP bias correction variables by sensor and product version.

	ACOS GOSAT v7.3	ACOS GOSAT v9	OCO-2 v9	OCO-2 v10	Variable Description
Land	(LandH only)	(LandH/LandM)	(nadir & glint)	(nadir & glint)	
	$\delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}, \delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}$	“CO <sub>2</sub> grad del” (see text)
	$dP$	–	–	–	Retrieved minus a priori surface pressure
	–	$dP_{\text{frac}}, dP_{\text{frac}}$	$dP_{\text{frac}}$	$dP_{\text{frac}}$	Elevation adjusted $dP$
					See equation 4 in (Kiel et al., 2019)
	DWS	$\text{DWS}, \sqrt{(\text{DWS})}$	DWS	$\log(\text{DWS})$	Combined aerosol optical depth of dust, water, and salt
	$\sqrt{\alpha_{\text{SCO}_2}}$	$\alpha_{\text{SCO}_2}, \sqrt{\alpha_{\text{SCO}_2}}$	–	–	Square-root of the retrieved albedo in the SCO <sub>2</sub> spectral band
	–	–	–	$\text{AOD}_{\text{fine}}$	Combined aerosol optical depth of sulfate and organic carbon
	–	$t_{\text{year}}$	–	–	Time in years
Ocean					
	$S_{32}$	–	–	–	Average signal in the WCO <sub>2</sub> and SCO <sub>2</sub> spectral bands
	$\delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}$	$\delta \nabla_{\text{CO}_2}$	“CO <sub>2</sub> grad del” (see text)
	$\log(\text{AOD}_{\text{dust}})$	–	–	–	Logarithm of dust AOD
	$H_{\text{ice}}$	–	–	–	Vertical height of ice cloud
	–	$\text{EOF}_{\text{SCO}_2}^3$	–	–	Third Empirical Orthogonal Function in the SCO <sub>2</sub> spectral band
	–	–	$dP$	$dP$	Retrieved minus a priori surface pressure
	–	$t_{\text{year}}$	–	–	Time in years

Table 7 summarizes the effect of the quality filtering and bias correction on the ACOS GOSAT XCO<sub>2</sub> for v7.3 and v9. For Ocean-Glint soundings, the v9 quality flag is substantially more restrictive compared to v7.3, i.e.  $\simeq 57\%$  pass rate compared to  $\simeq 78\%$ . This is mostly driven by the more extensive latitudinal coverage in the v9 record, which tends to include more soundings with high solar zenith angles (SZA) and low signal to noise ratio (SNR), which are more challenging for the L2FP. For H-gain land observations, the two versions have quite similar QF pass rates ( $\simeq 35\text{-}45\%$ ). The QF pass rate for v9 M-gain Land data is  $\simeq 39\%$  when compared against models, but  $\simeq 56\%$  against TCCON. In all cases there is a significant reduction in the scatter of the XCO<sub>2</sub> after application of the QF/BC; by a factor of  $\simeq 2$  for Ocean-Glint and Land M-gain, and a factor of 3 for Land H-gain. The QF/BC scatter is always slightly lower for v9 compared to v7.3, even though the number of soundings is greater by 1.5 to 10 times for the various scenarios.


**Table 7.** Comparison of ACOS GOSAT v7.3 versus v9 sounding throughput and XCO<sub>2</sub> scatter against truth metrics before and after filtering and bias correction.

Mode	Truth Metric	Version	N (soundings)	Throughput (%)	Sigma (ppm)	
					Unfiltered Raw	QF & BC
Ocean-Glint	Models	v7.3	82k	78%	1.7	1.0
		v9	1131k	56%	1.9	0.9
	TCCON	v7.3	2k	77%	2.0	1.2
		v9	15k	58%	2.4	1.1
Land H-gain	Models	v7.3	546k	37%	5.2	1.5
		v9	760k	37%	5.1	1.4
	TCCON	v7.3	5k	45%	4.3	1.7
		v9	56k	47%	4.4	1.6
Land M-gain	Models	v9	286k	39%	2.7	1.1
	TCCON	v9	9k	56%	2.8	1.1

Figure 2 shows the relative magnitudes of the bias correction on the good quality soundings by season, aggregated to 2.5° latitude by 5° longitude. The global median bias of -1.8 ppm has been removed for clarity. This highlights gradients and contrasts in the bias correction, which are of importance as gradients in CO<sub>2</sub> concentrations are the primary driver of CO<sub>2</sub> fluxes in atmospheric inversion systems. In general, the bias correction is necessary to remove spurious contrasts between land and ocean-glint XCO<sub>2</sub> values. The strongest relative bias corrections are positive adjustments over the bright land surfaces in M-gain viewing mode, specifically the Sahara in DJF and JJA, and Australia in DJF. The Land H-gain observations have a mix of relative bias correction values, ranging from mildly negative over high northern latitudes in JJA, to moderately positive over northern mid-latitudes in JJA in the western U.S. and the Middle East. Most of the Ocean-Glint observations have a mildly negative relative bias correction, with some mild positive values in the southern tropical oceans in DJF.

#### 4 Evaluation of ACOS GOSAT v9 XCO<sub>2</sub>


The ACOS GOSAT v9 XCO<sub>2</sub> record was characterized in five ways: (i) an analysis of the XCO<sub>2</sub> “good quality” data volume, (ii) a spatiotemporal analysis of the XCO<sub>2</sub> estimates, (iii) a validation against XCO<sub>2</sub> estimates from TCCON, (iv) a comparison to XCO<sub>2</sub> derived from models, and (v) a comparison with collocated XCO<sub>2</sub> estimates from the OCO-2 v10 product.



**Figure 2.** Maps of ACOS GOSAT v9 relative bias correction for DJF (A) and JJA (B) for good QF soundings in the eleven year data record at 2.5° by 5° latitude/longitude, after removal of the global, median bias of -1.8 ppm. Grid cells with less than 5 GOSAT soundings are not colored. The total number of soundings (N), and the mean bias ( $\mu$ ) and standard deviation of the bias ( $\sigma$ ) in each grid cell are given.

#### 315 4.1 ACOS GOSAT v9 “good quality” data volume


It is instructive to compare the ACOS GOSAT v9 product to the earlier v7.3 product to highlight similarities and differences in the quality filter screening. A time series histogram of the monthly throughput of the good quality filtered soundings for the v9 product compared to v7.3 is shown in Figure 3. The soundings have been binned by month, with the three GOSAT observation modes displayed by color. The v7.3 product did not contain any Land M-gain data in the L2Lite files (red in the figure) as the quality filtering and bias correction were not developed for that gain mode in v7.3 due to some unreconciled differences. An important feature of the v9 data record is the extension in time, which runs through June 2020, compared to a termination date of June 2016 for v7.3. Even for the overlapping v7.3 and v9 time period (2009 through mid 2016), there are some differences in the data volume for Land H-gain and Ocean-Glint observations. This is due to changes in both the details of the QF procedure, including changes in the variable thresholds used to assign QF=good/bad, and to some differences in the convergence characteristics of the L2FP retrieval. Generally, v9 is producing up to 60% more good-quality data than v7.3 near the end of the overlap period in 2016. There was a substantial increase in the number of good QF soundings from 2010 to 2019, due to the increased latitudinal range of the ocean observations as a result of improvements in the GOSAT pointing strategy, as well as improvements in the sounding selection for ACOS L2FP v9.



**Figure 3.** Time series histograms of the monthly number of good QF GOSAT soundings for v7.3 (top) and v9 (bottom) spanning the eleven year record. The large data gaps in early 2015 and 2019 were caused by a switch from the primary to secondary pointing mirror and a solar panel failure, respectively.



Figure 4 shows sounding density Hovmöller plots comparing ACOS GOSAT v7.3 (A) to v9 (B) with the three GOSAT observation modes combined. Again, the extended time period covered by v9 is evident. The increase in sounding density in the SH beginning in 2016 due to optimization of the GOSAT viewing strategy is prominent in the v9 product. This feature is also seen in the spatial maps showing the fraction of good quality soundings and the density per grid box, in panels C and D of Figure 1, which was introduced in Section 3.2. Persistently clear regions, such as the Sahara and western Australia, have as many as 30% of the observations assigned a good quality flag. Large regions of the tropical Pacific and Atlantic also contain a relatively high fraction of good quality soundings. On the other hand, tropical forests and high latitudes in general have low yields of good quality soundings. This is largely a combination of cloud contamination, dark surfaces at shortwave infrared wavelengths, and low solar illumination conditions, all three of which are problematic for retrieving CO<sub>2</sub> from space using reflected sunlight.




**Figure 4.** Sounding density comparing v7.3 (A) to v9 (B) good QF data as a function of time and latitude at 30 day by 15° latitude resolution for all viewing modes combined.

#### 4.2 ACOS GOSAT v9 XCO<sub>2</sub> spatiotemporal analysis

There has been a steady increase in the atmospheric burden of CO<sub>2</sub> since the onset of the industrial age due mainly to the burning of fossil fuels (e.g., Keeling et al., 1995). In May of 2009, at the beginning of the GOSAT mission, the mean global value of XCO<sub>2</sub> reported by the NOAA Global Monitoring Laboratory was 387.95 ppm, while by May of 2020, the mean global value had risen to 413.81 ppm (Dlugokencky and Tans, 2021). This yields a secular increase of  $\simeq 2.35$  ppm/yr. For comparison, Figure 5 shows the ACOS GOSAT v9 bias corrected and quality filtered XCO<sub>2</sub> as a function of latitude (15° increments) and time (30 d increments) for combined Land M and H-gain observations (A), and Ocean-Glint observations (B). Using the monthly mean XCO<sub>2</sub> values (combined land and ocean) for May 2009 (386.50 ppm) and May 2020 (411.82 ppm), the ACOS GOSAT v9 record has a secular increase of  $\simeq 2.30$  ppm/yr over the eleven year record. This small disagreement in secular trend

of approximately 2% is understandable, given the significant differences in the spatiotemporal sampling of the two data sets.

350 For the interested reader, a thorough comparison of satellite and surface-derived growth rates in atmospheric CO<sub>2</sub> is given in Buchwitz et al. (2018).



**Figure 5.** ACOS GOSAT v9 bias corrected and quality filtered XCO<sub>2</sub> as a function of latitude (15° increments) and time (30 d increments) for combined Land M-gain and H-gain observations (A), and Ocean-Glint observations (B). Grid cells with less than 10 GOSAT soundings are not colored.


The maps in Figure 6 show the spatial distribution of XCO<sub>2</sub> at 2.5° latitude by 5° longitude resolution for 2010 (top) and 2019 (bottom), for DJF (left) and JJA (right). The dynamic range of the color scale in each case is 6 ppm. However, due to the secular

355 increase in global CO<sub>2</sub> of  $\simeq 2.3$  ppm per year, the scale is centered  $\simeq 20$  ppm higher in 2019 compared to 2010. The strong latitudinal gradients in XCO<sub>2</sub> are similar in these two seasons, while the zonal gradient tends to be weakest in MAM (not shown), just before the summer draw down of CO<sub>2</sub> by the land biosphere begins. The increase in the number of Ocean-Glint soundings in the later part of the data record is also evident in these maps.

360 Qualitatively, the patterns in the maps look quite similar from 2010 to 2019, but with increased data coverage. In general, the highest concentrations of XCO<sub>2</sub> for the two selected seasons are observed by GOSAT in the northern hemisphere (NH) during DJF, especially over northern tropical Africa (between 0° and 15°N latitude), large portions of China, and the eastern United States. This stands to reason, as the atmospheric burden of CO<sub>2</sub> increases towards a peak during NH winter due to inactivity of the land biosphere, coupled with strong anthropogenic CO<sub>2</sub> emissions. During DJF the ACOS GOSAT v9 XCO<sub>2</sub> exhibits

365 relatively low concentrations across the entire SH, as would be expected if the Southern ocean were a strong carbon sink (e.g.,

Gruber et al., 2019)). In JJA, the XCO<sub>2</sub> is reduced over the mid-latitude and boreal forests, also expected behavior due to strong photosynthetic uptake of CO<sub>2</sub> during this season (e.g., Ciais et al., 2019).



**Figure 6.** ACOS GOSAT v9 bias corrected XCO<sub>2</sub> for the good QF soundings at 2.5° latitude by 5° longitude resolution for DJF 2009-2010 (A), JJA 2010 (B), DJF 2018-2019 (C), and JJA 2019 (D). The dynamic range of the color scale in each case is 6 ppm. However, due to the secular increase in global CO<sub>2</sub> of  $\simeq 2.2$  ppm per year, the scale is centered  $\simeq 20$  ppm higher in 2019 compared to 2010. Grid cells with less than 5 GOSAT soundings are not colored.

A quantification of differences in the bias corrected ACOS GOSAT v9 XCO<sub>2</sub> data product relative to the v7.3 product is given in Figure 7 for the overlapping period. The top row (panels A through C) show results for the Land H-gain observations, while the lower row (panels D through F) show results for the Ocean-Glint observations. Only soundings that were present in both data sets and assigned a good quality L2FP flag were used in this comparison. This restricts the analysis to April 2009 through June 2016, and also eliminates the v9 Land M-gain data, as no M-gain data exists in the v7.3 product. The mean and standard deviation of the  $\Delta$ XCO<sub>2</sub> for the approximately 311 k Land soundings at the single sounding level are -0.18 ppm and 0.72 ppm, respectively, as shown in Panel (A). When gridded and mapped at 2.5° latitude by 5° longitude resolution, as shown in Panel (B), the majority of the negative signal (v9 XCO<sub>2</sub> lower than v7.3) occurs at latitudes greater than approximately 45°. Most of the land mass at latitudes less than 45° have  $\Delta$ XCO<sub>2</sub> values closer to zero, with the largest positive signals appearing over equatorial forests. Furthermore, when the data is gridded and viewed versus time and latitude in 30 d by 15° increments, re-

spectively, as in Panel (C), we see that the  $\Delta\text{XCO}_2$  signal has an increasing tendency in time, i.e., the v9  $\text{XCO}_2$  increases more rapidly in time than v7.3. The cause of this effect is currently unknown, but is partially due to the implementation in v9 of a  
380 time dependent bias correction term of +0.05 ppm/yr for Land observations. This translates into an expected change of about 0.35 ppm in the v9 record over the 2009 to 2016 time span.

For the Ocean-Glint observations at the single sounding level (Panel D), the mean and standard deviation of the  $\Delta\text{XCO}_2$  are +0.28 ppm and 0.79 ppm, respectively. When gridded and mapped at  $2.5^\circ$  latitude by  $5^\circ$  longitude resolution (Panel E), the  
385 spatial distribution is fairly smooth, i.e., low variation in both latitude and longitude. Finally, when the data is gridded versus time and latitude (Panel F), the modest variation in latitude is confirmed, but a substantial time dependence is observed, with the  $\Delta\text{XCO}_2$  signal beginning negative in 2009 (v9  $\text{XCO}_2$  lower than v7.3), and switching to a positive  $\Delta\text{XCO}_2$  signal by 2016 (v9  $\text{XCO}_2$  higher than v7.3). The time dependent bias correction term for Ocean-Glint observations was +0.1 ppm/yr. This translates into an expected change of about 0.7 ppm over the 2009 to 2016 time span in the v9 record, accounting for some, but  
390 not all of this time dependent difference between v9 and v7.3.


This direct comparison between the v9 and v7.3  $\text{XCO}_2$  product only allows for statements as to their differences. It does not allow one to deduce which is closer to truth. Therefore, an analysis of the v9  $\text{XCO}_2$  data product against truth metrics follows. Furthermore, it is difficult to accurately determine the effect that the new v9  $\text{XCO}_2$  product will have on atmospheric inversion  
395 system results relative to v7.3 without further detailed study.

### 4.3 ACOS GOSAT v9 $\text{XCO}_2$ versus TCCON

A list of TCCON stations used in this work, including basic physical information and data citations, is given in Table 8. For the evaluation against the ACOS v9  $\text{XCO}_2$  data, the single sounding collocations described in Section 3.3, were aggregated into  
400 overpass mean values. Essentially the same TCCON dataset was used for both the QF/BC procedure as for the evaluation, as no hold-over data was maintained. Also, as described in Section 3.3, an averaging kernel correction was applied to the TCCON data in order to fairly compare to the satellite data. A one-to-one linear regression of the  $\text{XCO}_2$  provides a simple quantification of the agreement, as shown in Figure 8.

405 For Ocean-Glint observations (A), the mean ( $\mu$ ) of the differences in  $\text{XCO}_2$  ( $\Delta\text{XCO}_2^{\text{TCCON}} = \text{GOSAT} - \text{TCCON}$ ) is essentially zero: 0.00 ppm for the single-sounding (SS) results, and +0.01 ppm for the overpass mean (OPM) results. The corresponding standard deviations ( $\sigma$ ) are 1.08 ppm and 0.82 ppm for the SS and OPM results, respectively. This indicates that roughly half of the SS error variance is a result of instrument noise, or other random high frequency error sources ( $1.08^2=1.2$  ppm versus  $0.82^2=0.7$  ppm).


410



**Figure 7.** Analysis of the ACOS GOSAT calculated  $\Delta XCO_2$  (v9 minus v7.3) for the quality filtered and bias corrected soundings for the overlapping period spanning April 2009 through May 2016. Panels (A) through (C) show results for the Land H-gain observations, while Panels (D) through (F) show results for Ocean-Glint observations. Panels (A) and (D) shows the single sounding frequency distribution of  $\Delta XCO_2$  in 0.1 ppm bins. Panels (B) and (E) show the spatially gridded  $\Delta XCO_2$  at  $2.5^\circ$  latitude by  $5^\circ$  longitude resolution. Panels (C) and (F) show the  $\Delta XCO_2$  as a function of time and latitude in 30 d by  $15^\circ$  increments, respectively. The statistics in Panels (A) and (D) were calculated at the single sounding level, while those reported in Panels (B) and (E) were calculated on the grid box means.

**Table 8.** List of TCCON stations used in the BC/QF and XCO<sub>2</sub> evaluation of ACOS GOSAT v9, along with the data citations.

TCCON Station Name	Continent *=island	Latitude (degrees)	Altitude (meters)	Operational Date Range (YYYYMM - YYYYMM)	Data Citation
Eureka	North America	80.1 N	610	200608 – present	Strong et al. (2019)
Sodankylä	Europe	67.4 N	188	200901 – present	Kivi et al. (2014)
East Trout Lake	North America	54.4 N	502	201610 – present	Wunch et al. (2017a)
Bialystok	Europe	53.2 N	180	200903 – 201810	Deutscher et al. (2019)
Bremen	Europe	53.1 N	27	200407 – present	Notholt et al. (2019)
Karlsruhe	Europe	49.1 N	116	200909 – present	Hase et al. (2015)
Paris	Europe	48.8 N	60	201409 – present	Te et al. (2014)
Orleans	Europe	48.0 N	130	200908 – present	Warneke et al. (2019)
Garmisch	Europe	47.5 N	740	200707 – present	Sussmann and Rettinger (2018a)
Zugspitze	Europe	47.4 N	2960	201204 – present	Sussmann and Rettinger (2018b)
Park Falls	North America	45.9 N	440	200405 – present	Wennberg et al. (2017)
Rikubetsu	Asia	43.5 N	380	201311 – present	Morino et al. (2016)
Indianapolis	North America	39.9 N	270	201208 – 201212	Iraci et al. (2016b)
Lamont	North America	36.6 N	320	200807 – present	Wennberg et al. (2016b)
Four Corners	North America	36.8 N	1643	201103 – 201310	Dubey et al. (2014); Lindenmaier et al. (2014)
Anmyeondo	Asia	36.5 N	30	201408 – present	Goo et al. (2014)
Tsukuba	Asia*	36.1 N	30	200812 – present	Morino et al. (2018a)
Nicosia	Europe*	35.1 N	185	201908 – present	Petri et al. (2020)
Edwards	North America	35.0 N	699	201307 – present	Iraci et al. (2016a)
JPL	North America	34.2 N	390	201103 – 201307 201706 – 201805	Wennberg et al. (2016a) Wennberg et al. (2016a)
Caltech	North America	34.1 N	230	201209 – present	Wennberg et al. (2015)
Saga	Asia*	33.2 N	7	201106 – present	Kawakami et al. (2014)
Hefei	Asia	31.9 N	29	201509 – 201612	Liu et al. (2018)
Izana	Africa*	28.3 N	237	200705 – present	Blumenstock et al. (2017)
Burgos	Asia*	18.5 N	35	201703 – present	Morino et al. (2018b)
Ascension	Africa*	7.9 S	10	201205 – present	Feist et al. (2014)
Darwin	Australia	12.4 S	30	200508 – present	Griffith et al. (2014a)
Reunion	Africa*	20.9 S	87	201109 – present	De Mazière et al. (2017)
Wollongong	Australia	34.4 S	30	200805 – present	Griffith et al. (2014b)
Lauder	Australia*	45.0 S	370	200406 – present	Pollard et al. (2019)



**Figure 8.** Quality filtered and bias corrected ACOS GOSAT v9 vs collocated TCCON XCO<sub>2</sub> for Ocean-Glint (A), Land H-gain (B), and Land M-gain (C). Each point represents an overpass mean, which typically contain 5-10 GOSAT soundings per overpass. The legend in the lower right indicates the number of collocated overpass means for individual TCCON stations. Summary statistics for all stations combined are reported in the upper left of each panel for both single sounding and overpass means.

For Land H-gain observations (B),  $\mu=+0.10$  ppm and  $+0.14$  ppm for the SS and OPM, respectively. The Land H-gain  $\sigma$  are higher than for Ocean-Glint;  $1.60$  and  $1.14$  ppm for SS and OPM, respectively. Larger variations in  $\Delta XCO_2^{TCCON}$  are expected for Land H-gain due to variability in topography and surface brightness, as well as higher likelihood of contamination by cloud and aerosol, all of which are more challenging for the ACOS retrieval. Further, biology and atmospheric transport cause  $CO_2$  signals to vary more over land regions, and in addition, instrument noise is higher because the SNRs tend to be lower.


Land M-gain observations have near zero bias ( $\mu=-0.02$  ppm and  $+0.02$  ppm for SS and OPM, respectively), and scatter similar to that for Ocean-Glint ( $\sigma=1.09$  and  $0.84$  ppm for SS and OPM, respectively), likely driven by lower variability in surface topography and brightness compared to Land H-gain observations, as well as higher SNRs over these bright land surfaces.

The correlation in the  $XCO_2$  between the datasets in all observation modes is high, with Pearson  $R^2=0.98$ ,  $0.98$ , and  $0.99$  for Ocean-Glint, Land H-gain, and Land M-gain, respectively. Overall, these results indicate excellent agreement between the bias corrected and quality filtered ACOS GOSAT v9  $XCO_2$  product and collocated estimates from TCCON.

Figure 9 shows the mean absolute error (MAE) between the overpass mean collocated GOSAT and TCCON  $XCO_2$ , organized by latitude bins, season, and observation mode for v7.3 (Panels A and B) and v9 (Panels C, D, E). The calculation of the MAE and error bars follow the procedure reported in Chatterjee et al. (2013) (Equations 3 and 4). The error bars on the MAE represent the scatter around the mean. A smaller error bar, or a lower scatter, implies that the MAE values are more consistent across a group of TCCON stations within a latitude band and season. The MAEs tend to be lower for v9 compared to v7.3, with smaller error bars, and increased number of collocations. This is especially true for the SH Ocean-Glint data, where the MAE ranges from  $0.4$  to  $0.7$  ppm in v9 for all seasons, in contrast to v7.3, which had higher MAE ranging from  $0.5$  to  $0.85$  ppm in that region. In the v9 Land H-gain data, the MAE is roughly a function of latitude, with the highest values ( $\simeq 1.0$  ppm) seen between  $60^\circ N - 90^\circ N$ , and the lowest values ( $\simeq 0.7$  ppm) seen from  $30^\circ S - 60^\circ S$ . This stands to reason as lower variability of  $XCO_2$  in the SH tends to yield better agreement between satellite and ground based observations. The error bars on the v9 Land H-gain estimates in the  $30^\circ N - 60^\circ N$  latitude range are very small, due in part to the large number of collocations. There is very limited land data between  $0^\circ$  and  $30^\circ N$  (approximately 25% of Earth's surface), due to the sparsity of TCCON stations in this latitude band. Only Burgos ( $18.5^\circ N$ ) and Izana ( $28.3^\circ N$ ) are located in this range (reference Table 8), and many of the collocations from these sites are from GOSAT observations made in Ocean-Glint viewing.

Knowledge of the average  $XCO_2$  seasonal cycle can be used to disentangle the  $CO_2$  growth rate from the seasonal variability, as well as for quantifying potential seasonal biases between satellite and ground-based  $XCO_2$  estimates. Lindqvist et al. (2015) fitted a skewed sine wave (See Eq. 1 of Lindqvist et al., 2015) to the ACOS GOSAT v3.5  $XCO_2$  time series and the TCCON estimates of  $XCO_2$  at 16 stations, spanning April 2009 through December 2013. They found that ACOS GOSAT v7.3 captured the seasonal cycle within approximately 1 ppm of the TCCON estimates for all but the European sites, and that the satellite and ground-based  $CO_2$  growth rates agreed generally better than  $0.2$  ppm per year. Here, we provide an update to those results using






**Figure 9.** Mean Absolute Error (MAE; ppm) between ACOS GOSAT v9 XCO<sub>2</sub> and collocated TCCON observations for v7.3 (panels A and B) and v9 (panels C, D and E), binned by latitude for: Ocean-Glint (panels A and C), Land H-gain (panels B and D) and Land M-gain (panel E). Recall that there was no Land M-gain data generated for v7.3. The error bars on the MAE represent the standard error of the mean. The calculation of the MAE statistic and error bars follow the procedure reported in Chatterjee et al. (2013) (Equations 3 and 4). The total number of independent observations available within each latitude band and for each season are reported in the colored boxes. Latitude bands that do not have any collocated soundings are shaded in grey.

the eleven year ACOS GOSAT v9 XCO<sub>2</sub> data record. For this part of the analysis, a slightly more restrictive set of collocation criteria were implemented, compared to that described in Section 3.3 for the BC/QF procedure and to that used to generate Fig. 8. The seasonal cycle analysis required that the TCCON record spanned at least one contiguous year (a full seasonal cycle), and that a minimum of 20 collocations with GOSAT occurred. In addition, the three GOSAT observation modes (Ocean-Glint, Land M-gain, Land H-gain) were combined for each site, and satellite overpass means of XCO<sub>2</sub> were aggregated into daily means. This resulted in approximately 7700 daily averages at 26 TCCON stations over the 11 year GOSAT data record.

Figure 10 shows the results of the seasonal cycle fit for the Lamont, Oklahoma TCCON station. The one-to-one scatter of the 896 daily averaged XCO<sub>2</sub> values (A) indicates a bias of -0.27 ppm for the GOSAT product relative to TCCON, with a standard deviation of 1.25 ppm and a Pearson's R<sup>2</sup> of 0.99. The seasonal cycle fits (B) indicate excellent agreement in the

secular  $\text{CO}_2$  increase at this site; 2.34 ppm for both GOSAT and TCCON. The mean seasonal amplitudes indicate a slight disagreement of a few tenths of a ppm, with TCCON showing a slightly higher fitted peak  $\text{XCO}_2$  value during the spring maximum phase, compared to GOSAT. This is similar to the results for this site reported in (Fig. 4 of Lindqvist et al., 2015). The time series of the calculated difference in satellite and ground-based estimated  $\text{XCO}_2$  (GOSAT - TCCON), shown in (C), highlights the magnitude of the scatter about the mean bias, and suggests that there is no observable time-drift in the data at this site.



**Figure 10.** Seasonal cycle analysis of ACOS GOSAT v9  $\text{XCO}_2$  versus collocated TCCON for the Lamont, Oklahoma station, following the methodology of Lindqvist et al. (2015). Panel (A) shows the one-to-one scatter plot of the bias corrected daily mean  $\text{XCO}_2$  for GOSAT versus TCCON. Panel (B) shows the time series of GOSAT  $\text{XCO}_2$  (blue circles) with fit (blue line) and the TCCON  $\text{XCO}_2$  (grey triangles) with fit (orange line) over the eleven year data record. Panel (C) shows the time series of calculated  $\Delta\text{XCO}_2^{\text{TCCON}}$  (GOSAT - TCCON), with the mean difference (horizontal solid black line) and  $\pm 1$  standard deviation (gray shading). The three GOSAT observation modes have been combined into daily mean averages to provide the maximum number of collocations possible for the seasonal fits.

A summary of the data from each station that met the seasonal cycle collocation criteria is provided in Table 9. In addition, the full complement of plots are presented in Appendix A. Overall, the seasonal cycle analysis at most sites is in agreement, to within the estimated uncertainties. The standard deviation of the mean  $\text{XCO}_2$  biases for the 26 sites is 0.41 ppm for the ACOS GOSAT v9 record. This compares to a value of 0.51 ppm at 23 stations for ACOS GOSAT v7.3, suggesting an improvement in the quality of the v9  $\text{XCO}_2$  product.


#### 4.4 ACOS GOSAT v9 $\text{XCO}_2$ versus models

The collocation and calculation of the multi-model-mean (MMM) was described in Section 3.3. Although the model data used for evaluation was very similar to that used in the QF/BC procedure, some minor version updates and extensions in time were included, as indicated in Table 4. It is important to be aware that there can be a considerable delay between performing the QF/BC procedure and the full generation of the final product, during which time the models are often updated.

**Table 9.** Evaluation of the daily mean bias corrected ACOS GOSAT v9 XCO<sub>2</sub> (all viewing modes combined) against collocated TCCON estimates for individual stations. There were 7547 days total for the 25 stations. The following sites/instruments were excluded from this part of the analysis due to inadequate timeseries or seasonal cycle coverage; Eureka, Four Corners, Indianapolis (Influx), JPL2007, Lauder1, Lauder3, Manaus, and Ny Ålesund. The mean, standard deviation, and Pearson correlation coefficient ( $\mu$ ,  $\sigma$ ,  $R^2$ ) of the linear fit between GOSAT and TCCON are given in columns 3-5. The remaining columns quantify the seasonal cycle fit following the methodology described in Lindqvist et al. (2015). The bottom row provides mean summary statistics for the linear fit.

TCCON	N	$\Delta XCO_2$		Linear	Trend $\pm$ uncertainty		Amplitude $\pm$ uncertainty	
Station	(days)	(GOSAT - TCCON)		Correlation	(ppm/yr)		(ppm)	
Name		$\mu$ (ppm)	$\sigma$ (ppm)	( $R^2$ )	TCCON	GOSAT	TCCON	GOSAT
Sodankylä	166	0.91	1.57	0.98	2.32 $\pm$ 0.02	2.37 $\pm$ 0.04	9.83 $\pm$ 0.95	10.30 $\pm$ 2.76
East Trout Lake	86	0.69	1.52	0.94	2.19 $\pm$ 0.11	1.98 $\pm$ 0.23	10.30 $\pm$ 0.24	9.29 $\pm$ 0.54
Bialystok	252	0.01	1.26	0.98	2.29 $\pm$ 0.02	2.38 $\pm$ 0.04	8.97 $\pm$ 0.13	8.67 $\pm$ 0.26
Bremen	95	0.32	2.01	0.96	2.28 $\pm$ 0.04	2.42 $\pm$ 0.08	8.14 $\pm$ 0.27	8.48 $\pm$ 1.79
Karlsruhe	328	0.61	1.60	0.98	2.34 $\pm$ 0.02	2.43 $\pm$ 0.03	7.72 $\pm$ 0.14	7.19 $\pm$ 0.39
Paris	137	-0.28	1.95	0.92	2.10 $\pm$ 0.07	2.65 $\pm$ 0.10	7.89 $\pm$ 0.31	6.68 $\pm$ 0.42
Órleans	300	0.26	1.57	0.98	2.32 $\pm$ 0.02	2.41 $\pm$ 0.03	7.87 $\pm$ 0.13	6.96 $\pm$ 0.26
Garmisch	269	0.57	1.66	0.98	2.30 $\pm$ 0.02	2.28 $\pm$ 0.03	7.64 $\pm$ 0.16	7.52 $\pm$ 0.35
Zugspitze	107	-0.09	2.10	0.92	2.52 $\pm$ 0.04	2.66 $\pm$ 0.12	6.38 $\pm$ 0.23	8.27 $\pm$ 0.65
Park Falls	389	-0.12	1.35	0.98	2.32 $\pm$ 0.01	2.30 $\pm$ 0.03	8.94 $\pm$ 0.14	9.27 $\pm$ 0.27
Rikubetsu	58	-0.34	1.48	0.98	2.45 $\pm$ 0.07	2.85 $\pm$ 0.13	10.62 $\pm$ 0.41	12.19 $\pm$ 0.75
Lamont	896	-0.27	1.25	0.99	2.34 $\pm$ 0.01	2.34 $\pm$ 0.02	5.94 $\pm$ 0.09	5.66 $\pm$ 0.14
Anmyeondo	24	0.72	1.62	0.94	2.64 $\pm$ 0.22	2.93 $\pm$ 0.27	8.78 $\pm$ 0.68	10.25 $\pm$ 1.06
Tsukuba	389	0.75	1.71	0.97	2.54 $\pm$ 0.03	2.43 $\pm$ 0.04	6.84 $\pm$ 0.20	7.44 $\pm$ 0.30
Edwards	543	0.38	0.91	0.98	2.45 $\pm$ 0.01	2.46 $\pm$ 0.02	5.51 $\pm$ 0.08	5.63 $\pm$ 0.14
JPL	361	-0.12	1.31	0.98	2.44 $\pm$ 0.02	2.43 $\pm$ 0.03	5.30 $\pm$ 0.12	6.12 $\pm$ 0.22
Caltech	852	0.71	1.26	0.97	2.44 $\pm$ 0.02	2.49 $\pm$ 0.02	5.74 $\pm$ 0.10	5.85 $\pm$ 0.14
Saga	281	-0.03	1.50	0.97	2.31 $\pm$ 0.02	2.39 $\pm$ 0.04	6.59 $\pm$ 0.14	7.21 $\pm$ 0.27
Hefei	38	-0.22	1.77	0.78	3.22 $\pm$ 0.49	2.79 $\pm$ 0.72	6.60 $\pm$ 0.87	5.64 $\pm$ 0.88
Izana	180	-0.08	1.03	0.99	2.40 $\pm$ 0.01	2.26 $\pm$ 0.02	5.70 $\pm$ 0.11	5.76 $\pm$ 0.18
Burgos	80	-0.43	1.09	0.91	2.21 $\pm$ 0.07	2.38 $\pm$ 0.16	5.86 $\pm$ 0.23	5.31 $\pm$ 0.46
Ascension	310	0.49	0.73	0.98	2.37 $\pm$ 0.01	2.30 $\pm$ 0.00	0.32 $\pm$ 0.10	0.62 $\pm$ 0.10
Darwin	565	0.04	1.20	0.98	2.39 $\pm$ 0.01	2.32 $\pm$ 0.02	0.44 $\pm$ 0.09	3.20 $\pm$ 0.20
Reunion	309	0.11	0.84	0.99	2.38 $\pm$ 0.02	2.39 $\pm$ 0.02	1.57 $\pm$ 0.15	1.25 $\pm$ 0.13
Wollongong	532	-0.17	1.11	0.99	2.39 $\pm$ 0.01	2.36 $\pm$ 0.02	1.03 $\pm$ 0.09	1.34 $\pm$ 0.18
Lauder 2	194	-0.31	1.64	0.96	2.35 $\pm$ 0.01	2.23 $\pm$ 0.02	0.39 $\pm$ 0.10	0.30 $\pm$ 0.36
Mean	302	0.18	1.42	0.96				

Seasonal maps of  $\Delta\text{XCO}_2^{\text{MMM}}$  (GOSAT v9 minus MMM) are shown in Figure 11 for the eleven year data record binned at  $2.5^\circ$  latitude by  $5^\circ$  longitude. Generally, the agreement between the model derived values and the satellite estimates is quite good, with an annual mean difference of  $\simeq -0.15$  ppm, and binned scatter  $\simeq 0.5$  ppm. For Ocean-Glint observations, the  $\Delta\text{XCO}_2^{\text{MMM}}$  tends to be negative (positive) in the northern (southern) hemispheres. Land observations exhibit several distinct sub-continental scale disagreements, including a strong positive signal over northern tropical Africa in DJF (GOSAT  $\text{XCO}_2$  higher than MMM).




**Figure 11.** Seasonal maps of the mean  $\Delta\text{XCO}_2^{\text{MMM}}$  (GOSAT - MMM) spanning 2009 through 2018 for DJF (A), MAM (B), JJA (C), and SON (D) at  $2.5^\circ$  latitude by  $5^\circ$  longitude resolution. Grid boxes containing less than 10 collocations are colored white.

Figure 12 shows Hovmöller plots of the  $\Delta\text{XCO}_2^{\text{MMM}}$  Ocean-Glint data at 30 day by  $15^\circ$  latitude resolution for v7.3 (A) and v9 (B). The extension in time of v9 is evident, as well as the expansion in the latitude range of the Ocean-Glint observations since 2015. A direct comparison between the v7.3 and v9  $\Delta\text{XCO}_2^{\text{MMM}}$  values for the overlapping time period, April 2009 through June 2016, reveals a global mean bias and standard deviation of  $-0.54$  ppm and  $1.0$  ppm for the v7.3 product, and  $-0.20$  ppm and  $0.84$  ppm for v9, underscoring the improvement.

Of particular note are the strong positive  $\Delta\text{XCO}_2^{\text{MMM}}$  values in the v9 SH Ocean-Glint observations for the latter part of 2014, persisting through most of 2015. This feature is not seen in the v7.3 product, due to a paucity of SH Ocean-Glint data. It approximately coincides with the strong 2015-2016 El Niño event, where  $\Delta\text{XCO}_2^{\text{MMM}}$  signals were also seen in the OCO-2

v7 Ocean-Glint data, as reported in Chatterjee et al. (2017). It has been hypothesized that the 2015-2016 El Niño produced an anomalously strong carbon release from tropical land regions due to higher temperature and below average precipitation (Liu et al., 2017). In contrast to the positive SH signal, negative  $\Delta\text{XCO}_2^{\text{MMM}}$  values (GOSAT lower than MMM) are observed in the v9 NH oceans since 2016. It is unclear why the satellite and models disagree over such large spatial and temporal scales, but recent work by Müller et al. (2021) suggests that the ACOS v7.3 (and to a lesser extent v9)  $\text{XCO}_2$  are in fact biased low by approximately 1 to 1.5 ppm, as compared to a new independent evaluation data set generated from combined ship and aircraft measurements over the open oceans. Further investigation into the source of the ACOS GOSAT biases against models is warranted.




**Figure 12.** Time series of  $\Delta\text{XCO}_2^{\text{MMM}}$  (ACOS GOSAT v9 - MMM) versus latitude at 30 day by  $15^\circ$  resolution for Ocean-Glint observations for v7.3 (A) and v9 (B). Grid cells containing less than 10 collocations are colored white.

Figure 13 shows spatial maps of  $\Delta\text{XCO}_2^{\text{MMM}}$  for the truncated time span 2010 through 2015 comparing v7.3 (A and C) and v9 (B and D) for DJF (A and B) and JJA (C and D). There is significant decrease in scatter in the v9 product ( $\approx 0.5$  ppm) relative to v7.3 ( $\approx 0.75$  ppm). The low bias in DJF tropical Pacific vanishes in v9 and the positive bias in JJA extra-tropical land regions is reduced. The expanded latitudinal extent of the Ocean-Glint observations is evident in the v9 maps. One feature that is robust in both v7.3 and v9 is the large positive signal over northern tropical Africa in DJF. This feature was also observed in the OCO-2 v7 and v8 comparisons to a MMM (O'Dell et al., 2018), and in v10  $\text{XCO}_2$  anomaly maps (Hakkarainen et al., 2019).

#### 4.5 ACOS GOSAT v9 $\text{XCO}_2$ versus OCO-2

NASA's Orbiting Carbon Observatory-2 (OCO-2) has been collecting science data since September, 2014 from a near-polar low-Earth orbit (705 km altitude), with an afternoon equator crossing time of  $\approx 1:30$  PM (Crisp et al., 2017). Like GOSAT,



**Figure 13.** Maps of the mean  $\Delta XCO_2^{MMM}$  (GOSAT - MMM) spanning 2010 through 2015 for v7.3 DJF (A), v9 DJF (B), v7.3 JJA (C), and v9 JJA (D) at 2.5° by 5° latitude/longitude resolution. Only grid boxes with at least 10 collocations are shown.

OCO-2 takes measurements of reflected solar radiation in the Oxygen A-band (0.76  $\mu m$ ), and the weak and strong carbon dioxide bands (1.6 and 2.0  $\mu m$ , respectively), which are used to estimate XCO<sub>2</sub> using the ACOS L2FP retrieval algorithm (Eldering et al., 2017; O'Dell et al., 2018). However, due to differences in the orbit parameters of the two sensors, e.g., a 3 day repeat cycle for GOSAT versus a 16 day repeat cycle for OCO-2 (see Table 2 of Kataoka et al., 2017), the number of collocated soundings is somewhat limited. Therefore, some criteria must be defined in order to identify soundings that can be compared in a meaningful way. The underlying assumption of the collocation is that on scales of a few hundred kilometers and several hours, the natural variance in XCO<sub>2</sub> is not detectable in satellite derived estimates from the ACOS L2FP algorithm.

515

For this study, the coincidence criteria to match OCO-2 soundings to individual GOSAT soundings were those: (i) falling within 2° latitude and 3° longitude, (ii) with a maximum spatial separation of 300 km, and (iii) acquired within  $\pm 2$  hours. Due to the dense nature of the OCO-2 soundings relative to the sparseness of the GOSAT soundings, there are typically between zero and several hundred matched OCO-2 soundings per GOSAT footprint. A lower limit of 10 and an upper limit of 100 (randomly selected) OCO-2 soundings that meet the coincidence criteria were set in order to retain the GOSAT sounding for analysis. The individual L2FP quality flags are applied for both GOSAT and OCO-2 during the collocation procedure, and then the mean value of XCO<sub>2</sub> from the 10 to 100 collocated OCO-2 soundings is calculated and subtracted from the corresponding

520

GOSAT XCO<sub>2</sub> to produce  $\Delta\text{XCO}_2^{\text{OCO-2}}$ .

525 Here we compare ACOS GOSAT v9 against OCO-2 v10 (rather than to the deprecated v9), since we assume that science users will adopt the newest OCO-2 product. Major updates to the version 10 ACOS L2FP algorithm (Osterman et al., 2020), were discussed in Section 3.1.

One complexity in comparing ACOS GOSAT v9 and OCO-2 v10 is the fact that the two versions of the algorithm used  
530 different CO<sub>2</sub> priors. Typically, models which assimilate satellite CO<sub>2</sub> data take into account the unmeasured part of the prior CO<sub>2</sub> profile (specified via the retrieval’s averaging kernel) via an averaging kernel correction, as given in Eq. 1. Therefore, in order to fairly compare these two data sets as models would assimilate them, we need to remove their difference due to the unmeasured part of the CO<sub>2</sub> profile, as follows:


$$\text{XCO}_2' = \text{XCO}_2 + \sum_{i=1}^{20} h_i(1 - a_i) \cdot (u'_{a,i} - u_{a,i}), \quad (3)$$

535 where  $h$  is the XCO<sub>2</sub> pressure weighting function,  $a$  is the normalized XCO<sub>2</sub> averaging kernel,  $u_a$  is the ACOS v9 CO<sub>2</sub> prior profile used for GOSAT, and  $u'_a$  is the ACOS v10 CO<sub>2</sub> prior profile used for OCO-2. The summation takes place over the 20 vertical levels defined in the ACOS code. In summary, the total adjustment to the ACOS GOSAT XCO<sub>2</sub> value is calculated as the contribution of the difference in the vertical CO<sub>2</sub> priors at each level weighted by the one minus the averaging kernel at that level. The global mean adjustment due to the CO<sub>2</sub> prior correction was approximately 0.2 ppm, with 95% of corrections  
540 between -0.1 and +0.5 ppm.

Spatial maps of  $\Delta\text{XCO}_2^{\text{OCO-2}}$  (GOSAT v9 - OCO-2 v10) for the prior-corrected, collocated soundings are shown in Figures 14 and 15 for land and ocean, respectively. In each figure the maps are shown by season at 2.5° latitude by 5° longitude resolution. In all seasons, higher scatter in  $\Delta\text{XCO}_2^{\text{OCO-2}}$  is observed over land ( $\simeq 1$  ppm) than over ocean ( $< 0.7$  ppm), likely due  
545 to variability of land surface features and/or lower signal-to-noise ratios of the radiance measurements. The annual global mean  $\Delta\text{XCO}_2^{\text{OCO-2}}$  for land is near zero (0.06 ppm), and exhibits little variation with season. For Ocean-Glint, the global mean  $\Delta\text{XCO}_2^{\text{OCO-2}}$  is larger (-0.40 ppm) and varies more significantly by season from -0.2 (DJF) to -0.6 ppm (JJA). The disagreements in the Ocean-Glint data tend to be spatially coherent, with a notably large negative difference in the NH in most seasons. Currently, the underlying cause of these disagreements is unknown, and could stem from instrument calibration or sampling  
550 related issues, differences in retrieval algorithm versions, or even collocation issues.

The disagreement in XCO<sub>2</sub> for Ocean-Glint between ACOS GOSAT v9 and OCO-2 v10 is highlighted in Panel A of Figure 16, which shows  $\Delta\text{XCO}_2^{\text{OCO-2}}$  for the period September 2014 through December 2020 at 30 day by 15° latitude resolution for the Ocean-Glint observations. Panel B shows the number of collocated soundings in each bin. A large SH positive difference in  
555  $\Delta\text{XCO}_2^{\text{OCO-2}}$  (GOSAT higher than OCO-2 by  $\simeq 0.5$  ppm) is observed for the first two years of the time record. Then, in early





**Figure 14.** Spatial distribution of the bias corrected  $\Delta\text{XCO}_2^{\text{OCO-2}}$  (GOSAT v9 minus OCO-2 v10) for the good QF land soundings for DJF (A), MAM (B), JJA (C), and SON (D) for the overlapping period August 2014 through June 2020. The spatio-temporal requirements for matched soundings are a maximum separation of 300 km, and observation time within  $\pm 2$  hours. Maps are gridded at  $2.5^\circ$  latitude by  $5^\circ$  longitude resolution, and only grid boxes with at least 10 matched soundings are shown.


2016, there is what appears to be an abrupt jump to a large negative difference (GOSAT lower than OCO-2 by  $\simeq -0.5$  ppm) in the NH. From this point forward,  $\Delta\text{XCO}_2^{\text{OCO-2}}$  appears to be reasonably stable in time, although there is a persistent low difference in the NH for the remainder of the record.

560 Figure 17 shows the  $\Delta\text{XCO}_2^{\text{OCO-2}}$  data for the combined land H-gain and M-gain data, similar to Figure 16. The main feature here is that the overall variability is larger compared to the Ocean-Glint data, which we attribute to biases introduced by variations in both topography and surface albedo. A slightly positive (red) signal is observed during the September to December months in the SH, especially in 2014, 2018, and 2019. Although, additional investigation into such signals is warranted, it is beyond the scope of the current work.

565

A set of summary statistics for the ACOS GOSAT v9 versus OCO-2 v10  $\text{XCO}_2$  product is given in Table 10. The values reported here are on the individual collocations by year and season, rather than the spatially gridded averages as given in






**Figure 15.** Same as Figure 14, but for Ocean-Glint observations.

Figures 14 and 15. For the land observations, there has been a very slight upward trend in time of the  $\Delta XCO_2^{OCO-2}$  to slightly more positive values (GOSAT v9 larger than OCO-2 v10  $XCO_2$ ). On the other hand, for Ocean-Glint observations, the general trend has been an increasingly more negative  $\Delta XCO_2^{OCO-2}$  in time, as was seen in Figure 16. Additional investigation will be required to determine the root cause(s) of these differences.


## 5 Summary

The v9 ACOS GOSAT  $XCO_2$  product, spanning February 2009 through June 2020, has been compared to  $XCO_2$  estimates from TCCON, a suite of atmospheric inversion systems (models), and with collocated OCO-2 v10 data. The ACOS GOSAT v9 product is an improvement over ACOS GOSAT v7.3 relative to these standards. The v9 product provides a significant extension of the data record and contains data in M-gain viewing mode over bright land surfaces.

Of the 37.4 M estimates of  $XCO_2$  contained in the ACOS GOSAT v9 data record, approximately 80% were prefiltered due to contamination by cloud and/or aerosol, or due to insufficient SNR. Of the 7.0 M that were selected to run through the ACOS L2FP algorithm, approximately 6.1 M returned valid estimates of  $XCO_2$ . However, only  $\simeq 2$  M of those were identified as be-



**Figure 16.** Difference in  $XCO_2$  between ACOS GOSAT v9 and OCO-2 v10 ( $\Delta XCO_2^{OCO-2}$ ) as a function of time and latitude at 30 day by  $15^\circ$  latitude resolution for Ocean-Glint observations (panel A). Panel (B) shows the sounding density of the collocated soundings. Grid cells containing less than 10 collocations are colored white.



**Figure 17.** Same as Figure 16, but for land observations (combined H-gain and M-gain).

**Table 10.** A set of summary statistics for the comparison of the ACOS GOSAT v9 XCO<sub>2</sub> to the OCO-2 v10 product. Individual collocations for each year and season are given by N, while the mean  $\Delta$ XCO<sub>2</sub> and the standard deviation from the mean are given by  $\mu$  and  $\sigma$ , respectively, both in units ppm. The top portion of the table is for land observations, while the bottom is for Ocean-Glint (OceanH).

Land	DJF			MAM			JJA			SON		
Year	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$
2014	0	–	–	0	–	–	0	–	–	4564	0.11	1.37
2015	1963	0.02	1.36	3160	0.15	1.47	5631	-0.08	1.34	4108	-0.01	1.39
2016	4379	0.01	1.28	3672	-0.07	1.42	4701	-0.04	1.41	4923	0.02	1.43
2017	3610	0.10	1.40	4097	-0.04	1.37	3450	0.02	1.34	2892	-0.02	1.41
2018	3605	0.11	1.41	3904	-0.06	1.33	4738	0.06	1.36	4218	0.21	1.46
2019	2779	0.16	1.39	3917	0.06	1.41	4911	0.09	1.35	4823	0.24	1.40
2020	3422	0.17	1.37	3833	-0.04	1.34	1183	-0.03	1.22	0	–	–
OceanH	DJF			MAM			JJA			SON		
Year	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$	N	$\mu$	$\sigma$
2014	0	–	–	0	–	–	0	–	–	2603	0.10	0.76
2015	2373	0.29	0.74	3139	0.14	0.84	2982	-0.13	0.87	3103	0.14	0.81
2016	6029	0.07	0.80	5460	-0.45	0.90	5109	-0.66	0.89	7673	-0.26	0.85
2017	6847	-0.13	0.82	6235	-0.39	0.87	3523	-0.55	0.94	4332	-0.18	0.85
2018	5888	-0.05	0.86	5680	-0.47	0.89	2767	-0.65	0.90	5887	-0.32	0.85
2019	4815	-0.14	0.91	4887	-0.50	0.93	4511	-0.61	0.92	6972	-0.37	0.95
2020	5451	-0.34	0.94	4812	-0.60	1.01	385	-0.69	1.08	0	–	–

ing of “good” quality. This represents 5.4% of the total recorded soundings. The quality filtering and bias correction variables used for ACOS GOSAT v9 were similar to those used in previous product versions, and similar to those used for OCO-2 v9 and v10, but include for the first time, a correction to account for a small temporal drift in the data.

585 Comparisons with collocated estimates of XCO<sub>2</sub> from TCCON indicate overpass mean biases of  $\simeq$  0.1 to 0.2 ppm, and standard  
deviations of  $\simeq$  1.5 ppm. The mean squared error against TCCON is highest for land observations in the northern mid-latitudes  
(30 – 60°N), and lowest for Ocean-Glint and SH land M-gain observations. The statistics show improvement when compared  
to the results for v7.3, which spanned a shorter time period (April 2009 to June 2016). Specifically, the standard deviation of  
the mean station bias for the 26 sites is 0.41 ppm for the ACOS GOSAT v9 record, compared to 0.51 ppm at 23 stations for  
590 ACOS GOSAT v7.3.

Comparisons with collocated XCO<sub>2</sub> derived from a suite of 4 atmospheric inversion systems (models) suggests annual global  
mean differences of  $\simeq$  0.15 ppm and standard deviation of  $\simeq$  0.5 ppm. Hemispherical differences in XCO<sub>2</sub> estimates over

oceans were observed, as well as robust subcontinental scale land features. Results indicate better agreement with models  
595 in the ACOS v9 product ( $\mu=-0.20$  ppm,  $\sigma=0.8$  ppm) compared to v7.3 ( $\mu=-0.54$  ppm,  $\sigma=1.0$  ppm) for the overlapping period  
April 2009 through June 2016, but further investigation is required to explain the remaining disagreement over large spatial  
and temporal scales.

Comparisons with collocated OCO-2 v10 XCO<sub>2</sub> data show low bias but relatively high scatter for land observations ( $\mu=0.06$  ppm,  
600  $\sigma=1.0$  ppm, when averaged across seasons). Increased scatter over land is expected due to XCO<sub>2</sub> bias introduced by variability  
in topography and surface albedo. However, for Ocean-Glint observations, although the XCO<sub>2</sub> scatter is lower than that for  
land as expected ( $\sigma=0.7$  ppm), the global mean bias is relatively high ( $\mu=-0.4$  ppm, when averaged across seasons). These are  
issues that must be resolved in order for GOSAT v9 and OCO-2 v10 data to provide consistent information to atmospheric  
inversion systems for assessing fluxes of CO<sub>2</sub>.

605

Global estimates of CO<sub>2</sub> derived from satellite measurements provide coverage in traditionally data sparse regions where  
ground-based measurements are difficult. The assimilation of satellite XCO<sub>2</sub> into atmospheric inversion systems to quantify  
the spatiotemporal variations of carbon fluxes is a promising, but challenging, area of research. This research continues to  
benefit from various improvements in transport models, atmospheric inversion systems, and satellite retrievals. The role of the  
610 GOSAT record in this field remains unique due to its exceptional 11 year length and its coverage of nearly 5.5 years of the  
carbon cycle prior to the launch of OCO-2. The ACOS GOSAT v9 L2Std and L2Lite file products are both available on the  
NASA GES DISC (OCO-2 Science Team et al., 2019b, a).

## References

- 615 Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P., Langenfelds, R., Torn, M., Biraud, S., Stephens, B., Andrews, A., and Worthy, D.: Global CO<sub>2</sub> fluxes estimated from GOSAT retrievals of total column CO<sub>2</sub>, *Atmospheric Chemistry and Physics*, 13, 8695–8717, <https://doi.org/10.5194/acp-13-8695-2013>, <https://acp.copernicus.org/articles/13/8695/2013/>, 2013.
- Blumenstock, T., Hase, F., Schneider, M., Garcia, O. E., and Sepulveda, E.: TCCON data from Izana (ES), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.IZANA01.R1>, 2017.
- 620 Buchwitz, M., Reuter, M., Schneising, O., Noël, S., Gier, B., Bovensmann, H., Burrows, J. P., Boesch, H., Anand, J., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Kuze, A., Suto, H., Yoshida, Y., Crisp, D., and O'Dell, C.: Computation and analysis of atmospheric carbon dioxide annual mean growth rates from satellite observations during 2003–2016, *Atmospheric Chemistry and Physics*, 18, 17 355–17 370, <https://doi.org/10.5194/acp-18-17355-2018>, <https://acp.copernicus.org/articles/18/17355/2018/>, 2018.
- Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J., Tran, H., Kuze, A., Aleks, G. K., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO<sub>2</sub> and CH<sub>4</sub> observations from GOSAT, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL047888>, 2011.
- 625 Byrne, B., Liu, J., Lee, M., Baker, I., Bowman, K. W., Deutscher, N. M., Feist, D. G., Griffith, D. W. T., Iraci, L. T., Kiel, M., Kimball, J. S., Miller, C. E., Morino, I., Parazoo, N. C., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Velazco, V. A., Wennberg, P. O., and Wunch, D.: Improved constraints on northern extratropical CO<sub>2</sub> fluxes obtained by combining surface-based and space-based atmospheric CO<sub>2</sub> measurements, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032 029, <https://doi.org/10.1029/2019JD032029>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD032029>, 2020.
- CAMS: Copernicus Atmosphere Monitoring Service, [https://atmosphere.copernicus.eu/sites/default/files/2019-08/CAMS73\\_2018SC1\\_D73.1.4.1-2018-v1\\_201907\\_v1.pdf](https://atmosphere.copernicus.eu/sites/default/files/2019-08/CAMS73_2018SC1_D73.1.4.1-2018-v1_201907_v1.pdf), 2021.
- CarbonTracker: CarbonTracker, <https://carbontracker.noaa.gov>, 2021.
- 635 CarboScope: CarboScope, <http://www.bgc-jena.mpg.de/CarboScope>, 2021.
- Chatterjee, A., Engelen, R. J., Kawa, S. R., Sweeney, C., and Michalak, A. M.: Background error covariance estimation for atmospheric CO<sub>2</sub> data assimilation, *J. Geophys. Res. Atmos.*, 118, <https://doi.org/10.1002/jgrd.50654>, 2013.
- Chatterjee, A., Gierach, M. M., Sutton, A. J., Feely, R. A., Crisp, D., Eldering, A., Gunson, M. R., O'Dell, C. W., Stephens, B. B., and Schimel, D. S.: Influence of El Niño on atmospheric CO<sub>2</sub> over the tropical Pacific Ocean: Findings from NASA's OCO-2 mission, *Science*, 358, <https://doi.org/10.1126/science.aam5776>, <http://science.sciencemag.org/content/358/6360/eaam5776>, 2017.
- 640 Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morguí, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO<sub>2</sub> surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2010JD013887>, 2010.
- Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., and Bousquet, P.: Toward robust and consistent regional CO<sub>2</sub> flux estimates from in situ and spaceborne measurements of atmospheric CO<sub>2</sub>, *Geophysical Research Letters*, 41, 1065–1070, <https://doi.org/https://doi.org/10.1002/2013GL058772>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013GL058772>, 2014.

- Ciais, P., Tan, J., Wang, X., Roedenbeck, C., Chevallier, F., Piao, S.-L., Moriarty, R., Broquet, G., LeQuéré, C., Canadell, J., Peng, S., Poulter, B., Liu, Z., and Tans, P.: Five decades of northern land carbon uptake revealed by the interhemispheric CO<sub>2</sub> gradient, *Nature*, 568, 221–225, <https://doi.org/https://doi.org/10.1038/s41586-019-1078-6>, 2019.
- Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J.-F. L., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunch, D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/https://doi.org/10.1029/2012JD018087>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012JD018087>, 2012.
- Crisp, D., Fisher, B. M., O'Dell, C., Frankenberg, C., Basilio, R., Bösch, H., Brown, L. R., Castano, R., Connor, B., Deutscher, N. M., Eldering, A., Griffith, D., Gunson, M., Kuze, A., Mandrake, L., McDuffie, J., Messerschmidt, J., Miller, C. E., Morino, I., Natraj, V., Notholt, J., O'Brien, D. M., Oyafuso, F., Polonsky, I., Robinson, J., Salawitch, R., Sherlock, V., Smyth, M., Suto, H., Taylor, T. E., Thompson, D. R., Wennberg, P. O., Wunch, D., and Yung, Y. L.: The ACOS CO<sub>2</sub> retrieval algorithm—Part II: Global X<sub>CO<sub>2</sub></sub> data characterization, *Atmospheric Measurement Techniques*, 5, 687–707, <https://doi.org/10.5194/amt-5-687-2012>, 2012.
- Crisp, D., Pollock, H. R., Rosenberg, R., Chapsky, L., Lee, R. A., Oyafuso, F. A., Frankenberg, C., O'Dell, C. W., Brueffe, C. J., Doran, G. B., Eldering, A., Fisher, B. M., Fu, D., Gunson, M. R., Mandrake, L., Osterman, G. B., Schwandner, F. M., Sun, K., Taylor, T. E., Wennberg, P. O., and Wunch, D.: The on-orbit performance of the Orbiting Carbon Observatory-2 (OCO-2) instrument and its radiometrically calibrated products, *Atmospheric Measurement Techniques*, 10, 59–81, <https://doi.org/10.5194/amt-10-59-2017>, 2017.
- De Mazière, M., Sha, M. K., Desmet, F., Hermans, C., Scolas, F., Kumps, N., Metzger, J.-M., Duflot, V., and Cammas, J.-P.: TCCON data from Réunion Island (RE), Release GGG2014.R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.REUNION01.R1>, 2017.
- Deng, F., Jones, D. B. A., Henze, D. K., Bousseres, N., Bowman, K. W., Fisher, J. B., Nassar, R., O'Dell, C., Wunch, D., Wennberg, P. O., Kort, E. A., Wofsy, S. C., Blumenstock, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Heikkinen, P., Sherlock, V., Strong, K., Sussmann, R., and Warneke, T.: Inferring regional sources and sinks of atmospheric CO<sub>2</sub> from GOSAT XCO<sub>2</sub> data, *Atmospheric Chemistry and Physics*, 14, 3703–3727, <https://doi.org/10.5194/acp-14-3703-2014>, <https://acp.copernicus.org/articles/14/3703/2014/>, 2014.
- Deutscher, N. M., Notholt, J., Messerschmidt, J., Weinzierl, C., Warneke, T., Petri, C., and Grupe, P.: TCCON data from Bialystok (PL), Release GGG2014.R2, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.BIALYSTOK01.R2>, 2019.
- Dragokkenky, E. and Tans, P.: Trends in atmospheric carbon dioxide, NOAA/GML, [gml.noaa.gov/ccgg/trends/](http://gml.noaa.gov/ccgg/trends/), last accessed /today, 2021.
- Dubey, M. K., Lindenmaier, R., Henderson, B. G., Green, D., Allen, N. T., Roehl, C. M., Blavier, J.-F., Butterfield, Z. T., Love, S., Hamelmann, J. D., and Wunch, D.: TCCON data from Four Corners (US), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.FOURCORNERS01.R0/1149272>, 2014.
- Eldering, A., O'Dell, C. W., Wennberg, P. O., Crisp, D., Gunson, M. R., Viatte, C., Avis, C., Braverman, A., Castano, R., Chang, A., Chapsky, L., Cheng, C., Connor, B., Dang, L., Doran, G., Fisher, B., Frankenberg, C., Fu, D., Granat, R., Hobbs, J., Lee, R. A., Mandrake, L., McDuffie, J., Miller, C. E., Myers, V., Natraj, V., O'Brien, D., Osterman, G. B., Oyafuso, F., ayne, V. H., Pollock, H. R., Polonsky, I., Roehl, C. M., Rosenberg, R., Schwandner, F., Smyth, M., Tang, V., Taylor, T. E., To, C., Wunch, D., and Yoshimizu, J.: The Orbiting Carbon Observatory-2: first 18 months of science data products, *Atmospheric Measurement Techniques*, 10, 549–563, <https://doi.org/10.5194/amt-10-549-2017>, 2017.
- Feist, D. G., Arnold, S. G., John, N., and Geibel, M. C.: TCCON data from Ascension Island (SH), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.ASCENSION01.R0/1149285>, 2014.

- Feng, L., Palmer, P. I., Bösch, H., and Dance, S.: Estimating surface CO<sub>2</sub> fluxes from space-borne CO<sub>2</sub> dry air mole fraction observations using an ensemble Kalman Filter, *Atmospheric Chemistry and Physics*, 9, 2619–2633, <https://doi.org/10.5194/acp-9-2619-2009>, <https://acp.copernicus.org/articles/9/2619/2009/>, 2009.
- 690 Feng, L., Palmer, P. I., Parker, R. J., Deutscher, N. M., Feist, D. G., Kivi, R., Morino, I., and Sussmann, R.: Estimates of European uptake of CO<sub>2</sub> inferred from GOSAT XCO<sub>2</sub> retrievals: sensitivity to measurement bias inside and outside Europe, *Atmospheric Chemistry and Physics*, 16, 1289–1302, <https://doi.org/10.5194/acp-16-1289-2016>, <https://acp.copernicus.org/articles/16/1289/2016/>, 2016.
- Frankenberg, C., Platt, U., and Wagner, T.: Iterative maximum a posteriori (IMAP)-DOAS for retrieval of strongly absorbing trace gases: Model studies for CH<sub>4</sub> and CO<sub>2</sub> retrieval from near infrared spectra of SCIAMACHY onboard ENVISAT, *Atmospheric Chemistry and*  
 695 *Physics*, 5, 9–22, <https://doi.org/10.5194/acp-5-9-2005>, <https://www.atmos-chem-phys.net/5/9/2005/>, 2005.
- Frankenberg, C., Pollock, R., Lee, R. A. M., Rosenberg, R., Blavier, J.-F., Crisp, D., O'Dell, C. W., Osterman, G. B., Roehl, C., Wennberg, P. O., and Wunch, D.: The Orbiting Carbon Observatory (OCO-2): spectrometer performance evaluation using pre-launch direct sun measurements, *Atmospheric Measurement Techniques*, 8, 301–313, <https://doi.org/10.5194/amt-8-301-2015>, <https://amt.copernicus.org/articles/8/301/2015/>, 2015.
- 700 Goo, T.-Y., Oh, Y.-S., and Velazco, V. A.: TCCON data from Anmeyondo (KR), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.ANMEYONDO01.R0/1149284>, 2014.
- Griffith, D. W., Deutscher, N. M., Velazco, V. A., Wennberg, P. O., Yavin, Y., Aleks, G. K., Washenfelder, R. a., Toon, G. C., Blavier, J.-F., Murphy, C., Jones, N., Kettlewell, G., Connor, B. J., Macatangay, R., Roehl, C., Ryzek, M., Glowacki, J., Cullen, T., and Bryant, G.: TCCON data from Darwin (AU), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA,  
 705 <https://doi.org/10.14291/tcon.ggg2014.darwin01.R0/1149290>, 2014a.
- Griffith, D. W., Velazco, V. A., Deutscher, N. M., Murphy, C., Jones, N., Wilson, S., Macatangay, R., Kettlewell, G., Buchholz, R. R., and Riggensbach, M.: TCCON data from Wollongong (AU), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.wollongong01.R0/1149291>, 2014b.
- Gruber, N., Landschützer, P., and Lovenduski, N. S.: The variable southern ocean carbon sink, *Annual Review of Marine Science*,  
 710 11, 159–186, <https://doi.org/10.1146/annurev-marine-121916-063407>, <https://doi.org/10.1146/annurev-marine-121916-063407>, pMID: 30212259, 2019.
- Hakkarainen, J., Ialongo, I., Maksyutov, S., and Crisp, D.: Analysis of four years of global XCO<sub>2</sub> anomalies as seen by Orbiting Carbon Observatory-2, *Remote Sensing*, 11, <https://doi.org/10.3390/rs11070850>, <https://www.mdpi.com/2072-4292/11/7/850>, 2019.
- Hase, F., Blumenstock, T., Dohe, S., Gross, J., and Kiel, M.: TCCON data from Karlsruhe (DE), Release GGG2014R1, TCCON data archive,  
 715 hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.karlsruhe01.R1/1182416>, 2015.
- Heymann, J., Reuter, M., Hilker, M., Buchwitz, M., Schneising, O., Bovensmann, H., Burrows, J. P., Kuze, A., Suto, H., Deutscher, N. M., Dubey, M. K., Griffith, D. W. T., Hase, F., Kawakami, S., Kivi, R., Morino, I., Petri, C., Roehl, C., Schneider, M., Sherlock, V., Sussmann, R., Velazco, V. A., Warneke, T., and Wunch, D.: Consistent satellite XCO<sub>2</sub> retrievals from SCIAMACHY and GOSAT using the BESD algorithm, *Atmospheric Measurement Techniques*, 8, 2961–2980, <https://doi.org/10.5194/amt-8-2961-2015>, <https://amt.copernicus.org/articles/8/2961/2015/>, 2015.
- 720 Houweling, S., Baker, D., Basu, S., Boesch, H., Butz, A., Chevallier, F., Deng, F., Dlugokencky, E. J., Feng, L., Ganshin, A., Hasekamp, O., Jones, D., Maksyutov, S., Marshall, J., Oda, T., O'Dell, C. W., Oshchepkov, S., Palmer, P. I., Peylin, P., Poussi, Z., Reum, F., Takagi, H., Yoshida, Y., and Zhuravlev, R.: An intercomparison of inverse models for estimating sources and sinks of CO<sub>2</sub> using GOSAT

measurements, *Journal of Geophysical Research: Atmospheres*, 120, 5253–5266, <https://doi.org/https://doi.org/10.1002/2014JD022962>,  
725 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JD022962>, 2015.

Iraci, L. T., Podolske, J., Hillyard, P. W., Roehl, C., Wennberg, P. O., Blavier, J.-F., Allen, N., Wunch, D., Osterman, G. B.,  
and Albertson, R.: TCCON data from Edwards (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA,  
<https://doi.org/10.14291/tcon.ggg2014.edwards01.R1/1255068>, 2016a.

Iraci, L. T., Podolske, J. R., Hillyard, P. W., Roehl, C., Wennberg, P. O., Blavier, J.-F., Landeros, J., Allen, N., Wunch, D., Zavaleta, J.,  
730 Quigley, E., Osterman, G. B., Barrow, E., and Barney, J.: TCCON data from Indianapolis (US), Release GGG2014.R1, TCCON data  
archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.INDIANAPOLIS01.R1/1330094>, 2016b.

Jiang, F., Wang, H., Chen, J. M., Ju, W., Tian, X., Feng, S., Li, G., Chen, Z., Zhang, S., Lu, X., Liu, J., Wang, H., Wang, J., He,  
W., and Wu, M.: Regional CO<sub>2</sub> fluxes from 2010 to 2015 inferred from GOSAT XCO<sub>2</sub> retrievals using a new version of the  
Global Carbon Assimilation System, *Atmospheric Chemistry and Physics*, 21, 1963–1985, <https://doi.org/10.5194/acp-21-1963-2021>,  
735 <https://acp.copernicus.org/articles/21/1963/2021/>, 2021.

Kataoka, F., Crisp, D., Taylor, T. E., O'Dell, C. W., Kuze, A., Shiomi, K., Suto, H., Bruegge, C., Schwandner, F. M., Rosenberg, R., Chapsky,  
L., and Lee, R. A. M.: The cross-calibration of spectral radiances and cross-validation of CO<sub>2</sub> estimates from GOSAT and OCO-2, *Remote  
Sensing*, 9, <https://doi.org/10.3390/rs9111158>, <https://www.mdpi.com/2072-4292/9/11/1158>, 2017.

Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., and Sakashita, M.: TCCON data from Saga (JP), Release  
740 GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.saga01.R0/1149283>, 2014.

Keeling, C., Whorf, T., Wahlen, M., and van der Plichtt, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980,  
*Nature*, 375, 666–670, <https://doi.org/https://doi.org/10.1038/375666a0>, 1995.

Kiel, M., O'Dell, C. W., Fisher, B., Eldering, A., Nassar, R., MacDonald, C. G., and Wennberg, P. O.: How bias correction goes  
wrong: measurement of XCO<sub>2</sub> affected by erroneous surface pressure estimates, *Atmospheric Measurement Techniques*, 12, 2241–2259,  
745 <https://doi.org/10.5194/amt-12-2241-2019>, <https://www.atmos-meas-tech.net/12/2241/2019/>, 2019.

Kivi, R., Heikkinen, P., and Kyrö, E.: TCCON data from Sodankylä (FI), Release GGG2014.R0, TCCON data archive, hosted by Caltech-  
DATA, <https://doi.org/10.14291/tcon.ggg2014.sodankyla01.R0/1149280>, 2014.

Kulawik, S., Wunch, D., O'Dell, C., Frankenberg, C., Reuter, M., Oda, T., Chevallier, F., Sherlock, V., Buchwitz, M., Osterman, G., Miller,  
C. E., Wennberg, P. O., Griffith, D., Morino, I., Dubey, M. K., Deutscher, N. M., Notholt, J., Hase, F., Warneke, T., Sussmann, R., Robinson,  
750 J., Strong, K., Schneider, M., De Mazière, M., Shiomi, K., Feist, D. G., Iraci, L. T., and Wolf, J.: Consistent evaluation of ACOS-GOSAT,  
BESD-SCIAMACHY, CarbonTracker, and MACC through comparisons to TCCON, *Atmospheric Measurement Techniques*, 9, 683–709,  
<https://doi.org/10.5194/amt-9-683-2016>, <https://amt.copernicus.org/articles/9/683/2016/>, 2016.

Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform  
spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733,  
755 <https://doi.org/10.1364/AO.48.006716>, <http://ao.osa.org/abstract.cfm?URI=ao-48-35-6716>, 2009.

Kuze, A., Suto, H., Shiomi, K., Kawakami, S., Tanaka, M., Ueda, Y., Deguchi, A., Yoshida, J., Yamamoto, Y., Kataoka, F., Taylor, T. E., and  
Buijs, H. L.: Update on GOSAT TANSO-FTS performance, operations, and data products after more than 6 years in space, *Atmospheric  
Measurement Techniques*, 9, 2445–2461, <https://doi.org/10.5194/amt-9-2445-2016>, 2016.

Kuze, A., Kikuchi, N., Kataoka, F., Suto, H., Shiomi, K., and Kondo, Y.: Detection of Methane Emission from a Local Source Using GOSAT  
760 Target Observations, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12020267>, <https://www.mdpi.com/2072-4292/12/2/267>, 2020.



- Lindenmaier, R., Dubey, M. K., Henderson, B. G., Butterfield, Z. T., Herman, J. R., Rahn, T., and Lee, S.-H.: Multiscale observations of CO<sub>2</sub>, 13CO<sub>2</sub>, and pollutants at Four Corners for emission verification and attribution, *Proceedings of the National Academy of Sciences*, 111, 8386–8391, <https://doi.org/10.1073/pnas.1321883111>, <https://www.pnas.org/content/111/23/8386>, 2014.
- 765 Lindqvist, H., O'Dell, C. W., Basu, S., Boesch, H., Chevallier, F., Deutscher, N., Feng, L., Fisher, B., Hase, F., Inoue, M., Kivi, R., Morino, I., Palmer, P. I., Parker, R., Schneider, M., Sussmann, R., and Yoshida, Y.: Does GOSAT capture the true seasonal cycle of carbon dioxide?, *Atmospheric Chemistry and Physics*, 15, 13 023–13 040, <https://doi.org/10.5194/acp-15-13023-2015>, <https://www.atmos-chem-phys.net/15/13023/2015/>, 2015.
- Liu, C., Wang, W., and Sun, Y.: TCCON data from Hefei (PRC), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.HEFEI01.R0>, 2018.
- 770 Liu, J., Bowman, K. W., Schimel, D. S., Parazoo, N. C., Jiang, Z., Lee, M., Bloom, A. A., Wunch, D., Frankenberg, C., Sun, Y., O'Dell, C. W., Gurney, K. R., Menemenlis, D., Gierach, M., Crisp, D., and Eldering, A.: Contrasting carbon cycle responses of the tropical continents to the 2015–2016 El Niño, *Science*, 358, <https://doi.org/10.1126/science.aam5690>, <http://science.sciencemag.org/content/358/6360/eaam5690>, 2017.
- Morino, I., Yokozeki, N., Matzuzaki, T., and Horikawa, M.: TCCON data from Rikubetsu (JP), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.RIKUBETSU01.R1/1242265>, 2016.
- 775 Morino, I., Matsuzaki, T., and Horikawa, M.: TCCON data from Tsukuba (JP), 125HR, Release GGG2014R2, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.TSUKUBA02.R2>, 2018a.
- Morino, I., Velazco, V. A., Hori, A., Uchino, O., and Griffith, D. W. T.: TCCON data from Burgos, Ilocos Norte (PH), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.BURGOS01.R0>, 2018b.
- 780 Müller, A., Tanimoto, H., Sugita, T., Machida, T., Nakaoka, S., Patra, P. K., Laughner, J., and Crisp, D.: New approach to evaluate satellite-derived XCO<sub>2</sub> over oceans by integrating ship and aircraft observations, *Atmospheric Chemistry and Physics*, 21, 8255–8271, <https://doi.org/10.5194/acp-21-8255-2021>, <https://acp.copernicus.org/articles/21/8255/2021/>, 2021.
- Nelson, R. R. and O'Dell, C. W.: The impact of improved aerosol priors on near-infrared measurements of carbon dioxide, *Atmospheric Measurement Techniques*, 12, 1495–1512, <https://doi.org/10.5194/amt-12-1495-2019>, <https://www.atmos-meas-tech.net/12/1495/2019/>, 785 2019.
- Nguyen, H., Osterman, G., Wunch, D., O'Dell, C., Mandrake, L., Wennberg, P., Fisher, B., and Castano, R.: A method for colocating satellite XCO<sub>2</sub> data to ground-based data and its application to ACOS-GOSAT and TCCON, *Atmospheric Measurement Techniques*, 7, 2631–2644, <https://doi.org/10.5194/amt-7-2631-2014>, <https://amt.copernicus.org/articles/7/2631/2014/>, 2014.
- Notholt, J., Petri, C., Warneke, T., Deutscher, N. M., Palm, M., Buschmann, M., Weinzierl, C., Macatangay, R. C., 790 and Grupe, P.: TCCON data from Bremen (DE), Release GGG2014.R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.BREMEN01.R1>, 2019.
- OCO-2 Science Team, Gunson, M., and Eldering, A.: ACOS GOSAT/TANSO-FTS Level 2 bias-corrected XCO<sub>2</sub> and other select fields from the full-physics retrieval aggregated as daily files V9r, Goddard Earth Sciences Data and Information Services Center (GES DISC), Greenbelt, MD, USA, <https://doi.org/https://doi.org/10.5067/VWSABTO7ZII4>, last accessed /today, 2019a.
- 795 OCO-2 Science Team, Gunson, M., and Eldering, A.: ACOS GOSAT/TANSO-FTS Level 2 Full Physics Standard Product V9r, Goddard Earth Sciences Data and Information Services Center (GES DISC), Greenbelt, MD, USA, <https://doi.org/https://doi.org/10.5067/OSGTIL9OV0PN>, last accessed /today, 2019b.

- OCO-2 Science Team, Gunson, M., and Eldering, A.: OCO-2 Level 2 bias-corrected XCO<sub>2</sub> and other select fields from the full-physics retrieval aggregated as daily files V10r, Goddard Earth Sciences Data and Information Services Center (GES DISC), Greenbelt, MD, USA, <https://doi.org/https://doi.org/10.5067/E4E140XDMPO2>, last accessed /today, 2020.
- O'Dell, C., Osterman, G., Eldering, A., Cheng, C., Crisp, D., Frankenberg, C., and Fisher, B.: Orbiting Carbon Observatory-2 & 3 Retrievals of carbon dioxide from GOSAT using the atmospheric CO<sub>2</sub> observations from space (ACOS) algorithm; Level 2 standard product and lite data product data user's guide, v9, Tech. rep., Jet Propulsion Laboratory, [https://docsserver.gesdisc.eosdis.nasa.gov/public/project/OCO/ACOS\\_v9\\_DataUsersGuide.pdf](https://docsserver.gesdisc.eosdis.nasa.gov/public/project/OCO/ACOS_v9_DataUsersGuide.pdf), 2020.
- O'Dell, C. W., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Crisp, D., Eldering, A., Fisher, B., Gunson, M., McDuffie, J., Miller, C. E., Natraj, V., Oyafuso, F., Polonsky, I., Smyth, M., Taylor, T., Toon, G. C., Wennberg, P. O., and Wunch, D.: The ACOS CO<sub>2</sub> retrieval algorithm—Part 1: Description and validation against synthetic observations, *Atmospheric Measurement Techniques*, 5, 99–121, <https://doi.org/10.5194/amt-5-99-2012>, 2012.
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., Frankenberg, C., Kiel, M., Lindqvist, H., Mandrake, L., Merrelli, A., Natraj, V., Nelson, R. R., Osterman, G. B., Payne, V. H., Taylor, T. R., Wunch, D., Drouin, B. J., Oyafuso, F., Chang, A., McDuffie, J., Smyth, M., Baker, D. F., Basu, S., Chevallier, F., Crowell, S. M. R., Feng, L., Palmer, P. I., Dubey, M., García, O. E., Griffith, D. W. T., Hase, F., Iraci, L. T., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Sussmann, R., Te, Y., Uchino, O., and Velazco, V. A.: Improved Retrievals of Carbon Dioxide from the Orbiting Carbon Observatory-2 with the version 8 ACOS algorithm, *Atmospheric Measurement Techniques Discussions*, 2018, 1–57, <https://doi.org/10.5194/amt-2018-257>, <https://www.atmos-meas-tech-discuss.net/amt-2018-257/>, 2018.
- Osterman, G., O'Dell, C., Eldering, A., Fisher, B., Crisp, D., Cheng, C., Frankenberg, C., Lambert, A., Gunson, M., Mandrake, L., and Wunch, D.: Orbiting Carbon Observatory-2 & 3 Data Product User's Guide, v10, Tech. rep., Jet Propulsion Laboratory, [https://docsserver.gesdisc.eosdis.nasa.gov/public/project/OCO/OCO2\\_OCO3\\_B10\\_DUG.pdf](https://docsserver.gesdisc.eosdis.nasa.gov/public/project/OCO/OCO2_OCO3_B10_DUG.pdf), 2020.
- Oyafuso, F., Payne, V. H., Drouin, B. J., Devi, V. M., Benner, D. C., Sung, K., Yu, S., Gordon, I. E., Kochanov, R., Tan, Y., Crisp, D., Mlawer, E. J., and Guillaume, A.: High accuracy absorption coefficients for the Orbiting Carbon Observatory-2 (OCO-2) mission: Validation of updated carbon dioxide cross-sections using atmospheric spectra, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 203, 213–223, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2017.06.012>, <https://www.sciencedirect.com/science/article/pii/S0022407317301322>, *hITRAN2016 Special Issue*, 2017.
- Palmer, P. I., Feng, L., Baker, D., Chevallier, F., Bösch, H., and Somkuti, P.: Net carbon emissions from African biosphere dominate pan-tropical atmospheric CO<sub>2</sub> signal, *Nature Communications*, 10, 3344, <https://doi.org/10.1038/s41467-019-11097-w>, <https://doi.org/10.1038/s41467-019-11097-w>, 2019.
- Payne, V. H., Drouin, B. J., Oyafuso, F., Kuai, L., Fisher, B. M., Sung, K., Nemchick, D., Crawford, T. J., Smyth, M., Crisp, D., Adkins, E., Hodges, J. T., Long, D. A., Mlawer, E. J., Merrelli, A., Lunny, E., and O'Dell, C. W.: Absorption coefficient (ABSCO) tables for the Orbiting Carbon Observatories: Version 5.1, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 255, 107 217, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2020.107217>, <http://www.sciencedirect.com/science/article/pii/S0022407320302016>, 2020.
- Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Pétron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *Proceedings of the National Academy of Sciences*, 104, 18 925–18 930, <https://doi.org/10.1073/pnas.0708986104>, <https://www.pnas.org/content/104/48/18925>, 2007.

- 835 Petri, C., Vrekoussis, M., Rousogonous, C., Warneke, T., Sciare, J., and Notholt, J.: TCCON data from Nicosia (CY), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.nicosia01.R0>, 2020.
- Pollard, D. F., Robinson, J., and Shiona, H.: TCCON data from Lauder (NZ), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.LAUDER03.R0>, 2019.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K.,
- 840 Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *Journal of Climate*, 24, 3624 – 3648, <https://doi.org/10.1175/JCLI-D-11-00015.1>, <https://journals.ametsoc.org/view/journals/clim/24/14/jcli-d-11-00015.1.xml>, 2011.
- Rödenbeck, C.: Estimating CO<sub>2</sub> sources and sinks from atmospheric mixing ratio measurements using a global inversion of atmospheric
- 845 transport, Technical Report 6, Max Planck Institute for Biogeochemistry, Jena, 2005.
- Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO<sub>2</sub> data, *Biogeosciences*, 15, 2481–2498, <https://doi.org/10.5194/bg-15-2481-2018>, <https://bg.copernicus.org/articles/15/2481/2018/>, 2018.
- Strong, K., Roche, S., Franklin, J. E., Mendonca, J., Lutsch, E., Weaver, D., Fogal, P. F., Drummond, J. R., Batchelor, R.,
- 850 and Lindenmaier, R.: TCCON data from Eureka (CA), Release GGG2014.R3, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.EUREKA01.R3>, 2019.
- Sussmann, R. and Rettinger, M.: TCCON data from Garmisch (DE), Release GGG2014R2, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.GARMISCH01.R2>, 2018a.
- Sussmann, R. and Rettinger, M.: TCCON data from Zugspitze (DE), Release GGG2014.R1, TCCON data archive, hosted by CaltechDATA,
- 855 <https://doi.org/10.14291/TCCON.GGG2014.ZUGSPITZE01.R1>, 2018b.
- Taylor, T. E., O'Dell, C. W., Frankenberg, C., Partain, P. T., Cronk, H. Q., Savtchenko, A., Nelson, R. R., Rosenthal, E. J., Chang, A., Fisher, B. M., Osterman, G. B., Pollock, R. H., Crisp, D., Eldering, A., and Gunson, M. R.: Orbiting Carbon Observatory-2 (OCO-2) cloud screening algorithms: validation against collocated MODIS and CALIOP data, *Atmospheric Measurement Techniques*, 9, 973–989, <https://doi.org/10.5194/amt-9-973-2016>, 2016.
- 860 Tè, Y., Jeseck, P., and Janssen, C.: TCCON data from Paris (FR), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.paris01.R0/1149279>, 2014.
- Thompson, D. R., Benner, D. C., Brown, L. R., Crisp, D., Devi, V. M., Jiang, Y., Natraj, V., Oyafuso, F., Sung, K., Wunch, D., Castano, R., and Miller, C. E.: Atmospheric validation of high accuracy CO<sub>2</sub> absorption coefficients for the OCO-2 mission, *Journal Quant. Spect. Rad. Trans.*, 113, 2265–2276, <https://doi.org/10.1016/j.jqsrt.2012.05.021>, 2012.
- 865 UoE: UoE, <https://www.geos.ed.ac.uk/~lfeng/otool.pdf>, 2021.
- Wang, H., Jiang, F., Wang, J., Ju, W., and Chen, J. M.: Terrestrial ecosystem carbon flux estimated using GOSAT and OCO-2 XCO<sub>2</sub> retrievals, *Atmospheric Chemistry and Physics*, 19, 12 067–12 082, <https://doi.org/10.5194/acp-19-12067-2019>, <https://acp.copernicus.org/articles/19/12067/2019/>, 2019.
- Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N. M., Petri, C., and Grupe, P.: TCCON data from Orléans (FR), Release
- 870 GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.ORLEANS01.R1>, 2019.
- Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., and Allen, N.: TCCON data from Caltech (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.pasadena01.R1/1182415>, 2015.

- Wennberg, P. O., Roehl, C. M., Blavier, J.-F., Wunch, D., and Allen, N. T.: TCCON data from Jet Propulsion Laboratory (US), 2011, Release GGG2014.R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.JPL02.R1/1330096>, 2016a.
- 875 Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.: TCCON data from Lamont (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.lamont01.R1/1255070>, 2016b.
- Wennberg, P. O., Roehl, C., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R. a., Keppel-Aleks, G., Allen, N., and Ayers, J.: TCCON data from Park Falls (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, 880 <https://doi.org/10.14291/TCCON.GGG2014.PARKFALLS01.R1>, 2017.
- Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, <https://doi.org/10.1098/rsta.2010.0240>, 2011a.
- Wunch, D., Wennberg, P., Toon, G., Connor, B., Fisher, B., Osterman, G., Frankenberg, C., Mandrake, L., O'Dell, C., P.Ahonen, Biraud, 885 S., Castano, R., Cressie, N., Crisp, D., Deutscher, N., Eldering, A., Fisher, M., Griffith, D., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyro, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C., Morino, I., Notholt, J., Oyafuso, F., Rettinger, M., Robinson, J., Roehl, C., Salawitch, R., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D., Uchino, O., Warneke, T., and Wofsy, S.: A method for evaluating bias in global measurements of CO<sub>2</sub> total columns from space, *Atmospheric Chemistry and Physics*, 11, 12 317–12 337, <https://doi.org/10.5194/acp-11-12317-2011>, 2011b.
- 890 Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M., Liu, C., Feist, D. G., and Wennberg, P. O.: Documentation for the 2014 TCCON Data Release, Tech. rep., CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.DOCUMENTATION.R0/1221662>, 2015.
- Wunch, D., Mendonca, J., Colebatch, O., Allen, N. T., Blavier, J.-F., Springett, S., Neufeld, G., Strong, K., Kessler, R., and Worthy, D.: TCCON data from East Trout Lake, SK (CA), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.EASTTROUTLAKE01.R0/1348207>, 2017a.
- 895 Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., O'Dell, C., Mandrake, L., Viatte, C., Kiel, M., Griffith, D. W. T., Deutscher, N. M., Velazco, V. A., Notholt, J., Warneke, T., Petri, C., De Maziere, M., Sha, M. K., Sussmann, R., Rettinger, M., Pollard, D., Robinson, J., Morino, I., Uchino, O., Hase, F., Blumenstock, T., Feist, D. G., Arnold, S. G., Strong, K., Mendonca, J., Kivi, R., Heikkinen, P., Iraci, L., Podolske, J., Hillyard, P. W., Kawakami, S., Dubey, M. K., Parker, H. A., Sepulveda, E., García, O. E., Te, Y., Jeseck, P., Gunson, M. R., Crisp, D., and Eldering, A.: Comparisons of the Orbiting Carbon Observatory-2 (OCO-2) 900 X<sub>CO<sub>2</sub></sub> measurements with TCCON, *Atmospheric Measurement Techniques*, 10, 2209–2238, <https://doi.org/10.5194/amt-10-2209-2017>, <https://www.atmos-meas-tech.net/10/2209/2017/>, 2017b.
- Yoshida, Y., Ota, Y., Eguchi, N., Kikuchi, N., Nobuta, K., Tran, H., Morino, I., and Yokota, T.: Retrieval algorithm for CO<sub>2</sub> and CH<sub>4</sub> column abundances from short-wavelength infrared spectral observations by the Greenhouse Gases Observing Satellite, *Atmospheric Measurement Techniques*, 4, 717–734, <https://doi.org/10.5194/amt-4-717-2011>, 2011.
- 905 Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO<sub>2</sub> and XCH<sub>4</sub> and their validation using TCCON data, *Atmospheric Measurement Techniques*, 6, 1533–1547, <https://doi.org/10.5194/amt-6-1533-2013>, <https://amt.copernicus.org/articles/6/1533/2013/>, 2013.

The ACOS GOSAT v9 XCO<sub>2</sub> data are ~~publicly available via NASA's~~ [available on the NASA Goddard Earth Science Data and Information Services Center](https://doi.org/10.5067/OSGTH9OV0PN) (GES-DISC ~~at~~ <https://doi.org/10.5067/OSGTH9OV0PN> (L2Std files, OCO-2 Science Team et al. (2019b) ~~) and~~ [in both the per-orbit full format](https://doi.org/10.5067/OSGTH9OV0PN) (OCO-2 Science Team et al., 2019b, <https://doi.org/10.5067/OSGTH9OV0PN>) ~~and in~~ [the per-day lite format](https://doi.org/10.5067/VWSABTO7ZII4) (OCO-2 Science Team et al., 2019a, <https://doi.org/10.5067/VWSABTO7ZII4>). The monthly super-lite files, containing only the most essential variables for each satellite observation, are available at CaltechDATA (<https://doi.org/10.5067/OSGTH9OV0PN> (L2Lite files, OCO-2 Science Team et al. (2019a) [0000-0003-1080-9922](https://doi.org/10.5067/OSGTH9OV0PN))). The OCO-2 v10 L2Lite files containing the bias corrected and quality filtered XCO<sub>2</sub> data are [also](https://doi.org/10.5067/E4E140XDMPO2) available on the GES-DISC ~~at~~ <https://doi.org/10.5067/E4E140XDMPO2> (OCO-2 Science Team et al., 2020, <https://doi.org/10.5067/E4E140XDMPO2>). The TC-CON data for individual stations are available on the CaltechDATA site ~~at~~ (<https://data.caltech.edu/>). The CarbonTracker data are available on the NOAA GML site ~~at~~ (<https://carbontracker.noaa.gov>). The Carboscope model data are available at <http://www.bgc-jena.mpg.de/CarboScope>. The CAMS model data are available at <https://atmosphere.copernicus.eu/data>. The UoL model data are available at <https://www.geos.ed.ac.uk/~lfeng/>.


*Author contributions.* TET and CWO conceptualized the study and performed formal analysis. Additional formal analysis was performed by HL and AC. Funding acquisition was provided by DC, AK, POW, MG, and AE. Algorithm development was provided by TET, CWO, DC, POW, MG, AE, BF, MK, RRN, AM, and GO. Model data was provided by FC, PIP, and LF. TCCON stations were maintained and data provided by POW, NMD, MKD, DGF, OEG, DWTG, FH, LTI, RK, CL, MDM, IM, JN, YSO, HO, DFP, MR, CMR, MS, MKS, KS, RS, YT, VAV, MW, TW, and DW. Paper writing of the draft was performed by TET, CWO, and DC. All authors contributed to editing the final version of the manuscript.

*Competing interests.* We declare no competing interests in this work.


*Acknowledgements.* The CSU contribution to this work was supported by JPL subcontract 1439002. TET acknowledges assistance from Peter Somkuti and Heather Cronk at CSU/CIRA with Python map plotting. CarbonTracker results provided by NOAA ESRL, Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov>. The GEOS data used in this study were provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center. HL acknowledges support from the Academy of Finland (project 331829). AM's contributions to this work were supported by JPL subcontract 1577173. P.I.P. and L.F. acknowledge support from the UK National Centre for Earth Observation funded the National Environment Research Council (NE/R016518/1). The TCCON stations at Rikubetsu, Tsukuba, and Burgos are supported in part by the GOSAT series project. Local support for Burgos is provided by the Energy Development Corporation (EDC, Philippines). The TCCON site at Réunion Island is operated by the Royal Belgian Institute for Space Aeronomy with financial support since 2014 by the EU project ICOS-Inwire and the ministerial decree for ICOS (FR/35/IC1 to FR/35/C6) and local activities supported by LACy/UMR8105 and by OSU-R/UMS3365 – Université de La Réunion. The TCCON stations at Garmisch and Zugspitze

940 have been supported by the European Space Agency (ESA) under grant 4000120088/17/I-EF and by the German Bundesministerium für  
Wirtschaft und Energie (BMWi) via the DLR under grant 50EE1711D as well as by the Helmholtz Society via the research programme  
ATMO. The Paris TCCON site has received funding from Sorbonne Université, the French research center CNRS, the French space agency  
CNES, and Région Île-de-France. The Ascension Island TCCON station has been supported by the European Space Agency (ESA) un-  
der grant 4000120088/17/I-EF and by the German Bundesministerium für Wirtschaft und Energie (BMWi) under grants 50EE1711C and  
945 50EE1711E. We thank the ESA Ariane Tracking Station at North East Bay, Ascension Island, for hosting and local support. The Anmywondo  
TCCON station was funded by the Korea Meteorological Administration Research and Development Program "Development of Monitoring  
and Analysis Techniques for Atmospheric Composition in Korea." under Grant (KMA 2018-00522) NMD is supported by an Australian  
Research Council (ARC) Future Fellowship, FT180100327. The Darwin and Wollongong TCCON sites have been supported by a series  
of ARC grants, including DP160100598, DP140100552, DP110103118, DP0879468 and LE0668470, and NASA grants NAG5-12247 and  
950 NNG05-GD07G. The Eureka measurements were made at the Polar Environment Atmospheric Research Laboratory (PEARL) by the Cana-  
dian Network for the Detection of Atmospheric Change (CANDAC), primarily supported by the Natural Sciences and Engineering Research  
Council of Canada, Environment and Climate Change Canada, and the Canadian Space Agency. MKD thanks the LANL LDRD program  
for support operating the Four Corners TCCON site. The TCCON Nicosia site has received additional support from the European Union's  
Horizon 2020 research and innovation programme under grant agreement No. 856612 and the Cyprus Government, and by the University  
955 of Bremen. The Anmyeondo TCCON station is funded by the Korea Meteorological Administration Research and Development Program  
"Development of Monitoring and Analysis Techniques for Atmospheric Composition in Korea." under Grant (KMA 2018-00522).


## Appendix A: Seasonal cycle plots of ACOS GOSAT v9 XCO<sub>2</sub> versus TCCON for individual stations




**Figure A1.** Daily averaged bias-corrected ACOS GOSAT v9 versus collocated TCCON XCO<sub>2</sub> at Anmyeondo, Korea. Left panel (A) shows the one-to-one scatter plot of the daily mean XCO<sub>2</sub> for GOSAT versus TCCON. Middle panel (B) shows the time series of daily mean GOSAT XCO<sub>2</sub> (blue circles) with fit (blue line) and the TCCON XCO<sub>2</sub> (grey triangles) with fit (orange line) over the eleven year data record. Right panel (C) shows the time series of calculated  $\Delta XCO_2^{TCCON}$ , with the mean difference (horizontal solid black line) and  $\pm 1$  standard deviation (gray shading). In these plots, the three GOSAT observation modes have been combined in order to provide the maximum number of collocations possible for the seasonal fits.




**Figure A2.** Same as Figure A1, but for Ascension Island, located in the Pacific ocean off the west coast of Africa.



**Figure A3.** Same as Figure A1, but for Białystok, Poland.




**Figure A4.** Same as Figure A1, but for Bremen, Germany.




**Figure A5.** Same as Figure A1, but for Burgos, Philippines.







**Figure A6.** Same as Figure A1, but for Caltech, California.




**Figure A7.** Same as Figure A1, but for Darwin, Australia.




**Figure A8.** Same as Figure A1, but for Edwards, California.




**Figure A9.** Same as Figure A1, but for East Trout Lake, Canada.




**Figure A10.** Same as Figure A1, but for Garmisch, Germany.




**Figure A11.** Same as Figure A1, but for Hefei, China.




**Figure A12.** Same as Figure A1, but for Izaña, Tenerife, Spain.




**Figure A13.** Same as Figure A1, but for Jet Propulsion Laboratory (JPL), Pasadena, California. This site has been used occasionally for the simultaneous operation of a TCCON instrument during the thermal vacuum testing of OCO-2 (Frankenberg et al., 2015) and OCO-3.




**Figure A14.** Same as Figure A1, but for Karlsruhe, Germany.




**Figure A15.** Same as Figure A1, but for Lauder, New Zealand.




**Figure A16.** Same as Figure A1, but for Orleans, France.




**Figure A17.** Same as Figure A1, but for Paris, France.




**Figure A18.** Same as Figure A1, but for Park Falls, Wisconsin.




**Figure A19.** Same as Figure A1, but for Reunion Island, off the east coast of Madagascar.




**Figure A20.** Same as Figure A1, but for Rikubetsu, Japan.




**Figure A21.** Same as Figure A1, but for Saga, Japan.




**Figure A22.** Same as Figure A1, but for Sodankylä, Finland.



**Figure A23.** Same as Figure A1, but for Tsukuba, Japan.



**Figure A24.** Same as Figure A1, but for Wollongong, Australia.



**Figure A25.** Same as Figure A1, but for Zugspitze, Germany.