

High-resolution land-use land-cover change data for regional climate modelling applications over Europe - Part 1: The plant functional type basemap for 2015

Vanessa Reinhart^{1,2}, Peter Hoffmann^{1,2}, Diana Rechid¹, Jürgen Böhner³, and Benjamin Bechtel⁴

¹Helmholtz-Zentrum Hereon, Climate Service Center Germany (GERICS), Fischertwiete 1, 20095 Hamburg, Germany

²Universität Hamburg, Institute of Geography, Section Physical Geography, Bundesstraße 55, 20146 Hamburg, Germany

³Universität Hamburg, Institute of Geography, Cluster of Excellence “Climate, Climatic Change, and Society” (CLICCS), 805A Geomatikum, Bundesstraße 55, D-20146, Hamburg, Germany

⁴Ruhr-Universität Bochum, Department of Geography, Universitätsstraße 150/ Gebäude IA, 44801 Bochum, Germany

Correspondence: Vanessa Reinhart (vanessa.reinhart@hereon.de)

Abstract. The concept of plant functional types (PFTs) is shown to be beneficial in representing the complexity of plant characteristics in land use and climate change studies using regional climate models (RCMs). By representing land use and land cover (LULC) as functional traits, responses and effects of specific plant communities can be directly coupled to the lowest atmospheric layers. To meet the requirements of RCMs for realistic LULC distribution, we developed a PFT dataset for Europe (LANDMATE PFT Version 1.0; Reinhart et al., 2021b). The dataset is based on the high-resolution ESA-CCI land cover dataset and is further improved through the the additional use of climate information. Within the LANDMATE PFT dataset, satellite-based LULC information and climate data are combined to ~~achieve the best possible~~ create the representation of the diverse plant communities and their functions in the respective regional ecosystems while keeping the dataset most flexible for application in RCMs. Each LULC class of ESA-CCI is translated into PFT or PFT fractions including climate information by using the Holdridge Life Zone concept. Through ~~the~~ consideration of regional climate data, the resulting PFT map for Europe is regionally customized. A thorough evaluation of the LANDMATE PFT dataset is done using a comprehensive ground truth database over the European Continent. ~~A suitable evaluation method has been developed and applied to assess the quality of the new PFT dataset.~~ The assessment shows that the dominant LULC ~~groupstypes~~, cropland and woodland, are well represented within the dataset while uncertainties are found for some less represented LULC ~~groupstypes~~. The LANDMATE PFT dataset provides a realistic, high-resolution LULC distribution for implementation in RCMs and is used as basis for the LUCAS LUC dataset introduced in the companion paper by Hoffmann et al. (submitted) which is available for use as LULC change input for RCM experiment setups focused on investigating LULC change impact.

Copyright statement. CC BY 4.0

1 Introduction

20 Land use and land cover (LULC), including the vegetation type and function, ~~was declared an~~ were declared Essential Climate Variables (ECVs) by the Global Climate Observing System (GCOS) (Bojinski et al., 2014). Changes in ECVs are crucial factors of climate change and therefore need to be monitored and further represented in climate models to be able to assimilate and understand atmospheric processes and feedback effects on different scales. For LULC, anthropogenic modifications are the most important drivers of change. De- and reforestation and expansion of urban and cropland areas affect biogeophysical

25 (e.g., albedo, roughness, evapotranspiration, runoff) and biogeochemical (e.g., carbon emissions and sinks) surface properties and processes (Mahmood et al., 2014; Lawrence and Vandecar, 2015; Alkama and Cescatti, 2016; Perugini et al., 2017; Davin et al., 2020). Besides LULC changes, land management practices are being assessed regarding influence of related land surface modifications on regional climate, and also the potential of land management practices regarding climate change adaptation and mitigation efforts (Lobell et al., 2006; Kueppers et al., 2007; Burke and Emerick, 2016).

30 In order to represent impacts and feedbacks of LULC modifications as ~~realistic~~ realistically as possible, regional climate models (RCMs) require an accurate representation of LULC and its changes. In this context, the concept of plant functional types (PFTs) is ~~increasingly used~~ used frequently for the representation of LULC in RCMs ~~-(Davin et al., 2020).~~

PFTs are aggregated plant species groups that share comparable biophysical properties and functions. The aggregation makes it possible to represent these functionality groups within one single model grid unit as a mosaic. The main difference of the

35 PFT representation in comparison to the LULC class representation is the grouping of vegetation according to function instead of a descriptive definition. The function of a group is directly represented by the biophysical and biochemical properties that are prescribed or dynamically computed within the vegetation layer of an RCM.

A comprehensive review of the subsequent development of PFTs representing vegetation dynamics in climate models was done by Wullschleger et al. (2014). The need for applicable global PFT maps for vegetation models that are used with at-

40 mospheric models was already well emphasized by Box (1996). Moreover, the requirement that a climate model should include a vegetation model representing the biosphere was discussed by Lavorel et al. (2007). One criterion that is highly emphasized is the inter-regional applicability of a preferably simple PFT classification, which has the ability to capture key characteristics of the biosphere from biome to continental scale, regardless of climate zone and individual vegetation composition. A variety of PFT definitions and cross-walking procedures (CWPs), used for translating LULC products into global

45 or regional PFT maps ~~, are currently available. The European Space Agency Climate Change Initiative (ESA-CCI) and the United States Geological Service (USGS) provide the only two ready-to-use continuous global products to the community (Poulter et al., 2015; Sulla-Menashe and Friedl, 2018)~~ emerged in the last decades (Bonan et al., 2002; Poulter et al., 2011; Ottlé et al., 201

. The respective CWP documentation consist of the utilized input data, the translation table where each LULC class is assigned to PFT proportions and a description on how the input data is used to create the final product. However, the individual PFT

50 definitions and CWPs as well as the mostly satellite based input data differ greatly in complexity and temporal and horizontal resolution (Bonan et al., 2002; Winter et al., 2009; Lu and Kueppers, 2012). Moreover, inter-regional consistency cannot be achieved by products that origin from regionally constrained input data or regionally adapted CWPs. Therefore, the additional

use of climate information in the CWP from LULC to PFT is a highly useful step in order to create a dynamically customizable product, that can be adapted to various climate and vegetation characteristics (Poulter et al., 2011).

55 With the present work, we introduce a PFT map for the European Continent that specifically addresses the requirements of the RCM community (Bontemps et al., 2013). The land cover maps of the ESA-CCI are translated into 16 PFTs creating an updated version of the interactive MOsaic-based Vegetation (iMOVE) PFTs that were originally developed for the RCM REMO (Wilhelm et al., 2014). Climate information is implemented into the CWP employing the Holdridge ecosystem classification concept based on the Holdridge Life Zones (HLZs; Holdridge et al., 1967), which provide a global classification of climatic
60 zones in relation to potential vegetation cover. The HLZ concept is commonly used as a tool for ecosystem mapping from various overlapping research communities (Lugo et al., 1999; Yue et al., 2001; Khatun et al., 2013; Szelepcsényi et al., 2014; Tatli and Dalfes, 2021). This paper gives a detailed documentation on the preparation of the PFT map - hereinafter referred to as "LANDMATE PFT" - within the Helmholtz Institute for Climate Service Science (HICSS) project "Modelling human LAND surface Modifications and its feedbacks on local and regional cliMATE" (LANDMATE). The LANDMATE PFT map
65 is prepared in close collaboration with the EURO-CORDEX Flagship Pilot Study Land Use and Climate Across Scales (FPS LUCAS; Rechid et al., 2017). Within the FPS LUCAS, RCM experiments are coordinated among an RCM ensemble to investigate the impact of LULC change for past climate and future climate scenarios. Through creation of LANDMATE PFT and the time series LUCAS LUC (Hoffmann et al., submitted), the need for improved LULC and LULC change representation among the FPS LUCAS RCM ensemble is met. For the preparation of LANDMATE PFT, we developed a CWP for the
70 translation of LULC classes of ESA-CCI into 16 PFTs according to the needs of regional climate modellers from all over Europe (Bontemps et al., 2013). A key issue to address in the map development process is the accuracy of LULC representation in the final product (Hartley et al., 2017). In order to assess the quality of the product, we compared the LANDMATE PFT map to a comprehensive ground truth database for large parts of the European Continent. The quality information derived from the assessment supports the RCM community in addressing and interpreting uncertainties caused by LULC representation in
75 RCMs. The general workflow and subsequently all utilized datasets are summarized in section 2 while the major steps of the CWP are listed in section 3. Section 4 introduces in detail the accuracy assessment procedure followed directly by the results in section 5. All CWTs and figures corresponding to the CWP and the accuracy assessment can be found in Appendix A and B.

2 Methods and data

80 The LANDMATE PFT map (Reinhart et al., 2021b) is a combination of multiple datasets and concepts created using well-established methods and in addition, by considering the expertise of regional climate modellers from all over Europe within the FPS LUCAS.

2.1 General workflow

The workflow to generate the LANDMATE PFT map is summarized in fig. 1, which also includes the steps to **generated**
85 **generate** the LUCAS LUC dataset further described in the companion paper by Hoffmann et al. (submitted). First, **the-ESA-CCI**
a high-resolution land cover map ([ESA-CCI LC](#), Sect. 2.2.1), which has a native resolution of ~ 300 m, is aggregated to the 0.1°
target resolution using SAGA GIS (Conrad et al., 2015). The target resolution results from the FPS LUCAS ensemble resolution
(i.e., EURO-CORDEX domain EUR-11) that is used for **LULC change-LULCC** impact studies in FPS LUCAS Phase II. The
90 LULC type information from the original product is preserved in fractions per 0.1° grid cell which is advantageous to common
majority resampling methods. The sum of PFT fractions in the whole dataset remains the same in all target resolutions, only
the distribution of fractions per grid cell changes depending on the target resolution.




Figure 1. The general workflow to generate LANDMATE PFT 2015 Version 1.0. This workflow is part of the workflow to generate the LUCAS LUC time series as introduced in the companion paper by Hoffmann et al. (submitted)

The ~~A climate dataset for Europe~~ (E-OBS ~~gridded climate data~~ (, Sect. 2.2.2) is utilized for the preparation of the ~~HLZ a climate zone~~ map over Europe (Holdridge Life Zones, Sect. 2.2.4). From ~~E-OBS the climate dataset~~, the ensemble ~~mean means~~ 2-meter-temperature and annual precipitation from 1950-2020 are used to create the ~~HLZ climate zone~~ map of 0.1° horizontal resolution which is further implemented in the CWTs to prepare the final LANDMATE PFT maps. For regions that are not covered by E-OBS, the respective data of the CRU dataset (Sect. 2.2.3) is used.

For ~~A CWT (Sect. 3) is created for each of the the 37 ESA-CCI land cover classes, an individual CWT is created (Sect. 3) that includes a unique translation for each used HLZ.~~ LC classes. Since the table has three dimensions (Land cover class, HLZ and PFT), it was necessary to prepare the individual tables that include unique translations for each HLZ. For example, table 1 shows the CWT for LC class 40 (Mosaic natural vegetation (tree, shrub, herbaceous cover)(>50%) / cropland (<50%)). For each HLZ in the first column, the LC class 40 is translated into varying fractions of the LANDMATE PFTs. For the example class that means an increasing tree fraction from the boreal to the tropical HLZs. Note that the numbers of the HLZs are corresponding to the HLZ numbers in figure 2.

Table 1. Cross-walking table for ESA-CCI LC class 40 - Mosaic natural vegetation (tree, shrub, herbaceous cover)(>50%) / cropland(<50%) as shown in Appendix A

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Holdridge Life Zone	Tree tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	Shrub evergreen	deciduous	Grass C3	C4	Special vegetation Tundra	Swamps	Crops crops	Non-vegetated urban	bare ground	
1,2											35	30	35			
3-5											30	35	35			
6											25	40	35			
7									60				40			
8					10				50				40			
9,10					15				45				40			
11					20				40				40			
12					30		20		10				40			
13				10	10				30				40			
14,15				20	20			10	10				40			
16				25	20				15				40			
17				25	25				10				40			
18				30	30								40			
19									60				40			
20							35		25				40			
21			20		15		15		10				40			
22			25		10		15		10				40			
23,24			20		20		20						40			
25									60				40			
26							30		30				40			
27		10					50						40			
28		40					20						40			
29	40						20						40			
30	50						10						40			

The translation process is based on Wilhelm et al. (2014) where the translation of the Global Land Cover (GLC) 2006 to the 16 REMO-iMOVE PFTs is described. Since the nomenclature of GLC 2006 and ESA-CCI LC are similar and based on the same classification system some of the CWTs were initially adopted from (Wilhelm et al., 2014). For the more diverse ESA-CCI LC classes new CWTs need to be created. The new CWTs follow the translation of Poulter et al. (2015) (~~ESA-CCI PFTs~~ ESA POULTER) but were carefully revised and modified during the process. ~~This revision of the CWTs is supported by reference data and visual satellite image interpretation~~ After application of the CWP, an additional map of potential C3/C4 grass vegetation (NACP MsTMIP, Sect. 2.2.6 is used to divide the grass PFT fractions. The quality of the LANDMATE PFT

dataset is finally assessed by comparison to a comprehensive ground truth database ([LUCAS land use and land cover survey](#), Sect. 4).

2.2 Datasets & concepts

2.2.1 ESA-CCI LC

115 The European Space Agency Climate Change Initiative (ESA-CCI) provides continuous global land cover maps (ESA-CCI
 LC) on ~300 m horizontal grid resolution. The ESA-CCI LC maps are available for download in annual time steps for the
 years 1992-2018 (ESA, 2017). The classification of the LC maps follows the United Nations Land Cover Classification System
 (UN-LCCS) protocol (Di Gregorio, 2005) and consists of 22 level 1 classes and 14 additional level 2 classes, which include
 regional specifications. More information on ESA-CCI LC data processing can be found at [maps.elie.ucl.ac.be/CCI/viewer/
 120 download/ESACCI-LC-Ph2-PUGv2_2.0.pdf](http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf). An overview of the satellite missions involved in the production of ESA-CCI
 LC is given in table 2. Besides systematic global validation efforts (ESA, 2017; Hua et al., 2018), a few regional approaches
 investigated the quality of ESA-CCI LC over Europe (Vilar et al., 2019; Reinhart et al., 2021a).

Table 2. Satellite missions involved in the production of ESA-CCI LC according to ESA (2017)

Time period		Satellite product
Baseline	Production	MERIS FR/RR ¹ global SR ² composites
2003-2012		
1992-1999		Baseline 10-year global map; AVHRR ³ global SR composites for back-dating baseline
1999-2013		Baseline 10-year global map; SPOT-VGT ⁴ global SR composites for up and back-dating the baseline; PROBA-V ⁵ global SR composites at 300 m
2013-2015		Baseline 10-year global map; PROBA-V global SR composites at 1 km for years 2014 and 2015 for updating the baseline; PROBA-V time series at 300 m
Since 2016		Sentinel-3 OLCI and SLSTR ⁶ 7-day composites

¹MEdium Resolution Imaging Spectrometer Full Resolution/Reduced Resolution (ESA, 2002)

²Surface Reflectance

³Advanced Very-High-Resolution Radiometer (Hastings and Emery, 1992)

⁴SPOT Vegetation satellite program (Maisongrande et al., 2004)

⁵Project for On-Board Autonomy - Vegetation (Dierckx et al., 2014)

⁶Ocean and Land Colour Instrument (OLCI) and Sea and Land Surface Temperature Radiometer (SLSTR) (Donlon et al., 2012)

2.2.2 E-OBS Climate data

The E-OBS dataset (Cornes et al., 2018) is a daily gridded observational dataset, derived from station observations from European countries covering the period from 1950 to 2020. The point observations are interpolated using a spline method with random perturbations in order to produce an ensemble of realizations. For the creation of the HLZs that are used for the conversion of ESA-CCI LC classes to PFTs (Section 2.2.5), the ensemble mean of the 2-meter-temperature (TG) and precipitation (RR) on a regular 0.1° grid from E-OBS version 19.0e is used. It covers most of Europe, some parts of the Middle East and a narrow strip of Northern Africa.

2.2.3 CRU

The Climate Research Unit (CRU) TS 4.03 dataset is a global gridded high-resolution climate dataset based on station observations produced and maintained by the CRU of the University of East Anglia (Harris et al., 2014). The dataset provides global monthly means of climate parameters at 0.5° resolution from 1901 to 2019. In order to achieve the target resolution of 0.1° for the global LANDMATE PFT maps, the CRU climate data is downscaled using bilinear interpolation. Following Hoffmann et al. (2016), distance-weighted interpolation was applied to the atmospheric observation dataset CRU to extrapolate the climate data to the coastlines of the ESA-CCI LC maps in order to compensate for the different land-sea-masks of the products. The CRU climate dataset was used within this application for regions where E-OBS is not available. The bilinear interpolation of E-OBS caused minor issues on coastlines and small islands all over the research area, where this interpolation method was not able to correctly account for the resolution differences of the 0.1° EOBS and 0.018° LANDMATE PFT land-sea masks, respectively. The issue caused by the large resolution difference is fixed with a preceding extrapolation of the climate data along coastlines and islands in the LANDMATE PFT map Version 1.1 that is currently being prepared. Since the interpolation issue only affected a negligible amount of LANDMATE PFT cells the validation measures are not affected by this issue in a noticeable way.

2.2.4 Holdridge Life Zones

The Holdridge Life Zone (HLZ) concept was initially developed in 1967 (Holdridge et al., 1967) to define all divisions of the global biosphere, depending on the relation of biotemperature (average of monthly temperature above 0°C ; since plant activities are idle below freezing, all values below 0°C are adjusted to 0°C), mean annual precipitation and ratio of potential evapotranspiration to mean annual precipitation. By combining threshold values of biotemperature and annual rainfall, the 38 HLZs are created (Table 3). In the present analysis, the ~~tropical and subtropical~~ subtropical and warm temperate as well as the polar and subpolar HLZs are merged. Through the merging of the aforementioned HLZs, 30 individual HLZs in total are available for the creation of the European HLZ map (Fig. 2). The dynamic character of the specific quantitative ranges of the long-term means of the utilized climate parameters make the HLZ classification more flexible than other available global ecosystem classifications and therefore makes the HLZs most suitable for the application presented in this article. In addition the requirement for input data is relatively low.

Table 3. The Holdridge Life Zones following (Holdridge et al., 1967).

	Precipitation [mm]					
	<125	125 to <250	250 to <500	500 to <1000	1000 to <2000	>2000
<3	Subpolar dry tundra	Subpolar moist tundra	Subpolar wet tundra	Subpolar rain tundra	-	-
3 to <6	Boreal desert	Boreal dry shrub	Boreal moist forest	Boreal wet forest	Boreal rain forest	-
6 to <12	Cool temperate desert	Cool temperate desert shrub	Cool temperate steppe	Cool temperate moist forest	Cool temperate wet forest	Cool temperate rain forest
12 to <18	Warm temperate desert	Warm temperate desert scrub	Warm temperate thorn steppe/woodland	Warm temperate dry forest	Warm temperate moist forest	Warm temperate wet/rain forest
18 to <24	Subtropical desert	Subtropical desert shrub	Subtropical thorny steppe/woodland	Subtropical dry forest	Subtropical moist forest	Subtropical wet/rain forest
>24	Tropical desert	Tropical desert shrub	Tropical thorny woodland	Tropical very dry forest	Tropical dry forest	Tropical moist/wet/rain forest





Figure 2. Holdridge Life Zones map for the extent of LANDMATE PFT

155 In the past, the HLZ concept was not only found useful for global applications but successfully implemented especially for regional mapping approaches due to its ability to capture regional climate features with the support of bioclimatic variables (Daly et al., 2003; Tatli and Dalfes, 2016). Further, the HLZ concept was used for LULC change predictions, such as land use impact assessments, related to current and future climate change scenarios (Chen et al., 2003; Skov and Svenning, 2004; Yue et al., 2006; Saad et al., 2013; Szelepcsényi et al., 2018). With the implementation of climate data through the HLZ concept, 160 the resulting PFT maps become more detailed and can be customized to individual regions without losing global consistency.

2.2.5 Plant Functional Types

~~Table ??~~ [Figure 3](#) shows the LANDMATE PFTs that are based on the PFTs introduced by Wilhelm et al. (2014). The implementation of an irrigated cropland PFT (PFT 14) that is currently being developed within the HICSS project LANDMATE will be implemented in a later version of the dataset. In the initial version that is presented in this article, all cropland proportions are assigned to the cropland PFT (PFT 13). ~~LUCAS plant functional types based on Wilhelm et al. (2014) with modified crop types.~~

PFTs Names 1-Tropical broadleaf evergreen trees 2-Tropical deciduous trees 3-Temperate broadleaf evergreen trees 4-Temperate deciduous trees 5-Evergreen coniferous trees 6-Deciduous coniferous trees 7-Coniferous shrubs 8-Deciduous shrubs 9-C3-grass 10-C4 grass 11-Tundra 12-Swamp 13-Non-irrigated-croplands 14-Irrigated-croplands 15-Urban 16-Bare




Figure 3. LANDMATE PFT map for Europe for 2015 (a). Below a map section of the Alpine region shows an example of the resolution difference between LANDMATE PFT 0.1 (b) and LANDMATE PFT 0.018 (c). LANDMATE PFT 0.018 is used in the present accuracy assessment. For improved visualization all maps show the majority PFT per grid cell.

2.2.6 Potential C4 grass fraction NACP MsTMIP

The initial land cover map from the ESA-CCI LC does not provide a distinction between C3 and C4 grassland. Therefore, an additional product is used after applying the CWP. The map from the North American Carbon Program Multi-scale Synthesis and Terrestrial Model Intercomparison Project (NACP MsTMIP; Wei et al., 2014) is constructed based on the synergetic land cover product (SYNMAP) by Jung et al. (2006). SYNMAP is a combination of multiple high-resolution LULC products using a fuzzy agreement approach. The NACP MsTMIP map uses the grassland fractions from the SYNMAP product ~~and the~~. The potential C4 grass distribution is estimated-supported-by-generated by Wei et al. (2014) employing the well established method introduced by Still et al. (2003), which is based on the growing season temperature based-on-and rainfall, in combination with present climate conditions (Wei et al., 2014). The from the global CRU-NCEP dataset. The potential C4 grass map is provided in-on a 0.5° horizontal grid resolution-for the period from 1801 to 2010. For the preparation of LANDMATE PFT the NACP MsTMIP map of 2010 is used. The LANDMATE PFT grass fraction is split up into C4 and C3 grasses by multiplying the it with the potential C4 vegetation fraction and (1 - potential C4 vegetation fraction), respectively.

2.3 LUCAS - land use and land cover survey

The harmonized LUCAS *in situ* land cover and use database for field surveys from 2006 to 2018 (d'Andrimont et al., 2020) is the most consistent ground truth database for the European Continent. The survey was carried out at three-yearly intervals between 2006 and 2018. The systematic sampling design of the survey consists of a theoretical, regular grid over the European Continent with ~2 km grid size. The reference point locations are the corner points of the theoretical grid. Not all locations within the survey were easily accessible. Therefore, the survey is supported by *in situ* photo interpretation, *in-office* photo interpretation and satellite data in the latest time steps 2015 and 2018 (table 4). However, the main proportion of the reference points was recorded through location visits at all time steps, which makes this land survey the most reliable and consistent ground truth database for Europe.

Table 4. Number and recording method of reference points in the LUCAS land cover and use database per timestep.

Year	Reference points	<i>in situ</i>	<i>in situ</i> PI ⁷	<i>in-office</i> PI ⁸	GT ⁹ [%]
2006	168401	155238	13163		92.18
2009	234623	175029	59594		74.6
2012	270272	243603	26669		90.13
2015	340143	242823	25254	71970	71.39
2018	337854	215120	22894	99803	63.67

⁷Photo interpretation close to the reference location

⁸Photo interpretation with supporting data, such as satellite images

⁹Ground truth

The extent of the LUCAS survey was increased over time. The 2006 survey covered 11 countries while the 2018 map covers large parts of the European Continent with 28 countries. Throughout the survey, the ground truth data has been continuously checked for quality and plausibility. For the accuracy assessment of the LANDMATE PFT map the ground truth points of the year 2015 are employed (Sect. 4). In order to avoid confusion between the FPS LUCAS and the LUCAS ground truth dataset, the latter will be further referred to as **Ground Truth Survey** or **GT-SUR**.

3 Cross-walking procedure - ESA-CCI LC classes to PFTs

The CWP from ESA-CCI LC classes to PFTs presented in this article is based generally on (1) the translation introduced by Poulter et al. (2015) and (2) the translation by Wilhelm et al. (2014). Both translations are not just combined with each other but modified using additional data. The following sections introduce the PFTs of LANDMATE PFT aggregated into groups general LULC types and give an overview of the decisions on modifications that are made during the production process based on literature and additional data. ~~The final LANDMATE PFT map is shown in fig. 3.~~

3.1 Trees and shrubs, tropical and temperate | PFT 1-8

The LANDMATE PFTs are more diversified regarding tree-PFTs than the generic ~~ESA-CCI PFTs. The expansion of tree-PFTs to six in total was done at the expense of two~~ ESA POULTER PFTs. While the generic ESA POULTER PFTs have four shrub-PFTs, the LANDMATE dataset has only two while the tree-PFT count was increased to six. The increase of tree-PFT diversity is done in order to address the strong biogeophysical impacts of forested areas on regional and local climate, such as decreased albedo and increased roughness length (Bright et al., 2015). The effects of forested areas on near-surface climate are distinctively different to the effects of shrub or grass covered areas, and are also highly depending on tree species composition and latitudinal range (Bonan, 2008; Richardson et al., 2013). Another reason for the six tree-PFTs is the intended use of the PFT maps in RCMs. In the Land Surface Models (LSMs) of current generation RCMs, where a distinction is rather made between different tree or tree community types than between different shrub types. Therefore and with regard to the implementation process that needs to be done for each RCM individually, an increase in the number of tree-PFTs and a decrease in the number of shrub-PFTs is considered to be convenient. Accordingly, the tree and shrub proportions were distributed following both, the needleleaf and broadleaf definitions of the ESA-CCI LC classes as well as the HLZ map, where the HLZ map was decisive for an assignment of forest proportions to the temperate or tropical tree-PFT, respectively. Following a comparison with different forest datasets over Europe (not shown), the tree proportions in the translation of the mixed land cover classes (e.g. ~~lass-class~~ 61 - Tree cover, broadleaved, deciduous, closed (>40%)) are increased to be in line with the indicated overall forest amount over Europe.

3.2 Grassland | PFT 9 & 10

The generic ~~ESA-CCI~~ ESA POULTER PFTs include a natural grassland- and a managed grassland-PFT to include grassland and cropland respectively. The LANDMATE PFTs include two grassland-PFTs, distinguishing between C3 and C4 grass. The

contrasting photosynthetic pathways and therefore contrasting synthetic response to CO₂ and temperature determine specific ecosystem functions for both PFTs respectively. The main differences are found in global terrestrial productivity and water cycling (Lattanzi, 2010; Pau et al., 2013). The translation from the LULC classes that contain grassland proportions into C3 or C4 grass-PFTs respectively is supported by a map of potential C4 vegetation by Wei et al. (2014) where the potential global distribution of C4 is estimated using bioclimatic parameters (Sect. 2.2.6).

3.3 Tundra and swamps | PFT 11 & 12

The specific vegetation PFTs tundra and swamps are treated individually in LANDMATE PFT. Tundra is mostly used for the polar and subpolar HLZs, where the climatic conditions require a clear distinction of the land surface properties to the boreal and temperate regions regarding exchange and feedback processes with the atmosphere (Thompson et al., 2004). Chapin Iii et al. (2000) further suggest a differentiation of vegetation composition within these northern vegetation communities, which can also be realized using the introduced [translationCWP](#). The swamp-PFT is mostly used for translating the ESA-CCI LC mosaic tree/shrub/herbaceous classes and also partly for the flooded tree cover classes in most of the HLZs. Swamps occur mainly in the boreal and polar regions [in the European domain](#).

3.4 Cropland | PFT 13 & 14

Currently, two cropland-PFTs are defined in the LANDMATE PFT map. The cropland-PFT (PFT 13, see table ??) includes all managed, agricultural land surface proportions. The uncertainties of the translation of the ESA-CCI cropland classes and mixed cropland classes into the cropland-PFTs was investigated by Li et al. (2018) where the comparison of LULC change in the [ESA-CCI-ESA POULTER](#) PFT maps against other LULC products showed inconsistencies between global trends and geographical patterns between the products. However, Li et al. (2018) provide a modified CWT that was adjusted in regard to an improved knowledge base on how to translate LULC classes into PFTs for climate models. Particular focus is laid on mosaic classes and the sparsely vegetated classes of which appear numerous in ESA-CCI LC. Therefore, the [translation-CWP](#) from Li et al. (2018) for cropland is adopted into the present CWP.

The irrigated cropland-PFT (PFT 14, [see table ??fig 3](#)) is currently empty in the LANDMATE PFT map Version 1.0. This decision is made following intense research on available irrigation information. The ESA-CCI LC map that is used as initial input contains an "irrigated cropland" class but this information was not used in the process. The investigation on irrigated areas included the comparison of ESA-CCI LC to other products that are available, such as the irrigation map from the FAO (Siebert et al., 2005). Although the ESA-CCI LC quality assessment shows a very good agreement of the ESA-CCI LC irrigated cropland with the validation database (ESA, 2017), the comparison showed considerable differences between the products. The success of detection of irrigated areas is highly dependent on the correct detection of the crop types to infer the water needs of the respective crops, on atmospheric and environmental conditions and on the availability of multi-temporal, high resolution imagery (Bégué et al., 2018; Karthikeyan et al., 2020). Further, most remote sensing applications depend highly on ground truth data and local knowledge. Applications using different satellite imagery to detect agricultural management practices, such as irrigation, are only successfully tested and applied in local spatial units (Rufin et al., 2019; Ottosen et al., 2019). Therefore,

the irrigated cropland PFT remains unoccupied for now. Nevertheless, PFT 14 is defined within LANDMATE PFT Version 1.0 for the purpose of adding irrigated LULC fractions in the future. For the long term LUCAS LUC dataset (Hoffmann et al., submitted) which is extended backward and forward based on the LANDMATE PFT map for Europe 2015, irrigated cropland areas are already implemented following the irrigated area definition of the Land Use Harmonization (LUH2) dataset (Hurt et al., 2011).

3.5 Non-vegetated | PFT 15 & 16

The non vegetated-PFTs in the LANDMATE PFT dataset are urban and bare. The urban grid cells from ESA-CCI LC are directly translated into urban fractions for all HLZs in the CWP. The same applies for all bare ground proportions that are translated fully into the bare-PFT. In addition, the ESA-CCI LC mixed classes are split up and the bare ground proportions within the mixed classes are added to the bare-PFT. The explicit treatment of urban areas and especially differentiation from bare ground provides the possibility to resolve urban surface characteristics in RCMs. The treatment of urban areas as a slab surface or as an equal to rock surface as done in several RCM approaches cannot account for the complex geobiophysical processes associated with an urban agglomeration (Daniel et al., 2019; Belda et al., 2018). Due to the distinction of the two surface types, the LANDMATE PFT map can be used for impact studies with an urban focus.

3.6 Water, permanent snow & ice

The LANDMATE PFTs do not include individual PFT definitions for water and snow/ice respectively. Regarding the water representation, most currently used RCMs are utilizing a land-sea-mask to account for oceans and inland water areas. Therefore, an explicit definition of water as individual PFT has not been implemented. Consequently, ~~water grid cells~~ all water fractions, such as marine water, lakes and rivers are set to no data.^z In the present translation, the snow/ice grid cells from ESA-CCI land cover are translated into bare-PFT following Wilhelm et al. (2014). ~~LANDMATE PFT map for Europe for 2015 (a). Below a map section of the Alpine region shows an example of the resolution difference between LANDMATE PFT 0.1 (b) and LANDMATE PFT 0.018 (c). LANDMATE PFT 0.018 is used in the present accuracy assessment. For improved visualization all maps show the majority PFT per grid cell.~~

4 Quality assessment of the LANDMATE PFT map

The LANDMATE PFT map is based on the ESA-CCI LC map which was quality checked and compared to similar LULC products on a global (ESA, 2017; Yang et al., 2017; Hua et al., 2018; Li et al., 2018) and regional level (Reinhart et al., 2021a; Vilar et al., 2019). However, the translation from LULC classes to PFTs necessarily results in change of the map. The final product, the LANDMATE PFT map, is intended to be used in RCMs, which means the quality of the final product must be assessed in addition to the available quality assessments of the initial ESA-CCI LC map. In order to overcome the resolution difference, which is non negligible between LANDMATE PFT and the reference data GT-SUR, the LANDMATE PFT map is prepared on 0.018° horizontal resolution, which corresponds closely to the 2 km theoretical grid of GT-SUR.

The design of such a quality assessment of a large scale map product is not trivial, especially since the map product itself and the reference data are often different in structure and nomenclature, given that ground truth reference data is mostly collected as point data and independently from the assessed map product Foody (2002); Wulder et al. (2006); Olofsson et al. (2014). In order to produce reliable quality information for LANDMATE PFT, the present assessment follows closely the well established good-practice recommendations. Nevertheless, adjustments are done to account for the fractional structure of LANDMATE PFT. Section 4.2 provides additional information on the requirements of a "good practice" accuracy assessment, the key components and the selected sampling design and metrics.

290 **4.1 Research area**

The coverage of GT-SUR in the year 2015 includes 28 countries which are highlighted in dark grey in fig. 4.




Figure 4. Coverage of the Land Use and Coverage Area frame Survey (LUCAS) for reference year 2015 (top). The lower figure shows the points and LULC [group-type](#) representation within the grid cell highlighted in black color in the top map as an example for the whole research area.

The total number of GT-SUR points for 2015 is 340,143. Out of these points, 338,619 points (~99.55%) are covered with valid LANDMATE PFT grid cells of the assessed LULC [groups-types](#) and can be used in the analysis. Countries located within the contiguous area but missing in the assessment are Switzerland, Norway, the Russian Oblast Kaliningrad, Bosnia and Herzegovina, Montenegro, Albania, Serbia, Kosovo, North Macedonia, and Belarus. Figure 4 also shows the 2.5° grid that was used for the analysis of the accuracy assessment results (Sect. 5). Due to the fine scale and the high number of points over the whole research area, the visualization of the spatial analyses on continental scale is challenging. Therefore, the research area is split up through an overlay of a 2.5° grid (as shown in fig. 4). The overall and class-wise accuracy results for all points within

each 2.5° grid cell are aggregated in order to identify large scale spatial quality differences for the analyzed LULC groupstypes.
300 Additionally, the total number of points for each LULC group-type per grid cell are displayed in section 5.

4.2 Accuracy assessment - background & design

The key components of the accuracy assessment of a large-scale land cover product are **objective, sampling design, response design** and the final **analyses and estimation** (Wulder et al., 2006). All of the key components have great impact on the quality of the assessment and further, on the final metrics, especially in the present assessment, where reference and assessed
305 dataset differ widely in structure. LANDMATE PFT is a gridded dataset with fractional LULC classes but no information on the subgrid location within the grid cell. Other than that, the points of GT-SUR have fixed locations expressed through exact coordinates, but no (exact) information on the spatial extent of this class. Another challenge is the fractional structure of LANDMATE PFT itself, where one unit (grid cell) possibly contains multiple fractions. Therefore, the design of the accuracy assessment needs to be customized to the **objective**, which is to determine the overall quality of the LANDMATE PFT map for
310 Europe 2015 as well as the quality of individual LULC type representation within the map in order to derive recommendations for the use of LANDMATE PFTs in RCMs.

When it comes to the **sampling design**, sampling size, spatial distribution of the respective sample and the representation of each LULC group-type or class within the sample are crucial to produce reliable quality information about a LULC product (Stehman, 2009). ~~However, the~~ The collection of ground truth data is a rather expensive procedure regarding time and money,
315 which needs to be considered during the process. ~~The sample size is therefore a compromise size and cost. In~~ However, in the present assessment ~~we are able to rely on~~ an existing ground truth database containing over 340,000 records ~~is used as reference~~ , which eliminates the possible issue of a too small ~~sample size~~ reference database. It is also known that all assessed LULC groups-types are represented in a sufficiently high number (Table 6). Nevertheless, the present assessment is a special case situation with every unit of LANDMATE PFT containing more than one LULC group-type potentially. Therefore, the subsets
320 are selected through application of a filter to capture the map accuracy in a way that accounts for the fractional structure within the grid cells in the LANDMATE PFT map (see section 4.2.1).

The **response design** deals with the spatial support regions (SSR) and the labelling protocol or classification harmonization. The SSR is a buffer region around a sampling unit that is selected to account for small-scale landscape heterogeneity that is likely not captured by larger scale map products. In the present case, the sampling design is selected in a way that the grid
325 cells of LANDMATE PFT serve as SSR for each GT-SUR point. A fraction is not located precisely at one location within the respective grid cell but evenly distributed over the whole grid cell. Assuming, the uniformly distributed fraction can occur in small patches or in one large patch within the grid cell, the whole grid cell is defined as SSR for the respective LULC group-type. The labelling protocol needs to be determined to deal with the different legends of the reference and the assessed map. The harmonization of legends is selected in regard to the objective of the respective assessment, as in this case, to provide
330 information about the quality of representation of the most dominant LULC types in LANDMATE PFT. The labelling protocol used in the present assessment is summarized in table 5.

The **analyses and estimation** used are error matrices, that give an overview of the overall and LULC group-wise-type-wise accuracy of the LANDMATE PFT map. For both resolutions of LANDMATE PFT, the error matrices and the resulting accuracy measures overall accuracy (OA), producer's accuracy (PA) and user's accuracy (UA) are calculated, where PA and OA are calculated group-wise. The error matrix is a cross-tabulation between map and reference of the size $q \times q$, where q stands for the number of land cover classes or groups. The map classes are placed in the rows and the reference classes in the columns so that the diagonal of the matrix gives the sum of the correctly classified map units. The off-diagonal cell values represent the disagreement between the map and the reference. The overall accuracy is calculated according to equation 1:

$$OA_i = \frac{\sum_{i=1}^q n_{ii}}{n} * 100 \quad (1)$$

The sum of the agreeing diagonal elements n_{ii} of all LULC groups-types is divided by the number of all observations n . The PA represents the accuracy from the view of the map producer. The PA stands for the probability, that a LULC feature in the reference is classified as the respective feature by the map. The PA is calculated using equation 2 where the number of correctly classified units per LULC group-type n_{ii} is divided by the total number of LULC group-type occurrences of the reference n_{+i} :

$$PA_i = \frac{n_{ii}}{n_{+i}} * 100 \quad (2)$$

While the PA gives the proportion of features in the reference that are actually represented as those in the produced map, the UA is the accuracy from the perspective of the map user. It is the probability of a feature classified as such in the map is actually present in the reference. The UA is calculated using equation 3, where the number of correctly classified pixels n_{ii} per LULC group-type is divided by the row sum $n_{i+} = \sum_{j=1}^p n_{ji}$:

$$UA_i = \frac{n_{ii}}{n_{i+}} * 100 \quad (3)$$

4.2.1 Dataset harmonization & filter

The quality assessment is done assigning the PFT type with the maximum fraction per grid cell to the GT-SUR points located within the respective grid cell. The classifications of both datasets ~~need to be harmonized in~~ are harmonized as shown in table 5 where the focus is laid on the main LULC types in order to make the comparison as detailed as possible but also to be able to produce reliable and robust results for the RCM community. ~~For the analysis, the classifications of LANDMATE PFT and the GT-SUR are harmonized as shown in table 5-~~

Table 5. Classification harmonization between LANDMATE PFT map and GT-SUR

GT-SUR LC group	GT-SUR group name	LANDMATE PFT number	LANDMATE PFT name	Harmonization group number	Harmonization name
A	Artificial Land	15	Urban	1	URBAN
B	Cropland	13 14	Non-irrigated Crops Irrigated crops	2	CROPLAND
C	Woodland	1 2 3 4 5 6	Tropical broadleaf evergreen trees Tropical deciduous trees Temperate broadleaf evergreen trees Temperate deciduous trees Evergreen coniferous trees Evergreen deciduous trees	3	WOODLAND
D	Shrubland	7 8	Coniferous shrubs Deciduous shrubs	4	SHRUBLAND
E	Grassland	9 10	C3 Grass C4 Grass	5	GRASSLAND
F	Bare land	16	Bare	6	BARE AREAS
G	Water	11	Tundra	7	OTHER
H	Wetlands	12	Swamps		
Other	Marine areas				

The LULC ~~groups~~ types URBAN, CROPLAND, WOODLAND, SHRUBLAND, GRASSLAND, and BARE AREAS are harmonized without applying modifications to the classifications. The LANDMATE PFTs can easily be grouped or directly adopted while the GT-SUR level one classification (letters A-H) is completely adopted into the harmonized groups. ~~The LANDMATE PFT map is a product developed for the use in RCMs.~~ In general, RCMs implement a dedicated land-sea-mask to determine aquatic areas for both, inland and marine water. Therefore, the categories ~~WATER and MARINE areas are neglected in the analyses. The LANDMATE PFTs "Tundra" and "Swamp" can not be assigned with a~~ Water and Marine areas are not further analyzed. Since the LULC types Tundra, Swamps (LANDMATE PFT) and Wetlands (GT-SUR) cannot be harmonized with sufficient agreement to the GT-SUR ~~class definitions. Therefore, the LULC type definitions, the LULC types~~ are also not further analyzed in the assessment. Thus, the LULC types Water, Marine areas and Wetlands (GT-SUR ~~groups~~

~~water, wetlands and marine areas as well as the LANDMATE PFTs~~). Tundra and Swamps (LANDMATE PFT) are merged into the ~~group "OTHER" for the assessment~~LULC type OTHER. Although the group cannot be evaluated regarding the quality of the LANDMATE PFT map, the group needs to be involved in the assessment to keep the numbers in the assessment correct and reliable for all other groups. ~~However, as it is shown in table 6, only a minor amount of points/cells is affected.~~

370 Both datasets are provided in a regular Gaussian grid (WGS84 EPSG:4326) so that no reprojection of the datasets needs to be done for the comparison. ~~The descriptive statistics for each LULC group for the reference-~~




Figure 5. Overall cell count of LANDMATE PFT as function of the threshold for minimum coverage of the respective dominant LULC type.

The LANDMATE PFT dataset includes multiple LULC fractions per grid cell. Accordingly, the area proportion of the dominant LULC type varies widely and thus the likelihood that the ~~GT-SUR and the LANDMATE PFT dataset are summarized in table 6.~~ point sample falls within this area. The grid cells are grouped by minimum coverage of the dominant LULC type from 0.1 to 1 where 0.1 means a minimum coverage of 10 % and 1 means full coverage of the dominant LULC type. The coexisting fractions are not located in particular parts within a grid cell but equally distributed while the GT-SUR points have fixed locations on the map. With the applied grouping of the cells dependent on the minimum coverage of the dominant LULC type, the influence of grid cell heterogeneity on accuracy figures is investigated within the assessment. Figure 5 shows the number of LANDMATE PFT cells within the respective groups. While the groups 0.1-0.7 each contain over one third of the total LANDMATE PFT cells, the groups with a minimum coverage threshold of 0.8 and higher contain a number of cells that is rather not sufficient to represent the whole research area.

375
380
385
Beside the total number of LANDMATE PFT cells in the analysis, diversity among the represented LULC types is important. The right column of table 6 shows the number of LANDMATE PFT cells where the respective LULC type (left column) is dominant. The table is not grouped by minimum coverage but by LULC type and shows that each assessed LULC type is represented in a sufficiently high number when only the cells with dominant coverage (regardless the total proportion) are considered for each LULC type.

Table 6. General information on data in the comparison

LULC group type ¹⁰	GT-SUR ¹¹	LANDMATE PFT 0.018 ^{o12}	Dominant LAND- MATE PFT 0.018 ^o <u>dominant</u> ¹³
URBAN	14,393	65,000	7,577
CROPLAND	83,295	248,301	136,970
WOODLAND	124,374	277,290	124,437
SHRUBLAND	27,298	302,035	19,790
GRASSLAND	66,541	333,948	44,244
BARE AREAS	10,395	31,756	4,148
OTHER	12,340	28,823	1,470
Sums	338,636		338,636

The LANDMATE PFT dataset includes multiple LULC fractions per grid cell. Accordingly, the area proportion of the dominant LULC group varies widely and thus the likelihood that the GT-SUR point sample falls within this area. The filter applied is categorizing the grid cells regarding the proportion of the dominant LULC group, the higher the threshold, the stricter the filter and the more likely a specific sample falls into the subgrid fraction of the dominant class. The filter set numbers 1-10 are representing the cells containing a minimum of 10 – 100% of the dominant LULC group according to table ???. Filter sets with varying dominant LULC group share per grid cell from >10% to 100% Filter set number fraction-size of dominant LULC group LANDMATE PFT cells within filter set1 >10 % 338619 2 >20 % 338619 3 >30 % 336703 4 >40 % 311238 5 >50 % 259073 6 >60 % 203343 7 >70 % 137412 8 >80 % 74765 9 >90 % 26993 10 100 % 1449

4.3 Results

5 Results

In order to show the impact of the applied spatial filter, the spatial distribution of agreement and disagreement grid cell heterogeneity of LANDMATE PFT, the agreement of LANDMATE PFT with the reference GT-SUR is investigated. The point counts for each threshold for minimum coverage (0.1-1) of the dominant LULC type. For visualization of the spatial analysis, the point count and percentage agreement are aggregated and averaged, respectively, with the reference dataset are aggregated per 2.5° grid cell. After giving an overview over the overall accuracy measures the individual LULC group results

¹⁰LULC type analyzed in the quality assessment

¹¹Number of GT-SUR points per LULC type

¹²total number of grid cells in LANDMATE PFT that have a share >0% of the respective LULC type

¹³Number of cells where the LULC type is dominant in LANDMATE PFT 0.018°

are discussed in the following subsections. Note that due to the low overall point count as shown in table 6, the LULC groups SHRUBLAND and BARE AREAS are discussed together in section ??.

Figure 6 shows the spatial distribution of the filter sets over Europe while fig. ?? shows the overall accuracy for the filter sets. cell of the auxiliary grid, which was established as most useful for visualization of the results.

The distribution of the varying dominant LULC group filter sets over the research area in Europe. Since the >10% and the >20% filter set share the same number of points the >10% filter set is not shown.

Overall accuracy for the full domain of the 10 filter sets as introduced in table ?? as function of filter set size. Filter set numbers are shown in grey boxes.

In order to be able to capture the LULC group diversity and distribution characteristics of spatial distribution of the quality of the LULC type representation within LANDMATE PFT, the filter set assessed cells must be distributed well over the respective research area and contain a sufficiently large proportion of the total cells cell count of each LULC type. Figure 6 shows that the filter sets are distributed reasonably well up to filter set 7. Filter set 8 the distribution and count of cells grouped by threshold for minimum coverage according to fig. 5. The maps show that the groups with a threshold lower than 0.7 are distributed very well over the research area. Each region is covered with a sufficient amount of LANDMATE PFT grid cells that can be compared to the respective GT-SUR points. The 0.8 group shows a quite patchy pattern and a strongly decreasing sample number in Northern Europe. Within filter set 9 For the 0.9 group, the patchy pattern of low sample count and low number of cells per 2.5° grid cell spreads over the whole continent. While filter set 9 research area. While the 0.9 group could still be used for evaluation of LANDMATE PFT for limited regions in Europe, filter set 10 the group only containing cells with 100 % coverage of one LULC type (map 1) is clearly not evaluable due to the overall small sample cell count (< 1500). This pattern is also found for filter sets 9 & 10 of the individual LULC groups. The filter set sizes as well as the applied filter itself have direct impact on the OA shown in fig. ??.

The decreasing OA towards the higher sample count is an effect of the LANDMATE PFT grid cell heterogeneity representation in each filter set. Filter set 1-3 include all LANDMATE PFT grid cells where the dominant LULC group occupies a minimum of 10% to 30% respectively. Therefore, the probability that the GT-SUR point sample within the respective grid cell represents a location that is occupied by one of Figure 7 gives an overview of the cell count per group for each individual LULC type.




Figure 6. The distribution of the LANDMATE PFT cells grouped by threshold for minimum coverage of the respective dominant LULC type over the research area in Europe according to fig. 5. The same number of LANDMATE PFT cells falls into the groups with a minimum coverage of 0.1 and 0.2. Therefore, the 0.1 group is not shown in the figure.

For CROPLAND, WOODLAND and GRASSLAND, the threshold for minimum coverage of the respective dominant LULC type has a strong influence on the total cell count within each group while for URBAN and BARE AREAS, the cell count remains similar up to the 0.6 group. For SHRUBLAND, the cell count decreases strongly from the 0.4 group upwards. The curve characteristics suggest that the LULC types CROPLAND, WOODLAND, and GRASSLAND have a higher proportion of cells with a relatively low dominant coverage but since they are the three most populated LULC types overall (see tab. 6) the proportions are comparable to the other three groups. Figure 8 shows the PA for all LULC types dependent on the threshold for minimum coverage of the dominant LULC type including the overall accuracy for all LULC types together (dark grey line). While the overall accuracy is relatively independent of the threshold for minimum coverage, some LULC types are affected strongly. For WOODLAND, PA decreases rapidly for the 0.8 group. Considered that the non-dominant LULC groups is relatively high. The applied filter accounts for cell count for this group does decrease noticeably from 0.7 to 0.8 (fig. 7, the low PA is likely a result of this too low cell count. The PAs for GRASSLAND and SHRUBLAND remain almost constant but at a lower level compared to the impact of the structure difference of the two datasets. The higher probability of agreement is reflected in the increasing OA for the samples that include only grid cells with an occupation of 50% or more of the dominant LULC group other groups groups.




Figure 7. Cell count of LANDMATE PFT per LULC type as function of the threshold for minimum coverage of the respective dominant LULC type.




Figure 8. Producer's accuracy for LANDMATE PFT as function of the threshold for minimum coverage of the respective dominant LULC type.

Figure 9 shows the highest UA for WOODLAND and the lowest for SHRUBLAND while all other LULC types range in between. The threshold for minimum coverage of the individual LULC types has slightly more influence on the UA than on the PA of LANDMATE PFT, where the UA increases towards the groups with higher cell homogeneity.




Figure 9. User's accuracy for LANDMATE PFT as function of the threshold for minimum coverage of the respective dominant LULC type.

The spatial analysis for the six assessed LULC types for the 0.7 group is shown in fig. Sample 9 & 10 represent the LANDMATE PFT dataset not adequately regarding distribution and diversity while sample 8 shows a poor coverage in northern Europe. In order to include the largest proportion of the total sample in the analysis, the point count per LULC group as well as the PA per 2.5° grid cell for filter set 2 is analyzed in the results section (fig. B1)10. In order to give an overview of the spatial accuracy for the evaluable filter range, the agreement patterns for the range of evaluable groups, the respective figures for filter set 5 and 7 are shown in Appendix the 0.2 and 0.5 group are included in appendix B (tables B2 & 10B1 & B2).

5.0.1 URBAN

The urban representation in LANDMATE PFT for filter set 2 the 0.7 group is shown in fig. B1a and B1d. The PA for the filter sets 1-10 is shown in fig. ?? where an overall low PA for all filter sets is found 10a and 10d. Figure 8 shows that the PA for all groups is overall low and not majorly influenced by the threshold for minimum coverage. With increasing proportion coverage of the dominant LULC group type URBAN the PA increases slightly but is still lower than 40% for samples groups that include enough points to be considered representative for the research area. The overall low PA is reflected in the URBAN maps in fig. B1 as well as in fig. B2 and 10.

Producer's accuracy of the 10 filter sets (Filter set numbers in grey boxes) for the LULC group URBAN as a function of sample count per filter set.




Figure 10. Total count of GT-SUR-points-evaluated LANDMATE PFT grid cells per 2.5° grid cell (a-c; g-i) and producer's accuracy for the individual LULC groups-types (d-f;j-l) for filter-set-2-group.0.7 (dominant LULC group-type occupies > 2070% per LANDMATE PFT grid cell)

A visual-check A visualization of the map agreement between LANDAMTE-LANDMATE PFT and GT-SUR revealed reveals the issue that leads to the overall low PA. Figure 11 shows four large URBAN agglomerations in different areas of Europe where the red points represent GT-SUR urban points while the white points represent GT-SUR point representing non-

urban LULC [groupstypes](#). The grey-scaled squares represent the LANDMATE PFT URBAN fractions from zero (no coverage, white) to one (full coverage, black) within one grid cell.




Figure 11. Examples of URBAN representation in LANDMATE PFT (greyscale grid) and GT-SUR (points). Cities shown are Hamburg (a), London (b), Rome (c) and [Bucharest](#) (d).

The LANDMATE PFT grid cells with a large urban fraction [indicate](#) the respective city core [of the selected](#) [example cities](#) while the GT-SUR points that are located within the city core are mostly not classified as URBAN. However, the GT-SUR points do not fail to represent the structure of urban areas because [they](#) are characterized through a heterogeneous pattern of sealed surfaces, recreational areas (e.g. parks) and different building types and density, not through a homogeneous sealed area. The LANDMATE PFT map represents this heterogeneous structure through the varying fractions of non-urban PFTs within the grid cell. However, in order to make the impact of a larger city visible in an RCM simulation, it is beneficial for LANDMATE PFT to represent a larger city with a dense core structure. [In order to verify the representation of the large URBAN agglomerations in Europe, a comparison with the World Settlement Footprint for 2015 \(WSF, Marconcini et al., 2020\) dataset was done \(not shown\). The comparison showed that not only larger agglomerations but also smaller patches of settlements are represented well in LANDMATE PFT. Further, the URBAN fractions in LANDMATE PFT are directly adopted from the ESA-CCI LC dataset, which was thoroughly validated.](#) Therefore, despite the low agreement with GT-SUR in the present assessment, the URBAN PFT of LANDMATE PFT 2015 is [considered to be](#) of sufficiently good

quality and suitable to represent urban land cover in high resolution (~2-3 km) RCM simulations. Due to the ~~abovementioned~~ ~~aforementioned~~ comparability issues the UA of the LULC ~~group~~ ~~URBAN will not be further discussed~~ ~~type~~ ~~URBAN is not further discussed in this assessment~~.

5.0.1 CROPLAND

480 The CROPLAND representation in LANDMATE PFT shows, together with WOODLAND the highest PA for the research area. As shown in fig. ~~??~~8 the PA for all ~~filter-sets~~ ~~is~~ ~~ten~~ ~~groups~~ ~~is~~ ~~is~~ > 80% which is to be considered as a very good agreement with the reference.

~~Producer's accuracy of the 10 filter sets (Filter set numbers in grey boxes) for the LULC group CROPLAND as a function of sample count per filter set:~~

485 ~~Figure B1b~~ ~~Figure 10b~~ shows the distribution of CROPLAND points in GT-SUR over the research area. CROPLAND points are the second most frequent LULC ~~group~~ ~~type~~ in GT-SUR and are mainly distributed over middle and southern Europe. Although the northern European grid cells show a lower count of CROPLAND points, figure ~~B1e~~ ~~10e~~ shows that the PA is still very high in these areas. The PA increases with increasing ~~filter-set~~ ~~cell~~ homogeneity (Fig. B2 and 10). Regarding the UA for CROPLAND, LANDMATE PFT shows a strong overestimation, where ~~~51~~36% of the LANDMATE PFT CROPLAND
490 cells in ~~filter-set~~ ~~2~~ ~~the~~ ~~0.7~~ ~~group~~ are actually another LULC ~~group~~ ~~type~~ in the reference. ~~More than half of the LANDMATE PFT CROPLAND areas are mostly WOODLAND, GRASSLAND, and a mix of the other LULC groups in the reference, where 36 % are GRASSLAND and 13 % are WOODLAND.~~ The UA for CROPLAND increases rapidly towards the more homogeneous ~~filter-sets~~ (~~17~~ex61% for ~~filter-set~~ ~~7~~) ~~groups~~. However, the confusion with WOODLAND and GRASSLAND is non-negligible and will be discussed in section 7.

495 5.0.1 WOODLAND

For the representation of WOODLAND, the PA shows the second highest values with > 70% for all ~~filter-sets~~ ~~groups~~ with a reasonably high ~~point-count~~ (~~filter-sets~~ ~~1-7~~ ~~cell~~ ~~count~~ (~~groups~~ ~~0.1-0.7~~, fig. ~~??~~7). Similar to CROPLAND, the ~~sampling~~ ~~filter~~ ~~cell~~ ~~heterogeneity~~ does not have a large impact on PA. The highest PA is reached over the ~~northern~~ ~~Northern~~ European regions (Fig. ~~B1f~~ ~~10f~~). Deficits are visible over the southern British Isles, ~~some~~ ~~parts~~ ~~of~~ ~~France~~ ~~parts~~ ~~of~~ ~~Iberian~~ ~~Penninsula~~ and the coastline
500 along Belgium and the Netherlands. Further, ~~the~~ ~~Mediterranean~~ ~~Coast~~ ~~shows~~ a low PA ~~within~~ ~~grid~~ ~~is~~ ~~found~~ ~~for~~ cells that have an overall small ~~point-count~~ ~~cell~~ ~~count~~ ~~in~~ ~~the~~ ~~Mediterranean~~ (Fig. ~~B1e~~ ~~10c~~).

~~Producer's accuracy of the 10 filter sets (Filter set numbers in grey boxes) for the LULC group WOODLAND as a function of sample count per filter set:~~

The differences between northern and southern regions ~~tends~~ ~~tend~~ to increase towards the more homogeneous ~~filter-sets~~ ~~as~~ ~~shown~~ ~~in~~ ~~figures~~ ~~B2f~~ ~~and~~ ~~10f~~ ~~groups~~ (see fig. B1f and B2f for comparison). Agreement over the northern regions increases while agreement over the Iberian Peninsula decreases together with a rapid decrease of the ~~filter-set~~ ~~WOODLAND~~ ~~cell~~ count within the corresponding grid cells. The UA for WOODLAND is noticeably higher than for all other LULC ~~groups~~ ~~types~~ (> 70% for ~~filter-set~~ ~~2~~ ~~the~~ ~~0.2~~ ~~group~~ and increasing towards the more homogeneous ~~filter-sets~~ ~~groups~~) which emphasises the very good

quality of WOODLAND representation in LANDMATE PFT. ~~(The most confusion is found with LULC type GRASSLAND and the OTHER LULC types (not investigated in this assessment). Altogether, the UA of ~10% for filter set 2). Further, ~17x4% of the total LANDMATE PFT cells representing WOODLAND are actually CROPLAND or OTHER.~~

5.0.1 GRASSLAND

~~The GT-SUR sampling points show the highest GRASSLAND coverage in central Europe with the highest occurrence in Ireland and the southern part of France. 85% for group 0.7 is interpreted as a very good representation of the LULC type WOODLAND within LANDMATE PFT 2015.~~

The coverage of cells with the dominant LULC type GRASSLAND is well distributed except for the Northern European regions (Fig. B1h10h). The PA for LANDMATE PFT GRASSLAND according to fig. B1k is not noticeably higher in these areas but overall highest in the Southwest of the British Isles. For all filter sets 10k is very high on the British Isles and in some regions of Central Europe. For the remaining regions of the research area, the PA ranges between 32 and 34% which for GRASSLAND is considerably low (Fig. ??). This PA pattern remains similar throughout the range of evaluable groups (fig. B1k & B2k), with an average of 31-37%.

~~Producer's accuracy of the 10 filter sets (Filter set numbers in grey boxes) for the LULC group GRASSLAND as a function of sample count per filter set:~~

~~One~~ The main reason for this low accuracy of LANDMATE PFT regarding GRASSLAND can be found looking at the results of sections ?? and ?. The UAs the LULC types CROPLAND and WOODLAND. The UA of CROPLAND and WOODLAND reveal that ~2036% of the LANDMATE PFT CROPLAND cells and ~17x10% of the LANDMATE PFT WOODLAND cells are actually representing actually represent GRASSLAND in the reference, which adds up to over 60almost 55% of the total GT-SUR GRASSLAND points. Another reason is found in the dataset structure of LANDMATE PFT. A considerable amount of GRASSLAND is not part of the assessment because GRASSLAND does not make the dominant but the second dominant PFT in many grid cells (~45% of all LANDMATE PFT grid cells). Therefore, the seemingly weak GRASSLAND representation in LANDMATE PFT rather shows a weakness of the present assessment that is caused by the different dataset structures.

5.0.1 SHRUBLAND & BARE AREAS

The PA for SHRUBLAND and BARE AREAS is the lowest of all assessed LULC groups-types with < 20% for all filter sets groups of both LULC groups-types respectively (Fig. ?? and ??8). The low point overall cell count of both LULC groups-types might be one reason for the low PA. However, looking at the distribution of the SHRUBLAND and BARE AREA points in fig. B410i and ??, LANDMATE PFT is not able to capture the LULC groups-types even in grid cells with a relatively high point cell count. The GT-SUR shows includes ~27,000 SHRUBLAND points while LANDMATE PFT shows includes only ~19,000. 000 cells where SHRUBLAND is the dominant LULC type. Therefore, one reason for the poor SHRUBLAND representation lies within the base map (ESA-CCI LC) used for the creation of LANDMATE PFT, where the known small-low count of SHRUBLAND proportions was inherited by LANDMATE PFT. It must be noted, that a large proportion of SHRUBLAND in

545 ESA-CCI LC is part of the mixed LC classes, such as Shrubland/Cropland or Shrubland/Forest. The known deficit was partly compensated by the translation into the PFTs, where SHRUBLAND proportions were added to the total as proportions of the mixed ESA-CCI LC classes. Further SHRUBLAND makes the second dominant PFT in ~20% of the total LANDMATE PFT grid cells in the assessment. Just like for GRASSLAND, these SHRUBLAND proportions can not be addressed sufficiently within the present assessment.

~~Producer's accuracy of the 10 filter-sets (Filter-set numbers in grey-boxes) for the LULC group SHRUBLAND as a function of sample count per filter-set.~~

550 The overall BARE AREAS ~~sample-cell~~ count in LANDMATE PFT in ~~filter-set 2 is < 50%~~ the 0.7 group is only about 28 % of the actual BARE AREA points in GT-SUR. ~~Almost half~~ Further, within the 0.7 group, over 64 % of the GT-SUR BARE AREAS points are identified as CROPLAND while ~~17% are identified as WOODLAND or GRASSLAND~~. Only ~17% (~~< 21,000~~ points for filter-set 2 the 0.7 group) of the GT-SUR BARE AREAS are actually identified by LANDMATE PFT with the ~~largest agreement~~ highest PA in the Alps, Northern Great Britain, and Northern Scandinavia (Fig. ~~B-H10~~ B-H10). However, due to the comparably low ~~sample-cell~~ count the spatial assessment is ~~not robust~~ rather not reliable. Just like for SHRUBLAND, the 555 homogeneity of LANDMATE PFT cells does not have a large impact on the PA. UA is higher than PA with ~43% for group 0.2 and increasing towards ~~the more homogeneous filter-sets~~ more homogeneous groups (over 60 % for group 0.7). However, considering the rapidly decreasing ~~sample-cell~~ count for the more homogeneous ~~filter-sets~~ groups, the accuracy measures are becoming even less representative for the BARE AREA representation in LANDMATE PFT. Nevertheless, the ~~confusion with the other LULC groups~~ BARE AREA representation in LANDMATE PFT is further discussed in section 7.

560 5.1 Comparison to ESA POULTER PFT validation results

In order to compare the LANDMATE PFT map quality to the ESA POULTER PFT map quality, the validation workflow presented in this manuscript is also applied to the latter for the year 2015. The PA differences of LANDMATE PFT and ESA POULTER PFT are shown in fig. 12. The spatial PA differences vary between the assessed LULC types and groups. For URBAN, the differences are negligible. Since the ESA-CCI LC URBAN proportions are directly adopted in both PFT translations, this result was expected. The small differences in some grid cells of the map might result from the changes in the CWP within the LANDMATE PFT workflow that cause other LULC types to be of dominant coverage and therefore change the total cell counts per LULC type in the aggregation. LANDMATE PFT represents CROPLAND slightly worse than ESA POULTER PFT according to the differences seen in the maps. The threshold for minimum coverage does not have considerable influence on the CROPLAND representation, except for Northern Europe, where the highest minimum coverage threshold (0.7) shows the lowest PA for cropland. In contrast, the LANDMATE PFT WOODLAND representation is most improved for the highest minimum threshold (0.7) in comparison to the ESA POULTER PFTs. A similar result is found for the GRASSLAND representation. It is noticeable that the signal changes for the BARE AREAS representation. For the group with a minimum coverage of 0.2, ESA POULTER PFT shows the better BARE AREAS representation while for the 0.7 group, LANDMATE PFT shows the better quality.

570




Figure 12. Producer's accuracy of Comparison between the 10 filter sets (Filter set numbers in grey boxes) PA for LANDMATE PFT and ESA PFT. The plots show the LULC group BARE AREAS as differences between the PA where the positive values mean a function-higher PA of sample count per filter set LANDMATE PFT and the negative values a higher PA of ESA PFT. The differences were calculated per 2.5° cell of the auxiliary grid for each LULC type of the 0.2 group (left column), the 0.5 group (middle column) and the 0.7 group (right column)

575 6 Data availability

The LANDMATE PFT dataset for Europe 2015 is published with the Long Term Archiving Service (LTA) for large research datasets, which are relevant for climate or earth system research, of the German Climate Computing Service (DKRZ). As World Data Center for Climate (WDCC), the DKRZ LTA is accredited as regular member of the World Data System. The LANDMATE PFT dataset for Europe 2015 is available within the LANDMATE project data at https://cera-www.dkrz.de/WDCC/ui/cerasearch/entry?acronym=LM_PFT_LandCov_EUR2015_v1.0_af (Reinhart et al., 2021b). Within the LANDMATE project, a short documentation summarizes the technical information corresponding to LANDMATE PFT.

7 Discussion & conclusion

The present work introduces the preparation of the LANDMATE PFT map [2015](#) for the European Continent based on [several high-resolution](#) LULC datasets and climate data.

585 The LANDMATE PFT [map for 2015](#) Version 1.0 is prepared in order to provide realistic, high-resolution LULC representation for RCMs. The dataset includes LULC information from different, validated sources as well as regional climate information through involvement of the HLZs. ~~For each ESA-CCI land cover class, an individual CWT~~ [A cross-walking procedure \(CWP\)](#) is developed to translate the original LULC classes into PFTs. The various mixed LULC classes included in the base map ESA-CCI LC are extremely difficult to resolve within RCMs. Through the developed CWP, the mixed LULC classes can be disaggregated into PFT fractions, which improves the realistic representation of these [LULC](#) classes in RCMs. The involvement of the climate data further allows a customized translation of LULC classes for individual regions. The 16 LANDMATE PFTs are selected to provide simple transferability into various RCM families in order to be able to conduct coordinated RCM experiments where the implementation of a common, high quality LULC map provides minimum uncertainty for a multi-model ensemble.

595 The accuracy assessment of LANDMATE PFT is conducted in the form of a comparison with the ground truth dataset [LUCAS - land use and land cover survey \(GT-SUR\)](#). In order to account for the different structure of the reference GT-SUR and the assessed LANDMATE PFT map and further, ~~the fractional structure of the LANDMATE PFT grid cells, a filter is applied. All filtered LANDMATE PFT subsets are analyzed in terms of the grid cells are grouped by a threshold for minimum coverage of the dominant LULC type. All groups are analyzed regarding~~ agreement with the reference (i.e., GT-SUR). In order to investigate regional differences in accuracy measures, a spatial analysis supported by ~~gridded maps of an auxiliary grid over~~ the research area is done. The quality of the LANDMATE PFT map is assessed using the overall accuracy (OA) and the producer's and user's accuracy (PA and UA) for the individual LULC ~~groups-types~~. [The additional comparison to the generic ESA POULTER PFT map \(ESA POULTER PFT\) should give information on the regional improvement of LULC type representation in LANDMATE PFT.](#) Overall, the [assessment-validation](#) serves as recommendation and uncertainty information for regional climate modellers that use LANDMATE PFT, or the time series LUCAS LUC (Hoffmann et al., submitted), which is based on LANDMATE PFT, in RCMs.

Within the accuracy assessment, the OA does not change considerably between the evaluable filter-sets-groups of the respective LULC groups-types which shows that the dataset structure has no noticeable impact on that accuracy measure. The highest PA is found for CROPLAND and WOODLAND which are the dominant LULC groups-types in the research area. The lowest PA is found for SHRUBLAND and BARE AREAS, which are also the LULC groups-types with the lowest overall sample-cell count. The UA is found to be highest for WOODLAND, followed by CROPLAND, GRASSLAND and BARE AREAS. Both accuracy measures, PA and UA are highly-influenced-by-the-proportion-influenced-by-grid-cell-heterogeneity of the dominant LULC group-in-the-individual-type-within-a grid cell. The difference between the filter-sets-for-UA-of-the-LULC-groups-groups-for-UA is 10 to 20 % per group while the difference for PA is noticeable but considerably lower, which means that the applied filter-threshold-range has a higher influence on the former.

The URBAN representation in LANDMATE PFT represents a special case in the present assessment due to the heterogeneous structure of urban areas. Both datasets, GT-SUR and LANDMATE PFT are able to represent the LULC group-type URBAN very well for their respective purpose. Nevertheless, the PA for URBAN reflects the limitations of the present assessment method. The fine scale point data of GT-SUR represents the patchwork structure of recreational areas, building blocks, and other urban elements at the location of the respective points while LANDMATE PFT represents the urban area as an agglomeration of grid cells with URBAN as the dominant LULC group.~~The additional comparison with a high resolution dataset (WSF2015) showed that not only large but also small agglomerations of urban areas are represented well in LANDMATE PFT.~~ Therefore and despite of the accuracy assessment results for the LULC group-type URBAN, the LANDMATE PFT dataset can be recommended to be used in RCMs that resolve urban features over the European Continent.

A limitation of LANDMATE PFT is the overestimation of CROPLAND to the expense of ~~WOODLAND and GRASSLAND and GRASSLAND and WOODLAND and~~ the overestimation of WOODLAND to the expense of mostly GRASSLAND. This overestimation has a minor impact on the overall WOODLAND and CROPLAND representation but a major impact on the representation of GRASSLAND in LANDMATE PFT. The representation of GRASSLAND is comparably low due to the aforementioned reasons. Further, the LULC groups-types with the lowest point counts SHRUBLAND and BARE AREAS are not well represented, which happens due to the low overall sample size but also due to the overall too low representation in LANDMATE PFT, which is partly inherited from the base map ESA-CCI LC. The representation of these LULC groups-types needs to be considered when using LANDMATE PFT in RCM simulations using the supporting maps in fig. B1,B2 and 10. Nevertheless, the representation of SHRUBLAND and BARE AREAS is improved in some regions, compared to ESA POULTER PFT.

The representation of LULC groups-types in LANDMATE PFT is assessed through the comparison with ground truth data. The structural differences of the datasets, where gridded data is compared to point data, is a major weakness of this assessment. Although the fractional structure does not have a major influence on the OA, the LULC group-wise-type-wise PA and even more the UA is affected.

The present assessment takes into account the dominant LULC group-type per grid cell of LANDMATE PFT. Depending on the proportion of this LULC group-type, the second or third-most represented LULC group-type can occupy a considerable area of the respective grid cell. Therefore, a follow up assessment, where these LULC group-type proportions are also

considered and compared to the ground truth is needed in order to investigate, if the PA of the less dominant LULC ~~groups~~ types GRASSLAND, SHRUBLAND, and BARE AREAS is increased. The use of additional LULC data ~~, like it was done for URBAN in this assessment, would be an additional~~ specialized one one LULC type would be a useful step to validate the
645 quality of GRASSLAND, SHRUBLAND and BARE AREAS representation in LANDMATE PFT ~~-2015~~.

The results show that the LANDMATE PFT map is able to represent LULC over large parts of Europe in a sufficient quality. Especially the dominant LULC ~~groups~~ types are represented overall well which is highly beneficial for RCM experiments that require realistic, high-resolution LULC representation. Nevertheless, there are uncertainties found for the less represented LULC ~~group~~ types. When using LANDMATE PFT in an RCM it is crucial to consider these uncertainties when interpreting
650 simulation results. Especially the spatial distribution of uncertainties in LANDMATE PFT needs to be considered when comparing simulation results to observations because the input parameters in the employed land-surface schemes are influenced by the individual LULC, which subsequently considerably impacts on lower-atmosphere processes, such as the intensity of heat and moisture exchange. Thus, by carefully considering the issue of uncertainty introduced by the LULC input, misconclusions about RCM model performance and about small-scale interconnections can be ~~avoided~~ reduced (Ge et al., 2007; Sertel et al.,
655 2010; Santos-Alamillos et al., 2015; Reinhart et al., 2021a).

Beside the quality of the LULC product, the implementation process of each individual RCM is crucial for the realistic representation of LULC in regional climate model experiments. When translating a LULC product into the model specific LULC classes and structure, modifications are done that can change the map characteristics. When the LANDMATE PFT product is used in an RCM that only uses the dominant LULC fraction per grid cell, the overall LULC proportions can change.
660 The same applies when LANDMATE PFT is used in a model with limited fractions per grid cell or a different classification system. The present assessment gives a guideline on the quality of LANDMATE PFT (Version 1.0) when used unaltered. Through the involvement of the ground truth data, regional deficits of LANDMATE PFT are presented that can be compensated during the implementation process into the individual RCM or RCM family.

The findings of the present assessment support the identification of uncertainties within the LANDMATE PFT map for
665 Europe. Nevertheless, user feedback is crucial for the future overall improvement of LANDMATE PFT. The RCM community within the WCRP FPS LUCAS is already participating in the feedback process where implementation of LANDMATE PFT and the LUCAS LUC time series into different RCMs is comprehensively documented. The future work on LANDMATE PFT also includes the extension of the dataset to other CORDEX regions. Although, the dataset is based on various globally available datasets and therefore, can be created globally, the introduced quality assessment method must be performed for
670 each region individually, desirably using region-specific expert knowledge. Further, the assessment should be expanded in order to include the second or third-most represented LULC ~~group~~ type per grid cell to possibly achieve more accurate quality information about LANDMATE PFT.

Appendix A

Table A1. Cross-walking table for ESA-CCI LC class 10 - Cropland, rainfed and LC class 11 -Cropland, herbaceous cover. For LC class 10 and 11, no HLZ were assigned

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Tree							Shrub		Grass		Special vegetation					Non-vegetated
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	Crops crops		urban	bare ground
1-30									10				90			

Table A2. Cross-walking table for ESA-CCI LC class 12 - Cropland, tree or shrub cover. For LC class 12, no HLZ were assigned

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Tree							Shrub		Grass		Special vegetation					Non-vegetated
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	Crops crops		urban	bare ground
1-30								70					30			

Table A3. Cross-walking table for ESA-CCI LC class 20 - Cropland, irrigated or post flooding. For LC class 20, no HLZ were assigned

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Tree							Shrub		Grass		Special vegetation					Non-vegetated
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	Crops crops		urban	bare ground
1-30													100			

Table A4. Cross-walking table for ESA-CCI LC class 30 - Mosaic cropland (>50%) / natural vegetation (tree, shrub, herbaceous cover)(<50%).

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1-6											20	20	60			
7-9									40							
10					10				30							
11,12					30				10							
13,14									40							
15				5	5			20	10							
16				7.5	7.5			10	15							
17,18				20				10	10							
19									40							
20							20		20							
21,22				10	10				10							
23,24				10	10				20							
25									40							
26							20		20							
27		20					10		10							
28		10					15		15							
29	15						10		15							
30	20							10	10							

Table A5. Cross-walking table for ESA-CCI LC class 40 - Mosaic natural vegetation (tree, shrub, herbaceous cover)(>50%) / cropland(<50%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1,2											35	30	35			
3-5											30	35	35			
6											25	40	35			
7									60							
8					10				50							
9,10					15				45							
11					20				40							
12					30			20	10							
13				10	10			10	30							
14,15				20	20			10	10							
16				25	20				15							
17				25	25				10							
18				30	30											
19									60							
20							35		25							
21			20		15		15		10							
22			25		10		15		10							
23,24			20		20		20									
25									60							
26							30		30							
27		10					50									
28		40					20									
29	40						20									
30	50						10									

Table A6. Cross-walking table for ESA-CCI LC class 50 - Tree cover, broadleaved, evergreen, closed to open (>15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass	Special vegetation			Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6			12.5				12.5				75					
7-18			90	10												
19-24			100													
25-30	100															

Table A7. Cross-walking table for ESA-CCI LC class 60 - Tree cover, broadleaved, deciduous, closed to open (>15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass	Special vegetation			Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6								100								
7-24				70				15	15							
25-30		70						15	15							

Table A8. Cross-walking table for ESA-CCI LC class 61 - Tree cover, broadleaved, deciduous, closed (>40%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass	Special vegetation			Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6								85	15							
7-24				70				15	15							
25-30		70						15	15							

Table A9. Cross-walking table for ESA-CCI LC class 62 - Tree cover, broadleaved, deciduous, open (15-40%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass	Special vegetation			Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6								65	35							
7-24				30				25	45							
25-30		30						25	45							

Table A10. Cross-walking table for ESA-CCI LC class 70 - Tree cover, needleleaved, evergreen, closed to open (>15%) and LC class 71 - Tree cover, needleleaved, evergreen, closed (>40%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6					35	35	15		15							
7-18					70		10	5	15							
19-24			35		35		10	5	15							
25-30					70		10	5	15							

Table A11. Cross-walking table for ESA-CCI LC class 72 - Open (15-40%) needleleaved deciduous or evergreen forest (>5m)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6					15	15	25		45							
7-18					30		20	5	45							
19-24			15		15		20	5	45							
25-30					30		20	5	45							

Table A12. Cross-walking table for ESA-CCI LC class 80 - Tree cover, needleleaved, deciduous, closed to open (>15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-30						50	5	15	30							

Table A13. Cross-walking table for ESA-CCI LC class 81 - Treecover, needleleaved, deciduous, closed (>40%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-30						70		15	15							

Table A14. Cross-walking table for ESA-CCI LC class 82 - Tree cover, needleleaved, deciduous, open (15-40%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-30						30	5	20	45							

Table A15. Cross-walking table for ESA-CCI LC class 90 - Tree cover, mixed leaf type (broadleaved and needleleaved)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1-12				20	70				10							
13-24				70	20				10							
25-30	45	45							10							

Table A16. Cross-walking table for ESA-CCI LC class 100 - Mosaic tree and shrub (>50%) / herbaceous cover(<50%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1					30		30					30	10			
2,3					30		25					25	20			
4-6					30		20					20	30			
7-9				20	20		20		40							
10				25	25		20		30							
11				30	30		20		20							
12				30	30		25		15							
13				15	15			35	35							
14				20	20			30	30							
15				25	25			25	25							
16-18				25	25			30	20							
19,20			30				30		40							
21,22			35				35		30							
23,24			40				30		30							
25		20					50		30							
26		25					50		25							
27		30					45		25							
28		40					35		25							
29		60					20		20							
30		70					15		15							

Table A17. Cross-walking table for ESA-CCI LC class 110 - Mosaic herbaceous cover (>50%) / tree and shrub (<50%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1-6							50				45	5				
7				10	10		20		60							
8				10	20		10		60							
9				25	25				50							
10				30	30				40							
11,12				35	35				30							
13				15	15				70							
14,15				20	10				70							
16				30				10	60							
17,18				35				15	50							
19			15				15		70							
20			10				20		70							
21			20				10		70							
22			30				10		60							
23,24			35				15		50							
25		15					15		70							
26		20					10		70							
27		25					15		60							
28		30						10	60							
29		40						10	50							
30		50						10	40							

Table A18. Cross-walking table for ESA-CCI LC class 120 - Shrubland

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Holdridge Life Zone	Tree tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	Shrub evergreen	deciduous	Grass C3	C4	Special vegetation Tundra	Swamps	Crops crops	Non-vegetated urban	bare ground	
1-6							40				55	5				
7-12							10	50	40							
13							70		30							
14							40	30	30							
15							20	60	20							
16							20	70	10							
17,18							10	80	10							
19							10		90							
20							50		50							
21							90		10							
22							80	10	10							
23,24							100									
25							10	10	80							
26,27							20	60	20							
28							10	70	20							
29,30							10	80	10							

Table A19. Cross-walking table for ESA-CCI LC class 121 - Evergreen shrubland and LC class 122 - Deciduous Shrubland

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Holdridge Life Zone	Tree tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	Shrub evergreen	deciduous	Grass C3	C4	Special vegetation Tundra	Swamps	Crops crops	Non-vegetated urban	bare ground	
1-6							40				55	5				
7-12							60		40							
13,14							70		30							
15							80		20							
16-18							90		10							
19							10		90							
20							50		50							
21,22							90		10							
23,24							100									
25							20		80							
26-28							80		20							
29,30							90		10							

Table A20. Cross-walking table for ESA-CCI LC class 122 - Evergreen shrubland and LC class 122 - Deciduous Shrubland

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Holdridge Life Zone	Tree tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	Shrub evergreen	deciduous	Grass C3	C4	Special vegetation Tundra	Swamps	Crops crops	Non-vegetated urban	bare ground	
1-6											55	5				
7-12								60	40							
13,14								70	30							
15								80	20							
16-18								90	10							
19								10	90							
20								50	50							
21,22								90	10							
23,24								100								
25								80	20							
26-28								80	20							
29,30								90	10							

Table A21. Cross-walking table for ESA-CCI LC class 130 - Grassland

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6											90	10				
7-13									100							
14							5		95							
15								7.5	92.5							
16								10	90							
17								12.5	87.5							
18								15	85							
19									100							
20,21							5		95							
22							7.5		92.5							
23,24							10		90							
25									100							
26							5		95							
27							5	5	90							
28								10	90							
29								12.5	87.5							
30								15	85							

Table A22. Cross-walking table for ESA-CCI LC class 140 - Lichens and mosses

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6											90	10				
7-30									100							

Table A23. Cross-walking table for ESA-CCI LC class 150 - Sparse vegetation (tree, shrub, herbaceouscover)(<15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1-6											50	10				40
7-12							10		40							50
13				5	5		5		35							50
14				5	5		10		30							50
15				5	5			10	30							50
16				5	5				20							50
17,18				10	10			20	10							50
19							5		45							50
20,21			5				5		40							50
22			5				10		35							50
23			10				10		30							50
24			15				15		20							50
25							5	5	40							50
26,27		10					5	5	30							50
28,29		10							20							50
30	10								20							50

Table A24. Cross-walking table for ESA-CCI LC class 151 - Sparse tree (<15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
1-6											50	10				40
7-12							10		40							50
13				5	5				40							50
14,15				5	10				35							50
16				10	5				35							50
17,18				10	10				30							50
19-21			5						45							50
22			10						40							50
23			15						35							50
24			20						30							50
25		10							40							50
26-29		15							35							50
30	15								35							50

Table A25. Cross-walking table for ESA-CCI LC class 152 - Sparse shrub (<15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1							5				45	10				40
2-6							10				40	10				40
7-10							10		40							50
11,12							20		30							50
13,14							10		40							50
15,16								15	35							50
17,18								20	30							50
19							5		45							50
20,21							10		40							50
22,23							15		35							50
24							20		30							50
25							5		45							50
26							10		40							50
27							7.5	7.5	35							50
28,29									15							50
30									20							50

Table A26. Cross-walking table for ESA-CCI LC class 153 - Sparse herbaceous cover (<15%)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6											40	10				50
7-30									50							50

Table A27. Cross-walking table for ESA-CCI LC class 160 - Tree cover, flooded, fresh or brakish water

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6				10							45	45				
7-18				70								30				
19-24			70									30				
25-30	35	35										30				

Table A28. Cross-walking table for ESA-CCI LC class 170 - Tree cover, flooded, saline water

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops		Non-vegetated	
Holdridge Life Zone	tropical broadleaf evergreen	tropical broadleaf deciduous	temperate broadleaf evergreen	temperate broadleaf deciduous	evergreen coniferous	deciduous coniferous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops		urban	bare ground
1-6					40		30				10	20				
7-12				20				60				20				
13-18				30				50				20				
19-24			60				10	10				20				
25-30	80											20				

Table A29. Cross-walking table for ESA-CCI LC class 180 - Shrub or herbaceous cover, flooded, fresh / saline / brakish water

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical	tropical	temperate	temperate	evergreen	deciduous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
	broadleaf	broadleaf	broadleaf	broadleaf	coniferous	coniferous										
	evergreen	deciduous	evergreen	deciduous												
1-6											95	5				
7								10				90				
8							15	15	20			50				
9							20	20	20			40				
10-12							20	20	20			40				
13								20	20			60				
14								25	25			50				
15								30	30			40				
16								35	35			30				
17,18								45	15			40				
19,20							30	40	30							
21,22							40	40	20							
23							40	50	10							
24							30	60	10							
25							30	30	40							
26							30	40	30							
27							40	40	20							
28							40	50	10							
29							70	30								
30							90	10								

Table A30. Cross-walking table for ESA-CCI LC class 190 - Urban

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical	tropical	temperate	temperate	evergreen	deciduous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
	broadleaf	broadleaf	broadleaf	broadleaf	coniferous	coniferous										
	evergreen	deciduous	evergreen	deciduous												
1-30															100	

Table A31. Cross-walking table for ESA-CCI LC class 200 - Bare areas, LC class 201 - Consolidated bare areas and LC class 202 - Unconsolidated bare areas.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
	Tree						Shrub		Grass		Special vegetation		Crops	Non-vegetated		
Holdridge Life Zone	tropical	tropical	temperate	temperate	evergreen	deciduous	evergreen	deciduous	C3	C4	Tundra	Swamps	crops	urban	bare ground	
	broadleaf	broadleaf	broadleaf	broadleaf	coniferous	coniferous										
	evergreen	deciduous	evergreen	deciduous												
1-30															100	

Appendix B




Figure B1. Total count of GT-SUR points evaluated LANDMATE PFT grid cells per 2.5° grid cell (a-c; g-i) and producer's accuracy for the individual LULC groups types (d-f; j-l) for filter set 5 group 0.2 (dominant LULC group type occupies > 50.20% per LANDMATE PFT grid cell)




Figure B2. Total count of ~~GT-SUR points~~ evaluated LANDMATE PFT grid cells per 2.5° grid cell (a-c; g-i) and producer's accuracy for the individual LULC ~~groups types~~ (d-f; j-l) for filter-set-7-group 0.5 (dominant LULC ~~group-type~~ occupies > 7050% per LANDMATE PFT grid cell)

	1	2	3	4	5	6	7	SUM	UA
1	3234	806	1063	178	1769	120	407	7577	42.68
2	6625	67374	22298	5444	28559	4185	2485	136970	49.19
3	2414	5081	88064	8989	12818	1527	5544	124437	70.77
4	624	5316	4637	5498	1789	439	1487	19790	27.78
5	1411	4515	8063	6082	20763	1767	1643	44244	46.93
6	82	199	200	830	567	1810	460	4148	43.64
7	3	4	49	277	276	530	314	1453	21.61
SUM	14393	83295	124374	27298	66541	10378	12340		
PA	22.47	80.887	70.81	20.14	31.20	17.44	2.54	OA:	55.24

Table B1. Confusion matrix for LANDMATE PFT filter-set-1-group 0.1 - Dominant LULC group-type occupies a minimum of 10 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	3234	806	1063	178	1769	120	407	7577	42.68
2	6625	67374	22298	5444	28559	4185	2485	136970	49.19
3	2414	5081	88064	8989	12818	1527	5544	124437	70.77
4	624	5316	4637	5498	1789	439	1487	19790	27.78
5	1411	4515	8063	6082	20763	1767	1643	44244	46.93
6	82	199	200	830	567	1810	460	4148	43.64
7	3	4	49	277	276	530	314	1453	21.61
SUM	14393	83295	124374	27298	66541	10378	12340		
PA	22.47	80.887	70.81	20.14	31.20	17.44	2.54	OA:	55.24

Table B2. Confusion matrix for LANDMATE PFT filter-set-2-group 0.2 - Dominant LULC group-type occupies a minimum of 20 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	3221	793	1041	174	1748	117	404	7498	42.96
2	6596	67323	22210	5395	28488	4168	2457	136637	49.27
3	2377	5034	87838	8903	12750	1511	5483	123896	70.90
4	615	5280	4484	5363	1748	425	1401	19316	27.76
5	1401	4485	7961	5983	20716	1754	1559	43859	47.23
6	78	187	186	798	552	1799	452	4052	44.40
7	3	4	47	276	275	530	310	1445	21.45
SUM	14291	83106	123767	26892	66277	10304	12066		
PA	22.54	81.01	70.97	19.94	31.26	17.46	2.57	OA:	55.41

Table B3. Confusion matrix for LANDMATE PFT filter-set-3-group 0.3 - Dominant LULC group-type occupies a minimum of 30 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	3079	715	904	152	1597	109	364	6920	44.49
2	6263	66184	20069	4795	27209	4034	2304	130858	50.58
3	2061	4045	83073	7509	11168	1274	5030	114160	72.77
4	501	4813	3013	4235	1392	329	742	15025	28.19
5	1238	4031	6748	5091	19572	1571	1219	39470	49.59
6	54	123	122	606	469	1681	425	3480	48.30
7	2	2	40	254	258	517	252	1325	19.02
SUM	13198	79913	113969	22642	61665	9515	10336		
PA	23.33	82.82	72.89	18.70	31.74	17.67	2.44	OA:	57.22

Table B4. Confusion matrix for LANDMATE PFT filter-set-4-group 0.4 - Dominant LULC group-type occupies a minimum of 40 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	2632	499	676	117	1218	84	292	5518	47.70
2	5482	62499	15269	3772	23519	3737	1913	116191	53.79
3	1510	2215	71799	5277	7767	853	4284	93705	76.62
4	362	3865	1752	2689	915	206	350	10139	26.52
5	933	2992	4373	3605	16306	1227	893	30329	53.76
6	31	61	62	292	321	1375	392	2534	54.26
7	1	0	29	110	214	233	70	657	10.65
SUM	10951	72131	93960	15862	50260	7715	8194		
PA	24.03	86.65	76.41	16.95	32.44	17.82	0.85	OA:	60.74

Table B5. Confusion matrix for LANDMATE PFT filter-set-5-group 0.5 - Dominant LULC group-type occupies a minimum of 50 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	2123	284	464	85	844	67	231	4098	51.81
2	4436	56963	10802	2887	19016	3314	1556	98974	57.55
3	1025	978	57212	2949	4699	488	3345	70696	80.93
4	194	2459	967	1713	518	122	240	6213	27.57
5	628	1847	2584	2333	12497	798	630	21317	58.62
6	14	27	34	104	181	1022	339	1721	59.38
7	1	0	18	40	153	87	25	324	7.72
SUM	8421	62558	72081	10111	37908	5898	6366		
PA	25.21	91.06	79.37	16.94	32.97	17.33	0.39	OA:	64.70

Table B6. Confusion matrix for LANDMATE PFT filter-set-6-group 0.6 - Dominant LULC group-type occupies a minimum of 60 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	1684	167	311	53	568	44	185	3012	55.91
2	3288	49624	7217	2088	14351	2840	1145	80553	61.60
3	414	255	30158	806	1745	177	1910	35465	85.04
4	40	793	458	988	191	42	160	2672	36.98
5	410	1053	1363	1415	9113	478	425	14257	63.92
6	5	11	15	61	104	768	302	1266	60.66
7	1	0	9	19	99	50	9	187	4.81
SUM	5842	51903	39531	5430	26171	4399	4136		
PA	28.83	95.61	76.29	18.20	34.82	17.46	0.22	OA:	67.20

Table B7. Confusion matrix for LANDMATE PFT filter-set-7-group 0.7 - Dominant LULC group-type occupies a minimum of 70 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	1261	83	208	29	369	32	138	2120	59.48
2	2009	38997	4002	1296	9321	2239	745	58609	66.54
3	32	21	3201	54	195	8	108	3619	88.45
4	10	74	198	442	51	9	106	890	49.66
5	241	518	640	691	5957	240	229	8516	69.95
6	3	5	10	39	62	533	268	920	57.93
7	1	0	6	8	53	17	6	91	6.59
SUM	3557	39698	8265	2559	16008	3078	1600		
PA	35.45	98.23	38.73	17.27	37.21	17.32	0.38	OA:	67.41

Table B8. Confusion matrix for LANDMATE PFT filter-set-8-group 0.8 - Dominant LULC group-type occupies a minimum of 80 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	808	44	111	14	207	16	89	1289	62.68
2	592	17167	877	414	2601	1043	269	22963	74.76
3	1	1	47	1	1	0	14	65	72.31
4	2	7	28	74	11	1	10	133	55.64
5	40	81	108	181	1358	83	58	1909	71.14
6	3	1	7	20	28	338	230	627	53.91
7	0	0	1	2	2	1	1	7	14.29
SUM	1446	17301	1179	706	4208	1482	671		
PA	55.88	99.23	3.99	10.48	32.27	22.81	0.15	OA:	73.33

Table B9. Confusion matrix for LANDMATE PFT filter-set-9-group 0.9 - Dominant LULC group-type occupies a minimum of 90 % of a LANDMATE PFT grid cell

	1	2	3	4	5	6	7	SUM	UA
1	252	10	28	0	40	8	51	389	64.78
2	22	565	16	7	52	14	20	696	81.18
3	0	0	0	0	0	0	0	0	/
4	0	0	0	0	0	0	0	0	/
5	0	1	4	14	48	6	1	74	64.86
6	2	0	4	7	9	112	156	290	38.62
7	0	0	0	0	0	0	0	0	/
SUM	276	576	52	28	149	140	228		
PA	91.30	98.09	0.00	0.00	32.21	80.00	0.00	OA:	67.43

Table B10. Confusion matrix for LANDMATE PFT filter-set-10-group 1.0 - Dominant LULC group-type occupies 100 % of a LANDMATE PFT grid cell

675 *Author contributions.* VR conceptualized the paper outline and objective with the support of DR, PH and JB. VR and PH developed the cross-walking procedure and the corresponding cross-walking tables. PH developed the required translation software for the cross-walking procedure. VR developed the accuracy assessment design for the LANDMATE PFT map supported by BB. VR conducted the accuracy assessment and the visualization of results. VR wrote the original draft of the paper, VR, PH, DR, JB, and BB reviewed and edited the draft. VR wrote the final paper

680 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This work was financed within the framework of the Helmholtz Institute for Climate Service Science (HICSS), a co-operation between Climate Service Center Germany (GERICS) and Universität Hamburg, Germany and conducted as part of the project LANDMATE (Modelling human LAND surface modifications and its feedbacks on local and regional climate). We acknowledge the support of LUCAS by WCRP-CORDEX as a Flagship Pilot Study. We acknowledge the E-OBS dataset from the EU-FP6 project UERRA 685 (<http://www.uerra.eu>) and the Copernicus Climate Change Service, and the data providers in the ECA&D project (<https://www.ecad.eu>). We thank the European Space Agency (ESA) for making the Land cover products publicly available. Special thanks go to the FPS LUCAS partners for providing useful comments in order to improve the dataset.

References

- Alkama, R. and Cescatti, A.: Biophysical climate impacts of recent changes in global forest cover, *Science*, 351, 600–604, 2016.
- 690 Bégué, A., Arvor, D., Bellon, B., Betbeder, J., De Abelleira, D., PD Ferraz, R., Lebourgeois, V., Lelong, C., Simões, M., and R Verón, S.: Remote sensing and cropping practices: A review, *Remote Sensing*, 10, 99, 2018.
- Belda, M., Halenka, T., Huszar, P., Karlicky, J., and Nováková, T.: Do we need urban parameterization in high resolution regional climate simulations?, in: *AGU Fall Meeting Abstracts*, vol. 2018, pp. A21L–2878, 2018.
- Bojinski, S., Verstraete, M., Peterson, T. C., Richter, C., Simmons, A., and Zemp, M.: The concept of essential climate variables in support
695 of climate research, applications, and policy, *Bulletin of the American Meteorological Society*, 95, 1431–1443, 2014.
- Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *science*, 320, 1444–1449, 2008.
- Bonan, G. B., Levis, S., Kergoat, L., and Oleson, K. W.: Landscapes as patches of plant functional types: An integrating concept for climate and ecosystem models, *Global Biogeochemical Cycles*, 16, 5–1, 2002.
- Bontemps, S., Defourny, P., Radoux, J., Van Bogaert, E., Lamarche, C., Achard, F., Mayaux, P., Boettcher, M., Brockmann, C., Kirches,
700 G., et al.: Consistent global land cover maps for climate modelling communities: current achievements of the ESA's land cover CCI, in: *Proceedings of the ESA living planet symposium*, Edinburgh, pp. 9–13, 2013.
- Box, E. O.: Plant functional types and climate at the global scale, *Journal of Vegetation Science*, 7, 309–320, 1996.
- Bright, R. M., Zhao, K., Jackson, R. B., and Cherubini, F.: Quantifying surface albedo and other direct biogeophysical climate forcings of forestry activities, *Global Change Biology*, 21, 3246–3266, 2015.
- 705 Burke, M. and Emerick, K.: Adaptation to climate change: Evidence from US agriculture, *American Economic Journal: Economic Policy*, 8, 106–40, 2016.
- Chapin Iii, F., McGuire, A., Randerson, J., Pielke, R., Baldocchi, D., Hobbie, S. E., Roulet, N., Eugster, W., Kasischke, E., Rastetter, E., et al.: Arctic and boreal ecosystems of western North America as components of the climate system, *Global Change Biology*, 6, 211–223, 2000.
- 710 Chen, X., Zhang, X.-S., and Li, B.-L.: The possible response of life zones in China under global climate change, *Global and Planetary Change*, 38, 327–337, 2003.
- Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V., and Böhner, J.: System for automated geoscientific analyses (SAGA) v. 2.1. 4., *Geoscientific Model Development Discussions*, 8, 2015.
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J., and Jones, P. D.: An ensemble version of the E-OBS temperature and precipitation
715 data sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, 2018.
- Daly, C., Helmer, E. H., and Quiñones, M.: Mapping the climate of puerto rico, vieques and culebra, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 23, 1359–1381, 2003.
- Daniel, M., Lemonsu, A., Déqué, M., Somot, S., Alias, A., and Masson, V.: Benefits of explicit urban parameterization in regional climate modeling to study climate and city interactions, *Climate Dynamics*, 52, 2745–2764, 2019.
- 720 Davin, E. L., Rechid, D., Breil, M., Cardoso, R. M., Coppola, E., Hoffmann, P., Jach, L. L., Katragkou, E., de Noblet-Ducoudré, N., Radtke, K., et al.: Biogeophysical impacts of forestation in Europe: first results from the LUCAS (Land Use and Climate Across Scales) regional climate model intercomparison, *Earth System Dynamics*, 11, 183–200, 2020.
- Di Gregorio, A.: Land cover classification system: classification concepts and user manual: LCCS, vol. 2, Food & Agriculture Org., 2005.

- Dierckx, W., Sterckx, S., Benhadj, I., Livens, S., Duhoux, G., Van Achteren, T., Francois, M., Mellab, K., and Saint, G.: PROBA-V mission for global vegetation monitoring: standard products and image quality, *International Journal of Remote Sensing*, 35, 2589–2614, 2014.
- 725 Donlon, C., Berruti, B., Buongiorno, A., Ferreira, M.-H., Féménias, P., Frerick, J., Goryl, P., Klein, U., Laur, H., Mavrocordatos, C., et al.: The global monitoring for environment and security (GMES) sentinel-3 mission, *Remote Sensing of Environment*, 120, 37–57, 2012.
- d’Andrimont, R., Yordanov, M., Martinez-Sanchez, L., Eiselt, B., Palmieri, A., Dominici, P., Gallego, J., Reuter, H. I., Joebges, C., Lemoine, G., et al.: Harmonised LUCAS in-situ land cover and use database for field surveys from 2006 to 2018 in the European Union, *Scientific Data*, 7, 1–15, 2020.
- 730 ESA: Land Cover CCI Product User Guide Version 2, Tech. rep., European Space Agency, maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf, 2017.
- ESA: Land cover CCI product user guide version 2, Tech. Report, pp. p–105, 2017.
- ESA, E. A. P.: Available online: http://envisat.esa.int/handbooks/asar_CNTR.html (accessed on 27 January 2020), 2002.
- 735 Foody, G. M.: Status of land cover classification accuracy assessment, *Remote sensing of environment*, 80, 185–201, 2002.
- Ge, J., Qi, J., Lofgren, B. M., Moore, N., Torbick, N., and Olson, J. M.: Impacts of land use/cover classification accuracy on regional climate simulations, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Harris, I., Jones, P., Osborn, T., and Lister, D.: Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset, *International Journal of Climatology*, 34, 623–642, <https://doi.org/10.1002/joc.3711>, 2014.
- 740 Hartley, A., MacBean, N., Georgievski, G., and Bontemps, S.: Uncertainty in plant functional type distributions and its impact on land surface models, *Remote Sensing of Environment*, 203, 71–89, 2017.
- Hastings, D. A. and Emery, W. J.: The advanced very high resolution radiometer (AVHRR)-A brief reference guide, *Photogrammetric Engineering and Remote Sensing*, 58, 1183–1188, 1992.
- Hoffmann, P., Katzfey, J., McGregor, J., and Thatcher, M.: Bias and variance correction of sea surface temperatures used for dynamical downscaling, *Journal of Geophysical Research: Atmospheres*, 121, 12–877, 2016.
- 745 Hoffmann, P., Reinhart, V., Rechid, D., de Noblet-Ducoudré, N., Davin, E., Asmus, C., Bechtel, B., Böhner, J., Katrakgou, E., and Luysaert, S.: High-resolution land-use land-cover change data for regional climate modelling applications over Europe - Part 2: Historical and future changes, *ESSD*, submitted.
- Holdridge, L. R. et al.: Life zone ecology., *Life zone ecology.*, 1967.
- 750 Hua, T., Zhao, W., Liu, Y., Wang, S., and Yang, S.: Spatial consistency assessments for global land-cover datasets: A comparison among GLC2000, CCI LC, MCD12, GLOBCOVER and GLCNMO, *Remote Sensing*, 10, 1846, 2018.
- Hurt, G. C., Chini, L. P., Froking, S., Betts, R., Feddema, J., Fischer, G., Fisk, J., Hibbard, K., Houghton, R., Janetos, A., et al.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic change*, 109, 117, 2011.
- 755 Jung, M., Henkel, K., Herold, M., and Churkina, G.: Exploiting synergies of global land cover products for carbon cycle modeling, *Remote Sensing of Environment*, 101, 534–553, 2006.
- Karthikeyan, L., Chawla, I., and Mishra, A. K.: A review of remote sensing applications in agriculture for food security: Crop growth and yield, irrigation, and crop losses, *Journal of Hydrology*, 586, 124 905, 2020.
- Khatun, K., Imbach, P., and Zamora, J.: An assessment of climate change impacts on the tropical forests of Central America using the Holdridge Life Zone (HLZ) land classification system, *iForest-Biogeosciences and Forestry*, 6, 183, 2013.
- 760

- Kueppers, L. M., Snyder, M. A., and Sloan, L. C.: Irrigation cooling effect: Regional climate forcing by land-use change, *Geophysical Research Letters*, 34, 2007.
- Lattanzi, F. A.: C3/C4 grasslands and climate change, in: *Grassland Science in Europe*, pp. 3–13, 2010.
- Lavorel, S., Díaz, S., Cornelissen, J. H. C., Garnier, E., Harrison, S. P., McIntyre, S., Pausas, J. G., Pérez-Harguindeguy, N., Roumet, C.,
765 and Urcelay, C.: Plant functional types: are we getting any closer to the Holy Grail?, in: *Terrestrial ecosystems in a changing world*, pp. 149–164, Springer, 2007.
- Lawrence, D. and Vandecar, K.: Effects of tropical deforestation on climate and agriculture, *Nature climate change*, 5, 27–36, 2015.
- Li, W., MacBean, N., Ciais, P., Defourny, P., Lamarche, C., Bontemps, S., Houghton, R. A., and Peng, S.: Gross and net land cover changes in the main plant functional types derived from the annual ESA CCI land cover maps (1992–2015), 2018.
- 770 Lobell, D., Bala, G., and Duffy, P.: Biogeophysical impacts of cropland management changes on climate, *Geophysical Research Letters*, 33, 2006.
- Lu, Y. and Kueppers, L. M.: Surface energy partitioning over four dominant vegetation types across the United States in a coupled regional climate model (Weather Research and Forecasting Model 3–Community Land Model 3.5), *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- 775 Lugo, A. E., Brown, S. L., Dodson, R., Smith, T. S., and Shugart, H. H.: The Holdridge life zones of the conterminous United States in relation to ecosystem mapping, *Journal of biogeography*, 26, 1025–1038, 1999.
- Mahmood, R., Pielke Sr, R. A., Hubbard, K. G., Niyogi, D., Dirmeyer, P. A., McAlpine, C., Carleton, A. M., Hale, R., Gameda, S., Beltrán-Przekurat, A., et al.: Land cover changes and their biogeophysical effects on climate, *International journal of climatology*, 34, 929–953, 2014.
- 780 Maisongrande, P., Duchemin, B., and Dedieu, G.: VEGETATION/SPOT: an operational mission for the Earth monitoring; presentation of new standard products, *International Journal of Remote Sensing*, 25, 9–14, 2004.
- Marconcini, M., Metz-Marconcini, A., Üreyen, S., Palacios-Lopez, D., Hanke, W., Bachofer, F., Zeidler, J., Esch, T., Gorelick, N., Kakarla, A., et al.: Outlining where humans live, the World Settlement Footprint 2015, *Scientific Data*, 7, 1–14, 2020.
- Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., and Wulder, M. A.: Good practices for estimating area and
785 assessing accuracy of land change, *Remote Sensing of Environment*, 148, 42–57, 2014.
- Ottlé, C., Lescure, J., Maignan, F., Poulter, B., Wang, T., and Delbart, N.: Use of various remote sensing land cover products for plant functional type mapping over Siberia, *Earth System Science Data*, 5, 331–348, 2013.
- Ottosen, T.-B., Lommen, S. T., and Skjøth, C. A.: Remote sensing of cropping practice in Northern Italy using time-series from Sentinel-2, *Computers and Electronics in Agriculture*, 157, 232–238, 2019.
- 790 Pau, S., Edwards, E. J., and Still, C. J.: Improving our understanding of environmental controls on the distribution of C3 and C4 grasses, *Global Change Biology*, 19, 184–196, 2013.
- Perugini, L., Caporaso, L., Marconi, S., Cescatti, A., Quesada, B., de Noblet-Ducoudre, N., House, J. I., and Arneth, A.: Biophysical effects on temperature and precipitation due to land cover change, *Environmental Research Letters*, 12, 053002, 2017.
- Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N.: Plant functional type mapping for earth
795 system models, *Geoscientific Model Development*, 4, 993, 2011.
- Poulter, B., MacBean, N., Hartley, A., Khlystova, I., Arino, O., Betts, R., Bontemps, S., Boettcher, M., Brockmann, C., Defourny, P., et al.: Plant functional type classification for earth system models: results from the European Space Agency’s Land Cover Climate Change Initiative, *Geoscientific Model Development*, 8, 2315–2328, 2015.

- Rechid, D., Davin, E., de Noblet-Ducoudré, N., and Katragkou, E.: CORDEX Flagship Pilot Study" LUCAS-Land Use & Climate Across Scales"-a new initiative on coordinated regional land use change and climate experiments for Europe, EGUGA, p. 13172, 2017.
- 800 Reinhart, V., Fonte, C. C., Hoffmann, P., Bechtel, B., Rechid, D., and Böhner, J.: Comparison of ESA climate change initiative land cover to CORINE land cover over Eastern Europe and the Baltic States from a regional climate modeling perspective, *International Journal of Applied Earth Observation and Geoinformation*, 94, 102 221, 2021a.
- Reinhart, V., Hoffmann, P., and Rechid, D.: LANDMATE PFT land cover dataset for Europe 2015 (Version 1.0),
805 https://doi.org/10.26050/WDCC/LM_PFT_LandCov_EUR2015_v1.0, 2021b.
- Richardson, A. D., Keenan, T. F., Migliavacca, M., Ryu, Y., Sonnentag, O., and Toomey, M.: Climate change, phenology, and phenological control of vegetation feedbacks to the climate system, *Agricultural and Forest Meteorology*, 169, 156–173, 2013.
- Rufin, P., Frantz, D., Ernst, S., Rabe, A., Griffiths, P., Özdoğan, M., and Hostert, P.: Mapping cropping practices on a national scale using intra-annual landsat time series binning, *Remote Sensing*, 11, 232, 2019.
- 810 Saad, R., Koellner, T., and Margni, M.: Land use impacts on freshwater regulation, erosion regulation, and water purification: a spatial approach for a global scale level, *The International Journal of Life Cycle Assessment*, 18, 1253–1264, 2013.
- Santos-Alamillos, F., Pozo-Vázquez, D., Ruiz-Arias, J., and Tovar-Pescador, J.: Influence of land-use misrepresentation on the accuracy of WRF wind estimates: Evaluation of GLCC and CORINE land-use maps in southern Spain, *Atmospheric Research*, 157, 17–28, 2015.
- Sertel, E., Robock, A., and Ormeci, C.: Impacts of land cover data quality on regional climate simulations, *International Journal of Climatology*, 30, 1942–1953, 2010.
815
- Siebert, S., Döll, P., Hoogeveen, J., Faures, J.-M., Frenken, K., and Feick, S.: Development and validation of the global map of irrigation areas, *Hydrology and Earth System Sciences*, 9, 535–547, 2005.
- Skov, F. and Svenning, J.-C.: Potential impact of climatic change on the distribution of forest herbs in Europe, *Ecography*, 27, 366–380, 2004.
- 820 Stehman, S. V.: Sampling designs for accuracy assessment of land cover, *International Journal of Remote Sensing*, 30, 5243–5272, 2009.
- Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: Global distribution of C3 and C4 vegetation: carbon cycle implications, *Global biogeochemical cycles*, 17, 6–1, 2003.
- Sulla-Menashe, D. and Friedl, M. A.: User guide to collection 6 MODIS land cover (MCD12Q1 and MCD12C1) product, USGS: Reston, VA, USA, pp. 1–18, 2018.
- 825 Szelepcsényi, Z., Breuer, H., and Sümegei, P.: The climate of Carpathian Region in the 20th century based on the original and modified Holdridge life zone system, *Central European Journal of Geosciences*, 6, 293–307, 2014.
- Szelepcsényi, Z., Breuer, H., Kis, A., Pongrácz, R., and Sümegei, P.: Assessment of projected climate change in the Carpathian Region using the Holdridge life zone system, *Theoretical and applied climatology*, 131, 593–610, 2018.
- Tatli, H. and Dalfes, H. N.: Defining Holdridge's life zones over Turkey, *International Journal of Climatology*, 36, 3864–3872, 2016.
- 830 Tatli, H. and Dalfes, H. N.: Analysis of temporal diversity of precipitation along with biodiversity of Holdridge life zones, *Theoretical and Applied Climatology*, 144, 391–400, 2021.
- Thompson, C., Beringer, J., Chapin III, F. S., and McGuire, A. D.: Structural complexity and land-surface energy exchange along a gradient from arctic tundra to boreal forest, *Journal of Vegetation Science*, 15, 397–406, 2004.
- Vilar, L., Garrido, J., Echavarría, P., Martínez-Vega, J., and Martín, M. P.: Comparative analysis of CORINE and climate change initiative land cover maps in Europe: Implications for wildfire occurrence estimation at regional and local scales, *International Journal of Applied Earth Observation and Geoinformation*, 78, 102–117, 2019.
835

- Wei, Y., Liu, S., Huntzinger, D., Michalak, A., Viovy, N., Post, W., Schwalm, C., Schaefer, K., Jacobson, A., LU, C., Tian, H., Ricciuto, D., Cook, R., Mao, J., and Shi, X.: NACP MsTMIP: Global and North American Driver Data for Multi-Model Intercomparison, <https://doi.org/10.3334/ORNLDAAC/1220>, 2014.
- 840 Wilhelm, C., Rechid, D., and Jacob, D.: Interactive coupling of regional atmosphere with biosphere in the new generation regional climate system model REMO-iMOVE, *Geoscientific Model Development*, 7, 1093–1114, <https://doi.org/10.5194/gmd-7-1093-2014>, 2014.
- Winter, J. M., Pal, J. S., and Eltahir, E. A.: Coupling of integrated biosphere simulator to regional climate model version 3, *Journal of Climate*, 22, 2743–2757, 2009.
- Wulder, M. A., Franklin, S. E., White, J. C., Linke, J., and Magnussen, S.: An accuracy assessment framework for large-area land cover
845 classification products derived from medium-resolution satellite data, *International Journal of Remote Sensing*, 27, 663–683, 2006.
- Wullschleger, S. D., Epstein, H. E., Box, E. O., Euskirchen, E. S., Goswami, S., Iversen, C. M., Kattge, J., Norby, R. J., van Bodegom, P. M., and Xu, X.: Plant functional types in Earth system models: past experiences and future directions for application of dynamic vegetation models in high-latitude ecosystems, *Annals of botany*, 114, 1–16, 2014.
- Yang, Y., Xiao, P., Feng, X., and Li, H.: Accuracy assessment of seven global land cover datasets over China, *ISPRS Journal of Photogram-*
850 *metry and Remote Sensing*, 125, 156–173, 2017.
- Yue, T., Liu, J., Jørgensen, S. E., Gao, Z., Zhang, S., and Deng, X.: Changes of Holdridge life zone diversity in all of China over half a century, *Ecological Modelling*, 144, 153–162, 2001.
- Yue, T. X., Fan, Z. M., Liu, J. Y., and Wei, B. X.: Scenarios of major terrestrial ecosystems in China, *ecological modelling*, 199, 363–376, 2006.