

1 European pollen-based REVEALS land-cover reconstructions for the 2 Holocene: methodology, mapping and potentials

3
4 Esther Githumbi^{1,2}, Ralph Fyfe³, Marie-Jose Gaillard², Anna-Kari Trondman^{2,4}, Florence Mazier⁵, Anne-
5 Birgitte Nielsen⁶, Anneli Poska^{1,7}, Shinya Sugita⁸, ~~Martin Theuerkau⁹~~, Jessie Woodbridge³, Julien
6 Azuara^{9,10}, Angelica Feurdean^{10,11,12}, Roxana Grindean^{11,12,13}, Vincent Lebreton^{9,10}, Laurent Marquer^{13,14},
7 Nathalie Nebout-Combourieu^{9,10}, Migleĉ Staneĉikaitėĉ^{14,15}, Ioan Tanĉău^{11,12}, Spassimir Tonkov^{15,16},
8 Lyudmila Shumilovskikh^{16,17}, and LandClimII data contributors^{17,18+}.

9
10 ¹Department of Physical Geography and Ecosystem Science, University of Lund, 22362 Lund, Sweden

11 ²Department of Biology and Environmental Science, Linnaeus University, 39182 Kalmar, Sweden

12 ³School of Geography, Earth and Environmental Sciences, University of Plymouth, PL4 8AA Plymouth, United Kingdom

13 ⁴Division of Education Affairs, Swedish University of Agricultural Science (SLU), 23456 Alnarp, Sweden

14 ⁵Environmental Geography Laboratory, GEODE UMR 5602 CNRS, Université de Toulouse Jean Jaurès, 31058 Toulouse,
15 France

16 ⁶Department of Geology, Lund University, 22100 Lund, Sweden

17 ⁷Department of Geology, Tallinn University of Technology, 19086 Tallinn, Estonia

18 ⁸Institute of Ecology, Tallinn University of Technology, 10120 Tallinn, Estonia

19 ~~⁹Institute of Botany and Landscape Ecology, EMAU Greifswald, 1748 Greifswald, Germany~~

20 ^{9,10}Département Homme et Environnement, UMR 7194 Histoire Naturelle de l'Homme Préhistorique, 75013 Paris, France

21 ¹⁰Senckenberg Biodiversity and Climate Research Centre (BiK-F), 60325 Frankfurt am Main, Germany

22 ^{11,12}Department of Geology, Faculty of Biology and Geology, Babeş-Bolyai University, 400084 Cluj-Napoca, Romania

23 ^{12,13} Institute of Archaeology and History of Arts, Romanian Academy, Cluj-Napoca, 400015, Romania

24 ^{13,14}Department of Botany, University of Innsbruck, 6020 Innsbruck, Austria

25 ^{14,15}Institute of Geology and Geography, Vilnius University, Vilnius, LT-03101 Vilnius, Lithuania

26 ^{15,16}Department of Botany, Sofia University St. Kliment Ohridski, 1164 Sofia, Bulgaria

27 ^{16,17}Department of Palynology and Climate Dynamics, Georg-August-University, 37073 Göttingen, Germany

28 ^{17,18+}Team list

29 +A full list of authors appears at the end of the paper.

30

31 *Correspondence to:* Esther Githumbi (esther.githumbi@lnu.se)

Formatted: French (France)

32 **Abstract.** Quantitative reconstructions of past land-cover are necessary for research into the processes involved in climate-
33 human-land interactions. We present the first temporally continuous and most spatially extensive pollen-based land-cover
34 reconstruction for Europe over the Holocene (last 11,700 cal yr BP). We describe how vegetation cover has been quantified
35 from pollen records at a 1°x1° spatial scale using the 'Regional Estimates of VEgetation Abundance from Large Sites'
36 (REVEALS) model. REVEALS is a Landscape Reconstruction Algorithm (LRA) model calculates estimates of past regional
37 vegetation cover in proportions or percentages. REVEALS has been applied to 1128 pollen records across Europe and part of
38 the Eastern Mediterranean-Black Sea-Caspian-Corridor (30°-75°N, 25°W-50°E) to reconstruct the percentage cover of 31
39 plant taxa assigned to 12 plant functional types (PFTs) and three land-cover types (LCTs). A new synthesis of relative pollen
40 productivities (RPPs) available for European plant taxa was performed for this reconstruction. It includes > 1 RPP values for
41 39 taxa, and single values for 15 taxa (total of 54 taxa). As an illustration, we present maps of the results for five taxa (*Calluna*
42 *vulgaris*, *Cerealia-t*, *Picea abies*, ~~deciduous~~ *Quercus* ~~deciduous~~ and ~~evergreen~~ *Quercus* ~~evergreen~~) and three LCTs (open land
43 (OL), evergreen trees (ET) and summer-green trees (ST)) for 8 selected time windows. ~~We discuss the~~The reliability of the
44 REVEALS reconstructions and issues related to the interpretation of the results in terms of landscape openness and human-
45 induced vegetation change ~~and~~~~We then describe~~ the current use of this reconstruction and its future potential utility and
46 development ~~are discussed~~. The quality of input pollen count data (in terms of count size, pollen identification and chronology)
47 and number and type (lake or bog, large or small) of sites used for REVEALS analysis (for each grid cell) ~~Data quality primarily~~
48 ~~determined by amount and standard of data are the main~~ considerations ~~controls on~~ the data-reliability of the REVEALS dataset
49 ~~presented here. A large number of~~ sites with high quality pollen count data will produce more reliable land-cover estimates
50 ~~with lower standard errors compared to a low number of sites with lower quality pollen count data.~~ The REVEALS data
51 presented here can be downloaded from ~~https://doi.pangaea.de/10.1594/PANGAEA.937075?format=html#download~~
52 <https://doi.pangaea.de/10.1594/PANGAEA.937075> (Fyfe et al., 2022).

53

54 1 Introduction

55 The reconstruction of past land cover at global, continental and sub-continental scales is ~~necessary-essential~~ for the evaluation
56 of climate models, land-use scenarios and the study of past climate – land cover interactions. Vegetation plays a significant
57 role within the climate system through biogeochemical and biogeophysical feedbacks and forcings (Foley, 2005; Gaillard et
58 al., 2010, 2015, 2018; Strandberg et al., 2014). Land use has modified the land cover of Europe over Holocene timescales at
59 local, regional and continental scales (e.g. Roberts et al., 2019; Trondman et al., 2015; Woodbridge et al., 2018). Concerted
60 efforts have been made to model land-use and land-cover change (LULCC) over Holocene time scales (e.g. HYDE 3.2 (Klein
61 Goldewijk et al., 2017) and KK10 (Kaplan et al., 2011)). KK10 has been used to assess the impact of the scale of deforestation
62 between 6000 and 200 cal yr BP in Europe on the regional climate in the climate modelling study of Strandberg et al (2014).
63 The KK10-inferred land-cover change resulted in cooling or warming of the regional climate by 1° to 2° depending on the
64 season (winter or summer) and/or geographical location. Major changes in the forest cover of Europe over the Holocene may
65 therefore have had a significant impact on past regional climate, particularly those driven by deforestation since the start of
66 agriculture ~~during the Neolithic period, the timing of which varies slightly in different parts of Europe~~ (Fyfe et al., 2015;
67 Gaillard et al., 2015; Hofman-Kamińska et al., 2019; Nosova et al., 2018; Pinhasi et al., 2005; de Vareilles et al., 2021).
68 Estimating past land-cover change can enable quantification of the scale at which human impact on terrestrial ecosystems
69 perturbed the climate system. This in turn allows us to consider when environmental changes moved beyond the envelope of
70 natural variability (Ruddiman, 2003; Ruddiman et al., 2016). We focus here on the role of LULCC in the climate system;
71 anthropogenic land-cover change can have broader consequences on other processes and changes, such as erosion and fluvial
72 systems (Downs and Piégay, 2019), biodiversity loss (Barnosky et al., 2012), nutrient cycling (Guiry et al., 2018; McLauchlan
73 et al., 2013), habitat exploitation by megafauna (Hofman-Kamińska et al., 2019) and wider ecosystem functioning (Ellis, 2015;
74 Stephens et al., 2019).

75 The Earth System Modelling (ESM) community use LULCC model scenarios, along with dynamic vegetation models, to
76 understand interactions between different components of the earth system in the past (e.g. Gilgen et al., 2019; Smith et al.,
77 2016; He et al., 2014; Hibbard et al., 2010). Disagreement between LULCC scenarios suggests that their evaluation is needed
78 using independent, empirical datasets (Gaillard et al., 2010). Pollen-based reconstruction of past land cover represents probably
79 the best empirical data for this purpose, as ~~fossil pollen is~~ a direct proxy for past vegetation, and ~~thanks to~~ the ubiquity of
80 ~~pollen~~ data across the continent of Europe (e.g. Gaillard et al., 2010, 2018). The ~~Landscape Reconstruction Algorithm (LRA)~~
81 ~~with its two models REgional VEgetation Abundance from Large Sites (REVEALS) and LOVE models~~ (Sugita, 2007a) and
82 ~~LOcal VEgetation (Sugita, 2007b)~~ ~~is~~are the only current land-cover reconstruction approaches based on pollen data that
83 ~~incorporate assumptions that effectively~~ reduces the biases caused by the non-linear pollen-vegetation relationship, ~~and due to~~
84 differences in sedimentary archives, ~~basin size, inter-taxonomic differences in pollen productivity and dispersal characteristics,~~
85 and spatial scales. ~~REVEALS and LOVE are mechanistic models that transforms pollen count data to produce quantitative~~
86 ~~reconstructions of regional (spatial scale > 10⁴ km²) and local (spatial scale = relevant source area of pollen *sensu* Sugita~~

Field Code Changed

Formatted: French (France)

Formatted: French (France)

Field Code Changed

Field Code Changed

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Italic

87 (1993), > ca. 1-5 km radius) vegetation cover, respectively (Sugita, 2007a; 2007b). The REVEALS model was first tested and
88 validated in southern Sweden (Hellman et al., 2008a, 2008b) and later in other parts of Europe and the world (Mazier et al.,
89 2012; Soepboer et al., 2010; Sugita et al., 2010).
90 Moreover, the ‘Regional Estimates of VEgetation Abundance from Large Sites’ (REVEALS) model developed by Sugita
91 (2007a) makes it possible to quantify plant cover from pollen records at a regional spatial scale of ca. 100 km x 100 km (e.g.
92 Hellman et al., 2008a). The first pollen-based REVEALS reconstruction of plant cover over the Holocene covering a large part
93 of Europe (Trondman et al., 2015) was used for the assessment of LULCC scenarios (Kaplan et al., 2017), and helped to
94 evaluate climate model simulations using LULCC scenarios (Strandberg et al., 2014b). A comparison between REVEALS-
95 based open land cover from pollen records and Holocene deforestation-as simulated by HYDE 3.1 and KK10 showed that the
96 REVEALS reconstructions were more similar to the KK10 scenarios than the HYDE 3.1 ones (Kaplan et al., 2017). Therefore,
97 estimates of past plant cover from pollen are essential to both test and constrain LULCC models, and also provide alternative
98 inputs to Earth System Models (ESMs), Regional Climate Models (RCMs) and ecosystem models (Gaillard et al., 2018;
99 Harrison et al., 2020). This allows improved assessments of biogeophysical and biogeochemical forcings on climate due to
100 LULCC over the Holocene (Gaillard et al., 2010; Harrison et al., 2020; Ruddiman et al., 2016; Strandberg et al., 2014b).
101 Europe is of particular interest as one of the global regions of the globe that has experienced major human-induced land-cover
102 transformations. Europe has large N-S and W-E gradients in modern and historical climate and land use (Marquer et al., 2014,
103 2017). Early agriculture dates from the start of the Holocene in the SE Mediterranean region (Palmisano et al., 2019; Roberts
104 et al., 2019; Shennan, 2018), and human impact on vegetation across most of Europe is characterized by early land-cover
105 changes through agriculture (e.g. Marquer et al., 2014; Trondman et al., 2015). There is therefore a clear need to extend
106 quantitative vegetation reconstruction to the whole of Europe, including for the first time the Mediterranean region and
107 additional areas of Eastern Europe. The increase in the spatial coverage of sites, and the extension of temporal coverage to
108 cover the whole-entire Holocene to capture transient vegetation change at sub-millennial time scales is vital to capture
109 information on the transformation of the biosphere by human actions. increase the accuracy of the reconstructions. Europe has
110 a deep history of pollen data production (Edwards et al., 2017) and an open-access repository for pollen records (the European
111 Pollen Database (EPD)) as well as regional pollen repositories (list of databases and access links in sections 2.2 and the Data
112 Availability section). These data repositories result in, resulting in abundant pollen records that can be used for data-driven
113 reconstructions of past vegetation patterns at continental scales. Vegetation reconstructions for Europe based on pollen data
114 have used community-level approaches (Huntley, 1990), biomization methods (Davis et al., 2015; Prentice et al., 1996),
115 modern analogue techniques (MAT; Zanon et al., 2018), and pseudobiomization (Fyfe et al., 2010, 2015; Woodbridge et al.,
116 2014). These approaches capture the major trends in vegetation patterns over the course of the Holocene (Roberts et al., 2018;
117 Sun et al., 2020) and biomization methods have proved useful for evaluation of climate model results (e.g. Prentice and Webb
118 III, 1998). The results of these forms of pollen data manipulation either classify pollen data into discrete classes (e.g.
119 biomization, pseudobiomization) or are semi-quantitative, providing capturing relative change though time based on all pollen
120 taxa within a sample, at best rough estimates of the relationship between forest and open land cover (e.g. MAT). They cannot


Field Code Changed

Field Code Changed

121 achieve reconstructions of the cover of e.g. evergreen versus summer-green trees, or the cover of individual tree and herb taxa.
122 ~~Although useful in summarising palynological change over time based on entire pollen assemblages, such outputs are of limited~~
123 ~~use. They are thus of limited use~~-when differentiation of plant functional types (PFTs) is ~~essential-necessary~~ (e.g. Strandberg
124 et al., 2014). ~~Moreover,~~ Forest cover over the Holocene inferred from pollen records using these approaches differs from
125 forest cover obtained with REVEALS (Hellman et al., 2008a; Roberts et al., 2018); these differences indicate that REVEALS
126 ~~corrects~~ biases due to the non-linearity of the pollen-vegetation relationship ~~are not fully corrected by MAT or (pseudo-)~~
127 ~~biomization approaches.~~

128 In this paper we present the results of the second generation of REVEALS-based reconstruction of plant cover over the
129 Holocene in Europe, the first generation being the reconstruction published by Trondman et al. (2015). This second generation
130 reconstruction is, to date, the most spatially and temporally complete estimate of plant cover for Europe across the Holocene.
131 As with the Trondman et al. (2015) reconstruction, this new dataset is specifically designed to be used in climate modelling.
132 It is performed at a spatial scale of $1^\circ \times 1^\circ$ (ca. 100 km \times 100 km) across 30°-75°N, 25°W-50°E (Europe and part of the Eastern
133 Mediterranean-Black Sea-Caspian-Corridor) (Fig. 1). The number of pollen records used (1128), the area covered and time
134 length (entire Holocene) are a significant advance on the results presented in Trondman et al. (2015), which used 636 pollen
135 records covering NW Europe (including Poland and the Czech Republic and excluding western Russia and the Mediterranean
136 area), and produced estimates for five time windows (in cal yr BP, hereafter abbreviated BP): 6200-5700, 4200-3700, 700-
137 350, 350-100 BP and 100 BP to present. Marquer et al. (2014, 2017) produced continuous REVEALS reconstructions over the
138 entire Holocene, however only for transects of individual sites (19 pollen records) and groups of grid cells around them.

Formatted: English (United Kingdom)



139

140 **Figure 1: Study region showing site coverage. A.) Colours differences represent different modern biomes regions (purple = boreal,**
 141 **yellow = temperate, blue = Mediterranean) while size and colour of circle represents site type and size (see caption in panel A). B.)**
 142 **Grid cell reliability dependent on number of pollen records. black grid cells grey grid cells= reliable results, black grid cells grey grid**
 143 **cells = less reliable results. Reliable = 1 large lake or >>2 small lake(s) and/or small bog(s). less reliable = 1 bog (large or small) or 1**
 144 **small lake.**

Formatted: Font: 9 pt, Bold

145 The 1° × 1° scale corresponds approximatively to the spatial extent of pollen-based REVEALS reconstructions as evaluated
 146 by empirical studies in Europe (Hellman et al., 2008b; Soepboer et al., 2010): the REVEALS estimated abundances of plant
 147 taxa (in percentage cover) correspond most closely to the plant abundances in an area of ca. 100 km x 100 km or larger (see
 148 Li et al., 2020 for further discussion of the spatial scale of REVEALS reconstructions). This spatial scale is appropriate for
 149 climate models that typically use spatial scales of 0.25° to 1° (Gaillard et al., 2010). ~~REVEALS is a mechanistic model that
 150 transforms pollen count data to produce quantitative reconstructions of regional vegetation (Sugita, 2007a). The model was
 151 first tested and validated in southern Sweden (Hellman et al., 2008a, 2008b) and later in other parts of Europe and the world
 152 (Soepboer et al., 2010; Sugita et al., 2010).~~

Formatted: French (France)

Formatted: French (France)

Formatted: French (France)

Formatted: French (France)

Field Code Changed

Field Code Changed

Formatted: French (France)

Formatted: French (France)

Field Code Changed

Formatted: French (France)

Formatted: French (France)

153 2 Methods

154 2.1 REVEALS model and parameters

155 [The REVEALS model \(Sugita, 2007a\)](#) is a generalized version of the R-Value model of Davis ([Davis, 1963](#)). [The development](#)
 156 [of pollen-vegetation modelling from the R-Value model, via the ERV models of Andersen \(Andersen, 1970\) and Parsons and](#)
 157 [Prentice \(Parsons and Prentice, 1981\) through to the REVEALS model is described in detail in numerous earlier papers \(e.g.](#)
 158 [\(Broström et al., 2004; Bunting et al., 2013b; Sugita, 1993, 2007a\). Using simulations Sugita \(2007a\) showed that “large lakes”](#)
 159 [represent regional vegetation, i.e. between-lake differences in pollen assemblages are very small, which was the case for lakes](#)
 160 [≥50ha in the simulations \(Sugita, 2007a\). Tests using modern pollen data from surface lake sediments have shown that pollen](#)
 161 [assemblages from lakes ≥50ha are appropriate to estimate regional plant cover using the REVEALS model \(e.g. tests by](#)
 162 [Hellman et al. \(2008a and b\) in southern Sweden and by Sugita et al. \(2010\) in northern America\).](#)

163
 164 [Following the equation below, the](#)The REVEALS model, [represented by the model below \(equation 1\)](#), calculates estimates
 165 of regional vegetation abundance in proportions or percentage cover using fossil pollen counts from [“large lakes”](#) (Sugita,
 166 2007a).

Formatted: Line spacing: 1.5 lines

$$167 \hat{V}_i = \frac{n_{i,k} / \hat{\alpha}_i \int_R^{Z_{\max}} g_i(z) dz}{\sum_{j=1}^m \left(\frac{n_{j,k}}{\hat{\alpha}_j} \int_R^{Z_{\max}} g_j(z) dz \right)} = \frac{n_{i,k} / \hat{\alpha}_i K_i}{\sum_{j=1}^m (n_{j,k} / \hat{\alpha}_j K_j)} \quad (1)$$

- 168 • \hat{V}_i is the estimate of the regional vegetation abundance for taxon i (proportion or percentage).
- 169 • $n_{i,k}$ is the pollen count of taxon i at site k .

- 170 • $\hat{\alpha}_i$ is the estimate of pollen productivity (relative pollen productivity, RPP) for taxon i .
- 171 • z is the distance between the centre of the sedimentary basin and the pollen source.
- 172 • $g_i(z)$ is the pollen dispersal/deposition function for taxon i expressed as a function of distance z . Fall speed of pollen
- 173 (FSP), wind speed and atmospheric conditions are parameters needed to calculate this function.
- 174 • R is the radius of the sedimentary basin.
- 175 • Z_{\max} is the maximum distance within which most pollen originates (i.e. the maximum spatial extent of the regional
- 176 vegetation).
- 177 • m is the total number of taxa included,
- 178 • $K_i = \int_R^{Z_{\max}} g_i(z) dz$ is the “pollen dispersal-deposition coefficient” of taxon i from the border of the study site
- 179 (distance from the pollen sample corresponding to the radius R of the lake) to Z_{\max} .

181 ~~The REVEALS model was developed for pollen records from large lakes and~~ the assumptions of the REVEALS model are
 182 listed in Sugita (2007a). ~~The model was tested~~has been tested and validated in Europe (Hellman et al., 2008a; Mazier et al.,
 183 2012; Soepboer et al., 2010) and northern America (Sugita et al., 2010). Using simulations Sugita (2007a) demonstrated ~~using~~
 184 ~~simulations that, that~~ in theory, the model can ~~also~~ also be applied ~~on~~ to pollen records from multiple “small lakes” (< 50 ha),
 185 i.e. lakes for which between lake differences in pollen assemblages can be large; however, the REVEALS estimates ~~will~~
 186 ~~however using pollen records from “small lakes”~~ generally have larger standard errors (SE) than those based on pollen data
 187 from large lakes. ~~The latter was demonstrated for empirical pollen records from large lakes versus small sites (lakes and bogs)~~
 188 ~~by Trondman et al. (2016) in southern Sweden and Mazier et al. (2012) in the Czech Republic. Moreover, although~~ Although
 189 the application of the model on pollen data from bogs violates the model assumption that no plants grow on the basin,
 190 REVEALS can be applied using models of pollen dispersal and deposition for lakes or bogs. The Prentice’s model (Prentice,
 191 1985; 1988) describes deposition of pollen at a single point in a deposition basin and is suitable for pollen records from bogs.
 192 Sugita (1993) developed the “Prentice-Sugita model” that describes pollen deposition in a lake, i.e. on its entire surface with
 193 a subsequent mixing in the water body before deposition at the lake bottom. The original versions of both models use the
 194 Sutton model of pollen dispersal, i.e. a Gaussian plume model from a ground-level source under neutral atmospheric conditions
 195 (Sutton, 1953). A Lagrangian stochastic model of dispersion has also been introduced as an alternative for the description of
 196 pollen dispersal in models of the pollen-vegetation relationship in general, and in the REVEALS model in particular
 197 (Theuerkauf et al., 2013; 2016). It is difficult, in both theory and practice, to eliminate the effects of pollen coming from plants
 198 growing on sedimentary basins (e.g. Poaceae and Cyperaceae in bogs) on regional vegetation reconstruction. Previous studies
 199 have assessed the impacts of the violation of this assumption on ~~the~~ REVEALS outcomes (Mazier et al., 2012; Sugita et al.,
 200 2010; Trondman et al., 2016, 2015). An empirical study in southern Sweden (Trondman et al., 2016a) indicated that REVEALS
 201 estimates based on pollen records from multiple small sites (lakes and/or bogs) are similar to the REVEALS estimates based
 202 on pollen records from large lakes in the same region. The results also suggested that increasing the number of pollen records

Formatted: French (France)

203 significantly decreased the standard error of the REVEALS estimates, as expected based on simulations (Sugita, 2007a). It is
204 therefore appropriate to use pollen records from small bogs to increase the number of pollen records included in a REVEALS
205 reconstruction, following the protocol of the first generation REVEALS reconstruction for Europe (Mazier et al., 2012;
206 Trondman et al., 2015). However, REVEALS estimates of plant cover using pollen assemblages from large bogs only should
207 be interpreted with great caution (Mazier et al., 2012; see also section 4, Discussion).

208 The inputs needed to run the REVEALS model are: original pollen counts; relative pollen productivity estimates (RPPs) and
209 their standard deviation; fall speed of pollen (FSP); basin type (lake or bog); size of basin (radius); maximum extent of regional
210 vegetation; and wind speed (m/s) and atmospheric conditions. FSP can be calculated using measurements of the pollen grains
211 and the Stokes' law (Gregory, 1973). RPPs of major plant taxa can be estimated using datasets of modern pollen assemblages
212 and related vegetation and the Extended R-Value model (e.g. Mazier et al., 2008). RPPs exist for a large number of European
213 plant taxa, and syntheses of FSPs and RPPs were published earlier by ~~Broström in 2008 and Mazier in 2012~~ (Broström et al.,
214 ~~(2008) and~~ Mazier et al., (2012). The latter was used in the “first generation” REVEALS reconstruction (Trondman et al.,
215 2015). A new synthesis of European RPPs was performed for this “second generation” reconstruction (Appendices A, B, and
216 C). Preparation of data from individual pollen records, and the values of model parameters used, are described below (sections
217 2.2 and 2.3).

218 2.2 Pollen records – data compilation and preparation

219 1143 pollen records from 29 European countries and the Eastern Mediterranean-Black Sea-Caspian-Corridor were obtained
220 from databases and individual data contributors. The contributing databases include: the European Pollen Database (Fyfe et
221 al., 2009; Giesecke et al., 2014); the Alpine Palynological database (ALPADABA; Institute of Plant Sciences, University of
222 Bern; now also archived in EPD); the Czech Quaternary palynological database (PALYCZ; ~~Kuneš et al., 2009~~ Kuneš et al.,
223 2009); PALEOPYR (Lerigoleur et al., 2015); and datasets compiled within synthesis projects from the Mediterranean region
224 (Fyfe et al., 2018; Roberts et al., 2019) and the Eastern Mediterranean-Black Sea-Caspian-Corridor (EMBSecBIO project;
225 Marinova et al., 2018) (see Fig. 1 for map and; Data availability section for data location and team list for individual pollen
226 data contributors). We followed the protocols and criteria published in Mazier et al. (2012) and Trondman et al. (2015) for
227 selection of pollen records and application of the REVEALS model. Available pollen records were filtered based on criteria
228 including basin type (to exclude archaeological sites and marine records) and quality of chronological control (excluding sites
229 with poor age-depth models or fewer than three radiocarbon dates). This resulted in 1128 pollen records from lakes and bogs,
230 both small and large. The rationale behind the use of pollen records from small sites is based on the knowledge that REVEALS
231 estimates based on pollen records from multiple sites provide statistically validated approximations ~~reasonable approximations~~
232 of the regional cover of plant taxa (e.g. Trondman et al., 2016; see details under section 2.1 on the REVEALS model).
233 The taxonomy and nomenclature of pollen morphological types from the 1128 pollen records were harmonised. The pollen
234 morphological types were then consistently assigned to one of 31 RPP taxa (Table 1; see section 2.3 for details on the RPP
235 dataset used in this study), following the protocol outlined in Trondman et al. (2015: SI-2 with examples of harmonization

236 ~~between pollen-morphological types and RPP taxa~~. This process takes into account plant morphology, biology, and ecology
237 of the species that are included in each pollen morphological type ~~(see Trondman et al., 2015 for examples of harmonization~~
238 ~~between pollen-morphological types and RPP taxa)~~. ~~In this way~~Consequently, RPP-harmonized pollen count data were
239 produced for each of the 1128 pollen records. It should be noted that the EMBSecBIO data does not contain pollen counts
240 from cultivars, i.e. pollen from cereals and cultivated trees were deleted from the pollen records (Marinova et al., 2018).
241 Therefore, the cover of agricultural land (represented by ~~Cereals-cereals~~ in this reconstruction) will always be zero in the
242 Eastern Mediterranean-Black Sea-Caspian-Corridor in grid cells ~~including-with~~ only pollen records from EMBSecBIO,
243 ~~although-even though~~ agriculture did occur in the region from early Neolithic.
244 For application of REVEALS, an age-depth model (in cal yr BP) is required for each pollen record. We used the author's
245 original ~~published~~ model, the model available in the contributing database or, where necessary, a new age-depth model
246 constructed following the approach in Trondman et al. (2015). The age-depth model for each pollen record is used to aggregate
247 RPP-harmonised pollen count data into 25 time windows ~~aeross-throughout~~ the Holocene following a standard ~~time~~ division
248 used in Mazier et al. (2012) and Trondman et al. (2015), ~~which were~~ later adopted by the ~~Past Global Changes (PAGES)~~
249 LandCover6k working group (Gaillard et al., 2018). The first three time windows (present–100 BP (where present is the
250 ~~yeardate~~ of coring), 100–350 BP; 350–700 BP) capture the major human-induced land-cover changes since ~~the~~ Early Middle
251 Ages. Subsequent time windows are contiguous 500-year long intervals (e.g. 700–1200 BP, 1200–1700 BP, 1700–2200 BP,
252 etc.) with the oldest interval representing the start of the Holocene (11200–11700 BP). The use of 500-year long time windows
253 is motivated by the necessity to obtain sufficiently large pollen counts for reliable REVEALS reconstructions. Since the size
254 of the error on the REVEALS estimate partly depends on the size of the pollen count (Sugita, 2007a), the length of the time
255 window should be a reasonable compromise to ensure both a useful time resolution of the reconstruction and an acceptable
256 reliability of the REVEALS estimate of plant cover (Trondman et al., 2015).

257 Table 1: Land-cover types (LCTs) and Plant Functional Types (PFTs) according to Wolf et al. (2008) and their corresponding pollen
 258 morphological types. Fall speed of pollen (FSP) and the mean relative pollen productivity (RPP) estimates from using the new RPP
 259 synthesis (see section 2.3 and Appendices A-C for details) with their standard errors in brackets (see text for more explanations).
 260 *The FSP values of evergreen Quercus evergreen-t. and Mediterranean Ericaceae according to the original study (Mazier,
 261 unpublished) are 0.015 and 0.051, respectively (see Appendix B, Table B.3). The value of 0.035 (FSP of deciduous Quercus deciduous
 262 t.) and 0.038 (FSP of boreal-temperate Ericaceae) were used instead (see discussion in section 4.2 for explanation). t = type e.g.
 263 evergreen Quercus evergreen-t., RPP used in this study are relative to grass pollen productivity where Poaceae = 1 (indicated in bold).

Land-cover types (LCTs)	PFT	PFT definition	Plant taxa/Pollen-morphological types	FSP (m/s)	PPE-RPP (SD)
Evergreen trees (ET)	TBE1	Shade-tolerant evergreen trees	<i>Picea abies</i>	0.056	5.437 (0.097)
	TBE2	Shade-tolerant evergreen trees	<i>Abies alba</i>	0.12	6.875 (1.442)
	IBE	Shade-intolerant evergreen trees	<i>Pinus sylvestris</i>	0.031	6.058 (0.237)
	MTBE	Mediterranean shade-tolerant broadleaved evergreen trees	<i>Phillyrea</i>	0.015	0.512 (0.076)
			<i>Pistacia</i>	0.03	0.755 (0.201)
			<u>Evergreen Quercus evergreen-t.</u>	0.035*	11.043 (0.261)
	TSE	Tall shrub, evergreen	<i>Juniperus communis</i>	0.016	2.07 (0.04)
	MTSE	Mediterranean broadleaved tall shrubs, evergreen	Ericaceae [‡]	0.038*	4.265 (0.094)
<i>Buxus sempervirens</i>			0.032	1.89 (0.068)	
Summer green trees (ST)	IBS	Shade-intolerant summer-green trees	<i>Alnus glutinosa</i>	0.021	13.562 (0.293)
			<i>Betula</i>	0.024	5.106 (0.303)
	TBS	Shade-tolerant summer-green trees	<i>Carpinus betulus</i>	0.042	4.52 (0.425)
			<i>Carpinus orientalis</i>	0.042	0.24 (0.07)
			<i>Castanea sativa</i>	0.01	3.258 (0.059)
			<i>Corylus avellana</i>	0.025	1.71 (0.1)
			<i>Fagus sylvatica</i>	0.057	5.863 (0.176)
			<i>Fraxinus</i>	0.022	1.044 (0.048)
			<u>Deciduous Quercus deciduous-t.</u> [‡]	0.035	4.537 (0.086)
			<i>Tilia</i>	0.032	1.21 (0.116)
			<i>Ulmus</i>	0.032	1.27 (0.05)
TSD	Tall shrub, summer-green	<i>Salix</i>	0.022	1.182 (0.077)	
Open land (OL)	LSE	Low shrub, evergreen	<i>Calluna vulgaris</i>	0.038	1.085 (0.029)
	GL	Grassland - all herbs	<i>Artemisia</i>	0.025	3.937 (0.146)
			Amaranthaceae/Chenopodiaceae	0.019	4.28 (0.27)
			Cyperaceae	0.035	0.962 (0.05)
			<i>Filipendula</i>	0.006	3 (0.285)
			Poaceae	0.035	1 (0)
			<i>Plantago lanceolata</i>	0.029	2.33 (0.201)
			<i>Rumex acetosa-t</i>	0.018	3.02 (0.278)
	AL	Agricultural land - cereals	Cerealia-t	0.06	1.85 (0.380)
			<i>Secale cereale</i>	0.06	3.99 (0.320)

Formatted: Font: Bold

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold, Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold

Formatted: Font: Not Italic

Formatted: Font: Not Italic

266

267 2.3 Model parameter setting

268 For the purpose of this study, a new synthesis of the RPP values available for European plant taxa was performed in 2018-
269 2019 based on the latest synthesis by Mazier et al. (2012) and additional RPP studies published since then (Appendix A-C). It
270 provides new alternative RPP datasets for ~~the whole of~~ Europe, including or excluding plant taxa with dominant entomophily,
271 ~~and~~ with the important addition of plant taxa from the Mediterranean area (Table A1). ~~The location of studies included in the~~
272 ~~RPP synthesis are shown in Fig. C1 and related information is provided in Table C1.~~ The selection of RPP studies, RPP values
273 (shown in Appendix B, Tables B1 and B2) and calculation of mean RPP and their standard error (SD) for Europe are explained
274 in Appendix C. ~~The location of studies included in the RPP synthesis is shown in Fig. C1 and related information is provided~~
275 ~~in Table C1.~~ The synthesis includes a total of 54 taxa for which RPP values are available (Tables B1 and B2), 39 taxa from
276 studies in boreal and temperate Europe, and 15 taxa from studies in Mediterranean Europe of which seven include exclusively
277 sub-Mediterranean and Mediterranean taxa: *Buxus sempervirens*, *Carpinus orientalis*, *Castanea sativa*, Ericaceae
278 (Mediterranean species), *Phillyrea*, *Pistacia* and ~~evergreen Quercus evergreen~~-type. RPP values are available from both
279 boreal/temperate and Mediterranean Europe for seven taxa: i.e. Poaceae (reference taxon), *Acer*, *Corylus avellana*, Apiaceae,
280 *Artemisia*, *Plantago lanceolata* and Rubiaceae (Table B2). Table A1 presents the new RPP dataset for the 54 plant taxa and,
281 for comparison, the mean RPP values from Mazier et al. (2012) and from the recent synthesis by Wiczorek & Hertzschuh
282 (2020). Moreover, comparison with the RPP values of three studies not used in our synthesis is shown in Table A2. For the
283 REVEALS reconstructions presented in this paper, we excluded strictly entomophilous taxa, which resulted in a total of 31
284 taxa (Table 1). The excluded taxa are Compositae SF Cichorioideae (or Asteraceae subfamily Cichorioideae), *Leucanthemum*
285 (*Anthemis*-t., *Potentilla*-t., *Ranunculus acris*-t., and Rubiaceae. We included entomophilous taxa that are known to be
286 characterised by some anemophily, e.g. *Artemisia*, ~~Amaranthaceae~~/Chenopodiaceae, Rubiaceae, and *Plantago lanceolata*. We
287 excluded plant taxa with only one RPP value except Chenopodiaceae, *Urtica*, *Juniperus*, and *Ulmus*, and the seven exclusively
288 sub-Mediterranean and Mediterranean taxa mentioned above.

289 The FSP values (Tables 1 and A1) for boreal and temperate plant taxa were obtained from the literature (Broström et al., 2008;
290 Mazier et al., 2012); these values were in turn extracted from Gregory (1973) for trees, and calculated based on pollen
291 measurements and Stokes' law for herbs (Broström et al., 2004b). FSPs for Mediterranean taxa (*Buxus sempervirens*, *Castanea*
292 *sativa*, Ericaceae (Mediterranean species), *Phillyrea*, *Pistacia*, and *Quercus* evergreen type) were obtained by using pollen
293 measurements and Stokes' law (Mazier et al., unpublished); the FSP of *Carpinus betulus* (Mazier et al., 2012) was used for
294 *Carpinus orientalis* (Grindean et al., 2019).

295 The site radius was obtained from original publications where possible. Sites in the EMBSecBIO were classified as small
296 (0.01-1 km²), medium (1.1-50 km²) or large (50.1-500 km²). These were assigned radii of 399m, 2921m and 10000 m,
297 respectively. Where a site's radius could not be determined from publication, it was geolocated in Google Earth and the area
298 of the site was measured. A radius value was extracted assuming that a site shape is circular (Mazier et al., 2012). A constant

Field Code Changed

299 wind speed of 3 m/s, assumed to correspond approximatively to the modern mean annual wind speed in Europe, was used
300 following Trondman et al. (2015). Z_{\max} (maximum extent of the regional vegetation) was set to 100 km. Z_{\max} and wind speed
301 influence on REVEALS estimates has been evaluated earlier in simulation and empirical studies (Gaillard et al., 2008; Mazier
302 et al., 2012; Sugita, 2007a), [which support the values used for these parameters](#). Atmospheric conditions are assumed to be
303 neutral (Sugita, 2007a).

304 2.4 Implementation of REVEALS

305 REVEALS was implemented using the REVEALS function within the LRA R-package [of \(Abraham et al., 2014\)](#) (see Code
306 availability, section 6). The function enables the use of deposition models for bogs (Prentice's model) and lakes (Sugita's
307 model), and two dispersal models (a Gaussian plume model, and a Lagrangian stochastic model taken from the DISCOVER
308 package (Theuerkauf et al., 2016)). Within this study the Gaussian plume model was applied. The REVEALS model was run
309 on all pollen records within each $1^\circ \times 1^\circ$ grid cell across Europe. The REVEALS function [runs-is applied to](#) lake and bog sites
310 separately within each $1^\circ \times 1^\circ$ grid cell, and combines results (if [there is](#) more than one pollen record per cell) to produce a
311 single mean cover estimate (in proportion) and mean standard error (SE) for each taxon. The formulation of the SE [can-beis](#)
312 found in Appendix A of Sugita (2007a). The REVEALS SE [takes-into-accountaccounts for](#) the standard deviations on the
313 relative pollen productivities for the individual pollen taxa (Table 1) and the number of pollen grains counted in the sample
314 (Sugita, 2007a). The uncertainties of the averaged REVEALS estimates of plant taxa for a grid cell are calculated using the
315 delta method (Stuart and Ord., 1994), and expressed as the SEs derived from the sum of the within- and between-site variations
316 of the REVEALS results in the grid cell. The delta method is a mathematical solution to the problem of calculating the mean
317 of individual SEs (see Li et al., 2020, Appendix C, for [the](#) formula and further details). Results of the REVEALS function are
318 extracted by time window, producing 25 matrices of mean REVEALS [land-cover](#) estimates [of-cover](#) and 25 matrices of
319 corresponding mean SEs for each of the 31 RPP taxa and each grid cell. The 31 RPP taxa are also assigned to 12 plant functional
320 types (PFTs) and three land-cover types (LCTs) (Table 1) and their mean REVEALS estimates calculated. These PFTs follow
321 Trondman et al. (2015), with the addition of two PFTs for Mediterranean vegetation not reconstructed in earlier studies:
322 Mediterranean shade-tolerant broadleaved evergreen trees (MTBE) and Mediterranean broadleaved tall shrubs, evergreen
323 (MTSE). The mean SE for LCTs and PFTs including more than one plant taxon are calculated using the delta method (Stuart
324 and Ord., 1994), as [explained-described](#) above.

325 2.5 Mapping of the REVEALS estimates

326 To illustrate the information that the new REVEALS reconstruction provides, we present and describe [in \(section 3\)](#) maps of
327 the REVEALS estimates (% cover) and their associated SEs for the three LCTs (Fig. 2 to 4) and five taxa for eight selected
328 time windows: the five taxa are Cerealia-t and *Picea abies* (Fig. 5 and 6), ~~and~~ *Calluna vulgaris*, deciduous *Quercus* [type \(t\)](#),
329 and evergreen *Quercus* [t](#) (Fig. D1-D3). The selection of the five taxa and eight time windows is motivated essentially by
330 notable changes in spatial distribution of these taxa through time, with higher resolution for recent times characterised by the

331 largest and most rapid human-induced changes in vegetation cover. For visualisation purposes, the estimates are mapped in
332 nine % cover classes. These fractions are the same for the three LCTs (Figures 2-4), and the mapped output can therefore be
333 directly compared. In contrast, the colour scales used for the five taxa vary between maps depending on the abundance of the
334 PFT/taxon (Fig. 5 and 6, D1-D3). Different taxa thus have different scales and maps cannot be directly compared. We visualise
335 uncertainty in our data by plotting the SE as a circle inside each grid cell; it is the coefficient of variation (CV, i.e. the standard
336 error divided by the REVEALS estimate). Circles are scaled to fill the grid cell if the SE is equal or greater than the mean
337 REVEALS estimate (i.e. $CV \geq 1$). Grid-based REVEALS results that are based on pollen records from just 1 large bogs, or
338 single small bogs or lakes, provide lower quality results (see section 2.1 on the REVEALS model, and discussion section 4.1).
339 Grid cells quality are detailed in Table GC_quality_by_TW (see section 5, Data availability), by time window. ~~It should be~~
340 ~~stressed that the~~. The percentage scale ranges we use here are different from those used in the maps of Trondman et al. (2015)
341 and, therefore, the data visualisation we present cannot be directly compared with that of the 2015 study.

342 3 Results

343 ~~The The complete REVEALS land cover reconstruction full results, or REVEALS~~ dataset, includes mean REVEALS values
344 (in proportions) and their related mean SE for 31 individual tree and herb taxa, twelve PFTs and three LCTs for each grid cell
345 in 25 consecutive time windows of the Holocene (11.7 k BP to present) (~~see Data availability section~~). Here, results are
346 illustrated by maps of the three LCTs (Fig. 2-4) and five taxa (Fig. 5-6, D1-D3). The presented maps are not part of the
347 published dataset archived in [the PANGAEA Pangea online public database](#) (see Data availability, section 5), they are examples
348 of how the data can be [visually](#) presented and what they can be used for.


349 3.1 Land-cover types

350 The three land-cover types are evergreen trees (ET), summer-green trees (ST) and open land (OL). ET includes six PFTs which
351 are composed of nine pollen-morphological types (~~from~~ here after referred to as taxa). ST includes four PFTs which are
352 composed of eleven taxa while OL includes three PFTs that are in turn composed of nine taxa (Table 1).

353 3.1.1 Open Land (OL)

354 At the start of the Holocene, [open land \(OL\)](#) (Fig. 2) ~~exhibits a~~ higher [cover](#) in western Europe where it generally exceeds
355 80% ~~cover~~, compared with central Europe where it is more typically ~60%. There is a general decline in OL cover through the
356 early Holocene. At 5700-6200 BP most grid cells in central Europe have OL cover values between 10-50%. In western Europe,
357 whilst OL is generally reduced, several grid cells on the Atlantic fringe of northern Scotland persistently maintain 80-90% OL
358 cover. OL increases from the mid-Holocene, and by 2700-3200 BP the [British Isles](#) [United Kingdom](#), France, Germany and
359 the Mediterranean region have grid cells recording OL values >70%. In central, northern and eastern Europe grid cells OL


360 values vary between 10 - 70% at 2700-3200 BP. Time windows from the last two millennia show a consistent increase in OL
361 with values >60% across most of central, southern and western Europe and 20-70% in northern Europe.



363 Figure 12. Grid-based REVEALS estimates of Open Land (OL) cover for eight Holocene time windows. Percentage cover of open
364 land in 10% intervals represented by increasingly darker shades of green from 20%. Grey cells: cells without pollen data for the
365 time window, but with pollen data in other time windows. Circles in grid cells represent the coefficient of variation (CV; the standard
366 error divided by the REVEALS estimate). When $SE \geq REVEALS$ estimate, the circle fills the entire grid cell and the REVEALS
367 estimate is not different from zero. This occurs mainly where REVEALS estimates are low.

368 3.1.2 Evergreen Trees (ET)

369 The cover of eEvergreen tree (ET) cover (Fig. 3) at 9700-10200 BP is <30% across Europe, and by 7700-8200 BP fewer than
370 30 grid cells show ET >50%. ET percentage cover slowly increases through the early Holocene and at 5700-6200 BP groups
371 of grid cells in southern Europe record >80%, while in northern Europe ET cover ranges between 10% and 60%. There is a
372 consistent increase in ET cover over Europe during the mid- and late-Holocene with ET cover peaking at 2700-3200 BP before
373 starting to reduce. Across western parts of Europe, including the [British Isles](#) [United Kingdom](#), western France, Denmark, and
374 the Netherlands ET never exceeds 20% cover.




376 **Figure 23. Grid-based REVEALS estimates of Evergreen Tress (ET) cover for eight Holocene time windows.** See caption of Figure
377 2 for more explanations. Percentage cover of Evergreen Trees in 10% intervals represented by increasingly darker shades of green
378 from 20%. Grey cells: cells without pollen data for the time window, but with pollen data in other time windows. Circles in grid cells
379 represent the coefficient of variation (CV; the standard error divided by the REVEALS estimate). When SE > REVEALS estimate,
380 the circle fills the entire grid cell and the REVEALS estimate is not different from zero. This occurs mainly where REVEALS
381 estimates are low.

382

383 3.1.3 Summer-green Trees (ST)

384 The ~~cover estimate of cover of~~ summer-green trees (ST) ~~eovers~~ (Fig. 4) in the early Holocene at 9700-10200 BP is >40%
385 across Europe. A small number (<10) of grid cells in northern, western, central and southern Europe have cover >60%. This
386 significantly increases to 5700-6200 BP, at which time ST cover is >60% in central Europe, and 40-60% in northern Europe.
387 ST cover remains <20% in southern Europe. From 5700-6200 BP there is a steady decline in ST cover across Europe. At
388 2700-3200 BP only central Europe has ST cover >50% while ~~the rest of Europe exhibits values values are~~ <50% ~~for the rest of~~
389 ~~Europe~~. There is a consistent decline over the last two millennia BP. Most of Europe has ST cover <30% in the ~~two most last~~
390 ~~recent~~ time windows (100-350 ~~BP~~ and 100 BP-present), except for a group of grid cells in the southern Baltic states and
391 scattered records elsewhere.

Commented [EG1]: So that each figure and its caption is independent



393 Figure 34. Grid-based REVEALS estimates of Summer-green Trees (ST) cover for eight Holocene time windows. Percentage cover
394 of ST in 10% intervals represented by increasingly darker shades of green from 20%. Grey cells: cells without pollen data for the
395 time window, but with pollen data in other time windows. Circles in grid cells represent the coefficient of variation (CV; the standard
396 error divided by the REVEALS estimate). When SE ≥ REVEALS estimate, the circle fills the entire grid cell and the REVEALS
397 estimate is not different from zero. This occurs mainly where REVEALS estimates are low. See caption of Figure 2 for more
398 explanations.

399 3.2 Selected taxa

400 In terms of PFTs, *Cerealia-type (t)* is assigned to agricultural land (AL), *Picea abies* to shade tolerant evergreen trees (TBE1:
401 *Picea abies* is the only taxon in this PFT), *Calluna vulgaris* to low evergreen shrubs (LSE: *Calluna vulgaris* is the only taxon
402 in this PFT), deciduous *Quercus t.* to shade tolerant summer-green trees (TBS), and evergreen *Quercus* to Mediterranean
403 shade-tolerant broadleaved evergreen trees (MTBE) (Table 1).

404 3.2.1 *Cerealia-type*

405 *Cerealia-t.* (Fig. 5) is recorded throughout the Holocene with 10-15% as the maximum cover. *Cerealia-t.* is present in southern
406 Europe at 9700-10200 BP with several grid cells recording >5 to 10%. Whilst such values are rare, there are scattered grid
407 cells in central and western Europe recording the presence of *Cerealia-t.* at very low levels (0.5-1%). These values have high
408 SE (greater than the REVEALS estimate) and are therefore not different from zero; they correspond to single findings of
409 *Cerealia-t.* By 5700-6200 BP, grid cells in Estonia and France record 3-5% cover, and several regions within central and
410 western Europe record 0-5% (0.5-1%), although with high SEs. At 2700-3200 BP, *Cerealia-t.* is recorded across central and
411 western Europe in the British Isles United Kingdom, France, Germany, and Estonia with low values. In Norway, Sweden and
412 Finland it has 0-1% cover with high SEs. The highest cover (>5%) is observed across Europe from 1200 BP.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic


Formatted: Font: Not Italic



414 Figure 5. Grid-based REVEALS estimates of Cerealia \bar{t}_i cover for eight Holocene time windows. Percentage cover in 0.5%
415 intervals between 0 and 3%, 1% intervals between 3 and 5, and 5% interval between 5 and 10%. Percentage cover of Cerealia \bar{t}_i in
416 intervals represented by increasingly darker shades of green from 1-1.5%. Grey cells: cells without pollen data for the time window,
417 but with pollen data in other time windows. Circles in grid cells represent the coefficient of variation (CV; the standard error divided
418 by the REVEALS estimate). When $SE \geq REVEALS$ estimate, the circle fills the entire grid cell and the REVEALS estimate is not
419 different from zero. This occurs mainly where REVEALS estimates are low. See caption of Figure 2 for more explanations.

420 3.2.2 *Picea abies*

421 *Picea abies* (Fig. 6) cover (Fig. 6) is low (1-2%) at 9700-10200 BP, although a number of grid cells in central and eastern
422 Europe record values between 30 and 50%. By 7700-8200 BP, grid cells recording 30-50% cover are observed in more regions
423 of central and eastern Europe than earlier (Russia, Estonia, Romania, Slovakia and Austria). At 5700-6200 BP, almost all of OF
424 central Europe has consistent but low cover of *Picea abies*; values are higher towards northeast Europe (Russia, Estonia,
425 Latvia, Belarus and Lithuania), up to 30-50%. By 2700-3200 BP the cover of *Picea abies* has increased across central (ca.
426 10%) and northeast Europe (>30%). From 1200 BP, *Picea abies* is recorded in northern Europe, particularly in Norway and
427 Sweden with some grid cells recording 25-50% cover.



429 **Figure 6. Grid-based REVEALS estimates of *Picea* cover for eight Holocene time windows. Percentage cover in 1% interval between**
430 **0 and 2%, 3% interval between 2 and 5%, 5% intervals between 5 and 30%, and 20% interval between 30 and 50%. See caption of**
431 **Figure 2 for more explanations. Percentage cover of *Picea* in 1% intervals represented by increasingly darker shades of green from 5-**
432 **10%. Grey cells: cells without pollen data for the time window, but with pollen data in other time windows. Circles in grid cells**
433 **represent the coefficient of variation (CV; the standard error divided by the REVEALS estimate). When SE > REVEALS estimate,**
434 **the circle fills the entire grid cell and the REVEALS estimate is not different from zero. This occurs mainly where REVEALS**
435 **estimates are low.**

436 3.2.3 *Calluna vulgaris*

437 During the Holocene, *Calluna vulgaris* cover (Fig. D1) peaks at 50%, and is largely distributed in a central European belt from
438 the [British Isles/United Kingdom](#) across to the southern Baltic States. At 9700-10200 BP, it is recorded in only a few grid cells,
439 mostly in central and Western Europe, and at levels <10%. Cover slowly increases and by 7700-8200 BP, there are several
440 grid cells with cover >25% within the [British Isles/United Kingdom](#), and with 10-20% cover within Denmark. At 5700-6200
441 BP, grid cells in coastal locations in northwest Europe (particularly France, Germany and Denmark) have 50% *Calluna*
442 *vulgaris* cover. Cover steadily increases within the same grid cells and by 2700-3200 BP, cover has increased in northern and
443 Eastern Europe e.g. Norway, Estonia, with values up to 20% cover. The highest cover of *Calluna vulgaris* is recorded in the
444 last two millennia. Although some grid cells in southeast Europe record low cover values, these have high SE.

445 3.2.4 *Deciduous Quercus* [type \(t.\) deciduous](#)

446 [Deciduous Quercus](#) [t. deciduous](#) (Fig. D2) is recorded in central and western Europe at 9700-10200 BP at low levels (<10%),
447 while in southern Europe (Italy) ~~there are~~ several grid cells recording >20% cover. By 7700-8200 BP, cover in central and
448 western Europe is between 1-10% while in northern and eastern Europe grid cells it is <2% with high SEs. During the mid-
449 Holocene (5700-6200 BP) most of Europe, with the exception of some grid cells at the northern and southeast extremes, record
450 [deciduous Quercus](#) [deciduous](#) cover values between 2-15%. By 2700-3200 BP, ~~the %~~ cover in the same grid cells has decreased
451 to values between 2-10%. Thereafter, the number of grid cells recording [deciduous Quercus](#) [deciduous](#) cover remains similar;
452 however, the percentage cover slowly decreases and at 350-100 BP, the number of grid cells with [deciduous Quercus](#) [deciduous](#)
453 cover above 5% is very low.

454 3.2.5 *Evergreen Quercus* [type \(t.\) evergreen](#)

455 The spatial distribution of [evergreen Quercus](#) [t. evergreen](#) (Fig. D3) remains the same throughout the Holocene. Cover of
456 >30% is restricted to only a few grid cells and time windows. At the start of the Holocene, [evergreen Quercus](#) [t. evergreen](#) is
457 recorded with values <15% in southern Europe (Spain, Italy, Greece and Turkey) with high SEs. Cover of [evergreen Quercus](#)
458 [t. evergreen](#) does not exceed 15% until 6700-7200 BP (not shown), in grid cells located in Turkey, Greece and Italy. From
459 6700-7200 BP there is an increase in the number of grid cells recording [evergreen Quercus](#) [t. evergreen](#) in southern Europe
460 but most ~~exhibit show~~ low cover values (<15%), and have high SEs.

461 4 Discussion

462 The results presented here are the first full-Holocene grid-based REVEALS estimates of land-cover change for Europe
463 spanning the Mediterranean, temperate and boreal biomes, ~~and which highlighting~~ the spatial and temporal dynamics of 31
464 ~~plant~~ taxa, 12 PFTs and 3 LCTs across Europe over the last 11700 years. Previous studies have demonstrated major
465 differences between REVEALS results and pollen percentages (e.g. Marquer et al., 2014; Trondman et al., 2015), ~~and the~~
466 ~~differences between REVEALS results and other methods used to transform pollen data, including pseudobiomisation, and~~
467 ~~MAT (Roberts et al. 2018).~~ It is not the scope of this paper to evaluate the results in that context. This discussion focuses on
468 the reliability and potential of this “second generation” of REVEALS ~~land cover~~ reconstruction for Europe for use by the wider
469 science community.

470 4.1 Data reliability

471 The REVEALS results are reliant on the quality of the input datasets, namely pollen count data, chronological control for
472 sequences, and the number and reliability of RPP estimates used (~~further see~~ discussion on RPPs under 4.2). The standard
473 errors (SEs) can be considered a measure of the precision of the REVEALS results, and of reliability/quality (Trondman et al.,
474 2015). Where SEs are equal or greater than the REVEALS estimates (represented in the maps of Fig. 2-6 and D1-D3 as a circle
475 that fills the grid), caution should be applied in the use of the REVEALS estimates, as it implies that they are not different
476 from zero when taking the SEs into account. Whilst this is possible within an algorithmic approach that includes estimates of
477 uncertainty, it is conceptually impossible to have negative vegetation cover. If $SEs \geq \text{mean REVEALS value}$ it is therefore
478 uncertain whether the plant taxon has cover within the grid cell. Cover may either be very low or the taxon may be absent
479 within ~~the~~ region (grid cell in this case).

480 The size of pollen counts impacts on the size of REVEALS SEs (Sugita, 2007a); larger counts result in smaller SEs.
481 Aggregation of samples from pollen records to longer time windows results in larger count sizes and thus lower SEs (see
482 sections 2.2 above and 4.2 below). Our input dataset includes more than 59 million individual pollen identifications, organised
483 here into 16711 samples from 1128 sites, where a sample is an aggregated pollen count for RPP taxa for a time window at a
484 site. ~~77%Seventy-seven percent~~ of samples have count sizes in excess of 1000, which is deemed most appropriate for
485 REVEALS reconstructions (Sugita, 2007a). The mean count size across all samples is 3550. Samples with count sizes lower
486 than 1000 are still used, but result in higher SEs. More than half of the pollen records used in the study were sourced from
487 databases (see section 2.2). Note that the EMBSecBIO taxonomy has been pre-standardised, and the data compilers have
488 removed Cerealia-type ~~(t.)~~. This means that for grid cells within the Eastern Mediterranean-Black Sea-Caspian-Corridor,
489 caution is advised in the interpretation of ~~Cerealia-t~~ type. Nevertheless, pollen from ~~e.g.-ruderals is~~ often related to agriculture,
490 ~~such as for example~~. *Artemisia*, *Amaranthaceae/Chenopodiaceae*, and *Rumex acetosa* type are included in the land-cover type
491 open land (OL); therefore, changes in ~~OL cover of open land~~ in the Eastern Mediterranean-Black Sea-Caspian-Corridor may
492 be related to changes in agricultural land (see also discussion below, re agricultural, section 4.3).

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font: Not Italic

Formatted: Font: Not Italic

493 Aggregation of pollen counts to time windows depends on age-depth models. We have used the best age-depth models
494 available to us, based on the chronologies presented in Giesecke et al. (2014) for EPD sites, and through liaison with data
495 contributors. Nevertheless, future REVEALS runs may draw on improvements to age-depth modelling, which may result in
496 some original pollen count data being assigned to different time windows.

497 The REVEALS ~~presented~~ results ~~presented~~ here are provided for $1^\circ \times 1^\circ$ grid cells across Europe. The size and number of
498 suitable pollen records is an important factor in the quality of the REVEALS estimates for each grid cell. The REVEALS
499 model was developed for use with “large lakes” (≥ 50 ha; Sugita, 200a) (~~$\Rightarrow 100-500$ ha~~) that represent regional vegetation
500 (Sugita, 2007a). Grid cells with multiple large lakes will thus provide results with the highest level of certainty and reflect ~~best~~
501 the regional vegetation ~~most accurately~~. These grid cell results comprising of one or more large lakes are considered “high
502 quality” (dark grey grids in figure 1B). It has been shown both theoretically (Sugita, 2007a) and empirically (Fyfe et al., 2013;
503 Trondman et al., 2016a) that pollen records from multiple smaller (~~$<50+00$ ha~~) lakes will also provide REVEALS estimates
504 that reflect ~~the~~ regional vegetation. However, SEs may be larger if there is high variability in pollen composition between
505 records. We therefore also consider grid cells with multiple sites “high quality”. Application of REVEALS to pollen records
506 from large bogs violates assumptions of the model (see section 2.1 above). Therefore, REVEALS estimates for grid cells
507 including large bogs or single small sites (lake or bog) may not be representative of regional vegetation, particularly in areas
508 characterised by heterogeneous vegetation. We consider such estimates as “lower quality” (light grey grids in figure 1B),
509 although they may still provide first-order indications ~~on~~ of vegetation cover, and represent an improvement on pollen
510 percentage data (Marquer et al., 2014). Our results provide REVEALS estimates for a maximum of 420 grid cells per time
511 window. The number and type of pollen records in a grid cell can change between time windows: not all pollen records cover
512 the entire Holocene. It is therefore important to consider not just the number and type of pollen records in the total dataset, but
513 how this changes between time windows, to assess the reliability of individual results. Results for a maximum of 143 grid
514 cells are based on three or more sites, 65 on two sites, and a minimum of 212 grid cells on a single site. The results of a
515 maximum of 67 grid cells are based on single small bogs (<400 m radius), 68 on single small lakes (<400 m radius), and 82
516 on single large bogs. ~~It~~ This implies that about half the grid cells with REVEALS results should be considered as “lower
517 quality” results.

518 4.2 Role of RPPs and FSP in REVEALS results

519 A key assumption of the REVEALS model is that RPP values are constant within the region of interest, and through time
520 (Sugita, 2007a). Nevertheless, it has been suggested that RPPs may vary between regions, with the variation caused by
521 environmental variability (climate, land use), vegetation structure, or methodological design differences (Broström et al., 2008;
522 Hellman et al., 2008a; Mazier et al., 2012; Li et al., 2020; Wiczorek and Herzschuh, 2020). Wiczorek and Herzschuh (2020)
523 have shown that inter-taxon variability in RPP values is generally lower than intra-taxon variability, lending support to
524 application of the approach we used in the new synthesis of RPPs ~~in~~ for Europe (Appendix A-C), i.e. calculation of mean RPPs
525 using all available RPP values that can be considered as reliable. Nevertheless, some RPP taxa still present a challenge, for

526 example, Ericaceae, where Mediterranean tree forms have a greater number of inflorescences and hence may have a higher
527 RPP than low-growth form Ericaceae in central and northern Europe. ~~As we are using a single RPP dataset with the RPP of~~
528 ~~Ericaceae obtained in the Mediterranean region (more explanations below), the effect of higher pollen producing Ericaceae in~~
529 ~~the Mediterranean might result in underrepresentation of Ericaceae cover in eCentral nNorth Europe. Unfortunately, As we~~
530 have only unique RPP values for Ericaceae in both boreal-temperate Europe and Mediterranean Europe, and therefore the large
531 difference in RPP between the two biomes remains to be confirmed with more RPP studies.

532 Currently there is higher confidence in the boreal and temperate RPP values that are based on a wider set of studies increasing
533 the spread of values and hence reliability of the mean RPP values used (Mazier et al., 2012; Wiczorek and Herschuh, 2020),
534 whilst RPP values for Mediterranean taxa are based on fewer empirical RPP studies. The new RPP datasets for Europe
535 produced for this study (Appendix A-C) can be used in different ways. The RPPs provided in Table A1 can be used for entire
536 Europe, including ~~or excluding~~ entomophilous taxa ~~or not~~, and including all values from the Mediterranean area or only the
537 values for the strictly sub-Mediterranean and/or Mediterranean taxa. If one uses all RPPs from the Mediterranean area, there
538 will be taxa for which there is both a RPP value obtained in boreal/temperate Europe and a RPP value obtained in
539 Mediterranean Europe. Application of both RPP values in a single REVEALS reconstruction is not straightforward to achieve,
540 because the border between the two regions has shifted over the Holocene. In the REVEALS reconstruction presented in this
541 paper, we chose to use the RPPs from Mediterranean Europe only for the sub-Mediterranean and/or Mediterranean taxa
542 (including Ericaceae) (Table 1 and A1), and for all other taxa we used the RPPs from boreal/ temperate Europe. The major
543 issue with this choice is the RPP value of Ericaceae. Using only the large value from Mediterranean Europe may lead to an
544 under-representation of Ericaceae (*Calluna* excluded), in particular in boreal Europe, but perhaps also in temperate Europe.
545 Using only the small value from boreal/temperate Europe may lead to an over-representation of Ericaceae in Mediterranean
546 Europe.

547 Until we have more RPP values for each taxon, it is not possible to disentangle the effect of all factors influencing the
548 estimation of RPPs and to separate the effect of methodological factors from those of factors such as vegetation type, climate
549 and land use. The only way to evaluate the reliability of RPP datasets is to test them with modern or historical pollen
550 assemblages and related plant cover (Hellman et al., 2008a, 2008b). We argue that RPP values of certain taxa may not vary
551 substantially within some plant families or genera, while they might be variable within others, depending on the characteristics
552 of flowers and inflorescences that may be either very different or relatively constant within families or genera (see discussion
553 in (Li et al., (2018)). Therefore, we advise to use compilations of RPPs at continental or sub-continental scales rather than
554 compilations at multi-continental scales as the North Hemisphere dataset proposed by Wiczorek and Herzsuh (2020). We
555 consider the RPP selection used within this work as the most suitable for Europe to date, but expect revised and improved RPP
556 values as more RPP empirical studies are published. Moreover, experimentation in REVEALS applications will allow future
557 studies to evaluate the effects of using different RPP datasets on land-cover reconstructions (e.g. Mazier et al., 2012).

558 The role of FSP values in the pollen dispersal and deposition function ($g_i(z)$ in ~~the~~ equation (1) of the REVEALS model,
559 section 2.1) has been discussed by Theuerkauf et al. (2012). In this application of REVEALS we used the Gaussian Plume

560 Model (GPM) of dispersion and deposition as most existing RPP values have been estimated using this model. The GPM
561 approximates dispersal as a fast-declining curve with distance from the source plant, which implies short distances of transport
562 for pollen grain with high FSP compared to other models of dispersion and deposition (Theuerkauf, 2012). We have used the
563 FSP values obtained for deciduous Quercus type deciduous (0.035 m/s) and boreal-temperate Ericaceae (0.037 m/s) for
564 evergreen Quercus t. evergreen and Mediterranean Ericaceae, respectively, although the FSP values of those two taxa were
565 estimated to 0.015 and 0.051 in the Mediterranean study (Table 1 and A1). ~~Whether The possible effect of using a the lower~~
566 ~~FSP for evergreen Quercus t. evergreen (0.015 m/s) and at the higher FSP for Mediterranean Ericaceae (0.051 m/s) will have~~
567 ~~an effect on the REVEALS results is not known and may be lower cover of evergreen Quercus evergreen and higher cover of~~
568 ~~Mediterranean Ericaceae than our results suggest. This hypothesis however~~ requires further testing.

569 4.3 Use of the REVEALS land cover reconstructions results

570 The second generation dataset of pollen-based REVEALS land cover in Europe over the Holocene (this paper) is currently
571 used in two major research projects: LandClim, and PAGES LandCover6k. LandClim (a Swedish Research Council project;
572 e.g. Githumbi et al., 2019; Strandberg et al., 2014; Gaillard et al., 2010) studies the difference in the biogeophysical effect of
573 land-cover change on climate at 6000, 2500 and 200 BP (Githumbi et al., 2019; Strandberg et al., 2014), PAGES LandCover6k
574 focuses on providing datasets on past land-cover/land-use for climate modelling studies (e.g. Gaillard et al., 2018; Harrison
575 et al., 2020). The first generation REVEALS land-cover reconstruction (Marquer et al., 2014, 2017; Trondman et al., 2015) were
576 used to evaluate other pollen-based reconstructions of Holocene tree-cover changes in Europe (Roberts et al., 2018) and
577 scenarios of anthropogenic land-cover changes (ALCCs) (Kaplan et al., 2017) (see also section 1). The Trondman et al. (2015)
578 reconstructions were used to create continuous spatial datasets of past land cover using spatial statistical modelling
579 (Pirzamanbein et al., 2014, 2018, 2020). Spatially explicit datasets/maps based on the second generation of REVEALS
580 reconstruction are currently being produced within PAGES LandCover6k and used to evaluate and revise the HYDE (Klein
581 Goldewijk et al., 2017) and KK10 (Kaplan et al., 2009) ALCC scenarios. Moreover, LandCover6k archaeology-based
582 reconstructions of past land-use change (Morrison et al., 2021) will be integrated with the datasets of REVEALS land-cover.
583 Besides the uses listed above, the second generation of REVEALS reconstruction for Europe offers great potential for use in
584 a large range of studies on past European regional vegetation dynamics and changes in biodiversity over the Holocene (cf.
585 Marquer et al., 2014, 2017) and the relationship between regional plant cover, land use, and climate over millennial and
586 centennial time scales. Moreover the data can be used to create all sorts of maps of plant cover that can serve in various
587 contexts. We stress here that the reconstructions are of regional plant cover. They will thus have value in archaeological
588 research when impacts are expected at the regional level (e.g. the impact of early mining; Schauer et al., 2019) but
589 archaeological questions and research programmes that require information on local vegetation cover will require the full
590 application of the LRA (REVEALS and LOVE; Sugita, 2007a, b), such as the local vegetation estimates presented in Mehl et
591 al. (2015). The same approach of using the REVEALS results within the LOVE model is necessary for ecological questions
592 that require local vegetation estimates (e.g. Cui et al., 2013, 2014; Sugita et al., 2010).

Formatted: Heading 2

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

593 Several papers have discussed in depth the issues that need to be taken into account when interpreting REVEALS
594 reconstructions of past plant cover, in particular Trondman et al., (2015) and Marquer et al. (2017). The interpretation in terms
595 of human-induced vegetation change is one of the major challenges. The cover of open land (OL) may be used to assess
596 landscape openness, but is not a precise measure of human disturbance, as OL will include plant taxa characterizing both
597 naturally-open land and agricultural land that has been created by humans through the course of the Holocene with the
598 domestication of plants and livestock. Natural openness can occur in arctic and alpine areas, in wet regions, in river deltas and
599 around large lakes, as well as in eastern steppe areas. It is a particular challenge in the Mediterranean region where natural
600 vegetation openness represents a larger fraction of the land cover than in temperate or boreal Europe (Roberts et al., 2019).
601 Agricultural Land (AL; Trondman et al., 2015) is the only PFT that includes cultivars; nevertheless, it is restricted to cereal
602 cropping, and many other cultivated crop types that can be identified through pollen analysis do not yet have RPP values (e.g.
603 *Linum usitatissimum* (common flax), *Cannabis* (hemp), *Fagopyrum* (buckwheat), beans, etc.). Moreover, the Cerealia-t. pollen
604 morphological type includes pollen from wild species of Poaceae, especially when identification relies essentially on
605 measurements of the pollen grain and its pore and does not consider exine structure and sculpture (Beug, 2004; Dickson, 1988).
606 The maps presented and described in section 3 as an illustration of the results show similar changes in spatial distributions and
607 quantitative cover of plant taxa and land-cover types through time, between 6000 BP and present, as the results published in
608 Trondman et al., (2015). The much greater potential of the new REVEALS reconstruction resides in its larger spatial extent,
609 covering not only boreal and temperate Europe but also southern and eastern Europe, and its contiguous time windows across
610 the entire Holocene, from 11700 BP to present. The quality of results is also higher in a number of grid cells in comparison to
611 Trondman et al (2015), where new pollen records have been included, which may in several cases decrease the standard error
612 on the REVEALS estimates.

613 **5. Data availability**

614 All data files reported in this work, which were used for calculations, and figure [productions](#) are available for public download
615 at <https://doi.pangaea.de/10.1594/PANGAEA.937075> (Fyfe, Ralph M; Githumbi, Esther; Trondman, Anna-Kari; Mazier,
616 Florence; Nielsen, Anne Birgitte; Poska, Anneli; Sugita, Shinya; Woodbridge, Jessie; LandClimII contributors; Gaillard,
617 Marie-José (2024): A full Holocene record of transient gridded vegetation cover in Europe. PANGAEA,
618 <https://doi.org/10.1594/PANGAEA.931856>). The data and the DOI number are subject to future updates and only refer to this
619 version of the paper. The data available in Pangaea includes: 1) REVEALS reconstructions and their associated SE for the 25
620 time windows; 2) Metadata of the 1128 pollen records used; 3) LandClimII contributors listing the data
621 contributors/collectors/databases. 4) The list of FSP and RPP values used for the reconstructions and 5) Grid cell quality
622 information (in terms of available pollen data, which influences the result quality: mean REVEALS estimate of plant cover)
623 for all grid cells.

624 Pollen data were extracted from ALPADABA (<https://www.neotomadb.org/>), EMBSEC BIO
625 (<https://research.reading.ac.uk/palaeoclimate/embsecbio/>), EPD (<http://www.europeanpollendatabase.net/index.php>),
626 LandClimI, PALYCZ (<https://botany.natur.cuni.cz/palycz/>) and PALEOPYR (<http://paleopyr.univ-tlse2.fr/>). ~~The work of the~~
627 ~~work of the data contributors and the community is gratefully acknowledged.~~

628

629 6. Code availability

630 REVEALS was implemented using the REVEALS function within the LRA R-package (Abraham et al., 2014), available at
631 <https://github.com/petrkunes/LRA>.

632 Example code for data preparation and implementation of REVEALS, using two grid cells from SW Britain, is available at
633 <https://github.com/rmfyfe/landclimII>.

634 7. Conclusions

635 The application of the REVEALS model to 1128 pollen records distributed across Europe has produced the first full-Holocene
636 estimates of vegetation cover for 31 plant taxa in $1^\circ \times 1^\circ$ grid cells. These data are made available for use by the wider science
637 community, including aggregation of results to PFTs and LCTs. The REVEALS model assumptions are clearly stated to allow
638 interpretation and assessment of our results and several of the assumptions have been tested and validated. We can therefore
639 use the land-cover reconstructions to test the role of climate and humans on ~~the~~ Holocene vegetation at ~~the~~ regional scales in
640 terms of changes in plant cover over time and space. The overview of land-cover change across Europe over the Holocene can
641 be used to track the timing and rate of vegetation shifts, which is useful in discerning the drivers of the observed change
642 (Marquer et al., 2014; 2017). We can also study the effect of human-induced changes in regional vegetation cover on climate,
643 i.e. study land use as a climate forcing (e.g. Gaillard et al., 2010; Strandberg et al., 2014; Gaillard et al., 2018; Harrison et al.,
644 2020). Local reconstructions (LOVE) can be a complementary approach to archaeological surveys as fine-scale human use of
645 the landscape cannot be distinguished using REVEALS (regional estimates). The LOVE model requires that regional plant
646 cover is known: the REVEALS reconstructions are therefore needed for this purpose as well, and gridded reconstructions may
647 be a way to perform LOVE reconstructions, although other strategies can be chosen (e.g. Cui et al., 2013; Mazier et al., 2015).
648 Questions ~~such as~~ ~~aiming to~~ understand the degree of vegetation openness through the Holocene in Europe, or ~~on~~ ~~regarding~~
649 changes in the relationship between summer-green and evergreen tree cover through time can now and in the future be
650 answered and validated with fossil pollen data via the REVEALS approach. We expect that in the future imprecision can be
651 further reduced in terms of both the quality, and spatial extent, of REVEALS estimates, as more pollen records are
652 incorporated, and work on RPPs develops.

Formatted: Normal

Field Code Changed

653 Appendices

654 Appendix A - New RPP dataset for Europe

655 **A.1 Introduction**

656 The most common method to estimate RPPs involves the application of the Extended R-Value (ERV) model on datasets of
657 modern pollen assemblages and related vegetation cover. A summary of the ERV model and its assumptions, and an extensive
658 description of standardised field methods for the purpose of RPP studies are found in Bunting et al. (2013b). Estimation of
659 RPPs in Europe started with the studies by Sugita et al. (1999) and Broström et al. (2004) in Southern Sweden, and Nielsen et
660 al. (2004) in Denmark. The first tests of the RPP in pollen-based reconstructions of plant cover using the LRA's REVEALS
661 (REgional VEgetation Abundance from Large Sites) model (Sugita, 2007a) were published by Soepboer et al. (2007) in
662 Switzerland and Hellman et al. (2008a and b) in South Sweden. Over the last 15 years, a large number of RPP studies have
663 been undertaken in Europe North of the Alps, but it is only recently that RPP studies were initiated in the Mediterranean area
664 (Grindean et al., 2019; Mazier et al., unpublished). Two earlier syntheses of RPPs in Europe were published by Broström et
665 al. (2008) and Mazier et al. (2012). From 2012 onwards, these RPP values have been used in numerous applications of the
666 LRA's two models REVEALS and LOVE (LOCAL Vegetation Estimates) (Sugita, 2007a and b) to reconstruct regional and
667 local plant cover in Europe (e.g. Cui et al., 2013; Fyfe et al., 2013; Marquer et al., 2020; Mazier et al., 2015; Nielsen et al.,
668 2012; Nielsen and Odgaard, 2010; Trondman et al., 2015). Recently, Wiczorek and Herzschuh (2020) published a synthesis
669 of the RPPs available for the Northern Hemisphere; it includes new mean RPP values for Europe that were produced
670 independently from the synthesis we present here.

671 **A.1 New synthesis of European RPPs**

672 Table A1 is the result of the new synthesis of RPPs available in Europe we have performed for the REVEALS reconstruction
673 presented in the paper. It includes RPPs for 39 plant taxa from studies in boreal and temperate Europe of which 22 (Poaceae
674 included) are herbs or low shrubs, and for 22 plant taxa from studies in the Mediterranean area. The two regions have RPP
675 values for 7 plant taxa in common. These RPPs are compared to those from two syntheses published earlier, Mazier et al.
676 (2012) and Wiczorek and Herzschuh (2020). The number of selected RPP values (n) for Poaceae is larger than the total
677 number of RPP (tn), i.e. $n = tn + 1$. This is due to the fact that the study of Bunting et al. 2005 does not include a value for
678 Poaceae and the RPP values are related to *Quercus* (Bunting et al., 2005); therefore, RPPs related to Poaceae were calculated
679 by assuming the RPP value for *Quercus* (related to Poaceae; $Quercus_{(Poaceae)}$) was the same in this study region than the mean
680 of $Quercus_{(Poaceae)}$ RPPs from all other available studies.

681 The ranking of RPPs (relative to Poaceae, RPP=1) for 23 tree taxa (M: Mediterranean taxa), from the largest (13.56) to the
682 smallest (0.240), is as follows (Poaceae included for comparison ~~with herbs~~): *Alnus*> evergreen *Quercus* <evergreen-(M)>
683 *Abies alba*> *Pinus*> *Fagus sylvatica*> *Picea abies*> Ericaceae (M)> *Betula*> deciduous *Quercus* <t.> *Carpinus betulus*>
684 *Populus*> *Juniperus*> *Corylus avellana*> *Castanea sativa*> *Sambucus nigra*-t.> *Ulmus*> *Tilia*> *Salix*> *Fraxinus*> Poaceae

Formatted: Normal

Field Code Changed

685 (=1)> *Acer*> *Pistacia* (M)> *Phillyrea* (M)> *Carpinus orientalis* (M). All tree taxa have mean RPPs larger than 1 except *Acer*
686 (0.8), *Pistacia* (0.755), *Phillyrea* (0.512) and *Carpinus orientalis* (0.240). The ranking of RPPs for 24 herb and low shrub taxa,
687 from the largest (10.52) to the smallest (0.10), is as follows: *Urtica*> Chenopodiaceae> *Secale*> *Artemisia*> Rubiaceae> *Rumex*
688 *acetosa*-t.> *Filipendula*> *Plantago lanceolata*> *Trollius*> Ranunculaceae (M)> *Ranunculus acris*-t.> Cerealia-t.> *Potentilla*-
689 t.> *Plantago media*> *Calluna vulgaris*> Poaceae (=1)> Cyperaceae> *Plantago montana*> Fabaceae (M)> Rosaceae (M)>
690 Apiaceae> Compositae SF. Cichorioideae> *Empetrum*> *Leucanthemum* (*Anthemis*)-t.. ~~Of the taxa with Only six herb taxa have~~
691 RPPs larger than 3, ~~only six taxa are herbs while twelve are while 12 trees taxa have RPP > than 3.~~
692 The two studies in the Mediterranean area provide single RPP values for 16 taxa, five herb taxa (Poaceae included) and 11 tree
693 taxa of which six are sub-Mediterranean and/or Mediterranean, and three include both temperate and Mediterranean taxa
694 (Cupressaceae, Ericaceae, *Fraxinus*) (Table B2). The RPP of herb taxa are significantly different between the study of
695 Grindean et al. (2019) and our synthesis, except for *Artemisia* (5.89 and 3, 94, respectively). The RPP of *Corylus avellana*
696 from the study of Mazier et al. (unpublished) (3.44) is double as large as the mean RPP in our synthesis (1.71), and the mean
697 RPP of deciduous *Quercus* t. (deciduous species) in our synthesis (4.54) is four times as large as the RPP from the study of
698 Grindean et al. (2019) (1.10).
699

700 **Table A1: New synthesis of European RPPs: mean RPPs with their SDs in brackets, and mean RPPs from the**
701 **syntheses by Mazier et al. (2012) (St2 values) and Wieczorek and Herzsuh (2020), for comparison. This synthesis:**
702 **values in bold are new mean RPPs compared to Mazier et al. (2012). The RPP values from studies in the**
703 **Mediterranean area are indicated with “M” in the second column. The values emphasized in grey are the mean RPPs**
704 **used in the new REVEALS reconstruction for Europe (this paper). The values of fall speed of pollen (FSP) are from**
705 **Mazier et al. (2012) except those in italic, i.e. FSPs for Chenopodiaceae, *Urtica* and *Sambucus nigra-t.* (Abraham and**
706 **Kozáková, 2012), and *Populus* (Wieczorek and Herzsuh, 2020) and the new FSPs for Mediterranean taxa. For the**
707 **three syntheses, the number of selected RPP values (n) included in the calculation of the mean RPP estimate is**
708 **indicated with the total number of available RPP values (tn) in brackets. The reason why the number of selected**
709 **RPP values (n) for Poaceae is larger than the total number of RPP (tn) is provided in section A1. For explanation of**
710 **symbols, see captions below.**

711
712 * Separate mean RPP values for *Calluna vulgaris*, *Empetrum*, and Ericaceae (*Calluna* and *Empetrum* excluded) in this
713 synthesis, a single mean RPP values for all Ericales in Wieczorek and Herzsuh (2020)

714 ** Separate mean RPP values for Cerealia type (*Secale* excluded) and *Secale* in this synthesis, a single mean RPP for all cereals
715 in Wieczorek and Herzsuh (2020)

716 *** Separate mean RPP values for Compositae SF Cichorioideae and *Leucanthemum* (*Anthemis*) type in this synthesis, a single
717 mean RPP for all Asteraceae in Wieczorek and Herzsuh (2020). Note that there are no RPP for Asteraceae (Compositae SF
718 Cichorioideae and *Leucanthemum* (*Anthemis*) type excluded) in our synthesis

719 ^ Separate mean RPP values for *Filipendula* and *Potentilla* type in this synthesis, a single mean RPP for all Rosaceae in
720 Wieczorek and Herzsuh (2020); note that there are no RPP for Rosaceae (*Filipendula* and *Potentilla-t.* excluded) in our
721 synthesis; moreover *Filipendula* and *Potentilla-t.* are classified as herbs, while Rosaceae is classified as tree in Wieczorek and
722 Herzsuh (2020)

723 ^^ Separate mean RPP values for *Plantago lanceolata*, *P. media* and *P. montana* in this synthesis, a single mean RPP for all
724 Plantaginaceae in Wieczorek and Herzsuh (2020); note that there are no RPP for Plantaginaceae (*Plantago lanceolata*, *P.*
725 *media* and *P. montana* excluded) in our synthesis

726 ^^ Separate mean RPP values for *Ranunculus acris* type and *Trollius* in this synthesis, a single mean RPP for all
727 Ranunculaceae in Wieczorek and Herzsuh (2020); note that there are no RPP for Ranunculaceae (*Ranunculus acris-t* and
728 *Trollius* excluded) in our synthesis.

729

Field Code Changed

Formatted: Font: Bold

Formatted: Font: Bold

Field Code Changed

Study n (tn), FSP, RPP	This paper, synthesis			Mazier et al. 2012 St 3		Wieczorek & Herzsuh 2020 Europe version 2			
	n (tn)	FSP	RPP (SE)	n (tn)	RPP (SE)	n (tn)	RPP (SE)	Notes	
HERB TAXA	-	-	-	-	-	-	-	-	
Pbaceae (Reference taxon)	-	16(15)	0.035	1.00 (0.00)	9(8)	1.00 (0.00)	14(12)	1.00 (0.00)	
Herb taxa	-	-	-	-	-	-	-	-	
Amaranthaceae/Chenopodiaceae	-	1(1)	<u>0.019</u>	4.280 (0.270)	none	none	1(1)	4.28 (0.27)	Same value as in this synthesis
Apiaceae	-	1(1)	0.042	0.260 (0.010)	1(1)	0.26 (0.01)	3(3)	2.13 (0.41)	-
Apiaceae	M	1(1)	0.042	5.910 (1.230)	-	-	-	-	-
<i>Artemisia</i>	-	3(3)	0.025	3.937 (0.146)	1(1)	3.48 (0.20)	2(2)	4.33 (1.59)	-
<i>Artemisia</i>	M	1(1)	0.014	5.890 (3.160)	-	-	-	-	-
Comp. <i>Leucanth. (Anthemis)t</i> ,***	-	1(1)	0.029	0.100 (0.010)	1(1)	0.10 (0.01)	-	-	see Asteraceae all***
Comp. SF. Cichorioideae***	-	3(3)	0.051	0.160 (0.020)	3(3)	0.16 (0.02)	8(10)	0.22 (0.02)	Asteraceae all***
Comp. SF. Cichorioideae	M	1(1)	0.061	1.162 (0.075)	-	-	-	-	-
Comp. (Asteroideae + Cichorioideae)	M	1(1)	0.029	0.160 (0.100)	-	-	-	-	-
<i>Calluna vulgaris</i> *	-	2(4)	0.038	1.085 (0.029)	2(4)	1.09 (0.03)	-	-	see Ericales all*
Cerealia t.**	-	3(7)	0.060	1.850 (0.380)	2(4)	1.18 (0.04)	4(6)	2.36 (0.42)	Cereals all**
Cerealia t. (<i>Triticum</i> ., <i>Secale</i> , <i>Zea</i>)	M	1(1)	0.060	0.220 (0.120)	-	-	-	-	-
Cyperaceae	-	4(6)	0.035	0.962 (0.050)	4(6)	0.83 (0.04)	6(8)	0.56 (0.02)	-
<i>Empetrum</i> *	-	1(2)	0.038	0.110 (0.030)	1(2)	0.11 (0.03)	-	-	see Ericales all*
Ericaceae*	-	1(1)	0.038	0.070 (0.040)	1(1)	0.07 (0.04)	7(9)	0.44 (0.02)	Ericales all*
Fabaceae	M	1(1)	0.021	0.400 (0.070)	-	-	-	-	-
<i>Filipendula</i> ^	-	3(3)	0.006	3.000 (0.285)	2(3)	2.81 (0.43)	4(6)	0.97 (0.11)	Rosaceae all ^
<i>Pantago lanceolata</i> ^^	-	4(6)	0.029	2.330 (0.201)	3(4)	1.04 (0.09)	8(10)	2.49 (0.11)	Plantaginaceae all^^
<i>Pantago lanceolata</i>	M	1(1)	0.029	0.580 (0.320)	-	-	-	-	-
<i>Pantago media</i> ^^	-	1(1)	0.024	1.270 (0.180)	1(1)	1.27 (0.18)	-	-	see Plantaginaceae all^^
<i>Pantago montana</i> ^^	-	1(1)	0.030	0.740 (0.130)	1(1)	0.74 (0.13)	-	-	see Plantaginaceae all^^
<i>Potentilla</i> ^	-	2(3)	0.018	1.720 (0.200)	2(3)	1.72 (0.20)	-	-	see Rosaceae all^
Ranunculaceae	M	1(1)	0.020	2.038 (0.335)	-	-	-	-	-
<i>Ranunculus acris</i> ^^	-	2(2)	0.014	1.960 (0.360)	2(2)	1.96 (0.36)	3(5)	0.99 (0.12)	Ranunculaceae all^^^
Rosaceae (<i>Filipend.</i> , <i>Pot. t.</i> , <i>Sanguisorba</i>)	M	1(1)	0.018	0.290 (0.120)	-	-	-	-	-

Rubiaceae	-	2(3)	0.019	3.710 (0.340)	2(3)	3.71 (0.34)	3(5)	1.56 (0.12)	-
Rubiaceae	M	1(1)	0.019	0.400 (0.070)	-	-	-	-	-
<i>Rumex acetosat.</i>	-	3(4)	0.018	3.020 (0.278)	3(3)	0.85 (0.05)	3(4)	0.58 (0.03)	-
<i>Stacale**</i>	-	3(3)	0.060	3.990 (0.320)	1(1)	3.02 (0.05)	-	-	see Cereals all**
<i>Trollius^^^</i>	-	1(1)	0.013	2.290 (0.360)	1(1)	2.29 (0.36)	-	-	see Ranunculaceae all^^^
<i>Urtica</i>	-	1(1)	0.007	10.520 (0.310)	none	none	1(1)	10.52 (0.31)	Same value as in this synthesis
TREE TAXA	-	-	-	-	-	-	-	-	-
<i>Abies alba</i>	-	2(2)	0.120	6.875 (1.442)	2(2)	6.88 (1.44)	2(2)	6.88 (1.44)	Same value as in this synthesis
<i>Acer</i>	-	2(2)	0.056	0.800 (0.230)	2(2)	0.80 (0.23)	3(3)	0.23 (0.04)	-
<i>Acer</i>	M	1(1)	0.056	0.300 (0.090)	-	-	-	-	-
<i>Alnus</i>	-	5(7)	0.021	13.562 (0.293)	3(3)	9.07 (0.10)	4(6)	8.49 (0.22)	-
<i>Betula</i> (mainly <i>B. pubescens</i> , <i>B. pendula</i>)	-	7(9)	0.024	5.106 (0.303)	6(6)	3.99 (0.17)	6(8)	4.94 (0.44)	-
<i>Buxus sempervirens</i>	M	1(1)	0.032	1.890 (0.068)	-	-	-	-	-
<i>Carpinus betulus</i>	-	2(4)	0.042	4.520 (0.425)	2(2)	3.55 (0.43)	3(5)	3.09 (0.28)	-
<i>Carpinus orientalis</i>	M	1(1)	0.042	0.240 (0.070)	-	-	-	-	-
<i>Castanea sativa</i>	M	1(1)	0.010	3.258 (0.059)	-	-	-	-	-
<i>Corylus avellana</i>	-	4(4)	0.025	1.710 (0.100)	3(3)	1.99 (0.20)	3(4)	1.05 (0.33)	-
<i>Corylus avellana</i>	M	1(1)	0.025	3.440 (0.890)	-	-	-	-	-
Cupressaceae (<i>Juniperus</i> 3 species)	M	1(1)	0.020	1.618 (0.161)	-	-	-	-	See <i>Juniperus</i>
Ericaceae (<i>Arbutus unedo</i> , <i>Erica</i> 3 species)	M	1(1)	0.051	4.265 (0.094)	-	-	-	-	-
<i>Fagus sylvatica</i>	-	3(6)	0.057	5.863 (0.176)	4(4)	3.43 (0.09)	3(3)	2.35 (0.11)	-
<i>Fraxinus excelsior</i>	-	5(6)	0.022	1.044 (0.048)	3(3)	1.03 (0.11)	5(5)	2.97 (0.25)	-
<i>Fraxinus</i> (<i>F. excelsior</i> , <i>F. ornus</i>)	M	1(1)	0.022	2.990 (0.880)	-	-	-	-	-
<i>Juniperus communis</i>	-	1(2)	0.016	2.070 (0.040)	1(2)	2.07 (0.04)	1(1)	7.94 (1.28)	-
<i>Phillyrea</i>	M	1(1)	0.015	0.512 (0.076)	-	-	-	-	-
<i>Pistacia</i>	M	1(1)	0.030	0.755 (0.201)	-	-	-	-	-
<i>Picea abies</i>	-	4(8)	0.056	5.437 (0.097)	4(6)	2.62 (0.12)	4(6)	1.65 (0.15)	-
<i>Pinus</i> (mainly <i>P. sylvestris</i>)	-	6(9)	0.031	6.058 (0.237)	3(5)	6.38 (0.45)	4(6)	10.86 (0.80)	-
<i>Populus</i>	-	1(1)	0.025	2.660 (1.250)	none	none	1(1)	3.42 (1.60)	-
Dec. <i>Quercust.</i> (mainly <i>O. robur</i> , <i>O. petraea</i>)	-	6(8)	0.035	4.537 (0.086)	4(4)	5.83 (0.15)	5(7)	2.42 (0.10)	-

<u>Dec. Quercust. (mainly <i>O. peduncularis</i>)</u>	<u>M</u>	<u>1(1)</u>	<u>0.035</u>	<u>1.100 (0.350)</u>	-	-	-	-	-
<u>Evergreen Quercust. (<i>O. ilex</i>, <i>O. coccifera</i>)</u>	<u>M</u>	<u>1(1)</u>	<u>0.015</u>	<u>11.043 (0.261)</u>	-	-	-	-	-
<u>Salix</u>	-	<u>5(5)</u>	<u>0.022</u>	<u>1.182 (0.077)</u>	<u>3(4)</u>	<u>1.79 (0.16)</u>	<u>3(4)</u>	<u>0.39 (0.06)</u>	-
<u>Sambucus nigrat.</u>	-	<u>1(1)</u>	<u>0.013</u>	<u>1.300 (0.120)</u>	none	none	<u>1(1)</u>	<u>1.30 (0.12)</u>	- Same value as in this synthesis
<u>Tilia</u>	-	<u>4(5)</u>	<u>0.032</u>	<u>1.210 (0.116)</u>	<u>1(1)</u>	<u>0.80 (0.03)</u>	<u>3(4)</u>	<u>0.93 (0.09)</u>	-
<u>Ulmus</u>	-	<u>1(2)</u>	<u>0.032</u>	<u>1.270 (0.050)</u>	<u>1(1)</u>	<u>1.27 (0.05)</u>	none	-	-

730

Study n (tn), FSP, RPP	This paper, synthesis			Mazier et al. 2012 St 3		Wieczorek & Herzsuh 2020 Europe version 2		Notes
	n (tn)	FSP	RPP (SE)	n (tn)	RPP (SE)	n (tn)	RPP (SE)	
HERB TAXA								
Poaceae (Reference taxon)	16(15)	0.035	1.00 (0.00)	9(8)	1.00 (0.00)	14(12)	1.00 (0.00)	
Herb taxa								
Amaranthaceae/Chenopodiaceae	1(1)	0.019	4.280 (0.270)	none	none	1(1)	<u>4.28 (0.27)</u>	Same value as in this synthesis
Apiaceae	1(1)	0.042	0.260 (0.010)	1(1)	0.26 (0.01)	3(3)	2.13 (0.41)	
Artemisia	M 1(1)	0.042	5.910 (1.230)					
Artemisia	M 3(3)	0.025	3.937 (0.146)	1(1)	3.48 (0.20)	2(2)	4.33 (1.59)	
Asteraceae <i>Leucanth.</i> (<i>Anthemis</i>)-t***	M 1(1)	0.014	5.890 (3.160)					
Asteraceae Cichorioideae***	M 1(1)	0.029	0.100 (0.010)	1(1)	0.10 (0.01)			see Asteraceae all***
Asteraceae Cichorioideae	M 3(3)	0.051	0.160 (0.020)	3(3)	0.16 (0.02)	8(10)	0.22 (0.02)	Asteraceae all***
Asteraceae (Asteroideae + Cichorioideae)	M 1(1)	0.061	1.162 (0.075)					
<i>Calluna vulgaris</i> *	M 1(1)	0.029	0.160 (0.100)					see Ericales all*
Cerealia-t**	2(4)	0.038	<u>1.085 (0.029)</u>	2(4)	1.09 (0.03)			see Ericales all*
Cerealia-t (<i>Triticum</i> t., <i>Secale</i> , <i>Zea</i>)	3(7)	0.060	1.850 (0.380)	2(4)	1.18 (0.04)	4(6)	2.36 (0.42)	Cereals all**
Cyperaceae	M 1(1)	0.060	0.220 (0.120)					
<i>Empetrum</i> *	4(6)	0.035	0.962 (0.050)	4(6)	0.83 (0.04)	6(8)	0.56 (0.02)	
Ericaceae*	1(2)	0.038	0.110 (0.030)	1(2)	0.11 (0.03)			see Ericales all*
Fabaceae	M 1(1)	0.038	0.070 (0.040)	1(1)	0.07 (0.04)	7(9)	0.44 (0.02)	Ericales all*
<i>Filipendula</i> ^	M 1(1)	0.021	0.400 (0.070)					
<i>Plantago lanceolata</i> ^^	3(3)	0.006	3.000 (0.285)	2(3)	2.81 (0.43)	4(6)	0.97 (0.11)	Rosaceae all ^
<i>Plantago lanceolata</i>	4(6)	0.029	2.330 (0.201)	3(4)	1.04 (0.09)	8(10)	2.49 (0.11)	Plantaginaceae all^^
<i>Plantago media</i> ^^	M 1(1)	0.029	0.580 (0.320)					
<i>Plantago montana</i> ^^	1(1)	0.024	1.270 (0.180)	1(1)	1.27 (0.18)			see Plantaginaceae all^^
<i>Potentilla</i> -t^	1(1)	0.030	0.740 (0.130)	1(1)	0.74 (0.13)			see Plantaginaceae all^^
Ranunculaceae	2(3)	0.018	1.720 (0.200)	2(3)	1.72 (0.20)			see Rosaceae all^
<i>Ranunculus acris</i> -t^^^	M 1(1)	0.020	2.038 (0.335)					
Rosaceae (<i>Filipend.</i> , <i>Pot.</i> t., <i>Sanguisorba</i>)	2(2)	0.014	1.960 (0.360)	2(2)	1.96 (0.36)	3(5)	0.99 (0.12)	Ranunculaceae all^^^
Rubiaceae	M 1(1)	0.018	0.290 (0.120)					
Rubiaceae	2(3)	0.019	3.710 (0.340)	2(3)	3.71 (0.34)	3(5)	1.56 (0.12)	
<i>Rumex acetosa</i> -t	M 1(1)	0.019	0.400 (0.070)					
<i>Secale</i> **	3(4)	0.018	3.020 (0.278)	3(3)	0.85 (0.05)	3(4)	0.58 (0.03)	see Cereals all**
<i>Trallius</i> ^^	M 3(3)	0.060	3.990 (0.320)	1(1)	3.02 (0.05)			see Ranunculaceae all^^
<i>Urtica</i>	1(1)	0.013	2.290 (0.360)	1(1)	2.29 (0.36)	1(1)	<u>10.52 (0.31)</u>	Same value as in this synthesis
	1(1)	0.007	10.520 (0.310)	none	none			
TREE TAXA								
<i>Abies alba</i>	2(2)	0.120	6.875 (1.442)	2(2)	6.88 (1.44)	2(2)	<u>6.88 (1.44)</u>	Same value as in this synthesis
<i>Acer</i>	2(2)	0.056	0.800 (0.230)	2(2)	0.80 (0.23)	3(3)	0.23 (0.04)	
<i>Acer</i>	M 1(1)	0.056	0.300 (0.090)					
<i>Alnus</i>	5(7)	0.021	13.562 (0.293)	3(3)	9.07 (0.10)	4(6)	8.49 (0.22)	
<i>Betula</i> (mainly <i>B. pubescens</i> , <i>B. pendula</i>)	7(9)	0.024	5.106 (0.303)	6(6)	3.99 (0.17)	6(8)	4.94 (0.44)	
<i>Buxus sempervirens</i>	M 1(1)	0.032	1.890 (0.068)					
<i>Carpinus betulus</i>	2(4)	0.042	4.520 (0.425)	2(2)	3.55 (0.43)	3(5)	3.09 (0.28)	
<i>Carpinus orientalis</i>	M 1(1)	0.042	0.240 (0.070)					
<i>Castanea sativa</i>	M 1(1)	0.010	3.258 (0.059)					
<i>Corylus avellana</i>	4(4)	0.025	1.710 (0.100)	3(3)	1.99 (0.20)	3(4)	1.05 (0.33)	
<i>Corylus avellana</i>	M 1(1)	0.025	3.440 (0.890)					
Cupressaceae (<i>Juniperus</i> 3 species)	M 1(1)	0.020	1.618 (0.161)					See <i>Juniperus</i>
Ericaceae (<i>Arbutus unedo</i> , <i>Erica</i> 3 species)	M 1(1)	0.051	4.265 (0.094)					
<i>Fagus sylvatica</i>	3(6)	0.057	5.863 (0.176)	4(4)	3.43 (0.09)	3(3)	2.35 (0.11)	
<i>Fraxinus excelsior</i>	5(6)	0.022	1.044 (0.048)	3(3)	1.03 (0.11)	5(5)	2.97 (0.25)	
<i>Fraxinus</i> (<i>F. excelsior</i> , <i>F. ornus</i>)	M 1(1)	0.022	2.990 (0.880)					
<i>Juniperus communis</i>	1(2)	0.016	2.070 (0.040)	1(2)	2.07 (0.04)	1(1)	7.94 (1.28)	
<i>Phillyrea</i>	M 1(1)	0.015	0.512 (0.076)					
<i>Pistacia</i>	M 1(1)	0.030	0.755 (0.201)					
<i>Picea abies</i>	4(8)	0.056	5.437 (0.097)	4(6)	2.62 (0.12)	4(6)	1.65 (0.15)	
<i>Pinus</i> (mainly <i>P. sylvestris</i>)	6(9)	0.031	6.058 (0.237)	3(5)	6.38 (0.45)	4(6)	10.86 (0.80)	
<i>Populus</i>	1(1)	0.025	2.660 (1.250)	none	none	1(1)	3.42 (1.60)	
<i>Quercus</i> (mainly <i>Q. robur</i> , <i>Q. petraea</i>)	6(8)	0.035	4.537 (0.086)	4(4)	5.83 (0.15)	5(7)	2.42 (0.10)	
<i>Quercus</i> deciduous (mainly <i>Q. peduncul.</i>)	M 1(1)	0.035	1.100 (0.350)					
<i>Quercus</i> evergreen (<i>Q. ilex</i> , <i>Q. coccifera</i>)	M 1(1)	0.015	11.043 (0.261)					
<i>Salix</i>	5(5)	0.022	1.182 (0.077)	3(4)	1.79 (0.16)	3(4)	0.39 (0.06)	
<i>Sambucus nigra</i> -t	1(1)	0.013	1.300 (0.120)	none	none	1(1)	<u>1.30 (0.12)</u>	Same value as in this synthesis
<i>Tilia</i>	4(5)	0.032	1.210 (0.116)	1(1)	0.80 (0.03)	3(4)	0.93 (0.09)	
<i>Ulmus</i>	1(2)	0.032	1.270 (0.050)	1(1)	1.27 (0.05)	none		

732 A.2 Comparison of the new synthesis with two earlier syntheses (Table A1)

733 Of the 39 plant taxa for which we have a mean RPP in our new synthesis (**New**), 21 have a new mean RPP value compared to
734 the earlier synthesis of Mazier et al. (2012) (**Maz**), 18 taxa have the same mean RPPs in both syntheses. There are three new
735 taxa for which there were no RPP in **Maz**, i.e. **Amaranthaceae**/Chenopodiaceae, *Sambucus nigra*-t. and *Urtica*. The mean RPPs
736 are comparable between the two syntheses **New** and **Maz**, except for *Plantago lanceolata* (2.33 in **New**/1.04 in **Maz**), *Alnus*
737 (13.56/9.07), *Betula* (5.11/3.09), *Carpinus betulus* (4.52/3.55), *Fagus* (5.86/3.43), *Picea* (5.44/2.62) and *Quercus* (4.54/5.83).
738 *Abies alba* has the same RPP in all three syntheses. **Amaranthaceae**/Chenopodiaceae, *Sambucus nigra*-t. and *Urtica* have the
739 same single RPP values in the synthesis of Wieczorek and Hertzschuh (2020) (**W&H**) and **New**. **New** and **W&H** also have
740 comparable mean RPP values for *Artemisia*, Cereals (Cereals, *Secale* excluded in **New**, all Cereals in **W&H**), Compositae (SF
741 Cichorioideae in **N**, all **Compositae** (=Asteraceae) in **W&H**), Cyperaceae, *Plantago* (*P. lanceolata* in **New**, all Plantaginaceae
742 in **W&H**), *Betula*, *Corylus*, *Populus* and *Tilia*. There are relatively large differences in mean RPPs in **W&H** and **New** for 16
743 plant taxa, although the ranking of the plant taxa in terms of their mean RPPs is almost the same. Mean RPP is larger in **W&H**
744 than in **New** for Apiaceae (2.13/0.26), Ericales (0.44 in **W&H**) – *Empetrum* (0.11) and Ericaceae (0.07) in **New**, *Fraxinus*
745 (2.97/1.04), *Juniperus* (7.94/2.07), *Pinus* (10.86/6.06). Mean RPP is smaller in **W&H** than in **New** for *Filipendula* (0.97/3.00),
746 Rubiaceae (1.56/3.71), *Rumex acetosa* (0.58/2.02), *Acer* (0.23/0.80), *Alnus* (8.49/13.56), *Carpinus* (3.09/4.52), *Fagus*
747 (2.35/5.86)), *Picea* (1.65/5.44), *Quercus* (2.42/4.54) and *Salix* (0.39/1.18).
748 The larger differences between the mean RPPs in **New** and **W&H** than between **New** and **Maz** have not been examined in
749 detail. It is due to a slightly different selection of studies, i.e. the study of Theuerkauf et al. (2013) is not included in **W &H**
750 and we did not include in **New** (boreal and temperate Europe, Mediterranean area excluded) the studies of Bunting et al.
751 (2013a), Kuneš et al. (2019) and Grindean et al. (2019). Another important influencing factor is the selection of RPP values
752 for calculation of the mean RPP. Although the rules used to select RPP values are very similar between the syntheses, there
753 are obvious differences between **New** and **W&H** that are sometimes very significant (e.g. *Juniperus*).

754 A.3 Comparison of the new synthesis with three additional individual studies (Table A2)

755 The RPPs from Twiddle et al. (2012) (**Twi**) for *Pinus*, *Betula* and *Calluna* are considerably larger than the mean RPPs in our
756 synthesis (**New**). This is probably due to the assumption made on the RPP of *Picea* related to Poaceae. The RPP of *Picea*
757 varies greatly between the selected studies in **New**, from 0.57 to 8.43 (eight values available). If we assumed that the RPP of
758 *Picea* related to Poaceae in the study region of **Twi** was the mean RPP of the five smallest RPPs, i.e. 1.57, the RPP of the three
759 taxa would be 4.8 for *Pinus*, 3.4 for *Betula*, and 3.3 for *Calluna*, which is more comparable to the mean RPPs in **New**.
760 Three taxa in Bunting et al. (2013a) (**Bun**) have a RPP comparable to the mean RPP in **New**, i.e. for Cyperaceae, *Ranunculus*
761 *acris*-t., and *Rumex acetosa*-t. (*R. acetosa* in **Bun**). The other taxa have a RPP in **Bun** smaller than the mean RPP in **New**,
762 except *Plantago maritima* that has a larger RPP (5.8) in **Bun** than the mean RPP for *P. lanceolata* in **New**.

763 Of nine taxa, three have a RPP in Kuneš et al. (2019) ([Kun](#)) that is comparable to the mean RPP in [New](#), i.e. for *Plantago*
764 *lanceolata*, *Ranunculus acris*-t. and *Rumex acetosa*-t.. The other six taxa have a RPP larger than the mean RPP in [New](#)
765 (Compositae SF Cichorioideae, Cyperaceae and *Leucanthemum (Anthemis)*-t., or smaller ([Amaranthaceae](#)/Chenopodiaceae,
766 Rubiaceae) to considerably smaller (*Urtica*). Of the 14 tree taxa, only four have a RPP in [Kun](#) comparable to the mean RPP in
767 [New](#), i.e. for *Corylus*, *Fraxinus*, *Salix*, and *Ulmus*. For the other 10 tree taxa, the RPP in K is much smaller than the mean RPP
768 in N for *Abies alba*, *Alnus*, *Carpinus*, *Fagus*, *Picea*, *Pinus*, smaller for *Quercus*, and larger for *Acer* and *Tilia*.

769 Most of the RPP values of the three studies [Twi](#), [Bun](#) and [Kun](#) are in the range of the values selected from the studies included
770 in our synthesis ([New](#)) except for *Urtica*, *Abies alba*, *Carpinus*, and *Pinus* in [Kun](#). The Lagrangian Stochastic Model is used
771 in [Kun](#) instead of the Gaussian Plume Model in [New](#), which may be one of the factors behind the lower RPPs in [Kun](#), in
772 particular (but not only) for taxa with heavy pollen grains.

773

774 Table A2: Comparison of the mean RPPs in this synthesis with the RPP estimates from Britain (Twiddle, 2012),
 775 Greenland (Bunting et al., 2013a) and Czech Republic (Kuneš et al., 2019). Explanations for symbols in the taxa list,
 776 see caption below Table A4. + The original paper does not provide a RPP for Poaceae and values of standard deviations
 777 (SDs) for the RPPs. We extracted the RPP values related to *Picea* from Table 5 in Twiddle et al. (2012). RPPs related
 778 to Poaceae (1.00+) were then calculated by assuming that the RPP of *Picea* was equal to the mean RPP of *Picea* in
 779 Europe (this synthesis) (in bold). ++ The RPPs and their SDs are not listed in the original paper, we therefore
 780 extracted the values from Figure 4 in (Bunting et al., 2013a) and the decimals are approximate. +++ Kuneš et al.
 781 (2019): we chose the RPP values that were considered best by the authors, i.e. using the lake dataset (pollen from lake
 782 sediment), ERV sub-model 1 and the Lagrangian Stochastic Model (for details, see Discussion section, this paper). #
 783 value for *Plantago maritima* and ## two values for *Rumex acetosa* and *Rumex acetosella*, respectively (Bunting et al.,
 784 2013a), for comparison with *Plantago* spp. and *Rumex acetosa*-t. (this paper). Underlined RPPs are close to mean RPPs
 785 (this synthesis).
 786

Study	This paper synthesis RPP (SE)	Twiddle et al. (2012)+ RPP - ERV3 random GPM	Bunting et al. (2013)++ RPP (SE) - ERV1 GPM	Kunes et al (2019)+++ RPP (SE) - R ERV1 LSM
Information on analysis				
HERB TAXA				
Poaceae (Reference taxon)	1.000 (0.000)	1.00+	1.00 (0.00)	1.00 (0.00)
Herb taxa	-	-	-	-
Amaranthaceae/Chenopodiaceae	4.280 (0.270)	-	-	1.58 (0.74)
<i>Calluna vulgaris</i> *	1.085 (0.029)	11.42	-	-
Comp. <i>Leucanthemum</i> (<i>Anthemis</i>)t.***	0.10 (0.01)	-	-	0.94 (0.43)
Comp. SF. Cichorioideae***	0.160 (0.020)	-	-	1.04 (0.64)
Cyperaceae	0.962 (0.050)	-	0.95 (0.05)	2.10 (0.88)
<i>Plantago lanceolata</i> ^^	2.330 (0.201)	-	5.8 (0.3)#	2.24 (0.71)
<i>Potentilla</i> t.^	1.720 (0.200)	-	0.4 (0.03)	-
<i>Ranunculus acris</i> ^^	1.960 (0.360)	-	2.0 (0.1)	1.38 (1.13)
Rubiaceae	3.710 (0.340)	-	-	1.03 (0.74)
<i>Rumex acetos</i> t.	3.020 (0.278)	-	3.5 (0.3)/2.0 (0.1)##	1.94 (1.35)
<i>Urtica</i>	10.520 (0.310)	-	-	1.16 (0.52)
TREE TAXA				
<i>Abies alba</i>	6.875 (1.442)	-	-	1.08 (0.99)
<i>Acer</i>	0.800 (0.230)	-	-	1.25 (0.75)
<i>Alnus</i>	13.562 (0.293)	-	-	2.44 (0.73)
<i>Betula</i> (mainly <i>B. pubescens</i> , <i>B. pendula</i>)	5.106 (0.303)	13.16	3.75 (0.4)	2.53 (0.91)
<i>Carpinus betulus</i>	4.520 (0.425)	-	-	1.36 (0.36)
<i>Corylus avellana</i>	1.710 (0.100)	-	-	2.31 (1.13)
<i>Fagus sylvatica</i>	5.863 (0.176)	-	-	0.88 (0.25)
<i>Fraxinus excelsior</i>	1.044 (0.048)	-	-	0.79 (0.37)

Field Code Changed

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Field Code Changed

Field Code Changed

Formatted: Font: Bold

Formatted: Font: Bold

Field Code Changed

Field Code Changed

Formatted: Font: Bold

<i>Picea abies</i>	<u>5.437 (0.097)</u>	<u>5.44</u>	-	<u>2.39 (0.93)</u>
<i>Pinus</i> (mainly <i>P. sylvestris</i>)	<u>6.058 (0.237)</u>	<u>16.32</u>	-	<u>1.55 (0.44)</u>
<i>Dec. Quercust.</i> (mainly <i>Q. robur</i> , <i>Q. petraea</i>)	<u>4.537 (0.086)</u>	-	-	<u>2.08 (0.46)</u>
<i>Salix</i>	<u>1.182 (0.077)</u>	-	<u>0.7 (0.03)</u>	<u>1.43 (0.62)</u>
<i>Tilia</i>	<u>1.210 (0.116)</u>	-	-	<u>2.30 (1.24)</u>
<i>Ulmus</i>	<u>1.270 (0.050)</u>	-	-	<u>0.96 (0.77)</u>

787
788

Study	This paper, synthesis	<i>Twiddle et al. (2012)+</i>	<i>Bunting et al. (2013)++</i>	<i>Kunes et al (2019)+++</i>
<i>Information on analysis</i>	RPP (SE)	RPP - ERV3 random GPM	RPP (SE) - ERV1 GPM	RPP (SE) - R ERV1 LSM
HERB TAXA				
Poaceae (Reference taxon)	1.000 (0.000)	1.00+	1.00 (0.00)	1.00 (0.00)
Herb taxa				
Amaranthaceae/Chenopodiaceae	4.280 (0.270)			1.58 (0.74)
<i>Calluna vulgaris</i> *	1.085 (0.029)	11.42		
Comp. <i>Leucanthemum (Anthemis)</i> -t***	0.10 (0.01)			0.94 (0.43)
Comp. SF. Cichorioideae***	0.160 (0.020)			1.04 (0.64)
Cyperaceae	0.962 (0.050)		<u>0.95 (0.05)</u>	2.10 (0.88)
<i>Plantago lanceolata</i> ^^	2.330 (0.201)		5.8 (0.3)#	<u>2.24 (0.71)</u>
<i>Potentilla</i> -t^	1.720 (0.200)		0.4 (0.03)	
<i>Ranunculus acris</i> -t^^^	1.960 (0.360)		<u>2.0 (0.1)</u>	<u>1.38 (1.13)</u>
Rubiaceae	3.710 (0.340)			1.03 (0.74)
<i>Rumex acetosa</i> -t	3.020 (0.278)		<u>3.5 (0.3)/ 2.0 (0.1)##</u>	<u>1.94 (1.35)</u>
<i>Urtica</i>	10.520 (0.310)			1.16 (0.52)
TREE TAXA				
<i>Abies alba</i>	6.875 (1.442)			1.08 (0.99)
<i>Acer</i>	0.800 (0.230)			<u>1.25 (0.75)</u>
<i>Alnus</i>	13.562 (0.293)			2.44 (0.73)
<i>Betula</i> (mainly <i>B. pubescens</i> , <i>B. pendula</i>)	5.106 (0.303)	13.16	3.75 (0.4)	2.53 (0.91)
<i>Carpinus betulus</i>	4.520 (0.425)			1.36 (0.36)
<i>Corylus avellana</i>	1.710 (0.100)			<u>2.31 (1.13)</u>
<i>Fagus sylvatica</i>	5.863 (0.176)			0.88 (0.25)
<i>Fraxinus excelsior</i>	1.044 (0.048)			<u>0.79 (0.37)</u>
<i>Picea abies</i>	5.437 (0.097)	5.44		2.39 (0.93)
<i>Pinus</i> (mainly <i>P. sylvestris</i>)	6.058 (0.237)	16.32		1.55 (0.44)
<i>Quercus</i> (mainly <i>Q. robur</i> , <i>Q. petraea</i>)	4.537 (0.086)			2.08 (0.46)
<i>Salix</i>	1.182 (0.077)		0.7 (0.03)	<u>1.43 (0.62)</u>
<i>Tilia</i>	1.210 (0.116)			2.30 (1.24)
<i>Ulmus</i>	1.270 (0.050)			<u>0.96 (0.77)</u>

789

790 Appendix B - Selection of RPP values and calculation of the mean RPPs and their SDs

791 B.1 Methods

792 Tables B1 (Boreal and Temperate Europe) and B2 (Mediterranean Europe) list the RPP values from the 16 selected studies
793 according to the information on models used provided in Appendix C (Table C1) with further explanations on selection of
794 RPP studies. We followed similar procedures and rules as Mazier et al. (2012) and Li et al. (2018) to produce a new standard
795 RPP dataset for Europe. We consider that there are still too few RPP values per taxon to disentangle variability in the RPP
796 values for a particular taxon due to methodological issues, landscape characteristics, land use, or climate. We therefore use the
797 mean of selected RPP values for each taxon in the new standard RPP dataset, following Broström et al. (2008) and Mazier et
798 al. (2012). In boreal and temperate Europe, the number of RPP values per taxon varies between one and nine (*Betula*) (Table
799 B1), and in Mediterranean Europe, there is only one value per taxon (Table B2). In general, all three sub-models of the ERV
800 model were used in the RPP studies. We selected the RPP values obtained with the ERV sub-model considered by the authors
801 to have provided the best results (following the approach of Li et al., 2018). This is usually evaluated **from** by the shape of the
802 curve of likelihood function scores (LFS), or log likelihood (LL) (see e.g. Twiddle et al., 2012) and the LFS and LL values
803 themselves. All RPPs selected for this synthesis are expressed relative to Poaceae (RPP=1). In studies that used another
804 reference taxon and calculated a RPP for Poaceae, the RPPs were recalculated relative to Poaceae. In studies that did not
805 include a RPP value for Poaceae, it was assumed that the reference taxon had a RPP related to Poaceae equal to the mean of
806 the RPP values for that taxon in the other studies (e.g. Mazier et al., 2012). For simplicity, we used the value of *Quercus* (5.83)
807 calculated by Mazier et al. (2012) for the study by Bunting et al. (2005) (*Quercus* as reference taxon, no RPP value for
808 Poaceae). We could also have used the new mean RPP for *Quercus* (4.54) using our selected RPPs (five values, instead of
809 three in Mazier et al. (2012)). The latter would not have changed our results significantly; the mean RPP for *Quercus* would
810 have been 4.28 instead of 4.54 (Table A4). For the study by Baker et al. (2016), we used the RPP values obtained with Poaceae
811 as the reference taxon, given that the RPPs relative to *Quercus* or *Pinus* were almost identical when ERV submodel 3 was
812 used. The selection of RPP values in boreal and temperate Europe for the calculation of the mean RPP values of each taxon
813 (values emphasized **in green** in Table S1.2, A and B) is based on the following rules:

- 814 1. We excluded the RPP values that were not significantly different from zero considering the lower bound of its SE,
815 and values that were considered as uncertain by the authors of the original publications (e.g., *Vaccinium* for Finland
816 (Räsänen et al., 2007), *Pinus* for Central Sweden (von Stedingk et al., 2008)). Moreover, some RPP values were
817 excluded as they were assumed to be outliers or unreliable based on experts' knowledge on the plants involved, the
818 pollen-vegetation dataset, and the field characteristics of the related studies. For example, the RPPs for Cyperaceae,
819 *Potentilla*-t and Rubiaceae obtained in SW Norway (Hjelle, 1998) and those for *Salix* and *Calluna vulgaris* from
820 Central Sweden (von Stedingk et al., 2008) were assumed to be too low compared to the values obtained in other
821 study areas (Mazier et al., 2012).

Formatted: Highlight

822 2. (i) when five or more RPP estimates of pollen productivity ($N \geq 5$) were available for a pollen type, the largest and the
823 smallest RPP values (generally outlier values) were excluded, and the mean was calculated using the remaining three
824 or more RPP estimates; (ii) when $N=4$, the most deviating value was excluded, and the mean calculated using the
825 other three RPP values; (iii) when $N=3$, the mean was based on all values available except if one value was strongly
826 deviating from the other two; and (iv) when $N=2$, the mean was based on the two values available; an exception is
827 *Ulmus* for which we excluded the value from Germany (Theuerkauf et al. 2013) given that several of the RPPs in this
828 study are considerably higher than most values in the other available studies, i.e. for *Betula* (18.7), *Quercus* (17.85)
829 and *Tilia* (12.38). The latter values were also excluded from the mean RPP, as well as the unusually high values found
830 by Baker et al. (2016) for *Betula* (13.94), *Pinus* (23.12) and *Quercus* (18.47). Baker et al. (2016) argue that the high
831 RPP values might be characteristic of temperate deciduous forests that were little impacted by human activities. More
832 studies in this type of wooded environments would be needed to confirm this assumption. In the absence of such
833 studies we consider these values as outliers.

834 The SDs for the mean RPP values were calculated using the delta method (Stuart. and Ord., 1994), a mathematical solution to
835 the problem of calculating the mean of individual SDs (see e.g. Li et al. 2020 for more details).

836 **Table B1: Europe (Mediterranean area excluded): RPP estimates and their SDs (in brackets) with the total number of**
 837 **taxa per study indicated and in brackets the number of taxa with selected RPP estimates. (A) Studies using moss**
 838 **pollsters as pollen samples. (B) Studies using surface lake sediments as pollen samples. For explanation of symbols, see**
 839 **captions below Table B1 (B).**
 840 **(A)**

Type of pollen sample Region ERV submodel	Moss pollsters							
	Finland ERV 3	C Sweden ERV 3	S Sweden# ERV 3	Norway ERV 1	England## ERV 1	Swiss Jura ERV 1	Czech Rep* ERV 1	Poland** ERV 3
HERB TAXA								
Poaceae (Reference taxon)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)
Amaranthaceae/Chenopodiaceae							4.28 (0.27)	
Apiaceae				0.26 (0.009)				
Artemisia							2.77 (0.39)	
Calluna vulgaris		0.30 (0.03)	4.70 (0.69)	1.07 (0.03)				
Cerealia-t			3.20 (1.14)				0.0462 (0.0018)	
Comp. Leucanthemum (Anthemis)-t				0.10 (0.008)				
Comp. SF. Cichorioideae			0.24 (0.06)	0.06 (0.004)				
Cyperaceae	0.002 (0.0022)	0.89 (0.03)	1.00 (0.16)	0.29 (0.01)		0.73 (0.08)		
Empetrum	0.07 (0.06)	0.11 (0.03)						
Ericaceae		0.07 (0.04)						
Filipendula			2.48 (0.82)	3.39 (0.00)				
Plantago lanceolata			12.76 (1.83)	1.99 (0.04)			3.70 (0.77)	
Plantago media						1.27 (0.18)		
Plantago montana						0.74 (0.13)		
Potentilla-t			2.47 (0.38)	0.14 (0.005)		0.96 (0.13)		
Ranunculus acris-t			3.85 (0.72)	0.07 (0.004)				
Rubiaceae			3.95 (0.59)	0.42 (0.01)		3.47 (0.35)		
Rumex acetosa-t			4.74 (0.83)	0.13 (0.004)				
Secale			3.02 (0.05)					
Trollius						2.29 (0.36)		
Urtica							10.52 (0.31)	
Vaccinium	0.01 (0.01)							
TREE TAXA								
Abies						3.83 (0.37)		
Acer			1.27 (0.45)			0.32 (0.10)		
Alnus			4.20 (0.14)		8.74 (0.35)		2.56 (0.32)	15.95 (0.6622)
Betula	4.6 (0.70)	2.24 (0.20)	8.87 (0.13)		6.18 (0.35)			13.94 (0.2293)
Carpinus			2.53 (0.07)					4.48 (0.0301)
Corylus			1.40 (0.04)		1.51 (0.06)			1.35 (0.0512)
Fagus			6.67 (0.17)			1.20 (0.16)		
Fraxinus			0.67 (0.03)		0.70 (0.06)		1.11 (0.09)	
Juniperus		0.11 (0.45)	2.07 (0.04)					
Picea		2.78 (0.21)	1.76 (0.00)			8.43 (0.30)		
Pinus	8.40 (1.34)	21.58 (2.87)	5.66 (0.00)				6.17 (0.41)	23.12 (0.2388)
Quercus			7.53 (0.08)		5.83 (0.00)##		1.76 (0.20)	18.47 (0.1032)
Salix		0.09 (0.03)	1.27 (0.31)		1.05 (0.17)		1.19 (0.12)	
Sambucus nigra-t							1.30 (0.12)	
Tilia			0.80 (0.03)				1.36 (0.26)	0.98 (0.0263)
Ulmus			1.27 (0.05)					
Total number of taxa 39 (38)	6 (4)	10 (7)	26 (25)	12 (8)	7 (7)	11(10)	13(12)	8 (5)

841
842

Type of pollen sample	Moss polsters							
	Region	C	S	England	Swiss	Czech Rep*	Poland*	
	Finland	Sweden	Sweden#	Norway	##	Jura		*
ERV submodel	ERV 3	ERV 3	ERV 3	ERV 1	ERV 1	ERV 1	ERV 1	ERV 3
HERB TAXA	-	1.00	1.00	1.00	1.00	1.00	-	1.00
Poaceae (Reference taxon)	1.00 (0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	1.00 (0.00)	(0.00)
Amaranthaceae/Chenopodiaceae	-	-	-	0.26 (0.009)	-	-	4.28 (0.27)	-
Apiaceae	-	-	-	-	-	-	2.77 (0.39)	-
Artemisia	-	0.30 (0.03)	4.70 (0.69)	1.07 (0.03)	-	-	-	-
Calluna vulgaris	-	-	3.20 (1.14)	-	-	-	0.0462 (0.0018)	-
Cerealia t.	-	-	-	0.10 (0.008)	-	-	-	-
Comp.	-	-	0.24 (0.06)	0.06 (0.004)	-	-	-	-
Leucanthemum(Anthemis) t.	-	-	-	-	-	-	-	-
Comp. SF. Cichorioideae	-	0.89 (0.03)	1.00 (0.16)	0.29 (0.01)	-	0.73 (0.08)	-	-
Cyperaceae	0.002 (0.0022)	-	-	-	-	-	-	-
Empetrum	0.07 (0.06)	0.11 (0.03)	-	-	-	-	-	-
Ericaceae	-	0.07 (0.04)	-	-	-	-	-	-
Filipendula	-	-	2.48 (0.82)	3.39 (0.00)	-	-	-	-
Plantago lanceolata	-	-	12.76 (1.83)	1.99 (0.04)	-	-	3.70 (0.77)	-
Plantago media	-	-	-	-	-	1.27 (0.18)	-	-
Plantago montana	-	-	-	-	-	0.74 (0.13)	-	-
Potentilla t.	-	-	2.47 (0.38)	0.14 (0.005)	-	0.96 (0.13)	-	-
Ranunculus acrist.	-	-	3.85 (0.72)	0.07 (0.004)	-	-	-	-
Rubiaceae	-	-	3.95 (0.59)	0.42 (0.01)	-	3.47 (0.35)	-	-
Rumex acetosat.	-	-	4.74 (0.83)	0.13 (0.004)	-	-	-	-
Secale	-	-	3.02 (0.05)	-	-	-	-	-
Trollius	-	-	-	-	-	2.29 (0.36)	-	-
Urtica	-	-	-	-	-	-	10.52 (0.31)	-
Vaccinium	0.01 (0.01)	-	-	-	-	-	-	-

843

<u>TREE TAXA</u>								
	-	-	-	-	-	<u>3.83</u>	-	-
<i>Abies</i>	-	-	<u>1.27</u>	-	-	<u>(0.37)</u>	-	-
<i>Acer</i>	-	-	<u>(0.45)</u>	-	-	<u>0.32</u>	-	-
	-	-	<u>4.20</u>	-	-	<u>(0.10)</u>	-	-
<i>Alnus</i>	-	-	<u>(0.14)</u>	-	<u>8.74 (0.35)</u>	-	<u>2.56</u>	<u>15.95</u>
	-	-	<u>(0.13)</u>	-	-	-	<u>(0.32)</u>	<u>(0.6622)</u>
<i>Betula</i>	<u>4.6 (0.70)</u>	<u>2.24 (0.20)</u>	<u>8.87</u>	-	<u>6.18 (0.35)</u>	-	-	<u>13.94</u>
	-	-	<u>(0.07)</u>	-	-	-	-	<u>(0.2293)</u>
<i>Carpinus</i>	-	-	<u>2.53</u>	-	-	-	-	<u>4.48 (0.0301)</u>
	-	-	<u>(0.04)</u>	-	-	-	-	-
<i>Corylus</i>	-	-	<u>1.40</u>	-	<u>1.51 (0.06)</u>	-	-	<u>1.35 (0.0512)</u>
	-	-	<u>(0.04)</u>	-	-	-	-	-
<i>Fagus</i>	-	-	<u>6.67</u>	-	-	<u>1.20</u>	-	-
	-	-	<u>(0.17)</u>	-	-	<u>(0.16)</u>	-	-
<i>Fraxinus</i>	-	-	<u>0.67</u>	-	<u>0.70 (0.06)</u>	-	<u>1.11</u>	-
	-	-	<u>(0.03)</u>	-	-	-	<u>(0.09)</u>	-
<i>Juniperus</i>	-	<u>0.11 (0.45)</u>	<u>2.07</u>	-	-	-	-	-
	-	-	<u>(0.04)</u>	-	-	-	-	-
<i>Picea</i>	-	-	<u>1.76</u>	-	-	<u>8.43</u>	-	-
	-	<u>2.78 (0.21)</u>	<u>(0.00)</u>	-	-	<u>(0.30)</u>	-	-
<i>Pinus</i>	<u>8.40</u>	<u>21.58</u>	<u>5.66</u>	-	-	-	<u>6.17</u>	<u>23.12</u>
	<u>(1.34)</u>	<u>(2.87)</u>	<u>(0.00)</u>	-	-	-	<u>(0.41)</u>	<u>(0.2388)</u>
Deciduous <i>Quercus</i> .	-	-	<u>7.53</u>	-	<u>5.83</u>	-	<u>1.76</u>	<u>18.47</u>
	-	-	<u>(0.08)</u>	-	<u>(0.00)##</u>	-	<u>(0.20)</u>	<u>(0.1032)</u>
<i>Salix</i>	-	<u>0.09 (0.03)</u>	<u>1.27</u>	-	<u>1.05 (0.17)</u>	-	<u>1.19</u>	-
	-	-	<u>(0.31)</u>	-	-	-	<u>(0.12)</u>	-
<i>Sambucus nigrat.</i>	-	-	-	-	-	-	<u>1.30</u>	-
	-	-	-	-	-	-	<u>(0.12)</u>	-
<i>Tilia</i>	-	-	<u>0.80</u>	-	-	-	<u>1.36</u>	<u>0.98 (0.0263)</u>
	-	-	<u>(0.03)</u>	-	-	-	<u>(0.26)</u>	-
<i>Ulmus</i>	-	-	<u>1.27</u>	-	-	-	-	-
	-	-	<u>(0.05)</u>	-	-	-	-	-
Total number of taxa 39 (38)	6 (4)	10 (7)	26 (25)	12 (8)	7 (7)	11(10)	13(12)	8 (5)

844

845 (B)

Type of pollen sample Region ERV submodel	lake surface sediment				
	Estonia ERV 3	Denmark ERV 1	Swiss Plateau	Germany*** ERV 3	Germany ****
HERB TAXA					
Poaceae (Reference taxon)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)
<i>Artemisia</i>	3.48 (0.20)				5.56 (0.020)
<i>Calluna vulgaris</i>		1.10 (0.05)	0.00076 (0.0019)		
Cerealia-t	1.60 (0.07)	0.75 (0.04)	0.17 (0.03)	9.00 (1.92)	0.08 (0.001)
Compositae <i>Leucanthemum (Anthemis)-t</i>			0.24 (0.15)		
Cyperaceae	1.23 (0.09)				
<i>Filipendula</i>	3.13 (0.24)				
<i>Plantago lanceolata</i>		0.90 (0.23)			2.73 (0.043)
<i>Rumex acetosa -t</i>		1.56 (0.09)			2.76 (0.022)
<i>Secale</i>				4.08 (0.96)	4.87 (0.006)
TREE TAXA					
<i>Alnus</i>	13.93 (0.15)		9.92 (2.86)	15.51 (1.25)	13.68 (0.049)
<i>Betula</i>	1.81 (0.02)		2.42 (0.39)	9.62 (1.92)	19.70 (0.117)
<i>Carpinus</i>			4.56 (0.85)	9.45 (0.51)	
<i>Corylus</i>			2.58 (0.39)	0.76 (0.17)	
<i>Fagus</i>		5.09 (0.22)	1.39 (0.21)	5.83 (0.45)	9.63 (0.008)
<i>Fraxinus</i>				6.74 (0.68)	1.35 (0.012)
<i>Juniperus</i>			0.57 (0.16)		
<i>Picea</i>	4.73 (0.13)	1.19 (0.42)	1.35 (0.45)	1.58 (0.28)	5.81 (0.007)
<i>Pinus</i>	5.07 (0.06)			5.66 (0.00)	5.39 (0.222)
<i>Populus</i>			2.56 (0.39)	2.66 (1.25)	
<i>Quercus</i>	7.39 (0.20)			2.15 (0.17)	17.85 (0.049)
<i>Salix</i>	2.31 (0.08)				
<i>Tilia</i>				1.47 (0.23)	12.38 (0.101)
<i>Ulmus</i>					11.51 (0.101)
Total number of taxa (selected values) 23 (22)	11 (11)	7 (7)	13 (9)	13 (10)	15 (11)

846

847 B)

Type of pollen sample Region ERV submodel	lake surface sediment				
	Estonia ERV 3	Denmark ERV 1	Swiss Plateau	Germany** ERV 3	Germany ****
HERB TAXA					
Poaceae (Reference taxon)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)	1.00 (0.00)
<i>Artemisia</i>	3.48 (0.20)				5.56 (0.020)
<i>Calluna vulgaris</i>		1.10 (0.05)	0.00076 (0.0019)		
Cerealia t.	1.60 (0.07)	0.75 (0.04)	0.17 (0.03)	9.00 (1.92)	0.08 (0.001)
Compositae <i>Leucanthemum(Anthemis) t.</i>			0.24 (0.15)		
Cyperaceae	1.23 (0.09)				

<i>Filipendula</i>	3.13 (0.24)	-	-	-	-
<i>Plantago lanceolata</i>	-	0.90 (0.23)	-	-	2.73 (0.043)
<i>Rumex acetosa</i> t.	-	1.56 (0.09)	-	-	2.76 (0.022)
<i>Secale</i>	-	-	-	4.08 (0.96)	4.87 (0.006)
TREE TAXA	-	-	9.92 (2.86)	-	-
<i>Alnus</i>	13.93 (0.15)	-	2.42 (0.39)	15.51 (1.25)	13.68 (0.049)
<i>Betula</i>	1.81 (0.02)	-	4.56 (0.85)	9.62 (1.92)	19.70 (0.117)
<i>Carpinus</i>	-	-	2.58 (0.39)	9.45 (0.51)	-
<i>Corylus</i>	-	-	0.76 (0.17)	-	-
<i>Fagus</i>	-	5.09 (0.22)	1.39 (0.21)	5.83 (0.45)	9.63 (0.008)
<i>Fraxinus</i>	-	-	-	6.74 (0.68)	1.35 (0.012)
<i>Juniperus</i>	-	-	0.57 (0.16)	-	-
<i>Picea</i>	4.73 (0.13)	1.19 (0.42)	1.35 (0.45)	1.58 (0.28)	5.81 (0.007)
<i>Pinus</i>	5.07 (0.06)	-	-	5.66 (0.00)	5.39 (0.222)
<i>Populus</i>	-	-	2.56 (0.39)	2.66 (1.25)	-
Deciduous <i>Quercus</i> t.	7.39 (0.20)	-	-	2.15 (0.17)	17.85 (0.049)
<i>Salix</i>	2.31 (0.08)	-	-	-	-
<i>Tilia</i>	-	-	-	1.47 (0.23)	12.38 (0.101)
<i>Ulmus</i>	-	-	-	-	11.51 (0.101)
Total number of taxa (selected values) 23 (22)	11 (11)	7 (7)	13 (9)	13 (10)	15 (11)

848

849
850 # RPPs for herbs from Broström et al. (2004); RPPs for trees from Sugita et al. (1999) (reference taxon *Juniperus*), converted
851 to Poaceae as reference taxon by Broström et al. (2004).
852 ## Bunting et al. (2005), reference taxon *Quercus* and no RPP for Poaceae; RPPs relative to Poaceae calculated by Mazier et
853 al. (2012) assuming that the RPP of *Quercus* relative to Poaceae is the same as the mean RPP of *Quercus* from three other
854 studies in NW Europe.
855 * New RPPs from the Czech Republic (Abraham and Kozáková, 2012).
856 ** New RPPs from Poland. Poaceae as reference taxa (see text for more details)
857 *** New RPPs from Germany (Matthias et al., 2012), reference taxon *Pinus*. RPPs converted to Poaceae as reference
858 taxon. We selected the RPP estimates obtained with the dataset of vegetation cover including only the trees that had reached
859 their flowering age (allFIDage) (for more information, see Matthias et al., 2012).
860 **** New RPPs from Germany (Theuerkauf et al., 2012); in the original publication, the ERV analysis was performed with
861 the Lagrangian Stochastic Model (LSM) for dispersal of pollen and with *Pinus* as reference taxon. For this synthesis, Martin
862 Theuerkauf redid the analysis with the Gaussian Plume Model for dispersal of pollen (Parsons and Prentice, 1981; Prentice
863 and Parsons, 1983) and with Poaceae as reference taxon.
864 **Green:** selected RPP estimates to be included in the mean RPP values.
865 **Red:** RPP estimates excluded because $SE \geq RPP$.
866 **Orange:** RPP estimates excluded because of a too large difference with the other available estimates and their mean (less than
867 half or more than double the mean RPP).
868 **Light blue:** RPP estimates excluded due to its extreme high value compared to the other available estimates (much over double
869 the mean of the other RPPs), i.e. from the study at Bialowice forest (Poland, Baker et al., 2016) for *Betula*, *Pinus* and *Quercus*,
870 Central Sweden (von Stedingk et al., 2008) for *Pinus*, and Germany**** (Theuerkauf et al., 2013) for *Betula*, *Quercus*, *Tilia*,
871 and *Ulmus*.

Formatted: Highlight

872 **Table B2: Mediterranean area: RPP estimates and their SDs from two available studies, and mean RPPs for northern**
873 **and temperate Europe (Table A1, Appendix A), for comparison. The single RPPs emphasized in green were used in the**
874 **REVEALS reconstruction for Europe (this paper). The plant taxa emphasized in bold are sub-Mediterranean and/or**
875 **Mediterranean plant species and genera. The values emphasized with grey shadow are the mean RPPs that were used**
876 **in the REVEALS reconstruction (this paper) for entire Europe (Mediterranean area included). See Appendix AB for**
877 **more details. FSP values: from Mazier et al. (2012) except (') new values from Mazier et al. (unpubl.), (") value from**
878 **Abraham and Kózáková (2012), ("" value from (Commerford et al., 2013). *, **FSP from Mazier et al. (2012) used in**
880 **the REVEALS reconstruction (this study) for Ericaceae (Medit)* and evergreen Quercus t. evergreen** instead of the**
new FSP values from Mazier et al. (unpubl.); for more explanations, see Discussion section, this paper.

Region	France Medit. (ERV3)			Romania (ERV3)			Europe. Medit. excluded		
Study reference	Mazier et al. (unpubl.)			Grindean et al. (2019)			This paper (Tables A1)		
	RPP	SD	FSP	RPP	SD	FSP	RPP	SD	FSP
HERB TAXA	-	-	-	-	-	-	-	-	-
Poaceae (reference taxon)	1.000	0.000	0.035	1.00	0.00	0.035	1.00	0.00	0.035
<i>Apiaceae</i>	-	-	-	5.91	1.23	0.042	0.26	0.01	0.042
<i>Artemisia</i>	-	-	-	5.89	3.16	0.014"	3.937	0.146	0.014"
Compositae (Asteroideae + Cichorioideae)	-	-	-	0.16	0.10	0.029	-	-	-
Comp. SF. Asteroideae (<i>Anthemis</i> t., <i>Leucanthemum</i>)	-	-	-	-	-	-	0.10	0.01	0.029
Comp. SF. Cichorioideae	1.162	0.675	0.061'	-	-	-	0.16	0.02	0.05
Cerealia (Cerealia t. + <i>Triticum</i> t. + <i>Secale</i> + <i>Zea</i>)	-	-	-	0.22	0.12	0.060	-	-	-
Cerealia t. (Cerealia t., <i>Secale</i> excluded)	-	-	-	-	-	-	1.85	0.38	0.060
Cerealia - <i>Secale cereale</i>	-	-	-	-	-	-	3.99	0.33	0.060
Fabaceae	-	-	-	0.40	0.07	0.021""	-	-	-
<i>Plantago lanceolata</i>	-	-	-	0.58	0.32	0.029	2.33	0.20	0.029
Ranunculaceae	2.038	0.335	0.020'	-	-	-	-	-	-
Ranunculaceae - <i>Ranunculus acris</i> t.	-	-	-	-	-	-	1.96	0.36	0.014
Ranunculaceae - <i>Trollius</i>	-	-	-	-	-	-	2.29	0.36	0.013
Rosaceae (<i>Filipendula</i> , <i>Potentilla</i> t., <i>Sanguisorba</i>)	-	-	-	0.29	0.12	0.018	-	-	-
Rosaceae - <i>Filipendula</i>	-	-	-	-	-	-	3.00	0.28	0.006
Rosaceae - <i>Potentilla</i> t.	-	-	-	-	-	-	1.72	0.20	0.018
Rubiaceae	-	-	-	0.40	0.07	0.019	3.71	0.34	0.019
TREE/SHRUB TAXA	-	-	-	-	-	-	-	-	-
<i>Acer</i>	-	-	-	0.30	0.09	0.056	0.80	0.23	0.056
<i>Buxus sempervirens</i>	1.890	0.068	0.032'	-	-	-	-	-	-
<i>Carpinus betulus</i>	-	-	-	-	-	-	4.52	0.43	0.042
<i>Carpinus orientalis</i>	-	-	-	0.24	0.07	0.042	-	-	-
<i>Castanea sativa</i>	3.258	0.059	0.010'	-	-	-	-	-	-

Commented [MG2]: Replace green by another way to emphasize these values; DELETE orange!!! Not needed!

Formatted: Highlight

Formatted: Highlight

Field Code Changed

Formatted: Font: Bold

<i>Corylus avellana</i>	3.440	0.890	0.025	-	-	-	1.71	0.10	0.025
Cupressaceae (<i>Juniperus communis</i> , <i>J. phoenicea</i> , <i>J. oxycedrus</i>)	1.618	0.161	0.020'	-	-	-	-	-	-
Cupressaceae - <i>Juniperus communis</i>	-	-	-	-	-	-	2.07	0.04	0.016
Ericaceae (<i>Arbutus unedo</i> , <i>Erica arborea</i> , <i>E. cinerea</i> , <i>E. multiflora</i>)	4.265	0.094	0.051'	-	-	-	-	-	-
Ericaceae (<i>Vaccinium</i> dominant, <i>Calluna</i> excluded)	-	-	-	-	-	-	0.07	0.04	0.038*
<i>Fraxinus excelsior</i>	-	-	-	-	-	-	1.04	0.02	0.022
<i>Fraxinus</i> (<i>F. excelsior</i> , <i>F. ornus</i>)	-	-	-	2.99	0.88	0.022	-	-	-
<i>Phillyrea</i>	0.512	0.076	0.015'	-	-	-	-	-	-
<i>Pistacia</i>	0.755	0.201	0.030'	-	-	-	-	-	-
Evergreen <i>Quercus</i> t. (<i>Q. ilex</i> , <i>Q. coccifera</i>)	11.04	3	0.261	0.015'	-	-	-	-	-
Deciduous <i>Quercus</i> t. (<i>Q. spp.</i> , <i>Q. pedunculata</i> dominant)	-	-	-	1.10	0.35	0.035	-	-	-
Deciduous <i>Quercus</i> t. (<i>Q. petraea</i> + <i>Q. rubra</i>)	-	-	-	-	-	-	4.54	0.09	0.035**
Total number of taxa	11	-	-	13	-	-	-	-	-

881
882

Region	France Medit. (ERV3)			Roumania (ERV3)			Europe, Medit. excluded		
Study reference	Mazier et al. (unpubl.)			Grindean et al. (2019)			This paper (Tables 2A, 2B)		
	RPP	SD	FSP	RPP	SD	FSP	RPP	SD	FSP
HERB TAXA									
Poaceae (reference taxon)	1.000	0.000	0.035	1.00	0.00	0.035	1.00	0.00	0.035
Apiaceae				5.91	1.23	0.042	0.26	0.01	0.042
<i>Artemisia</i>				5.89	3.16	0.014''	3.937	0.146	0.014''
Asteraceae (Asteroideae + Cichorioideae)				0.16	0.10	0.029			
Asteraceae Asteroideae (<i>Anthemis</i> t., <i>Leucanthemum</i>)							0.10	0.01	0.029
Asteraceae Cichorioideae	1.162	0.675	0.061'				0.16	0.02	0.05
Cerealia (Cerealia t. + <i>Triticum</i> t. + <i>Secale</i> + <i>Zea</i>)				0.22	0.12	0.060			
Cerealia (Cerealia t., <i>Secale</i> excluded)							1.85	0.38	0.060
Cerealia - <i>Secale cereale</i>							3.99	0.33	0.060
Fabaceae				0.40	0.07	0.021'''			
<i>Plantago lanceolata</i>				0.58	0.32	0.029	2.33	0.20	0.029
Ranunculaceae	2.038	0.335	0.020'						
Ranunculaceae - <i>Ranunculus acris</i> t.							1.96	0.36	0.014
Ranunculaceae - <i>Trollius</i>							2.29	0.36	0.013
Rosaceae (<i>Filipendula</i> , <i>Potentilla</i> t., <i>Sanguisorba</i>)				0.29	0.12	0.018			
Rosaceae - <i>Filipendula</i>							3.00	0.28	0.006
Rosaceae - <i>Potentilla</i> t.							1.72	0.20	0.018
Rubiaceae				0.40	0.07	0.019	3.71	0.34	0.019
TREE/SHRUB TAXA									
<i>Acer</i>				0.30	0.09	0.056	0.80	0.23	0.056
<i>Buxus sempervirens</i>	1.890	0.068	0.032'						
<i>Carpinus betulus</i>							4.52	0.43	0.042
<i>Carpinus orientalis</i>				0.24	0.07	0.042			
<i>Castanea sativa</i>	3.258	0.059	0.010'						
<i>Corylus avellana</i>	3.440	0.890	0.025				1.71	0.10	0.025
Cupressaceae (<i>Juniperus communis</i> , <i>J. phoenicea</i> , <i>J. oxycedrus</i>)	1.618	0.161	0.020'						
Cupressaceae - <i>Juniperus communis</i>							2.07	0.04	0.016
Ericaceae (<i>Arbutus unedo</i> , <i>Erica arborea</i> , <i>E. cinerea</i> , <i>E. multiflora</i>)	4.265	0.094	0.051'						
Ericaceae (<i>Vaccinium</i> dominant, <i>Calluna</i> excluded)							0.07	0.04	0.038*
<i>Fraxinus excelsior</i>							1.04	0.02	0.022
<i>Fraxinus</i> (<i>F. excelsior</i> , <i>F. ornus</i>)				2.99	0.88	0.022			
<i>Phillyrea</i>	0.512	0.076	0.015'						
<i>Pistacia</i>	0.755	0.201	0.030'						
<i>Quercus</i> evergreen (<i>Q. ilex</i> , <i>Q. coccifera</i>)	11.043	0.261	0.015'						
<i>Quercus</i> deciduous (<i>Q. spp.</i> , <i>Q. peduncularis</i> dominant)				1.10	0.35	0.035			
<i>Quercus</i> deciduous (<i>Q. petraea</i> + <i>Q. rubra</i>)							4.54	0.09	0.035**
Total number of taxa	11			13					

883

884 **Appendix C - Selection of RPP studies**

885 **C.1 Introduction**

886 The most common method to estimate RPPs involves the application of the Extended R-Value (ERV) model on datasets of
887 modern pollen assemblages and related vegetation cover. A summary of the ERV model and its assumptions, and an extensive
888 description of standardised field methods for the purpose of RPP studies are found in Bunting et al. (2013). Estimation of RPPs
889 in Europe started with the studies by Sugita et al. (1999) and Broström et al. (2004) in Southern Sweden, and Nielsen et al.
890 (2004) in Denmark. The first tests of the RPP in pollen-based reconstructions of plant cover using the LRA's REVEALS
891 (REgional VEgetation Abundance from Large Sites) model (Sugita, 2007a) were published by Soepboer et al. (2007) in
892 Switzerland and Hellman et al. (2008a and b) in South Sweden. Over the last 15 years, a large number of RPP studies have
893 been undertaken in Europe North of the Alps, but it is only recently that RPP studies were initiated in the Mediterranean area
894 (Grindean et al., 2019; Mazier et al., unpublished). Two earlier syntheses of RPPs in Europe were published by Broström et al.
895 al. (2008) and Mazier et al. (2012). From 2012 onwards, these RPP values have been used in numerous applications of the
896 LRA's two models REVEALS and LOVE (LOcal VEgetation Estimates) (Sugita, 2007a and b) to reconstruct regional and
897 local plant cover in Europe (Cui et al., 2013; Fyfe et al., 2013; Marquer et al., 2020; Mazier et al., 2015; Nielsen et al., 2012;
898 Nielsen and Odgaard, 2010; Trondman et al., 2015). Recently, Wiczorek and Herzschuh (2020) published a synthesis of the
899 RPPs available for the Northern Hemisphere; it includes new mean RPP values for Europe that were produced independently
900 from the synthesis we present here.



901 **C.2 Selection of RPP studies and related information on methods used**

902 The synthesis of mean RPPs presented here was produced in 2018 and applied in REVEALS reconstructions 2018-2020. Of
903 nineteen RPP studies available (in July 2021), we selected fifteen published between 1998 and 2018 and one unpublished
904 study in 2018 (Grindean et al., 2019). The sixteen study regions are distributed in twelve European countries (Figure C1) and
905 detailed in Table C1. Three studies are not included in our synthesis: Britain (Twiddle et al., 2012) because of the absence of
906 Poaceae in the calculated RPPs, curves of likelihood function scores exhibiting departures from theoretically correct curves,
907 and doubts expressed by the authors on the reliability of the values; Greenland (Bunting et al., 2013) because this land area
908 was not included in the REVEALS reconstruction of Holocene plant cover in Europe presented in this paper; and Czech
909 Republic (Kuneš et al., 2019) because the study was not ready when we finalized our synthesis. However, we compare the
910 RPP values from these three studies with the mean RPP values in this synthesis (Appendix A, Table A2).
911 All studies used the ERV model to calculate RPPs, and all but one study used modern pollen assemblages and vegetation; only
912 Nielsen et al. (2004; Denmark) used historical pollen and vegetation data. Eleven studies used pollen assemblages from moss
913 pollsters, five studies from lake sediments. Grindean et al. (2019; Romania) also used some pollen assemblages from surface

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Normal



914

915
916
917
918
919
920
921
922

Figure C1: Location of the selected studies of relative pollen productivities (RPP) in Europe. 1. Britain, (Bunting et al., 2005); 2. Czech Republic, (Abraham and Kozáková, 2012); 3. Denmark, (Nielsen, 2004); 4. Estonia, (Poska et al., 2011); 5. Finland, (Räsänen et al., 2007); 6. France, Mazier et al. unpublished; 7. Germany, (Matthias et al., 2012); 8. Germany, (Theuerkauf et al., 2012); 9. Norway, (Hjelle, 1998); 10. Poland, (Baker et al., 2016); 11. Romania, (Grindean et al., 2019); 12. Sweden, (von Stedingk et al., 2008); 13. Sweden, (Sugita et al., 1999); 14. Sweden, (Broström et al., 2004); 15. Switzerland, (Soepboer et al., 2007); 16. Switzerland, (Mazier et al., 2008).

Formatted: Centered
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold, French (France)
Formatted: French (France)
Formatted: Font: Bold, French (France)
Field Code Changed
Formatted: French (France)
Field Code Changed
Formatted: Font: Bold, French (France)
Formatted: French (France)
Field Code Changed
Formatted: Font: Bold
Formatted: French (France)
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold
Field Code Changed
Formatted: Font: Bold

923 soil samples. All studies used distance-weighted vegetation except two, Hjelle et al. (1998; SW Norway) and Sugita et al.
924 (1999; S Sweden). The Gaussian Plume Model (GPM) was used for pollen dispersal and deposition to distance-weight
925 vegetation, i.e. the Prentice's bog model (Parsons and Prentice, 1981; Prentice and Parsons, 1983) in studies using pollen from
926 moss pollsters, and the Sugita's lake model (Sugita, 1993) in studies using pollen from lake sediments (see also caption of
927 Table C1). In the case of the study by Theuerkauf et al. (2013), the published RPP values were calculated using the Lagrangian
928 Stochastic Model. For the purpose of this synthesis, Theuerkauf recalculated the RPPs using the GPM bog model in the
929 application of the ERV model. The distribution of sites for collection of pollen samples and vegetation data within the study
930 regions is random or random stratified in seven of the eleven studies using moss pollsters; the five remaining studies used
931 selected sites (or systematic distribution). Studies using lake sediments normally result in a systematic site distribution.
932 Broström et al. (2005) and Twiddle et al. (2012) showed that random distribution of sites provided better estimates of "relevant
933 source area of pollen" (RSAP; *sensu* Sugita, 1994) and thus of RPPs, given that the reliable RPPs are those obtained at the
934 RSAP distance and beyond. Both studies indicated that systematic distribution of sites have the tendency to result in curves of
935 likelihood function scores that do not follow the theoretical behaviour, i.e. an increase of the scores with distance until the
936 values reach an asymptote. However, the difference in RPPs between systematic and random sampling is generally not very
937 large. Nonetheless, systematic sampling may lead to uncertainty in terms of reliability of RPPs and random distribution of
938 sites is recommended and has generally been used in studies using moss pollsters or soil samples published from 2008 and
939 onwards.

940 Table C1: Selection of studies for the synthesis of relative pollen productivity (RPP) estimates. Emphasized in bold: additional, new
941 studies compared to the studies included in the synthesis of Mazier et al. (2012). Symbols: ¹L=lakes; M=moss pollsters; S=surface
942 soil; ²Other distance-weighting models were used in most studies, including the Gaussian Plume Model (GPM), 1/d, 1/d² (d=distance)
943 and the Lagrangian Stochastic Model (LSM). The GPM is used in both the model developed for bogs (Parsons and Prentice, 1981;
944 Prentice and Parsons, 1983) and lakes (Sugita, 1993). For this RPP synthesis, we chose the results from the analyses using GPM
945 rather than 1/d or 1/d². Note: In the study of Theuerkauf et al. (2013) the LSM was used. For this synthesis, Theuerkauf recalculated
946 his RPPs using the lake model developed by Sugita (1993); ³ Number of plant taxa for which RPP was estimated, including the
947 reference taxon. Note: In the study by Theuerkauf et al. (2013) RPPs were estimated for 17 taxa using LSM. The RPPs were
948 recalculated using the lake model (Sugita, 1993) for 15 taxa (see note under ² above) for this synthesis. In the study of Sugita et al.
949 (1999) RPPs were calculated for 14 trees and 3 herbs. We used only the values for the 14 trees in this synthesis, following the syntheses
950 by Broström et al. (2008) and Mazier et al. (2012); [^] Britain: the study includes two areas (a and b) in which RPP estimates were
951 calculated for different sets of taxa and the two areas have different numbers of sites: a. Calthorpe (34), 5 taxa; b. Wheatfen (17,
952 same 5 taxa and *Corylus* (6 taxa in total); ^{^^} random distribution restricted to areas of the study region with existing vegetation
953 maps (therefore no sites outside these areas); i.e. study region including separate areas (Mazier et al., 2008). ^{*} Vegetation data from
954 historical maps around 1800 CE; ⁺ lake sediments dated to ca. 1800; ^{*} The reference taxon used in the original study is different
955 from Poaceae. For this synthesis the RPPs were converted to values relative to Poaceae; ^{**} The study of Bunting et al. (2005) does
956 not include a RPP for Poaceae. In order to calculate the RPPs relative to Poaceae, it was assumed that the RPP of *Quercus* was equal
957 to the mean of RPPs from three other studies in Europe (see Mazier et al., 2012 for details). Although we have included new RPP
958 values for *Quercus* in this synthesis, we did not recalculate the RPPs from Bunting et al. (2005) with a new mean value for *Quercus*,
959 but used the same values as in Mazier et al. (2012). For comparison, the mean value for *Quercus* using the RPPs of the additional
960 studies included in this synthesis is 4.28 (instead of 5.83 in Mazier et al., 2012). This would imply slightly lower RPPs in Britain also
961 for *Alnus*, *Betula*, *Corylus*, *Fraxinus* and *Salix*. # no distance weighting used for vegetation data because there was no information
962 about vegetation with increasing distance from the pollen sample (Hjelle et al., 1998; Sugita et al., 1999). In the Swedish study,
963 vegetation data within a 10² m² (herb taxa) and 10³ m² quadrat (tree taxa) centred on the pollen sample was used (Sugita et al., 1999).
964

965

Country	Region	No sites	Site distrib.	Pollen sample ¹	ERV sub-model	Distance weighting model ²	Reference taxon	No taxa ³	Reference
Britain	East Anglian; Norfolk woodlands	(34 + 19) [^]	selected	M	1	GPM Prentice's bog	<i>Quercus</i> Poaceae**	6	Bunting et al. 2005
Czech Republic	Central Bohemia: agricultural landscape	54	stratified random	M	1	GPM Prentice's bog	Poaceae	13	Abraham & Kózáková 2012
Denmark	Ancient agricultural landscape ⁺	30	selected	L ⁺⁺	1	GPM Sugita's lake	Poaceae	7	Nielsen et al. 2004
Estonia	Hemiboreal forest zone: mixed woodland - agricultural landscape	40	selected	L	3	GPM Sugita's lake	Poaceae	10	Poska et al. 2011
Finland	N Finland	24	stratified random	M	3	GPM Prentice's bog	Poaceae	6	Räsänen et al. 2007
France	Mediterranean region	23	random	M	3	GPM Prentice's bog	Poaceae	11	Mazier et al. unpubl.
Germany	Eastern Germany: Brandenburg, agricultural landscape	49	selected	L	3	GPM Sugita's lake	<i>Pinus</i> Poaceae*	16	Matthias et al. 2012
	NE Germany: agricultural landscape	27	selected	L	3	LSM GPM Sugita's Lake ²	<i>Pinus</i> Poaceae*	11 (15) ³	Theuerkauf et al. 2013
Norway	SW Norway: Hordaland and Sogn og Fjordane, mown or grazed grass-land and heath	39	selected	M	1	None [#]	Poaceae	17	Hjelle 1998
Poland	NE Poland: Bialowieza Forest	18	stratified random	M	3	GPM Prentice's bog	Poaceae	8	Baker et al. 2016
Romania	SE Romania: Forest-steppe region	26	random	M & S	3	GPM Prentice's bog	Poaceae	13	Grindean et al. 2019
Sweden	West- Central Sweden: Forest-tundra ecotone	30	random	M	3	GPM Prentice's bog	Poaceae	10	von Stedingk et al. 2008
	S Sweden: ancient cultural landscapes	114	selected	M	3	None [#]	<i>Juniperus</i> Poaceae*	14 (17) ³	Sugita et al. 1999
	S Sweden: unfertilized mown or grazed grasslands	42	selected	M	3	GPM Prentice's bog	Poaceae	11	Broström et al. 2004

966 **Table C1: Selection of studies for the synthesis of relative pollen productivity (RPP) estimates. Emphasized in bold:**
 967 **additional, new studies compared to the studies included in the synthesis of Mazier et al. (2012). For explanation of**
 968 **symbols, see captions below the Table.**

Country	Region	No sites	Site distrib.	Pollen sample [†]	ERV sub-model	Distance weighting model [‡]	Reference taxon	No taxa [§]	Reference
Britain	East Anglian: Norfolk woodlands	(34 + 19) ^Δ	selected	M	1	GPM Prentice's bog	<i>Quercus</i> Poaceae ^{*,§}	6	Bunting et al. 2005
Czech Republic	Central Bohemia: agricultural landscape	54	stratified random	M	1	GPM Prentice's bog	Poaceae	13	Abraham & Kóžaková 2012
Denmark	Ancient agricultural landscape [~]	30	selected	L ⁺⁺	1	GPM Sugita's lake	Poaceae	7	Nielsen et al. 2004
Estonia	Hemiboreal forest zone: mixed-woodland –agricultural landscape	40	selected	L	3	GPM Sugita's lake	Poaceae	10	Poska et al. 2011
Finland	N-Finland	24	stratified random	M	3	GPM Prentice's bog	Poaceae	6	Räsänen et al. 2007
France	Mediterranean region	23	random	M	3	GPM Prentice's bog	Poaceae	11	Mazier et al. unpubl.
Germany	Eastern Germany: Brandenburg: agricultural landscape	49	selected	L	3	GPM Sugita's lake	<i>Pinus</i> Poaceae ^{*,§}	16	Matthias et al. 2012
	NE Germany: agricultural landscape	27	selected	L	3	LSM GPM Sugita's Lake [‡]	<i>Pinus</i> Poaceae ^{*,§}	11 (+5) [‡]	Theuerkauf et al. 2013
Norway	SW-Norway: Hordaland and Sogn og Fjordane, mown or grazed grass-land and heath	39	selected	M	1	None [#]	Poaceae	17	Hjelle 1998
Poland	NE Poland: Białowieża Forest	18	stratified random	M	3	GPM Prentice's bog	Poaceae	8	Baker et al. 2016
Romania	SE Romania: Forest-steppe region	26	random	M & S	3	GPM Prentice's bog	Poaceae	13	Grindean et al. 2019
Sweden	West-Central Sweden:	30	random	M	3	GPM Prentice's	Poaceae	10	von Stedingk et al. 2008

	Forest-tundra ecotone					bog			
	S-Sweden: ancient cultural landscapes	14	selected	M	3	None [#]	<i>Juniperus</i> Poaceae [‡]	14 (17) ³	Sugita et al. 1999
	S-Sweden: unfertilized mown or grazed grasslands	42	selected	M	3	GPM Prentice's bog	Poaceae	11	Broström et al. 2004
Switzerland	Lowland: agricultural landscape	20	selected	L	3	GPM Prentice's bog	Poaceae	13	Soepboer et al. 2007
	Jura Mountain: pasture woodlands	20	(stratified) random ^{△△}	M	1	GPM Prentice's bog	Poaceae	11	Mazier et al. 2008

969

970 [†]L=lakes; M=moss pollsters; S=surface soil

971 [‡]Other distance-weighting models were used in most studies, including the Gaussian Plume Model (GPM), 1/d, 1/d²

972 (d=distance) and the Lagrangian Stochastic Model (LSM). The GPM is used in both the model developed for bogs (Parsons

973 and Prentice, 1981; Prentice and Parsons, 1983) and lakes (Sugita, 1993). For this RPP synthesis, we chose the results from

974 the analyses using GPM rather than 1/d or 1/d². Note: In the study of Theuerkauf et al. (2013) the LSM was used. For this

975 synthesis, Theuerkauf recalculated his RPPs using the lake model developed by Sugita (1993).

976 ³Number of plant taxa for which RPP was estimated, including the reference taxon. Note: In the study by Theuerkauf et al.

977 (2013) RPPs were estimated for 17 taxa using LSM. The RPPs were recalculated using the lake model (Sugita, 1993) for 15

978 taxa (see note under ² above) for this synthesis. In the study of Sugita et al. (1999) RPPs were calculated for 14 trees and 3

979 herbs. We used only the values for the 14 trees in this synthesis, following the syntheses by Broström et al. (2008) and Mazier

980 et al. (2012).

981 [△]Britain: the study includes two areas (a and b) in which RPP estimates were calculated for different sets of taxa and the two

982 areas have different numbers of sites: a. Calthorpe (34), 5 taxa; b. Wheatfen (17), same 5 taxa and *Corylus* (6 taxa in total).

983 ^{△△} random distribution restricted to areas of the study region with existing vegetation maps (therefore no sites outside these

984 areas); i.e. study region including separate areas (Mazier et al., 2008).

985 ^{††}Vegetation data from historical maps around 1800 CE.

986 ^{†††}lake sediments dated to ca. 1800.

987 [‡]The reference taxon used in the original study is different from Poaceae. For this synthesis the RPPs were converted to values

988 relative to Poaceae.

989 ^{‡‡}The study of Bunting et al. (2005) does not include a RPP for Poaceae. In order to calculate the RPPs relative to Poaceae,

990 it was assumed that the RPP of *Quercus* was equal to the mean of RPPs from three other studies in Europe (see Mazier et al.,

991 2012 for details). Although we have included new RPP values for *Quercus* in this synthesis, we did not recalculate the RPPs


992 from Bunting et al. (2005) with a new mean value for *Quercus*, but used the same values as in Mazier et al. (2012). For

993 comparison, the mean value for *Quercus* using the RPPs of the additional studies included in this synthesis is 4.28 (instead of
994 5.83 in Mazier et al., 2012). This would imply slightly lower RPPs in Britain also for *Alnus*, *Betula*, *Corylus*, *Fraxinus* and
995 *Salix*.

996 # no distance weighting used for vegetation data because there was no information about vegetation with increasing distance
997 from the pollen sample (Hjelle et al., 1998; Sugita et al., 1999). In the Swedish study, vegetation data within a 10²-m² (herb
998 taxa) and 10³-m² quadrat (tree taxa) centred on the pollen sample was used (Sugita et al., 1999).

999 **Appendix D Maps of REVEALS cover for three plant taxa (*Calluna vulgaris*, deciduous *Quercus* deciduous and
1000 evergreen *Quercus* evergreen)**

1001



1002

1003

1004

1005


1006

1007

1008

Figure D1. Grid-based REVEALS estimates of *Calluna vulgaris* cover for eight Holocene time windows. Percentage cover in 2% interval between 0 and 2%, 3% interval between 2 and 5%, 5% intervals between 5 – 35% and 15% interval between 35 and 50%. **Intervals represented by increasingly darker shades of green from 5-10%.** Grey grid cells have no data (pollen) for *Calluna vulgaris* in the mapped time window. The circles represent the coefficient of variation (CV; the standard error divided by the REVEALS estimate). When $SE \geq REVEALS$ estimate, the circle fills the entire grid cell and the REVEALS estimate is not different from zero. This occurs mainly where REVEALS estimates are low.

1009




1011

1012

1013 **Figure D2.** Grid-based REVEALS estimates of *deciduous Quercus deciduous* cover in eight Holocene time windows.
 1014 Percentage cover in 1% interval between 0 and 2%, 3% interval between 2 and 5%, 5% intervals between 5 and 30%
 1015 and 20% interval between 30 and 50%. **Intervals represented by increasingly darker shades of green from 2-5%.** See
 1016 **caption of Figure A1 for more explanations.** Grey grid cells have no data (pollen) for *Calluna vulgaris* in the mapped
 1017 **time window.** The circles represent the coefficient of variation (CV; the standard error divided by the REVEALS
 1018 **estimate).** When $SE > REVEALS$ estimate, the circle fills the entire grid cell and the REVEALS estimate is not
 1019 **different from zero.** This occurs mainly where REVEALS estimates are low.

1020



1021
1022
1023
1024
1025
1026
1027
1028

Figure D3. Grid-based REVEALS estimates of evergreen *Quercus* evergreen cover for eight Holocene time windows. Percentage cover in 0.5% intervals between 0 and 1%, 1% intervals between 1 and 5%, 5% intervals between 5 and 15 and 15% interval between 15 and 30%. See caption of Figure A1 for more explanations. Intervals represented by increasingly darker shades of green from 1-2%. Grey grid cells have no data (pollen) for *Calluna vulgaris* in the mapped time window. The circles represent the coefficient of variation (CV; the standard error divided by the REVEALS estimate). When $SE \geq REVEALS$ estimate, the circle fills the entire grid cell and the REVEALS estimate is not different from zero. This occurs mainly where REVEALS estimates are low.

1029 **+Team list**

1030 Åkesson Christine (School of Geography & Sustainable Development, University of St. Andrews, UK), Balakauskas Lauras
1031 (Department of Geology and Mineralogy, Vilnius University, Vilnius, Lithuania), Batalova Vlada (Lomonosov Moscow State
1032 University, Department of Physical geography and Landscape science, Moscow, Russia), Birks H.J.B. (Department of
1033 Biological Sciences and Bjerknes Centre for Climate Research, University of Bergen, Norway), Bjune Anne. E. (Department
1034 of Biological Sciences and Bjerknes Centre for Climate Research, University of Bergen, Norway), Borisova Olga (Insitute of
1035 Geography, Russian Academy of Sciences, Moscow, Russia), Bozilova Elissaveta (Department of Botany, Sofia University
1036 St. Kliment Ohridski, Sofia, Bulgaria), Burjachs Francesc (ICREA Barcelona, Catalonia, Spain; Rovira i Virgili University
1037 (URV), Tarragona, Catalonia, Spain; Institut Català de Paleoecologia Humana i Evolució Social (IPHES), Campus Sescelades
1038 URV, W3, 43007 Tarragona, Spain), Cheddadi Rachid (Institut des Sciences de l'Evolution de Montpellier, Université de
1039 Montpellier, CNRS-UM-IRD, Montpellier, France), Christiansen Jörg (Department of Palynology and Climate Dynamics,
1040 Georg-August University, Göttingen, Germany), David Remi (Archeosciences Laboratory, UMR 6566 CReAAH, CNRS,
1041 Rennes I University, Rennes, France), de Klerk Pim (State Museum of Natural History, Karlsruhe, Germany), Di Rita Federico
1042 (Dipartimento di Biologia Ambientale, Università di Roma "La Sapienza", Piazzale Aldo Moro, 5, 00185, Roma, Italia),
1043 Dörfler Walter (Institute für Ur- und Frühgeschichte, Christian-Albrechts University, Kiel, Germany), Doyen Elise
1044 ([Paleobotanab, Bureau d'étude spécialisé en reconstitution des paléoenvironnements à partir de vestiges botaniques, 01300](#)
1045 [Nattages](#))[Laboratoire Chrono-Environnement, Franche-Comté University, Besançon, France](#)), Eastwood Warren (School of
1046 Geography, Earth and Environmental Sciences, University of Birmingham B15 2TT, UK), Etienne David (Savoie Mont Blanc
1047 University, Chambéry, France), Feeser Ingo (Institut für Ur- und Frühgeschichte, Christian-Albrechts University, Kiel,
1048 Germany), Filipova-Marinova Mariana (Museum of Natural History, Varna, Bulgaria), Fischer E. (Institute für Ur- und
1049 Frühgeschichte, Christian-Albrechts University, Kiel, Germany), Galop Didier (GEODE UMR 5602, Toulouse University,
1050 Toulouse, France), Garcia Jose Sebastian Carrion (Departamento de Biología Vegetal, Facultad de Biología, Universidad de
1051 Murcia, 30100 Murcia, Spain), [Gauthier Emilie \(Laboratoire Chrono-Environnement, UMR 6249 CNRS-Franche-Comté](#)
1052 [University, Besançon, France\)](#), [Giesecke Thomas \(Department of Physical Geography, Utrecht University, Utrecht, The](#)
1053 [Netherlands\)](#), Herking Christa (Institute of Botany and Landscape Ecology, EMAU, Greifswald, Germany), Herzsuh Ulrike
1054 (Alfred-Wegener-Institut Potsdam, Germany), Jouffroy-Bapicot Isabelle (Laboratoire Chrono-Environnement, [UMR 6249](#)
1055 [CNRS, Franche-Comté University, Besançon, France](#)), Kasianova Alisa (Department of Palynology and Climate Dynamics,
1056 Georg-August-University, Göttingen, Germany), Kouli Katerina (Department of Geology and Geoenvironment, National and
1057 Kapodistrian University of Athens, Panepistimioupolis, 15784 Ilissia, Greece), Kuneš Petr (Department of Botany, Charles
1058 University, Prague, Czech RepublicCzech), Lagerås Per (The Archaeologists, National Historical Museums, Lund, Sweden),
1059 Latałowa Małgorzata (Department of Plant Ecology, University of Gdansk, Poland), Lechterbeck Jutta (State Office for
1060 Cultural Heritage Baden-Wuerttemberg, Germany), Leroyer Chantal (Archeosciences Laboratory, UMR 6566 CReAAH,

Field Code Changed

Formatted: English (United Kingdom)

1061 CNRS, Rennes1 University, Rennes, France), Leydet Michelle (European Pollen Database, IMBE, Aix-Marseille Université,
1062 Avignon Université, IRD, Aix-en-Provence, France), Lisystina Olga ([Department of Geology, Tallinn University of](#)
1063 [Technology, 19086 Tallinn, Estonia](#))~~Department of Postglacial Geology, Tallinn University of Technology, Tallinn, Estonia~~,
1064 Lukanina Ekaterina (Department of Palynology and Climate Dynamics, Georg-August-University, Göttingen, Germany),
1065 Magyari Enikő (Department of Environmental and Landscape Geography, Eötvös Loránd University, Budapest, Hungary),
1066 Marguerie Dominique (Archeosciences Laboratory, UMR 6566 CReAAH, CNRS, Rennes1 University, Rennes, France),
1067 ~~Marta~~-Mariotti [Lippi Marta](#) (Dipartimento di Biologia, Università di Firenze, Via G. La Pira, 4, 50121 Firenze, Italy), Mensing
1068 Scott (Department of Geography, University of Nevada, Reno, NV 89557, USA), Mercuri Anna Maria (Laboratorio di
1069 Palinologia e Paleobotanica, Dipartimento di Scienze della Vita, Università di Modena e Reggio Emilia, Italy), Miebach
1070 Andrea (Steinmann Institute for Geology, Mineralogy, and Paleontology, University of Bonn, Bonn, Germany), [Milburn Paula](#)
1071 ([College of Science and Engineering, University of Edinburgh, Edinburgh, Scotland](#)), [Miras Yannick \(CNRS HNHP UMR](#)
1072 [7194, Museum National d'Histoire Naturelle, Paris, France](#)), [Morales del Molino César \(Alpine Pollen Database, Institute of](#)
1073 [Plant Sciences, Bern University, Switzerland\)](#), -Mrotzek Almut (Institute of Botany and Landscape Ecology, EMAU
1074 ,Greifswald, Germany), ~~Milburn Paula (College of Science and Engineering, University of Edinburgh, Edinburgh, Scotland)~~,
1075 Nosova Maria (Main Botanical Garden, Russian Academy of Sciences, Moscow, Russia), [Odgaard Bent Vad \(Department of](#)
1076 [Geoscience, Aarhus University, Denmark\)](#), Overballe-Petersen Mette (Forest & Landscape, Faculty of Life Sciences,
1077 University of Copenhagen, Frederiksberg, Denmark), Panajiotidis Sampson (Aristotle University of Thessaloniki, Department
1078 of Forestry and Natural Environment, PO Box: 270, GR54124 Thessaloniki, Greece), Pavlov Danail (Society of Innovative
1079 Ecologists of Bulgaria, Varna, Bulgaria), Persson⁺ Thomas[‡] (Department of Geology, Lund University, Lund, Sweden), Pinke
1080 Zsolt (Department of Physical Geography, Eötvös Loránd University, Budapest, Hungary), Ruffaldi Pascale (Laboratoire
1081 Chrono-Environnement, [UMR 6249 CNRS, Franche-Comté University, Besançon, France](#)), Sapelko Tatyana (Institute of
1082 Limnology, Russian Academy of Sciences, St. Petersburg, Russia), [Schmidt Monika \(Department of Palynology and Climate](#)
1083 [Dynamics, Georg-August-University, Göttingen, Germany\)](#), -Schult Manuela (Institute of Botany and Landscape Ecology,
1084 EMAU, Greifswald, Germany), ~~-Schmidt Monika (Department of Palynology and Climate Dynamics, Georg August~~
1085 ~~University, Göttingen, Germany)~~, ~~Stancikaite Migele (Institute of Geology and Geography, Nature Research Centre,~~
1086 ~~Akademijos Str. 2, LT-08412, Vilnius, Lithuania)~~~~Institute of Geology and Geography, Vilnius University, Vilnius, Lithuania~~,
1087 Stivrins Normunds (Department of Geography, Faculty of Geography and Earth Sciences, University of Latvia, Jelgavas iela
1088 1, Riga, 1004, Latvia), Tarasov Pavel E. (Institute of Geological Sciences, Free University of Berlin, Germany), ~~Theuerkauf~~
1089 ~~Martin (Institute of Botany and Landscape Ecology, EMAU Greifswald, 1748 Greiswald, Germany)~~-Tonkov Spassimir
1090 (~~Department of Botany, Sofia University St. Kliment Ohridski, Sofia, Bulgaria~~), Veski Siim (Department of Geology, Tallinn
1091 University of Technology, Tallinn, Estonia), Wick Lucia (IPNA, University of Basel, Basel, Switzerland), Wiethold Julian
1092 (INRAP, Direction interrégionale Grand-Est Nord, Laboratoire archéobotanique, Metz, France), Woldring Henk (Groningen
1093 Institute of Archaeology, University of Groningen, The Netherlands), Zernitskaya Valentina (Institute for Nature Management,
1094 National Academy of Sciences of Belarusk, Minsk, Republic of Belarus).

Formatted: Superscript

Formatted: Font: (Default) +Headings (Times New Roman)

Formatted: Not Superscript/ Subscript

1095 **Author Contribution**

1096 MJG coordinated the study as part of LandClim II and PAGES LandCover6k, two research projects for which she is the overall
1097 coordinator and administrator. MJG, AKT, EG, FM, RF, ABN, AP and SS conceptualised the study and methodology. SS
1098 developed the REVEALS model and helped with all issues related to the application of the model and interpretation of results.
1099 EG, AKT, RF, FM, ABN, and AP collected new pollen records from individual authors. JW provided part of the pollen records
1100 from the Mediterranean area (collected earlier for a separate project). MS and ST provided unpublished pollen records. EG
1101 and AKT had the major responsibility of handling the pollen data files and collecting all related metadata. AKT collected new
1102 values of relative pollen productivity estimates (RPPs) in Europe. MT provided unpublished RPP values for Germany and FM
1103 for the Mediterranean area. FM, JA, VL, LM, and NNC were all involved in the unpublished RPP study in southern France,
1104 and AF, RG, ABN and IT performed the RPP study in Romania. MJG performed the selection of RPP values for the new RPP
1105 synthesis used in this paper, EG made the calculations of mean RPPs, and MJG wrote Appendices A, B, and C, and prepared
1106 the Figures and Tables therein. RF performed the REVEALS model runs and created Figure 1 and the maps of REVEALS-
1107 based plant cover (Figures 2-6 and D1-D3). EG, RF and MJG designed the manuscript, EG prepared the first draft of the
1108 manuscript and all Tables, and the final manuscript for submission, RF and MJG wrote parts of the text and edited the full
1109 manuscript. All the co-authors were involved in commenting the manuscript.

1110 **Competing interests**

1111 The authors declare that they have no conflict of interest.

1112 **Figures entirely compiled by the manuscript authors:** Since such figures are part of the manuscript, they will receive the
1113 same distribution licence as the entire manuscript, namely a CC BY License. No citation is needed and no reproduction rights
1114 must be obtained.

1115 **Acknowledgements**

1116 This study was funded by a research project financed by the Swedish Research Council VR (Vetenskapsrådet) on
1117 “Quantification of the bio-geophysical and biogeochemical forcings from anthropogenic de-forestation on regional Holocene
1118 climate in Europe, LandClim II”. Financial support from the Linnaeus University’s Faculty of Health and Life Science is
1119 acknowledged for Marie-José Gaillard, Anna-Kari Trondman, and Esther Githubi. This is a contribution to the strategic
1120 research areas MERGE (Modelling the Regional and Global Earth system) and the Past Global Change (PAGES) project and
1121 its working group LandCover6k (<http://pastglobalchanges.org/landcover6k>), which in turn received support from the Swiss
1122 National Science Foundation, the Swiss Academy of Sciences, the US National Science Foundation, and the Chinese Academy
1123 of Sciences. Anneli Poska was supported by the ESF project number PRG323. We thank Sandy Harrison (University of
1124 Reading, UK) for providing the pollen records from the EMBSeCBIO project. [The work of the data contributors to - and the](#)
1125 [database managers of ALPADABA \(https://www.neotomadb.org/\), EMBSECBIO](#)
1126 [\(https://research.reading.ac.uk/palaeoclimate/embsecbio/\), EPD \(http://www.europeanpollendatabase.net/index.php\),](#)
1127 [LandClimI \(Trondman et al., 2015\), PALYCZ \(https://botany.natur.cuni.cz/palycz/\), and PALEOPYR \(http://paleopyr.univ-](#)
1128 [tlse2.fr/\) is gratefully acknowledged.](#)

1129 **References**

- 1130 Abraham, V. and Kozáková, R.: Relative pollen productivity estimates in the modern agricultural landscape of Central
1131 Bohemia (Czech Republic), *Rev. Palaeobot. Palynol.*, 179, 1–12, doi:10.1016/j.revpalbo.2012.04.004, 2012.
- 1132 Abraham, V., Oušková, V. and Kuneš, P.: Present-Day Vegetation Helps Quantifying Past Land Cover in Selected Regions of
1133 the Czech Republic, edited by B. Bond-Lamberty, *PLoS One*, 9(6), e100117, doi:10.1371/journal.pone.0100117, 2014.
- 1134 Andersen, S. T.: The relative pollen productivity and pollen representation of north European trees, and correction factors for
1135 tree pollen spectra, *Danmarks Geol. Undersogelse II*, 96, 99, 1970.
- 1136 Baker, A. G., Zimny, M., Keczyński, A., Bhagwat, S. A., Willis, K. J. and Latalowa, M.: Pollen productivity estimates from
1137 old-growth forest strongly differ from those obtained in cultural landscapes: Evidence from the Białowieża National Park,
1138 Poland, *The Holocene*, 26(1), 80–92, doi:10.1177/0959683615596822, 2016.
- 1139 Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., Getz, W. M., Harte, J., Hastings, A.,
1140 Marquet, P. A., Martinez, N. D., Mooers, A., Roopnarine, P., Vermeij, G., Williams, J. W., Gillespie, R., Kitzes, J., Marshall,
1141 C., Matzke, N., Mindell, D. P., Revilla, E. and Smith, A. B.: Approaching a state shift in Earth’s biosphere, *Nature*, 486(7401),
1142 52–58, doi:10.1038/nature11018, 2012.
- 1143 Beug, H. J.: Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete (Guide to pollen determination for
1144 Central Europe and neighboring areas), Verlag Dr. Friedrich Pfeil., 2004.
- 1145 Broström, A., Sugita, S. and Gaillard, M.-J.: Pollen productivity estimates for the reconstruction of past vegetation cover in
1146 the cultural landscape of southern Sweden, *The Holocene*, 14(3), 368–381, doi:10.1191/0959683604hl713rp, 2004.

1147 Broström, A., Nielsen, A. B., Gaillard, M.-J., Hjelle, K., Mazier, F., Binney, H., Bunting, J., Fyfe, R., Meltsov, V., Poska, A.,
1148 Räsänen, S., Soepboer, W., von Stedingk, H., Suutari, H. and Sugita, S.: Pollen productivity estimates of key European plant
1149 taxa for quantitative reconstruction of past vegetation: a review, *Veg. Hist. Archaeobot.*, 17(5), 461–478, doi:10.1007/s00334-
1150 008-0148-8, 2008.

1151 Bunting, M. J., Armitage, R., Binney, H. A. and Waller, M.: Estimates of 'relative pollen productivity' and 'relevant source
1152 area of pollen' for major tree taxa in two Norfolk (UK) woodlands, *The Holocene*, 15(3), 459–465,
1153 doi:10.1191/0959683605hl821rr, 2005.

1154 Bunting, M. J., Schofield, J. E. and Edwards, K. J.: Estimates of relative pollen productivity (RPP) for selected taxa from
1155 southern Greenland: A pragmatic solution, *Rev. Palaeobot. Palynol.*, 190, 66–74, doi:10.1016/j.revpalbo.2012.11.003, 2013a.

1156 Bunting, M. J., Farrell, M., Broström, A., Hjelle, K. L., Mazier, F., Middleton, R., Nielsen, A. B., Rushton, E., Shaw, H. and
1157 Twiddle, C. L.: Palynological perspectives on vegetation survey: a critical step for model-based reconstruction of Quaternary
1158 land cover, *Quat. Sci. Rev.*, 82, 41–55, doi:10.1016/j.quascirev.2013.10.006, 2013b.

1159 Commerford, J. L., McLauchlan, K. K. and Sugita, S.: Calibrating Vegetation Cover and Grassland Pollen Assemblages in the
1160 Flint Hills of Kansas, USA, *Am. J. Plant Sci.*, 04(07), 1–10, doi:10.4236/ajps.2013.47A1001, 2013.

1161 Cui, Q.-Y., Gaillard, M.-J., Lemdahl, G., Sugita, S., Greisman, A., Jacobson, G. L. and Olsson, F.: The role of tree composition
1162 in Holocene fire history of the hemiboreal and southern boreal zones of southern Sweden, as revealed by the application of the
1163 Landscape Reconstruction Algorithm: Implications for biodiversity and climate-change issues, *The Holocene*, 23(12), 1747–
1164 1763, doi:10.1177/0959683613505339, 2013.

1165 Davis, B. A. S., Collins, P. M. and Kaplan, J. O.: The age and post-glacial development of the modern European vegetation: a
1166 plant functional approach based on pollen data, *Veg. Hist. Archaeobot.*, 24(2), 303–317, doi:10.1007/s00334-014-0476-9,
1167 2015.

1168 Davis, M. B.: On the theory of pollen analysis, *Am. J. Sci.*, 261(10), 897–912, doi:10.2475/ajs.261.10.897, 1963.

1169 Dickson, C.: Distinguishing cereal from wild grass pollen: some limitations, *Circaea*, 5, 67–71, 1988.

1170 Downs, P. W. and Piégay, H.: Catchment-scale cumulative impact of human activities on river channels in the late
1171 Anthropocene: implications, limitations, prospect, *Geomorphology*, 338, 88–104, doi:10.1016/j.geomorph.2019.03.021, 2019.

1172 Edwards, K. J., Fyfe, R. and Jackson, S. T.: The first 100 years of pollen analysis, *Nat. Plants*, 3(2), 17001,
1173 doi:10.1038/nplants.2017.1, 2017.

1174 Ellis, E. C.: Ecology in an anthropogenic biosphere, *Ecol. Monogr.*, 85(3), 287–331, doi:10.1890/14-2274.1, 2015.

1175 Foley, J. A.: Global Consequences of Land Use, *Science (80-.)*, 309(5734), 570–574, doi:10.1126/science.1111772, 2005.

1176 Fyfe, R., de Beaulieu, J.-L., Binney, H., Bradshaw, R. H. W., Brewer, S., Le Flao, A., Finsinger, W., Gaillard, M.-J., Giesecke,
1177 T., Gil-Romera, G., Grimm, E. C., Huntley, B., Kunes, P., Kühl, N., Leydet, M., Lotter, A. F., Tarasov, P. E. and Tonkov, S.:
1178 The European Pollen Database: past efforts and current activities, *Veg. Hist. Archaeobot.*, 18(5), 417–424,
1179 doi:10.1007/s00334-009-0215-9, 2009.

1180 Fyfe, R., Roberts, N. and Woodbridge, J.: A pollen-based pseudobiomisation approach to anthropogenic land-cover change.,

1181 <http://dx.doi.org.proxy.lnu.se/10.1177/0959683610369509>, 20(7), 1165–1171, doi:10.1177/0959683610369509, 2010.

1182 Fyfe, R., Twiddle, C., Sugita, S., Gaillard, M. J., Barratt, P., Caseldine, C. J., Dodson, J., Edwards, K. J., Farrell, M., Froyd,

1183 C., Grant, M. J., Huckerby, E., Innes, J. B., Shaw, H. and Waller, M.: The Holocene vegetation cover of Britain and Ireland:

1184 Overcoming problems of scale and discerning patterns of openness, *Quat. Sci. Rev.*, 73, 132–148,

1185 doi:10.1016/j.quascirev.2013.05.014, 2013.

1186 Fyfe, R. M., Woodbridge, J. and Roberts, N.: From forest to farmland: pollen-inferred land cover change across Europe using

1187 the pseudobiomization approach, *Glob. Chang. Biol.*, 21(3), 1197–1212, doi:10.1111/gcb.12776, 2015.

1188 Fyfe, R. M., Woodbridge, J. and Roberts, C. N.: Trajectories of change in Mediterranean Holocene vegetation through

1189 classification of pollen data, *Veg. Hist. Archaeobot.*, 27(2), 351–364, doi:10.1007/s00334-017-0657-4, 2018.

1190 Fyfe, R. M., Githumbi, E., Trondmann, A.-K., Mazier, F., Nielsen, A. B., Poska, A., Sugita, S., Woodbridge, J., Contributors,

1191 L. and Gaillard, M.-J.: A full Holocene record of transient gridded vegetation cover in Europe, *Pangaea*,

1192 doi:<https://doi.pangaea.de/10.1594/PANGAEA.937075>, 2022.

1193 Gaillard, M.-J., Sugita, S., Bunting, M. J., Middleton, R., Broström, A., Caseldine, C., Giesecke, T., Hellman, S. E. V., Hicks,

1194 S., Hjelle, K., Langdon, C., Nielsen, A.-B., Poska, A., von Stedingk, H. and Veski, S.: The use of modelling and simulation

1195 approach in reconstructing past landscapes from fossil pollen data: a review and results from the POLLANDCAL network,

1196 *Veg. Hist. Archaeobot.*, 17(5), 419–443, doi:10.1007/s00334-008-0169-3, 2008.

1197 Gaillard, M.-J., Sugita, S., Mazier, F., Trondman, A.-K., Broström, A., Hickler, T., Kaplan, J. O., Kjellström, E., Kokfelt, U.,

1198 Kuneš, P., Lemmen, C., Miller, P., Olofsson, J., Poska, A., Rundgren, M., Smith, B., Strandberg, G., Fyfe, R., Nielsen, A. B.,

1199 Alenius, T., Balakauskas, L., Barnekow, L., Birks, H. J. B. B., Bjune, A., Björkman, L., Giesecke, T., Hjelle, K., Kalnina, L.,

1200 Kangur, M., van der Knaap, W. O., Koff, T., Lagerås, P., Latalowa, M., Leydet, M., Lechterbeck, J., Lindbladh, M., Odgaard,

1201 B., Peglar, S., Segerström, U., von Stedingk, H. and Seppä, H.: Holocene land-cover reconstructions for studies on land cover-

1202 climate feedbacks, *Clim. Past*, 6(4), 483–499, doi:10.5194/cp-6-483-2010, 2010.

1203 Gaillard, M.-J., Kleinen, T., Samuelsson, P., Nielsen, A. B., Bergh, J., Kaplan, J. O., Poska, A., Sandström, C., Strandberg,

1204 G., Trondman, A.-K. and Wramneby, A.: Second Assessment of Climate Change for the Baltic Sea Basin, edited by The

1205 BACC II Author Team, Springer International Publishing, Cham., 2015.

1206 Gaillard, M. J., Morrison, K. D., Madella, M. and Whitehouse, N.: Editorial: Past land-use and land-cover change: the

1207 challenge of quantification at the subcontinental to global scales, *Past Glob. Chang. Mag.*, 26(1), 3–3,

1208 doi:10.22498/pages.26.1.3, 2018.

1209 Giesecke, T., Davis, B., Brewer, S., Finsinger, W., Wolters, S., Blaauw, M., de Beaulieu, J.-L., Binney, H., Fyfe, R. M.,

1210 Gaillard, M.-J., Gil-Romera, G., van der Knaap, W. O., Kuneš, P., Kühl, N., van Leeuwen, J. F. N. N., Leydet, M., Lotter, A.

1211 F., Ortu, E., Semmler, M. and Bradshaw, R. H. W. W.: Towards mapping the late Quaternary vegetation change of Europe,

1212 *Veg. Hist. Archaeobot.*, 23(1), 75–86, doi:10.1007/s00334-012-0390-y, 2014.

1213 Githumbi, E., Fyfe, R., Kjellström, E., Lindström, J., Lu, Z., Mazier, F., Nielsen, A. B., Poska, A., Smith, B., Strandberg, G.,

1214 Sugita, S., Zhang, Q. and Gaillard, M.-J.: Holocene quantitative pollen-based vegetation reconstructions in Europe for climate

1215 modelling: LandClim II, in INQUA 2019: Life on the Edge, Dublin. [online] Available from:
1216 [https://portal.research.lu.se/portal/en/publications/holocene-quantitative-pollenbased-vegetation-reconstructions-in-europe-](https://portal.research.lu.se/portal/en/publications/holocene-quantitative-pollenbased-vegetation-reconstructions-in-europe-for-climate-modelling-landclim-ii(46cc8471-f51c-4117-a7c6-ccff00638e82)/export.html)
1217 [for-climate-modelling-landclim-ii\(46cc8471-f51c-4117-a7c6-ccff00638e82\)/export.html](https://portal.research.lu.se/portal/en/publications/holocene-quantitative-pollenbased-vegetation-reconstructions-in-europe-for-climate-modelling-landclim-ii(46cc8471-f51c-4117-a7c6-ccff00638e82)/export.html) (Accessed 9 August 2021), 2019.

1218 Gregory, P.: Spores: their properties and sedimentation in still air. Microbiology of the atmosphere. A plant science
1219 monograph, 1973.

1220 Grindean, R., Nielsen, A. B., Tanțău, I. and Feurdean, A.: Relative pollen productivity estimates in the forest steppe landscape
1221 of southeastern Romania, *Rev. Palaeobot. Palynol.*, 264, 54–63, doi:10.1016/j.revpalbo.2019.02.007, 2019.

1222 Guiry, E., Beglane, F., Szpak, P., Schulting, R., McCormick, F. and Richards, M. P.: Anthropogenic changes to the Holocene
1223 nitrogen cycle in Ireland, *Sci. Adv.*, 4(6), eaas9383, doi:10.1126/sciadv.aas9383, 2018.

1224 Harrison, S. P., Gaillard, M. J., Stocker, B. D., Vander Linden, M., Klein Goldewijk, K., Boles, O., Braconnot, P., Dawson,
1225 A., Fluet-Chouinard, E., Kaplan, J. O., Kastner, T., Pausata, F. S. R., Robinson, E., Whitehouse, N. J., Madella, M. and
1226 Morrison, K. D.: Development and testing scenarios for implementing land use and land cover changes during the Holocene
1227 in Earth system model experiments, *Geosci. Model Dev.*, 13(2), 805–824, doi:10.5194/gmd-13-805-2020, 2020.

1228 He, F., Vavrus, S. J., Kutzbach, J. E., Ruddiman, W. F., Kaplan, J. O. and Krumhardt, K. M.: Simulating global and local
1229 surface temperature changes due to Holocene anthropogenic land cover change, *Geophys. Res. Lett.*, 41(2), 623–631,
1230 doi:10.1002/2013GL058085, 2014.

1231 Hellman, S., Gaillard, M.-J., Broström, A. and Sugita, S.: The REVEALS model, a new tool to estimate past regional plant
1232 abundance from pollen data in large lakes: validation in southern Sweden, *J. Quat. Sci.*, 23(1), 21–42, doi:10.1002/jqs.1126,
1233 2008a.

1234 Hellman, S. E. V., Gaillard, M., Broström, A. and Sugita, S.: Effects of the sampling design and selection of parameter values
1235 on pollen-based quantitative reconstructions of regional vegetation: a case study in southern Sweden using the REVEALS
1236 model, *Veg. Hist. Archaeobot.*, 17(5), 445–459, doi:10.1007/s00334-008-0149-7, 2008b.

1237 Hibbard, K., Janetos, A., van Vuuren, D. P., Pongratz, J., Rose, S. K., Betts, R., Herold, M. and Feddema, J. J.: Research
1238 priorities in land use and land-cover change for the Earth system and integrated assessment modelling, *Int. J. Climatol.*, 30(13),
1239 2118–2128, doi:10.1002/joc.2150, 2010.

1240 Hjelle, K. L.: Herb pollen representation in surface moss samples from mown meadows and pastures in western Norway, *Veg.*
1241 *Hist. Archaeobot.*, 7(2), 79–96, doi:10.1007/BF01373926, 1998.

1242 Hofman-Kamińska, E., Bocherens, H., Drucker, D. G., Fyfe, R. M., Gumiński, W., Makowiecki, D., Pacher, M., Piličiauskienė,
1243 G., Samojlik, T., Woodbridge, J. and Kowalczyk, R.: Adapt or die—Response of large herbivores to environmental changes
1244 in Europe during the Holocene, *Glob. Chang. Biol.*, 25(9), 2915–2930, doi:10.1111/gcb.14733, 2019.

1245 Huntley, B.: European vegetation history: Palaeovegetation maps from pollen data - 13 000 yr BP to present, *J. Quat. Sci.*,
1246 5(2), 103–122, doi:10.1002/jqs.3390050203, 1990.

1247 Kaplan, J., Krumhardt, K., Gaillard, M.-J., Sugita, S., Trondman, A.-K., Fyfe, R., Marquer, L., Mazier, F. and Nielsen, A.:
1248 Constraining the Deforestation History of Europe: Evaluation of Historical Land Use Scenarios with Pollen-Based Land Cover

1249 Reconstructions, *Land*, 6(4), 91, doi:10.3390/land6040091, 2017.

1250 Kaplan, J. O., Krumhardt, K. M. and Zimmermann, N.: The prehistoric and preindustrial deforestation of Europe, *Quat. Sci.*
1251 *Rev.*, 28(27–28), 3016–3034, doi:10.1016/j.quascirev.2009.09.028, 2009.

1252 Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C. and Goldewijk, K. K.: Holocene carbon emissions
1253 as a result of anthropogenic land cover change, *The Holocene*, 21(5), 775–791, doi:10.1177/0959683610386983, 2011.

1254 Klein Goldewijk, K., Beusen, A., Doelman, J. and Stehfest, E.: Anthropogenic land use estimates for the Holocene – HYDE
1255 3.2, *Earth Syst. Sci. Data*, 9(2), 927–953, doi:10.5194/essd-9-927-2017, 2017.

1256 Kuneš, P., Abraham, V., Kovářik, O., Kopecký, M., Břizová, E., Dudová, L., Jankovská, V., Knipping, M., Kozšková, R.,
1257 Nováková, K., Petr, L., Pokorný, P., Roszková, A., Rybničková, E., Svobodová-Svitavská, H. and Wacnik, A.: Czech
1258 quaternary palynological Database - Palycz: review and basic statistics of the data, *Preslia*, 81(3), 209–238, 2009.

1259 Kuneš, P., Abraham, V., Werchan, B., Plesková, Z., Fajmon, K., Jamrichová, E. and Roleček, J.: Relative pollen productivity
1260 estimates for vegetation reconstruction in central-eastern Europe inferred at local and regional scales, *The Holocene*, 29(11),
1261 1708–1719, doi:10.1177/0959683619862026, 2019.

1262 Lerigoleur, E., Mazier, F., Jégou, L., Perret, M. and Galop, D.: PALEOPYR: un système d'information pour la gestion et
1263 l'exploitation de données palaeoenvironnementales sur le massif nord-pyrénéen. *Ingénierie des Systèmes d'Information* 3.
1264 [online] Available from: <http://paleopyr.univ-tlse2.fr/%0A>, 2015.

1265 Li, F., Gaillard, M.-J., Xu, Q., Bunting, M. J., Li, Y., Li, J., Mu, H., Lu, J., Zhang, P., Zhang, S., Cui, Q., Zhang, Y. and Shen,
1266 W.: A Review of Relative Pollen Productivity Estimates From Temperate China for Pollen-Based Quantitative Reconstruction
1267 of Past Plant Cover, *Front. Plant Sci.*, 9(September), doi:10.3389/fpls.2018.01214, 2018.

1268 Li, F., Gaillard, M.-J., Cao, X., Herzsuh, U., Sugita, S., Tarasov, P. E., Wagner, M., Xu, Q., Ni, J., Wang, W., Zhao, Y., An,
1269 C., Beusen, A. H. W., Chen, F., Feng, Z., Goldewijk, C. G. M. K., Huang, X., Li, Y., Li, Y., Liu, H., Sun, A., Yao, Y., Zheng,
1270 Z. and Jia, X.: Towards quantification of Holocene anthropogenic land-cover change in temperate China: A review in the light
1271 of pollen-based REVEALS reconstructions of regional plant cover, *Earth-Science Rev.*, 203(February), 103119,
1272 doi:10.1016/j.earscirev.2020.103119, 2020.

1273 Marinova, E., Harrison, S. P., Bragg, F., Connor, S., Laet, V., Leroy, S. A. G., Mudie, P., Atanassova, J., Bozilova, E., Caner,
1274 H., Cordova, C., Djamali, M., Filipova-Marinova, M., Gerasimenko, N., Jahns, S., Kouli, K., Kotthoff, U., Kvavadze, E.,
1275 Lazarova, M., Novenko, E., Ramezani, E., Röpke, A., Shumilovskikh, L., Tanțău, I. and Tonkov, S.: Pollen-derived biomes
1276 in the Eastern Mediterranean–Black Sea–Caspian–Corridor, *J. Biogeogr.*, 45(2), 484–499, doi:10.1111/jbi.13128, 2018.

1277 Marquer, L., Gaillard, M.-J., Sugita, S., Trondman, A.-K., Mazier, F., Nielsen, A. B., Fyfe, R., Odgaard, B. V., Alenius, T.,
1278 Birks, H. J. B., Bjune, A. E., Christiansen, J., Dodson, J., Edwards, K. J., Giesecke, T., Herzsuh, U., Kangur, M., Lorenz,
1279 S., Poska, A., Schult, M. and Seppä, H.: Holocene changes in vegetation composition in northern Europe: why quantitative
1280 pollen-based vegetation reconstructions matter, *Quat. Sci. Rev.*, 90, 199–216, doi:10.1016/j.quascirev.2014.02.013, 2014.

1281 Marquer, L., Gaillard, M.-J., Sugita, S., Poska, A., Trondman, A.-K., Mazier, F., Nielsen, A. B., Fyfe, R., Jönsson, A. M.,
1282 Smith, B., Kaplan, J. O., Alenius, T., Birks, H. J. B., Bjune, A. E., Christiansen, J., Dodson, J., Edwards, K. J., Giesecke,

1283 T., Herzsuh, U., Kangur, M., Koff, T., Latalowa, M., Lechterbeck, J., Olofsson, J. and Seppä, H.: Quantifying the effects of
1284 land use and climate on Holocene vegetation in Europe, *Quat. Sci. Rev.*, 171, 20–37, doi:10.1016/j.quascirev.2017.07.001,
1285 2017.

1286 Marquer, L., Mazier, F., Sugita, S., Galop, D., Houet, T., Faure, E., Gaillard, M.-J., Haunold, S., de Munnik, N., Simonneau,
1287 A., De Vleeschouwer, F. and Le Roux, G.: Pollen-based reconstruction of Holocene land-cover in mountain regions:
1288 Evaluation of the Landscape Reconstruction Algorithm in the Vicdessos valley, northern Pyrenees, France, *Quat. Sci. Rev.*,
1289 228, 106049, doi:10.1016/j.quascirev.2019.106049, 2020.

1290 Matthias, I., Nielsen, A. B. and Giesecke, T.: Evaluating the effect of flowering age and forest structure on pollen productivity
1291 estimates, *Veg. Hist. Archaeobot.*, 21(6), 471–484, doi:10.1007/s00334-012-0373-z, 2012.

1292 Mazier, F., Broström, A., Gaillard, M.-J., Sugita, S., Vittoz, P. and Buttler, A.: Pollen productivity estimates and relevant
1293 source area of pollen for selected plant taxa in a pasture woodland landscape of the Jura Mountains (Switzerland), *Veg. Hist.*
1294 *Archaeobot.*, 17(5), 479–495, doi:10.1007/s00334-008-0143-0, 2008.

1295 Mazier, F., Gaillard, M. J., Kunes, P., Sugita, S., Trondman, A.-K. and Brostrom, A.: Testing the effect of site selection and
1296 parameter setting on REVEALS-model estimates of plant abundance using th Czech Quaternary Palynological database
1297 Testing the effect of site selection and parameter setting on REVEALS-model estimates of plant abunda, *Rev. Palaeobot.*
1298 *Palynol.*, 187, 38–49 [online] Available from: <https://halshs.archives-ouvertes.fr/halshs-00959845>, 2012.

1299 Mazier, F., Broström, A., Bragée, P., Fredh, D., Stenberg, L., Thiere, G., Sugita, S. and Hammarlund, D.: Two hundred years
1300 of land-use change in the South Swedish Uplands: comparison of historical map-based estimates with a pollen-based
1301 reconstruction using the landscape reconstruction algorithm, *Veg. Hist. Archaeobot.*, 24(5), 555–570, doi:10.1007/s00334-
1302 015-0516-0, 2015.

1303 McLauchlan, K. K., Williams, J. J., Craine, J. M. and Jeffers, E. S.: Changes in global nitrogen cycling during the Holocene
1304 epoch, *Nature*, 495(7441), 352–355, doi:10.1038/nature11916, 2013.

1305 Morrison, K. D., Hammer, E., Boles, O., Madella, M., Whitehouse, N., Gaillard, M.-J., Bates, J., Vander Linden, M., Merlo,
1306 S., Yao, A., Popova, L., Hill, A. C., Antolin, F., Bauer, A., Biagetti, S., Bishop, R. R., Buckland, P., Cruz, P., Dreslerová, D.,
1307 Dusseldorp, G., Ellis, E., Filipovic, D., Foster, T., Hannaford, M. J., Harrison, S. P., Hazarika, M., Herold, H., Hilpert, J.,
1308 Kaplan, J. O., Kay, A., Klein Goldewijk, K., Kolář, J., Kyazike, E., Laabs, J., Lancelotti, C., Lane, P., Lawrence, D., Lewis,
1309 K., Lombardo, U., Lucarini, G., Arroyo-Kalin, M., Marchant, R., Mayle, F., McClatchie, M., McLeester, M., Mooney, S.,
1310 Moskal-del Hoyo, M., Navarrete, V., Ndiema, E., Góes Neves, E., Nowak, M., Out, W. A., Petrie, C., Phelps, L. N., Pinke, Z.,
1311 Rostain, S., Russell, T., Sluyter, A., Styring, A. K., Tamanaha, E., Thomas, E., Veerasamy, S., Welton, L. and Zanon, M.:
1312 Mapping past human land use using archaeological data: A new classification for global land use synthesis and data
1313 harmonization, edited by J. Freeman, *PLoS One*, 16(4), e0246662, doi:10.1371/journal.pone.0246662, 2021.

1314 Nielsen, A. B.: Modelling pollen sedimentation in Danish lakes at c.ad 1800: an attempt to validate the POLLSCAPE model,
1315 *J. Biogeogr.*, 31(10), 1693–1709, doi:10.1111/j.1365-2699.2004.01080.x, 2004.

1316 Nielsen, A. B. and Odgaard, B. V.: Quantitative landscape dynamics in Denmark through the last three millennia based on the

1317 Landscape Reconstruction Algorithm approach, *Veg. Hist. Archaeobot.*, 19(4), 375–387, doi:10.1007/s00334-010-0249-z,
1318 2010.

1319 Nielsen, A. B., Giesecke, T., Theuerkauf, M., Feeser, I., Behre, K.-E., Beug, H.-J., Chen, S.-H., Christiansen, J., Dörfler, W.,
1320 Endtmann, E., Jahns, S., de Klerk, P., Kühl, N., Latalowa, M., Odgaard, B. V., Rasmussen, P., Stockholm, J. R., Voigt, R.,
1321 Wiethold, J. and Wolters, S.: Quantitative reconstructions of changes in regional openness in north-central Europe reveal new
1322 insights into old questions, *Quat. Sci. Rev.*, 47, 131–149, doi:10.1016/j.quascirev.2012.05.011, 2012.

1323 Nosova, M. B., Novenko, E. Y., Severova, E. E. and Volkova, O. A.: Vegetation and climate changes within and around the
1324 Polistovo-Lovatskaya mire system (Pskov Oblast, north-western Russia) during the past 10,500 years, *Veg. Hist. Archaeobot.*,
1325 28(2), 123–140, doi:10.1007/s00334-018-0693-8, 2018.

1326 Palmisano, A., Woodbridge, J., Roberts, C. N., Bevan, A., Fyfe, R., Shennan, S., Cheddadi, R., Greenberg, R., Kaniewski, D.,
1327 Langgut, D., Leroy, S. A. G., Litt, T. and Miebach, A.: Holocene landscape dynamics and long-term population trends in the
1328 Levant, *The Holocene*, 29(5), 708–727, doi:10.1177/0959683619826642, 2019.

1329 Parsons, R. W. and Prentice, I. C.: Statistical approaches to R-values and the pollen—vegetation relationship, *Rev. Palaeobot.*
1330 *Palynol.*, 32(2–3), 127–152, doi:10.1016/0034-6667(81)90001-4, 1981.

1331 Pinhasi, R., Fort, J. and Ammerman, A. J.: Tracing the origin and spread of agriculture in Europe, *PLoS Biol.*, 3(12), 1–9,
1332 doi:10.1371/journal.pbio.0030410, 2005.

1333 Pirzamanbein, B., Lindström, J., Poska, A., Sugita, S., Trondman, A.-K., Fyfe, R., Mazier, F., Nielsen, A. B., Kaplan, J. O.,
1334 Bjune, A. E., Birks, H. J. B., Giesecke, T., Kangur, M., Latalowa, M., Marquer, L., Smith, B. and Gaillard, M.-J.: Creating
1335 spatially continuous maps of past land cover from point estimates: A new statistical approach applied to pollen data, *Ecol.*
1336 *Complex.*, 20, 127–141, doi:10.1016/j.ecocom.2014.09.005, 2014.

1337 Pirzamanbein, B., Lindström, J., Poska, A. and Gaillard, M. J.: Modelling Spatial Compositional Data: Reconstructions of past
1338 land cover and uncertainties, *Spat. Stat.*, 24, 14–31, doi:10.1016/j.spasta.2018.03.005, 2018.

1339 Pirzamanbein, B., Poska, A. and Lindström, J.: Bayesian Reconstruction of Past Land Cover From Pollen Data: Model
1340 Robustness and Sensitivity to Auxiliary Variables, *Earth Sp. Sci.*, 7(1), doi:10.1029/2018EA000547, 2020.

1341 Poska, A., Meltsov, V., Sugita, S. and Vassiljev, J.: Relative pollen productivity estimates of major anemophilous taxa and
1342 relevant source area of pollen in a cultural landscape of the hemi-boreal forest zone (Estonia), *Rev. Palaeobot. Palynol.*, 167(1–
1343 2), 30–39, doi:10.1016/j.revpalbo.2011.07.001, 2011.

1344 Prentice, C.: Records of vegetation in time and space: the principles of pollen analysis, in *Vegetation history*, pp. 17–42,
1345 Springer Netherlands, Dordrecht., 1988.

1346 Prentice, C., Guiot, J., Huntley, B., Jolly, D. and Cheddadi, R.: Reconstructing biomes from palaeoecological data: a general
1347 method and its application to European pollen data at 0 and 6 ka, *Clim. Dyn.*, 12(3), 185–194, doi:10.1007/BF00211617, 1996.

1348 Prentice, I. C.: Pollen Representation, Source Area, and Basin Size: Toward a Unified Theory of Pollen Analysis, *Quat. Res.*,
1349 23(1), 76–86, doi:10.1016/0033-5894(85)90073-0, 1985.

1350 Prentice, I. C. and Parsons, R. W. A.: Maximum Likelihood Linear Calibration of Pollen Spectra in Terms of Forest

1351 Composition, *Biometrics*, 39(4), 1051–1057, doi:10.2307/2531338, 1983.

1352 Prentice, I. C. and Webb III, T.: BIOME 6000: reconstructing global mid-Holocene vegetation patterns from palaeoecological
1353 records, *J. Biogeogr.*, 25(6), 997–1005, doi:10.1046/j.1365-2699.1998.00235.x, 1998.

1354 Räsänen, S., Suutari, H. and Nielsen, A. B.: A step further towards quantitative reconstruction of past vegetation in
1355 Fennoscandian boreal forests: Pollen productivity estimates for six dominant taxa, *Rev. Palaeobot. Palynol.*, 146(1–4), 208–
1356 220, doi:10.1016/j.revpalbo.2007.04.004, 2007.

1357 Roberts, C. N., Woodbridge, J., Palmisano, A., Bevan, A., Fyfe, R. and Shennan, S.: Mediterranean landscape change during
1358 the Holocene: Synthesis, comparison and regional trends in population, land cover and climate, *The Holocene*, 29(5), 923–
1359 937, doi:10.1177/0959683619826697, 2019.

1360 Roberts, N., Fyfe, R. M., Woodbridge, J., Gaillard, M.-J., Davis, B. A. S. S., Kaplan, J. O., Marquer, L., Mazier, F., Nielsen,
1361 A. B., Sugita, S., Trondman, A.-K. and Leydet, M.: Europe’s lost forests: a pollen-based synthesis for the last 11,000 years,
1362 *Sci. Rep.*, 8(1), 716, doi:10.1038/s41598-017-18646-7, 2018.

1363 Ruddiman, W. F.: The Anthropogenic Greenhouse Era Began Thousands of Years Ago, *Clim. Change*, 61(3), 261–293,
1364 doi:10.1023/B:CLIM.0000004577.17928.fa, 2003.

1365 Ruddiman, W. F., Fuller, D. Q., Kutzbach, J. E., Tzedakis, P. C., Kaplan, J. O., Ellis, E. C., Vavrus, S. J., Roberts, C. N., Fyfe,
1366 R., He, F., Lemmen, C. and Woodbridge, J.: Late Holocene climate: Natural or anthropogenic?, *Rev. Geophys.*, 54(1), 93–
1367 118, doi:10.1002/2015RG000503, 2016.

1368 Shennan, S.: *The First Farmers of Europe An Evolutionary Perspective*, Cambridge University Press, Cambridge. [online]
1369 Available from: [https://www-cambridge-org.proxy.lnu.se/se/academic/subjects/archaeology/archaeology-europe-and-near-
1370 and-middle-east/first-farmers-europe-evolutionary-perspective?format=HB&isbn=9781108422925](https://www-cambridge-org.proxy.lnu.se/se/academic/subjects/archaeology/archaeology-europe-and-near-and-middle-east/first-farmers-europe-evolutionary-perspective?format=HB&isbn=9781108422925), 2018.

1371 Soepboer, W., Sugita, S., Lotter, A. F., van Leeuwen, J. F. N. and van der Knaap, W. O.: Pollen productivity estimates for
1372 quantitative reconstruction of vegetation cover on the Swiss Plateau, *The Holocene*, 17(1), 65–77,
1373 doi:10.1177/0959683607073279, 2007.

1374 Soepboer, W., Sugita, S. and Lotter, A. F.: Regional vegetation-cover changes on the Swiss Plateau during the past two
1375 millennia: A pollen-based reconstruction using the REVEALS model, *Quat. Sci. Rev.*, 29(3–4), 472–483,
1376 doi:10.1016/j.quascirev.2009.09.027, 2010.

1377 von Stedingk, H., Fyfe, R. M. and Allard, A.: Pollen productivity estimates from the forest—tundra ecotone in west-central
1378 Sweden: implications for vegetation reconstruction at the limits of the boreal forest, *The Holocene*, 18(2), 323–332,
1379 doi:10.1177/0959683607086769, 2008.

1380 Stephens, L., Fuller, D., Boivin, N., Rick, T., Gauthier, N., Kay, A., Marwick, B., Armstrong, C. G., Barton, C. M., Denham,
1381 T., Douglass, K., Driver, J., Janz, L., Roberts, P., Rogers, J. D., Thakar, H., Altaaweel, M., Johnson, A. L., Sampietro Vattuone,
1382 M. M., Aldenderfer, M., Archila, S., Artioli, G., Bale, M. T., Beach, T., Borrell, F., Braje, T., Buckland, P. I., Jiménez Cano,
1383 N. G., Capriles, J. M., Diez Castillo, A., Çilingiroğlu, Ç., Negus Cleary, M., Conolly, J., Coutros, P. R., Covey, R. A.,
1384 Cremaschi, M., Crowther, A., Der, L., di Lernia, S., Doershuk, J. F., Doolittle, W. E., Edwards, K. J., Erlandson, J. M., Evans,

1385 D., Fairbairn, A., Faulkner, P., Feinman, G., Fernandes, R., Fitzpatrick, S. M., Fyfe, R., Garcea, E., Goldstein, S., Goodman,
1386 R. C., Dalpoim Guedes, J., Herrmann, J., Hiscock, P., Hommel, P., Horsburgh, K. A., Hritz, C., Ives, J. W., Junno, A., Kahn,
1387 J. G., Kaufman, B., Kearns, C., Kidder, T. R., Lanoë, F., Lawrence, D., Lee, G.-A., Levin, M. J., Lindsoug, H. B., López-
1388 Sáez, J. A., Macrae, S., Marchant, R., Marston, J. M., McClure, S., McCoy, M. D., Miller, A. V., Morrison, M., Motuzaite
1389 Matuzeviciute, G., Müller, J., Nayak, A., Noerwidi, S., Peres, T. M., Peterson, C. E., Proctor, L., Randall, A. R., Renette, S.,
1390 Robbins Schug, G., Ryzewski, K., Saini, R., Scheinsohn, V., Schmidt, P., Sebillaud, P., Seitsonen, O., Simpson, I. A.,
1391 Soltysiak, A., Speakman, R. J., Spengler, R. N., Steffen, M. L., et al.: Archaeological assessment reveals Earth's early
1392 transformation through land use, *Science* (80-.), 365(6456), 897–902, doi:10.1126/science.aax1192, 2019.
1393 Strandberg, G., Kjellström, E., Poska, A., Wagner, S., Gaillard, M.-J., Trondman, A.-K., Mauri, A., Davis, B. A. S. S., Kaplan,
1394 J. O., Birks, H. J. B. B., Bjune, A. E., Fyfe, R., Giesecke, T., Kalnina, L., Kangur, M., van der Knaap, W. O., Kokfelt, U.,
1395 Kuneš, P., Latašová, M., Marquer, L., Mazier, F., Nielsen, A. B., Smith, B., Seppä, H. and Sugita, S.: Regional climate
1396 model simulations for Europe at 6 and 0.2 k BP: sensitivity to changes in anthropogenic deforestation, *Clim. Past*, 10(2), 661–
1397 680, doi:10.5194/cp-10-661-2014, 2014.
1398 Stuart, A. and Ord, J. K.: *Kendall's advanced theory of statistics, Distrib. theory*, 1 [online] Available from:
1399 <https://ci.nii.ac.jp/naid/10004597057> (Accessed 2 July 2021), 1994.
1400 Sugita, S.: A Model of Pollen Source Area for an Entire Lake Surface, *Quat. Res.*, 39(2), 239–244,
1401 doi:10.1006/qres.1993.1027, 1993.
1402 Sugita, S.: Theory of quantitative reconstruction of vegetation I: pollen from large sites REVEALS regional vegetation
1403 composition, *The Holocene*, 17(2), 229–241, doi:10.1177/0959683607075837, 2007a.
1404 Sugita, S.: Theory of quantitative reconstruction of vegetation II: all you need is LOVE, *The Holocene*, 17(2), 243–257,
1405 doi:10.1177/0959683607075838, 2007b.
1406 Sugita, S., Gaillard, M. J. and Broström, A.: Landscape openness and pollen records: A simulation approach, *Holocene*, 9(4),
1407 409–421, doi:10.1191/095968399666429937, 1999.
1408 Sugita, S., Parshall, T., Calcote, R. and Walker, K.: Testing the Landscape Reconstruction Algorithm for spatially explicit
1409 reconstruction of vegetation in northern Michigan and Wisconsin, *Quat. Res.*, 74(2), 289–300,
1410 doi:10.1016/j.yqres.2010.07.008, 2010.
1411 Sun, A., Luo, Y., Wu, H., Chen, X., Li, Q., Yu, Y., Sun, X. and Guo, Z.: An updated biomization scheme and vegetation
1412 reconstruction based on a synthesis of modern and mid-Holocene pollen data in China, *Glob. Planet. Change*, 192(May 2019),
1413 103178, doi:10.1016/j.gloplacha.2020.103178, 2020.
1414 Sutton, O. : *Micrometeorology*, Q. J. R. Meteorol. Soc., 79(341), 457–457, doi:10.1002/qj.49707934125, 1953.
1415 Theuerkauf, M., Kuparinen, A. and Joosten, H.: Pollen productivity estimates strongly depend on assumed pollen dispersal,
1416 *The Holocene*, 23(1), 14–24, doi:10.1177/0959683612450194, 2012.
1417 Theuerkauf, M., Couwenberg, J., Kuparinen, A. and Liebscher, V.: A matter of dispersal: REVEALSinR introduces state-of-
1418 the-art dispersal models to quantitative vegetation reconstruction, *Veg. Hist. Archaeobot.*, 25(6), doi:10.1007/s00334-016-

1419 0572-0, 2016.

1420 Trondman, A.-K., Gaillard, M.-J., Sugita, S., Björkman, L., Greisman, A., Hultberg, T., Lagerås, P., Lindbladh, M. and Mazier,
1421 F.: Are pollen records from small sites appropriate for REVEALS model-based quantitative reconstructions of past regional
1422 vegetation? An empirical test in southern Sweden, *Veg. Hist. Archaeobot.*, 25(2), 131–151, doi:10.1007/s00334-015-0536-9,
1423 2016.

1424 Trondman, A. K., Gaillard, M. J., Mazier, F., Sugita, S., Fyfe, R., Nielsen, A. B., Twiddle, C., Barratt, P., Birks, H. J. B.,
1425 Bjune, A. E., Björkman, L., Broström, A., Caseldine, C., David, R., Dodson, J., Dörfler, W., Fischer, E., van Geel, B., Giesecke,
1426 T., Hultberg, T., Kalnina, L., Kangur, M., van der Knaap, P., Koff, T., Kuneš, P., Lagerås, P., Latalowa, M., Lechterbeck, J.,
1427 Leroyer, C., Leydet, M., Lindbladh, M., Marquer, L., Mitchell, F. J. G., Odgaard, B. V., Peglar, S. M., Persson, T., Poska, A.,
1428 Rösch, M., Seppä, H., Veski, S. and Wick, L.: Pollen-based quantitative reconstructions of Holocene regional vegetation cover
1429 (plant-functional types and land-cover types) in Europe suitable for climate modelling, *Glob. Chang. Biol.*, 21(2), 676–697,
1430 doi:10.1111/gcb.12737, 2015.

1431 Twiddle, C. L.: Pollen Analysis : Not Just a Qualitative Tool, *Geomorphol. Tech.*, 1–11, 2012.

1432 de Vareilles, A., Pelling, R., Woodbridge, J. and Fyfe, R.: Archaeology and agriculture: plants, people, and past land-use,
1433 *Trends Ecol. Evol.*, 36(10), 943–954, doi:10.1016/j.tree.2021.06.003, 2021.

1434 Wiczorek, M. and Herzschuh, U.: Compilation of relative pollen productivity (RPP) estimates and taxonomically harmonised
1435 RPP datasets for single continents and Northern Hemisphere extratropics, *Earth Syst. Sci. Data*, 12(4), 3515–3528,
1436 doi:10.5194/essd-12-3515-2020, 2020.

1437 Wolf, A., Callaghan, T. V. and Larson, K.: Future changes in vegetation and ecosystem function of the Barents Region, *Clim.*
1438 *Change*, 87(1–2), 51–73, doi:10.1007/s10584-007-9342-4, 2008.

1439 Woodbridge, J., Fyfe, R. M. and Roberts, N.: A comparison of remotely sensed and pollen-based approaches to mapping
1440 Europe’s land cover, edited by M. Bush, *J. Biogeogr.*, 41(11), 2080–2092, doi:10.1111/jbi.12353, 2014.

1441 Zanon, M., Davis, B. A. S. S., Marquer, L., Brewer, S. and Kaplan, J. O.: European Forest Cover During the Past 12,000
1442 Years: A Palynological Reconstruction Based on Modern Analogs and Remote Sensing, *Front. Plant Sci.*, 9(March), 1–25,
1443 doi:10.3389/fpls.2018.00253, 2018.

1444