

1 **A multi-dimensional dataset of Ordovician to Silurian graptolite**
2 **specimens for virtual examination, global correlation and shale gas**
3 **exploration**

4
5 Hong-He Xu ^{1*}, Zhi-Bin Niu ^{1,2*}, Yan-Sen Chen ¹, Xuan Ma ¹, Xiao-Jing Tong ¹,
6 Yi-Tong Sun ¹, Xiao-Yan Dong ¹, Dan-Ni Fan ¹, Shuang-Shuang Song ¹, Yan-
7 Yan Zhu ¹, Ning Yang ¹, Qing Xia ¹

8 ¹ State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of
9 Geology and Palaeontology and Center for Excellence in Life and
10 Paleoenvironment, Chinese Academy of Sciences, 210008 Nanjing, China

11 ² College of Intelligence and Computing, Tianjin University, 300354 Tianjin,
12 China

13 *The first authors.

14 **Correspondence:** Hong-He Xu (hwxu@nigpas.ac.cn), or Zhi-Bin Niu
15 (zniu@tju.edu.cn)

16
17 **Abstract**

18 Multi- elemental and -dimensional data are more and more important in
19 the development of data-driven research, as is the case in modern
20 palaeontology, in which examination by experts, or someday artificial
21 intelligence, every fossil specimen plays a fundamental role. We here release
22 a dataset of 1,550 graptolite specimens representing 113 Ordovician to
23 Silurian graptolite species or subspecies that are significant in global
24 stratigraphic correlation and shale gas exploration. The dataset contains
25 2,951 high-resolution images and a data table of each specimen's scientific
26 information, e.g., taxonomic, geologic, and geographic information, and
27 references. Our dataset provides images for specialists or laymen worldwide,
28 is supported by the tool, FSIDvis (Fossil Specimen Image Dataset Visualizer),
29 which we developed to facilitate the human-interactive exploration of the rich-
30 attribution image dataset, and also a nonlinear dimension reduction
31 technique, t-SNE (t-Distributed Stochastic Neighbor Embedding), to project
32 image data into two-dimensional space to visualize and explore similarities.
33 Our dataset potentially contributes to virtual examinations of specimens
34 (VES), global bio-stratigraphic correlation, and improvement of the shale gas
35 exploration efficiency. All data, images and the spreadsheet file, are

36 available from <https://doi.org/10.5281/zenodo.5205215> (Xu, 2022).

37

38 **1. Introduction**

39 Fossils are direct evidence of prehistoric life and are probably the most
40 important research object of palaeontology and stratigraphy, during which
41 fossils are collected, sampled, illustrated, described, curated, and deposited
42 as permanent specimens in museum or institution for further investigation
43 (Shute and Foster, 1999). Examinations to fossil specimens is a key and
44 indispensable part of descriptive palaeontology. Such, however, can be
45 partially achieved in a convenient and low-cost way, with the aid of multi-
46 dimensional fossil specimen dataset as in this study.

47 Graptolites are an extinct group of marine, colonial, organic-walled
48 hemichordates and have over 210 genera/3,000 species in worldwide fossil
49 records from the Cambrian to Carboniferous (c. 510~320 Ma) shales (Maletz,
50 2017). Graptolites extensively diversified in the Ordovician Period and
51 witnessed the second-largest mass extinction in geological life history, i.e., the
52 end-Ordovician mass extinction (Goldman et al., 2020). Graptolites evolved
53 quickly and spread globally in the Paleozoic (Fig. 1), and its species are
54 widely used as significant index fossils for determining rock ages and regional
55 bio-stratigraphic correlation. Bio-zones based on graptolite species dividing
56 the Ordovician and Silurian Periods are generally less than one million years
57 in duration; such a short geological interval makes possible a precise
58 understanding of life evolution in geological history (Chen et al., 2012; 2018).
59 Up to 102 Ordovician and Silurian graptolite species were selected as global
60 bio-zones for dating sediments and understanding the evolutionary pattern of
61 palaeobiology; and 13 global stratotype sections and points (GSSPs) are
62 defined by the first appearance datum (FAD) of graptolite species from the
63 Cambrian, Ordovician, and Silurian systems (Goldman et al., 2020). (Fig. 2).

64 Additionally, bio-zones or indication zones based on graptolite species
65 assist with identifying mining beds for shale gas exploration (Fig. 1).
66 Graptolitic shale yields a significant volume of shale gas and comprises more
67 than 9% global hydrocarbons rocks (Klemme and Ulmishek, 1991;
68 Podhalańska, 2013). In China, over 61.4% of natural gas is yielded from
69 Ordovician and Silurian graptolitic shale of southern China (Zou et al., 2019).
70 Identification of graptolite species helps to locate shale gas mining beds;

71 especially, 16 graptolite species were chosen as “gold callipers” to locate
72 favourable exploration beds (FEBs) of shale gas from China (Zou et al., 2015)
73 (Fig. 2).

74 In this paper, we describe a multi-dimensional and integrated dataset of
75 graptolite specimens. The dataset potentially contributes to a range of
76 scientific activities and provides 1) easy access to and virtual examination of
77 fossil specimens through high-resolution images and detailed scientific
78 information for teaching and training in paleontology and geologic survey; 2) a
79 standard fossil specimen image dataset for use in bio-stratigraphic correlation
80 and to improve exploration efficiency in the shale gas industry, and 3) a
81 potential aid of developing image-based automated classification.

82 83 **2. Materials and methods**

84 All images in our dataset were taken from graptolite specimens that are
85 preserved in shale and were collected from China. These specimens are
86 housed at the Nanjing Institute of Geology and Palaeontology (NIGP),
87 Chinese Academy of Sciences (CAS), with serial numbers and prefix NIGP.

88 We spent over two years to photograph every specimen using a single-
89 lens reflex camera Nikon D800E with Nikkor 60 mm macro-lens and a Leica
90 M125 or M205C microscope equipped with Leica cameras (Fig. 3). Every
91 image is well focused and shows the morphology of the graptolite. In total, we
92 took 40,597 images, including 20,644 camera photos (each with a resolution
93 of 4,912 × 7,360) and 19,953 microscope photos (each with a resolution of
94 2,720 × 2,048). Photos of low contrast or bad focus were removed from the
95 whole collection. We selected only photos that show the morphology of the
96 specimen and the diagnostic characters of each graptolite species that the
97 specimen represents (Fig. 4). We selected one or two images for each
98 specimen as the final dataset, uploaded to, and stored in our cloud server
99 (Fig. 3).

100 101 **3. Data description**

102 Our final dataset consists of 2,951 high-resolution images and a related
103 spreadsheet file. Every image is a high-resolution photo taken from a
104 collection of 1,550 graptolite specimens. These specimens were formally
105 published between 1958 and 2020. They belong to 113 graptolite species or

106 subspecies of 41 genera and 16 families of the Order Graptoloidea (see the
107 spreadsheet file, Fig 5). The geological age of these graptolite species ranges
108 from the Middle Ordovician (467.3 Ma) to the Telychian (433.4 Ma) Stage of
109 the Silurian Period (Fig. 5).

110 These graptolite species have relatively abundant fossil records and are
111 significant in regional and global bio-stratigraphic correlation. They are
112 commonly used in geological age determination and shale gas FEB
113 indication, including 32 graptolite bio-zones from the Darriwilian Stage of the
114 Ordovician Period (467.3 Ma) to the Telychian Stage of the Silurian Period
115 (433.4 Ma) and 16 “gold callipers” of shale gas FEBs for the case of 20 m to
116 80 m thick graptolite shale in China (Fig. 6). These species also include two
117 “golden spike” graptolite species for the two GSSPs in southern China (i.e.,
118 bases of the Darriwilian Stage in the Middle Ordovician System and the
119 Hirnantian Stage in the Upper Ordovician System)(Goldman et al., 2020;
120 Zhang et al., 2020).

121 The name of the individual image file is initialled by the specimen’s unique
122 number and taxonomical species name. Every specimen was photographed
123 with scale bar. The scale is attached to an image of the entire rock specimen.
124 The other image is a close-up of the fossil within the coloured loop drawn on
125 the whole specimen. Occasionally in the large images, the scale bar is
126 embedded and beside the fossil specimen. For example, in the file named
127 ‘9721Cardiograptus_amplus_S.jpg’, the genus name and species name are
128 connected by the underline symbol, avoiding the space symbol. ‘9721’ is the
129 specimen number, ‘Cardiograptus_amplus’ means the species name is
130 *Cardiograptus amplus* and ‘_S’ means it is a photo with scale bar. In all scale
131 bars, the minimum unit is one millimetre.

132 The image files are in JPG format. The single JPG file size ranges from
133 822 KB to 7.055 MB. The whole volume of the dataset is 10.4 GB. The quality
134 of specimen images in our dataset is much better than that in any previous
135 publications because most specimens were first studied many years ago and
136 their illustrations were in black and white, in low-resolution and/or printed on
137 paper publications only. Most of these specimens were illustrated only once,
138 or never clearly photographed. The image collection of our dataset provides
139 necessary complement for these specimens and, furthermore, unfolds their
140 scientific value to experts or anyone who is interested in fossils.

141 Every of specimen is tagged with scientific information, including genus
142 and species names, nominator, nomination year, specimen number, collection
143 number, locality (province, city, county), geological horizon and section,
144 collector name, collecting time, identifier, identifying time, related references,
145 and illustration labels. Specimens can be indexed and located in their detailed
146 housing drawers and cabinets using any of the above information. Their
147 detailed geologic information can also be obtained from the geological
148 section-based database, the Geobiodiversity Database (Xu et al., 2020) and
149 forms key elements of fossil specimen metadata (Xu et al., in press). All
150 related information is collected and recorded in a separate spreadsheet file
151 released with our image dataset (Xu et al., 2022).

152 Some specimens of our collection have a long research history, since
153 1958, and their taxonomical status might have changed in the light of
154 graptolite systematic studies (Maletz, 2017; Zhang et al., 2020). We invited
155 graptolite palaeontologists to curate every specimen to make sure that its
156 scientific information is updated and widely accepted. The spreadsheet file
157 includes following fields: species ID, Phylum, Class, Order, Suborder,
158 Infraorder, Family, Subfamily, Genus, Revised species name, tagged species
159 name, total number of specimens, specimen serial number, image file name,
160 microscope photo number, SLR photo number, Stage, age from, age to, mean
161 age value, locality, longitude, latitude, horizon, and first published reference. It
162 is noted that the 'Revised species name' of every specimen reflect the
163 emendation and correction study in Ma (2020), with help of graptolite expert
164 Prof Zhang Y-D (NIGP, CAS), which might need further study or peer-
165 reviewed. One can always search specimens according to tagged species
166 names.

167 Our dataset, with the image collection and comprehensive information of
168 a large batch of fossil specimens, supports virtual examination of specimens
169 in a convenient and low-cost way. Experts or laymen can look through,
170 examine, and even measure fossil specimens without need for
171 regional/international travel and formalities. Such greatly benefits
172 palaeontology in research, teaching, and science communication (Rahman et
173 al., 2012).

174

175 **4. Data visualization**

176 We have developed an interactive web exploration tool, FSIDvis (Fossil
177 Specimen Image Dataset Visualizer), to assist users to examine better the
178 scientific contents of our data (Fig. 7).

179 We further explore the distribution of these graptolite images and
180 visualize the feature embedding of our graptolite dataset (Fig. 8) using
181 different colors to denote different specimens. In detail, for each annotated
182 image, we first resized it into 448×448 pixels and fed it into the trained
183 Convolutional Neural Network (CNN) model. The output 1×1×2048 feature
184 map from the last average pooling layer is flattened and projected to a 113
185 (number of species) dimensional fully connected layer to represent an image
186 embedding. After that, we use t-SNE (t-Distributed Stochastic Neighbor
187 Embedding), a nonlinear dimension reduction technique for high-dimensional
188 data, to project the image embeddings into two-dimensional space for
189 visualization. Finally, we indicate the image data distribution by a scatter plot,
190 we use 15 colors to represent 15 families of the order Graptoloidea, covering
191 42 genera and 113 species. The distribution of the images in this figure is
192 based on species, showing a potential of automatic classifying graptolite
193 species using artificial intelligence (Niu and Xu, 2020).

194 195 **5. Conclusions**

196 A multi-dimensional, integrated dataset based on 1,550 pieces of
197 graptolite specimens is released. It contains 2,951 high-resolution images and
198 a spreadsheet file showing structured records of every specimen's scientific
199 information. During the preparation of the dataset, 113 Ordovician to Silurian
200 graptolite species or subspecies were selected for their significances in
201 stratigraphic correlation and shale gas exploration, and these specimens were
202 carefully photographed and taxonomically curated.

203 Our dataset provides experts or laymen with a mean of virtual
204 examination of a batch of fossil specimens in a convenient and low-cost way.
205 It potentially contributes to global bio-stratigraphic correlation, especially with
206 those bio-zone graptolite species, and in the shale gas industry to
207 improvement of exploration efficiency. A fossil specimen database needs to
208 fulfil the purpose and requirement of virtual examination of specimens. This
209 greatly benefits palaeontologic research and science communication. The
210 whole dataset is visualized by the tool FSIDvis (Fossil Specimen Image Data

211 Visualizer) and a nonlinear dimension reduction technique, t-SNE (t-
212 Distributed Stochastic Neighbor Embedding).

213

214 **Data availability.** The dataset is archived and publicly available from
215 <https://doi.org/10.5281/zenodo.5205215>. Visualized version is available at
216 <http://fsidvis.fossil-ontology.com:8089/>

217

218 **Author contributions.** H.-H.X. and Z.-B.N. equally designed the project,
219 developed the model, and performed the simulations. H.-H.X. prepared and
220 revised the manuscript. Y.-S.C. gave technical support. X.M. curated fossil
221 specimens. Others contributed to specimen photography.

222

223 **Competing interests.** The authors declare that they have no conflict of
224 interest.

225

226 **Acknowledgments.** We thank Prof. Zhang Yuandong and Dr. Chen Qing,
227 NIGP, CAS, for careful curating and examining of the graptolite specimens;
228 Prof. Peter M Sadler, University of California (USA), for comment and
229 improving the manuscript; Dr. Pan Zhaohui, Institute of Vertebrate
230 Palaeontology and Paleoanthropology, CAS; Mr. Pan Yaohua and Mr. Wu
231 Junqi, College of Intelligence and Computing, Tianjin University, for
232 constructive suggestions and help.

233

234 **Financial support.** This research has been supported by the Strategic
235 Priority Research Program of the Chinese Academy of Sciences (Grants
236 XDA19050101 and XDB26000000) and National Natural Science Foundation
237 of China (61802278).

238

239 **References**

240 Chen, X., Chen, Q., Zhen, Y., Wang, H., Zhang, L., Zhang, J. and Xiao, Z.:
241 Circumjacent distribution pattern of the Lungmachiean graptolitic black
242 shale (early Silurian) on the Yichang Uplift and its peripheral region.
243 Science China Earth Sciences, 61, 1195–1203, 2018.

244 Chen, X., Zhang, Y., Li, Y., Fan, J., Tang, P., Chen, Q. and Zhang, Y.:
245 Biostratigraphic correlation of the Ordovician black shales in Tarim Basin

246 and its peripheral regions. *Science China Earth Sciences*, 55, 1230–1237,
247 2012.

248 Goldman, D., Sadler, P.M. and Leslie, S.A.: The Ordovician Period, in
249 *Geologic Time Scale 2020*. Elsevier. p. 631–694, 2020.

250 Klemme, H.D. and Ulmishek, G.F.: Effective petroleum source rocks of the
251 world: stratigraphic distribution and controlling depositional factors. *AAPG*
252 *Bulletin*, 75, 1809–1851. 1991.

253 Ma, X.: Palaeontology, biostratigraphy and palaeoecology of the graptolite
254 from the Hulo Formation (Darriwilian – Sandbian) in northwestern
255 Zhejiang Province, East China. A Ph.D dissertation submitted to University
256 of Chinese Academy of Sciences (supervised by Prof. Zhang Y-D). 1-301.
257 2020. Maletz, J.: Part V, Second Revision, Chapter 13: The history of
258 graptolite classification. *Treatise Online*, 88:1–11, 2017.

259 Niu, Z.-B. and Xu, H.-H.: AI-based graptolite identification improve shale gas
260 exploration. *bioRxiv*. doi: <https://doi.org/10.1101/2022.01.17.476477>

261 Peters, S. E. and McClennen, M.: The Paleobiology Database application
262 programming interface. *Paleobiology*, 42, 1–7, 2016.

263 Podhalańska, T.: Graptolites–stratigraphic tool in the exploration of zones
264 prospective for the occurrence of unconventional hydrocarbon deposits.
265 *Przegląd Geologiczny*, 61, 621–629, 2013.

266 Rahman, I.A., Adcock, K. and Garwood, R.J.: Virtual fossils: a new resource
267 for science communication in paleontology. *Evolution: Education and*
268 *Outreach*. 5, 635–641, 2012.

269 Shute, C.H. and Foster, T.S.: Curation in museum collections. In: Jones, T.P.,
270 Rowe, N.P., eds, *Fossil plants and spores: modern techniques*. Geological
271 Society of London. 184–186, 1999.

272 Xu H.H, Nie T., Guo W., Chen Y.-S. and Yuan W.-W.: Palaeontological fossil
273 specimen metadata standard. *Acta Palaeotologica Sinica*, 61, 280–290.
274 DOI: 10.19800/j.cnki.aps.2022007. 2022.

275 Xu, H.-H., Niu, Z.-B. and Chen, Y.-S.: A status report on a section-based
276 stratigraphic and palaeontological database–the Geobiodiversity
277 Database. *Earth System Science Data*, 12, 3443–3452, 2020.


278 Xu, H.-H.: High-resolution images of 1550 Ordovician to Silurian graptolite
279 specimens for global correlation and shale gas exploration.
280 <https://doi.org/10.5281/zenodo.5205215>. 2022.

281 Zhang, Y.D. Zhan, R.B., Wang, Z.H., Yuan, W., Fang., Liang, Y.,Yan, Wang, Y.,
282 Liang, K. et al.: 2020. Illustrations of index fossils from the Ordovician
283 strata in China. Zhejiang University Press. 1–575, 2020.

284 Zou, C.N., Dong, D., Wang, Y., Li, J., Huang., Wang, S., Guan, Q. et al.: Shale
285 gas in China: Characteristics, challenges and prospects (I). Petroleum
286 Exploration and Development. 42, 689–701, 2015.

287 Zou, C.N., Gong, J., Wang, H.Y. and Shi, Z.: Importance of graptolite evolution
288 and biostratigraphic calibration on shale gas exploration. China Petroleum
289 Exploration. 24, 1–6, 2019.

290



291

292

293

294


295

296

297

298

Figure 1. Global distribution of graptolite shale and shale gas production region. Most graptolite fossils were yielded from these shale sediments and their distribution is based on their occurrence records in global Ordovician and Silurian sediments. All data are from Peters and McClennen (2016) and Xu et al. (2020). The map is from © OpenStreetMap contributors 2021. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.



299

300 **Figure 2.** Graptolite species of our dataset are significant to biostratigraphy
 301 and dating of Ordovician and Silurian sediments. These graptolites also
 302 witnessed several macro-evolution events, including the great Ordovician
 303 biodiversity event (GOBE), Late Ordovician mass extinction (LOME).
 304 Radiation of several graptolite groups (bold verticle lines) occurs in this
 305 geological time. Two global stratotype sections and points (GSSPs), based on
 306 graptolite species record, are in southern China (the spike marks in left figure)
 307 (data from Goldman et al., 2020). Bio- or indication zones based on graptolite
 308 species assist with identifying mining beds for shale gas exploration in
 309 southern China. 16 graptolite indicator-zones are used in the shale gas
 310 exploration in China (Zou et al., 2015) (right part in the figure).
 311




312

313 **Figure 3.** The process of creating the graptolite specimen image dataset.


314 The graptolite specimens were carefully curated and revised to select the
 315 species with biostratigraphy and application significances. Every image was
 316 obtained from specimens that were macro-photographed using a single-lens
 317 reflex camera and microscope. After professional revision and cleaning, the
 318 whole dataset was uploaded to and stored in our cloud server.

319



320
 321
 322
 323
 324
 325
 326

Figure 4. Typical images of graptolite specimens in our dataset. Every image was taken from a unique graptolite specimen. Our dataset only selected the photos that well show morphology of every specimen and diagnostic character of each graptolite species that the specimens represent. The scientific species name of every specimen is given on each image.



327


328 **Figure 5.** Geographic distribution (A) and geologic range (B) of graptolite
 329 species of our dataset. Each graptolite specimen locality is represented by a
 330 pie chart where each colour is encoded as one graptolite family of the Order
 331 Graptoloidea. The sector size is proportional to the specimen number for
 332 every family. The radius of the pie chart is proportional to the total number of
 333 specimens from the same locality. The dashed-lines circle the main areas of
 334 shale gas production. The map is from © OpenStreetMap contributors 2021.
 335 Distributed under the Open Data Commons Open Database License (ODbL)
 336 v1.0.

337

System	Series	Stage	Graptolite biozone (22)	Stage	Graptolite indicator zone for shale gas FEB (16)
Silurian	Wenlock	Homerian	<i>Colonograptus deubeli</i>	Telychian	<i>Spirograptus turriculatus</i> (N2)
		Sheinwoodian	<i>Colonograptus praedeubeli</i>		<i>Spirograptus guerichi</i> (N1)
		Telychian	<i>Spirograptus turriculatus</i>		<i>Stimulograptus sedgwickii</i> (LM8)
	Llandovery	Aeronian	<i>Lituigraptus convolutus</i>	Aeronian	<i>Lituigraptus convolutus</i> (LM7)
			<i>Demirastrites triangulatus</i>		<i>Demirastrites triangulatus</i> (LM6)
		Rhuddanian		<i>Coronograptus cyphus</i>	<i>Coronograptus cyphus</i> (LM5)
				<i>Cystograptus vesiculosus</i>	<i>Cystograptus vesiculosus</i> (LM4)
				<i>Parakidograptus acuminatus</i>	<i>Parakidograptus acuminatus</i> (LM3)
				<i>Akidograptus ascensus</i>	<i>Akidograptus ascensus</i> (LM2)
				<i>Metabolograptus persculptus</i>	<i>Metabolograptus persculptus</i> (LM1)
Ordovician	Upper	Hirnantian	<i>Metabolograptus persculptus</i>	Hirnantian	<i>Metabolograptus extraordinarius</i> (WF4)
			<i>Metabolograptus extraordinarius</i>		<i>Dicellograptus mirus</i> (WF3c)
		Katian		<i>Paraorthograptus pacificus</i>	<i>Tangyagraptus typicus</i> (WF3b)
				<i>Dicellograptus complexus</i>	<i>Paraorthograptus pacificus</i> (WF3a)
				<i>Dicellograptus ornatus</i>	<i>Dicellograptus complexus</i> (WF2)
	Middle	Sandbian	<i>Dicellograptus complanatus</i>	Katian	<i>Dicellograptus complanatus</i> (WF1)
		Darrivilian			<i>Orthograptus calcaratus</i>
					<i>Hustedograptus teretiusculus</i>
					<i>Archiclimacograptus riddellensis</i>
					<i>Pterograptus elegans</i>
Dapingian	<i>Nicholsonograptus fasciculatus</i>				
	<i>Levisograptus dentatus</i>				
	<i>Levisograptus austrodentatus</i>				


338

339 **Figure 6.** Graptolite species selected as global bio-zone (left) and indicator
340 zone (right) for shale gas favourable exploration beds (FEBs) of our dataset.
341 Among our dataset of 113 graptolite species, there are 22 graptolite index
342 species from global correlation from the Middle Ordovician to (470.0 Ma) to
343 the Wenlock of the Silurian Period (427.4 Ma), and 16 graptolite species as
344 ‘gold callipers’ to locate FEBs of shale gas in China. Note that some graptolite
345 species are duplicate in the two lists.
346



347

348 **Figure 7.** FSIDvis (Fossil Specimen Image Dataset Visualizer) system
 349 interface. a) Fossil on geographic distribution view, showing fossil specimen
 350 location on the map. The lens (a.1) is a tailor-designed specimens' picker that
 351 facilitates users to collect interest fossils of a region where the inner ring and
 352 outer ring represent the family and genus. When the user chooses a genus,
 353 the corresponding detailed species with images will be listed in the fossil list
 354 view (a.2), where the detailed information and further high-resolution image if
 355 the specimens are given. Hit the space bar for locking the selection. b)
 356 Geological age scale view, providing the geologic age selection ability; the top
 357 one is the chronostratigraphic age scale, and the bottom one is an age slider
 358 that facilitates the users to choose a specific age slot interactively. The web
 359 exploration tool of graptolite is provided at [http://fsidvis.fossil-](http://fsidvis.fossil-ontology.com:8089/)
 360 [ontology.com:8089/](http://fsidvis.fossil-ontology.com:8089/). The map is from © OpenStreetMap contributors 2021.
 361 Distributed under the Open Data Commons Open Database License (ODbL)
 362 v1.0.
 363



364

365 **Figure 8.** t-SNE embeded visualization of our graptolite specimen images.

366 Individual specimens are denoted and grouped by different colors. These

367 groups match different graptolite families (blocks with several small images).