

1           **A multi-dimensional dataset of Ordovician to Silurian graptolite**  
2           **specimens for virtual examination, global correlation and shale gas**  
3           **exploration**

4  
5   Hong-He Xu <sup>1\*</sup>, Zhi-Bin Niu <sup>1,2\*</sup>, Yan-Sen Chen <sup>1</sup>, Xuan Ma <sup>1</sup>, Xiao-Jing Tong <sup>1</sup>,  
6   Yi-Tong Sun <sup>1</sup>, Xiao-Yan Dong <sup>1</sup>, Dan-Ni Fan <sup>1</sup>, Shuang-Shuang Song <sup>1</sup>, Yan-  
7           Yan Zhu <sup>1</sup>, Ning Yang <sup>1</sup>, Qing Xia <sup>1</sup>

8   <sup>1</sup> State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of  
9   Geology and Palaeontology and Center for Excellence in Life and  
10   Paleoenvironment, Chinese Academy of Sciences, 210008 Nanjing, China

11   <sup>2</sup> College of Intelligence and Computing, Tianjin University, 300354 Tianjin,  
12   China

13   \*The first authors.

14   **Correspondence:** Hong-He Xu ([hwxu@nigpas.ac.cn](mailto:hwxu@nigpas.ac.cn)), or Zhi-Bin Niu  
15   ([zniu@tju.edu.cn](mailto:zniu@tju.edu.cn))

16  
17   **Abstract**

18       Multi- elemental and -dimensional data are more and more important in  
19   the development of data-driven research, as is the case in modern  
20   palaeontology, in which examination by experts, or someday artificial  
21   intelligence, every fossil specimen plays a fundamental role. We here release  
22   a dataset of 1,550 graptolite specimens representing 113 Ordovician to  
23   Silurian graptolite species or subspecies that are significant in global  
24   stratigraphic correlation and shale gas exploration. The dataset contains  
25   2,951 high-resolution images and a data table of each specimen's scientific  
26   information, e.g., taxonomic, geologic, and geographic information, comment,  
27   and references. Our dataset provides images for specialists or laymen  
28   worldwide, is supported by the tool, FSIDvis (Fossil Specimen Image Dataset  
29   Visualizer), which we developed to facilitate the human-interactive exploration  
30   of the rich-attribution image dataset, and also a nonlinear dimension reduction  
31   technique, t-SNE (t-Distributed Stochastic Neighbor Embedding), to project  
32   image data into two-dimensional space to visualize and explore similarities.  
33   Our dataset potentially contributes to virtual examinations of specimens  
34   (VES), global bio-stratigraphic correlation, and improvement of the shale gas  
35   exploration efficiency. All data, images and the spreadsheet file, are available

36 from <https://doi.org/10.5281/zenodo.5205215> (Xu, 2022).

37

38

## 39 **1. Introduction**

40 Fossils are direct evidence of prehistoric life and are probably the most  
41 important research object of palaeontology and stratigraphy, during which  
42 fossils are collected, sampled, illustrated, described, curated, and deposited  
43 as permanent specimens in museum or institution for further investigations  
44 (Shute and Foster, 1999). Examinations of fossil specimens are a key and  
45 indispensable part of descriptive palaeontology. Such, however, can be  
46 partially achieved in a convenient and low-cost way, with the aid of multi-  
47 dimensional fossil specimen dataset as in this study.

48 Graptolites are an extinct group of marine, colonial, organic-walled  
49 hemichordates and have over 210 genera/3,000 species in worldwide fossil  
50 records from the Cambrian to Carboniferous (c. 510~320 Ma) shales (Maletz,  
51 2017). Graptolites extensively diversified in the Ordovician Period and  
52 witnessed the second-largest mass extinction in geological life history, i.e., the  
53 end-Ordovician mass extinction (Goldman et al., 2020). Graptolites evolved  
54 quickly and spread globally in the Palaeozoic (Fig. 1), and its species are  
55 widely used as significant index fossils for determining rock ages and regional  
56 bio-stratigraphic correlation. Bio-zones based on graptolite species dividing  
57 the Ordovician and Silurian Periods are generally less than one million years  
58 in duration; such a short geological interval makes possible a precise  
59 understanding of life evolution in geological history (Chen et al., 2012; 2018).  
60 Up to 102 Ordovician and Silurian graptolite species were selected as global  
61 bio-zones for dating sediments and understanding the evolutionary pattern of  
62 palaeobiology; and 13 global stratotype sections and points (GSSPs) are  
63 defined by the first appearance datum (FAD) of graptolite species from the  
64 Cambrian, Ordovician, and Silurian systems (Goldman et al., 2020) (Fig. 2).

65 Additionally, bio-zones or indication zones based on graptolite species  
66 assist with identifying mining beds for shale gas exploration (Fig. 1).  
67 Graptolitic shale yields a significant volume of shale gas and comprises more  
68 than 9% global hydrocarbons rocks (Klemme and Ulmishek, 1991;  
69 Podhalańska, 2013). In China, over 61.4% of natural gas is yielded from  
70 Ordovician and Silurian graptolitic shale of southern China (Zou et al., 2019).

71 Identification of graptolite species helps to locate shale gas mining beds;  
72 especially, 16 graptolite species were chosen as “gold callipers” to locate  
73 favourable exploration beds (FEBs) of shale gas from China (Zou et al., 2015)  
74 (Fig. 2).

75 In this paper, we describe a multi-dimensional and integrated dataset of  
76 graptolite specimens. The dataset potentially contributes to a range of  
77 scientific activities and provides 1) easy access to and virtual examination of  
78 fossil specimens through high-resolution images and detailed scientific  
79 information for teaching and training in paleontology and geologic survey; 2) a  
80 standard fossil specimen image dataset for use in bio-stratigraphic  
81 correlations and to improve exploration efficiency in the shale gas industry,  
82 and 3) a potential aid of developing image-based automated classification.

83

## 84 **2. Materials and methods**

85 All images in our dataset were taken from graptolite specimens that are  
86 preserved in shale and were collected from China. These specimens are  
87 housed at the Nanjing Institute of Geology and Palaeontology (NIGP),  
88 Chinese Academy of Sciences (CAS), with serial numbers and the prefix  
89 NIGP.

90 We spent over two years to photograph every specimen using a single-  
91 lens reflex camera Nikon D800E with Nikkor 60 mm macro-lens and a Leica  
92 M125 or M205C microscope equipped with Leica cameras (Fig. 3). Every  
93 image is well focused and shows the morphology of the graptolite. In total, we  
94 took 40,597 images, including 20,644 camera photos (each with a resolution  
95 of 4,912 × 7,360) and 19,953 microscope photos (each with a resolution of  
96 2,720 × 2,048). Photos of low contrast or bad focus were removed from the  
97 whole collection. We only selected photos that show the morphology of the  
98 specimen and the diagnostic characters of each graptolite species that the  
99 specimen represents (Fig. 4). We selected one or two images for each  
100 specimen as the final dataset, uploaded to, and stored in our cloud server  
101 (Fig. 3).

102

## 103 **3. Data description**

104 Our final dataset consists of 2,951 high-resolution images and a related  
105 spreadsheet file. Every image is a high-resolution photo taken from a

106 collection of 1,550 graptolite specimens. These specimens were formally  
107 published between 1958 and 2020. They belong to 113 graptolite species or  
108 subspecies of 41 genera and 16 families of the Order Graptoloidea (see the  
109 spreadsheet file, Fig 5). The geological age of these graptolite species ranges  
110 from the Middle Ordovician (467.3 Ma) to the Telychian Stage of the Silurian  
111 Period (433.4 Ma) (Fig. 5).

112 These graptolite species have relatively abundant fossil records and are  
113 significant in regional and global bio-stratigraphic correlations. They are  
114 commonly used in geological age determination and shale gas FEB  
115 indication, including 32 graptolite bio-zones from the Darriwilian Stage of the  
116 Ordovician Period (467.3 Ma) to the Telychian Stage of the Silurian Period  
117 (433.4 Ma) and 16 “gold callipers” of shale gas FEBs for the case of 20 ~ 80  
118 m thick graptolite shale in China (Fig. 6). These species also include two  
119 “golden spike” graptolite species for the two GSSPs in southern China (i.e.,  
120 bases of the Darriwilian Stage in the Middle Ordovician Series and the  
121 Hirnantian Stage in the Upper Ordovician Series) (Goldman et al., 2020;  
122 Zhang et al., 2020).

123 The name of the individual image file is initialled by the specimen’s unique  
124 number and taxonomical species name. Every specimen was photographed  
125 with scale bar. The scale is attached to an image of the entire rock specimen.  
126 The other image is a close-up of the fossil within the coloured loop drawn on  
127 the whole specimen. Occasionally in the large images, the scale bar is  
128 embedded and beside the fossil specimen. For example, in the file named  
129 ‘9721Cardiograptus\_amplus\_S.jpg’, the genus name and species name are  
130 connected by the underline symbol, avoiding the space symbol. ‘9721’ is the  
131 specimen number, ‘Cardiograptus\_amplus’ means the species name is  
132 *Cardiograptus amplus* and ‘\_S’ means it is a photo with scale bar. In all scale  
133 bars, the minimum unit is one millimetre.

134 The image files are in JPG format. The single JPG file size ranges from  
135 822 KB to 7.055 MB. The whole volume of the dataset is 10.4 GB. The quality  
136 of specimen images in our dataset is much better than that in any previous  
137 publications because most specimens were first studied many years ago and  
138 their illustrations were in black and white, in low-resolution and/or printed on  
139 paper publications only. Most of these specimens were illustrated only once,  
140 or never clearly photographed. The image collection of our dataset provides

141 necessary complement for these specimens and, furthermore, unfolds their  
142 scientific value to experts or anyone who is interested in palaeontology.

143 Every of specimen is tagged with scientific information, including genus  
144 and species names, nominator, nomination year, specimen number, collection  
145 number, locality (province, city, county), geologic horizon and section, collector  
146 name, collecting time, identifier, identifying time, related references, and  
147 illustration labels. Specimens can be indexed and located in their detailed  
148 housing drawers and cabinets using any of above field element. Their detailed  
149 geologic information can also be obtained from the geological section-based  
150 database, the Geobiodiversity Database (Xu et al., 2020) and forms key  
151 elements of fossil specimen metadata (Xu et al., 2022). All related information  
152 is collected and recorded in a separate spreadsheet file released with our  
153 image dataset (Xu, 2022).

154 Some specimens of our collection have a long research history, since  
155 1958, and their taxonomical status might have changed in the light of  
156 graptolite systematic studies (Maletz, 2017; Zhang et al., 2020). We invited  
157 graptolite palaeontologists to curate every specimen to make sure that its  
158 scientific information is updated and widely accepted. The spreadsheet file  
159 includes following fields: species ID, Phylum, Class, Order, Suborder,  
160 Infraorder, Family, Subfamily, Genus, Revised species name, tagged species  
161 name, total number of specimens, specimen serial number, image file name,  
162 microscope photo number, SLR photo number, Stage, age from, age to, mean  
163 age value, locality, longitude, latitude, horizon, and first published reference. It  
164 is noted that the 'Revised species name' of every specimen reflect the  
165 emendation and correction study in Ma (2020), with comments of graptolite  
166 experts Prof Zhang Y-D and Dr Chen Q (NIGP, CAS), which differs from  
167 formal synonyms and might need further study or peer-reviewing. One can  
168 always search specimens according to tagged species names and examine  
169 specimens through our dataset, which, with the image collection and  
170 comprehensive information of a large batch of fossil specimens, supports  
171 virtual examination of specimens in a convenient and low-cost way. Experts or  
172 laymen can look through, examine, and even measure fossil specimens  
173 without need for regional/international travel and formalities. Such greatly  
174 benefits palaeontology in research, teaching, and science communication  
175 (Rahman et al., 2012).

176

#### 177 **4. Data visualization**

178 We have developed an interactive web exploration tool, FSIDvis (Fossil  
179 Specimen Image Dataset Visualizer), to assist users to examine better the  
180 scientific contents of our data (Fig. 7).

181 We further explore the distribution of these graptolite images and  
182 visualize the t-SNE feature embedding of our graptolite dataset (Fig. 8) using  
183 different colors to denote different families. In detail, for each annotated  
184 image, we first resized it into 448×448 pixels and fed it into the trained  
185 convolutional neural network (CNN) model. The output 1×1×2048 feature map  
186 from the last average pooling layer is flattened and projected to a 113  
187 (number of species) dimensional fully connected layer to represent an image  
188 embedding. After that, we use t-SNE (t-Distributed Stochastic Neighbor  
189 Embedding), a nonlinear dimension reduction technique for high-dimensional  
190 data, to project the image embeddings into the two-dimensional space for  
191 visualization. Finally, we indicate the image data distribution by a scatter plot,  
192 we use 15 colours to represent 15 families of the Order Graptoloidea,  
193 covering 42 genera and 113 species. The distribution of the images in this  
194 figure is based on species, showing a potential of automatic classifying  
195 graptolite species using CNN of the artificial intelligence (Niu and Xu, 2020).

196

#### 197 **5. Conclusions**

198 A multi-dimensional, integrated dataset based on 1,550 pieces of  
199 graptolite specimens is released. It contains 2,951 high-resolution images and  
200 a spreadsheet file showing structured records of every specimen's scientific  
201 information. During the preparation of the dataset, 113 Ordovician to Silurian  
202 graptolite species or subspecies were selected for their significances in  
203 stratigraphic correlation and shale gas exploration, and all specimens were  
204 carefully photographed and taxonomically curated.

205 Our dataset provides experts or laymen with a mean of virtual examination  
206 of a batch of fossil specimens in a convenient and low-cost way. It potentially  
207 contributes to global bio-stratigraphic correlation, especially with those bio-  
208 zone graptolite species, and in the shale gas industry to improvement of  
209 exploration efficiency. A fossil specimen database needs to fulfil the purpose  
210 and requirement of virtual examination of specimens. This greatly benefits

211 palaeontologic research and science communication. The whole dataset is  
212 visualized by the tool FSIDvis (Fossil Specimen Image Data Visualizer) and a  
213 nonlinear dimension reduction technique, t-SNE (t-Distributed Stochastic  
214 Neighbor Embedding).

215

216 **Data availability.** The dataset is archived and publicly available from  
217 <https://doi.org/10.5281/zenodo.5205215>. The visualization tool FSIDvis is  
218 available at <http://fsidvis.fossil-ontology.com:8089/>

219

220 **Author contributions.** H.-H.X. and Z.-B.N. equally designed the project,  
221 developed the model, and performed the simulations. H.-H.X. prepared and  
222 revised the manuscript. Y.-S.C. gave technician supports. X.M. revised and  
223 curated fossil specimens. Others contributed in specimen photography.

224

225 **Competing interests.** The authors declare that they have no conflict of  
226 interest.

227

228 **Acknowledgments.** We thank Prof. Zhang Yuandong and Dr. Chen Qing,  
229 NIGP, CAS, for careful curating and examining to graptolite specimens; Prof.  
230 Peter M Sadler, University of California (USA), for comment and improving the  
231 manuscript; Dr. Pan Zhaohui, Institute of Vertebrate Palaeontology and  
232 Paleoanthropology, CAS; Mr. Pan Yaohua and Mr. Wu Junqi, College of  
233 Intelligence and Computing, Tianjin University, for constructive suggestions  
234 and help.

235

236 **Financial support.** This research has been supported by National Key R&D  
237 Program of China (2022YFF0800200), Special Project on Network and  
238 Informatization of the Chinese Academy of Sciences (CAS-WX2021SF-0205),  
239 the Strategic Priority Research Program of the Chinese Academy of Sciences  
240 (Grants XDA19050101 and XDB26000000), and National Natural Science  
241 Foundation of China (61802278).

242

## 243 **References**

244 Chen, X., Chen, Q., Zhen, Y. et al.: Circumjacent distribution pattern of the  
245 Lungmachie graptolitic black shale (early Silurian) on the Yichang Uplift

246 and its peripheral region. *Science China Earth Sciences*, 61, 1195–1203,  
247 2018.

248 Chen, X., Zhang, Y., Li, Y. et al.: Biostratigraphic correlation of the Ordovician  
249 black shales in Tarim Basin and its peripheral regions. *Science China*  
250 *Earth Sciences*, 55, 1230–1237, 2012.

251 Goldman, D., Sadler, P. M. and Leslie, S. A.: The Ordovician Period, in  
252 *Geologic Time Scale 2020*. Elsevier. p. 631–694, 2020.

253 Klemme, H.D. and Ulmishek, G.F.: Effective petroleum source rocks of the  
254 world: stratigraphic distribution and controlling depositional factors. *AAPG*  
255 *Bulletin*, 75, 1809–1851. 1991.

256 Ma, X.: Palaeontology, biostratigraphy and palaeoecology of the graptolite  
257 from the Hulo Formation (Darriwilian – Sandbian) in northwestern  
258 Zhejiang Province, East China. A Ph.D dissertation submitted to University  
259 of Chinese Academy of Sciences (supervised by Prof. Zhang Y-D). 1-301.  
260 2020. [for formal publication of academic using, please contact the  
261 present author H.-H. Xu]

262 Maletz, J.: Part V, Second Revision, Chapter 13: The history of graptolite  
263 classification. *Treatise Online*, 88:1–11, 2017.

264 Niu, Z.-B. and Xu, H.-H.: AI-based graptolite identification improve shale gas  
265 exploration. *bioRxiv*. Doi: <https://doi.org/10.1101/2022.01.17.476477>

266 Peters, S. E. and McClennen, M.: The Paleobiology Database application  
267 programming interface. *Paleobiology*, 42, 1–7, 2016.

268 Podhalańska, T.: Graptolites–stratigraphic tool in the exploration of zones  
269 prospective for the occurrence of unconventional hydrocarbon deposits.  
270 *Przegląd Geologiczny*, 61, 621–629, 2013.

271 Rahman, I. A., Adcock, K. and Garwood, R.J.: Virtual fossils: a new resource  
272 for science communication in paleontology. *Evolution: Education and*  
273 *Outreach*. 5, 635–641, 2012.

274 Shute, C. H. and Foster, T. S.: Curation in museum collections. In: Jones, T.P.  
275 and Rowe, N.P., eds, *Fossil plants and spores: modern techniques*.  
276 *Geological Society of London*. 184–186, 1999.

277 Xu, H.H, Nie, T., Guo, W. et al.: Palaeontological fossil specimen metadata  
278 standard. *Acta Palaeotologica Sinica*, 61, 280–290. DOI:  
279 10.19800/j.cnki.aps.2022007. 2022.

280 Xu, H.-H., Niu, Z.-B. and Chen, Y.-S.: A status report on a section-based



281 stratigraphic and palaeontological database—the Geobiodiversity  
282 Database. Earth System Science Data, 12, 3443–3452, 2020.


283 Xu, H.-H.: High-resolution images of 1550 Ordovician to Silurian graptolite  
284 specimens for global correlation and shale gas exploration.  
285 <https://doi.org/10.5281/zenodo.5205215>. 2022.

286 Zhang, Y. D. Zhan, R. B., Wang, Z. H. et al.: 2020. Illustrations of index fossils  
287 from the Ordovician strata in China. Zhejiang University Press. 1–575,  
288 2020.

289 Zou, C. N., Dong, D., Wang, Y. et al.: Shale gas in China: Characteristics,  
290 challenges and prospects (I). Petroleum Exploration and Development.  
291 42, 689–701, 2015.

292 Zou, C. N., Gong, J., Wang, H. Y. et al. : Importance of graptolite evolution  
293 and biostratigraphic calibration on shale gas exploration. China Petroleum  
294 Exploration. 24, 1–6, 2019.

295



296

297

298

299


300

301

302

303

**Figure 1.** Global distribution of graptolite shale and shale gas production region. Most graptolite fossils were yielded from these shale sediments and their distribution is based on their occurrence records in global Ordovician and Silurian sediments. All data are from Peters and McClennen (2016) and Xu et al. (2020). The map is from © OpenStreetMap contributors 2021. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.



305


306 **Figure 2.** Graptolite species of our dataset are significant to biostratigraphy  
 307 and dating of Ordovician and Silurian sediments. These graptolites also  
 308 witnessed several macro-evolution events, including the great Ordovician  
 309 biodiversity event (GOBE), Late Ordovician mass extinction (LOME).  
 310 Radiation of several graptolite groups (bold vertical lines) occurs in this  
 311 geological time. Two global stratotype sections and points (GSSPs), based on  
 312 graptolite species record, are in southern China (the spike marks in left figure)  
 313 (data from Goldman et al., 2020). Bio- or indication zones based on graptolite  
 314 species assist with identifying mining beds for shale gas exploration in  
 315 southern China. 16 graptolite indicator-zones are used in the shale gas  
 316 exploration in China (Zou et al., 2015) (right part in the figure).  
 317



318

319 **Figure 3.** The process of creating the graptolite specimen image dataset. The  
 320 graptolite specimens were carefully curated and revised to select the species  
 321 with biostratigraphy and application significances. Every image was obtained  
 322 from specimens that were macro-photographed using a single-lens reflex  
 323 camera and microscope. After professional revision and cleaning, the whole  
 324 dataset was uploaded to and stored in our cloud server.

325



326

327 **Figure 4.** Typical images of graptolite specimens in our dataset. Every image


328 was taken from a unique graptolite specimen. Our dataset only selected the

329 photos that well show morphology of every specimen and diagnostic

330 character of each graptolite species that the specimens represent. The

331 scientific species name of every specimen is given on each image.

332



334

335

**Figure 5.** Geographic distribution (A) and geologic range (B) of graptolite

336

species of our dataset. Each graptolite specimen locality is represented by a

337

pie chart where each colour is encoded as one graptolite family of the Order

338

Graptoloidea. The sector size is proportional to the specimen number for

339

every family. The radius of the pie chart is proportional to the total number of

340

specimens from the same locality. The dashed-lines circle the main areas of

341

shale gas production. The map is from © OpenStreetMap contributors 2021.

342

Distributed under the Open Data Commons Open Database License (ODbL)

343

v1.0.

344




System	Series	Stage	Graptolite biozone (22)	Stage	Graptolite indicator zone for shale gas FEB (16)	
Silurian	Wenlock	Homerian	<i>Colonograptus deubeli</i>	Telychian	<i>Spirograptus turriculatus</i> (N2)	
		Sheinwoodian	<i>Colonograptus praedeubeli</i>		<i>Spirograptus guerichi</i> (N1)	
	Llandovery	Telychian	<i>Spirograptus turriculatus</i>	Aeronian	<i>Stimulograptus sedgwickii</i> (LM8)	
		Aeronian	<i>Lituiograptus convolutus</i>		<i>Lituiograptus convolutus</i> (LM7)	
			<i>Demirastrites triangulatus</i>		<i>Demirastrites triangulatus</i> (LM6)	
		Rhuddanian		<i>Coronograptus cyphus</i>	Rhuddanian	<i>Coronograptus cyphus</i> (LM5)
				<i>Cystograptus vesiculosus</i>		<i>Cystograptus vesiculosus</i> (LM4)
				<i>Parakidograptus acuminatus</i>		<i>Parakidograptus acuminatus</i> (LM3)
				<i>Akidograptus ascensus</i>		<i>Akidograptus ascensus</i> (LM2)
				<i>Metabolograptus persculptus</i>		<i>Metabolograptus persculptus</i> (LM1)
Ordovician	Upper	Hirnantian	<i>Metabolograptus persculptus</i>	Hirnantian	<i>Metabolograptus extraordinarius</i> (WF4)	
			<i>Metabolograptus extraordinarius</i>			
		Katian	<i>Paraorthograptus pacificus</i>	Katian	<i>Dicellograptus mirus</i> (WF3c)	
			<i>Dicellograptus complexus</i>		<i>Tangyagraptus typicus</i> (WF3b)	
	Middle	Sandbian	<i>Dicellograptus ornatus</i>	<i>Paraorthograptus pacificus</i> (WF3a)		
			<i>Dicellograptus complanatus</i>	<i>Dicellograptus complexus</i> (WF2)		
			<i>Orthograptus calcaratus</i>	<i>Dicellograptus complanatus</i> (WF1)		
		Darrivilian	<i>Hustedograptus teretiusculus</i>			
			<i>Archiclimacograptus riddellensis</i>			
			<i>Pterograptus elegans</i>			
Dapingian	<i>Nicholsonograptus fasciculatus</i>					
	<i>Levisograptus dentatus</i>					
		<i>Levisograptus austrodentatus</i>				

346

347 **Figure 6.** Graptolite species selected as global bio-zone (left) and indicator  
 348 zone (right) for shale gas favourable exploration beds (FEBs) of our dataset.  
 349 Among our dataset of 113 graptolite species, there are 22 graptolite index  
 350 species from global correlation from the Middle Ordovician to (470.0 Ma) to  
 351 the Wenlock of the Silurian Period (427.4 Ma), and 16 graptolite species as  
 352 'gold callipers' to locate FEBs of shale gas in China. Note that some graptolite  
 353 species are duplicate in the two lists.


354



355

356 **Figure 7.** FSIDvis (Fossil Specimen Image Dataset Visualizer) system  
 357 interface. a) Fossil on geographic distribution view, showing fossil specimen  
 358 location on the map. The lens (a.1) is a tailor-designed specimens' picker that  
 359 facilitates users to collect interest fossils of a region where the inner ring and  
 360 outer ring represent the family and genus. When the user chooses a genus,  
 361 the corresponding detailed species with images will be listed in the fossil list  
 362 view (a.2), where the detailed information and further high-resolution image if  
 363 the specimens are given. Hit the space bar for locking the selection. b)  
 364 Geological age scale view, providing the geologic age selection ability; the top  
 365 one is the chronostratigraphic age scale, and the bottom one is an age slider  
 366 that facilitates the users to choose a specific age slot interactively. The map is  
 367 from © OpenStreetMap contributors 2021. Distributed under the Open Data  
 368 Commons Open Database License (ODbL) v1.0.  
 369





370

371 **Figure 8.** t-SNE embedding visualization of our graptolite specimen images.

372 Individual specimens are denoted by different colors and grouped in the

373 visualization. These groups also taxonomically match different graptolite

374 families (blocks with several small images).