

1 **A multi-dimensional dataset of Ordovician to Silurian graptolite**
2 **specimens for virtual examination, global correlation and shale gas**
3 **exploration**

4
5 Hong-He Xu ^{1*}, Zhi-Bin Niu ^{1,2*}, Yan-Sen Chen ¹, Xuan Ma ¹, Xiao-Jing Tong ¹,
6 Yi-Tong Sun ¹, Xiao-Yan Dong ¹, Dan-Ni Fan ¹, Shuang-Shuang Song ¹, Yan-
7 Yan Zhu ¹, Ning Yang ¹, Qing Xia ¹

8 ¹ State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of
9 Geology and Palaeontology and Center for Excellence in Life and
10 Paleoenvironment, Chinese Academy of Sciences, 210008 Nanjing, China

11 ² College of Intelligence and Computing, Tianjin University, 300354 Tianjin,
12 China

13 *The first authors.

14 **Correspondence:** Hong-He Xu (hwxu@nigpas.ac.cn), or Zhi-Bin Niu
15 (zniu@tju.edu.cn)

16
17 **Abstract**

18 Multi- elemental and -dimensional data are more and more important in
19 the development of data-driven research, as is the case in modern
20 palaeontology, in which examination by experts, or someday artificial
21 intelligence, every fossil specimen plays a fundamental role. We here release
22 a dataset of 1,550 graptolite specimens representing 113 Ordovician to
23 Silurian graptolite species or subspecies that are significant in global
24 stratigraphic correlation and shale gas exploration. The dataset contains
25 2,951 high-resolution images and a data table of each specimen's scientific
26 information, e.g., taxonomic, geologic, and geographic information, comment,
27 and references. Our dataset provides images for specialists or laymen
28 worldwide, is supported by the tool, FSIDvis (Fossil Specimen Image Dataset
29 Visualizer), which we developed to facilitate the human-interactive exploration
30 of the rich-attribution image dataset, and also a nonlinear dimension reduction
31 technique, t-SNE (t-Distributed Stochastic Neighbor Embedding), to project
32 image data into two-dimensional space to visualize and explore similarities.
33 Our dataset potentially contributes to virtual examinations of specimens
34 (VES), global bio-stratigraphic correlation, and improvement of the shale gas
35 exploration efficiency. All data, images and the spreadsheet file, are available

36 from <https://doi.org/10.5281/zenodo.6688670> (Xu, 2022).

37

38

39 **1. Introduction**

40 Fossils are direct evidence of prehistoric life and are probably the most
41 important research object of palaeontology and stratigraphy, during which
42 fossils are collected, sampled, illustrated, described, curated, and deposited
43 as permanent specimens in museum or institution for further investigations
44 (Shute and Foster, 1999). Examinations of fossil specimens are a key and
45 indispensable part of descriptive palaeontology. Such, however, can be
46 partially achieved in a convenient and low-cost way, with the aid of multi-
47 dimensional fossil specimen dataset as in this study.

48 Graptolites are an extinct group of marine, colonial, organic-walled
49 hemichordates and have over 210 genera/3,000 species in worldwide fossil
50 records from the Cambrian to Carboniferous (c. 510~320 Ma) shales (Maletz,
51 2017). Graptolites extensively diversified in the Ordovician Period and
52 witnessed the second-largest mass extinction in geological life history, i.e., the
53 end-Ordovician mass extinction (Goldman et al., 2020). Graptolites evolved
54 quickly and spread globally in the Palaeozoic (Fig. 1), and its species are
55 widely used as significant index fossils for determining rock ages and regional
56 bio-stratigraphic correlation. Bio-zones based on graptolite species dividing
57 the Ordovician and Silurian Periods are generally less than one million years
58 in duration; such a short geological interval makes possible a precise
59 understanding of life evolution in geological history (Chen et al., 2012; 2018).
60 Up to 102 Ordovician and Silurian graptolite species were selected as global
61 bio-zones for dating sediments and understanding the evolutionary pattern of
62 palaeobiology; and 13 global stratotype sections and points (GSSPs) are
63 defined by the first appearance datum (FAD) of graptolite species from the
64 Cambrian, Ordovician, and Silurian systems (Goldman et al., 2020) (Fig. 2).

65 Additionally, bio-zones or indication zones based on graptolite species
66 assist with identifying mining beds for shale gas exploration (Fig. 1).
67 Graptolitic shale yields a significant volume of shale gas and comprises more
68 than 9% global hydrocarbons rocks (Klemme and Ulmishek, 1991;
69 Podhalańska, 2013). In China, over 61.4% of natural gas is yielded from
70 Ordovician and Silurian graptolitic shale of southern China (Zou et al., 2019).

71 Identification of graptolite species helps to locate shale gas mining beds;
72 especially, 16 graptolite species were chosen as “gold callipers” to locate
73 favourable exploration beds (FEBs) of shale gas from China (Zou et al., 2015)
74 (Fig. 2).

75 In this paper, we describe a multi-dimensional and integrated dataset of
76 graptolite specimens. The dataset potentially contributes to a range of
77 scientific activities and provides 1) easy access to and virtual examination of
78 fossil specimens through high-resolution images and detailed scientific
79 information for teaching and training in paleontology and geologic survey; 2) a
80 standard fossil specimen image dataset for use in bio-stratigraphic
81 correlations and to improve exploration efficiency in the shale gas industry,
82 and 3) a potential aid of developing image-based automated classification.

83

84 **2. Materials and methods**

85 All images in our dataset were taken from graptolite specimens that are
86 preserved in shale and were collected from China. These specimens are
87 housed at the Nanjing Institute of Geology and Palaeontology (NIGP),
88 Chinese Academy of Sciences (CAS), with serial numbers and the prefix
89 NIGP.

90 We spent over two years to photograph every specimen using a single-
91 lens reflex camera Nikon D800E with Nikkor 60 mm macro-lens and a Leica
92 M125 or M205C microscope equipped with Leica cameras (Fig. 3). Every
93 image is well focused and shows the morphology of the graptolite. In total, we
94 took 40,597 images, including 20,644 camera photos (each with a resolution
95 of 4,912 × 7,360) and 19,953 microscope photos (each with a resolution of
96 2,720 × 2,048). Photos of low contrast or bad focus were removed from the
97 whole collection. We only selected photos that show the morphology of the
98 specimen and the diagnostic characters of each graptolite species that the
99 specimen represents (Fig. 4). We selected one or two images for each
100 specimen as the final dataset, uploaded to, and stored in our cloud server
101 (Fig. 3).

102

103 **3. Data description**

104 Our final dataset consists of 2,951 high-resolution images and a related
105 spreadsheet file. Every image is a high-resolution photo taken from a

106 collection of 1,550 graptolite specimens. These specimens were formally
107 published between 1958 and 2020. They belong to 113 graptolite species or
108 subspecies of 41 genera and 16 families of the Order Graptoloidea (see the
109 spreadsheet file, Fig 5). The geological age of these graptolite species ranges
110 from the Middle Ordovician (467.3 Ma) to the Telychian Stage of the Silurian
111 Period (433.4 Ma) (Fig. 5).

112 These graptolite species have relatively abundant fossil records and are
113 significant in regional and global bio-stratigraphic correlations. They are
114 commonly used in geological age determination and shale gas FEB
115 indication, including 32 graptolite bio-zones from the Darriwilian Stage of the
116 Ordovician Period (467.3 Ma) to the Telychian Stage of the Silurian Period
117 (433.4 Ma) and 16 “gold callipers” of shale gas FEBs for the case of 20 ~ 80
118 m thick graptolite shale in China (Table 1). These species also include two
119 “golden spike” graptolite species for the two GSSPs in southern China (i.e.,
120 bases of the Darriwilian Stage in the Middle Ordovician Series and the
121 Hirnantian Stage in the Upper Ordovician Series) (Goldman et al., 2020;
122 Zhang et al., 2020).

123 The name of the individual image file is initialled by the specimen’s unique
124 number and taxonomical species name. Every specimen was photographed
125 with scale bar. The scale is attached to an image of the entire rock specimen.
126 The other image is a close-up of the fossil within the coloured loop drawn on
127 the whole specimen. Occasionally in the large images, the scale bar is
128 embedded and beside the fossil specimen. For example, in the file named
129 ‘9721Cardiograptus_amplus_S.jpg’, the genus name and species name are
130 connected by the underline symbol, avoiding the space symbol. ‘9721’ is the
131 specimen number, ‘Cardiograptus_amplus’ means the species name is
132 *Cardiograptus amplus* and ‘_S’ means it is a photo with scale bar. In all scale
133 bars, the minimum unit is one millimetre.

134 The image files are in JPG format. The single JPG file size ranges from
135 822 KB to 7.055 MB. The whole volume of the dataset is 10.4 GB. The quality
136 of specimen images in our dataset is much better than that in any previous
137 publications because most specimens were first studied many years ago and
138 their illustrations were in black and white, in low-resolution and/or printed on
139 paper publications only. Most of these specimens were illustrated only once,
140 or never clearly photographed. The image collection of our dataset provides

141 necessary complement for these specimens and, furthermore, unfolds their
142 scientific value to experts or anyone who is interested in palaeontology.

143 Every of specimen is tagged with scientific information, including genus
144 and species names, nominator, nomination year, specimen number, collection
145 number, locality (province, city, county), geologic horizon and section, collector
146 name, collecting time, identifier, identifying time, related references, and
147 illustration labels. Specimens can be indexed and located in their detailed
148 housing drawers and cabinets using any of above field element. Their detailed
149 geologic information can also be obtained from the geological section-based
150 database, the Geobiodiversity Database (Xu et al., 2020) and forms key
151 elements of fossil specimen metadata (Xu et al., 2022). All related information
152 is collected and recorded in a separate spreadsheet file released with our
153 image dataset (Xu, 2022).

154 Some specimens of our collection have a long research history, since
155 1958, and their taxonomical status might have changed in the light of
156 graptolite systematic studies (Maletz, 2017; Zhang et al., 2020). We invited
157 graptolite palaeontologists to curate every specimen to make sure that its
158 scientific information is updated and widely accepted. The spreadsheet file
159 includes following fields: species ID, Phylum, Class, Order, Suborder,
160 Infraorder, Family, Subfamily, Genus, Revised species name, tagged species
161 name, total number of specimens, specimen serial number, image file name,
162 microscope photo number, SLR photo number, Stage, age from, age to, mean
163 age value, locality, longitude, latitude, horizon, and first published reference. It
164 is noted that the 'Revised species name' of every specimen reflect the
165 emendation and correction study in Ma (2020), with comments of graptolite
166 experts Prof Zhang Y-D and Dr Chen Q (NIGP, CAS), which differs from
167 formal synonyms and might need further study or peer-reviewing. One can
168 always search specimens according to tagged species names and examine
169 specimens through our dataset, which, with the image collection and
170 comprehensive information of a large batch of fossil specimens, supports
171 virtual examination of specimens in a convenient and low-cost way. Experts or
172 laymen can look through, examine, and even measure fossil specimens
173 without need for regional/international travel and formalities. Such greatly
174 benefits palaeontology in research, teaching, and science communication
175 (Rahman et al., 2012).

176

177 **4. Data visualization**

178 We have developed an interactive web exploration tool, FSIDvis (Fossil
179 Specimen Image Dataset Visualizer), to assist users to examine better the
180 scientific contents of our data (Fig. 6).

181 We further explore the distribution of these graptolite images and
182 visualize the t-SNE feature embedding of our graptolite dataset (Fig. 7) using
183 different colors to denote different families. In detail, for each annotated
184 image, we first resized it into 448×448 pixels and fed it into the trained
185 convolutional neural network (CNN) model. The output 1×1×2048 feature map
186 from the last average pooling layer is flattened and projected to a 113
187 (number of species) dimensional fully connected layer to represent an image
188 embedding. After that, we use t-SNE (t-Distributed Stochastic Neighbor
189 Embedding), a nonlinear dimension reduction technique for high-dimensional
190 data, to project the image embeddings into the two-dimensional space for
191 visualization. Finally, we indicate the image data distribution by a scatter plot,
192 we use 15 colours to represent 15 families of the Order Graptoloidea,
193 covering 42 genera and 113 species. The distribution of the images in this
194 figure is based on species, showing a potential of automatic classifying
195 graptolite species using CNN of the artificial intelligence (Niu and Xu, 2020).

196

197 **5. Conclusions**

198 A multi-dimensional, integrated dataset based on 1,550 pieces of
199 graptolite specimens is released. It contains 2,951 high-resolution images and
200 a spreadsheet file showing structured records of every specimen's scientific
201 information. During the preparation of the dataset, 113 Ordovician to Silurian
202 graptolite species or subspecies were selected for their significances in
203 stratigraphic correlation and shale gas exploration, and all specimens were
204 carefully photographed and taxonomically curated.

205 Our dataset provides experts or laymen with a mean of virtual examination
206 of a batch of fossil specimens in a convenient and low-cost way. It potentially
207 contributes to global bio-stratigraphic correlation, especially with those bio-
208 zone graptolite species, and in the shale gas industry to improvement of
209 exploration efficiency. A fossil specimen database needs to fulfil the purpose
210 and requirement of virtual examination of specimens. This greatly benefits

211 palaeontologic research and science communication. The whole dataset is
212 visualized by the tool FSIDvis (Fossil Specimen Image Data Visualizer) and a
213 nonlinear dimension reduction technique, t-SNE (t-Distributed Stochastic
214 Neighbor Embedding).

215

216 **Data availability.** The dataset is archived and publicly available from
217 <https://doi.org/10.5281/zenodo.6688670>. The visualization tool FSIDvis is
218 available at <http://fsidvis.fossil-ontology.com:8089/>

219

220 **Author contributions.** H.-H.X. and Z.-B.N. equally designed the project,
221 developed the model, and performed the simulations. H.-H.X. prepared and
222 revised the manuscript. Y.-S.C. gave technician supports. X.M. revised and
223 curated fossil specimens. Others contributed in specimen photography.

224

225 **Competing interests.** The authors declare that they have no conflict of
226 interest.

227

228 **Acknowledgments.** We thank Prof. Zhang Yuandong and Dr. Chen Qing,
229 NIGP, CAS, for careful curating and examining to graptolite specimens; Prof.
230 Peter M Sadler, University of California (USA), for comment and improving the
231 manuscript; Dr. Pan Zhaohui, Institute of Vertebrate Palaeontology and
232 Paleoanthropology, CAS; Mr. Pan Yaohua and Mr. Wu Junqi, College of
233 Intelligence and Computing, Tianjin University, for constructive suggestions
234 and help.

235

236 **Financial support.** This research has been supported by National Key R&D
237 Program of China (2022YFF0800200), Special Project on Network and
238 Informatization of the Chinese Academy of Sciences (CAS-WX2021SF-0205),
239 the Strategic Priority Research Program of the Chinese Academy of Sciences
240 (Grants XDA19050101 and XDB26000000), and National Natural Science
241 Foundation of China (61802278).

242

243 **References**

244 Chen, X., Chen, Q., Zhen, Y. et al.: Circumjacent distribution pattern of the
245 Lungmachiean graptolitic black shale (early Silurian) on the Yichang Uplift

246 and its peripheral region. *Science China Earth Sciences*, 61, 1195–1203,
247 2018.

248 Chen, X., Zhang, Y., Li, Y. et al.: Biostratigraphic correlation of the Ordovician
249 black shales in Tarim Basin and its peripheral regions. *Science China*
250 *Earth Sciences*, 55, 1230–1237, 2012.

251 Goldman, D., Sadler, P. M. and Leslie, S. A.: The Ordovician Period, in
252 *Geologic Time Scale 2020*. Elsevier. p. 631–694, 2020.

253 Klemme, H.D. and Ulmishek, G.F.: Effective petroleum source rocks of the
254 world: stratigraphic distribution and controlling depositional factors. *AAPG*
255 *Bulletin*, 75, 1809–1851. 1991.

256 Ma, X.: Palaeontology, biostratigraphy and palaeoecology of the graptolite
257 from the Hulo Formation (Darriwilian – Sandbian) in northwestern
258 Zhejiang Province, East China. A Ph.D dissertation submitted to University
259 of Chinese Academy of Sciences (supervised by Prof. Zhang Y-D). 1-301.
260 2020. DOI:10.5281/zenodo.7827023.

261 Maletz, J.: Part V, Second Revision, Chapter 13: The history of graptolite
262 classification. *Treatise Online*, 88:1–11, 2017.

263 Niu, Z.-B. and Xu, H.-H.: AI-based graptolite identification improve shale gas
264 exploration. *bioRxiv*. DOI:[10.1101/2022.01.17.476477](https://doi.org/10.1101/2022.01.17.476477)

265 Peters, S. E. and McClennen, M.: The Paleobiology Database application
266 programming interface. *Paleobiology*, 42, 1–7, 2016.

267 Podhalańska, T.: Graptolites–stratigraphic tool in the exploration of zones
268 prospective for the occurrence of unconventional hydrocarbon deposits.
269 *Przegląd Geologiczny*, 61, 621–629, 2013.

270 Rahman, I. A., Adcock, K. and Garwood, R.J.: Virtual fossils: a new resource
271 for science communication in paleontology. *Evolution: Education and*
272 *Outreach*. 5, 635–641, 2012.

273 Shute, C. H. and Foster, T. S.: Curation in museum collections. In: Jones, T.P.
274 and Rowe, N.P., eds, *Fossil plants and spores: modern techniques*.
275 *Geological Society of London*. 184–186, 1999.

276 Xu, H.H, Nie, T., Guo, W. et al.: Palaeontological fossil specimen metadata
277 standard. *Acta Palaeotologica Sinica*, 61, 280–290. DOI:
278 [10.19800/j.cnki.aps.2022007](https://doi.org/10.19800/j.cnki.aps.2022007). 2022.

279 Xu, H.-H., Niu, Z.-B. and Chen, Y.-S.: A status report on a section-based
280 stratigraphic and palaeontological database–the Geobiodiversity

281 Database. *Earth System Science Data*, 12, 3443–3452, 2020.


282 Xu, H.-H.: High-resolution images of 1550 Ordovician to Silurian graptolite
283 specimens for global correlation and shale gas exploration.
284 <https://doi.org/10.5281/zenodo.6688670>. 23 June, 2022.

285 Zhang, Y. D. Zhan, R. B., Wang, Z. H. et al.: 2020. Illustrations of index fossils
286 from the Ordovician strata in China. Zhejiang University Press. 1–575,
287 2020.

288 Zou, C. N., Dong, D., Wang, Y. et al.: Shale gas in China: Characteristics,
289 challenges and prospects (I). *Petroleum Exploration and Development*.
290 42, 689–701, 2015.

291 Zou, C. N., Gong, J., Wang, H. Y. et al.: Importance of graptolite evolution and
292 biostratigraphic calibration on shale gas exploration. *China Petroleum*
293 *Exploration*. 24, 1–6, 2019.

294



295

296

297

298


299

300

301

302

Figure 1. Global distribution of graptolite shale and shale gas production region. Most graptolite fossils were yielded from these shale sediments and their distribution is based on their occurrence records in global Ordovician and Silurian sediments. All data are from Peters and McClennen (2016) and Xu et al. (2020). The map is from © OpenStreetMap contributors 2021. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.




303
 304 **Figure 2.** Graptolite species of our dataset are significant to biostratigraphy
 305 and dating of Ordovician and Silurian sediments. These graptolites also
 306 witnessed several macro-evolution events, including the great Ordovician
 307 biodiversity event (GOBE), Late Ordovician mass extinction (LOME).
 308 Radiation of several graptolite groups (bold vertical lines) occurs in this
 309 geological time. Two global stratotype sections and points (GSSPs), based on
 310 graptolite species record, are in southern China (the spike marks in left figure)
 311 (data from Goldman et al., 2020). Bio- or indication zones based on graptolite
 312 species assist with identifying mining beds for shale gas exploration in
 313 southern China. 16 graptolite indicator-zones are used in the shale gas
 314 exploration in China (Zou et al., 2015) (right part in the figure).
 315



316


317 **Figure 3.** The process of creating the graptolite specimen image dataset. The
 318 graptolite specimens were carefully curated and revised to select the species
 319 with biostratigraphy and application significances. Every image was obtained
 320 from specimens that were macro-photographed using a single-lens reflex
 321 camera and microscope. After professional revision and cleaning, the whole
 322 dataset was uploaded to and stored in our cloud server.

323



324
 325
 326
 327
 328
 329
 330


Figure 4. Typical images of graptolite specimens in our dataset. Every image was taken from a unique graptolite specimen. Our dataset only selected the photos that well show morphology of every specimen and diagnostic character of each graptolite species that the specimens represent. The scientific species name of every specimen is given on each image.



331

332 **Figure 5.** Geographic distribution (A) and geologic range (B) of graptolite
 333 species of our dataset. Each graptolite specimen locality is represented by a
 334 pie chart where each colour is encoded as one graptolite family of the Order
 335 Graptoloidea. The sector size is proportional to the specimen number for
 336 every family. The radius of the pie chart is proportional to the total number of
 337 specimens from the same locality. The dashed-lines circle the main areas of
 338 shale gas production. The map is from © OpenStreetMap contributors 2021.
 339 Distributed under the Open Data Commons Open Database License (ODbL)
 340 v1.0.

341




342

343 **Figure 6.** FSIDVis (Fossil Specimen Image Dataset Visualizer) system
 344 interface. a) Fossil on geographic distribution view, showing fossil specimen
 345 location on the map. The lens (a.1) is a tailor-designed specimens' picker that
 346 facilitates users to collect interest fossils of a region where the inner ring and
 347 outer ring represent the family and genus. When the user chooses a genus,
 348 the corresponding detailed species with images will be listed in the fossil list
 349 view (a.2), where the detailed information and further high-resolution image if
 350 the specimens are given. Hit the space bar for locking the selection. b)

351 Geological age scale view, providing the geologic age selection ability; the top
 352 one is the chronostratigraphic age scale, and the bottom one is an age slider
 353 that facilitates the users to choose a specific age slot interactively. The map is
 354 from © OpenStreetMap contributors 2021. Distributed under the Open Data
 355 Commons Open Database License (ODbL) v1.0.

356



357

358 **Figure 7.** t-SNE embedding visualization of our graptolite specimen images.

359 Individual specimens are denoted by different colors and grouped in the

360 visualization. These groups also taxonomically match different graptolite

361 families (blocks with several small images).

362

363

364 **Table 1.** Graptolite species selected as global biozone and indicator zone

365 (right) for shale gas favourable exploration beds (FEBs) of our dataset.

Period	Epoch	Age	Graptolite biozone	Graptolite indicator zone for shale gas FEB
Silurian	Llandovery	Telychian	<i>Cyrtograptus centrifugus</i>	
			<i>Cyrtograptus insectus</i>	
			<i>Cyrtograptus lapworthi</i>	
			<i>Monoclimacis griestoniensis</i>	
			<i>Monoclimacis crispus</i>	
			<i>Spirograptus turriculatus</i>	<i>Spirograptus turriculatus</i> (N2)
		<i>Spirograptus guerichi</i>	<i>Spirograptus guerichi</i> (N1)	
		Aeronian	<i>Stimulograptus sedgwickii</i>	<i>Stimulograptus sedgwickii</i> (LM8)
			<i>Lituigraptus convolutus</i>	<i>Lituigraptus convolutus</i> (LM7)
Rhuddanian	<i>Demirastrites triangulatus</i>	<i>Demirastrites triangulatus</i> (LM6)		

Ordovician			<i>Coronograptus cyphus</i>	<i>Coronograptus cyphus</i> (LM5)	
			<i>Cystograptus acuminatus</i>	<i>Cystograptus vesiculosus</i> (LM4)	
			<i>Parakidograptus acuminatus</i>	<i>Parakidograptus acuminatus</i> (LM3)	
	Upper	Hirnatian	<i>Akidograptus ascensus</i>	<i>Akidograptus ascensus</i> (LM2)	
			<i>Normalograptus persculptus</i>	<i>Metabolograptus persculptus</i> (LM1)	
			<i>Normalograptus extraordinarius</i>	<i>Metabolograptus extraordinarius</i> (WF4)	
			Katian	<i>Paraorthograptus pacificus</i>	<i>Dicellograptus mirus</i> (WF3c)
				<i>Dicellograptus complexus</i>	<i>Tangyagraptus typicus</i> (WF3b)
				<i>Dicellograptus complanatus</i>	<i>Paraorthograptus pacificus</i> (WF3a)
		<i>Dicellograptus elegans</i>		<i>Dicellograptus complexus</i> (WF2)	
		<i>Geniculograptus pygmaeus</i>		<i>Foliomena - Nankinolithus</i> (WF1)	
		Sandbian	<i>Diplacanthograptus spiniferus</i>		
			<i>Diplacanthograptus caudatus</i>		
			<i>Climacograptus bicornis</i>		
			<i>Nemagraptus gracilis</i>		
		Middle	Darriwilian	<i>Jiangxigraptus vagus</i>	
				<i>Didymograptus murchisoni</i>	
				<i>Pterograptus elegans</i>	
				<i>Nicholograptus fasciculatus</i>	
				<i>Acrograptus ellesae</i>	
Dapingian	<i>Undulograptus austrodentatus</i>				
	<i>Exigraptus clavus</i>				