



A 30 year monthly 5 km gridded surface elevation time series for the Greenland Ice Sheet from multiple satellite radar altimeters

Baojun Zhang^{1*}, Zemin Wang^{1*}, Jiachun An^{1†}, Tingting Liu^{1†}, Hong Geng²

¹Chinese Antarctic Center of Surveying and Mapping, Wuhan University, Wuhan, 430079, China

5 ²School of Resource and Environment Science, Wuhan University, Wuhan, 430079, China

*These authors contributed equally to this work.

†Correspondence to: Jiachun An (jcan@whu.edu.cn) and Tingting Liu (tliu23@whu.edu.cn)

Abstract. A long-term time series of ice sheet surface elevation change (SEC) is important for study of ice sheet variation and its response to climate change. In this study, we used an updated plane-fitting least-squares regression strategy to generate a
10 30 year surface elevation time series for the Greenland Ice Sheet (GrIS) at monthly temporal resolution and 5×5 km grid spatial resolution using ERS-1, ERS-2, Envisat, and CryoSat-2 satellite radar altimeter observations obtained between August 1991 and December 2020. The accuracy and reliability of the time series are effectively guaranteed by application of sophisticated corrections for intermission bias and interpolation based on empirical orthogonal function reconstruction. Validation using both airborne laser altimeter observations and the European Space Agency GrIS Climate Change Initiative
15 (CCI) product indicated that our merged surface elevation time series is reliable. The accuracy and dispersion of errors of SECs of our results were 19.3% and 8.9% higher, respectively, than those of CCI SECs, and even 30.9% and 19.0% higher, respectively, in periods from 2006–2010 to 2010–2014. Further analysis showed that our merged time series could provide detailed insight into GrIS SEC on multiple temporal (up to 30 years) and spatial scales, thereby providing opportunity to explore potential associations between ice sheet change and climatic forcing. The merged surface elevation time series data
20 are available at <http://dx.doi.org/10.11888/Glacio.tpdc.271658> (Zhang et al., 2021).

1 Introduction

Over recent decades, the Greenland Ice Sheet (GrIS) has experienced increasing substantial imbalance. Driven by atmospheric and oceanic warming (Straneo and Heimbach, 2013; Hanna et al., 2012), this imbalance has become a leading driver of global sea level change, whose contribution to which is almost 1.4 times that of the Antarctic Ice Sheet (Shepherd et al., 2018; Shepherd et al., 2020). As a result of changes in surface mass balance (SMB) and ice dynamics, ice sheet elevation change (EC) is a direct indicator of climate change. Furthermore, with an appropriate density model of ice firn layers, ECs can be used
25 to monitor variation in ice sheet mass balance. Thus, a long-term time series of GrIS EC is essential for assessing climate change (Sørensen et al., 2018). Since 1991, various satellite altimetry missions have made continuous observation of ice sheet ECs a reality (Shepherd et al., 2019; Simonsen et al., 2021). This approach, which uses measurements of surface EC (SEC)



30 derived from satellite altimetry to monitor ice sheet mass balance, provides an unprecedented advantage in terms of
spatiotemporal resolution in comparison with two other satellite-based techniques: gravimetric mass balance derived from
satellite gravimetry and input–output balance derived from remotely sensed ice flow (Shepherd et al., 2019; Simonsen et al.,
2021).

The effective life of a single satellite mission is limited, which means reconstruction of a long-term ice sheet elevation time
35 series requires observations from multiple altimeter missions to be combined. In such a process, the method adopted to
eliminate system biases is a crucial factor. System biases include intermission bias, ascending–descending bias, and time-
variable penetration effects. It is generally believed that intermission bias is derived mainly from orbital errors, and differences
in the centre of gravity and phase of antennae between satellites (Frappart et al., 2016). Owing to its distinct spatial pattern
(Zwally et al., 2005; Frappart et al., 2016), intermission bias is generally corrected for each grid cell using an estimate
40 calculated from observations over overlapping epochs (Paolo et al., 2016; Sørensen et al., 2018; Adusumilli et al., 2018;
Schröder et al., 2019; Shepherd et al., 2019; Simonsen et al., 2021). The ascending–descending bias can be considered to
comprise both intra-mission ascending–descending bias and inter-mission ascending–descending bias (Zhang et al., 2020).
Both are related to the angle between radar polarization and wind-induced features of the firm (Armitage et al., 2014; Remy et
al., 2006). The former can be corrected by introducing a term for satellite flight direction into a regression model (Simonsen
45 and Sorensen, 2017; Mcmillan et al., 2014; Yang et al., 2019), or reduced by re-tracking the radar return waveform using a
threshold re-tracker (Helm et al., 2014; Schröder et al., 2019). No specific treatment has been proposed for handling the latter,
except that it is accounted for by the introduction of estimations of a series of parameters into a regression model (Zhang et
al., 2020). It has been proven that using a large amount of data to fine-tune the correction of intermission bias and ascending–
descending bias can ensure better self-consistency and reliability of the combined time series of elevation (Zhang et al., 2020).
50 Unfortunately, this method is unsuitable for combining data from multiple satellite missions simultaneously, because
introduction into the fitting model of additional parameters and the increasingly complicated topological relationships between
them will lead to regression failure. For mitigating time-variable penetration effects, there are currently three common
approaches: including waveform parameters in the regression model (Flament and Remy, 2012; Simonsen and Sorensen, 2017),
re-tracking the radar return waveforms with a threshold re-tracker (Nilsson et al., 2016; Helm et al., 2014; Schröder et al.,
55 2017), or applying a waveform deconvolution model to the radar return waveforms (Arthern et al., 2001; Mcmillan et al., 2016;
Slater et al., 2019). However, none of these approaches can account completely for the time-variable penetration effects.
Therefore, construction of a long-term elevation time series through combining data from sensors with similar characteristics
might represent a more reasonable approach.

The ground tracks of polar orbiting altimetry satellites over the polar ice sheets are irregular. Additionally, certain outliers
60 exist in altimeter observations, especially in relation to the early altimetry missions, e.g., ERS-1 (Schröder et al., 2019). These
problems will result in lack of available data values in certain cells of a joint elevation time series. Thus, to estimate the volume
or mass change over a basin or an entire ice sheet, gridding methods such as kriging (e.g., Bamber et al., 2009; Slater et al.,
2018), tension continuous curvature splines (e.g., Zhang et al., 2017), or inverse distance weight (e.g., Chuter and Bamber,



2015) are usually employed to interpolate or even extrapolate the results for unobserved cells. However, such straightforward
65 interpolations are unable to reflect the true patterns of elevation or ECs in steep and very active areas across ice sheet margins
(Hurkmans et al., 2012), not to mention the accuracy of the extrapolation results where there are insufficient constraints.
Assuming that the spatial distribution pattern of the variation of ice sheet SECs is very small temporally, then orthogonal
spatial maps of surface elevation (SE) variability can be extracted using empirical orthogonal function (EOF) decomposition
from a sufficiently long elevation time series. Together with sparse observations, orthogonal spatial maps can be used to realize
70 interpolation (reconstruction) of a time series of early satellite-derived ECs. Actually, EOF reconstruction has already been
used for reconstruction of sea surface temperature (e.g., Smith et al., 1996) and sea level change (e.g., Chambers et al., 2002;
Jin et al., 2012) time series. The high-quality observations of Envisat and CryoSat-2, especially the higher-resolution coverage
of Cryosat-2, provide potential for the use of EOF reconstruction for interpolation of an early elevation time series.

Here, we improve a previously proposed algorithm (Zhang et al., 2020) that requires a large volume of observations in an
75 integrated adjustment model for simultaneous correction of intermission bias and ascending–descending bias. While retaining
its advantages, we develop a 30 year (1991–2020) monthly 5×5 km gridded SE time series for the GrIS by merging ERS-1,
ERS-2, Envisat, and CryoSat-2 radar altimetry observations. Then, to facilitate the use of the elevation time series, we use the
EOF reconstruction method for more reliable interpolation of data for grid cells with missing values. In this paper, the details
of the data processing are presented. The final merged SE time series dataset is freely available at
80 <http://dx.doi.org/10.11888/Glacio.tpd.271658> (Zhang et al., 2021).

2 Material and methodology

2.1 Satellite radar altimetry data

In this study, we used ice sheet SE observations from four European Space Agency (ESA) satellite radar altimeter missions:
ERS-1, ERS-2, Envisat, and CryoSat-2. Since the launch of ERS-1 in 1991, satellite radar altimeter missions have continuously
85 collected SE observations for 30 years using similar sensors. Currently, following the retirement of the first three missions,
only CryoSat-2 remains in operation.

For ERS-1 and ERS-2, we used the Level 2 (L2) Geophysical Data Record (GDR) product from the REAPER project version
RP01, which has been reprocessed to align both the ERS and Envisat datasets (Brockley et al., 2017). For Envisat, we
downloaded the latest L2 GDR product version 3.0 from ESA, which is better than the previous version (ver. 2.1) in terms of
90 coverage and performance at cross-overs. For CryoSat-2, we used the latest Baseline D L2 GDR data provided by ESA. Over
land ice, Baseline D improves the ascending–descending crossover statistics to 0.1 m from 1.9 m achieved with the previous
version Baseline C (Meloni et al., 2020). Here, heights re-tracked by the ICE-1 re-tracker for ERS-1, ERS-2, and Envisat, the
OCOG re-tracker for the CryoSat-2 LRM, and the Wingham/Wallis model fit re-tracker for the CryoSat-2 SRAIn were used.
Before performing combined calculations, all erroneous height records were eliminated using standard quality flags.



95 2.2 Airborne laser altimetry data

To bridge the gap in observations between the ICESat and ICESat-2 laser altimeter missions, the Operation IceBridge project (OIB) implemented more than 1000 airborne surveys during 2009–2020. During the OIB campaign, the airborne laser altimeter payload (i.e., the Airborne Topographic Mapper (ATM)) recorded a large number of high-precision ice sheet SE observations. Additionally, prior to OIB, several Pre-IceBridge airborne ATM surveys were conducted between 1993 and 2008. The OIB and Pre-IceBridge ATM SE datasets can both be download from the National Snow and Ice Data Center. Because their accuracy is 10 cm or better (Krabill et al., 2004), we used these ATM elevation measurements (i.e., ATM L2 product) to validate our merged SE time series. Additionally, SECs derived from OIB and Pre-IceBridge ATM measurements (i.e., ATM L4 product) were also used.

2.3 Generation of surface elevation time series

Our previous study (Zhang et al., 2020) demonstrated that using a large amount of data to fine-tune the correction of intermission bias and ascending–descending bias can develop a more self-consistent and reliable combined elevation time series. However, use of the updated plane-fitting least-squares regression model of Zhang et al. (2020) to merge data from three or more satellite missions is not straightforward. As the number of satellite altimetry missions involved in the calculation increases, additional terms of system bias and the increasingly complex topological relationships between them must be considered in the least-squares regression model, making the model overly complicated and ultimately incomprehensible. However, it is possible to divide the calculation into several individual steps, reducing the complexity of the model, while retaining its advantages.

First, the intra-mission ascending–descending bias b_{AD} at a grid cell for each radar altimeter mission can be estimated as follows:

$$\begin{aligned} h(lon_i, lat_i, t_i) = & h_0 + a_0(lon_i - lon_0) + a_1(lat_i - lat_0) + a_2(lon_i - lon_0)^2 + a_3(lat_i - lat_0)^2 \\ & + a_4(lon_i - lon_0)(lat_i - lat_0) + a_5(t_i - t_0) + a_6 \cos(2\pi(t_i - t_0)) + a_7 \sin(2\pi(t_i - t_0)) \\ & + a_8(bs_i - \overline{bs}) + b_{AD}(-1)^{AD} + res(lon_i, lat_i, t_i), \end{aligned} \quad (1)$$

where $h(lon_i, lat_i, t_i)$ denotes the surface height measured at longitude (lon_i), latitude (lat_i), and time (t_i); the reference epoch t_0 was set to 2010.0 in this study; lon_0 , lat_0 , and h_0 represent the longitude, latitude, and height (at t_0), respectively, of the centre of a grid cell; a_0 – a_4 are the quadratic expansion for surface topography; a_5 – a_7 denote the linear and seasonal signals for temporal changes of SE; a_8 is a parameter to mitigate the time-variable penetration effects of the radar signal using the anomaly of backscattered power ($bs_i - \overline{bs}$); and $res(lon_i, lat_i, t_i)$ denotes the residuals of the regression. Note that the SARIn and LRM observations of CryoSat-2 should be calculated separately here.

Second, the intermission bias b_{im} between Envisat and SARIn (or LRM) of CryoSat-2 can be calculated from the corrected heights $h^c(lon_i, lat_i, t_i)$ by applying b_{AD} :



$$\begin{aligned}
 h^c(lon_i, lat_i, t_i) = & h_0 + a_0(lon_i - lon_0) + a_1(lat_i - lat_0) + a_2(lon_i - lon_0)^2 + a_3(lat_i - lat_0)^2 \\
 & + a_4(lon_i - lon_0)(lat_i - lat_0) + a_5(t_i - t_0) + a_6 \cos(2\pi(t_i - t_0)) + a_7 \sin(2\pi(t_i - t_0)) \\
 & + a_8(bs_i - \bar{bs}) + b_{im}(-1)^{im} + res(lon_i, lat_i, t_i).
 \end{aligned} \tag{2}$$

The above implies that Envisat is taken as reference, which means that the bias between SARIn and LRM will be corrected in this step. After applying b_{im} , Envisat and CryoSat-2 data will be consistent, and subsequently ERS-2 and then ERS-1 data can also be corrected to be consistent with them using Eq. (2).

Third, all the consistent heights $h^{cc}(lon_i, lat_i, t_i)$ can be used in the final least-squares regression to construct the SE time series:

$$\begin{aligned}
 h^{cc}(lon_i, lat_i, t_i) = & h_0 + a_0(lon_i - lon_0) + a_1(lat_i - lat_0) + a_2(lon_i - lon_0)^2 + a_3(lat_i - lat_0)^2 \\
 & + a_4(lon_i - lon_0)(lat_i - lat_0) + a_5(t_i - t_0) + a_6 \cos(2\pi(t_i - t_0)) + a_7 \sin(2\pi(t_i - t_0)) \\
 & + a_8(bs_i - \bar{bs}) + res(lon_i, lat_i, t_i).
 \end{aligned} \tag{3}$$

The elevation anomaly can be derived as follows:

$$\begin{aligned}
 \nabla h(lon_i, lat_i, t_i) = & h^{cc}(lon_i, lat_i, t_i) \\
 & - \left(a_0(lon_i - lon_0) + a_1(lat_i - lat_0) + a_2(lon_i - lon_0)^2 + a_3(lat_i - lat_0)^2 \right. \\
 & + a_4(lon_i - lon_0)(lat_i - lat_0) + a_5(t_i - t_0) + a_6 \cos(2\pi(t_i - t_0)) + a_7 \sin(2\pi(t_i - t_0)) \\
 & \left. + a_8(bs_i - \bar{bs}) \right).
 \end{aligned} \tag{4}$$

Note that the removal of h_0 is to facilitate study of ECs, and that h_0 can be used to generate an independent digital elevation model (DEM). The DEM and the corresponding surface slope and aspect are shown in Fig. 1. When necessary, h_0 can be added back. Then, the monthly SE time series for a grid cell can be obtained as follows:

$$\overline{\Delta h}(lon_0, lat_0, t^j) = \frac{1}{n} \sum_{i=1}^n \Delta h(lon_0, lat_0, t_i), \tag{5}$$

where n is the number of corrected elevations in month t^j .

To generate a robust time series of 5 km gridded elevations, the above least-squares fitting is first performed on a 2 km polar-stereographic grid over the GrIS. For each grid cell, all observations within 2.5 km of its centre are used for the iterative least-squares estimation under the constraints of 3σ outlier rejection criteria and the same thresholds as in Zhang et al. (2020). Then, a 40 km floating median low-pass filter, similar to Schröder et al. (2019), and the same spatiotemporal median filter as used by Zhang et al. (2020), are implemented to generate the final merged robust 5 km gridded time series.

2.4 Interpolation based on EOF reconstruction

Assuming that the spatial patterns of GrIS SECs are stationary in time, the three-dimensional GrIS SE anomaly time series $\nabla H(lon, lat, t)$ can be represented as a linear combination of the EOF modes $eof_i(lon, lat)$ and principal components $pc_i(t)$ (Chambers et al., 2002; Jin et al., 2012):



$$\nabla H(lon, lat, t) = \sum_{i=1}^N eof_i(lon, lat)pc_i(t), \quad (6)$$

where N is the total number of EOF modes; and lon , lat , and t denote the temporal and spatial position of a certain SE anomaly. The purpose of solving EOF modes is to supplement the sparse monthly gridded data attributable to poor observations in the early years. Therefore, we use the gridded time series during 2003–2020 obtained from the higher quality observations of Envisat and CryoSat-2 here. To mitigate errors caused by extrapolation, only grid cells with at least 100 elevation anomalies in 216 months are retained. The missing values in the retained gridded time series are interpolated using ordinary kriging, a technique usually employed to generate a DEM (Bamber et al., 2009; Slater et al., 2018).

Then, for the monthly grid that needs interpolation, the following equation can be established:

$$v_t = \sum_{i=1}^M eof_i(lon, lat)PC_i(t) - T(lon, lat, t), \quad (7)$$

where $T(lon, lat, t)$ denotes the values already in this monthly grid; M means choosing the first M EOF modes; v_t is the interpolation (reconstruction) error; and $PC_i(t)$ is the principal components to be estimated corresponding to each of the M modes for this monthly grid, which can be estimated to minimize v_t using a linear least-squares estimator. To determine M , we experimented by adjusting it from 1 to 216 modes. We found that both the percentage of the explained variance and the root mean square (RMS) difference between the reconstructed time series of monthly EC and that of the observations become insensitive after 30 modes with only minor changes, as can be seen in Fig. 2. Thereby, the elevation anomalies missing from grid cells during the period of the ERS missions are interpolated using EOF reconstruction. Note that we also first deduct the seasonal signals before performing the EOF reconstruction, and then return them to the reconstruction results here. The performance of both EOF reconstruction and ordinary kriging is shown in Fig. 3, illustrating the superiority of the former in comparison with the latter. Especially in extrapolation, there are many obvious over-interpolations in the ordinary kriging result.

2.5 Uncertainty for surface elevation time series

As described in Sect. 2.3, the elevation anomaly in a grid cell of the merged SE time series is derived using a median estimator. Thus, to obtain a realistic estimation of error, we also use the scaled median absolute deviation MAD_S as a metric of its uncertainty following Ewert et al. (2012):

$$MAD_S = k \text{median}(|H - \text{median}(H)|). \quad (8)$$

The scale factor k is set to 1.4826 to make MAD_S a consistent estimator similar to the standard deviation.



As for those interpolated elevation anomalies in the gridded time series, because of the complicated interpolation methods adopted, it is difficult to estimate their uncertainty using formal error propagation techniques. Here, we use the scaled RMS of the residuals ε derived from the elevation anomalies h in a grid cell as follows:

$$185 \quad h = b_0 + b_1 t + b_2 t^2 + b_3 \cos(2\pi(t - t_0)) + b_4 \sin(2\pi(t - t_0)) + \varepsilon, \quad (9)$$

where b_0 is a constant; and b_1 – b_4 denote the linear, quadratic, and seasonal signals of the temporal changes of SE, respectively.

A scale factor of 1.05 is used to compensate for the reduced RMS error due to fitting (Wahr et al., 2006).

It should be noted that when using our elevation time series to estimate the volume change for individual drainage basins and major sectors, the correlated uncertainties between adjacent grid cells should also be considered. According to Schröder et al.
190 (2019), applying a scaling factor to the squared uncertainties can account for the autocorrelation over an area.

3 Results

3.1 Surface elevation anomaly time series

The average rate of SECs in a certain time interval can be calculated from SE time series using a least-squares fitting model with a first-order polynomial and a sine wave with a 1 year period. The additional annual items are used to avoid the bias
195 caused when the entire annual cycle is not fully covered.

The average SEC rates and their uncertainties over the periods 1991–2000, 2001–2010, 2011–2020, and 1991–2020 from our merged elevation anomaly time series are shown in Fig. 4. As reported by Shepherd et al. (2020), the GrIS has been losing ice throughout most of the intervening period. Thus, maps of these long-term average SEC rates show signals of continuous thinning in many areas along the coast. Overall, the most notable signals of GrIS thinning are concentrated on the west coast
200 of Greenland, especially along Melville Bay and near Jakobshavn Isbræ. Comparison of the average rates in the different periods reveals significantly accelerated and expanded thinning in many outlet glaciers, e.g., Jakobshavn Isbræ and Upernavik Isstrøm on the west coast of the GrIS, Zachariæ Isstrøm and Nioghalvfjordsfjorden glacier in the northeast of the GrIS, Kangerdlugssuaq Gletscher and Helheimgletscher in the southeast of the GrIS, and Petermann Gletscher and Humboldt Gletscher in the northwest of the GrIS. The main contributor to the significant thinning detected in these regions is ice dynamics.
205 The volume of solid ice being discharged into the ocean is increasing because of warmer air and ocean temperatures (Mouginot et al., 2015; Aschwanden et al., 2016; Shepherd et al., 2020; Wood et al., 2021). Signs of thickening are evident mainly in accumulation areas with higher elevation in central and northwestern parts of the GrIS, e.g., the area near King Christian X Land. These weak signals of thickening mainly reflect the increase in SMB derived from a combination of high snowfall and low surface melting (Simonsen et al., 2021).

210 Irrespective of whether thickening or thinning, it can be seen that the rates of SEC in different periods vary. To gain insight into the spatiotemporal changes of average SEC rates, the average SEC rates and their uncertainties at 5 year intervals during 1991–2020 for our time series are illustrated in Fig. 5. Considerable variation in the mean SEC rates is evident, e.g., abnormal



accumulation during 1996–2000, gradually increasing loss from 2000–2005 to 2011–2015, and deceleration of thinning during 2015–2020. Benefitting from the higher temporal and spatial resolution of our combined time series, the small-scale
215 spatiotemporal evolution of the average rates of SECs can be analysed in detail. Taking Jakobshavn Isbræ as an example, Fig. 5 clearly reveals its evolution from thinning in the early 1990s, to equilibrium in the late 1990s, to accelerated thinning in the first decade of the 2000s, and then to deceleration of thinning in recent years. Similarly, the evolution of other glaciers can be obtained from our time series.

Our 5 km gridded time series can also provide a more detailed evolution of SEC characteristics on temporal scales of up to 30
220 years. The four examples presented in Fig. 6 illustrate that our results have the capability to pinpoint such signals of GrIS SEC. Jakobshavn Isbræ is the largest and fastest outlet glacier on the west coast of Greenland; however, its thinning throughout the observational period since 1991 is not continuous. For example, short-term deceleration of thinning and thickening during 1996–2001 and since 2014, caused by deceleration of the ice flow (Joughin et al., 2004; Khazendar et al., 2019), can be seen in Fig. 6(a). A rapid drop in the surface elevation of Jakobshavn Isbræ is evident during 2003–2013. The rate of surface
225 lowering increases with increasing distance from the grounding line. During this period, the mean rate at position A (Fig. 6(a)) is up to -2.85 ± 0.04 m yr⁻¹. Upernavik Isstrøm consists of five glaciers, all of which flow into the same fjord. Zachariæ Isstrøm and Nioghalvfjærdsfjorden drain the majority of the Northeast Greenland Ice Stream. Unlike Jakobshavn Isbræ, their surfaces have lowered consistently since 1991 (Fig. 6(b) and (c)). The average rates of thinning at A, B, and C in Upernavik Isstrøm are -1.34 ± 0.03 , -1.01 ± 0.01 , and -0.60 ± 0.01 m yr⁻¹, respectively. The thinning rates of Zachariæ Isstrøm and
230 Nioghalvfjærdsfjorden are slower, i.e., mean SEC rates of -0.67 ± 0.02 , -0.28 ± 0.01 , and -0.21 ± 0.01 m yr⁻¹ at A, B, and C, respectively. Similarly, the closer to the grounding line, the faster the rate of thinning of the ice of those glaciers. King Christian X Land, located in the northeast of the GrIS, is a highly representative accumulation area. The ice velocity in this area is very small and there is no outflow glacier, and its change is driven mainly by SMB (Aschwanden et al., 2016; Velicogna et al., 2014). It shows weak continuous thickening over the entire observational period since 1991 (Fig. 6(d)). The rate of thickening
235 at A, B, and C is 0.13 ± 0.01 , 0.11 ± 0.01 , and 0.06 ± 0.01 m yr⁻¹, respectively. Whether continuous thinning or thickening, the rates do not remain constant, i.e., there are always periods of acceleration, deceleration or short-term fluctuation that are evident traces of the driving force of climate change on ice sheet change. For example, abnormal melting in 2012 and accumulation since 2016–2017, both driven by the North Atlantic Oscillation (NAO), are clearly visible in the time series of the above regions (Wood et al., 2021; Simonsen et al., 2021).

240 3.2 Ice sheet volume time series

Volume change of an ice sheet is an important parameter for determining the response of the ice sheet to the effects of climate change. A GrIS altimetric volume time series can be derived from our merged time series. In this study, the ICE-6G_D model (Peltier et al., 2018) and a scale factor (Groh et al., 2012) were used to correct for the effects of Glacial Isostatic Adjustment and elastic solid earth rebound. See Sect. 2.5 in Zhang et al. (2020) for the detailed calculation. The volume time series of the
245 entire GrIS and certain sub-regions estimated from our time series are shown in Figs. 7 and 8, respectively. Linear and quadratic



trend estimates can be inferred from the volume time series using a least-squares fitting model with a second-order polynomial and a sine wave with a 1 year period.

Over the entire GrIS, we detect an overall volume loss of $53.8 \pm 4.5 \text{ km}^3 \text{ yr}^{-1}$ with an acceleration in loss of $2.2 \pm 0.3 \text{ km}^3 \text{ yr}^{-2}$ during 1991–2020. Six of eight ice drainage systems show volume loss. The largest volume loss ($19 \pm 1.4 \text{ km}^3 \text{ yr}^{-1}$) and greatest acceleration in loss ($0.9 \pm 0.1 \text{ km}^3 \text{ yr}^{-2}$) are both from the ice sheet along the northwestern coast. Drainage systems located in central western and southwestern parts of the GrIS are the other two largest contributors to ice loss with volumes and rates of acceleration of $-10.2 \pm 1.3 \text{ km}^3 \text{ yr}^{-1}$ and $-0.5 \pm 0.1 \text{ km}^3 \text{ yr}^{-2}$, and $-10.2 \pm 1.9 \text{ km}^3 \text{ yr}^{-1}$ and $-0.6 \pm 0.1 \text{ km}^3 \text{ yr}^{-2}$, respectively. The only two drainage systems to show volume accumulation are located in central eastern and northeastern parts of the GrIS. However, their trends of volume accumulation are very weak, i.e., $1.3 \pm 0.8 \text{ km}^3 \text{ yr}^{-1}$ and $0.1 \pm 0.1 \text{ km}^3 \text{ yr}^{-2}$, and $-1.6 \pm 2.6 \text{ km}^3 \text{ yr}^{-1}$ and $0.5 \pm 0.2 \text{ km}^3 \text{ yr}^{-2}$, respectively.

In addition to studying the long-term trend of altimetric volume change of the ice sheet, our merged time series also provides detailed insight into small-scale fluctuations in volume change that reflect the effects of climate change on a temporal scale of up to 30 years. The evolution of ice sheet volume for the entire GrIS and certain sub-regions can be divided into different processes, as shown in Figs. 7 and 8, respectively. Before 1997, because of the contribution of various drainage systems in western Greenland (Fig. 8(a) and (e)–(h)), the GrIS presented rapid volume loss. Simonsen et al. (2021) thought that these ice losses were attributable mainly to the main outflow glaciers along the west coast. Then, the overall volume of the GrIS was alleviated, as also confirmed by the changes in the 5-year average SEC rates (see Fig. 6). This is attributed to increase of the SMB in the northeastern and central eastern drainage systems (Fig. 8(b) and (c)), and to reduction of ice discharge of outlet glaciers along the west coast (Fig. 8(g) and (h)). Subsequently, the GrIS entered a period of rapid ice loss because of the reduced SMB that was mostly attributable to meltwater runoff and increased ice discharge (Fig. 8(a) and (e)–(h)) (Simonsen et al., 2021; Shepherd et al., 2020; Velicogna et al., 2014). Then, all drainage systems entered another period of slowdown in ice loss. In fact, these processes are full of the traces of the effects of climate change. The rapid ice loss since 2003 was driven by the transition of the NAO from a high positive phase to a low-to-negative phase, which reduced SMB by enhancing melting and reducing snowfall, and accelerated ice discharge of outlet glaciers by driving warmer subsurface waters on the continental shelf (Bevis et al., 2019; Wood et al., 2021). The subsequent slowdown was because the NAO transitioned back to a more positive phase. It came from the anomalous increase in snowfall and anomalously low surface melting due to NAO-driven shifts in atmospheric forcing since 2016–2017 (Shepherd et al., 2020; Simonsen et al., 2021) and the slowed ice discharge attributable to NAO-driven shifts in oceanic forcing since 2010 (Wood et al., 2021). The weak signal of volume accumulation of drainage systems located in central eastern and northeastern parts of the GrIS (Fig. 8(b) and (c)) was also attributed to the two short-term abnormally increased snowfalls driven by the shift of the NAO, one in the early 2000s (Shepherd et al., 2020) and the other in the late 2010s (Simonsen et al., 2021). The volume of accumulated low-density snow exceeded that of lost high-density ice.



4 Uncertainty analysis

4.1 Error sources

280 In Sect.2.5, we described how to estimate the uncertainty of our results. In fact, the uncertainties of the SE time series (Fig. 6),
altimetric volume time series (Figs. 7 and 8), and average rates of SEC (Figs. 4 and 5) from our combined time series have
been given in the figures mentioned. Here, we discuss the sources of errors in our merged time series. The SE time series
developed in this study was based on ERS-1, ERS-2, Envisat, and CryoSat-2 radar altimeter observations. According to the
characteristics of the radar altimetry satellite data, the following factors will introduce errors into our results.

285 The first is the penetration of the signal into the surface snow, which causes a radar altimeter not to observe the actual surface
height of the ice sheet. Furthermore, surface processes such as melting, refreezing, and firn compaction might produce a new
reflecting surface that could result in errors. In this study, the time-variable penetration effects of the radar signal were
mitigated using the common strategy of including corrections for waveform parameters into the least-squares regression model
(see Eq. (1)), as performed in many previous studies (Flament and Remy, 2012; Sørensen et al., 2018; Zhang et al., 2020).

290 However, surface penetration suppresses noise induced by seasonal snowfall, making radar altimetric measurements more
relevant to mass change than those obtained from laser altimetry (Sørensen et al., 2018). Therefore, our time series obtained
by combining data from multiple altimetry missions is more suited to track dynamical processes and inter-annual or long-term
surface processes (Zhang et al., 2020; Simonsen et al., 2021).

The beam-limited footprint of a radar altimeter that has radius with scale of up to kilometres is another source of error. A
295 footprint with such a large radius makes it difficult for the radar altimeter to accurately measure ice surface height in areas
with complex terrain or drastic changes in elevation. Thus, the uncertainties of average SEC rates (Figs. 4 and 5) for marginal
areas with complex terrain are larger than those for the central ice sheet. This effect is also reflected in the estimation of
intermission bias and ascending–descending bias (Frappart et al., 2016; Zhang et al., 2020). In this study, official relocation
results in data products relocated by the point of closest approach were used to suppress this influence.

300 Additionally, cross-calibration between observations of different missions and interpolation or extrapolation of unobserved
cells will also introduce errors into the combined results. For the former, we used large amounts of data to fine-tune the
correction of intermission bias and ascending–descending bias for each grid cell, which is an approach proven to ensure better
self-consistency and reliability of the combined elevation time series (Zhang et al., 2020). For the latter, we used the EOF
reconstruction method to reduce interpolation error, because it can incorporate more temporal and spatial information to
305 constrain the interpolation results.

4.2 Comparison with airborne laser altimetry elevation

To validate our merged results, we first used the high-precision ATM L2 surface heights. Before performing a comparison, a
40 km floating median low-pass filter was applied to the ATM L2 data to eliminate outliers. Moreover, the mean height (at t_0)
of the centre of each grid cell h_0 was first added back into our merged GrIS SE anomaly time series to match the surface



310 heights. Then, we searched for all ATM L2 observations located within 2.5 km and a 10 d interval of each of the grid nodes
of our three-dimensional time series. The result of subtracting the elevation value of a grid node from the median of the ATM
L2 observations represents the difference for that location. The results of the validation are shown in Fig. 9(a). It can be seen
that the larger differences are concentrated primarily in steeper areas at the margins of the GrIS. This might be due to the poor
observation accuracy of radar altimeters in areas of complex terrain (Zhang et al., 2020). Another possible reason is that many
315 of them are interpolations or extrapolations. Over the GrIS, the median, RMS error, and the 10th and 90th percentile ranges
($P_{90}-P_{10}$) are -2.82 , 99.43 , and 145.25 m, respectively (Table 1), i.e., better than obtained for the 5 km interpolated grid cells
of the DEM of Slater et al. (2018), which are comparable to our calculations in terms of strategy and resolution (their median
and RMS error values were 25.4 and 138.6 m, respectively).

4.3 Comparison with airborne laser altimetry elevation changes

320 We also used ATM L4 SECs to evaluate our merged results. Similarly, a 40 km floating median low-pass filter was also used
to eliminate outliers in the ATM L4 SECs before performing the validation. The ATM L4 SECs are derived from every two
coincident ATM elevation measurements. We compared the ATM L4 data points with grids in our merged gridded time series
that lay within a 2.5 km radius and a 15 d interval of the observation instants of that point. Subsequently, the SE differences
and SEC differences between the ATM L4 observations and our merged time series at the same epochs were obtained, as
325 shown in Fig. 9(b) and (c). The spatial distribution patterns of SE differences and SEC differences are similar to those of the
SE differences mentioned in Sect. 4.2. The larger differences are distributed in areas with complex terrain at the margins of
the GrIS. The median, RMS error, and $P_{90}-P_{10}$ over the GrIS are also listed in Table 1. Overall, the median values of these
difference are both near 0, and the two $P_{90}-P_{10}$ values are both relatively small. Thus, although the RMS error of the SE
differences is larger than that of both Schröder et al. (2019) and Zhang et al. (2020) for the Antarctic Ice Sheet, our result is
330 still considered reliable. Furthermore, the integrity of ATM L4 data covering only the outlet glaciers of the West Antarctic Ice
Sheet and the Antarctic Peninsula Ice Sheet is limited.

4.4 Comparison with ESA GrIS Climate Change Initiative elevation changes

The ESA GrIS Climate Change Initiative (CCI) project has provided a dataset of SEC over the GrIS with a 5 year mean during
1992–2020 derived from ESA’s Ku-band radar satellite level-2 data products, which can be downloaded for free from
335 http://products.esa-icesheets-cci.org/products/details/cci_sec_2020.tar.gz/. Here, our results are verified through
intercomparison with that dataset. For consistent comparison, 5 year average SEC rates for the same observation epochs were
estimated from our time series using a least-squares fitting model with a first-order polynomial and a sine wave with a 1 year
period. Then, we compared the CCI SECs or our SECs with ATM L4 SECs at each grid node located within a 2.5 km radius.
To remove the influence of interpolation using EOF reconstruction, we only compared results that were not interpolated. The
340 median was also used to eliminate the influence of outliers.



Figure 10 shows the median, RMS error, and $P_{90}-P_{10}$ of the results of the intercomparison over the GrIS at 5 year intervals. It can be seen that our results are better in most periods, especially in those time intervals across the period of overlapping observations of Envisat and CryoSat-2 (i.e., from 2006–2010 to 2010–2014). Statistics of validation with GrIS CCI SECs listed in Table 2 also confirm this assertion. In comparison with the CCI SECs, the accuracy (RMS error) and dispersion of errors ($P_{90}-P_{10}$) of our results are improved by 19.3% and 8.9%, respectively, over all periods. In all periods from 2006–2010 to 2010–2014, the accuracy (RMS error) and dispersion of errors ($P_{90}-P_{10}$) of our results are improved by 30.9% and 19.0%, respectively. It might indicate that the effectiveness of our method for inter-mission bias correction for ERS-1 and ERS-2 has been reprocessed to align with Envisat by REAPER (Brockley et al., 2017).

5 Data availability

The surface elevation time series of the GrIS can be downloaded from the Big Earth Data Platform for Three Poles at <http://dx.doi.org/10.11888/Glacio.tpd.271658> (Zhang et al., 2021). In this repository, the time series is provided in NetCDF (.nc) format.

6 Conclusions

In this study, we developed a 30 year SE time series over the GrIS by combining ERS-1, ERS-2, Envisat, and CryoSat-2 satellite radar altimeter observations. A large number of operations, especially an updated plane-fitting least-squares regression strategy and an EOF reconstruction method, were performed to ensure that the time series has higher accuracy with monthly time resolution and 5×5 km spatial grid resolution. Validations with airborne laser altimetry observations and ESA GrIS CCI SECs indicated that our merged SE time series is reliable. In terms of the 5 year average SEC rates, the accuracy and dispersion of errors of our results were 19.3% and 8.9% higher than those of the CCI SECs, respectively. Benefiting from the finer correction of the inter-mission bias, the accuracy and dispersion of errors in our results were improved by up to 30.9% and 19.0%, respectively, in periods from 2006–2010 to 2010–2014.

The SECs and volume changes of the ice sheet are important variables that reflect the effects of climate change. As shown in Sect. 3, our data series can be used not only for studying long-term changes in the elevation and volume of the GrIS, but also for studying their temporal and spatial evolutions in detail on a temporal scale of up to 30 years. In particular, benefiting from the high temporal and spatial resolutions of our time series, the temporal and spatial evolution processes of ice loss from the main outflow glaciers in the GrIS can also be described in detail. These evolution processes are the response of the GrIS to oceanic and atmospheric changes driven by climate change. Thus, our merged time series provides an opportunity to examine the potential associations between ice sheet changes and climate forcing. The spatiotemporal patterns of accelerating or decelerating SEC of the GrIS, caused by shifts in atmospheric forcing and oceanic forcing driven by NAO phase transformation, reveal the sensitivity of the GrIS to climate forcing.



375 The mass balance of an ice sheet is a climate-related variable that has greater scientific value than elevation change. If combined with an appropriate ice density model, we could obtain a mass balance time series from our merged time series with much higher spatial resolution and longer temporal coverage than that of either GRACE or GRACE-FO. This could have advantages for studying mass change in small basins, especially the mass balance of outflow glaciers, thereby improving the estimation accuracy of the mass balance of the GrIS and reducing the uncertainty of projections of future sea level change.

Author Contributions.

BZ performed the calculation and wrote the manuscript. ZW contributed to the conception of the study. JA advised on validation and revised the manuscript. TL supervised the work. HG contributed to discussions and analysis of the results. All authors contributed to improvement of the manuscript.

380 **Competing interests.**

The authors declare that they have no conflict of interest.

Acknowledgements.

385 We would like to thank the organizations that shared their datasets and software for use in this study. The ERS-1, ERS-2, Envisat, and CryoSat-2 observations and the GrIS CCI SECs were provided by the European Space Agency and the airborne elevation data were provided by the National Snow and Ice Data Center. All geographical plots were produced using Generic Mapping Tools. We thank James Buxton MSc, from Liwen Bianji (Edanz) (www.liwenbianji.cn), for editing the English text of a draft of this manuscript.

Financial support.

390 This work was supported by the National Key Research and Development Program of China (2018YFC1406102), National Natural Science Foundation of China (42006184, 41941010), and Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA19070100).

References

- Adusumilli, S., Fricker, H. A., Siegfried, M. R., Padman, L., Paolo, F. S., and Ligtenberg, S. R. M.: Variable Basal Melt Rates of Antarctic Peninsula Ice Shelves, 1994–2016, *Geophys Res Lett*, 45, 4086–4095, doi:10.1002/2017gl076652, 2018.
- 395 Armitage, T. W. K., Wingham, D. J., and Ridout, A. L.: Meteorological Origin of the Static Crossover Pattern Present in Low-Resolution-Mode CryoSat-2 Data Over Central Antarctica, *IEEE Geoscience and Remote Sensing Letters*, 11, 1295–1299, doi:10.1109/lgrs.2013.2292821, 2014.



- Arthern, R. J., Wingham, D. J., and Ridout, A. L.: Controls on ERS altimeter measurements over ice sheets: Footprint-scale topography, backscatter fluctuations, and the dependence of microwave penetration depth on satellite orientation, *J Geophys Res-Atmos*, 106, 33471–33484, doi:10.1029/2001jd000498, 2001.
- Aschwanden, A., Fahnestock, M. A., and Truffer, M.: Complex Greenland outlet glacier flow captured, *Nature Communications*, 7, 10524, doi:10.1038/ncomms10524, 2016.
- Bamber, J. L., Gomez-Dans, J. L., and Griggs, J. A.: A new 1 km digital elevation model of the Antarctic derived from combined satellite radar and laser data - Part 1: Data and methods, *The Cryosphere*, 3, 101–111, doi:10.5194/tc-3-101-2009, 2009.
- Bevis, M., Harig, C., Khan, S. A., Brown, A., Simons, F. J., Willis, M., Fettweis, X., van den Broeke, M. R., Madsen, F. B., Kendrick, E., Caccamise, D. J., van Dam, T., Knudsen, P., and Nylen, T.: Accelerating changes in ice mass within Greenland, and the ice sheet's sensitivity to atmospheric forcing, *Proceedings of the National Academy of Sciences*, 116, 1934, doi:10.1073/pnas.1806562116, 2019.
- Brockley, D. J., Baker, S., Femenias, P., Martinez, B., Massmann, F.-H., Otten, M., Paul, F., Picard, B., Prandi, P., Roca, M., Rudenko, S., Scharroo, R., and Visser, P.: REAPER: Reprocessing 12 Years of ERS-1 and ERS-2 Altimeters and Microwave Radiometer Data, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 5506–5514, 10.1109/tgrs.2017.2709343, 2017.
- Chambers, D. P., Mehlhoff, C. A., Urban, T. J., Fujii, D., and Nerem, R. S.: Low-frequency variations in global mean sea level: 1950–2000, *Journal of Geophysical Research: Oceans*, 107, 1-1-1-10, doi:10.1029/2001JC001089, 2002.
- Ewert, H., Groh, A., and Dietrich, R.: Volume and mass changes of the Greenland ice sheet inferred from ICESat and GRACE, *J Geodyn*, 59–60, 111–123, 10.1016/j.jog.2011.06.003, 2012.
- Flament, T. and Remy, F.: Dynamic thinning of Antarctic glaciers from along-track repeat radar altimetry, *J Glaciol*, 58, 830–840, 10.3189/2012JG11J118, 2012.
- Frappart, F., Legrésy, B., Niño, F., Blarel, F., Fuller, N., Fleury, S., Birol, F., and Calmant, S.: An ERS-2 altimetry reprocessing compatible with ENVISAT for long-term land and ice sheets studies, *Remote Sensing of Environment*, 184, 558–581, doi:10.1016/j.rse.2016.07.037, 2016.
- Groh, A., Ewert, H., Scheinert, M., Fritsche, M., Rülke, A., Richter, A., Rosenau, R., and Dietrich, R.: An investigation of Glacial Isostatic Adjustment over the Amundsen Sea sector, West Antarctica, *Global Planet Change*, 98-99, 45–53, 10.1016/j.gloplacha.2012.08.001, 2012.
- Hanna, E., Mernild, S. H., Cappelen, J., and Steffen, K.: Recent warming in Greenland in a long-term instrumental (1881–2012) climatic context: I. Evaluation of surface air temperature records, *Environmental Research Letters*, 7, 045404, doi:10.1088/1748-9326/7/4/045404, 2012.
- Helm, V., Humbert, A., and Miller, H.: Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8, 1539–1559, doi:10.5194/tc-8-1539-2014, 2014.
- Hurkmans, R. T. W. L., Bamber, J. L., Sørensen, L. S., Joughin, I. R., Davis, C. H., and Krabill, W. B.: Spatiotemporal interpolation of elevation changes derived from satellite altimetry for Jakobshavn Isbræ, Greenland, *Journal of Geophysical Research: Earth Surface*, 117, doi:10.1029/2011JF002072, 2012.
- Jin, T., Li, J., Jiang, W., and Chu, Y.: Low-frequency sea level variation and its correlation with climate events in the Pacific, *Chinese Science Bulletin*, 57, 3623–3630, doi:10.1007/s11434-012-5231-y, 2012.
- Joughin, I., Abdalati, W., and Fahnestock, M.: Large fluctuations in speed on Greenland's Jakobshavn Isbræ glacier, *Nature*, 432, 608–610, doi:10.1038/nature03130, 2004.
- Khazendar, A., Fenty, I. G., Carroll, D., Gardner, A., Lee, C. M., Fukumori, I., Wang, O., Zhang, H., Seroussi, H., Moller, D., Noël, B. P. Y., van den Broeke, M. R., Dinardo, S., and Willis, J.: Interruption of two decades of Jakobshavn Isbræ acceleration and thinning as regional ocean cools, *Nat Geosci*, 12, 277–283, doi:10.1038/s41561-019-0329-3, 2019.
- Krabill, W., Hanna, E., Huybrechts, P., Abdalati, W., Cappelen, J., Csatho, B., Frederick, E., Manizade, S., Martin, C., Sonntag, J., Swift, R., Thomas, R., and Yungel, J.: Greenland Ice Sheet: Increased coastal thinning, *Geophys Res Lett*, 31, L24402, 10.1029/2004GL021533, 2004.
- McMillan, M., Shepherd, A., Sundal, A., Briggs, K., Muir, A., Ridout, A., Hogg, A., and Wingham, D.: Increased ice losses from Antarctica detected by CryoSat-2, *Geophys Res Lett*, 41, 3899–3905, doi:10.1002/2014gl060111, 2014.
- McMillan, M., Leeson, A., Shepherd, A., Briggs, K., Armitage, T. W. K., Hogg, A., Kuipers Munneke, P., van den Broeke, M., Noël, B., van de Berg, W. J., Ligtenberg, S., Horwath, M., Groh, A., Muir, A., and Gilbert, L.: A high-resolution record of Greenland mass balance, *Geophys Res Lett*, 43, 7002–7010, doi:10.1002/2016GL069666, 2016.
- Meloni, M., Bouffard, J., Parrinello, T., Dawson, G., Garnier, F., Helm, V., Di Bella, A., Hendricks, S., Ricker, R., Webb, E., Wright, B., Nielsen, K., Lee, S., Passaro, M., Scagliola, M., Simonsen, S. B., Sandberg Sørensen, L., Brockley, D., Baker, S., Fleury, S., Bamber, J., Maestri, L., Skourup, H., Forsberg, R., and Mizzi, L.: CryoSat Ice Baseline-D validation and evolutions, *The Cryosphere*, 14, 1889–1907, doi:10.5194/tc-14-1889-2020, 2020.
- Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A., and Paden, J.: Fast retreat of Zachariae Isstrøm, northeast Greenland, *Science*, 350, 1357, doi:10.1126/science.aac7111, 2015.
- Nilsson, J., Gardner, A., Sørensen, L. S., and Forsberg, R.: Improved retrieval of land ice topography from CryoSat-2 data and its impact for volume-change estimation of the Greenland Ice Sheet, *The Cryosphere*, 10, 2953–2969, doi:10.5194/tc-10-2953-2016, 2016.




- Paolo, F. S., Fricker, H. A., and Padman, L.: Constructing improved decadal records of Antarctic ice shelf height change from multiple satellite radar altimeters, *Remote Sensing of Environment*, 177, 192–205, doi:10.1016/j.rse.2016.01.026, 2016.
- 455 Peltier, W. R., Argus, D. F., and Drummond, R.: Comment on “An Assessment of the ICE-6G_C (VM5a) Glacial Isostatic Adjustment Model” by Purcell et al, *Journal of Geophysical Research: Solid Earth*, 123, 2019–2028, doi:10.1002/2016JB013844, 2018.
- Remy, F., Legresy, B., and Benveniste, J.: On the Azimuthally Anisotropy Effects of Polarization for Altimetric Measurements, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 3289–3296, doi:10.1109/tgrs.2006.878444, 2006.
- 460 Schröder, L., Horwath, M., Dietrich, R., Helm, V., van den Broeke, M. R., and Ligtenberg, S. R. M.: Four decades of Antarctic surface elevation changes from multi-mission satellite altimetry, *The Cryosphere*, 13, 427–449, doi:10.5194/tc-13-427-2019, 2019.
- Schröder, L., Richter, A., Fedorov, D. V., Eberlein, L., Brovkvov, E. V., Popov, S. V., Knöfel, C., Horwath, M., Dietrich, R., Matveev, A. Y., Scheinert, M., and Lukin, V. V.: Validation of satellite altimetry by kinematic GNSS in central East Antarctica, *The Cryosphere*, 11, 1111–1130, doi:10.5194/tc-11-1111-2017, 2017.
- 465 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophys Res Lett*, 46, 8174–8183, doi:10.1029/2019GL082182, 2019.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlström, A., Babonis, G., Barletta, V., Blazquez, A., Bonin, J., Csatho, B., Cullather, R., Felikson, D., Fettweis, X., Forsberg, R., Gallee, H., Gardner, A., Gilbert, L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mouginit, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noel, B., Otosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg-Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem, M., Vishwakarma, B. D., Wiese, D., Wouters, B., and The, I. t.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, doi:10.1038/s41586-018-0179-y, 2018.
- 475 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlström, A., Babonis, G., Barletta, V. R., Björk, A. A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., Cullather, R., Engdahl, M. E., Felikson, D., Fettweis, X., Forsberg, R., Hogg, A. E., Gallee, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P. L., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., 480 Mottram, R., Mouginit, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Otosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem, M., Vishwakarma, B. D., Wiese, D., Wilton, D., Wagner, T., Wouters, B., Wuite, J., and The, I. T.: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, doi:10.1038/s41586-019-1855-2, 2020.
- 485 Simonsen, S. B. and Sorensen, L. S.: Implications of changing scattering properties on Greenland ice sheet volume change from Cryosat-2 altimetry, *Remote Sensing of Environment*, 190, 207–216, doi:10.1016/j.rse.2016.12.012, 2017.
- Simonsen, S. B., Barletta, V. R., Colgan, W. T., and Sørensen, L. S.: Greenland Ice Sheet Mass Balance (1992–2020) From Calibrated Radar Altimetry, *Geophys Res Lett*, 48, e2020GL091216, doi:10.1029/2020GL091216, 2021.
- 490 Slater, T., Shepherd, A., Mcmillan, M., Armitage, T. W. K., Otosaka, I., and Arthern, R. J.: Compensating Changes in the Penetration Depth of Pulse-Limited Radar Altimetry Over the Greenland Ice Sheet, *IEEE Transactions on Geoscience and Remote Sensing*, 57, 9633–9642, doi:10.1109/TGRS.2019.2928232, 2019.
- Slater, T., Shepherd, A., McMillan, M., Muir, A., Gilbert, L., Hogg, A. E., Konrad, H., and Parrinello, T.: A new digital elevation model of Antarctica derived from CryoSat-2 altimetry, *The Cryosphere*, 12, 1551–1562, 10.5194/tc-12-1551-2018, 2018.
- 495 Smith, T. M., Reynolds, R. W., Livezey, R. E., and Stokes, D. C.: Reconstruction of Historical Sea Surface Temperatures Using Empirical Orthogonal Functions, *J Climate*, 9, 1403–1420, doi:10.1175/1520-0442(1996)009<1403:ROHSST>2.0.CO;2, 1996.
- Sørensen, L. S., Simonsen, S. B., Forsberg, R., Khvorostovsky, K., Meister, R., and Engdahl, M. E.: 25 years of elevation changes of the Greenland Ice Sheet from ERS, Envisat, and CryoSat-2 radar altimetry, *Earth and Planetary Science Letters*, 495, 234–241, doi:10.1016/j.epsl.2018.05.015, 2018.
- 500 Straneo, F. and Heimbach, P.: North Atlantic warming and the retreat of Greenland's outlet glaciers, *Nature*, 504, 36–43, doi:10.1038/nature12854, 2013.
- Velicogna, I., Sutterley, T. C., and van den Broeke, M. R.: Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data, *Geophys Res Lett*, 41, 8130–8137, 10.1002/2014GL061052, 2014.
- Wahr, J., Swenson, S., and Velicogna, I.: Accuracy of GRACE mass estimates, *Geophys Res Lett*, 33, doi:10.1029/2005GL025305, 2006.
- 505 Wood, M., Rignot, E., Fenty, I., An, L., Björk, A., van den Broeke, M., Cai, C., Kane, E., Menemenlis, D., Millan, R., Morlighem, M., Mouginit, J., Noël, B., Scheuchl, B., Velicogna, I., Willis, J. K., and Zhang, H.: Ocean forcing drives glacier retreat in Greenland, *Science Advances*, 7, eaba7282, doi:10.1126/sciadv.aba7282, 2021.
- Yang, Y. D., Li, F., Hwang, C., Ding, M. H., and Ran, J. J.: Space-Time Evolution of Greenland Ice Sheet Elevation and Mass From Envisat and GRACE Data, *J Geophys Res-Earth*, 124, 2079–2100, doi:10.1029/2018jf004765, 2019.




- 510 Zhang, B., Wang, Z., An, J., Liu, T., and Geng, H.: Surface elevation time series over the Greenland Ice Sheet (1991–2020), National Tibetan Plateau Data Center [dataset], doi:10.11888/Glacio.tpd.c.271658, 2021.
- Zhang, B., Wang, Z., Li, F., An, J., Yang, Y., and Liu, J.: Estimation of present-day glacial isostatic adjustment, ice mass change and elastic vertical crustal deformation over the Antarctic ice sheet, *J Glaciol*, 63, 703–715, doi:10.1017/jog.2017.37, 2017.
- Zhang, B., Wang, Z., Yang, Q., Liu, J., An, J., Li, F., Liu, T., and Geng, H.: Elevation Changes of the Antarctic Ice Sheet from Joint Envisat and CryoSat-2 Radar Altimetry, *Remote Sensing*, 12, doi:10.3390/rs12223746, 2020.
- 515 Zwally, H. J., B., G. M., Beckley, M. A., and Saba, J. L.: Antarctic and Greenland Drainage Systems, GSFC Cryospheric Sciences Laboratory, http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php, 2012.
- Zwally, H. J., Giovinetto, M. B., Li, J., Cornejo, H. G., Beckley, M. A., Brenner, A. C., Saba, J. L., and Yi, D. H.: Mass changes of the Greenland and Antarctic ice sheets and shelves and contributions to sea-level rise: 1992–2002, *J Glaciol*, 51, 509–527, doi:10.3189/172756505781829007, 2005.


520




525 **Figure 1:** (a) SE of the GrIS DEM at reference time 2010.0, and (b) surface slope and (c) aspect derived from the DEM.




530 **Figure 2:** Percentage of variance explained (red line) and cumulative variance explained (blue line) by each EOF mode; RMS error (green line) and its derivative (purple line) of the difference between the reconstructed time series of monthly elevation change and that of observation in different EOFs.




535 **Figure 3:** Interpolation performance of EOF reconstruction and ordinary kriging: (a), (b), and (c) are the results for March 1992 observation, EOF reconstruction, and ordinary kriging interpolation, respectively, and (d), (e), and (f) are the same for April 1992.



540 **Figure 4:** Maps of long-term SECs and their uncertainties from the combined elevation time series over the periods of (a) and (e) 1991–2000, (b) and (f) 2001–2010, (c) and (g) 2011–2020, and (d) and (h) 1991–2020.



545 **Figure 5:** Maps of long-term SECs and their uncertainties from the combined elevation time series over the periods of (a) and (g), 1991–1995, (b) and (h) 1996–2000, (c) and (i) 2001–2005, (d) and (j) 2006–2010, (e) and (k) 2011–2015, and (f) and (l) 2016–2020.



550 **Figure 6:** SE anomaly time series near (a) Jakobshavn Isbræ, (b) Upernavik Isstrøm, (c) Zachariae Isstrøm and
 Nioghalvfjærdsfjorden glacier, and (d) King Christian X Land. The locations of the selected points (A, B, and C) are marked
 in the left-hand maps of elevation change over 1991–2020. The time series and the 1σ uncertainty ranges for each point are
 given in the right-hand plots with the time series of points A and B shifted along the SE anomaly axis for better visibility.

555





Figure 7: Volume change of the entire GrIS south of 81.5°N from our merged altimetric time series (green dots) and its corresponding result after removing seasonal oscillations using a 13 month moving average (blue solid curve). The solid red line is the best-fit quadratic curve for the linear and quadratic trend estimates of volume change. The grey error bars show the 1 σ uncertainty range of the altimetry data. The red shaded area in the inset indicates the coverage of the GrIS with reference to the Greenland drainage system boundaries in Zwally et al. (2012).



565 **Figure 8:** Volume change of sub-regions south of 81.5°N from our merged altimetric time series (green dots) and their corresponding results after removing seasonal oscillations using a 13 month moving average (blue solid curves). The solid red lines are the best-fit quadratic curves for the linear and quadratic trend estimates of volume change. The grey error bars show the 1σ uncertainty range of the altimetry data. The red shaded area in the inset of each panel indicates the coverage of the sub-region with reference to the Greenland drainage system boundaries in Zwally et al. (2012).

570




Figure 9: Validation with ATM laser altimeter observations: (a) differences between SE derived from ATM L2 and our merged time series, (b) SE differences, and (c) SEC differences between ATM L4 observations and our merged time series at the same epochs.

575




Figure 10: Validation with GrIS CCI SECs: (a) median, (b) RMS error, and (c) $P_{90}-P_{10}$ of the SEC differences between CCI (orange) and those derived from our merged time series (green) at 5 year intervals during 1992–2019.

580 **Table 1.** Statistics of the results of validation with ATM laser altimeter observations. The median, RMS error, and $P_{90}-P_{10}$ of the biases are given.



Validation data		Statistics of the comparison with validation data		
Data	Variable	Median	RMS	P ₉₀ – P ₁₀
ATM L2	SE (m)	-2.82	99.43	145.25
ATM L4	SE differences (m)	-0.02	8.84	3.78
ATM L4	SECs (m yr ⁻¹)	-0.00	3.63	0.95

Table 2. Statistics of the results of validation with GrIS CCI SECs. The median, RMS error, and P90–P10 of the biases are given.

	Statistics of the comparison with ATM L4 SECs					
	All Periods			Periods across the Envisat and CryoSat-2 connections		
	Median (m yr ⁻¹)	RMS (m yr ⁻¹)	P ₉₀ – P ₁₀ (m yr ⁻¹)	Median (m yr ⁻¹)	RMS (m yr ⁻¹)	P ₉₀ – P ₁₀ (m yr ⁻¹)
GrIS CCI	-0.05	0.57	0.79	-0.04	0.81	1.16
This study	-0.03	0.46	0.72	-0.09	0.56	0.94

585