

Reply to reviewer 1 of 'EMO-5: A high-resolution multi-variable gridded meteorological data set for Europe' by Vera Thiemig et al.

Dear Reviewer,

we are please to share with you that we have gone through all your comments and addressed them. Please find below our reply to each major point you raised.

Major points

1. The introduction is too short. Greater effort is required on scene setting including a discussion of the role and limitations of both reanalysis and satellite observations to provide the reader with necessary context to then properly interpret the paper. Indeed, there is in general a lack of sufficient acknowledgement of prior and ongoing valuable work in this area across the submitted manuscript as a whole to provide the reader with necessary context to properly and fairly evaluate the value of your work. For example, ECA&D and the associated E-OBS gridded product served via the C3S CDS should be noted much more prominently and far earlier and calls into substantial question your assertion on lines 39-41. I would be hard pressed to defend the statement you make there given the availability of the well documented and highly utilised E-OBS product. Similarly, the efforts under C3S to collate station data as documented in e.g. Noone et al., 2021 warrants more attention than is currently given. Failing to acknowledge these – Copernicus funded – efforts just gives an impression of dysfunctionality across Copernicus program activities as well as – for those with the knowledge – giving the distinct impression of over-selling what you have produced and doing so in glorious isolation. This is disappointing as a knowledgeable reviewer and equally does not bode well for realising programmatic synergies across Copernicus services in the new Copernicus budgetary cycle activities.

The paragraph starting at line 55 should come earlier and be expanded. Then in numerous places comparisons to existing products / approaches should be made. For example, the QC method in 3.1 should make reference to QC procedures of EOBS, GHCND and HadISD at a minimum.

Reply to point 1:

Despite that we do not think that the review of a scientific paper is a discussion platform for Copernicus internal management matters, we recognise that this is truly a concern for the reviewer. Therefore we would like to reassure him (her) that there is good collaboration ongoing between the various Copernicus core services and that we seek synergies and collaborations wherever possible. In general, the dialogue between CEMS and C3S has always existed, and efforts on mutual-beneficial solutions have been a focus. In fact, this very manuscript was shared with the entrusted entity of C3S some months ago, and was

only submitted after addressing all their comments. Hence, there is transparency and collaboration established between C3S and CEMS, including in this matter.

It might on first glance appear like a service duplication within Copernicus, however the meteorological data collection services of C3S and CEMS serve two very different needs, which principal their requirements and result into two very different products, each with their rightfulness to exist. On the one hand, CEMS needs an operational real-time dataset that contains as much valid information as available at near real-time to compute the initial state of the system, and in addition some decades of historical data created with the same method as the real-time data to allow calibration and validation of the operational system and consistency between the calibration of the model and the operational running of the forecast. On the other hand, C3S serves the community with a climatological dataset, and hence their data products need to be of sufficient length, consistency, and continuity to determine climate variability and climate change. It appears that this distinction was not clear and had caused the reviewer's concerns.

Some of the reviewer's requests, such as e.g. the wish for mentioning ECA&D and E-OBS earlier than in line 39 and more prominently as well as include those in product/method comparison seemed arbitrary and at times hints a conflict of interest.

Nevertheless, we have put great effort into the rewriting of the introduction to improve the scene setting as the reviewer calls it to assure that the reason, distinction and use of EMO-5 is crystal clear.

1 Introduction

The (timely) availability of meteorological data and their quality determine in many cases the capacity of environmental modelling. Easy accessibility to quality-controlled data, with good coverage and spatial resolution, can provide a solid foundation for various environmental modelling applications, such as e.g. hydrological or agricultural models.

One prominent example that relies heavily on good quality meteorological input data provided in near real-time is the European Flood Awareness System (EFAS), which is part of the Emergency Management Service (EMS) of Copernicus, the EU's Earth observation programme. EFAS provides a flood monitoring and forecast service for riverine and flash floods across the whole of Europe. The forecasts of EFAS are calculated using the semi-distributed hydrological rainfall-runoff model LISFLOOD (<https://ec-jrc.github.io/lisflood/>) which relies heavily on quality-controlled, (sub-)daily meteorological information on precipitation, temperature, wind speed, solar radiation and water vapour pressure for a) model calibration and validation as well as for b) the computation of initial conditions during the operational running. In 2006, during the set-up phase of EFAS for pre-operational running there was an imminent need for a pan-European, quality-controlled, high-resolution, multi-variable, near real-time as well as historical meteorological data set. However, despite the existence of a fairly good coverage network of in situ stations across Europe, at that time, there existed no overarching service that collected in near real-time all those meteorological in situ data across the entire European domain. For this reason, the Joint

Research Centre started in 2006 with the collection, quality control and gridding of real-time and historic (from 1970) meteorological data across Europe and neighbouring regions, a service that became in 2012 known as the Copernicus EMS Meteorological Data Collection Centre (MDCC).

The CEMS MDCC runs around the clock and produces daily near real-time meteorological grids which are used in the operational running of not only EFAS, but also by two other major CEMS services, namely the European Forest Fire Information System (EFFIS; <https://effis.jrc.ec.europa.eu/>; San-Miguel, J. et al., 2019) and the European Drought Observatory (EDO; <https://edo.jrc.ec.europa.eu/>; Spinoni et al., 2016; Cammalleri, C. et al., 2020). At the same time, the MDCC collects also historical data in an offline mode and feeds those into the MDCC data collection for the production of historical meteorological grids. This is important as historical data (produced with the same method) are indispensable for the calibration of the various models and indicators. While the service produces daily updated grids (EMO-5 operational grids), we have re-run our archive from 1990-2019 to produce a new long-term dataset, which we refer to as version 1 of the EMO-5. EMO-5 (version 1) comprises daily 5x5 km grids for six variables - **precipitation, minimum and maximum air temperature, wind speed, solar radiation and water vapour pressure** - and additional 6-hourly grids for precipitation and **mean air temperature**. The underlying data comes from 21 data providers that contributed to the MDCC data collection, by sharing data from a total of about 18,694 in situ stations across Europe and five gridded data sets over selected areas. For the in situ stations, it should be noted that while some stations measure multiple variables, others measure only one or two. Furthermore, some stations provided data for the entire archive period from 1970, while others only for a limited time-period. The gridded data sets have been added to the MDCC data collection, in order to improve the quality of the resulting meteorological grids by increasing the information density over selected areas, mainly data scarce areas and areas with complex topography.

Throughout the last decade, many other observational meteorological data grids emerged, such as: E-OBS (Haylock et al., 2008; Cornes et al., 2018) for the whole of Europe; seNorge2 and seNorge2018 for Norway (Lussana et al., 2018); SPREAD for Spain (Serrano-Notivolli et al., 2017); ZAMG-INCA for Austria (Haiden et al., 2011); CombiPrecip for Switzerland (Sideris et al., 2014); CarpatClim for Hungary, Serbia, Romania, Ukraine, Slovakia, Poland, Czech Republic and Croatia (Antolović et al., 2013; Spinoni et al., 2015); EURO4M-APGD for the European Alps and adjacent flatland regions (Isotta et al. 2014); SAFRAN for France, Spain and Tunisia (Quintana-Seguí et al., 2008; Vidal et al., 2010; Quintana-Seguí et al., 2017; Trambly et al., 2019). However, despite the availability of these data sets and their immense value each of them holds, EMO-5 represents a uniquely valuable resource due to a combination of its pan-European coverage, near real-time production, high temporal (6-hourly and daily) and spatial (5x5 km) resolution, large amount of input data (18,694 in situ stations and five high resolution regional observational grids), seven different variables (i.e. precipitation, minimum, maximum, and mean air temperature, wind speed, solar radiation and water vapour pressure), and a long historical data record (from 1 January 1990). These characteristics combine to make EMO-5, to our knowledge, the most complete gridded

multi-variable observational near-real time meteorological data set covering the whole of Europe (and peripheral areas).

The aim of this paper is to present EMO-5 (version 1) and its potential usage, by providing an insight into its data sources, the applied methods and the quality of the resulting data set. In this paper, the evaluation of the resulting grid quality is focused mainly on the gridded precipitation data, as precipitation is the most crucial driver for hydrological modelling (our main focus). However, we invite the scientific community to expand the evaluation exercise beyond EMO-5 gridded precipitation.

Based on the information provided in this paper, the scientific community will understand that EMO-5 (version 1 as well as the operational grids) is not a climatological dataset (unlike e.g. the Essential Climate variables produced by C3S, see Noone et al., 2021), but an operational data set based on the maximum amount of quality-controlled information available at any given time. The implications of this are, that we do not advice to use this dataset to determine climate variability and climate change, however any other environmental application, especially those with real-time, high spatial resolution or multi-variable needs is likely to benefit from this data set. Hence, by making the EMO-5 (version 1) data publicly available, we aim to support many other environmental applications and services that would benefit from using those data, such as e.g. hydrological, agricultural or other environmental applications.

The remainder of this paper is organised as outlined in the following. The source data are described in Section 2. The entire workflow of the grid creation, including the quality control criteria applied during the data collection and an evaluation of various interpolation methods, are described in Section 3. Data access information are given in Section 4. An evaluation of the grid quality for precipitation is described in Section 5, and finally, some conclusions are presented in Section 6, followed by a future outlook.

2. The reason why the analysis is limited to 1970 needs to be made far more explicit. There are many data that extend back further than 1970 and the choice of 1970 is arbitrary with no clear rationale given to justify the choice. From an application perspective there would be considerable value to extending the analysis to earlier periods to the extent possible.

Given that the product is reputedly available from 1970 onwards it is problematic that Figures 1-3 only show availability post-1990. This gives the impression that you may be wishing to hide / obscure availability of data in the first 20 years of the series. The figures should show 1970 onwards or the product should be cut at 1990 and published as such. Later at the end of section 3 you suggest this is the case. Please clarify whether the product extends 1970 to date or 1990 to date and ensure consistently stating so throughout the paper.

Reply to point 2:

The frame in which the meteorological data are collected is described in the introduction (Section 1, lines 32-54): to serve the need of the CEMS, and in particular EFAS. EFAS requires a good coverage of observations for the most recent decades (to calibrate and validate our hydrological model) as well as real time data (for the initialisation of the hydrological model used for operational forecasting). For this reason, we are not collecting earlier data (before 1970). Also, earlier data is already covered by C3S, and hence would be a duplication within the Copernicus service.

With regards to the second paragraph of your comment: In line 48 it is specified that the MDCC is collecting meteorological data from 1970 onwards. This is specifying the time frame of the collection of the in situ meteorological data under CEMS, but not the starting time of the product (EMO-5), which is explicitly mentioned throughout the manuscript as 1990, see:

- abstract (lines 23-24): *“EMO-5 (release 1) covers the time period from 1990 to 2019,[...]”*
- introduction (line 65): *“[...] (from 1 January 1990).”*
- Section 2 (line 123): *“For each of the seven EMO-5 meteorological variables, the location (and hence density) of the input data, as well as the record length per station and the number of input stations over time (1990-2019),”*
- Figure 1-3
- Section 3.3.6 (lines 322-323): *“The EMO-5 version which will be available on the JRC Data Catalogue has been produced for the whole length of the archive from 1990 till 2019, and it is foreseen to extend this dataset on a monthly base.”*
- Section 6 (line 467): *“EMO-5 covers the time period from 1990 onwards [...]”*

Despite the consistent reporting of the starting time of EMO-5, we recognise that the sentence in Section 3.3.6 (lines 322-323) and in particular *“for the whole length of the archive”* might have caused the confusion, for which reason we modified it to: *“The EMO-5 (version 1) dataset has been produced for the time period from 1990 till 2019.”*

3. The methodology in almost all aspects needs to be strengthened and much more explicit. The method must be outlined to the extent that the analysis could be to first order independently reproduced based upon the description. This is necessary and a central tenet of the scientific method itself. Please revise the method to be considerably more comprehensive such that based upon the method description an independent researcher may reasonably be able to recreate your approach.

Reply to point 3:

We have checked this point, but we fail to see where the methodology is not described in enough detail to be reproducible. Obviously the input data are not provided and neither the code, but beside those we have described all that is needed to reproduce the methodology including the variable-specific data validation rules, the interpolation algorithm, the station selection criteria, the variable specific aggregation periods and any other variable specific information that is needed.

It has also not been a comment of the other two reviewers, so it is difficult for us to understand where an expansion would be required in the opinion of the reviewer.

4. The validation activity would be stronger if it could be shown in more detail for all variables and would, undoubtedly, serve to increase user uptake. However, the only real validation in the strict scientific sense is actually the leave-one-out analysis which is not even contained within the validation section at all. What is presently classed as being validation is actually characterisation of the product. Validation requires a value that is of known quality to compare against and none of the three sub-sections in the section purportedly to do with validation actually undertake such a comparison. That section should be retitled to Characterisation accordingly. Then the leave-one-out validation should be considerably expanded, shown for numerous parameters, and aggregated across well-sampled and sparsely-sampled regions separately.

Reply to point 4:

The leave-one-out validation for all variables is in Table 4 and is explicitly mentioned in line 159. It is referenced in Section 3.2 in which we determine the optimal interpolation scheme for this operational, real-time data set. We have considered moving it under the validation section, but as the choice of the interpolation scheme is part of the methodology we see benefits for leaving it in Section 3.2.

As it wasn't obvious that Table 3 contained the leave-one-out analysis we extended the table caption to: *"Summary of the error measures for the three interpolation schemes based on the leave one out analysis. (Best values are in bold)"*

We do not have a section called "Validation", only called "Evaluation" in which we evaluate the product from various angles, hence we don't think renaming is necessary.

5. In section 2 the input data is outlined to a very perfunctory level – greater detail here would help including a breakdown by source, data policy etc and links to sources where available publicly by elevating Table A1 to the main text and augmenting with additional information. Also, the maps in Figures 1 to 3 conflate station and gridded input in ways that are really inaccessible. Maps of just primary station source by time and / or variable would be more accessible and avoid proverbially mixing apples and oranges together.

Reply to point 5:

As the reviewer pointed out correctly, specifications of the input data for EMO-5 can be found in Section 2 and in Appendix 1. The reviewer can find the requested breakdown of sources in Table A1 (line 705), which shows not only the full list of data providers (as requested by the reviewer), but also specifies the variables and the number of stations being collected from each data provider. The table does not contain web links to the data provider as we do not consider it a good practise, as for ones those web links outdate quickly and each data provider can easily be found through using any search engine. Also, CEMS MDCC does not use weblinks for collecting those data, as those are not scraped from

the web, but rather provided directly to us by the data providers listed in Table A1. As the reviewer suggested under point 15 to transfer multi-page figures in the Appendix for their length, we believe that with the same reasoning also Table A1 as a multi-page table should remain in the Appendix. However, if there is a strong preference by the reviewer we have no problems with transferring it to the main text.

We have added information on the data policy at the end of Section 2: *“All data collected by the MDCC, are covered by the EEA-EUMETNET Public Duty License Agreement or the EEA non-EUMETNET Partner license Agreement, which means those data are share-able between the Copernicus services.”*

To Figure 1-3: As specified in the Figure captions and in the main text (lines 121ff), they show for each of the EMO-5 variable, the station location, record length per station and the number of stations over time. It is a common practise to show multiple data (sources) in one image as we have done. The differentiation between those was done by colour coding the different data sources, hence there is no mixing of apple and pears.

Worse, Section 2 is also completely silent on sharing of these station data onwards to e.g. the World Data Centre for Meteorology at NOAA NCEI or the C3S in-situ database effort or ECA&D and thus E-OBS. This limits the potential utility of this data collection activity if you are not actively trying to share data where you can with the activities being undertaken to improve access globally to meteorological holdings, many of which are funded by Copernicus. Where is the joined up thinking? How does this serve Copernicus if you are not actively sharing across Copernicus Services?

As CEMS is not the owner of those in situ station data we do not have the mandate to distribute them further, with the only exception to other Copernicus services, as the data has been collected under the EUMETNET data licence agreement, which permits that. On that note, we can assure the reviewer that there is an open and strong collaboration between CEMS and the other Copernicus core services, and in particular with C3S to streamline efforts and to avoid duplications. There have been multiple meetings between CEMS-C3S on this and related topics in the past and are pleased to share also with the reviewer that we do support the sharing of those data within the Copernicus services. We do not need to add this information to the manuscript, as the fact that those data are collected under the EUMETNET data licence agreement, does tell the relevant parties (if they don't already know) that those data is sharable within the Copernicus Services.

6. The quality control is a little limited, in particular in not including any form of neighbour buddy checking as is state of the art in e.g. GHCND or HadISD. Either that or, between the text and the table, the presence of buddy checks is obscured. You should at a minimum be explicit that the applied checks consist of a minimum set of record-based, logic and spike-based checks and do not include a number of other checks including repeat strings, distributional checks, frequent value checks and neighbour-based checks as is the case in state-of-the-art QC procedures such as GHCND or HadISD. You should aim to incorporate a broader array of such checks in future.

Reply to point 6:

All quality control checks that are currently performed by the CEMS MDCC on the operationally collected meteorological data (and hence also EMO-5) are explicitly mentioned in the “Quality control on input data” section (Section 3.1). We can confirm to the reviewer that there have been no additional checks obscured.

As also to the knowledge of the reviewer, an expansion of the quality control framework of the MDCC has been foreseen and is currently under development, and will be part of the next version of EMO-5. In fact, this is mentioned in Section 6 under “Conclusions and future work” (see lines 500ff): “[...] to improve the current quality control framework, new data validation rules, such as spatial comparison with neighbouring stations or additional statistical checks, will be implemented.”

The reviewer has to remember that EMO-5 is created in an operational service to serve a particular purpose. Due to the potential of this data set, we have decided to make it publically available as we see it our duty given that it is a Copernicus product and hence shall be free and open. The QC might be in the opinion of the reviewer limited, but it shall not be forgotten that this data set has been used very successfully for the past decade for the calibration and initialisation of the hydrological model underlying the European Flood Awareness System.

Furthermore, I would expect the discussion of QC in section 3.1 to include some summary of the frequency with which different values are flagged and some consideration of heterogeneity of flagging of values across sources and regionally. Does the frequency of QC test failure look reasonable? Does it raise any flags for particular sources etc. etc. are all questions I would expect this paper to address if it is to build user confidence in applicability of the resulting product and yet it is presently silent on these issues.

This is indeed something of interest to the reader. CEMS MDCC publishes every year an analysis on the data collection containing more in detail information, an excellent source for the potential EMO-5 user to receive further information on data providers and provision, database, post-processing, improvements to the data flow and post-processing and gap analysis. We will make sure that next years report contains also statistics on the QC flagging. We have added this source of information in the Conclusion section:

“Lastly, the CEMS Meteorological Data Collection Centre publishes every year an annual report on the CEMS meteorological data collection, with updated information on e.g. data providers and provision, database, post-processing, improvements to the data flow and post-processing and gap analysis. All reports can be found on the EFAS website, while last years report is referenced here: Rehfeldt, et al., 2021.”

7. The interpolation scheme discussion in 3.2 requires substantive additional detail. No discussion is forthcoming around why the different schemes have different skill per parameter. This must, intrinsically, be to do with the spatio-temporal correlation

structures of the different parameters and how effectively the three schemes can handle these. In particular the parameters that vary more smoothly in space and time clearly are better suited to IDW whereas the much smaller scale correlation structure precipitation is better suited to SPHEREMAP. The lack of geophysical interpretation of the findings does not build the necessary user confidence that you understand what underlies your methods.

The analysis should also remark upon the similarity in skill measures for most variables and diagnostics as this implies that interpolation choice isn't a first order effect presumably? Furthermore, is there any gradation in skill between well sampled and sparsely sampled regions? One would presume so but your present aggregation just lumps all cases together. It would be considerably more informative to split this analysis and repeat for well-sampled and sparsely-sampled regions. I would expect no impact of choice in well-sampled regions where interpolation choice makes little difference but much larger impacts in sparsely sampled regions. Can you repeat that analysis but splitting out by regions of greater and lesser station density? What does that tell you?

Reply to point 7:

This is a very interesting comment and these are all valid. But the aim of the comparison was to find the one optimal interpolation scheme for all parameters in terms of computational effort, computational time and lowest uncertainty. As precipitation is of highest importance in hydrologic modelling, a higher weight was given to the performance in precipitation than the other parameters. It was driven by the application side and not to characterize interpolation schemes.

The proposed comparison of the interpolation schemes considering the station density is an additional study and beyond the scope of this paper. A similar study was presented by GPCC in a paper from 2013 at the EGU General Assembly only for precipitation. Briefly, the interpolation uncertainty decreases with increasing station availability. The interpolation schemes didn't depict the same sensitivity, a scheme with the 'lowest' uncertainty at low station density had the 'highest' uncertainty at high station density and vice versa.

8. Given that ERA-Interim/Land values may be biased in terms of variance and mean state isn't the time-varying use of this product to infill in data sparse regions potentially problematic for long-term product homogeneity? Also, ERA5-Land has replaced ERA-Interim -land having higher resolution and improved quality. Should the algorithm not use this product instead? In particular because ERA-Interim land is no longer produced in CDAS mode introducing potentially a major discontinuity into your product upon cessation of ERA-Interim land production?

Reply to point 8:

We do agree with the reviewer on his/her view on the quality of ERA-Interim/Land and ERA5-Land. As specified in Table 1 (line 665) the import of ERA-Interim/Land data as virtual stations has been limited to peripheral areas for which the coverage with in situ stations in

the MDCC database is very low, such as Iceland, North Africa and the eastern part of the EFAS domain (Near East, Caucasus, Russia); visible in Figure 1 (top, left panel). After the discontinuity of ERA-Interim/Land an import of ERA5/Land had been discussed intensively, with the final decision of not including ERA-5/Land. Reason for that decision was that since the start of importing ERA-Interim/Land for data sparse areas more and more stations in those areas were added to the MDCC data base, which diminished the potential gain of adding ERA-5/Land.

With regards to the product homogeneity concern: As specified in the introduction (first and second paragraph) the prime purpose behind the creation of the meteorological grids is to serve the European Flood Awareness System. EFAS requires a good coverage of observations for the most recent decades (to calibrate and validate our hydrological model) as well as real time data (for the initialisation of the hydrological model used for operational forecasting). The more accurate the meteorological data, the better for the application, hence it is not the homogeneity over time that is prime concern, but to have the best possible representation of the meteorological situation for each time step. We are aware that this requirement differs from applications that focus on climate research. In order to give that credit, we added it as limitation in the Introduction as well as in the Conclusion section:

In the introduction:

“[...] Based on the information provided in this paper, the scientific community will understand that EMO-5 (version 1 as well as the operational grids) is not a climatological dataset (unlike e.g. the Essential Climate variables produced by C3S, see Noone et al., 2021), but an operational data set based on the maximum amount of quality-controlled information available at any given time. The implications of this are, that we do not advice to use this dataset to determine climate variability and climate change, however any other environmental application, especially those with real-time, high spatial resolution or multi-variable needs is likely to benefit from this data set. [...]”

In the Conclusion section:

“ The presented EMO-5 (version 1) is the result of an operational real-time service, which utilizes for every grid realisation the maximum amount of valid information available. This is very different to climate data sets, which need to be of sufficient length, consistency, and continuity to determine climate variability and climate change. EMO-5 favours the maximum amount of available information over long-term product homogeneity and is therefore not suitable for climate studies i.e. trend analysis. However, EMO-5 is suitable for environmental application which do not have this homogeneity requirement, but value a data set that uses the maximum amount of information available for each grid realisation.”

9. The assumptions around 06-18 and 18-06 for Tn and Tx will miss many true Tx and Tn values particularly in the winter half season when daily maxima or minima often occur outside these windows. Particularly so in the higher latitudes of the domain. They also risk introducing a discontinuity with the mean temperatures calculated

over the full 24 hours. At a minimum caveats need to be added but, ideally, you would calculate Tx, Tn and Tm consistently for full 24 hour periods to enable cases where Tx and / or Tn fall outside of nominal day / night.

We assume that the reviewer is familiar with the WMO guidelines on the calculation of daily minimum and maximum temperature (WMO-No. 306) and hence recognise that we are following the WMO standard, which we regard as good practise, especially considering that we are an operational service.

Nevertheless, we are aware about the mentioned implications and have added a paragraph in the conclusions on this topic:

“Further, there might be some reservations towards using the EMO-5 minimum night-time and maximum day-time temperature as the daily minimum and maximum temperature respectively. To recall, the daily minimum and maximum temperature within EMO-5 are calculated following the WMO guideline (WMO-No. 306), which assumes the minimum temperature to occur between 18:00 and 06:00, and the maximum temperature between 06:00 and 18:00. However, particularly in the winter half season and in the higher latitudes, the daily maxima or minima temperature occur sometimes outside these windows. For this reason, some data sets, such as E-OBS, calculate the daily minimum and maximum temperature over the full 24 hour period. Lavaysse et al. (2018) who investigated the minimum and maximum temperature of EMO-5 (referred to as LisFlood in their paper) and E-OBS with regards to their suitability in the frame of temperature extremes in Europe (heat- and cold waves), came to the conclusion that the two observational datasets showed only minor differences in heat and cold waves occurrences and intensities, which according to Lavaysse is probably due to the good agreement in representing both, the minimum and maximum temperature.”

10. More details are required upon the land-sea mask used and how it is aggregated to the 5km grid. Are stations omitted based upon whether land or ocean at the native e.g. 5 minute resolution of the mask but then the gridded product produced for all 5km gridcells that contain any land? The method is not reproducible or understandable absent of such details.

Reply to point 10:

Thank you, this paragraph could indeed be misunderstood. The decision about the inclusion or exclusion of a station from gridding is done independently from the land-sea-mask. Per default, all stations within the EFAS domain (European window, plus neighbouring countries) are used for gridding, even if these are at very small islands or drilling platforms. If a station reports very irregular, low quality or no data, then this station is excluded from gridding. It is now reworded as:

“A land-sea mask is used to exclude sea surfaces from the gridding procedure, as EMO-5 originates from the need for near real-time information on observed meteorological conditions over land surface areas. The land-sea-mask is not used to define stations to be excluded from gridding.”

11. Section 3.3.4 (Implications of altitude for temperature and water vapour pressure) is a little basic compared to state-of-the-art spline fitting techniques as used e.g. in HadEX or CRUTS. Why has such a relatively speaking simple approach been used as is described in 3.3.4 and how does it compare to spline fitting or other techniques?

Reply to point 11:

EMO-5 is an operational, real-time data set. Hence the entire methodology needs to be robust and executable in a couple of hours. We tested some techniques to fit parameters for the interpolation to the available data for the actual time stamp, but the quality of the fitted parameters was often below well-developed climatological means. Sometimes the fitting routine even didn't converged.

12. The mean temperature discussion in 3.3.5 would benefit from discussing various pieces of literature about how to create mean temperatures and the difference between the mean of a 24 hour period, the mean of max / min and various other alternative methods. There have been several papers on the different ways to calculate means and the random and systematic effects they can have. For the purpose of your dataset the key issue is whether the different ways you have done so may impart systematic or random effects. This can be ascertained from a representative (climatologically) set of well-sampled stations being deliberately degraded to calculate the mean from the 24(+) instantaneous values and then various approximations. This can then at least bound your uncertainties.

Reply to point 12:

We have added a reference to Luedeling (2018), who interpolated hourly time series from min-max values. We have also done a simple cross-validation of a set of simulated 6-hourly times series and compared with the observations, both for the entire time series, and by looking at the 6-hourly averages for the different times of the day separately. We agree that both the methodology and the data set would make it possible to extend both analyses and the discussion on this topic considerably, but we see this as out of scope for this manuscript due to length issues.

We have extended Section 3.3.5 as follows:

“The goodness of this approximation was analysed in a simple cross-validation procedure. We picked around 800 stations that already had 6-hourly observations from the start of the period (1990). For these stations, we compared the 6-hourly observations with the simulated 6-hourly dataset based on maximum and minimum temperatures. A hexbin-plot is shown in Figure 7, where darker colors indicate a high density of points indicate a temperature with a large number of observed and simulated values. The line shows where simulations are equal to the observations, whereas the dashed lines indicate where the simulated values are 2 degrees higher or lower than the observations. We can see that a large majority of the

simulated values are within 2 degrees of the observations, although there are cases with larger deviations.

The correlation between observations and simulations range from 0.91-0.99. The simulated values are on the average unbiased, with mean and median around 0.1 degrees below the observations. However, when looking at the simulated temperatures for different times of the day, it can be noted that the method has a tendency to underestimate the average night temperature, and overestimate the afternoon temperature. The mean and median of the underestimation is 1.8 degrees for the night temperature and the overestimation 1.5 degrees for the afternoon, i.e., the simulated 6-hour periods are slightly more extreme than the observations. The root mean squared error (RMSE) for each station ranged from 1-4 degrees, with mean and median RMSE of 2.2 degrees. The difference was seen as acceptable for our purposes, and is quite similar to what was observed by e.g. Luedeling (2018)."

13. I would expect considerably more in the data availability section. It is grossly insufficient to just point to a website. You must describe things such as the data format(s) (ideally there should be at least two to cater to a range of users), any software tools available, version control, version retention policies, whether files are available as spatial or temporal aggregations, whether the files contain uncertainty information, whether they are univariate or multivariate. Otherwise no user is going to go to that site on the off chance. You are not really helping yourselves to advertise the availability if all you do is point to a URL. Users need to know what they are being offered and how well it is ultimately managed.

Reply to point 13:

We followed the official instructions for submission provided by ESSD with regards to what should be in the individual sections (see <https://www.earth-system-science-data.net/submission.html>). For the data availability section it reads as follows:

Data availability: this section should contain all necessary information on data access. Upon acceptance for publication, this includes at least the data set DOI(s) and their data-citation(s). If more than 5 DOIs are necessary, please create a table including DOIs and citations and reference this table here instead. This is also the section that holds review links or other access tokens to your data set if necessary for the review. Optionally, overarching project URLs or other meta-information can be included. The reader must gain direct access to the data by the means described in this section.

Despite being regarded as “grossly insufficient” by the reviewer, the data availability section is in line with the journal’s content guideline. Nevertheless, we agree with the reviewer on the value of that information for which reason we have extended the data availability section, and have created a README.txt which has been uploaded to the data repository.

4 Data availability

EMO-5 (Thiemig et al., 2021) is a Copernicus product and as such free and open to everyone. It can be accessed through the Data Catalogue of the European Commission’s Joint Research Centre at <https://doi.org/10.2905/0BD84BE4-CEC8-4180-97A6-8B3ADAAC4D26>.

The repository contains a CF-1.6 compliant NetCDF stack files for each variable, as well as a README file with detailed product specifications, which are briefly summarised in Table 5 below.

Table 5: Product specification of EMO-5.

<i>Data set description</i>	
Name of the data set	EMO-5 (EMO stands for “European Meteorological Observations”, whereas the 5 denotes the spatial resolution of 5 km)
Short description	EMO-5 (version 1) is a European high-resolution, (sub-)daily, multi-variable meteorological data set built on historical and real-time observations obtained by integrating data from 18,964 ground weather stations, four high-resolution regional observational grids (i.e. CombiPrecip, ZAMG - INCA, EURO4M-APGD and CarpatClim) as well as one global reanalysis (ERA-Interim/Land). EMO-5 includes at daily resolution: total precipitation, temperatures (minimum and maximum), wind speed, solar radiation and water vapour pressure. In addition, EMO-5 also makes available 6-hourly precipitation and mean temperature. The raw observations from the ground weather stations underwent a set of quality controls, before SPHEREMAP and Yamamoto interpolation methods were applied in order to estimate for each 5x5 km grid cell the variable value and its affiliated uncertainty, respectively.
Created by	Copernicus Emergency Management Service
Horizontal coverage	Europe (EFAS domain)
Horizontal resolution	5 km
Spatial gaps	Only land areas are covered by this data set
Temporal coverage	1990-01-01 till 2019-12-31
Temporal resolution	daily and 6-hourly
Temporal gaps	No gaps
Number of available variables	7
Variables available at daily resolution	total precipitation, temperatures (minimum and maximum), wind speed, solar radiation and water vapour pressure
Variables available at 6-hourly resolution	total precipitation and mean temperatures
Units	precipitation [mm], temperature [°C], vapor pressure [hPa], solar

	<i>radiation [J/m²], wind speed [m/s]</i>
<i>Projection</i>	<i>Lambert Azimuthal Equal-Area (5km)</i>
<i>Data type</i>	<i>5*5 km grids</i>
<i>Available version</i>	<i>version 1</i>
<i>DOI of dataset</i>	<i>https://doi.org/10.2905/OBD84BE4-CEC8-4180-97A6-8B3ADAAC4D26</i>
<i>PID of dataset</i>	<i>http://data.europa.eu/89h/0bd84be4-cec8-4180-97a6-8b3adaac4d26</i>

14. seNorge2 and seNorge2018 are as I understand it consecutive versions of a single dataset with 2018 replacing 2. You should use just the 2018 version as the other version has been deprecated. Having also reviewed that paper and in particular considering their discussion it would furthermore not be appropriate to treat that product as truth given the substantial caveats made around interpolation in data sparse areas of complex topography. Therefore the seNorge2018 validation must be noted to be conditional on the verity of that product which cannot be assured a priori. You cannot treat either seNorge version as 'truth' which significantly inhibits their value as a means of validation. There are many other national gridded products and comparison to a range of such products e.g. the UK analysis from the Met Office may be instructive and reduce the dependency upon a single product which may itself contain substantive systematic biases.

Reply to point 14:

In the absence of knowing the actual "truth" as neither in situ measurements nor high-resolution do represent the truth under all circumstances, we do not treat either of those as truth, and have neither suggested it in the manuscript. What we did however do is an intercomparison, and we do learn from those through identifying similarities and differences in the products, especially if we do understand their underlying production methodologies. That said, we do find value in not only comparing it to seNorge2018, but also to seNorge2, as the newer dataset is not a simple successor. Apart from the number of stations in seNorge2018, there were very large differences and evolution in the methodology used to create it, which are very well documented in the literature (referred to as Lussana et al., 2018 and Lussana, 2018b in the paper). The use of both datasets has the advantage to provide ideas on how to improve the EMO-5 dataset in the future and what influence the different approaches have (topography, use of reanalyses, ...). The changes and limitations are very well documented, which is very helpful for the interpretation of the results. For this very reason, the authors consider it useful to include both data sets.

Of course, a comparison with many different high-resolution regional datasets would be interesting. However considering already the length of the manuscript out of scope. At the same time, the comparison done with seNorge data is without doubt interesting and useful,

as the seNorge datasets as an example fulfils many considerations: high-resolution datasets, which is very important for a high-resolution dataset like EMO-5 as a comparison; the Norwegian topography is very challenging; there are data sparse and dense areas; and it's documentation. In addition, as described above, there are versions with very different methodologies, which the authors see as a positive. Because this helps to understand the EMO-5 dataset with both strengths and limitations.

15. I cannot make head or tail of Figure 11 from the caption and the individual images. The caption needs expanding and the panels given better titles. I cannot understand why the panels in each case vary in the manner they do and it is non-intuitive to me and therefore will be so to your readers. A figure extending over several pages is also barely legible. You would probably be better to concentrate upon 1 or 2 events that could be given more space in the main text and place the remaining cases in supplemental materials.

Reply to point 15:

We do agree that our analysis of extreme precipitation events is quite extended with a long figure. However, knowing that we are interested in using the data set for reproducing floods the capacity of the data set to capture high precipitation events is of particular interest/ importance to us, and hence worth investigating. Also, we have included numerous events as we did not want to choose a well-working one and then call it a 'representative' example. We believe this has a much higher value to the reader than 1-2 cherry-picked examples.

16. Lacking in entirety is any meaningful discussion section which may highlight caveats, future work avenues, future priorities, synergies with other Copernicus and global activities (such as WMO) that may yield improvements etc. etc. I would expect a substantive discussion section which covered such points.

Reply to point 16:

Future plans for EMO-5 are mentioned under Section "6 Conclusions and future work", see last paragraph (l 498ff), which reads as follows:

"As part of the operational Copernicus EMS, the number of stations (historical and near real-time) that are used for gridding in EMO-5 will be continuously increased, through adding new data providers and the integration of new, high-resolution regional observational grids, where available. In addition, to improve the current quality control framework, new data validation rules, such as spatial comparison with neighbouring stations or additional statistical checks, will be implemented. Finally, as it is foreseen to increase the spatial resolution of EFAS from the current 5 km grid to a 1 arc minute grid (approximately equal to 1.8 km at the equator), also EMO-5 will increase the spatial resolution in its next version."

Synergies with other Copernicus services is a continuous endeavour and takes place not through discussing/stating those in scientific publications, but via direct dialogues and collaborations, which we practice both.

As already mentioned under point 8, we have added a paragraph to the conclusion section which highlights caveats and limitations of EMO-5. It reads as follows:

“The presented EMO-5 (version 1) is the result of an operational real-time service, which utilizes for every grid realisation the maximum amount of valid information available. This is very different to climate data sets, which need to be of sufficient length, consistency, and continuity to determine climate variability and climate change. EMO-5 favours the maximum amount of available information over long-term product homogeneity and is therefore not suitable for climate studies i.e. trend analysis. However, EMO-5 is suitable for environmental application which do not have this homogeneity requirement, but value a data set that uses the maximum amount of information available for each grid realisation.

Further, there might be some reservations towards using the EMO-5 minimum night-time and maximum day-time temperature as the daily minimum and maximum temperature respectively. To recall, the daily minimum and maximum temperature within EMO-5 are calculated following the WMO guideline (WMO-No. 306), which assumes the minimum temperature to occur between 18:00 and 06:00, and the maximum temperature between 06:00 and 18:00. However, particularly in the winter half season and in the higher latitudes, the daily maxima or minima temperature occur sometimes outside these windows. For this reason, some data sets, such as E-OBS, calculate the daily minimum and maximum temperature over the full 24 hour period. Lavaysse et al. (2018) who investigated the minimum and maximum temperature of EMO-5 (referred to as LisFlood in their paper) and E-OBS with regards to their suitability in the frame of temperature extremes in Europe (heat- and cold waves), came to the conclusion that the two observational datasets showed only minor differences in heat and cold waves occurrences and intensities, which according to Lavaysse is probably due to the good agreement in representing both, the minimum and maximum temperature.”