

MIS 5e sea-level history along the Pacific Coast of North America

Daniel R. Muhs¹

¹U.S. Geological Survey, MS 980, Box 25046, Federal Center, Denver, Colorado 80225 USA

Correspondence to: Daniel R. Muhs (dmuhs@usgs.gov)

Abstract. The primary last interglacial, marine isotope substage (MIS) 5e records on the Pacific Coast of North America, from Washington (USA) to Baja California Sur (Mexico), are found in the deposits of erosional marine terraces. Warmer coasts along the southern Golfo de California host both erosional marine terraces and constructional coral reef terraces. Because the northern part of the region is tectonically active, MIS 5e terrace elevations vary considerably, from a few meters above sea level to as much as 70 m above sea level. The primary paleo-sea level indicator is the shoreline angle, the junction of the wave-cut platform with the former sea cliff, which forms very close to mean sea level. Most areas on the Pacific Coast of North America have experienced uplift since MIS 5e time, but the rate of uplift varies substantially as a function of tectonic setting. Chronology in most places is based on uranium-series ages of the solitary coral *Balanophyllia elegans* (erosional terraces) or the colonial corals *Porites* and *Pocillopora* (constructional reefs). In areas lacking corals, correlation to MIS 5e often can be accomplished using amino acid ratios of fossil mollusks, compared to similar ratios in mollusks that also host dated corals. U-series analyses of corals that have experienced largely closed-system histories range from ~124 to ~118 ka, in good agreement with ages from MIS 5e reef terraces elsewhere in the world. There is no geomorphic, stratigraphic, or geochronology evidence for more than one high-sea stand during MIS 5e on the Pacific Coast of North America. However, in areas of low uplift rate, the outer parts of MIS 5e terraces apparently were re-occupied by the high-sea stand at ~100 ka (MIS 5c), evident from mixes of coral ages and mixes of molluscan faunas with differing thermal aspects. This sequence of events took place because glacial isostatic adjustment processes acting on North America resulted in regional high-sea stands at ~100 ka and ~80 ka that were higher than is the case in far-field regions, distant from large continental ice sheets. During MIS 5e time, sea surface temperatures (SST) off the Pacific Coast of North America were higher than is the case at present, evident from extralimital southern species of mollusks found in dated deposits. Apparently, no wholesale shifts in faunal provinces took place, but in MIS 5e time, some species of bivalves and gastropods

25 lived hundreds of kilometers north of their present northern limits, in good agreement with SST estimates derived from
26 foraminiferal records and alkenone-based reconstructions in deep-sea cores. Because many areas of the Pacific Coast of
27 North America have been active tectonically for much or all of the Quaternary, many earlier interglacial periods are recorded
28 as uplifted, higher elevation terraces. In addition, from southern Oregon to northern Baja California, there are U-series-dated
29 corals from marine terraces that formed at ~80 ka, during MIS 5a. In contrast to MIS 5e, these terrace deposits host
30 molluscan faunas that contain extralimital northern species, indicating cooler SST at the end of MIS 5. Here I present a
31 review and standardized database of MIS 5e sea-level indicators along the Pacific Coast of North America and the
32 corresponding dated samples. The database is available in Muhs et al. (2021a) [<https://doi.org/10.5281/zenodo.5903285>].

33 **1 Introduction**

34 Because of the prospect of future sea-level rise, there has been an increasing interest in past, but geologically recent times
35 of higher than present sea level. One of the best studied of these is the last interglacial (LIG), recognized in terrestrial geologic
36 records as the Sangamon (North America) or Eemian (Europe) periods. Within the deep-sea sediment core record, Arrhenius
37 (1952) initiated the widely accepted practice of numbering Quaternary interglacial and glacial stages, which was encouraged
38 with the pioneering work on oxygen isotopes in such cores by Emiliani (1955). Interglacial periods have odd numbers and
39 glacial periods have even numbers. Thus, the last interglacial (*sensu lato*) in deep-sea cores is known as marine isotope stage
40 (MIS) 5. Shackleton (1969) recognized five major substages of the MIS 5 complex (5e, 5d, 5c, 5b, 5a, from oldest to youngest),
41 and those substages are now widely recognized by marine stratigraphers and paleoclimatologists. Another nomenclature
42 suggested by Martinson et al. (1987) is followed by some investigators, with the peaks of these substages referred to as “events”
43 MIS 5.5, 5.4, 5.3, 5.2, and 5.1, from oldest to youngest. MIS 5e or 5.5 is considered to be the period of peak global warmth
44 and minimal global ice of the late Quaternary (see review in Murray-Wallace and Woodroffe, 2014). Some investigators also
45 consider that MIS 5e alone is the last interglacial (*sensu stricto*).

46 The Pacific Coast of North America contains a rich record of Quaternary sea level history, particularly the peak of the LIG,
47 MIS 5e, generally considered to date from ~130 ka to ~115 ka. Part of the richness of this sea level record is due to the tectonic
48 setting of North America (Fig. 1). Most of the continent is situated on the North America lithospheric plate. However, Baja
49 California and part of westernmost California are both located on the Pacific plate, and the southern part of Central America
50 is on the Caribbean plate. The boundaries between the two major (Pacific and North America) and smaller lithospheric plates
51 are the tectonic controls on the crustal blocks that form the Pacific Coast of North America. In southwestern Canada and the
52 northwestern USA, the Cascadia subduction zone occurs where the southeast-moving Juan de Fuca and Gorda plates meet the
53 North America plate (Fig. 2). Farther south, from northern California to the Golfo de California, the dominant structural

54 control is the San Andreas Fault, a major right-lateral (dextral) strike-slip system, with many smaller, subparallel faults
55 associated with it. Still farther south, at the head of the Golfo de California, the structural style changes again, with the
56 boundary between the Pacific and North America plates taking the form of a spreading center, the northernmost part of the
57 East Pacific Rise (Fig. 2). Finally, the structural style changes to the south once more, back to a subduction zone, in southern
58 Mexico and Central America. Here the Cocos plate is being subducted under the North America plate (in the northern part)
59 and under the Caribbean plate in the southern part (Fig. 1).

60 The importance of tectonic setting for studies of past shorelines, such as that of MIS 5e, is due to its influence on vertical
61 movement of coastal crustal blocks. In collisional zones, such as the Cascadia subduction zone, it could be expected that some
62 vertical movement might be found in the crust of the overriding plate. Indeed, a classical study by Uyeda and Kanamori (1979)
63 proposed that where the dip of the subducting plate is shallow, rapid uplift should be seen in the overriding plate. However,
64 detailed studies of marine terraces in northern California and Oregon by H.M. Kelsey and his colleagues (Kelsey, 1990;
65 McInelly and Kelsey, 1990; Kelsey and Bockheim, 1994; Kelsey et al., 1994, 1996; Polenz and Kelsey, 1999) have shown
66 convincingly that it is actually local structures (faults and folds) *within* the upper plate that control the rates of marine terrace
67 uplift seen along much of the Cascadia subduction zone. Farther south, within the San Andreas Fault zone, rates of uplift are
68 highly variable (see summary in Muhs et al., 2014b). Along much of the coast bordering this fault zone, uplift rates are modest,
69 likely (though not yet proven) because movements along faults that have a predominantly strike-slip (horizontal) sense of
70 movement have a small vertical component. Exceptions to this occur where there are restraining bends in these faults, the
71 most famous of which is the "big bend" area of the San Andreas Fault zone (Fig. 2). Here, crustal compression results in
72 extremely high rates of uplift. Away from the zone of maximum uplift south of the big bend in the San Andreas Fault, Shaw
73 and Suppe (1994) proposed that uplift of the Santa Cruz Island and Anacapa Island shelf area of southern California is due to
74 movement on an underlying, blind thrust fault. High rates of uplift can also be found on coastlines adjacent to triple junctions,
75 such as the Mendocino triple junction (Fig. 2) and the Panama triple junction ("PTJ" in Fig. 1). Along coastlines bordering a
76 spreading-center plate boundary, such as that in the Golfo de California, crustal blocks are moving away from each other and
77 accommodation space is increasing, so uplift rates are not expected to be particularly high. This simplified picture is to a great
78 extent borne out by field studies (e.g., Ortlieb, 1991), although local structures can again play a role in generating uplift over
79 limited parts of such a coastline. Uplift in Central America is rapid in places, due to subduction of seamounts on the Cocos
80 and Nazca plates.

81 Marine terraces along the Pacific Coast of North America have been studied for more than a century. Lawson (1893)
82 considered that emergent terraces formed by episodic (and presumably rapid) uplift, what would now be referred to as
83 coseismic uplift. Smith (1900), studying terraces on the California islands, concurred with this hypothesis, reasoning that
84 episodic uplift must alternate with periods of "comparative quiescence." Interestingly, the concept of episodic rapid uplift is
85 now known to have validity for some parts of the Pacific Coast, in diverse tectonic settings (see discussion below on Holocene
86 shorelines).

87 Grant and Gale (1931) also considered emergent terraces to have a tectonic origin, but also pointed out the possibility of a
88 eustatic component. It was Davis (1933), however, studying marine terraces in the Malibu, California region, who was likely
89 the first to point out explicitly that although uplift was obviously a factor in the formation of a flight of marine terraces, a
90 eustatic component was important as well. Davis (1933) considered that uplift rates were likely to have geographic variability,
91 but he noted that eustatic records ought to be the same everywhere. Despite the publication of this important paper, there was
92 a return to the idea of terraces being dominantly of a tectonic origin in later studies by Putnam (1942), Woodring et al. (1946),
93 and Upson (1951). Woodring et al. (1946) thought that eustatic effects were either obliterated or obscured in the geologic
94 record of marine terraces. Although Upson (1951) considered that terraces were formed principally by episodic uplift, he
95 recognized that there were problems with this explanation and thought that a eustatic component was present.

96 Interestingly, it was a master's degree thesis at the University of California at Berkeley that articulated our current concepts
97 of marine terraces, uplift, and sea level clearly for the first time. Alexander (1953), working on both marine and stream terraces
98 in the Capitola-Watsonville area of central California, measured the maximum elevations of marine terraces and the tops of
99 stream-fill terraces, noting their similar elevations, and reasoning that they must have a common, eustatic control. He also
00 noted that in between times of stream terrace formation, there were episodes of valley cutting, which indicated periods of
01 eustatically lowered sea level, during glacial periods. On the other hand, multiple marine terraces indicated long-term tectonic
02 uplift. He (Alexander, 1953, p. 36) concluded that "Thus, the marine terraces of the Capitola-Watsonville area are regarded
03 as having originated under conditions of a slowly and continuously rising coast against which occurred at least three complete
04 cycles of eustatic changes in sea level." This is a remarkable conclusion, reached before any modern methods of
05 geochronology were in common use, and based only on sound field mapping, elevation measurements, and geomorphic
06 reasoning. It was this concept, along with uranium-series geochronology, that allowed Broecker et al. (1968), Mesolella et al.
07 (1969), and Veeh and Chappell (1970) to infer that coral reef terraces on the uplifting coasts of Barbados and New Guinea
08 recorded interglacial periods that supported the Milankovitch or orbital theory of climate change. In California, Alexander's
09 (1953) concept was accepted explicitly or implicitly by subsequent workers in the following decades (e.g., Vedder and Norris,
10 1963; Birkeland, 1972; Bradley and Griggs, 1976; Wehmiller et al., 1977a), and his contribution is now recognized in one of
11 the leading textbooks on geomorphology (Anderson and Anderson, 2010).

12 Dating of marine terraces on the Pacific Coast of North America had a development similar to that for other coastlines.
13 Early use of uranium-series (U-series) analyses of corals was reported by Veeh and Valentine (1967), Valentine and Veeh
14 (1969), and Ku and Kern (1974). In these investigations and most subsequent studies, the taxon analyzed is the solitary coral
15 *Balanophyllia elegans* (Gerrodette, 1979), which is by far the most common coral found in Oregon and California marine
16 terrace deposits. These early studies permitted an interpretation that low-elevation terraces at Cayucos, San Nicolas Island,
17 and Point Loma could all date to MIS 5e. Other studies attempted U-series analyses of fossil mollusks (e.g., Bradley and
18 Addicott, 1968; Szabo and Rosholt, 1969; Szabo and Vedder, 1971), but a seminal study by Kaufman et al. (1971), with
19 extensive data from California terraces, showed that mollusks are inappropriate materials for U-series geochronology.

20 A new development in geochronology, however, brought mollusks back to the forefront in dating marine terraces on the
21 Pacific Coast. Using the Cayucos, San Nicolas Island, and Point Loma U-series coral ages as calibration points, Wehmiller et
22 al. (1977a), Wehmiller (1982), and Kennedy et al. (1982) showed that terraces from Baja California Sur to Oregon could be
23 correlated to MIS 5e on the basis of amino acid ratios in fossil mollusks, a profound finding that demonstrated the extensive
24 nature of the last interglacial record on the Pacific Coast of North America. In addition, these studies also showed that uplift
25 rates on the Pacific Coast are variable, overturning a long-held concept that the lowest marine terrace is everywhere of the
26 same age. Indeed, terraces estimated to be as young as ~50 ka were found in areas of high uplift rate.

27 Concerted efforts to find corals yielded more U-series ages of marine terraces. Rockwell et al. (1989) mapped 14 terraces
28 on Punta Banda, Baja California, the lowest 3 of which have shoreline angle elevations of 15-17 m, 22 m, and 27-43 m. The
29 1st or "Lighthouse" terrace has U-series ages (by alpha spectrometry) of corals and hydrocorals of ~80 ka, the 2nd terrace is
30 undated, and the 3rd or "Sea Cave" terrace has ages of ~120 ka. This was the first study on the Pacific Coast to provide
31 definitive geochronologic evidence of both MIS 5e and MIS 5a (as well as a likely MIS 5c at ~22 m) terraces. Muhs et al.
32 (1990, 1992, 1994) reported additional U-series ages, again by alpha spectrometry, for MIS 5e terraces at Cayucos, Point San
33 Luis, San Nicolas Island, San Clemente Island, and Point Loma (all in California), and Punta Banda, Isla Guadalupe, and Cabo
34 Pulmo (in Baja California and Baja California Sur). Terraces dating to MIS 5a were reported from Coquille Point, near
35 Bandon, Oregon; Point Arena, San Nicolas Island, and Point Loma (all in California); and Punta Banda (Baja California).

36 The development of U-series dating of corals by thermal ionization mass spectrometry (TIMS) led to a new level of
37 complexity in the understanding of the Pacific Coast marine terrace record. Stein et al. (1991) redated corals from the Cayucos
38 and Point Loma areas, confirming that fossils dating to MIS 5e were present, but also showing the possibility that some corals
39 dated to MIS 5c (~100 ka). Muhs et al. (2002a) confirmed these results for both Cayucos and Point Loma. Kennedy et al.
40 (1982), in their amino acid study along the Pacific Coast, reported that terraces correlated to MIS 5a had molluscan faunas
41 with cool-water aspects, whereas those correlated to MIS 5e hosted molluscan faunas with warm-water aspects. Cool-water
42 faunas were confirmed with corals dated to ~80 ka using TIMS by Muhs et al. (2006) in a later study, in partial support of
43 Kennedy et al. (1982). However, Muhs et al. (2002a) showed that the terraces at Cayucos and Point Loma, containing both
44 MIS 5e and MIS 5c corals, hosted molluscan faunas with a mix of both warm-water taxa (thought to date from MIS 5e) and
45 cool-water taxa (thought to date from MIS 5c). This idea was explored in more detail on San Nicolas Island, where the lowest
46 three terraces (1, 2b, and 2a, in ascending elevation order) were mapped in detail, terraces elevations were measured precisely
47 with differential GPS methods, corals from all three terraces were dated with TIMS, and the faunas were characterized (Muhs
48 et al., 2012). Terrace 1 dates to ~80 ka and hosts a cool-water fauna, terrace 2b has both 100 ka and 120 ka corals and hosts a
49 mix of cool-water and warm-water taxa, and terrace 2a has only ~120 ka corals, no cool-water taxa, but several warm-water
50 taxa. This finding raised the possibility that the MIS 5c high-sea stand in this region had a paleo-sea level elevation higher
51 than what would have been inferred from the classic records on Barbados and New Guinea, and that this high stand overtook
52 at least the outer part of the MIS 5e terrace, reworking and mixing its fossils (with warm-water taxa) with shells dating to MIS
53 5c (with cool-water taxa). Subsequent studies have shown that other terraces dating to MIS 5e (by TIMS U-series on corals)

54 or correlated to MIS 5e (by amino acids on mollusks) also contain mixes of warm-water and cool-water taxa (Muhs et al.,
55 2014a, 2014b; Muhs and Groves, 2018).

56 The main aim of this paper is to serve as a description to accompany a standardized database of MIS 5e sea-level indicators
57 compiled following the WALIS template (Rovere et al., 2020). From the published papers in the area of interest, I extracted
58 sea level indicators and standardized the quantification of their elevation and indicative meaning (Shennan, 1982; Shennan et
59 al., 2015; Rovere et al., 2016), along with appropriate metadata. Each sea level indicator was then associated with one or more
60 samples, dated with U-series or amino acid racemization (AAR) methods, that were also added to the database. In some cases,
61 U-series dated samples were already present in the WALIS database from the compilation of Chutcharavan and Dutton, 2021).

62 **2 Sea level indicators**

63 As pointed out by Rovere et al. (2016), critical to reconstructing past sea level during MIS 5e (or any past high-sea stand,
64 for that matter) is an accurate assessment of paleo-sea level indicators (Table 1). For the vast majority of MIS 5e geomorphic
65 records along the Pacific Coast of North America, the best relative sea level (RSL) indicator is what is called the *shoreline*
66 *angle*, a term that goes back to the classic study of terraces in the Malibu, California area by Davis (1933). The shoreline
67 angle is the junction of the marine platform (or "wave cut bench"), formed in the surf zone and the sea cliff, when viewed in
68 cross section (Fig. 3a). Davis (1933) and virtually all investigators who have followed him have generally regarded the
69 shoreline angle as the best overall RSL, because it is considered to form at or near sea level. Kelsey (2015) points out that
70 depending on bedrock type, structures within the local bedrock, orientation of the coast with respect to wave exposure, and
71 other factors, shore-parallel variability of the shoreline angle elevation on modern coastlines can vary by as much as 1-4 m.
72 In the San Diego area, however, measurements made by Kern (1977) indicate that modern shoreline angles typically form
73 within a meter of modern sea level. Whether shoreline angles on the Pacific Coast form closest to mean sea level or high-tide
74 level is probably not known with any certainty. In any case, however, the range of variability of shoreline angle elevations
75 noted by Kelsey (2015) is typically greater than the mean tidal range. In southern and central California, from San Diego to
76 San Francisco Bay, mean tidal range is typically only 1.1 to 1.2 m; in northern California, it increases to about 1.2 to 1.5 m;
77 and in Oregon, it is 1.6 to 1.8 m (data from: https://tidesandcurrents.noaa.gov/tide_predictions.html). In most places that I
78 have studied along the Pacific Coast, marine platforms or wave-cut benches are typically only visible at low tide (Fig. 4) and
79 are not visible at high tide. In most of California, therefore, with a mean tidal range (low to high tide) of only about a meter,
80 these observations suggest that Kern's (1977) observations have general validity, and shoreline angles approximate mean sea
81 level.

82 For the field geomorphologist studying marine terraces, a much greater challenge lies in mapping shorelines accurately and
83 finding good exposures of ancient shoreline angles. After terrace emergence, the wave-cut platform and the marine sediments
84 covering it become the locus of deposition of terrestrial deposits, including alluvium, colluvium, and eolian sand (Fig. 5).
85 Such deposits obscure the precise location of the *inner edge* of a marine terrace. The term "inner edge" is often used

86 interchangeably with the term shoreline angle, but here it is meant to express the spatial extent of a shoreline, i.e., viewed
87 planimetrically, in a shore-parallel sense. Put another way, it is the mapped expression of where the shoreline angle is situated,
88 marking the former junction of land and sea. Terrestrial deposits that cover inner edges of marine terraces not only make
89 mapping of a given terrace difficult, but also can be extensive enough that they cover two or more discrete terraces. Alluvial
90 and eolian deposition can sometimes generate a rather smooth surface that gives the impression of being an actual marine
91 platform surface, which may in reality be many meters below (Fig. 3b). In the example shown in Figure 3b, the unwary
92 researcher might assume that there is only one terrace here, and also could easily assume that the "apparent inner edge" is
93 where the actual shoreline angle is situated, when in fact it is seaward of this and at a much lower elevation.

94 Even where shoreline angles are well exposed, an additional complication can arise, particularly in areas where uplift rates
95 are low or no uplift at all is occurring. In such areas, successions of sea-level high stands that have similar paleo-sea levels
96 may reoccupy a terrace. Examples of this with reoccupation of the MIS 5e terrace by the MIS 5c high-sea stand were noted
97 earlier. Where this has occurred, it may sometimes be difficult or impossible to estimate paleo-sea level during MIS 5e time,
98 even when a shoreline angle is well exposed, as it would be unclear which high-sea stand produced that feature.

99 In the database, the upper and lower limit of the indicative range for a shoreline angle were set as the Mean Lower Low
00 Water (MLLW) and the Mean Higher High Water (MHHW) reported for the nearest NOAA tide station. In studies reviewed
01 here, whenever the shoreline angle elevation was not reported by the original authors, the dated sample elevation was used,
02 and readers are cautioned that such elevations are therefore minimum-limiting for an estimate of paleo-sea level.

03 Because most Pacific Coast marine terraces develop on a high-energy, erosive coastline, biological indicators of RSL are
04 rare. Typically, marine fossils are found in a poorly sorted mix of sand and gravel. As a consequence, the fossils in marine
05 terrace deposits, even those near the former shoreline, have been transported there by waves, sometimes from depths of 20 m
06 or more. Exceptions to this, while uncommon, do occur and most often take the form of rock-boring mollusks in growth
07 position, particularly bivalves in the Pholadidae family. A good example of this is the species *Penitella penita*. This taxon
08 typically occurs in the mid-intertidal zone, based on modern specimen collections in the Santa Barbara Museum of Natural
09 History (P. Valentich-Scott, written communication, March 2020). Only rarely is *P. penita* found below depths of ~10 m
10 (Coan et al., 2000). Thus, if fossil *P. penita* is found in growth position in bored holes of a wave-cut bench (Fig. 6), it is likely
11 that one is within 10 m of paleo-sea level. While this criterion is not as specific an RSL indicator as a shoreline angle, it is
12 often a complementary tool for paleo-sea level. Other species of bivalves can potentially serve as paleo-sea level indicators if
13 they are articulated, a characteristic not possible with gastropods. For example, the large bivalve *Saxidomus nuttalli*, which is
14 presently found from northern California to Baja California Sur, typically lives in muddy sediments within the intertidal zone
15 to ~10 m depth (Coan et al., 2000), a range similar to *Penitella penita*. Thus, if an articulated fossil specimen of *S. nuttalli* is
16 found, it is *possible* that it is close to where it was situated when it was living, because wave transport commonly will
17 disarticulate shells. However, *S. nuttalli* is not a rock-boring mollusk, so without occurrence in a hole that it has bored, one
18 can never be certain, even with articulated shells, that one is near the position where the specimen lived.

19 Farther south, along both shores of the Golfo de California and the Pacific Coast of mainland Mexico and Central America,
20 ocean water temperatures are higher than farther north, and hermatypic (reef-building) corals are found (Fig. 7). Although
21 hermatypic corals can be found throughout much of this region, true coral reefs are far less common. For example, within the
22 Golfo de California, although corals can be found along almost all of the Baja California coast and much of the Sonoran coast,
23 true coral reefs have been documented only at a few localities. The region from the upper Golfo de California to Panama does,
24 however, host a surprising diversity of coral species (Reyes-Bonilla and López-Pérez, 1998; Glynn and Ault, 2000; Glynn et
25 al., 2017; Toth et al., 2017). Some of the most important genera are *Porites* (7 species), *Pocillopora* (6 species), *Psammocora*
26 (4 species), and *Pavona* (5 species). *Porites panamensis* (formerly *P. californica* in some studies) is found from the upper
27 Golfo south to Panama, but also has a disjunct distribution, with colonies of this taxon also found in Bahía Magdalena, on the
28 Pacific coast of Baja California Sur (Squires, 1959). According to Glynn and Ault (2000), maximum shelf depths where coral
29 colonies or reefs have been observed, from the Golfo de California to Panama, are ~10 m or less. This important observation
30 provides a third relative sea level indicator; where fossil hermatypic corals are found in growth position, sea level was likely
31 no higher than ~10 m above that elevation.

32 **3 Elevation measurements and geochronology**

33 **3.1 Elevation measurements**

34 Virtually all of the studies cited herein provide measurements of the elevations of the RSL indicators. In most studies that
35 were conducted before approximately 2010, measurements were typically made using contours on topographic maps, hand
36 level and/or metered tape, transit and stadia rod, or barometric altimeter. For these studies, unless uncertainties are reported
37 in the original manuscript (or where the shoreline angle elevation range is given), elevation uncertainties are assumed to be
38 20% of the original elevation. This procedure assumes that higher elevation shoreline angles will have greater uncertainties
39 and attribution to an appropriate sea level datum. After approximately 2010, most studies provide elevation measurements
40 done by either handheld or differential Global Positioning System (GPS) instruments (Table 2). Where elevation
41 measurements were made with a handheld GPS instrument, uncertainties can be substantial, and here it is assumed that
42 measurement errors are within ± 3 m of the reported value. For measurements made with a differential GPS instrument,
43 uncertainties are those given in the original study; if not reported, measurement errors are assumed to be within ± 0.5 m.

44 **3.2 Geochronology**

45 All of the RSL indicators that represent MIS 5e on the Pacific Coast of North America considered here have
46 geochronological constraints based on either direct numerical dating using uranium-series (U-series) methods on corals or the
47 correlated-age method of amino acid geochronology, with ties to nearby U-series-dated (coral) localities. As a result, each
48 RSL data point in the database is associated with one or more fossil samples dated with either U-series or amino acid
49 geochronology. Luminescence methods have not been widely applied in this region, although the study by Grove et al. (1995)

50 in the Tomales Bay area provides an important exception. U-series dating of mollusks was once considered a promising
51 method for dating marine terrace fossils in California, but the study by Kaufman et al. (1971) has shown convincingly that
52 mollusks do not take up U during growth, and frequently behave as open systems with respect to U and its daughter products.
53 Thus, early studies that have attempted to date marine terraces by this method are not considered reliable. More recently,
54 cosmogenic isotopes have been attempted in developing chronologies for marine terraces in California (Perg et al., 2001).
55 This method, while promising in theory, requires careful discrimination of which sediments are sampled for analysis. In a
56 study by Perg et al. (2001), ages derived for the terraces near Santa Cruz, California, do not agree with U-series ages on marine
57 terrace corals from the same area (Muhs et al., 2006). The latter investigators speculated that the sediments analyzed by Perg
58 et al. (2001) were likely taken from the terrestrial deposits overlying the marine terrace deposits, which explains the younger
59 than expected cosmogenic ages. Finally, the unique altitudinal-spacing method of Bull (1985) has been applied to marine
60 terraces on the Pacific Coast of North America. Terraces correlated to MIS 5e using this method are not considered in the
61 present review, because Bull's (1985) method assumes that the sea level history derived from the Huon Peninsula of New
62 Guinea is a faithful representation of sea level history on all coastlines around the world (this issue is reviewed in more detail
63 below).

64 3.3.1 Uranium-series dating

65 Uranium-series dating is based on the fortunate characteristic of corals (Fig. 8) to take up small amounts of U (^{238}U , ^{235}U ,
66 ^{234}U) from seawater into their aragonite skeletons during growth. The U assimilated by corals is in isotopic equilibrium with
67 seawater. In contrast, Th and Pa are very insoluble elements, and therefore ocean water contains essentially no dissolved Th
68 or Pa. Thus, ^{230}Th and ^{231}Pa atoms, absent in living corals, accumulate in a fossil, due to decay of ^{234}U and ^{235}U , respectively.
69 These two "daughter-deficient" methods utilize daughter/parent activity ratios ($^{230}\text{Th}/^{234}\text{U}$ and $^{231}\text{Pa}/^{235}\text{U}$) that begin with 0 in
70 living corals and continue to increase in a fossil until equilibrium values of 1.0 are reached. In addition, ^{234}U is present in
71 seawater with an ~14-16% greater activity than ^{238}U (i.e., the $^{234}\text{U}/^{238}\text{U}$ activity value in seawater is ~1.15). In a fossil coral,
72 the $^{234}\text{U}/^{238}\text{U}$ activity value decreases down to an equilibrium value of 1.0 over time, resulting in a third clock, a "daughter-
73 excess" method.

74 Both solitary and colonial corals take up U from seawater during growth, usually in amounts ranging from 2-3 ppm,
75 although some genera of corals (notably species of *Acropora*) take up U in amounts ranging from 3-4 ppm. Along the northern
76 part of the Pacific Coast of North America, from Oregon to Baja California, the most common species used for U-series dating
77 is the solitary coral *Balanophyllia elegans* (Fig. 8). Based on studies of living and dead-collected modern specimens, *B.*
78 *elegans* takes up some additional U after death, but apparently does so from seawater while still submerged and in isotopic
79 equilibrium with U in the ocean (Muhs et al., 2002a, 2006). Farther south, where colonial, hermatypic corals are found, species
80 of the genera *Pocillopora* (Fig. 8) and *Porites* are the taxa most commonly used for U-series dating. In practice, the two clocks
81 used most commonly in U-series dating are $^{230}\text{Th}/^{234}\text{U}$ and $^{234}\text{U}/^{238}\text{U}$. Because of the laboratory challenges in using a Pa spike,
82 few laboratories measure $^{231}\text{Pa}/^{235}\text{U}$. It is a common practice to assess $^{230}\text{Th}/^{234}\text{U}$ ages by plotting measured $^{230}\text{Th}/^{238}\text{U}$ values
83 against measured $^{234}\text{U}/^{238}\text{U}$ values, along with the expected isotopic evolution pathways, assuming initial $^{234}\text{U}/^{238}\text{U}$ values in

84 seawater. In Figure 9a, such an array is shown for corals from Bermuda and the pathways expected from initial seawater
85 values (for $^{234}\text{U}/^{238}\text{U}$ 1.140-1.155; shown here from 1.140-1.160; for $^{230}\text{Th}/^{234}\text{U}$, the initial value is 0.0). Corals that follow
86 these expected isotopic evolution pathways yield ages that likely have minimal bias and can be considered to have had mostly
87 closed-system histories with respect to U-series nuclides. In Figure 9b, what is shown is a much more common situation with
88 corals from the Pacific Coast of North America, with examples from 1st and 2nd terraces on San Nicolas Island, California.
89 While some corals that indicate a closed-system history, similar to Bermuda, others plot above the closed-system evolution
90 pathways. This indicates an open-system history with respect to U-series isotopes in these corals, likely due to recoil-derived
91 additions of ^{230}Th and ^{234}U from dissolved U in water passing through the host sediment. An alternative method of assessing
92 degree of closed-system history of fossil corals is to plot the apparent $^{230}\text{Th}/^{238}\text{U}$ age as a function of its back-calculated initial
93 $^{234}\text{U}/^{238}\text{U}$ value, using the measured $^{234}\text{U}/^{238}\text{U}$ value and the apparent age. Examples of this approach are given in Figure 10,
94 where samples, if they have experienced a closed-system history, should fall within the blue-shaded bands that define the range
95 of variability of modern seawater. As is evident from the plots shown in Figure 10, both solitary corals from the Pacific Coast
96 of North America, and colonial corals from Barbados are prone to open-system histories, but some corals show good evidence
97 of a likely closed-system history. In the examples shown here, it would appear that those corals with closed-system histories
98 on the Pacific Coast have an age range of ~124 ka to ~114 ka.

99 In examining U-series data from corals of reef terraces on Barbados, Gallup et al. (1994) noted that even with open-system
00 histories on isotope evolution plots, a roughly linear trend was observed, with corals that plotted farther above the closed-
01 system pathway showing a bias to older apparent ages. On the Pacific Coast of North America, the same kind of trend is seen
02 as that on Barbados (see Fig. 9a), indicating that this may be a general condition in the near-surface environment where fossil
03 corals are found, despite substantial differences in climate, soil and groundwater hydrology, and composition of surrounding
04 terrains. Nevertheless, noting this typically linear trend on Barbados, Gallup et al. (1994) suggested that extrapolation of linear
05 trends back to a closed-system composition could yield an approximate age for a given terrace. This is also part of the basis
06 of the open-system method of U-series age correction devised by Thompson et al. (2003).

07 Because of the analytical challenges in determining $^{231}\text{Pa}/^{235}\text{U}$ ages, it has become a common practice within the U-series
08 geochronology community to assess the reliability of $^{230}\text{Th}/^{234}\text{U}$ ages with use of the back-calculated $^{234}\text{U}/^{238}\text{U}$ values and a
09 comparison to modern seawater. Although in principle this is an appropriate cross-check, it is not completely reliable. Studies
10 by Gallup et al. (2002) and Cutler et al. (2003) on corals from Barbados and New Guinea showed that some corals that
11 demonstrated concordant $^{231}\text{Pa}/^{235}\text{U}$ and $^{230}\text{Th}/^{234}\text{U}$ ages did not show back-calculated $^{234}\text{U}/^{238}\text{U}$ values within the range of
12 modern seawater. Conversely, some corals that did show back-calculated $^{234}\text{U}/^{238}\text{U}$ values within the range of modern seawater
13 did not have concordant $^{231}\text{Pa}/^{235}\text{U}$ and $^{230}\text{Th}/^{234}\text{U}$ ages.

14 Marine terrace corals dated by U-series methods are found within the WALIS database and/or within the compilation of
15 Chutcharavan and Dutton (2021). Generalized information about each U-series-dated locality can be found in Table S1.

16 **3.2.2 Amino acid geochronology**

17 In the absence of corals in a marine terrace deposit or emergent reef, mollusks, both bivalves and gastropods, can be used
18 for amino acid geochronology. For marine terraces on the Pacific Coast of North America, amino acid geochronology was
19 pioneered by John F. Wehmiller and his colleagues (Wehmiller et al., 1977a; Lajoie et al., 1980; Kennedy et al., 1982;
20 Wehmiller, 1982, 1992, 2013a, 2013b). The method is based on the observation that living organisms contain only amino
21 acids with the "L" (*levo*, or left-handed) configuration. Upon death of an organism, amino acids of the L configuration convert
22 to amino acids of the "D" (*dextro*, or right-handed) configuration, a reaction called racemization. Racemization is a reversible
23 process that results in increased D/L ratios in a fossil until an equilibrium ratio of 1.0 is reached. A related process, called
24 epimerization, is conversion of the amino acid L-isoleucine (found in living organisms) to D-alloisoleucine (not found in living
25 organisms). Epimerization, like racemization, begins with D-alloisoleucine/L-isoleucine values of 0.0 in a fossil, but this ratio
26 increases over time until an equilibrium value of 1.25-1.30 is reached (Miller and Mangerud, 1985). Some of the fossils that
27 have been most commonly used on the Pacific Coast of North America are the bivalves *Saxidomus* and *Chione* (in protected,
28 sandy or muddy, bay environments) and *Tegula* (in high-energy, rocky-shore environments), shown in Figure 8.

29 Amino acid values in fossil mollusks can be used for lateral correlation of marine terrace deposits, exploiting the fact that
30 both racemization and epimerization rates increase with higher diagenetic temperature histories. This means that D/L values
31 in shells reach equilibrium values more quickly in warmer climates than they do in cooler climates. Thus, on north-south-
32 trending coastlines in the Northern Hemisphere, such as the Pacific Coast of North America, shells in terrace deposits at more
33 southerly localities are expected to have higher D/L values than shells of the same genus but of similar age in cooler, northerly
34 localities. When D/L values are arrayed on a latitudinal plot or a plot of mean annual air temperatures, there should be a south-
35 to-north decrease in D/L values in shells of the same age. In practice, some localities along such an array have independent
36 age control from U-series dating of corals. If so, then a shore-parallel correlation of locality to locality, from south to north,
37 can be accomplished, yielding an "aminozone" corresponding to the age of the independently dated localities. Shells from
38 younger terraces would define an aminozone below such a zone and older terraces would define an aminozone above it.

39 The first major attempts at aminostratigraphic correlation along the Pacific Coast using the approach just described were
40 those by Wehmiller et al. (1977a), Kennedy et al. (1982), and Wehmiller (1982). The north-to-south correlation of terraces
41 from Kennedy et al. (1982) is shown in Figure 11a, along with three U-series-dated localities that serve as calibration points.
42 Kennedy et al. (1982) also noted that most localities correlated to either MIS 5a or MIS 3 hosted terrace faunas with cool-
43 water aspects, whereas those correlated to MIS 5e had warm-water faunas, or at least faunas that were "neutral," lacking cool
44 or water-water taxa. In the time since the Kennedy et al. (1982) study was conducted, more U-series ages on coral have been
45 reported (~120 ka, ~80 ka, and ~47 ka), many of which support the original aminostratigraphic correlations (Fig. 11b).
46 Nevertheless, some localities are now known to host mixes of warm and cool faunas and at least two of these have mixes of
47 ~120 ka (MIS 5e) and ~100 ka (MIS 5c) corals (Fig. 11b). This issue is discussed in more detail below.

48 Even with some concerns, amino acid geochronology has been shown to be a very powerful coast-parallel correlation tool.
49 Even within the limited geographic range of central California to northern Baja California, there is enough of an air temperature

50 gradient that aminostratigraphic correlation can be accomplished. At a given locality where two terraces are found (one at a
51 low elevation, one at a higher elevation), MIS 5a and MIS 5e terrace deposits can usually be distinguished from one another
52 (Fig. 12). Furthermore, lateral correlation of MIS 5e and MIS 5a deposits from central California to northern Baja California
53 can be made, anchored by localities with U-series ages on corals.

54 Similar to U-series-dated marine terrace corals, those terrace localities correlated to MIS 5e with amino acid racemization
55 or epimerization methods are found within the WALIS database, along with linkage to the U-series-dated localities that served
56 as calibration. Generalized information about each locality correlated to MIS 5e with amino acid geochronology can be found
57 in Table S2.

58 **3.2.3 Zoogeographic aspects of terrace faunas**

59 In a pioneering study of marine terraces on the Pacific Coast of North America, Kennedy et al. (1982) used the
60 aminostratigraphic approach described above to extend earlier work by Wehmiller et al. (1977a). Both studies established
61 that the lowest marine terrace along the Pacific Coast of North America is not the same age at all localities, due to varying
62 rates of uplift from one reach of coast to another. In addition, Kennedy et al. (1982) noted that localities dated (by U-series
63 on coral) to or correlated with MIS 5e host either zoogeographically "neutral" molluscan fossil faunas or faunas that contain
64 extralimital southern species. In contrast, localities that were either dated or correlated to the ~80 ka MIS 5a host molluscan
65 fossil faunas with several extralimital northern species (Fig. 11). Extralimital species (or northward or southward-ranging
66 species) are those that, while extant, do not live at a particular locality at present, but are found either entirely or mostly to the
67 north (cool waters in this region) or to the south (warmer waters in this region). An example of a locality, dated to ~130 ka by
68 thermoluminescence (Grove et al., 1995), is the marine deposit in Tomales Bay, north of San Francisco, California. This
69 deposit contains many "neutral" species, i.e., those that still live in the area at present, but also host a large number of
70 extralimital southern and southward-ranging species (Fig. 13). In contrast, the Davenport terrace in the Santa Cruz, California
71 area, dated to ~80 ka by U-series methods on corals (Muhs et al., 2006), hosts only one southward-ranging species, but several
72 extralimital northern and northward-ranging species. Warmer waters off California during MIS 5e and cooler waters during
73 MIS 5a are consistent with the zoogeographic aspects of planktonic foraminiferal faunas found in deep-sea cores (Kennett and
74 Venz, 1995) and with sea surface temperatures (SST) derived from alkenones (Herbert et al., 2001; Yamamoto et al., 2007).

75 **4 Relative Sea Level indicators**

76 Relative sea level indicators from the Pacific Coast of North America for MIS 5e and all pertinent data related to them are
77 given in Table S1 and Figure 14 (U-series-dated coral-bearing localities) and Table S2 and Figure 15 (localities correlated to
78 MIS 5e using aminostratigraphy). In the sections that follow, the regions these localities are from are discussed with respect
79 to the nature of the sea level record, as this differs from region to region. Within the course of these discussions, previous
80 studies are examined and the basis for the age assignments is discussed critically. For simplicity, the review of the regions is
81 taken from north to south. In the text that follows, there is an indication near each site discussed of what the unique RSL
82 identification is, corresponding to the WALIS database.

83 4.1 Southwestern Canada

84 Records of marine deposits dating to MIS 5e are difficult to find on the coast of British Columbia. Erosion by repeated
85 advances of the Cordilleran ice sheet has likely removed much of the potential record. Furthermore, the sedimentary record
86 that does exist is highly complex, due to rapid sedimentation rates, active tectonics, and glacial isostatic adjustment (GIA)
87 effects. Mollusk-bearing glaciomarine sediments were deposited in lowland areas adjacent to coastal British Columbia or in
88 Puget Sound when isostatic depression of these areas allowed inflow of ocean waters. Thus, at least some of the marine record
89 that is now emergent is not strictly "interglacial," but likely occurred at the transition between a glacial period and the following
90 interglacial period.

91 In southwestern Canada, most investigators have hypothesized, from the stratigraphic sections that have been studied, that
92 the main record of MIS 5e is the Muir Point Formation (Hicock and Armstrong, 1983; Alley and Hicock, 1986; Hicock,
93 1990; Clague et al., 1992). On Vancouver Island in British Columbia (Fig. 16), the Muir Point Formation consists of gravel,
94 sand, and silt, with abundant peat and wood layers, suggesting a mostly terrestrial origin, but Hicock and Armstrong (1983)
95 hypothesize an alluvial fan-to-floodplain-to-coastal plain-to-delta sequence, based on the sediment facies. Indeed, Alley and
96 Hicock (1986) and Hicock (1990) report minor amounts of marine dinoflagellate cysts in a part of the Muir Point Formation,
97 implying tidal or estuarine conditions, and these investigators infer a paleo-sea level of at least +10 m, relative to present. A
98 last interglacial origin for the Muir Point Formation was hypothesized early in the study of this formation by its stratigraphic
99 position: it has organic materials that date to >40 ka and has normal polarity, but is underlain by older till and overlain by mid-
00 Wisconsin (MIS 3) Cowichan Head Formation sediments, in turn overlain by Vashon Till dating to the Fraser Glaciation (=late
01 Wisconsin, or MIS 2) (Alley and Hicock, 1986; Hicock, 1990). Vegetation evidence also suggests a climate at least as warm
02 as today's, based primarily on the abundance of thermophyllous *Pseudotsuga* (Douglas fir) pollen, implying interglacial
03 conditions (Hicock and Armstrong, 1983; Alley and Hicock, 1986; Hicock, 1990). An MIS 5e age is permitted by optically
04 stimulated luminescence (OSL) ages of 119 ± 9 ka and 112 ± 11 ka from the Muir Point Formation at and near its type section
05 (Lian et al., 1995). Because more study is needed for assessment of the age of the Muir Point Formation, no specific entry in
06 the WALIS database was attempted here.

07 4.2 Washington, USA

08 Only two fossil-bearing localities are candidates for MIS 5e deposits in the State of Washington, one in Puget Sound and
09 the other on the outer coast, at Willapa Bay (Fig. 16). Both have had a confusing and/or controversial history of study.

10 4.2.1 Whidbey Island, Puget Sound

11 As is the case with British Columbia, the southern Puget Sound area, within the boundaries of Washington State, has been
12 subjected to rapid sedimentation rates, active tectonics, and GIA effects, as well as removal of much of the geologic record,
13 due to advances and retreats of the Cordilleran ice sheet. Also similar to British Columbia, the main geologic unit that most
14 investigators agree records the last interglacial period (MIS 5e) is not primarily a marine deposit at all, but a terrestrial deposit
15 called the Whidbey Formation. Hansen and Mackin (1949) were among the first to study the formation, noting that it occurred

16 stratigraphically below deposits dating to the last glacial period (i.e., MIS 4 through MIS 2), and that it hosted pollen indicating
17 an interglacial vegetation similar to that of the present. Easterbrook et al. (1967) were the first investigators to apply the formal
18 name Whidbey Formation to the pollen-bearing unit studied by Hansen and Mackin (1949) and designated the type locality
19 on coastal bluffs of southwestern Whidbey Island (Fig. 16). At the type section, Easterbrook et al. (1967) and Easterbrook
20 (1968, 1969) noted that the Whidbey Formation is underlain by what is called Double Bluff Drift, consisting of till and
21 glaciomarine sediments. At this locality, the Whidbey Formation is overlain by glacial deposits of Possession (MIS 4?) and
22 Vashon (MIS 2) age. Easterbrook et al. (1967) conducted pollen analyses of Whidbey Formation sediments and concluded
23 that the vegetation implied an interglacial climate similar to the present. They also reported ages that showed the unit was
24 beyond the range of radiocarbon dating. More detailed pollen work was conducted by Heusser and Heusser (1981), who
25 reached the same conclusions about past climate conditions. Karrow et al. (1995) reported on nonmarine fossils in the Whidbey
26 Formation, including mollusks, ostracodes, insects, fish, vertebrates, and plant macrofossils. Their interpretations are similar
27 to those of Hansen and Mackin (1949), Easterbrook et al. (1967), and Heusser and Heusser (1981), that the deposit likely
28 represents an interglacial period with a degree of warmth similar to that of the present. It is important to note that in all of the
29 studies just cited, the Whidbey Formation is described as a *terrestrial* deposit, likely formed as floodplain sediments. None
30 of the studies cited here mention the presence of marine fossils within the deposit. Later studies have all confirmed a likely
31 MIS 5 age for the Whidbey Formation, based on thermoluminescence (TL) dating (151 ± 43 ka to 102 ± 38 ka; Berger and
32 Easterbrook, 1993), optically stimulated luminescence (OSL) dating (107 ± 8 ka; Lian et al., 1995), and $^{40}\text{Ar}/^{39}\text{Ar}$ dating of
33 plagioclase from pumice within the formation (128 ± 9 ka; Dethier et al., 2008).

34 With the advent of amino acid geochronology, several studies presented data on some of the marine-shell-bearing deposits
35 of Whidbey Island. Most of these studies focus on a shell-bearing deposit along Admiralty Bay, on the west coast of Whidbey
36 Island. This deposit is visible in an ~17 m thick coastal exposure (~2 m above sea level) of diamicton and/or glaciomarine
37 sediment composed of gravel, sand, and silt, with a layer of marine fossils, dominated by *Saxidomus gigantea*, in its uppermost
38 part (Polenz et al., 2009). The upper contact of this complex deposit is obscured by a recent landslide, but a short distance
39 inland and at higher (~40 m to ~60 m) elevations, glaciomarine deposits of the Everson Interstade, outwash of the Fraser
40 Glaciation, and till of the Vashon Stade (all of MIS 2 age) are mapped (Polenz et al., 2009) and likely overlie the shell-bearing
41 deposits exposed at lower elevation. Kvenvolden et al. (1980) reported that this fossiliferous deposit lies stratigraphically
42 between the Whidbey Formation and "middle Wisconsin sediments" and used amino acid ratios in *Saxidomus gigantea* to
43 estimate an age of ~80 ka. Their study, however, presents no stratigraphic evidence of the Whidbey Formation being exposed
44 at Admiralty Bay. Blunt (1982) analyzed shells from the same locality as Kvenvolden et al. (1980) and another locality ~150
45 m to the north. He used kinetic modeling to derive an age range of 77 ka to 99 ka for the locality studied by Kvenvolden et
46 al. (1980) and 75-110 ka for the newer locality. In a later study derived primarily from data in Blunt (1982), Blunt et al. (1987,
47 p. 331-332) described the Admiralty Bay deposit as belonging to the Possession Glaciation (which postdates the Whidbey
48 Formation), but later in the same paper (p. 340 and p. 346) said that the deposits are correlated with the Whidbey Formation.
49 These investigators also used *Saxidomus gigantea* and kinetic modeling to estimate ages of ~96 ka and ~107 ka for the deposit,

50 apparently pooling the two localities. Using these data, Easterbrook correlated the deposit at Admiralty Bay with the Whidbey
51 Formation. Kennedy et al. (1982) used the *Saxidomus gigantea* single-shell analysis in Kvenvolden et al. (1980) to estimate
52 an aminozone-derived, correlated age of 80 ka, and apparently used the pooled *Saxidomus gigantea* data in Blunt (1982) to
53 estimate an aminozone-derived, correlated age of 120 ka. In addition, Kennedy et al. (1982) reported that the 80 ka locality
54 hosts a cool-water fauna and the ~120 ka locality hosts a warm-water fauna. This is puzzling, because no faunal data are given
55 in Kvenvolden et al. (1980) or Kennedy et al. (1982), although Blunt (1982) reports a single extralimital northern or at least
56 northward-ranging species, *Mya truncata*, in the fossil deposit at Admiralty Bay. Furthermore, the two localities are only ~150
57 m apart and occur at roughly the same elevation according to Blunt (1982). Finally, Polenz et al. (2009) presented
58 sedimentological data indicating that the deposits at Admiralty Bay have a glaciomarine origin. These investigators correlated
59 the deposit either to the pre-last interglacial Double Bluff Glaciation (their favored option) or the post-last interglacial
60 Possession Glaciation. In my own examinations of the deposits at Admiralty Bay, I have seen no evidence for more than one
61 stratigraphic unit. I also agree with Polenz et al. (2009) that the shell-bearing deposit exposed there is likely glaciomarine
62 drift, dating to the transition between the penultimate glacial period (MIS 6), represented by the Double Bluff unit, and MIS
63 5e. It is likely an older equivalent of the shell-bearing glaciomarine drift of the late, last glacial Fraser glaciation, a unit called
64 the Everson glaciomarine deposits. For this reason, this locality has not been entered into the WALIS database.

65 **4.2.2 Willapa Bay**

66 The only other emergent, fossil-bearing locality that is a candidate for an MIS 5e deposit in Washington State is along the
67 inner shores of Willapa Bay (**RSL ID 3684**) (Fig. 16). Near Bay Center, sea cliffs expose marine sediments that are richly
68 fossiliferous (Fig. 17). Addicott (1966) reported the fossil fauna from this locality, which consists mostly of bivalves, and no
69 taxa are extralimital, or even northward or southward ranging. Kvenvolden et al. (1979) provided the first published amino
70 acid data from this area. They recognized four stratigraphic units (I, intertidal; II, subtidal; III, subaerial; and IV, subtidal,
71 from oldest to youngest). Their unit IV is the thickest and apparently the most extensive deposit, interpreted to have an
72 estuarine origin; the top of this unit defines a marine terrace surface, at an elevation of ~13 m. Almost all of the *Saxidomus*
73 *gigantea* specimens they analyzed are from this youngest deposit. Using assumed calibration ages of ~68 ka and ~100 ka for
74 the lowest terrace at Santa Cruz, California (Bradley and Addicott, 1968), which also hosts fossil *Saxidomus gigantea*,
75 Kvenvolden et al. (1979) used linear kinetic modeling (taking temperature differences into account) to generate age estimates
76 of 190 ± 40 ka for units I and II, and 120 ± 40 ka for unit IV, which they correlated to MIS 7 (I and II) and MIS 5 (IV). Their
77 terrestrial unit III was interpreted to have formed when sea level lowered during MIS 6. It is now known that U-series ages
78 on mollusks, including the ~68 ka and ~100 ka ages for Santa Cruz reported by Bradley and Addicott (1968), are not reliable
79 (Kaufman et al., 1971). Nevertheless, reliable U-series ages on corals from the same terrace yielded ages in between these,
80 averaging about 80 ka (Muhs et al., 2006). Thus, the newer ages, if used as calibration, would not change the original kinetic
81 model ages for the Willapa Bay deposits. In any case, Wehmiller (1981) challenged Kvenvolden et al.'s (1979) age estimates,
82 arguing that nonlinear kinetic modeling is more appropriate for numerical ages using amino acid data. Using nonlinear kinetic
83 modeling, Wehmiller (1981) recalculated the ages of units I/II and IV at Willapa Bay to be 300 ± 50 ka and 70 ± 15 ka,

84 respectively, suggesting correlation with MIS 9 and MIS 5a. Kvenvolden et al. (1981) countered that nonlinear kinetics could
85 be applied to amino acid values within the ranges of what their samples yielded, and also noted that Wehmiller's (1981) age
86 estimates would require a much more complex geologic history than their age estimates. Kennedy et al. (1982) reported new
87 amino acid values in *Saxidomus* from unit IV and using a lateral correlation (aminozone) approach, considered that the unit
88 IV deposits at Willapa Bay were of MIS 5a age, in agreement with Wehmiller (1981). They also reported that the fauna at the
89 Bay Center locality of unit IV hosted cool-water forms, although Addicott (1966) reported no extralimital species or
90 northward-ranging species. The cool-water aspect of the fauna at Bay Center is apparently based on the identification of *Mya*
91 *japonica* in these deposits, reported by Kennedy (1978). Although *M. japonica* was once considered to range only in the
92 Arctic seas, from Japan to Nome, Alaska (Abbott, 1974), Coan et al. (2000) consider that *M. japonica* does not have differences
93 with *M. arenaria* that are sufficient to merit specific status. If so, then there are no extralimital northern species in the fauna
94 of unit IV at Willapa Bay and the assemblage as a whole can be considered zoogeographically "neutral." Given all the
95 uncertainties in what has been reported thus far for Willapa Bay, it seems likely that unit IV of Kvenvolden et al. (1979) could
96 date to MIS 5e, but more geochronological information is needed to be certain of this.

97 **4.3 Oregon, USA**

98 Moving south from Washington, coastal Oregon is where the dominant geomorphic expression of MIS 5e shorelines as
99 erosional marine terraces begins. The coast of Oregon is within the Cascadia subduction zone (Fig. 2) and most of it can be
00 characterized as a high-wave-energy environment. Thus, erosional marine terraces are common landforms along a substantial
01 amount of the coast, particularly in the central and southern parts of Oregon (Fig. 16). A pioneering study by Griggs (1945)
02 involved the mapping and naming of the lowest three marine terraces in southern Oregon and the terrace names are still in use
03 today. More recently, detailed mapping of marine terraces along the Oregon coast has been conducted primarily by H.M.
04 Kelsey and his colleagues and students (Kelsey, 1990; McNelly and Kelsey, 1990; Bockheim et al., 1992; Kelsey and
05 Bockheim, 1994; Kelsey et al., 1996). Candidate landforms for some or all substages of MIS 5 are the lowest three terraces
06 found along much of the central and southern Oregon coast. Kennedy et al. (1982) inferred that the lowest of these, the Whisky
07 Run terrace near Coquille Point (Figs. 18, 19) likely correlated to MIS 5a because of a U-series age of ~72 ka on a coral from
08 its deposits, as well as relatively low D/L leucine values in *Saxidomus gigantea*, and a cool-water aspect to the terrace fauna
09 (Zullo, 1969; Kennedy, 1978). Later, both alpha-spectrometry and TIMS U-series ages of corals from the Whisky Run terrace
10 confirmed an age of ~80 ka, and a more extensive cool-water fauna was reported (Muhs et al., 1990, 2006). Higher terraces
11 are present in this area and north to Cape Arago, named the Pioneer, Seven Devils, and Metcalf terraces (lowest to highest),
12 mapped by McNelly and Kelsey (1990). Based on the ~80 ka age of the Whisky Run terrace, McNelly and Kelsey (1990)
13 inferred that MIS 5e is represented by the Seven Devils terrace, with the Pioneer terrace correlated to MIS 5c.

14 Farther south, the lowest terrace at Cape Blanco also hosts a cool-water fauna (Addicott, 1964a), but based on amino acid
15 values, Kennedy et al. (1982) interpreted this terrace to be of post-MIS 5 age, possibly as young as MIS 3. In a later study,
16 Kelsey (1990) remapped the terraces in this area and named this the Cape Blanco terrace (Fig. 20). He correlated this with the
17 Whisky Run terrace at Coquille Point, supported by new amino acid and oxygen isotope values in *Saxidomus gigantea* from

18 Cape Blanco (Muhs et al., 1990). Kelsey also mapped and named higher landforms above the Cape Blanco terrace, the Pioneer,
19 Silver Butte, and Indian Creek, from lowest to highest (Figs. 18, 19). He considered the Pioneer terrace to represent the MIS
20 5c high-sea stand and the Silver Butte terrace to represent the MIS 5e high stand. Amino acid data given by Muhs et al. (1990)
21 support the correlation of the Pioneer terrace to MIS 5c, but no fossils have yet been found on the Silver Butte terrace.

22 North of Coquille Point, near Newport, Oregon, Kennedy et al. (1982) reported amino acid values in *Saxidomus gigantea*
23 from a low marine terrace near Newport jetty and a higher terrace at Yaquina Bay. On the basis of these amino acid ratios and
24 a cool-water fauna (lower terrace) and a warm-water fauna (higher terrace), Kennedy et al. (1982) correlated these terraces
25 with MIS 5a and MIS 5e, respectively. Later mapping by Kelsey et al. (1996) identified these as the Newport (lower) and
26 Yachats (higher) terraces, respectively, with an intermediate-elevation landform they named the Wakonda terrace. They
27 correlated the Newport, Wakonda, and Yachats terraces with MIS 5a, 5c, and 5e, respectively.

28 Summarizing all these studies, U-series, amino acid, oxygen isotope, and faunal data all support a correlation of the lowest
29 marine terrace at Newport, Coquille Point, and Cape Blanco to MIS 5a (Fig. 19). At Cape Blanco, similar amino acid ratios
30 and oxygen isotope data correlate the Pioneer terrace with MIS 5c. At Newport-Yaquina Bay, the Yachats terrace (**RSL ID**
31 **3685**) is correlated to MIS 5e by amino acids and faunal data. Lack of fossils precludes correlation of intermediate and higher
32 terraces at all these localities. To address this problem, Kelsey and Bockheim (1994) used degree of soil development to
33 correlate undated terraces in all three areas, plus a fourth area in southernmost Oregon, near Cape Ferrelo (Fig. 16), where all
34 terraces lack fossils. With the generation of a soil development index that utilizes time-dependent soil properties (e.g., Bt
35 horizon thickness, color, texture, clay content), they identified, from north to south, the Yachats, Seven Devils, Silver Butte,
36 and Gowman terraces as the likely candidates for records of the MIS 5e high-sea stand.

37 **4.4 Northern California, USA**

38 **4.4.1 Crescent City coastal plain**

39 Surprisingly few studies of marine terraces have been undertaken in northern California, in part because fossil-bearing
40 occurrences that would permit dating are rare. Northernmost California is within the Cascadia subduction zone, similar to
41 coastal Oregon (Fig. 21). About 25 km south of the Oregon border, marine terraces have been studied for decades on the
42 Crescent City coastal plain. Maxon (1933) named all the marine terrace deposits in this area collectively the Battery Formation,
43 and he also noted the presence of fossil invertebrates in the deposits. Similarly, Delattre and Rosinski (2012) mapped deposits
44 of the entire Crescent City coastal plain as the Battery Formation. The first attempt at dating these deposits was by Kennedy
45 et al. (1982), who presented amino acid data from *Saxidomus gigantea* from low-elevation (~7 m) sea cliff exposures in
46 southern Crescent City. These investigators also reported a cool-water fauna from this low-elevation terrace and on the basis
47 of D/L leucine values, correlated the terrace with MIS 5a.

48 In contrast to Maxon (1933) and Delattre and Rosinski (2012), Polenz and Kelsey (1999) recognized three marine terraces
49 in this area (Qpm3, Qpm2, and Qpm1, from youngest to oldest), differentiated by subtle elevation changes and differing
50 degrees of soil development, following the approach used by Kelsey and Bockheim (1994) in southern Oregon. Polenz and
51 Kelsey (1999) correlated the three terraces they mapped (Qpm3, Qpm2, Qpm1) with MIS 5a, 5c, and 5e, respectively, although

52 they noted that Qpm1 could correlate with MIS 7. The localities studied by Kennedy et al. (1982) are situated on what Polenz
53 and Kelsey (1999) mapped as Qpm2, the terrace they correlated with MIS 5c. It should be noted, however, that it is
54 questionable whether amino acid ratios can distinguish ~80 ka deposits from ~100 ka deposits, and cool-water faunas are
55 expected from terraces of either age, based on alkenone paleotemperature data from a nearby deep-sea core (ODP 1020) studied
56 by Herbert et al. (2001). Thus, the best evidence for a possible MIS 5e shoreline in this area is the Qpm1 terrace mapped by
57 Polenz and Kelsey (1999), found mostly inland of the younger terraces. This terrace has maximum platform elevations of ~29
58 m to ~15 m.

59 **4.4.2 Trinidad Head area**

60 Marine terraces are scarce between Crescent City and along a coastal reach ~60 km to the south. However, in the Trinidad
61 Head area (Fig. 21), there are multiple marine terraces, well expressed geomorphically. This area, like the Crescent City
62 coastal plain, is also within the Cascadia subduction zone. Carver (1992) mapped seven terraces in this area, with additional
63 undifferentiated higher elevation terraces. Based on ages assigned from the oxygen isotope record and an untested assumption
64 of a constant uplift rate, Carver (1992) gave estimated terrace ages that were also followed by Delattre and Rosinski (2012).
65 The rationale for these age assignments is reported to be from degree of soil development and thermoluminescence (TL) ages
66 reported by Berger et al. (1991). However, the method by which ages from degree of soil development are derived is not
67 described and all of the TL ages reported by Berger et al. (1991) are either not consistent with Carver's (1992) mapping or date
68 younger deposits that overlie the marine terrace deposits. Although not mapped, McCrory (2000) also presented shore-parallel
69 terrace profiles for seven marine terraces in this area, with shoreline angle elevations ranging from ~15 m to ~255 m. No
70 numerical ages are available, but McCrory (2000) used a graphical method of estimating terrace ages, as described by Lajoie
71 (1986). Finally, Padgett et al. (2019) remapped the terraces in this area and assigned ages based on degree of soil development
72 and an assumption that the terrace with the most prominent inner edge (their "Surface 3") dates to MIS 5e. They further
73 assumed that the lower-elevation "Surface 1" and "Surface 2" terraces date to MIS 5a and 5c, respectively. Thus, the mapping
74 and age assignments of Carver (1992) and Delattre and Rosinski (2012) disagree with those of Padgett et al. (2019), but it is
75 important to emphasize that none of these studies have any supporting numerical ages. Interestingly, pre-MIS 5e ages are
76 given by Kennedy et al. (1982) for marine deposits in this area, based on D/L values in *Saxidomus gigantea*, but because no
77 geomorphic or stratigraphic data are given, it is not known how these aminostratigraphic data can be linked to the other studies.
78 Much more work needs to be done on dating the terraces in this area, and although it seems likely that an MIS 5e record is
79 present, it cannot be determined at this time which shoreline is representing it.

80 **4.4.3 Eureka-Cape Mendocino area**

81 The Eureka area is situated within the southernmost part of the Cascadia subduction zone, but Cape Mendocino is close to
82 the Mendocino triple junction, where the North America, Gorda, and Pacific plates intersect (Fig. 21). McLaughlin et al.
83 (2000) mapped the geology of the Eureka, California area, as well as the Cape Mendocino area to the south of Eureka. Within
84 the Eureka area itself, these investigators mapped a unit simply called "Qt," which is primarily nonmarine, fluvial terrace
85 deposits, but which also includes shallow marine deposits, including an informally named deposit called the "Hookton marine"

86 unit (Ogle, 1953). Wehmiller et al. (1977b) and Kennedy et al. (1982) reported leucine D/L values in *Saxidomus gigantea*
87 from two localities in the Eureka area (one of which is at an elevation of 15-17 m, **RSL ID 3686**). Both localities are within
88 the "Qt" unit of McLaughlin et al., 2000) that would permit correlation of the host deposits with MIS 5e. One of these localities
89 is reported to host a warm-water fauna and the other is reported to host a zoogeographically "neutral" fauna, but stratigraphic
90 and faunal details are not given. It seems likely that the Eureka area hosts marine deposits that correlate to MIS 5e, but more
91 detailed work is required to confirm this.

92 Between Cape Mendocino and Point Delgada, the region is within the influence of the tectonically active Mendocino triple
93 junction (Fig. 21). Along this rugged part of the coast, numerous terraces have been identified, including some dating to the
94 Holocene and even historic time, described later. For the Pleistocene, terrace elevation transects were reported by Merritts
95 and Bull (1989). No numerical ages are given, but terraces correlated to MIS 5e were reported at elevations of ~150 m (a short
96 distance north of Point Delgada) and at ~250 m (near Punta Gorda). These correlations were made using a graphical correlation
97 method described by Bull (1985). Bull's (1985) method was developed before much was known about the importance of GIA
98 effects, which have been shown to affect the California coast, and this issue is discussed later. Thus, whether the correlations
99 of Merritts and Bull (1989) of the terraces in the Punta Gorda and Point Delgada areas to MIS 5e are valid or not will have to
00 await independent dating.

01 **4.4.4 Laguna Point to Point Arena**

02 Between Laguna Point and Point Arena, marine terraces form the coastal plain area of this part of northern California. This
03 area is south of the Mendocino triple junction and is within the strike-slip tectonic region of the San Andreas Fault zone (Fig.
04 21). Although detailed mapping has not been conducted in this area, general terrace maps are available and a portion of
05 terraced coastline is shown here (Fig. 22). Merritts and Bull (1989) have reported elevations of terrace inner edges on shore-
06 normal transects in this area as well. These investigators report as many as six marine terraces in the Cabrillo Point area, from
07 ~10 m to ~130 m above sea level. Unfortunately, only one locality, thus far, has yielded fossils in this reach of coastline. On
08 the lowest terrace (shoreline angle of ~10 m) at Laguna Point, within MacKerricher State Park (Fig. 14), Kennedy et al. (1982)
09 reported D/L leucine ratios on *Saxidomus gigantea* fragments, as well as a fauna with extralimital northern species. The amino
10 acid ratios on these shells plotted above their ~80 ka aminozone, creating a dilemma: the amino acid data suggested correlation
11 with MIS 5e but the fauna was typical of MIS 5a deposits. Dorothy J. Merritts of Franklin and Marshall College returned to
12 this locality and recovered two solitary corals (*Balanophyllia elegans*) that she submitted to laboratories at the U.S. Geological
13 Survey. For this review, she kindly allowed use of these previously unpublished data. One coral has a low U content and an
14 apparent age of ~156 ka, too old to be considered to be of MIS 5e age, but likely biased old because of U loss. However, the
15 other coral has a U content of 4.82 ±0.10 ppm, a ²³²Th content of 0.06 ppm, a ²³⁰Th/²³²Th value of 180, a ²³⁴U/²³⁸U value of
16 1.0976 ±0.0016, a ²³⁰Th/²³⁸U value of 0.7771 ±0.0024, and an age of 130.4 ±0.9 ka. The back-calculated initial ²³⁴U/²³⁸U
17 value of 1.1411 ±0.0022 is within the range of modern seawater, giving the age of ~130 ka a high degree of confidence. This
18 age is consistent with the age implied by the amino acid data reported by Kennedy et al. (1982). The issue of an MIS 5e terrace

19 such as this, containing a cool-water fauna, is contrary to the general model proposed by Kennedy et al. (1982). These
20 apparently contradictory observations can be reconciled, however, in areas of low uplift rate, by formation of an MIS 5e
21 terrace, followed by reoccupation (and fossil reworking) during the MIS 5c high-sea stand, a topic that is explored in more
22 detail later.

23 Marine terraces at Point Arena and south of it have been mapped by Muhs et al. (2003, 2006), who report three terraces in
24 this area. Corals from their "Qt1," the lowest terrace (~20-25 m), were dated by alpha-spectrometric methods to ~80 ka, or
25 MIS 5a (Muhs et al., 1990, 1994). Later dating of coral from this terrace by TIMS also gave an age of ~80 ka, and
26 paleontological studies yielded a fauna with extralimital northern forms (Muhs et al., 2006). All these results are in good
27 agreement with those of Kennedy et al. (1982), who correlated the terrace to MIS 5a on the basis of D/L leucine values in
28 *Saxidomus gigantea*. The two higher terraces mapped by Muhs et al. (2003) have elevations of ~40-45 m (Qt2) and ~60-65
29 m (Qt3). A reasonable working hypothesis is that Qt3 represents the last interglacial peak (MIS 5e) high-sea stand and that
30 the intermediate Qt2 terrace formed during the MIS 5c high-sea stand. Unfortunately, fossils on both of these terraces have
31 yet to be found, so testing of this hypothesis is not yet possible.

32 4.4.5 Tomales Bay

33 Tomales Bay (RSL ID 3794) has been of interest to geologists because of its unusual configuration, conditioned largely by
34 the fact that the San Andreas Fault zone is situated within the bay, parallel to the bay's long axis (Fig. 23). At a few points on
35 the eastern side of Tomales Bay, there are exposures of a marine deposit called the Millerton Formation, long considered to
36 be of Pleistocene age. On the most recent geologic map of the area, the formation is simply included within what is mapped
37 as "marine terrace deposits" (Graymer et al., 2006). At Toms Point, it is a fossil-rich bed (with abundant *Saxidomus* and
38 *Chione* shells), ~0.5 m thick, overlying a bench cut on Franciscan rocks, and overlain by nonmarine terrestrial deposits. The
39 shell-rich bed is ~8-9 m above sea level at Toms Point. At Millerton Point, the beds are gravelly, ~1.0 m thick, and are rich
40 in *Ostrea* and *Leukomca* (formerly *Protothaca*) shells, all exposed just above modern beach level. Johnson (1962), who
41 conducted the most thorough study of the fossils from the Millerton Formation, noted that several extralimital southern and
42 southward-ranging species are present (Fig. 13), with no northern species, implying water temperatures much warmer than
43 those at Tomales Bay today. Kennedy et al. (1982) noted the warm-water aspect of the fauna in the Millerton Formation, and
44 presented D/L leucine data in *Saxidomus* that fall slightly above their ~120 ka, MIS 5e aminozone. These investigators did not
45 specifically accept or reject a possible MIS 5e age for the formation. Grove et al. (1995) studying the tectonics of the area,
46 reported a TL age of 134 ± 12 ka for the Millerton Formation (analyzed by G.W. Berger). Unfortunately, no analytical data
47 are given for further consideration of this TL age. However, Muhs and Groves (2018) presented D-alloisoleucine/L-isoleucine
48 data for *Chione* from the Millerton Formation, collected at Toms Point. Their data fall into a last interglacial (MIS 5e)
49 aminozone when compared with a south-to-north transect of similar data from *Chione* in Baja California and California. More
50 dating of this important formation would be highly desirable, but the available information indicates that the Millerton
51 Formation along the northeast shores of Tomales Bay represents deposits that can be correlated to MIS 5e.

52 4.5 Central California, USA

53 4.5.1 Point Año Nuevo-Santa Cruz area

54 South of San Francisco, marine terraces dominate many parts of the coast of central California. As noted earlier, it was in
55 this area, east of Santa Cruz (Fig. 21), that Alexander (1953) formulated the modern concept of how marine terraces form on
56 tectonically active coastlines, specifically as landforms cut during interglacial sea-level high stands superimposed on crustal
57 blocks experiencing steady uplift. Just west of where Alexander (1953) worked, Bradley and Griggs (1976) mapped six
58 prominent terraces in the Santa Cruz-Point Año Nuevo area (Fig. 24). The lowest of these six marine terraces, between Santa
59 Cruz and extending to at least just north of Point Año Nuevo, is called the Santa Cruz terrace. Detailed seismic profiling and
60 examination of outcrops carried out by Bradley and Griggs (1976) show that the Santa Cruz terrace actually consists of three
61 distinct platforms cut on bedrock. However, the three wave-cut platforms are covered with marine and, importantly, nonmarine
62 deposits that have smoothed over the subaerial surface topographically into a single, broad landform. Bradley and Griggs
63 (1976) referred to the three buried platforms as the Greyhound level (~45 m), Highway 1 level (~35 m), and Davenport level
64 (~20 m). These elevations are rough averages, as the shoreline angle elevations vary as a function of where they are situated
65 with respect to active geologic structures. Too few exposures allow for these terraces to be mapped separately, but isolated
66 outcrops where the Davenport platform can be identified are present along the coast between Santa Cruz and north of Point
67 Año Nuevo (Fig. 24). Above the Santa Cruz terrace complex, higher terraces are found at ~55-60 m (Cement terrace), ~90-
68 100 m (Western terrace), ~120-140 m (Wilder terrace), ~180-195 m (Black Rock terrace), and ~240-260 m (Quarry terrace).

69 For years, there has probably been no greater speculation about the age of a coastal landform in California than that of the
70 Davenport platform between Point Año Nuevo and Santa Cruz. Anderson and Menking (1994) reviewed many of the previous
71 age estimates made for this terrace. Bradley and Addicott (1968) reported U-series ages of mollusks from this terrace that
72 ranged from ~100 ka to ~60 ka, but it is now well known that U-series analyses of mollusks are not reliable (Kaufman et al.,
73 1971). In a later study, Bradley and Griggs (1976) recognized this problem and suggested instead that the Highway 1 platform,
74 present just above the Davenport platform, was cut during the ~120 ka (MIS 5e) high-sea stand. They interpreted the Davenport
75 platform, although at a lower elevation, to have been cut during a hypothesized lower, ~140 ka sea stand, such as that seen at
76 reef VIIa on New Guinea (Bloom et al., 1974). Based on amino acid ratios in fossil mollusks and the faunal zoogeographic
77 aspect, Kennedy et al. (1982) concluded that the fossils on the Davenport platform dated to the ~80 ka high stand of sea during
78 MIS 5a. Assuming an age of ~120,000 yr for the Highway 1 terrace, Lajoie et al. (1991) estimated an age of ~100 ka (MIS
79 5c) for the Davenport terrace. Perg et al. (2001) used cosmogenic isotopes to estimate an age of 65 ka for the Highway 1
80 terrace where it is found just northwest of Santa Cruz, which would correlate the terrace with MIS 3. These workers offered
81 cosmogenic isotope ages for the higher terraces as well, including ~92 ka (Western terrace), correlated to MIS 5a; ~137 ka
82 (Wilder terrace), correlated to MIS 5c; ~139 ka (Black Rock terrace), correlated to MIS 5e; and ~226 ka (Quarry terrace),
83 correlated to MIS 7. The age estimates of Perg et al. (2001) combined with the elevations given above would characterize this
84 reach of coastline as having one of the highest rates of uplift along the California coast, exceeded only by the coast near the
85 Mendocino triple junction and south of the "big bend" of the San Andreas Fault, discussed later.

86 Muhs et al. (2006) reported U-series ages corals on from Green Oaks Creek, Point Año Nuevo, and Point Santa Cruz, all
87 from deposits of the Davenport level of the Santa Cruz terrace (Fig. 24). Twelve corals gave ages ranging from ~84 ka to ~76
88 ka and all 11 corals collected near Green Oaks Creek have back-calculated initial $^{234}\text{U}/^{238}\text{U}$ values ranging from 1.154 to
89 1.1460. These values fall well within the range of modern seawater, giving a high degree of confidence that the corals have
90 experienced closed-system histories with respect to U-series isotopes. In addition, deposits of the Davenport terrace have
91 faunas containing a large number of extralimital northern or northward-ranging species of mollusks, consistent with dated,
92 ~80 ka terraces elsewhere in Oregon and California (Addicott, 1966; Kennedy, 1978; Muhs et al., 2006). The U-series ages
93 of ~80 ka agree with the amino acid age estimate reported earlier by Kennedy et al. (1982).

94 Although Perg et al. (2001) did not analyze sediments from the Davenport platform directly, their ages for the higher
95 terraces imply that the Davenport terrace should correlate with one of the interstadial high stands of sea, recorded as uplifted,
96 coral-reef terraces on New Guinea. These terraces date to ~50,000, ~40,000 or ~30,000 yr B.P. (Chappell et al., 1996; Cutler
97 et al., 2003). However, based on the U-series ages for corals from the Davenport terrace, it is very likely that the cosmogenic
98 ages for the older Santa Cruz terraces are underestimates. A reasonable explanation is that the ages reported by Perg et al.
99 (2001) reflect the ages of alluvium that overlies the marine deposits. The terrestrial sedimentary cover in this area is typically
00 much thicker than the marine cover and marine sediments are rarely, if ever, exposed at the ground surface. For example, at
01 Point Año Nuevo, the sea cliff exposes the Davenport platform at ~7.8 m above sea level, overlain by ~0.5 to ~0.2 m of marine
02 deposits with fossils. However, above the marine deposits are ~9.8 m of alluvial sands and gravels, interbedded with silts and
03 clays. A well-developed soil, with an A/E/Bt [Bts]/C profile developed in these nonmarine deposits, indicates a substantial
04 age for this alluvium.

05 In light of the ~80 ka (MIS 5a) age for the Davenport terrace, a reasonable working hypothesis is that the other platforms
06 in the Santa Cruz terrace complex date to earlier high-sea stands of MIS 5. Thus, the Highway 1 platform could date to MIS
07 5c and the Greyhound platform could date to MIS 5e. If this correlation is correct, then the much of the MIS 5e shoreline
08 (represented hypothetically by the Greyhound platform) has been eroded away. Based on the shore-parallel elevation profiles
09 of Bradley and Griggs (1976), less than ~4 km of the shore-parallel extent of this terrace still exists along ~32 km of coastline
10 that they mapped. In contrast, the Highway 1 platform occurs nearly continuously from the city of Santa Cruz northwest to
11 Point Año Nuevo. If this platform was cut during MIS 5c, then much of the MIS 5e (Greyhound) terrace must have been
12 removed before much uplift could take place, contrary to the high uplift rates implied by Perg et al. (2001). The challenge
13 here, as in many places, is to devise a method of dating terraces that lack fossils.

14 **4.5.2. San Luis Obispo County, California**

15 In northern San Luis Obispo County, five marine terraces have been mapped by Hanson et al. (1994) in the area around
16 San Simeon (Fig. 25). These terraces are, in ascending order, the Point (7-8 m), San Simeon (4-23 m), Tripod (23-38 m), Oso
17 (29-47), and La Cruz (53-79 m) terraces. Terrace elevations vary as a function of proximity to the northwest-trending San
18 Simeon Fault zone. Unfortunately, fossils are apparently lacking in this area, so there is little age control for any of these
19 terraces. Hanson et al. (1994) correlated the Tripod terrace to MIS 5e, based on a simple lateral correlation to the low-elevation

20 marine terrace exposed near Cayucos, to the south (see discussion below). However, this correlation is currently only a
21 working hypothesis, as the Cayucos terrace is ~35 km distant, and it is not certain that such a long, shore-parallel correlation
22 can be justified.

23 West of the town of Cayucos (**RSL ID 3688**) a broad marine terrace extends along the coast for several kilometers, as
24 discussed earlier (Fig. 6a). This terrace and its deposits are well known, in part because the terrace sediments host a rich
25 molluscan fauna (Valentine, 1958), but also because it is the first terrace in California where a coral was dated by U-series
26 (Veeh and Valentine, 1969). The latter workers reported an age of ~130 ka for this coral, which is recalculated here to ~122
27 ka, using more recent estimates of the half-lives of U-series nuclides. With this correlation to the MIS 5e, the Cayucos terrace
28 has been an important calibration point for many aminostratigraphic studies (Wehmiller et al., 1977a; Kennedy et al., 1982;
29 Wehmiller, 1982; Muhs et al., 2014b). The terrace is broad, with a shore-normal extent of up to ~600 m. Although it has
30 been reported that the shoreline angle of the terrace, at around 7-8 m above sea level, is exposed near the town of Cayucos
31 itself (Stein et al., 1991), this measurement is actually of the wave-cut bench behind a paleo-sea stack. However, the
32 measurement is not greatly different from that made by extending a shore-normal topographic profile of the wave-cut bench
33 landward and finding its possible intersection with an extension of the paleo-sea cliff topographic profile downward (Fig. 6b),
34 which yields a possible shoreline angle elevation of ~8 m (**RSL IDs 3776, 3801**).

35 Using TIMS U-series methods, Stein et al. (1991) analyzed 12 corals from near LACMIP loc. 10731 (**RSL ID 3688**), near
36 the town of Cayucos (Figs. 6a, 25). Eleven corals from this locality gave ages ranging from ~125 ka to ~113 ka, with somewhat
37 elevated initial $^{234}\text{U}/^{238}\text{U}$ values, indicating that the ages are probably biased old to some degree, but still likely correlative to
38 MIS 5e. However, one coral gave an apparent age of ~101 ka, with an initial $^{234}\text{U}/^{238}\text{U}$ value that was only slightly elevated,
39 indicating minimal age bias. Muhs et al. (2002a) revisited the same locality and analyzed seven corals, with four yielding ages
40 of ~123 ka to ~116 ka, in broad agreement with Stein et al. (1991), and with three giving ages of ~110 ka to ~108 ka. Muhs
41 et al. (2002a) interpreted these data to indicate the possibility that during MIS 5c at ~100 ka, the high-sea stand overtook at
42 least the outer part of the terrace created during MIS 5e at ~120 ka, with the result that fossils of two ages were mixed together.
43 This interpretation is supported by a reexamination of the fossil fauna reported by Valentine (1958), using updated modern
44 zoogeography. Although the fauna contains several extralimital southern species of mollusks, as well as some southward-
45 ranging mollusks, it also contains some extralimital northern and northward-ranging species. This mix of warm-water (~120
46 ka?) and cool-water (~100 ka?) molluscan forms was interpreted by Muhs et al. (2002a) to be consistent with the apparent mix
47 of ~120 ka and ~100 ka corals.

48 South of Cayucos, in the Diablo Canyon (**RSL ID 3808**)-Point San Luis (**RSL ID 3777**)-Shell Beach (**RSL ID 3807**) area
49 (Fig. 25), Hanson et al. (1994) mapped a sequence of at least 12 marine terraces, up to an elevation of at least ~200 m. The
50 lowest two terraces, Q1 (12-4 m) and Q2 (34-12 m) were correlated with MIS 5a and 5e, respectively, using a variety of dating
51 methods. A U-series age, recalculated here using updated half-lives, on corals from the Q2 at Point San Luis (done by alpha
52 spectrometry) is ~118 ka and supports this correlation (Muhs et al., 1994), as do ages (by TIMS) on corals from Shell Beach
53 that range from ~127 ka to ~122 ka (Stein et al., 1991). However, at other localities of the Q2 terrace, Hanson et al. (1994)

54 noted inconsistencies between amino acid age estimates of the terrace mollusks and zoogeographic aspects of the faunas. For
55 example, at some localities, amino acid ratios would imply correlation to MIS 5a, but faunas have a warm-water aspect,
56 implying a correlation to MIS 5e. At other localities, such as Shell Beach, both amino acid data and faunal data would indicate
57 an MIS 5a age (Hanson et al., 1994), but U-series data by Stein et al. (1991) imply an MIS 5e age. Some of these issues were
58 also discussed by Wehmiller (1992) and Kennedy (2000), but details of the faunas and amino acid data do not allow for
59 complete resolution of the problem. At one locality on the Q2 terrace, near Diablo Canyon, Muhs et al. (1994) reported a U-
60 series age on coral (by alpha spectrometry) of ~108 ka, implying correlation with MIS 5c. If that age is correct, then it is
61 possible that, like Cayucos, the Q2 terrace in this area contains fossils of both MIS 5c and 5e age, which would explain the
62 occurrence of both cool-water (~100 ka) and warm-water (~120 ka) forms on the same terrace. More work is needed at
63 localities on both the Q1 and Q2 terraces to resolve this problem.

64 **4.6 Southern California, USA**

65 Coastal southern California is defined here as that part of the coast that is east or south of Point Conception (Fig. 25). This
66 point is a major geographic feature, because here the California coast changes from a north-south orientation to an east-west
67 one. The change in coastal orientation is structurally controlled by the orientation of the San Andreas Fault, which has a major
68 restraining bend (informally referred to as the "Big Bend") inland of, and to the northeast of Point Conception. Point
69 Conception also marks a major faunal boundary in the marine invertebrate communities of the Pacific Coast of North America,
70 long recognized by marine zoogeographers (Valentine, 1966). This has relevance to the interpretation of marine terrace faunas,
71 discussed in more detail below.

72 **4.6.1 Point Conception to Arroyo Hondo**

73 Marine terraces form the coastal plain between Point Conception and Arroyo Hondo (Fig. 25). Upson (1951) was the first
74 to study these terraces and map them. He also noted the presence of marine fossils in some of the deposits and reported
75 identifications of the taxa, based on an examination of his collections by W.P. Woodring. Preliminary age assignments for
76 low-elevation terraces in this area, based on amino acid ratios, were made by Kennedy et al. (1982). Later, however, Rockwell
77 et al. (1992) mapped five terraces (I, II, III, IV, and V, from lowest to highest) in the area and Kennedy et al. (1992) provided
78 new amino acid data and faunal lists from terraces I, II, and III. Terrace I, also called the Cojo terrace has shoreline angle
79 elevations varying from ~17 m to ~10 m, depending on proximity to local structures. Two bone specimens from deposits
80 overlying this terrace gave concordant $^{230}\text{Th}/^{238}\text{U}$ and $^{231}\text{Pa}/^{235}\text{U}$ ages of ~70 ka and ~87 ka, leading Rockwell et al. (1992) to
81 conclude that these were close minimum-limiting ages and permitted correlation with MIS 5a. Two localities on terrace I have
82 D/L leucine and valine values in *Saxidomus* that are significantly lower than those of this genus from deposits from terrace III
83 (~40-30 m), which Rockwell et al. (1992) correlated with MIS 5e (**RSL ID 3687**). The intermediate-elevation terrace II is
84 correlated with MIS 5c. Faunas in deposits of both terraces I and II have a cool-water aspect, whereas the fauna of terrace III
85 has a warm-water aspect, supporting these age assignments (Kennedy et al., 1992). Thus far, no corals have been reported in
86 terrace deposits in this area.

87 **4.6.2 Malibu**

88 Marine terrace deposits along the Malibu coast of Los Angeles County have been studied since the landmark paper of Davis
89 (1933), who named the two most prominent landforms the Malibu (higher) and Dume (lower) terraces. These two terraces,
90 plus an intermediate one called the Corral terrace, were mapped in detail by Birkeland (1972), who also measured the shoreline
91 angle elevations. These elevations vary in a shore-parallel sense, ranging from 76-61 m (Malibu), 54-46 m (Corral), and 40-
92 15 m (Dume). Szabo and Rosholt (1969) analyzed mollusks from the Corral (called "Terrace C" in their paper) and Dume
93 terraces for U-series isotopes, including ^{231}Pa and ^{235}U . They recognized that mollusks are open systems with regard to U-
94 series isotopes but devised an open-system model of age determination. From this model, they proposed ages of ~154 ka to
95 ~115 ka (average of ~131 ka) for the Corral terrace and ~112 ka to ~95 ka (average of ~104 ka) for the Dume terrace. Szabo
96 and Rosholt (1969) correlated the Corral terrace to the ~120 ka Rendezvous Hill ("Barbados III") terrace of Barbados and the
97 Dume terrace to the ~105 ka Ventnor ("Barbados II") terrace of the same island. The open-system model received considerable
98 criticism from Kaufman et al. (1971), who concluded that mollusks were not suitable for U-series dating, using either closed-
99 system or open-system approaches. Interestingly, Simms et al. (2016) nevertheless accepted the ~131 ka and ~104 ka mollusk
00 ages and correlated the Corral and Dume terraces with MIS 5e and 5c, respectively. Kennedy et al. (1982), however, reported
01 amino acid data from the Dume terrace that suggested correlation with MIS 5e, supported by the presence of a warm-water
02 fauna, studied earlier by Addicott (1964b). Although more work needs to be conducted on these terraces, it seems likely that
03 the lowest, Dume terrace (**RSL IDs 3689, 3690**) is the most probable representative of MIS 5e.

04 **4.6.3 Palos Verdes Hills-San Pedro**

05 The Palos Verdes Hills, also in Los Angeles County, is an uplifted crustal block with at least a dozen marine terraces (Fig.
06 26). The crustal block is bounded by faults on its southeast and northern sides. Based on mapping of terraces in a now-classic
07 study by Woodring et al. (1946), the Palos Verdes Hills was likely an island during some point or points in its history, most
08 recently during the last interglacial period, or MIS 5e. Woodring et al. (1946) numbered the terraces, from "1" (the lowest
09 terrace within the city of San Pedro) to "13" (the highest in the Palos Verdes Hills). The marine deposits overlying terrace 1
10 in San Pedro were referred to as the Palos Verdes Sand (**RSL IDs 3691, 3772, 3773, 3795**) and this unit was regarded as being
11 of the same age throughout its mapped extent, although it was recognized that there are substantial differences in the faunal
12 character from place to place (Woodring et al., 1946). To the west, on the Palos Verdes Hills, deposits of all the terraces were
13 considered by Woodring et al. (1946) to be older than the Palos Verdes Sand, so the lowest elevation terrace there was referred
14 to as terrace 2. Aminostratigraphic work by Muhs et al. (1992) showed that the Palos Verdes Sand in northern San Pedro is
15 correlative with terrace 4 (~72 m) on the Palos Verdes Hills and the Palos Verdes Sand in southern San Pedro is correlative
16 with terrace 2 (~47 m) on the Palos Verdes Hills (Fig. 26). These investigators correlated terrace 4 with MIS 5e and terrace 2
17 with MIS 5a, based on both aminostratigraphy and terrace faunas (warm-water faunas on terrace 4; cool-water faunas on
18 terrace 2). Because of these correlations, Muhs et al. (2006) considered that the terrace numbering system of Woodring et al.
19 (1946) was misleading and instead named terrace 2 the "Paseo del Mar" terrace and terrace 4 the "Gaffey" terrace (Figs. 26,
20 27). These correlations remained untested until some years later, when corals were recovered from both terraces 2 (Paseo del
21 Mar terrace) and 4 (Gaffey terrace, **RSL IDs 3771, 3784**). U-series ages by TIMS gave ages of ~80 ka (MIS 5a) from three

22 localities on terrace 2, and ages of ~119 ka to ~113 ka (MIS 5e) from a single locality on terrace 4 (Muhs et al., 2006). It is
23 likely that intermediate-elevation terrace 3, found only on the west side of the Palos Verdes Hills, correlates with MIS 5c, but
24 fossil corals or mollusks from this terrace have not yet been found.

25 **4.6.4 Newport Bay area**

26 The Newport Bay area of Orange County, California, south of Los Angeles, has long been known for its highly fossiliferous
27 marine terrace deposits (**RSL IDs 3692, 3693, 3796, 3820, 3821**). Terraces were mapped by Vedder et al. (1957) and then
28 remapped by Vedder et al. (1975). The most extensive of these is the area locally referred to as Newport Mesa (Fig. 28).
29 Grant et al. (1999) measured the shoreline angle elevations of eight terraces in this area and Newport Mesa corresponds to
30 their "Terrace 2." The shoreline angle elevations of this terrace range from ~32-36 m. A lower-elevation surface (their
31 "Terrace 1") has shoreline angle elevations ranging from ~19-22 m. From a fossil locality in the eastern part of Terrace 2,
32 Kanakoff and Emerson (1959) reported what is likely the most abundant marine invertebrate fauna of Pleistocene age on the
33 Pacific Coast of North America, with at least 500 species of mollusks, corals, bryozoans, brachiopods, echinoids, crabs,
34 barnacles, and worms. When the suitability of U-series dating of mollusks was still in a stage of assessment, Szabo and Vedder
35 (1971) attempted dating fossils by this method from the lowest three terraces in the area. As is usually the case with mollusks,
36 virtually all the specimens analyzed had evidence of open-system histories. Wehmiller et al. (1977a) and later Kennedy et al.
37 (1982) analyzed mollusks from the area for amino acid geochronology, primarily using the genera *Saxidomus* and *Leukoma*.
38 They showed that mollusks from half a dozen localities on Terrace 2 likely date to MIS 5e, but at least three localities have
39 evidence of older, pre-MIS 5e fossils, one of which is the main locality studied by Kanakoff and Emerson (1959). Grant et al.
40 (1999) conducted TIMS U-series analyses of corals from both Terrace 2 (two localities) and Terrace 1 (one locality). One of
41 their localities on Terrace 2 is close to the main locality studied by Kanakoff and Emerson (1959) and three analyses of one
42 *Paracyathus pedroensis* coral colony gave ages of ~124-120 ka, with minimal likely age bias, permitting correlation to MIS
43 5e. Three *Balanophyllia elegans* samples from the same terrace at another locality gave older apparent ages, but with clear
44 evidence of an open system history. A single *Paracyathus pedroensis* coral colony from Terrace 1 gave an apparent age of
45 ~106 ka, allowing correlation to MIS 5c. At least two of the localities correlated to MIS 5e by Wehmiller et al. (1977a) are
46 on what Grant et al. (1999) later mapped as Terrace 1 and correlated to MIS 5c, based on their U-series age from this terrace,
47 highlighting the need for additional study of these terraces.

48 **4.6.5 San Diego County**

49 In the San Diego area (**RSL IDs 3694 to 3701, 3785, 3822**), multiple marine terraces have been documented (Kern and
50 Rockwell, 1992). The lowest two terraces, mapped by Kern (1977), have received the most attention. These are the Bird Rock
51 terrace (shoreline angle elevation of ~8 m) and the Nestor terrace (shoreline angle elevation of ~23 m), best exposed along the
52 west coast of Point Loma (Fig. 29). Both terraces host deposits and fossils that are thought to represent high-energy, rocky
53 intertidal environments (Kern, 1977). A more quiet-water, "bay" fauna characterizes what is called the Bay Point Formation
54 at somewhat more protected localities. This formation is considered to be correlative to deposits of the Nestor terrace
55 (Valentine, 1959; Kern, 1971). Ku and Kern (1974) reported three alpha-spectrometric U-series ages of corals from the Nestor

56 terrace (~109 ka, ~131 ka, and ~124 ka). Of these, the age of ~109 ka is the only analysis that yielded an initial $^{234}\text{U}/^{238}\text{U}$
57 value within the range of modern seawater. Nevertheless, an "average" age of ~121 ka (correlated to MIS 5e) for the Nestor
58 terrace has been assumed by many subsequent investigators who have used this terrace as a calibration point for amino acid
59 geochronology (Wehmiller et al., 1977a; Wehmiller and Belknap, 1978; Wehmiller and Emerson, 1980; Emerson et al., 1981;
60 Kennedy et al., 1982; Wehmiller, 1982; Keenan et al., 1987). Stein et al. (1991) reported somewhat older U-series ages of
61 individual corals, analyzed by TIMS, ranging from ~145 ka to ~133 ka, and offered the possibility that the Nestor terrace was
62 not cut during the MIS 5e high-sea stand. The same investigators also reported an age of ~97 ka for the Bird Rock terrace.
63 Muhs et al. (1994) redated corals from both the Nestor and Bird Rock terraces using alpha-spectrometric U-series analyses,
64 and reported ages of 126 ± 6 ka and 85 ± 4 ka, respectively.

65 In some amino acid studies that have used the Nestor terrace fossils as calibration points (Wehmiller et al., 1977a; Kennedy
66 et al., 1982), the fauna has been reported to be one characterized by warm-water forms, although no taxa are specifically
67 mentioned. Using the detailed fauna presented by Valentine and Meade (1961) and Kern (1977), however, Muhs et al. (2002a)
68 challenged the idea that the Nestor terrace hosts predominantly warm-water species. Although some extralimital southern
69 forms are present, there are a larger number of northward-ranging species. Muhs et al. (2002a) also reported new TIMS U-
70 series analyses of individual corals from deposits of the Nestor terrace. Nine of these have ages ranging from ~128 ka to ~113
71 ka, but three corals have ages ranging from ~109 ka to ~98 ka, similar to what was reported for Cayucos, California, discussed
72 earlier. These investigators interpreted the results from both localities to indicate that the deposits at Cayucos and on the
73 Nestor terrace contain fossils representing MIS 5e (with warm-water mollusks) and MIS 5c (with cool-water mollusks).

74 Elsewhere in the San Diego area, at Torrey Pines State Park (Fig. 25), and near the Mexican border, Wehmiller et al. (1977a)
75 and Kennedy et al. (1982) correlated low-elevation marine terrace deposits to the Nestor terrace using amino acid
76 geochronology. These investigators correlated the Torrey Pines and "border locality" deposits with MIS 5e based on both
77 amino acid ratios and reports of faunas with warm-water aspects (Emerson and Addicott, 1953; Valentine, 1960). There have
78 been, however, no U-series or amino acid studies of the quiet-water Bay Point Formation fossils, so assumed correlations to
79 MIS 5e for these deposits remain hypothetical.

80 **4.6.6 Channel Islands**

81 The eight islands off the coast of southern California are called the Channel Islands, because of the proximity of the northern
82 chain of four islands to Santa Barbara Channel (Fig. 25). Of the eight islands, all but Santa Catalina Island are characterized
83 by geomorphically well expressed marine terraces. In addition, some of the most fossiliferous and best-preserved terraces
84 along the entire coast of North America are found on the Channel Islands. Five islands are preserved either in Channel Islands
85 National Park or by The Nature Conservancy and two (San Nicolas Island and San Clemente Island) are owned by the U.S.
86 Navy. Thus, the urban development that has obscured much of the marine terrace geomorphology on mainland California is
87 absent on the islands. Fourteen terraces have been mapped on San Nicolas Island (Vedder and Norris, 1963), to an elevation
88 of ~240 m, and San Clemente Island hosts at least 20 terraces, the highest at an elevation of almost 600 m. Even tiny Santa
89 Barbara Island, which has an area of less than 3 km², hosts at least five marine terraces (Muhs and Groves, 2018).

90 The majority of work on last interglacial marine terrace records has been done on the southern islands. Muhs et al. (1994)
91 reported alpha-spectrometric ages of the lowest marine terraces on San Clemente Island (**RSL IDs 3755 and 3800**) and San
92 Nicolas Island (**RSL IDs 3775 and 3813 to 3817**). Their study showed that the 2nd emergent terraces on both islands have
93 deposits hosting fossils that likely date to MIS 5e and the 1st terrace on San Nicolas Island dates to ~80 ka, or MIS 5a. Later,
94 higher precision TIMS U-series analyses confirmed these ages for both San Clemente Island and San Nicolas Island (Muhs et
95 al., 2002a, 2006). More detailed work on San Nicolas Island, however, with both new terrace mapping and new TIMS U-
96 series ages (Muhs et al., 2012), showed that although the 1st terrace is a single landform with deposits dating to ~80 ka, the
97 2nd terrace is a composite feature, with a broad, lower elevation surface (terrace "2b") and a narrow, higher elevation surface
98 (terrace "2a"). Fossils from terrace 2a date only to MIS 5e and do not contain cool-water mollusks, but fossils from terrace 2b
99 date to both MIS 5e (~120 ka) and MIS 5c (~100 ka) and contain a mix of mollusks with both warm-water and cool-water
00 aspects, similar to what was reported for Cayucos and the Nestor terrace at Point Loma. Muhs et al. (2012) interpreted these
01 results to indicate that the MIS 5c high-sea stand was high enough and the uplift rate on San Nicolas Island was low enough
02 that much of the MIS 5e terrace (2a) was removed by sea cliff retreat at ~100 ka, and fossils from both the ~120 ka and ~100
03 ka sea stands were mixed into the deposits of terrace 2b. With these new findings in mind, Muhs et al. (2014a) examined the
04 faunal record of the MIS 5e terrace on San Clemente Island, which is also a composite landform (i.e., two platforms, 2a and
05 2b, as on San Nicolas Island). This investigation showed that the "MIS 5e" terrace deposits on this island also contain a mix
06 of both warm-water and cool-water fossils, chiefly mollusks, that also imply a mix of MIS 5e (warm) and MIS 5c (cool) taxa.
07 This interpretation also explains a previously enigmatic molluscan oxygen isotope record, implying cooler waters in what had
08 been thought to be solely ~120 ka deposits (Muhs and Kyser, 1987).

09 The largest of the northern Channel Islands (San Miguel, Santa Rosa, and Santa Cruz) also have marine terraces that date
10 to MIS 5e. TIMS U-series analyses give ages of ~120 ka for the 2nd emergent terraces (shoreline angle elevations of ~20-24
11 m) on San Miguel (**RSL ID 3778**) and Santa Rosa (**RSL IDs 3779 to 3782**) Islands (Muhs et al., 2014b). As is the case on
12 San Nicolas and San Clemente Islands, the fossil faunas from the deposits of these terraces host both warm-water and cool-
13 water taxa (Orr, 1960; Muhs et al., 2014b), although currently there is only sparse geomorphic evidence of two high sea stands
14 (i.e., both MIS 5e and 5c). At one locality on Santa Rosa Island, there is a marine terrace with an outer edge at ~7 m above
15 sea level with an uncertain shoreline angle elevation, as the inner part of the terrace is covered by eolian sand. Apparent TIMS
16 U-series ages of corals from this terrace range from ~113 ka to ~110 ka and all have slightly elevated initial $^{234}\text{U}/^{238}\text{U}$ values,
17 indicating at least some bias to older ages (Muhs et al., 2015). Further, mollusks from this terrace (SRI-1 on Fig. 12) have
18 lower amino acid ratios than those in the 24-m-high terrace (SRI-5F on Fig. 12) dated to ~120 ka elsewhere on the island.
19 Thus, it is possible that this isolated terrace fragment represents an MIS 5c record, but more work is needed to confirm this.
20 On both islands, there is a lower elevation terrace with a shoreline angle elevation of ~3 m (San Miguel Island) and ~7 m
21 (Santa Rosa Island). TIMS U-series analyses give an age of ~80 ka for this terrace on Santa Rosa Island and amino acid ratios
22 in mollusks indicate a similar age for the 3-m-high terrace on San Miguel Island (Muhs et al., 2015, 2018). The lowest-
23 elevation terrace on Santa Cruz Island (**RSL IDs 3789, 3790, 3811, 3812**) has shoreline angle elevations ranging from ~6 m

24 to ~17 m and both U-series ages of corals and amino acid ratios in mollusks indicate that it dates to MIS 5e (Pinter et al., 1998;
25 Muhs and Groves, 2018). Thus far, there is no evidence of terraces dating to MIS 5c or 5a on Santa Cruz Island, suggesting
26 that the long-term uplift rate on this island is relatively low.

27 The two smallest of the Channel Islands, Anacapa Island (**RSL ID 3791**) and Santa Barbara Island (**RSL IDs 3792, 3793**),
28 both have low-elevation terraces, with shoreline angle elevations of ~10-11 m above sea level. No coral ages are yet available
29 for these terraces, but amino acid ratios indicate a mixed population of mollusk ages, correlated to MIS 5e and either MIS 5c
30 or MIS 5a (Fig 12). The terrace fauna on Santa Barbara Island is diverse, with a large number of warm-water forms and a
31 smaller number of cool-water forms (Lipps et al., 1968; Muhs and Groves, 2018). The much more sparse fauna on Anacapa
32 Island hosts some warm-water forms, but only one northward-ranging species.

33 Santa Catalina Island's marine terrace record, or to put it more accurately, its apparent lack of a record, has been an enigma
34 for decades (see Smith, 1933, for one of the earliest discussions). The island is situated between crustal blocks to the north
35 (Palos Verdes Hills) and south (San Clemente Island) that both host abundant marine terraces. Bedrock is not a limiting factor,
36 because the Catalina Schist that characterizes much of the island is similar to that of the Franciscan rocks that host marine
37 terraces elsewhere in central and northern California (e.g., Cayucos, Fig. 6). In support of this, Emery (1958) showed that
38 submarine terraces are found off Santa Catalina Island. At a minimum, even with no uplift, with the likelihood of a higher
39 than present sea level during MIS 5e, one should expect to see some evidence of a terrace that dates to ~120 ka. One hypothesis
40 that has been offered is that the island is subsiding and that the terrace record is largely submerged (Castillo et al., 2018).
41 However, there is no a priori reason to suppose that in a tectonic setting similar to those elsewhere in southern California that
42 Santa Catalina Island should be subsiding when all adjacent areas are uplifting. Schumann et al. (2012) presented evidence
43 that in fact the opposite is true, i.e., Santa Catalina Island is experiencing uplift, possibly at a high enough rate that fluvial
44 erosion has removed most evidence of any terraces. This remains the most viable explanation to date, but more work is needed
45 to confirm this.

46 **4.7 Baja California, Pacific Coast**

47 South of the city of Ensenada, Baja California, there is a prominent peninsula called Punta Banda (Fig. 30). Rockwell et
48 al. (1989) mapped several marine terraces on this peninsula and provided alpha-spectrometric U-series ages of corals and
49 hydrocorals from the lowest terraces. The 3rd terrace, called the Sea Cave terrace (**RSL IDs 3786, 3802**), has a shoreline
50 angle elevation that varies between ~34 m and ~40 m (Fig. 31). The 1st terrace, called the Lighthouse terrace, has a shoreline
51 angle elevation that varies between ~15 m and ~18 m. An intermediate, unnamed terrace at ~22 m elevation occurs only in a
52 small area on the outer part of the peninsula. U-series ages indicate that the Sea Cave terrace is ~120 ka (MIS 5e) and that the
53 Lighthouse terrace is ~80 ka (MIS 5a). The intermediate terrace at ~22 m could represent MIS 5c, but no corals were found
54 on the terrace. Muhs et al. (2002a) reported new, TIMS U-series analyses of corals from both the Sea Cave and Lighthouse
55 terraces that confirm the earlier ages generated by alpha spectrometry. Many of the corals from Punta Banda show minimal
56 age bias, based on back-calculated initial $^{234}\text{U}/^{238}\text{U}$ values (Fig. 10c). Corals from the Sea Cave terrace that show mostly

57 closed-system histories have ages ranging from ~124 ka to ~118 ka, and those from the Lighthouse terrace range from ~83 ka
58 to ~80 ka.

59 Isla Guadalupe is situated ~260 km southwest of the Pacific coast of Baja California (Fig. 2). The island is attractive as a
60 reference locality for estimating paleo-sea level during the last interglacial period because it is one of the few localities adjacent
61 to North America that can be considered, *a priori*, to be tectonically stable. The island is distant from any plate boundary, has
62 no active faults nearby, has no active volcanoes on it or near it, is bounded on its eastern side by a seafloor with undisturbed
63 marine sediment, and has no history of recent earthquakes (Gonzalez-Garcia et al., 2003). Lindberg et al. (1980) reported that
64 emergent marine deposits are found on the southern and eastern of the coasts of the island. These deposits have elevations of
65 ~1 m to ~8 m above sea level, with most localities described as ~1 m to ~6 m above sea level (**RSL IDs 3803 to 3805**). From
66 these deposits, Muhs et al. (2002a) reported ages of ~123 ka to ~118 ka for *Pocillopora* corals and most show closed-system
67 histories. An extensive faunal list by Lindberg et al. (1980) indicates that the ~120 ka deposits host a large number of
68 extralimital species of mollusks. In addition, Isla Guadalupe marks, thus far, the northernmost occurrence of hermatypic corals
69 along the Pacific Coast of North America during MIS 5e (Durham, 1980).

70 South of Punta Banda, marine terraces are prominent landforms all along the coast of Baja California. In a coastal reach
71 from Punta Banda south for at least 300 km, multiple marine terraces are present, mapped by Orme (1980) (Fig. 30). Many
72 low-elevation marine terrace deposits along this reach of coast are highly fossiliferous (Emerson, 1956, 1960; Emerson and
73 Addicott, 1958; Addicott and Emerson, 1959; Valentine, 1960a, 1961) and are candidates for records as MIS 5e shorelines.
74 Unfortunately, little work has been done on age determinations for most of these terraces. What geochronologic work has
75 been done along the Pacific coast of both Baja California and Baja California Sur is aminostratigraphic correlation, using U-
76 series-dated localities, such as the Nestor terrace to the north, and a single locality to the south, Bahía Magdalena (see
77 Wehmiller and Emerson, 1980). Thus, before reviewing the results of the aminostratigraphic correlations, the work done at
78 Bahía Magdalena is discussed first.

79 Bahía Magdalena (**RSL IDs 3707, 3798**) is situated on the Pacific side of Baja California Sur (Fig. 32) and marine deposits
80 there have long been famous for their extensive Pleistocene fauna (Jordan, 1936). Near the village of Puerto Magdalena on
81 the peninsula, one of Jordan's (1936) fossil sites (California Academy of Sciences [CAS] locality 754) contains fragments of
82 the colonial coral *Porites californica* (now considered to be *P. panamensis*). These corals occur in marine terrace deposits
83 that have a *maximum* elevation of ~6 m and have been dated to ~118 ka to ~116 ka by alpha-spectrometric uranium-series
84 methods (Omura et al., 1979). When the data of Omura et al. (1979) are recalculated using the more recent estimates of half-
85 lives (Cheng et al., 2013), the coral ages from Bahía Magdalena are ~114.8 ka to ~114.0 ka.

86 The northernmost locality on the Pacific coast of Baja California that has been examined for paleontology and
87 aminostratigraphic correlation is a low-elevation (outer edge elevation of ~5 m to ~10 m) terrace at Camalú (**RSL ID 3702**,
88 Fig. 30), studied by Valentine (1980). Amino acid ratios in *Leukoma* shells from the deposits of this terrace led Valentine
89 (1980) to conclude that the terrace dates to MIS 5e. It is important to note here that although Valentine (1980) did not report
90 actual amino acid ratios for Camalú, his interpretation is supported by data presented graphically by Keenan et al. (1987).

91 Faunal data make an MIS 5e interpretation for the terrace at Camalú complicated, however, as there are several northward-
92 ranging species of mollusks and only one southward-ranging species (Valentine, 1980). Approximately 300 km south of
93 Camalú, at Punta Santa Rosalíllita (**RSL ID 3962**), Woods (1980) mapped three emergent marine terraces, named Tomatal
94 (shoreline angle of ~7 m), Andres (~25-30 m), and Aeropuerto (~50-60 m). Although there are no U-series ages on corals
95 from these terraces, Woods (1980) reported amino acid data from fossil mollusks that correlate the Tomatal terrace with the
96 peak of the LIG at ~120 ka. As is the case at Camalú, data presented graphically by Keenan et al. (1987) support this
97 interpretation. Amino acid data from these localities are also reported in Wehmiller and Pellerito (2015).

98 Farther south on the Pacific coast of northern Baja California Sur, at Bahía Tortugas and Bahía Asunción (Fig. 33), Emerson
99 et al. (1981) and Keenan et al. (1987) reported amino acid data for low elevation terraces that they correlated to MIS 5e. Two
00 terraces are present at Bahía Tortugas (**RSL IDs 3703, 3704**), one at ~27-24 m and the other at ~12 m, although it is not clear
01 if these elevations refer to shoreline angles or simply the fossil localities that were studied. In any case, amino acid ratios
02 clearly distinguish the two terrace deposits, with the higher elevation terrace attributed to MIS 5e, based on aminostratigraphic
03 correlation between Bahía Magdalena to the south and the Nestor terrace to the north. The lower terrace, with lower ratios, is
04 considered to be "~95 ka," but correlation to either MIS 5c or MIS 5a is possible. Deposits of both terraces contain *both* warm-
05 water and cool-water species, but the upper terrace contains substantially more warm-water forms than the lower terrace.

06 At Bahía Asunción (**RSL IDs 3705, 3706**), Keenan et al. (1987) identified three age groups of marine terrace deposits
07 based on amino acid ratios. The youngest of these is correlated with MIS 5e, based on aminostratigraphic correlation to Bahía
08 Magdalena. The deposits correlated to MIS 5e are situated ~6 m above sea level at one locality and ~11-12 m above sea level
09 at another locality. However, deposits hosting what these investigators considered to be older deposits, based on amino acid
10 ratios, have elevations that fall within the same general range as those correlated to MIS 5e, so additional work in this area is
11 warranted to clarify the age-elevation relations.

12 **4.8 Golfo de California coasts of Baja California, Baja California Sur, and Sonora, Mexico**

13 As discussed above, waters in the Golfo de California are distinctly warmer than those in the Pacific Ocean along the west
14 coast of Baja California and Baja California Sur (Mitchell et al., 2002). Thus, the potential for finding coral-bearing marine
15 deposits or even true coral reefs is greater in this region than on the outer coast of Baja California and Baja California Sur.
16 Cabo Pulmo is located in the southernmost part of Baja California Sur (Fig. 32), adjacent to the Golfo de California. Ortlieb
17 (1987) mapped emergent marine terraces near Cabo Pulmo, as well as to the southwest, towards Cabo San Lucas, and to the
18 north. Squires (1959) described a coral-bearing marine terrace deposit near Cabo Pulmo (**RSL ID 3806**), which Ortlieb (1987)
19 reported as having a shoreline angle elevation of ~6 m. Muhs et al. (2002a) reported three TIMS U-series analyses of *Porites*
20 and *Pocillopora* corals from this deposit. The *Pocillopora* colony gave an apparent age of ~140 ka, but is clearly biased old,
21 based on an elevated initial $^{234}\text{U}/^{238}\text{U}$ value. On the other hand, *Porites* corals from these deposits gave ages of ~127 ka and
22 ~120 ka and have only slightly elevated initial $^{234}\text{U}/^{238}\text{U}$ values. Thus, it is clear that this deposit represents MIS 5e.

23 North of Cabo Pulmo, Isla Cerralvo (**RSL ID 3832**) is situated off the eastern coast of Baja California Sur. Tierney and
24 Johnson (2012) studied a section 8.7 m thick on southwestern tip of the island, composed of alternating layers of growth-

25 position corals (Fig. 32) and cobbles, the latter interpreted to be from storm transport. Five coral-cobble cycles are represented
26 by the layers in this section, all interpreted to represent a single interglacial period. Tierney and Johnson (2012) report a U-
27 series age of ~126 ka from one of the *Porites* coral colonies found in the section. The highest growth-position reef layer is at
28 an elevation of ~3.9 m, overlain by sands interpreted to be from a prograding beach, up to an elevation of 7.1 m. Coral-bearing
29 sediments (interpreted here to be from storm deposits) occur as high as ~8.7 m above sea level.

30 Marine terrace deposits are exposed on both sides of Punta Coyote (**RSL IDs 3828 to 3831**), Baja California Sur (Fig. 32).
31 U-series analyses, done by alpha spectrometry, have been conducted on both *Porites* and *Pocillopora* corals recovered in low-
32 elevation terrace deposits here, reported by Sirkin et al. (1990) and Szabo et al. (1990). Apparent ages of corals from each of
33 five localities would allow correlation of the deposits to MIS 5e, but one coral has a back-calculated initial $^{234}\text{U}/^{238}\text{U}$ value that
34 is higher than modern seawater and three have initial $^{234}\text{U}/^{238}\text{U}$ values that are *lower* than modern seawater, an unusual
35 situation, and how this affects apparent ages is not known. Based on what general information is given, it appears that these
36 deposits have inner edge elevations that may be on the order of ~8 to ~10 m above sea level. Farther north, at Bahía Coyote
37 (**RSL IDs 3826 and 3827**), DeDiego-Forbis et al. (2004) reported U-series ages of *Porites* corals from terraces exposed along
38 the coast. The highest elevations of what appear to be growth-position *Porites* colonies are estimated to be ~18 m to ~22 m
39 above modern sea level. At least four of the corals analyzed appear to have experienced gain of bulk U, but two corals analyzed
40 have acceptable U contents and apparent ages of ~138 ka. Because both of these corals have initial $^{234}\text{U}/^{238}\text{U}$ values that are
41 higher than modern seawater, both are biased old by some amount, but permit correlation of the terraces to MIS 5e.

42 Studies by Johnson (2002) and Johnson et al. (2007) provide data on last interglacial coral reefs at two localities in Baja
43 California Sur, Isla Coronado, and Punta Chivato (Fig. 32a, b, c). TIMS U-series analyses were conducted on these corals in
44 laboratories of the U.S. Geological Survey, and complete analytical data are given in Muhs et al. (2014b). At Isla Coronado
45 (**RSL ID 3818**), Johnson et al. (2007) report that a *Porites panamensis* colony formed one of the largest fossil structures yet
46 reported in the Golfo de California. The top of the coral reef surface is ~12 m above sea level. Analysis of coral from this
47 reef gave an age of ~127 ka, with an initial $^{234}\text{U}/^{238}\text{U}$ value higher than modern seawater. Thus, the age is likely biased old by
48 some amount, but still allows correlation to MIS 5e. At Punta Chivato (**RSL ID 3819**), a *Porites panamensis* colony ~15 cm
49 high, in growth position (Fig. 32), is situated on Pleistocene river gravels at a present elevation of 7.5 m to 10 m above sea
50 level (Johnson, 2002; Johnson et al., 2007). U-series analysis of a *Porites* sample from this colony gave an age of 117.7 ka,
51 with an initial $^{234}\text{U}/^{238}\text{U}$ value indistinguishable from modern seawater, giving a high degree of confidence that this deposit
52 correlates with MIS 5e. Just south of Punta Chivato, a low-elevation marine terrace is present along the coast near Mulegé
53 (**RSL IDs 3823 and 3824**, Fig. 32). Rather than a constructional coral reef, this landform appears to be a California-style
54 marine terrace, with a wave-cut platform and overlying deposits that contain corals. The terrace has a shoreline angle elevation
55 of ~12 m above sea level and two alpha-spectrometric U-series analyses gave ages of ~146 and ~124 ka, with initial $^{234}\text{U}/^{238}\text{U}$
56 values only slightly higher than modern seawater (Ashby et al., 1987). Both corals have somewhat lower than optimum
57 $^{230}\text{Th}/^{232}\text{Th}$ values, suggesting the possibility of some inherited ^{230}Th , which would bias the apparent ages older. Nevertheless,
58 it is likely that the terrace correlates with MIS 5e, as concluded by Ashby et al. (1987). Between Punta Chivato and Mulegé,

59 ~15 km north of the latter locality, Libbey and Johnson (1997) listed an extensive (>40 species) fossil molluscan fauna from
60 a terrace that they report can be traced to, or nearly to Punta Chivato. Many of these taxa have modern ranges that extend
61 from the upper part of the Golfo de California to southern Mexico, Panama, Ecuador, or Peru, indicating a likely marine
62 paleotemperature range at least as warm as that of the present.

63 North of Punta Chivato, coral-bearing marine terrace deposits have not been reported on either coast of the Golfo de
64 California. Nevertheless, molluscan-rich terrace deposits are common and permit the possibility of amino acid geochronology.
65 By far the most extensive studies of these deposits are those by Ortlieb (1987, 1991). Because of the relatively high mean
66 annual air temperatures in the Golfo de California (~20°C in the north ranging to ~23°C in the south), even amino acid
67 geochronology becomes problematic in identifying deposits that correlate to MIS 5e. The reason for this is that many species
68 of mollusks will have reached, or be close to, racemic equilibrium for most amino acids after ~120 ka. Ortlieb's (1987, 1991)
69 approach to this problem was to consider that shells (the bivalves *Chione* and *Dosinia*) that were beyond radiocarbon range,
70 but yielded amino acid ratios not yet at equilibrium (but close to it) could be interpreted to be of MIS 5e age. This method is
71 supported by his analyses of shells from both Bahía Magdalena, on the Pacific side of Baja California Sur and at Bahía San
72 Nicolas (RSL IDs 3741, 3742), on the Golfo de California side of Baja California (Figs. 32, 33), where U-series ages on corals
73 have been obtained (Omura et al., 1979; Ortlieb, 1987). His interpretations are supported by more recent amino acid data on
74 *Chione* reported by Umhoefer et al. (2014), calibrated to U-series data reported by DeDiego-Forbis et al. (2004) from Bahía
75 Coyote (Fig. 34). Also shown in this figure are D-alloisoleucine/L-isoleucine values in radiocarbon-dated *Chione* shells of
76 late Holocene age, from Cholla Bay, Sonora (Martin et al., 1996) and a blue-shaded band that defines the equilibrium range
77 for D-alloisoleucine/L-isoleucine (1.25-1.35; Miller and Mangerud, 1985). When all data are considered, it is apparent that
78 shells falling within a range of ~0.70 to ~1.00 can be correlated to U-series-dated 120 ka localities. This range of values is
79 substantially lower than the equilibrium range of 1.25-1.35, but considerably higher than the range of values in Holocene
80 shells, ~0.02 to ~0.12. Ortlieb's (1987, 1991) results, along with those by Umhoefer et al. (2014) show, therefore, that an MIS
81 5e shoreline can be traced from the uppermost Golfo de California, south along the coasts of Baja California and Sonora, for
82 at least ~850 km. Most of the MIS 5e shorelines studied by Ortlieb (1987, 1991) have relatively low elevations, ranging from
83 ~2 m to ~8 m above sea level (Table S2).

84 **4.9 Pacific Coast of southern Mexico**

85 Although corals presently flourish along the Pacific coast of southern Mexico, south of the Golfo de California (López-
86 Pérez, 1998), there are only scattered reports of fossil corals as reefs or in emergent terrace deposits along this reach of
87 coastline. Emergent marine terraces or fossil coral reefs have been reported from both mainland Mexico, south of Oaxaca
88 (Palmer, 1928a,b; Squires, 1959) and offshore Tres Marias Islands (Fig. 7) (Hertlein and Emerson, 1959; Foose, 1962). There
89 do not appear to have been any recent studies of these deposits, nor are any geochronological data available. Given the
90 elevations that are described in these studies, however, as well as the fossil records, they are candidates as MIS5e shoreline
91 records and deserve further study.

92 **4.10 Central America**

93 The tectonic setting of the Pacific coast of Central America differs from that of Mexico to the north. Here, both the Cocos
94 and Nazca plates are being actively subducted under either the North America plate or the Caribbean plate (Figs. 1, 35a).
95 Furthermore, the Panama triple junction is situated offshore, just south of the Costa Rica-Panama border (Fig. 35a), making
96 this area tectonically and structurally complex.

97 Ocean temperatures are, in principle, warm enough to support hermatypic coral reef growth off the Pacific coast of Central
98 America from Guatemala to Panama. Toth et al. (2017) point out, however, that between Mexico and Nicaragua, there are
99 very few if any true coral reefs on the Pacific coast, in what is referred to as the "Central American faunal gap" (Fig. 7). These
00 investigators hypothesize that the lack of modern coral reefs along this reach of coast may be a function of a lack of hard
01 substrate available for larval settlement, although severe and prolonged upwelling has also been offered as a contributing factor
02 (Glynn et al., 2017). In all likelihood, this also limits the potential for finding emergent coral reefs that date to MIS 5e as well.

03 Farther south, along the coasts of Costa Rica and Panama, modern coral reefs are much more common and many developed
04 as early as ~7,000 yr ago (see summary in Toth et al., 2017). This is also an area where subduction of the Cocos plate beneath
05 the Caribbean plate is currently active and uplift rates are high (Gardner et al., 1992; Marshall and Anderson, 1995). Much of
06 the work on emergent marine terraces here has been focused on the Nicoya Peninsula and Osa Peninsula of Costa Rica, and
07 the Burica Peninsula of Panama (Fig. 35a). Gardner et al. (1992) reported marine terraces of Holocene age (~7000 yr to 980
08 yr) at elevations of ~3 to ~9 m on the Osa Peninsula and Marshall and Anderson (1995) report Holocene marine terraces with
09 ages of ~4700 yr to ~500 yr at elevations of ~4 m to ~16 m on the Nicoya Peninsula, demonstrating that the uplift rate in this
10 region of active subduction is relatively high. Fisher et al. (1998), Gardner et al. (2001), and Sak et al. (2004) all pointed out
11 that uplift in this region is controlled primarily by roughness of the subducting plate: forearc uplift on the Caribbean plate
12 corresponds to the position of migrating seamounts on the northeastward-moving Cocos plate.

13 With such high uplift rates, based on the elevations of Holocene marine terraces, any MIS 5e marine terraces on the Pacific
14 coast of Costa Rica would have to be at relatively high elevations now. On the Nicoya Peninsula, Marshall and Anderson
15 (1995) recognized two marine terraces. The younger of these is the suite of Holocene marine deposits, called the "Cabuya"
16 terrace. The higher terrace, called the "Cobano" terrace, is a broad coastal mesa, situated at an average elevation of ~180 m,
17 and is hypothesized to have formed during MIS 5e, although no geochronologic data are presented in support of this (Marshall
18 and Anderson, 1995). Using Holocene uplift and rotation rates, Gardner et al. (2001) estimated the Cobano terrace to have
19 formed between ~200 ka and ~100 ka, also permitting an interpretation of an MIS 5e age.

20 On the Osa Peninsula of Costa Rica, Gardner et al. (2013) mapped Quaternary marine deposits of three ages, from youngest
21 to oldest, the Jiménez (Holocene), Tigre (MIS 3?), and Rincón (MIS 5?) members of what they called the Marengo Formation
22 (Fig. 35b). They reported an OSL age of 109 ± 28 ka for deposits of the Rincón member and correlated this unit to MIS 5e.
23 Based on marine terrace shell radiocarbon ages, the Jiménez member dates to the Holocene. Gardner et al. (2013) also dated
24 shells from the Tigre member, which, when calibrated, range from ~31 ka to ~48 ka with a few samples yielding apparently
25 infinite ages. They correlated the Tigre member with MIS 3.

26 Gardner et al.'s (2013) correlation of the Tigre member to MIS 3, based on their radiocarbon ages, requires some scrutiny.
27 A critical examination applies to similar radiocarbon ages reported by Gardner et al. (1992) and Sak et al. (2004), also on the
28 Osa Peninsula, as do ages reported by Morell et al. (2011) for terrace shells on the Burica Peninsula of Panama (Fig. 35a).
29 Indeed, some of the marine terrace radiocarbon ages reported by Morrell et al. (2011) date not only to MIS 3, but actually give
30 apparent ages dating to the late last glacial period (MIS 2), at a time when sea level was several tens of meters below present.

31 Emergent marine deposit shells giving apparent radiocarbon ages of MIS 3 age have been reported on coastlines in various
32 parts of the globe for decades, with some investigators claiming that such ages require a paleo-sea level close to, or even above
33 present sea level during this interstadial period. Periodically, there have been critiques of such claims (Thom, 1973; Bloom,
34 1983; Colman et al., 1989), but many investigators continue to regard shell radiocarbon ages of ~30 ka to ~45 ka as truly finite
35 and accurate. The problem is that modern carbon is nearly everywhere and has considerable mobility. This means that old
36 shells are notorious for incorporating at least small amounts of modern carbon. Thus, even very small amounts of modern
37 carbon can make an infinitely old shell yield an apparently “finite” radiocarbon age (Pigati et al., 2007). An 80-ka sample, for
38 example, with a very small amount of modern carbon, can easily yield an apparent radiocarbon age of ~40 ka to ~45 ka. More
39 work needs to be done on the ages of the older marine terrace fossils of both the Osa Peninsula of Costa Rica and the Burica
40 Peninsula before any inferences about terraces of either MIS 3 or MIS 5 age can be made.

41 **5 Last Interglacial sea level fluctuations**

42 One of the issues that has been actively debated in the past few decades is whether MIS 5e was characterized by a single
43 sea-level high stand or multiple high stands. Some of the original evidence for more than one high stand came from the Huon
44 Peninsula of New Guinea, where reefs VIIa and VIIb were interpreted to represent early and later high-sea stands of MIS 5e,
45 respectively (Bloom et al., 1974; Chappell, 1974). U-series dating of corals from these two terraces using TIMS methods
46 confirmed that both terraces likely date to MIS 5e (Stein et al., 1993). Such a record on New Guinea does not, however,
47 require that this was a global phenomenon, because coseismic uplift has been well documented for this coast, with as many as
48 six coral reefs emerging in the Holocene alone (Ota et al., 1993). On Barbados, the uplift rate is much lower than that on New
49 Guinea, but more than one high stand of sea during MIS 5e has been proposed here as well (Schellmann and Radtke, 2004;
50 Thompson and Goldstein, 2005). Unlike New Guinea, however, multiple Holocene terraces have not been reported on
51 Barbados, and coseismic uplift is a less likely explanation for possible multiple LIG terraces.

52 In addition to tectonically active coastlines, there have been claims of multiple sea stands during MIS 5e from deep-sea
53 records and reefs on tectonically stable coastlines. Rohling et al. (2008), studying the oxygen isotope record in planktonic
54 foraminifera recovered from Red Sea sediment cores, suggested that there could have been as many as four separate high
55 stands of sea during MIS 5e. Thompson et al. (2011) reported TIMS U-series ages of corals from San Salvador Island and
56 Great Inagua Island in the Bahamas, proposing at least two high stands during MIS 5e, and possibly as many as four high
57 stands, similar to the Red Sea record of Rohling et al. (2008).

58 Modeling efforts have also addressed the question of a dual high-sea stand during the LIG. Kopp et al. (2009) conducted
59 a statistical analysis of a database generated from many reported MIS 5e deposits worldwide, from both tectonically active

60 and stable coastlines. These investigators concluded that early within MIS 5e there was a sea-level high, followed by a drop
61 of ~4 m, succeeded by another sea-level high. Unfortunately, some of the hypothesized MIS 5e sites used by Kopp et al.
62 (2009) are either poorly dated or not dated at all, rendering this reconstruction uncertain. In a more recent review of both field
63 and modeling evidence, Barlow et al. (2018) concluded that there is no evidence of more than one high-sea stand during MIS
64 5e. Along the Pacific Coast of North America, there has been, thus far, no evidence of more than one high-sea stand during
65 MIS 5e, along either tectonically stable or uplifting coasts.

66 **6 Other interglacials**

67 **6.1 Interglacials prior to MIS 5e**

68 Because of ongoing tectonic processes during the Quaternary, multiple marine terraces are recorded along much of the
69 Pacific Coast of North America, from southern Oregon to Baja California. As noted earlier, Woodring et al. (1946) mapped
70 13 marine terraces in the Palos Verdes Hills (Fig. 26), the highest of which is at an elevation of ~400 m. Vedder et al. (1957,
71 1975) and Grant et al. (1999) documented at least six marine terraces above the Newport Mesa terrace, correlated to MIS 5e,
72 in the Newport Beach area. Vedder and Norris (1963) mapped 14 marine terraces on San Nicolas Island (Fig. 36a), with the
73 highest at an elevation of ~270 m. Fossils are found in deposits of all 14 terraces. On this island, amino acid ratios in fossil
74 *Tegula* specimens show a steady increase with terrace elevation (Fig. 36b). By the time the 8th and 10th terraces are reached,
75 D-alloisoleucine/L-isoleucine values in *Tegula* are at equilibrium values of ~1.25, indicating considerable antiquity. San
76 Clemente Island hosts more than 20 marine terraces, and these landforms show superb geomorphic preservation (Fig. 37).
77 Fossil-bearing marine terrace deposits are found as high as ~265 m (Cockerell, 1939), similar to San Nicolas Island, and the
78 highest marine terrace is found at an elevation of nearly 600 m. If the late Quaternary uplift rate has been steady over the
79 history of the island (Muhs et al., 2014a), the highest terrace on San Clemente Island could be ~3 Ma. Even some of the
80 smallest islands off the California coast host a long-term history of interglacial high-sea stands superimposed on steady uplift.
81 Santa Barbara Island has an area of only ~2.6 km², yet it hosts at least five marine terraces, up to an elevation of ~100 m (Fig.
82 38). On many of the California islands, pre-MIS 5e marine terraces are distinguished from younger terraces by the presence
83 of the extinct fossil gastropod *Pusio fortis* (formerly *Calicantharus fortis*). For example, on Santa Barbara Island, this taxon
84 is found in deposits of the 2nd, 3rd, and 4th terraces (Fig. 38), but is not found in deposits of the 1st terrace, which appears to
85 contain a mix of fossils dating to MIS 5e and MIS 5c (Muhs and Groves, 2018). Multiple marine terraces are found along the
86 Pacific coast of Mexico as well. Rockwell et al. (1989) recognized 14 marine terraces on Punta Banda, in northern Baja
87 California, with the highest at an elevation of ~347 m. Farther south, Orme (1980) mapped multiple marine terraces, with the
88 highest between Cabo San Quintin and Punta Baja (Fig. 30), at an elevation of ~300 to ~357 m.

89 Unfortunately, there are few data on the possible ages of pre-MIS 5e terraces on the Pacific Coast of North America.
90 Indeed, numerical ages of terraces dating from MIS 7, 9, and 11 have yet to be confirmed for any part of the Pacific coast of

91 the continent, although it is likely that marine terraces representing these high-sea stands are preserved. Although corals are
92 present in deposits of several higher elevation terraces, open-system histories have likely prevailed in many of these fossils.
93 For example, Muhs et al. (2004) presented U-series data for corals from the 10th terrace (elevation ~236 m) on San Nicolas
94 Island, indicating possible ages of ~600 ka to ~450 ka. These apparent ages are, however, not consistent with the late
95 Quaternary uplift rate, nor are they consistent with amino acid ratios at equilibrium values in fossil mollusks from this terrace
96 (Fig. 36b). A more promising isotopic method of age determination for fossils of pre-MIS 5e terraces on the Pacific Coast is
97 Sr isotope stratigraphy, a calibrated method of geochronology. Early experiments with this method in California showed
98 promise (Ludwig et al., 1992) and since that time, better calibration curves have been developed (Howarth and McArthur,
99 1997).

00 Latitudinal, north-south-trending aminozones parallel or subparallel to MIS 5a and MIS 5e aminozones, show the potential
01 for at least lateral correlation of older, pre-MIS 5e marine terraces. Wehmiller (1982) used such an approach on the Pacific
02 Coast of North America, from southern Baja California Sur to Oregon. His data showed the possibility for marine records
03 prior to MIS 5e, including high-sea stands associated with MIS 7, 9, 11, 13, and 15.

04 Another possibility for dating older terraces is the use of kinetic modeling with amino acid ratios. In this approach, a
05 theoretical kinetic pathway is used with a calibrated amino acid ratio for shells from a deposit that is independently dated (such
06 as by U-series on coral). Clarke and Murray-Wallace (2006) review the various mathematical expressions for different kinetic
07 pathways. One of the most widely used method is the parabolic kinetic model, derived from heating experiments that simulate
08 long periods of geologic time (Mitterer and Kriausakul, 1989). When this method is applied to the terrace sequence on San
09 Nicolas Island (Fig. 36b), using the ~120 ka age for terrace 2a and its D-alloisoleucine/L-isoleucine value of 0.52, the higher
10 amino acid ratios for the older terraces yield apparent ages of ~375 ka (terrace 4), ~480 ka (terrace 5), ~510 ka (terrace 6), and
11 ≥ 680 ka for terraces 8 and 10. If these ages are correct, they would permit correlation of terrace 4 with MIS 11 and terraces 5
12 and 6 with MIS 13. Along the terrace transect shown in Figure 36b, this would also imply that terraces that formed during
13 MIS 7 and MIS 9 were likely removed by erosion during MIS 5e. Interestingly, the ages and uplift rates derived from older
14 terraces in this exercise are similar to the uplift rate derived from the MIS 5e terrace. While all of these implied results seem
15 reasonable geologically, it is important to remember that kinetic modeling of amino acid racemization and epimerization is
16 still theoretical and age estimates derived from such an approach are simply possibilities for additional testing.

17 Still another method to address the question of ages of older marine terrace deposits is the use of cosmogenic isotopes. In
18 the San Diego region, one of the oldest marine terrace deposits is called the Clairemont terrace, part of a larger complex of
19 marine terrace and beach ridge deposits called the Lindavista terrace sequence. Early data on *Tivela stultorum* valves from
20 the Lindavista unit showed D/L values at or near equilibrium for all amino acids (Wehmiller et al., 1977a), implying
21 considerable antiquity. Based on data in Lajoie et al. (1991), as many as 13 terraces occur above the MIS 5e terrace, each with

22 wave-cut benches and prominent beach ridges. The Clairemont terrace is found at an elevation of ~96 m above sea level, and
23 Simms et al. (2020) used cosmogenic nuclides at two localities to estimate an age of ~1.48 Ma for this terrace.

24 **6.2 High-sea stands after MIS 5e**

25 Whereas ages of pre-MIS 5e marine terraces on the Pacific Coast of North America are rare, there are several marine terrace
26 ages that postdate the peak of the last interglacial period, mostly for the relatively high-sea stands of MIS 5c (~100 ka) and
27 MIS 5a (~80 ka). With regard to MIS 5c, TIMS U-series ages of corals dating to this high-sea stand have been confirmed, but
28 mixed with MIS 5e deposits at Cayucos, Point Loma, and San Nicolas Island, as discussed above (Stein et al., 1993; Muhs et
29 al., 2002a, 2012). In two other areas, both of which have somewhat higher late Quaternary uplift rates, there are terraces that
30 are good candidates for MIS 5c records, although both are as yet undated. On the Palos Verdes Hills, what Woodring et al.
31 (1946) mapped as the "2nd" and "4th" terraces have been dated to ~80 ka and ~120 ka, respectively (Muhs et al., 2006), as
32 noted earlier. To avoid confusion with terrace numbering that is inconsistent from this area to nearby San Pedro, Muhs et al.
33 (2006) named these (informally) as the Paseo del Mar (2nd) and Gaffey (4th) terraces. In the western part of the Palos Verdes
34 Hills (Fig. 26), there is an intermediate-elevation terrace that Woodring et al. (1946) mapped as the "3rd" terrace. Because it
35 occupies a morphostratigraphic position between the ~80 ka (2nd) Paseo del Mar and ~120 (4th) Gaffey terraces, it is very
36 likely that this terrace records the MIS 5c high-sea stand. Woodring et al. (1946) did not report any fossil localities on this
37 terrace, and 30 years of periodic searches by the present author have not resulted in any either, so the terrace remains undated.

38 The other locality that provides morphostratigraphic evidence of a possible MIS 5c record is Punta Banda, in northern Baja
39 California. At this locality (Fig. 31), Rockwell et al. (1989) reported a small terrace fragment at ~22 m above sea level above
40 the Lighthouse (1st) terrace at ~15 m and the Sea Cave (3rd) terrace at ~34 m. Similar to the Palos Verdes Hills, the Lighthouse
41 and Sea Cave terraces are dated to ~80 ka and ~120 ka, respectively, by both alpha-spectrometric U-series (Rockwell et al.,
42 1989) and TIMS U-series methods (Muhs et al., 2002a). Unfortunately, as with the Palos Verdes Hills, no corals have yet
43 been found on the 2nd, ~22 m terrace on Punta Banda.

44 While marine terraces dated to MIS 5c are rare on the Pacific Coast of North America, terraces dated to MIS 5a are abundant
45 (Fig. 39 and Table S3). Corals have been acquired and dated by TIMS U-series methods at Coquille Point (Oregon), Point
46 Arena (northern California), three localities between Point Año Nuevo and Santa Cruz (central California), Santa Rosa and
47 San Nicolas Islands, the Palos Verdes Hills, and Point Loma (all in southern California), and Punta Banda (northern Baja
48 California). Analytical and faunal data for these terraces are given in Addicott (1966), Zullo (1969), Kern (1977), Kennedy

49 (1978), Rockwell et al. (1989), and Muhs et al. (2002a, 2006, 2012). In all cases, the faunas are characterized by cool-water
50 forms, with several extralimital northern and northward-ranging species.

51 In addition to U-series-dated localities, a large number of localities lack corals, but have mollusks that permit
52 aminostratigraphic correlation to MIS 5a, following the approach pioneered by Wehmiller et al. (1977a) and Kennedy et al.
53 (1982). These localities can be found from near Newport, Oregon, into northern California, and to the Channel Islands of
54 southern California (Fig. 39). Like their U-series-dated counterparts, these terraces host faunas with extralimital northern or
55 northward-ranging species of mollusks.

56 It is very likely that there are terraces dating to MIS 5a and/or MIS 5c in Baja California, Baja California Sur, and Sonora
57 as well, based on amino acid and faunal studies by Emerson et al. (1981) and Ortlieb (1987). In addition, along the Pacific
58 coast of northern Baja California, numerous fossil localities, shown earlier in Figure 30, contain mixes of warm-water and
59 cool-water molluscan faunas. Although none of these terraces have either U-series or aminostratigraphic data for age control
60 (with the exception of Camalú, as noted earlier), their low elevations allow for the possibility that they record some part of
61 MIS 5. The mixes of cool-water and warm-water mollusks invite comparison to similar mixes of faunas with contrasting
62 thermal aspects, found at Cayucos, Point Loma, and San Nicolas Island, that have U-series ages on corals that include both
63 MIS 5e and 5c.

64 Two localities in southern California have U-series and amino acid evidence for emergent marine terraces dating to MIS 3.
65 Both localities are south of the "big bend" in the San Andreas Fault (Figs. 1, 2) where this large constraining bend brings about
66 a shift from fault lateral movement to predominantly crustal compression between the Pacific and North America plates. The
67 result is unusually high rates of uplift, such that terraces formed when sea level was substantially lower than present are now
68 emergent. Isla Vista, a university community (Fig. 25), is built on a marine terrace whose outer edge is at an elevation of ~7
69 m. Based on amino acid ratios in *Saxidomus* valves, Wehmiller et al. (1977a), Wehmiller (1982), and Kennedy et al. (1982)
70 thought that this terrace likely postdated MIS 5a. This conclusion was supported by the fact that the fauna within the deposits
71 of this terrace contains a large number of extralimital northern species (Wright, 1972), consistent with very cold waters off the
72 California coast at this time. This inference is expected for the time period postdating MIS 5a, based on independent evidence
73 (Kennett and Venz, 1995). Gurrola et al. (2014) reported U-series ages of ~49 ka and ~47 ka for corals from this terrace,
74 which support the original age interpretations (see Muhs et al., 2014b, for isotopic data for one of these specimens). All these
75 investigators correlated this terrace with MIS 3.

76 Also in southern California, there is a terrace that has been correlated to MIS 3 near a locality simply called "Sea Cliff,"
77 northwest of Ventura (Fig. 25). Along an ~6 km reach of coastline here, there are two marine terraces, a low-elevation surface
78 dated to the Holocene (see discussion below) and a higher elevation terrace of Pleistocene age above it. The Pleistocene
79 terrace has a variable elevation in a shore-parallel sense, from just over ~100 m to just over ~200 m above sea level (Wehmiller

80 et al., 1978). Although no corals have yet been found in the deposits of the Pleistocene terrace, amino acid ratios indicate that
81 it is likely ~50 ka, similar to the terrace at Isla Vista (Wehmiller et al., 1978; Kennedy et al., 1982; Wehmiller, 1982). The
82 elevation of this terrace, along with its young age and formation at a time of relatively low sea level indicates that this reach
83 of coastline has experienced an extremely high rate of uplift.

84 In northern California, near the Mendocino triple junction of the Gorda, Pacific, and North America plates, there is a third
85 locality with a marine terrace correlated to MIS 3 (McLaughlin et al., 1983a, b). This terrace is found near Point Delgada (Fig.
86 21) and has a maximum elevation of ~7 m above sea level. Correlation of this terrace to MIS 3 is based on a radiocarbon age
87 of ~45 ka from fossil wood found in terrestrial deposits that overlie the marine terrace deposits. Although radiocarbon ages
88 on wood are usually reliable, this apparent age is near or at the limit of the method and in addition is found within overlying
89 deposits, not the marine terrace deposits themselves. Thus, the cautions discussed earlier with regard to modern carbon
90 contamination would apply here as well. Although the age from Point Delgada is interpreted to be a close, minimum-limiting
91 age, it is in fact just a minimum-limiting age and the terrace itself could be older.

92 Based on early results of amino acid geochronology in Kennedy et al. (1982), it was originally thought that marine terrace
93 deposits at a fourth locality, Cape Blanco, Oregon (Figs. 18-20), could correlate with MIS 3. Later amino acid studies, linked
94 with a nearby U-series-dated, coral-bearing locality (Coquille Point, Oregon), showed that the low terrace at Cape Blanco
95 likely dates to MIS 5a (Muhs et al., 1990). The fauna at Cape Blanco, with its cool-water species, is similar to that at Coquille
96 Point (Muhs et al., 2006). Furthermore, oxygen isotope ratios in fossil *Saxidomus gigantea* and *Mya truncata* collected from
97 the two localities do not have significant differences (Muhs et al., 1990).

98 **6.3 Holocene sea level indicators**

99 Emergent Holocene marine deposits are found at several localities along the Pacific Coast of North America. Within the
00 southern Puget Sound area of Washington State (Fig. 16), emergent marine terraces or peat-covered tidal flats are found at
01 five localities, as much as ~7 m above sea level, with ages ranging between ~1,000 and 1,100 yr B.P. (Bucknam et al., 1992).
02 It has long been recognized (e.g., Kelsey, 1990) that there is the potential for coseismic uplift along the zone where the Juan
03 de Fuca plate is being subducted beneath the North America plate (Fig. 2). What is interesting about the Holocene terraces in
04 the Puget Sound area, however, is that they are some distance inland from this subduction zone. Bucknam et al. (1992) attribute
05 Holocene uplift here to reverse slip along an inferred fault within the crust of the North America plate.

06 Near the Mendocino triple junction area of northern California (Figs. 2, 21), Holocene marine terraces have also been
07 documented (Lajoie et al., 1991). The most recent of these produced 1.4 m of uplift associated with the Ms 7.1 earthquake at
08 Cape Mendocino in 1992 (Carver et al., 1994). Merritts (1996) reported that earlier Holocene, coseismic uplift events had
09 occurred prior to the A.D. 1992 earthquake. Based primarily on radiocarbon ages of marine shells, at least four such events
10 occurred between ~7 ka and ~0.6 ka.

11 South of the “big bend” area of the San Andreas Fault (Figs. 2, 25), crustal compression is the dominant tectonic style.
12 Thus, in the area to the south, uplift rates are very high. Sarna-Wojcicki et al. (1987) mapped two marine terraces in this area,
13 between Ventura and Santa Barbara. The higher of the two terraces ranges in elevation from ~120 to ~210 m, and amino acid
14 data in mollusks reported by Wehmiller et al. (1978) indicate that it is likely ~45 ka (MIS 3), as discussed above. The lower
15 of the two terraces has elevations that range from ~6 to ~35 m. Radiocarbon ages of marine mollusks from this terrace range
16 from ~5 ka to ~1.8 ka (Sarna-Wojcicki et al., 1987). More recent detailed work by Rockwell et al. (2016) identified four
17 Holocene terraces in this area, with radiocarbon ages of ~6.7 ka, ~4.4 ka, ~2.1 ka, and ~0.95 ka. Each terrace represents a
18 separate coseismic uplift event.

19 Still farther south, the coast of Central America is adjacent to the subduction zone, where the northeast-moving Cocos plate
20 is being subducted beneath the Caribbean plate (Fig. 35). In addition, the Panama triple junction is situated just south of the
21 Costa Rica-Panama border, where the Cocos, Caribbean, and Nazca plates intersect. On the Nicoya Peninsula of Costa Rica,
22 Marshall and Anderson (1995) reported marine terraces at elevations of ~4 to ~16 m above sea level, with radiocarbon ages
23 of ~4.1 ka to ~0.4 ka. Gardner et al. (2001), working the same general area, reported similar elevations and ages for two
24 terraces, in agreement with, but adding detail to the study of Marshall and Anderson (1995). Fisher et al. (1998) ascribed uplift
25 in this region to subduction of seamount chains on the Cocos plate. Marine terraces of Holocene age have also been reported
26 for the nearby Osa Peninsula of Costa Rica (Gardner et al., 2013) and the Burica Peninsula of Panama (Fig. 35) by Morell et
27 al. (2011).

28 **6.4 Implications for paleozoogeography**

29 There has been considerable interest in MIS 5e not only for its implications for future sea-level rise, but also for warming
30 of the oceans. Indeed, sea-level rise is linked to ocean warming due to the possibility of thermal expansion of the world's
31 oceans. In addition, however, possible ocean warming during MIS 5e has importance for understanding how modern marine
32 ecosystems might respond to future warming.

33 Global-scale studies of MIS 5e have been carried out using proxy paleoclimate data from deep-sea cores with the goal of
34 estimating sea surface temperatures (SST). Results of these investigations have not been entirely consistent. CLIMAP Project
35 Members (1984) concluded that overall, the last interglacial ocean was not significantly different from the modern ocean. It
36 is important to note, however, that for many regions of the world, including much of the ocean around Australia, the
37 Mediterranean Basin, the Bering Sea, the central Pacific Ocean, and the eastern Pacific Ocean off North America, there were
38 few cores available. Using a larger dataset, Turney and Jones (2010) concluded that MIS 5e global temperatures were on
39 average ~1.5°C warmer than present, although part of this conclusion is based on ice and terrestrial records. In yet another
40 compilation, McKay et al. (2011) concluded that on a global scale, SST during MIS 5e was not significantly different from
41 the present. From this, these investigators inferred that thermal expansion likely played only a minor role, if any, in the higher
42 than present sea level during MIS 5e. However, as the same investigators also pointed out, some regions are exceptions to this
43 generalization.

44 In some regions where core data are sparse, shallow invertebrate marine fossil faunas serve as an important record of SST.
45 For example, extralimital species of mollusks and corals, indicating warmer SST during MIS 5e, have been documented in the
46 Indian Ocean along the western coast of Australia (Kendrick et al., 1991), around New Zealand and the southern coast of
47 Australia (Murray-Wallace et al., 2000), along the Bering Sea and Arctic Ocean coasts of Alaska (Brigham-Grette and
48 Hopkins, 1995), in the eastern Atlantic Ocean off Africa and the Mediterranean Basin (Cuerda, 1975, 1987, 1989; Cuerda and
49 Sacarès, 1992; Hearty et al., 1986; Meco et al., 2002, 2006; Muhs et al., 2014c), in the western Atlantic Ocean around Bermuda
50 (Richards et al., 1969; Muhs et al., 2002b), and in the Pacific Ocean along the shores of Oahu in the Hawaiian Islands (Kosuge,
51 1969; Muhs et al., 2002b; Groves, 2011). The sites studied in Australia, the central Pacific, the western Atlantic Ocean, the
52 eastern Atlantic Ocean off Africa, the Mediterranean, and the central Pacific are all localities anchored by reliable MIS 5e U-
53 series ages on corals.

54 In the context of both deep-sea core proxy climate data and shallow-water marine invertebrate records from around the
55 globe, it is interesting to consider what the SST off the Pacific Coast of North America was during MIS 5e. Herbert et al.
56 (1998) showed that alkenone unsaturation indices, derived from modern core-top samples, correlate in a linear fashion with
57 modern SST. Using this relation, Herbert et al. (2001) generated both oxygen isotope values in foraminifera (to identify MIS
58 5e) and alkenone unsaturation indices to estimate SST in five cores, taken off northern California to south of Cabo San Lucas,
59 Baja California Sur. In all cases, SST during MIS 5e is substantially higher than at present or during earlier parts of the
60 Holocene. One of the cores examined (Ocean Drilling Project, or ODP 893) is from Santa Barbara Basin, the same locality
61 studied for temperature-sensitive foraminiferal species by Kennett and Venz (1995). The latter workers found that MIS 5e
62 was the only time, other than the Holocene, when warm-water foraminifera were present in Santa Barbara Basin, in good
63 agreement with the alkenone unsaturation index data. Two other cores, one off central California and one off southern
64 California, studied by Yamamoto et al. (2007), also gave alkenone-based SST indicating substantially warmer waters off the
65 Pacific Coast during MIS 5e compared to present.

66 Given these findings, it is pertinent to evaluate the shallow-water marine terrace records of mollusks and other invertebrates
67 of MIS 5e age from the Pacific Coast of North America. As noted earlier, pioneering amino acid studies by Wehmiller et al.
68 (1977a) and Kennedy et al. (1982) considered that marine terrace deposits correlated to MIS 5a (~80 ka) had cool-water faunas
69 and those correlated to MIS 5e (~120 ka) had warm-water faunas. While the cool-water forms that are so prominent in terraces
70 correlated to MIS 5a by amino acid geochronology (Kennedy et al., 1982) have been largely confirmed to indeed be ~80 ka,
71 based on TIMS U-series ages of corals (Muhs et al., 2002a, 2006), the GIA-related fossil mixing of ~120 ka and ~100 ka (MIS
72 5c) fossils into single terrace deposits complicates matters. Nevertheless, using those localities where there is good evidence
73 for MIS 5e age fossils and assuming that the cool-water forms represent MIS 5c, the shallow-water marine invertebrate record
74 still allows some inferences about SST during the peak of the last interglacial period. Here, examples of bivalves, gastropods
75 and corals are examined from dated deposits to illustrate what can be inferred about ocean temperatures during MIS 5e.

76 Two species of bivalves that live dominantly in tropical waters off the Pacific Coast of North America are *Chione undatella*
77 and *Dosinia ponderosa*. *C. undatella* is one of two species of *Chione* (*C. californiensis* is the other) that presently live only

78 south of the Point Conception area (Fig. 40 a). Although *C. undatella* is found only as far north as Goleta (near Santa Barbara),
79 California, it ranges south along the coast of Mexico, including the Golfo de California, to Peru, and is also found on the
80 Galapagos Islands (Coan and Valentich-Scott, 2012). In fossil form, *C. undatella* is found in several marine terrace deposits
81 either directly dated to MIS 5e or correlated to it on the basis of amino acids, from Bahía Magdalena, Baja California Sur,
82 north to Potrero Canyon near Los Angeles, California (Fig. 40a). In addition, however, it has also been reported from a terrace
83 correlated to MIS 5e from near San Luis Obispo Bay by Kennedy (2000) and also has been found at Tomales Bay, California.
84 The presence of *C. undatella* at Tomales Bay is particularly significant, because this locality is ~500 km northwest of its
85 modern northern limit. *Dosinia ponderosa* at present ranges only as far north as Laguna Ojo de Liebre, on the Pacific coast
86 of Baja California Sur, just east of Punta Eugenia (Fig. 40b). Like *C. undatella*, *D. ponderosa* ranges south along the coast of
87 Baja California Sur, including the Golfo de California, all the way to Peru and including the Galapagos Islands (Coan and
88 Valentich-Scott, 2012). In fossil form, it is found at several localities dated or correlated to MIS 5e in Baja California Sur and
89 Sonora, all within its present range, but also as far north as Potrero Canyon near Los Angeles, California, and at Newport Bay
90 (Fig. 40b). These California localities are ~750 km northwest of the modern northern limit of *D. ponderosa*.

91 The gastropod fossil records from MIS 5e terrace deposits also show that what are now southern species lived farther north
92 during the last interglacial period. *Mexacanthina lugubris* is a gastropod only rarely found as far north as San Diego. Bertsch
93 and Aguilar Rosas (2016) report that on the Pacific coast, *M. lugubris* presently lives from San Diego to Cabo San Lucas and
94 on the eastern Golfo de California coast, the species is found from Bahía Kino, Sonora to Mazatlán, Sinaloa. Fossil occurrences
95 of *Mexacanthina lugubris* in deposits dated to ~120 ka are found all along this taxon's modern Pacific Coast distribution, from
96 Bahía de Magdalena, Baja California Sur, to Point Loma, near San Diego. However, there are also some occurrences reported
97 in MIS 5e deposits, well north of the modern range endpoint for *Mexacanthina lugubris* (Fig. 41a). Although warmer waters
98 during the last interglacial period allowed *Chione undatella* to migrate north of its modern range by several hundred kilometers,
99 Point Conception was apparently a barrier to northward migration of *Mexacanthina lugubris* beyond the Santa Barbara region.
00 Another gastropod, *Stramonita biserialis*, presently lives from Cedros Island, just north of Punta Eugenia, south along Baja
01 California Sur, throughout the Golfo de California, and all the way to Chile, as well as being on the Galapagos Islands (Keen,
02 1971). In MIS 5e deposits, it is found at localities in the Golfo de California and along the coast of Baja California Sur, all
03 within its modern range (Fig. 41b). However, it is also found in MIS 5e deposits on Isla Guadalupe, along the northwestern
04 coast of Baja California, and in some southern California localities as far north as San Pedro, California, near Los Angeles
05 (Fig. 41b). The occurrence of *S. biserialis* in San Pedro is a northward extension of its modern range by nearly 700 km.

06 One particularly interesting locality, with U-series ages of ~120 ka on corals, is the low-elevation marine deposit on Isla
07 Guadalupe, off the Pacific coast of Baja California (Figs. 42, 43). Isla Guadalupe is interesting zoogeographically even at
08 present, because its modern marine invertebrate fauna has elements of both the Californian and Panamanian faunal provinces
09 (Lindberg et al., 1980), making it a particularly sensitive area. The MIS 5e marine deposits here have a fauna that has been
10 reported by Lindberg et al. (1980) and Durham (1980). Muhs et al. (2002a) summarized the modern geographic ranges of this
11 fauna, showing that it has a substantial number of extralimital southern species of taxa, along with what was then thought to

12 be two northward-ranging species. With new data on ranges of species that have been published since that time (Coan and
13 Valentich-Scott, 2012; Berschauer and Clark, 2018), that paleozoogeographic analysis has been redone here. Results indicate
14 that the fauna contains no northward-ranging species, but hosts 13 extralimital southern species (Fig. 44). All but three of
15 these taxa have southern range endpoints south of the equator, and six species have northern range endpoints no farther north
16 than Bahía Magdalena, which is over 700 km southeast of Isla Guadalupe.

17 In addition to the extralimital southern species of bivalves and mollusks within the MIS 5e fauna of Isla Guadalupe, there
18 are two other taxa which merit additional discussion for the paleozoogeographic significance. The MIS 5e deposits of Isla
19 Guadalupe host the North American Pacific Coast's northernmost MIS 5e occurrence of a hermtypic colonial coral, *Pocillopora*
20 *guadalupensis* (Durham, 1980). This species is not known to be living in the eastern Pacific at present (Reyes-Bonilla and
21 López-Pérez, 1998), nor has it been found in other Pleistocene or Pliocene marine deposits in the region (López Pérez, 2008).
22 The eastern Pacific region at present hosts five species of *Pocillopora*, with one species (*P. verrucosa*) found as far north as
23 Isla San Marcos, on the eastern coast of Baja California Sur, and four species currently living offshore near Cabo San Lucas
24 (Reyes-Bonilla and López-Pérez, 1998). However, Isla Guadalupe is ~1000 km northwest of Cabo San Lucas (Fig. 42).
25 Durham (1980) pointed out that *P. guadalupensis* more closely resembles *Pocillopora* species of the central and western
26 Pacific Ocean (such as *P. ligulata*) than it does to species of this genus found in the southeastern Pacific, a relationship he
27 described as "strange." However, it is now known that living *P. ligulata* is found not only in the central Pacific Ocean
28 (including the Hawaiian Islands) but is also found off the coasts of Colombia and Ecuador (Glynn et al., 2017). Thus, one
29 could hypothesize that perhaps there is a last interglacial evolutionary link between *P. ligulata* from tropical waters of
30 northwestern South America and *P. guadalupensis* of Isla Guadalupe.

31 The other particularly noteworthy fossil reported by Lindberg et al. (1980) from the MIS 5e fauna of Isla Guadalupe is the
32 cowry *Cypraea (Erosaria) cernica*, now called *Naria cernica*. This species is not known to occur anywhere along the Pacific
33 coasts of the Americas, and currently lives in the tropical waters of the Indo-West Pacific province (Burgess, 1985). The
34 closest living populations of this species to Isla Guadalupe are in the Hawaiian Islands (Severns, 2011), although interestingly,
35 the species has not yet been reported as a fossil within MIS 5e (or older) deposits on the Hawaiian Islands (Groves, 2011). In
36 any interpretation, however, Isla Guadalupe is thousands of kilometers away from any present location where *Naria cernica*
37 can be found, making its presence on this island during MIS 5e a remarkable find.

38 Collectively, the MIS 5e fossil record for bivalves (*Chione undatella*, *Dosinia ponderosa*), gastropods (*Mexacanthina*
39 *lugubris*, *Stramonita biserialis*, *Naria cernica*), and coral (*Pocillopora guadalupensis*) from several localities, from Baja
40 California to northern California, indicates that water temperatures off the Pacific Coast of North America were substantially
41 warmer than present. It does not appear that there were wholesale shifts of entire faunal provinces, such as the present-day
42 Californian province being replaced entirely by Panamic species (Fig. 43). Lindberg et al. (1980) point out that of the modern
43 fauna of Isla Guadalupe, ~75% are from the Californian province, ~6% are Panamic, and ~19% are biprovincial. In contrast,
44 the MIS 5e fauna of Isla Guadalupe consists of ~39% Californian species, 32% Panamic species, and ~29% biprovincial
45 species. Thus, while it is clear that greater numbers of warm-water species lived in more northerly locations than is the case

46 today, certain species existed within each faunal province as they do today. Furthermore, it appears that some of the physical
47 geographic barriers that define provincial boundaries also served as barriers during MIS 5e, despite northward migrations. A
48 good example of this is the migration of *Mexacanthina lugubris* north of where its present northern limit is situated, but
49 apparently Point Conception prevented this taxon from migration farther north, into what is now the Oregonian province (Fig.
50 43). Still, while it may be difficult to quantify the degree of ocean warming off the Pacific Coast during MIS 5e, the presence
51 of numerous extralimital species in terrace deposits at many localities is consistent with the alkenone and foraminiferal data
52 from deep sea cores that eastern Pacific Ocean SST were higher than present, from northern California to southern Baja
53 California.

54 **6.5 Controversies**

55 In some of the earliest studies of late Quaternary sea level history, supported by what was then the relatively new U-series
56 dating method, there was general agreement that sea level stands during MIS 5c (~100 ka) and MIS 5a (~80 ka) were
57 substantially below modern sea level, by as much as 10 to 20 m (Broecker et al., 1968; Meselella et al., 1969; Veeh and
58 Chappell, 1970; Bloom et al., 1974; Chappell, 1974). These early studies were on Barbados and New Guinea, two areas far
59 apart from one another, unrelated tectonically, and having quite different long-term uplift rates. The broad agreement in
60 paleo-sea level estimates for MIS 5c and 5a at both localities seemed to provide support for a "global" eustatic sea-level
61 history for the late Quaternary. Later, a third locality with emergent reef terraces, the northwest coast of Haiti, showed
62 general agreement with Barbados and New Guinea for paleo-sea levels at MIS 5c and 5a (Dodge et al., 1983), which
63 reinforced the concept of a global eustatic sea-level curve. For New Guinea, the original paleo-sea level estimates were
64 refined by Chappell and Shackleton (1986).

65 Because marine terraces on many coastlines lack materials suitable for dating, a number of graphical methods emerged in
66 an attempt to compare the elevation spacing of a suite of undated marine terraces with a "global" sea level curve. Most of
67 these schemes assumed that the detailed paleo-sea level record of the Huon Peninsula of New Guinea (Chappell and
68 Shackleton, 1986) is a faithful representation of global, eustatic sea level change. Lajoie (1986) even ventured the opinion
69 that dating a suite of marine terraces was simply a matter of correlating the undated landforms with the appropriate peaks on
70 the New Guinea sea level curve. Similarly, Bull (1985) proposed that dating of an entire suite of otherwise undated marine
71 terraces could be accomplished solely by graphical means. In Bull's (1985) method, a given terrace was assigned an age and
72 paleo-sea level corresponding to a possible correlative terrace on New Guinea. The resultant uplift rate, along with the New
73 Guinea sea level curve, was used to plot inferred amounts of uplift for other terraces. The process was repeated for different
74 assumed ages of the original terrace chosen and different uplift rates. Whichever of the resultant plots yielded the best-fit
75 linear array of points on an inferred uplift vs. age plot was interpreted to be the correct correlation and uplift rate. In this
76 method, once the "correct" uplift rate was identified, all terraces in the suite were dated simultaneously. The technique and
77 variations of it have been applied to undated or partially dated terrace sequences in New Zealand (Bull and Cooper, 1986),

78 northern California (Merritts and Bull, 1989; McCrory, 2000), central California (Hanson et al., 1994), southern California
79 (Trecker et al., 1998), Mexico (Mayer and Vincent, 1999), and Italy (Calanchi et al., 2002).

80 Despite the apparent agreement for a global eustatic sea level curve, there were always localities with marine terrace
81 elevations that did not seem to fit the Barbados-New Guinea sea level curve for the late Quaternary. On the Atlantic Coastal
82 Plain of the USA, emergent marine deposits, a few meters above sea level, gave U-series ages on coral of ~80 ka (Cronin et
83 al., 1981), unexpected on a passive continental margin, given the sea level estimates at this time from Barbados and New
84 Guinea. Similar results were obtained on tectonically stable Bermuda, where the marine facies of the Southampton
85 Formation, at 1-2 m above sea level, yielded U-series ages on coral averaging ~80 ka (Harmon et al., 1983). Later studies on
86 both Bermuda and the Atlantic Coast Plain, with more elevation measurements and precise TIMS U-series dating, gave the
87 same results as these early studies (Muhs et al., 2002b; Wehmiller et al., 2004). On the tectonically active Ryukyu Islands of
88 Japan, where reef terraces dating to ~120 ka, ~100 ka, and ~80 ka are all present, elevations yield paleo-sea levels at MIS 5c
89 and 5a that are close to present (Ota and Omura, 1992). U-series ages and terrace elevations from the Pacific Coast of North
90 America (California and Mexico) also give paleo-sea level estimates for MIS 5c and 5a that are much closer to present sea
91 level than what would be expected from the Barbados-New Guinea records (Muhs et al., 1994). Thus, despite the
92 attractiveness of being able to date, using graphical techniques, an entire suite of marine terraces with no independent age
93 control, it is in fact a hazardous practice.

94 The explanation for the disagreement between some paleo-sea level estimates and the Barbados-New Guinea sea-level
95 history is likely due to GIA processes. Indeed, GIA effects can and should be expected in high-latitude and mid-latitude
96 regions of the Northern Hemisphere where large, continental ice sheets were found during glacial periods. Thus, using the
97 Barbados-New Guinea sea level curve for terrace correlations via elevation spacing in such regions will likely yield spurious
98 results. In some far-field regions, distant from the Laurentide, Cordilleran and Fennoscandian ice sheets, the elevation-
99 spacing method of terrace correlation might be applicable, but in virtually all mid-latitude and high-latitude regions, the
00 approach is untenable. Creveling et al. (2015) modeled apparent sea levels around the world, assuming a true "eustatic"
01 high-sea level of +6 m. These investigators showed that relative sea level, at the end of MIS 5e, could have varied from
02 ~5.3-5.7 m above present (in far-field regions such as Australia and South Africa) to as much as ~9-11 m above present (on
03 coastlines and islands of North America or around it). Further efforts along these lines by Dendy et al. (2017) confirm these
04 differences and provide additional insights on how ice sheet configuration during the penultimate glaciation (MIS 6)
05 influenced sea levels during MIS 5e. A combined field and modeling study on San Nicolas Island, California showed that
06 simulations of GIA processes over the period since MIS 5e yielded a sea-level history that matched the elevation spacing of
07 marine terraces dating to MIS 5c and 5a (Muhs et al., 2012). Later modeling by Creveling et al. (2017) refined paleo-sea
08 level estimates for both of these time periods and extended the concept of differing sea level histories to much of the globe.
09 Simms et al. (2016, 2020) conducted GIA modeling specifically on the Pacific Coast of North America and also confirmed
10 MIS 5c and 5a paleo-sea levels higher than what would be predicted by the Barbados-New Guinea terrace records.

11 GIA processes may also help explain what in the past had been an enigmatic observation about MIS 5e marine terrace
12 faunas and controversy about their origins. It was noted above that there is no persuasive field evidence of more than one
13 high-sea stand during MIS 5e on the Pacific Coast of North America. What *has* been documented, however, is evidence that
14 in areas of low uplift rate, marine terraces that formed during MIS 5e were reoccupied by the high-sea stand that followed it,
15 MIS 5c (~100 ka). The evidence of this sequence of events has actually been in existence for more than a century, with the
16 recognition of "thermally anomalous" faunas, i.e., those fossil faunas with *both* extralimital northern and extralimital southern
17 species of mollusks within the same deposit. Many hypotheses have been proposed to account for this (see review in Muhs
18 and Groves, 2018), but TIMS U-series dating finally demonstrated that corals of both MIS 5e and MIS 5c age exist in the same
19 marine terrace deposits at localities in central and southern California (see data in Table S1). At Cayucos (central California)
20 and Point Loma (southern California), both ~120 ka and ~100 ka corals exist within the same terrace deposits and terrace
21 deposits at both localities also host a mix of warm-water (~120 ka?) and cool-water (~100 ka?) mollusks (Muhs et al., 2002a).
22 On San Nicolas Island, the same mix of ~120 ka and ~100 ka corals and warm and cool mollusks is present in what Muhs et
23 al. (2012) called terrace 2b. Remnants of a slightly higher elevation terrace ("2a") have only ~120 ka corals and no cool-water
24 mollusks. Where corals are lacking from other Channel Islands marine terrace deposits, amino acid data show the likelihood
25 of two ages of shells, along with a mix of warm-water and cool-water mollusks (see Muhs and Groves, 2018, for examples).
26 Muhs et al. (2012) showed that the likely explanation for these observations is a low uplift rate combined with GIA processes
27 that resulted in a higher sea level during MIS 5c (Fig. 45).

28 Despite the general agreement between GIA models and field evidence from the Pacific Coast, there is an unresolved issue.
29 As noted, GIA modeling for MIS 5a and MIS 5c conducted by Muhs et al. (2012), Creveling et al. (2017) and Simms et al.
30 (2016) fit the elevation differences seen in terraces of these ages in California. A problem that remains, however, is the relative
31 elevation of the MIS 5e sea level on the Pacific Coast. Simms et al. (2016) modeled relative sea level at ~119 ka, including a
32 correction for the eustatic component of sea-level rise (taken to be 6 ± 3 m, relative to present), from Washington State to
33 southernmost Baja California Sur. Their results indicate a paleo-sea level as high as +13 m relative to present along much of
34 this coast from Washington to the southern Channel Islands, decreasing to +12 m in northern Baja California, and ultimately
35 decreasing to +10 m in southernmost Baja California Sur. Along with their modeled paleo-sea levels for MIS 5a and 5c, there
36 is a good match to the elevational spacing of a number of terrace sequences along the coast. In addition, there is good
37 agreement with the Simms et al. (2016) modeling for MIS 5e and similar modeling done for selected sites on the Pacific Coast
38 by Creveling et al. (2015). Despite these promising results, Muhs et al. (2021b) pointed out that there are several localities,
39 from central California to southernmost Baja California Sur, where elevations of MIS 5e terraces do not agree with the GIA
40 model results. These sites include Cayucos, Point San Luis, Santa Cruz Island, Anacapa Island, and Santa Barbara Island,
41 California, as well as Isla Guadalupe, Baja California. Older, higher elevation terraces at many of these localities preclude an
42 explanation of subsidence, as these higher terraces indicate a trend of steady, long-term uplift in the Quaternary. The reason
43 for the differences between the field data and these well conceptualized GIA models is not understood and needs more study.

44 It is interesting to note that GIA processes and their effect on relative sea levels may not be limited to MIS 5 paleo-sea
45 levels. Returning to San Nicolas Island, it was noted that there are 14 terraces on this island (Fig. 37). Vedder and Norris
46 (1963) reported faunal data from most of these terraces. Deposits of the 5th, 8th, and 10th terraces all contain mixes of both
47 warm-water and cool-water species of mollusks, suggesting a similar sequence of events as that described above for MIS 5e
48 and MIS 5c. Higher, older terraces elsewhere in California have not yet been investigated for this same kind of record but
49 would be a worthwhile effort.

50 A controversy that exists for MIS 5e along the Pacific Coast of North America is the amount of sea surface warming during
51 the last interglacial period derived from the fossil record compared to that from climate modeling. As noted earlier, faunal
52 evidence from a variety of marine terrace localities, from southern Baja California Sur to north of San Francisco Bay, indicates
53 substantial warming during MIS 5e, relative to present. Northward migration of what are now subtropical or tropical species
54 into mid-latitudes is documented at several localities (Figs. 13, 40, 41, 42, 44). These observations from the marine terrace
55 record are mirrored in the foraminiferal and alkenone records found in deep-sea cores off the Pacific Coast of North America
56 (Kennett and Venz, 1995; Herbert et al., 2001; Yamamoto et al., 2007), as summarized earlier. Modeling of SST (as well as
57 land surface temperatures) by Otto-Bliesner et al. (2013) indicates, however, that there was little annual surface temperature
58 change during MIS 5e, compared to pre-industrial modern time, on a global basis. In addition, for the Pacific Coast of North
59 America specifically, their model results mirror that of the global simulation, i.e., very little difference in MIS 5e time
60 compared to pre-industrial modern time. Otto-Bliesner et al. (2013) noted that their modeling was not able to reproduce many
61 proxy paleoclimate records that indicate greater warmth during MIS 5e. The reason for the disagreement between the model
62 results and the marine terrace faunal records (as well as those from deep-sea cores) is not understood at present and needs
63 more investigation.

64 **7 Future research directions**

65 In examining the work done to date on the Pacific Coast of North America, several topics that could merit additional work
66 have been mentioned. However, some specific needs that would be particularly useful are described here. Terrace mapping
67 can certainly be improved in many areas that have not received much attention. Particularly needed are good maps of marine
68 terraces in certain parts of northern California, central California, Baja California, and Sonora. More U-series ages on corals
69 are needed, particularly in Oregon, northern California, and northern Baja California. Preliminary field studies that I have
70 conducted indicate that the solitary coral *Balanophyllia elegans* is present in marine terrace deposits in some parts of northern
71 California where geochronology has yet to be conducted. Similarly, around the Golfo de California, colonial corals are likely
72 present in many marine terrace deposits that have not yet been studied, both on the Baja California peninsula, and on the coast
73 of mainland Mexico. More characterization of fossil faunas is needed. Very little work has yet been done on the coast of the
74 Golfo de California but could be carried out in concert with new U-series dating of colonial corals. Continued refinement of
75 GIA models and testing of those models is needed, particularly in view of the growing appreciation that paleo-sea levels around
76 the world during MIS 5e, 5c, and 5a are going to differ from coast to coast. Higher relative sea levels during MIS 5c and 5a,
77 with faunal mixing, is confirmed at only three localities at present in coastal California, but should be investigated at other

78 localities, particularly to see if there is a GIA gradient, which some models suggest should have existed (Simms et al., 2016).
79 Examination of fossil faunas and dating, perhaps with Sr isotopes, of pre-MIS 5e interglacials is an endeavor that could be
80 very usefully pursued. Records of older interglacials are found on the Palos Verdes Hills (Woodring et al., 1946), in the
81 Newport Bay area (Vedder et al., 1957, and on San Nicolas Island (Vedder and Norris, 1963). Such studies could test the
82 degree to which GIA effects were active during the middle and early Quaternary. Development of new dating methods for
83 marine terrace deposits that lack corals or even mollusks is encouraged. Cosmogenic and luminescence methods have promise,
84 but need to be investigated more thoroughly, particularly in those areas where there is independent geochronologic control
85 using U-series methods on corals. Finally, it would be useful to continue exploration of why climate modelling has shown
86 very little evidence for warming of the eastern Pacific Ocean off North America during MIS 5e, whereas the fossil records
87 (terrace faunas, foraminifera, alkenones) all point to substantial warming during this period. Because of the expectation of
88 future warming of the eastern Pacific Ocean, understanding the cause of different reconstructions from models and geologic
89 records is a worthy goal.

90 **8 Data availability**

91 Data from this study are open access and available at the following link: <https://doi.org/10.5281/zenodo.5557355>. Data
92 were exported from the WALIS database on 14 April 2021 and database descriptions can be found at the following link:
93 <https://doi.org/10.5281/zenodo.3961543> [Rovere et al., 2020]. Further information about the database can be examined here:
94 <https://warmcoasts.eu/world-atlas.html> (last access: 14 April 2021).

95 **9 Author contribution**

96 The manuscript was written by D.R. Muhs, who also drew all figures and compiled Tables S1, S2, and S3. D.R. Muhs, A.
97 Rovere, D. Ryan, and J.F. Wehmiller all contributed to the WALIS database for this review.

98 **10 Competing interests**

99 There are no competing interests of which the author is aware.

00 **11 Acknowledgments**

01 I thank the Climate Research and Development Program of the U.S. Geological Survey for supporting my research on
02 sea level history through the "Geologic Records of High Sea Levels" project. Much appreciation goes to Alessio Rovere,
03 who kindly invited me to contribute this review and provided an excellent review and much editorial guidance. Many thanks
04 go to Deidre Ryan, who ably assisted with the details of compiling amino acid data. A big thank-you goes to John
05 Wehmiller (University of Delaware), who helped me sort through amino acid data from nearly 40 years ago. Thanks go to
06 Markes Johnson (Williams College) and Lauren Toth (U.S. Geological Survey) for contributing beautiful photos of fossil
07 and modern corals, respectively. Laura Brothers and Janet Slate (U.S. Geological Survey), Jessica (JC) Creveling (Oregon
08 State University), John Wehmiller, Hayley C. Cawthra (Council for Geoscience Western Cape office and Nelson Mandela

09 University), and Barbara Mauz (University of Liverpool and University of Salzburg) read an earlier version of the paper and
10 made many helpful comments for its improvement, which I appreciate very much.

11 **References**

- 12 Abbott, R.T.: *American Seashells: The Marine Mollusca of the Atlantic and Pacific Coasts of North America* (2nd ed.), Van
13 Nostrand Reinhold Company, New York, 663 pp., 1974.
- 14 Addicott, W.O.: Interpretation of the invertebrate fauna from the upper Pleistocene Battery Formation near Crescent City,
15 California, *California Academy of Science Proceedings (Series 4)*, 31, 341-347, 1963.
- 16 Addicott, W.O.: A late Pleistocene invertebrate fauna from southwestern Oregon, *Journal of Paleontology*, 38, 650-661, 1964a.
- 17 Addicott, W.O.: Pleistocene invertebrates from the Dume terrace, western Santa Monica Mountains, California, *Bulletin of*
18 *the Southern California Academy of Sciences*, 63, 141-150, 1964b.
- 19 Addicott, W.O.: Late Pleistocene marine paleoecology and zoogeography in central California, U.S. Geological Survey
20 Professional Paper, 523-C, C1-C21, 1966.
- 21 Addicott, W.O., and Emerson, W.K.: Late Pleistocene invertebrates from Punta Cabras, Baja California, Mexico, *American*
22 *Museum Novitates*, 1925, 1-33, 1959.
- 23 Alexander, C.S.: The marine and stream terraces of the Capitola-Watsonville area, *University of California Publications in*
24 *Geography*, 10, 1-44, 1953.
- 25 Alley, N.F., and Hicock, S.R.: The stratigraphy, palynology, and climatic significance of pre-middle Wisconsin Pleistocene
26 sediments, southern Vancouver Island, British Columbia, *Canadian Journal of Earth Sciences*, 23, 369-382, 1986.
- 27 Alvarado, J.J., Reyes-Bonilla, H., Álvarez del Castillo Cardenas, P.A., Buitrago, F., and Aguirre-Rubí, J.: Coral reefs of the
28 Pacific coast of Nicaragua, *Coral Reefs*, 29, 201, 2010.
- 29 Anderson, R.S., and Anderson, S.P.: *Geomorphology: The Mechanics and Chemistry of Landscapes*, Cambridge University
30 Press, Cambridge, 637 pp., 2010.
- 31 Anderson, R.S., and Menking, K.M.: The Quaternary marine terraces of Santa Cruz, California: Evidence for coseismic
32 uplift on two faults, *Geological Society of America Bulletin*, 106, 649-664, 1994.
- 33 Arrhenius, G.: Sediment cores from the east Pacific, *Swedish Deep-Sea Expedition 1947-1948, Reports* 5, fasc. 1, 227 pp.,
34 1952.
- 35 Ashby, J.R., Ku, T.L., and Minch, J.A.: Uranium-series ages of corals from the upper Pleistocene Mulege terrace, Baja
36 California Sur, Mexico, *Geology*, 15, 139-141, 1987.
- 37 Barlow, N.L.M., McClymont, E.L., Whitehouse, P.L., Stokes, C.R., Jamieson, S.S.R., Woodroffe, S.A., Bentley, M.J., Callard,
38 S.L., Ó Cofaigh, C., Evans, D.J.A., Horrocks, J.R., Lloyd, J.M., Long, A.J., Margold, M., Roberts, D.H., and Sanchez-

- 39 Montes, M.L.: Lack of evidence for a substantial sea-level fluctuation within the Last Interglacial, *Nature Geoscience*,
40 11, 627-634, 2018.
- 41 Berger, G.W., and Easterbrook, D.J.: Thermoluminescence dating tests for lacustrine, glaciomarine, and floodplain
42 sediments from western Washington and British Columbia, *Canadian Journal of Earth Sciences*, 30, 1815-1828,
43 1993.
- 44 Berger, G.W., Burke, R.M., Carver, G.A., and Easterbrook, D. J.: Test of thermoluminescence dating with coastal sediments
45 from northern California, *Chemical Geology*, 87, 21-37, 1991.
- 46 Berschauer, D.P., and Clark, R.N.: *Sea Shells of Southern California: Marine Shells of the Californian Province*, San Diego
47 Shell Club, Encinitas, California, 131 pp., 2018.
- 48 Bertsch, H., and Aguilar Rosas, L.E.: *Marine Invertebrates of Northwest Mexico*, Instituto de Investigaciones
49 Oceanológicas, UABC, Ensenada, Mexico, 432 pp., 2016.
- 50 Birkeland, P.W.: Late Quaternary eustatic sea-level changes along the Malibu coast, Los Angeles County, California,
51 *Journal of Geology*, 80, 432-448, 1972.
- 52 Bloom, A.L.: Sea level and coastal morphology of the United States through the late Wisconsinan glacial maximum, in: *Late*
53 *Quaternary Environments of the United States, Volume 1, The Late Pleistocene*, edited by: Porter, S.C., University
54 of Minnesota Press, Minneapolis, Minnesota, pp. 215-229, 1983.
- 55 Bloom, A.L., Broecker, W.S., Chappell, J.M.A., Matthews, R.K., and Mesolella, K.J.: Quaternary sea level fluctuations on a
56 tectonic coast: New $^{230}\text{Th}/^{234}\text{U}$ dates from the Huon Peninsula, New Guinea, *Quaternary Research*, 4, 185-205, 1974.
- 57 Bockheim, J.G., Kelsey, H.M., and Marshall, J.G., III: Soil development, relative dating, and correlation of late Quaternary
58 marine terraces in southwestern Oregon, *Quaternary Research*, 37, 60-74, 1992.
- 59 Blunt, D.J.: *Geochemistry of amino acids in mollusks and wood, Pacific Northwest, United States*, M.S. thesis, California
60 State University, Hayward, California, 1982.
- 61 Blunt, D.J., Easterbrook, D.J., and Rutter, N.W.: Chronology of Pleistocene sediments in the Puget Lowland, Washington,
62 *Washington Division of Geology and Earth Resources Bulletin*, 77, 321-353, 1987.
- 63 Bradley, W.C., and Addicott, W.O.: Age of first marine terrace near Santa Cruz, California, *Geological Society of America*
64 *Bulletin*, 79, 1203-1210, 1968.
- 65 Bradley, W.C., and Griggs, G.B.: Form, genesis, and deformation of central California wave-cut platforms, *Geological Society*
66 *of America Bulletin*, 87, 433-449, 1976.
- 67 Brigham-Grette, J., and Hopkins, D.M.: Emergent marine record and paleoclimate of the last interglaciation along the
68 northwest Alaskan coast, *Quaternary Research*, 43, 159–173, 1995.
- 69 Broecker, W.S., Thurber, D.L., Goddard, J., Ku, T.-L., Matthews, R.K., and Mesolella, K.J.: Milankovitch hypothesis
70 supported by precise dating of coral reefs and deep-sea sediments, *Science*, 159, 297-300, 1968.
- 71 Bucknam, R.C., Hemphill-Haley, E., and Leopold, E.B.: Abrupt uplift within the past 1700 years at southern Puget Sound,
72 *Washington, Science*, 258, 1611-1614, 1992.

- 73 Bull, W.B.: Correlation of flights of global marine terraces, in: Proceedings of the 15th Annual Geomorphology Symposium,
74 edited by: Morisawa, M., and Hack, J.T., Allen and Unwin, Boston, 129-152, 1985.
- 75 Bull, W.B., and Cooper, A.F.: Uplifted marine terraces along the Alpine Fault, New Zealand, *Science*, 234, 1225-1228, 1986.
- 76 Burgess, C.M.: *Cowries of the World*, Gordon Verhoef Seacomber Publications, Cape Town, South Africa, 289 pp, 1985.
- 77 Calanchi, N., Lucchi, F., Pirazzoli, P.A., Romagnoli, C., Tranne, C.A., Radtke, U., Reyss, J.L., and Rossi, P.L.: Late
78 Quaternary relative sea-level changes and vertical movements at Lipari (Aeolian Islands), *Journal of Quaternary*
79 *Science*, 17, 459-467, 2002.
- 80 Carver, G.A.: Late Cenozoic tectonics of coastal northern California, in: *Field Guide to the Late Cenozoic Subduction*
81 *Tectonics & Sedimentation of North Coastal California*, Pacific Section, American Association of Petroleum
82 Geologists, pp. 1-9, 1992.
- 83 Carver, G.A., Jayko, A.S., Valentine, D.W., and Li, W.H.: Coastal uplift associated with the 1992 Cape Mendocino earthquake,
84 northern California, *Geology*, 22, 195-198, 1994.
- 85 Castillo, C.M., Klemperer, S.L., Ingle, J.C., Jr., Powell, C.L., II, Legg, M.L., and Francis, R.D.: Late Quaternary subsidence
86 of Santa Catalina Island, California Continental Borderland, demonstrated by seismic-reflection data and fossil
87 assemblages from submerged marine terraces, *Geological Society of America Bulletin*, 131, 21-42, 2018.
- 88 Chappell, J.: Geology of coral terraces, Huon Peninsula, New Guinea: A study of Quaternary tectonic movements and sea-
89 level changes, *Geological Society of America Bulletin*, 85, 553-570, 1974.
- 90 Chappell, J., and Shackleton, N.J.: Oxygen isotopes and sea level, *Nature*, 324, 137-140, 1986.
- 91 Chappell, J., Omura, A., Esat, T., McCulloch, M., Pandolfi, J., Ota, Y., and Pillans, B.: Reconciliation of late Quaternary sea
92 levels derived from coral terraces at Huon Peninsula with deep sea oxygen isotope records, *Earth and Planetary*
93 *Science Letters*, 141, 227-236, 1996.
- 94 Cheng, H., Edwards, R.L., Shen, C.-C., Polyak, V.J., Asmerom, Y., Woodhead, J., Hellstrom, J., Wang, Y., Kong, X., Spötl,
95 C., Wang, X., and Alexander, E.C., Jr.: Improvements in ^{230}Th dating, ^{230}Th and ^{234}U half-life values, and U-Th
96 isotopic measurements by multi-collector inductively coupled plasma mass spectrometry, *Earth and Planetary*
97 *Science Letters*, 371-372, 82-91, 2013.
- 98 Chutcharavan, P.M., and Dutton, A.: A global compilation of U-series dated fossil coral sea-level indicators for the last
99 interglacial period (MIS 5e), *Earth System Science Data Discussions*, 1–41, <https://doi.org/10.5194/essd-2020-381>,
00 2021.

- 01 Clague, J.J., Easterbrook, D.J., Hughes, O.L., and Matthews, J.V., Jr.: The Sangamonian and early Wisconsinan stages in
02 western Canada and northwestern United States, Geological Society of America Special Paper, 270, 253-268, 1992.
- 03 Clarke, S.J., and Murray-Wallace, C.V.: Mathematical expressions used in amino acid racemisation geochronology—A review,
04 Quaternary Geochronology, 1, 261-278, 2006.
- 05 CLIMAP Project Members: The last interglacial ocean, Quaternary Research, 21, 123-224, 1984.
- 06 Coan, E.V., and Valentich-Scott, P.: Bivalve seashells of tropical west America, Santa Barbara Museum of Natural History,
07 Santa Barbara, 1258 pp., 2012.
- 08 Coan, E.V., Valentich-Scott, P., and Bernard, F.R.: Bivalve seashells of western North America: marine bivalve mollusks from
09 Arctic Alaska to Baja California, Santa Barbara Museum of Natural History, Santa Barbara, 764 pp., 2000.
- 10 Cockerell, T.D.A.: Pleistocene shells from San Clemente Island, California, The Nautilus 53, 22-23, 1939.
- 11 Colman, S.M., Mixon, R.B., Rubin, M., Bloom, A.L., and Johnson, G.H.: Comments and reply on "Late Pleistocene barrier-
12 island sequence along the southern Delmarva Peninsula: Implications for middle Wisconsin sea levels," Geology, 17,
13 84-88, 1989.
- 14 Creveling, J.R., Mitrovica, J.X., Hay, C.C., Austermann, J., and Kopp, R.E.: Revisiting tectonic corrections applied to
15 Pleistocene sea-level highstands, Quaternary Science Reviews, 111, 72-80, 2015.
- 16 Creveling, J.R., Mitrovica, J.X., Clark, P.U., Waelbroeck, C., and Pico, T.: Predicted bounds on peak global mean sea level
17 during marine isotope stages 5a and 5c, Quaternary Science Reviews, 163, 193-208, 2017.
- 18 Cronin, T.M., Szabo, B.J., Ager, T.A., Hazel, J.E., and Owens, J.P.: Quaternary climates and sea levels of the U.S. Atlantic
19 Coastal Plain, Science, 211, 233-240, 1981.
- 20 Cuerda, J.: Los tiempos Cuaternarios en Baleares, Inst. de Estudios Baleáricos de la Diputación Provincial de Baleares, Palma,
21 304 pp., 1975.
- 22 Cuerda, J.: Moluscos marinos y salobres del Pleistoceno Balear, La Caja de Baleares, "Sa Nostra," Palma, 421 pp., 1987.
- 23 Cuerda, J.: Los tiempos Cuaternarios en Baleares (2nd ed.), Dir. Gral. Cultura. Conselleria de Cultura, Educació i Esports,
24 Govern Balear, Mallorca, Spain, 356 pp., 1989.
- 25 Cuerda, J., and Sacarès, J.: El Quaternari al Migjorn de Mallorca, Direcció General de Cultura, Conselleria de Cultura.
26 Educació i Esports, Govern Balear Spain, 130 pp, 1992.
- 27 Cutler, K.B., Edwards, R.L., Taylor, F.W., Cheng, H., Adkins, A., Gallup, C.D., Cutler, P.M., Burr, G.S., and Bloom, A.L.:
28 Rapid sea-level fall and deep-ocean temperature change since the last interglacial period, Earth and Planetary
29 Science Letters, 206, 253-271, 2003.
- 30 Davis, W.M.: Glacial epochs of the Santa Monica Mountains, California, Geological Society of America Bulletin, 44, 1041-
31 1133, 1933.

- 32 DeDiego-Forbis, T., Douglas, R., Gorsline, D., Nava-Sanchez, E., Mack, L., and Banner, J.: Late Pleistocene (Last
33 Interglacial) terrace deposits, Bahia Coyote, Baja California Sur, México, *Quaternary International*, 120, 29-40,
34 2004.
- 35 Delanghe, D., Bard, E., and Hamelin, B.: New TIMS constraints on the uranium-238 and uranium-234 in seawaters from the
36 main ocean basins and the Mediterranean Sea, *Marine Chemistry*, 80, 79-93, 2002.
- 37 Delattre, M., and Rosinski, A.: Preliminary geologic map of portions of the Crescent City and Orick quadrangles, California,
38 California Geological Survey, scale 1:100,000, 2012.
- 39 Dendy, S., Austermann, J., Creveling, J.R., and Mitrovica, J.X.: Sensitivity of Last Interglacial sea-level high stands to ice
40 sheet configuration during Marine Isotope Stage 6, *Quaternary Science Reviews*, 171, 234-244, 2017.
- 41 Dethier, D.P., Dragovich, J.D., Sarna-Wojcicki, A.M., and Fleck, R.J.: Pumice in the interglacial Whidbey Formation at
42 Blowers Bluff, central Whidbey Island, WA, USA, *Quaternary International*, 178, 229-237, 2008.
- 43 Dodge, R.E., Fairbanks, R.G., Benninger, L.K., and Maurrasse, F.: Pleistocene sea levels from raised coral reefs of Haiti,
44 *Science*, 219, 1423-1425, 1983.
- 45 Durham, J.W.: A new fossil *Pocillopora* (coral) from Guadalupe Island, Mexico, in: *The California Islands: Proceedings of a*
46 *Multidisciplinary Symposium*, edited by: Power, D.M., Santa Barbara, Santa Barbara Museum of Natural History,
47 pp. 63-70, 1980.
- 48 Dutton, A., and Lambeck, K.: Ice volume and sea level during the last interglacial, *Science*, 337, 216-219, 2012.
- 49 Easterbrook, D.J.: Pleistocene stratigraphy of Island County, State of Washington, Department of Water Resources, *Water*
50 *Supply Bulletin* 25, Part I, 1-34, 1968.
- 51 Easterbrook, D.J.: Pleistocene chronology of the Puget Lowland and San Juan Islands, Washington, *Geological Society of*
52 *America Bulletin*, 80, 2273-2286, 1969.
- 53 Easterbrook, D.J., Crandell, D.R., and Leopold, E.B.: Pre-Olympia Pleistocene stratigraphy and chronology in the central
54 Puget Lowland, Washington, *Geological Society of America Bulletin*, 78, 13-20, 1967.
- 55 Edwards, R.L., Cheng, H., Murrell, M.T., and Goldstein, S.J.: Protactinium-231 dating of carbonates by thermal ionization
56 mass spectrometry: implications for Quaternary climate change, *Science* 276, 782-786, 1997.
- 57 Emerson, W.K.: Pleistocene invertebrates from Punta China, Baja California, Mexico, *Bulletin of the American Museum of*
58 *Natural History*, 111, 313-342, 1956.
- 59 Emerson, W.K.: Pleistocene invertebrates from near Punta San José, Baja California, Mexico, *American Museum Novitates*,
60 2002, 1-7, 1960.
- 61 Emerson, W.K.: Invertebrate faunules of late Pleistocene age, with zoogeographic implications, from Turtle Bay, Baja
62 California Sur, Mexico, *The Nautilus*, 94, 67-89, 1980.

- 63 Emerson, W.K., and Addicott, W.O.: A Pleistocene invertebrate fauna from the southwest corner of San Diego County,
64 California, Transactions of the San Diego Society of Natural History, 11, 429-444, 1953.
- 65 Emerson, W.K., and Addicott, W.O.: Pleistocene invertebrates from Punta Baja, Baja California, Mexico, American Museum
66 Novitates, 1909, 1-11, 1958.
- 67 Emerson, W.K., Kennedy, G.L., Wehmiller, J.F., and Keenan, E.: Age relations and zoogeographic implications of late
68 Pleistocene marine invertebrate faunas from Turtle Bay, Baja California Sur, Mexico, The Nautilus, 95, 105-116,
69 1981.
- 70 Emery, K.O.: Shallow submerged marine terraces of southern California, Geological Society of America Bulletin, 69, 39-60,
71 1958.
- 72 Emiliani, C.: Pleistocene temperatures, Journal of Geology, 63, 538-578, 1955.
- 73 Fisher, D.M., Gardner, T.W., Marshall, J.S., Sak, P.B., and Protti, M.: Effect of subducting sea-floor roughness on fore-arc
74 kinematics, Pacific coast, Costa Rica, Geology, 26, 467-470, 1998.
- 75 Foose, R.M.: Reconnaissance geology of Maria Cleopha Island, Tres Marias Islands, Mexico, Bulletin of the American
76 Association of Petroleum Geologists, 46, 1740-1745, 1962.
- 77 Gallup, C.D., Edwards, R.L., and Johnson, R.G.: The timing of high sea levels over the past 200,000 years, Science, 263,
78 796-800, 1994.
- 79 Gallup, C.D., Cheng, H., Taylor, F.W., and Edwards, R.L.: Direct determination of the timing of sea level change during
80 Termination II, Science, 295, 310-313, 2002.
- 81 Gardner, T., Marshall, J., Merritts, D., Bee, B., Burgette, R., Burton, E., Cooke, J., Kehrwald, N., Protti, M., Fisher, D., and
82 Sak, P.: Holocene forearc block rotation in response to seamount subduction, southeastern Península de Nicoya,
83 Costa Rica, Geology, 29, 151-154, 2001.
- 84 Gardner, T.W., Verdonck, D., Pinter, N.M., Slingerland, R., Furlong, K.P., Bullard, T.F., and Wells, S.G.: Quaternary uplift
85 astride the aseismic Cocos Ridge, Pacific coast, Costa Rica, Geological Society of America Bulletin, 104, 219-232,
86 1992.
- 87 Gardner, T.W., Fisher, D.M., Morrell, K.D., and Cupper, M.L.: Upper-plate deformation in response to flat slab subduction
88 inboard of the aseismic Cocos Ridge, Osa Peninsula, Costa Rica, Lithosphere, 5, 247-264, 2013.
- 89 Gerrodette, T.: Equatorial submergence in a solitary coral, *Balanophyllia elegans*, and the critical life stage excluding the
90 species from shallow water in the south, Marine Ecology-Progress Series 1, 227-235, 1979.
- 91 Glynn, P.W., and Ault, J.S.: A biogeographic analysis and review of the far eastern Pacific coral reef region, Coral Reefs,
92 19, 1-23, 2000.
- 93 Glynn, P.W., Alvarado, J.J., Banks, S., Cortés, J., Feingold, J.S., Jiménez, C., Maragos, J.E., Martínez, P., Maté, J.L.,
94 Moanga, D.A., Navarrete, S., Reyes-Bonilla, H., Riegl, B., Rivera, F., Vargas-Ángel, B., Wieters, E.A., and Zapata,
95 F.A.: Eastern Pacific coral reef provinces, coral community structure and composition: An overview, in: Coral

96 Reefs of the Eastern Tropical Pacific, edited by: Glynn, P.W., Manzello, D.P., and Enochs, I.C., Springer,
97 Dordrecht, Netherlands, pp. 107-176, 2017.

98 Gonzalez-Garcia, J.J., Prawirodirdjo, L., Bock, Y., and Agnew, D.: Guadalupe Island, Mexico as a new constraint for Pacific
99 plate motion, *Geophysical Research Letters*, 30 (16), 1872, doi:10.1029/2003GL017732, 2003.

100 Grant, L. B., Mueller, K. J., Gath, E. M., Cheng, H., Edwards, R. L., Munro, R., and Kennedy, G. L.: Quaternary uplift and
101 earthquake potential of the San Joaquin Hills, southern Los Angeles Basin, California, *Geology*, 27, 1031-1034, 1999.

102 Grant, U.S., IV, and Gale, H.R.: Catalogue of the marine Pliocene and Pleistocene of California and adjacent regions,
103 *Memoirs of the San Diego Society of Natural History*, 1, 1-1036, 1931.

104 Graymer, R.W., Moring, B.C., Saucedo, G.J., Wentworth, C.M., Brabb, E.E., and Knudsen, K.L.: Geologic map of the San
105 Francisco Bay region, U.S. Geological Survey Scientific Investigations Map, 2918, scale 1:275,000, 2006.

106 Griggs, A.B.: Chromite-bearing sands of the southern part of the coast of Oregon, U.S. Geological Survey Bulletin, 945E,
107 113-150, 1945.

108 Grove, K., Colson, K., Binkin, M., Dull, R., and Garrison, C.: Stratigraphy and structure of the late Pleistocene Olema Creek
109 Formation, San Andreas fault zone north of San Francisco, California, in: *Recent Geologic Studies in the San*
110 *Francisco Bay Area*, edited by: Sanginés, E.M., Andersen, D.W., and Busing, A.V., Pacific Section of the Society
111 of Economic Paleontologists and Mineralogists, 76, 55-76, 1995.

112 Groves, L.T.: Fossil marine molluscs of the Hawaiian Islands, in: *Shells of the Hawaiian Islands—The Sea Shells*, edited by:
113 Severns, M., ConchBooks, Hackenheim, Germany, pp. 536-546, 2011.

114 Gurrola, L.D., Keller, E.A., Chen, J.H., Owen, L.A., and Spencer, J.Q.: Tectonic geomorphology of marine terraces: Santa
115 Barbara fold belt, California, *Geological Society of America Bulletin*, 126, 219-233, 2014.

116 Hansen, H.P., and Mackin, J.H.: A pre-Wisconsin forest succession in the Puget Lowland, Washington, *American Journal of*
117 *Science*, 247, 833-855, 1949.

118 Hanson, K.L., Wesling, J.R., Lettis, W.R., Kelson, K.I., and Mezger, L.: Correlation, ages, and uplift rates of Quaternary
119 marine terraces: South-central coastal California, *Geological Society of America Special Paper*, 292, 45-71, 1994.

120 Harmon, R.S., Mitterer, R.M., Kriausakul, N., Land, L.S., Schwarcz, H.P., Garrett, P., Larson, G.J., Vacher, H.L., and Rowe,
121 M.: U-series and amino-acid racemization geochronology of Bermuda: Implications for eustatic sea-level fluctuation
122 over the past 250,000 years, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 44, 41-70, 1983.

123 Hearty, P.J., Miller, G.H., Stearns, C.E., and Szabo, B.J.: Aminostratigraphy of Quaternary shorelines in the Mediterranean
124 basin, *Geological Society of America Bulletin*, 97, 850-858, 1986.

125 Herbert, T.D., Schuffert, J.D., Thomas, D., Lange, C., Weinheimer, A., Peleo-Alampay, A., and Herguera, J.-C.: Depth and
126 seasonality of alkenone production along the California margin inferred from a core top transect, *Paleoceanography*,
127 13, 263-271, 1998.

- 28 Herbert, T.D., Schuffert, J.D., Andreasen, D., Heusser, L., Lyle, M, Mix, A., Ravelo, A.C., Stott, L.D., and Herguera, J.C.:
29 Collapse of the California Current during glacial maxima linked to climate change on land, *Science*, 293, 71-76, 2001.
- 30 Hertlein, L.G., and Emerson, W.K.: Results of the Puritan-American Museum of Natural History expedition to western
31 Mexico. 5. Pliocene and Pleistocene megafossils from the Tres Mariás Islands, *American Museum Novitates*, 1940,
32 1-15, 1959.
- 33 Heusser, C.J., and Heusser, L.E.: Palynology and paleotemperature analysis of the Whidbey Formation, Puget Lowland,
34 Washington, *Canadian Journal of Earth Sciences*, 18, 136-149, 1981.
- 35 Hicock, S.R.: Last interglacial Muir Point Formation, Vancouver Island, British Columbia, *Géographie Physique et*
36 *Quaternaire*, 44, 337-340, 1990.
- 37 Hicock, S.R., and Armstrong, J.E.: Four Pleistocene formations in southwest British Columbia: their implications for patterns
38 of sedimentation of possible Sangamonian to early Wisconsin age, *Canadian Journal of Earth Sciences*, 20, 1232-
39 1247, 1983.
- 40 Howarth, R.J., and McArthur, J.M.: Statistics for strontium isotope stratigraphy: A robust LOWESS fit to the marine Sr-
41 isotope curve for 0 to 206 Ma, with look-up table for derivation of numeric age, *Journal of Geology*, 105, 441-456,
42 1997.
- 43 Jennings, C.W.: Fault activity map of California and adjacent areas with locations and ages of recent volcanic eruptions,
44 California Division of Mines and Geology, Geologic Data Map no. 6, scale 1:750 000, 1994.
- 45 Jennings, C.W., and Strand, R.G.: Geologic map of California, Ukiah sheet, State of California, Department of Natural
46 Resources, Division of Mines, scale 1:250,000, 1960.
- 47 Johnson, M.E.: *Discovering the Geology of Baja California: Six Hikes on the Southern Gulf Coast*, University of Arizona
48 Press, Tucson, Arizona, 220 pp., 2002.
- 49 Johnson, M.E., López-Pérez, R.A., Ranson, C.R., and Ledesma-Vázquez, J.: Late Pleistocene coral-reef development on Isla
50 Coronados, Gulf of California, *Ciencias Marinas*, 33, 105-120, 2007.
- 51 Johnson, R.G.: Mode of formation of marine fossil assemblages of the Pleistocene Millerton Formation of California,
52 *Geological Society of America Bulletin*, 73, 113-130, 1962.
- 53 Jordan, E.K.: The Pleistocene fauna of Magdalena Bay, Lower California, *Contributions from the Department of Geology of*
54 *Stanford University*, 1(4), 103-174, 1936.
- 55 Kanakoff, G.P., and Emerson, W.K.: Late Pleistocene invertebrates of the Newport Bay area, California, Los Angeles
56 County Museum Contributions in Science, 31, 1-47, 1959.
- 57 Karrow, P.F., Ceska, A., Hebda, R.J., Miller, B.B., Seymour, K.L., and Smith, A.J.: Diverse nonmarine biota from the
58 Whidbey Formation (Sangamonian) at Point Wilson, Washington, *Quaternary Research*, 44, 434-437, 1995.
- 59 Kaufman, A., Broecker, W.S., Ku, T.-L., and Thurber, D.L.: The status of U-series methods of mollusk dating, *Geochimica*
60 *et Cosmochimica Acta*, 35, 1155-1183, 1971.

- 61 Kaufman, D.S., and Manley, W.F.: A new procedure for determining DL amino acid ratios in fossils using reverse phase
62 liquid chromatography, *Quaternary Science Reviews*, 17, 987-1000, 1998.
- 63 Keen, A.M.: *Sea Shells of Tropical West America: Marine Mollusks from Baja California to Peru* (2nd edition), Stanford
64 University Press, 1064 pp., Stanford, 1971.
- 65 Keenan, E.M., Ortlieb, L., and Wehmiller, J.F.: Amino acid dating of Quaternary marine terraces, Bahia Ascunción, Baja
66 California Sur, Mexico, *Journal of Coastal Research*, 3, 297-305, 1987.
- 67 Kelsey, H.M.: Late Quaternary deformation of marine terraces on the Cascadia subduction zone near Cape Blanco, Oregon,
68 *Tectonics*, 9, 983-1014, 1990.
- 69 Kelsey, H.M.: Geomorphological indicators of past sea levels, in: *Handbook of Sea-Level Research*, edited by: Shennan, I.,
70 Long, A.J., and Horton, B.P., John Wiley & Sons, Ltd., pp. 66-82, 2015.
- 71 Kelsey, H.M., and Bockheim, J.G.: Coastal landscape evolution as a function of eustasy and surface uplift rate, Cascadia
72 margin, southern Oregon, *Geological Society of America Bulletin*, 106, 840-854, 1994.
- 73 Kelsey, H.M., Engebretson, D.C., Mitchell, C.E., and Ticknor, R.L.: Topographic form of the Coast Ranges of the Cascadia
74 Margin in relation to coastal uplift rates and plate subduction, *Journal of Geophysical Research*, 99, 12,245-12,255,
75 1994.
- 76 Kelsey, H.M., Ticknor, R.L., Bockheim, J.G., and Mitchell, C.E.: Quaternary upper plate deformation in coastal Oregon,
77 *Geological Society of America Bulletin*, 108, 843-860, 1996.
- 78 Kendrick, G.W., Wyrwoll, K.-H., and Szabo, B.J.: Pliocene-Pleistocene coastal events and history along the western margin
79 of Australia, *Quaternary Science Reviews*, 10, 419-439, 1991.
- 80 Kennedy, G.L.: Pleistocene paleoecology, zoogeography and geochronology of marine invertebrate faunas of the Pacific
81 Northwest Coast (San Francisco Bay to Puget Sound), Unpublished Ph.D. thesis, University of California-Davis,
82 824 pp., 1978.
- 83 Kennedy, G.L.: Zoogeographic correlation of marine invertebrate faunas, in: *Quaternary Geochronology*, edited by: Noller,
84 J.S., Sowers, J.M., and Lettis, W.R., American Geophysical Union Reference Shelf 4, 413-424, Washington, D.C.,
85 2000.
- 86 Kennedy, G.L., Lajoie, K.R., and Wehmiller, J.F.: Aminostratigraphy and faunal correlations of late Quaternary marine
87 terraces, Pacific Coast, USA, *Nature*, 299, 545-547, 1982.
- 88 Kennedy, G.L., Wehmiller, J.K., and Rockwell, T.K.: Paleoecology and paleozoogeography of late Pleistocene marine-terrace
89 faunas of southwestern Santa Barbara County, California, in: *Quaternary Coasts of the United States: Marine and*
90 *Lacustrine Systems*, edited by: Fletcher, C.H. III and Wehmiller, J.F., SEPM (Society for Sedimentary Geology)
91 Special Publication, no. 48, 343-361, 1992.
- 92 Kennett, J.P., and Venz, K.: Late Quaternary climatically related planktonic foraminiferal assemblage changes: Hole 893A,
93 Santa Barbara Basin, California, *Proceedings of the Ocean Drilling Program, Scientific Results 146 (Part 2)*, 281-
94 293, 1995.

- 95 Kern, J.P.: Paleoenvironmental analysis of a late Pleistocene estuary in southern California, *Journal of Paleontology*, 45, 810-
96 823, 1971.
- 97 Kern, J.P.: Origin and history of upper Pleistocene marine terraces, San Diego, California, *Geological Society of America*
98 *Bulletin*, 88, 1553-1566, 1977.
- 99 Kern, J.P., and Rockwell, T.K.: Chronology and deformation of Quaternary marine shorelines, San Diego County,
00 California, in: *Quaternary Coasts of the United States: Marine and Lacustrine Systems*, edited by: Fletcher, C.H. III
01 and Wehmiller, J.F., SEPM (Society for Sedimentary Geology) Special Publication, no. 48, 377-382, 1992.
- 02 Kopp, R.E., Simons, F.J., Mitrovica, J.X., Maloof, A.C., and Oppenheimer, M.: Probabilistic assessment of sea level during
03 the last interglacial stage, *Nature*, 462, 863-868, 2009.
- 04 Kosuge, S.: Fossil mollusks of Oahu, Hawaii Islands, *Bulletin of the National Science Museum [Tokyo, Japan]*, 12, 783-
05 794, 1969.
- 06 Ku, T.-L., and Kern, J.P.: Uranium-series age of the upper Pleistocene Nestor terrace, San Diego, California, *Geological*
07 *Society of America Bulletin*, 85, 1713-1716, 1974.
- 08 Kvenvolden, K.A., Blunt, D.J., and Clifton, H.E.: Amino-acid racemization in Quaternary shell deposits at Willapa Bay,
09 Washington, *Geochimica et Cosmochimica Acta*, 43, 1505-1520, 1979.
- 10 Kvenvolden, K.A., Blunt, D.J., McMenamin, M.A., and Straham, S.E.: Geochemistry of amino acids in shells of the clam
11 *Saxidomus*, *Physics and Chemistry of the Earth*, 12, 321-332, 1980.
- 12 Kvenvolden, K.A., Blunt, D.J., and Clifton, H.E.: Age estimations based on amino acid racemization: reply to comments of
13 J.F. Wehmiller, *Geochimica et Cosmochimica Acta*, 45, 265-267, 1981.
- 14 Lajoie, K.R.: Coastal tectonics, in: *Active Tectonics*, edited by: Wallace, R., National Academy Press, Washington, D.C., pp.
15 95-124, 1986.
- 16 Lajoie, K.R., Wehmiller, J.F., and Kennedy, G.L.: Inter- and intragenetic trends in apparent racemization kinetics of amino
17 acids in Quaternary mollusks, in: *Biogeochemistry of Amino Acids*, edited by: Hare, P.E., John Wiley and Sons,
18 pp. 305-340, 1980.
- 19 Lajoie, K.R., Ponti, D.J., Powell, C.L. II, Mathieson, S.A., and Sarna-Wojcicki, A.M.: Emergent marine strandlines and
20 associated sediments, coastal California; a record of Quaternary sea-level fluctuations, vertical tectonic movements,
21 climatic changes, and coastal processes, in: Morrison, R.B., ed., *Quaternary nonglacial geology; Conterminous U.S.*,
22 edited by: Morrison, R.B., Geological Society of America, Boulder, Colorado, *The Geology of North America*, v. K-
23 2, pp. 190-203, 1991.
- 24 Lawson, A.C.: The post-Pliocene diastrophism of the coast of southern California, *University of California Bulletin of the*
25 *Department of Geology*, 1, 115-160, 1893.
- 26 Lian, O.B., Hu, J., Huntley, D.J., and Hicock, S.R.: Optical dating studies of Quaternary organic-rich sediments from

- 27 southwestern British Columbia and northwestern Washington State, *Canadian Journal of Earth Sciences*, 32, 1194-
28 1207, 1995.
- 29 Libbey, L.K., and Johnson, M.E.: Upper Pleistocene rocky shores and intertidal biotas at Playa La Palmita (Baja California
30 Sur, Mexico), *Journal of Coastal Research*, 13, 216-225, 1997.
- 31 Lindberg, D.R., Roth, B., Kellogg, M.G., and Hubbs, C.L.: Invertebrate megafossils of Pleistocene (Sangamon interglacial)
32 age from Isla de Guadalupe, Baja California, Mexico, in: *The California Islands: Proceedings of a Multidisciplinary*
33 *Symposium*, edited by: Power, D.M., Santa Barbara Museum of Natural History, Santa Barbara, California, pp. 41-
34 62, 1980.
- 35 Lipps, J.H., Valentine, J.W., and Mitchell, E.: Pleistocene paleoecology and biostratigraphy, Santa Barbara Island, California,
36 *Journal of Paleontology*, 42, 291-307, 1968.
- 37 López Pérez, R.A.: Fossil corals from the Gulf of California, México: Still a depauperate fauna, but it bears more species than
38 previously thought, *Proceedings of the California Academy of Sciences*, 59, 503-519, 2008.
- 39 Ludwig, K.R., Muhs, D.R., Simmons, K.R., and Moore, J.G.: Sr-isotope record of Quaternary marine terraces on the California
40 coast and off Hawaii, *Quaternary Research*, 37, 267-280, 1992.
- 41 Mann, P.: Global catalogue, classification and tectonic origins of restraining- and releasing bends on active and ancient strike-
42 slip fault systems, *Geological Society of London Special Publications*, 290, 13-142, 2007.
- 43 Marshall, J.S., and Anderson, R.S.: Quaternary uplift and seismic cycle deformation, Península de Nicoya, Costa Rica,
44 *Geological Society of America Bulletin*, 107, 463-473, 1995.
- 45 Martin, R.E., Wehmiller, J.F., Harris, M.S., and Liddell, W.D.: Comparative taphonomy of bivalves and foraminifera from
46 Holocene tidal flat sediments, Bahía la Choya, Sonora, Mexico (Northern Gulf of California): taphonomic grades and
47 temporal resolution, *Paleobiology*, 22, 80-90, 1996.
- 48 Martinson, D.G., Pisias, N.G., Hays, J.D., Imbrie, J., Moore, T.C., Jr., and Shackleton, N.J.: Age dating and the orbital theory
49 of the ice ages: Development of a high-resolution 0 to 300,000-year chronostratigraphy, *Quaternary Research*, 27,
50 1-29, 1987.
- 51 Maxson, J.H.: Economic geology of portions of Del Norte and Siskiyou counties, northwesternmost California, California
52 *Journal of Mines and Geology*, 29, 123-160, 1933.
- 53 Mayer, L., and Vincent, K.R.: Active tectonics of the Loreto area, Baja California Sur, Mexico, *Geomorphology*, 27, 243-255,
54 1999.
- 55 McCrory, P.A.: Upper plate contraction north of the migrating Mendocino triple junction, northern California: Implications
56 for partitioning of strain, *Tectonics*, 19, 1144-1160, 2000.
- 57 McInelly, G.W., and Kelsey, H.M.: Late Quaternary tectonic deformation in the Cape Arago-Bandon region of coastal Oregon
58 as deduced from wave-cut platforms, *Journal of Geophysical Research*, 95, 6699-6713, 1990.

- 59 McKay, N.P., Overpeck, J.T., and Otto-Bliesner, B.L.: The role of ocean thermal expansion in Last Interglacial sea level rise,
60 *Geophysical Research Letters*, 38, L14605, doi:10.1029/2011GL048280, 2011.
- 61 McLaughlin, R.J., Lajoie, K.R., Sorg, D.H., Morrison, S.D., and Wolfe, J.A.: Tectonic uplift of a middle Wisconsin marine
62 platform near the Mendocino triple junction, California, *Geology*, 11, 35-39, 1983a.
- 63 McLaughlin, R.J., Lajoie, K.R., Sorg, D.H., Morrison, S.D., and Wolfe, J.A.: Comment and reply on 'Tectonic uplift of a
64 middle Wisconsin marine platform near the Mendocino triple junction, California,' *Geology*, 11, 621-622, 1983b.
- 65 McLaughlin, R.J., Ellen, S.D., Blake, M.C., Jr., Jayko, A.S., Irwin, W.P., Aalto, K.R., Carver, G.A., and Clarke, S.H., Jr.:
66 *Geology of the Cape Mendocino, Eureka, Garberville, and southwestern part of the Hayfork 30 x 60 minute*
67 *quadrangles and adjacent offshore area, northern California, U.S. Geological Survey Miscellaneous Field Studies,*
68 *MF-2336, scales 1:100,000 (2 sheets) and 1:137,000 (1 sheet), 2000.*
- 69 Meco, J., Guillou, H., Carracedo, J.C., Lomoschitz, A., Ramos, A.J.G., and Rodríguez Yáñez, J.J.: The maximum warmings
70 of the Pleistocene world climate recorded in the Canary Islands, *Palaeogeography, Palaeoclimatology, Palaeoecology*,
71 185, 197–210, 2002.
- 72 Meco, J., Ballester, J., Betancourt, J.F., Cilleros, A., Scaillet, S., Guillou, H., Carracedo, J.C., Lomoschitz, A., Petit-Maire, N.,
73 Ramos, A.J.G., Perera, N., and Meco, J.M.: *Paleoclimatología del Neógeno en las Islas Canarias: Geliense,*
74 *Pleistoceno y Holoceno, Ministerio de Medio Ambiente, Universidad de las Palmas de Gran Canaria, 203 pp., 2006.*
- 75 Merritts, D.J.: The Mendocino triple junction: Active faults, episodic coastal emergence, and rapid uplift, *Journal of*
76 *Geophysical Research*, 101, 6051-6070, 1996.
- 77 Merritts, D., and Bull, W.B.: Interpreting Quaternary uplift rates at the Mendocino triple junction, northern California, from
78 uplifted marine terraces, *Geology*, 17, 1020-1024, 1989.
- 79 Mesolella, K.J., Matthews, R.K., Broecker, W.S., and Thurber, D.L.: The astronomical theory of climatic change: Barbados
80 data, *Journal of Geology*, 77, 250-274, 1969.
- 81 Miller, G.H., and Mangerud, J.: Aminostratigraphy of European marine interglacial deposits, *Quaternary Science Reviews*, 4,
82 215-278, 1985.
- 83 Mitchell, D.L., Ivanova, D., Rabin, R., Brown, T.J., and Redmond, K.: Gulf of California sea surface temperatures and the
84 North American monsoon: Mechanistic implications for observations, *Journal of Climate*, 17, 2261-2281, 2002.
- 85 Mitterer, R.M., and Kriauskaul, N.: Calculation of amino acid racemization ages based on apparent parabolic kinetics,
86 *Quaternary Science Reviews*, 8, 353-357, 1989.
- 87 Morrell, K.D., Fisher, D.M., Gardner, T.W., La Femina, P., Davidson, D., and Teletzke, A.: Quaternary outer fore-arc
88 deformation and uplift inboard of the Panama Triple Junction, Burica Peninsula. *Journal of Geophysical Research*,
89 116, doi: 10.1029/2010JB007979, 2011.
- 90 Muhs, D.R.: Quaternary sea-level events on northern San Clemente Island, California, *Quaternary Research*, 20, 322-341,
91 1983.

- 92 Muhs, D.R.: Amino acid age estimates of marine terraces and sea levels, San Nicolas Island, California, *Geology*, 13, 58-61,
93 1985.
- 94 Muhs, D.R., and Groves, L.T.: Little islands recording global events: Late Quaternary sea-level history and paleozoogeography
95 of Santa Barbara and Anacapa Islands, Channel Islands National Park, California, *Western North American*
96 *Naturalist*, 78, 540-589, 2018.
- 97 Muhs, D.R., and Kyser, T.K.: Stable isotope compositions of fossil mollusks from southern California: Evidence for a cool
98 last interglacial ocean, *Geology*, 15, 119-122, 1987.
- 99 Muhs, D.R., and Simmons, K.R.: Taphonomic problems in reconstructing sea-level history from the late Quaternary marine
00 terraces of Barbados, *Quaternary Research*, 88, 409-429, 2017.
- 01 Muhs, D.R., Simmons, K.R., and Steinke, B.: Timing and warmth of the last interglacial period: New U-series evidence
02 from Hawaii and Bermuda and a new fossil compilation for North America, *Quaternary Science Reviews*, 21, 1355-
03 1383, 2002b.
- 04 Muhs, D.R., Kelsey, H.M., Miller, G.H., Kennedy, G.L., Whelan, J.F., and McInelly, G.W.: Age estimates and uplift rates for
05 late Pleistocene marine terraces: Southern Oregon portion of the Cascadia forearc, *Journal of Geophysical Research*,
06 95, 6685-6698, 1990.
- 07 Muhs, D.R., Miller, G.H., Whelan, J.F., and Kennedy, G.L.: Aminostratigraphy and oxygen isotope stratigraphy of marine-
08 terrace deposits, Palos Verdes Hills and San Pedro areas, Los Angeles County, California, in: *Quaternary Coasts of*
09 *the United States: Marine and Lacustrine Systems*, edited by: Fletcher, C.H. III and Wehmiller, J.F., SEPM (Society
10 for Sedimentary Geology) Special Publication, no. 48, 363-376, 1992.
- 11 Muhs, D. R., Kennedy, G. L., and Rockwell, T. K.: Uranium-series ages of marine terrace corals from the Pacific coast of
12 North America and implications for last-interglacial sea level history, *Quaternary Research*, 42, 72-87, 1994.
- 13 Muhs, D.R., Simmons, K.R., Kennedy, G.L., and Rockwell, T.K.: The last interglacial period on the Pacific Coast of North
14 America: Timing and paleoclimate, *Geological Society of America Bulletin*, 114, 569-592, 2002a.
- 15 Muhs, D.R., Prentice, C., and Merritts, D.J.: Marine terraces, sea level history and Quaternary tectonics of the San Andreas
16 fault on the coast of California, in: *Quaternary Geology of the United States*, INQUA 2003 Field Guide Volume,
17 edited by: Easterbrook, D., Desert Research Institute, Reno, Nevada, pp. 1-18, 2003.
- 18 Muhs, D.R., Wehmiller, J.F., Simmons, K.R., and York, L.L.: Quaternary sea level history of the United States, in: *The*
19 *Quaternary Period in the United States*, edited by: Gillespie, A.R., Porter, S.C., and Atwater, B.F., Elsevier,
20 Amsterdam, pp. 147-183, 2004.
- 21 Muhs, D.R., Simmons, K.R., Kennedy, G.L., Ludwig, K.R., and Groves, L.T.: A cool eastern Pacific Ocean at the close of
22 the last interglacial complex, *Quaternary Science Reviews*, 25, 235-262, 2006.
- 23 Muhs, D.R., Simmons, K.R., Schumann, R.R., Groves, L.T., Mitrovica, J.X., and Laurel, D.: Sea-level history during the last
24 interglacial complex on San Nicolas Island, California: Implications for glacial isostatic adjustment processes,
25 paleozoogeography and tectonics, *Quaternary Science Reviews*, 37, 1-25, 2012.

- 26 Muhs, D.R., Groves, L.T., and Schumann, R.R.: Interpreting the paleozoogeography and sea level history of thermally
27 anomalous marine terrace faunas: A case study from the last interglacial complex of San Clemente Island, California,
28 Monographs of the Western North American Naturalist, 7, 82-108, 2014a.
- 29 Muhs, D.R., Simmons, K.R., Schumann, R.R., Groves, L.T., DeVogel, S.B., Minor, S.A., and Laurel, D.: Coastal tectonics on
30 the eastern margin of the Pacific Rim: Late Quaternary sea-level history and uplift rates, Channel Islands National
31 Park, California, USA, Quaternary Science Reviews, 105, 209-238, 2014b.
- 32 Muhs, D.R., Meco, J., and Simmons, K.R.: Uranium-series ages of corals, sea level history, and palaeozoogeography, Canary
33 Islands, Spain: An exploratory study for two Quaternary interglacial periods, Palaeogeography, Palaeoclimatology,
34 Palaeoecology, 394, 99-118, 2014c.
- 35 Muhs, D.R., Simmons, K.R., Groves, L.T., McGeehin, J.P., Schumann, R.R., and Agenbroad, L.D.: Late Quaternary sea-
36 level history and the antiquity of mammoths (*Mammuthus exilis* and *Mammuthus columbi*), Channel Islands
37 National Park, California, USA, Quaternary Research, 83, 502-521, 2015.
- 38 Muhs, D.R., Pigati, J.S., Schumann, R.R., Skipp, G.L., Porat, N., and DeVogel, S.B.: Quaternary sea-level history and the
39 origin of the northernmost coastal aeolianites in the Americas: Channel Islands National Park, California, USA,
40 Palaeogeography, Palaeoclimatology, Palaeoecology, 491, 38-76, 2018.
- 41 Muhs, D.R., Wehmiller, J.F., Ryan, D.D., and Rovere, A.: MIS 5e relative sea-level index points along the Pacific coast of
42 North America (1.1) [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.5903285>, 2021a.
- 43 Muhs, D.R., Schumann, R.R., Groves, L.T., Simmons, K.R., and Florian, C.R.: The marine terraces of Santa Cruz Island,
44 California: Implications for glacial isostatic adjustment models of last-interglacial sea-level history, Geomorphology,
45 389, 107826, <https://doi.org/10.1016/j.geomorph.2021.107826>, 2021b.
- 46 Murray-Wallace, C.V., and Woodroffe, C.D.: Quaternary Sea Level Changes: A Global Perspective, Cambridge University
47 Press, Cambridge, 484 pp, 2014.
- 48 Murray-Wallace, C.V., Beu, A.G., Kendrick, G.W., Brown, L.J., Belperio, A.P., and Sherwood, J.E.: Palaeoclimatic
49 implications of the occurrence of the arcoid bivalve *Andadara trapezia* (Deshayes) in the Quaternary of Australasia,
50 Quaternary Science Reviews, 19, 559-590, 2000.
- 51 Ogle, B.A.: Geology of the Eel River Valley area, Humboldt County, California, Bulletin of the California Division of Mines
52 and Geology, 164, 1-128, 1953.
- 53 Omura, A., Emerson, W.K., and Ku, T.L.: Uranium-series ages of echinoids and corals from the upper Pleistocene Magdalena
54 terrace, Baja California Sur, Mexico, The Nautilus, 94, 184-189, 1979.

- 55 Orme, A.R.: Marine terraces and Quaternary tectonism, northwest Baja California, Mexico, *Physical Geography*, 1, 138-161,
56 1980.
- 57 Orr, P.C.: Late Pleistocene marine terraces on Santa Rosa Island, California, *Geological Society of America Bulletin*, 71,
58 1113-1120, 1960.
- 59 Ortlieb, L.: Néotectonique et variations du niveau marin au Quaternaire dans la région du Golfe de Californie, Mexique, Institut
60 Français de Recherche Scientifique pour le Développement en Coopération, Paris, France, Collection Études et
61 Thèses, 1036 pp, 1987.
- 62 Ortlieb, L.: Quaternary shorelines along the northeastern Gulf of California; geochronological data and neotectonic
63 implications, *Geological Society of America Special Paper*, 254, 95-120, 1991.
- 64 Ota, Y., and Omura, O.: Contrasting styles and rates of tectonic uplift of coral reef terraces in the Ryukyu and Daito Islands,
65 southwestern Japan, *Quaternary International*, 15/16, 17-29, 1992.
- 66 Ota, Y., Chappell, J., Kelley, R., Yonekura, N., Matsumoto, E., Nishimura, T., and Head, J.: Holocene coral reef terraces and
67 coseismic uplift of Huon Peninsula, Papua New Guinea, *Quaternary Research*, 40, 177-188, 1993.
- 68 Otto-Bliesner, B.L., Rosenbloom, N., Stone, E.J., McKay, N.P., Lunt, D.J., Brady, E.C., and Overpeck, J.T.: How warm was
69 the last interglacial? New model-data comparisons, *Philosophical Transactions of the Royal Society A*, 371,
70 20130097, <http://dx.doi.org/10.1098/rsta.2013.0097>, 20 pp, 2013.
- 71 Padgett, J.S., Kelsey, H.M., and Lamphear, D.: Upper-plate deformation of Late Pleistocene marine terraces in the Trinidad,
72 California, coastal area, southern Cascadia subduction zone, *Geosphere*, 15, 1323-1341, 2019.
- 73 Palmer, R.H.: Fossil and recent corals and coral reefs of western Mexico, *Proceedings of the American Philosophical Society*,
74 67, 21-31, 1928.
- 75 Perg, L.A., Anderson, R.S., and Finkel, R.C.: Use of a new ¹⁰Be and ²⁶Al inventory method to date marine terraces, Santa
76 Cruz, California, USA, *Geology*, 29, 879-882, 2001.
- 77 Pigati, J.S., Quade, J., Wilson, J., Jull, A.J.T., and Lifton, N.A.: Development of low-background vacuum extraction and
78 graphitization systems for ¹⁴C dating of old (40-60 ka) samples, *Quaternary International*, 166, 4-14, 2007.
- 79 Pindell, J.L., and Kennan, L.: Tectonic evolution of the Gulf of Mexico, Caribbean and northern South America in the mantle
80 reference frame: an update, *Geological Society of London Special Publications*, 328, 1-55, 2009.
- 81 Pinter, N., Lueddecke, S.B., Keller, E.A., and Simmons, K.R.: Late Quaternary slip on the Santa Cruz Island fault,
82 California, *Geological Society of America Bulletin*, 110, 711-722, 1998.
- 83 Polenz, M., and Kelsey, H.M.: Development of a late Quaternary marine terraced landscape during on-going tectonic
84 contraction, Crescent City coastal plain, California, *Quaternary Research*, 52, 217-228, 1999.
- 85 Polenz, M., Schasse, H.W., Kalk, M.L., and Petersen, B.B.: Geologic map of the Camano 7.5-minute quadrangle, Island
86 County, Washington, Washington Department of Natural Resources, Division of Geology and Earth Resources
87 Geologic Map, GM-68, scale 1:24,000, 2009.
- 88 Putnam, W.C.: Geomorphology of the Ventura region, California, *Geological Society of America Bulletin*, 53, 691-754, 1942.

- 89 Reyes-Bonilla, H., and López-Pérez, A.: Biogeography of the stony corals (Scleractinia) of the Mexican Pacific, *Ciencias*
90 *Marinas*, 24, 211-224, 1998.
- 91 Reyes-Bonilla, H., and López-Pérez, A.: Corals and coral-reef communities in the Gulf of California, in: *Atlas of Coastal*
92 *Ecosystems in the Western Gulf of California*, edited by: Johnson, M.E., and Ledesma-Vázquez, J., University of
93 Arizona Press, Tucson, Arizona, pp. 45-57, 2009.
- 94 Richards, H.G., Abbott, R.T., and Skymer, T.: The marine Pleistocene mollusks of Bermuda, *Notulae Naturae*, 425, 1-10,
95 1969.
- 96 Rockwell, T.K., Muhs, D.R., Kennedy, G.L., Hatch, M.E., Wilson, S.H., and Klinger, R.E.: Uranium-series ages, faunal
97 correlations and tectonic deformation of marine terraces within the Agua Blanca fault zone at Punta Banda, northern
98 Baja California, Mexico, in: *Geologic Studies in Baja California*, edited by: Abbott, P.L., Pacific Section, Society of
99 Economic Paleontologists and Mineralogists, Los Angeles, California, Book 63, 1-16, 1989.
- 00 Rockwell, T.K., Nolan, J., Johnson, D.L., and Patterson, R.H.: Ages and deformation of marine terraces between Point
01 Conception and Gaviota, Western Transverse Ranges, California, in: *Quaternary Coasts of the United States:*
02 *Marine and Lacustrine Systems*, edited by: Fletcher, C.H. III and Wehmiller, J.F., SEPM (Society for Sedimentary
03 Geology) Special Publication, no. 48, 333-341, 1992.
- 04 Rockwell, T.K., Clark, K., Gamble, L., Oskin, M.E., Haaker, E.C., and Kennedy, G.L.: Large Transverse Range earthquakes
05 cause coastal upheaval near Ventura, southern California, *Bulletin of the Seismological Society of America*, 106,
06 2706-2720, 2016.
- 07 Rohling, E.J., Grant, K., Hemleben, Ch., Siddall, M., Hoogakker, B.A.A., Bolshaw, M., and Kucera, M.: High rates of sea-
08 level rise during the last interglacial period, *Nature Geoscience*, 1, 38-42, 2008.
- 09 Rovere, A., Raymo, M.E., Vacchi, M., Lorscheid, T., Stocchi, P., Gómez-Pujol, L., Harris, D.L., Casella, E., O'Leary, M.J.,
10 and Hearty, P.J.: The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in
11 a warmer world, *Earth-Science Reviews*, 159, 404-427, 2016.
- 12 Rovere, A., Ryan, D., Murray-Wallace, C., Simms, A., Vacchi, M., Dutton, A., Lorscheid, T., Chutcharavan, P., Brill, D.,
13 Bartz, M., Jankowski, N., Mueller, D., Cohen, K., and Gowan, E.: Descriptions of database fields for the World Atlas
14 of Last Interglacial Shorelines (WALIS), Zenodo, [https://doi:10.5281/zenodo.3961544](https://doi.org/10.5281/zenodo.3961544), 2020.
- 15 Sak, P.B., and Fisher, D.M.: Effects of subducting seafloor roughness on upper plate vertical tectonism: Osa Peninsula, Costa
16 Rica, *Tectonics*, 23, doi: 10.1029/2002TC001474, 2004.
- 17 Sarna-Wojcicki, A.M., Lajoie, K.R., and Yerkes, R.F.: Recurrent Holocene displacement on the Javon Canyon fault--A
18 comparison of fault-movement history with calculated recurrence intervals, *U.S. Geological Survey Professional*
19 *Paper*, 1339, Chapter 8, 125-135, 1987.
- 20 Schellmann, G., and Radtke, U.: A revised morpho- and chronostratigraphy of the Late and Middle Pleistocene coral reef
21 terraces on southern Barbados (West Indies), *Earth-Science Reviews*, 64, 157-187, 2004.

- 22 Schumann, R.R., Minor, S.A., Muhs, D.R., Groves, L.T., and McGeehin, J.P.: Tectonic influences on the preservation of
23 marine terraces: Old and new evidence from Santa Catalina Island, California, *Geomorphology*, 179, 208-224, 2012.
- 24 Severns, M.: *Shells of the Hawaiian Islands—The Sea Shells*, ConchBooks, Hackenheim, Germany, 562 pp., 2011.
- 25 Shackleton, N.J., 1969. The last interglacial in the marine and terrestrial records, *Proceedings of the Royal Society of London*
26 *Series B*, 174, 135-154, 1969.
- 27 Shaw, J.H., and Suppe, J.: Active faulting and growth folding in the eastern Santa Barbara Channel, California, *Geological*
28 *Society of America Bulletin*, 106, 607-626, 1994.
- 29 Shennan I.: Interpretation of Flandrian Sea-level data from the fenland, *Proceedings of the Geologists' Association*, 83, 53–
30 63, 1982.
- 31 Shennan, I., Long, A.J., and Horton, B.P. eds.: *Handbook of sea-level research*, John Wiley & Sons, 2015.
- 32 Simkin, T., Tilling, R.I., Vogt, P.R., Kirby, S.H., Kimberly, P., and Stewart, D.B.: This dynamic planet; world map of
33 volcanoes, earthquakes, impact craters, and plate tectonics, U.S. Geological Survey Geologic Investigations Series
34 Map, I-2800, scale 1:30,000,000, 2006.
- 35 Simms, A.R., Rouby, H., and Lambeck, K.: Marine terraces and rates of vertical tectonic motion: The importance of glacio-
36 isostatic adjustment along the Pacific coast of central North America, *Geological Society of America Bulletin*, 128,
37 81-93, 2016.
- 38 Simms, A.R., Rood, D.H., and Rockwell, T.K.: Correcting MIS5e and 5a sea-level estimates for tectonic uplift, an example
39 from southern California, *Quaternary Science Reviews*, 248, doi.org/10.1016/j.quascirev.2020.106571, 2020.
- 40 Sirkin, L., Szabo, B.J., Padilla A., G., Pedrin A., S., and Diaz R., E.: Uranium-series ages of marine terraces, La Paz
41 Peninsula, Baja California Sur, Mexico, *Coral Reefs*, 9, 25-30, 1990.
- 42 Smith, W.S.T.: A topographic study of the islands of southern California, *University of California Bulletin of the*
43 *Department of Geology*, 2, 179-230, 1900.
- 44 Smith, W.S.T.: Marine terraces on Santa Catalina Island, *American Journal of Science* 25, 123–136, 1933.
- 45 Squires, D.F.: Results of the Puritan–American Museum of Natural History expedition to western Mexico. 7. Corals and
46 coral reefs in the Gulf of California, *Bulletin of the American Museum of Natural History*, 118, 367–432, 1959.
- 47 Stein, M., Wasserburg, G.J., Lajoie, K.R., and Chen, J.H.: U-series ages of solitary corals from the California coast by mass
48 spectrometry, *Geochimica et Cosmochimica Acta*, 55, 3709-3722, 1991.
- 49 Stein, M., Wasserburg, G.J., Aharon, P., Chen, J.H., Zhu, Z.R., Bloom, A., and Chappell, J.: TIMS U-series dating and
50 stable isotopes of the last interglacial event in Papua New Guinea, *Geochimica et Cosmochimica Acta*, 57, 2541-
51 2554, 1993.

- 52 Szabo, B.J., and Rosholt, J.N.: Uranium-series dating of Pleistocene molluscan shells from southern California—An open
53 system model, *Journal of Geophysical Research*, 74, 3253-3260, 1969.
- 54 Szabo, B.J., and Vedder, J.G.: Uranium-series dating of some Pleistocene marine deposits in southern California, *Earth and*
55 *Planetary Science Letters*, 11, 283-290, 1971.
- 56 Szabo, B.J., Hausback, B.P., and Smith, J.T.: Relative inactivity during the past 140,000 years of a portion of the La Paz
57 fault, southern Baja California Sur, Mexico, *Environmental Geology and Water Sciences*, 15, 119-122, 1990.
- 58 Thom, B.G.: The dilemma of high interstadial sea levels during the last glaciation, *Progress in Geography*, 5, 167-246, 1973.
- 59 Thompson, W.G., and Goldstein, S.L.: Open-system coral ages reveal persistent suborbital sea-level cycles, *Science*, 308,
60 401-404, 2005.
- 61 Thompson, W.G., Spiegelman, M.W., Goldstein, S.L., and Speed, R.C.: An open-system model for U-series age
62 determinations of fossil corals, *Earth and Planetary Science Letters*, 210, 365-381, 2003.
- 63 Tierney, P.W., and Johnson, M.E.: Stabilization role of crustose coralline algae during late Pleistocene reef development on
64 Isla Cerralvo, Baja California Sur (Mexico), *Journal of Coastal Research*, 28, 244-254, 2012.
- 65 Toth, L.T., Macintyre, I.G., and Aronson, R.B.: Holocene reef development in the eastern tropical Pacific, in: *Coral Reefs of*
66 *the Eastern Tropical Pacific*, edited by: Glynn, P.W., Manzello, D.P., and Enochs, I.C., Springer, Dordrecht,
67 Netherlands, pp. 177-201, 2017.
- 68 Trecker, M.A., Gurrola, L.D., and Keller, E.A.: Oxygen-isotope correlation of marine terraces and uplift of the Mesa Hills,
69 Santa Barbara, California, USA, *Geological Society of London Special Publications*, 146, 57-69, 1998.
- 70 Turney, C.S.M., and Jones, R.T.: Does the Agulhas Current amplify global temperatures during super-interglacials? *Journal*
71 *of Quaternary Science*, 25, 839-843, 2010.
- 72 Umhoefer, P.J., Maloney, S.J., Buchanan, B., Arrowsmith, J.R., Martinez-Gutiérrez, G., Kent, G., Driscoll, N., Harding, A.,
73 Kaufman, D., and Rittenour, T.: Late Quaternary faulting history of the Carrizal and related faults, La Paz region,
74 Baja California Sur, Mexico, *Geosphere*, 10, 476-504, 2014.
- 75 Upson, J.E.: Former marine shore lines of the Gaviota quadrangle, Santa Barbara County, California, *Journal of Geology*, 59,
76 415-446, 1951.
- 77 Uyeda, S., and Kanamori, H.: Back-arc opening and the mode of subduction, *Journal of Geophysical Research*, 84, 1049-1061,
78 1979.
- 79 Valentine, J.W.: Upper Pleistocene mollusca from Potrero Canyon, Pacific Palisades, California, *Transactions of the San*
80 *Diego Society of Natural History*, 12, 181-205, 1956.
- 81 Valentine, J.W.: Late Pleistocene megafauna of Cayucos, California and its zoogeographic significance, *Journal of*
82 *Paleontology*, 32, 687-696, 1958.

- 83 Valentine, J.W.: Pleistocene molluscan notes, 3. Rocky coast faunule, Bahia San Quintin, Mexico, *The Nautilus*, 74, 18-23,
84 1960a.
- 85 Valentine, J.W.: Habitats and sources of Pleistocene mollusks at Torrey Pines Park, California, *Ecology*, 41, 161-165,
86 1960b.
- 87 Valentine, J.W.: Paleoecologic molluscan geography of the Californian Pleistocene, *University of California Publications in*
88 *Geological Sciences*, 34, 309-442, 1961.
- 89 Valentine, J.W.: Pleistocene molluscan notes. 4. Older terrace faunas from Palos Verdes Hills, California, *Journal of Geology*,
90 70, 92-101, 1962.
- 91 Valentine, J.W.: Numerical analysis of marine molluscan ranges on the extra-tropical northeastern Pacific shelf, *Limnology*
92 *and Oceanography*, 11, 198-211, 1966.
- 93 Valentine, J.W.: Camalú: A Pleistocene terrace fauna from Baja California, *Journal of Paleontology*, 54, 1310-1318, 1980.
- 94 Valentine, J.W., and Meade, R.F.: Californian Pleistocene paleotemperatures, *University of California Publications in*
95 *Geological Sciences*, 40, 1-45, 1961.
- 96 Valentine, J.W., and Veeh, H.H.: Radiometric ages of Pleistocene terraces from San Nicolas Island, *Geological Society of*
97 *America Bulletin*, 80, 1415-1418, 1969.
- 98 Vedder, J.G., and Norris, R.M.: *Geology of San Nicolas Island California*, U.S. Geological Survey Professional Paper, 369,
99 1-65, 1963.
- 00 Vedder, J.G., Yerkes, R.F., and Schoellhamer, J.E.: *Geologic map of the San Joaquin Hills-San Juan Capistrano area*, Orange
01 *County, California*, U.S. Geological Survey Oil and Gas Investigations Map, OM 193, scale, 1:24,000, 1957.
- 02 Vedder, J.G., Yerkes, R.F., and Schoellhamer, J.E.: *Revised geologic map, structure sections and well table, San Joaquin Hills-*
03 *San Juan Capistrano area, California*, U.S. Geological Survey Open-File Report, 75-552, 1975.
- 04 Veeh, H.H., and Valentine, J.W.: Radiometric ages of Pleistocene fossils from Cayucos, California, *Geological Society of*
05 *America Bulletin*, 78, 547-550, 1967.
- 06 Veeh, H.H., and Chappell, J.: *Astronomical theory of climatic change: Support from New Guinea*, *Science*, 167, 862-865,
07 1970.
- 08 Weber, G.E.: Late Pleistocene slip rates on the San Gregorio fault zone at Point Año Nuevo, San Mateo County, California,
09 *in: Geology and Tectonics of the Central California Coast Region, San Francisco to Monterey*, edited by: Greene,
10 H.G., Weber, G.E., and Wright, T.L., *Pacific Section of the American Association of Petroleum Geologists*,
11 *Bakersfield, California*, pp. 193–203, 1990.
- 12 Weber, G.E., Lajoie, K.R., and Griggs, G.B.: *Coastal tectonics & coastal geologic hazards in Santa Cruz and San Mateo*
13 *Counties, California*, *Geological Society of America (Cordilleran Section), Field Trip Guide*, 187 pp., 1979.

- 14 Wehmiller, J.F.: Kinetic model options for interpretation of amino acid enantiomeric ratios in Quaternary mollusks: Comments
15 on a paper by Kvenvolden et al. (1979), *Geochimica et Cosmochimica Acta*, 45, 261-264, 1981.
- 16 Wehmiller, J.F.: A review of amino acid racemization studies in Quaternary mollusks: Stratigraphic and chronologic
17 applications in coastal and interglacial sites, Pacific and Atlantic coasts, United States, United Kingdom, Baffin
18 Island, and tropical islands, *Quaternary Science Reviews*, 1, 83-120, 1982.
- 19 Wehmiller, J.F.: Aminostratigraphy of southern California Quaternary marine terraces, in: *Quaternary Coasts of the United*
20 *States: Marine and Lacustrine Systems*, edited by: Fletcher, C.H. III and Wehmiller, J.F., SEPM (Society for
21 *Sedimentary Geology*) Special Publication, no. 48, 317-321, 1992.
- 22 Wehmiller, J.F.: United States Quaternary coastal sequences and molluscan racemization geochronology--What have they
23 meant for each other over the past 45 years? *Quaternary Geochronology*, 16, 3-20, 2013a.
- 24 Wehmiller, J.F.: Amino acid racemization, coastal sediments, in: *Encyclopedia of Scientific Dating Methods*, edited by:
25 Rink, W.J., and Thompson, J., Springer-Verlag, Berlin Heidelberg, pp. 1-10, 2013b.
- 26 Wehmiller, J.F., and Belknap, D.F.: Alternative kinetic models for the interpretation of amino acid enantiomeric ratios in
27 Pleistocene mollusks: Examples from California, Washington, and Florida, *Quaternary Research*, 9, 330-348, 1978.
- 28 Wehmiller, J.F., and Emerson, W.K.: Calibration of amino acid racemization in late Pleistocene mollusks: Results from
29 Magdalena Bay, Baja California Sur, Mexico, with dating applications and paleoclimatic implications, *The*
30 *Nautilus*, 94, 31-36, 1980.
- 31 Wehmiller, J.F., and Pellerito, V.: Database of Quaternary coastal geochronologic information for the Atlantic and Pacific
32 coasts of North America, Delaware Geological Survey Open File Report, 50, 2015.
- 33 Wehmiller, J.F., Lajoie, K.R., Kvenvolden, K.A., Peterson, E., Belknap, D.F., Kennedy, G.L., Addicott, W.O., Vedder, J.G.,
34 and Wright, R.W.: Correlation and chronology of Pacific coast marine terrace deposits of continental United States
35 by fossil amino acid stereochemistry--technique evaluation, relative ages, kinetic model ages, and geologic
36 implications, U.S. Geological Survey Open-File Report, 77-680, 1-196, 1977a.
- 37 Wehmiller, J.F., Kennedy, G.L., and Lajoie, K.R.: Amino acid racemization age estimates for Pleistocene marine deposits in
38 the Eureka-Fields Landing area, Humboldt County, California, U.S. Geological Survey Open-File Report, 77-517,
39 25 pp., 1977b.
- 40 Wehmiller, J.F., Lajoie, K.R., Sarna-Wojeicki, A.M., Yerkes, R.F., Kennedy, G.L., Stephens, T.A., and Kohl, R.F.: Amino-
41 acid racemization dating of Quaternary mollusks, Pacific Coast United States, U.S. Geological Survey Open-File
42 Report, 78-701, pp. 445-448, 1978.
- 43 Wehmiller, J.F., Simmons, K.R., Cheng, H., Edwards, R.L., Martin-McNaughton, J., York, L.L., Krantz, D.E., and Shen, C.-
44 C.: Uranium-series coral ages from the U.S Atlantic Coastal Plain--the "80 ka problem" revisited, *Quaternary*
45 *International*, 120, 3-14, 2004.

- 46 Woodring, W.P., Bramlette M.N., and Kew, W.S.W.: Geology and paleontology of Palos Verdes Hills, California, U.S.
47 Geological Survey Professional Paper, 207, 1-145, 1946.
- 48 Woods, A.J.: Geomorphology, deformation, and chronology of marine terraces along the Pacific Coast of central Baja
49 California, Mexico, Quaternary Research, 13, 346-364, 1980.
- 50 Wright, R.H.: Late Pleistocene marine fauna, Goleta, California, Journal of Paleontology, 46, 688-695, 1972.
- 51 Yamamoto, M., Yamamuro, M., and Tanaka, Y.: The California current system during the last 136,000 years: response of
52 the North Pacific High to precessional forcing, Quaternary Science Reviews, 26, 405-414, 2007.
- 53 Zullo, V.A.: A late Pleistocene marine invertebrate fauna from Bandon, Oregon, Proceedings of the California Academy of
54 Science, 36, 347-361, 1969.

55 **FIGURE CAPTIONS**

- 56 **Figure 1: Tectonic setting of North America showing lithospheric plates, plate boundaries, and features referred to in**
57 **the text. Redrawn in simplified form from Simkin et al. (2006). PTJ, Panama Triple Junction.**
- 58 **Figure 2: Tectonic setting of the Pacific Coast of North America, from southern Canada to southern Baja California**
59 **Sur, Mexico, showing plates, plate boundaries, structures, and localities referred to in the text. Redrawn in**
60 **simplified form from Simkin et al. (2006). CSZ, Cascadia Subduction Zone; MTJ, Mendocino Triple**
61 **Junction; SAF, San Andreas Fault.**
- 62 **Figure 3. Diagrams showing the terminology used for marine terraces: (a) simple case of a modern wave-cut bench**
63 **or platform in the surf zone with marine gravels and modern shells (blue symbols), shoreline angle, and**
64 **single emergent marine terrace above it, with a colluvial cover masking most of the marine terrace deposits**
65 **with their fossils (red symbols). Such an emergent terrace could have formed from a higher-than-present sea**
66 **level or from uplift since the time of initial terrace formation; (b) more complex case on an uplifting coast**
67 **with the features described above, but an additional (older) terrace above the lower one. Note that in (b),**
68 **colluvial deposits cover both of the emergent terraces, making them appear as one landform, with a single**
69 **inner edge that is at a higher elevation than the shoreline angles of both emergent terraces.**
- 70 **Figure 4: (a) Modern wave-cut bench exposed at low tide, shoreline angle, sea cliff, and wave-cut bench of emergent,**
71 **~80 ka marine terrace, Cormorant Rock, San Nicolas Island, California, USA. (b) Modern wave-cut bench**
72 **exposed at low tide, overlying marine gravels, shoreline angle, and sea cliff, San Pedro, California, USA.**
73 **Photographs by D.R. Muhs.**
- 74 **Figure 5: Examples of exposures of ancient shoreline angles: (a) north coast of Santa Cruz Island, California, just**
75 **east of Prisoners Harbor; (b) west side of Santa Barbara Island, California, showing benches and shoreline**
76 **angles of three of the four lowest marine terraces. Photographs by D.R. Muhs.**
- 77 **Figure 6: Two methods of estimating paleo-sea level when shoreline angles are not exposed, Cayucos, California. (a)**
78 **Map showing extent of marine terrace dated to ~120 ka and fossil localities from Muhs et al. (2002a). (b)**
79 **Topographic profile of a shore-normal transect in the vicinity of fossil localities LACMIP. 11923, 11762, and**
80 **11922, showing measured bench elevations and paleo-sea cliff elevations, where they are exposed;**
81 **intersection of extrapolated wave-cut bench slope landward and paleo-sea cliff slope downward yields an**
82 **estimated shoreline angle elevation of ~8 m. (c) Photograph of outer edge of terrace at fossil locality**
83 **LACMIP 11923, showing wave-cut bench 3 m above sea level with *Penitella penita* fossils (rock-boring**
84 **bivalves) in growth position (see enlargement in (d)). *P. penita* lives in waters 10 m deep or shallower, so**
85 **bench elevation (3 m) plus maximum depth of growth (10 m) yields a *maximum-limiting* paleo-sea level of ~13**
86 **m above present. Photographs by D.R. Muhs.**
- 87 **Figure 7. (a) Map showing the distribution of living hermatypic corals and coral reefs along the Golfo de California**
88 **coasts of Mexico, the Pacific coast of Mexico, and the Pacific coast of Central America (compiled from Reyes-**
89 **Bonilla and López-Pérez, 2009; Alvarado et al., 2010; Glynn et al., 2017). (b), (c) Examples of modern**

hermatypic corals along the Pacific coast of Central America (photographs courtesy of Lauren Toth, U.S. Geological Survey).

Figure 8. Examples of fossil marine organisms used for geochronology of marine terrace deposits on the Pacific coast of North America: (a) solitary corals *Balanophyllia elegans* (fossil), San Nicolas Island, California (U-series dating); (b) *Porites panamensis* (fossil), Isla Carmen, Baja California Sur (U-series dating); (c) *Saxidomus* (fossil), San Nicolas Island, California (amino acid geochronology); (d) *Tegula* (fossil), San Clemente Island, California (amino acid geochronology); (e) *Chione* (modern, upper row; fossil, lower row), Cholla Bay, Sonora, Mexico (amino acid geochronology). All photographs by D.R. Muhs.

Figure 9. Isotopic evolution diagrams for (a) colonial corals from the Southampton Formation (~80 ka) and Devonshire marine member of the Rocky Bay Formation (~120 ka) of Bermuda and (b) solitary corals from terrace 1 (~80 ka) and terrace 2 (~120 ka) of San Nicolas Island, California. Bermuda data are from Muhs et al. (2002a), but do not include two samples that have evidence of U loss. San Nicolas Island data are from Muhs et al. (2006). Blue bands show isotopic evolution pathways for corals having mostly closed-system history and initial $^{234}\text{U}/^{238}\text{U}$ activity values of 1.16 to 1.14, which bracket measured values in modern seawater (Chen et al., 1986; Delanghe et al., 2002) and modern corals (Muhs et al., 2002b).

Figure 10. Plots of apparent $^{230}\text{Th}/^{238}\text{U}$ ages vs. back-calculated initial $^{234}\text{U}/^{238}\text{U}$ values in solitary corals from (a) Eel Point terrace, San Clemente Island, California, (b) Terrace 2, west end of San Nicolas Island, California [same as in Fig. 9b, but with different scales], (c) Sea Cave terrace, Punta Banda, Baja California, Mexico, and (d) colonial corals (mostly *Acropora palmata*) from the Rendezvous Hill terrace, north end of Barbados, West Indies. San Clemente Island and Punta Banda data are from Muhs et al. (2002b), San Nicolas Island data are from Muhs et al. (2006), and Barbados data are from Muhs and Simmons (2017). Also shown (blue bands) is the range of $^{234}\text{U}/^{238}\text{U}$ activity values in modern seawater (Chen et al., 1986; Delanghe et al., 2002). Note that in both solitary corals and colonial corals, samples plotting above seawater values tend to be biased to older apparent ages, but the degree of bias varies from locality to locality.

Figure 11. (a) Plot showing D-leucine/L-leucine in fossil *Saxidomus* shells (or equivalent values converted from *Leukoma staminea* shells; see Lajoie et al., 1980) from marine terrace deposits of the Pacific Coast of the USA, from Kennedy et al. (1982). Localities are arranged from north (left) to south (right), parallel to latitudinal trend of mean annual air temperatures increasing to the south. Samples plotting along the pink line are correlated with MIS 5e (~120 ka) based on calibration to U-series-dated corals from Cayucos and Point Loma; samples plotting along the blue line are correlated to MIS 5a (~80 ka), based on U-series-dated corals from Coquille Point, Oregon. Calibration points used are the only ones that were available at the time of the original study. Samples plotting below these lines are correlated with MIS 3 or to Holocene-dated deposits (gray line). Not included from the original study are data points from Whidbey Island, Washington, which are interpreted to be from glaciomarine deposits (Polenz et al., 2009). Colors of circles indicate molluscan fauna thermal aspects (see discussion of Fig. 13). (b) Plot of same data as in (a), except new U-series ages of corals, generated since 1982, have been added and thermal aspects of some faunas have been modified (Muhs et al., 2002b, 2006, 2014b).

Figure 12. (a) Plot of mean D/L values in glutamic acid (vertical axis) in fossil *Tegula* from dated (filled circles) and undated (open circles) marine terraces on the California and Baja California coast, shown as a function of latitude (horizontal axis) as a proxy for long-term temperature history, cooler in the northwest, warmer in the southeast. Error bars are ± 1 standard deviation, based on D/L values in 3 to 6 individual shells from the same deposit. Colored bands (“aminozones”) indicate correlation between fossil localities of the same age, anchored by U-series dating of corals. Terrace name abbreviations: SMI, San Miguel Island; SRI, Santa Rosa Island; SCRZI, Santa Cruz Island; N, Nestor; BR, Bird Rock; PDM, Paseo del Mar; G, Gaffey; SC, Sea Cave; L, Lighthouse; see Muhs et al. (1994, 2002b, 2006, 2014b, 2015) for terrace stratigraphic names and U-series ages. Data from Santa Cruz Island-West, Santa Cruz Island-South, Santa Barbara Island, and Anacapa Island are from Muhs and Groves (2018); all other data are from Muhs et al. (2014b). (b) Same as in (a), but for mean D/L values in valine.

Figure 13. Modern geographic ranges of extralimital and northward or southward-ranging fossil mollusks found in ~80,000 yr B.P. marine terrace deposits at Green Oaks Creek, Point Año Nuevo and Santa Cruz, California

40 and the ~130,000 yr B.P. Millerton Formation at Toms Point, Tomales Bay, California. Ages are from Grove
41 et al. (1995) for the Millerton Formation at Toms Point and Muhs et al. (2006) for the other localities. Fossil
42 data for the Millerton Formation are from Johnson (1962); Davenport terrace fossil data are from Addicott
43 (1966) and Muhs et al. (2006). Modern species names and geographic ranges updated by the author from
44 Abbott and Haderlie (1980), O'Clair and O'Clair (1998), and Coan et al. (2000).

45 **Figure 14.** Map of the Pacific Coast of North America with structural features as shown in Figure 2, but also plotted
46 are localities (filled red circles) where U-series ages of corals dating to MIS 5e (~120 ka) have been reported.
47 Abbreviations are keyed to Table S1 and are as follows: CP, Cayucos Point; C, Cayucos; DC, Diablo Canyon;
48 PSL, Point San Luis; SB, Shell Beach; SMI, San Miguel Island; SRI, Santa Rosa Island; SCRZI, Santa Cruz
49 Island; SNI, San Nicolas Island; PV, Palos Verdes Hills; NB, Newport Beach; SCI, San Clemente Island; PL,
50 Point Loma; PB, Punta Banda; IG, Isla Guadalupe; BM, Bahía Magdalena; CP, Cabo Pulmo; CE, Isla
51 Cerralvo; LP, La Paz; PCO, Punta Coyote; IC, Isla Coronado; BSN, Bahía San Nicolas; MU, Mulegé; PC,
52 Punta Chivato; CSZ, Cascadia Subduction Zone; MTJ, Mendocino Triple Junction; SAF, San Andreas Fault.

53 **Figure 15.** Map of the Pacific Coast of North America with structural features as shown in Figure 2, but also plotted
54 are localities (open red circles) where amino acid geochronology has permitted correlation of marine deposits
55 to MIS 5e (~120 ka). Abbreviations are keyed to Table S2 and are as follows: WB, Willapa Bay; YB, Yaquina
56 Bay; E, Eureka; TB, Tomales Bay; AH, Arroyo Hondo; SRI, Santa Rosa Island; SCRZI, Santa Cruz Island;
57 ANA, Anacapa Island; SBI, Santa Barbara Island; M, Malibu; PP, Pacific Palisades; PVH, Palos Verdes Hills;
58 SP, San Pedro; NM, Newport Mesa; LB, Laguna Beach; SO, San Onofre; TP, Torrey Pines; PL, Point Loma;
59 BL, Border locality; CM, Camalú; PSR, Punta Santa Rosalíllita; BT, Bahía Tortuga; BA, Bahía Asunción;
60 BSN, Bahía San Nicolas North; BC, Bahía Concepción; BSI, Bahía Santa Inés; SL, San Lucas; SR, Santa
61 Rosalia; CSM, Caleta Santa Maria; SB, Salina la Borrascosa; PG, Punta Gorda; ESB, East of Salina la
62 Borrascosa; PLO, Puerto Lobos; PLI, Puerto Libertad; PC, Punta Cuevas; SPC, Southeast of Punta Cuevas;
63 PT, Punta Tepopa; IT, Isla Tiburón; PK, Punta Kino; CSZ, Cascadia Subduction Zone; MTJ, Mendocino
64 Triple Junction; SAF, San Andreas Fault.

65 **Figure 16:** Map of southwestern Canada, Washington, and Oregon, showing localities referred to in the text

66 **Figure 17:** Photographs of possible MIS 5e marine deposits in the Willapa Bay area, near Bay Center, Washington:
67 (a), (b) location of shell-bearing layer relative to modern sea level; (c) (d) closeup views showing shell bed with
68 *Ostrea conchaphilia*, *Saxidomus gigantea*, and coniferous wood fragments. All photographs by D.R. Muhs.

69 **Figure 18:** Maps of marine terraces in the Coquille Point (a) and Cape Blanco (b) areas of southwestern Oregon. (a)
70 Qwr, Whisky Run terrace deposits; Qp, Pioneer terrace deposits; Qsd, Seven Devils terrace deposits
71 (correlated to MIS 5e); Qm, Metcalf terrace deposits. (b) Qcb, Cape Blanco terrace deposits; Qp, Pioneer
72 terrace deposits; Qsb, Silver Butte terrace deposits (correlated to MIS 5e); Qic, Indian Creek terrace deposits;
73 Qpr, Poverty Ridge terrace deposits. Redrawn from terrace maps in McNelly and Kelsey (1990) for (a) and
74 Kelsey (1990) for (b).

75 **Figure 19:** Cross sections of marine terrace deposits in coastal Oregon in the Newport-Yaquina Bay area (a), Cape
76 Arago-Coquille Point area (b), and Cape Blanco area (c), and estimated ages based on U-series dating of corals,
77 amino acid geochronology of mollusks, and degree of soil development. Cross sections from Kelsey et al. (1996)
78 for (a), McNelly and Kelsey (1990) for (b), and Kelsey (1990) for (c); geochronological data from Kennedy et
79 al. (1982), Muhs et al. (1990, 2006), and Kelsey et al. (1996). Deposit abbreviations as defined in Figure 18
80 caption.

81 **Figure 20.** Photographs of the Pioneer (~100 ka, MIS 5c?) and Cape Blanco (~80 ka, MIS 5a?) marine terraces (a) in
82 the Cape Blanco area, with closeups wave-cut bench on Miocene sandstone, shell bed, and terrace sediments
83 (b, c). All photographs by D.R. Muhs.

84 **Figure 21.** Map of coastal northwestern California showing geographic and structural features and locations discussed
85 in the text. CSZ, Cascadia Subduction Zone; MFZ, Mendocino Fault Zone; SAF, San Andreas Fault; SGH,
86 San Gregorio fault zone; H, Hayward Fault; M, Maacama Fault Zone; GV, Green Valley Fault; H-R,
87 Healdsburg-Rodgers Fault.

88 **Figure 22:** (a) Map showing the distribution of marine terraces in the Laguna Point-Fort Bragg-Cabrillo Point area
89 (redrawn from Jennings and Strand, 1960); (b) photograph of 10-m-high marine terrace, correlated to MIS

- 90 5e (Merritts and Bull, 1989); (c) pholad holes in outer edge of 10-m-high, MIS 5e terrace. Photographs by D.R.
91 Muhs.
- 92 **Figure 23:** (a) Landsat band 5 image (from U.S. Geological Survey) of the Tomales Bay, California area, with fossil
93 localities of the Millerton Formation (filled red circles) from Johnson (1962); the formation is correlated to
94 MIS 5e by thermoluminescence (Grove et al., 1995) and amino acid geochronology (Muhs and Groves, 2018).
95 (b) Ground photograph from Toms Point, looking south to Tomales Bay. (c) Marine terrace deposits with
96 fossils exposed on wave-cut bench at Toms Point. (d) Marine deposits with fossils exposed at Millerton Point.
97 Photographs by D.R. Muhs.
- 98 **Figure 24.** Map of a portion of the central coast of California, from Santa Cruz to just north of Point Año Nuevo,
99 showing marine terraces, fossil localities, and location of the San Gregorio fault zone (solid gray lines; dashed
00 where uncertain). Marine terrace inner edges redrawn from Bradley and Griggs (1976) and Weber et al.
01 (1979); location of the San Gregorio fault zone from Weber et al. (1979) and Weber (1990).
- 02 **Figure 25.** Map of geographic features in southern California and localities referred to in the text. Grey lines are faults
03 from Jennings (1994). AI, Anacapa Island; SBI, Santa Barbara Island.
- 04 **Figure 26.** Map of marine terrace deposits (brown shades), terrace inner edges (black lines), fossil localities (open red
05 circles), landslide deposits (gray shades), and faults in the Palos Verdes Hills area, Los Angeles County,
06 California. Redrawn from Woodring et al. (1946). PVS, Palos Verdes Sand, part of which is of MIS 5e age
07 (Muhs and Groves, 2018); PDM, Paseo del Mar terrace (~80 ka, MIS 5a); G, Gaffey terrace (~120 ka, MIS 5e;
08 Muhs et al., 2006). LACMIP, Natural History Museum of Los Angeles County Invertebrate Paleontology fossil
09 locality numbers; WBK, fossil localities of Woodring et al. (1946).
- 10 **Figure 27:** View of outer edges of the Paseo del Mar (~80 ka) and Gaffey (~120 ka) terraces, looking northwest from
11 Point Fermin (see Fig. 26). On left side of photograph, a small fragment of the 12th terrace is visible
12 (=LACMIP loc. 1304 on Fig. 26). Photograph by D.R. Muhs.
- 13 **Figure 28:** (a) Photograph of Newport Mesa (terrace 2) of MIS 5e age, view to the west, upper Newport Bay in the
14 foreground. (b) Map of marine terraces in the Newport Bay area, redrawn from Vedder et al. (1957, 1975)
15 and Grant et al. (1999); terrace elevations and ages increase with terrace numbering (terrace 1=youngest;
16 terrace 8=oldest). Filled red circles are fossil localities with U-series ages from Grant et al. (1999); open red
17 circles are amino acid geochronology fossil localities of Wehmiller et al. (1977a). Photograph in (a) by D.R.
18 Muhs.
- 19 **Figure 29:** (a) Map showing marine terrace inner edges (redrawn from Kern, 1977) and LACMIP fossil localities where
20 U-series ages are reported by Muhs et al. (1994; 2002b); corals from the Nestor terrace date to ~120 ka (MIS
21 5e) and ~100 ka (MIS 5c); those from the Bird Rock terrace date to ~80 ka (MIS 5a). (b) Photograph showing
22 outer edges of the Nestor and Bird Rock terraces on the west coast of Point Loma, looking north (see location
23 in (a)). Photograph by D.R. Muhs.
- 24 **Figure 30:** Map of northwestern Baja California, showing marine terrace deposits (brown shades), terrace inner edges
25 (black lines), and fossil localities (red/orange circles). Punta Banda fossil localities are shown in Figure 31.
26 South of Punta Banda, most fossil localities are undated, but based on elevations shown here, many likely date
27 to some part of MIS 5 and could contain mixes of fossils of two ages, with cool-water (blue dots) and warm-
28 water (red dots) faunas (see text for discussion). Marine terrace deposits and inner edge mapping redrawn
29 from Orme (1980).
- 30 **Figure 31:** (a) Map of the Punta Banda area, south of Ensenada, Baja California (see Fig. 30 for location). Lighthouse
31 terrace is dated to MIS 5a (~80 ka) and Sea Cave terrace is dated to MIS 5e (~120 ka), both by U-series on
32 corals (Rockwell et al., 1989; Muhs et al., 2002b). Terrace mapping is redrawn from Rockwell et al. (1989).
33 (b) Photograph of the ~120 ka Sea Cave terrace on the northeast side of Punta Banda, showing thick alluvial
34 cover. Photograph by D.R. Muhs.
- 35 **Figure 32:** (a) Satellite image of southernmost Baja California Sur and parts of Sonora and Sinaloa, acquired on 27
36 November 2011 using the MODIS instrument on the Aqua satellite (courtesy of the NASA Rapid Response
37 Team). Red circles indicate marine terrace localities where U-series ages on corals have yielded MIS 5e ages
38 (see Table S1). (b), (c), (d), (e) Photographs of growth-position corals dated to MIS 5e from Punta Chivato,

- 39 Isla Coronado, and Isla Cerralvo (see Johnson, 2002; Johnson et al., 2007; Tierney and Johnson, 2012). All
40 coral photographs are courtesy of Markes Johnson, Williams College.
- 41 **Figure 33:** Satellite image of northwestern Mexico (Baja California, Baja California Sur, Sonora, and parts of adjacent
42 areas) showing localities (red circles) where amino acid geochronology of fossil mollusks has yielded MIS 5e
43 ages (Ortlieb, 1987, 1991; Valentine, 1980; Woods, 1980; Emerson et al., 1981; Keenan et al., 1981). Also
44 shown for reference are two U-series-dated MIS coral localities, Punta Banda (PB) and Isla Guadalupe (IG).
45 Image acquired 28 May 2006 by the Medium Resolution Imaging Spectrometer (MERIS) onboard the Envisat
46 satellite, courtesy of the European Space Agency. Abbreviations of amino acid localities (see Table S2): CM,
47 Camalú; PSR, Punta Santa Rosalillita; BT, Bahía Tortuga; BA, Bahía Asunción; BSN, Bahía San Nicolas
48 North; BC, Bahía Concepción; BSI, Bahía Santa Inés; SL, San Lucas; SR, Santa Rosalia; CSM, Caleta Santa
49 Maria; SB, Salina la Borrascosa; PG, Punta Gorda; ESB, East of Salina la Borrascosa; PLO, Puerto Lobos;
50 PLI, Puerto Libertad; PC, Punta Cuevas; SPC, Southeast of Punta Cuevas; PT, Punta Tepopa; IT, Isla
51 Tiburón; PK, Punta Kino.
- 52 **Figure 34:** Plot showing ratios of D-alloisoleucine to L-isoleucine in fossil *Chione* (red squares) and *Dosinia* (blue
53 squares) as a function of latitude in Baja California, Baja California Sur, and Sonora (localities listed in Table
54 S2). Amino acid data from Ortlieb (1987, 1991), and Umhoefer et al. (2014). Also shown (solid green squares)
55 are ratios of D-alloisoleucine to L-isoleucine in late Holocene, radiocarbon-dated *Chione* shells from the upper
56 Golfo de California (data from Martin et al., 1996) and ratios of D-alloisoleucine to L-isoleucine (blue shade)
57 at equilibrium (from Miller and Mangerud, 1985). Pink shade defines aminozone correlated to MIS 5e, based
58 on calibration with U-series-dated deposits that also contain *Chione* shells (gold squares).
- 59 **Figure 35:** (a) Map of Central America, showing structural features (redrawn from Mann, 2007, and Pindell and
60 Kennan 2009), lithospheric plates, directions of present plate movements (arrows), and localities referred to in
61 text. (b) Map showing marine terrace deposits on the Osa Peninsula of Costa Rica (redrawn from Gardner et
62 al., 2013); legend shows possible correlation to marine isotope stages (MIS).
- 63 **Figure 36:** (a) Map of marine terraces on San Nicolas Island, California (from Vedder and Norris, 1963; Muhs et al.
64 2012); orange dots are fossil localities. (b) Topographic profile across the lowest 11 terraces in the Celery Creek
65 Canyon area (blue boxed area in (a)), showing terrace numbers, shoreline angle elevations (from Muhs et al.,
66 2018), and (green lettering) ratios of the amino acids D-alloisoleucine to L-isoleucine in the fossil gastropod
67 *Tegula* (data mostly from Muhs, 1985). Assuming a constant rate of uplift, terrace 4 could be ~390 ka (~MIS
68 11). Note that large jump in elevation between terraces 2a and 4 is mirrored by substantial increase in amino
69 acid ratios, indicating the likelihood that terraces representing MIS 9 (~300 ka) and MIS 7 (~200 ka) were
70 removed by sea cliff retreat during MIS 5e.
- 71 **Figure 37:** View of the lowest six marine terraces on the west coast of San Clemente Island, California, looking south.
72 Terrace 2b has been dated to MIS 5e (~120 ka) by TIMS U-series methods on coral (Muhs et al., 2002b);
73 terrace 1 is estimated to be ~80 ka (MIS 5a) based on amino acid ratios in mollusks (Muhs, 1983). Terrace 6
74 could be as old as ~600 ka or more, if the rate of uplift has been constant over time. SLA=shoreline angle
75 elevation, in meters above modern sea level. Photograph by D.R. Muhs.
- 76 **Figure 38:** Use of an extinct species as a biostratigraphic marker. (a) Map of marine terraces and fossil localities on
77 Santa Barbara Island, California (from Muhs and Groves, 2018), along with (b) topographic profile of terraces
78 (location shown in (a)). Also shown in both (a) and (b) are terrace deposits that host the extinct fossil gastropod
79 *Pusio fortis* (formerly *Calicantharus fortis*), shown in (c). Note that in (a) and (b), this taxon is not found on
80 terrace 1, whose deposits contain a mix of fossils dating to MIS 5e and 5c, but the species is found on higher
81 terraces, indicating that it likely became extinct before MIS 5e. Photograph in (c) is from 69-m-high terrace
82 deposits on San Miguel Island, by D.R. Muhs.
- 83 **Figure 39:** Map of the Pacific Coast of North America with structural features as shown in Figure 2, but also plotted
84 are localities (filled blue circles) where U-series ages of corals dating to MIS 5a (~80 ka) have been reported
85 and localities (open blue circles) where amino acid ratios in mollusks permit correlation to MIS 5a.
86 Abbreviations are keyed to Table S3 and are as follows: BC, Bay Center, Willapa Bay; NJ, Newport Jetty; FP,
87 Five Mile Point; CP Coquille Point; CB, Cape Blanco; CC, Crescent City; PA, Point Arena; HB, Half Moon
88 Bay; GO, Green Oaks Creek; AN, Point Año Nuevo; SC, Santa Cruz; G, Gaviota; AH, Arroyo Hondo; SMI,

89 San Miguel Island; SRI, Santa Rosa Island; SNI, San Nicolas Island, PV, Palos Verdes Hills; SP, San Pedro;
90 SCI, San Clemente Island; PL, Point Loma; PB, Punta Banda; CSZ, Cascadia Subduction Zone; MTJ,
91 Mendocino Triple Junction; SAF, San Andreas Fault.

92 **Figure 40:** Examples of extralimital southern species of bivalves indicating sea surface temperatures higher than
93 present during MIS 5e (filled and open circles); also shown, in light purple shading, are the modern ranges of
94 these taxa (from Coan and Valentich-Scott, 2012): (a) *Chione undatella* and (b) *Dosinia ponderosa*. Age and
95 fossil data compiled from sources as follows: Jordan (1936), Valentine (1956, 1960), Kanakoff and Emerson
96 (1959), Johnson (1962), Omura et al. (1979), Emerson (1980), Emerson et al. (1981), Kennedy et al. (1982),
97 Ashby et al. (1987), Keenan et al. (1987), Ortlieb (1987, 1991), Grove et al. (1995), Grant et al. (1999), Kennedy
98 (2000), and Muhs and Groves (2018).

99 **Figure 41:** Examples of extralimital southern species of gastropods indicating sea surface temperatures higher than
00 present during MIS 5e (filled and open circles); also shown, in light purple shading, are the modern ranges of
01 these taxa (from Bertsch and Aguilar Rosas, 2016 and Keen, 1971): (a) *Mexacanthina lugubris* and (b)
02 *Stramonita biserialis*. Age and faunal data compiled from Jordan (1936), Kanakoff and Emerson (1959),
03 Valentine (1962, 1980), Lipps et al. (1968), Kern (1977), Ashby et al. (1979), Omura et al. (1979), Lindberg et
04 al. (1980), Emerson et al. (1981), Ortlieb (1987), Rockwell et al. (1989), Muhs et al. (1983, 1992, 2002b, 2014a),
05 Grant et al. (1999), Wehmiller and Pellerito (2015). Note that the presence of *Stramonita biserialis* on San
06 Clemente Island is from a recent discovery of this taxon by L.T. Groves, Natural History Museum of Los
07 Angeles County, from the NOTS Pier terrace (Muhs, 1983), correlated to the Eel Point terrace (~120 ka; Muhs
08 et al., 2002b) by amino acid ratios in mollusks.

09 **Figure 42:** (a) Map showing the modern distribution of species of the hermatypic coral genus *Pocillopora* (purple
10 shading; from Reyes-Bonilla and López-Pérez, 1998), and U-series-dated, MIS 5e occurrences of this genus
11 elsewhere in the region. MIS 5e age and faunal data from: Durham (1980), Lindberg et al. (1980), Sirkin et al.
12 (1990), Szabo et al. (1990), Muhs et al. (2002b), DeDiego-Forbis et al. (2004), Johnson et al. (2007), and Tierney
13 and Johnson (2012). Occurrence of fossil *Pocillopora* on the Mexican coast south of Oaxaca is undated, but
14 could be of MIS 5e age, and is from Palmer (1928). (b) Photograph of fragments of *Pocillopora guadalupensis*
15 from Isla Guadalupe, Mexico, dated to ~120 ka (Muhs et al., 2002b). Photograph by D.R. Muhs.

16 **Figure 43:** Map of a part of the Pacific Coast of North America showing marine invertebrate faunal zones (from
17 Valentine, 1966) and their correlation with mean annual sea surface temperatures (temperature data from
18 U.S. National Oceanic and Atmospheric Administration).

19 **Figure 44:** Extralimital southern species of gastropods and bivalves (from Lindberg et al., 1980) in marine deposits of
20 Isla Guadalupe, Mexico (see Fig. 42 for location) dated to MIS 5e by Muhs et al. (2002b), and their modern
21 latitudinal distribution (from Keen, 1971 for gastropods; from Coan and Valentich-Scott, 2012 for bivalves).
22 Note also that the deposits also contain the extralimital southern genus of coral *Pocillopora* (see Fig. 42), and
23 the Indo-Pacific gastropod *Naria cernica* (see text for discussion).

24 **Figure 45:** Cross section of the lowest marine terraces on San Nicolas Island, California and modeled sea level curves
25 for this island, the Florida Keys, and Barbados (from Muhs et al., 2012), showing link between sea level
26 history, uplift rate, and terrace geomorphology, influenced by glacial isostatic adjustment (GIA) processes.
27 Qmt, Quaternary marine terrace deposits.
28

Name of RSL indicator	Description of RWL	Description of IR
Marine terrace elevation	Junction of the wave-cut platform and sea cliff	Shoreline angles typically are within a meter or less of mean sea level.
Pholadidae elevation	Elevation of rock-boring mollusks in growth position	Bivalves in the Pholadidae family typically live within the intertidal zone and usually at depths of ~10 m or less, providing a minimum paleo-sea level elevation
Coral reef terrace elevation	Uppermost growth-position coral reef colony is a minimum elevation; habitat depth measurement needs to be added for better accuracy	In southern Baja California Sur and adjacent parts of mainland Mexico and Central America, colonial hermatypic corals in growth position (e.g., <i>Porites</i> and <i>Pocillopora</i>) typically live in the intertidal zone and are usually found at depths of ~10 m or less, providing a minimum paleo-sea level elevation

29 **Table 1: different types of RSL indicators, reference water level (RWL) and indicative range (IR) on the Pacific**
30 **Coast of North America.**

Measurement technique	Description	Typical accuracy
Hand level and tape	Hand level and metered tape and/or use of transit and stadia	On the order of tens of centimeters
Aneroid altimeter	American Paulin System aneroid altimeter	On the order of 0.5 meter
Global Positioning System (GPS)	Uses satellite array and triangulation with post-processing to increase accuracy	On the order of tens of centimeters if satellite geometry is favorable

31 **Table 2: measurement techniques used to establish the elevation of MIS 5e shorelines on the Pacific Coast of North**
32 **America.**




FIGURE 1




FIGURE 2






FIGURE 3

(a)



(b)




FIGURE 4





FIGURE 5










FIGURE 6




(b) *Porites*, Granitos del Oro reef, Gulf of Chiriquí



(c) *Pocillopora*, Contadora reef, Gulf of Panamá

(a) *Balanophyllia elegans*, coral for U-series dating



Middle Pleistocene, San Nicolas Island, California

(c) *Saxidomus*, bivalve for amino acid geochronology



Last interglacial, San Nicolas Island, California

(b) *Porites*, coral for U-series dating



Last interglacial, Isla Carmen, Baja California Sur

(d) *Tegula*, gastropod for amino acid geochronology



Middle Pleistocene, San Clemente Island, CA

(e) *Chione*, bivalve for amino acid geochronology



Modern *Chione*, Cholla Bay, Sonora

Last-interglacial (?) *Chione*, Cholla Bay, Sonora





FIGURE 9




FIGURE 10





FIGURE 11





FIGURE 12




FIGURE 13




FIGURE 14



FIGURE 15




FIGURE 16






FIGURE 17







FIGURE 18

(a) Newport-Yaquina Bay area



(b) Cape Arago-Coquille Point area



(c) Cape Blanco area






FIGURE 19

(a)



(b)



(c)






FIGURE 21









FIGURE 22



(b) Tomales Bay from Toms Point



(c) Terrace deposits at Toms Point



(d) Fossils in terrace deposits at Millerton Point




FIGURE 24




FIGURE 25




FIGURE 26



SOUTHWEST



**12th terrace,
(Early or middle Pleistocene)**
Elevation: ~396 m



**Gaffey terrace,
~120,000 years (MIS 5e)**
(Terrace "4" of Woodring
et al., 1946)
Outer edge: ~60 m



**Paseo Del Mar terrace,
~80,000 years (MIS 5a)**
(Terrace "2" of Woodring
et al., 1946)
Outer edge: ~45 m



**Monterey Formation
(Miocene)**



*Pacific
Ocean*

FIGURE 27

(a)



(b)




FIGURE 28






FIGURE 29



- Marine terrace deposits (brown) and terrace inner edges (black lines)
- Marine terrace fossil localities
- Warm-water species present
- Cool-water species present
- 5-10 m* Fossil locality elevation

FIGURE 30







FIGURE 32



FIGURE 33

NORTH

SOUTH

Uppermost
Golfo de California

Puerto Lobos

Puerto Libertad

Punta Cuevas

Isla Tiburon

Punta Kino

Caleta Santa Maria

Bahía San Nicolas

Bahía Magdalena

Bahía Coyote

Punta Coyote

← Amino acid ratios at equilibrium →




FIGURE 34





FIGURE 35





FIGURE 36




FIGURE 37





FIGURE 38






FIGURE 39



Chione undatella
in 120 ka deposits


- U-series, coral
- Aminostratigraphy



Dosinia ponderosa
in 120 ka deposits


- U-series, coral
- Aminostratigraphy

FIGURE 40



Distribution of *Mexacanthina lugubris* at ~120 ka:

- U-series, coral
- Aminostratigraphy



***Stramonita biserialis* in 120 ka deposits**

- U-series, coral
- Aminostratigraphy

FIGURE 41





FIGURE 42

To Dixon Entrance, Canada
~54°N to 55°N



To northern Peru,
~5°S to ~6°S




FIGURE 44




FIGURE 45