

1 **Moment tensor catalogue of earthquakes in West Bohemia from 2008** 2 **to 2018**

3
4 Václav Vavryčuk¹, Petra Adamová¹, Jana Doubravová¹, Josef Horálek¹

5
6 ¹Institute of Geophysics, Boční II/1401, 14100 Praha 4, Czech Republic

7
8 **Correspondence:** Václav Vavryčuk (vv@ig.cas.cz)

9 10 **Abstract**

11 We present a unique catalogue of full moment tensors (MTs) of earthquakes with M_L between 0.5 and 4.4 that
12 occurred in West Bohemia, Czech Republic, in the period from 2008 to 2018. The MTs were calculated from
13 vertical components of P-wave amplitudes. The MT inversion was based on the principal component analysis
14 applied to optimally filtered velocity records of local seismic stations deployed in the West Bohemia area. The
15 minimum number of inverted stations is 15 and the RMS between theoretical and observed amplitudes is lower
16 than 0.5. The catalogue is exceptional in several aspects: (1) it represents an extraordinary extensive dataset of
17 more than 5.100 MTs, (2) it covers a long period of seismicity in the studied area, during which several
18 prominent earthquake swarms took place, (3) the locations and retrieved MTs of earthquakes are of a high
19 accuracy. Additionally, we provide three-component records at the West Bohemia (WEBNET) seismic stations,
20 the velocity model in the region, and the technical specification of the stations. The dataset is ideal for being
21 utilized by a large community of researchers for various seismological purposes, e.g., for studies of (1) the
22 migration of foci and the spatiotemporal evolution of seismicity, (2) redistribution of stress during periods of
23 intense seismicity, (3) the interaction of faults, (4) the Coulomb stress along the faults and local stress
24 anomalies connected to fault irregularities, (5) diffusivity of fluids along the activated faults, or (6) the time-
25 dependent seismic risk due to the migration of seismicity in the region. In addition, the dataset is optimum for
26 developing and testing new inversions for MTs and for tectonic stress. Since most of the earthquakes are non-
27 shear, the dataset can contribute to studies of non-double-couple components of MTs and their relation to shear-
28 tensile fracturing and/or seismic anisotropy in the focal zone.

29 30 31 **1 Introduction**

32 The seismic moment tensor (MT) describes equivalent body forces acting at an earthquake source (Knopoff
33 and Randall, 1970). It is a basic quantity evaluated for earthquakes that informs us about their moment
34 magnitude, focal mechanism and type of faulting. It can be separated into double-couple (DC), isotropic (ISO)
35 and compensated linear vector dipole (CLVD) components (Jost and Hermann, 1998; Vavryčuk, 2015). The
36 DC component is produced by shear faulting in isotropic media; the ISO and CLVD components reflect
37 complexities in the earthquake source, e.g., irregularly shaped faults, seismic anisotropy, shear-tensile faulting

38 induced by fluid injection in volcanic or geothermal areas, or the presence of a material interface in the focal
39 zone (Frohlich, 1994; Julian et al. 1998; Miller et al. 1998; Růžek et al., 2003; Šílený and Milev, 2008;
40 Vavryčuk 2005, 2006, 2011a, 2013, 2015; Vavryčuk and Hrubcová 2017).

41

42 Since earthquakes do not occur separately but in sequences, it is necessary to compile high-quality MT
43 catalogues for understanding origins of seismicity, tectonic stress regime and seismic energy release of any
44 region under study. In this way, we can identify prominent periods of seismicity, trace faults and fault
45 segments, monitor migration of earthquake foci, analyse interactions of nearby or intersecting faults, and map
46 the fluid flow along the fault systems in the focal zone (Vavryčuk et al., 2021). Hence, MT catalogues are
47 fundamental sources of information for all detailed studies of seismicity on the local, regional or global scale.

48

49 In this paper, we present recordings, locations and high-quality moment tensors of 5134 earthquakes that
50 occurred in the West Bohemia geothermal region, Czech Republic in the period from 2008 to 2018. The
51 earthquakes were monitored by the West Bohemia local seismic network WEBNET (Horálek et al., 2000;
52 Fischer et al., 2010). Their locations were calculated by the double-difference location method and the moment
53 tensors were determined using the moment tensor inversion of P waves based on the principal component
54 analysis. Because of its extent and quality, the presented dataset is unique and represents an extraordinary
55 dataset, which might find exciting applications in numerous future studies.

56

57 **2 West-Bohemia seismoactive region**

58 The region of West Bohemia is located in the western part of the Bohemian Massif, where three major tectonic
59 units are merged: the Saxothuringian, the Teplá-Barrandian and the Moldanubian. The region is
60 geodynamically active exposed to the Tertiary and Quaternary volcanism associated with CO₂ emanations,
61 dry and wet mofettes, and numerous mineral springs (Kämpf et al., 2013; Hrubcová et al., 2017; Bräuer et al.,
62 2018). Two major fault systems are identified in the area: the Mariánské Lázně fault striking in the NW-SE
63 direction and the Ore-Mountain fault striking in the WSW-ENE direction (Figure 1b). The recently most active
64 fault is, however, a left-lateral strike-slip fault in the N-S direction, situated at the eastern boundary of the Cheb
65 Basin filled by up to 300 m thick Tertiary and Quaternary sediments. The seismically active faults were
66 identified at depth by clustering of hypocentres and by focal mechanisms (Vavryčuk et al., 2013), but they also
67 have some geological evidence on the surface (Bankwitz et al., 2003).

68

69 The seismic energy in the West Bohemia region is typically released in the form of earthquake swarms. The
70 occurrence of the earthquake swarms has been well documented in the region since the beginning of the 19th
71 century. A significant increase of the earthquake activity was observed at the turn of the 19th and 20th century,
72 when several larger swarms were observed. There were earthquake swarms in 1897, 1900, 1903 and 1908.
73 During the last 40 years, the seismicity occurs in the area of 40 x 50 square kilometres, but the most intense
74 seismicity is focused in the Nový Kostel zone with size of 3 x 12 square kilometres (Fischer et al., 2014;
75 Čermáková and Horálek, 2015). Foci of earthquakes in this zone are clustered along a fault striking in the
76 roughly N-S direction (Figure 1a) with depths ranging from 6 to 11 km. The duration of the earthquake swarms

77 varies; it lasts from several days for micro-swarms up to 2-3 months for the most prominent swarms. The
 78 swarms may consist of several thousands of earthquakes. The local magnitudes M_L of the earthquakes rarely
 79 exceed a value of 4.0. The strongest instrumentally recorded swarm activity occurred in 1985/86 with two
 80 main shocks having magnitudes of M_L 4.6 and 4.2.

81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124




Figure 1. (a) The map view of earthquake foci in the period from 2008 to 2018, (b) topographic map with tectonic faults (black dashed lines) and positions of stations (blue triangles), and (c) the magnitude-time plot with the colour-coded time. The red dots in (b) show the earthquake foci. The dashed rectangle around the foci in (b) defines the area shown in panel (a). The red full arrows mark the orientation of the maximum and minimum principal stress axes. The dashed-dotted line marks the boundary between the Czech Republic and Germany. The position of West Bohemia in Europe is indicated in the inset. The focal mechanisms typical for the area are also indicated.

125 **3 Monitoring system**

126 The seismic activity in the region is monitored by the local seismic network WEBNET (Figure 1b, Table 1).
 127 The network is operating since 1994 and the number of stations gradually increased (Horálek et al., 2000;
 128 Fischer et al., 2010). After its major upgrade in 2008, the WEBNET network consists of 23 seismic stations
 129 within the epicentral distance of 25 km. The stations cover the area rather uniformly with a minor azimuthal
 130 gap of 45° to the south. The three component ground-velocity records are sampled at 250 Hz and the frequency
 131 response is flat at least between 1 and 80 Hz. Until September 2014, all data were processed based on triggered
 132 records. Since the beginning of September 2014, the recordings are processed by using automatic pre-
 133 processing of continuous recordings. Another major upgrade of the network was realized in 2015. Originally,
 134 the stations were equipped by the Le-3DLite and SM3 seismometers; some of them were lately upgraded using
 135 the Guralp CMG-3ESP seismometers. The station with the nearest epicentral distance (station NKC) is
 136 additionally equipped with the broadband STS-2 seismometer. For a technical specification of the WEBNET
 137 seismic stations, see Table 1. Full information on stations is provided in files Webnet.xml (formatted according
 138 to the FDSN StationXML standard, <http://docs.fdsn.org/projects/stationxml/en/latest/>) and Webnet.dataless
 139 (formatted according to the Dataless SEED format ([https://ds.iris.edu/ds/nodes/dmc/data/formats/dataless-
 140 seed/](https://ds.iris.edu/ds/nodes/dmc/data/formats/dataless-seed/)) that are included in the dataset.

141
 142 **Table 1.** Location and instrumentation of the WEBNET seismic stations
 143

Code	Site name	Latitude (°N)	Longitude (°E)	<i>h</i> (m)	Sensor before 2015	Digitizer before 2015	Sensor after 2015	Digitizer after 2015	Note
BUBD	Bublava	50.38174	12.51362	746	LE-3DLite	Gaia	LE-3DLite	Gaia	
HOPD	Horní Paseky	50.22378	12.26547	731	LE-3DLite	Gaia	LE-3DLite	Gaia	
HRC	Hrádek	50.19348	12.53660	596	LE-3DLite	Gaia	LE-3DLite	Gaia	Out of order from 2015
HRED	Hřebeny	50.21425	12.56491	589	LE-3DLite	Gaia	LE-3DLite	Gaia	Timing problems in 2011 and 2014
HUC	Komorní Hůrka	50.09997	12.33612	480	-	-	CMG-3ESPC	Taurus	Installed in 2016, anomalous site effects
KAC	Kaceřov	50.14361	12.51708	548	SM-3	Janus-Trident	SM-3	Janus-Trident	
KOC	Kopaniny	50.26417	12.23288	621	SM-3	5800 PCM	CMG-3ESPC	Centaur	
KOPD	Kopanina	50.20319	12.47473	536	LE-3DLite	Gaia	LE-3DLite	Gaia	

KRC	Kraslice	50.33069	12.52950	806	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
KVC	Květná	50.20496	12.51134	666	SM-3	5800 PCM	CMG-3ESPC	Centaur	
LAC	Lazy	50.04967	12.62396	884	SM-3	5800 PCM	CMG-3ESPC	Centaur	
LBC	Luby	50.26461	12.41123	684	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
LOUD	Loučná	50.27753	12.57449	692	LE-3DLite	Gaia	LE-3DLite	Gaia	
NKC	Nový Kostel	50.23234	12.44706	610	SM-3 CMG-40T	5800 PCM Janus-Trident	CMG-3ESPC STS-2	Centaur	
PLED	Plesná	50.20890	12.33767	556	LE-3DLite	Gaia	LE-3DLite	Gaia	
POC	Počátky	50.31997	12.42662	841	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
POLD	Polná	50.15603	12.23497	556	LE-3DLite	Gaia	LE-3DLite	Gaia	
SKC	Skalná	50.16911	12.36050	501	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
SNED	Sněžná	50.31088	12.50131	756	LE-3DLite	Gaia	LE-3DLite	Gaia	
STC	Studenec	50.25794	12.51849	712	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
TRC	Trojmezí	50.30344	12.14466	612	LE-3DLite	Gaia	CMG-3ESPC	Centaur	
VAC	Vackov	50.23450	12.37634	581	SM-3	Janus-Trident	CMG-3ESPC	Centaur	
ZHC	Zelená Hora	50.06984	12.30810	677	CMG-40T	Janus-Trident	CMG-3ESPC	Centaur	
MAC	Chlum sv. Maří	50.14429	12.53516	609	-	-	CMG-3ESPC	Centaur	Installed in 2017


144 Quantity h means the altitude of the stations. Recording systems: Taurus – Nanometrics digitizer; Janus-Trident
145 – Nanometrics communications controller-digitizer; Centaur – Nanometrics digitizer; Gaia – Vistec digitizer;
146 5800 PCM – Lennartz digitizing system. Seismometers: SM-3 – SP sensor; LE-3DLite – Lennartz SP sensor;
147 CMG-40T – Guralp BB sensor; CMG-3ESPC – Guralp BB sensor. Station HUC has anomalous site effects
148 and it was not used in the MT inversion. Time in station HRED is erroneously shifted by 0.45 s in 2011 and
149 by 2 s in 2014.

150

151 **4 Seismicity in 2008-2018**

152 The West Bohemia region is characterized by a continuous background seismicity scattered over the whole
153 region interrupted by earthquake swarm sequences located mostly in the Nový Kostel focal zone. The most
154 intense periods of seismicity are in 2008, 2011, 2014, 2017 and 2018 (Figures 1c and 2). All these sequences

155 are typical earthquake swarms except for the seismic activity in 2014, which was exceptional. This sequence
 156 resembled a mainshock-aftershock sequence rather than the earthquake swarm (Hainzl et al., 2016; Jakoubková
 157 et al., 2018; Vavryčuk and Adamová, 2018) being formed by three pronounced activity periods. The strongest
 158 events in these periods reached magnitude significantly larger than the other events (Figure 2c). The seismic
 159 sequences differ in the earthquake productivity, in the duration, and in the number of periods of the intense
 160 seismicity (Figure 2). The strongest event in the period from 2008 to 2018 reached magnitude M_L of 4.2 and
 161 it occurred in 2014.



199 **Figure 2.** Magnitude-time plots of the major seismic sequences in the period from 2008 to 2018. According to
 200 the Bath law (Bath, 1965), the 2014 activity resembles rather a mainshock-aftershock sequence, because the
 201 difference in magnitudes ΔM_L between two strongest events in individual seismicity phases exceeds 1. In other
 202 seismic sequences, the magnitude gaps between two strongest events are not so prominent.

203

204 **5 Magnitudes and foci locations**

205 The local magnitude of earthquakes is computed from the velocity records according to the formula of Horálek
 206 et al. (2000). The locations are computed in two steps. First, initial locations were calculated by the NonLinLoc
 207 code (Lomax et al., 2009) in a layered velocity model (see Table 2) developed by Málek et al. (2005). For the
 208 locations, manual picks of the P and S arrivals were used. Second, we applied the double-difference location
 209 algorithm developed by Waldhauser and Ellsworth (2000) to differential times calculated from manual picks.
 210 The relative precision of hypocentres was less than ± 20 m within the cluster (Bouchaala et al., 2013). The
 211 absolute location of the cluster was determined with the accuracy of about ± 100 m in the horizontal plane and
 212 ± 350 m in depth (see Bouchaala et al., 2013).

213

214 The locations of foci point to complex geometry of the fault system in the focal area (Figure 3). The seismicity
 215 migrated from south to north in time and the individual seismic sequences occurred along different subfaults
 216 (Fischer et al., 2010; Bouchaala et al., 2013; Vavryčuk et al., 2013; Jakoubková et al., 2017). For example, the
 217 2008, 2011 and 2017 swarms activated three similarly oriented subfaults separated with gaps and offsets
 218 between them. The barrier between the fault segments activated in 2008 and 2011 was broken in 2014 (Hainzl
 219 et al., 2016; Vavryčuk and Adamová, 2018), and the gap between the fault segments activated in 2011 and
 220 2017 was broken during the 2018 swarm (Bachura et al., 2021; Vavryčuk et al., 2021). The overall direction
 221 of the whole fault system is defined by strike of 170° and dip of 75° . However, some fault segments may
 222 deviate from this overall direction significantly. For example, small echelon faults located at the deepest part
 223 of the fault system have strike of 305° and dip of 65° (see Figure 3c, blue dots at the depth range of 10.5-11
 224 km).

225

226

227

Table 2. The layered velocity model

Depth (km)	0.0	0.2	0.5	1.0	2.0	4.0	6.0	10.0	20.0	32.0
v_P (km/s)	4.30	5.06	5.33	5.60	5.87	6.09	6.35	6.74	7.05	7.25
Q_P	30	40	50	60	80	100	150	200	300	400

228

229

Ratio v_P/v_S is 1.70 and ratio Q_P/Q_S is 2.

230

231

232

233

234

235

236

237

238




Figure 3. The earthquakes foci with local magnitude $M_L \geq 0.5$ in (a) 3D view, (b) cross-section vertical view, and (c) in-plane vertical view. The foci are colour coded according to time: dark blue – 2008, light blue – 2011, yellow – 2014, red – 2017, and brown – 2008.

6 Moment tensors

6.1 MT inversion of earthquakes

The MT inversion requires accurate locations of earthquakes, an accurate crustal velocity model, dense coverage of stations on the focal sphere and low seismic noise (Šílený, 2009; Ford et al., 2010; Stierle, Bohnhoff, et al., 2014; Stierle, Vavryčuk, et al., 2014). We can invert amplitudes of seismic phases, amplitude ratios or full waveforms (Dreger and Woods, 2002; Cesca et al., 2006; Sokos and Zahradník, 2008; Cesca and Dahm, 2008; Vavryčuk et al., 2008; Zahradník et al., 2008; Fojtíková et al., 2010; Kwiątek et al., 2016;

288 Jechumtálová and Šílený, 2005; Vavryčuk and Kühn, 2012; Kühn and Vavryčuk, 2013; Yu et al., 2018, 2019).
289 The applicability of the individual MT inversions is specific, depending on the magnitude of analysed
290 earthquakes, predominant wave frequencies and epicentral distances of stations. MTs of moderate or large
291 earthquakes are usually calculated from full waveforms recorded at regional or global seismic networks. By
292 contrast, MTs of small earthquakes and microearthquakes are commonly calculated from amplitudes of P
293 and/or S waves picked in short-period seismograms recorded at local networks. In this way, the sensitivity of
294 the MT inversion to small-scale complexities of the local geological structure are suppressed and a
295 computationally demanding modelling of high-frequency full waveforms is avoided.

296

297 The inversion for MTs of small earthquakes and microearthquakes is challenging for several reasons: (1) the
298 waveforms are complex due to high frequencies and noise, and (2) the datasets are extensive with thousands
299 of events, which require a semi- or fully-automated processing. Here, the MT inversion developed by Vavryčuk
300 et al. (2017) is applied. The inversion is based on the principal component analysis (PCA), which transforms
301 correlated waveforms into a set of the so called principal components (see Figure 4). The first component has
302 the highest variance and reproduces a so-called ‘common wavelet’, i.e., a wavelet with the highest similarity
303 with all analysed traces. This common wavelet physically represents a signal radiated by the earthquakes
304 source, which can be distorted during its propagation from the source to the receiver by inhomogeneities in
305 the geological structure, site effects or seismic noise.

306

307 Subsequently, the common wavelet is correlated with individual recorded traces and the effective P-wave
308 amplitudes are calculated as the amplification factors applied to the common wavelet, in order to optimally
309 reproduce the recorded traces. The obtained amplitudes are inverted for the MTs using the generalized linear
310 inversion (Lay and Wallace, 1995). The Green’s function amplitudes are computed by the ray method
311 (Červený, 2001; Vavryčuk, 1999, 2008) and incorporated the effects of the Earth’s surface. An inhomogeneous
312 medium with a vertical gradient obtained by smoothing the layered model of Málek et al. (2005) was applied
313 for computing the rays by the ray-tracing algorithm. The inversion is robust, fast and insensitive to noise in
314 data.

315


316

317 **6.2 Individual steps of the MT inversion**

318 The MT inversion consists of data pre-processing, alignment of traces, computation of the effective amplitudes
319 using the PCA method and the MT inversion. The data are not corrected for the frequency response of sensors,
320 because the response is flat for all sensors at least from 2 to 60 Hz. The individual steps of the inversion are as
321 follows (see Figure 5):

- 322 1. Data pre-processing, which comprises: (a) an oversampling of records in order to perform an accurate
323 alignment of waveforms, (b) band-pass filtering to enhance the signal-to-noise ratio, and (c) a rough
324 alignment of waveforms using manual picks, if available, or using an automatic picking algorithm called
325 the Suspension Bridge Picking (SBPx), see FeedMeImATroll (2021).

326 2. Two-step accurate alignment of waveforms, which comprises: (a) an alignment of waveforms using the
 327 cross-correlation with the waveform of the highest signal-to-noise ratio, (b) calculation of the first principal
 328 component from the aligned waveforms, (c) another alignment of waveforms using the cross-correlation
 329 with the computed first principal component, and (d) calculation of the refined first principal component
 330 from the aligned waveforms.



365 **Figure 4.** Example of the MT inversion of the earthquake on 24 May 2014 at 16:14:30 with ML 2.1. (a) Whole
 366 velocity records; (b) window with aligned P waves; (c) the common wavelet (red line) together with the P-
 367 wave traces at individual stations (blue lines); (d) the common wavelet represented by the first principal
 368 component (red line) and noise in waveforms represented by the second principal component (black line). The
 369 polarities of the P-wave in panel (c) are switched to be consistent with the polarity of the common wavelet.
 370 The inset in plot (a) shows the focal mechanism and positions of stations on the focal sphere (red circles mark
 371 negative polarities, and blue plus signs mark the positive polarities).

- 372 3. Calculation of the PCA amplitudes and weights in the MT inversion, which comprises: (a) calculation of
 373 the PCA coefficients of the first principal component, which serve as the effective amplitudes used in the
 374 MT inversion, (b) calculation of the correlation coefficients between individual traces and the first principal
 375 component, which serve as the weights in the linear MT inversion scheme (in this way, a station with a
 376 waveform significantly different from the common wavelet suppressed in the inversion),
- 377 4. Repeated MT inversion for several alternative band-pass filters and time windows, in order to adapt the
 378 inversion to earthquakes with a varying frequency content. The inversion is firstly run with the whole set
 379 of stations, and secondly with eliminating two stations producing the largest misfits in the inversion.

380 In this way, we obtain a set of candidate MTs. The optimum MT is that with the minimum root-mean-squares
 381 (RMS) of differences between the synthetic amplitudes A^{synth} and the observed amplitudes A^{obs}

$$382 \text{ RMS} = \frac{\sqrt{\sum_{i=1}^N (A_i^{synth} - A_i^{obs})^2}}{\sqrt{\sum_{i=1}^N (A_i^{synth})^2}}, \quad (1)$$

383 where N is the number of stations. The optimum MT is normalized and expressed in a relative scale, because
 384 it is computed from wave amplitudes but not from full displacement records. The scalar moment is obtained
 385 by integrating the common (displacement) wavelet. The optimum MTs were further decomposed into the DC,
 386 ISO and CLVD components according to Equations (6-10) of Vavryčuk (2015).




Figure 5. Flowchart of the PCA moment tensor inversion.

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465




Figure 6. Examples of focal mechanisms and histograms of the CLVD and ISO errors. (a) Earthquake on 1 September, 2011 at 12:54:05.7 with $ML = 0.6$, and (b) earthquake on 11 May, 2018 at 06:26:09.0 with $ML = 2.3$.




Figure 7. Inversion for the full MT solution ('Non-DC MT' and 'Non-DC misfits') and for the DC solution ('DC MT' and 'DC misfits') for earthquakes in Figure 6. The mean amplitude misfits for the full MT and DC solutions are shown by blue and red dashed lines, respectively.

466
 467 In order to estimate errors of the MTs, the inversion is performed for each MT repeatedly 100 times using
 468 amplitudes distorted by noise characterized by a flat probability distribution. The level of noise ranges from
 469 -25% to 25% of the inverted amplitude at each trace. The scatter of the solutions served for estimating: (1) the
 470 mean errors in the P/T axes directions calculated as the mean of deviations between the directions of the P/T
 471 axes of noise-free solution and the noisy solutions, (2) the mean errors in the percentages of the DC, ISO and
 472 CLVD components calculated as the standard deviations of the DC, ISO and CLVD values of noisy MT
 473 solutions.

474
 475 Figure 6 exemplifies the MT inversion for two earthquakes, which display significant non-DC components.
 476 The histograms of the CLVD and ISO errors indicate that the ISO component is always better constrained than
 477 the CLVD component. Nevertheless, despite the numerical errors produced by the inversion, the histograms
 478 prove that both the events contain also true non-DC components. This is also confirmed by a comparison of
 479 fits for the full MTs and for the DC solutions for the events shown in Figure 7. The figure indicates that the
 480 misfits for the full MT solutions are almost twice lower than those for the DC solutions. This proves that at
 481 least some part of the non-DC components retrieved by the MT inversion should be of physical origin.

482
 483

484 **7 Basic characteristics of the MT catalogue**

485 Firstly, we processed all events with the local magnitude larger than 0.5. After that, we checked manually the
 486 quality of input data and the retrieved MT and we excluded earthquakes: (1) recorded at a low number of
 487 stations ($N < 15$), (2) with extremely low signal-to-noise ratio, (3) produced unstable moment tensors with
 488 anomalously high RMS ($RMS > 0.5$). In this way, we obtained a dataset of 5134 earthquakes listed in the
 489 catalogue. Table 3 summarizes the numbers of events in individual years. The magnitude-frequency
 490 distribution of the analysed events is shown in Figure 8.

491
 492
 493
 494

Table 3. Number of reported events for each year.

Year	2008	2009	2010	2011	2012	2013
Number of events	990	38	24	1211	69	201
Year	2014	2015	2016	2017	2018	2008-2018
Number of events	831	36	32	580	1122	5134

495
 496
 497
 498
 499

500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545




Figure 8. Cumulative (blue) and non-cumulative (red) magnitude-frequency distribution of the analysed earthquakes.

The earthquakes inverted for MTs were recorded mostly by 20 or more stations (Figure 9, middle column). The RMS varied during the whole period and ranged mostly from 0 to 0.5 (Figure 9, right column). The MTs with RMS higher than 0.5 were considered as unreliable. The variation of the RMS in time is probably produced by varying station coverage due to the foci migration. The P/T axes form compact and non-overlapping clusters for all seismic sequences in the studied time period (Figure 9). The position of clusters slightly differs in individual years and indicates some stress variation in the focal zone. Directions of the P/T axes are well resolved with the mean standard deviation less than 2° (Figure 10, two left columns). The errors of the ISO and CLVD components are mostly about 1.5-2% and 5-6%, respectively (Figure 10, two right columns). Comparing these errors for individual activity periods, we see that the errors tend to slightly decrease with time. This might be due to a continuously increasing quality of the WEBNET network. The histograms of the standard deviations of the P/T axes and the ISO and CLVD errors for the whole period from 2008-2018 are shown in Figure 11.

546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595




Figure 9. The P/T axes (left-hand plots), histograms of the RMS of the number of stations used in the MT inversion (middle plots), and histograms of the RMS of the retrieved MTs (right-hand plots) for seismic activities in 2008 (a), 2011(b), 2014 (c), 2017 (d) and 2018 (e). N denotes the number of stations, which recorded the individual earthquakes.

596
597


598
599
600
601
602
603
604
605

606
607
608
609
610
611
612
613
614

615
616
617
618
619
620
621
622
623

624
625
626
627
628
629
630
631
632

633
634
635
636
637
638
639
640
641
642



643 **Figure 10.** Histograms of mean deviations of the P/T axes and histograms of the ISO and CLVD standard
644 errors for MTs of earthquakes from individual prominent seismic activities: in 2008 (a), 2011(b), 2014 (c),
645 2017 (d) and 2018 (e). The mean P/T deviations and the ISO and CLVD standard errors were calculated for
646 each event from 100 MTs inverted using randomly generated noisy data.




Figure 11. Histograms of mean deviations of the P/T axes (a-b) and histograms of the standard ISO and CLVD errors (c-d) for the 5134 reported MTs. The mean P/T deviations and the ISO and CLVD standard errors were calculated for each event from 100 MTs inverted using randomly generated noisy data.

8 Description of the dataset

The dataset consists of the following directories:

- Waveforms – this directory is further structured into subdirectories according to individual years and earthquakes. Three-component velocity records are stored in ASCII files with four columns (time + 3 components: Z,N,E) individually for each station and each earthquake. The first line of the files contains time of the first sample. The pre-event time before the P-wave arrival is 2s.
- Model – this directory contains the ASCII file ‘model.crust’, which defines the layered velocity model for the West Bohemia region (depth in km, P-wave velocity in km/s, v_P/v_S ratio, P-wave quality factor Q_P and ratio Q_P/Q_S of P-wave and S-wave quality factors).
- Stations – this directory contains the ASCII file ‘Webnet_coordinates.dat’ with coordinates of stations (site, name of the station, latitude, longitude, elevation), file ‘Webnet.xml’ with a full technical specification of stations formatted according to the FDSN StationXML standard (<http://docs.fdsn.org/projects/stationxml/en/latest/>) and file ‘Webnet.dataless’ with a full technical

691 specification of stations formatted according to the Dataless SEED format
692 (<https://ds.iris.edu/ds/nodes/dmc/data/formats/dataless-seed/>).

- 693 • Moments – this directory contains the ASCII file ‘catalogue_2008-2018.dat’ with double difference
694 locations, magnitudes, moment tensors and their errors, RMS and the numbers of inverted stations.
- 695 • Figures – this directory is further structures into subdirectories according to individual years. Four
696 figures are provided for each earthquake in the .pdf format (see Figure 12): complete waveforms of
697 vertical components, a detail of the P-waveforms, the focal mechanism with positions of stations, and
698 the RMS at individual stations.

699

700 File ‘catalogue_2008-2018.dat’ lists the following quantities for each earthquake:

- 701 • Event identification (composed form year and the sequential number of the event in the year, e.g.
702 2008-216)
- 703 • Double-difference locations
 - 704 • Origin time (year, day, hour, minute, second)
 - 705 • Latitude (°N)
 - 706 • Longitude (°E)
 - 707 • Depth (km)
- 708 • Local magnitude M_L (calculated according to Horálek et al., 2000)
- 709 • N – number of stations used in the MT inversion
- 710 • Frequencies f_1 a f_2 (in Hz) – low and high corner frequencies of the optimum Butterworth 4th-order
711 band-pass filter
- 712 • RMS – for its definition, see Equation (1)
- 713 • Moment magnitude M_w
- 714 • Components of the normalized moment tensor: $M_{11}, M_{12}, M_{13}, M_{22}, M_{23}, M_{33}$ (x_1 – North, x_2 – East, x_3
715 – down). The moment tensor is normalized using the Euclidean norm (see Equation 17 of Vavryčuk,
716 2015)
- 717 • Strike1, dip1, rake1, strike2, dip2, rake2 (in °)
- 718 • DC, CLVD, ISO (in %, calculated according to Equations 6-10 of Vavryčuk, 2015)
- 719 • Errors of DC, CLVD, ISO (in %, for the definition of errors, see the text)
- 720 • Deviations of the P/T axes (in °, for the definition of errors, see the text)

721

722

723

724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770




Figure 12. Example of plots provided for each earthquake in the dataset. (a) Vertical components of complete waveforms recorded at the WEBNET stations and aligned according to the arrival time of the P wave. Stations are sorted according to their epicentral distance. (b) Vertical components of the P waves aligned according to their arrival time and with a polarity switched according to the polarity of the common wavelet. (c) The focal mechanism with positions of the stations on the focal sphere (negative polarities – red circles, positive polarities – blue plus signs). (d) Root-mean-squares (RMS) of the differences between the theoretical and observed amplitudes of the P waves.

9 Discussion and conclusions

We publish a unique catalogue of moment tensors of earthquakes that occurred in the West Bohemia in the period from 2008 to 2018. The catalogue is exceptional in several aspects: (1) it represents an extraordinary extensive dataset of more than 5.000 MTs, (2) it covers a long period of seismicity in the studied area, during which several prominent earthquake swarms took place, (3) the foci locations and retrieved MTs are of a very high accuracy. In addition, the three-component velocigrams recorded at the WEBNET stations together with the velocity model in the region and the technical specification of stations are provided. This predetermines the dataset to be utilized by a large community of researchers for various seismological purposes.

771

772 The great potential of the dataset or its subsets has so far been proved in studies of origins of the swarm activity
773 in this area (Horálek and Fischer, 2008; Fischer et al., 2010; Fischer et al., 2014), migration of seismicity in
774 time due to fluid flow and/or stress redistribution in the focal zone (Hainzl et al, 2012, 2016; Vavryčuk and
775 Hrubcová, 2017), changes of the v_p/v_s ratio in the focal zone (Dahm & Fischer, 2014; Bachura & Fischer,
776 2016), identification of fault segments and their mutual interaction (Vavryčuk and Adamová, 2018; Vavryčuk
777 et al., 2021), the fault instability (Vavryčuk, 2011b, 2014), differences in the seismic energy release in
778 earthquake swarms and mainshock-aftershock sequences (Čermáková and Horálek, 2015; Vavryčuk and
779 Adamová, 2018), the efficiency of new moment tensor inversion algorithms such as the MT inversion based
780 on the PCA (Vavryčuk et al., 2017), the MT inversion using the empirical Green's functions (Vavryčuk and
781 Adamová, 2020) or the MT inversion using the network calibration (Davi and Vavryčuk, 2012; Davi et al.,
782 2013). The provided records were also utilized in a study of seismic anisotropy based on the analysis of shear-
783 wave splitting (Vavryčuk and Boušková 2008), identification of shallow discontinuities in the Earth's crust
784 (Hrubcová et al., 2016), lateral variation of depth of the Moho discontinuity (Hrubcová et al., 2013, 2017), and
785 for detailed mapping of the non-DC components of MTs and shear-tensile fracturing in the Nový Kostel focal
786 zone (Vavryčuk, 20011a; Vavryčuk et al., 2021).

787

788 The dataset is ideal for being utilized in many other studies in future, e.g., for studies of (1) the interaction
789 between the scattered background regional seismicity and the swarm seismicity focused in the Nový Kostel
790 zone, (2) the Coulomb stress and local stress anomalies connected to fault irregularities, (3) diffusivity of fluids
791 along the activated faults, or (4) time-dependent seismic risk due to the migration of seismicity in the region.
792 In addition, the dataset is optimum for developing and testing new MT inversions (Šílený and Vavryčuk, 2000,
793 2002), stress inversions, and for the spatiotemporal evolution of tectonic stress. Since most of the earthquakes
794 are non-shear, the dataset can contribute to studies of the non-DC components and their relation to shear-tensile
795 fracturing and/or seismic anisotropy in the focal zone (Vavryčuk, 1997; Vavryčuk and Boušková, 2008).

796

797

798 **Data availability**

799 The MT catalogue and waveforms are available at the Mendeley Data Repository
800 <https://doi.org/10.17632/9pwy7rgzkt.2> (Vavryčuk et al., 2022a) and at the International Seismological
801 Centre (ISC) Dataset Repository <https://doi.org/10.31905/H212Z6OX> (Vavryčuk et al., 2022b).

802

803

804 **Acknowledgements**

805 The study was supported by the Grant Agency of the Czech Republic, Grant No. 19-06422S. We thank
806 Grzegorz Kwiatek and one anonymous reviewer for their helpful reviews. The moment tensor decomposition
807 was performed using public open Matlab code MT_DECOMPOSITION ([https://www.ig.cas.cz/en/mt-
808 decomposition/](https://www.ig.cas.cz/en/mt-decomposition/)).

810 **References**

- 811 Bachura, M., and Fischer, T.: Detailed velocity ratio mapping during the aftershock sequence as a tool to
812 monitor the fluid activity within the fault plane, *Geophys. J. Int.*, 453, 215-222, 2016.
- 813 Bachura, M., Fischer, T., Doubravová, J., and Horálek, J.: From earthquake swarm to a main shock-
814 aftershocks: the 2018 activity in West Bohemia/Vogtland, *Geophys. J. Int.*, 224(3), 1835-1848, 2021.
- 815 Bankwitz, P., Schneider, G., Kämpf, H., and Bankwitz, E.: Structural characteristics of epicentral areas in
816 Central Europe: study case Cheb Basin (Czech Republic), *J. Geodyn.*, 35, 5-32.
817 [https://doi.org/10.1016/S0264-3707\(02\)00051-0](https://doi.org/10.1016/S0264-3707(02)00051-0), 2003.
- 818 Bath, M.: Lateral inhomogeneities in the upper mantle, *Tectonophysics*, 2, 483–514, 1965.
- 819 Bouchaala, F., Vavryčuk, V., and Fischer, T.: Accuracy of the master-event and double-difference locations:
820 Synthetic tests and application to seismicity in West Bohemia, Czech Republic, *J. Seismol.*, 17(3), 841-
821 859, <https://doi.org/10.1007/s10950-013-9357-4>, 2013.
- 822 Bräuer, K., Kämpf, H., Niedermann, S., and Strauch, G.: Monitoring of helium and carbon isotopes in the
823 western Eger Rift area (Czech Republic): Relationships with the 2014 seismic activity and indications for
824 recent (2000-2016) magmatic unrest, *Chem. Geol.*, 482, 131-145,
825 <https://doi.org/10.1016/j.chemgeo.2018.02.017>, 2018.
- 826 Čermáková, H., and Horálek, J.: The 2011 West Bohemia (Central Europe) earthquake swarm compared with
827 the previous swarms of 2000 and 2008, *J. Seismol.*, 19, 899–913. <https://doi.org/10.1007/s10950-015-9502-3>, 2015.
- 829 Červený, V.: *Seismic Ray Theory*. Cambridge University Press, Cambridge, 2001.
- 830 Cesca, S., and Dahm, T.: A frequency domain inversion code to retrieve time-dependent parameters of very
831 long period volcanic sources, *Comput. Geosci.*, 34(3), 235-246, 2008.
- 832 Cesca, S., Buforn, E., and Dahm, T.: Amplitude spectra moment tensor inversion of shallow earthquakes in
833 Spain, *Geophys. J. Int.*, 166, 839-854, 2006.
- 834 Dahm, T., and Fischer, T.: Velocity ratio variations in the source region of earthquake swarms in NW Bohemia
835 obtained from arrival time double- differences, *Geophys. J. Int.*, 196(2), 957-970, 2014.
- 836 Davi, R., and Vavryčuk, V.: Seismic network calibration for retrieving accurate moment tensors, *Bull. Seism.*
837 *Soc. Am.*, 102(6), 2491-2506, <https://doi.org/10.1785/0120110344>, 2012.
- 838 Davi, R., Vavryčuk, V., Charalampidou, E.-M., and Kwiątek, G.: Network sensor calibration for retrieving
839 accurate moment tensors of acoustic emissions, *Int. J. Rock Mech. Mining Sci.*, 62, 59-67,
840 <https://doi.org/10.1016/j.ijrmms.2013.04.004>, 2013.
- 841 Dreger, D., and Woods, B.: Regional distance seismic moment tensors of nuclear explosions, *Tectonophysics*
842 356, 139-156, 2002.
- 843 Fischer, T., Horálek, J., Michálek, J., and Boušková, A.: The 2008 West Bohemia earthquake swarm in the
844 light of the WEBNET network, *J. Seismol.*, 14, 665–682, 2010.
- 845 Fischer, T., Horálek, J., Hrubcová, P., Vavryčuk, V., Bräuer, K., and Kämpf, H.: Intra-continental earthquake
846 swarms in West-Bohemia and Vogtland: a review, *Tectonophysics*, 611, 1-27.
847 <https://doi.org/10.1016/j.tecto.2013.11.001>, 2014.
- 848 Fojtíková, L., Vavryčuk, V., Cipciar, A., and Madarás, J.: Focal mechanisms of micro-earthquakes in the Dobrá
849 Voda seismoactive area in the Malé Karpaty Mts. (Little Carpathians), Slovakia, *Tectonophysics*, 492,
850 213-229, <https://doi.org/10.1016/j.tecto.2010.06.007>, 2010.
- 851 Fojtíková, L., and Zahradník, J.: A new strategy for weak events in sparse networks: The first-motion polarity
852 solutions constrained by single-station waveform inversion, *Seismol. Res. Lett.*, 85(6), 1265-1274,
853 <https://doi.org/10.1785/0220140072>, 2014.
- 854 Ford, S.R., Dreger, D.S., and Walter, W.R.: Network sensitivity solutions for regional moment-tensor
855 inversions, *Bull. Seism. Soc. Am.*, 100(5A), 162-1970, 2010.

856 FeedMeImATroll: Suspension Bridge Picking Algorithm (SBPx)
857 ([https://www.mathworks.com/matlabcentral/fileexchange/51996-suspension-bridge-picking-algorithm-](https://www.mathworks.com/matlabcentral/fileexchange/51996-suspension-bridge-picking-algorithm-sbpx)
858 [sbpx](https://www.mathworks.com/matlabcentral/fileexchange/51996-suspension-bridge-picking-algorithm-sbpx)), MATLAB Central File Exchange, Retrieved September 21, 2021.

859 Frohlich, C.: Earthquakes with non-double-couple mechanisms, *Science*, 264(5160), 804-809, 1994.

860 Hainzl, S., Fischer, T., and Dahm, T.: Seismicity-based estimation of the driving fluid pressure in the case of
861 swarm activity in Western Bohemia, *Geophys. J. Int.*, 191(1), 271-278, 2012.

862 Hainzl, S., Fischer, T., Čermáková, H., Bachura, M., and Vlček, J.: Aftershocks triggered by fluid intrusion:
863 Evidence for the aftershock sequence occurred 2014 in West Bohemia/Vogtland, *J. Geophys. Res.*, 121(4),
864 2575–2590, 2016.

865 Horálek J., Fischer T., Boušková A., and Jedlička P.: The Western Bohemia/Vogtland region in the light of the
866 WEBNET network, *Stud. Geophys. Geod.*, 44(2), 107–125, 2000.

867 Horálek, J., and Fischer, T.: Role of crustal fluids in triggering the West Bohemia/Vogtland earthquake swarms:
868 just what we know (a review), *Stud. Geophys. Geod.*, 52, 455–478, 2008.

869 Hrubcová, P., Vavryčuk, V., Boušková, A., and Horálek, J.: Moho depth determination from waveforms of
870 microearthquakes in the West Bohemia/Vogtland swarm area, *J. Geophys. Res.*, 118, 1–17,
871 <http://dx.doi.org/10.1029/2012JB009360>, 2013.

872 Hrubcová, P., Vavryčuk, V., Boušková, A., and Bohnhoff, M.: Shallow crustal discontinuities inferred from
873 waveforms of microearthquakes: Method and application to KTB drill site and West Bohemia swarm area,
874 *J. Geophys. Res., Solid Earth*, 121, 881-902, <https://doi.org/10.1002/2015JB012548>, 2016.

875 Hrubcová, P., Geissler, W.H., Bräuer, K., Vavryčuk, V., Tomek, Č., and Kämpf, H.: Active magmatic
876 underplating in western Eger Rift, Central Europe, *Tectonics*, 36, <https://doi.org/10.1002/2017TC004710>,
877 2017.

878 Jakoubková, H., Horálek, J., and Fischer, T.: 2014 mainshock-aftershock activity versus earthquake swarms in
879 West Bohemia, Czech Republic, *Pure Appl. Geophys.*, 175(1), 109-131, [https://doi.org/10.1007/s00024-](https://doi.org/10.1007/s00024-017-1679-7)
880 [017-1679-7](https://doi.org/10.1007/s00024-017-1679-7), 2018.

881 Jechumtálová, Z., and Šílený, J.: Amplitude ratios for complete moment tensor retrieval, *Geophys. Res. Lett.*,
882 32, L22303, 2005.

883 Jost, M.L., and Hermann, R.B.: A student's guide to and review of moment tensors, *Seismol. Res. Lett.*, 60,
884 37-57, 1989.

885 Julian, B.R., Miller, A.D., and Foulger, G.R.: Non-double-couple earthquakes 1: Theory, *Rev. Geophys.*, 36,
886 525–549, 1998.

887 Kämpf, H., Bräuer, K., Schumann, J., Hahne, K., and Strauch, G.: CO₂ discharge in an active, non-volcanic
888 continental rift area (Czech Republic): Characterisation (δ C-13, He-3/He-4) and quantification of
889 diffuse and vent CO₂ emissions, *Chem. Geol.*, 339, 71-83,
890 <https://doi.org/10.1016/j.chemgeo.2012.08.005>, 2013.

891 Knopoff, L., and Randall, M.J.: The compensated linear vector dipole: A possible mechanism for deep
892 earthquakes, *J. Geophys. Res.*, 75, 4957-4963, 1970.

893 Kühn, D., and Vavryčuk, V.: Determination of full moment tensors of microseismic events in a very
894 heterogeneous mining environment, *Tectonophysics*, **589**, 33-43,
895 <https://doi.org/10.1016/j.tecto.2012.12.035>, 2013.

896 Kwiatek, G., Martínez-Garzón, P., and Bohnhoff, M.: HybridMT: A MATLAB/Shell environment package for
897 seismic moment tensor inversion and refinement, *Seismol. Res. Lett.*, 87(4), 964-976, 2016.

898 Lay, T., and Wallace, T.C.: *Modern Global Seismology*, Academic Press, New York, 1995.

899 Lomax, A., Michelini, A., and Curtis, A.: Earthquake location, direct, global-search methods, in *Complexity*
900 *In Encyclopedia of Complexity and System Science, Part 5*, Springer, New York, pp. 2449-2473,
901 <https://doi.org/10.1007/978-0-387-30440-3>, 2009.

902 Málek, J., Horálek, J., and Janský, J.: One-dimensional qP-wave velocity model of the upper crust for the West
903 Bohemia/Vogtland earthquake swarm region, *Stud. Geophys. Geod.*, 49, 501-524, 2005.

904 Miller, A.D., Foulger, G.R., and Julian, B.R.: Non-double-couple earthquakes 2: Observations, *Rev. Geophys.*,
905 36, 551-568, 1998.

906 Růžek, B., Vavryčuk, V., Hrubcová, P., Zedník, J. and Celebration Working Group: Crustal anisotropy in the
907 Bohemian Massif, Czech Republic: Observations based on Central European Lithospheric Experiment
908 Based on Refraction (CELEBRATION) 2000, *J. Geophys. Res.*, 108, B8, art. no. 2392, doi:
909 10.1029/2002JB002242, 2003.

910 Šílený, J.: Resolution of non-double-couple mechanisms: Simulation of hypocenter mislocation and velocity
911 structure mismodeling, *Bull. Seism. Soc. Am.*, 99(4), 2265-2272, 2009.

912 Šílený, J., and Milev, A.: Source mechanism of mining induced seismic events - Resolution of double couple
913 and non double couple models, *Tectonophysics*, 456(1-2), 3-15, 2008.

914 Šílený, J., and Vavryčuk, V.: Approximate retrieval of the point source in anisotropic media: numerical
915 modelling by indirect parametrization of the source, *Geophys. J. Int.*, 143, 700-708.
916 <https://doi.org/10.1046/j.1365-246X.2000.00256.x>, 2000.

917 Šílený, J., and Vavryčuk, V.: Can unbiased source be retrieved from anisotropic waveforms by using an
918 isotropic model of the medium? *Tectonophysics*, 356, 125-138, [https://doi.org/10.1016/S0040-](https://doi.org/10.1016/S0040-1951(02)00380-3)
919 1951(02)00380-3, 2002.

920 Sokos, E., and Zahradník, J.: ISOLA — A Fortran code and Matlab GUI to perform multiple point source
921 inversion of seismic data, *Comput. Geosci.*, 34, 967–977, 2008.

922 Stierle, E., Vavryčuk, V., Šílený, J., and Bohnhoff, M.: Resolution of non-double-couple components in the
923 seismic moment tensor using regional networks: 1. A synthetic case study, *Geophys. J. Int.*, 196(3),
924 1869-1877, <https://doi.org/10.1093/gji/ggt502>, 2014.

925 Stierle, E., Bohnhoff, M., and Vavryčuk, V.: Resolution of non-double-couple components in the seismic
926 moment tensor using regional networks: 2. Application to aftershocks of the 1999 Mw 7.4 Izmit
927 earthquake, *Geophys. J. Int.*, 196(3), 1878-1888, <https://doi.org/10.1093/gji/ggt503>, 2014.

928 Vavryčuk, V.: Elastodynamic and elastostatic Green tensors for homogeneous weak transversely isotropic
929 media, *Geophys. J. Int.*, 130(3), 786-800. <https://doi.org/10.1111/j.1365-246X.1997.tb01873.x>, 1997.

930 Vavryčuk, V.: Weak-contrast reflection/transmission coefficients in weakly anisotropic elastic media: *P*-wave
931 incidence, *Geophys. J. Int.*, 138, 553-562, doi:10.1046/j.1365-246X.1999.00890.x, 1999. Erratum,
932 *Geophys. J. Int.*, 140 (2000), 248, 1999.

933 Vavryčuk, V.: Spatially dependent seismic anisotropy in the Tonga subduction zone: a possible contributor to
934 the complexity of deep earthquakes, *Phys. Earth Planet. Inter.*, 155, 63-72.
935 <https://doi.org/10.1016/j.pepi.2005.10.005>, 2006.

936 Vavryčuk, V.: Real ray tracing in anisotropic viscoelastic media, *Geophys. J. Int.*, 175, 617-626, doi:
937 10.1111/j.1365-246X.2008.03898.x, 2008.

938 Vavryčuk, V.: Tensile earthquakes: Theory, modeling, and inversion, *J. Geophys. Res., Solid Earth*, 116(B12),
939 B12320. <https://doi.org/10.1029/2011JB008770>, 2011a.

940 Vavryčuk, V.: Principal earthquakes: Theory and observations from the 2008 West Bohemia swarm, *Earth
941 Planet. Sci. Lett.*, 305, 290-296, <https://doi.org/10.1016/j.epsl.2011.03.002>, 2011b.

942 Vavryčuk, V.: Is the seismic moment tensor ambiguous at a material interface? *Geophys. J. Int.*, 194(1), 395-
943 400, <https://doi.org/10.1093/gji/ggt084>, 2013.

944 Vavryčuk, V.: Iterative joint inversion for stress and fault orientations from focal mechanisms, *Geophys. J. Int.*,
945 199(1), 69-77, <https://doi.org/10.1093/gji/ggu224>, 2014.

946 Vavryčuk, V.: Moment tensor decompositions revisited, *J. Seismol.*, 19(1), 231-252,
947 <https://doi.org/10.1007/s10950-014-9463-y>, 2015.

948 Vavryčuk, V., and Adamová, P.: Detection of stress anomaly produced by interaction of compressive fault steps
949 in the West Bohemia swarm region, Czech Republic, *Tectonics*, 37, 4212-4225.
950 <https://doi.org/10.1029/2018TC005163>, 2018.

951 Vavryčuk, V., Adamová, P., Doubravová, J., and Jakoubková, H.: Moment tensor inversion based on the
952 principal component analysis of waveforms: Method and application to microearthquakes in West
953 Bohemia, Czech Republic, *Seismol. Res. Lett.*, 88(5), 1303-1315, <https://doi.org/10.1785/0220170027>,
954 2017.

955 Vavryčuk, V., Adamová, P., Doubravová, J., and Ren, Y.: Mapping stress and fluids on faults by nonshear
956 earthquakes, *J. Geophys. Res.: Solid Earth*, 126, e2020JB021287, <https://doi.org/10.1029/2020JB021287>,
957 2021.

958 Vavryčuk, V., Adamová, P., Doubravová, J., and Horálek, J.: WEBNET moment tensor catalogue 2008-2018,
959 *Mendeley Data*, V2, <https://doi.org/10.17632/9pwy7rgzkt.2>, 2022a.

960 Vavryčuk, V., Adamová, P., Doubravová, J., and Horálek, J.: WEBNET moment tensor catalogue 2008-2018,
961 *ISC Seismological Dataset Repository*, <https://doi.org/10.31905/H212Z6OX>, 2022b.

962 Vavryčuk, V., and Boušková, A.: S-wave splitting from records of local micro-earthquakes in West
963 Bohemia/Vogtland: An indicator of complex crustal anisotropy, *Stud. Geophys. Geod.*, 52, 631-650,
964 <https://doi.org/10.1007/s11200-008-0041-z>, 2008.

965 Vavryčuk, V., Bouchaala, F., and Fischer, T.: High-resolution fault image from accurate locations and focal
966 mechanisms of the 2008 swarm earthquakes in West Bohemia, Czech Republic, *Tectonophysics*, 590,
967 189-195, <https://doi.org/10.1016/j.tecto.2013.01.025>, 2013.

968 Vavryčuk, V., and Hrubcová, P.: Seismological evidence of fault weakening due to erosion by fluids from
969 observations of intraplate earthquake swarms, *J. Geophys. Res.*, 122,
970 <https://doi.org/10.1002/2017JB013958>, 2017.

971 Vavryčuk, V., and Kühn, D.: Moment tensor inversion of waveforms: a two-step time-frequency approach,
972 *Geophys. J. Int.*, 190, 1761-1776, <https://doi.org/10.1111/j.1365-246X.2012.05592.x>, 2012.

973 Waldhauser F. and Ellsworth W. L.: A double-difference earthquake location algorithm: Method and
974 application to the northern Hayward fault, California, *Bull. Seismol. Soc. Am.*, 90/6:1353–1368,
975 <https://doi.org/10.1785/0120000006>, 2000.

976 Yu, Ch., Vavryčuk, V., Adamová, P., and Bohnhoff, M.: Moment tensors of induced microearthquakes in The
977 Geysers geothermal reservoir from broadband seismic recordings: Implications for faulting regime, stress
978 tensor and fluid pressure, *J. Geophys. Res., Solid Earth*, 123, 8748-8766,
979 <https://doi.org/10.1029/2018JB016251>, 2018.

980 Yu, Ch., Vavryčuk, V., Adamová, P., and Bohnhoff, M.: Frequency-dependent moment tensors of induced
981 microearthquakes, *Geophys. Res. Lett.*, 46, 6406-6414, <https://doi.org/10.1029/2019GL082634>, 2019.

982 Zahradník, J., Sokos, E., Tselentis, G.-A., and Martakis, N.: Non-double-couple mechanism of moderate
983 earthquakes near Zakynthos, Greece, April 2006; explanation in terms of complexity, *Geophys. Prospect.*,
984 56, 341-356, 2008.

985
986