

Last interglacial—(MIS 5e)—sea-level proxies in the glaciated Northern Hemisphere

April S. Dalton¹, Evan J. Gowan^{2,3,4}, Jan Mangerud⁵, Per Möller⁶, Juha P. Lunkka⁷, Valery Astakhov^{8,9}

5

¹Department of Physical Geography and Geoecology, Charles University, Prague, Czech Republic

²Department of Earth and Environmental Sciences, Kumamoto University, Kumamoto, Japan

³Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, Germany

⁴MARUM, University of Bremen, Bremen, Germany

10 ⁵Department of Earth Science, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway

⁶Department of Geology, Quaternary Sciences, Lund University, Lund, Sweden

⁷Geology Research Group, Oulu Mining School, University of Oulu, P.O. Box 9000, Oulu FI-90014 Finland

⁸Institute of Earth Sciences, St. Petersburg University, Universitetskaya 7/9, 199034, St. Petersburg, Russia

⁹A. P. Karpinsky Russian Geological Research Institute (VSEGEI), Sredny pr. 74, 199178, St. Petersburg, Russia

15 *Correspondence to:* April S. Dalton (aprls.dalton@gmail.com) and Evan J. Gowan (evangowan@gmail.com)

Abstract. Because global sea level during the last interglacial (LIG; 130–115 ka) was higher than today, the LIG is a useful approximate analogue for improving predictions of future sea-level rise. Here, we synthesize sea-level proxies for the LIG in the glaciated Northern Hemisphere for inclusion in the World Atlas of Last Interglacial Shorelines (WALIS) database. We describe 82 sites from Russia, northern Europe, Greenland and North America from a variety of settings, including 20 boreholes, riverbank exposures and along coastal cliffs. Marine sediments at these sites were constrained to the LIG using a variety of radiometric methods (radiocarbon, ²³⁰U-Series/uranium-thorium-dating, potassium-argon-argon-dating), non-radiometric methods (amino acid dating, luminescence methods, and electron spin resonance, tephrochronology) as well as various stratigraphic and palaeoenvironmental approaches. In general, the sites reported in this paper do not offer constraint on the global LIG highstand, but rather evidence of glacial isostatic adjustment (GIA)-influenced sea-level positions following the Marine Isotope Stage 6 glaciation (MIS 6; 191–130 ka). As the areas in this database were covered by ice sheets from the penultimate glaciation and were affected by glacial isostatic adjustment (GIA), most of the proxies suggest how that sea level was much higher during the LIG than at the present time during the LIG. Moreover, many of the sites show evidence of regression due to sea-level fall (owing to due to glacial isostatic GIA-uplift), and some also show fluctuations that may reflect regrowth of continental ice or increased influence of the global sea-level signal. In addition to documenting LIG sea level sites in a large swath of the Northern Hemisphere, this compilation is highly relevant for reconstructing the size of MIS 6 ice sheets through GIA modelling. The database is available at <https://doi.org/10.5281/zenodo.5602212> (Dalton et al., 2021).

1 INTRODUCTION

During the last interglacial (LIG), between 130 and 115 ka (peak interglacial at 123 ka; Lisiecki and Raymo, 2005),
35 temperatures were warmer than today by up to 5°C in some regions of the northern Hemisphere (Dahl-Jensen et al., 2013),
and global sea levels were ~~up to 5–m to 10–m~~ higher (Dutton and Lambeck, 2012; Dyer et al., 2021). Like today, Greenland
and Antarctica were the predominant global ice stores, as large continental ice sheets that grew repeatedly during the
Quaternary over North America and Eurasia were absent at that time (see Batchelor et al., 2019). The LIG therefore
represents a useful analogue for understanding the behavior of large continental ice sheets in a warming world, which is key
40 for improving predictions of future melting of the Greenland and Antarctica ice sheets and concomitant sea-level rise (Slater
et al., 2021). The LIG is known regionally as the Kazantsevo interglacial (Siberia; here however redefined as Karginsky, per
Astakhov 2013), the Mikulino interglacial (Russia), the Eemian (western, central and northern Europe), the Ipswichian
(United Kingdom), the Langelandsey interglaciation (Greenland) and the Sangamonian (North America).

45 The World Atlas of Last Interglacial Shorelines (WALIS) is a standardized database that has been created to archive global
sea level sites constrained to the LIG. Here, we contribute 82 sites from the formerly glaciated Northern Hemisphere to
WALIS. We focus on sites that were covered by ice during the Marine Isotope Stage 6 (MIS 6) glaciation (191–ka to 130
ka; Lisiecki and Raymo, 2005; Fig. 1), including Russia, Finland, Estonia, Poland, Sweden, Norway, Svalbard, Iceland,
Greenland, Canada, and the United States. Sea-level proxies in the glaciated regions of the southern North Sea, Jutland
50 Peninsula and Great Britain are the subject of separate studies in this issue (Cohen et al., in preparation 2021). To
standardize the presentation of LIG proxy sites, we use marine isotope stages (MIS), as presented in Lisiecki and Raymo
(2005) and consider that LIG corresponds with MIS 5e (130–to 115 ka) and the penultimate glaciation to be MIS 6. Most of
the sediments described herein were deposited in isostatically depressed land immediately following the retreat of major
55 continental ice sheets from the MIS 6 glaciation (Figs. 21–2). Isostatic recovery is sometimes preserved as a sequence of
deep water, followed by shallowing and shoreline, deltaic, and estuarine depositional settings. Such LIG sites were subject
to considerable erosion from subsequent glaciations – (notably, during MIS 5d (peaked at 109 ka), MIS 4 (71–57 ka) and
MIS 2 (29–14 ka; see Batchelor et al., 2019 and Fig. 2) and are therefore only sporadically preserved. There are also many
more sites preserved

60 In the first part of this paper (Sections 2–5), we define the types of sea-level proxies, elevation measurements, dating
techniques and quality assessment, all of which are technical aspects of entering the LIG data into the WALIS database. In
Section 6, we describe each LIG site in detail, paying particular attention to the elevation of marine sediments and any
geochronological constraints. In Section 7, we present sites containing LIG marine sediments that are not *in situ* and have
been transported and/or glaciotectonised following deposition. These sites are unsuitable as precise indicators of relative sea
65 level (RSL) and are therefore excluded from the WALIS database. However, they contribute to the general picture of LIG

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, English (United States), Pattern: Clear

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, English (United States), Pattern: Clear

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, English (United States), Pattern: Clear

sea level and are therefore included here. In the Discussion (Section 8) we provide an overview of the LIG sites compiled for this paper, as well as examples of marine deposits of other ages in the glaciated region, notably MIS 7 (243 ka to 191 ka), MIS 5c (peak 96 ka), MIS 5a (peak 82 ka) MIS 3 (57–29 ka) and Holocene (11.5 ka to present-day; Fig. 2). We conclude with suggestions for future research. As shown in Figure 2, the LIG is known regionally as the Kazantsevo interglacial
70 (Siberia; here however redefined as Karginsky, per Astakhov 2013), the Mikulino interglacial (Russia), the Eemian (western, central and northern Europe), the Ipswichian (United Kingdom), the Langelandslev interglaciation (Greenland) and the Sangamonian (North America).

2 SEA-LEVEL PROXIES

Our approach to describing sea-level proxies differs from the standard approach used in most of the studies in WALIS.

75 Since most of the LIG sites are located in places that were undergoing rapid sea-level changes dominantly due to glacial isostatic adjustment (GIA)GIA rather than global sea-level change, it is essentially impossible to pinpoint when sea level was at a particular elevation (especially given the large uncertainty in the dating methods). In many locations, there are indications of regression from an often indeterminate highstand position at deglaciation, to a position below the elevation of the outcrop. For many sites, there is clear evidence of coarse grained, wave influenced deposits that show that sea level was
80 near the elevation of the investigated the deposit. The indicative meaning of these deposits, as defined by Rovere et al. (2016), is not sufficiently clear to deduce a precise sea-level position.

We regard it as more useful to describe the sea-level proxies in terms of how much information a site provides in terms of changes in sea level at each site during the LIG. Accordingly, The primary target for this compilation is mainly intended for
85 researchers for those who are interested in inferringinferring the size of the MIS 6 ice sheets through GIA modelling (e.g. Lambeck et al., 2006). In the database descriptions (Dalton et al., 2021), we have indicated the relative water depth based on the geological descriptions at the sites (i.e. deep water, shallow water, near sea level, above sea level). From this information, it should be possible to test the reliability of MIS 6 ice sheet reconstructions and deduce LIG sea level.

90 In our database (Dalton et al., 2021), the vast majority of the sea-level proxies are denoted as "marine limiting". The entered elevation marks the highest elevation of marine sediments at a site. This usually marks an unconformity between the marine sediments and overlying younger sediments that date to between MIS 5d and MIS 1 (started at 14 ka; see Fig. 2). In a few cases, the contact is a conformable transition from marine to terrestrial (often fluvial or lacustrine) sedimentation. At sites where there is reasonably high confidence of the indicative meaning (generally where there is confidence of the sea-level
95 highstand), we have defined them as sea-level indicators using the standard approach of WALIS (i.e. Rovere et al., 2016), and note that sea level likely never exceeded that elevation. At sites where there is evidence of a regression and transgression

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, English (United States), Pattern: Clear

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, English (United States), Pattern: Clear

within the LIG, we have created two entries in the database. Based on the amount of information on sea-level position and variations at a site, we have assigned a quality score, which will be elaborated on in Section 5.

3 ELEVATION MEASUREMENTS

100 A summary of elevation measurement techniques and datums, as used in cited research, are found in Tables 1 and 2.
However, The techniques used to measure the elevation(s) were often not stated in many of the studies covered in our database and were instead extracted from stated elevations and section diagrams in the original publications. In these cases, we applied a nominal uncertainty of 20% of the stated elevation, as recommended by Rovere et al. (2016), or 10 m, which ever was smaller. The later constraint is It is unlikely the elevation uncertainty will be worse than the contour

105 intervals of typical topographic maps (10–20 m), provided the authors were precise in the pinpointing the location of their site. For studies that involved the authors of this paper, we were able to provide the details of the elevation measurements and provide narrower uncertainties. The datum used was not stated in most of the studies presented here and are is assumed to be referenced to present day mean sea level. The tidal range in most locations covered in our database is presently relatively small (*i.e.* < 1–2 m), so this is unlikely to add significant uncertainty.

Formatted: Not Highlight

110 4 OVERVIEW OF DATING TECHNIQUES

A large number of dating techniques have been applied to LIG marine deposits covered in our database. These dating techniques include absolute (luminescence, electron spin resonance, U/Th), minimum limiting (radiocarbon), and relative approaches (amino acid racemization, stratigraphy, environmental conditions). Generally, The absolute dating techniques have relatively large uncertainties and cannot be used to give a precise timing of deposition within the LIG. When combined 115 with paleo-environmental conditions, it can usually be concluded that the deposit has a LIG age, rather than being part of another period of high sea level (*e.g.* Holocene, MIS 3, MIS 5a/c, MIS 7; see Fig. 2). Consideration of quality of the age control in the database is elaborated in Section 5.

4.1.1 Amino acid racemization dating

120 Amino acid geochronology measures the racemisation of amino acids. For the LIG, the epimerization of D-allo-isoleucine to L-isoleucine is most used (known as the D/L ratio; Oldale et al., 1982; Miller and Mangerud, 1985). Older shells have a higher isoleucine epimerization ratio than younger shells. However, this is a relative dating technique, with the epimerization controlled by regional diagenetic temperature, among other factors (Andrews et al., 1983). Therefore, this technique can only be used for correlation between sites or to differentiate between different marine incursions in each region, rather than give precise ages.

125 4.1.2 Radiocarbon dating

Radiocarbon dating measures the amount of radioactive carbon (^{14}C) remaining in organic material after death of the dated animal/plant. The time since death can be approximated by consideration of the mean half-life (5,730 years ka) of ^{14}C (Stuiver and Polach, 1977) and then converted to calendar year via calibration (Reimer et al., 2020). However, this chronological method is only useful for samples less than ~45,000 years old because the remaining ^{14}C in old samples is too scarce to be reliably measured beyond that point, and the sample becomes increasingly susceptible to modern-day carbon contamination (Douka et al., 2010). Thus, for the purposes of identifying MIS 5e/LIG marine sites, radiocarbon ages offer only minimum constraint.

4.1.3 Stratigraphic inferences

In many cases, the stratigraphic position of a particular marine unit ~~has provided~~ evidence for its age-of-deposition. When 135 the marine unit is overlain by tills that are independently assigned to MIS 5d/b (peak MIS 5d conditions were at 87 ka), MIS 4 or MIS 2 glaciation, ~~this is stratigraphic context is~~ used as evidence to support a LIG n MIS 5e age assignment (e.g., at Pöhja-Uhtju and Peski; Miettinen et al., 2002; [Sections 6.22-6.23](#)). ~~Conversely, As another example, sometimes~~ the presence 140 of a till directly underneath the marine sediments ~~that~~ suggests significant isostatic depression (often related to the MIS 6 glaciation), ~~which, and along with evidence of~~ rapid marine inundation into the isostatically depressed landscape and shallowing of marine waters, ~~is used as evidence of an LIG age assignment owing to subsequent rebound~~ (e.g., Ile aux Coudres; Occhietti et al., 1995; [Section 6.45](#)).

4.1.4 Palaeoenvironmental inferences

Climate during [the LIG](#) was several degrees warmer than present-day temperatures in the Northern Hemisphere (Rasmussen et al., 2003; Sánchez Goñi et al., 2012). As a result, palaeo-indicators of warmer-than-present-day conditions are 145 often used as support for a LIG age assignment. Marine-based palaeoecological indicators commonly preserved in the stratigraphic record include dinoflagellate cysts, foraminifera, Coelenterata, Bryozoa and Mollusca, diatoms, and marine gastropods (Bergsten et al., 1998; Mangerud et al., 1981). In the terrestrial sediments that often overlie the [LIG](#) marine unit, pollen [assemblage zones \(PAZ\)](#) and Coleoptera are some of the most used markers for determining palaeo-temperature (Dredge et al., 1992; Miettinen et al., 2002). Some caution is needed when considering these as correlating to 150 [the LIG](#) in the absence of other numerical dating methods, as it is possible that these deposits could be from an older interglacial period. In our database, we refer to any site with environmental conditions supporting LIG assignment as an "Eemian interglacial deposit", as defined by Mangerud et al. (1979).

4.1.5 Thermoluminescence dating

A common dating method for ~~MIS-5eLIG~~ marine sediments is thermoluminescence (TL), although in recent years it ~~is-has~~
155 ~~been~~ largely replaced by optically stimulated luminescence (OSL; described below in Section 4.1.7). ~~This-The~~ TL technique measures the last exposure of a sediment to sunlight via the resetting of electrons (Huntley et al., 1985; Lamothe and Huntley, 1988). In a laboratory setting, these changes are measured by heat stimulation. Either quartz or feldspar can be used, although feldspar is susceptible to anomalous fading, which can lead to large uncertainties in the age estimation ~~because the electrons trapped in the crystal lattice 'leak' over time~~ (Godfrey-Smith et al., 1988; Huntley et al., 1985).

160 4.1.6 Infrared stimulated luminescence (IRSL)dating

~~This-The~~ infrared stimulated luminescence (IRSL) method uses wavelengths in the infrared range to induce luminescence in feldspar, which in some cases has been shown to reduce uncertainties in age estimates compared with TL (Godfrey-Smith et al., 1988).

4.1.7 Optically stimulated luminescence (OSL)dating

165 Similar to TL dating, OSL measures the refilling of shallow electron traps in sediment grains that occur during burial following exposure of the sediment grain to sunlight; therefore, the time since burial can be obtained (Duller, 2008; Huntley et al., 1985). In a laboratory setting, the release of these electrons is induced by light stimulation and ~~measured-the~~ escaping dose ~~is measured~~. A key consideration in OSL dating is the former depositional context (largely shallow marine settings in our dataset) and its impact on bleaching (zeroing) of the sediments by sunlight, as well as the burial history of the sediment
170 ~~(multiple ages in Svalbard; Alexanderson and Landvik, 2018; Alexanderson et al., 2011a)~~. Generally, OSL is considered to be more reliable than TL and IRSL.

4.1.8 Electron spin resonance (ESR)dating

~~Electron spin resonance (ESR) dating~~ estimates the time since deposition of certain materials (largely molluscs in our database) by measuring the trapping of electrons within the material's crystal lattice. A key factor is the radiation
175 occurring from the enclosing sediment, as well as the radioactivity of the sample (Grün, 1989). The precision of the age from ESR dates on mollusc shells is complicated, since the shells have an open system to uranium (Schellmann and Radtke, 1999). As a result, care must be taken in interpreting ESR ages, and the uncertainty can be larger than the stated analytical uncertainty. The analytical techniques used to determine ESR ages for many of the sites in our database are described in detail in Molodkov (1988) and Molodkov et al. (1998).

180 4.1.9 **Uranium-thorium dating**

As with ESR dating, uranium-thorium (U/Th) dates ~~of from~~ mollusc shells are complicated to interpret since the shells have an open system to uranium, and uranium and relies on assumptions on the exchange of the element with the surrounding environment (Radke et al., 1985). As a result, this technique is not commonly applied to LIG deposits. U/Th dating of ~~molluscs~~ This method ~~shells~~ was used to constrain the age of deposits at two sites in our database (Miller et al., 1977, 185 Israelson et al., 1994). For the later, the ages U/Th age provided only minimum ages constraint.

4.1.10 **Tephrochronology**

At one site in our database (Galtalækur site in Iceland; Vliet-Lanoë et al., 2018; Section 6.40), tephrochronology is used to support the MIS 5e LIG age assignment. Contained within the marine sediments is a tephra layer that was linked via geochemical analyses to a specific eruption (Grimsvötn volcano) which was, in turn, constrained to MIS 5e the LIG based on 190 the position of the Grimsvötn tephra in North Atlantic marine sediment cores (Davies et al., 2014).

4.1.11 **Potassium–argon K–Ar dating**

At one site in our database (Galtalækur site in Iceland; Vliet-Lanoë et al., 2018) potassium-argon (K/Ar) dating is used to constrain the age of a glacio-volcanic unit that underlies the MIS 5e LIG marine unit. This method measures ~~the~~ the rate of decay from Kpotassium to Arargon and was possible at this specific site owing to given the presence of volcanic rocks. It 195 and is otherwise not a common dating method in common to Quaternary research.

5 QUALITY ASSESSMENT

At each LIG marine site documented in the WALIS database there is a quality assessment rating for the RSL proxies. In the WALIS documentation, the standard RSL rating is designed for far-field sea-level indicators, in which a precise assignment of sea-level position can often be determined. For glaciated areas where sea level 200 was rapidly changing due to glacial isostatic rebound after the end of the MIS 6 glaciation, this type of quality assessment is not as useful, especially since the dating techniques are not precise enough to pinpoint when the marine sediments were deposited. As one of the primary uses of this database will be GIA modelling, we have devised a rating scheme to assess the usefulness for this purpose (Table 3). The best quality RSL proxies are sedimentary sequences that have well documented elevation measurements, and in which there is a clear transition from deep marine to shallow 205 marine, beach, and terrestrial environments. A high rating is also assigned if it can be proved that sea-level position remained above a threshold for a long period of time, *i.e.* for most or all of the LIG. The rating decreases when there is less geological evidence of sea-level position changes and proximity to sea level.

210 The chronological rating for each LIG site follows the WALIS standard (Table 4) and is determined by how precise the age of the marine deposit can be constrained. In the standard WALIS documentation, higher chronological scores are generally only assigned for U/Th dated corals that can have precision of less than a few thousand years, which is not the case in the techniques applied to glaciated region sea-level proxies. Despite this, based on stratigraphy and environmental proxies, there is reasonable confidence that most of the proxies documented in our database fall either within the LIG, during the latest part of MIS 6 when the ice sheets were retreating, or during the early parts of MIS 5d when ice sheets began to grow again.

215 Without environmental indicators, it can be difficult to distinguish deposits that might instead date to the sea-level highstand that happened after the MIS 5d glaciation (e.g. Mangerud et al., 1998). For North American sites, a high rating assignment from environmental conditions alone is not applied, as the LIG stratigraphy is not as well understood as in Europe (Otvos, 2015). Sites that have multiple age determinations increase the confidence in our LIG age assignment. However, the precision of the techniques is never high enough to elevate any site rating to the highest level.

220

As a result of this, almost all of the proxies are set to have a score of 2 (not numerically age constrained and based on relative ages or environmental conditions) or 3 (numerically age constrained). Higher scores in WALIS are generally only assigned for U/Th dated corals that can have precision of less than a few thousand years, which is not the case in the techniques applied to glaciated region sea-level proxies. Despite this, based on stratigraphy and environmental proxies, there is reasonable confidence that all of the proxies documented in our database fall either in the LIG, the latest part of MIS 6 when the ice sheets were retreating, or during the early parts of MIS 5d when the ice sheets began to grow again. Without the environmental indicators, it can be difficult to distinguish deposits that might instead date to the sea-level highstand that happened after the MIS 5d glaciation (e.g. Mangerud et al., 1998). The primary challenge of assessing past sea level in formerly glaciated areas is that the sea-level position likely did not remain fixed for any length of time. Moreover, the dating techniques that can be applied to LIG deposits lack the precision to determine when sea level was at a specific elevation (with the possible exception of the correlation with European pollen records, if such long-distance correlations are valid). Therefore, we chose to describe the proxies in terms of how much information they give to show sea-level variations in the LIG. Most of the studies we looked at lack information on how elevation was determined. However, we were able to include this information in the database for sites that involved the authors of this study. Since a large portion of sites described here were overridden by ice sheets during the last glacial period (MIS 2; see Fig. 1), most of the sections are incomplete due to erosion.

230

Here, we describe sites from the formerly glaciated Northern Hemisphere that contain *in situ* LIG marine sediments, ordered roughly from east to west, and sorted by country. Geochronological results are reported for all the sites where data are available. Sediments overlying or underlying the marine strata are generalized unless they provide additional context for the LIG marine event. We offer no interpretation of tectonics, glacio-isostatic adjustmentGIA or eustacy. Elevation measurements are in MASL (meters above sea level (MASL)) or MBSL (meters below sea level (MBSL)). Owing to the

235 **Formatted:** Strikethrough

global scale of this database, it is not possible to map all features/locations [described in the text \(especially rivers\)](#) and the reader is referred to the original publications for [specific local site information](#). [Additional details \(including site coordinates, 245 elevation of marine sediments, quality scores for both RSL and age determinations\)](#) [are](#) summarized in Table 5 and detailed in the database of Dalton et al. (2021).

[7.16.1 Novorybnoye 2, Taimyr Peninsula, Russia \(72.83, 105.79\)](#)

On the southernmost Taimyr Peninsula, ~30 m-high river bluffs close to the small settlement of Novorybnoye, on the south shore of the Khatanga River, expose a complex Mid- to Late Pleistocene stratigraphy ([Fig. 3B](#)). As described in Kind and Leonov (1982), this record encompasses three glacial till units on top of Cretaceous sandstone; the till beds are described as interbedded with two marine sediment successions, the lowermost to be, in their terminology, [Kazantsevo \(i.e. MIS 5e, LIG\) LIG](#) in age and the upper one sandwiched between two Early Zyryanka tills (MIS 5d–5a). This chronostratigraphy was, however, not substantiated by any numerical ages. The Novorybnoye bluffs were reinvestigated by Möller et al. (2019a; 2019b), resulting in two more observed marine units and a very different chronology, in which the lowermost marine unit 255 probably dates to MIS 9–11 ([424–337 ka](#)). Relevant to the present study, the LIG marine sediments span 14.8–24 MASL and are divided into two units: F1 and F2. Marine unit F1 (spanning 14.8–21.5 MASL) is at the base a glaciomarine clayey silt with numerous occurrences of ice-rafted debris. Above these marine sediments are stratified and normally graded sandy shoreface sediment (F2) with an abundance of *Hiatella arctica* and *Astarte* sp. shells from 21.5–24 MASL. An OSL age of 124 ka is supported by a mollusc ESR date at 131 ka; Möller et al. (2019a) thus concluded that 260 this entire marine unit (F1–F2) represents marine inundation following deglaciation and sediment deposition during isostasy-driven shore regression at the transition from MIS 6 into the [Karginsky interglacial \(MIS 5e\) LIG](#). The absolute height of unit F sediments is ≥25 MASL, which is a minimum deglacial sea-level altitude.

Formatted: Font: (Default) +Body (Times New Roman)

[7.26.2 Bol'shaya Balakhnya River \(BBR 17\), Taimyr Peninsula, Russia \(73.62, 105.36\)](#)

In the lower reaches of the Bol'shaya Balakhnya River ([Fig. 3B](#)), a sediment succession shows marine silty clay with dispersed ice-rafted debris situated between 6.2–7 MASL (site BBR 17B) and 7–13 MASL (site BBR 17A). At BBR 17B, there is an upper erosional contact to fluvial sediment at 7 MASL (Möller et al., 2019a, 2019b; Der Sarkissian et al., 2020). The mollusc fauna in the marine sediments is dominated by arctic *Portlandia arctica*, but there is also an abundant occurrence of subarctic taxa: *Buccinum undatum*, *Mytilus edulis* and *Macoma baltica*, suggesting higher than present influx of Atlantic water. [Electron spin resonance](#) ages on *P. arctica* are 101–105 ka (n = 3) and 270 indicate an MIS 5c age (known locally as Early Zyryanka). However, two molluscs in the above-lying fluvial sediments (OSL-dated to a MIS 3 age; the molluscs redeposited from erosion of the [underlying](#) marine sediment), yield ESR ages of 122 and 123 ka. These dates, together with the interglacial-type mollusc fauna composition, clearly set

the marine sediments into the LIG, however with poor indication of sea level at deposition.

7.36.3 Kamennaya River, Taimyr Peninsula, Russia (76.53, 103.52)

275 Around the Kamennaya River (Fig. 3B), which is a tributary to the Leningradskaya River, Gudina et al. (1983) described several sites exposing marine and nearshore marine sediments forming regressive terraces from 133 ~~down to~~ 40 MASL (site no. 373 located at the highest altitude of with highest altitude, 133 MASL, site no 373, coordinates as above). The sediment successions are divided into a lower coarsening upwards member, from silts and clays to cross-bedded sand, and an upper member of sand and gravel. All sediments are rich in molluscs, predominantly *Astarte borealis*, *A. crenata*, *A. montuagi*, *Macoma calcarea*, *Hiatella arctica* and *Mya truncata*, (i.e. species that are ~~A~~rtic to non-conclusive in their biogeography). As opposed to the molluscs, the foraminifera association (48 species detected) has a dominance of subarctic species, thus suggesting warmer than present sea temperatures. Based on the latter, Gudina et al. (1983) suggested that the marine sediments in the Leningradskaya basin were deposited at shore regression during the ~~Kazantsevo interglacial (Karginsky in terms of Astakhov (2013); i.e. MIS 5e) LIG. Gudina et al. (1983) do not present any numerical age data.~~

280

7.46.4 Kratnaya River sections, Taimyr Peninsula, Russia (77.5, 103.2)

285 Three river-cut sections along the Kratnaya River (denoted KR1, KR2 and KR3; Fig. 3B), expose a thick basal till on top of which are marine sediments with a slight upwards-coarsening trend with off-shore silt and sand grading into shoreface-deposited sand with gravel stringers (Möller et al., 2008; Möller et al., 2015). The base of these marine sediments is situated at 37 MASL and the uppermost logged sediments reach ~43 MASL, but the marine sediment succession can locally 290 be followed upslope to at least 50 MASL. The sediments host a variable abundance of ~~the~~ molluscs with *Hiatella arctica*, *Mya truncata* and *Astarte borealis* as dominating taxa (all ~~A~~rtic to non-conclusive in their biogeography). Six ESR ages on 295 molluscs from the marine sediment form an age cluster between 111 ka and 142 ka, while two OSL dates gave younger ages of 84 ka and 100 ka (Möller et al., 2008; Möller et al., 2015). The ESR ages firmly suggest a transition from ~~the MIS 6 glaciation (known locally as the Late Taz) (MIS 6 glaciation)~~ into the ~~Karginsky (MIS 5e) interglacial LIG~~ for the emplacement of this ~~regressive~~ marine sediment succession.

7.56.5 Anjeliko River sections, Taimyr Peninsula, Russia

300 Based on the composite stratigraphy of five river-cut sections (three of which have LIG sediments and are included in the database) along and in the vicinity of the Anjeliko River (Fig. 3B), Möller et al. (2008) and Möller et al. (2015) report three marine units intercalated with glacial tills and suggested that two of these marine events, together with underlying tills, represent two full glacial cycles coupled with marine inundation and regression following deglaciations. Relevant to the present study, the intermediate marine unit consists of glaciomarine debris flow sediments followed by an upwards coarsening sediment succession of off-shore clay and silt into shore-face/fore-shore sand. These marine sediments are situated between 55 and 58 MASL. Outside of logged sediment successions, the marine sediments could be followed to ~80

MASL and, if observations from investigations by Mirošnikov (1959) and Šnejder (1989) from other nearby locations (notably on Cape Chelyuskin) refer to the same sediments, they might have a highest altitude of up to 140 MASL. The sediments host a variable abundance of the molluscs *Hiatella arctica*, *Mya truncata* and *Astarte borealis* as dominating taxa, all Arctic to non-conclusive in their biogeography. Three ESR ages on molluscs from the marine sediment yielded ages of 143, 145 and 156 ka, while two OSL ages suggested 79 and 135 ka (see Möller et al., 2015). The age envelope (disregarding the 79 ka OSL age) suggests a transition from the MIS 6 glaciation (known locally as the Late Taz) into the LIG for this marine sediment succession. The uppermost glacial/marine sediment succession in the Anjeliko River area dates from a glaciation followed by marine inundation during MIS 5d-5c (known locally as the Early Zyryanka; see Möller et al., 2008).

7.66.6 Ozernaya River, October Revolution Island, Russia (79.12, 96.92)

October Revolution Island is located offshore of the Taimyr Peninsula, in the Severnaya Zemlya archipelago, Arctic Russia (Fig. 3B). On this island, along the north to south-flowing Ozernaya River, are occasional exposures of up to 50-m thick paleo-valley fill sediment successions, ordered in a pancake-like stratigraphy. First briefly described by Bolshiyanov and Makeyev (1995), a more in-depth description was made in Möller et al. (2007), showing four till beds (Till I-IV) interbedded with three marine units (Marine I-III). Relevant to the present study, Marine unit III is situated between 75.5 and 80.5 MASL (sites Oz1b and 1d). These marine sediments can laterally be followed to higher ground with no covering till (sites Oz2 and 3; see Fig. 5A in Möller et al., 2015), eventually terminating in sets of beach ridges, the highest at ~140 MASL. In the database (Dalton et al., 2021), we have treated the highest elevation beach ridge as a sea-level indicator. The marine sediments show a general coarsening upward trend with off-shore to shore-face deposited silt and clay with varying frequency of ice-dropped clasts, continuing into massive or vaguely stratified sand (site Oz1) and finally beach-face gravels (sites Oz 2-3). Besides Arctic molluscs as *Hiatella arctica*, *Mya truncata* and *Astarte borealis*, the marine sediments also host biogeographically subarctic species such as *Chlamys islandica* and *Buccinum undatum*. The foraminifera fauna is also mainly Arctic, but a warmer-water indicator is *Trifarina* cf. *angulosa* and the also occurring *Elphidium ustulatum* and *Islandiella inflata* that are rarely found in deposits younger than the LIG in Europe (Möller et al., 2007).

The age of the Marine unit III is not straight-forward, as discussed in Möller et al. (2007, 2015) and Möller et al. (2015). Electron spin resonance (ESR) ages on the mollusc fauna have an age envelope of 77–105 ka (n = 8), while ESR ages on nearby found *Chlamys islandica* is 105 and 120 ka (Bolshiyanov and Makeyev, 1995). *GSL* (Optically stimulated luminescence) ages on the sediment show an age envelope of 143–176 ka (n = 11). Based on the stratigraphic position in a regional context, the interglacial fauna elements in the marine sediments and an evaluation of the numerical ages from performed dating attempts, Möller et al. (2007) favoured an interpretation that the Marine unit III sediments in the Ozernaya River valley were deposited at the transition between the MIS 6 glaciation (known locally as the Late Taz) and the

LIG, and that the deglacial sea had a highstand of at least 140 MASL.

7.76.7 Lower Agapa River, Taimyr Peninsula, Russia (71.6, 88.3)

In the lower reaches of the Agapa River (Fig. 3B), several river sections display three marine units (Gudina et al., 1968; later 340 reinvestigated by Sukhorukova, 1998). The lowermost unit is interbedded fine sand and silt with the boreal bivalve *Cyprina islandica* (now *Arctica islandica*) followed by silt and clay (Unit 2), 30–35 m thick, spanning 30 to ≥63 MASL. Unit 2 hosts a rich mollusc fauna (16 subarctic (arcto-boreal) and 20 arctic species), however with no species list given. The foraminifera fauna also suggests a component of Atlantic water inflow. Marine Unit 2 with interglacial-type marine fauna reaches ≥63 MASL, which is a minimum sea-level altitude for these deeper marine sediments. ~~No numerical age data are presented in~~ 345 ~~Gudina et al. (1968), nor in Sukhorukova (1998). However, the~~ ~~The~~ presented biostratigraphy strongly suggests that Unit 2, with a warmer-water fauna, represents interglacial conditions, most ~~probably MIS 5e (Karginsky)~~ likely the LIG.

7.86.8 Karginsky Cape, West Siberian Plain, Russia (69.95, 83.57)

At Karginsky Cape (Fig. 3B), Western Siberian Plain, Kind (1974) described a ~16-m thick sequence of marine sands and silts situated between 5 and 21 MASL, sandwiched between two tills. This ~~site~~ is the LIG marine stratotype ~~for this region~~ 350 ~~(Karginsky Formation; see stratigraphy in Fig. 4)~~. These marine sediments contain shells of the biogeographically ~~Arctic-to-non-conclusive~~ molluscs *Astarte borealis*, *Macoma calcarea*, *Mya truncata* and *Ciliatocardium ciliatum*, as well as the subarctic *Mytilus edulis*. Shells of a typical boreal mollusc, *Arctica islandica*, were found only on the beach. The sequence also contains remains of plants presently growing some 3–5° ~~(approx. 300–500 km)~~ to the south, suggesting it was deposited during a warmer interval. Conventional radiocarbon dates obtained in this section 355 ~~were~~ 42, 42, 46 and ≥52 ka. However, the first ESR date on an *Arctica islandica* shell yielded an age of 121.9 ka (Katzenberger and Grün, 1985; Arkhipov, 1989). The LIG age of this marine formation was later confirmed by 6 OSL dates in the range of 117–97 ka (Astakhov and Nazarov, 2010b; Nazarov, 2011; Nazarov et al., 2018, 2020).

7.96.9 Tanama River, Western Siberian Plain, Russia (2 sites)

360 Along the Tanama River (Fig. 3B), Nazarov et al., (2021) describe marine sediments overlying till from the Taz (MIS 6) glaciation, ~~below which are -and-~~ marine sediments from an earlier interglacial (interpreted as MIS 7 based on OSL dates). The upper unit of marine sediments (known locally as the Payuta marine formation) consist of sands and silts, with numerous shells of *Arctica islandica*, which indicate~~s~~ warmer water conditions. The unit is associated with terraces that reach an elevation of 60–70 MASL. The change in slope at 70 m took on the appearance of strandlines. Overlying the 365 marine sediments are lacustrine and alluvial sediments that ~~gave yielded ages in the range of~~ MIS 4–3 ages.

7.9.16.9.1 Tanama site 1

The unit is associated with terraces that reach an elevation of between 60 and 70 MASL. ~~The change in slope at 70 m took on the appearance of strandlines.~~

7.9.26.9.2 Tanama site 2 (69.83, 79.00)

370 The unit is associated with terraces that reach an elevation of between 60 and 70 MASL. ~~The change in slope at 70 m took on the appearance of strandlines.~~

7.106.10 Bol'shaya Kheta, West Siberian Plain, Russia (4 sites)

In the Western Siberia Plain, there are LIG outcrops along the Bol'shaya Kheta River (Nazarov et al., 2020, 2021; Astakhov and Semionova, 2021). First described by Volkova (1958), these sites contain a similar ~~identical~~ stratigraphic record that can

375 be traced along the river sections (Fig. 3), with LIG marine sediments generally ranging from 5 ~~to~~ 40 MASL (~~stratigraphy and ages summarized in Fig. 4~~). The upper marine formation was assigned to the ~~Kazantsevo interglacial~~^{LIG} by Sachs (1953), ~~in modern terminology corresponding to the Karginsky interglacial (MIS 5e; LIG)~~. This marine unit contains only the extant species *Cyrtodaria siliqua* and *C. kurriana* and a rare occurrence of the boreal *Arctica islandica*. ~~These sites are mapped in Fig. 3B.~~

Formatted: Font: Not Italic

7.10.16.10.1 Site 7251, Bol'shaya Kheta River (68.47, 83.12)

At this site, the upper marine sand- and clay ~~beds~~ ~~s (constrained to MIS 5e; LIG)~~ are situated between 20 and 30 MASL (Fig. 3). Two OSL ages on these marine sediments yielded ~~ages of~~ 124 \pm 31 ka and 121 \pm 11 ka (Nazarov et al., 2020, 2021; Astakhov and Semionova, 2021). These marine sediments directly overlie a till of MIS 6 age.

7.10.26.10.2 Site 7248, Bol'shaya Kheta River

385 Located 55 km south from ~~the Northern River sections~~^{site 7251}, marine sand- and clay ~~beds~~ ~~s (constrained to MIS 5e; LIG)~~ are located between 5 ~~to~~ ~~and~~ 30 MASL (Fig. 3). Three OSL attempts on these marine sediments yielded ~~ages of~~ 110 \pm 16 ka, 127 \pm 20 ka and 114 \pm 12 ka (Nazarov et al., 2020, 2021; Astakhov and Semionova, 2021).

7.10.36.10.3 Site 7249, Bol'shaya Kheta River (68.00, 83.13)

390 At this site, ~~m~~Marine sand- and ~~e~~lays (~~LIG~~)clay are located between 5 ~~to~~ ~~and~~ 30 MASL (Fig. 3; Nazarov et al., 2021). No chronological constraints are available for this site, but the marine unit can be traced along the ~~riverbank~~ ~~shoreline~~ to the other sites dated to the LIG.

7.10.46.10.4 Site 7246, Bol'shaya Kheta River

Marine sand and clay beds (LIG) are located between 5 to 30 MASL (Fig. 3; Nazarov et al., 2021). A single OSL age of 132 ± 11 ka suggests deposition during the LIG (Nazarov et al., 2021).

395 7.116.11 Observations Cape, West Siberian Plain, Russia (68.97, 76.10)

On the tip of the Taz Peninsula, Western Siberian Plain, the Observations Cape site (Fig. 3B) contains parallel laminated sand, silt and clay with an abundance of boreal molluscs as *Modiolus* sp and *Zirfaea crispata* *Zirphaea crispata* and subarctic species as *Buccinum undatum*, *Macoma balthica* and *Mytilus*. Described and dated by Astakhov and Nazarov (2010b), these marine sediments are situated between 3 and 35 MASL and yielded 6 OSL dates with a mean age of 135 ka (stratigraphy and ages summarized in Fig. 43), leading to a LIG MIS 5e (LIG) age assignment. These marine sediments are underlain by Middle Pleistocene glacial till and overlain by fluvial sand with OSL ages at of 77 and 74 ka BP.

Formatted: Font: Italic

7.126.12 Sula, Pechora Lowland, Russia (67, 50.34)

In the Pechora Lowland (Fig. 5B), the best dated LIG site is the succession of shoreline sands along the Sula River, a left tributary to the Pechora River (denoted sites 21/22). The sand formation was originally described as lying on top of marine clay with a cool-indicating mollusc fauna (Lavrova, 1949), the latter, however, was not confirmed by later descriptions. The well-exposed marine sand spans the interval 31–41 MASL, starting with thin foreshore gravel in tabular foresets, containing paired shells of subarctic *Mytilus edulis* (stratigraphy and ages summarized in Fig. 4). The fining upwards and bioturbated sand contains abundant shells of boreal molluscs *Arctica islandica* and rare shells of *Cerastoderma edule* and *Zirphaea crispata* – all species not presently living east of the Kola Peninsula. This mollusc fauna is typical for a shallow sea with positive bottom temperatures and occurs through the sand formation, topped by a cross-bedded beach gravel. The marine unit, attributed to the LIG, is overlain by fluvial sand and cryoturbated black silty clay of glaciolacustrine origin, topped by aeolian silt (Mangerud et al., 1999). An approximate LIG age was later confirmed by numerous OSL dates; altogether 16 ages in the range of 90–128 ka with a mean age of 112 ± 2 ka (see Fig. 3, and Murray et al., 2007).

415 7.136.13 River Yangarei, Pechora Lowland, Russia (68.7, 61.83)

In addition to the Sula site, marine sediments of a MIS 5e LIG age are also found at much higher elevations within the Pechora Lowland (Fig. 5B). As an example, marine sand with mollusc shells occur along the Yangarei River at 70 MASL, the sediments yielding OSL ages at of 121.6 ± 9.2 ka and 114.8 ± 8.9 ka (Astakhov and Semionova, 2021).

7.146.14 Vorga-Yol Section, Pechora Lowland, Russia

420 Also in the Pechora Lowland, OSL ages of 126 ± 8 ka, 131 ± 8 ka and 149 ± 10 ka were obtained from sand with shell
fragments -at 90 MASL, directly underlying the terminal glaciofluvial delta at Vorga-Yol section (Astakhov and Semionova,
2021; [Fig. 5B](#)).

7.156.15 Pyoza River, Arkhangelsk district, Russia (11 sites)

425 Marine sediments assigned to the [Mikulinian](#) (LIG ([known locally as the Mikulinian](#))) were first noted in stratigraphic
records along the Pyoza River, [in the](#) Arkhangelsk district, [Russia](#), in the early-mid 20th century and examined more recently
by Houmark-Nielsen et al., (2001) and Grøsfjeld et al., (2006). We [compile describe](#) 11 sites below, [which are all mapped in](#)
[Figs. 5B-C](#).

7.15.16.15.1 Zaton site (65.58, 44.63)

430 First discovered by Ramsay (1904), the Zaton site (Site 0 of Houmark-Nielsen et al., 2001) was examined by Devyatova and
Loseva (1964) and Devyatova (1982). As described most recently by Grøsfjeld et al. (2006), the entire LIG marine
sequence spans 2 [MASL to](#) 11 MASL. At the base of the section, from 2 [to](#) 7.5 MASL are marine clays with a gradual
transition into silty sands, interpreted as an offshore to shoreface sediment succession (>45 m to <12 m water depth). At the
435 top of the section, from 7.5 [to](#) 10 MASL are laminated sand and silt, separated by gravel horizons. These sediments are
interpreted as deposited in a [strong higher-energy](#) coastal/tidal environment with channel erosion and infilling (foreshore
environment; <12 m depth). Capping this sediment succession, from 10 [to](#) 11 MASL, [are](#) cross-bedded fluvial sand. [Marine](#)
molluscs, dinoflagellate cysts and benthic foraminifera are present throughout (Grøsfjeld et al., 2006). The
stratigraphy observed at this site is laterally continuous for ~800 m.

440 Several geochronological attempts have been made at the Zaton site. Early TL dating on sediments overlying the marine
[unit](#) yielded an age of 93 ka, which supports a LIG-age for the underlying marine sediments (Hütt et al., 1985).
Amino acid dating on shells from the marine unit yielded mean [a](#) D/L ratio of 0.051 ± 0.006 , which is [slightly higher than](#)
[expected](#) but [probably Eemian](#) according to Miller and Mangerud (1985). Subsequent ESR dating confirmed a LIG
age for this marine deposit (120 [82](#) ka; Molodkov and Raukas, 1988; Molodkov and Bolikhovskaya, 2002). A LIG-age is
also suggested by pollen analyses (Devyatova, 1982); according to the established pollen-based [LIG](#) climate for western
445 Europe (Zagwijn, 1996), marine sediments at this site span the entire interval from ~133 to ~119.5 ka (Grøsfjeld et al.,
2006). Moreover, molluscs from this site suggest warmer-than-present day conditions (notably, *Corbula gibba* and *Balanus*
improvisus), which also supports a LIG age (Grøsfjeld et al., 2006).

7.15.26.15.2 Bychye

The Bychye site (sometimes spelled “Bychie”; Site 1 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) was first described by Devyatova and Loseva (1964) and again by Devyatova (1982). As described most recently by Grøsfjeld et al. (2006), the stratigraphy at this site is similar to the Zaton site. The entire LIG-marine sequence spans from 12.5 to 23 MASL. At the modern-day river level (~12 MASL) is a till unit, which is overlain by marine clays which gradually coarsen into clayey silt (12.5 to 19 MASL; >45 m water depth). These sediments are terminated by an erosional horizon (at 19 MASL), indicating falling sea level (Grøsfjeld et al., 2006), followed by laminated sand and silt separated by gravel horizons with channel incisions (from 19 to 23 MASL), interpreted as a regression sequence (shoreface/tidal environment; <12 m water depth). Marine molluscs, dinoflagellate cysts and benthic foraminifera (Grøsfjeld et al., 2006) are present throughout. The stratigraphy at this site is laterally continuous for ~500 m.

The marine sediments at the Bychye site are assigned to the LIG based on palaeoenvironmental and stratigraphic inferences. Pollen data (examined by Devyatova, 1982) suggest correlation to the ~133 to ~124-ka interval (Grøsfjeld et al., 2006) LIG climate for western Europe (Zagwijn, 1996). Various marine molluscs, dinoflagellate cysts and benthic foraminifera suggest a transition from cooler-than-present to warmer-than-present temperatures, which supports the capturing of the MIS 6 deglaciation, followed by establishment of LIG warmer marine conditions (Grøsfjeld et al., 2006).

7.15.36.15.3 Viryuga W (65.82, 46.00)

Located on the northern side of the Pyoza River, the Viryuga W site (Site 4 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) was first described by Devyatova and Loseva (1964). Our descriptions are derived from the more recent examination by Houmark-Nielsen et al. (2001) and Grøsfjeld et al. (2006). At the base is a marine unit (spanning 21 to 39 MASL) containing stratified sands with shells. A till is present between 39 and 45 MASL. Capping the stratigraphic section is an upper marine unit (spanning 46 to 49 MASL), consisting of a clayey diamict containing abundant *Mya truncata* and *Macoma calcarea* shells, often paired and thus suggesting *in-situ* preservation. Dinoflagellate cysts and benthic foraminifera are present throughout (Grøsfjeld et al., 2006).

The lower marine unit yielded OSL ages ranging between 237–194 ka, suggesting an ice-free interval during MIS 7 (Houmark-Nielsen et al., 2001). Accordingly, Grøsfjeld et al. (2006) interpreted the upper marine sediments as the earliest part of the LIG. Correlation of pollen data from this site with the climate for western Europe (Zagwijn, 1996) place the upper marine sediment unit between ~133 and ~130 ka (Grøsfjeld et al., 2006). Marine molluscs and benthic foraminifera suggest cooler-than-present conditions during this time, which suggests deposition began during the MIS 6 deglaciation at the beginning of the LIG (Grøsfjeld et al., 2006). However, because the upper marine sediments themselves remain undated and

the stratigraphic context is unclear, earlier workers (Devyatova and Loseva, 1964; Houmark-Nielsen et al., 2001) interpreted
480 this upper unit as Weichselian-aged (e.g. late MIS 5).

7.15.46.15.4 Viryuga E-(65.80, 45.99)

The Viryuga E site (Site 6 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) was first described by Devyatova and Loseva (1964) and again by Grøsfjeld et al. (2006). Following Grøsfjeld et al. (2006), the basal interval at this site is a clayey till between 56 to 58 MASL, suggested inferred to have been laid down emplaced during the MIS 6 glaciation. The
485 till is, with a sharp boundary, overlain by clayey glaciomarine sediment between 58 and 60 MASL, interpreted as deposited during LIG marine inundation and sea-level high-stand following collapse of the MIS 6 ice sheet collapse. Capping the stratigraphy is marine, mollusc-bearing cross-bedded sand with gravel and clay bedding (60 to 63 MASL), interpreted as shoreface deposits. Marine molluscs, dinoflagellate cysts and benthic foraminifera are present throughout (Grøsfjeld et al., 2006). Marine molluscs and benthic foraminifera suggest cooler-than-present conditions during deposition, followed by
490 warmer-than present molluscs; together, these data suggest deposition during the deglacial phase at the beginning end of MIS 6 of the LIG and then into the early LIG (Grøsfjeld et al., 2006). Water depths were likely >45 m at time of deposition. No direct dates are available at this site. However, correlation of pollen data from this site with climate for western Europe (Zagwijn, 1996), along with the stratigraphic context, place the marine sediments between ~133 and ~130 ka (Grøsfjeld et al., 2006).

495 7.15.56.15.5 Kalinov-(65.79, 46.22)

At the base of the The Kalinov site (Site 8 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) displays at the base marine clays are interbedded with sands (situated from 28 and 37 MASL). +The sediment succession was interpreted as deposited in a lower shoreface environment (Grøsfjeld et al., 2006). The marine sediments are capped by fluvial sands from 38 to 40 MASL. The marine clay interval contains marine molluscs, dinoflagellate cysts and benthic foraminifera
500 (Grøsfjeld et al., 2006). The Ecorrelation of pollen data from this site with climate for western Europe (Zagwijn, 1996) places the marine sediment succession between ~133 ka and ~130 ka (Grøsfjeld et al., 2006), although there are no direct dates at this site. Marine molluscs and benthic foraminifera suggest cooler-than-present conditions during this time, which suggest deposition during the deglacial phase at the end of MIS 6 and into the early LIG (Grøsfjeld et al., 2006)during the deglacial phase at the beginning of the LIG (Grøsfjeld et al., 2006).

505 7.15.66.15.6 Yatsevets

At the base of the Yatsevets site (Site 10 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) is a till situated between 32 and 34 MASL. Overlying this till is a sequence of marine clay between 33 and 38 MASL, containing marine molluscs, dinoflagellate cysts and benthic foraminifera (Grøsfjeld et al., 2006). Molluscs and benthic foraminifera suggest cooler-than-present conditions during sediment deposition and with an increasing water depth (>45 m depth), which support deposition

510 during the deglacial phase at the beginning of the LIG (Grøsfjeld et al., 2006). Correlation of pollen data (Devyatova and Loseva, 1964) from this site with the climate for western Europe (Zagwijn, 1996) place the marine sediment deposition between ~133 and ~130 ka (Grøsfjeld et al., 2006).

7.15.76.15.7 Site 11 Orlovets (65.71, 46.84)

The Orlovets site ~~displays contains~~ almost entirely laminated marine silts that coarsen upward (from 38 ~~and~~ 43.5 MASL), 515 the sediment succession interpreted as deposited in a lower shoreface environment (Grøsfjeld et al., 2006). ~~T~~The laminated silts are capped by sand and gravel, interpreted as deposited in a foreshore environment ~~(Grøsfjeld et al., 2006)~~. Numerous marine molluscs, dinoflagellate cysts and benthic foraminifera are present throughout ~~(Grøsfjeld et al., 2006)~~, all suggesting slightly warmer marine conditions than present-day ~~conditions~~ (Grøsfjeld et al., 2006). ~~No direct dates are available for the marine sediments; however, a~~ According to the established ~~Eemian LIG~~ climate for western Europe (Zagwijn, 1996), marine 520 sediments at this site ~~are suggested to potentially span an age interval~~ from ~128 to ~124 ka (Grøsfjeld et al., 2006).

7.15.86.15.8 Site 12 Orlovets (65.69, 46.93)

The stratigraphic record at Site 12 Orlovets is identical to that of Site 11 Orlovets. Laminated marine silts that coarsen upward (from 38 ~~to~~ 43.5 MASL) were interpreted as deposited in a lower shoreface environment (Grøsfjeld et al., 2006). At this site, however, the marine interval is capped by sand, interpreted to be deposited in a fluvial environment (Grøsfjeld et 525 al., 2006).

7.15.96.15.9 Site 13 Yolkino (65.68, 47.60)

Site 13 Yolkino was first described by Devyatova and Loseva (1964) and ~~again later~~ by Houmark-Nielsen et al. (2001) and Grøsfjeld et al. (2006). At the base of the section is marine sand containing molluscs, dinoflagellate cysts and benthic foraminifera, between 48 and 51 MASL. ~~There are no direct age constraints on the sediments at this site. However,~~ 530 ~~e~~Correlation of pollen data from this site with the climate for western Europe (Zagwijn, 1996) place marine sediment deposition between ~130 and ~128 ka, ~~and~~ ~~(Grøsfjeld et al., 2006)~~. ~~T~~he marine molluscs suggest warmer than present-day conditions (Grøsfjeld et al., 2006). The marine sediments are overlain by a series of organic-bearing lacustrine sediments, dated by OSL to 89 ka (Houmark-Nielsen et al., 2001), as well as tills and fluvial sediments.

7.15.106.15.10 Site 14 Yolkino

535 The stratigraphic record at Site 14 Yolkino is identical to that of Site 13 Yolkino. Importantly, marine sand containing molluscs, dinoflagellate cysts and benthic foraminifera are situated at the base of this stratigraphic section between 48 and 51 MASL. An OSL age on these marine sediments suggests deposition at 124 ka (Houmark-Nielsen et al., 2001).

7.15.116.15.11 Burdui

Marine sand at the Burdui site (Site 24 of Houmark-Nielsen et al., 2001 and Grøsfjeld et al., 2006) is situated between 59 and 60 MASL. These sediments are overlain by a 1-m interval of proglacial sand, dated by OSL to 97 ka (Houmark-Nielsen et al., 2001). Marine molluscs, dinoflagellate cysts and benthic foraminifera are present throughout (Grøsfjeld et al., 2006). Analysis of the marine molluscs suggest warmer conditions than present-day (Grøsfjeld et al., 2006). According to the established Eemian LIG climate for western Europe (Zagwijn, 1996), the marine sediments at this site seemingly span an age interval from ~131 to ~130 ka (Grøsfjeld et al., 2006).

545 7.166.16 Ponoi River, Kola Peninsula, Russia (67.078, 41.131)

The Ponoi site is in the Ponoi River ~~v~~Valley on the eastern part of the Kola Peninsula (Fig. 6B). The most complete section in the area is about 20-m thick and ~~has~~was~~been~~ studied by Lavrova (1960), Nikonov (1966), Gudina and Yevzerov (1973), Ikonen and Ekman (2001) and Korsakova et al. (2016). According to the latter two studies, a till at the base of the section (interpreted as deposited during ~MIS 6) ~~at the base of the section~~ is overlain by a marine clay unit (denoted unit 2) with sand and gravel interbeds. This clay unit is situated between 7 and 11 MASL and contains sporadic unbroken mollusc shells and shell fragments. The mollusc, foraminifera, and diatoms, together with palynological and lithostratigraphical evidence, ~~indicate~~suggest ~~that~~that the marine clay can be correlated with the Eemian interglacial stage~~LIG~~ (i.e. LIG; MIS 5-e). Overlying this marine unit are stratified sands and gravels with shell detritus and unbroken shells. This upper unit is 8~~~~~m thick and has a sharp lower contact with the underlying marine clays (Korsakova et al., 2016). Mollusc shells and sand in this upper unit have been dated ~~with~~via ESR and OSL to between 96 ± 8 ka (ESR) and 71.9 ± 8.2 ka (OSL) indicating that this upper sand was deposited during approximately MIS 5 a-c (Early Weichselian substage; ~~see~~ Korsakova et al., 2016). the Early Weichselian substage (approximately MIS 5 a-c; ~~see~~ Korsakova et al., 2016). This age assignment for the uppermost sediments supports and earlier LIG age for the underlying marine clays. Capping the entire stratigraphic section is a glacial unit consisting of glaciolacustrine silt with clast and a till unit (Korsakova et al. 2016).

560 7.176.17 Svyatoi Nos, Kola Peninsula, Russia (68.016, 39.874)

The Svyatoi Nos site is located on the northeastern coast of the Kola Peninsula (Fig. 6B). As described by Korsakova (2019, 2021), the MIS 6 (known locally as the Moscovian glaciation) glaciomarine sediments are overlain by marine mollusc-bearing silty sands that are present between 11 and 16 MASL. The mollusc species indicate sublittoral faunal assemblage (e.g. Ikonen and Ekman, 2001) and temperate saline water conditions (e.g. Korsakova, 2019). Based mainly on mollusc, foraminifera, and pollen results, the silty sands are thought to have been deposited during the LIG (Gudina and Yevzerov, 1973; Ikonen and Ekman, 2001; Korsakova, 2009, 2019) despite the slightly younger IRSL age of 109.9 ± 10.9 ka obtained from these sands.

7.186.18 Chapoma, Kola Peninsula, Russia

In the southeastern part of the Kola Peninsula (Fig. 6B), exposures at the Chapoma site occur in the river terraces on the 570 bank of the River Chapoma, about 3.4 km from the river mouth (Gudina and Yevzerov, 1973). The exposure is approximately 25-m high. A till bed at the base is suggested to have been deposited during MIS 6 (known locally as the Moscovian glaciation the Moseovian glaciation (–MIS 6–); see Gudina and Yevzerov, 1973; Korsakova et al., 2004; Korsakova, 2019). Relevant to the present study, the till bed is overlain by clay, silt and sandy silt beds that spans 6.5–10 575 MASL, and with abundant mollusc shells (Korsakova, 2019). The foraminifera fauna identified from these marine sediments include a rich and relatively warmwater-indicating fauna with species such as *Bulimina aculeata*, *Bolivina pseudoduplicata* and *Hyalina baltica* (e.g. Gudina and Yevzerov, 1973; Ikonen and Ekman, 2001). Pollen data from the marine sediments indicate a succession from a closed to open *Betula–Pinus* forests. Electron spin resonanceSR dating obtained from molluscs at ~~around~~ 9 MASL yielded ages of 128 ± 7.5 ka and 138.5 ± 9.6 ka, which supports deposition during the LIG. An additional marine bed at this section post-dates the LIG and was likely deposited during MIS 5a (see Discussion).

580 7.196.19 Strelna River, Kola Peninsula, Russia (66.911, 33.637)

Located just south of the Kola Peninsula, the Strelna site is situated approximately 7 km upstream from the mouth of the River Strelna (Fig. 6B). Many ~~workers~~researchers, for example Grave et al. (1969) have studied the litho- and biostratigraphy for this >16-m thick sediment succession. The most recent summary of ~~the~~ previous work is given in Korsakova (2019). At the base of this site is a 2.5-m-thick unit of mollusc-bearing fine sand and silt situated between 33 and 585 35.5 MASL. The ESR-dated molluscs from this unit yielded an age of 111.5 ± 11.2 ka. Taken together, the mollusc and diatom assemblages suggest that the basal marine sediments were deposited in a coastal environment, possibly indicating a ~~contemporary~~ sea level above 36 MASL (Korsakova et al., 2016). Pollen evidence suggest that a *Betula/Pinus* ~~birch and pine~~ forests existed in the area (Grave et al., 1969). Accordingly, this unit is most likely correlative with the LIG (–MIS 5e–) (e.g. Korsakova et al., 2004; Korsakova, 2019).

Formatted: Font: Italic

590 7.206.20 Varzuga, Kola Peninsula, Russia (66.399, 36.641)

Exposures along the banks of the Varzuga River between the Koytolov and Kletnoy rapids (south of Varzuga village) have been studied by Lavrova (1960), Gudina and Yevzerov (1973), Apukthin (1978) and Lunkka et al. (2018). An exposure located on the right bank of the Kletnoy rapid (Fig. 6B) consists of a marine mollusc-bearing clayey silt at the base of the section (situated between 10 and 14 MASL), conformably overlain by glaciolacustrine 595 silt and sand-rich silt (Site 'S1' of Lunkka et al., 2018). Pollen, diatom and foraminifera indicate that the marine silt was deposited in a sublittoral zone at water depths of 40–50 m during an interglacial stage, correlated with the LIG (Ikonen and Ekman, 2001, and references therein; Lunkka et al., 2018). Although the sedimentary sequence along the banks of the Varzuga River is glaciotectonized in places (Apukthin, 1978), the marine clay/silt situated between 10 and 14 MASL is in

600 *situ*, where it is conformably overlain by glaciolacustrine silt at around 14 MASL. These overlying glaciolacustrine silts are thought to have been deposited during a later stage of MIS 5, prior to 88 ka (Lunkka et al., 2018). Capping the stratigraphic sections are two till units interbedded with sand and silt (Lunkka et al., 2018).

7.216.21 Petrozavodsk, Western Russia (61.77, 34.40)

605 Marine sediments assigned to the LIG were first noted in a borehole record from Petrozavodsk, western Russia (Fig. 6B), described by Wollosovich (1908). These sediments are located at 40 MASL and conformably overlie glaciolacustrine clays (Wollosovich, 1908; Lukashov, 1982). A marine origin for these sediments is confirmed by saline diatom taxa (Ikonen and Ekman, 2001) and several mollusc species, suggesting saline conditions on the order of 10 to 15 % (Funder et al., 2002). ~~No geochronological data are available for these sediments; instead, the~~ The LIG-age is based on correlation ~~of to~~ to the local pollen record ~~to an established Fennian (MIS 5e; LIG) pollen assemblage~~ (Lukashov, 1982; Ikonen and Ekman, 2001). Based on the pollen taxa and its correlation to the saline diatom taxa, Ikonen and Ekman (2001) showed that the marine phase 610 prevailed in the area for a long time during the LIG, i.e. from ~~the Pinus-Betula PAZ to the Picea-Alnus-Carpinus PAZ~~ (possibly between 130 ~~ka to~~ 124 ka).

7.226.22 Peski, Western Russia (60.15, 29.29)

615 Miettinen et al. (2002) described a 32-m borehole record from Peski, western Russia (Fig. 6B). Sediments between 9 and 13.5 MASL were correlated to the LIG. The base of the borehole is a till (interpreted as deposited during the MIS 6 glaciation), followed by a dark bluish, organic-bearing clay and silt deposit containing *Portlandia arctica* molluscs between 10.5 and 11.5 MASL, interpreted as deposited during relatively deep-water marine conditions. The marine deposit is assigned to the LIG based on palaeoecological and stratigraphic context. Pollen data from this interval suggest warm conditions, especially the occurrence of *Corylus* and *Carpinus*, which are associated with the climatic optimum of the LIG (Miettinen et al., 2002). Diatoms contained in the marine unit suggest relatively 620 deep-water, planktonic conditions, possibly representing the maximum of the marine transgression (Miettinen et al., 2002). Marine conditions were confirmed by the subsequent identification of dinoflagellate cysts and foraminifera in these sediments (Miettinen et al., 2014). The LIG unit gradually transitions to later-stage MIS 5 deposition, which is distinguished by a change to grey colour and cooler conditions deduced from pollen and diatoms. This transition 625 likely indicates that sea level remained above 13.5 m throughout the entire LIG. Stratigraphically, the marine unit is overlain by a 16-m thick till associated with the advance of MIS 5d/b ice sheets (Miettinen et al., 2002).

7.236.23 Põhja-Uhtju, Estonia (59.68, 26.51)

Miettinen et al. (2002) described a Quaternary sediment sequence in a 70-m deep borehole from the Põhja-Uhtju island in Estonia (Fig. 6B). The base of the recorded stratigraphy is composed of sands and silty clay, interpreted as deposited during the MIS 6 deglaciation. Overlying these sediments, between 51–49 MBSL, is clay and silt, interpreted as marine in origin

Formatted: Font: Italic
Formatted: Font: Italic
Formatted: Font: Italic
Formatted: Font: Italic
Formatted: Font: Italic

630 and suggesting a sea level above the present at that time. Above these are ~35 m of silty clays and tills, associated with later ice advances. The 51–49 MBSL interval is assigned to a LIG marine incursion, an interpretation based on their palaeoecological and stratigraphic context. Pollen data, especially the occurrence of *Corylus* and *Carpinus*, suggest warm conditions at the climatic optimum of the LIG (Miettinen et al., 2002). The diatom assemblage in these sediments suggest shallow marine to brackish water conditions.

635 Miettinen et al. (2002) describe a Quaternary sediment sequence in a 70-m deep borehole from the Põhja-Uhtju island in Estonia. The base of the record showed sands and silty clay, interpreted as deposited during the deglaciation of MIS 6 ice. Next, located at 51 to 49 MBSL, was clay and silt, interpreted as marine in origin, suggesting that contemporary sea level could have been above present sea level. Overlying these marine sediments were ~35 m of silty clays and tills associated with later ice advances. The interval of 51 to 49 MBSL is assigned to the MIS 5e marine incursion based on palaeoecological and stratigraphic context. Pollen data suggest warm conditions, especially *Corylus* and *Carpinus*, which are associated with the climatic optimum of MIS 5e (Miettinen et al., 2002). The diatom assemblage in these sediments suggest shallow, marine to brackish water conditions. No geochronological attempts were made at this site.

640 Similar to the nearby island of Põhja-Uhtju, a borehole at Suur-Prangli, Estonia (Fig. 6B) records marine sediments assigned to the LIG. Liivrand (1987) reported a silt/clay sediment succession between 61 and 75 MBSL that is both overlain and underlain by tills. The diatom record suggests brackish-water conditions, followed by a shallow marine environment (Liivrand, 1991). This assignment to the LIG was based on stratigraphic position (bracketed by tills assigned to MIS 6 and MIS 4) and paleoclimate succession recorded in pollen composition, notably a maximum *Picea* and *Carpinus* interval (Liivrand, 1991). As further justification for the LIG age assignment, the pollen assemblage observed at Suur-Prangli show a similar succession to nearby pollen successions (Forsström and Punkari, 1997) and a well-dated LIG pollen record from 645 Germany (Field et al., 1994).

650 Similar to the nearby island of Põhja-Uhtju, marine sediments assigned to MIS 5e have also been reported from a borehole record at Suur-Prangli, Estonia. Liivrand (1987) reported a silt/clay sediment succession between 61 and 75 MBSL that is both overlain and underlain by tills. The diatom record suggests brackish conditions followed by a shallow marine environment (Liivrand, 1991). These marine sediments are assigned to MIS 5e based on stratigraphic position (bracketed by 655 tills) and pollen assemblage (Liivrand, 1987). The MIS 5e pollen assemblage is dominated by *Picea* and *Carpinus*.

Extensive late Pleistocene stratigraphic records in the Lower Vistula region of Poland were first described by Roemer (1864). We describe two representative sites that later have been subject to extensive stratigraphic, sedimentological and palaeoecological analyses. However, we note that other additional occurrences of the MIS 5e LIG marine records from this area are presented in are also present in this area (see Makowska (–1986).

660 **7.27.16.25.1 Obrzynowo**

A 212-m borehole record from Obrzynowo, Poland (Fig. 6B), was first described in an unpublished report from the Polish Geological Institute at Warsaw University (Janczyk-Kopikowa and Marks, 2002), and was subsequently published in Knudsen et al. (2012). This borehole record (ground surface is at 104.5 MASL) contains a sequence of glacial and interglacial deposits and intersects marine sediments between 108 and 115 m depth, which situates the entire marine sequence between 10.5 and 3.5 MBSL (Janczyk-Kopikowa and Marks, 2002; Knudsen et al., 2012). This record captures a beach/shoreline environment from 10.5~~to~~6.5 MBSL (fine-grained sand and silt with mollusc shells) that gradually transitions into deeper-water marine deposit from 6.5~~to~~3.5 MBSL (clays, silts and very fine sands). Dinoflagellate cysts and diatoms confirm saline conditions (Knudsen et al., 2012). These marine sediments are both overlain and underlain by tills, fluvial sediments, varved clays and organic-rich sediments and this site likely documents several glacial-interglacial transitions over the late Quaternary. The marine sequence at Obrzynowo has been assigned to MIS 5e the LIG based on its stratigraphic position and a comparison with the Licze core (discussed next; Head et al., 2005), as well as comparison to other regional pollen successions that have been assigned to the Eemian-LIG (an interpretation especially based on the presence of *Picea* and *Carpinus*; Mamakowa, 1989, 1988).

7.27.26.25.2 Licze (53.75, 19.13)

675 A borehole record at the Licze site (ground surface at 87 MASL; Fig. 6B) records a transition from freshwater to marine conditions between 8–14.5 MBSL (Makowska et al., 2001; Zawidzka, 1997) in which there is freshwater molluscs that are gradually replaced by a marine assemblage. The first marine molluscs are present at 13.5 MBSL and they remain common until 8 MBSL (Mamakowa, 1989).

Dinoflagellate cysts and diatoms confirm marine conditions through this stratigraphic interval and suggest warmer waters
680 Dinoflagellate cysts and diatoms confirm marine conditions through this stratigraphic interval and suggest warmer waters
Dinoflagellate cysts and diatoms confirm marine conditions through this stratigraphic interval and suggest warmer waters
Dinoflagellate cysts and diatoms confirm marine conditions through this stratigraphic interval and suggest warmer waters
than present-day conditions (Head et al., 2005; Knudsen et al., 2012). Throughout this interval with sediment transition from
685 silty sand to silt and clay suggests an environmental change (transgression) from a beach environment to deeper marine
conditions (Makowska et al., 2001). These marine sediments are both overlain and underlain by extensive Quaternary
deposits (tills, fluvial deposits, varved clays and organic-rich sediments). These marine sediments are assigned to the LIG
owing to their stratigraphic position, along with a comparison to the pollen record at the Obrzynowo site (Head et al., 2005)
and to other regional pollen successions, assigned to the LIG (especially due to the presence of *Picea* and *Carpinus*;
Mamakowa, 1989, 1988)

690 **7.286.26 Rewal coastline, Poland (3 sites)**

Marine sediments bracketed by tills were described in several borehole records along the Rewal coastline, Poland, by Krzyszkowski et al. (1999) as well as by Krzyszkowski (2010). The marine sediments consist of fine-grained sand (known as the Rewal Sand) containing a rich assemblage of plant detritus and marine shells (largely boreal *Cardium* sp. (now *Cerastoderma*), but also *Astarte borealis*, and the boreal *Thracia popyracea*; Krzymińska, 1996). Krzyszkowski et al. 695 (1999) suggested that these sediments represent deposition either a lagoon or a beach setting. They were environment and they are assigned to MIS 5e LIG based on their stratigraphic position. Both sites are overlain by the Ninikowo Till (assigned to the last glacial cycle) and underlain by the Pustkowo Till (assigned to MIS 6 glaciation; Krzyszkowski et al., 1999). Here, we describe the position of LIG marine sediments in three representative boreholes from this region (sites mapped in Fig. 6B).

700 **7.28.16.26.1 Rewal borehole (54.09, 15.03)**

The Rewal borehole (WH5) encountered a marine sand unit at an elevation between 5.5 and 13 MBSL (Krzyszkowski et al., 1999).

7.28.26.2 Ciećmierz borehole (53.99, 15.03)

The Ciećmierz borehole intersected a thin lens of the Rewal Sands situated between 6.5 and 8 MBSL. Additional pollen 705 work at this site suggests boreal conditions at the time of deposition of this marine unit (Krzyszkowski, 2010).

7.28.36.26.3 Sliwin borehole (54.08, 15.01)

The Sliwin borehole (WH4) encountered the marine unit between 6.3 and 9 MBSL (Krzyszkowski et al., 1999).

7.296.27 Ollala, Finland

Forsström et al. (1988) described a sediment exposure and four boreholes at Ollala in central Finland (Fig. 6B) on their 710 lithology, pollen, plant macrofossil and diatom content. At borehole F, there is a till (>4 m thick) at the base above the Proterozoic crystalline bedrock. Overlying this till are < 1 m of glaciofluvial sands, which then change into silt (situated between 116.25 and 116.5 MASL) and then thick greenish gyttja (situated from 116.5 to 117.5 MASL). The silt and gyttja intervals (together comprising 116.25 to 117.5 MASL) contain diatom taxa indicating a shallow marine (saline and brackish water) basin passing upwards into fresh-water diatom taxa, typical for a lake basin. This change from saline to fresh-water 715 taxa takes place in the lower part of the gyttja layer at 117 MASL, indicating that the fresh-water lake was isolated from the marine basin during the latter part of the MIS 5e LIG (LIG), likely the result of glacio-isostatic GIA uplift. Pollen data from this interval also clearly suggest that these sediments were deposited during interglacial conditions (Grönlund, 1991a, 1991b; Nenonen, 1995). There are no finite absolute dates from the Ollala sediments. The gyttja unit is overlain by 0.5-m of

glacially sheared and compressed organic-rich material that is, in turn, overlain by up to 5-m thick Weichselian till
720 (Forssström et al., 1987; Grönlund, 1991a). This marine gytta unit is also present between 116.6 and 116.8 MASL at nearby borehole B (Forssström et al., 1987; Grönlund, 1991a).

7.306.28 Ukonkangas, Finland (63.916, 25.851)

The Ukonkangas site is a till-covered esker, located 8 km southeast of Kärsämäki, central Finland (Fig. 6B). According to Grönlund (1991b) the Ukonkangas gravel pit exposed gravel at its base, overlain by ~0.5-m-thick bluish, organic-bearing silt (situated between 105.4 and 105.95 MASL). This silt bed is laterally continuous for 15 meters in the exposure, suggesting it is *in situ*. Grönlund (1991b) interpreted the silt bed as deposited in a littoral zone of the Eemian (LIG) LIG White Sea (Fig. 13), with time becoming more shallow and less saline. Pollen content indicates that the regional vegetation was forested and composed of broad-leaved trees and other pollen (e.g. *Osmunda*) indicating Eemian interglacial temperate climatic conditions (Eriksson, 1993). The diatom taxa are typical to the Eemian LIG White/Baltic Sea taxa, which 730 also occur at the Eemian-LIG sites in adjacent areas (Grönlund 1991a, b). The silt unit is overlain by parallel bedded sand, interpreted as beach sand (situated between 105.95 and 108 MASL; Grönlund 1991b). The top 2 metres of the section is composed of till (MIS 2?). No geochronological data are available for this site.

7.316.29 Viitala, Finland (62.598, 23.002)

At the Viitala site, located in western Finland (Fig. 6B), three borehole records documented a clay beneath the MIS 2 till (Nenonen et al., 1991). The clay unit is ~5 m thick. Diatom taxa indicate that the lower ~2 m of the clay was deposited in a large cool and freshwater basin during the early part of the LIG, and that the upper ~3 m was deposited in a brackish/saline littoral zone of the LIG White/Baltic Sea (e.g. Grönlund, 1991a; see Fig. 1). These marine sediments are situated between 81.5 and 84.5 MASL. Pollen data from the entire clay unit suggest a lower birch-dominated zone (~2 m; freshwater sediments) and an upper birch–pin–oak–hazel-dominated zone (~3m; marine 740 sediments). The interglacial-type pollen taxa and the stratigraphical position of the clay suggest that the clay unit was deposited during the LIG (Nenonen et al., 1991; Eriksson, 1993).

7.326.30 Mertuanaja, Finland (64.113, 24.586)

The Mertuanaja region is in the outskirts of Ylivieska, Finland (Fig. 6B), and is the stratotype area for the Eemian interglacial LIG (LIG; MIS 5e) of southern Finland (Nenonen, 1995; Eriksson et al., 1999). We present a composite of several sediment exposures studied in the Mertuanaja area. At the base of the stratigraphy is a till, deposited during the Saalian glaciation (MIS 6) (e.g. Nenonen, 1995; Lunkka et al., 2016). Overlying this till is the Mertuanaja Clay (average

750 thickness ~1.6 m), consisting of laminated silt and organic-bearing silty clay in its basal part overlain by a sand layer (2–20 cm thick), which is, in turn, overlain by laminated silt. Diatom data suggest deposition in a fresh-water environment in the basal part of the Mertuanaja Clay, changing to marine/brackish water, and then back to fresh-water conditions at deposition of the laminated silt and clay above the sand layer (Nenonen, 1995; Eriksson et al., 1999). Relevant to the present study is the transition from the fresh-water diatom taxa at the basal part of the Mertuanaja Clay into the marine taxa that takes place
755 at around 59 MASL. Coastal marine conditions are recorded in this stratigraphic record until the sand layer at 60–61 MASL, interpreted as littoral sand. The pollen spectra from the Mertuanaja Clay suggest that it was deposited during interglacial conditions correlated to the LIG (e.g. Grönlund, 1991a; Nenonen, 1995). A sand and gravel unit overlying the Mertuanaja Clay was OSL dated to 110 ± 7 ka (Lunkka et al., 2019), followed by tills and an upper-most sand and gravel unit dated by radiocarbon methods to between 70 and 35 ka (Lunkka et al., 2016).

760 **7.336.31 Norra Sannäs, Sweden (61.7830, 16.6931)**

Sub-till marine sediments were intersected in boreholes in the vicinity of Lake Dellen, central Sweden (Fig. 6B), during field work carried out in the late 1980s and early 1990s. Robertsson et al. (1997) described a representative borehole record from Norra Sannäs (surface at 45 MASL) that shows a marine (littoral) sequence at 17.35-m depth in the core, which translates to 27.65 MASL. The diatom assemblage in clay sediments suggests deposition during marine to brackish conditions, which
765 abruptly change to fresh-water diatoms at 17.1-m depth, suggesting the end of marine conditions (Robertsson et al., 1997). Stratigraphically, the marine sediments are overlain and underlain by tills that have been assigned to MIS 4 and MIS 6, respectively, thus suggesting a LIG age for the marine sediments. This age assignment is also supported by pollen data (Robertsson et al., 1997), which bear a similar succession to other LIG records from the region. The sole geochronological attempt at this site was a radiocarbon dating of the marine sediments that
770 gave an age of 25.48 ka (Ua-1666); Robertsson et al. (1997) considered this age unreliable owing to a very low organic content at the sampled interval.

7.346.32 Fjøsanger, Norway (60.3431, 5.330)

The Fjøsanger site, located in the outskirts of Bergen City (Fig. 6B), is situated along a small fjord, well inside the extent of the Scandinavian Ice Sheet during large Quaternary glaciations, including MIS 6 and MIS 2. The marine limit was 55 MASL
775 during the last deglaciation, dated to 11.5 ka 11,500 cal years BP (Mangerud et al., 2019). Relevant to the present study, at Fjøsanger there is a continuous sequence of shallow marine sediments, presently situated between 0 and 15 MASL, where molluscs, foraminifera, and pollen show that the climate changed from cold to warmer-than-present conditions and back to cold (Mangerud et al., 1981). The sequence is covered by a till. The fauna, especially the boreal *Littorina littorea*, and the coarse-grained sediments (gravel) suggest very shallow-water sediments, but beach sediments were not found; thus, sea level
780 was has been slightly higher. Correlation of the pollen stratigraphy with sites in the Netherlands, Denmark and Germany shows clearly that the warm period represents the Eemian LIG, a conclusion supported by the occurrence of the marine

gastropod *Bitium reticulatum*, which is known only from the LIG sediments in Europe (Mangerud et al., 1981), and by TL ages (Hütt et al., 1983). Amino-acid stratigraphical correlation with classical European LIG sites yielded slightly high D/L values for Fjøsanger, but within uncertainty (Miller and Mangerud, 1985). A surprising, and indeed important, conclusion is 785 that the RSL at Fjøsanger was above 15 MASL from late MIS 6 through the entire LIG and into MIS 5d (Fig. 2) (Mangerud et al., 1981). We consider this as a secure conclusion because the Fjøsanger sequence shows a complete LIG succession and 15 MASL is the elevation of the top of the *in situ* marine beds. Fjøsanger was one of the first sites where it was demonstrated that the LIG (Eemian) should be correlated with MIS 5e, and not with the entire MIS 5 (Mangerud et al., 1979).

7.356.33 Bø, Norway (59.3622, 5.276)

790 The Bø site is located on the eastern side of the large island Karmøy, on the extreme south-west coast of Norway (Fig. 6B). Like the Fjøsanger site, Bø is situated well inside the extent of the Scandinavian Ice Sheet during large Quaternary glaciations, including MIS 6 and MIS 2. The marine limit was about 16 MASL during the last deglaciation, dated to 17.5 ka (Vasskog et al., 2019). Relevant to the present study, interglacial sediments were found between 1 and 6 MBSL in an excavation, but only the upper 3 m could be sampled (Andersen et al., 1983). The interglacial sequence 795 was covered by till and other sediments. The pollen stratigraphy (Andersen et al., 1983; Høeg, 1999) and amino-acid stratigraphy (Miller and Mangerud 1985; Sejrup, 1987) clearly shows that this site is LIG in age. The very early part of the interglacial is missing in the samples, but molluscs, foraminifera (Sejrup, 1987) and pollen show that the warmest part is present. Andersen et al. (1983) concluded that LIG RSL was more than 15–20 MASL because they postulate an open sea-connection across the island. Sejrup (1987) 800 described paired shells of the boreal molluscs (*Lucinoma borealis* and *Pecten maximus*); these presently live-in water depths of 20–50 m and he postulated that sea level was this high during the LIG, however at lowest at the end of the interglacial. While we cannot determine a precise RSL at Bø, the results support the conclusion from Fjøsanger, namely that sea level in western Norway during the entire LIG was at least 15 m higher than the current sea level.

7.366.34 Hidalen, Svalbard, Norway (78.90, 28.13)

805 The Hidalen site is in eastern Svalbard, on the island of Kongsøya (Fig. 8B). The marine limit from the last deglaciation is about 100 MASL (Salvigsen, 1981). The stratigraphy at Hidalen contains three coarsening-upward sequences from marine silt to littoral gravel, separated by till beds (Ingólfsson et al., 1995). The youngest coarsening-upward sequence (Unit F) is of Holocene age, whereas the two older have non-finite ¹⁴radiocarbon ages. The oldest (Unit B) obtained a TL age of 148 ka and combined with the amino acid D/L ratio, Ingólfsson et al. (1995) considered Unit B to be of pre-Eemian LIG age. Based 810 on the amino acid D/L values they argued that the younger Unit D is of Eemian-LIG (MIS 5e) age. Mangerud et al. (1998) presented two alternative interpretations, the first being the one by Ingólfsson et al. (1995). However, they argued that it was more probable that the oldest Unit B is of Eemian-LIG age. Kongsøya is a national park with strong regulations, and no Quaternary geologist has later been allowed to go ashore to solve this problem. For the present interest, we state that MIS

815 LIG marine sediments are present on Kongsøya and they span ~50 to >60 MASL. If Unit D is indeed of LIG age, then sea level was up to ~80 MASL. In our database, we have added both units, and note the controversy in age.

7.376.35 Kapp Ekholm, Svalbard, Norway (78.55, 16.55)

The Kapp Ekholm site ([mapped in Fig. 58B](#); [field photographs in Fig. 9](#)) is in central Svalbard, ~14 km outside the large Nordenskiöldbreen glacier, which occupies the head of Isfjorden-Billefjorden. This implies that if Kapp Ekholm was ice free, glaciers on Svalbard were not much larger than at present. The marine limit during the last deglaciation was 90 ~~m~~ 820 ~~as.1~~ MASL (Salvigsen, 1984) and [the stratigraphy at Kapp Ekholm suggests that](#) this site was repeatedly covered by ice during the Quaternary. The site was first described by Lavrushin (1967, 1969) who found the subarctic mollusc *Mytilus edulis* within one of the marine formations (Fig. 610). *Mytilus* requires warmer climate on Svalbard than during the 20th century (Mangerud and Svendsen, 2018). Kapp Ekholm is the only pre-LGM site on Svalbard where *Mytilus* or other “warm-water” molluscs are found.

825 Several marine and glacial units are found in superposition at Kapp Ekholm ([stratigraphy shown in Fig. 9](#)). The entire exposure is described in detail by Mangerud and Svendsen (1992); here we will only describe the formation containing *Mytilus* (‘Formation B’), which is situated between 5 and 22 MASL. The lower ~5 m of Formation B contains thick marine mud with numerous floating stones (called ‘marine diamicton’). The stones have probably rolled down the steep slope from 830 the shore. Molluscs, including thick *Mya truncata*, in living position, are common and paired shells of subarctic *Mytilus edulis* are also found in this unit (Fig. 6). The mud is overlain by thin and almost horizontal sand beds, capped by 12 m thick, steeply dipping gravel foresets. The latter are interpreted as formed by long-shore drift, and the top (22 MASL) therefore reflects sea level during the formation. The entire Formation B suggests a shallowing during deposition and thus sea level was considerably higher than 22 MASL when the mud and sand were deposited. Chronologically, Formation B is dated with 835 five OSL dates that yielded an average of 118 ka (Mangerud et al., 1998), suggesting a LIG age. Moreover, the frequent occurrence of the “warm-water” demanding *Mytilus edulis*, suggests that Formation B should be correlated with an interglacial in northern Europe when sea water was warmer than at present. Amino acid D/L values also strongly suggest that Formation B cannot stem from an older interglacial.

7.386.36 Skilvika, Svalbard, Norway (77.57, 14.44)

840 In southwestern Svalbard, Skilvika is located on [the shore of Bellsund](#), western Spitsbergen ([Fig. 8B](#)). The site was first described by Semevskij (1967), and later studied by several scientists. A detailed description was provided by Landvik et al. (1992). The entire Quaternary exposure is ~~one-1~~-km long and stretches up to 35 MASL. Most of the exposure consists of 845 Formations 3 and 4, which are the candidates for [MIS 5e-5e LIG](#) age and together span 15 to 30 MASL ([stratigraphy summarize in Fig. 711](#)). Formation 3 is a coarsening upward sequence from glaciomarine silt, through sand (shoreface) to cross-bedded gravel (foreshore). The overlying Formation 4 consists of a strange sediment, namely foresets of boulders and

containing marine fossils in sand lenses between the up to 2-m large boulders. The foresets of Formation 4 interfinger with the beds in Formation 3 (Figs 11 and 12). The interpretation is that Formations 3 and 4 together represent a prograding beach showing a sea level about 30 m above the present (Landvik et al., 1992). Formation 4 was formed by an advance of the local glacier. There were found neither foraminifera (Lycke et al., 1992) nor molluscs (Landvik et al., 1992) that require 850 as warm water as at present and based on TL ages and amino acid D/L values Landvik et al. (1992) concluded a MIS 5c age, which was also accepted by Mangerud et al. (1998), although with the comment that “an Eemian age cannot be ruled out”. Alexanderson and Landvik (2018) then performed an extensive dating program, obtaining 20 OSL ages from Formation 3. The result was large spread in ages from 66–263 ka, although when excluding two outliers the range was 99 ± 7 to 149 ± 17 ka, with a mean of 119 ± 5 ka (n=18). After another quality screening, they obtained a mean age of 118 ± 7 ka 855 (n=10). The conclusion is that Formations 3 and 4 probably are of LIG age, although warm-water species are missing, and a MIS 5c age cannot be excluded. The RSL was about 30 MASL.

7.396.37 Kongsfjordhallet, Svalbard, Norway (79.03, 11.88)

The site Kongsfjordhallet is located on the north shore of Kongsfjorden, northwestern Spitsbergen (Fig. 8B). Marine limit during the last glaciation was about 40 MASL (Lehman and Forman, 1992). These exposures have been studied by 860 several scientists, apparently first by Boulton (1979). The oldest Quaternary marine units are probably about 1 million years old (Houmark-Nielsen and Funder, 1999). In this review we mainly rely on results reported by Alexanderson et al. (2018). Relevant to the present study, units 4 and 5 are considered to be of LIG age. The altitude of these units varies along the ~700 m studied section of the coastline, but they are largely present from 27–34 MASL. The base of these sediments is ~1.5 m thick glacimarine mud. This is overlain by up to 3-m-thick littoral sand and gravel, suggesting a 865 shallowing due to glacial isostatic uplift. The assumed LIG age is based on two OSL dates with an average of 132 ± 7 ka, supported by a diverse foraminifera fauna (Alexanderson et al., 2018). This age is also consistent with the chronology of the entire sequence. We find it probable that these units are of LIG age, although based on two OSL ages only. The RSL was about 34 MASL when the littoral sands and gravels were deposited. However, the units are parts of an emergence cycle; thus, sea level was initially higher than indicated by the littoral sediments.

Formatted: Not Highlight

870 7.406.38 Poolepynten, Svalbard, Norway (78.45, 11.66)

The Poolepynten site is located on the island Prins Karls Forland on the west coast of Spitsbergen (Fig. 8B). The marine limit from the last deglaciation is about 40 MASL (Forman, 1990). The site was first described by Miller (1982) and later in more detail by Bergsten et al. (1998), Andersson et al. (1999) and Alexanderson et al. (2013). The site has attracted much interest because the oldest known bones of polar bear (*Ursus maritimus*) was discovered here (Ingólfsson and Wiig, 875 2009; Lindqvist et al., 2010). In the present review, we mainly rely on results reported by Alexanderson et al. (2013). The exposure shows a sequence of several marine and glacial units with ages between 10–130 ka BP as dated with different geochronological methods. The lowermost unit (named A1 in Alexanderson et al., 2013) is considered to be of MIS-5eLIG

age, and is the only unit discussed here. It is exposed between 2 and 4 MASL and consists of thin sand beds with scattered pebbles, shell fragments and kelp, and it is interpreted as shallow marine or sublittoral deposits. The polar bear jaw was
880 found in this unit.

The MIS-5eLIG age of unit A1 is based on several lines of evidence: the stratigraphical position, amino acid D/L values (Miller, 1982), slightly warm foraminifera fauna (Bergsten et al., 1998), the phylogenetic position of the polar bear mandible (Lindqvist et al., 2010), and two robust OSL ages giving of 118 ± 13 and 105 ± 10 ka (Alexanderson et al., 2013). We
885 consider it very probable that Unit A1 is of MIS-5eLIG age and that it indicates a relative sea level RSL of 2–4 MASL. However, as with all other sites on Svalbard, the unit is interpreted as part of an emergence cycle caused by glacio-isostatic GIA uplift. Thus, the mentioned sea level does not show the MIS-5eLIG high-stand and possibly neither the low stand.

7.416.39 Leinstranda, Svalbard, Norway (78.88, 11.55)

890 The Leinstranda site is located on the northwest coast of Spitsbergen (Fig. 8B). The site was discovered by Troitsky et al. (1979) and later studied by several scientists, in most detail by Miller et al. (1989) and
895 Alexanderson et al. (2011b). A synthesis and critical review was provided by Alexanderson et al. (2011a). The marine limit from the last deglaciation is 46 MASL (Forman, 1990). The sequence consists of several formations of marine, beach and glacial sediments dated to 200–10 ka with different geochronological methods. The units considered to be of LIG age are found between 16 and 19 MASL. The lower part consists of 2-m-thick glaciomarine silt considered to reflect a sea level about 80 m higher than at present. The sediments are coarsening upwards to shore-face sand reflecting a sea level of 20 MASL at its formation. Almost all dated samples are collected from the shore-face sand. Marine mollusc shells, some being paired, are found in all units. All species are common on Svalbard today.

900

Miller et al. (1989) found a foraminifera fauna that suggested slightly warmer conditions than at present and named the unit for the Leinstranda interglacial. Their amino-acid results suggested an Eemian LIG age, which subsequently was supported by IRSL dating (Forman, 1999). We consider four recent OSL dates, with an average of 129 ± 10 ka, to provide the most reliable age (Alexanderson et al., 2011a; Alexanderson et al., 2011b); but we appreciate that several methods agree on a MIS-905 5e age. The conclusion is that Leinstranda represents a MIS-5eLIG site. The sea level was above 20 MASL when the dated bed was deposited and sea level was thus considerably higher before. Whether sea level later in the MIS-5eLIG dropped even lower due to glacial isostatic uplift is unknown.

Formatted: Not Highlight

7.426.40 Galtalækur site in Jökulhlaup valley, Iceland

Van Vliet-Lanoë et al. (2018) describe, in central southern Iceland, marine sediments exposed in deeply incised valleys (Rangá, Þjórsá and Hvítá rivers) that formed because of late glacial jökulhlaups (flooding events) from nearby glaciers (Fig. 13B). The Rangá Formation is around 30-m thick and records two separate marine transgressions following the MIS 6 deglaciation of the region. The marine unit (represented by the regional stratigraphic unit R-C2) consists largely of sands draped by thin silt sediment, representing a coastal fluvial setting, where a tidal delta formed in an estuary (Van Vliet-Lanoë et al., 2018). It is best preserved at the Galtalækur site, where it is located at approximated 120 MASL (extracted from a stratigraphic plot; exact measurements at that site were not provided). The Rangá Formation sediments are assigned to MIS 5e the LIG based on stratigraphic context, absolute dating and tephrachronology. From a stratigraphic standpoint, the Rangá Formation is overlain by extensive deglaciation deposits (tills, glaciofluvial and glacial marine sediments) that are dated to the Holocene via tephrachronology (presence of Vedde ash; Van Vliet-Lanoë et al., 2018). Contained in these marine sediments is a tephra layer from the Grimsvötn volcano which is dated to MIS 5e the LIG based on its position in North Atlantic marine cores (Davies et al., 2014). Both underlying and overlying units that bracket the Rangá Formation have absolute ages. Van Vliet-Lanoë et al. (2018) presented three K-Ar ages on the glacio-volcanic unit that underlies the Rangá Formation that yielded ages between 155 ka and 129 ka. Two other ages from this unit (using the $^{39}\text{Ar}/^{40}\text{Ar}$ method) also yielded ages in this range (Clay et al., 2015; Flude et al., 2008).

7.436.41 Region of Scoresby Sund, East Greenland (8 sites)

Last Interglacial sites in the region of Scoresby Sund, eastern Greenland, were the subjects of intensive research in the early 1990s as part of the Polar North Atlantic Margins; Late Cenozoic Evolution (PONAM) project, which resulted in several papers published in the mid-1990s. A common indicator of LIG age is the presence of warmer-water than today fauna, e.g. *Balanus crenatus*. Present are also members of the Astarte genus, some arctic and some subarctic to their biogeography (Funder et al., 2002). The presence of *Astarte borealis*, generally regarded as an Arctic species, is, however, thought only possible in this region under conditions of increased advection of Atlantic water, which occurred only during the LIG and the Holocene (Vosgrea et al., 1994). Some sites are also constrained to the LIG based on >20 luminescence ages (Mejdahl and Funder, 1994). Below, we present 8 locations where these LIG sediments were documented in the vicinity of Scoresby Sund. The coordinates for these sites are estimated from maps in the original publications.

7.43.16.41.1 Kikiakajik section (70.038, -22.2467)

At Kikiakajik (Fig. 13C) there is a 700-m long and 10–15-m high coastal cliff, but it is partly covered by perennial snowbanks. Mangerud and Funder (1994) cleaned and described two exposures. The first exposure has a coarsening-upwards sequence from horizontally bedded marine silts at 3.5 MASL through sand to gravel foresets, which are cut by a till at 7.5 MASL. The second exposure consists only of the gravel foresets at 9–13 MASL, covered by till. The silt and sand

contain a molluscs fauna similar to the one described at Kap Hope, including *Balanus crenatus* and the warmer than today-
940 indicating *Astarte borealis*. A single shell was radiocarbon dated to >44 ka, and the sand was dated with TL
to 132 ± 10 ka. The distinct fauna strongly suggests a correlation between Kap Hope and Kikiakajik,
and the warm water requirements of the fauna together with the dates from both sites indicate a LIG age.

7.43.26.41.2 **Kap Hope (70.459, -22.318)**

At Kap Hope there is a 20-m high coastal section where Mangerud and Funder (1994) described a marine silt between 2 and
945 3 MASL, directly overlying bedrock (Fig. 13C). The silt contains a rich *Astarte* fauna including the warmer-water species
Balanus crenatus and *Astarte borealis*, which are not living in East Greenland today. The fauna suggests an offshore
environment warmer than at present. From the silt there is a coarsening up-wards sequence through sand (well-sorted, ripple
laminations with gravel lenses) to gravel (includes well-rounded boulders), representing a regression from off-shore through
shoreface to beach environment at 10 m MASL. The beach sediments are covered by a 5-m thick sandy gravel interpreted
950 to suggest a rising relative sea-levelRSL, eventually capped by a till. Two radiocarbon-¹⁴C-dates from individual shells gave
non-finite ages (>42 ka) and two TL/OSL dates from the shoreface sand yielded 97 ± 10 ka and 75 ± 8 ka, respectively.
Mangerud and Funder (1994) assigned the entire clay-sand-gravel sequence to the LIG based on the fauna and supported by
the TL/OSL dates, although the latter yielded too young ages. In the database, we have added separate entries for both the
regressive and transgressive units.

955 7.43.36.41.3 **Kap Stewart composite (70.44, -22.78)**

Marine sediments assigned to the LIG are present from 0.5-40 MASL around Kap Stewart (Tveranger et al., 1994, Fig.
13C). In their study, the authors examined several sites to better understand the sedimentology and to find as many
fossils as possible. However, ultimately, all sedimentary structures, dates and fossils were assumed to represent one single
site, which we follow here. To provide additional context on the LIG marine event, we provide a brief overview of each
960 individual site below.

At Loc. 471 marine sediments interpreted as deposited in a tidal, shallow marine delta environment were found between 9
and 12 MASL and at Loc. 473 similar sediments were found at 0.5-to-8 MASL (Tveranger et al., 1994). These sediments
contain marine molluscs suggesting warmer-than-present water. Based on these molluscs and the stratigraphic position
965 below diamicton and more recent sediments, these deposits were assigned a LIG (MIS-5e)-age. From Loc. 473 the
interglacial unit could be mapped on the surface some 500 m inland (towards the northN) to Loc. 468 where the unit
consists of tidally influenced channels and estuarine/lagunal deposits up to about 40 MASL. The unit was mapped further
through Loc. 470 where it consists of trough cross-bedded sand and gravel located between 30 and 38 MASL and
interpreted as fluvial deposits with paleocurrents towards the fjord. The interpretation is that the transition from marine to a
970 coarsening upwards fluvial sequence reflects a progradation of the paleo-Ostraev river delta. The vertical stacking of tidal

channels and the thickness of fluvial deposits indicate that progradation took place during a RSL rise up to about 40 MASL.

7.43.46.41.4 Hesteelv composite (70.44, -23.10)

Similar to the Kap Stewart site, in the Hesteelv area (Fig. 13C), workers examined several sites to better understand the stratigraphic record (Tveranger et al., 1994). Ultimately, all sediment successions, dates and fossils from studied sections were assumed to represent the same marine unit, situated between 5–35 MASL, and tied to the LIG ~~Ultimately, all sedimentary structures, dates and fossils were assumed to represent one single site, with the LIG marine unit situated between 5 and 35 MASL. For additional context on the LIG marine sediment, we provide below a brief overview of some individual sites. Below, to provide additional context on the LIG marine event, we provide a brief overview of some individual sites.~~

At Hesteelv 13 outcrops, located in an area stretching 5 km along the fjord and 1 km inland, were cleaned and sedimentological logged in detail (Tveranger et al., 1994). Sediment units could (at several places) be traced on the surface between the outcrops and the correlations between the outcrops mentioned here are considered reliable. At Loc. 153 a continuous outcrop was cleaned from 5–71 MASL and this is a key exposure for the full stratigraphy of the area but here we will only describe the LIG sequence (Unit A in Tveranger et al., 1994), which is found 5–35 m MASL. The lower 15 m consist of 2–50-cm thick sandy turbidites interbedded with massive or weakly laminated mud layers containing molluscs, dropstones and synsedimentary slumps. The turbidites are overlain by low-angle, cross-stratified sand, with layers of gravel, and sand with climbing ripples and planar cross-bedding. Depositional direction was towards the fjord (i.e. south) throughout the unit. The unit is interpreted as a prograding delta sequence with transition from offshore turbidites to shoreface deposits with migrating bars. The altitude and facies indicate a sea-level between 35 and 40 MASL. The interglacial sediments are overlain by a till, interstadial marine sediments and another till and deglacial sediments, not described here. The LIG unit can be followed from Loc. 153 to Loc. 152 (18–30 MASL) and Loc. 154 (9–33 MASL) by surface mapping. Both localities have a coarsening upward sequence like that at Loc. 153 and thus support the interpretation there. Last interglacial sediments were described also from some of the other outcrops and fossils collected. At some places erosional, regressive facies are mapped and interpreted to stem from the subsequent falling RSL. Several terrestrial plant macrofossils (e.g. *Betula pubescens*), beetles and marine molluscs (e.g. *Mytilus edulis*) collected from the LIG unit in different outcrops show warmer-than-at-present climate, and mean July temperature was estimated to have been about 5 °C (Tveranger et al., 1994). Thus, an interglacial origin is clear. The average of 9 OSL ages is 115 ± 29 ka and suggests a LIG age, which is consistent with the full stratigraphy at the site and indeed by correlation with other sites along Scoresby Sund (Vosgerau et al., 1994).

7.43.56.41.5 Site 443d, Fynselv area

At Site 443d (Fig. 13C), marine sediments were identified between 12 and 21.5 MASL, dated by TL/OSL to 120 ± 10 ka, 121 ± 10 ka, and 122 ± 10 ka (Hansen et al., 1994). The marine sediment succession s-were was interpreted as deposited in a deltaic to shallow marine environments~~suecession~~ (units Ib, Ic, Id). These marine sediments -and- are overlain by a till ~~followed by~~ and marine sediments, representing a later incursion.

7.43.66.41.6 Langelandselv composite (70.546, -23.644)

Like the Kap Stewart and Hesteelv sites, workers in the ~~6.40.7~~ Langelandselv region (Fig. 13C) examined several sites to better understand the stratigraphic record (Landvik et al., 1994). The LIG marine unit (amalgamated from several sites) spans 0 ~~to~~ 70 MASL. For additional context on the LIG marine sediment, we provide below a brief overview of some ~~individual sites. Below, to provide additional context on the LIG marine event, we provide a brief overview of some individual sites.~~

At sites 77A and 77D, a layer of marine silts and sands ~~are-is~~ situated between 0 and 5 MASL (Landvik et al., 1994). OSL/TL dating of the uppermost silt sediments yielded 122 ka and 121 ka (Mejdahl and Funder, 1994). Nearby, at sites 74 and 76A, marine silts and sands were documented between 4 and 11 MASL. These marine sediments are overlain by ~10m of sand, interpreted to be of fluvial origin. These were OSL/TL dated to 100 ka, 106 ka, 117 ka and 129 ka, the whole sediment succession suggesting an environmental change from marine conditions into a fluvial setting (Landvik et al., 1994).

At four sites in the Langelandselv region (sites 95, 96, 97 and 113), LIG marine sediments are situated at an anomalously high elevation. Landvik et al. (1994) suggested these marine sediments were deposited early during the LIG during isostatic recovery from the MIS 6 glaciation and were believed to represent an earlier interval than the other sites from the Scoresby Sund area, typically situated much closer to present-day sea level. At sites 95, 96 and 97, Landvik et al. (1994) described a sand unit containing ripple and herring-bone cross-beds, situated between 44 and 57 MASL. These sands were interpreted as a shallow marine environment and were OSL dated to 117 ka (Landvik et al., 1994). The mollusc assemblage in these sands suggests deposition during a warm-water interval, which is also indicative of the LIG (Vosgreaux et al., 1994). Finally, at site 113, Landvik et al. (1994) described marine sand (Unit A₀) situated between 67 and 70 MASL.

7.43.76.41.7 Location 72, Aucellaev River (70.585, -23.764)

On the north shore of Scoresby Sund, along the Aucellaev River (Fig. 13C), are marine sediments at the base of the stratigraphic record which consist of mollusc-bearing (largely *Astarte borealis*) sand and silt, situated between 12 and 16 MASL, and interpreted as deposited in a sublittoral shallow marine environment (Unit 2; see Israelson et al., 1994). The sediments were dated via OSL and TL ~~dated to~~ 122 ka and 144 ± 15 ka (sample R911004). A broad LIG age is also

suggested by the presence of warm-water mollusc fauna. The marine sediment unit is both underlain and overlain by till and on top is sediment from a later marine event.

7.43.86.41.8 Lollandselv-Falsterselv region, Greenland (70.873, -24.179)

1035 Ingólfsson et al. (1994) noted LIG marine sediments at 23 stratigraphic sites between the Lollandselv and Falsterselv rivers (Fig. 13C). A composite diagram of these sites shows shell-bearing marine silts situated between 0 and 8 MASL in this region (Ingólfsson et al., 1994). The maximum elevation of marine sediments in this location is 20 MASL. The authors interpreted these marine silts as deposited during the LIG, representing a transition from glaciomarine to tidal and eventually a littoral depositional setting. Macrofossils (plants, molluscs) suggest temperatures warmer than the Holocene. These LIG
1040 marine sediments are overlain by a series of cross-bedded sands (fluvial), till (ice advance), topped by sediments suggesting a later marine incursion.

7.44.6.42 Thule, Western Greenland (7 sites)

1045 Kelly et al., (1999) described three separate marine events in sediment sequences from western Greenland (Fig. 13D). The middle marine sediment succession (known as the Qarmat Eevent) contains mollusc shells from *Mya truncata*, *Hiatella arctica* and *Chlamys islandica*. The age of these marine sediments is assigned to the LIG, based on TL ages between 91~~–~~to 154 ka (mean age at 127 ka). The Qarmat sediments coarsen upwards, and the uppermost sediments have wave influenced sedimentation. We outline 8~~–~~7 sites below (locations mapped in Fig. 13D), as described by Kelly et al. (1999). Some of the chronological data ~~were are from described in~~ Sejrup (1990).

7.44.16.42.1 Iterlak K (76.710, -69.410)

1050 ~~At this site, s~~Sands and muds are situated from 25~~–~~to 27.5 MASL. Marine shells are present, and two amino acid dates suggest deposition during the LIG, which is supported by a non-finite radiocarbon age~~±~~.

7.44.26.42.2 Iterlak L

~~At this site, s~~Sands and muds are situated from 11~~–~~to 14 MASL. No marine shells or foraminifera were noted, but a TL age suggests deposition at 118 ka.

1055 7.44.36.42.3 Saunders Ø B (76.597, -69.736)

~~At this site, t~~The Qarmat marine event is situated between 3~~–~~to 20 MASL. Glaciomarine muds are present from 3~~–~~to 6 MASL. Overlying the glaciomarine sediments is a ~1-m interval of sands, followed by clast-supported gravels with varying amounts of mud and sand, which was interpreted as a beach deposit. Marine shells are present. Three TL ages from the lower-most sediments yielded 154 ka, 153 ka and 119 ka.

1060 **7.44.46.42.4** Saunders Ø C

~~At this site, t~~The marine interval spans 5~~–~~to 12 MASL. The lower part of the section consists largely of sands which gradually coarsens to sandy gravels at ~8 MASL, which was interpreted as a beach deposit. Marine shells are present and the uppermost sediments, at ~12 MASL, were dated with a TL to 91 ka.

7.44.56.42.5 Narsaarsuk D (76.453, -69.287)

1065 Sand with occasional coarse gravel beds are situated from 0~~–~~to 6 MASL. The sediments have signs of being wave influenced. Marine shells and foraminifera are present, and amino acid dating of shells suggest deposition during the LIG.

7.44.66.42.6 Narsaarsuk E (76.453, -69.287)

1070 Clay and silt, transitioning to sands are situated from 11~~–~~to 17 MASL. Marine shells and foraminifera are present. A TL age from the upper sediments suggests deposition at 119 ka, which is supported by amino acid dating suggesting a LIG timeframe, as do two non-finite radiocarbon ages.

7.44.76.42.7 Narsaarsuk F+G (76.453, -69.287)

At this site, sand with occasional coarse gravel beds is situated from 3~~–~~to 9 MASL. Marine shells, foraminifera and plant matter are present. Amino acid dating suggests deposition of these sediments broadly within the LIG, which is supported by a non-finite radiocarbon age as well as a TL age of 123 ka.

1075 **7.45.6.43** Iles de la Madeleine, Quebec, Canada (3 sites)

In eastern Canada, the Iles de la Madeleine contains sediment exposures bearing an extensive stratigraphic record that details the interplay between sea level and Pleistocene ice sheets. Marine sediments assigned to ~~MIS 5~~the LIG have been described at three sites, shortly described below.

7.45.16.43.1 Camping site

1080 Rémillard et al. (2017) described marine sediments (Fig. 14B) along the cliffs of the Iles de la Madeleine that overlie local bedrock. These sediments consist of fine to medium red sand with gravel beds, moderately to poorly-sorted, and 10~~–~~to 30 cm in thickness. This unit was located at 14 MASL with a maximum thickness of 4 m. The sediments were interpreted as a barrier beach sediment displaying alternating low and high energy fluctuations and were assigned to the early part of MIS 5 (likely 5e), based on an IRSL age of 115 \pm 8 ka (OSL51; Rémillard et al., 2017). The

1085 **7.45.26.43.2 Portage du Cap**

The Portage du Cap site (Fig. 14B) was first described by Prest et al. (1976) and subsequently by Dredge et al. (1992). At this site, a sequence of sub-till silts, pebbles and gravel are exposed within a gravel pit. The entire marine sequence spans 14~~–~~to 17 MASL (Dredge et al., 1992). Underlying the entire marine sequence is sandstone bedrock containing borings from *Zirphella* sp., a species that lives in the modern intertidal zone in the region. The lowest unit, situated at 14 MASL, is a 1090 grey marine silt, rich in dinoflagellate cysts, containing a pollen, beetle and diatom assemblage suggesting conditions warmer than present-day in the region (Dredge et al., 1992; Prest et al., 1976). Within this unit there are clast-supported beach gravels with well-rounded pebbles. Overlying the gravel unit is a 1~~–~~to 2 m thick organic sand unit with a woody organic horizon. For this reason, the lowermost section of this stratigraphic sequence has been assigned to a MIS-5eLIG marine incursion that immediately followed the removal of MIS 6 ice from this area. On top of this unit is a gravel unit with 1095 a pollen assemblage similar to present-day, likely the result of relative climatic cooling following peak MIS-5eLIG conditions, paired with local sea-level rise and beach formation (Dredge et al., 1992). Three radiocarbon ages are available at this site. Plant detritus and wood have been dated to >38 ka and >35 ka~~BP~~ (GSC-2313 and BGS-259; Prest et al., 1976) and a marine shell was dated to 42.9 ± 0.72 ka ~~BP~~ (GSC-4633 HP; McNeely and McCuaig, 1991). The finite age on the marine shell is possibly beyond the reliability limit for dating marine shells (Pigati, 2002), and all radiocarbon ages 1100 considered as minimum such.

7.45.36.43.3 Le Bassin site (47.2333, -61.8983)

Dredge et al. (1992) described a stratigraphic section of sub-till peat overlain by clays, sands and gravels at the Bassin site, Iles de la Madeleine, Canada (Fig. 14B). At the base of the section overlying bedrock is a 0.5-m sand interval, followed by a 0.2-m peat layer. This was interpreted as formed in nearshore fluvial and lacustrine environments. Overlying 1105 the peat layer is a 0.9-m silt and clay unit, followed by 0.1 m of sand, both interpreted as marine in origin based on the presence of warm-water oyster shells (*Ostrea virginica*). The altitude of these marine sediments is unclear; we assume the base of the stratigraphic section is 0 MASL ('just below tide level' according to Dredge et al., 1992), which would situate the marine silt, clay and sands between 0.7 and 1.7 MASL. Pollen data from these sediments suggest conditions similar to present-day (Mott and Grant, 1985). The sub-till sediments suggest that, following deglaciation of this region, the local area 1110 became a peatland and, as sea levels rose, it transitioned to a lagoon environment, with beach ridge and coastal barrier formation (Dredge et al., 1992). Three U/Th ages on these sediments range from 89~~–~~106 ka (UQT-182, UQT-183, UQT-184) and were interpreted as minimum age constraints by the original authors. Two diamictons (interpreted as tills deposited during ice advances) truncate the sand unit. Based on pollen and stratigraphic position, Dredge et al. (1992) suggested

a LIG age assignment

7.46.44 Clyde Foreland, Baffin Island, Canada (3 sites)

First described by Løken (1966) and Feyling-Hanssen (1967), sub-till mollusc-bearing marine sands are present along cliffs of the Clyde Foreland, Baffin Island (Fig. 14C). These are known as the Cape Christian marine sediments. Several radiocarbon dating attempts on organic material from this unit yielded infinite ages or finite ages (e.g. QL-188) that can be considered as minimum ages. Feyling-Hanssen (1967) suggested that these sediments were deposited during MIS 5 or the LIG owing to its content of relatively warm-water-indicating foraminifera (*Cassidulina teretis*). Warm conditions were confirmed via pollen analyses. Later, these sites were re-visited by Miller et al. (1977) who reported and a ^{230}Th age of 130 ka (Miller et al., 1977). Below, we document 3 of these sites, as described by Miller et al. (1977). These sites are correlated to each other on the basis of based on stratigraphic position, and amino acid ratios.

1125 7.46.44.1 Profile 6 (70.694, -68.946)

At this site, the Cape Christian marine sediments are situated between 27 and 28 MASL. They are underlain by a series of sands and tills and overlain by a thin layer of buried soil along with till from later Pleistocene ice advance.

7.46.44.2 Profile 9 (70.596, -68.413)

At this site, Cape Christian marine sediments (coarse sand and cobbles) are found between 9 and 11 MASL. These marine sediments are overlain by two tills and younger marine sediments. This is the type section for the Cape Christian marine sediments.

7.46.44.3 Profile 10 (70.575, -68.347)

Miller et al. (1977) documented coarse sands and gravels between 1.8 MASL and 2.5 MASL, which they correlate to the LIG Cape Christian marine sediments.

1135 7.47.6.45 Ile aux Coudres, Quebec, Canada

Occhietti et al. (1995) described a core taken on Ile aux Coudres, located in the Saint Lawrence Estuary, Canada, that showed a succession of Pleistocene-aged deposits (Fig. 14B). A till was noted at the base of this record (known as the Baie-Saint-Paul glacial complex), followed by a series of clays, rhythmites and deltaic sediments, known as the “Guettard Sea” sediments, present from 125 to 2 MBSL. Occhietti et al. (1995) described, at the base of this stratigraphic unit, a very compact, finely stratified grey clay between 125 and 102 MBSL. A dinoflagellate cyst at 119 MBSL suggests a marine origin for these sediments, and the authors interpreted it as a transgressive prodeltaic (deep marine) deposit. Next, between 102 and 71 MBSL, are grey rhythmites, hypothesized to be a high-level prodeltaic deposit resulting from the gradual shallowing of marine waters. Lastly, the sediment core shows silts and sandy silts between 71 and 2 MBSL, interpreted as prodeltaic, transitioning to deltaic sediments. This uppermost unit contains benthic foraminifera common to brackish marine

1145 water. Based on stratigraphic context (i.e. heavily isostatic depression following large-scale glaciation), Occhietti et al. (1995) aligned this record with the transition period between MIS 6 and [the LIG](#). They [inferred](#) a large-scale, long-lasting glaciation followed by ice recession, rapid marine inundation into the isostatically depressed landscape, and shallowing of marine waters owing to subsequent rebound. This accumulation of marine sediments may represent [3.5 kyrs](#) of deposition (Occhietti et al., 1995). Since the age of the deposit is based on regional correlations and 1150 environmental conditions from pollen assemblage composition, rather than direct dating, we assign the lowest age quality score.

[7.486.46 Bridgehampton, New York, United States \(40.9374, -72.3064\)](#)

1155 Marine sediments, typically sandy clay that is brown or grey/green in colour, were first encountered at depth on Long Island, New York in the early 1900s (Fuller, 1914; [Fig. 14B](#)). The so-called “Gardiners Clay” is present at depth in several borehole records between 20 and 45 MBSL in various well records in this region (Nemickas and Koszalka, 1982). Paleontological work revealed a variety of warm-water fauna, including foraminifera, coelenterata, bryozoa and mollusca (Gustavson, 1972), confirming a marine origin for these sediments. At the Bridgehampton site, the Gardiners Clay was encountered between 23 and 46 MBSL (Nemickas and Koszalka, 1982).

1160 The Gardiners Clay is provisionally assigned to [the LIG](#), based on stratigraphic context and amino acid [dating](#). Stratigraphically, this marine unit is overlain by a series of Pleistocene deposits, notably the Montauk till, associated with the most recent (MIS 2) [glaciation](#) (Nemickas and Koszalka, 1982). Moreover, at some sites, the Gardiners Clay is underlain by a gravel deposit (Jameco Gravel; Fuller, 1914) that has been associated with fluvial or glaciofluvial deposition immediately following retreat of the MIS 6 ice sheet and prior to the 1165 marine incursion. The interpretation of the age is complicated by the fact that no amino acid racemization had been conducted on shells from the Bridgehampton site. However, shells contained in the Gardiners Clay from a nearby outcrop (40002; the stratigraphy of which was not described in detail) yielded an age assignment of early MIS 5 (Wehmiller et al., 1988; Wehmiller and Pellerito, 2015). The significant difference in elevation and depositional context (at-depth in a drill core vs. sub-aerial outcrop) implies the Gardiners Clay might represent two marine deposits of different ages.

1170 **[7.496.47 Kwatabohegan River, Ontario, Canada \(51.1389, -82.1167\)](#)**

The Hudson Bay Lowlands contain a rich stratigraphic record consisting of tills interspersed by non-glacial sediments. Along the Kwatabohegan River ([Fig. 14B](#)), *in situ* marine sediments were first discovered underlying till by Bell (1904), and later described in detail by Skinner (1973). These so-called “Bell Sea” sediments were first reported to be at 75 MASL (Skinner, 1973), however subsequent geochronological work on samples originally obtained by Skinner (1973) report the elevation as 1175 90 MASL (McNeely, 2002). The reason for this discrepancy is unknown. These marine sediments consist of 0.8 m compact

bluish grey sand and silt that are deformed owing to subsequent Wisconsinan ice advance over the region. An undulating bed of marine mollusc shells is present near the mid-point of the sediment package (Skinner, 1973).

The Bell Sea sediments are assigned to MIS 5e the LIG based on stratigraphic context, amino acid dating and minimum 1180 limiting radiocarbon ages. Stratigraphically, this marine deposit is overlain by tills that are assigned to the Wisconsin Glaciation MIS 4 and MIS 2 glaciations, thus it preceeded the build-up of ice during the Wisconsinan. Amino acid techniques dating also supports a MIS 5e LIG assignment for the Bell Sea sediments; Andrews et al. (1983) show that isoleucine epimerization ratios from shells in the Bell Sea sediments (average of ~0.2) are significantly older than ratios from assumed mid-Wisconsinan marine deposits (~MIS 5-3) in this region (average of ~0.14), which are even older than shells from the 1185 post-LGM marine incursion (average of ~0.03). In calculating the amino acid dates, the average regional diagenetic temperature (0.6°C) can only be reconciled with the isoleucine epimerization ratios if the smallest isoleucine epimerization ratio (Bell Sea sediments), is assigned to MIS 5e the LIG (Andrews et al., 1983). For that reason, the Bell Sea sediments were assigned to ~130 ka (Andrews et al., 1983). Finally, two radiocarbon ages on marine shells (*Hiatella arctica*) from the Bell Sea deposit yielded >37 ka (GSC-1475; Blake, 1988) and 47.85 ± 1.09 ka (TO-2503; McNeely, 2002). The finite age is 1190 unreliable because shells samples are highly susceptible to modern-day contamination, and the date is very close to the limit of radiocarbon dating (McNeely, 2002). Thus, both are considered as minimum limiting radiocarbon ages.

7.506.48 East of Nicholson Peninsula, Northwest Territories, Canada (69.894, -128.521)

Along the westernmost coastline of the Northwest Territories, Canada, sediments representing the Liverpool Bay 1195 Interglaciation are exposed as part of the Ikpisuguk Formation (Fig. 14D). As described by Rampton (1988), a layer of organic-bearing marine sand and silt is present between 0 and 2 MASL (Locality VH-83-050), which was interpreted as an intertidal beach complex. Fossils from a nearby exposure of the Ikpisuguk Formation (VH-83-040; does not contain a marine unit) suggest a climate warmer-than-present, which supports LIG deposition (Rampton, 1988). Radiocarbon age attempts on driftwood contained in this unit were non-finite (GSC-3722) and amino acid ratios 0.1– 1200 0.15, both supporting a LIG age assignment based on the amino acid framework for nearby Banks Island (Vincent, 1982, 1983).

87 OTHER LIG MARINE SITES

During our search for MIS 5e/LIG marine sites to include in this database, we located several LIG sites containing 1205 marine sediments that are not *in situ*, mostly due to post-depositional, i.e. later, glacial deformation and/or relocation. These sites are thus unsuitable as indicators of RSL but merit inclusion in this manuscript paper. We describe several ten of these such sites below, ordered east to west.

8.17.1 Lower Ob sites, West Siberian Plain, Russia

As shown in Fig. 34, the stratigraphic record present at the Lower Ob sites, West Siberian Plain, contains a well-dated MIS 5eLIG section consisting of alluvial sedimentum with OSL ages of 125_–to–138 ka and thick peat with U/Th ages of 133 and 141 ka. These terrestrial sediments occur at the same altitudes or slightly higher than the corresponding marine sediments of the Arctic. However, marine sediments at the base of this stratigraphic sequence (separated from the overlying terrestrial beds by a till) yielded a youngest age of 153 ± 15 (sand with rich fauna of boreal foraminifers of the Kazantsevo-LIG assemblage in Hashgort borehole at lat/long 65.42, 65.67; Arkhipov et al., 1992). This marine unit possibly represents an earlier marine incursion and is therefore excluded from our database.

8.27.2 More-Yu, Pechora Lowland, Russia

At the More-Yu site, Pechora Lowland (lat/long 67.867, 60.183), OSL ages of 112 ka and 120 ka were obtained from large sand blocks with shells of boreal molluscs included into the glacially deformed diamict sequence (Fig. 3). The uranium/thorium/U/Th date of 130 ± 8 ka from this section on boreal *Arctica islandica* shell confirms an Eemian LIG age of the detached marine sand and a Weichselian age of the encompassing till with fossil glacial ice (Astakhov and Svendsen, 2002). We exclude this site from our database owing to the likely glacial translocation of the marine unit. Another site in the Pechora Lowland is Vastiansky Kon, which was first described in 1938 and revisited by Tveranger et al. (1998). The stratigraphic record, along with the presence of warmer-water marine shells, suggest a LIG age for marine sediments at this site. However, it has been glaciotectonized and is therefore inappropriate for inclusion in our database.

8.37.3 Malaya Kachkovka, Kola Peninsula, Russia

The Malaya Kachkovka site is in the tributary valley of the Malaya Kachkovka River on the eastern coast of the Kola Peninsula (lat/long 67.40, 40.90; 140 MASL). The entire sediment exposure is ~10 m thick (Gudina and Yevzerov, 1973). The mollusc and foraminifera fauna, as well as the pollen content of a marine sediment interval between 127.5–134 MASL were studied by Lavrova (1960) and Gudina and Yevzerov (1973). The marine part of the section comprises a 0.5-m thick clay unit at the base of the exposure, overlain by mollusc-bearing sands. Lavrova (1960) considered that the mollusc fauna contained in the sediments represents the upper sublittoral zone, and according to Gudina and Yevzerov (1973) the faunal assemblage represents the warmest fauna of interglacial marine sediment faunas discovered on the Kola Peninsula. Uranium/thorium ages on *Astarte borealis* shells from coarse and medium marine sands at around 132 MASL yielded ages of 102 ± 4 and 114 ± 4 ka (Arslanov et al., 1981), suggesting that the sediments were deposited during the LIG. However, Gudina and Yevzerov (1973) pointed out that the characteristics of the faunal assemblages and the high altitude of the marine sediments might suggest that they do not represent the LIG but rather an older interglacial. Even if this is not the case (i.e. from an earlier interglacial), the till on top of the marine sediments might indicate that the marine sequence is not in an

in situ position and has been glacially thrust as a block to its present high-altitude position. The site is thus excluded from our LIG database.

The Malaya Kachkovka site is in the tributary valley of the Malaya Kachkovka River on the eastern coast of the Kola Peninsula, Russia (67.40, 40.90; 140 MASL). The entire sediment exposure is ~10 m thick (Gudina and Yevzerov, 1973).

1240 Mollusc fauna, foraminifera and pollen content of the marine sediment interval (127.5 to 134 MASL) were studied by Lavrova (1960) and Gudina and Yevzerov (1973). The marine part of the section comprises 0.5 m thick clay unit at the base of the exposure overlain by mollusc-bearing sands. Lavrova (1960) considered that the mollusc fauna in the marine sequence represents the upper sublittoral zone, and according to Gudina and Yevzerov (1973) the faunal assemblage represents the warmest fauna of the interglacial marine sediment faunas discovered on the Kola Peninsula. $^{230}\text{Th}/^{234}\text{U}$ ages on *Astarte borealis* shells from coarse and medium marine sands at around 132 MASL yielded ages of 102±4 ka and 114±4 ka (Arslanov et al., 1981) suggesting that the sediments formed during the Eemian interglacial (MIS 5e). However, Gudina and Yevzerov (1973) pointed out that the characteristics of faunal assemblages and the high altitude of the marine interglacial sediments might suggest that the sediments at the Malaya Kachkovka site do not represent the Eemian interglacial sediments but rather an older interglacial. Even if that was not the case, the marine sequence is overlain by glacial till and therefore, it is probable that the marine sequence is not *in situ* and has been glacially thrust as a block to its present high altitude position. Grave et al. (1969) described a riverbank section of the Ludyanoi creek (lat/long 66.33, 39.92; 70 MASL), a small tributary to Pulonga River, southeastern costal area of the Kola Peninsula. They described a sediment exposure where the top 10.7 m of the section is glaciofluvial sands and gravels overlying a 1.5-m thick till. Below the till is a 3.7-m thick greenish grey marine clay with shells and shell fragments recognised as *Astarte borealis*, *A. elliptica*, *A. crenata* and *A. montaqui*, of which the latter two are considered subarctic to their biogeography. The top of the marine clay unit is at around 60 M~~ASL~~SL, but it is not precisely known if this marine unit is *in situ*. It is therefore excluded from our database.

Formatted: Font: Italic

8.67.5 Lovozero, Kola Peninsula, Russia

There is a thick sediment sequence at the Lovozero site located in the inner part of the Kola Peninsula, Russia (lat/long 67.08, 38.970; 210 MASL). The lower part of the drill hole sections contains a 44-m thick sand and silt unit between 140 and 1260 184 MASL. A proportion of the diatom flora in the lower part of the sediment core is marine and the pollen taxa is dominated with pine and birch (Ikonen and Ekman 2001). However, it is generally thought that diatoms and pollen are mostly reworked in this lower sand-rich sediment unit and therefore, the correlation of this unit to *Eemian interglacial* the LIG (MIS 5e) is not possible and the position of the sea level cannot be reconstructed (e. g. Ikonen and Ekman, 2001). We do not include the Lovozero site in our database.

1265 8.77.6 Evijärvi, Finland

The Evijärvi site is situated in central Ostrobothnia, western Finland (lat/long 63.434, 23.322; 67 MASL). At this site, the LIG sediments occur on the proximal flank of a drumlin (Eriksson et al., 1980). In the top 6.4 m of the borehole record,

several till and sand beds occur. The interglacial sediments consist of silt (9.0–9.5 m depth) and gyttja (6.4–7.4 m depth) interbedded with till. Samples for pollen and diatoms were analysed from borehole sediments (Grönlund, 1991; Eriksson, 1993). The pollen spectra of the silt layer (9.0–9.5 m depth) and gyttja layer (6.4–7.4 m depth) indicate only one local pollen assemblage zone consisting of *Betula*, *Alnus*, *Picea* and *Corylus*, which is correlated to the LIG (Erikson, 1993). Diatoms in the silt bed (9.0–9.5 depth) above the till at the base of the core are exclusively marine while the diatom taxa in the gyttja is dominated with marine lagoonal types (Grönlund, 1991a). We do not include the Evijärvi site in our database because these LIG marine sediments are not at their original position of deposition but have been transported for an unknown distance and 1275 altitude by ice (e.g. Eriksson, 1993).

8.87.7 Norinkylä, Finland

The Norinkylä site is situated in southern Ostrobothnia, western Finland (lat/long 62.58, 22.020, 114 MASL). This site exposes till on top of the flanks of an esker, and in between are glacially deformed gyttja, organic-rich silt and sand, suggested to be in the LIG in western Finland. Boreholes made in the Rahkaneva mire next to the till-covered esker display silt, clay and gyttja in between two till beds with a maximum thickness of 4.5 m. The inter-till sediments have been studied for their lithology and pollen and diatom content (Grönlund, 1991b; Erikson, 1993). The pollen assemblage shows a succession typical for the LIG in western Finland (Eriksson, 1993). Fresh-water diatom taxa dominate the lowermost part of the inter-till sediments (biostratigraphically belonging to the early LIG 1280 Betula regional zone; Grönlund, 1991b) while there is a transition to marine diatom taxa that takes place at around 97 MASL. Saline diatom taxa dominate the interglacial sediment between 97 and 102 MASL (Grönlund, 1991 a, b). We do not include the Norinkylä LIG sediments in our database because these sediments are glaciectonised and have been transported an unknown distance from their original place of deposition.

Formatted: Font: Italic

8.97.8 Svartenhuk Halvø, west-central Greenland

1290 Kelly (1986) and Bennike et al. (1994) describe a raised marine deposit in west-central Greenland, which was correlated to the LIG based on AAamino acid dating. They found no evidence of a Holocene highstand in this location. However, more recent work at this site suggests it may not have been deposited during the LIG (Lane et al., 2015). We were unable to add this site to

8.107.9 Nantucket Island, Massachusetts, United States

1295 Marine sediments, bracketed by two tills, on Nantucket Island, Massachusetts, United States, were first described in the 1800s (Desor and Cabot, 1849; Verrill, 1875). These marine sediments are located between 0–20 MASL (Oldale et al., 1982). These sediments consist of a gravel base, overlain by several meters of cross-bedded so-called “Sankaty Sands” that contain stratified silt and clay along with abundant marine shells. Oldale et al. (1982) assigned the Sankaty Sands to MIS

1300 the LIG based on stratigraphic context, palaeoenvironmental data, absolute chronology (U/Th, minimum radiocarbon ages) and relative age determinations (amino acid racemization). The lowermost till was assigned to MIS 6 and the upper till to MIS 2 ice advances, and the Sankaty Sands were stratigraphically constrained to the time interval immediately following retreat of the MIS 6 ice sheet. Palaeoenvironmentally, well-preserved warm-water oyster shells (*Crassostrea virginica*) and clam shells (*Mercenaria mercenaria*) are present at the base of this marine unit, and cooler-water clams (*Mercenaria campechiensis*) and mussels (*Mytilus edulis*) are present near the top (Verrill, 1875). These marine shells capture the 1305 transition from warm-water LIG conditions to more temperate MIS 5d conditions (Oldale et al., 1982). In terms of absolute chronology, corals located within the Sankaty Sands were dated via U/Th methods to 133 ± 7 ka, which the authors believed to be a maximum age for this deposit (SHIO/80-9; Oldale et al., 1982). Moreover, seven radiocarbon ages were obtained from wood and shells located at this site. Four yielded non-finite ages and three were finite. 1310 However, the latter were suspected to be affected by modern carbon contamination (Oldale et al., 1982). Finally, amino acid racemization analyses (isoleucine epimerization ratios) of 10 shells from this marine unit suggest deposition between 140 ka and 120 ka (Oldale et al., 1982). In summary, the palaeoenvironmental data, stratigraphic context and available age assignments all suggest a LIG age for the Sankaty Sands. Despite the clear evidence of LIG age assignment, the section is glaciectonized, and is therefore unreliable as an estimate of palaeo-sea level.

8.117.10 Western Banks Island, Canada

1315 Lakeman and England (2014) describe a glaciectonized marine deposit on Phillips Island, located off the west coast of Banks Island, which dated to the LIG based on OSL dates and non-finite radiocarbon dates. The exposure has interbedded sand, silt and clay with abundant mollusc fossils. The OSL samples were taken at 7 MASL. Several other raised in-situ marine

98 DISCUSSION

1320 This manuscript brings together 82 MIS 5e/LIG sea-level proxies from the formerly glaciated Northern Hemisphere. In general, the sites reported in this paper do not offer constraint on the global MIS 5e/LIG high-stand, but rather any evidence of GIAe-influenced sea-level positions after the end of the MIS 6 glaciation. The motivation behind some of the research described herein was to constrain the extent of the Barents Sea, and the (possible) connection between the White Sea and the Baltic Sea caused by the glacio-isostatic depression of western Russia during the MIS 6 glaciation and into the 1325 LIG (Ikonen and Ekman, 1991; Funder et al., 2002; Miettinen et al., 2014). Last Interglacial sites in Svalbard, Iceland and Greenland are largely located along the modern-day coastline. Comparatively fewer LIG sites are present in North America, likely owing to more glacial erosion or their sub-surface position in only borehole records.

~~The primary challenge of assessing past sea level in formerly glaciated areas is that the sea level position likely did not remain at a fixed position for any length of time. The dating techniques that can be applied to LIG deposits lack the precision~~

1330 Northern Hemisphere ice sheets waxed and waned (Batchelor et al., 2019) many times during the Quaternary (Batchelor et al., 2019), allowing for marine incursions prior to the LIG (e.g. during MIS 7) and during later intervals (e.g., during MIS 5c, MIS 5a, MIS 3 and the Holocene), with maximum high-stands preferentially during the transitional stages between glacials and following interglacials/interstadials. However, given difficulties inherent to dating sediments beyond the range of
1335 radiocarbon dating, chronological constraints remain a key challenge for these deposits. Here, we highlight some locations where pre-LIG and post-LIG sea-level oscillations are recorded. This list is not exhaustive; rather, it is meant to provide examples of other sea-level oscillations that are preserved in the stratigraphic record, and to provide context for the difficulties in assigning ages to marine deposits in the glaciated Northern Hemisphere.

1340 Several pre-LIG marine units have been described at Kongsfjordhallet, Svalbard (Houmark-Nielsen and Funder, 1999). The oldest marine unit is probably as much as 1 million years old. Marine sediments dating to MIS 7 have been documented at several sites in Russia. At the Bol'shaya Kheta River sections (lat/long 67.97, 83.10), in addition to LIG marine sediments (see Section 6.10), a lower marine bed contains shells of the extinct
1345 mollusk *Cyrtodaria jenisseae* Sachs (today generally considered as *Cyrtodaria angusta*, see e.g. Möller et al., 2019a) and OSL ages at 225 ± 16 , 226 ± 21 , 162 ± 23 , 231 ± 17 ka indicate a MIS 7 age (Nazarov et al., 2020; Astakhov and Semionova, 2021). A pre-LIG age is also suggested for numerous sites on the Taimyr Peninsula. Kind and Leonov (1982) claimed that >60 outcrops on central Taimyr Peninsula, south of Lake Taimyr, expose LIG-aged marine sediments
1350 . In the Bol'shaya Balakhnya river valley such sediment should reach 30–40 MASL, while further to the NE, in the Bol'shaya Rassokha river valley, such sediment should reach 60–70 MASL. However, as demonstrated in Möller et al. (2019a) from a large number of sections along the Bol'shaya Balakhnya River, these sediments are of an older age as suggested from numerous ESR and OSL dates (Möller et al., 2019a, 2019b). The sediments that Kind and Leonov (1982) presumed were LIG in age are often overlain by a till; as the sediment sections are south of the Severokokora ice marginal zone, marking the southern maximum boundary of the MIS 5d (known locally as the Early Zyryanka) glaciation over Taimyr (Möller et al., 2019a), this till must
1355 predate the MIS 5d glaciation (MIS 6 or older) and the marine sediments are consequently of a pre-LIG age.

Post-LIG marine sediments are far more commonly preserved in the stratigraphic record. For example, at the Chapoma site on the Kola Peninsula, in addition to an LIG marine sediments (see Section 6.186), an upper marine silt is present between 11 and 13 MASL (Korsakova, 2019). Two ESR dates on shells at around 12 MASL in this marine unit yielded ages of $85.5 \pm$
1360 3.2 ka and 86.0 ± 3.9 ka (Arslanov 1981), placing it into MIS 5a. Multiple sea-level oscillations are also present at the Lower Agapa River sections (Gudina et al., 1968) as well as the Strelna River site (Section 6.197) where a sequence of upper marine sediments were was dated via IRSL and ESR to between 102 ka and 84 ka (Korsakova et al., 2004; Korsakova,

2019). Möller et al. (2015, 2019a) describe several marine sequences in the central and southern parts of the Taimyr Peninsula, Russia. All these marine sediment successions were dated with ESR and OSL; with six outliers excluded the
1365 mean age is ~86 ka ($n = 62$) with an age scatter of ± 15 ka, which firmly puts these sediments into MIS 5c–b (known locally as the Early Zyryanka). Finally, marine sediments at the Anjeliko River site (Section 6.5) have been constrained to an ESR age cluster of 80–93 ka (mean age 86 ka), suggesting deposition at marine inundation during MIS 5a following a Kara Sea Ice Sheet glaciation during MIS 5d–5c.

1370 Marine sediments dating to MIS 3 have been described in the glaciated region of Russia and Europe. For example, at the Ozernaya River on October Revolution Island, offshore of Taimyr Peninsula (section 6.6), Marine unit IV sediments were exposed in two sections on opposite sides of the river valley, ~2.5 km apart (Oz 4 and Oz 5; Möller et al., 2007). These sediments contained a rhythmical sedimentation pattern, several *in situ*-positioned paired mollusc shells and almost complete skeletons of at least 9 narwhals (*Monodon Monoceros*) that were in the process of eroding out at the top of the sediment
1375 succession. The Marine IV sediments were interpreted as formed in a marine setting at a water level >65 MASL within a deeply embayed estuary behind a valley-mouth bar/barrier system in the Ozernaya valley (Möller et al., 2007). Radiocarbon ages-dating on of a narwhal tusk and molluscs yielded ages of ~50, 45.4 and 46.8 cal. ka BP-ka (all at the upper limit of radiocarbon dating), ages supported by a single mollusc shell ESR dated to 59 ka BP. The dating of Marine IV sediments thus suggests that they are of a MIS 3 age, formed at a marine inundation of the island that probably was glaciated during the
1380 whole time-span of MIS 5d–to MIS 4.

Marine sediments dating to MIS 3 are uncommon in North America and are often less reliably constrained than similar sites in Russia and Europe. Nevertheless, shallow marine and beach sediments in Eastern Canada constrained to MIS 3, using OSL and radiocarbon dating, are situated at 30, 37 and ~15 MASL (Rémillard et al., 2016, 2017). Also, in the Saint
1385 Lawrence Lowlands, marine sediments dated by radiocarbon to MIS 3 are situated at 30 MASL (Dionne and Occhietti, 1996). Finally, there is evidence for a marine incursion between the LIG and the Holocene the Severn Marine site in the Hudson Bay Lowlands, Canada was dated to MIS 3, using OSL techniques (Severn River marine unit: 55 MASL). These marine sediments have been dated via amino acid techniques (Andrew et al., 1983), TL age attempts (Forman et al., 1987) and OSL attempts (Dalton Dalton et al., 2016). However, there is no clear consensus on the age of this marine incursion as
1390 most attempts spanned MIS 5a–3 and were of relatively low precision (e.g. ± 10 ka). However, whether Hudson Bay was ice-free or remained glaciated through MIS 3 remains a topic of ongoing discussion (Dalton et al., 2019; Miller et al., 2019).

9.58.2 Holocene sea-level databases

Holocene marine sediments are widely preserved in the glaciated Northern Hemisphere and are documented in several databases, including the Russian Arctic (Baranskaya et al., 2018), the Baltic Sea (Rosentau et al., 2021), European west coast
1395 (García-Artola et al., 2018), Greenland (Long et al., 2011) and eastern Canada (Vacchi et al., 2018).

109 CONCLUSIONS AND FUTURE RESEARCH

Reconstructing sea-level change through the MIS 5e/LIG is critical for understanding the sensitivity of the Earth System to future change. We here ~~here~~ contribute 82 LIG sea-level proxies from the formerly glaciated Northern Hemisphere to the *World Atlas of Last Interglacial Shorelines* (WALIS) database. Given their position in the envelope of Northern Hemisphere ice sheets, these data are useful for testing the reliability of MIS 6 ice sheet reconstructions, and deducing LIG sea level should also be useful for inferring the relative size of MIS 6 ice sheets. Obtaining an accurate chronology remains one of the most significant challenges for LIG deposits in the glaciated region. When geochronological data are lacking, often the only way to distinguish between interglacial deposits is via palaeoecological inferences, which have their own set of uncertainties. Therefore, key areas of future research should focus on revisiting these sites to vet the stratigraphic record and testing new geochronological methods (especially U/Th, and OSL techniques, and the potential for cosmogenic nuclide methods to constrain the age of buried sediments, e.g. Balco and Rovey, 2008). Future work to locate new LIG sites (particularly in North America, which has a relative scarcity of LIG sites) should be focussed on coastal regions that are known to contain extensive stratigraphic records (e.g. the Saint Lawrence valley).

1410 DATA AVAILABILITY

The database on LIG sites in the glaciated Northern Hemisphere is available here: <https://doi.org/10.5281/zenodo.5602212> (Dalton et al., 2021). A detailed description of database fields in the WALIS database is available here: <https://doi.org/10.5281/zenodo.3961544> (Rovere et al., 2020).

1415 AUTHOR CONTRIBUTIONS

ASD and EJG conceptualised this paper. ASD wrote the first draft with substantial input from JM, PM, JPL and VA. EJG managed data entry into the WALIS database and developed the methodological framework. All authors reviewed/edited subsequent drafts of the manuscript.

COMPETING INTERESTS

1420 The authors declare no competing interests.

DISCLAIMER

The information contained in this database was the result of studies from many scientists over the course of several decades. Please cite the original sources of the data in addition to this database.

ACKNOWLEDGEMENTS

1425 [Funding for E.J.G. was provided through a Japan Society for the Promotion of Science Postdoctoral Fellowship, Helmholtz Exzellenznetzwerks "The Polar System and its Effects on the Ocean Floor \(POSY\)" and Helmholtz Climate Initiative REKLIM \(Regional Climate Change\), a joint research project at the Helmholtz Association of German research centres \(HGF\).](#) Fieldwork for J.M. was funded by several consecutive grants from the Research Council of Norway. Specific project funding for research on Taimyr/Russia was provided through grants from the Swedish Natural Science Research Council

1430 (VR) to P.M. (contract nos. G-650-199815671/2000 and 621-2008-3759) and logistics were mainly arranged and, to a large extent, funded by the Swedish Polar Research Secretariat (SPRS). [Fieldwork for J.P.L. was funded by the Academy of Finland and the Finnish Academy of Science and Letters.](#)

Overview of Russian sites by V.A. was performed according to the research plan of the St. Petersburg State University, Russia.

1435 This study was also supported by the PACES-II program at the Alfred Wegener Institute and the Bundesministerium für Bildung und Forschung funded project, PalMod. The WALIS database was developed by the ERC Starting Grant "Warmcoasts" (ERC-StG-802414) and PALSEA. PALSEA is a working group of the International Union for Quaternary Sciences (INQUA) and Past Global Changes (PAGES), which in turn received support from the Swiss Academy of Sciences and the Chinese Academy of Sciences. The database structure was designed by A. Rovere, D. Ryan, T. Lorscheid, A.

1440 Dutton, P. Chutcharavan, D. Brill, N. Jankowski, D. Mueller, M. Bartz, E.J. Gowan and K. Cohen. Thanks go to Svend Funder for helpful discussions, [as well as two anonymous Reviewers.](#)

REFERENCES

1445 Alexanderson, H., Landvik, J. Y., Molodkov, A., and Murray, A. S.: A multi-method approach to dating middle and late Quaternary high relative sea-level events on NW Svalbard – A case study, *Quat. Geochronol.*, 6, 326–340, 2011a.

Alexanderson, H., Landvik, J. Y., and Ryen, H. T.: Chronology and styles of glaciation in an inter-fjord setting, northwestern Svalbard, *Boreas*, 40, 175–197, 2011b.

Alexanderson, H., Ingólfsson, Ó., Murray, A. S., and Dudek, J.: An interglacial polar bear and an early Weichselian glaciation at Poolepynten, western Svalbard, *Boreas*, 42, 532–543, 2013.

1450 Alexanderson, H., Henriksen, M., Ryen, H. T., Landvik, J. Y. and Peterson, G.: 200 ka of glacial events in NW Svalbard: an emergence cycle facies model and regional correlations, *Arktos*, 4, 1–25, 2018.

Alexanderson, H. and Landvik, J. Y.: Sediments of MIS 5e age suggested by new OSL dates from the Skilvika section, west Svalbard, *Polar Res.*, 37, 1503907, 2018.

1455 Andersen, B. G., Sejrup, H.-P., Kirkhus, Ø.: Eemian and Weichselian Deposits at Bø on Karmøy, SW Norway: A Preliminary Report. *Norges geologiske undersøkelse* 380, 189–201, 1983.

Andersson, T., Forman, S. L., Ingólfsson, Ó., and Manley, W. F.: Late Quaternary environmental history of central Prins Karls Forland, western Svalbard, *Boreas*, 28, 292–307, 1999.

Andrews, J. T., Shilts, W. W., and Miller, G. H.: Multiple deglaciations of the Hudson Bay Lowlands, Canada, since deposition of the Missinaibi (Last-Interglacial?) Formation, *Quat. Res.*, 19, 18–37, 1983.

1460 Apukhtin, N. I.: Novye dannye po stratigraphii chetvertichnykh otlozheniy yuzhnoy chasti Kolskogo poluostrova [New data on Quaternary stratigraphy of the southeastern part of the Kola Peninsula]. Trudy VSEGEI, Leningrad: chetvertisnaya geologiya i geomorfologiya. In: Proceeding of the All-Russian Geological Institute: Quaternary Geology and Geomorphology, vol. 297, pp. 53–65 [in Russian], 1978.

Arkhipov, S. A.: A chronostratigraphic scale of the glacial Pleistocene of the West Siberian North. In: Skabichevskaya, N. A. (Ed.), *Pleistotsen Sibiri. Stratigrafia i mezhregionalnye korrelatsii*. Nauka, Novosibirsk, 19–30, 1989.

1465 Arkhipov, S. A., Levchuk, L. K. and Shelkopyas, V. N.: Marine Quaternary sediments of the Lower Ob. In: Murzayeva, V.E., Punning, J.-M. K., Chichagova, O.A., eds. *Geokronologiya chetvertichnogo perioda*, Nauka, Moscow, 90–101, 1992.

Arslanov, KhA., Yevzerov, V. Y., Tertichnyi, N. I., Gerasimov, S. A. and Lokshin, N. V.: K voprosu o vozraste otlozhenii borealnoi transgressii (ponoiskikh sloeov) na Kolskom poluostrove [On the age of deposits (Ponoi Beds) Boreal transgression in the Kola Peninsula]. In: Velichko, A.A., Faustova, M.A. (Eds.), *Pleistotsenovye Oledeneniya Vostochno-Evropeiskoi Ravniny* [Pleistocene Glaciations on the East European Plain]. Nauka Press, Moscow, 28–37 [in Russian], 1981.

Astakhov, V. I.: Age of remnants of a Pleistocene glacier in Bol'shezemel'skaya Tundra. *Doklady Earth Sciences* 384(4), 468–472. Translated from *Doklady Akademii Nauk* 384, 534–538, 2002.

1470 Astakhov, V. I., Arslanov, Kh.A. and Nazarov, D. V.: The age of mammoth fauna on the Lower Ob. *Doklady Earth Sciences* 396 (4), 538–542. Translated from *Doklady Akademii Nauk*, 396, 253–257, 2004.

Astakhov, V. I., Arslanov, Kh.A., Maksimov, F. E., Kuznetsov, V. Yu., Razina, V. V. and Nazarov, D. V.: The age of interglacial peat on the Lower Ob, Doklady Earth Sciences 401, 298-302. Translated from Doklady Akademii Nauk 2005, 400 (1), 95-99, 2005.

1480 Astakhov, V., Nazarov, D. V.: Stratigraphy of the Upper Pleistocene in northern West Siberia and its geochronometric foundation. Regionalnaya Geologiya i Metallogeniya 43, 36-47 [in Russian], 2010a.

Astakhov, V., Nazarov, D. V.: Correlation of Upper Pleistocene sediments in northern West Siberia, Quat. Sci. Rev., 29, 3616-3629, 2010b.

Astakhov, V. I.: Pleistocene glaciations of northern Russia – a modern view. Boreas 42, 1-24, 2013.

1485 Astakhov, V. I. and Semionova, L.: The penultimate interglaciation of northern Russia. Quaternary International, 605-606, 142-154, 2021

Balco, G., and Rovey, C. W: An isochron method for cosmogenic nuclide dating of buried soils and sediments, American Journal of Science, 308, 1083-1114, 2008.

Baranskaya, A. V., Khan, N. S., Romanenko, F. A., Roy, K., Peltier, W. R., Horton, B. P.: A postglacial relative sea-level database for the Russian Arctic coast, Quat. Sci. Rev., 199, 188-205, 2018.

1490 Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., Stokes, C. R., Murton, J. B., and Manica, A.: The configuration of Northern Hemisphere ice sheets through the Quaternary, Nat. Commun., 10, 3713, 2019.

Bell, J. M.: Economic Resources of Moose River Basin, Report of the Bureau of Mines Part I, 1904. 135-191, 1904.

Bennike, O. and Böcher, J.: Early Weichselian interstadial land biotas at Thule, Northwest Greenland, Boreas, 21, 111-117, 1992.

1495 Bennike, O. and Böcher, J.: Land biotas of the last interglacial/glacial cycle on Jameson Land, East Greenland, Boreas, 23, 479-487, 1994.

Bennike, O., Hansen, K.B., Knudsen, K.L., Penney, D. N., and Rasmussen, K.L.: Quaternary marine stratigraphy and geochronology in central West Greenland. Boreas, 23, 194-215, 1994.

1500 Bergsten, H., Andersson, T., and Ingólfsson, Ó.: Foraminiferal stratigraphy of raised marine deposits, representing isotope stage 5, Prins Karls Forland, western Svalbard, Polar Res., 17, 81-89, 1998.

Blake, W. J.: Geological Survey of Canada Radiocarbon Dates XXVII, Geological Survey of Canada, Paper 87-7, 100, 1988.

Bolshiyanov, D. Y. and Makeyev, V. M.: Archipelago of Severnaya Zemlya: Environmental History, Gidrometeoizdat, St. Petersburg, 1995.

1505 Bolshiyanov, D. and Molodkov, A.: Marine Pleistocene deposits of the Taymyr Peninsula and their age from ESR dating. In Kassens, H., Bauch, I., Dmitrenko, H., Eicken, H., Hubberten, H.-W., Melles, M., Thiede, J., Timokhov, L. (Eds.), Land-ocean Systems in the Siberian Arctic: Dynamics and History. Springer-Verlag, Berlin, 469-475, 1998.

Boulton, G. S.: Glacial history of the Spitsbergen archipelago and the problem of a Barents Shelf ice sheet, Boreas, 8, 31-57, 1979.

1510 Clay, P. L., Busemann, H., Sherlock, S. C., Barry, T. L., Kelley, S. P., and McGarvie, D. W.: *40Ar/39Ar ages and residual volatile contents in degassed subaerial and subglacial glassy volcanic rocks from Iceland*, *Chem. Geol.*, 403, 99-110, 2015.

[Cohen, K. M., Barnett, R., Cartelle, V., Busschers, F. S., and Barlow, N. L. M.: Last Interglacial sea-level data points for NW Europe. Earth System Science Data, in preparation](#)
[Cohen, K. M., Cartelle, V., Barnett, R., Busschers, F. S., and Barlow, N. L. M.: Last Interglacial sea-level data points from Northwest Europe, Earth Syst. Sci. Data Discuss. \[preprint\]](#)
<https://doi.org/10.5194/essd-2021-390>, in review, 2021.

Formatted: Font: (Default) Times New Roman, 10 pt, Forecolor: Auto, English (United States), Pattern: Clear

Formatted: Font color: Auto

1515 Dahl-Jensen, D., Albert, M. R., Aldahan, A., Azuma, N., Balslev-Clausen, D., Baumgartner, M., Berggren, A. M., Bigler, M., Binder, T., Blunier, T., Bourgeois, J. C., Brook, E. J., Buchardt, S. L., Buizert, C., Capron, E., Chappellaz, J., Chung, J., Clausen, H. B., Cvijanovic, I., Davies, S. M., Ditlevsen, P., Eicher, O., Fischer, H., Fisher, D. A., Fleet, L. G., Gfeller, G., Gkinis, V., Gogineni, S., Goto-Azuma, K., Grinsted, A., Gudlaugsson, H., Guillemin, M., Hansen, S. B., Hansson, M., Hirabayashi, M., Hong, S., Hur, S. D., Huybrechts, P., Hvidberg, C. S., Iizuka, Y., Jenk, T., Johnsen, S. J., Jones, T. R., Jouzel, J., Karlsson, N. B., Kawamura, K., Keegan, K., Kettner, E., Kipfstuhl, S., Kjaer, H. A., Koutnik, M., Kuramoto, T., Köhler, P., Laepple, T., Landais, A., Langen, P. L., Larsen, L. B., Leuenberger, D., Leuenberger, M., Leuschen, C., Li, J., Lipenkov, V., Martinerie, P., Maselli, O. J., Masson-Delmotte, V., McConnell, J. R., Miller, H., Mini, O., Miyamoto, A., Montagnat-Rentier, M., Mulvaney, R., Muscheler, R., Orsi, A. J., Paden, J., Panton, C., Pattyn, F., Petit, J. R., Pol, K., Popp, T., Possnert, G., Prié, F., Prokopiou, M., Quiquet, A., Rasmussen, S. O., Raynaud, D., Ren, J., Reutenauer, C., Ritz, C., Röckmann, T., Rosen, J. L., Rubino, M., Rybak, O., Samyn, D., Sapart, C. J., Schilt, A., Schmidt, A. M. Z., Schwander, J., Schüpbach, S., Seierstad, I., Severinghaus, J. P., Sheldon, S., Simonsen, S. B., Sjolte, J., Solgaard, A. M., Sowers, T., Sperlich, P., Steen-Larsen, H. C., Steffen, K., Steffensen, J. P., Steinhage, D., Stocker, T. F., Stowasser, C., Sturevik, A. S., Sturges, W. T., Sveinbjörnsdóttir, A., Svensson, A., Tison, J. L., Uetake, J., Vallelonga, P., van de Wal, R. S. W., van der Wel, G., Vaughn, B. H., Vinther, B., Waddington, E., Wegner, A., Weikusat, I., White, J. W. C., Wilhelms, F., Winstrup, M., Wittrant, E., Wolff, E. W., Xiao, C., and Zheng, J.: *Eemian interglacial reconstructed from a Greenland folded ice core*, *Nature*, 493, 489-494, 2013.

Dalton, A. S., Finkelstein, S. A., Barnett, P. J., and Forman, S. L.: *Constraining the Late Pleistocene history of the Laurentide Ice Sheet by dating the Missinaibi Formation, Hudson Bay Lowlands, Canada*, *Quat. Sci. Rev.*, 146, 288-299, 2016.

1535 Dalton, A. S., Gowan, E. J., Mangerud, J., Möller, P., Lunkka, J. P., and Astakhov V: *Last interglacial (MIS 5e) sea level proxies in the glaciated Northern Hemisphere (1.0)* [Data set]. Zenodo: <https://doi.org/10.5281/zenodo.5602212>, 2021.

1540 Davies, S. M., Abbott, P. M., Meara, R. H., Pearce, N. J. G., Austin, W. E. N., Chapman, M. R., Svensson, A., Bigler, M., Rasmussen, T. L., Rasmussen, S. O., and Farmer, E. J.: *A North Atlantic tephrostratigraphical framework for 130–60 ka b2k: new tephra discoveries, marine-based correlations, and future challenges*, *Quat. Sci. Rev.*, 106, 101-121, 2014.

Der Sarkissian, C., Möller, P., Hofman, C. A., Ilsøe, P., Rick, T. C., Schiøtte, T., Sørensen, M. V., Dalén, L., and Orlando, L.: *Unveiling the ecological applications of ancient DNA from mollusk shells*, *Front. Ecol. Evol.*, 8, 2020.

Desor, M. E. and Cabot, E. C.: *On the Tertiary and more recent deposits in the Island of Nantucket*, *Geological Society of London Quarterly Journal*, 5, 340-344, 1849.

1545 Devyatova, E. I. and Loseva, E. I.: *Stratigraphy and Palaeogeography of the Quaternary Period in the Mezen River Basin*, Akademiya Nauk SSSR, Izdalstvo Nauka, Leningrad, 105 p. [in Russian], 1964.

1550 Devyatova, E. I.: *Late Pleistocene Environments as Related to Human Migrations in the Dvina River Basin and in Karelia*, Karelia, Petrozavodsk [in Russian], 156 p., 1982.

Dionne, J.-C. and Occhietti, S.: Aperçu du Quaternaire à l'embouchure du Saguenay, Québec, *Geogr. Phys. Quatern.*, 50, 5-34, 1996.

1555 Donner, J.: The Eemian site of Norinkylä compared with other interglacial and interstadial sites in Ostrobothnia, Western Finland, *Annales Academiae Scientiarum Fennicae A III*, 149, 31, 1988.

Douka, K., Higham, T. F. G., and Hedges, R. E. M.: Radiocarbon dating of shell carbonates: old problems and new solutions, *Munibe Suplemento*, 31, 18-27, 2010.

1560 Dredge, L. A., Mott, R. J., and Grant, D. R.: Quaternary stratigraphy, paleoecology, and glacial geology, Îles de la Madeleine, Québec, *Can. J. Earth Sci.*, 29, 1981-1996, 1992.

Duller, G. A. T.: Single-grain optical dating of Quaternary sediments: why aliquot size matters in luminescence dating, *Boreas*, 37, 589-612, 2008.

Dutton, A. and Lambeck, K.: Ice Volume and Sea Level During the Last Interglacial, *Science*, 337, 216-219, 2012.

1565 Dyer, B., Austermann, J., D'Andrea, W. J., Creel, R. C., Sandstrom, M. R., Cashman, M., Rovere, A., Raymo, M. E.: Sea-level trends across The Bahamas constrain peak last interglacial ice melt, Proc. Natl. Acad. Sci. U.S.A., 118, e2026839118, 2021.

Eriksson, B., Grönlund, T. and Kujansuu, R.: Interglasiaalikerrostuma Evijärvellä, Pohjanmaalla. (An interglacial deposit at Evijärvi, in the Pohjanmaa region, Finland). *Geologi*, 32, 65-71, [in Finnish with English summary], 1980.

1570 Eriksson, B.: The Eemian pollen stratigraphy and vegetational history of Ostrobothnia, Finland. *Bulletin of the Geological Survey of Finland* 372, 36 p., 1993.

Eriksson, B., Grönlund, T. and Uutela A.: Biostratigraphy of Eemian sediments at Mertuanjoa, Pohjanmaa (Ostrobothnia), western Finland, *Boreas* 28, 274-291, 1999.

Feyling-Hanssen, R. W.: The Clyde Foreland, Field Report North-Central Baffin Island 1966. Department of Energy, Mines & Resources, Geographic Branch, Ottawa, 35-55, 1967.

1575 Field, M. H., Huntley, B., and Müller, H.: Eemian climate fluctuations observed in a European pollen record, *Nature*, 371, 779-783, 1994.

Flude, S., Burgess, R., and McGarvie, D. W.: Silicic volcanism at Ljósufjöll, Iceland: Insights into evolution and eruptive history from Ar-Ar dating, *J. Volcanol. Geotherm. Res.*, 169, 154-175, 2008.

1580 Forman, S. L., Wintle, A. G., Thorleifson, L. H., and Wyatt P. H.: Thermoluminescence properties and age estimates for Quaternary raised marine sediments, Hudson Bay Lowland, Canada, Can. J. Earth Sci., 24, 2405-2411, 1987.

Forman, S. L.: Post-glacial relative sea-level history of northwestern Spitsbergen, Svalbard, *Geol. Soc. Am. Bull.*, 102, 1580-1590, 1990.

Forman, S. L.: Infrared and Red Stimulated Luminescence Dating of Late Quaternary Nearshore Sediments from Spitsbergen, Svalbard, *Arct. Antarct. Alp. Res.*, 31, 34-49, 1999.

1585 Forsström, L., Eronen, M. and Grönlund, T.: On marine phases and shoreline levels of the Eemian interglacial and Weichselian interstadials on the coast of Ostrobothnia, Finland. Geological Survey of Finland, Special Paper 2, 37-42, 1987.

Forsström L., Aalto M., Eronen M. and Grönlund T.: Stratigraphic evidence for Eemian crustal movements and relative sea-level changes in eastern Fennoscandia, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 68, 317-335, 1988.

Forsström, L. and Punkari, M.: Initiation of the last glaciation in northern Europe, *Quat. Sci. Rev.*, 16, 1197-1215, 1997.

1590 Fuller, M. L.: The geology of Long Island, New York, U. S. Geological Survey Professional Paper 82, 1914. 231 p., 1914.

Funder, S., Demidov, I., and Yelovicheva, Y.: Hydrography and mollusc faunas of the Baltic and the White Sea–North Sea seaway in the Eemian, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 184, 275-304, 2002.

Gao, C., Huot, S., McDonald, A. M., Crabtree, D. C., and Turton, C. L.: Subtill nonglacial deposits and their climatic implications for the Last Interglacial (MIS 5e), Hudson Bay Lowlands, Canada, *Quat. Sci. Rev.*, 248, 106590, 2020.

1595 García-Artola, A., Stéphan, P., Cearreta, A., Kopp, R. E., Khan, N. S., and Horton, B. P.: Holocene sea-level database from the Atlantic coast of Europe, *Quat. Sci. Rev.*, 196, 177-192, 2018.

Godfrey-Smith, D. I., Huntley, D. J., and Chen, W. H.: Optical dating studies of quartz and feldspar sediment extracts, *Quat. Sci. Rev.*, 7, 373-380, 1988.

1600 Grave, M.K., Gunova, V.S., Devyatova, E.I., Lavrova, M.A., Lebedeva, R.M., Samsonova, L.Ya and Cheremisinova, E.A.: Mikulinskoye mezhlednikoviyne na yugo-vostoke Kolskogo poluostrova [Mikulinian interglacial in the southeastern Kola Peninsula]. In: Grave, M.K., Koshechkin, B.I. (Eds.), *Main Problems of Geomorphology and Anthropogen Stratigraphy of the Kola Peninsula*. Nauka Press, Leningrad, pp. 25-56 [in Russian], 1969.

Grönlund, T.: The diatom stratigraphy of the Eemian Baltic Sea on the basis of sediment discoveries in Ostrobothnia, Finland. Geological Survey of Finland, Report of Investigation 112, 26 pp, 1991a.

1605 Grönlund, T.: New Cores from Eemian interglacial deposits in Ostrobothnia, Finland. Geological Survey of Finland Bulletin 352, 31 pp, 1991b.

Grøsfjeld, K., Funder, S., Seidenkrantz, M. S., and Glaister, C.: Last Interglacial marine environments in the White Sea region, northwestern Russia, *Boreas* 35, 493-520, 2006.

Grün, R.: Electron spin resonance (ESR) dating, *Quat. Int.*, 1, 65-109, 1989.

1610 Gudina, V. I., Nuzhdina, N. A., and Troitskiy, S. L.: New data on the marine Pleistocene of the Taymyr Lowland, Akademiya Nauk SSSR, Sibirskoe Otdelenie. *Geologia i Geofizika*, 1968. 40-48 [in Russian], 1968.

Gudina, V. I., Yevzerov, V. Ya.: The Stratigraphy and Foraminifera of the Upper Pleistocene in the Kola Peninsula. The British Library Board, p. 1981. <https://doi.org/10.1017/S0016756800026224>. Translated by Lees, E., Edited by Hughes, M, 1973.

1615 Gudina, V. I., Kryukov, V. D., Levchuk, L. K., and Sudkov, L. A.: Upper-Pleistocene sediments in north-eastern Taymyr, *Bulletin of Commission on Quaternary Researches*, 52, 90-97 [in Russian], 1983.

Gusskov, S.A. and Levchuk, L.K.: Foraminiferal complexes and palaeoceanographic reconstructions of the Middle and Late Pleistocene interglacial basins in the North of Siberia. *Antropozoikum, Quaternary of Siberia. Journal of Geological Sciences* 23, Czech Geological Survey, Prague, 125–132, 1999.

1620 Gustavson, T. C.: A Warm-Water Pleistocene Fauna from the Gardiners Clay of Eastern Long Island. *J. Paleontol.*, 46, 447-449, 1972.

Hansen, L. A., Ørgensen, M. E., Houmark-Nielsen, M., and Kronborg, C.: Late Pleistocene stratigraphy and depositional environments of the Fynselv area, Jameson Land, East Greenland, *Boreas*, 23, 385-397, 1994.

1625 Head, M. J., Seidenkrantz, M.-S., Janczyk-Kopikowa, Z., Marks, L., and Gibbard, P. L.: Last Interglacial (Eemian) hydrographic conditions in the southeastern Baltic Sea, NE Europe, based on dinoflagellate cysts, *Quat. Int.*, 130, 3-30, 2005.

Helmens, K.F., Räsänen, M., Johansson, P.W., Jungner, H. and Korjonen, K.: The Last Interglacial – Glacial cycle in NE Fennoscandia: a nearly continuous record from Sokli (Finnish Lapland), *Quat. Sci. Rev.*, 19, 1605-1623, 2000.

Hirvas, H.: Pleistocene stratigraphy of Finnish Lapland. *Geological Survey of Finland Bulletin* 354, 123 pp, 1991.

1630 Høeg, H. I.: Pollenanalytiske undersøkelser i Rogaland og Ersdal i Vest-Agder [Pollen analytical investigations in Rogaland and in Ersdal in Vest-Agder]. Artikksamling til Kerstin Griffin på 60-årsdagen [Museum landscape. A collection of papers for the 60th birthday of Kerstin Griffin]. AmS-Rapport 12A, 1999. 145–226, 1999.

Houmark-Nielsen, M. and Funder, S.: Pleistocene stratigraphy of Kongsfjordhallet, Spitsbergen, Svalbard, *Polar Res.*, 18, 39-50, 1999.

1635 Houmark-Nielsen, M., Demidov, I., Funder, S., Grøsfjeld, K., Kjær, K. H., Larsen, E., Lavrova, N., Lyså, A. and Nielsen, J. K.: Early and Middle Valdaian glaciations, ice-dammed lakes and periglacial interstadials in northwest Russia: new evidence from the Pyoza River area, *Glob. Planet. Change* 31, 215-237, 2001.

Huntley, D. J., Godfrey-Smith, D. I., and Thewalt, M. L. W.: Optical dating of sediments, *Nature*, 313, 105-107, 1985.

1640 Hütt, G., Punning, J.-M., Mangerud, J.: Thermoluminescence dating of the Eemian-Early Weichselian sequence at Fjøsanger, western Norway, *Boreas* 12, 227-231, 1983.

Hütt, G., Molodkov, A., Kessel, H., and Raukas, A.: ESR dating of subfossil Holocene shells in Estonia, *Nucl. Tracks Radiat. Meas.*, 10, 891-898, 1985.

1645 Ikonen, L., Ekman, I.: Biostratigraphy of the Mikulino interglacial sediments in NW Russia: the Petrozavodsk site and a literature review. *Annales Academiae Scientiarum Fennicae A III Geologica-Geographica* 161. 88 pp, 2001.

Ingólfsson, Ó., Lyså, A., Funder, S., Möller, P., Björck, S.: Late Quaternary glacial history of the central west coast of Jameson Land, East Greenland, *Boreas*, 23, 447-458, 1994.

1650 Ingólfsson, Ó., Rögnvaldsson, F., Bergsten, H., Hedenäs, L., Lemdahl, G., Lirio, J. M., and Sejrup, H. P.: Late Quaternary glacial and environmental history of Kongsøya, Svalbard, *Polar Res.*, 14, 123-139, 1995.

Ingólfsson, Ó. and Wiig, Ø.: Late Pleistocene fossil find in Svalbard: the oldest remains of a polar bear (*Ursus maritimus* Phipps, 1744) ever discovered, *Polar Res.*, 28, 455-462, 2009.

Janczyk-Kopikowa, Z. and Marks, L.: Eemian marine sediments in Poland, Unpublished Report, Polish Geological Institute, Warsaw University, Warsaw, 2002. 5, 2002.

1655 Jørgensen, T., Haile, J., Möller, P., Andreev, A., Boessenkool, S., Rasmussen, M., Kienast, F., Coissac, E., Taberlet, P., Brochman, C., Bigelow, N. H., Andersen, K., Orlando, L., Gilbert, M. P. T. and Willerslev, E.: A comparative study of ancient sedimentary DNA, pollen and macrofossils from permafrost sediments in Northern Siberia reveals long-term vegetational stability, *Mol. Ecol.*, 21, 1989-2003, 2012.

1660 Kelly, M.: Quaternary pre-Holocene marine events of western Greenland. *Rapport Grønlands Geologiske Undersøgelse*, 131, 1-23, 1986.

Kelly, M., Funder, S., Houmark-Nielsen, M., Knudsen, K., L., Kronborg, C., Landvik, J., and Sorby, L.: Quaternary glacial and marine environmental history of northwest Greenland: a review and reappraisal, *Quat. Sci. Revs.*, 18, 373-392, 1999.

Kind, N. V.: Geokronologija pozdniego antropogena po izotopnym dannym [Late Quaternary geochronology according to isotope data]. Moscow, Nauka, 255 p. [in Russian], 1974.

1665 Kind, N. V. and Leonov, B. N.: *Antropogen Taimyra* (The Anthropocene of Taymyr Peninsula), Nauka, Moscow, 1982.

Knudsen, K. L., Jiang, H., Gibbard, P. L., Kristensen, P., Seidenkrantz, M.-S., Janczyk-Kopikowa, Z., and Marks, L.: Environmental reconstructions of Eemian Stage interglacial marine records in the Lower Vistula area, southern Baltic Sea, *Boreas*, 41, 209-234, 2012.

1670 Korsakova, O.P., Molodkov, A.N., Kolka, V.V.: Geological-stratigraphic position of Upper Pleistocene marine sediments in the Southern Kola Peninsula: evidence from geochronological and geological data. *Dokl. Earth Sci.* 398, 908-912, 2004.

Korsakova, O. P.: Pleistocene marine deposits in the coastal areas of Kola Peninsula (Russia), *Quaternary International*, 206, 3-15, 2009.

Korsakova, O., Kolka, V., Semenova, L.: Late Pleistocene stratigraphy according to the sediments sequence from eastern Kola Peninsula, Ponoi River Valley (North-Western Russia), *Quat. Int.*, 420, 280-293, 2016.

1675 Korsakova, O.: Formal stratigraphy of the Neopleistocene (Middle and Upper/Late Pleistocene) in the Kola region, NW Russia, *Quat. Int.*, 534, 42-59, 2019.

Korsakova, O.: Upper Pleistocene and Holocene stratigraphy in the Kola Peninsula and Northern Karelia (N-W Russia): Marine and glacial units. *Quat. Int.*, [605-606, 108-125, 2021, #press](#).

1680 Krzmińska, J.: Analiza mikrofauny i malakofauny z osadów strefy brzegowej – czesc zachodnia, *Arch. Oddz. Geologii Morza. PIG-PIB*, Gdańsk, 1996.

Krzyszkowski, D., Dobrakowa, E., Dobrakci, R., Czerwonka, J. A., and Kuszell, T.: Stratigraphy of Weichselian deposits in the cliff sections between Lukecin and Niechorze, Baltic Coast, northwestern Poland, *Quaternary Studies in Poland*, 16, 27-45, 1999.

1685 Krzyszkowski, D.: Till stratigraphy, petrography and palaeogeography along the northwestern coastal region of Poland, *Buletyn państwowego instytutu geologicznego*, 438, 51-92, 2010.

Lakeman, T. R. and England, J. H.: Late Wisconsinan glaciation and postglacial relative sea-level change on western Banks Island, Canadian Arctic Archipelago, *Quaternary Research*, 80, 99-112, 2014.

Lambeck, K., Purcell, A., Funder, S., Kjær, K. H., Larsen, E. and Möller, P.: Constraints on the Late Saalian to early Middle Weichselian ice sheet of Eurasia from field data and rebound modelling, Boreas, 35, 539-575, 2006.

1690 Lamothé, M. and Huntley, D. J.: Thermoluminescence Dating of Late Pleistocene Sediments, St. Lawrence Lowland, Québec, *Geogr. Phys. Quatern.*, 42, 33, 1988.

Landvik, J. Y., Bolstad, M., Lycke, A. K., Mangerud, J., and Sejrup, H. P.: Weichselian stratigraphy and palaeoenvironments at Bellsund, western Svalbard, *Boreas*, 21, 335-358, 1992.

1695 Landvik, J. Y., Lyså, A., Funder, S. and Kelly, M.: The Eemian and Weichselian stratigraphy of the Langelandselv area, Jameson Land, East Greenland, *Boreas*, 23, 412-423, 1994.

Lane, T. P., Roberts, D. H., Ó Cofaigh, C., Vieli, A., and Moreton, S. G.: The glacial history of the southern Svartenhuk Halvø, West Greenland, *Arktos*, 19, 2015.

1700 Lavrova, M. A.: Chetvertichnaya Geologiya Kolskogo Poluostrova [Quaternary Geology of the Kola Peninsula]. Academy of Sciences of USSR Press, Moscow- Leningrad [in Russian], 1960.

Lavrova, M. A. and Troitsky, S. L.: Interglacial transgressions in northern Europe and Siberia. In: Gromov, V.I. (Ed.) *Khronologiya i klimaty chetvertichnogo perioda*. [Chronology and climates of the Quaternary]. Academy of Sciences Press, Moscow, 124-135 [in Russian], 1960.

1705 Lavrushin, J. A.: Pleistocene of Northern Spitsbergen [in Russian, internal English translation, The Norwegian Polar Research Institute], *Doklady Akademii Nauka, SSR*, 176(I), 167-170, 1967.

Lavrushin, J. A.: Quaternary Deposits of Spitsbergen [in Russian, internal English translation, The Norwegian Polar Research Institute], Nauka, Moscow, 1969.

1710 Lehman, S. J. and Forman, S. L.: Late Weichselian glacier retreat in Kongsfjorden, west Spitsbergen, Svalbard, *Quat. Res.*, 37, 139-154, 1992.

Liivrand, E.: Regional type section of the Eemian marine deposits on Suur-Prangli [In Russian with an English Summary], *P. Est. Acad. Sci. Geol.*, 36, 20-26, 1987.

Liivrand, E.: Biostratigraphy of the Pleistocene deposits in Estonia and correlations in the Baltic region, Stockholm University, Stockholm, 114 pp., 1991.

1715 Lindqvist, C., Schuster, S. C., Sun, Y., Talbot, S. L., Qi, J., Ratan, A., Tomsho, L. P., Kasson, L., Zeyl, E., Aars, J., Miller, W., Ingólfsson, Ó., Bachmann, L., and Wiig, Ø.: Complete mitochondrial genome of a Pleistocene jawbone unveils the origin of polar bear, *Proc. Natl. Acad. Sci. U. S. A.*, 107, 5053-5057, 2010.

Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, PA1003, 2005.

1720 Løken, O. H.: Baffin Island refugia older than 54 000 years, *Science*, 153, 1378-1380, 1966.

Long, A. J., Woodroffe, S. A., Roberts, D. H., Dawson, S.: Isolation basins, sea-level changes and the Holocene history of the Greenland Ice Sheet, *Quat. Sci. Rev.*, 30, 3748-3768, 2011.

Lukashov, A. D.: Guidebook for Excursions A-4, C-4, Karelia, INQUA XI Congress, Moscow, 57 pp., 1982.

Lunkka, J. P., Lintinen, P., Nenonen, K. and Huhta, P.: Stratigraphy of the Koivusaarenneva exposure and its correlation across central Ostrobothnia, Finland. *Bulletin of the Geological Society of Finland* 88, 53–67, 2016.

Lunkka, J. P., Kaparulina, E. Putkinen, N. and Saarnisto, M.: Late Pleistocene palaeoenvironments and the last deglaciation on the Kola Peninsula, Russia, *Arktos* 4, 18, 2018.

Lunkka J. P., Sarala P. and Gibbard P. L.: Morphostratigraphy and its application to correlation of multiple till sequences in formerly glaciated terrains – an example from the eastern flank of the Scandinavian Ice Sheet. 3rd International Congress of Stratigraphy - Strati 2019, 2nd-9th July 2019 Milano. Abstracts p. 347, 2019.

Lycke, A. K., Mangerud, J., and Sejrup, H. P.: Late Quaternary foraminiferal stratigraphy from western Svalbard, *Boreas*, 21, 271-288, 1992.

Makowska, A.: Pleistocene seas in Poland-sediments, age and palaeogeography, *Prace Instytutu Geologicznego*, 120, 1-74, 1986.

1735 Makowska, A., Kabulski, P., and Uniejewska, M.: The Tychnowy sea deposits from new boreholes against the structure of the Pleistocene complex in the Prabuty-Susz area, lower Vistula region, *Bulletyn Panstwowego Instytutu Geologicznego*, 398, 69-84, 2001.

Mamakowa, K.: Pollen stratigraphy of the Eemian and adjoining glacial deposits based on continuous sequences in Poland, *B. Pol. Acad. Sci.*, 36, 299-307, 1988.

1740 Mamakowa, K.: Late Middle Polish glaciation, Eemian and Early Vistulian vegetation at Imbramowice near Wroclaw and the pollen stratigraphy of this part of the Pleistocene in Poland, *Acta Palaeobot.*, 29, 11-176, 1989.

Mangerud, J., Sønstegaard, E., and Sejrup, H.-P.: Correlation of the Eemian (interglacial) Stage and the deep-sea oxygen-isotope stratigraphy, *Nature* 277, 189-192, 1979.

1745 Mangerud, J., Sønstegaard, E., Sejrup, H.-P. and Haldorsen, S.: A continuous Eemian-Early Weichselian sequence containing pollen and marine fossils at Fjøsanger, western Norway, *Boreas* 10, 137-208, 1981.

Mangerud, J. and Svendsen, J. I.: The last interglacial-glacial period on Spitsbergen, Svalbard, *Quat. Sci. Rev.*, 11, 633-664, 1992.

Mangerud, J. and Funder, S.: The interglacial-glacial record at the mouth of Scoresby Sund, East Greenland, *Boreas*, 23, 349-358, 1994.

1750 Mangerud, J., Dokken, T., Hebbeln, D., Heggen, B., IngÓlfsson, Ó., Landvik, J. Y., Mejdahl, V., Svendsen, J. I., and Vorren, T. O.: Fluctuations of the Svalbard-Barents Sea ice sheet during the last 150 000 years, *Quat. Sci. Rev.*, 17, 11-42, 1998.

Mangerud, J., Svendsen, J. I. and Astakhov, V. I.: Age and extent of the Barents and Kara Sea ice sheets in Northern Russia, *Boreas*, 28, 46-80, 1999.

1755 Mangerud, J. and Svendsen, J. I.: The Holocene Thermal Maximum around Svalbard, Arctic North Atlantic; molluscs show early and exceptional warmth, *Holocene*, 28, 65-83, 2018.

Mangerud, J., Hughes, A.L.C., Sæle, T.H. and Svendsen, J.I.: Ice-flow patterns and precise timing of ice sheet retreat across a dissected fjord landscape in western Norway, *Quat. Sci. Rev.*, 214, 139-163, 2019.

1760 Mauz, B., Vacchi, M., Green, A., Hoffmann, G., and Cooper, A.: Beachrock: a tool for reconstructing relative sea level in the far-field, *Mar. Geol.*, 362, 1-16, 2015.

McNeely, R. and McCuaig, S.: Geological Survey of Canada Radiocarbon Dates XXIX, Geological Survey of Canada, Paper 89-7, 1-134, 1991.

McNeely, R.: Geological Survey of Canada Radiocarbon Dates XXXIII, Geological Survey of Canada, Current Research 1765 2001, 1-51, 2002.

Mejdahl, V. and Funder, S. Luminescence dating of Late Quaternary sediments from East Greenland, *Boreas* 23, 525-535, 1994.

Miettinen, A., Rinne, K., Haila, H., Hyvärinen, H., Eronen, M., Delusina, I., Kadastik, E., Kalm, V., and Gibbard, P.: The 1770 marine Eemian of the Baltic: new pollen and diatom data from Peski, Russia, and Põhja-Uhtju, Estonia, *J. Quat. Sci.*, 17, 445-458, 2002.

Miettinen, A., Head, M. J., and Knudsen, K. L.: Eemian sea-level highstand in the eastern Baltic Sea linked to long-duration White Sea connection, *Quat. Sci. Rev.*, 86, 158-174, 2014.

Miller, G. H., Andrews, J. T. and Short, S. K.: The last interglacial-glacial cycle, Clyde foreland, Baffin Island, N.W.T.: 1775 stratigraphy, biostratigraphy, and chronology, *Canadian Journal of Earth Sciences*, 14, 2824-2857, 1977.

Miller, G. H.: Quaternary Depositional Episodes, Western Spitsbergen, Norway: Aminostratigraphy and Glacial History, *Arct. Antarct. Alp. Res.*, 14, 321-340, 1982.

Miller, G. H. and Mangerud, J.: Aminostratigraphy of European marine interglacial deposits, *Quat. Sci. Rev.*, 4, 215-278, 1985.

Miller, G. H., Sejrup, H. P., Lehman, S. J., and Forman, S. L.: Glacial history and marine environmental change during the 1780 last interglacial-glacial cycle, western Spitsbergen, Svalbard, *Boreas*, 18, 273-296, 1989.

Miller, G. H. and Andrews, J. T.: Hudson Bay was not deglaciated during MIS 3, *Quat. Sci. Rev.*, 225, 105944, 2019.

Möller, P., Bolshiyanov, D. Yu. and Bergsten, H.: Weichselian geology and palaeoenvironmental history of the central Taymyr Peninsula, Siberia, indicating no glaciation during the last global glacial maximum, *Boreas*, 28, 92-114, 1999.

1785 Möller, P., Lubinski, D. J., Ingólfsson, Ó., Forman, S. L., Seidenkrantz, M.-S., Bolshiyanov, D. Y., Lokrantz, H., Antonov, O., Pavlov, M., Ljung, K., Zeeberg, J., and Andreev, A.: Erratum to: Severnaya Zemlya, Arctic Russia: a nucleation area for Kara Sea ice sheets during the Middle to Late Quaternary, *Quat. Sci. Rev.*, 26, 1149-1191, 2007.

Möller, P., Fedorov, G., Pavlov, M., Seidenkrantz, M.-S., and Sparrenbom, C.: Glacial and palaeoenvironmental history of the Cape Chelyuskin area, Arctic Russia, *Polar Res.*, 27, 222-248, 2008.

1790 Möller, P., Alexanderson, H., Funder, S., and Hjort, C.: The Taimyr Peninsula and the Severnaya Zemlya archipelago, Arctic Russia: a synthesis of glacial history and palaeo-environmental change during the Last Glacial cycle (MIS 5e–2), *Quat. Sci. Rev.*, 107, 149–181, 2015.

Möller, P., Benediktsson, Í. Ö., Anjar, J., Bennike, O., Bernhardson, M., Funder, S., Håkansson, L. M., Lemdahl, G., Licciardi, J. M., Murray, A. S., and Seidenkrantz, M.-S.: Glacial history and palaeo-environmental change of southern Taimyr Peninsula, Arctic Russia, during the Middle and Late Pleistocene, *Earth-Sci. Rev.*, 196, 102832, 2019a.

1795 Möller, P., Benediktsson, I. O., Anjar, J., Bennike, O., Bernhardson, M., Funder, S., Håkansson, L. M., Lemdahl, G., Licciardi, J. M., Murray, A. S., and Seidenkrantz, M. S.: Data set on sedimentology, palaeoecology and chronology of Middle to Late Pleistocene deposits on the Taimyr Peninsula, Arctic Russia, *Data Brief*, 25, 104267, 2019b.

Molodkov, A.: ESR dating of Quaternary shells: recent advances, *Quat. Sci. Revs.*, 7, 477–484, 1988.

1800 Molodkov, A. N. and Raukas, A.: The age of Upper Pleistocene marine deposits of the Boreal transgression on the basis of electron-spin resonance (ESR) dating of subfossil mollusc shells, *Boreas* 17, 267–272, 1988.

Molodkov, A., Dreimanis, A., ĀBoltiņš, O. and Raukas, A.: The ESR age of *Portlandia arctica* shells from glacial deposits of Central Latvia: an answer to a controversy on the age and genesis of their enclosing sediments, *Quat. Sci. Revs.*, 17, 1077–1094, 1998.

1805 Molodkov, N. A. and Bolikhovskaya, N. S.: Eustatic sea-level and climate changes over the last 600 ka as derived from mollusc-based ESR-chronostratigraphy and pollen evidence in Northern Eurasia, *Sediment. Geol.* 150, 185–201, 2002.

Mott, R. J., and Grant, D. R.: Pre-Wisconsinan paleoenvironments in Atlantic Canada, *Geographie physique et Quaternaire*, 39: 239–254, 1985.

Murray, A. S., Svendsen, J.I., Mangerud, J. and Astakhov, V.I.: Testing the accuracy of quartz OSL dating using a known age Eemian site on the Sula river, northern Russia, *Quat. Geochron.*, 2, 102–109, 2007.

1810 Nazarov, D.V.: New data on Quaternary sediments in the central part of the West Siberian Arctic. *Regionalnaya Geologiya i Metallogeniya* 30/31, 213–221 [in Russian], 2007.

Nazarov, D.V.: Quaternary deposits of the Central part of the West Siberian Arctic. Abstract of the thesis. diss. Cand. geol-min. sciences. SPb. [in Russian], 2011

1815 Nazarov, D.V., Pushka, D.V., Nikolskaya, O.A. and Gerasicheva, A.V.: State geological map of the Russian Federation. Scale 1: 1,000,000 (third generation). Series West Siberian. Map of Quaternary formations. Sheet R-44 - Gydan Peninsula. Cartographic factory VSEGEI. [in Russian], 2018.

Nazarov, D.V., Nikolskaya, O.A., Gladysheva, A.S., Zhigmanovsky, I.V., Ruchkin, M.V. and Merkuljev, A.V.: Pleistocene marine deposits of the Lower Yenisei basin: OSL age, facies and paleontological characteristics. *Regionalnaya Geologiya i Metallogeniya* 82, 16–34 [in Russian], 2020.

1820 Nazarov, D. V. Nikolskaia, O. A., Gladysheva, A. S., Zhigmanovskiy, I. V., Ruchkin, M., V., Merkuljev, A. V., Thomsen, K., J.: Evidence for the intrusion of marine Atlantic waters into the West Siberian Arctic during the Middle Pleistocene, *Boreas*, <https://doi.org/10.1111/bor.12558>, 2021.

Nenonen, K., Eriksson, B. and Grönlund, T.: The Till Stratigraphy of Ostrobothnia, Western Finland, with Reference to New Eemian Interglacial Sites. *Striae* 34, 65–76, 1991.

1825 Nenonen, K.: Pleistocene Stratigraphy and Reference Sections in Southern and Western Finland. Geological Survey of Finland, Regional Office for Mid-Finland. 94 pp, 1995.

Nemickas, B. and Koszalka, E. J.: Geohydrologic Appraisal of Water Resources of the South Fork, Long Island, New York, Geological Survey water-supply paper 2073, 1982. 55 p. + maps, 1982.

1830 Niemelä, J. and Tynni, R.: Interglacial and interstadial sediments in the Pohjanmaa region, Finland, Bulletin of the Geological Survey of Finland 302, 48 pp, 1979.

Nikonov, A.A.: Stratigrafiya i paleogeografiya antropogena Kolskogo poluostrova i prilegayushchikh oblastei [Quaternary stratigraphy and paleogeography of the Kola Peninsula and adjacent areas]. In: Grichuk, V.P. (Ed.), Verhniy Pleistocen. Stratigrafiya I Absolyutnaya Geohronologiya [Upper Pleistocene. Stratigraphy and Geochronology]. Nauka Press, Moscow, pp. 92–105 [in Russian], 1966.

1835 Occhietti, S., Long, B., Clet, M., Boespflug, X., and Sabeur, N.: Séquence de la transition Illinoien-Sangamonien: forage IAC-91 de l'Île aux Coudres, estuaire moyen du Saint-Laurent, Québec, Can. J. Earth Sci., 32, 1950–1964, 1995.

Oldale, R. N., Valentine, P. C., Cronin, T. M., Spiker, E. C., Blackwelder, B. W., Belknap, D. F., Wehmiller, J. F., and Szabo, B. J.: Stratigraphy, structure, absolute age, and paleontology of the upper Pleistocene deposits at Sankaty Head, Nantucket Island, Massachusetts, Geology, 10, 246–252, 1982.

1840 Ottos, E. G.: Beach ridges—definitions and significance, Geomorphology, 32, 83–108, 2000.

Pigati, J. S.: On correcting ^{14}C ages of gastropod shell carbonate for fractionation, Radiocarbon, 44, 755–760, 2002.

Prest, V. K., Terasmae, J., Matthews, J. V. J., and Lichti-Federovich, S.: Late-Quaternary History of Magdalen Islands, Quebec, Maritime Sediments, 12, 39–59, 1976.

1845 Radtke, U., Mangini, A. and Grün, R.: ESR dating of marine fossil shells. Nuclear Tracks and Radiation Measurements (1982), 10, 879–884, 1985.

Rampton, V. N.: Quaternary Geology of the Tuktoyaktuk Coastlands, Northwest Territories, Geological Survey of Canada Memoir 423, 98 p., 1988.

Ramsay, W.: Beiträge zur Geologie der recenten und pleistocänen Bildungen der Halbinsel Kanin. Bulletin de Société de Géographie de Finlande, Fennia 21, 66 p., 1904.

1850 Rasmussen, T. L., Thomsen, E., Kuijpers, A., and Wastegård, S.: Late warming and early cooling of the sea surface in the Nordic seas during MIS 5e (Eemian Interglacial), Quat. Sci. Rev., 22, 809–821, 2003.

1855 Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capone, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP), Radiocarbon, 62, 725–757, 2020.

Rémillard, A. M., St-Onge, G., Bernatchez, P., Hétu, B., Buylaert, J.-P., Murray, A. S., and Vigneault, B.: Chronology and stratigraphy of the Magdalen Islands archipelago from the last glaciation to the early Holocene: new insights into the glacial and sea-level history of eastern Canada, Boreas, 45, 604–628, 2016.

Rémillard, A. M., St-Onge, G., Bernatchez, P., Hétu, B., Buylaert, J.-P., Murray, A. S., and Lajeunesse, P.: Relative sea-level changes and glacio-isostatic adjustment on the Magdalen Islands archipelago (Atlantic Canada) from MIS 5 to the late Holocene, *Quat. Sci. Rev.*, 171, 216-233, 2017.

1865 Robertsson, A.-M., Svedlund, J.-O., Andrén, T., and Sundh, M.: Pleistocene stratigraphy in the Dellen region, central Sweden, *Boreas*, 26, 237-260, 1997.

Roemer, F.: Notizuber das Vorkommen von *Cardium edule* und *Buccinum (Nassa) reticulatum* im Diluvial-Kies bei Bromberg im Grossherzogthum Posen, *Zeitschrift für Deutschen Geologischen Gesellschaft*, 16, 611-614, 1864.

1870 Rosentau, A., Kleemann, V., Bennike, O., Steffen, H., Wehr, J., Latinović, M., Bagge, M., Ojala, A., Berglund, M., Becher, G.P. and Schoning, K.: A Holocene relative sea-level database for the Baltic Sea. *Quaternary Science Reviews*, 266, p.107071, 2021.

Rovere, A., Raymo, M.E., Vacchi, M., Lorscheid, T., Stocchi, P., Gomez-Pujol, L., Harris, D.L., Casella, E., O'Leary, M.J. and Hearty, P.J.: The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world. *Earth-Sci. Rev.*, 159, 404-427, 2016.

1875 Rovere, A., Ryan, D., Murray-Wallace, C., Simms, A., Vacchi, M., Dutton, A., Lorscheid, T., Chutcharavan, P., Brill, D., Bartz, M., Jankowski, N., Mueller, D., Cohen, K., and Gowan, E.: Descriptions of database fields for the World Atlas of Last Interglacial Shorelines (WALIS) (Version 1.0), Zenodo, <https://doi.org/10.5281/zenodo.3961544>, 2020

Saarnisto, M., Eriksson, B. and Hirvas, H.: Tepsankumpu revisited – pollen evidence of stable Eemian climates in Finnish Lapland. *Boreas* 28, 12–22, 1999.

1880 Salvigsen, O.: Radiocarbon Dated Raised Beaches in Kong Karls Land, Svalbard, and their Consequences for the Glacial History of the Barents Sea Area, *Geografiska Annaler: Series A, Physical Geography*, 63, 283-291, 1981.

Salvigsen, O.: Occurrence of pumice on raised beaches and Holocene shoreline displacement in the inner Isfjorden area, Svalbard, *Polar Res.*, 2, 107-113, 1984.

1885 Sánchez Goñi, M. F., Bakker, P., Desprat, S., Carlson, A. E., Van Meerbeeck, C. J., Peyron, O., Naughton, F., Fletcher, W. J., Eynaud, F., Rossignol, L., and Renssen, H.: European climate optimum and enhanced Greenland melt during the Last Interglacial, *Geology*, 40, 627-630, 2012.

Schellmann, G. and Radtke, U.: Problems encountered in the determination of dose and dose rate in ESR dating of mollusc shells, *Quat. Sci. Revs.*, 18, pp.1515-1527, 1999.

1890 Sejrup, H.-P.: Molluscan and foraminiferal biostratigraphy of an Eemian-Early Weichselian section on Karmøy, southwestern Norway. *Boreas* 16, 27-42, 1987.

Sejrup, H.-P.: Amino acid geochronology. In S. Funder, (Ed.), *Late Quaternary stratigraphy and glaciology in the Thule area. Meddeleser om Grønland*, Geoscience, 22, 36-39, 1990.

Semevskij, D. V.: Pleistocene deposits in Spitsbergen [In Russian, Internal English translation, Norsk Polarinstitutt], 1967.

1895 Skinner, R. G.: Quaternary stratigraphy of the Moose River Basin, Ontario, *Geological Survey of Canada Bulletin*, 225, 1-77, 1973.

Slater, T., Lawrence, I. R., Otosaka, I. N., Shepherd, A., Gourmelen, N., Jakob, L., Tepes, P., Gilbert, L., and Nienow, P.: Review article: Earth's ice imbalance, *Cryosphere*, 15, 233-246, 2021.

Šnejder, G. V.: Stratigraphy of Cenozoic deposits of some topographic features of the Taymyr Peninsula, north-eastern extremity, *PGO Sevmorgeologija* 1989, Leningrad, 35-48, 1989.

1900 Stuiver, M. and Polach, H. A.: Discussion: reporting of ^{14}C data, *Radiocarbon*, 19, 355-363, 1977.

Sukhorukova, S. S.: The key sequence of marine sediments and climate fluctuations of the Kazantsevo (Eemian) interglacial (northern Siberia), *Geologia i geofizika*, 39, 74-84, 1998.

1905 Troitsky, L., Punning, J.-M., Hütt, G., and Rajamä, R.: Pleistocene glaciation chronology of Spitsbergen, *Boreas*, 8, 401-407, 1979.

Troitsky, S.L. and Kulakov, A.P.: Sea level oscillations and topography of the coastlands. In: Timofeyev, D.A., ed. *Problemy ekzogenного рельефообразования* 1, Nauka, Moscow, 351-426 [in Russian], 1976.

1910 Tveranger, J., Houmark-Nielsen, M., Løvberg, K., and Mangerud, J.: Eemian-Weichselian stratigraphy of the Flakkerhuk ridge, southern Jameson Land, East Greenland, *Boreas*, 23, 359-384, 1994.

[Tveranger, J., Astakhov, V., Mangerud, J., and Svendsen, J. I.: Signature of the last shelf-centered glaciation at a key section in the Pechora basin, Arctic Russia, *J. Quat. Sci.*, 13, 189-203, 1998.](#)

1915 Vasskog, K., Svendsen, J.-I., Mangerud, J., Agasøster Haaga, K., Svean, A. and Lunnan, E.M.: Evidence of early deglaciation (18 000 cal a bp) and a postglacial relative sea-level curve from southern Karmøy, south-west Norway, *J. Quat. Sci.*, 34, 410-423, 2019.

Verrill, A. E.: On the post-Pliocene fossils of Sankaty Head, Nantucket Island, with a note on the geology by S. H. Scudder, *American Journal of Science*, 10, 364-375, 1875.

[Vincent, J-S.: The Quaternary history of Banks Island, N.W.T., Canada, *Géogr. Phys. Quat.*, 36, 209-232, 1982.](#)

1920 [Vincent, J-S.: La géologie du Quaternaire et la geomorphologie de Pile Banks, Arctique Canadien, Commission géologique du Canada, Mémoire 405 \[in French\]](#), 1983.

Volkova, V. S: Sediments of Boreal transgression on Bolshaya and Malaya Kheta rivers (left tributaries of the Yenissei). *Doklady pervichnoi organizatsii naychno-tehnicheskogo Gornogo obshchestva* 1,43-49 [in Russian], 1958.

1925 Vosgerau, H., Funder, S., Kelly, M., Knudsen, K. L., Kronborg, C., Madsen, H. B. and Sejrup, H. P.: Palaeoenvironment and changes in relative sea level during the last interglaciation at Langelandselv. Jameson Land. East Greenland, *Boreas* 23, 398-411, 1994.

Wehmiller, J. F., Belknap, D. F., Boutin, B. S., Mirecki, J. E., Rahaim, S. D., York, L. L., and Easterbrook, D. J.: A review of the aminostratigraphy of Quaternary mollusks from United States Atlantic Coastal Plain sites. In: *Dating Quaternary Sediments*, Geological Society of America, 1988.

1930 Wehmiller, J. F. and Pellerito, V.: An evolving database for Quaternary aminostratigraphy, *GeoResJ*, 6, 115-123, 2015.

[Willerslev, E., Davison, J., Moora, M., Zobel, M., Coissac, E., Edward, M. E., Lorenzen, E. D., Vestergård, M., Gussarova, G., Haile, J., Craine, J., Gielly, L., Boessenkool, S., Epp, S. L., Pearman, P. B., Cheddadi, R., Murray, D., Bräthen, K. A.,](#)

Zagwijn, W. H.: An analysis of Eemian climate in western and central Europe, Quat. Sci Revs. 15, 451-469, 1996.

1935 Zawidzka, K.: Badania petrograficzno-litologiczne osadów czwartorzędowych, Szczegolowa. Szczegolowa Mapa Geologiczna Polski w skali 1:50,000. CAG PIG, Warszawa, 1997.

FIGURES

1940




Figure 1. Location of Last Interglacial (Last interglacial (LIG)4G) marine sites in the formerly glaciated Northern Hemisphere, along with the extent of MIS 6 and Last Glacial Maximum ice sheets (Batchelor et al., 2019). Inset map shows extent of the LIG White Sea that inundated the isostatically depressed landscape of western Russia and northwestern Europe.

1945




Figure 2. The timescales covered in this [manuscript](#), [paper](#), along with regional nomenclature. The benthic $\delta^{18}\text{O}$ curve is the LR04 stack, from which the Marine Isotope Stages are derived (Lisiecki and Raymo, 2005).

Formatted: Superscript




Figure 3. The location of LIG marine sites in the Taimyr Peninsula and Western Siberian Plain, Russia. Panel A situates the sites in the Northern Hemisphere, whereas Panel B provides a regional map. Sites plotted on this map are detailed in Sections 6.1–6.11. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al., (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.




Figure 34. An overview of the best dated sections of the last interglacial (LIG) in the Russian Arctic mainland. Numbers along the columns are geochronometric ages in kan-kilo years. Relative positions of samples are shown by black dots: large ones – dated by optical luminescence, smaller ones – by radiocarbon; black triangles indicate dates by uranium-seriesU/Th and electron-spin-resonanceESR methods. Sources of information: Sula (Murray et al., 2007), More-Yu (Astakhov and Svendsen, 2002), Nadym Ob (Astakhov et al., 2004, 2005), Observations Cape (Astakhov and Nazarov, 2010a, b), Bol. Kheta (Nazarov et al., 2020), Karginsky Cape (Astakhov, 2013; Nazarov et al., 2020). These stratigraphic sections are sometimes summaries of several sites; see text for details. Comprehensive documentation of each site (including detailed sea level measurements and chronological data) is available at <https://doi.org/10.5281/zenodo.5602212> (Dalton et al., 2021).




Figure 5. The location of LIG marine sites in the Arkhangelsk district and Pechora Lowland, Russia. Panel A situates the sites in the Northern Hemisphere, whereas Panels B and C provide regional maps. Sites plotted on this map are detailed in Sections 6.12–6.15. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al., (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.





Figure 6. The location of LIG marine sites in western Russia, Estonia, Poland, Finland, Sweden and Norway. Panel A situates the sites in the Northern Hemisphere, whereas Panel B provides a regional map. Sites plotted on this map are detailed in Sections 6.16–6.33. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al., (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.



1985 **Figure 47.** Relative sea-level curve for Fjøsanger during the entire (Eemian \approx) MIS 5eLIG, shown as a dotted uncertainty band. The dashed part of the curve is considered most uncertain. The line at 25 m.a.s.l.MASL shows the threshold in the Bergen Valley; when sea level was higher than this threshold the fjord turned into a sound that was open in both ends. The dotted line at 15 m.a.s.l.MASL shows the top of the marine MIS 5eLIG marine sediments and thus the minimum altitude of the relative sea-levelRSL during the entire MIS 5eLIG. Modified from Mangerud et al., (1981).

1990




Figure 8. The locations of LIG marine sites in Svalbard, Norway. Panel A situates the sites in the Northern Hemisphere, whereas Panel B provides a regional map. Sites plotted on this map are detailed in Sections 6.34–6.39. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al., (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.





Figure 59. Photograph of Section II at Kapp Ekholm, Svalbard. Panel A is original photograph, and panel B shows stratigraphic units. Solid white line indicates a confirmed stratigraphic contact; dashed line indicates inferred stratigraphic contact (largely due to sediment slumping). The LIG MIS 5 sediments (indicated by the white text arrowbox) are about 10 m in thickness. Most *Mytilus edulis* were found in the lower part of the marine diamicton. Photo J. Mangerud 1988.



Figure 10. Preserved *Mytilus edulis* in the [LIG](#) sediments at Kapp Ekholm, Svalbard. This mollusc requires warmer water than [the conditions](#) around Svalbard for the last millennia, until it immigrated in 2004. Photo J. Mangerud 1988.



2010
Figure 711. Composite log of the Skilvika exposure with luminescence (TL and OSL in ka) and radiocarbon ages (in cal. ka BP for ages > 40 ka BP, un-calibrated for younger ages). Formations (Fm) 3 and 4 are the candidates for MIS-5e/the LIG. [From Alexanderson and Landvik \(2018\)](#) ([from Alexanderson and Landvik, 2018](#)).



2015

Figure 812. The upper half of the photo shows the extremely coarse foreset in Formation 4 at Skilvika. Note how the foresets interfinger with the sand and gravel beds in Formation 3. Photo J. Mangerud 1984.





Figure 13. The locations of LIG marine sites in Iceland and Greenland. Panel A situates the sites in the Northern Hemisphere, whereas Panels B, C and D provide regional maps. Sites plotted on this map are detailed in Sections 6.40–6.42. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al., (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.

2020

2025




Figure 14. The locations of LIG marine sites in North America. Panel A situates the sites in the Northern Hemisphere, whereas Panels B, C and D provide regional maps. Sites plotted on this map are detailed in Sections 6.43–6.48. Extent of MIS 6 and Last Glacial Maximum ice sheets after Batchelor et al. (2019). Scale calculated at centre of mapped area. Base layer: ArcGIS World Imagery.




Figure 9.15. Schematic diagram showing each site with quality scores. Summary of quality scores for the RSL data (horizontal axis) (upper panel) and chronology data (vertical axis). (lower panel).

TABLES

2040

Table 1: Measurement techniques used to establish the elevation of MIS-5eLIG marine deposits (Rovere et al., 2020). GPS= Global positioning system.

Formatted: Font: Not Bold

Measurement technique	Description	Typical accuracy
Barometric altimeter	Difference in barometric pressure between a point of known elevation (often sea level) and a point of unknown elevation. Not accurate and used only rarely in sea-level studies	Up to $\pm 20\%$ of elevation measurement
Cross-section from publication	The elevation was extracted from a published sketch/topographic section.	Variable, depending on the scale of the sketch or topographic section.
Differential GPS	GPS positions acquired in the field and corrected either in real time or during post-processing with respect to the known position of a base station or a geostationary satellite system (e.g. Omnistar). Accuracy depends on satellite signal strength, distance from base station, and number of static positions acquired at the same location.	$\pm 0.02/\pm 0.08$ m, depending on survey conditions and instruments used (e.g., single-band vs dual-band receivers)
Distance from top of drill core	Distance from the top of drill core.	Depending on coring technique and sampling procedures
Handheld GPS	Commercial hand-held GPS	Dependent on model and satellite coverage but could be as low as 1-2 m.
Inclinometer	Elevation measured with inclinometer starting from a point of known altitude.	Variable depending on the distance between reference and measured point.
Metered tape or rod	The end of a tape or rod is placed at a known elevation point, and the elevation of the unknown point is calculated using the metered scale and, if necessary, clinometers to calculate angles.	Up to $\pm 10\%$ of elevation measurement
Not reported	The elevation measurement technique was not reported, most probably hand level or metered tape.	20% of the original elevation reported added in root mean square to the sea level datum error
Theodolite and rod	Elevation derived from triangulation with a theodolite.	Usually very precise, centimetric accuracy, depending on distance.
Topographic map and digital elevation models	Elevation derived from the contour lines on topographic maps. Most often used for large-scale landforms (i.e. marine terraces). Several meters of error are possible, depending on the scale of the map or the resolution of the DEM	Variable with scale of map and technique used to derive DEM.

2045

Table 2. Sea level datums reviewed in this study (Rovere et al., 2020).

Datum name	Datum description
Mean Sea Level / General definition	General definition of MSL, with no indications on the datum to which it is referred to. A datum uncertainty can be established on a case-by-case basis.

Table 3. Quality scores for sea level proxies.

Description	Quality rating
Sedimentary sequence confidently shows a regression from deep (> 10 m) marine to terrestrial conditions, or the deposit provides, via age constraints, a limiting elevation for a long portion of the interglacial. Elevation measurements are well documented to be within 3 m.	5 (excellent)
Same as above, but details of elevation measurements are not well documented.	4 (good)
Sedimentary sequence is regressive, but it is less certain that sea level fell below the elevation of the top of the sequence (e.g. beach sediments).	3 (average)
Sedimentary sequence is marine limiting, but there is evidence that sea level was within 10 m of the top of the sequence (e.g. shoreface and wave influenced littoral sediments).	2 (poor)
Sedimentary sequence is marine limiting, but there is evidence of sea level fluctuations.	1 (very poor)
The sequence is marine or terrestrial limiting, and there is no information on potential sea level position.	0 (minimal)

Formatted: Font: Not Bold

Table 4. Quality scores for age, from the WALIS documentation (Rovere et al., 2020)

Description	Quality rating
Very narrow age range, e.g. a few thousand ² s of years, that allow the attribution to a specific timing within the LIG (MIS 5e) (e.g. 117 ± 2 ka)	5 (excellent)
Narrow age range, confidently dated to the LIG allowing the attribution to a specific substage of MIS 5 (e.g., MIS 5e)	4 (good)
The RSL data point can be attributed only to a generic interglacial (e.g. MIS 5)	3 (average)
Only partial information or minimum age constraints are available	2 (poor)
Different age constraints point to different interglacials	1 (very poor)
Not enough information to attribute the RSL data point to any Pleistocene interglacial.	0 (rejected)

Formatted: Font: Not Bold

2050

2055

Table 5. Summary of LIG sites from the glaciated Northern Hemisphere that contain RSL indicators. Further details are available in Dalton et al., (2021). AAR = Amino Acid Racemization; Lum.= Luminescence; ESR = Electron Spin Resonance; ChrStrat.= Chronostratigraphy. Comprehensive documentation of each site (including detailed sea level measurements and chronological data) is available at <https://doi.org/10.5281/zenodo.5602212> (Dalton et al., 2021).

Formatted: Font: Not Bold

Formatted: Font: 9 pt, Font color: Black, English (United States)

Section in manuscript	WALIS RSL ID	Site name	Subsite	Country ^{Nat} ion	Lat. (DD)	Long. (DD)	RSL indicator elevation (m)	RSL indicator elevation error (m)	Age attribution	Quality of RSL data	Quality of age data
6.1	4123	Novorybnoye 2	Unit F	Russia	72.83	105.79	24	2	Lum.; ESR	2	4
6.2	4124	Bol'shaya Balakhnya River (BBR 17)	Unit A	Russia	73.62	105.36	13	2	ESR; ChrStrat.	0	3
6.3	4091	Kamennaya River		Russia	76.53	103.52	132.6	10	ChrStrat.	4	4
6.4	4088	Kratnaya River	KR1	Russia	77.51	103.21	43.3	2	Lum.; ESR; Other dating	1	3
6.4	4089	Kratnaya River	KR2	Russia	77.51	103.20	39.1	2	Lum.; ESR; Other dating	1	3
6.4	4090	Kratnaya River	KR3	Russia	77.50	103.20	36.2	2	Lum.; ESR; Other dating	1	3
6.5	4085	Anjeliko River	AR3	Russia	77.35	102.73	58.5	2	Lum.; ESR; Other dating	3	3
6.5	4086	Anjeliko River	AR4	Russia	77.36	102.68	59.2	2	Lum.; ESR; Other dating	3	3
6.5	4087	Anjeliko River	Bolotniy River BR1	Russia	77.39	102.66	48.8	2	Lum.; ESR; Other dating	4	3
6.6	4000	October Revolution Island	Ozemaya River, highest beach ridge	Russia	79.12	96.92	140	5	AAR; Lum.; ESR; Other dating	4	3
6.7	4139	Lower Agapa River		Russia	71.60	88.30	63	5	ChrStrat.	1	4
6.8	4140	Karginsky Cape		Russia	69.95	83.57	21	5	Lum.; ESR; ChrStrat.	0	4
6.9.1	4147	Tanama	Tanama 1	Russia	70.24	79.76	65	5	Lum.; ChrStrat.	3	4
6.9.2	4148	Tanama	Tanama 2	Russia	69.83	79.00	65	5	Lum.; ChrStrat.	3	4
6.10.1	4255	Bol'shaya Kheta	Site 7251	Russia	68.47	83.12	30	0.5	Lum.; ChrStrat.	2	4
6.10.2	4256	Bol'shaya Kheta	Site 7248	Russia	67.97	83.10	30	0.5	Lum.; ChrStrat.	2	4
6.10.3	4257	Bol'shaya Kheta	Site 7249	Russia	68.00	83.13	30	0.5	Lum.; ChrStrat.	2	4
6.10.4	4258	Bol'shaya Kheta	Site 7246	Russia	67.96	83.21	30	0.5	Lum.; ChrStrat.	2	4
6.11	4151	Observation Cape		Russia	68.97	76.10	35	5	Lum.	2	3
6.12	4152	Sula	Sula 21/22	Russia	67.00	50.34	50	5	Lum.; ChrStrat.	4	4
6.13	4153	River Yangarei	Yangarei-1	Russia	68.70	61.83	70.5	5	Lum.; ChrStrat.	0	4
6.14	4154	Vorga-Yol		Russia	66.70	56.75	91	5	Lum.	2	4
6.15.1	4188	Pyoza River	Zaton site	Russia	65.58	44.63	10	2	AAR; ESR; ChrStrat.	4	4
6.15.2	4189	Pyoza River	Bychye site	Russia	65.79	45.06	23	4.6	ChrStrat.	2	4
6.15.3	4190	Pyoza River	Viryuga W.	Russia	65.82	46.00	49	9.8	ChrStrat.	0	3
6.15.4	4191	Pyoza River	Viryuga E.	Russia	65.82	46.00	63	10	ChrStrat.	2	4
6.15.5	4192	Pyoza River	Kalinov	Russia	65.79	46.22	37	7.4	ChrStrat.	3	4

6.15.6	4193	Pyoza River	Yatsevets	Russia	65.70	46.52	38	7.6	ChrStrat.	1	4
6.15.7	4194	Pyoza River	Site 11 Orlorets	Russia	65.71	46.84	43.5	8.7	ChrStrat.	3	4
6.15.8	4195	Pyoza River	Site 12 Orlorets	Russia	65.69	46.93	43.5	8.7	ChrStrat.	3	4
6.15.9	4196	Pyoza River	Site 13 Yolkino	Russia	65.68	47.60	51	10	ChrStrat.	2	4
6.15.10	4197	Pyoza River	Site 14 Yolkino	Russia	65.68	47.60	51	10	Lum.; ChrStrat.	2	4
6.15.11	4198	Pyoza River	Burdui	Russia	65.67	48.06	60	10	ChrStrat.	2	4
6.16	4155	Ponoi River	Unit 2	Russia	67.08	41.13	11.5	2.3	Lum.; ESR; ChrStrat.	2	4
6.17	4156	Svyatoi Nos		Russia	68.02	39.87	16	3.2	Lum.; ChrStrat.	0	4
6.18	4157	Chapoma		Russia	66.11	38.97	10	2	ESR; ChrStrat.	4	4
6.19	4158	Strelna River		Russia	66.010	383.52 64	35.5	7.1	ESR; ChrStrat.	2	4
6.2	4159	Varzuga	S1	Russia	66.40	36.64	14	2.8	ChrStrat.	0	4
6.21	4160	Petrozavodsk		Russia	61.77	34.40	40	8	ChrStrat.	1	4
6.22	3712	Peski		Russia	60.15	29.29	13.5	8	ChrStrat.	4	4
6.23	3711	Pöhja-Uhtju		Estonia	59.68	26.51	-49	1	ChrStrat.	2	4
6.24	3985	Suur-Prangli		Estonia	59.62	25.01	-61	1	ChrStrat.	2	4
6.25.1	3987	Lower Vistula Region	Obrzynowo	Poland	53.78	19.27	-3.5	10	ChrStrat.	4	4
6.25.2	3986	Lower Vistula Region	Licze	Poland	53.75	19.13	-8	10	ChrStrat.	4	4
6.26.1	3990	Rewal coastline	Rewal borehole	Poland	54.09	15.03	-5.5	0.4	ChrStrat.	3	4
6.26.2	3991	Rewal coastline	Ciećmierz borehole	Poland	53.99	15.03	-6.5	4	ChrStrat.	3	4
6.26.3	3989	Rewal coastline	Sliwin borehole	Poland	54.08	15.01	6.3	0.8	ChrStrat.	3	4
6.27	4161	Ollala	Borehole F	Finland	64.18	25.35	116.5	1	ChrStrat.	4	4
6.28	4162	Ukonkangas		Finland	63.91	25.85	105.7	1	ChrStrat.	5	4
6.29	4163	Viitala		Finland	62.60	23.00	84.5	1	ChrStrat.	2	4
6.3	4164	Mertuanaja		Finland	34.11	24.59	60	1	Lum.; ChrStrat.	4	4
6.31	3988	Norra Sannäs		Sweden	61.78	16.69	27.65	9	ChrStrat.; Other dating	4	4
6.32	3708	Fjøsanger		Norway	60.34	5.33	15	0.1	AAR; Lum.	5	4
6.33	3709	Bø		Norway	59.36	5.28	-1	0.1	AAR; ChrStrat.	5	4
6.34	3833	Hidalen	Unit D	Norway (Svalbard) and Jan Mayen	78.90	28.13	83	2	AAR	3	2
6.34	3834	Hidalen	Unit B	Norway (Svalbard) Svalbard and Jan Mayen	78.90	28.13	64	2	AAR; Lum.	2	2
6.35	3787	Kapp Ekholm	Formation B	Norway (Svalbard) Svalbard and Jan Mayen	78.55	16.55	22	0.1	AAR; Lum.; ChrStrat.	3	4
6.36	3810	Skilvika	Formation 3	Norway (Svalbard) Svalbard and Jan Mayen	77.57	14.44	28	0.5	Lum.	3	3
6.37	3799	Kongsfjordhallet		Norway (Svalbard) Svalbard and Jan Mayen	79.03	11.88	34	1	Lum.	3	3

6.38	3809	Poolepynten	Unit A1	Norway (Svalbard)& valbard and Jan Mayen	78.45	11.66	5	2	AAR; Lum.; Other dating	3	4
6.39	3788	Leinstranda		Norway (Svalbard)& valbard and Jan Mayen	78.88	11.56	19.2	0.5	Lum.; ChrStrat.	3	4
6.4	3710	Galtalækur		Iceland	63.99	-19.96	120	10	ChrStrat.; Other dating	2	3
6.41.1	4168	Kikiakajik	Beach	Greenland	70.04	-22.25	9.75	3.4	Lum.; ChrStrat.; Other dating	3	4
6.41.2	4166	Kap Hope	Regressive sequence	Greenland	70.46	-22.32	8	2.23	Lum.; ChrStrat.; Other dating	5	4
6.41.2	4167	Kap Hope	Transgressive sequence	Greenland	70.46	-22.32	15	1	Lum.; ChrStrat.; Other dating	3	4
6.41.3	4169	Kap Stewart		Greenland	70.44	-22.78	40	8	Lum.; ChrStrat.	4	4
6.41.4	4170	Hesteevl		Greenland	70.44	-23.10	35	7	Lum.; ChrStrat.	3	4
6.41.5	4172	Fynselv	443d	Greenland	70.46	-23.32	21.5	4.3	Lum.; ChrStrat.	3	4
6.41.6	4254	Langelandselv	composite section	Greenland	70.55	-23.64	70	10	AAR; Lum.; ChrStrat.	4	4
6.41.7	4181	Aucellaev River	Location 72	Greenland	70.59	-23.76	16	3.2	AAR; Lum.; ChrStrat.	1	4
6.41.8	4182	Lollandselv- Falsterselv		Greenland	70.87	-24.18	20	4	ChrStrat.	4	2
6.42.1	4199	Thule	Iterlak K	Greenland	76.71	-69.41	27.5	5.5	AAR; ChrStrat.; Other dating	2	4
6.42.2	4200	Thule	Iterlak L	Greenland	76.71	-69.42	14	2.8	Lum.; ChrStrat.	2	4
6.42.3	4201	Thule	Saunders Ø B	Greenland	76.60	-69.74	20	4	Lum.; ChrStrat.	3	4
6.42.4	4202	Thule	Saunders Ø C	Greenland	76.60	-69.74	12	2.4	Lum.; ChrStrat.	3	4
6.42.5	4203	Thule	Narsaarsuk D	Greenland	76.45	-69.29	6	1.2	AAR; ChrStrat.	2	4
6.42.6	4204	Thule	Narsaarsuk E	Greenland	76.45	-69.29	6	1.2	AAR; Lum.; ChrStrat.	2	4
6.42.7	4205	Thule	Narsaarsuk F+G	Greenland	76.45	-69.29	9	1.8	AAR; Lum.; ChrStrat.	2	4
6.43.1	3661	Iles de la Madeleine	Camping	Canada	47.35	-61.88	14	4	Lum.	2	3
6.43.2	3640	Iles de la Madeleine	Portage du Cap	Canada	47.25	-61.91	17	4.88	ChrStrat.; Other dating	3	2
6.43.3	4183	Iles de la Madeleine	Le Bassin	Canada	47.23	-61.90	2	1	ChrStrat.; Other dating	4	2
6.44.1	4185	Clyde Foreland	Profile 6	Canada	70.69	-68.95	28	5.6	ChrStrat.; Other dating	1	4
6.44.2	4186	Clyde Foreland	Profile 9	Canada	70.60	-68.41	11	5.6	U-Series; AAR; ChrStrat.	0	4
6.44.3	4187	Clyde Foreland	Profile 10	Canada	70.58	-68.35	2.5	0.5	AAR; ChrStrat.	1	4
6.45	3639	Ile aux Coudres	Site du Forage	Canada	47.41	-70.42	-2	0.4	ChrStrat.; Other dating	2	2

6.46	3680	Long Island	Bridgehampton - core S59793	United States of America	40.94	-72.31	-23	1.4	AAR; ChrStrat.	1	2
6.47	3637	Kwatabohegan River	Marine unit	Canada	51.14	-82.12	90	18	AAR; Other dating	0	2
6.48	4184	East of Nicholson Peninsula	VH-83-050	Canada	69.89	-128.52	2	0.4	AAR; ChrStrat.; Other dating	3	2