

# 1 Pre- and post-production processes increasingly dominate 2 greenhouse gas emissions from agri-food systems

3 Francesco N. Tubiello<sup>1</sup>, Kevin Karl<sup>1,2</sup>, Alessandro Flammini<sup>1,3</sup>, Johannes Gütschow<sup>4</sup>, Griffiths  
4 Obli-Laryea<sup>1</sup>, Giulia Conchedda<sup>1</sup>, Xueyao Pan<sup>1</sup>, Sally Yue Qi<sup>2</sup>, Hörn Halldórudóttir  
5 Heiðarsdóttir<sup>1</sup>, Nathan Wanner<sup>1</sup>, Roberta Quadrelli<sup>5</sup>, Leonardo Rocha Souza<sup>6</sup>, Philippe Benoit<sup>2</sup>,  
6 Matthew Hayek<sup>7</sup>, David Sandalow<sup>2</sup>, Erik Mencos-Contrera<sup>8,9</sup>, Cynthia Rosenzweig<sup>9,8</sup>, Jose'  
7 Rosero Moncayo<sup>1</sup>, Piero Conforti<sup>1</sup> and Maximo Torero<sup>1</sup>

8 <sup>1</sup> Food and Agriculture Organization, Rome, Italy

9 <sup>2</sup> Center on Global Energy Policy, Columbia University, New York, USA

10 <sup>3</sup> United Nations Industrial Development Organization, Department of Environment, Vienna, Austria

11 <sup>4</sup> Potsdam Institute for Global Climate Research, Potsdam, Germany

12 <sup>5</sup> International Energy Agency, Paris, France

13 <sup>6</sup> United Nations Statistics Division, New York, USA

14 <sup>7</sup> Department of Environmental Science, New York University, New York, USA

15 <sup>8</sup> Center of Global Climate Research, Columbia University, New York, USA

16 <sup>9</sup> NASA Goddard Institute for Space Studies, New York, USA

17 \*Corresponding author: Francesco N. Tubiello, [francesco.tubiello@fao.org](mailto:francesco.tubiello@fao.org)

18

19 **Abstract.** We present results from the FAOSTAT *Emissions shares* database, covering emissions from agri-food  
20 systems and their shares to total anthropogenic emissions for 196 countries and 40 territories, for the period 1990-  
21 2019. We find that in 2019, global agri-food systems emissions were 16.5 (95% CI range: 11-22) billion metric  
22 tonnes (Gt CO<sub>2eq</sub> yr<sup>-1</sup>), corresponding to 31% (range: 19-43%) of total anthropogenic emissions. Of the agri-food  
23 systems total, global emissions within the farm gate— from crop and livestock production processes including on-  
24 farm energy use— were 7.2 Gt CO<sub>2eq</sub> yr<sup>-1</sup>; emissions from land use change, due to deforestation and peatland  
25 degradation, were 3.5 Gt CO<sub>2eq</sub> yr<sup>-1</sup>; and emissions from pre- and post-production processes— manufacturing of  
26 fertilizers, food processing, packaging, transport, retail, household consumption and food waste disposal— were  
27 5.8 Gt CO<sub>2eq</sub> yr<sup>-1</sup>. Over the study period 1990-2019, agri-food systems emissions increased in total by 17%, largely  
28 driven by a doubling of emissions from pre- and post-production processes. Conversely, the FAOSTAT data show  
29 that since 1990 land use emissions decreased by 25%, while emissions within the farm gate increased 9%. In 2019,  
30 in terms of individual greenhouse gases (GHGs), pre- and post- production processes emitted the most CO<sub>2</sub> (3.9  
31 Gt CO<sub>2</sub> yr<sup>-1</sup>), preceding land use change (3.3 Gt CO<sub>2</sub> yr<sup>-1</sup>) and farm-gate (1.2 Gt CO<sub>2</sub> yr<sup>-1</sup>) emissions. Conversely,  
32 farm-gate activities were by far the major emitter of methane (140 Mt CH<sub>4</sub> yr<sup>-1</sup>) and of nitrous oxide (7.8 Mt N<sub>2</sub>O  
33 yr<sup>-1</sup>). Pre-and post-production processes were also significant emitters of methane (49 Mt CH<sub>4</sub> yr<sup>-1</sup>), mostly  
34 generated from the decay of solid food waste in landfills and open-dumps. One key trend over the 30-year period  
35 since 1990 highlighted by our analysis is the increasingly important role of food-related emissions generated  
36 outside of agricultural land, in pre- and post-production processes along the agri-food system, at global, regional

1 and national scales. In fact, our data show that by 2019, pre- and post-production processes had overtaken farm-  
2 gate processes to become the largest GHG component of agri-food systems emissions in Annex I parties (2.2 Gt  
3 CO<sub>2eq</sub> yr<sup>-1</sup>). They also more than doubled in non-Annex I parties (to 3.5 Gt CO<sub>2eq</sub> yr<sup>-1</sup>), becoming larger than  
4 emissions from land-use change. By 2019 food supply chains had become the largest agri-food system component  
5 in China (1100 Mt CO<sub>2eq</sub> yr<sup>-1</sup>); USA (700 Mt CO<sub>2eq</sub> yr<sup>-1</sup>) and EU-27 (600 Mt CO<sub>2eq</sub> yr<sup>-1</sup>). This has important  
6 repercussions for food-relevant national mitigation strategies, considering that until recently these have focused  
7 mainly on reductions of non-CO<sub>2</sub> gases within the farm gate and on CO<sub>2</sub> mitigation from land use change. The  
8 information used in this work is available as open data with DOI 10.5281/zenodo.5615082 (Tubiello et al., 2021d).  
9 It is also available to users via the FAOSTAT database (FAO, 2021a), with annual updates.

10 **Keywords:** Agri-food systems, GHG emissions, farm gate, land use change, supply chains

11

## 1 **1. Introduction**

2 Agriculture is a significant contributor to climate change as well as one of the economic sectors most at risk from  
3 it. Greenhouse gas (GHG) emissions generated within the farm gate by crop and livestock production and related  
4 land use change contribute about one-fifth to one-quarter of total emissions from all human activities, when  
5 measured in CO<sub>2</sub> equivalents (Mbow et al., 2019; Smith et al., 2014; Vermeulen et al., 2012). The impacts are even  
6 starker in terms of individual GHG emissions. Agriculture contribute nearly 50% of global anthropogenic methane  
7 (CH<sub>4</sub>) and 75% of the total nitrous oxide (N<sub>2</sub>O) emissions (FAO, 2021b; Gütschow et al., 2021; Sauniois, et al.,  
8 2020). Once pre- and post-production activities along agri-food systems supply chains are included, food and  
9 agriculture activities generate up to one-third of all anthropogenic emissions globally (Crippa et al., 2021a,b;  
10 Rosenzweig et al., 2020; Tubiello et al., 2021a). This larger food systems perspective expands the potential for  
11 designing GHG mitigation strategies across the entire food system, i.e., over and above the more traditional focus  
12 on agricultural production and land use management that is currently found within countries' Nationally  
13 Determined Contributions (FAO, 2019).

14 Significant progress has recently resulted in the development of novel databases with global coverage of country-  
15 level data on agri-food systems emissions (Crippa et al., 2021a,b; Tubiello et al., 2021a). Tubiello et al. (2021a),  
16 in particular, provided a mapping of emission categories of the Intergovernmental Panel on Climate Change  
17 (IPCC)—used by countries for reporting their national GHG inventories (NGHGI) to the United Nations  
18 Framework Convention on Climate Change (UNFCCC)—unto internationally accepted food and agriculture  
19 concepts that are more easily understood by farmers and planners in countries, including in Ministries of  
20 Agriculture. By providing a correspondence between IPCC and FAO terminology, we seek to help countries to  
21 more adequately capture important aspects of food and agriculture activities within existing climate reporting, so  
22 that they can better identify effective climate actions across their agri-food systems (Fig. 1, adapted from Tubiello  
23 et al., 2021a). Firstly, the correspondence mapping expands the IPCC “agriculture” definition to include, in  
24 addition to non-CO<sub>2</sub> emissions from the farm, also the CO<sub>2</sub> generated in drained peatlands on agricultural land  
25 (Conchedda and Tubiello, 2020; Drösler et al., 2014) and by energy use in farm operations (FAO, 2020b; Flammini  
26 et al., 2021; Sims and Flammini, 2014). Secondly, it usefully disaggregates the ‘Land Use, land use change and  
27 forestry’ (LULUCF) of IPCC (2003) by separating out the emissions directly linked to food and agriculture  
28 activities, such as those generated by deforestation (Curtis et al., 2020; Tubiello et al., 2021c) and peat fires  
29 (Prosperi et al., 2020), from carbon removals, which are largely associated to processes in managed forests rather  
30 than on agricultural land (Grassi et al., 2021).

31 We present herein and discuss results from the first agri-food systems emissions database in FAOSTAT. The new  
32 database covers, as in previous versions (Tubiello et al., 2013) agriculture production activities within the farm  
33 gate and associated land use and land use change emissions on agricultural land. Importantly, it also includes  
34 estimates of emissions from pre- and post-production processes along food supply chains, including: fertilizer  
35 manufacturing, energy use within the farm gate, food processing, domestic and international food transport, retail,  
36 packaging, household consumption and food systems waste disposal. The database provides emissions data for  
37 four main GHG gases/categories (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and fluorinated gases) and their combined CO<sub>2</sub>eq levels. Data  
38 are available by country, over the period 1990-2019, as well as by regional and other relevant aggregations.

1 Importantly, data are provided in both IPCC and FAO classifications, facilitating the identification of national  
2 mitigation strategies across agri-food systems in countries, regionally and globally.

## 3 **2. Materials and methods**

4 Recent work (Rosenzweig et al., 2021; Tubiello et al., 2021a) helped to characterize agri-food systems emissions  
5 into three components: 1) Farm Gate; 2) Land Use Change; and 3) Pre- and Post-Production. Emissions estimates  
6 from the first two—generated by crop and livestock production activities within the farm gate and by the  
7 conversion of natural ecosystems to agriculture, such as deforestation and peatland degradation—are well  
8 established (IPCC, 2019). In particular, FAO disseminates annual updates in FAOSTAT (FAO, 2021; Tubiello,  
9 2019). This paper expands the available FAOSTAT data to include estimates of emissions from pre- and post-  
10 production processes, s., including energy use in fertilizer manufacturing; food processing; packaging; transport;  
11 retail; household consumption; and waste disposal.

### 12 **2.1 Mapping Agri-food Systems Components**

13 The new FAOSTAT data are provided, for each country, in both IPCC and FAO classifications. Specifically, on  
14 the one hand, data can be downloaded using IPCC emissions categories: *Energy; Industrial Processes and Product*  
15 *Use* (IPPU, henceforth referred to as Industry); *Waste; Agriculture; Land Use, Land Use Change and Forestry*  
16 *(LULUCF); and Other*. Both the total emissions from IPCC sectors are provided, as well as the portion directly  
17 related to agri-food systems. On the other hand, through the IPCC to FAO mapping discussed above and extending  
18 previous work (Tubiello, 2021a), data can also be downloaded in relevant FAO categories, covering emissions  
19 from: *Farm Gate, Land Use Change; and Pre- and Post-Production* processes (Fig. 1).

20 The FAOSTAT emissions estimates follow the IPCC (2006) “territorial approach,” i.e., they are assigned to the  
21 countries where they occur, independently of production or consumption considerations. For example, CO<sub>2</sub>  
22 emissions from energy use in fertilizers manufacturing are accounted for in the producing country, while the N<sub>2</sub>O  
23 emissions from fertilizer used on a country’s agricultural land for crop production are accounted for in that country.  
24 Similarly, emissions from energy use in agri-food systems activities are accounted for in countries where fuel  
25 combustion for that particular activity occurs, including electricity generation. The methods applied herein do not  
26 cover additional, upstream emissions associated with fuel supply chains, which are therefore not assigned to agri-  
27 food systems. More details on the scope of this work are found in section 2.3.

### 28 **2.2 Emissions Estimates**

29 FAO regularly disseminates emissions data for fifteen sub-domains in relation to the farm gate and land use change  
30 components of agri-food systems emissions, with published methodologies and results (i.e., Tubiello et al., 2021a).  
31 This manuscript relies in addition on new methods for computing emissions from pre- and post-production  
32 processes. Specifically, methods for emissions from energy use in fertilizers manufacturing, food processing, retail  
33 and household consumption, as well as refrigeration in retail are presented in Tubiello et al, (2021b), while Karl  
34 and Tubiello (2021 a,b) presented methods for estimating agri-food systems emissions in transport and waste  
35 disposal. Finally, emissions from on-farm energy use were developed by Flammini *et al.*, 2021). We refer the  
36 interested reader to those original publications for full details, while for completeness we also provide a sufficiently  
37 detailed summary of methods and coefficients as Supplementary Material of this manuscript.

1 More generally, a step-wise approach was followed for the estimation of agri-food systems emissions, as follows  
2 *Step 1*: identify, for each food systems component, the relevant international statistics needed to characterize  
3 country-level activity data (AD);  
4 *Step 2*: determine the food-related shares of the activity data ( $AD_{food}$ ) and assigns relevant GHG emission factors  
5 (EF) to each activity;  
6 *Step 3*: implement the generic IPCC method for estimating GHG emissions ( $E_{food}$ ), using inputs of activity data  
7 and emission factors from the first two steps, as follows:

$$8 \quad E_{food} = EF * AD_{food} \quad (1)$$

9 *Step 4*: Impute missing agri-food systems GHG emissions data by component. This step was limited to pre- and  
10 post-production processes, and applied where country-specific activity data were lacking. The imputation method  
11 used PRIMAP, a complete dataset of emissions estimates for all IPCC sectors, by country, covering the period  
12 1990-2019 (Gütschow et al., 2021). The PRIMAP dataset is already available in FAOSTAT for the computation  
13 of emissions shares of agriculture to the total anthropogenic total (FAO, 2019; Tubiello et al., 2021a). It compiles  
14 all available information on GHG emissions by country, including from official reporting. It was used  
15 internationally as the basis for an early, first-order estimate of agri-food systems shares in total GHG emissions  
16 (IPCC, 2019). Additionally, it was recently used in a UNFCCC Synthesis Report (UNFCCC, 2021) to assess world  
17 GHG emissions from all sectors in preparation of a stock take exercise that will be undertaken in 2022-2023 to  
18 assess countries' performance against their mitigation commitments under the Paris Agreement. The imputations  
19 in equation (1) were performed by applying to the PRIMAP sectoral emissions country-specific food system  
20 emissions shares (Tubiello et al., 2021b for more details).

## 21 **2.3 Global Warming Potentials Used**

22 The estimated emissions data expressed in CH<sub>4</sub> and N<sub>2</sub>O gases were converted to CO<sub>2</sub>-equivalents by using the  
23 100-year global warming potentials (GWP) of the IPCC (2014) Fifth Assessment Report, and specifically:

24  $GWP-CH_4 = 28$ ;  $GWP-N_2O = 265$ ;  $GWP-Fgases = 5195$ . The value for F-gases was obtained as an average of  
25 several distinct products (Tubiello et al., 2021b).

## 26 **2.3 Data uncertainty and limitations**

### 27 **2.3.1 Boundaries**

28 The processes covered herein do not span all processes attributable to agri-food systems. In particular, the scope  
29 of this work does not include, by design, upstream GHG emissions in the fuel chain, such as petroleum refining,  
30 as well as methane leaks during extraction processes and piping. These are expected to be not negligible if  
31 considered. While emissions from such sources can be estimated using a fixed fuel chain coefficient for certain  
32 fuel supply chains (see Crippa et al., 2021a), the authors do not consider such sources to be within scope of this  
33 work. GHG emissions attributable to electricity generation are included in the scope of this work, which itself  
34 excludes upstream GHG emissions in the fuel chain used to generate electricity (Flammini et al., 2021; Tubiello  
35 et al., 2021b).

1 Conversely, emissions of fluorinated gases (f-gases) from household refrigeration and from climate-controlled  
2 transportation were not included for lack of available country-level data for disaggregated cold chain elements.  
3 However, one estimate suggests that the majority (over 60%) of global food-related F-gas emissions occur in the  
4 retail stage, which is accounted for here in this work (International Institute of Refrigeration, 2021). Emissions  
5 from pesticide manufacturing were also not included due to the paucity of information and methodologies for their  
6 estimation at country level, in contrast to advanced work in fertilizers manufacturing (Brentrup et al., 2016;  
7 Brentrup et al., 2018; IFS, 2019). Bellardy et al. (2008) estimated global emissions from pesticides manufacturing  
8 to be roughly 72 (range: 3-140) Mt CO<sub>2</sub>eq yr<sup>-1</sup>, roughly 1-2% of the pre- and post-production total estimated in  
9 this work.

### 10 **2.3.2 Uncertainty**

11 Uncertainties in FAOSTAT farm gate and land use change emissions estimates have been characterized elsewhere,  
12 and computed in line with IPCC (2006) guidelines as ranging 30—70% across component processes. For the  
13 purpose of this analysis, we assigned uncertainties of 30% and 50% respectively to the farm gate and land use  
14 change components of the FAOSTAT agri-food systems emissions, in line with previous work (i.e., Tubiello et  
15 al., 2013; 2021b). The uncertainties in the estimates of pre- and post-production activities described herein are by  
16 contrast less documented. On the one hand, uncertainties in underlying energy activity data and emissions factors  
17 are typically lower than for the other two components, ranging 5-20% (Flammini et al., 2022). On the other, the  
18 relative novelty in estimating food systems shares for a range of activity data across many processes makes our  
19 estimates more uncertain, with heavy reliance on literature results from a subset of countries and regions that are  
20 necessarily extended to the rest of the world (Karl and Tubiello, 2021a). For this reason, we assigned an overall  
21 uncertainty of 30% to the pre- and post-production component. This is higher than the uncertainty of the underlying  
22 energy processes, but more in line with values used in recent work (Crippa et al., 2021a). As shown below,  
23 considering a roughly equal, one-third contribution of the three components and their assigned uncertainties, an  
24 overall uncertainty of 40% was estimated for the agri-food systems emissions totals, applicable to countries and  
25 regional aggregates.

26 The above uncertainties are meant only as first rough estimates, useful to determine tentative 95% confidence  
27 intervals for the overall agri-food system component of FAOSTAT emissions. Significantly more research is  
28 needed for further refinements in future studies, in particular on better characterizing sub-regional and regional  
29 activity data and emissions coefficients, given the diversity in agri-food system typology and their dependence on  
30 physical geography and national socio-economic drivers. These limitations nonetheless reflect the paucity of  
31 activity data available to describe agri-food systems components and their trends, globally and regionally. While  
32 knowledge and data exist for regions and countries such as the EU, USA China, and India, much remains to be  
33 done in terms of regional and country specific coverage.

### 34 **2.3.3 Areas for Advancement**

35 Work towards estimating agri-food systems emissions at the country level can be advanced in several ways. The  
36 present approach could be expanded on by including other country- and region-specific studies that estimate trends  
37 in energy consumption across a range of similar activities as proxies— whether or not they are distinctly related  
38 to food. Furthermore, other data sources could help explain and estimate variations in agri-food systems between  
39 countries, such as: GDP per capita, urbanization levels, proxies for infrastructure and industrial development, and

1 geographic and climate considerations. The development of a methodology to estimate emissions from pesticides  
2 could be explored, as it would help complement the understanding of emissions associated with chemical use in  
3 agriculture, in addition to fertilizers. Emissions from machinery manufacturing and from upstream GHG emissions  
4 in the fuel chain could also be added to further refine the analysis. This work could be further expanded by focusing  
5 on specific food commodities— requiring an additional focus on international trade and on supply and demand  
6 patterns (Dalin and Rodríguez-Iturbe, 2016). Such analysis would ultimately enable consumers to understand the  
7 full carbon footprint of particular commodities across global supply chains, which can facilitate GHG mitigation  
8 actions taken at the consumer level (Poore and Nemecek, 2018). Furthermore, it would be also useful to further  
9 investigate the increasing role of bioenergy and renewables as important mitigation opportunities in the food sector  
10 (Clark et al., 2020, JRC, 2015; Pablo-Romero et al., 2017; Wang, 2014).

## 11 **Data availability**

12 The GHG emission data presented herein cover the period 1990-2019, at the country level, with regional and global  
13 aggregates. They are available as open data, with DOI 10.5281/zenodo.5615082 (Tubiello et al., 2021d) and via  
14 the FAOSTAT emissions shares database (FAO, 2021a).

## 15 **3 Results**

### 16 **3.1 Global trends**

17 The FAOSTAT dataset considered in this study estimates in 2019 total anthropogenic emissions at 52 Gt CO<sub>2eq</sub>  
18 yr<sup>-1</sup> without land use, land use change and forestry emissions (LULUCF), and 54 Gt CO<sub>2eq</sub> yr<sup>-1</sup> with LULUCF—  
19 consistently with recent estimates (IPCC, 2019). We use the latter figure to compute emissions shares. In 2019  
20 world-total agri-food systems emissions, expressed in terms of 95% confidence intervals (CI) determined using an  
21 overall uncertainty of 40%, were 16.5 (CI range: 10-23) billion metric tonnes (Gt CO<sub>2eq</sub> yr<sup>-1</sup>), corresponding to  
22 31% (range: 19%-42%) of total anthropogenic emissions (Tab. 1). Of the food systems total, global emissions  
23 within the farm gate –from crop and livestock production processes including on-farm energy use—were 7.2  
24 (range: 5-9) Gt CO<sub>2eq</sub> yr<sup>-1</sup>; emissions from land use change, due to deforestation and peatland degradation, were  
25 3.5 (range: 2-5) Gt CO<sub>2eq</sub> yr<sup>-1</sup>; and emissions from pre- and post-production processes –manufacturing of fertilizers,  
26 food processing, packaging, transport, retail, household consumption and food waste disposal—were 5.8 (range:  
27 4-8) Gt CO<sub>2eq</sub> yr<sup>-1</sup>. Over the study period 1990-2019, agri-food systems emissions increased in total by 17%, though  
28 they have remained rather constant since about 2006 (Fig. 2). These trends were largely driven by a doubling of  
29 emissions from pre- and post-production processes, while land use emissions decreased by 25% and farm gate  
30 increased only 9%. In terms of single GHG, pre- and post- production processes emitted the most CO<sub>2</sub> (3.9 Gt CO<sub>2</sub>  
31 yr<sup>-1</sup>) in 2019, preceding land use change (3.3 Gt CO<sub>2</sub> yr<sup>-1</sup>) and farm-gate (1.2 Gt CO<sub>2</sub> yr<sup>-1</sup>) emissions. Conversely,  
32 farm-gate activities were by far the major emitter of methane (140 Mt CH<sub>4</sub> yr<sup>-1</sup>) and of nitrous oxide (7.8 Mt N<sub>2</sub>O  
33 yr<sup>-1</sup>). Pre-and post- processes were also significant emitters of methane (49 Mt CH<sub>4</sub> yr<sup>-1</sup>), mostly generated from  
34 the decay of solid food waste in landfills and open-dumps.

35 Emissions from within the farm gate and those due to related land use processes, including details of their sub-  
36 components, have been discussed in Tubiello et al. (2021a) and are regularly presented within FAOSTAT statistical  
37 briefs (e.g., FAO, 2020a; 2021b). Here we provide a detailed discussion of the components of agri-food systems  
38 emissions from pre- and post-production activities along supply chains and their relative contribution to the food

1 system totals (Fig. 3). Considering that the uncertainties used above are rough estimates, we will not report  
2 uncertainties in the following analysis. Our data show that in 2019 emissions from deforestation were the single  
3 largest emission component of agri-food systems, at 3.1 Gt CO<sub>2</sub> yr<sup>-1</sup>, having decreased 30% since 1990. The second  
4 most important component were non-CO<sub>2</sub> emissions from enteric fermentation (2.8 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), with increases  
5 of 13%. These were followed by emissions from livestock manure (1.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>) and several pre- and post-  
6 production emissions, including CO<sub>2</sub> from household consumption (1.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), CH<sub>4</sub> from food waste  
7 disposal (1.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), mostly CO<sub>2</sub> from fossil-fuel combustion for on-farm energy use (1.0 Gt CO<sub>2</sub>eq yr<sup>-1</sup>),  
8 and CO<sub>2</sub> and F-gases emissions from food retail (0.9 Gt CO<sub>2</sub>eq yr<sup>-1</sup>). Importantly, our data show that growth in  
9 pre- and post-production components was particularly strong, with emissions from retail increasing from 1990 to  
10 2019 by more than seven-fold, while emissions from household consumption more than doubled over the same  
11 period.

12 Finally, while emissions from agri-food systems increased globally by 16 percent between 1990 and 2019, their  
13 share in total emissions decreased, from 40 percent to 31 percent, as did the per capita emissions, from 2.7 to  
14 2.1 tonnes CO<sub>2</sub>eq per capita (Fig 2.)

### 15 **3.2 Regional Trends**

16 Our results indicate significant regional variation in terms of the composition of agri-food systems emissions by  
17 component (Fig. 4). Specifically, in terms of total agri-food systems emissions in 2019, Asia had the largest  
18 contribution, at 7 Gt CO<sub>2</sub>eq yr<sup>-1</sup>, followed by Africa (2.7 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), South America (2.4 Gt CO<sub>2</sub>eq yr<sup>-1</sup>) and  
19 Europe (2.1 Gt CO<sub>2</sub>eq yr<sup>-1</sup>). North America (1.5 Gt CO<sub>2</sub>eq yr<sup>-1</sup>) and Oceania (0.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>) were the smallest  
20 emitters among regions (Fig. 4). Focusing on GHG emissions beyond agricultural land, pre- and post-production  
21 emissions in 2019 were largest in Asia (2.9 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), followed by Europe and North America (0.8-1.1 Gt  
22 CO<sub>2</sub>eq yr<sup>-1</sup>). Regions also varied in terms of how agri-food systems components contributed to the total (Tab. 2).  
23 In 2019, pre- and post- production emissions were the largest food systems contributor in Europe (55%), North  
24 America (52%) and Asia (42%). Conversely, they were smallest in Oceania (23%), Africa (14%) and South  
25 America (12%). Additionally, the contribution of pre- and post-production processes along food supply chains  
26 significantly increased since 1990, when in no region they were the dominant emissions component. Since then,  
27 they doubled in all regions except in Africa—where it remained below 15%.

28 The data show which pre- and post-production process was most important by region (Tab. 2). In 2019, food  
29 household consumption was the dominant process outside of agricultural land emissions in Asia (0.9 Gt CO<sub>2</sub>eq yr<sup>-1</sup>)  
30 and Africa (0.2 Gt CO<sub>2</sub>eq yr<sup>-1</sup>). Conversely, Europe, Oceania and North America pre- and post-production  
31 processes were led by emissions from food retail (0.3-0.4 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), while South America was dominated by  
32 emissions from food waste disposal (0.2 Gt CO<sub>2</sub>eq yr<sup>-1</sup>).

### 33 **3.3 Country Trends**

34 Our estimates show a marked variation among countries in terms of total emissions as well as the composition of  
35 contributions across farm gate, land use change and pre- and post-processing components (Fig. 5). China had the  
36 most emissions (1.9 Gt CO<sub>2</sub>eq yr<sup>-1</sup>), followed by India, Brazil, Indonesia and the USA (1.2-1.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>).  
37 Democratic Republic of Congo (DRC) and Russian Federation followed with 0.5-0.6 Gt CO<sub>2</sub>eq yr<sup>-1</sup>, followed by  
38 Pakistan, Canada and Mexico with 0.2-0.3 Gt CO<sub>2</sub>eq yr<sup>-1</sup>. The contribution of the three main agri-food systems



1 components to the national total differed among countries significantly (Fig. 5). For instance, China and India had  
2 virtually no contribution from land use change to agri-food systems emissions. The land use contribution was also  
3 minor in the USA, Russian Federation and Pakistan. Conversely, the latter was the dominant emissions component  
4 in Brazil, Indonesia and the DRC. Additionally, the new database allowed for an in-depth analysis by country of  
5 pre- and post-production emissions along the agri-food chain, highlighting a significant variety in most relevant  
6 sub-process contribution (Tab. 3). For the year 2019, pre- and post-production emissions were dominated in China  
7 by food household consumption processes (463 Mt CO<sub>2</sub>eq yr<sup>-1</sup>), whereas food waste disposal was the dominant  
8 pathway in Brazil, Indonesia (77 Mt CO<sub>2</sub>eq yr<sup>-1</sup>), DRC (8 Mt CO<sub>2</sub>eq yr<sup>-1</sup>), Pakistan (33 Mt CO<sub>2</sub>eq yr<sup>-1</sup>) and Mexico,  
9 (56 Mt CO<sub>2</sub>eq yr<sup>-1</sup>). Emissions from food retail dominated the pre- and post-production component in the USA  
10 (292 Mt CO<sub>2</sub>eq yr<sup>-1</sup>), Russian Federation (177 Mt CO<sub>2</sub>eq yr<sup>-1</sup>) and Canada (20 Mt CO<sub>2</sub>eq yr<sup>-1</sup>). Finally, on-farm  
11 energy use was the largest pre- and post-production component in India (205 Mt CO<sub>2</sub>eq yr<sup>-1</sup>).

## 12 4 Discussion

### 13 4.1 Comparisons with previous work

14 The overall assessment of total agri-food systems emissions found in this work confirms recent previous findings  
15 by the IPCC (2019) and Crippa et al. (2021). With regards to pre- and post-production, the FAOSTAT estimates  
16 were consistent (Tab. 4) with previous findings (i.e., Crippa et al., 2021a, b; Vermuelen et al., 2012; Poore and  
17 Nemecek, 2018). In particular, emissions estimates for food transport, processing, waste and retail were consistent  
18 with EDGAR-FOOD (Karl and Tubiello, 2021b) and estimates for fertilizers manufacturing were in line with  
19 previous work by Vermeulen (2012). Conversely, FAOSTAT estimates were higher than EDGAR-FOOD for  
20 household consumption and lower for food packaging, the latter possibly linked to FAOSTAT estimates excluding  
21 indirect emissions from fuel supply chains, which were instead included in previously published estimates. Finally,  
22 our estimates of F-gas emissions from retail agreed well with those published in EDGAR-FOOD.

23 The most important disagreement with previous work was observed in relation to household consumption  
24 emissions. FAOSTAT estimates in this work, 1.2 Gt CO<sub>2</sub>eq, were nearly three times those of EDGAR-FOOD  
25 (with reference to 2015, the last year for which EDGAR data was available). While much more research is needed  
26 to refine estimates in this important agri-food systems component, our estimates were in fact well aligned with  
27 earlier FAO (2011) work (Figure 4), as well as more consistent with observed population growth, an important  
28 determinant of household consumption trends.

### 29 4.2 Trends

30 One notable trend over the 30-year period since 1990 is the increasingly important role of food-related emissions  
31 generated outside of agricultural land, in pre- and post-production processes along food supply chains, at all global,  
32 regional and national scales. Our data show that by 2019, pre- and post-production processes had overtaken farm-  
33 gate processes to become the largest GHG component of agri-food systems emissions in Annex I parties (2.2 Gt  
34 CO<sub>2</sub>eq yr<sup>-1</sup>). While farm gate emissions still dominated food-systems processes in non-Annex I parties, emissions  
35 from pre- and post-production were closing the gap in 2019, surpassing land use change, and having doubled since  
36 1990 to 3.5 Gt CO<sub>2</sub>eq yr<sup>-1</sup>. By 2019, pre- and post-production processes had become the largest agri-food system  
37 component in China (1.1 Gt CO<sub>2</sub>eq yr<sup>-1</sup>); USA (0.7 Gt CO<sub>2</sub>eq yr<sup>-1</sup>) and EU-27 (0.6 Gt CO<sub>2</sub>eq yr<sup>-1</sup>). This has important  
38 repercussions for food-relevant national mitigation strategies, such as those included in countries' NDCs,

1 considering that until recently these have focused mainly on reductions of non-CO<sub>2</sub> gases within the farm gate and  
2 on CO<sub>2</sub> mitigation from land use change (Hönle et al., 2019).

3 Importantly, the FAOSTAT database presented here allows for an estimation of the percentage share contribution  
4 of food systems emissions in total anthropogenic emissions, by country as well as at regional and global levels,  
5 over the period 1990-2019. A number of important issues can be highlighted to this end (Tab. 5 and Fig. 6). First,  
6 in terms of CO<sub>2</sub>eq, the share of world total agri-food systems emissions decreased from 40% in 1990 to 31% in  
7 2019. Thus while it is important to note that one-third of all GHG emissions today are generated by agri-food  
8 systems, their shares in total emissions may continue to decrease in the near future. This decreasing trend was  
9 driven by trends in large regions, consistently with transformations in their agri-food systems and land use change  
10 patterns. For instance, in South America, the region with the highest food systems share over the entire study  
11 period (Fig. 6), food shares decreased from 96% in 1990 to 72% in 2019. In Africa, from 67% to 57%, in Asia  
12 from 49% to 24% and in Oceania from 57% to 39%. In contrast to these trends, our data suggested that in regions  
13 dominated by modern agri-food systems, such as Europe and North America, the overall share of agri-food systems  
14 emissions in fact increased from 1990 to 2019, specifically from 24% to 31% in Europe and from 17% to 21% in  
15 North America. Such increases could be explained by increases in absolute emissions from pre- and post-  
16 production activities (Tab. 5), re-enforced by concomitant emissions decreases in non-food sector, especially  
17 energy systems (Lamb et al., 2022). The noted increase in absolute emissions from pre- and post-production  
18 activities was in fact present in all regions, leading to increases in the relative contributions to agri-food systems  
19 of this component, except for Africa.

20 An analysis on agri-food systems impacts on total GHG emissions would not be complete without a focus on  
21 component gases in addition to quantities expressed in CO<sub>2</sub>eq. The FAOSTAT data confirm the trends from 1990  
22 to 2019 seen for total CO<sub>2</sub>eq emissions, with important features (Tab. 6). First, the impact of agri-food systems on  
23 world total CO<sub>2</sub> emissions was 21% in 2019 (down from 31% in 1990), a respectable share considering the  
24 importance of carbon dioxide in any effective long-term mitigation strategy. While most regions had contributions  
25 around this value, ranging 13%-23% for North America, Oceania, Europe and Asia, the CO<sub>2</sub> contribution of agri-  
26 food systems was highest in Africa (52%) and South America (70%), largely in relation to land use change  
27 emissions, still significant therein. Europe and North America were the only regions where the CO<sub>2</sub> share of agri-  
28 food systems actually increased from 1990 to 2019, confirming the growing weight of pre- and post-production  
29 processes, which typically involve fossil-fuel energy use and thus emissions of CO<sub>2</sub> gas through combustion.

30 Second, the data highlight the significant contribution of agri-food systems to 2019 world total emissions of CH<sub>4</sub>  
31 (53%) and N<sub>2</sub>O (78%), also confirmed at regional levels (Tab. 6), linked to farm gate production processes  
32 (Tubiello, 2019).

33 Finally, the data highlight a very large increase in agri-food systems contributions of F-gas emissions, which went  
34 from near zero in 1990 to more than one-quarter of the world total in 2019 –with larger contributions in many  
35 regions. Such a marked increase is consistent with the growth in use of hydrofluorocarbons (HFCs) as refrigerants  
36 in the food retail and other sectors, following the banning of CFCs in 1990 (Hart et al., 2020; International Institute  
37 of Refrigeration, 2021; Tubiello et al., 2021b). Our findings are furthermore consistent with the strong growth in  
38 F-gas emissions reported in recent studies (Minx et al., 2021; Park et al., 2021).

1 An important aspect of the dataset presented in this study is its provision of information mapped across IPCC and  
2 FAO categories alike. Specific IPCC sectors include *Agriculture* and *Land use, land use change and forestry*  
3 (*LULUCF*). The IPCC further considers the *Agriculture, Forestry and Other Land Use (AFOLU)*. While countries  
4 report their agriculture and food emissions to the UNFCCC within National GHG Inventories, our findings  
5 highlight the importance to expand that reporting to a fuller agri-food systems view, one that properly weights the  
6 contribution of food to the global economy. Indeed, our results show that agri-food systems emissions in 2019  
7 were one-third of total anthropogenic emissions. This important picture does not emerge from NGHGI reporting  
8 aligned to IPCC categories, according to which for instance, *LULUCF* and *AFOLU* emissions contributed  
9 respectively 4% and 15% of the total.

## 10 **5 Conclusions**

11 This paper provided details of a new FAOSTAT database on GHG emissions along the entire agri-food systems  
12 chain, including crop and livestock production processes on the farm, land use change activities from the  
13 conversion of natural ecosystems to agricultural land, and processes along food supply chains, from input  
14 manufacturing to food processing, transport and retail, including household consumption and waste disposal.

15 The data are provided in open access mode to users worldwide and are available by country over the time period  
16 1990-2019, with plans for annual updates. The major trends identified in this work help locate GHG emissions  
17 hotspots in agri-food systems at the country, regional and global level. This can inform the process of designing  
18 effective mitigation actions in food and agriculture. This work adds to knowledge on GHG emissions from  
19 agriculture and land use—generally well established in the literature—by adding critical information on emissions  
20 from a range of pre- and post-production processes. The new data highlight the increasingly important role that  
21 pre- and post-production processes along supply chains play in the overall GHG footprint of agri-food systems,  
22 globally and in most countries, providing new insights into food and agriculture development trends and future  
23 mitigation options.

24 The granularity of the dataset allows, for the first time, to highlight specific processes of importance in specific  
25 countries or group of countries with similar characteristics. The relevance of the information being provided cuts  
26 across several national and international priorities, specifically those aiming at achieving more productive and  
27 sustainable food systems, including in relation to climate change. To this end, the work presented herein completes  
28 a mapping of IPCC categories, used by countries for reporting to the climate convention, to food and agriculture  
29 categories that are more readily understandable by farmers and ministries of agriculture in countries. This helps  
30 better identify agri-food systems entry points within existing and future national determined contributions. Finally,  
31 the methodological work underlying these efforts complements and extends recent pioneering efforts by FAO and  
32 other groups in characterizing technical coefficients to enable quantifying the weight of agri-food systems within  
33 countries' emissions profiles. The next steps in such efforts would need the involvement of interested national and  
34 international experts in compiling a first set of coefficients for agri-food systems as a practical 'agri-food systems  
35 annex' to the existing guidelines of the Intergovernmental Panel on Climate Change, providing guidance to  
36 countries on how to better characterize food and agriculture emissions within their national GHG inventories.

## 37 **6. Disclaimer**

1 The views expressed in this paper are the authors' only and do not necessarily reflect those of FAO, UNSD,  
2 UNIDO and IEA.

3

#### 4 **7. Acknowledgements**

5 FAOSTAT is supported by the FAO regular budget, funded by its member countries. We acknowledge the efforts  
6 of national experts who provide the statistics on food and agriculture as well as on energy use that are at the basis  
7 of this effort. All authors contributed critically to the drafts and gave final approval for the publication. We are  
8 grateful for overall support by the Food Climate Partnership at Columbia University.

9

## 1 6 References

- 2 Bellarby, J., Foereid, B., & Hastings, A. 2008. *Cool Farming: Climate Impacts of Agriculture and Mitigation*  
3 *Potential*.
- 4 Brentrup, F., Hoxha, A., and Christensen, B. 2016. Carbon footprint analysis of mineral fertilizer production in  
5 Europe and other world regions. In *10th International Conference on Life Cycle Assessment of Food*.
- 6 Brentrup, F., Lammel, J., Stephani, T., and Christensen, B. 2018. Updated carbon footprint values for mineral  
7 fertilizer from different world regions. In *11th International Conference on Life Cycle Assessment of Food*.  
8 <https://www.researchgate.net/publication/329774170> Updated carbon footprint values for mineral fertilizer f  
9 [rom different world regions](https://www.researchgate.net/publication/329774170)
- 10 Clark, M. A., Domingo, N. G. G., Colgan, K., Thakrar, S. K., Tilman, D., Lynch, J., Azevedo, I. L., & Hill, J. D.  
11 2020. Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets. *Science*,  
12 *370*(6517), 705–708. <https://doi.org/10.1126/science.aba7357>
- 13 Crippa, M., Solazzo, E., Guizzardi, D., Monforti-Ferrario, F., Tubiello, F. N., & Leip, A. 2021a. Food systems are  
14 responsible for a third of global anthropogenic GHG emissions. *Nature Food*, 1–12.  
15 <https://doi.org/10.1038/s43016-021-00225-9>
- 16 Crippa, M., Guizzardi, D., Schaaf, E., Solazzo, E., Muntean, M., Monforti-Ferrario, F., Olivier, J.G.J., Vignati, E.  
17 2021b. Emissions Database for Global Atmospheric Research (EDGAR) v6.0. Available from:  
18 [https://edgar.jrc.ec.europa.eu/index.php/dataset\\_ghg60](https://edgar.jrc.ec.europa.eu/index.php/dataset_ghg60)
- 19 Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A., & Hansen, M. C. 2018. Classifying drivers of global forest  
20 loss. *Science*, *361*(6407), 1108–1111. <https://doi.org/10.1126/science.aau3445>
- 21 Dalin, C., and Rodríguez-Iturbe, I. 2016. Environmental impacts of food trade via resource use and greenhouse  
22 gas emissions. *Environmental Research Letters*, *11*(3), 035012. <https://doi.org/10.1088/1748-9326/11/3/035012>
- 23 Drösler, M., Verchot, L. V., Freibauer, A., Pan, G., Evans, C. D., Borroniere, R. A., ... Couwenberg,  
24 J. 2014. Chapter 2: Drained inland organic soils. In T. Hiraishi, T. Krug, K. Tanabe, N. Srivastava, B.  
25 Jamsranjav, M. Fukuda, & T. Troxler (Eds.), *2013 Supplement to the 2006 IPCC Guidelines for National*  
26 *Greenhouse Gas Inventories: Wetlands* (pp. 1–79). IPCC: Switzerland.
- 27 Fang, X., Ravishankara, A. R., Velders, G. J. M., Molina, M. J., Su, S., Zhang, J., Hu, J., & Prinn, R. G. 2018.  
28 Changes in Emissions of Ozone-Depleting Substances from China Due to Implementation of the Montreal  
29 Protocol. *Environmental Science & Technology*, *52*(19), 11359–11366. <https://doi.org/10.1021/acs.est.8b01280>
- 30 FAO, 2011. Energy-smart food for people and climate. Rome. Available at:  
31 <http://www.fao.org/3/i2454e/i2454e00.pdf>
- 32 FAO, 2014. Opportunities for Agri-Food Chains to become Energy-Smart. Available online at:  
33 <http://www.fao.org/3/a-i5125e.pdf>
- 34 FAO. 2020a. Emissions due to agriculture. Global, regional and country trends 2000–2018. FAOSTAT Analytical  
35 Brief Series No 18. Rome
- 36 FAO. 2020b. FAOSTAT: Energy Use. In: *FAO.org* [online]. <https://www.fao.org/faostat/en/#data/GN>
- 37 FAO. 2021a. FAOSTAT: Emissions Shares. In: *FAO.org* [online]. <https://www.fao.org/faostat/en/#data/EM>
- 38 FAO. 2021b. FAOSTAT: Emissions Totals. In: *FAO.org* [online]. <https://www.fao.org/faostat/en/#data/GT>
- 39 Flammini, A., Pan, X., Tubiello, F.N., Qiu, S.Y., Rocha Souza, L., Quadrelli, R., Bracco, S., Benoit, P. and Sims,  
40 R., 2021. Emissions of greenhouse gases from energy use in agriculture, forestry and fisheries: 1970–2019. *Earth*  
41 *System Science Data Discussions*. Available at <https://essd.copernicus.org/preprints/essd-2021-262/>
- 42 Gütschow J., Jeffery L. and R. Gieseke. 2021. The PRIMAP-hist national historical emissions time series v2.3  
43 (1850-2017). GFZ Data Services. Available from: <https://doi.org/10.5880/pik.2019.001>
- 44 Hart, M., Austin, W., Acha, S., Le Brun, N., Markides, C. N., & Shah, N. 2020. A roadmap investment strategy to  
45 reduce carbon intensive refrigerants in the food retail industry. *Journal of Cleaner Production*, *275*, 123039.  
46 <https://doi.org/10.1016/j.jclepro.2020.123039>
- 47 Hönle, S. E., Heidecke, C., & Osterburg, B. 2019. Climate change mitigation strategies for agriculture: An analysis  
48 of nationally determined contributions, biennial reports and biennial update reports. *Climate Policy*, *19*(6), 688–  
49 702. <https://doi.org/10.1080/14693062.2018.1559793>

- 1 International Fertiliser Society. 2019. The carbon footprint of fertiliser production: regional reference values.  
2 Available from: [https://www.fertilizerseurope.com/wp-content/uploads/2020/01/The-carbon-footprint-of-](https://www.fertilizerseurope.com/wp-content/uploads/2020/01/The-carbon-footprint-of-fertilizer-production_Regional-reference-values.pdf)  
3 [fertilizer-production\\_Regional-reference-values.pdf](https://www.fertilizerseurope.com/wp-content/uploads/2020/01/The-carbon-footprint-of-fertilizer-production_Regional-reference-values.pdf)
- 4 International Institute of Refrigeration. 2021. *The Carbon Footprint of the Cold Chain, 7<sup>th</sup> Informatory Note on*  
5 *Refrigeration and Food*. [https://iifiir.org/en/fridoc/the-carbon-footprint-of-the-cold-chain-7-lt-sup-gt-th-lt-sup-gt-](https://iifiir.org/en/fridoc/the-carbon-footprint-of-the-cold-chain-7-lt-sup-gt-th-lt-sup-gt-informatory-143457)  
6 [informatory-143457](https://iifiir.org/en/fridoc/the-carbon-footprint-of-the-cold-chain-7-lt-sup-gt-th-lt-sup-gt-informatory-143457)
- 7 IPCC. 1996. Climate Change 1995 - The Science of Climate Change: Contribution of Working Group I to the  
8 Second Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge. Available at:  
9 [https://www.ipcc.ch/ipccreports/sar/wg\\_I/ipcc\\_sar\\_wg\\_I\\_full\\_report.pdf](https://www.ipcc.ch/ipccreports/sar/wg_I/ipcc_sar_wg_I_full_report.pdf)
- 10 IPCC. 2000. Good practice guidance and uncertainty management in national greenhouse gas inventories. In: J.  
11 Penman *et al.* (Eds.), IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama,  
12 Japan. Available at: [http://www.ipcc-nggip.iges.or.jp/public/gp/english/gp\\_gaum\\_en.html](http://www.ipcc-nggip.iges.or.jp/public/gp/english/gp_gaum_en.html)
- 13 IPCC. 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National  
14 Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. & Tanabe K. (eds).  
15 Published: IGES, Japan. Available at: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/vol5.html>
- 16 IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth  
17 Assessment Report of the Intergovernmental Panel on Climate Change. Available at:  
18 [https://www.ipcc.ch/site/assets/uploads/2018/02/SYR\\_AR5\\_FINAL\\_full.pdf](https://www.ipcc.ch/site/assets/uploads/2018/02/SYR_AR5_FINAL_full.pdf)
- 19 JRC. European Commission. 2015. Monforti, F., Dallemand, J., Pascua, I., Motola, V., Banja, M., Scarlat, N.,  
20 Medarac, H., Castellazzi, L., Labanca, N., Bertoldi, P., Pennington, D., Goralczyk, M., & Schau, E., Saouter, E.,  
21 & Sala, S., Notarnicola, B., Tassielli, G., and Renzulli, P. Energy use in the EU food sector: State of play and  
22 opportunities for improvement.
- 23 Karl, K. and Tubiello, F.N. 2021a. *Methods for Estimating Greenhouse Gas Emissions from Food Systems:*  
24 *Domestic Food Transport*. Rome.
- 25 Karl, K. and Tubiello, F.N. 2021b. *Methods for Estimating Greenhouse Gas Emissions from Food Systems: Food*  
26 *Systems Waste Disposal*. Rome.
- 27 Lamb, W. F., Grubb, M., Diluiso, F., & Minx, J. C. 2022. Countries with sustained greenhouse gas emissions  
28 reductions: An analysis of trends and progress by sector. *Climate Policy*, 22(1), 1–17.  
29 <https://doi.org/10.1080/14693062.2021.1990831>
- 30 Mbow, C., C. Rosenzweig, L.G. Barioni, T.G. Benton, M. Herrero, M. Krishnapillai, E. Liwenga, P. Pradhan,  
31 M.G. Rivera-Ferre, T. Sapkota, F.N. Tubiello, Y. Xu, 2019: Food Security. In: *Climate Change and Land: an*  
32 *IPCC special report on climate change, desertification, land degradation, sustainable land management, food*  
33 *security, and greenhouse gas fluxes in terrestrial ecosystems* [P.R. Shukla, et al. (eds.)].
- 34 Minx, J. C., Lamb, W. F., Andrew, R. M., Canadell, J. G., Crippa, M., Döbbling, N., Forster, P. M., Guizzardi,  
35 D., Olivier, J., Peters, G. P., Pongratz, J., Reisinger, A., Rigby, M., Saunio, M., Smith, S. J., Solazzo, E., & Tian,  
36 H. 2021. A comprehensive and synthetic dataset for global, regional, and national greenhouse gas emissions by  
37 sector 1970–2018 with an extension to 2019. *Earth System Science Data*, 13(11), 5213–5252.  
38 <https://doi.org/10.5194/essd-13-5213-2021>
- 39 Pablo-Romero, María del P., Rafael Pozo-Barajas, and Rocío Yñiguez. 2017. “Global Changes in Residential  
40 Energy Consumption.” *Energy Policy* 101. Available at: <https://doi.org/10.1016/j.enpol.2016.10.032>.
- 41 Park, W. Y., Shah, N., Vine, E., Blake, P., Holuj, B., Kim, J. H., & Kim, D. H. 2021. Ensuring the climate benefits  
42 of the Montreal Protocol: Global governance architecture for cooling efficiency and alternative refrigerants.  
43 *Energy Research & Social Science*, 76, 102068. <https://doi.org/10.1016/j.erss.2021.102068>
- 44 Poore, J. & Nemecek, T. 2018. Reducing Food’s Environmental Impacts through Producers and Consumers.  
45 *Science* 360, no. 6392: 987–92. <https://doi.org/10.1126/science.aag0216>.
- 46 Ritchie H., 2019. Food production is responsible for one-quarter of the world’s greenhouse gas emissions.  
47 Available online at <https://ourworldindata.org/food-ghg-emissions>
- 48 Rosenzweig, C., Tubiello, F.N., Sandalow, D., Benoit, P. and Hayek, M.N., 2021. Finding and fixing food system  
49 emissions: the double helix of science and policy. *Environmental Research Letters*, 16(6).
- 50 Rosenzweig, C., Mbow, C., Barioni, L. G., Benton, T. G., Herrero, M., Krishnapillai, M., Liwenga, E. T., Pradhan,  
51 P., Rivera-Ferre, M. G., Sapkota, T., Tubiello, F. N., Xu, Y., Mencos Contreras, E., & Portugal-Pereira, J. 2020.

1 Climate change responses benefit from a global food system approach. *Nature Food*, 1(2), 94–97.  
2 <https://doi.org/10.1038/s43016-020-0031-z>

3 Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A.,  
4 Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D.  
5 R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., ... Zhuang, Q. 2020. The Global Methane Budget  
6 2000–2017. *Earth System Science Data*, 12(3), 1561–1623. <https://doi.org/10.5194/essd-12-1561-2020>

7 Sims R.E.H. and Flammini A., 2014. Energy-smart food – technologies, practices and policies. Chapter 6.  
8 Sustainable Energy Solutions in Agriculture. Taylor & Francis Group, London, UK

9 Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsidig, E. A., Haberl, H., Harper, R., House, J.,  
10 Jafari, M., Masera, O., Mbow, C., Ravindranath, N. J., Rice, C. W., Robledo Abad, C., Romanovskaya, A.,  
11 Sperling, F., and Tubiello, F. N. 2014. Agriculture, Forestry and Other Land Use (AFOLU), in Working Group III  
12 contribution to the IPCC 5th Assessment Report, Climate Change 2014: Mitigation of Climate Change, edited by:  
13 Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner,  
14 S., Eickemeier, P., Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., Zwickel, T., and Minx, J. C.,  
15 Cambridge University Press, Cambridge, UK, New York, NY, USA

16 Tao, S., M. Y. Ru, W. Du, X. Zhu, Q. R. Zhong, B. G. Li, G. F. Shen, *et al.* 2018. “Quantifying  
17 the Rural Residential Energy Transition in China from 1992 to 2012 through a Representative National  
18 Survey.” *Nature Energy* 3, no. 7: 567–73. <https://doi.org/10.1038/s41560-018-0158-4>.

19 Tassou, S. A., Ge, Y., Hadawey, A., & Marriott, D. 2011. Energy consumption and conservation in food retailing.  
20 *Applied Thermal Engineering*, 31(2), 147–156. <https://doi.org/10.1016/j.applthermaleng.2010.08.023>

21 Tubiello, F. N., Salvatore, M., Rossi, S., Ferrara, A., Fitton, N., & Smith, P. 2013. *The FAOSTAT database of*  
22 *greenhouse gas emissions from agriculture*. 8(1), 015009. <https://doi.org/10.1088/1748-9326/8/1/015009>

23 Tubiello, F.N., Rosenzweig, C., Conchedda, G., Karl, K., Gütschow, J., Xueyao, P., Obli-Laryea, G., Wanner, N.,  
24 Qiu, S.Y., De Barros, J. and Flammini, A., 2021a. Greenhouse gas emissions from food systems: building the  
25 evidence base. *Environmental Research Letters*, 16(6), p.065007.

26 Tubiello, F.N., Flammini, A., Karl, K., Obli-Laryea, G., Qiu, S.Y., Heiðarsdóttir, H., Pan, X., and Conchedda, G.,  
27 2021b. *Methods for estimating greenhouse gas emissions from food systems. Part III: energy use in fertilizer*  
28 *manufacturing, food processing, packaging, retail and household consumption*. FAO Statistics Working Paper 21-  
29 Rome.

30 Tubiello, F. N., Conchedda, G., Wanner, N., Federici, S., Rossi, S., & Grassi, G. 2021c. Carbon emissions and  
31 removals from forests: New estimates, 1990–2020. *Earth System Science Data*, 13(4), 1681–1691.  
32 <https://doi.org/10.5194/essd-13-1681-2021>

33 Tubiello, F.N., Karl, K., Flammini, A., Conchedda, G., and Obli-Laryea, G., 2021d. Food Systems Emissions  
34 Shares, 1990-2019. Open access dataset, available at <https://zenodo.org/record/5615082>

35 UNFCCC. 2021. Nationally determined contributions under the Paris Agreement. *Synthesis report by the*  
36 *secretariat*. Available at: <https://unfccc.int/documents/306848>

37 Wang, Lijun. 2014. “Energy Efficiency Technologies for Sustainable Food Processing.” *Energy Efficiency*, vol. 7,  
38 no. 5, pp. 791–810. *Springer Link*. doi:10.1007/s12053-014-9256-8.

39 Vermeulen, Sonja J., Bruce M. Campbell, and John S.I. Ingram. 2012. “Climate Change and Food Systems.”  
40 *Annual Review of Environment and Resources* 37, no. 1: 195–222. <https://doi.org/10.1146/annurev-environ-020411-130608>.

41  
42  
43  
44  
45

1 TABLES

2

3

Activity	Category	1990	2019	Change
Net Forest conversion	Land Use Change	4,392	3,058	-30%
Enteric Fermentation	Farm-Gate	2,494	2,823	13%
Livestock Manure	Farm-Gate	1,101	1,315	19%
Household Consumption	Pre- and Post- Production	541	1,309	142%
Waste Disposal	Pre- and Post- Production	984	1,278	30%
On-farm energy use	Farm-Gate	757	1,021	35%
Food Retail	Pre- and Post- Production	128	932	631%
Drained organic soils	Pre- and Post- Production	736	833	13%
Rice Cultivation	Farm-Gate	621	674	9%
Fires	Land Use Change	558	654	17%
Synthetic Fertilizers	Farm-Gate	422	601	42%
Food Transport	Pre- and Post- Production	327	586	79%
Food Processing	Pre- and Post- Production	421	510	21%
Fertilizers Manufacturing	Pre- and Post- Production	152	408	168%
Food Packaging	Pre- and Post- Production	166	310	87%
Crop Residues	Farm-Gate	161	226	40%
Forestland	N/A	-3,391	-2,571	-24%

4

5 **Table 1.** GHG emissions (Mt CO<sub>2</sub>eq) by agri-food systems component for all processes considered in this work.  
6 Data on forestland removals are provided for completeness of land-based emissions available in FAOSTAT.  
7 Uncertainties (not shown) are estimated at 30% for farm gate and pre- and post-production components and at 50%  
8 for land use change processes.

9



1

<i>Region</i>	<i>Farm Gate</i>	<i>LUC</i>	<i>PPP</i>	<i>Total</i>	<i>%PPP</i>	<i>%PPP (1990)</i>	<i>Highest PPP</i>	<i>note</i>
<i>Asia</i>	3.2	0.9	2.9	7.0	42%	24%	0.9	Household
<i>Africa</i>	1.1	1.2	0.4	2.7	14%	16%	0.2	Household
<i>South America</i>	1.0	1.1	0.3	2.4	12%	6%	0.1	Waste
<i>Europe</i>	0.9	0.1	1.1	2.1	55%	26%	0.4	Retail
<i>Northern America</i>	0.6	0.2	0.8	1.5	52%	35%	0.3	Retail
<i>Oceania</i>	0.2	0.0	0.1	0.3	23%	11%	0.0	Retail

2

3

4 **Table 2.** Regional GHG emissions (Gt CO<sub>2</sub>eq) by agri-food systems component, showing farm gate, land use  
5 change (LUC), pre- and post-production processes (PPP) and total emissions Percentage contribution of PPP  
6 shown for the year 1990 and 2019. The last two columns show the largest estimated contributing PPP activity by  
7 region. Uncertainties are estimated to be 30% for farm gate and PPP activities, 50% for land use change.

8

<b>Country</b>	<b>Farm-gate</b>	<b>LUC</b>	<b>PPP</b>	<b>Total</b>	<b>Main PPP</b>	<b>Main PPP Name</b>
<b>China</b>	792	0	1102	1894	469	Household Consumption
<b>India</b>	768	0	618	1386	205	On-farm
<b>Brazil</b>	553	663	144	1360	79	Waste Disposal
<b>Indonesia</b>	491	658	132	1281	76	Waste Disposal
<b>USA</b>	477	60	696	1232	292	Retail
<b>DRC</b>	28	624	9	660	8	Waste Disposal
<b>Russian Federation</b>	146	35	362	542	177	Retail
<b>Pakistan</b>	205	7	71	283	33	Waste Disposal
<b>Canada</b>	97	96	81	274	20	Retail
<b>Mexico</b>	115	15	116	246	56	Waste Disposal

1  
2 **Table 3.** Top ten country GHG emissions (Mt CO<sub>2</sub>eq) by agri-food systems component and total food systems  
3 emissions, 2019. The last two columns show the dominant sub-component of pre- and post-production processes.  
4 Agri-food system GHG emissions from the top 10 countries represent 55% of global agri-food system emissions.  
5 Country level uncertainties those used for global and regional estimates.

6  
7

Food system component	FAO (2011) <sup>1</sup>	Vermeulen <i>et al.</i> (2012) <sup>2</sup>	Poore & Nemecek (2018) <sup>3</sup>	Ritchie (2019) <sup>4</sup>	Tubiello <i>et al.</i> (2021a) <sup>5</sup>	Crippa <i>et al.</i> (2021) EDGAR-FOOD <sup>6</sup>	This analysis <sup>6</sup>	
Reference year	Mid-2000s	2004–2007	2009–2011	2017	2019	2015	2019	
Fertilizer manufacturing	-	0.3–0.6	-	-	-	-	0.4	
Food processing	2.1	0.2	0.6	0.5	4.3 (incl. retail and household consumption)	0.5	0.5	
Food packaging		0.4	0.6	0.7		1.0	0.3	
Food transport			0.8	0.8		0.5	0.9	0.6
Food retail		0.7	0.4	0.4			0.8	0.9
Food household consumption	1.2	0.2	-	-		0.5	1.3	
Waste disposal	-	0.1	-	-	1.0	1.6	1.3	
On-farm electricity generation	-	-	-	-	-	-	0.5	
<b>TOTAL</b>	<b>3.3</b>	<b>1.9–2.2</b>	<b>2.4</b>	<b>2.4</b>	<b>5.8</b>	<b>5.3</b>	<b>5.8</b>	

2 <sup>1</sup> Includes emissions from indirect energy inputs (e.g. manufacturing of machinery). Global estimate based on  
3 literature.

4 <sup>2</sup> Global estimate based on Chinese and British emission patterns and literature.

5 <sup>3</sup> Meta-analysis of life-cycle assessments

6 <sup>4</sup> Global estimate based on literature

7 <sup>5</sup> Global estimate largely based on country-level (bottom-up) analysis (relying on FAOSTAT and EDGAR-FOOD)

8 <sup>6</sup> Global estimate largely based on country-level (bottom-up) analysis

9

10 **Table 4.** Overview of pre- and post-food production GHG emission estimates from selected studies, Gt CO<sub>2</sub>eq.

11 Adapted from Tubiello *et al.* (2021b).

12

13

1

	Farm gate		Land Use Change		Pre- and Post-Production		Agri-food Total	Systems
	1990	2019	1990	2019	1990	2019	1990	2019
<b>Africa</b>	705	1139	1017	1220	323	388	2045	2747
	23%	24%	33%	26%	11%	8%	67%	57%
<b>Asia</b>	2595	3250	1273	865	1223	2930	5091	7044
	25%	11%	12%	3%	12%	10%	49%	24%
<b>Europe</b>	1603	854	88	83	589	1140	2280	2077
	16%	13%	1%	1%	6%	17%	23%	31%
<b>North America</b>	538	574	175	156	376	777	1089	1507
	8%	8%	3%	2%	6%	11%	17%	21%
<b>South America</b>	728	982	1974	1106	176	281	2878	2369
	23%	30%	64%	34%	6%	9%	93%	72%
<b>Oceania</b>	267	223	65	16	42	71	374	309
	40%	28%	10%	2%	6%	9%	57%	39%
<b>World</b>	6604	7214	4676	3503	2886	5827	14165	16544
	19%	13%	13%	6%	8%	11%	40%	31%

2

3

4

5

6

7

8

9

8

9

**Table 5.** Regional GHG emissions (Mt CO<sub>2</sub>eq) by agri-food systems component and total food systems emissions, 2019. The last two columns show the dominant sub-component of pre- and post-production processes. Uncertainties (not shown) are estimated at 30% for farm gate and pre- and post-production components and at 50% for land use change processes.

1

	1990	2019	1990	2019	1990	2019	1990	2019	1990	2019
	CO <sub>2</sub> eq		CO <sub>2</sub>		CH <sub>4</sub>		N <sub>2</sub> O		F-gases	
World	40	31	31	21	60	53	79	78	0	27
Africa	67	57	65	52	63	58	90	87	0	20
Northern America	17	21	11	13	36	42	60	70	0	56
South America	93	72	97	70	82	75	94	92	0	6
Asia	49	24	38	16	66	49	84	80	0	9
Europe	23	31	13	23	46	47	70	74	0	28
Oceania	57	39	38	22	76	64	93	77	0	63

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

## 1 **FIGURE LEGENDS**

2

3 **Figure 1.** Mapping of emissions across agri-food systems. Left-hand panel: IPCC sectors and processes used in  
4 national GHG emissions inventories. Right-hand panel: food and agriculture sectors and categories aligned to  
5 FAO's definitions.

6

7 **Figure 2.** World-total GHG emissions from agri-food systems, 1990-2019. Color bars show contributions by  
8 emissions within the farm gate (yellow); land use change (green) and pre- and post- production along food supply  
9 chains (blue). Source: FAOSTAT (FAO, 2021). Also shown are emissions per capita (authors' own calculations).

10

11 **Figure 3.** World total 2019 GHG emission from agri-food systems, showing contributions on agricultural land  
12 (left panel) and from pre- and post- production along food supply chains (right panel). Net removals on forest land  
13 are also shown, for completeness. The sum of emissions from agricultural land and forest land correspond to the  
14 IPCC AFOLU category. Source: FAOSTAT (FAO, 2021).

15

16 **Figure 4.** Total GHG emission from agri-food systems by FAO regions, 2019. Color bars show contributions by  
17 emissions within the farm gate (yellow); land use change (green) and pre- and post- production along food supply  
18 chains (blue). Source: FAOSTAT (FAO, 2021).

19

20 **Figure 5.** Total GHG emission from agri-food systems by country, top ten emitters, 2019. Color bars show  
21 contributions by emissions within the farm gate (yellow); land use change (green) and pre- and post- production  
22 along food supply chains (blue). Source: FAOSTAT (FAO, 2021).

23


24 **Figure 6.** Top panel: Agri-food systems emissions (GtCO<sub>2</sub>eq yr<sup>-1</sup>); Bottom panel: shares of agri-food systems in  
25 total anthropogenic emissions (%). Data shown by region, 1990-2019. Color bars show contributions component:  
26 farm gate (yellow); land use change (green) and pre- and post- production along food supply chains (blue). Source:  
27 FAOSTAT (FAO, 2021).

28

IPCC	Food Systems Activity	GHG			FAO			
		CH <sub>4</sub>	N <sub>2</sub> O	CO <sub>2</sub>				
<b>AFOLU</b>	<b>LULUCF</b>	Net Forest Conversion	x	x	x	<b>LAND USE CHANGE</b>	<b>AGRICULTURAL LAND</b>	<b>FOOD SYSTEMS</b>
		Tropical Forest Fires	x	x	x			
		Peat Fires	x		x			
		Drained Organic Soils	x		x			
	<b>AGRICULTURE</b>	Burning - Crop residues	x	x		<b>FARM GATE</b>		
		Burning - Savanna	x	x				
		Crop Residues		x				
		Drained Organic Soils		x				
		Enteric Fermentation	x					
		Manure Management	x	x				
		Manure Applied to Soils		x				
		Manure Left on Pasture		x				
		Rice Cultivation	x					
		Synthetic Fertilizers		x				
<b>ENERGY</b>	On-farm Energy Use	x	x	x	<b>PRE AND POST PRODUCTION</b>			
	Transport	x	x	x				
	Processing	x	x	x				
	Packaging	x	x	x				
	Fertilizer manufacturing	x	x	x				
	Household consumption	x	x	x				
	Retail –Energy Use	x	x	x				
<b>Industry</b>	Retail –Refrigeration	x	x	x				
<b>WASTE</b>	Solid Food Waste	x						
	Incineration			x				
	Industrial Wastewater	x	x					
	Domestic Wastewater	x	x					

Figure 1. Mapping of emissions across agri-food systems. Left-hand panel: IPCC sectors and processes used in national GHG emissions inventories. Right-hand panel: food and agriculture sectors and categories aligned to FAO’s definitions


1  
2



3  
4  
5  
6  
7


**Figure 2. World-total GHG emissions from agri-food systems, 1990-2019.** Color bars show contributions by emissions within the farm gate (yellow); land use change (green) and pre- and post- production along food supply chains (blue). Source: FAOSTAT (FAO, 2021). Also shown are emissions per capita (authors' own calculations).





1  
 2 **Figure 3. World total 2019 GHG emission from agri-food systems.** Contributions on agricultural land are displayed on the  
 3 left, and from pre- and post- production along food supply chains on the right. Net removals on forest land are also shown in  
 4 the center for completeness. The sum of emissions from agricultural land and forest land correspond to the IPCC AFOLU  
 5 category. Source: FAOSTAT (FAO, 2021).

6



1

2


3 **Figure 4. Total GHG emission from agri-food systems by FAO regions, 2019.** Color bars show contributions by emissions

4 within the farm gate (yellow); land use change (green) and pre- and post- production along food supply chains (blue). Source:

5 FAOSTAT (FAO, 2021).


6

7



1  
2  
3  
4  
5

**Figure 5. Total GHG emission from agri-food systems by country, top ten emitters, 2019.** Color bars show contributions by emissions within the farm gate (yellow); land use change (green) and pre- and post- production along food supply chains (blue). Source: FAOSTAT (FAO, 2021).



1  
2  
3  
4  
5  
6  
7

**Figure 6. Top panel: Agri-food systems emissions (GtCO<sub>2</sub>eq yr<sup>-1</sup>).** Bottom panel: shares of agri-food systems in total anthropogenic emissions (%). Data shown by region, 1990-2019. Color bars show contributions component: farm gate (yellow); land use change (green) and pre- and post- production along food supply chains (blue). Source: FAOSTAT (FAO, 2021).