

Historical and Future Weather Data for Dynamic Building Simulations in Belgium using the MAR model: Typical & Extreme Meteorological Year and Heatwaves

5 Sebastien Doutreloup¹, Xavier Fettweis¹, Ramin Rahif², Essam Elnagar³, Mohsen S. Pourkiaei⁴, Deepak Amaripadath², Shady Attia²

¹: Laboratory of Climatology and Topoclimatology, Department de Geography, UR SPHERES, University of Liège, Belgium

²: Sustainable Building Design Lab, Dept. UEE, Faculty of Applied Sciences, University of Liege, Belgium

10 ³: Thermodynamics Laboratory, Aerospace and Mechanical Engineering Department, Faculty of Applied Sciences, University of Liège, Belgium

⁴: Atmospheres and Monitoring lab (SAM), UR Spheres, University of Liege, Arlon Campus Environment, Avenue de Longwy 185, 6700 Arlon, Belgium.

Correspondence to: Doutreloup S. (s.doutreloup@uliege.be)

Abstract. Increasing temperature due to global warming will influence building, heating and cooling practices. Therefore
15 this dataset aims to provide formatted and adapted meteorological data for specific users who work in building designing, architecture, building energy management systems, modeling renewable energy conversion systems, or other people interested in this kind of projected weather data. These meteorological data are produced from the regional climate MAR model simulations. This regional model adapted and validated over Belgium is forced firstly by the ERA5 reanalysis which represents the closest climate to reality, and secondly by 3 ESMS from the [CMIP6 Sixth Coupled Model Intercomparison Project](#)
20 [Project](#) database, namely BCC-CSM2-MR, MPI-ESM.1.2, and MIROC6. The main advantage of using the MAR model is that the generated weather data have a high resolution ([hourly data and 5km](#)) and are spatially and temporally homogeneous. The generated weather data follow two protocols. On one hand, the Typical Meteorological Year (TMY) and eXtreme Meteorological Year (XMY) files are generated [following largely inspired by](#) the method proposed by the standard ISO15927-4, allowing the reconstruction of typical and extreme years while keeping a plausible variability of the
25 meteorological data. On the other hand, the HeatWaves Event (HWE) meteorological data are generated according to [the method proposed by Ouzeau et al. \(2016\)](#) used to detect the heatwave events and to classify them according to three criteria of the heatwave (the most intense, the longest duration and the highest temperature). All generated weather data are freely available on the open online repository Zenodo (<https://doi.org/10.5281/zenodo.5606983> ; (Doutreloup, S. and Fettweis X., 2021)) [and these data are produced within the framework of the research project OCCuPANt](#)
30 (<https://www.occupant.uliege.be/> - ULiège).

Keywords. Climate simulation; Future weather files; Typical year; Extreme year; Heatwave; Regional Modeling; Belgium

1 Introduction

On a global scale, the warmest (SSP5-8.5) scenario from the IPCC last assessment report (IPCC et al., 2021) suggests a temperature increase of +5°C by 2100. However, the regions of the world will not warm up at the same speed or intensity. Some regions, such as the poles, ~~will warm faster and larger than equatorial regions will warm faster and to higher levels than equatorial regions~~ (Lee, J. Y. et al., 2021). Over the temperate oceanic regions such as Western Europe, the temperature is expected to increase between +1°C and +5°C in 2100 depending on the climate models and greenhouse gas emission scenarios used (Termonia et al., 2018; RMI, 2020; IPCC et al., 2021).

Moreover, IPCC et al. (2021) affirm that extreme events will become more probable and more intense. In particular, the maximum temperature is expected to increase faster (sometimes up to twice) than the mean temperature (Seneviratne, S. I. et al., 2021). Over Western-Central Europe, maximum temperatures are projected to increase up to +7°C for a global increase of +5°C (Seneviratne, S. I. et al., 2021).

More concretely, during the summer season, hot extremes (including heatwaves) are already increasing and will continue to strengthen with global warming both in intensity and frequency (Suarez-Gutierrez et al., 2020; Seneviratne, S. I. et al., 2021; Dunn et al., 2020). The consequences of these heatwaves will affect human health (Fouillet et al., 2006), agriculture, the comfort and health of life inside buildings (Bruffaerts et al., 2018; Sherwood and Huber, 2010; Buysse et al., 2010), and the energy demand especially for cooling systems (Larsen et al., 2020). This is what motivated some previous studies in Belgium to represent the energy needs for heating and cooling under average and extreme weather conditions (Ramon et al., 2019).

~~This comfort of life. The energy efficiency and living comfort areis~~ precisely what motivates the ULiège OCCuPANT project (Impacts Of Climate Change on the indoor environmental and energy PerformAnce of buildiNGs in Belgium during summer, <https://www.occupant.uliege.be/>) in which climate data are involved. ~~The OCCuPANT project aims to evaluate the vulnerability of building inhabitants in a warmer climate context and more particularly during heatwaves. The OCCuPANT project aims to evaluate the energy performance and vulnerability of building inhabitants in the context of climate change. Acquiring reliable current and future climate data is vital in any study related to climate change and defines its quality (Pérez-Andreu and al., 2018).~~

The purpose of this dataset is to propose meteorological data coming from a fine resolution regional climate model over Belgium and the neighbouring regions. ~~These data will then be used to anticipate future climate changes which will influence both the production of heating and cooling demands as well as the electrical grids in larger scales. These changes will require innovations in building design and systems which will necessarily take time. Thus, the more they are anticipated, the better we will find solutions.~~ The use of a regional model allows building spatially and temporally continuous and homogeneous past and future meteorological data according to different warming scenarios for some Belgian cities. This regional model is fed by the ERA-5 reanalysis model (Hersbach et al., 2020) to simulate the past climate (1980-2020) and also by 3 different Earth System Models (ESMs) from the [Sixth Coupled Model Intercomparison Project \(CMIP6\)](#) database

65 (Wu et al., 2019; Tatebe et al., 2019; Gutjahr et al., 2019) to obtain different future projections and associated uncertainties for the same scenario SSP5-8.5.

70 The future climate data are very useful to predict the variations in heating and cooling demands in buildings. The characterization of minimum outdoor temperature under future scenarios is necessary to estimate the heating and cooling season needs. This can result in rethinking the building designs and make them resilient against the impact of climate change. For instance, in heating-dominated region of Belgium, the concept of building design is more on heat retention to decrease the heating energy use during the winter. However, by warming weather conditions in the last decades, such a design concept caused significant overheating problems during the summer. Therefore, it is necessary to predict the future performance of buildings and adapt them to the variations in the outdoor weather conditions. Designing cooling systems that can last for 100 years is challenging. However, it is possible to increase the preparedness of buildings to climate change through passive design strategies, the use of active heating/cooling systems, or both. Both active and passive solutions may need regular replacements over the building life span.

75 For each city, considered period, model, and scenario, two synthetic files (in CSV format) are generated following largely inspired by the method in ISO-15927-4: Typical Meteorological Year file (TMY) and eXtreme Meteorological Year file (XMY). In addition to these synthetic files, files focused on heatwaves are also generated, namely a file for the most intense heatwave event, one for the warmest heatwave, one for the longest heatwave, and one containing all the heatwaves detected within a specific period. Finally, these files are described in detail in the methodology section.

2. Methodology

2.1 MAR model and area of interest

85 The regional climate model used in this study is the “Modèle Atmosphérique Régional” model (hereafter called “MAR”) in its version 3.11.4 (Kittel, 2021). The main role of MAR is to downscale a global model or reanalysis in order to get weather outputs at a finer spatial and temporal resolution (Fig. 1). As shown in Fig. 1, MAR is a three-dimensional atmospheric model coupled to a one-dimensional transfer scheme between the surface, vegetation, and atmosphere (Ridder and Gallée, 1998). The MAR model also includes an urban island module which modifies the city grid points to simulate urban heat island through a modification of the surface albedo (fixed to 0.1) and an absence of vegetation which influence the thermal and humidity exchanges between soil and atmosphere. Initially, the MAR model was developed for both Greenland (Fettweis et al., 2013) and Antarctica ice sheets (Agosta et al., 2019; Kittel et al., 2021). However, it has recently successfully adapted to temperate regions such as Belgium (Fettweis X. et al., 2017 ; Doutrélop et al., 2019; Wyard et al., 2021; Fettweis X. et al., 2017). In the framework of this study, MAR is initially forced every 6 hours at its lateral boundaries (temperature, wind, and specific humidity) by the reanalysis ERA5 (called hereafter MAR-ERA5) which is available at a horizontal resolution of ~31 km (Hersbach et al., 2020). Different kinds of observations (*in situ* weather observation, radar

data, satellites, *etc*) are 6-hourly assimilated into ERA5 to be the closest to the observed climate. In this way, the simulations of MAR-ERA5 can be considered as ~~a reconstruction the closest simulation of to~~ the current observed climate.

Then, the MAR model has been forced every 6 hours by 3 ESMs from the CMIP6 database (Eyring et al., 2016). These ESMs do not contain any observational data and represent only ~~the mean an~~ evolution of the average and interannual variability of the climatic parameters over long periods. These models contain two characteristic periods: one in the past, from 1980 to 2014 (hereafter called "historical" scenario), and another in the future, from 2015 to 2100 according to different SSP scenarios (SSP5_8_5, SSP3_7_0, and SSP2_4_5). The selection and description of each of these ESMs as well as a comparison with MAR-ERA5 over the historical period are presented in the next section.

The atmospheric variables used to force MAR every 6 hours at each ~~MAR~~-vertical level are temperature, surface pressure, wind, and specific humidity, as well as the sea surface temperature over the North Sea from both ERA5 reanalysis (from 1980 to 2020) and the three ESMs (from 1980 to 2100). The spatial resolution of MAR is 5km over an integration domain (120 x 90 grid cells) centered over Belgium as shown in Fig. 2 to build hourly outputs.

The choice of 12 cities is motivated, on the one hand, by the size of the cities which must be sufficient to show a temperature increase compared to the neighbouring countryside and, on the other hand, to best represent climate spatial variability observed in Belgium. For example, the city of Ostend is strongly influenced by the thermal inertia of the sea, while the city of Arlon has a more continental climate.

2.2 Forcing models and MAR simulations

2.2.1 Choice of representative ~~ESMs~~Earth System Models

The Sixth Coupled Model Intercomparison Project (CMIP6; Eyring et al. (2016)) database contains about 30 ESMs from many scientific institutes around the world. For practical reasons, we can't regionalize all these ESMs. Thus, we had to select a few representative ESMs for our region of interest, West-Europe.

Our choice was based on two criteria. The first criterion is that the ESM should represent (with the lowest possible bias) the main atmospheric circulation in the free atmosphere over western Europe by evaluating the geopotential height at 500hPa and the temperature at 700hPa during summer and winter with respect to ERA5 over 1980-2014 based on the skill score method developped by Connolley and Bracegirdle (2007). After selecting the ESMs that meet this first criterion, the second criterion is to choose 3 ESMs representing the CMIP6 models spread in 2100 for the same scenario (SSP5-8.5 here). Namely, we keep only 3 ESMs: BCC-CSM2-MR (Wu et al., 2019), MPI-ESM.1.2 (Gutjahr et al., 2019), and MIROC6 (Tatebe et al., 2019). The ESM BCC-CSM2-MR simulates a warming close to the averageensemble mean of ~~all the~~ 30 ESMs for the 2100 horizon withusing the SSP5-8.5 scenario, the ESM MIROC6 simulates a larger warming than averageensemble mean, and the ESM MPI-ESM.1.2 simulates a lower warming than average-the ensemble mean by 2100. The use of these 3 models allows us to obtain a first approximation of the uncertainty from ESMs without having to downscale all the 30 available models of CMIP6.

2.2.2 Future socioeconomic scenarios

130 Shared Socioeconomic Pathways (SSPs; Riahi et al. (2017)) are scenarios of global socio-economic evolution projected to 2100. These SSPs are used to develop greenhouse gas emission (GHG) scenarios associated with different climate policies and are used to force each future ESMs. There are 3 main scenarios, namely SSP2-4.5 (intermediate GHG), SSP3-7.0 (high GHG), and SSP5-8.5 (very high GHG), ~~which are with their~~ respectively increasingly warming for 2100. For more details about these scenarios, we refer to Riahi et al. (2017).

135 For the same practical reasons that led us to choose only 3 ESMs out of the 30 available models, we cannot afford to calculate every SSP scenario of each ESM. Thus, since the ESMs do not simulate general atmospheric circulation changes (Eyring, V. et al., 2021) leading that only temperature drives the evolution of the climate. Hence, only the SSP5-8.5 scenario has been calculated to save computational time. Then, scenarios SSP3-7.0 and SSP2-4.5 are reconstructed from the MAR simulations forced by the ESMs with the SSP5-8.5 scenario since the warming rates from the lower scenarios are included in this scenario SSP5-8.5 but for a different earlier period.

140 Therefore, this method involves determining the period of the SSP5-8.5 scenario that corresponds to the warming rate projected by the other scenario for a given period. For this purpose, we use the raw data of each ESM and each scenario has been aggregated over Belgium to the horizon 2100. Then, for each ESM and for the SSP3-7.5 and SSP2-4.5 scenarios, we identify the corresponding period (the most equivalent mean and variability) in the SSP5-8.5 scenario ~~for each decade~~. Then, we extract the weather data of this corresponding period for SSP3-7.5 and SSP2-4.5 out of the SSP5-8.5 forced MAR simulations (i.e., weather data of MAR-MPI with scenario SSP5-8.5 during the period 2066-2085 represent MAR-MPI with scenario SSP3-7.0 for the period 2081-2100). This method is open to discussion for several reasons, firstly because climate does not react in a linear way to an increase of GHG flowing in different SSP scenarios. Moreover, with an equal warming rate but different periods, the Earth, Atmosphere and Ocean systems will not have the same (spatial and temporal) responses due to their inertia. Despite these precautions, the method used here allows on the one hand a very quick estimation without

145 additional computer time. On the other hand, it remains valid as a first approximation, especially since the most interesting weather variable in this study is temperature, which is, therefore, the least sensitive to these issues.

150

2.2.3 Evaluation of the MAR simulations

To verify that these ESMs forced MAR simulations can be used to anticipate future periods, it is necessary to evaluate them over the present period (namely 1980-2014 which corresponds to the period of the historical scenario). The aim is to determine if they are able to represent the average and the climate variability over the Belgian territory for this period as observed (i.e. ERA5 in your case). So, we compare the 3 MAR simulations forced by the 3 ESMs with the MAR simulation forced by the ERA5 reanalyses. In this case, as the two most important variables for this database are the temperature at 2m above ground level (a.g.l.) and the incident solar radiation, the means and standard deviation of these data over the period

1980-2014 and over the Belgian territory are compared in Tab. 1 and Fig. 3. These values are compared on an annual scale as well as on a summer scale since the OCCuPANt project focuses on heatwaves.

The results of this comparison presented in Tab. 1 indicate that the incident solar radiations and temperature at 2m a.g.l. values proposed by the 3 MAR simulations forced by the ESMs are ~~mostly very~~ close to the average simulated by MAR-ERA5 with (not statistically significant) biases between MAR forced by the ESMs and MAR-ERA5 lower than the standard deviation (i.e. the interannual variability) of MAR-ERA5. We can also note that the interannual variability of the MAR simulations forced by the ESMs is close to the interannual variability of the MAR-ERA5 simulation. We can then conclude that the MAR simulations forced by the ESMs are able to represent the ~~current mean~~ climate simulated by MAR-ERA5 and its interannual variability with success, except MAR-MIR which significantly overestimates temperature and solar radiation in summer. Knowing that MAR-MIR simulated the largest warming in 2100, this simulation needs to be considered as the extreme climates we could have.

2.3 Generating the Typical Meteorological Year TMY and eXtreme Meteorological Year XMY files

The Typical Meteorological Year (TMY) and the eXtreme Meteorological Year (XMY) are datasets that are widely used by building designers and others for modeling renewable energy conversion systems (Wilcox and Marion, 2008). The TMYs are the synthetic years (on an hourly basis) constructed by representative typical months (Barnaby and Crawley, 2011) which are selected by comparing the distribution of each month within the long-term (minimum 10 years) distribution of that month for the available observations or modeled data (using Finkelstein-Schafer statistics (Finkelstein and Schafer, 1971)). The XMY is the extension of the TMY weather data and is formed by selecting the most deviating (i.e., extreme) months over a certain dataset instead of typical months (Ferrari and Lee, 2008). There are many methods to reconstruct this kind of weather file (Ramon et al., 2019), but for ~~the purpose of~~ this study, a protocol for the construction of these typical years has been developed based on the ISO15927-4 (European Standard, 2005) and is described briefly below.

The method consists of reconstituting each month of the typical (resp. extreme) year with the most typical (resp. extreme) month present in the considered period for a considered city. The comparison is essentially based on two variables, namely temperature at 2m a.g.l. and incident solar radiation. The choice of these two climatic parameters (called “*para*” hereafter) is related to the fact that they both influence the comfort inside the buildings. Therefore, we generate files with the typical year according to the temperature at 2m a.g.l. and, a typical year according to the incident solar radiation.

Here are the steps to find the most typical (resp. extreme) month for each weather *para*:

- 1 Converting hourly file in daily file: From all the hourly data ~~offrom~~ all the same calendar months available within the selected period, the daily mean of *para* is ~~ealeulated~~computed;
- 2 Selecting the typical (extreme) month: For each calendar month, the percentile 50 (resp. 95) of *para* is calculated over the studied period ~~eovered~~ to find the month which is the closest to the 50 percentile (resp. 95) of *para*;

190 | 3 Extracting hourly data of this typical (extreme) month: Finally, the hourly weather values of this typical (resp. extreme) month are stored in the file of the typical (resp. extreme) year.

| The hourly weather variables available in the TMY and XMY files are described in Table 2:

- 1 Dry bulb temperature at 2m a.g.l. (°C)
- 195 | 2 Relative humidity at 2m a.g.l. (%)
- 3 Global horizontal radiation (Wh/m²)
- 4 Diffuse solar radiation (Wh/m²)
- 5 Direct normal radiation (Wh/m²)
- 6 Wind speed at 10m a.g.l. (m/s)
- 200 | 7 Wind direction (north degrees)
- 8 Dew point temperature at 2m a.g.l. (°C)
- 9 Atmospheric Pressure (Pa)
- 10 Cloudiness (in tenths)
- 11 Sky temperature (K) according to (Duffie and Beckman, 2013)
- 205 | 12 Specific humidity at 2m a.g.l. (kg/kg)
- 13 Precipitation (mm)

| 2.4 Definition of a heatwave and generating the HWEHeatWave Event files

In Belgium, heatwaves are officially defined by two definitions, a retrospective one, and a prospective one. The retrospective heatwave is defined as periods of at least 5 consecutive days with a maximum temperature higher than 25°C, of which at least 3 days within this period with a maximum temperature higher than 30°C (RMI, 2020). The prospective heatwave is a period with a predicted minimum temperature of 18.2°C or more and a maximum temperature of 29.6°C or higher both for 3 consecutive days (Tsachoua, L. and Reynders, D., 2016).

210 | However, these definitions are static and do not consider the local climate of each region. For example, onin the highlands where it is on average colder, these heatwave criteria are not necessarily met, even though this region also experiences a heatwave. Moreover, when comparing the different ESMs, this fixed heatwave definition criterion could induce artifacts since each ESM has its own variability and biases over the current climate. For these reasons, we have used the statistical definition of a heatwave from (Ouzeau et al., 2016) computed for each MAR pixel regardless of its basic climate, and each ESM independently of its own internal variability.

| The calculation method, according to (Ouzeau et al., 2016), is as follows and it is illustrated byin Fig. 4:

220 ▪ For the period 1980-2014 (which corresponds to the "historical" scenario in the ESMS), for each pixel and each MAR simulation, we calculate three thresholds defined by 3 percentiles of the daily mean temperature: Sint = 95th percentile, Sdeb = 97.5th percentile and Spic = 99.5th percentile ;

225 ▪ A heatwave is detected when the daily mean temperature reaches Spic. The duration of this event is the number of days between the first day when the daily mean temperature is equal to or greater than Sdeb and either when the daily mean temperature falls below Sint or when the daily mean temperature falls below Sdeb for three consecutive days at least.

230 ▪ In this dataset, we add a condition compared to (Ouzeau et al., 2016) which is that the minimum duration of the heatwave must be at least 5 consecutive days otherwise the heatwave event is not considered.

Once the heatwaves events (called "HWE" hereafter) are detected, we can characterize them according to three criteria:

- 1 the duration which is the number of consecutive days of the HWE;
- 230 2 the maximal daily mean temperature reached during the HWE;
- 3 the global intensity which is calculated by the cumulative difference between the temperature and the Sdeb threshold during the HWE, divided by the difference between Spic and Sdeb.

The hourly data provided in each HWE file are the same as for the TMY and XMY files (see section above). For each period, each city, each scenario, and each forcing, 4 files are created:

- 235 ▪ a file containing hourly weather data for the longest HWE;
- a file containing the hourly weather data of the HWE characterized by the highest daily average temperature;
 - a file containing the hourly weather data of the HWE characterized by the highest intensity;
 - a file that concatenates all the hourly weather data of all the HWEs present in a period, for each city, each scenario, and each MAR model.

240 Finally, the HWE files contain only the period corresponding to a heatwave event. However, a heatwave can also be dependent on the period preceding and/or following it (depending on the use of the files and the purpose of the users). Thus, a suggestion for users could be to combine HWE files with files of typical or extreme years in order to obtain simulations of a normal year with one or more heatwaves. We have not built these files in order to leave it up to the users to combine or not these files and to prepare them according to their own constraints and interests.-

245 3. The dataset

3.1 Data availability

The data described in this manuscript can be freely accessed on the Zenodo open-access repository: <https://doi.org/10.5281/zenodo.5606983> (Doutreloup, S. and Fettweis X., 2021). As the files are numerous, each zipped folder contains all the data for each city concerned by this dataset.

250 3.2 Structure of each file

All files are in CSV format comma-separated. The file names are formatted in such a way that they contain all the information about the origin of the file. Each file name is composed as follows:

- For TMY or XMY: CityName_Type_Period_Scenario_MARmodel_Para.csv
- For Heatwave: CityName_Type_Period_Scenario_MARmodel.csv

255 with :

- CityName: the name of the city considered in this file;
- Type: the nature of this file, namely TMY (typical meteorological year), XMY (extreme meteorological year), HWE-HI (most intense heatwave event), HWE-HT (warmest heatwave event), HWE-LD (longest heatwave event), and HWE-all (all heatwave events);
- 260 - Period: the time period considered;
- Scenario: the IPCC scenario of the ESM considered, namely “hist”, “SSP5-8.5”, “SSP3-7.0”, and “SSP2-4.5” (note that for MAR-ERA5 the name of the scenario is set by default to "hist" even if the ERA5 forcing does not contain any scenario);
- MAR model: the version of MAR used, namely: MAR-ERA5, MAR-BCC, MAR-MPI or MAR-MIR;
- 265 - Para: is the climatic parameter used to generate the TMY or XMY file, namely the temperature at 2m a.g.l. (TT) or the incident solar radiation (SWD).

For TMY and XMY files, the header is composed of the task number (which is a reference number for internal use in the OCCuPANt project), the name of the city, and the name of the different variables accompanied by their unit. For HWE files, the header is composed of the task number (same remark as above), the name of the city, the characterization of the

270 heatwave (longest, warmest, and most intense) accompanied by its period, and the name of the different variables accompanied by their unit. Then comes the weather data with the time variable in the first column. Weather data are in universal time and leap year mode for the MAR-ERA5 and in no-leap year mode for MAR-BCC, MAR-MPI, and MAR-MIR.

A short example of how to select the desired data and files, as well as an example of how to use it, is provided in Appendix

275 A for Liege-City.

4. Conclusion

The goal of this dataset is to provide formatted and adapted meteorological data for specific users who work in building designing, architecture, building energy management systems, modeling renewable energy conversion systems, or other people interested in this kind of data. These weather data are derived from the regional climate MAR model. This regional

280 model adapted and validated over Belgium ~~is forced by 4 ESMs~~ is forced by a reanalysis and 3 ESMs. On one hand, MAR is forced by the ERA5 reanalysis to represent the closest climate to reality. On the other hand, MAR is forced by 3 ESMs from

the CMIP6 database, namely BCC-CSM2-MR, MPI-ESM.1.2, and MIROC6. The main advantage of using the MAR model is that the generated weather data at high resolution are spatially and temporally homogeneous.

285 The generated weather data follow two protocols. Firstly, the TMY and XMY meteorological data are generated
accordingly largely inspired by the method proposed by the standard ISO15927-4, allowing the reconstruction of typical and
extreme years while keeping the plausible variability of the meteorological data. Secondly, the meteorological data
concerning heatwaves (HWE) are generated according to the method proposed by Ouzeau et al. (2016) to detect the
heatwave events and to classify them according to three criteria of the heatwave. The OCCuPANT project, in which this
290 paper is included, aims to identify the highest temperature, the longest heatwave duration, and the most intense heatwave
event. Finally, all generated weather data are open source and freely available on the open repository Zenodo
(<https://doi.org/10.5281/zenodo.5606983> ; (Doutreloup, S. and Fettweis X., 2021).

5 Appendices

5.1 Appendix A: Example for Liege city

295 The Liege folder contains 596 files. This is a huge amount of files, so to find your way around we suggest the
following method for TMY and XMY files :

- 1 Choosing a typical or extreme year, i.e. select XMY for an extreme year;
- 2 Choosing the parameter that determines the typical or extreme year ([TT for temperature and SWD for incident solar
radiation](#)), i.e. if you want a year with extreme incident solar radiation (i.e. sunshine), you should select SWDbased.
- 300 3 Choosing the reference period, i.e. 2085-2100 for the end of the century;
- 4 Choosing the socio-economic scenario, i.e. “ssp585” for a world that continues to use fossil fuels;
- 5 Choosing one of the 3 MAR models, i.e. MAR-BCC for the MAR model forced by BCC-CSM2-MR;
- 6 Finally, the choice is made for the file:

Liege-City_XMY2085-2100_ssp585_MAR-BCC_SWDbased.csv

305 This file selection method is a proposal but each user is free to develop his own method and of course you can use several
files to compare them. For example, if we want to compare typical and extreme temperatures for the end of the century and
for the two selected parameters with MAR-BCC, we get Fig. A-1. Fig. A-1 clearly shows that the extreme year temperature
based on temperature is much higher than the typical year temperature. On the other hand, the extreme year temperature
based on incident solar radiation is much lower, especially in January, than the typical year temperature because extreme
310 radiation in winter usually means cold weather.

The method is almost identical for the heatwave files except that there are no parameters to determine the typical and
extreme years but we have to choose if we want to look at the longest, the most intense, the hottest, or all heatwaves present

within the period. Therefore, if we want to compare the longest heatwave (HWE-LD) over the same period, for the same model and scenario as the example above, we choose the file :

315 Liege-City_HWE-LD_2085-2100_ssp585_MAR-BCC.csv

Fig. A-2 compares the temperature evolution during the warmest heatwave obtained from each of the MAR-BCC, MAR-MPI, and MAR-MIR simulations. The header of each file shows the warmest daily average temperature, namely 37.2°C for MAR-BCC, 35.3°C for MAR-MPI, and 37.8°C for MAR-MIR. However, it can be seen in Fig. A-2 that the hourly
320 temperatures obviously rise much higher than these daily average values and in this case rise to ~44°C for MAR-BCC and MAR-MIR and ~42°C for MAR-MPI.

6 Author contribution

All authors participated in the conceptualization of this paper. SD and XF designed the experiments and methodology and SD carried them out. SD prepared and created figures and data. XF oversaw the research. SD wrote the initial manuscript
325 and all authors participated in the revision.

7 Competing interests

The authors declare that they have no conflict of interest.

8 Acknowledgment

This research was partially funded by the Walloon Region under the call “Actions de Recherche Concertées 2019 (ARC)”
330 and the project OCCuPANT, on the Impacts Of Climate Change on the indoor environmental and energy PerformAnce of buildiNgs in Belgium during summer. The authors would like to gratefully acknowledge the Walloon Region and Liege University for funding. Computational resources have been provided by the Consortium des Équipements de Calcul Intensif (CÉCI), funded by the Fonds de la Recherche Scientifique de Belgique (F.R.S.-FNRS) under Grant No. 2.5020.11 and by the Walloon Region.

335 9 Financial support

This research was partially funded by the Walloon Region under the call “Actions de Recherche Concertées 2019 (ARC)” and the project OCCuPANT, on the Impacts Of Climate Change on the indoor environmental and energy PerformAnce of buildiNgs in Belgium during summer.

10 References


- 340 Agosta, C., Amory, C., Kittel, C., Orsi, A., Favier, V., Gallée, H., van den Broeke, M. R., Lenaerts, J. T. M., van Wessem, J. M., van de Berg, W. J., and Fettweis, X.: Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979–2015) and identification of dominant processes, *The Cryosphere*, 13, 281–296, <https://doi.org/10.5194/tc-13-281-2019>, 2019.
- Barnaby, C. S. and Crawley, U. B.: Weather data for building performance simulation, in: *Building Performance Simulation*
- 345 *for Design and Operation*, Routledge, 2011.
- Bruffaerts, N., De Smedt, T., Delcloo, A., Simons, K., Hoebeke, L., Verstraeten, C., Van Nieuwenhuysse, A., Packeu, A., and Hendrickx, M.: Comparative long-term trend analysis of daily weather conditions with daily pollen concentrations in Brussels, Belgium, *Int. J. Biometeorol.*, 62, 483–491, <https://doi.org/10.1007/s00484-017-1457-3>, 2018.
- Buysse, D. J., Cheng, Y., Germain, A., Moul, D. E., Franzen, P. L., Fletcher, M., and Monk, T. H.: Night-to-night sleep
- 350 variability in older adults with and without chronic insomnia, *Sleep Med.*, 11, 56–64, <https://doi.org/10.1016/j.sleep.2009.02.010>, 2010.
- [Conolley, W. M.; Bracegirdle, T. J. An Antarctic assessment of IPCC AR4 coupled models. *Geophys. Res. Lett.* 2007, 34, L22505, doi:10.1029/2007GL031648.](https://doi.org/10.1029/2007GL031648)
- Doutreloup, S. and Fettweis X.: Typical & Extreme Meteorological Year and Heatwaves for Dynamic Building Simulations
- 355 in Belgium based on MAR model Simulations (version 1.0.0) [Data set], <https://doi.org/10.5281/zenodo.5606983>, 2021.
- Doutreloup, S., Wyard, C., Amory, C., Kittel, C., Erpicum, M., and Fettweis, X.: Sensitivity to Convective Schemes on Precipitation Simulated by the Regional Climate Model MAR over Belgium (1987–2017), *Atmosphere*, 10, 34, <https://doi.org/10.3390/atmos10010034>, 2019.
- Duffie, J. A. and Beckman, W. A.: *Solar Engineering of Thermal Processes*, John Wiley&Sons, Inc., Hoboken, New Jersey,
- 360 2013.
- Dunn, R. J. H., Alexander, L. V., Donat, M. G., Zhang, X., Bador, M., Herold, N., Lippmann, T., Allan, R., Aguilar, E., Barry, A. A., Brunet, M., Caesar, J., Chagnaud, G., Cheng, V., Cinco, T., Durre, I., Guzman, R. de, Htay, T. M., Ibadullah, W. M. W., Ibrahim, M. K. I. B., Khoshkam, M., Kruger, A., Kubota, H., Leng, T. W., Lim, G., Li-Sha, L., Marengo, J., Mbatha, S., McGree, S., Menne, M., Skansi, M. de los M., Ngwenya, S., Nkrumah, F., Oonariya, C., Pabon-Caicedo, J. D.,
- 365 Panthou, G., Pham, C., Rahimzadeh, F., Ramos, A., Salgado, E., Salinger, J., Sané, Y., Sopaheluwakan, A., Srivastava, A., Sun, Y., Timbal, B., Trachow, N., Trewin, B., Schrier, G. van der, Vazquez-Aguirre, J., Vasquez, R., Villarroel, C., Vincent, L., Vischel, T., Vose, R., and Yussof, M. N. B. H.: Development of an Updated Global Land In Situ-Based Data Set of Temperature and Precipitation Extremes: HadEX3, *J. Geophys. Res. Atmospheres*, 125, e2019JD032263, <https://doi.org/10.1029/2019JD032263>, 2020.
- 370 [European Standard: ISO 15927-4 Hygrothermal performance of buildings — calculation and presentation of climatic data — Part 4: Hourly data for assessing the annual energy use for heating and cooling, 2005. European Standard: EN ISO 15927-](https://www.iso.org/standard/54548.html)

4:2005 Hygrothermal performance of buildings — calculation and presentation of climatic data — Part 4: Hourly data for assessing the annual energy use for heating and cooling (ISO 15927-4:2005)

- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Eyring, V., N.P. Gillett, K.M. Achuta Rao, R. Barimalala, M. Barreiro Parrillo, N. Bellouin, C. Cassou, P.J. Durack, Y. Kosaka, S. McGregor, S. Min, O. Morgenstern, Y. Sun, and Eyring, V., N.P. Gillett, K.M. Achuta Rao, R. Barimalala, M. Barreiro Parrillo, N. Bellouin, C. Cassou, P.J. Durack, Y.: Human Influence on the Climate System., in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)], 2021.
- Ferrari, D. and Lee, T.: Beyond TMY: Climate data for specific applications, in: *Proceedings 3rd International Solar Energy Society conference—Asia Pacific region (ISES-AP-08)*, 12, 2008.
- Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., and Gallée, H.: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, 7, 469–489, <https://doi.org/10.5194/tc-7-469-2013>, 2013.
- Fettweis X., Wyard C., Doutreloup S., and Belleflamme A.: Noël 2010 en Belgique: neige en Flandre et pluie en Haute-Ardenne, *Bull. Société Géographique Liège*, 68, 97–107, 2017.
- Finkelstein, J. M. and Schafer, R. E.: Improved goodness-of-fit tests, *Biometrika*, 58, 641–645, <https://doi.org/10.1093/biomet/58.3.641>, 1971.
- Fouillet, A., Rey, G., Laurent, F., Pavillon, G., Bellec, S., Guihenneuc-Jouyau, C., Clavel, J., Jouglu, E., and Hémon, D.: Excess mortality related to the August 2003 heat wave in France, *Int. Arch. Occup. Environ. Health*, 80, 16–24, <https://doi.org/10.1007/s00420-006-0089-4>, 2006.
- Gutjahr, O., Putrasahan, D., Lohmann, K., Jungclaus, J. H., von Storch, J.-S., Brüggemann, N., Haak, H., and Stössel, A.: Max Planck Institute Earth System Model (MPI-ESM1.2) for the High-Resolution Model Intercomparison Project (HighResMIP), *Geosci. Model Dev.*, 12, 3241–3281, <https://doi.org/10.5194/gmd-12-3241-2019>, 2019.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G. D., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P. de, Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- IPCC, Masson Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I.


- Gomis, M., and Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2021.
- 410 Kittel, C.: Present and future sensitivity of the Antarctic surface mass balance to oceanic and atmospheric forcings: insights with the regional climate model MAR, PhD Thesis, Université de Liège, Liège, Belgique, Liège, 2021.
- Kittel, C., Amory, C., Agosta, C., Jourdain, N. C., Hofer, S., Delhasse, A., Doutreloup, S., Huot, P.-V., Lang, C., Fichet, T., and Fettweis, X.: Diverging future surface mass balance between the Antarctic ice shelves and grounded ice sheet, *The Cryosphere*, 15, 1215–1236, <https://doi.org/10.5194/tc-15-1215-2021>, 2021.
- 415 Larsen, M. A. D., Petrović, S., Radoszynski, A. M., McKenna, R., and Balyk, O.: Climate change impacts on trends and extremes in future heating and cooling demands over Europe, *Energy Build.*, 226, 110397, <https://doi.org/10.1016/j.enbuild.2020.110397>, 2020.
- Lee, J. Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J. P. Dunne, F. Engelbrecht, E. Fischer, J. C. Fyfe, C. Jones, A., and Maycock, J. Mutemi, O. Ndiaye, S. Panickal, T. Zhou: Future Global Climate: Scenario-Based Projections and Near-Term Information, *Camb. Univ. Press, Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the*
- 420 *Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)], 2021.
- Ouzeau, G., Soubeyroux, J.-M., Schneider, M., Vautard, R., and Planton, S.: Heat waves analysis over France in present and future climate: Application of a new method on the EURO-CORDEX ensemble, *Clim. Serv.*, 4, 1–12, <https://doi.org/10.1016/j.cliser.2016.09.002>, 2016.
- 425 <https://doi.org/10.1016/j.cliser.2016.09.002>, 2016.
- [Pérez-Andreu V., Aparicio-Fernández C., Martínez-Ibernón A. and Vivancos J.-L.: Impact of climate change on heating and cooling energy demand in a residential building in a Mediterranean climate, *Energy*, vol. 165, pp. 63–74, Dec. 2018, doi: 10.1016/j.energy.2018.09.015.](https://doi.org/10.1016/j.energy.2018.09.015)
- Ramon, D., Allacker, K., van Lipzig, N. P. M., De Troyer, F., and Wouters, H.: Future Weather Data for Dynamic Building
- 430 Energy Simulations: Overview of Available Data and Presentation of Newly Derived Data for Belgium, in: *Energy Sustainability in Built and Urban Environments*, edited by: Motoasca, E., Agarwal, A. K., and Breesch, H., Springer, Singapore, 111–138, https://doi.org/10.1007/978-981-13-3284-5_6, 2019.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K.,
- 435 Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Glob. Environ. Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.

- 440 Ridder, K. D. and Gallée, H.: Land Surface–Induced Regional Climate Change in Southern Israel, *J. Appl. Meteorol. Climatol.*, 37, 1470–1485, [https://doi.org/10.1175/1520-0450\(1998\)037<1470:LSIRCC>2.0.CO;2](https://doi.org/10.1175/1520-0450(1998)037<1470:LSIRCC>2.0.CO;2), 1998.
- RMI: Rapport climatique 2020 de l’information aux services climatiques, Royal Meteorological Institute of Belgium, Brussels, 2020.
- Seneviratne, S. I., Zhang X., M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S., and
- 445 Lewis, F. Otto, I. Pinto, M. Satoh, S. M. Vicente-Serrano, M. Wehner, B. Zhou: Weather and Climate Extreme Events in a Changing Climate, *Clim. Change 2021 Phys. Sci. Basis Contrib. Work. Group Sixth Assess. Rep. Intergov. Panel Clim. Change Masson-Delmotte V P Zhai Pirani Connors C Péan Berg. N Caud Chen Goldfarb M Gomis M Huang K Leitzell E Lonnoy J B R Matthews T K Maycock T Waterfield O Yelekçi R Yu B Zhou Eds*, 2021.
- Sherwood, S. C. and Huber, M.: An adaptability limit to climate change due to heat stress, *Proc. Natl. Acad. Sci.*, 107, 9552–
- 450 9555, <https://doi.org/10.1073/pnas.0913352107>, 2010.
- Suarez-Gutierrez, L., Müller, W. A., Li, C., and Marotzke, J.: Hotspots of extreme heat under global warming, *Clim. Dyn.*, 55, 429–447, <https://doi.org/10.1007/s00382-020-05263-w>, 2020.
- Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O’ishi, R., Yamazaki, D.,
- 455 Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geosci. Model Dev.*, 12, 2727–2765, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.
- Termonia, P., Van Schaeybroeck, B., De Cruz, L., De Troch, R., Caluwaerts, S., Giot, O., Hamdi, R., Vannitsem, S., Duchêne, F., Willems, P., Tabari, H., Van Uytven, E., Hosseinzadehtalaei, P., Van Lipzig, N., Wouters, H., Vanden
- 460 Broucke, S., van Ypersele, J.-P., Marbaix, P., Villanueva-Birriel, C., Fettweis, X., Wyard, C., Scholzen, C., Doutreloup, S., De Ridder, K., Gobin, A., Lauwaet, D., Stavrakou, T., Bauwens, M., Müller, J.-F., Luyten, P., Ponsar, S., Van den Eynde, D., and Pottiaux, E.: The CORDEX.be initiative as a foundation for climate services in Belgium, *Clim. Serv.*, 11, 49–61, <https://doi.org/10.1016/j.cliser.2018.05.001>, 2018.
- Tsachoua, L. and Reynders, D.: Hittegolf en Ozonpiekenplan,
- 465 https://www.belspo.be/belspo/organisation/publ/pub_ostc/agora/ragjj146_en.pdf, 2016.
- Wilcox, S. and Marion, W.: Users Manual for TMY3 Data Sets, 2008.
- Wu, T., Lu, Y., Fang, Y., Xin, X., Li, L., Li, W., Jie, W., Zhang, J., Liu, Y., Zhang, L., Zhang, F., Zhang, Y., Wu, F., Li, J., Chu, M., Wang, Z., Shi, X., Liu, X., Wei, M., Huang, A., Zhang, Y., and Liu, X.: The Beijing Climate Center Climate System Model (BCC-CSM): the main progress from CMIP5 to CMIP6, *Geosci. Model Dev.*, 12, 1573–1600,
- 470 <https://doi.org/10.5194/gmd-12-1573-2019>, 2019.
- Wyard, C., Scholzen, C., Doutreloup, S., Hallot, É., and Fettweis, X.: Future evolution of the hydroclimatic conditions favouring floods in the south-east of Belgium by 2100 using a regional climate model, *Int. J. Climatol.*, 41, 647–662, <https://doi.org/10.1002/joc.6642>, 2021.




475


Figure 1: Workflow of the MAR model. The MAR model (black box) needs to be forced by a reanalysis model or global model (upper orange box) and by static data (lower orange box) and the result of the MAR simulation gives meteorological variables in two or three dimensions (yellow box).




480 **Figure 2: Model topography (color background, in meters), localization of cities (black) used in this dataset for Belgium, and localization of neighboring countries (in red).**



485 **Figure 3: Annual mean temperature (in C, lower lines) and annual summer temperature (in C, upper lines) of MAR forced by ERA5 reanalysis (in black bold lines), MAR forced by BCC-CSM2-MR (in red lines), MAR forced by MPI-ESM1.2 (in blue lines), and MAR forced by MIROC6 (in green lines) between 1980 and 2100 according to the scenario SSP5-8.5. The average is computed here over the whole integration domain excluding the ocean.**




490 **Figure 4 (figure and legend coming from Ouzeau et al., 2016): Characterization of a heatwave from a daily mean temperature indicator over France (example of a time series from June 30 to August 5, 1983): Duration (start and end), maximal temperature, and global intensity (red area of the plot). Temperatures above (below) climatological line (1981–2010 reference period) are represented by the pink (blue) area. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)**



495 **Figure A-1: Typical (black) and extreme (red) temperature for Liege city simulated by MAR forced by BCC-CSM2-MR with the scenario *sspSSP5-8.5* for the period 2085-2100. The typical/extreme months are based on temperature (left) and incident solar radiation (right). These data are extracted from these files :**

- Liege-City_TMY2085-2100_ssp585_MAR-BCC_TTbased.csv
- Liege-City_XMY2085-2100_ssp585_MAR-BCC_TTbased.csv
- Liege-City_TMY2085-2100_ssp585_MAR-BCC_SWDbased.csv
- 500 - Liege-City_XMY2085-2100_ssp585_MAR-BCC_SWDbased.csv



505 **Figure A-2: Temperature during warmest heatwave events for Liege city simulated by MAR-BCC (i.e., MAR forced by BCC-CSM2-MR), MAR-MPI (i.e., MAR forced by MPI-ESM.1.2), and MAR-MIR (i.e., MAR forced by MIROC6) with the scenario *sspSSP5-8.5* for the period 2085-2100. These data are extracted from these files :**

- Liege-City_HWE-HT_2085-2100_ssp585_MAR-BCC.csv

- Liege-City_HWE-HT_2085-2100_ssp585_MAR-MPI.csv
- Liege-City_HWE-HT_2085-2100_ssp585_MAR-MIR.csv

1980-2014 over Belgium territory	Time scale	MAR-ERA5	MAR-BCC	MAR-MPI	MAR-MIR
Mean annual temperature at 2m a.g.l. (in C)	annual	10.1 +/- 6.2	10.3 +/- 6.0	10.1 +/- 6.4	10.4 +/- 7.2
	summer	17.7 +/- 1.3	17.6 +/- 1.7	18.6 +/- 1.2	19.9 +/- 1.1
Mean incident solar radiation (in W/m ²)	annual	119 +/- 78	122 +/- 79	116 +/- 83	134 +/- 85
	summer	213 +/- 20	221 +/- 20	224 +/- 22	239 +/- 25

510 Table 1: Comparison of MAR-ERA5 (i.e., MAR forced by the reanalysis ERA5) with MAR-BCC (i.e., MAR forced by BCC-CSM2-MR), MAR-MPI (i.e., MAR forced by MPI-ESM1.2), and MAR-MIR (i.e., MAR forced by MIROC6) to mean temperature at 2m **above ground level** and mean incident solar radiation over Belgium territory during 1980-2014 (**mean value +/- standard error**).

<u>Weather variable name</u>	<u>Level</u>	<u>Units</u>
<u>Dry bulb temperature</u>	<u>2m a.g.l.</u>	<u>°C</u>
<u>Relative humidity</u>	<u>2m a.g.l.</u>	<u>%</u>
<u>Global horizontal radiation</u>	<u>Ground (horizontal surface)</u>	<u>Wh/m²</u>
<u>Diffuse solar radiation</u>	<u>Ground (horizontal surface)</u>	<u>Wh/m²</u>
<u>Direct normal radiation</u>	<u>Ground (horizontal surface)</u>	<u>Wh/m²</u>
<u>Wind speed</u>	<u>10m a.g.l.</u>	<u>m/s</u>
<u>Wind direction</u>	<u>10m a.g.l.</u>	<u>north degrees</u>
<u>Dew point temperature</u>	<u>2m a.g.l.</u>	<u>°C</u>
<u>Atmospheric Pressure</u>	<u>ground</u>	<u>Pa</u>
<u>Cloudiness</u>	<u>All troposphere</u>	<u>tenths</u>
<u>Sky temperature</u>	<u>All troposphere</u>	<u>K</u>

(according to (Duffie and Beckman,

<u>Specific humidity</u>	<u>2m a.g.l.</u>	<u>2013)</u> <u>kg/kg</u>
<u>Precipitation</u>	<u>Ground</u>	<u>mm</u>

515 Table 2: name, level and units of all weather variables used in the produced weather files.