
New gridded dataset of rainfall erosivity (1950–2020) on the Tibetan Plateau

Yueli Chen¹, Xingwu Duan², Minghu Ding¹, Wei Qi¹, Ting Wei¹, Jianduo Li^{3,1}, Yun Xie⁴

5 ¹State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

²Institute of International Rivers and Eco-security, Yunnan University, Kunming, 650091, China

³CMA Earth System Modeling and Prediction Centre, Beijing, 100081, China

10 ⁴State Key Laboratory of Earth Surface Processes and Resources Ecology, Faculty of Geographic Science, Beijing Normal University, Beijing, 100875, China

Correspondence to: Xingwu Duan (xwduan@ynu.edu.cn), Minghu Ding (dingminghu@foxmail.com)

Abstract. The risk of water erosion on the Tibetan Plateau (TP), a typical fragile ecological area, is increasing with climate change. Rainfall erosivity map is useful for understanding the spatial-temporal pattern of rainfall erosivity and identifying hot spots of soil erosion. This study generate an annual gridded rainfall erosivity dataset on a 0.25° grid for the TP in 1950–2020. 1-min precipitation observations at 1787 weather stations for 7 years and 0.25° hourly European Center for Medium-Range Weather Forecasts Reanalysis 5 (ERA5) precipitation data for 71 years are employed in this study. Our results indicate that the ERA5-based estimates have a marked tendency to underestimate annual rainfall erosivity when compared to the station-based estimates, because of the systematically biases of ERA5 precipitation data including the large underestimation of the maximum contiguous 30-min peak intensity and relatively slight overestimation of event erosive precipitation amount. The multiplier factor map over the TP, which was generated by Inverse Distance Weighted method based on the relative changes between the available station-based annual rainfall erosivity grid values and the corresponding ERA5-based values, was employed to correct the ERA5-based annual rainfall erosivity and then reconstruct the annual rainfall erosivity dataset. The multi-year average correction coefficient over the TP between the station-based annual rainfall erosivity values and the newly released data is 0.67. In addition, the probability density and various quantile values of the new data are generally consistent with the station-based values. The data offers a view of large-scale spatial-temporal

30 variability in the rainfall erosivity and addresses the growing need for the information to predict rainfall-induced hazards over the TP. The dataset are available at <http://data.tpsc.ac.cn/en/data/37c34046-3c2a-4737-b3c9-35af398da62a/>.

1 Introduction

Precipitation is the main driver of water erosion because it directly affects the detachment of soil particles, breakdown of aggregates, and transport of eroded particles via runoff (Wischmeier and Smith, 1965, 1978). The *R* factor, that is, the multi-year average rainfall erosivity, which is described by the Universal Soil Loss Equation (USLE; Wischmeier and Smith, 1965, 1978) and Revised USLE (RUSLE; Renard, 1997), is an indicator of the potential ability of rainfall and runoff to affect soil erosion. Generally, the *R* factor is calculated by using the classical (Wischmeier and Smith, 1965) or statistical algorithms (e.g., Liu et al., 2002) according to the temporal resolution of the available precipitation data.

The classical algorithm for rainfall erosivity requires a continuous precipitation data series with <15-min temporal resolution (Angulo-Martínez and Beguería, 2009). As networks of weather stations and observation platforms have matured considerably in the past two decades, rainfall erosivity has been calculated using the classical algorithm at the local scale (Agnese et al., 2006; Ma et al., 2014; Wang et al., 2017), and the application of the algorithm has been gradually extended to the national (Panagos et al., 2015; Kim et al., 2020; Yue et al., 2021) and global scale (Panagos et al., 2017; Liu et al., 2020). Despite substantial progress, it is still notable that the relative error of the estimated rainfall erosivity increases rapidly with increasing time interval of the precipitation data. For example, the relative error based on hourly data was more than 80%, compared with the results based on 1-min data (Lobo and Bonilla, 2015; Yin et al., 2015; Shin et al., 2019). In addition, the accuracy of the rainfall erosivity map is greatly reduced by inadequate weather station coverage, especially in areas with complex climates and terrains (Yue et al., 2021). Therefore, the accuracy of rainfall erosivity estimation depends strongly on both of the temporal and spatial resolution of the precipitation data (Panagos et al., 2017; Kim et al., 2020).

Compared with in-situ observations, gridded precipitation data (e.g. satellite-based, reanalysis and fused datasets) are not subjected to topographical limitations and could supply continuous precipitation data (Beck et al., 2017). These data have been widely used to estimate the rainfall erosivity in China, especially in the regions with scarce in-situ observations (Teng et al., 2018), Germany (Risal et al., 2018), Africa (Vrieling et al., 2010), the United States (Kim et al., 2020), and other regions. They have contributed greatly to our knowledge of the spatiotemporal patterns of rainfall erosivity; however, the uncertainties in rainfall erosivity directly calculated by using gridded precipitation data have not been

quantified, although obvious biases between gridded and observed precipitation values have been demonstrated (Freitas et al., 2020).

65 The Tibetan Plateau (TP) referred to as the Third Pole is one of the highest plateaus worldwide and has an average altitude of more than 4000 m (Yao et al., 2012). Since the mid-1950s, the TP has experienced significant warming exceeding that of other regions in the same latitude zone (Liu and Chen, 2000). Owing to increasing snowmelt and more frequent heavy precipitation events, which may cause more severe soil erosion, knowledge of the rainfall erosivity on the TP is highly important for
70 soil sustainability and thus water and food security. The accuracy of rainfall erosivity estimation depends mainly on the spatiotemporal accuracy of the precipitation data, especially on the TP, where the seasonal and regional precipitation patterns exhibit significant variability owing to westerly winds, the Indian monsoon, and land–atmosphere interaction.

Many efforts have been made to study the rainfall erosivity on the TP (Table 1). Most studies employed
75 the empirical methods, however, our study has demonstrated that these empirical methods always resulted to obvious biases over the TP, when compared with the values based on the 1-min precipitation data by using the standard method (paper submitted). In the term of the type of the precipitation data, dozens of station-based precipitation data were commonly used to calculate the rainfall erosivity (e.g., Qin et al., 2016; Gu et al., 2020). Yue et al. (2022) has reported that the scarce weather stations can
80 significantly reduce the estimation accuracy of the rainfall erosivity in the regions with complex terrain and climate, especially on the TP. Therefore, the accuracy of the estimated rainfall erosivity on the TP are largely reduced by the current empirical estimation models and the scarcity of the historical weather stations. In other words, the precipitation data with high spatial-temporal resolution are essential to calculate the rainfall erosivity on the TP.

Table 1. *R* factor of TP in previous studies

Study scale	Type of precipitation data	Number of weather stations	Temporal resolution	Calculation Method	Spatial characteristics	<i>R</i> factor (MJ·mm·ha ⁻¹ ·h ⁻¹ ·yr ⁻¹)	Reference
China	Weather station	China: 2381 TP: < 100	1 hour	Standard	Kriging	TP: 273	Yue et al., 2022
Southwestern China	CRU_TS4		Monthly	Empirical	Grid, no bias correction	Tibet: 3407	Cao et al., 2018
China	TRMM, weather station	China: 650 TP: < 50	Daily	Empirical	Grid, bias correction of TRMM precipitation	No value	Teng et al., 2017
China	Weather station	China: 756 TP: < 50	Daily	Empirical	Kriging	TP: 408	Qin et al., 2016
China's dryland region	Weather station	China's dryland region 298	Daily	Empirical	HASM interpolation	Most of TP: 1–500	Yang et al., 2015
China	Weather station	China: 590 TP: < 50	Daily	Empirical	Kriging	Central and eastern TP: 147	Liu et al., 2013
China	Weather station	China: 564	Daily	Empirical	Kriging	Cold zone of TP: 368 Sub-cold zone of TP: 427	Zhang et al., 2003
Tibet	Weather station	Tibet: 38	Daily	Empirical	Station-averaged	Tibet: 714	Gu et al., 2020
Tibet	TRMM 3B42 gridded		Daily	Empirical	Grid, no bias correction	Tibet: 768	Yan et al., 2010

Note: CRU_TS4: Climatic Research Unit Time Series 4 gridded precipitation product. TRMM: Tropical Rainfall Measuring Mission gridded precipitation product. Empirical method means the rainfall erosivity values are calculated by using the empirical equations based on daily or monthly precipitation data. The standard method is proposed by USLE or RUSLE. The boundaries of the TP used in these studies has some slight differences.

To expand the spatial coverage and extend the time series of rainfall erosivity over the TP, the various
90 gridded precipitation datasets, for example, satellite-based Tropical Rainfall Measuring Mission
(TRMM) and station-based Climatic Research Unit Time Series 4 gridded precipitation datasets
(CRU_TS4), are also introduced into the soil erosion study of the TP in recent decade (Yan et al., 2010;
Teng et al., 2017; Gao et al., 2018). The performances of these gridded precipitation mainly depend on
the spatial-temporal accuracy of the gauge observations, and thus these datasets always present obvious
95 biases, due to insufficient density of the weather station network over the TP (Yuan et al., 2021). It is
notable that the significant biases of various gridded precipitation data have also been widely identified
(Sun et al., 2018), and their impacts on the rainfall erosivity estimation have not been assessed.

In recent researches, the model-based gridded precipitation datasets begin to be concerned (Li et al.,
2020; Zhou et al., 2021), because they could resolve the complex topography and climate effects over
100 the TP and provide long-term data by setting simulation period. The European Center for
Medium-Range Weather Forecasts Reanalysis 5 (ERA5) as the newly generation is one of the most
widely used precipitation datasets in the world (Hersbach et al. 2019). Compared with other gridded
precipitation datasets, ERA5 succeeded in reproducing the inter-annual and decadal variabilities of
precipitation and reflecting the spatial-temporal patterns (Yuan et al., 2021), and performed marginally
105 better in detecting daily precipitation over the whole TP for the long-term periods (Jiang et al., 2021),
despite the bias in precipitation amount was also reported (Jiang et al., 2021; Jiao et al., 2021).
Therefore, this study aims to reconstruct the historical annual rainfall erosivity with 0.25° spatial
resolution in 1950–2020 over the TP, by employing the 0.25° hourly ERA5 precipitation data for 71
years to generate a long-term background values and utilizing the 1-min precipitation observations at
110 1787 weather stations for 7 years to identify and correct the biases of the estimates. In detail, this paper
describes (1) the performance of ERA5 precipitation data at the weather stations; (2) the performance
of the ERA5-based annual rainfall erosivity calculated by using the standard method recommended by
the USLE model; (3) the correction of the ERA5-based annual rainfall erosivity and the validation of
the newly generated dataset.

115 **2 Study Area and Source Data**

2.1 Tibetan Plateau

The study area is the TP (26–40°N, 73–105°E), which is located in Southwestern China and covers an

area of approximately 2.5 million km². The elevation of the TP ranges from 84 to 8246 m, with an average value of 4379 m. Precipitation in the southeastern TP is influenced by warm, humid Indian monsoons, whereas in the western TP, it is influenced more strongly by the mid-latitude westerlies (Yao et al., 2012). The annual precipitation is concentrated from May to October (Gu et al., 2020), and shows a spatial pattern of a wet east and west with a dry middle (Li et al., 2020). Along with the significant climate change and a very fragile ecological environment, the TP has high potential for soil loss, especially in the eastern TP and Hengduan Mountains, which are among the most severely eroded areas in China (Teng et al., 2019).

2.2 Precipitation data

Previous studies of the TP have used in-situ precipitation observations with <50 stations and coarse temporal resolution, e.g., hourly (Yue et al., 2021), daily (Wang et al., 2017), or half-monthly (Teng et al., 2018; Gu et al., 2020; Liu et al., 2020). By contrast, this study estimated the rainfall erosivity on the TP using precipitation observations at 1-min intervals in 2013–2020 at 1787 weather stations obtained from the National Meteorology Information Center of the China Meteorological Administration [Figure 1(a)].

To ensure the accuracy of the in situ precipitation data, we evaluated their quality. The data integrity of each station was first checked using quality control codes at 1-min intervals by month. Because precipitation on the TP occurs mainly from May to September, observed data with an integrity of >90% from May to September in a year can be used to calculate the annual rainfall at the station. The number of stations with data suitable for calculating the annual rainfall erosivity for each year is shown in the lower left corner of Figure 1(a); it ranges from 628 to 1472, with an average of 1114 stations for 2013–2020 (excluding 2017, because a disruption in data reception caused the loss of precipitation observations in August 2017). Moreover, we examined the station density in each 0.25° grid, which is consistent with the spatial resolution of the ERA5 data [Figure 1(b)]. The number of stations in each grid varies from 1 to 29, and the mean value is 2.1. A total of 836 grids (20% of the grids covering the TP) have observed precipitation values. Because the data quality varies, the available grids with observations change annually; on average, there are 589 available grids with observation records for 2013–2020, excluding 2017.

The hourly 0.25° ERA5 data represent the most recent generation of ECMWF global atmospheric reanalysis and offer higher spatial resolution than ERA-Interim and other improvements since 1979 (Hersbach et al., 2019). The precipitation data are the sum of large-scale precipitation and convective precipitation consisting of rain and snow, as determined by the ECMWF Integrated Forecasting System.




Figure 1. (a) Spatial distribution of weather stations on TP; the inset shows the number of available weather stations by year. (b) Number of available weather stations in each grid with 0.25° spatial resolution; the inset shows the number of available weather stations by year.

3 Methodology

Figure 2 shows the overall algorithm for generating the annual rainfall erosivity dataset with the 0.25° spatial resolution over the TP in 1950–2020. We firstly calculated the annual rainfall erosivity by using the standard method of rainfall erosivity based on the 1-min in-situ precipitation observations and 0.25° hourly ERA5 precipitation data, respectively. Secondly, the performances of the ERA5 were systematically assessed in the terms of the detecting accuracy of the precipitation for erosive events and the estimation accuracy of ERA5-based annual rainfall erosivity. Finally, the historical annual rainfall erosivity data for the TP was produced after correcting the ERA5-based annual rainfall erosivity.




Figure 2. Schematic representation of algorithm for generating annual rainfall erosivity dataset for 1950–2020. r_{obs_year} and r_{ERA5_year} represent the station-based and ERA5-based annual rainfall erosivity values, respectively. r_{cor_year} means the corrected values of the r_{ERA5_year} by using the multiplier factor map.

170

3.1 Algorithm of annual rainfall erosivity

A rainfall event is defined following Wischmeier and Smith (1978) as having measurable rainfall with no interruption or at most a 6-h interruption. If a rainfall event is interrupted for more than 6 h, subsequent rainfall is considered to belong to a new rainfall event. Rainfall events of more than 12 mm are selected as erosive events following Xie et al. (2000), and the EI_{30} index of the erosive event is calculated. Specifically, the rainfall erosivity of an erosive rainfall event is calculated as follows (Brown and Foster, 1987):

$$e_r = 0.29[1 - 0.72\exp(-0.05i_r)] \quad (1)$$

$$E = \sum_{r=1}^n (e_r \cdot P_r) \quad (2)$$

$$r_{event} = E \cdot I_{30} \quad (3)$$

180

where E ($\text{MJ} \cdot \text{ha}^{-1}$) is the total energy of the erosive event, and r_{event} ($\text{MJ} \cdot \text{mm} \cdot \text{ha}^{-1} \cdot \text{h}^{-1}$) is the event rainfall erosivity of the event. For the 1-min precipitation data (ERA5 data), i_r (mm/h) is the rainfall intensity for the r^{th} minute (hour), e_r ($\text{MJ} \cdot \text{ha}^{-1} \cdot \text{mm}^{-1}$) is the unit energy for the r^{th} minute (hour), P_r

(mm) is the rainfall amount for the r^{th} minute (hour), n is the rainfall duration, and I_{30} (mm/h) is the maximum contiguous 30-min (1-h) peak intensity. After the event rainfall erosivity at all stations was calculated, we identified and removed extreme outliers of the event rainfall erosivity at each site, which resulted from temporary abnormalities in the automatic observation equipment and were not identified during quality control of the precipitation data. We used boxplots to detect extreme outliers. The lower and upper quartiles were defined as the 25th percentile of event rainfall erosivity (Q1) and the 75th percentile (Q2); the difference (Q2 – Q1) is called the interquartile range (IQR). Event rainfall erosivity data at a station outside the lower and upper bounds (Q1 – 3IQR, Q2 + 3IQR) are considered extreme outliers.

The observed annual rainfall erosivity values ($r_{\text{station_year}}$) were obtained by summing the rainfall erosivity for all erosive events per year by station. Next, the ERA5-based annual rainfall erosivity ($r_{\text{ERA5_year}}$) for all the grids in the TP were calculated. Notably, for easy comparison of $r_{\text{station_year}}$ and $r_{\text{ERA5_year}}$, the $r_{\text{station_year}}$ values were upscaled to the grid values ($r_{\text{obs_year}}$) with 0.25° spatial resolution by averaging the station-based values in the same grid. Figure 1(b) shows the spatial distribution of the available grids with $r_{\text{obs_year}}$. Steps 2 to 3 in Figure 2 are all based on $r_{\text{obs_year}}$ and $r_{\text{ERA5_year}}$ data at grid scale.

3.2 Assessment of the performance of the ERA5 precipitation data

The performance of the ERA5 precipitation data were assessed at 280 grid cells, which corresponded to 7% of all the grids over the TP. Given the importance of erosive rainfall events to soil erosion, we focused on the performance of the ERA5 precipitation data in detecting characteristics of erosive rainfall event, including multi-year averaged annual erosive precipitation amount and frequency, and mean erosive event precipitation amount and I_{30} .

The mean values of $r_{\text{ERA5_year}}$ for 2013–2020 were compared with those of $r_{\text{obs_year}}$ by station. The absolute bias (AB) and correction coefficient (r) were used to evaluate the accuracy of annual rainfall erosivity estimation using ERA5 data. The AB is calculated as shown in Eq. 4.

$$AB = \sum_{i=1}^n (r_{\text{ERA5_year}_i} - r_{\text{obs_year}_i}) / n \quad (4)$$

where i is the i^{th} annual rainfall erosivity value, $r_{\text{ERA5_year}_i}$ is the ERA5-based annual rainfall erosivity in the i^{th} year, $r_{\text{obs_year}_i}$ is the observed annual rainfall erosivity in the i^{th} year, and n is the number of years of data. Moreover, the empirical orthogonal function (EOF) was employed to assess the

spatiotemporal pattern of annual rainfall erosivity revealed by the ERA5 reanalysis precipitation data by comparing it with the pattern revealed by the observed values.

215 3.3 Reconstruction and validation of annual rainfall erosivity

For the soil erosion process, it is known that not all the precipitation events but the erosive events have close relationship with the water erosion process. Our study indicated the precipitation characteristics derived from ERA5 data for erosive events showed high correction with those from in-situ precipitation observations over the TP (Figure 4). In addition, there was a high correlation between the station-based
220 and ERA5-based annual rainfall erosivity (Figure 5), and their spatiotemporal distribution patterns also showed well agreement (Figure 7). These findings have demonstrated that it is reasonable to generate the rainfall erosivity dataset for the TP by using the ERA5 precipitation data, and meanwhile, the correction is also essential because of the obvious biases identified in the ERA5-based rainfall erosivity values.

225 Relative changes between the in-situ and modeled precipitation are always used to correct the modeled precipitation for accuracy improvement, such as the global precipitation data from WorldClim (Fick et al., 2017), the gridded precipitation data of the China Meteorological Forcing Dataset (He et al., 2020) and the bias adjusted ERA5 precipitation data (Cucchi et al., 2020). Given the close correlation between the precipitation and rainfall erosivity, the relative changes were also employed to correct the
230 ERA5-based annual rainfall erosivity in this study. Here, we have used a hypothesis that the bias of the ERA5-based annual rainfall erosivity resulted from ERA5 precipitation data at each grid keeps steady by year. In detail, the correction process can be divided into three steps. Firstly, the r_{obs_year} values were divided by r_{ERA5_year} for each year, and then the calculated results, i.e., the multiplier factor values, were averaged for years. Secondly, inverse distance weighted (IDW) interpolation was used to generate a
235 multiplier factor map of the TP with 0.25° spatial resolution. Thirdly, the corrected annual rainfall erosivity dataset (r_{cor_year}) was obtained as the product of r_{ERA5_year} and the multiplier factor for each grid.

Specifically, there are 373 grids with observed annual rainfall erosivity values from 2014 to 2020. The r_{obs_year} and r_{ERA5_year} values in these grids were used to generate the multiplier factor map. The r_{obs_year}
240 and r_{ERA5_year} values in other grids for 2014–2020, which were not used before, are available for assessing the accuracy. Moreover, the year 2013 was regarded as a complete verification year, in which

the assessment of the $r_{\text{cor_year}}$ was conducted in all the TP grids with $r_{\text{obs_year}}$ values. Table 2 lists the number of validation grids at each year, and Figure 3 shows the spatial distribution of the validation grids for 2013–2020 (excluding 2017).

245

Table 2. Numbers of grids used in this study

Year	Total number of grids with observations	Number of validation grids	Percentage of validation data in total data (%)
2013	381	381	100
2014	477	104	22
2015	504	131	26
2016	562	189	34
2018	712	339	48
2019	745	372	50
2020	742	369	50




Figure 3. Spatial distribution of validation grids covering the TP for 2013–2020 (excluding 2017).

250

4 Results

4.1 Detecting accuracy of ERA5 for erosive rainfall events

Figure 4 compared the multi-year average annual erosive precipitation amount and frequency, and mean erosive event precipitation amount and I_{30} derived from ERA5 precipitation data with those from
 255 in-situ observations. In detail, the ERA5-based multi-year average annual erosive precipitation amount is three times more than the station-based value across the TP. The overestimation of the multi-year average annual precipitation amount was also reported by Jiao et al. (2021). The ERA5 overestimated the annual erosive precipitation frequency by 1.6 times. For the erosive event rainfall amount, ERA5 was almost twice as much as the station-based value, which differed from the finding of Jiao et al.
 260 (2021) that the daily precipitation amount with more than 10 mm are underestimated by ERA5. This result demonstrates that the erosive rainfall events in the TP cannot be simply equivalent to the daily

precipitation events (Chen et al., 2022). In addition, the mean I_{30} of ERA5 for erosive events are only one ninth of the station-based value. Because the relatively slight overestimation of ERA5 precipitation data in the erosive event precipitation amount could not offset the substantial underestimation in I_{30} , the ERA5-based estimates showed a marked tendency to underestimate the rainfall erosivity when compared to the station-based estimates. Overall, the comparison between the two data sources indicated that there were significant biases of ERA5 data in detecting precipitation characteristics for erosive events in the TP, however, also presented high correlations with correlation coefficient being 0.33 – 0.84.





Figure 4. Scatterplots of the station-based multi-year average (a) annual erosive precipitation amount (Ptot), (b) annual erosive precipitation frequency (PF), (c) mean erosive event precipitation amount (Pevent), (d) mean I_{30} for erosive events (I_{30}) vs those derived from ERA5 data at the corresponding grid cells in 2013–2020.

4.2 Evaluation of rainfall erosivity estimation using ERA5 data

The accuracy of annual rainfall erosivity estimation using the ERA5 precipitation data for 2013–2020

was assessed and compared with the $r_{\text{obs_year}}$ values in 280 grids covering the TP. The correlation coefficient of the mean annual rainfall erosivity based on the observed and ERA5 precipitation data is 0.71. For most stations, the ERA5-based values were significantly underestimated (Figure 5).




280

Figure 5. Comparison of mean annual rainfall erosivity based on observed and ERA5-based results for seven years (2013–2020, excluding 2017). The dotted line is the result of an optimal model (with an intercept of 0 and regression coefficient of 1). The red line is the regression result. Colors of dots represent the grid density. Unit: $\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$.

285

To further evaluate the quality of mean annual rainfall erosivity estimation using ERA5 data, the performance of the ERA5 data in each grid was evaluated, as shown in Figure 6. The spatial pattern of the ERA5-based mean annual rainfall erosivity is consistent with that of the observed values. Specifically, areas with large annual rainfall erosivity are located mainly in the southeastern part of the plateau, especially at the southeast edge, whereas the mean annual values in the northwestern part of the plateau are relatively small. However, the observed mean annual rainfall erosivity on the TP is $344 \text{ MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$, and the ERA5-based results underestimate this value by 47%. Moreover, except for most of the grids in the northwest corner and individual grids in the southeastern part of the plateau, the mean annual rainfall erosivity values in most grids in the TP are lower than the observed values.

290



295

Figure 6. Mean annual rainfall erosivity in 2013–2020 (excluding 2017) based on (a) in situ precipitation observations and (b) ERA5 reanalysis precipitation data. (c) *AB* between the values based on ERA5 reanalysis data and precipitation observations. Unit: $\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$.

300

The accuracy of the spatiotemporal variability of the mean annual rainfall erosivity on the TP obtained using the ERA5 dataset is also crucial for determining whether ERA5 is suitable for rainfall erosivity calculation. We used the first three EOF modes, which are considered to provide most of the valuable information, for evaluation. The spatial pattern of the first three EOFs of the observed values accounts for 77% of the total variance, and that of the first three EOFs of the ERA5-based values accounts for

305 84% of the total variance (Figure 7). Clearly, ERA5 successfully captured the spatial pattern of the EOF modes, especially the first two EOF modes, revealed by the observed values. In addition, the corresponding principal components of the EOF modes of the ERA5-based values are also consistent

with the temporal variation trend of the observed values. Therefore, it can be concluded that the ERA5-based mean annual rainfall erosivity generally reproduces the spatiotemporal patterns of the rainfall erosivity on the TP.

310




Figure 7. First three EOF modes of observed and ERA5-based mean annual rainfall erosivity on the TP in 2013–2020 (excluding 2017).

315

4.2 Reconstruction and validation of corrected annual rainfall erosivity

Using the observed and ERA5-based annual rainfall erosivity, we calculated the multiplier factors for 373 grids [Figure 8(a)]. The multiplier factors for the TP range from 0 to 23, with a mean value of 2.4. Multiplier factors of <1 indicate that the ERA5-based annual rainfall erosivity is overestimated, and conversely, the annual rainfall erosivity in the grid is underestimated. Most of the areas with overestimated ERA5-based mean annual rainfall erosivity are located in the Tarim, Qaidam, and Yarlung Zangpo basins. In other areas, the annual rainfall erosivity is typically underestimated, and areas with greater underestimation appear east of the Qaidam basin and in the source area of the Yellow River. We also produced a multiplier factor map of the TP by IDW interpolation based on the multiplier factors of 373 grids [Figure 8(b)].

320
325




Figure 8. (a) Spatial distribution of multiplier factors of 373 grids, (b) multiplier factor map of TP generated by IDW interpolation.

330 The corrected annual rainfall erosivity in 2013–2020 (excluding 2017) was then calculated in the validation grids as the product of the ERA5-based annual values and multiplier factors from the map. Figure 9 compares the observed and ERA5-based annual rainfall erosivity in the validation grids by year. In 2014–2020 (excluding 2017), the multi-year averaged correction coefficient between r_{obs_year} and r_{cor_year} is 0.67, which is 0.13 larger than the value between r_{obs_year} and r_{ERA5_year} . Moreover, all of

335 the data for 2013, which were not used to produce the multiplier factor map, were used to conduct an independent assessment. The results show that the correction coefficient also increases, from 0.53 to 0.67, after the ERA5-based annual rainfall erosivity is corrected, indicating significant improvement.





Figure 9. Comparison of ERA5-based annual rainfall erosivity ($\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$) with observed values in validation grids for 2013–2020 (excluding 2017). The dotted line is the result of an optimal model (with an intercept of 0 and a regression coefficient of 1). The black solid lines are the regression result. Colors of dots represent the grid density.

Violin plots are an alternative method of synthetically evaluating the accuracy of the corrected annual rainfall erosivity. Figure 10 compares the observed, ERA5-based, and corrected annual rainfall erosivity in the validation grids for 2013–2020 (excluding 2017). The corrected annual rainfall erosivity values for 2014–2020 are better than the ERA5-based values in terms of both the probability density and the values corresponding to different quantiles. Even in 2013, a completely independent verification year, the accuracy of the corrected annual rainfall erosivity is greatly improved. Specifically, the observed grid-averaged multi-year mean annual rainfall erosivity is 329

MJ·mm·ha⁻¹·h⁻¹·yr⁻¹ in 2013–2020 (excluding 2017), where the ERA5-based value is 190 MJ·mm·ha⁻¹·h⁻¹·yr⁻¹, and the corrected value is 374 MJ·mm·ha⁻¹·h⁻¹·yr⁻¹. The relative error is significantly reduced, from -42% to 14%, by multiplier factor correction.



355

Figure 10. Violin plots of observed, ERA5-based, and corrected annual rainfall erosivity in validation grids for 2013–2020 (excluding 2017). Y axis shows annual rainfall erosivity in MJ·mm·ha⁻¹·h⁻¹. The boxplot diagram of the median of the violin plots shows the maximum value, 75% quantile value, 50% quantile value, 25% quantile value, and minimum value. The horizontal lines represent average values.

360

4.3 Rainfall erosivity in the TP and related uncertainties

Because of the large variability of the spatiotemporal patterns of precipitation, the R factor, an essential input for soil loss estimation, must be calculated using a minimum of 20 years of precipitation data (Renard et al., 1997). In this study, the annual rainfall erosivity values of the TP for 71 years based on the 0.25° hourly ERA5 precipitation data were calculated by the algorithm shown in Section 3.1. Next, after correction by the multiplier factor map, the new annual rainfall erosivity dataset for 1950–2020 and R factor map were produced.

The annual rainfall erosivity fluctuates considerably within a range of 239 to 408 $\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$ (Figure 10). However, no obvious increasing or decreasing trend appears in the past 71 years across the TP. Regarding the spatial distribution, the R factor generally shows a decreasing trend from southeast to northwest. The areas with R factors below 200 $\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$ are concentrated in the northwestern part of the TP, whereas regions with high R factors appear mainly in the southeastern TP, especially in the Bomi–West Sichuan and Dawang–Chayu areas. The TP-averaged R factor is 307 $\text{MJ}\cdot\text{mm}\cdot\text{ha}^{-1}\cdot\text{h}^{-1}\cdot\text{yr}^{-1}$, which is obviously lower than those from previous studies (e.g. Qin et al., 2016; Cao et al., 2018), excluding Liu et al. (2013) and Yue et al. (2022).




Figure 11. R factor map of TP with the 0.25° spatial resolution for 1950–2020. Inset represents the yearly change in annual rainfall erosivity.

380 Compared with the previous studies, there are two essential improvements by using the data-driven
approach in this study. On the one hand, the 1-min precipitation observations from 1787 weather
stations are firstly used to calculate the accuracy rainfall erosivity values by employing the standard
algorithm. With the densely-spaced rainfall erosivity values, it is able to yield realistic spatial
distribution and identifying the high spatial heterogeneity of the rainfall erosivity over the TP. On the
385 other hand, not only a *R* factor map, we also produced a high-precision time series of annual rainfall
erosivity for 71 years after correcting the ERA5-based estimations, which may offer great help to
reveal the spatial-temporal evolution over the TP under the climate change.

It is also notable that some uncertainties are also unavoidably involved in the newly reconstructed
dataset. As the biases of the ERA5 precipitation data in detecting the characteristics of the erosive
390 rainfall events have been revealed, we intended to use multiple factors to correct the ERA5-based
rainfall erosivity values by grid, to reduce the biases resulting from the ERA5 data. Limited by the
scarcity of the in-situ precipitation observations from weather stations before 2013 (less than 100
weather stations), it is hardly to yield realistic spatial distribution of the multiple factor map by year.
Here, we made a hypothesis that the biases of the ERA5-based annual rainfall erosivity always kept
395 steady in various years, and thus the multi-year average annual multiple factor map from 2014–2020 is
used in correcting process. With the improvement of the weather/climate forecast models in the future,
the biases of the estimated rainfall erosivity by using gridded precipitation data will continue to reduce.

5 Data availability

400 The new gridded annual rainfall erosivity dataset for the TP for 1950–2020 is available at
<http://data.tpdc.ac.cn/en/data/37c34046-3c2a-4737-b3c9-35af398da62a/> (Chen et al., 2021).

6 Conclusions

This study presents a new gridded dataset of annual rainfall erosivity over the TP based on the 1-min
in-situ precipitation data from 1787 weather stations and the long-term ERA5 precipitation data. The
405 annual rainfall erosivity data are available over 71 years (from 1950 to 2020) on a 0.25° grid. The
TP-averaged correction coefficient between the station-based annual rainfall erosivity and the newly
released data is 0.67. In addition, the probability density and various quantile values of the new data are
generally consistent with the station-based values across the TP.

This dataset offers a unique view of large- to local-scale features in rainfall erosivity variability over
410 the TP, where it is hardly to obtain the long-term in-situ precipitation data with sufficient
spatial-temporal resolution. This new data availability opens up many interesting applications in soil
erosion study and disaster research, including:

- (1) providing input data of the R factor for the TP, which is needed for soil erosion modelling;
- (2) understanding the present processes of water erosion over the TP and improving future
415 projections;
- (3) identifying the hot spots at high risk of the landslide and flood hazards.

The data are available in Network Common (NC) Data Format that can be readily imported into
standard geographical information system software (e.g. ArcGIS) or accessed programmatically (e.g.
MATLAB, Python).

420

Acknowledgments

This research was jointly supported by the Second Tibetan Plateau Scientific Expedition and Research
(STEP) Program (Grant No. 2019QZKK0106, Grant No. 2019QZKK0307), and the Basic Research
Special Project of the Chinese Academy of Meteorological Sciences (Grant No. 2020Z003).

425 Author contributions

YC designed the study and wrote the manuscript. XD and MD contributed to the manuscript
preparation and dataset generation. WQ, TW and JL contributed to the analysis method used in this
study. YX contributed to the suggestions for manuscript revision and visualization.

Declaration of interests

430 The authors declare that they have no competing financial interests or personal relationships that could
have influenced the work reported in this paper.

References

Agnese, C., Bagarello, V., Corrao, C., D'Agostino, L., and D'Asaro, F.: Influence of the rainfall
measurement interval on the erosivity determinations in the Mediterranean area, *J. Hydrol.*, 329(1),

-
- 435 39-48, doi: doi.org/10.1016/j.jhydrol.2006.02.002, 2006.
- Angulo-Martínez, M., and Beguería, S.: Estimating rainfall erosivity from daily precipitation records: A comparison among methods using data from the Ebro Basin (NE Spain), *J. Hydrol.*, 379(1), 111-121, doi: doi.org/10.1016/j.jhydrol.2009.09.051, 2009.
- Beck, H.E., Vergopolan, N., Pan, M., Levizzani V., Dijk, A., Weedon, G., Brocca, L., Pappenberger, F.,
440 Huffman, G.J., Wood, E.F.: Global-scale evaluation of 23 precipitation datasets using gauge observations and hydrological modeling, 21, 6201-6217, <https://doi.org/10.5194/hess-21-6201-2017>, 2017.
- Brown, C.L., and Foster, R.G.: storm Erosivity Using Idealized Intensity Distributions, *Trans. ASAE.*, 30(2), 379, doi: doi.org/10.13031/2013.31957, 1987.
- 445 Cao, Y., Wang, S.J., Bai, X.Y., and Li, H.W.: Inversion calculation and spatial-temporal pattern of rainfall erosivity in southwestern China over one hundred years, *Acta Ecologica Sinica*, 38(24), 8766-8773, DOI: 10.5846/stxb201805241142, 2018. In Chinese.
- Chen, Y., Duan, X., Ding, M, Qi, W., Wei, T., and Li, J.: 2021, A new gridded dataset of rainfall erosivity (1950-2020) in the Tibetan Plateau [dataset], National Tibetan Plateau Data Center, DOI: 10.11888/ Terre.tpdc.271833, 2021.
450
- Cucchi, M., Weedon, G. P., Amici, A., Bellouin, N., Lange, S., Müller Schmied, H., Hersbach, H., and Buontempo, C.: WFDE5: bias-adjusted ERA5 reanalysis data for impact studies, *Earth Syst. Sci. Data*, 12, 2097–2120, <https://doi.org/10.5194/essd-12-2097-2020>, 2020.
- Fick, S.E. and R.J. Hijmans: WorldClim 2: new 1km spatial resolution climate surfaces for global land
455 areas. *Int. J. Climatol.*, 37 (12): 4302-4315, <https://doi.org/10.1002/joc.5086>, 2017.
- Freitas, E.D.S., Coelho, V.H.R., Xuan, Y., Melo, D.D.C.D., Gadelha, A.N., Santos, E.A., Galvão, C.D.O., Ramos Filho, G.M., Barbosa, L.R., Huffman, G.J., Petersen, W.A., and Almeida, C.D.N.: The performance of the IMERG satellite-based product in identifying sub-daily rainfall events and their properties, *J. Hydrol.*, 589, 125128, doi: doi.org/10.1016/j.jhydrol.2020.125128, 2020.
- 460 Gu, Z., Feng, D., Duan, X., Gong, K., Li, Y., and Yue, T.: Spatial and temporal patterns of rainfall erosivity in the Tibetan Plateau, *Water*, 12, 200, doi: 10.3390/w12010200, 2020.
- He, J., Yang, K., Tang, W., Lu, H., Qin, J., Chen, Y., and Li, X.: The first high-resolution meteorological forcing dataset for land process studies over China, *Sci. Data.*, 7(1), 25, doi: 10.1038/s41597-020-0369-y, 2020.

-
- 465 Hersbach, H., and Coauthors: Global reanalysis: goodbye ERA-Interim, hello ERA5,
doi:10.21957/vf291hehd7, <https://www.ecmwf.int/node/19027>, 2019.
- Jiang Q., Li W., Fan Z., He X., Sun W., Chen S., Wen J., Gao J., and Wang J.: Evaluation of the ERA5
reanalysis precipitation dataset over Chinese Mainland. *J. Hydrol.*, 595,125660,
<https://doi.org/10.1016/j.jhydrol.2020.125660>, 2021.
- 470 Jiao D., Xu N., Yang F., and Xu K.: Evaluation of spatial-temporal variation performance of ERA5
precipitation data in China. *Sci. Rep.*, 11,17956, <https://doi.org/10.1038/s41598-021-97432-y>, 2021.
- Kim, J., Han, H., Kim, B., Chen, H., and Lee, J.: Use of a high-resolution-satellite-based precipitation
product in mapping continental-scale rainfall erosivity: A case study of the United States, *Catena*,
193, 104602, doi: doi.org/10.1016/j.catena.2020.104602, 2020.
- 475 Li, D., Yang, K., Tang, W., Li, X., Zhou, X., and Guo, D.: Characterizing precipitation in high altitudes
of the western Tibetan plateau with a focus on major glacier areas, *Int. J. Climatol.*, 40(12),
5114-5127, doi: doi.org/10.1002/joc.6509, 2020.
- Li J.: Hour station-based precipitation characteristics over the Tibetan Plateau. *Int. J. Climatol.*, 38:
1560-1570, <https://doi.org/10.1002/joc.52812>, 2018.
- 480 Liu, B., Xie, Y., Li, Z., Liang, Y., Zhang, W., Fu, S., Yin, S., Wei, X., Zhang, K., Wang, Z., Liu, Y.,
Zhao, Y., and Guo, Q.: The assessment of soil loss by water erosion in China, *Int. Soil Water Conse.*,
8(4), 430, doi: [10.1016/j.iswcr.2020.07.002](https://doi.org/10.1016/j.iswcr.2020.07.002), 2020.
- Liu, X., and Chen, B.: Climatic warming in the Tibetan Plateau during recent decades, *Int. J. Climatol.*,
20(14), 1729-1742, doi: [10.1002/1097-0088\(20001130\)20:14<1729::AID-JOC556>3.0.CO;2-Y](https://doi.org/10.1002/1097-0088(20001130)20:14<1729::AID-JOC556>3.0.CO;2-Y),
485 2000.
- Liu, B., Tao, H., Song, C., Guo, B., Shi, Z., Zhang, C., Kong, F., and He, B.: Temporal and spatial
variations of rainfall erosivity in China during 1960 to 2009, *Geographical Research*, 32(2): 245-256,
doi: [10.11821/yj2013020005](https://doi.org/10.11821/yj2013020005), 2013. In Chinese.
- Liu, B.Y., Zhang K.L., and Xie Y.: An empirical soil loss equation. In Proceedings of the 12th
490 International Soil Conservation Organization Conference, Beijing, China, 26-31 May 2002; Tsinghua
University Press: Beijing, China, 2002.
- Lobo, G.P., and Bonilla, C.A.: Sensitivity analysis of kinetic energy-intensity relationships and
maximum rainfall intensities on rainfall erosivity using a long-term precipitation dataset, *J. Hydrol.*,
527, 788-793, doi: doi.org/10.1016/j.jhydrol.2015.05.045, 2015.

-
- 495 Ma, X., He, Y., Xu, J., van Noordwijk, M., and Lu, X.: Spatial and temporal variation in rainfall erosivity in a Himalayan watershed, *Catena*, 121, 248-259, doi: doi.org/10.1016/j.catena.2014.05.017, 2014.
- Ma Y., Yang Y., Han Z., et al.: Comprehensive evaluation of ensemble multi-satellite precipitation dataset using the dynamic Bayesian model averaging scheme over the Tibetan Plateau. *J. Hydrol.*, 500 556: 634-644, <https://doi.org/10.1016/j.jhydrol.2017.11.050>, 2018.
- Panagos, P., Borrelli, P., Meusburger, K., Yu, B., Klik, A., Jae Lim, K., Yang, J.E., Ni, J., Miao, C., Chattopadhyay, N., Sadeghi, S.H., Hazbavi, Z., Zabihi, M., Larionov, G.A., Krasnov, S.F., Gorobets, A.V., Levi, Y., Erpul, G., Birkel, C., Hoyos, N., Naipal, V., Oliveira, P.T.S., Bonilla, C.A., Meddi, M., Nel, W., Al Dashti, H., Boni, M., Diodato, N., Van Oost, K., Nearing, M., and Ballabio, C.: Global 505 rainfall erosivity assessment based on high-temporal resolution rainfall records, *Sci. Rep.*, 7(1), 4175, doi: doi.org/10.1038/s41598-017-04282-8, 2017.
- Panagos, P., Meusburger, K., Ballabio, C., Borrelli, P., Beguería, S., Klik, A., Rymaszewicz, A., Michaelides, S., Olsen, P., Tadić, M.P., Aalto, J., Lakatos, M., Dumitrescu, A., Rousseva, S., Montanarella, L., and Alewell, C.: Reply to the comment on “Rainfall erosivity in Europe” by 510 Auerswald et al., *Sci. Total Environ.*, 532, 853-857, doi: doi.org/10.1016/j.scitotenv.2015.05.020, 2015.
- Qi W., Liu J., and Chen D.: Evaluations and improvements of GLDAS2.0 and GLDAS2.1 forcing data’s applicability for basin scale hydrological simulations in the Tibetan Plateau. *J. Geophys. Res-atmos*, 123: 13128-13148, <https://doi.org/10.1029/2018JD029116>, 2018.
- 515 Qin, W., Guo, Q., Zuo, C., Shan, Z., Ma, L., and Sun, G.: Spatial distribution and temporal trends of rainfall erosivity in mainland China for 1951–2010, *Catena*, 147, 177-186, doi:doi.org/10.1016/j.catena.2016.07.006, 2016.
- Renard, K.G., Foster, G.A., Weesies, D.K., McCool, D.K., and Yoder, D.C.: Predicting soil erosion by water: a guide to conservation planning with the revised universal soil loss equation (RUSLE). U.S. 520 Department of Agriculture, Agriculture Handbook No. 703, 1997.
- Risal, A., Lim, K.J., Bhattarai, R., Yang, J.E., Noh, H., Pathak, R., and Kim, J.: Development of web-based WERM-S module for estimating spatially distributed rainfall erosivity index (EI30) using RADAR rainfall data, *Catena*, 161, 37-49, doi: doi.org/10.1016/j.catena.2017.10.015, 2018.
- Shin, J., Kim, T., Heo, J., and Lee, J.: Spatial and temporal variations in rainfall erosivity and erosivity

-
- 525 density in South Korea, *Catena*, 176, 125-144, doi: doi.org/10.1016/j.catena.2019.01.005, 2019.
- Sun, Q., Miao C., Duan Q., Ashouri H., Sorooshian S., and Hsu K.: A Review of Global Precipitation Data Sets: Data Sources, Estimation, and Intercomparisons, *Rev. Geophys.*, 56, 79-107, <https://doi.org/10.1002/2017RG000574>, 2018.
- Teng, H., Hu, J., Zhou, Y., Zhou, L., and Shi, Z.: Modelling and mapping soil erosion potential in China, *J. Integr. Agr.*, 18(2), 251-264, doi: doi.org/10.1016/S2095-3119(18)62045-3, 2019.
- 530 Teng, H., Liang, Z., Chen, S., Liu, Y., Viscarra Rossel, R.A., Chappell, A., Yu, W., and Shi, Z.: Current and future assessments of soil erosion by water on the Tibetan Plateau based on RUSLE and CMIP5 climate models, *Sci. Total Environ.*, 635, 673-686, doi: doi.org/10.1016/j.scitotenv.2018.04.146, 2018.
- 535 Vrieling, A., Sterk, G., and de Jong, S.M.: Satellite-based estimation of rainfall erosivity for Africa, *J. Hydrol.*, 395(3), 235-241, doi: doi.org/10.1016/j.jhydrol.2010.10.035, 2010.
- Wang, Y., Cheng, C., Xie, Y., Liu, B., Yin, S., Liu, Y., and Hao, Y.: Increasing trends in rainfall-runoff erosivity in the Source Region of the Three Rivers, *Sci. Total. Environ.*, 592, 639-648, <https://doi.org/10.1016/j.scitotenv.2017.02.235>, 2017.
- 540 Wischmeier, W. H., and Smith, D. D., 1965. Predicting rainfall-erosion losses from cropland east of the Rocky Mountains: Guide for selection of practices for soil and water conservation. US 434 Department of Agriculture.
- Wischmeier, W. H., and Smith, D. D., 1978. Predicting rainfall erosion losses: a guide to conservation planning. Department of Agriculture, Science and Education Administration. U. S. Department of
- 545 Agriculture, Agriculture Handbook No. 537.
- Xie, Y., Liu, B. Y., and Zhang, W. B.: Study on standard of erosive rainfall. *Journal of Soil and Water Conservation*, 14(4), 6-11, doi: 10.3321/j.issn:1009-2242.2000.04.002, 2000. In Chinese.
- Yan, D., Fan, J., Guo, F., Guo, X., and Gong, K.F.: Spatiotemporal distribution of precipitation erosivity in Tibet autonomous region, *Bulletin of Soil and Water Conservation*, 30(3): 17-21, doi: 10.13961/j.cnki.stbctb.2010.04.025, 2010. In Chinese.
- 550 Yang, F., and Lu, C.: Spatiotemporal variation and trends in rainfall erosivity in China's dryland region during 1961-2012, *Catena*, 133, 362-372, doi: dx.doi.org/10.1016/j.catena.2015.06.005, 2015.
- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D.B., and Joswiak, D.: Different glacier status with atmospheric

-
- 555 circulations in Tibetan Plateau and surroundings, *Nat. Clim. Change*, 2, 663-667, doi:
doi.org/10.1038/nclimate1580, 2012.
- Yin, S., Xie, Y., Liu, B., and Nearing, M.: Rainfall erosivity estimation based on rainfall data collected
over a range of temporal resolutions, *Hydrol. Earth Syst. Sc.*, 19, doi: 10.5194/hess-19-4113-2015,
2015.
- 560 Yuan X., Yang K., Lu H., He, J., Sun J., and Wang Y.: Characterizing the features of precipitation for
the Tibetan Plateau among four gridded datasets: Detection accuracy and spatio-temporal
variabilities. *Atmos. Res.*, 264, 105875, <https://doi.org/10.1016/j.atmosres.2021.105875>, 2021.
- Yue, T., Yin, S., Xie, Y., Yu, B., and Liu, B.: Rainfall erosivity mapping over mainland China based on
high density hourly rainfall records, *Earth Syst. Sci. Data.*, 14, 665-682,
565 doi.org/10.5194/essd-2020-370, 2022.
- Zhang, W., Xie, Y., and Liu, B.: Spatial distribution of rainfall erosivity in China. *Journal of Mountain
Science*, 21(1): 33-40, doi: 10.16089/j.cnki.1008-2786, 2003. In Chinese.
- Zhou, X., Yang, K., Ouyang, L., Wang, Y., Jiang, Y., Li, X., Chen, D., Prein, A.:
Added value of kilometer-scale modeling over the third pole region: a CORDEXCPTP pilot study,
570 *Clim. Dyn.*, 57, 1673-1687, <https://doi.org/10.1007/s00382-021-05653-8>, 2021.