

# Optical and biogeochemical properties of diverse Belgian inland and coastal waters

Alexandre Castagna<sup>1</sup>, Luz Amadei Martínez<sup>1</sup>, Margarita Bogorad<sup>1</sup>, Ilse Daveloose<sup>1</sup>, Renaat Dasseville<sup>1</sup>, Heidi Dierssen<sup>2</sup>, Matthew Beck<sup>3</sup>, Jonas Mortelmans<sup>4</sup>, Héloïse Lavigne<sup>3</sup>, Ana Dogliotti<sup>5</sup>, David Doxaran<sup>6</sup>, Kevin Ruddick<sup>3</sup>, Wim Vyverman<sup>1</sup>, and Koen Sabbe<sup>1</sup>

<sup>1</sup>Protistology and Aquatic Ecology, Ghent University, Ghent, Belgium

<sup>2</sup>Department of Marine Sciences, University of Connecticut, Groton, CT, United States

<sup>3</sup>Operational Directorate Natural Environment, Royal Belgian Institute of Natural Sciences, Brussels, Belgium

<sup>4</sup>Marine Observation Centre (MOC), Flanders Marine Institute (VLIZ), Oostende, Belgium

<sup>5</sup>Instituto de Astronomía y Física del Espacio (IAFE), CONICET-Universidad de Buenos Aires, Buenos Aires, Argentina

<sup>6</sup>Laboratoire d'Océanographie de Villefranche, CNRS-Sorbonne University, Villefranche, France

**Correspondence:** Alexandre Castagna (alexandre.castagna@ugent.be)

**Abstract.** From 2017 to 2019, an extensive sampling campaign was conducted in Belgian inland and coastal waters, aimed at providing paired data of optical and biogeochemical properties to support research into optical monitoring of aquatic systems. The campaign was focused on inland waters, with sampling of four lakes and a coastal lagoon along the growth season, in addition to samples of opportunity of other four lakes. Campaigns also included the Scheldt estuary over a tidal cycle and two sampling campaigns in the Belgian coastal zone. Measured parameters include inherent optical properties (absorption, scattering and beam attenuation coefficients, near-forward volume scattering function, turbidity), apparent optical properties (Secchi disk depth, substrate and water-leaving Lambert-equivalent bi-hemispherical reflectance), and biogeochemical properties (suspended particulate matter, mineral fraction of particle mass, particle size distribution, pigment concentration, DNA metabarcoding, flow microscopy counts, and bottom type classification). The diversity of water bodies and environmental conditions covered a wide range of system states. The chlorophyll *a* concentration varied from 0.63 mg m<sup>-3</sup> to 382.72 mg m<sup>-3</sup>, while the suspended particulate matter concentration varied from 1.02 g m<sup>-3</sup> to 791.19 g m<sup>-3</sup>, with mineral fraction varying from 0 to 0.95. Depending on system and season, phytoplankton assemblages were dominated by cyanobacteria, green algae (Mamiellophyceae, Pyramimonadophyceae) or diatoms. The dataset is available from Castagna et al. (2022), <https://www.pangaea.de/tok/c67200d99ea9bbbeadd9edec9690f937b5bacbff>.

## 15 1 Introduction

Datasets of paired optical and biogeochemical properties are essential for developing and validating the interpretation of optical signals captured with *in situ* instrumentation or remote sensors. Though the data gathered in the last 50 years provide a large collection of conditions across a diverse set of environments, three major caveats are observed in the freely accessible datasets (*e.g.*, SeaBASS at <https://seabass.gsfc.nasa.gov/> and license categories A and B of LIMNADES at <https://limnades.stir.ac.uk/>):

20 1. The majority of the data concerns open oceans and coastal waters, with little representation of inland water systems;

2. The majority of the data concerns multispectral measurements, particularly at wavebands typical of ocean colour sensors;
3. The majority of the paired biogeochemical data includes only broad features (*e.g.*, chlorophyll *a* concentration) deemed suitable for operational retrievals with multispectral instruments.

In order to fully exploit the potential of hyperspectral satellite missions, hyperspectral datasets paired with detailed com-  
25 position information of aquatic systems are required (Dierssen et al., 2020). In addition, more extensive data is necessary to develop and validate regional optical retrievals over complex optical systems such as lakes, lagoons, estuaries and rivers.

The data presented here were gathered and processed during three projects funded by the Belgian Science Policy Of-  
fice (BELSPO) and one project funded by the Research Foundation – Flanders (FWO). The PONDER project (BELSPO  
SR/00/325) focused on developing tools for spaceborne remote sensing of inland water systems using high spatial resolution  
30 ( $\leq 30$  m) sensors. During the course of the project, global coverage and open access data for high spatial resolution sensors was only available for multispectral missions. The HYPERMAQ project (BELSPO SR/00/335) focused on exploring hyperspectral data for detailed biogeochemical retrievals in support of the new generation of hyperspectral spaceborne remote sensing missions. The PHYTOBEL project (BELSPO SR/02/213) provided funding for data curation and publication. The Flemish part of the LifeWatch BE, a long-term research project funded by FWO, provided biogeochemical data and infrastructure for  
35 sampling the Belgian coastal zone (BCZ). Samples were taken from 2017 to 2019 and cover eight lakes, the Spuikom lagoon, the Scheldt estuary and the BCZ.

The goal of this data report is to provide a detailed description and validation of the methods used during the research, present a summary of the observations and briefly discuss aspects of the data that might be relevant for potential users.

## 2 Methods and data description

40 Measurements are presented in three groups: (1) Inherent optical properties (IOPs) consisting of absorption, scattering and beam attenuation coefficients, near-forward volume scattering function and turbidity; (2) Apparent optical properties (AOPs) consisting of Secchi disk depth, water-leaving and substrate Lambert-equivalent bi-hemispherical reflectance; and (3) biogeochemical properties consisting of suspended particulate matter concentration, mineral fraction of particle mass, particle size distribution, pigment concentration, DNA metabarcoding, flow imaging microscopy counts, and bottom type classifica-  
45 tion. Table 1 presents the description of relevant acronyms, symbols, constants and subscripts used in this study. The studied aquatic systems are listed in Table 2. The measured parameters and the number of samples are listed in Tables 3, 4 and 5. All coordinates are relative to the WGS84 datum and times are reported in UTC.

Linear models used for consistency check between parameters were fitted using robust linear regression (iterated reweighted least squares with Huber’s loss function; Venables and Ripley, 2002). For selected parameters, the data summary is provided  
50 in the form of a violin plot, which in addition to the median point, 25th and 75th percentiles, presents the empirical distribution of the data as the variable width of the planar shape. All analyses were performed in R (version 3.6.3, R Core Team, 2020) with aid of packages ‘MASS’ (version 7.3-51.5, Venables and Ripley, 2002), ‘vioplot’ (version 0.3.6, Adler and Kelly, 2020), ‘dada2’ (version 1.12.1, Callahan et al., 2016) and ‘decontam’ (version 1.6.0, Davis et al., 2018).

## 2.1 Study sites

55 The majority of the measurements were performed in Belgian inland waters, in particular four lakes and a coastal lagoon. The sampling frequency along the growth season (spring to autumn) varied with the sampling year. The field campaigns were performed seasonally throughout 2017 (fortnightly) and 2018 (monthly), spanning a broad range of conditions. Data for four additional lakes with sporadic sampling during 2017 are also included in the dataset. Those systems were previously described in Castagna et al. (2020) and are reintroduced here for convenience. A summary of the study sites is presented in Table 2.

60 The Spuikom (51°13'41.0"N 2°57'09.5"E) is a shallow, brackish-marine coastal lagoon (typical salinity between 27 PSU and 33 PSU), with a surface area of 0.82 km<sup>2</sup> and an average depth of 1.5 m. It is connected to the sea through the Ostend harbor by a lock system. The lagoon exchanges water with the harbor continuously, resulting in median daily renewal of 3.7 % of its volume. It is used for recreational activities and shellfish aquaculture. During the observation period (2017 to 2018) it experienced a cycle of diatom blooms in spring followed by a transition to transparent waters in autumn, with extensive  
65 bottom coverage by macroalgae. It is subject to strong coastal winds that can cause sediment resuspension, adding bottom sediments and benthic diatoms to the water column. The chlorophyll *a* (Chl *a*) concentration typically varied from 2 mg m<sup>-3</sup> to 25 mg m<sup>-3</sup>, with the exception of an intense nanoflagellate bloom that reached a Chl *a* concentration of ≈130 mg m<sup>-3</sup>. The Secchi disk depth ranged from 0.62 m to bottom.

The Hazewinkel (51°03'57.4"N 4°23'27.1"E) is a mesotrophic lake with an area of 0.66 km<sup>2</sup> and a maximum depth of  
70 20 m. It is used for recreational and sport activities. During the observation period (2017) it presented a peak of phytoplankton abundance in spring, with Chl *a* concentration reaching 20 mg m<sup>-3</sup>, after which the concentration stayed around 5 mg m<sup>-3</sup>. Secchi disk depth varied from 1 m to 6 m. Submerged macrophytes were confined to near-shore locations due to a steep basin slope.

The Donkmeer (51°02'23.4"N 3°58'47.2"E) is the second largest lake in Flanders, with a surface area of 0.86 km<sup>2</sup> and  
75 an average depth of 2 m. It experiences recurrent cyanobacterial blooms of *Anabaena spp.* and *Planktothrix agardhii* (Descy et al., 2011). It is used for recreational activities and fishing. During the observation period (2017 to 2018) it experienced cyanobacterial blooms from summer to autumn, reaching a Chl *a* concentration of ≈400 mg m<sup>-3</sup> and a Secchi disk depth of 0.2 m. Its northern and southern portions are connected through a narrow and shallow passage, with the northern portion experiencing a shorter period of cyanobacterial blooms due to management actions.

80 The Dikkebus (50°49'07.4"N 2°50'39.5"E; 0.36 km<sup>2</sup>, 2.5 m deep) and the Zillebeke (50°50'15.4"N 2°54'34.3"E; 0.28 km<sup>2</sup>, 2 m deep) are two lakes created in the 13th century for water supply, a function that remains to date along with recreational activities. Both lakes exhibit annual blooms of the cyanobacteria *Microcystis aeruginosa* and *Aphanizomenon flos-aquae* (Descy et al., 2011). During the observation period (2018), intense cyanobacterial blooms were observed with the Chl *a* concentration reaching 105 mg m<sup>-3</sup> and in the Zillebeke those blooms were associated with fish mortality. The Secchi disk depth ranged  
85 from 0.5 m to 2 m. Additional stations were included from four other Flemish lakes: Bocht (51°04'24.0"N 4°23'19.8"E; 0.35 km<sup>2</sup> and 18 m deep), Nieuwdonk (51°02'04.1"N 3°58'32.2"E; 0.26 km<sup>2</sup> and 22 m deep) and two unnamed adjacent

lakes (51°06'06.1"N 4°01'44.1"E;  $\approx 0.16 \text{ km}^2$  each and 10 m deep), here referred to as “DelMare”, a sand extraction lake with water sports activities, and “Molsbroek”, a nature reserve.

Other inland water systems included were the Scheldt estuary and the Leuven-Dijle canal. The Scheldt is a rain-fed low-land river, with an estuary environment subjected to tides from the mouth at Vlissingen (The Netherlands) to Ghent (Belgium, 160 km upstream), where a system of locks prevent further propagation of the tide (Meire et al., 2005). It has a large economical importance as a transport waterway, connecting the harbours of Antwerp and Ghent to the North Sea. The Scheldt estuary was sampled in mid October 2019 at two locations near the city of Sint-Amands (51°03'18.0"N 4°11'59.6"E and 51°04'24.1"N, 4°11'24.0"E), with a time series including a full tidal cycle (tidal range of 6 m). During the observation period the Chl *a* concentration reached  $55.7 \text{ mg m}^{-3}$  at high tide, while the suspended particulate matter (SPM) concentration reached  $791.2 \text{ g m}^{-3}$ . In the same campaign performed in the Scheldt, two samples were taken at the Leuven-Dijle canal, a highly transparent artificial waterway running parallel to the Dijle river, connecting Leuven to the Zenne-Dijle confluence, and ultimately to the Scheldt. The sampling position was close to the lock of Zennegat (51°03'46.7"N 4°25'50.0"E).

The stations sampled in the BCZ are part of the regular LifeWatch sampling campaigns (Mortelmans et al., 2019). The April and July 2018 campaigns were augmented to include spectroscopic measurements. The BCZ is a shallow part of the North Sea (1 m to 40 m), experiencing high tidal fluctuations (average of 4 m) and strong tidal currents ( $1 \text{ m s}^{-1}$ ). Those conditions, combined with limited freshwater discharge into the region, result in a well mixed-water column (van Beusekom and Diel-Christiansen, 1993). It experiences high turbidity (SPM concentration from  $1 \text{ g m}^{-3}$  to  $200 \text{ g m}^{-3}$ ) with large influence from particulate material imported through the Strait of Dover (Fettweis and Van den Eynde, 2003). The BCZ develops a turbidity maximum zone near Zeebrugge, also influenced by the decreasing magnitude of the residual transport vectors from the East border (Fettweis and Van den Eynde, 2003). The inflow of the Yser and Scheldt rivers influence the availability of nutrients, with an increase of nitrogen and phosphorus since the second half of the 20th century, followed by a de-eutrophication phase during which nitrogen and especially phosphorus decreased (Desmit et al., 2020). The phytoplankton seasonal dynamics are well described (e.g., Reid et al., 1990), with an early spring diatom bloom followed by a mixed bloom of the haptophyte *Phaeocystis globosa* and diatoms. A recent review of the phytoplankton seasonal dynamic was provided by Castagna et al. (2021). The IOPs in this region were extensively studied by Astoreca et al. (2006, 2009, 2012).

## 2.2 Sampling

Sampling was performed from a diverse set of platforms. For the inland water campaigns, samples were taken from pontoons or an inflatable boat, depending on the system and date. The water was sampled just below the surface, taking care not to draw in materials floating at the surface (e.g., pollen, debris, etc). Field samples were stored in 5 L semi-transparent plastic carboys, kept in the dark and cold during the transport to the laboratory, and processed within 4 to 6 hours from sampling. Sampling for the coastal campaigns was performed from the Research Vessel (RV) Simon Stevin (Flanders Marine Institute, VLIZ). For most marine samples, water was sampled just below the surface using Niskin bottles (General Oceanics, Inc.), attached to a rosette system. The exception were subsurface samples for flow imaging microscopy, taken with a bucket from the side of

120 the ship. Filtrations were performed on board and water subsamples were kept in the dark and cold during transport to the laboratory for spectrophotometric measurements.

Macrophytes, sediment and biofilm were sampled in the Spuikom lagoon. Macrophytes were sampled during 2017 and 2018, by collecting floating specimens or recovering specimens from the bottom using a rake. Specimens were stored in transparent plastic bags containing water from the lagoon, and kept in the dark and cold during transport to the laboratory. Sediments and  
125 biofilm were sampled in April 2018 using polymethyl methacrylate (PMMA) tubes attached to a short corer. The cores were retrieved with care not to disturb the surface of the sediment, sealed and transported to the laboratory in a vertical position. In July 2018, biofilm patches had detached from the bottom of the Spuikom and were sampled floating at the surface and stored in plastic bags for microscopic examination. The cores and macrophytes were stored in a climate room at 4°C for up to 3 days until analysis.

### 130 2.3 Inherent Optical Properties

Most of the IOP measurements were performed *ex situ*, with a benchtop spectrophotometer (Lambda 650S, PerkinElmer) equipped with an integrating sphere (150 mm internal diameter). The interior of the integrating sphere is made of highly reflective (nominal 99 %) sintered polytetrafluoroethylene (PTFE). The spectral IOP measurements made with the spectrophotometer include particle absorption ( $a_p$ ), chemically decomposed into *in vivo* pigment absorption ( $a_\phi$ ) and depigmented particle absorp-  
135 tion ( $a_d$ ), chromophoric dissolved organic matter absorption ( $a_g$ ), particle beam attenuation ( $c_p$ ) and scattering ( $b_p$ ) Napierian coefficients. Methods followed recommendations from Pegau et al. (2002) and IOCCG (2018). Measurements covered the range from 250 nm to 850 nm in 1 nm steps, with a 2 nm integration slit and 0.24 s integration time (250 nm min<sup>-1</sup>). Data is provided in the range of 380 nm to 850 nm, and include a single pass of a smoothing function to reduce noise (rectangular filter, 10 nm window moving average).

140 Turbidity, defined as the side-scattering at 860 nm relative to Formazin standards (ISO 7027:1999; Dogliotti et al., 2015; Boss et al., 2009a), was measured in discrete samples with a portable turbidimeter (2100P ISO, HACH). Additional IOP data were measured from *in situ* instrumentation for a subset of water systems and stations. These include the diffraction peak ( $\psi < 15^\circ$ ) of the particle volume scattering function (VSF;  $\beta_p(670)$ ) and non-water beam attenuation at an acceptance angle of 0.018° measured at 670 nm ( $c_{nw}(670, 0.018)$ ; LISST, Sequoia Scientific).

#### 145 2.3.1 Absorption coefficient from dissolved components (< 0.45 μm)

Absorption due to chromophoric dissolved organic matter (CDOM) was determined in the laboratory from fresh and undiluted filtered subsamples, using a 5 cm pathlength quartz cuvette. Subsamples were filtered with 0.45 μm polyamide (nylon) fiber syringe filters. Between samples, the syringe and cuvette were rinsed with deionized water. The filters were first rinsed by filtrating 15 mL of deionized water before use with samples. The first sample filtration volume was used for final rinse of  
150 the quartz cuvette and the volume discarded. The cuvette was then gently filled with the second filtration volume to avoid the formation of bubbles, and allowed to rest for at least 3 min before measurements. The blank was determined with deionized water, using the same cuvette and at the same temperature (room temperature, ≈20°C). Quality control evaluated the presence

of absorption peaks at 676 nm as an indication of cell leakage and the offset from zero absorption at 750 nm as a indication of the presence of hydrosols.


155 The use of deionized water as blank for all measurements resulted in a mismatch of salinity for marine and brackish samples. Dissolved salts change the complex refractive index of water (Quan and Fry, 1995; Röttgers et al., 2014b), affecting reflection and refraction interactions with the cuvette wall, water scattering and absorption. In the wavelength range reported here, the scattering effects and the absorption of salts are negligible, however the dissolved salts form what is known as salt-solvated water, mixed with pure water (Max and Chapados, 2001). Salt-solvated water can have higher or lower absorption than pure  
160 water, depending on the wavelength, with a magnitude defined by the salt type (Max and Chapados, 2001; Röttgers et al., 2014b). This difference is particularly important in the NIR range, due to the spectral shape of  $a_g$ . The effect of the mismatch in salinity was investigated by comparing the effects of salinity on blank readings for  $a_g$  determinations from deionized water and artificial seawater at a salinity of 35 (NaCl at 35 g kg<sup>-1</sup>). The results are shown in Fig. S1 (supplementary material) and the resulting difference spectrum was subtracted from  $a_g$  of the Spuikom lagoon and the BCZ for an approximate correction.  
165 Dissolved molecular oxygen (O<sub>2</sub>) also contributes to absorption in the UV range (Jonaz and Fournier, 2007, and references therein), however in the wavelength range reported here CDOM dominates the signal.

The Lambda 650S was not available in 2017 and  $a_g$  was measured for a subset of samples using another spectrophotometer (UV-1601, Shimadzu). Sample preparation was the same, except that  $a_g$  was only measured at selected wavelengths (380 nm, 400 nm, 433 nm, 550 nm and 750 nm). To retrieve the full spectrum for these samples, an hyperbolic model was fitted to the  
170 available wavelengths (Twardowski et al., 2004). The hyperbolic model fit was also applied to all measurements in the range of 380 nm to 850 nm to provide a smooth, fitted version of  $a_g$  and to calculate the CDOM absorption spectral hyperbolic slope coefficient,  $S_{h_g}$ :

$$a_g(\lambda) = a_g(443) \left( \frac{\lambda}{443} \right)^{-S_{h_g}}. \quad (1)$$

Most samples had negligible absorption in the NIR ( $< |0.1| \text{ m}^{-1} \text{ nm}^{-1}$ ), fluctuating around zero due to instrument noise  
175 or residual blank salinity difference effect. Four stations had positive or negative offsets that were larger than  $|0.1|$ , and the negative offsets were attributed to the presence of bubbles in the blank of the day. All spectra were subtracted by the average  $a_g$  between 750 nm and 850 nm. Five freshwater samples had a pronounced Chl *a* absorption peak at 676 nm, and since NIR signals were close to zero, those peaks likely result from cell leakage. The  $a_g$  for those stations were flagged.

The magnitudes of  $a_g(412)$  varied between 0.547 m<sup>-1</sup> nm<sup>-1</sup> and 4.185 m<sup>-1</sup> nm<sup>-1</sup>, with values above 2.25 m<sup>-1</sup> nm<sup>-1</sup>  
180 generally observed only in freshwater systems. The exception was a sample in the Spuikom collected during a nanoflagellate bloom, when  $a_g(412)$  reached 3.160 m<sup>-1</sup> nm<sup>-1</sup>. The range of the  $S_{h_g}$  was between 5.60 and 8.81. The observed values of  $a_g$  and fitted  $S_{h_g}$  are presented in Fig. 1.



**Figure 1.** Chromophoric dissolved organic matter (CDOM) absorption in the Belgian water systems. (A) CDOM spectral absorption coefficient,  $a_g$ ; and (B) hyperbolic slope of  $a_g$ ,  $S_{hg}$ , fitted in the range of 380 nm to 850 nm.

### 2.3.2 Absorption coefficient from particulate components ( $> 0.7 \mu\text{m}$ )

Particle absorption was measured using the filterpad method, with particles concentrated onto a glass fiber filter (GF/F, effective mesh of  $0.7 \mu\text{m}$ ). To improve the homogeneity of particle deposition over the filtration area, two stacked filters were used and samples with high abundance of particles were diluted in the filtration funnel with deionized water. Immediately after filtration,

filters were transferred to PetriSlides (Merck), wrapped in aluminum foil and frozen in liquid nitrogen. Filters were then stored at -80 °C until analysis.

Before analysis, the filters were allowed to thaw to room temperature and kept hydrated with deionized water. To avoid dislocating large particles deposited on the filter fibers, hydration was performed by raising the filter, adding a droplet of water to the PetriSlide base and gently lowering the filter onto the droplet, resulting in water spreading by capillarity. The “inside sphere” variant of the quantitative filterpad method was used (Stramski et al., 2015), using a center mount coated with PTFE. Filters were read twice, with a 90° rotation between reads to average small deviations from homogeneous deposition. This was especially important to account for large cyanobacteria colonies when these occurred in low densities (Fig. S2, supplementary material). Pigments were then oxidized by treating the filters with sodium hypochlorite (NaClO; Ferrari and Tassan, 1999), using the same approach as for hydration. Filters were read after 15 min or complete oxidation, following the same orientations as of the pigmented readings. The pathlength amplification correction was taken from Stramski et al. (2015) as recommended in IOCCG (2018). The *in vivo* pigment absorption coefficient ( $a_\phi$ ) was calculated as the total particulate absorption coefficient ( $a_p$ ) subtracted by the depigmented particle absorption coefficient ( $a_d$ ). The chemical oxidation step was not performed for the Scheldt samples.

Sodium hypochlorite has an absorption peak at  $\approx 300$  nm, with residual absorption up to 500 nm. One method to remove the oxidant’s signal is to rinse the filter with pure water before the determinations of  $a_d$  and this method was used for a subset of samples to evaluate the impact of sodium hypochlorite and rinsing. We observed that some samples had an apparent loss of  $a_d$  observed between determinations with sodium hypochlorite and after rinsing (Fig. 2). The magnitude of this effect varied across the lakes, with Spuikom, Donkmeer, Hazewinkel and Bocht suffering little to no loss between 550 nm and 850 nm, while Dikkebus and Zillebeke showed large reductions in  $a_d$  after rinsing. This effect might provide additional information on the nature of the particles in a given system, but needs to be further explored to understand its causes. A direct consequence for measurements of  $a_d$  and  $a_\phi$  is that rinsing might underestimate  $a_d$ .


Therefore we developed a statistical method to remove the oxidant’s signal, instead of applying rinsing. We observed that when there was no loss of signal with rinsing, the minimum difference between  $a_p$  and rinsed  $a_d$  in the UV was at 305 nm, with  $a_d(305)$  typically between 80 % and 90 % of  $a_p(305)$  (Fig. 2A). Based on these observations, we fitted an exponential function to measured  $a_d$  with sodium hypochlorite in the range of 550 nm to 850 nm, and included a point-estimate of rinsed  $a_d$  at 305 nm as  $0.8a_p(305)$ . The additional data point at 305 nm helps to set the curvature of the exponential model. Finally, an offset was included as necessary to match the  $a_p$  absorption in the NIR end. The exponential model with an offset was:

$$a_d(\lambda) = \alpha_d + (a_d(550) - \alpha_d)e^{-S_{e_d}(\lambda-550)}, \quad (2)$$

where  $\alpha_d$  ( $\text{m}^{-1} \text{nm}^{-1}$ ) is a spectrally flat offset (Estapa et al., 2012) and  $S_{e_d}$  ( $\text{nm}^{-1}$ ) is the spectral exponential slope of  $a_d$ .


The treatment with sodium hypochlorite can also introduce another artifact to the measurements. For a set of samples we observed a baseline offset between  $a_p$  and  $a_d$  in the NIR even before rinsing, propagating to increased baseline absorption of  $a_\phi$ . This likely results from sodium hypochlorite removing the adsorbed organic layer over particles (Binding et al., 2008),





**Figure 2.** Evaluation of the treatment of the filters with sodium hypochlorite (NaClO) and subsequent rinse on the determinations of  $a_d$ . (A) A sample from the Bocht (BL\_08) and (B) a sample from the Dikkebus (DK\_01). The sample in (B) show loss of absorbing material after rinsing.


220 with the magnitude of the effect proportional to the concentration of particles and organic matter. In our samples this effect was larger in the maximum turbidity zone of the BCZ. This loss of absorbing material was not observed in a study by Röttgers et al. (2014a) including samples from a diverse set of environments, though the authors did not apply NaClO to the North Sea or Baltic Sea samples. The baseline effect was compensated for by adding the minimum offset necessary to match  $a_p$  and  $a_d$  in the



**Figure 3.** Depigmented particle absorption in the Belgian water systems. (A) Fitted depigmented particle absorption coefficient,  $a_d$ ; and (B) spectral exponential slope of  $a_d$ ,  $S_{e_d}$ , fitted in the range of 380 nm to 850 nm. Note that for the Scheldt samples, the exponential model was fitted directly to  $a_p$ .

range of 800 nm to 850 nm without generating negative  $a_\phi$  values. This constant offset likely underestimates the contribution  
 225 of organic absorption in the blue end due to the exponential spectral shape of this component (*e.g.*, Cael and Boss, 2017).

Similar to what was observed for  $a_g(412)$ , the magnitudes of  $a_d(412)$  varied between  $0.055 \text{ m}^{-1} \text{ nm}^{-1}$  and  $4.570 \text{ m}^{-1} \text{ nm}^{-1}$ , with values above  $2.79 \text{ m}^{-1} \text{ nm}^{-1}$  generally observed only in freshwater systems. The baseline absorption coefficient in the NIR ( $a_p$  or  $a_d$ , and equal to  $a_\alpha$ ) ranged from  $0.003 \text{ m}^{-1} \text{ nm}^{-1}$  to  $7.933 \text{ m}^{-1} \text{ nm}^{-1}$ , with only Scheldt samples presenting val-



**Figure 4.** Particle and *in vivo* pigment absorption in the Belgian water systems. (A) Particle absorption coefficient,  $a_p$ ; and (B) *In vivo* pigment absorption coefficient,  $a_\phi$ . The  $a_\phi$  presented here was calculated with the fitted  $a_d$ .

ues higher than  $0.565 \text{ m}^{-1} \text{ nm}^{-1}$ . The range of the  $S_{e_d}$  was between  $0.0064 \text{ nm}^{-1}$  and  $0.0194 \text{ nm}^{-1}$ . The fitted values of  $a_d$  and  $S_{e_d}$  are presented in Fig. 3. Since particle absorption samples from the Scheldt were not depigmented but were dominated by  $a_d$  (Fig. 4A), an estimate of  $S_{e_d}$  is provided by fitting Eq. 2 directly to  $a_p$  (Fig. 3B).

The integral-normalized  $a_\phi$  presented a diversity of pigment absorption peaks (Fig. 4B), showing spectral shapes associated with dominance of cyanobacteria, green and red lineages of algae. Considering that  $a_\phi$  was retrieved from fitted  $a_d$ , an independent validation was performed against the total Chl *a* concentration (described later). In the blue end of the spectrum, we used

235 Gaussian decomposition (Hoepffner and Sathyendranath, 1991; Chase et al., 2013) to extract the magnitude associated with the  
 Soret band of Chl *a in vivo* (435 nm; Fig. 5A). In the red end of the spectrum, we compared the  $a_{\phi}(676)$  against models fitted  
 to global datasets (Nardelli and Twardowski, 2016, Fig. 5B). We note that our observations are similar to those reported by  
 Hoepffner and Sathyendranath (1991), while the value of Nardelli and Twardowski (2016) are lower likely due a combination  
 of their lower Chl *a* concentration range and their estimation of  $a_{\phi}(676)$  from the  $a_{nw}(650, 676, 715)$  line height (Roesler and  
 240 Barnard, 2013). The median spectral mass-specific *in vivo* pigment absorption coefficient at 676 nm,  $*a_{\phi}(676)$ , was equal to  
 $0.022 \text{ m}^2 \text{ mg}^{-1} \text{ nm}^{-1}$  (Fig. 5C), similar to the value of  $0.020 \text{ m}^2 \text{ mg}^{-1} \text{ nm}^{-1}$  presented by Hoepffner and Sathyendranath  
 (1991). The comparisons in the blue and red spectral regions we observe a linearity in log scale, with larger spread in the  
 comparison  $a_{\text{Gauss}}(435)$  likely due to a combination of the uncertainty in the estimate of the fitted  $a_d$  and the variable pigment  
 packaging effect (Morel and Bricaud, 1981; Latimer, 1983), both larger in the blue spectral range.


### 245 2.3.3 Near-forward ( $< 15^\circ$ ) particle volume scattering function

The near-forward particle volume scattering function (VSF;  $\beta_p$ ) at 670 nm was measured with two Laser In-Situ Scattering  
 and Transmissometry (LISST, Sequoia Scientific Inc.) instruments, models 200X and 100X, equipped with a detector type C.  
 The LISST-100X has a set of 32 concentric detectors measuring  $\beta_p$  between  $0.038^\circ$  and  $7.519^\circ$ , while the LISST-200X has  
 a set of 36 concentric detectors, the first 32 detectors measuring  $\beta_p$  between  $0.081^\circ$  and  $15.394^\circ$ . Single depth measurements  
 250 were taken at 1 m depth, with the instrument deployed horizontally (LISST-200X) or vertically (LISST-100X), having the  
 optical sensor end oriented down. The vertical deployment, preferentially made on the shady side of the platform, was chosen  
 to reduce the interference of environmental light since the model 100X does not compensate for external light. Considering  
 those deployment conditions and that our systems were mostly turbid, we expect negligible influence of environmental light  
 on the measurements. The instruments were used in a subset of stations. Calibrations with deionized water were performed  
 255 monthly and the most recent calibration used for blank values. The VSF was retrieved from the raw LISST data file, using the  
 procedures described in Agrawal (2005) and the instrument calibration files.

Quality control was performed by flagging entries in which the single scattering transmittance ( $T_b$ ) was lower than 30 %, to  
 avoid artifacts produced by multiple scattering:

$$T_b(670) = e^{-b_t(670, 0.018)l_{\text{LISST}}}, \quad (3)$$

260 where  $b_t$  is the total scattering coefficient and  $l_{\text{LISST}}$  is the pathlength of the LISST instrument. The non-water beam attenuation  
 coefficient at 670 nm measured by the LISST transmittance sensor with an acceptance angle of  $0.018^\circ$ ,  $c_{nw}(670, 0.018)$ , was  
 subtracted by the  $a_{nw}(670)$  measured with a benchtop spectrophotometer to calculate  $b_t(670, 0.018)$ . The VSF is an important  
 parameter for a tentative correction of the spectral particle beam attenuation and scattering coefficients measured with the  
 benchtop spectrophotometer.



**Figure 5.** Relation between the total chlorophyll *a* (monovinyl chlorophyll *a* + chlorophyllide *a*) and *in vivo* pigment absorption. (A) Magnitude of the Gaussian peak centered at 435 nm; (B) the *in vivo* pigment absorption at 676 nm,  $a_{\phi}(676)$ ; and (C) mass-specific *in vivo* pigment absorption at 676 nm,  $*a_{\phi}(676)$ . The  $a_{\phi}(676)$  for the Scheldt samples were estimated as the  $a_{\text{nw}}(650, 676, 715)$  line height (Roesler and Barnard, 2013).

#### 265 2.3.4 Beam attenuation and scattering from particulate components ( $> 0.45 \mu\text{m}$ )

In addition to the  $c_{\text{nw}}(670, 0.018)$  measured by the LISST instruments, the spectral particle beam attenuation ( $c_{\text{p}}$ ) and scattering ( $b_{\text{p}}$ ) coefficients were calculated from the  $c_{\text{nw}}$  measured on fresh samples with the Lambda 650S spectrophotometer, by subtracting  $a_{\text{g}}$  and  $a_{\text{g}}+a_{\text{p}}$ , respectively. To reduce the acceptance angle of the spectrophotometer (Boss et al., 2009b; Leymarie et al., 2010), a black barrier with a central circular aperture of 2.4 mm diameter was placed in the entry port of the integrating  
270 sphere. With a distance of 69.5 cm from the center of the sample cell to the integrating sphere (due to system of mirrors extending the pathlength), the detector acceptance angle in water for this configuration is  $0.074^\circ$ . The integration time of the instrument was increased to 0.4 s ( $150 \text{ nm min}^{-1}$ ) to compensate for the reduced signal, but noise levels were still noticed throughout the spectra. On later samples, the reference beam was partially blocked with a 1 % transmittance filter to reduce the dynamic range imposed by the lack of a similar aperture in the reference beam.

275 As with the  $a_{\text{g}}$ ,  $c_{\text{nw}}$  was measured in undiluted water subsamples, using 1 cm, 5 cm or 10 cm quartz cuvettes, depending on the turbidity. Blank readings were measured with deionized water, resulting in mismatched salinity for brackish and marine samples. The effect of the mismatched salinity was evaluated by comparing the beam attenuation between deionized water and artificial seawater at a salinity of 35 (NaCl at  $35 \text{ g L}^{-1}$ ) in the same experimental conditions from sample measurements. In the range from 380 nm to 850 nm, the difference in beam attenuation coefficient was  $\approx 0.12 \text{ m}^{-1} \text{ nm}^{-1}$  (Fig. S3, supplementary  
280 material). The difference spectrum was subtracted from all brackish and marine samples for an approximate correction.

The measurements of  $c_{\text{p}}(670, 0.074)$  made with the benchtop spectrophotometer were tested against the LISST determinations of  $c_{\text{p}}(670, 0.018)$ , with a similar method as described in Boss et al. (2009b). Measurements of  $c_{\text{nw}}$  from both instruments were converted to  $c_{\text{p}}$  by removing the absorption by CDOM at 670 nm. The difference between  $c_{\text{p}}(670)$  estimated by each instrument should be accounted for by the additional scattering signal captured by the larger acceptance angle of the spec-  
285 trophotometer, given by the integration of the VSF between  $0.018^\circ$  and  $0.074^\circ$ . The VSF between the lowest measured angle by the LISST ( $0.038^\circ$  or  $0.081^\circ$ ) and its transmittance acceptance angle ( $0.018^\circ$ ) was calculated with linear extrapolation in  $\log_{10}$  space from the two lowest measured angles. A detailed demonstration for sample DN\_29 is presented in Fig. S4 (supplementary material). Fig. 6A shows the result for all samples with paired data available, with data closely following the 1:1 line, and variation possibly caused by artifacts related to *in situ* versus *ex situ* measurements. This analysis validates the  $c_{\text{p}}$   
290 measurements with the spectrophotometer.

The additive error due to the contamination with the near-forward scattered signal within the acceptance angle can be accurately corrected for only when the VSF of each sample is known. However, an approximate correction is possible. Fig. 6B shows that the scattering signal in the range of  $0.018^\circ$  to  $0.074^\circ$  represents an average of 43.2 % of the particle scattering coefficient, with a stable relation across systems and seasons. Assuming that the angular shape of the VSF presents minor  
295 spectral variability and that the VSFs of different systems and seasons in our dataset are well represented by the subset for which LISST data is available, it is possible to use the approximate correction:

$$b_{\text{p}}(\lambda, 0.074) = c_{\text{p}}(\lambda, 0.074) - a_{\text{p}}(\lambda), \quad (4)$$

$$b_p(\lambda, 0.018) = \frac{b_p(\lambda, 0.074)}{1 - 0.432}, \quad (5)$$

300

$$c_p(\lambda, 0.018) = a_p(\lambda) + b_p(\lambda, 0.018). \quad (6)$$


The LISST-equivalent (acceptance angle of  $0.018^\circ$ ) version of the spectral  $b_p$  and  $c_p$  are also provided in conjunction with the measured values (acceptance angle of  $0.074^\circ$ ), as recommended by IOCCG (2018).

Quality control included the flagging of  $c_p$  spectra showing irregular behavior. Particle beam attenuation is a smooth spectral  
305 function (Roesler and Boss, 2003) due to anomalous diffraction that cause a complementary pattern between scattering and absorption (Zaneveld and Kitchen, 1995). Disturbances can be present in the form of small peaks caused by pigment absorption or oscillations caused by anomalous dispersion when there is a large contribution of small particles to the beam attenuation, but their shape and position are characteristic. Large, abrupt and otherwise irregular spectral behavior likely arises from motion of large particles in the beam cross sectional area. Six samples from Belgian lakes were flagged by this procedure, all with  
310 presence of large cyanobacteria aggregates. All BCZ samples of April, taken during a major *Phaeocystis* bloom, were also discarded due to similar effects of the large colonies. Beam attenuation remains elusive for standard bentchtop spectrophotometric analysis (sequential spectral scanning) under these conditions. As an example, cyanobacteria aggregates were observed floating on the surface of the cuvette at the end of the spectral run, even though the samples were mixed at the start of the measurement.

315 Another source of bias related to beam attenuation measurements in a spectrophotometer is that total internal reflection will occur for a fraction of the scattered light, dependent on the incident angles on the water side of the quartz wall. Some fraction of the internally reflected light could scatter back into directions in the detector's acceptance angle and artificially decrease beam attenuation measurements (IOCCG, 2018). Our experimental procedure did not include a dark baffle inside the cuvette to reduce this potential effect. Finally, our measurements show an oscillation centered at  $\approx 500$  nm, regardless of the system,  
320 particle type or concentration. This suggests that this oscillation is an artifact of the measurement procedure, though we could not identify its source. The spectral (LISST-equivalent) beam attenuation and particle scattering coefficients are presented in Fig. 7

### 2.3.5 Turbidity

Turbidity was quantified as formazin nephelometric units (FNU), a NIR side-scattering ( $90^\circ$ ) VSF magnitude relative to  
325 formazin standards. For the BCZ and Scheldt campaigns, turbidity was measured just after water sampling, while for the lakes and lagoon the turbidity was measured after transportation to the laboratory. The transportation might result in changes in particle aggregation due to changes in turbulence. However, we found a single linear relation between turbidity and suspended particulate matter (described later) across all systems and seasons.




**Figure 6.** Evaluation of  $c_p$  measurement bias due to the finite acceptance of the spectrophotometer. (A) Relation between the difference of the  $c_p$  measured with the spectrophotometer and the LISST, and the VSF integral in the range of solid angles defined by the difference of their acceptance angles. (B) Relation between the scattering coefficient ( $b_p = c_p - a_p$ ) and the magnitude of scattering between  $0.018^\circ$  and  $0.074^\circ$  scattering angles ( $\psi$ ).

## 2.4 Apparent Optical Properties

330 The AOP measured *in situ* were the Secchi disk depth ( $d_{\text{Secchi}}$ ) and the Lambert-equivalent water-leaving bi-hemispherical reflectance ( $\rho_{\text{wl}}^{\text{L}}$ ).  $\rho_{\text{wl}}^{\text{L}}$  was measured with the above water or on-water protocols, depending on the system. The Lambert-





**Figure 7.** LISST-equivalent (acceptance angle of  $0.018^\circ$ ) spectral particle beam attenuation coefficient,  $c_p$  (A), and integral-normalized particle scattering coefficient,  $b_p$  (B).

equivalent bi-hemispherical reflectance ( $\rho_s^L$ ) measurements of substrate samples (sediments, macroalgae) were performed *ex situ*, underwater and in air, using natural illumination.

The “Lambert-equivalent” qualification indicates that the bi-directional reflectance distribution function (BRDF) of the targets is assumed to be well represented by the Lambert model, and the hemispherical-directional measurement is converted to bi-hemispherical by scaling it with the cosine-weighted solid angle integral of an hemisphere,  $\pi$  sr. The water-leaving signal is not strictly Lambertian, however this approximation is commonly used for remote sensing purposes (*cf.* Frouin et al., 2019).

### 2.4.1 Secchi disk depth

A standard quadrant Secchi disk (black and white, 30 cm diameter) was used to measure transparency. The Secchi disk was  
340 deployed from the shady side of the sampling platform, recording the depths of disappearance and reappearance. The Secchi  
disk depth ( $d_{\text{Secchi}}$ ) was recorded as the average of the two depths. A tentative correction for the effect of sun zenith angle  
(described later) at the time of measurement (Lee et al., 2015) was applied following the formulation of Verschuur (1997).  
This correction normalizes the Secchi disk depth measurements to the Sun in the zenith. Measured and corrected values are  
provided. A comparison between turbidity the inverse Secchi disk depth, corrected for the Sun zenith angle, shows a log-linear  
345 relation and is presented in Fig. S5 (supplementary material).

### 2.4.2 Lambert-equivalent water-leaving bi-hemispherical reflectance

Reflectance spectroscopy measurements were performed in Belgian lakes and the coastal lagoon using the “on-water” method,  
also known as the skylight-blocked approach (Lee et al., 2013, 2019; Ruddick et al., 2019b). Measurements were made with  
a hand held spectrometer (FieldSpec HH, Analytical Spectral Devices) equipped with a 7.5° Field-of-View (FOV) foreoptics.  
350 The instrument has bands with 3.6 nm full width at half maximum (FWHM) and a spectral sampling of 1.6 nm, covering the  
range from 325 nm to 1075 nm. Water-leaving radiance ( $L_{\text{wl}}$ ) was recorded at 0.5 m horizontal distance from the deployment  
platform, aligned with the Sun azimuth, and with the opening of the lens’ cylindrical shield at 2.5 cm below the water surface.  
The global downwelling plane irradiance just above the surface ( $E_{\text{d}}(0^+)$ ) was estimated from near-coincident measurements  
of the exitant radiance of a sintered PTFE reference target (nominal reflectivity of 12 %), held parallel to the surface (Castagna  
355 et al., 2019; Ruddick et al., 2019a). An example of the measurement approach is presented in Fig. S6 (supplementary material).  
A total of 10  $L_{\text{wl}}$  spectra and 5 plaque exitant radiance spectra were averaged per station, with the measurement sequence  
completed within 2 min. The Lambert-equivalent water-leaving bi-hemispherical reflectance ( $\rho_{\text{wl}}^{\text{L}}$ ) was estimated according to:

$$\rho_{\text{wl}}^{\text{L}}(\theta, \Delta\phi, \lambda) = \pi \frac{L_{\text{wl}}^{\text{Meas}}(\theta, \Delta\phi, \lambda)}{(1 - \epsilon) E_{\text{d}}(\lambda)}, \quad (7)$$

where  $\epsilon$  is the estimated fractional shadowing error from the instrument and platform,  $\theta$  is the nadir angle ( $0^\circ$ ) and  $\Delta\phi$  here  
360 refers to the platform-sensor system relative azimuth to the Sun ( $0^\circ$ ). The superscript “Meas” for  $L_{\text{wl}}$  in Eq. 7 refers to the  
measured water-leaving signal, biased due to shadowing of the water system. Measurements were resampled to a regular 1 nm  
interval before Eq. 7 was applied. The shadowing error  $\epsilon$  was calculated for the range of IOPs observed in the lakes and  
lagoon and for the deployment from the inflatable boat and sensor as used in the campaigns, using a backward Monte Carlo  
radiative transfer code (unpublished). Formally,  $\epsilon$  is a function of the deployment setup, Sun zenith angle, diffuse fraction of  
365  $E_{\text{d}}$ ,  $a_{\text{t}}$  and  $c_{\text{t}}$ . However, the main IOP defining  $\epsilon$  is  $a_{\text{t}}$  (Fig. S7, supplementary material). The correction was applied using  
field observations of the required parameters. For stations without measurements of IOPs, spectral  $c_{\text{p}}$  was estimated from a  
multivariate linear regression against turbidity calculated over all available data and  $a_{\text{g}}$  was estimated as the median  $a_{\text{g}}$  of given  
water system. The diffuse fraction of  $E_{\text{d}}$  was simulated from the Sun zenith angle for clear skies (Castagna et al., 2019) and  
was set to 1 when field observations recorded clouds covering the direct Sun illumination.

370 The same spectrometer used in the lakes and lagoon campaigns was used in the Scheldt campaign, with the above water approach (Mobley, 1999; Ruddick et al., 2019b, a). For this measurement,  $\Delta\phi$  refers to the view direction relative to the Sun, set at  $135^\circ$ . The water-system radiance ( $L_{ws}$ ) was recorded at a nadir angle of  $40^\circ$ , the sky radiance ( $L_{sky}$ ) was measured at the nominal specular angle from  $L_{ws}$ ,  $140^\circ$ , and  $E_d$  estimated from the exitant radiance of the reference sintered PTFE plaque.  $\rho_{wl}^L$  was estimated according to:

$$375 \rho_{wl}^L(\theta, \Delta\phi, \lambda) = \pi \frac{L_{ws}(\theta, \Delta\phi, \lambda) - \tilde{\rho}_f L_{sky}(180^\circ - \theta, \Delta\phi, \lambda)}{E_d(\lambda)}, \quad (8)$$


where  $\tilde{\rho}_f$  is the effective Fresnel reflectance.

For the BCZ campaigns of 2018, reflectance spectroscopy measurements were also made with the above water approach, using a set of three spectroradiometers (VIS-ARC RAMSES, TriOS) fixed on the bow of the RV Simon Stevin (Castagna et al., 2021). The effective Fresnel reflectance was estimated from wind speed and  $L_{sky}$  according to Ruddick et al. (2006).  
 380 Further details of processing and quality control are described in Ruddick et al. (2006). The RAMSES instruments have a typical bandwidth of 10 nm FWHM with spectral sampling every  $\approx 3$  nm. Radiance and irradiance measurements were resampled to a regular 2.5 nm interval before Eq. 8 was applied, and  $\rho_{wl}^L$  was linearly interpolated to 1 nm interval to match the other reflectance data. The above water measurements were not corrected for possible disturbances by the platform in the spectroscopic measurements (Shang et al., 2020) and were not corrected for the non-nadir viewing geometry (Gleason et al.,  
 385 2012).

The available  $\rho_{wl}^L(0^\circ)$  and  $\rho_{wl}^L(40^\circ, 135^\circ)$  are presented in Fig. 8. As expected from the wide range of SPM concentrations,  $\rho_{wl}^L$  at 810 nm varied over three orders of magnitude, from 0.00036 to 0.10560. Similarly, the large diversity in terms of particle composition (*e.g.*, mineral fraction, taxonomy, pigment) and relative contribution of  $a_\phi$  translate to a diversity of  $\rho_{wl}^L$  spectral shapes. A validation of  $\rho_{wl}^L$  and the shadowing correction for the on water approach is provided in Fig. 9, by estimating turbidity  
 390 from  $\rho_{wl}^L(730)$  following the algorithm proposed by Nechad et al. (2010). The Nechad et al. (2010) algorithm was calibrated with data between  $\approx 1 \text{ g m}^{-3}$  and  $\approx 100 \text{ g m}^{-3}$  and the comparison is restricted to that range, covering all systems with the exception of the Scheldt.


### 2.4.3 Lambert-equivalent bi-hemispherical reflectance of bottom substrates

Measurements of reflectance spectroscopy of the surface of the sediment cores were made with the portable spectrometer  
 395 described previously (FieldSpec HH, Analytical Spectral Devices). The PMMA tubes were cut 5 cm above the sediment's surface level, and measurements made in nadir view, with 2.5 cm of water above the sediment. Measurements were performed with the on-water approach as described previously, *i.e.* an extension to the shield of the foreoptics was submerged to 2.5 cm below the water surface. A circular NIST-traceable Munsel card was used as a submerged reference, gently placed over the sediment, receiving the same illumination as the sediment's surface. An example of the measurement setup is shown in Fig. S8  
 400 (supplementary material). The bottom depth at the sampled stations varied between 1.3 m to 1.93 m and all but the station SP\_43 (1.3 m deep) had the Secchi disk depth above the bottom (average Secchi disk depth of 1.4 m). In July 2018, reflectance



**Figure 8.** Lambert-equivalent water-leaving bi-hemispherical reflectance,  $\rho_{wl}^L$ , estimated on Belgian water systems. The view geometry is dependent on the measurement approach, being nadir view ( $\theta = 0^\circ$ ) for lakes and the Spuikom, and  $\theta = 40^\circ$   $\Delta\phi = 135^\circ$  for the Scheldt and the BCZ. (A) Magnitude of  $\rho_{wl}^L$  and (B) its integral-normalization with respect to wavelength.  $\rho_{wl}^L(0^\circ)$  values measured with the on water approach presented here were corrected for shadowing.

spectroscopy measurements were also made of floating biofilm mats that had detached from the bottom of the Spuikom, presumably due to enhanced buoyancy caused by bubbles trapped in the mucilage matrix following several days of high irradiance and heat. Measurements were performed from a boat, without disturbing the floating biofilm mats. A sample taken




**Figure 9.** Assessment of  $\rho_{wl}^L$  data quality by comparing the turbidity estimated from reflectance using the algorithm of Nechad et al. (2010) to the measured turbidity.


405 for microscopy examination, revealed an assemblage composed of representatives of the benthic diatom genera *Pleurosigma*, *Gyrosigma* and *Navicula* (Fig. S9, supplementary material).

Reflectance spectroscopy measurements of the most conspicuous macrophyte species occurring in the Spuikom (*Cladophora glomerata*, *Ulva sp.* and *Sargassum muticum*) were performed in air, with an hyperspectral camera (SOC710-VP, Surface Optics Cooperation). The macrophyte reflectance measurements were made in nadir view with natural illumination. The  
 410 hyperspectral camera was set up at 1 m above the samples, using a f/2.8 aperture following the observation of lower spatial variability in the visible to red-edge spectral region. The 12 % (nominal) reflectivity sintered PTFE plaque was used to estimate  $E_d$  and a spectrally flat 5 % reflectivity sheet was used as background. Specimens were folded over a supporting petridish to avoid signal from the background and the reflectance averaged over circular areas to average over the three dimensional structures (Fig. S10, supplementary material). As the macroalgae were washed free of the sediment before the measurements,  
 415 the determined  $\rho_s^L$  represent pure end members of substrate reflectance.

The average  $\rho_s^L$  spectra of macroalgae and the spectra of sediment samples are presented in Fig. 10.  $\rho_s^L$  spectra of macroalgae reached values as high as 0.4 in the NIR range, while sediments, independent of biofilm thickness, had an average  $\rho_s^L$  in the NIR of  $\approx 0.11$ . In the visible range, the spectral shape of *Sargassum muticum* seems indistinguishable from the sediments with thickest biofilms, as expected due to similar pigment composition of brown algae and diatoms. The similarity in pigment com-  
 420 position also explains the similarity between the average  $\rho_s^L$  of healthy specimens of the green algae *Ulva sp.* and *Cladophora glomerata* (cf. Kotta et al., 2014).



**Figure 10.** Lambert-equivalent substrate bi-hemispherical reflectance,  $\rho_s^L$ , estimated from samples on the Spuikom. The view geometry was nadir view ( $\theta = 0^\circ$ ,  $\Delta\phi$  undefined) for all measurements. (A) The magnitude of  $\rho_s^L$  in the visible to NIR spectral range and (B) the magnitude in the visible spectral range. In (B) only the sediment samples with the thinnest and thickest biofilm mats are shown, as judged by the reflectance magnitude at 676 nm.



**Figure 11.** Relation between suspended particulate matter (SPM) concentration and turbidity in Belgian water systems.


## 2.5 Biogeochemical data

### 2.5.1 Suspended particulate matter and mineral fraction

Determinations of suspended particulate matter (SPM) concentration ( $\text{g m}^{-3}$ ) were performed gravimetrically, after filtration  
 425 of the suspensions with pre-treated glass fiber filters (GF/F, Whatman, nominal mesh size of  $0.7 \mu\text{m}$ ). For brackish and marine  
 samples, the filters were rinsed with distilled water to remove salt from the filtration area and rim of the filter (Strickland and  
 Parsons, 1968). The filters were then dried overnight at  $60^\circ\text{C}$  and cooled to room temperature before mass determinations.  
 The pre-treatment of filters involved combustion for one hour at  $450^\circ\text{C}$  to eliminate organic components, followed by washing  
 to remove loose glass fibers and blank mass determinations. To calculate the mineral fraction ( $f_m$ ) of the SPM, the filters  
 430 were heated to  $500^\circ\text{C}$  for one hour for thermal oxidation of organic matter and cooled to room temperature before new mass  
 determinations. For some filters, the last mass determinations were lower than the blank mass, indicating a combination of  
 very low mineral fraction and loss of glass fibers during manipulation. Those samples had the mineral fraction set to zero. The  
 observed range of SPM across all systems was from  $1.02 \pm 0.09 \text{ g m}^{-3}$  to  $791.19 \pm 0.10 \text{ g m}^{-3}$  and the range of mineral fraction  
 was from  $0 \pm 0.00$  to  $0.95 \pm 0.08$ . A comparison between SPM and turbidity is presented in Fig. 11.

### 435 2.5.2 Particle size distribution

The particle size distribution (PSD) was inverted from the LISST measurements of VSF, using the random shaped particle  
 VSF kernels provided by the manufacturer. The LISST-100X with random shaped particle inversion provides the PSD between  
 $1.8 \mu\text{m}$  and  $381 \mu\text{m}$  in 32 log-spaced intervals, while the LISST-200X provides the PSD between  $1 \mu\text{m}$  and  $500 \mu\text{m}$  in 36



**Figure 12.** Particle size distribution (PSD) in the Belgian water systems. (A) Retrieved with the LISST-100X; and (B) retrieved with the LISST-200X. The continuous particle number concentration PSD was calculated from the volume concentration PSD assuming spherical particles of median diameter in each size bin and normalizing by the width of each logarithmically spaced size bin.

log-spaced intervals. The LISST software automatically flags measurements with transmittance lower than 30 %. For further details of the measurements with the LISST, see Sec. 2.3.3. Values are provided as volume concentration PSD (cm<sup>3</sup> m<sup>-3</sup>, ppm). The PSDs converted to particle number concentration, using the volume of a sphere with the median diameter of each retrieved bin size range (Buonassissi and Dierssen, 2010), are shown in Fig. 12. Assessing differences in the PSD measured with the two instruments is under further investigation, but beyond the scope of this data report.




### 2.5.3 Pigment concentration

445 Pigment mass concentrations were determined using High Performance Liquid Chromatography (HPLC), following the method of Van Heukelem and Thomas (2001). Cells were broken with sonication and the suspension was cleared by filtration through a 0.22  $\mu\text{m}$  syringe filter. The HPLC was equipped with a reverse-phase column (Eclipse XDB C<sub>8</sub>) and the detection was performed with spectral absorption (Agilent 1100 series, Diode Array Detector). Pigment standards were acquired from the Danish Hydrographic Institute (DHI) and quantified pigments are listed in Table 6. For the BCZ samples, the measurements  
450 are part of the regular marine LifeWatch BE sampling campaigns (Flanders Marine Institute, 2021a), described in Mortelmans et al. (2019). The observed range of total Chl *a* (Chl *a* + Chlide *a*) across all systems was from 0.63 mg m<sup>-3</sup> to 387.53 mg m<sup>-3</sup>. Fig. 13 presents the distributions of total Chl *a* per system and the spread of Chl *b* to total Chl *c* (Chl *c*<sub>1</sub>*c*<sub>2</sub> + Chl *c*<sub>3</sub>).

### 2.5.4 DNA metabarcoding

DNA metabarcoding (amplicon sequencing) was performed only for samples from the Spuikom and the BCZ. The molecular  
455 analysis was based on replicate filters collected for pigment analysis. DNA extraction was performed with the DNeasy Plant Mini Kit (Qiagen), with the polymerase chain reaction (PCR) amplification targeting the variable region 4 (V4) of the nuclear 18S ribosomal RNA gene. The 18S rRNA V4 primers were the TAREuk454FWD1 (5' CCAGCASCYGGCGTAATTCC 3') and the TAREukREV3 (5' ACTTTCGTTCTTGATYRA 3'; Stoeck et al., 2010). Paired-end (2 x 300 base pairs) sequencing was performed with the Illumina MiSeq technology (Illumina, San Diego, US) by Genewiz (Leipzig, Germany). The primers were  
460 trimmed from the sequenced reads using the FASTX-Toolkit (Gordon and Hannon, 2010). The resulting base-pair sequences were processed with the DADA2 algorithm (Callahan et al., 2016) to resolve amplicon single variants (ASVs). Probable contaminant sequences were removed using negative controls, following the method of Davis et al. (2018). Taxonomic assignment to the ASVs was based on the Protist Ribosomal Reference database (PR<sup>2</sup> version 4.12; Guillou et al., 2012). The raw molecular data can be found at the Sequence Read Archive (SRA) of the National Center for Biotechnology Information (NCBI)  
465 under the accession number PRJNA778668 (<https://www.ncbi.nlm.nih.gov/sra/PRJNA778668>).


The raw data was further processed to an aggregation level that is relevant for optical monitoring. The assigned taxonomy with PR<sup>2</sup> was updated to follow the taxonomy of the World Register of Marine Species (WoRMS Editorial Board, 2021), aggregated to species rank and filtered to remove non-pigmented organisms. Heterotrophic organisms were filtered at division rank, with the exception of exclusively heterotrophic dinoflagellates, which were filtered at the lowest rank possible based  
470 on reference sources (Hasle et al., 1997; WoRMS Editorial Board, 2021). The data was further annotated to indicate: (1) the pigmentation group (*sensu* Jeffrey et al., 2011) of each species based on our pigment ratio database; and (2) the toxicity, based on the IOC-UNESCO HAB reference list (Moestrup et al., 2021). More than 200 different species from 19 classes of phytoplankton were identified.



**Figure 13.** Pigment concentrations in Belgian water systems. (A) Total chlorophyll *a* (TChl *a*) concentration; and (B) relation between monovinyl chlorophyll *b* (Chl *b*) and total chlorophyll *c* (TChl *c*) concentrations. TChl *a* = monovinyl chlorophyll *a* + chlorophyllide *a*, TChl *c* = chlorophyll *c*<sub>1c2</sub> + divinyl chlorophyll *c*<sub>3</sub> (cf. Table 6). A value of 0.01 mg m<sup>-3</sup> was added to represent in the log-log plot the samples with concentrations below the detection limit.

### 2.5.5 Flow imaging microscopy

475 As part of the regular LifeWatch BE samples in the BCZ, organisms in the range of 55 μm to 300 μm were counted and identified to the lowest taxonomic unit possible using flow imaging microscopy (FlowCam VS IV, Fluid Imaging Technologies,




**Figure 14.** Relative abundance of *Rhizosolenia spp.* cells as estimated by DNA metabarcoding (amplicon sequencing) and microscopy (FlowCam). Data normalized using the counts of common species of diatoms in each dataset.

Inc.; Amadei Martínez et al., 2020; Flanders Marine Institute, 2021b). Fresh water samples were fixed with lugol and stored in the dark until analysis. The images were analyzed with the VisualSpreadsheet software (Fluid Imaging Technologies, Inc.) for automatic taxonomic identification. Data was visually inspected to validate the classification. The fixation with lugol results in the disaggregation of *Phaeocystis globosa* colonies and single-cell *P. globosa* were not detected due to the lower size range of the instrument configuration. In total, 30 different species were detected, the majority of which were diatoms. A comparison of the spatio-temporal pattern of relative abundance of *Rhizosolenia spp.* cells between amplicon sequencing and FlowCam is shown in Fig. 14.

### 2.5.6 Bottom cover

For systems in which the bottom can be visible from the surface, the bottom type was described. Bottom cover was classified based on visual inspection from the surface or from images taken with a submersible camera (Fig. 15). For some stations the bottom cover was classified based on sampling of bottom material. The discrete classes were: (1) Sediment (potentially with biofilm); (2) Shells; (3) *Cladophora*; (4) *Sargassum*; (5) Brown algae (not specified); and (6) Heterogeneous. The last class was used when the bottom was covered by a complex mixture of other classes. This classification is qualitative in the sense that describes the major composition of the bottom at a given station, but does not provide fractional cover.



**Figure 15.** Example of bottom cover classes in the Spuiikom. (A) Sediment (with biofilm); (B) Sediment, shells and brown algae; (C) Heterogeneous.

## 2.6 Ancillary parameters

In addition to the optical and biogeochemical parameters, a series of ancillary parameters were also determined: Time, position, local depth, sampling platform, Sun zenith angle, and visual descriptions of the sky and water system.

For the inland water campaigns, time and position of each station were recorded from a handheld Global Navigation Satellite System (GNSS) receiver (GPSmap 62s, Garmin), which is enabled to receive corrections from the European Geostationary Navigation Overlay Service (EGNOS). The local depth at the time of sampling was estimated with a handheld single beam echosounder (Echotest II, Plastino) at the start of each station. The exception was one very shallow station (0.2 m), where a folding ruler was used. For marine stations, position, time and local depth were taken from the ship's navigation data, acquired with a differential GPS and a mounted single beam ecosounder (JFE 380-25, Japan Radio Co., Ltd.). Sun zenith angles were

500 calculated with the HORIZONS system (Jet Propulsion Laboratory, NASA; <https://ssd.jpl.nasa.gov/horizons/>). All ancillary parameters, with the exception of the Sun zenith angle, were combined into a single metadata file.

### 3 Conclusions and recommendations

This study described in detail the first open dataset of paired optical and biogeochemical measurements in a diverse set of water systems located in Belgium. The wide range of observed conditions and the relative scarcity of similar open datasets in inland waters make this a relevant contribution to the community. Potential users of this dataset are encouraged to contact the authors for further inquiries concerning the data and an updated status of studies in development using the dataset.

Although the raw, measured IOPs and AOPs are provided, their fitted or corrected versions are recommended for use. For example, the fitted  $a_g$  removes noise and oscillation in the NIR due to salinity mismatch between blanks and brackish or marine samples. The fitted  $a_d$  was found as the least biased way to remove the absorption signal of NaClO, providing the best estimate of  $a_d$  and  $a_\phi$ . The  $b_p$ , and consequentially  $c_p$ , corrected to an acceptance angle of  $0.018^\circ$  (equivalent to the LISST instruments) provides a more accurate estimate of the scattering and beam attenuation. The  $d_{\text{Secchi}}$  was corrected for the Sun zenith angle, normalizing all measurements to the Sun in the zenith, and  $\rho_{\text{wl}}^{\text{L}}$  measured with the on-water method were corrected for platform and instrument shadowing. The results of the fitting and corrections were evaluated through consistency checks between different data types and instruments as presented in the text.

### 515 4 Data availability

Data is available from Castagna et al. (2022), hosted at PANGAEA (<http://www.pangaea.de>) under the doi: <https://www.pangaea.de/tok/c67200d99ea9bbbeadd9edec9690f937b5bacbff>

*Author contributions.* AC, RD, and ID performed field campaigns in Belgian lakes. AC, HL, KR, MB, AD, and DD participated on field campaigns in the Spuikom, Scheldt and BCZ. LA joined the Sheldt and BCZ campaigns, JM joined BCZ campaigns and HD joined Spuikom campaigns. AC performed laboratory measurements of IOPs, field and laboratory measurements of AOP, processing of LISST data, measurements of turbidity, SPM and mineral fraction, classification of the bottom cover, curated the data and wrote the manuscript. MB performed reflectance spectroscopy measurements in the BCZ. ID and RD performed HPLC analysis of pigments. RB and AC performed analysis of DNA metabarcoding. LA performed flow microscopy counts. HD, KS and WV provided infrastructure, planning and methodological discussion. All authors reviewed the manuscript.

525 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research was funded by BELSPO Stereo III projects PONDER (SR/00/325), HYPERMAQ (SR/00/335) and and PHYTOBEL (SR/02/213). H.D. acknowledges funding by NASA Ocean Biology and Biochemistry through the Plankton, Aerosol, Cloud, ocean Ecosystem (PACE) project (NNX15AC32G). We are thankful to the RV Simon Stevin crew for support during LifeWatch BE campaigns, to Darja Belišová, Heidi Tantu, Kathryn Morrissey, Reinhoud de Blok, and Tine Verstraete for help in selected field campaigns, to 530 Sofie D'hondt for the extraction and amplification of the bulk plankton DNA, to Willem Stock for identification of benthic diatoms in the floating biofilms, to Olivier De Clerck for identification of macroalgae, to Koen De Rycker for operating the short corer, to Dieter Vansteewegen and André Cattrijsse for support during the BCZ and two extended campaigns in the Spuikom, and to the Café Zates for support during the Scheldt campaigns. We are appreciative of all researchers and steering committee members of the PONDER and HYPERMAQ projects for the interesting discussions along the development of this research. We acknowledge the R Core team and the authors of the R 535 packages for developing and maintaining the free software used in this research. BCZ data provided as part of the Flemish contribution to the LifeWatch ESFRI by the Flanders Marine Institute (VLIZ).

## References

- Adler, D. and Kelly, S. T.: vioplot: violin plot, <https://github.com/TomKellyGenetics/vioplot>, r package version 0.3.6, 2020.
- 540 Agrawal, Y. C.: The optical volume scattering function: Temporal and vertical variability in the water column off the New Jersey coast, *Limnology and Oceanography*, 50, 1787–1794, <https://doi.org/10.4319/lo.2005.50.6.1787>, 2005.
- Amadei Martínez, L., Mortelmans, J., Dillen, N., Debusschere, E., and Deneudt, K.: LifeWatch observatory data: phytoplankton observations in the Belgian Part of the North Sea, *Biodiversity Data Journal*, 8, e57 236, <https://doi.org/10.3897/BDJ.8.e57236>, 2020.
- Astoreca, R., Ruddick, K., Rousseau, V., Mol, B., Parent, J.-Y., and Lancelot, C.: Variability of the inherent and apparent optical properties in a highly turbid coastal area: impact on the calibration of remote sensing algorithms, *EARSeL eProceedings*, 5, 1–17, 2006.
- 545 Astoreca, R., Rousseau, V., and Lancelot, C.: Coloured dissolved organic matter (CDOM) in Southern North Sea waters: Optical characterization and possible origin, *Estuarine, Coastal and Shelf Science*, 85, 633–640, <https://doi.org/10.1016/j.ecss.2009.10.010>, 2009.
- Astoreca, R., Doxaran, D., Ruddick, K., Rousseau, V., and Lancelot, C.: Influence of suspended particle concentration, composition and size on the variability of inherent optical properties of the Southern North Sea, *Continental Shelf Research*, 35, 117–128, <https://doi.org/10.1016/j.csr.2012.01.007>, 2012.
- 550 Binding, C. E., Jerome, J. H., Bukata, R. P., and Booty, W. G.: Spectral absorption properties of dissolved and particulate matter in Lake Erie, *Remote Sensing of Environment*, 112, 1702–1711, <https://doi.org/10.1016/j.rse.2007.08.017>, 2008.
- Boss, E., Taylor, L., Gilbert, S., Gundersen, K., Hawley, N., Janzen, C., Johengen, T., Purcell, H., Robertson, C., Schar, D. W. H., Smith, G. J., and Tamburri, M. N.: Comparison of inherent optical properties as a surrogate for particulate matter concentration in coastal waters, *Limnology and Oceanography: Methods*, 7, 803–810, <https://doi.org/10.4319/lom.2009.7.803>, 2009a.
- 555 Boss, E. S., Slade, W. H., Behrenfeld, M. J., and Dall’Olmo, G.: Acceptance angle effects on the beam attenuation in the ocean, *Optics Express*, 17, 1535–1550, <https://doi.org/10.1364/OE.17.001535>, 2009b.
- Buonassissi, C. J. and Dierssen, H. M.: A regional comparison of particle size distributions and the power law approximation in oceanic and estuarine surface waters, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2010JC006256>, 2010.
- Cael, B. B. and Boss, E. S.: Simplified model of spectral absorption by non-algal particles and dissolved organic materials in aquatic 560 environments, *Optics Express*, 25, 25 486, <https://doi.org/10.1364/OE.25.025486>, 2017.
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon data, *Nature Methods*, 13, 581–583, <https://doi.org/10.1038/nmeth.3869>, 2016.
- Castagna, A., Carol Johnson, B., Voss, K. J., Dierssen, H. M., Patrick, H., Germer, T. A., Sabbe, K., and Vyverman, W.: Uncertainty in 565 global downwelling plane irradiance estimates from sintered polytetrafluoroethylene plaque radiance measurements, *Applied Optics*, 58, 4497–4511, <https://doi.org/10.1364/AO.58.004497>, 2019.
- Castagna, A., Simis, S. G. H., Dierssen, H., Vanhellefont, Q., Sabbe, K., and Vyverman, W.: Extending Landsat 8: Retrieval of an Orange contra-Band for Inland Water Quality Applications, *Remote Sensing*, 12, 637, <https://doi.org/10.3390/rs12040637>, 2020.
- Castagna, A., Dierssen, H., Organelli, E., Bogorad, M., Mortelmans, J., Vyverman, W., and Sabbe, K.: Optical Detection of Harmful Algal Blooms in the Belgian Coastal Zone: A Cautionary Tale of Chlorophyll  $c_3$ , *Frontiers in Marine Science*, 8, 1892, 570 <https://doi.org/10.3389/fmars.2021.770340>, 2021.
- Castagna, A., Amadei Martínez, L., Bogorad, M., Daveloose, I., Dasseville, R., Dierssen, H. M., Beck, M., Mortelmans, J., Lavigne, H., Dogliotti, A., Doxaran, D., Ruddick, K., Vyverman, W., and Sabbe, K.: Dataset of Optical and biogeochemical properties of diverse Belgian inland and coastal waters, PANGAEA, <https://doi.org/10.1594/PANGAEA.940240>, dataset, 2022.

- Chase, A., Boss, E. S., Zaneveld, R., Bricaud, A., Claustre, H., Ras, J., Dall'Olmo, G., and Westberry, T. K.: Decomposition of in situ  
575 particulate absorption spectra, *Methods in Oceanography*, 7, 110–124, <https://doi.org/10.1016/j.mio.2014.02.002>, 2013.
- Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., and Callahan, B. J.: Simple statistical identification and removal of contaminant  
sequences in marker-gene and metagenomics data, *Microbiome*, 6, 226, <https://doi.org/10.1186/s40168-018-0605-2>, 2018.
- Descy, J.-P., Pirlot, S., Verniers, G., Viroux, L., Lara, Y., Wilmotte, A., Vyverman, W., Vanormelingen, P., Van Wichelen, J., Van Gremberghe,  
I., Triest, L., Peretyatko, A., Everbecq, E., and Codd, G.: B-BLOOMS 2 - Cyanobacterial blooms: toxicity, diversity, modeling and  
580 management, Tech. rep., Belgian Science Policy, Brussels, Belgium, 2011.
- Desmit, X., Nohe, A., Borges, A. V., Prins, T., De Cauwer, K., Lagring, R., Van der Zande, D., and Sabbe, K.: Changes in chlorophyll  
concentration and phenology in the North Sea in relation to de-eutrophication and sea surface warming, *Limnology and Oceanography*,  
65, 828–847, <https://doi.org/10.1002/lno.11351>, 2020.
- Dierssen, H., Bracher, A., Brando, V., Loisel, H., and Ruddick, K.: Data Needs for Hyperspectral Detection of Algal Diversity Across the  
585 Globe, *Oceanography*, 33, 74–79, <https://doi.org/10.5670/oceanog.2020.111>, 2020.
- Dogliotti, A. I., Ruddick, K. G., Nechad, B., Doxaran, D., and Knaeps, E.: A single algorithm to retrieve turbidity from remotely-sensed data  
in all coastal and estuarine waters, *Remote Sensing of Environment*, 156, 157–168, <https://doi.org/10.1016/j.rse.2014.09.020>, 2015.
- Estapa, M. L., Boss, E., Mayer, L. M., and Roesler, C. S.: Role of iron and organic carbon in mass-specific light absorption by particulate mat-  
ter from Louisiana coastal waters, *Limnology and Oceanography*, 57, 97–112, <https://doi.org/https://doi.org/10.4319/lo.2012.57.1.0097>,  
590 2012.
- Ferrari, G. M. and Tassan, S.: A method using chemical oxidation to remove light absorption by phytoplankton pigments, *Journal of Phycol-  
ogy*, 35, 1090–1098, <https://doi.org/10.1046/j.1529-8817.1999.3551090.x>, 1999.
- Fettweis, M. and Van den Eynde, D.: The mud deposits and the high turbidity in the Belgian–Dutch coastal zone, southern bight of the North  
Sea, *Continental Shelf Research*, 23, 669–691, [https://doi.org/10.1016/S0278-4343\(03\)00027-X](https://doi.org/10.1016/S0278-4343(03)00027-X), 2003.
- 595 Flanders Marine Institute: LifeWatch observatory data: nutrient, pigment, suspended matter and secchi measurements in the Belgian Part of  
the North Sea, <https://doi.org/10.14284/441>, accessed: 2021-08-31, 2021a.
- Flanders Marine Institute: LifeWatch observatory data: phytoplankton observations by imaging flow cytometry (FlowCam) in the Belgian  
Part of the North Sea, <https://doi.org/10.14284/527>, accessed: 2021-08-31, 2021b.
- Frouin, R. J., Franz, B. A., Ibrahim, A., Knobelspiesse, K., Ahmad, Z., Cairns, B., Chowdhary, J., Dierssen, H. M., Tan, J., Dubovik, O.,  
600 Huang, X., Davis, A. B., Kalashnikova, O., Thompson, D. R., Remer, L. A., Boss, E., Coddington, O., Deschamps, P.-Y., Gao, B.-C.,  
Gross, L., Hasekamp, O., Omar, A., Pelletier, B., Ramon, D., Steinmetz, F., and Zhai, P.-W.: Atmospheric Correction of Satellite Ocean-  
Color Imagery During the PACE Era, *Frontiers in Earth Science*, 7, 145, <https://doi.org/10.3389/feart.2019.00145>, 2019.
- Gleason, A. C., Voss, K. J., Gordon, H. R., Twardowski, M., Sullivan, J., Trees, C., Weidemann, A., Berthon, J.-F., Clark, D.,  
and Lee, Z.: Detailed validation of the bidirectional effect in various Case I and Case II waters, *Optics Express*, 20, 7630,  
605 <https://doi.org/10.1364/OE.20.007630>, 2012.
- Gordon, A. and Hannon, G.: FASTX-Toolkit, [http://hannonlab.cshl.edu/fastx\\_toolkit/index.html](http://hannonlab.cshl.edu/fastx_toolkit/index.html), version 0.0.13, 2010.
- Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de Vargas, C., Decelle, J., del Campo, J.,  
Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H., Lara, E., Le Bescot, N., Logares, R., Mahé, F., Massana, R.,  
Montresor, M., Morard, R., Not, F., Pawlowski, J., Probert, I., Sauvadet, A.-L., Siano, R., Stoeck, T., Vaulot, D., Zimmermann, P., and  
610 Christen, R.: The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with  
curated taxonomy, *Nucleic Acids Research*, 41, D597–D604, <https://doi.org/10.1093/nar/gks1160>, 2012.



- Hasle, G. R., Steidinger, K. A., Syvertsen, E. E., Jansen, K., Jhrondsen, J., and Heimdahl, B. R.: Identifying Marine Phytoplankton, Elsevier, San Diego, California, <https://doi.org/10.1016/B978-0-12-693018-4.X5000-9>, 1997.
- 615 Hoepffner, N. and Sathyendranath, S.: Effect of pigment composition on absorption properties of phytoplankton, *Marine Ecology Progress Series*, 73, 11–23, <https://doi.org/10.3354/meps073011>, 1991.
- IOCCG: Inherent Optical Property Measurements and Protocols: Absorption Coefficient, vol. 1 of *IOCCG Ocean Optics and Biogeochemistry Protocols for Satellite Ocean Colour Sensor Validation*, IOCCG, Dartmouth, NS, Canada, 2018.
- ISO 7027:1999: Water quality – Determination of turbidity, Standard, International Organization for Standardization, Geneva, CH, 1999.
- Jeffrey, S. W., Wright, S. W., and Zapata, M.: Microalgal classes and their signature pigments, in: *Phytoplankton Pigments: Characterization, Chemotaxonomy, and Applications in Oceanography*, edited by Roy, S., Llewellyn, C. A., Egeland, E. S., and Johnsen, G., chap. 1, pp. 3–77, Cambridge University Press, Cambridge, UK, 2011.
- 620 Jonaz, M. and Fournier, G. R.: *Light Scattering by Particles in Water: Theoretical and Experimental Foundations*, Elsevier, Amsterdam, The Netherlands, <https://doi.org/10.1016/B978-0-12-388751-1.X5000-5>, 2007.
- Kotta, J., Remm, K., Vahtmäe, E., Kutser, T., and Orav-Kotta, H.: In-air spectral signatures of the Baltic Sea macrophytes and their statistical separability, *Journal of Applied Remote Sensing*, 8, 1 – 14, <https://doi.org/10.1117/1.JRS.8.083634>, 2014.
- 625 Latimer, P.: The deconvolution of absorption spectra of green plant materials -Improved corrections for the sieve effect, *Photochemistry and Photobiology*, 38, 731–734, <https://doi.org/10.1111/j.1751-1097.1983.tb03608.x>, 1983.
- Lee, Z., Pahlevan, N., Ahn, Y.-H., Greb, S., and O'Donnell, D.: Robust approach to directly measuring water-leaving radiance in the field, *Applied Optics*, 52, 1693–1701, <https://doi.org/10.1364/AO.52.001693>, 2013.
- 630 Lee, Z., Shang, S., Hu, C., Du, K., Weidemann, A., Hou, W., Lin, J., and Lin, G.: Secchi disk depth: A new theory and mechanistic model for underwater visibility, *Remote Sensing of Environment*, 169, 139–149, <https://doi.org/10.1016/j.rse.2015.08.002>, 2015.
- Lee, Z., Wei, J., Shang, Z., Garcia, R., Dierssen, H. M., Ishizaka, J., and Castagna, A.: On-Water Radiometry Measurements: Skylight-Blocked Approach and Data Processing (Appendix to IOCCG Protocol Series 2019), Tech. Rep. December, 2019.
- Leymarie, E., Doxaran, D., and Babin, M.: Uncertainties associated to measurements of inherent optical properties in natural waters, *Applied Optics*, 49, 5415–5436, <https://doi.org/10.1364/AO.49.005415>, 2010.
- 635 Max, J.-J. and Chapados, C.: IR spectroscopy of aqueous alkali halide solutions: Pure salt-solvated water spectra and hydration numbers, *The Journal of Chemical Physics*, 115, 2664–2675, <https://doi.org/10.1063/1.1337047>, 2001.
- Meire, P., Ysebaert, T., Van Damme, S., Van Den Bergh, E., Maris, T., and Struyf, E.: The Scheldt estuary: A description of a changing ecosystem, *Hydrobiologia*, 540, 1–11, <https://doi.org/10.1007/s10750-005-0896-8>, 2005.
- 640 Mobley, C. D.: Estimation of the remote-sensing reflectance from above-surface measurements, *Applied Optics*, 38, 7442, <https://doi.org/10.1364/AO.38.007442>, 1999.
- Moestrup, Ø., Akselmann-Cardella, R., Churro, C., Fraga, S., Hoppenrath, M., Iwataki, M., Larsen, J., Lundholm, N., and Zingone, A.: IOC-UNESCO Taxonomic Reference List of Harmful Micro Algae, <http://www.marinespecies.org/hab>, <https://doi.org/10.14284/362>, accessed on 2021-07-26, 2021.
- 645 Morel, A. Y. and Bricaud, A.: Theoretical results concerning light absorption in a discrete medium, and application to specific absorption of phytoplankton, *Deep Sea Research Part A. Oceanographic Research Papers*, 28, 1375–1393, [https://doi.org/10.1016/0198-0149\(81\)90039-X](https://doi.org/10.1016/0198-0149(81)90039-X), 1981.
- Mortelmans, J., Deneudt, K., Cattrijsse, A., Beauchard, O., Daveloose, I., Vyverman, W., Vanaverbeke, J., Timmermans, K., Peene, J., Roose, P., Knockaert, M., Chou, L., Sanders, R., Stinchcombe, M., Kimpe, P., Lammens, S., Theetaert, H., Gkritzalis, T., Hernandez, F.,

- 650 and Mees, J.: Nutrient, pigment, suspended matter and turbidity measurements in the Belgian part of the North Sea, *Scientific Data*, 22, <https://doi.org/10.1038/s41597-019-0032-7>, 2019.
- Nardelli, S. C. and Twardowski, M. S.: Assessing the link between chlorophyll concentration and absorption line height at 676 nm over a broad range of water types, *Optics Express*, 24, A1374, <https://doi.org/10.1364/OE.24.0A1374>, 2016.
- Nechad, B., Ruddick, K. G., and Park, Y.: Calibration and validation of a generic multisensor algorithm for mapping of total suspended matter in turbid waters, *Remote Sensing of Environment*, 114, 854–866, <https://doi.org/10.1016/j.rse.2009.11.022>, 2010.
- 655 Pegau, W. S., Zaneveld, J. R. V., Mitchell, B. G., Mueller, J. L., Kahru, M., Wieland, J., and Stramska, M.: *Inherent Optical Properties: Instruments, Characterizations, Field Measurements and Data Analysis Protocols*, vol. IV, NASA, 2002.
- Quan, X. and Fry, E. S.: Empirical equation for the index of refraction of seawater, *Applied Optics*, 34, 3477, <https://doi.org/10.1364/AO.34.003477>, 1995.
- 660 R Core Team: *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2020.
- Reid, P. C., Lancelot, C., Gieskes, W. W. C., Hagmeier, E., and Weichart, G.: Phytoplankton of the North Sea and its dynamics: A review, *Netherlands Journal of Sea Research*, 26, 295–331, [https://doi.org/10.1016/0077-7579\(90\)90094-W](https://doi.org/10.1016/0077-7579(90)90094-W), 1990.
- Roesler, C. S. and Barnard, A. H.: Optical proxy for phytoplankton biomass in the absence of photophysiology: Rethinking the absorption line height, *Methods in Oceanography*, 7, 79–94, <https://doi.org/10.1016/j.mio.2013.12.003>, 2013.
- 665 Roesler, C. S. and Boss, E. S.: Spectral beam attenuation coefficient retrieved from ocean color inversion, *Geophysical Research Letters*, 30, 1468, <https://doi.org/10.1029/2002GL016185>, 2003.
- Röttgers, R., Dupouy, C., Taylor, B. B., Bracher, A., and Woźniak, S. B.: Mass-specific light absorption coefficients of natural aquatic particles in the near-infrared spectral region, *Limnology and Oceanography*, 59, 1449–1460, <https://doi.org/10.4319/lo.2014.59.5.1449>, 2014a.
- 670 Röttgers, R., McKee, D., and Utschig, C.: Temperature and salinity correction coefficients for light absorption by water in the visible to infrared spectral region, *Optics Express*, 22, 25 093, <https://doi.org/10.1364/OE.22.025093>, 2014b.
- Roy, S., Llewellyn, C. A., Egeland, E. S., and Johnsen, G., eds.: *Phytoplankton Pigments: Characterization, Chemotaxonomy, and Applications in Oceanography*, Cambridge University Press, Cambridge, UK, 2011.
- 675 Ruddick, K. G., Cauwer, V. D., Park, Y.-J., and Moore, G.: Seaborne measurements of near infrared water-leaving reflectance : The similarity spectrum for turbid waters, *Limnology and Oceanography*, 51, 1167–1179, <https://doi.org/10.4319/lo.2006.51.2.1167>, 2006.
- Ruddick, K. G., Voss, K. J., Banks, A., Boss, E., Castagna, A., Frouin, R., Hieronymi, M., Jamet, C., Johnson, B., Kuusk, J., Lee, Z., Ondrusek, M., Vabson, V., and Vendt, R.: A review of protocols for Fiducial Reference Measurements of downwelling irradiance for the validation of satellite remote sensing data over water, *Remote Sensing*, 11, <https://doi.org/10.3390/rs11151742>, 2019a.
- 680 Ruddick, K. G., Voss, K. J., Banks, A. C., Boss, E. S., Castagna, A., Frouin, R., Hieronymi, M., Jamet, C., Johnson, B. C., Kuusk, J., Lee, Z., Ondrusek, M., Vabson, V., and Vendt, R.: A Review of Protocols for Fiducial Reference Measurements of Downwelling Irradiance for the Validation of Satellite Remote Sensing Data over Water, *Remote Sensing*, 11, 1742, <https://doi.org/10.3390/rs11151742>, 2019b.
- Shang, Z., Lee, Z., Wei, J., and Lin, G.: Impact of ship on radiometric measurements in the field: a reappraisal via Monte Carlo simulations, *Optics Express*, 28, 1439, <https://doi.org/10.1364/OE.28.001439>, 2020.
- 685 Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D., Breiner, H. W., and Richards, T. A.: Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water, *Molecular Ecology*, 19, 21–31, <https://doi.org/10.1111/j.1365-294X.2009.04480.x>, 2010.

- Stramski, D., Reynolds, R. A., Kaczmarek, S., Uitz, J., and Zheng, G.: Correction of pathlength amplification in the filter-pad technique for measurements of particulate absorption coefficient in the visible spectral region, *Applied Optics*, 54, 6763, <https://doi.org/10.1364/AO.54.006763>, 2015.
- 690 Strickland, J. D. H. and Parsons, T. R.: *A Practical Handbook of Seawater Analysis*, Fisheries Research Board of Canada, Ottawa, Canada, 1968.
- Twardowski, M. S., Boss, E., Sullivan, J. M., and Donaghay, P. L.: Modeling the spectral shape of absorption by chromophoric dissolved organic matter, *Marine Chemistry*, 89, 69–88, <https://doi.org/10.1016/j.marchem.2004.02.008>, 2004.
- 695 van Beusekom, J. and Diel-Christiansen, S.: *A synthesis of phyto and zooplankton dynamics in the North Sea environment*, Tech. rep., 1993.
- Van Heukelem, L. and Thomas, C. S.: Computer-assisted high-performance liquid chromatography method development with applications to the isolation and analysis of phytoplankton pigments, *Journal of Chromatography A*, 910, 31–49, [https://doi.org/10.1016/S0378-4347\(00\)00603-4](https://doi.org/10.1016/S0378-4347(00)00603-4), 2001.
- Venables, W. N. and Ripley, B. D.: *Modern Applied Statistics with S*, Springer, New York, fourth edn., <http://www.stats.ox.ac.uk/pub/MASS4>, ISBN 0-387-95457-0, 2002.
- 700 Verschuur, G. L.: Transparency Measurements in Garner Lake, Tennessee: The Relationship between Secchi Depth and Solar Altitude and a Suggestion for Normalization of Secchi Depth Data, *Lake and Reservoir Management*, 13, 142–153, <https://doi.org/10.1080/07438149709354305>, 1997.
- WoRMS Editorial Board: *World Register of Marine Species (WoRMS)*, <https://doi.org/10.14284/170>, accessed: 2021-06-02, 2021.
- 705 Zaneveld, J. R. V. and Kitchen, J. C.: The variation in the inherent optical properties of phytoplankton near an absorption peak as determined by various models of cell structure, *Journal of Geophysical Research*, 100, 13 309, <https://doi.org/10.1029/95JC00451>, 1995.

**Table 1.** Description of relevant acronyms, symbols, constants and subscripts used in this manuscript.

|            | Description  |   |
|------------|--|---|
| Acronyms   | AOP  | Apparent optical properties   |
|            | BCZ  | Belgian coastal zone  |
|            | BRDF   | Bidirectional reflectance distribution function   |
|            | CDOM   | Chromophoric dissolved organic matter   |
|            | FNU  | Formazin nephelometric units  |
|            | FWHM   | Full-Width-at-Half-Maximum  |
|            | IOP  | Inherent optical properties   |
|            | NIR  | Near-infrared range of the electromagnetic spectrum (700 nm to 1000 nm)   |
|            | PMMA   | Polymethyl methacrylate   |
|            | PSD  | Particle size distribution  |
|            | PTFE   | Polytetrafluoroethylene   |
|            | SPM  | Suspended particulate matter  |
|            | UV   | Ultraviolet range of the electromagnetic spectrum (10 nm to 400 nm)   |
| Symbols    | $a$  | Spectral Napierian absorption coefficient ( $\text{m}^{-1} \text{nm}^{-1}$ )  |
|            | $b$  | Spectral Napierian scattering coefficient ( $\text{m}^{-1} \text{nm}^{-1}$ )  |
|            | $c$  | Spectral Napierian beam attenuation coefficient ( $\text{m}^{-1} \text{nm}^{-1}$ )  |
|            | $d_{\text{Secchi}}$                                  | Secchi disk depth (m)   |
|            | $E_d(0^+)$   | Spectral global (direct + diffuse) downwelling plane irradiance just above the water surface ( $\text{W m}^{-2} \text{nm}^{-1}$ ) |
|            | $f_m$  | Mineral fraction of SPM (1)   |
|            | $l$  | Metric pathlength (m)   |
|            | $L$  | Spectral radiance ( $\text{W m}^{-2} \text{sr}^{-1} \text{nm}^{-1}$ )   |
|            | $S_h$  | Spectral hyperbolic slope coefficient (1)   |
|            | $S_e$  | Spectral exponential slope coefficient ( $\text{nm}^{-1}$ )   |
|            | $T$  | Transmittance (1)   |
|            | $\alpha_d$   | Spectral Napierian depigmented particle absorption coefficient offset ( $\text{m}^{-1} \text{nm}^{-1}$ )                          |
|            | $\beta$  | Spectral Napierian volume scattering function ( $\text{m}^{-1} \text{sr}^{-1} \text{nm}^{-1}$ )                                   |
|            | $\Delta\phi$   | Relative azimuth angle ( $^\circ$ )   |
|            | $\epsilon$   | Fractional error caused by shadowing of the downwelling light for on water spectroscopy measurements (1)                          |
|            | $\theta$   | Nadir angle ( $^\circ$ )  |
|            | $\lambda$  | Wavelength (nm)   |
|            | $\rho^L$   | Lambert-equivalent bi-hemispherical reflectance (1)   |
|            | $\tilde{\rho}_r$                                     | Effective Fresnel reflectance (1)   |
|            | $\psi$   | Polar scattering angle ( $^\circ$ or rad)   |
| Constants  | $e$  | Euler's number ( $\approx 2.718282$ )   |
|            | $\pi$  | The ratio of a circle's circumference to its diameter ( $\approx 3.141593$ )  |
| Subscripts | b  | Relative to backscattering  |
|            | d  | Relative to the depigmented particle components   |
|            | g  | Relative to the CDOM component  |
|            | nw   | Relative to all non-water components  |
|            | p  | Relative to the particle components   |
|            | s  | Relative to the substrate (macroalgae, sediments)   |
|            | t  | Relative to all components (total)  |
|            | wl   | Relative to water-leaving   |
| ws         | Relative to water-system (water-leaving + interface) |   |
| $\phi$     | Relative to the <i>in vivo</i> pigments component    |   |

**Table 2.** Summary of Belgian water systems sampled in this study.

| Type     | System       | Platform        | Period    |
|----------|--------------|-----------------|-----------|
| Fresh    | Bocht        | Boat, Pontoon   | 2017      |
|          | Donkmeer     | Boat, Pontoon   | 2017-2018 |
|          | DelMare      | Boat            | 2017      |
|          | Dikkebus     | Pontoon         | 2018      |
|          | Hazewinkel   | Boat, Pontoon   | 2017      |
|          | Leuven-Dijle | Pontoon         | 2019      |
|          | Molsbroek    | Boat            | 2017      |
|          | Nieuwdonk    | Pontoon         | 2017      |
|          | Zillebeke    | Boat            | 2018      |
| Brackish | Scheldt      | Pontoon         | 2019      |
|          | Spuikom      | Pontoon, Boat   | 2017-2019 |
| Marine   | BCZ          | RV Simon Stevin | 2018      |

**Table 3.** Summary of measured and derived inherent optical properties per water system. Values indicate the number of valid, quality checked, measurements.

| Type         | System       | $a_g$      | $a_p$      | $a_d$              | $a_\phi$   | $c_p$     | $b_p$     | $\beta_p$                  | Turb       |
|--------------|--------------|------------|------------|--------------------|------------|-----------|-----------|----------------------------|------------|
|              |              |            |            | $(m^{-1} nm^{-1})$ |            |           |           | $(m^{-1} sr^{-1} nm^{-1})$ | (FNU)      |
| Fresh        | Bocht        | 1          | 7          | 7                  | 7          | -         | -         | 2                          | 8          |
|              | Donkmeer     | 18         | 33         | 32                 | 32         | 8         | 8         | 10                         | 38         |
|              | DelMare      | -          | 2          | 2                  | 2          | -         | -         | 1                          | 3          |
|              | Dikkebus     | 7          | 7          | 7                  | 7          | 3         | 3         | 5                          | 7          |
|              | Hazewinkel   | 1          | 17         | 17                 | 17         | -         | -         | 1                          | 18         |
|              | Leuven-Dijle | 2          | 2          | -                  | -          | 2         | 2         | -                          | 2          |
|              | Molsbroek    | -          | 1          | 1                  | 1          | -         | -         | -                          | 1          |
|              | Nieuwdonk    | -          | 1          | 1                  | 1          | -         | -         | -                          | 2          |
|              | Zillebeke    | 8          | 8          | 8                  | 8          | 3         | 3         | 6                          | 8          |
| Brackish     | Scheldt      | 21         | 21         | -                  | -          | 21        | 21        | 2                          | 21         |
|              | Spuikom      | 35         | 51         | 46                 | 46         | 21        | 21        | 15                         | 60         |
| Marine       | BCZ          | 9          | 19         | 19                 | 19         | -         | -         | 19                         | 19         |
| <b>Total</b> |              | <b>102</b> | <b>169</b> | <b>140</b>         | <b>140</b> | <b>58</b> | <b>58</b> | <b>61</b>                  | <b>187</b> |

**Table 4.** Summary of measured apparent optical properties per water system.

| <b>Type</b>  | <b>System</b> | $\rho_{wt}^L$<br>(1) | $\rho_s^L$<br>(1) | $d_{Secchi}$<br>(m) |
|--------------|---------------|----------------------|-------------------|---------------------|
|              | Bocht         | 7                    | -                 | 8                   |
|              | Donkmeer      | 34                   | -                 | 37                  |
|              | DelMare       | -                    | -                 | 4                   |
|              | Dikkebus      | 7                    | -                 | 7                   |
| Fresh        | Hazewinkel    | 20                   | -                 | 21                  |
|              | Leuven-Dijle  | -                    | -                 | -                   |
|              | Molsbroek     | -                    | -                 | 1                   |
|              | Nieuwdonk     | -                    | -                 | 2                   |
|              | Zillebeke     | 7                    | -                 | 7                   |
| Brackish     | Scheldt       | 8                    | -                 | -                   |
|              | Spuikom       | 62                   | 14                | 68                  |
| Marine       | BCZ           | 19                   | -                 | 18                  |
| <b>Total</b> |               | <b>164</b>           | <b>14</b>         | <b>173</b>          |

**Table 5.** Summary of measured and derived biogeochemical parameters per water system.

| <b>Type</b>  | <b>System</b> | <b>SPM</b><br>(g m <sup>-3</sup> ) | <b>f<sub>m</sub></b><br>(1) | <b>PSD</b><br>(cm <sup>3</sup> m <sup>-3</sup> ) | <b>Pigments</b><br>(mg m <sup>-3</sup> ) | <b>Amplicon</b><br>(counts) | <b>FlowCam</b><br>(counts) | <b>Bottom</b><br>(type) |
|--------------|---------------|------------------------------------|-----------------------------|--|--|-----------------------------|----------------------------|-------------------------|
| Fresh        | Bocht         | 8                                  | 8                           | 2  | 8  | -                           | -                          | -                       |
|              | Donkmeer      | 34                                 | 34                          | 10   | 36                                       | -                           | -                          | -                       |
|              | DelMare       | 2                                  | 2                           | 1  | 3  | -                           | -                          | -                       |
|              | Dikkebus      | 7                                  | 7                           | 5  | 7  | -                           | -                          | -                       |
|              | Hazewinkel    | 18                                 | 18                          | 1  | 18                                       | -                           | -                          | -                       |
|              | Leuven-Dijle  | 2                                  | 2                           | -  | -  | -                           | -                          | -                       |
|              | Molsbroek     | 1                                  | 1                           | -  | 1  | -                           | -                          | -                       |
|              | Nieuwdonk     | 2                                  | 2                           | -  | 2  | -                           | -                          | 1                       |
|              | Zillebeke     | 8                                  | 8                           | 6  | 8  | -                           | -                          | -                       |
| Brackish     | Scheldt       | 21                                 | 21                          | 2  | 12                                       | -                           | -                          | -                       |
|              | Spuikom       | 56                                 | 48                          | 15   | 53                                       | 48                          | -                          | 26                      |
| Marine       | BCZ           | 19                                 | 19                          | 19   | 19                                       | 15                          | 13                         | -                       |
| <b>Total</b> |               | <b>178</b>                         | <b>170</b>                  | <b>61</b>  | <b>167</b>                               | <b>63</b>                   | <b>13</b>                  | <b>27</b>               |

**Table 6.** List of quantified pigments. Names and abbreviations follow Roy et al. (2011). Known co-elutions according to Van Heukelem and Thomas (2001) are described in the table footnotes.

| <b>Name</b>  | <b>Abbreviation</b>  | <b>Name</b>                              | <b>Abbreviation</b>         |
|--|--|--|-----------------------------|
| (monovinyl) Chlorophyll <i>a</i>                                     | Chl <i>a</i>   | Diadinoxanthin                           | Diadino                     |
| (monovinyl) Chlorophyll <i>b</i>                                     | Chl <i>b</i>   | Diatoxanthin                             | Diato                       |
| Chlorophyll <i>c</i> <sub>1</sub> <i>c</i> <sub>2</sub> <sup>†</sup> | Chl <i>c</i> <sub>1</sub> <i>c</i> <sub>2</sub> <sup>†</sup> | Echinenone                               | Echin                       |
| (divinyl) Chlorophyll <i>c</i> <sub>3</sub>                          | Chl <i>c</i> <sub>3</sub>                                    | Fucoxanthin                              | Fuco                        |
| Chlorophyllide <i>a</i>  | Chlide <i>a</i>  | 19'-Butanoyloxyfucoxanthin               | But-fuco                    |
| Pheophorbide <i>a</i>  | Pheide <i>a</i>  | 19'-Hexanoyloxyfucoxanthin               | Hex-fuco                    |
| Pheophytin <i>a</i>  | Phe <i>a</i>   | Lutein                                   | Lut                         |
| α-Carotene   | βε-Car   | Mutatoxanthin                            | Mutato                      |
| β-Carotene   | ββ-Car   | Myxoxanthophyll                          | Myxo                        |
| Lycopene   | Lyc  | 9'- <i>cis</i> -Neoxanthin <sup>††</sup> | <i>c</i> -Neo <sup>††</sup> |
| Alloxanthin  | Allo   | Peridinin                                | Peri                        |
| Antheraxanthin   | Anth   | Prasinoxanthin <sup>†††</sup>            | Pras <sup>†††</sup>         |
| Astaxanthin  | Asta   | Violaxanthin                             | Viola                       |
| Canthaxanthin  | Cantha   | Zeaxanthin                               | Zea                         |
| Crocoxanthin   | Croco  |  |                             |
| α-Cryptoxanthin  | α-Cypt   |  |                             |
| β-Cryptoxanthin  | β-Cypt   |  |                             |

<sup>†</sup> Chl *c*<sub>1</sub> *c*<sub>2</sub> = Chl *c*<sub>1</sub> + Chl *c*<sub>2</sub> + magnesium 2,4-divinylpheoporphyryin *a*<sub>5</sub> monomethyl ester (MgDVP). MgDVP is present in trace concentrations in most algae and cyanobacteria, but is a major component of PRASINO 3A and 3B pigmentation groups (*sensu* Jeffrey et al., 2011);

<sup>††</sup> *c*-Neo and 19'-Hexanoyloxy-4-ketofucoxanthin (Hex-kfuco) co-elute and are not known to co-occur in the same organism or pigmentation group (*sensu* Jeffrey et al., 2011);

<sup>†††</sup> Pras and micromonol (Microl) co-elute and co-occur in the PRASINO 3B pigmentation group (*sensu* Jeffrey et al., 2011).