

# Global Covenant of Mayors, a dataset of GHG emissions for 6,200 cities in Europe and the Southern Mediterranean Countries

Albana Kona<sup>1</sup>, ~~Paolo Bertoldi<sup>1</sup>~~, Fabio ~~Ferrario~~ Monforti-Ferrario<sup>1</sup>, ~~Paolo Bertoldi<sup>1</sup>~~,  
5 Marta Giulia Baldi<sup>1</sup>, ~~Georgia Kakoulaki<sup>1</sup>~~, ~~Eleonora Lo Vullo<sup>1</sup>~~, ~~Georgia Kakoulaki<sup>1</sup>~~,  
Nadja Vetter<sup>1</sup>, Christian Thiel<sup>1</sup>, Giulia Melica<sup>1</sup>, ~~Eleonora Lo Vullo<sup>1</sup>~~, Alessandra  
Sgobbi<sup>2</sup>, Christofer Ahlgren<sup>2</sup>, Brieuc Posnic<sup>3</sup>

<sup>1</sup>European Commission – Joint Research Centre - Ispra

<sup>2</sup>European Commission – Directorate General for Climate Action

<sup>3</sup>European Commission – Directorate General for Energy

Correspondence to: [Fabio.MONFORTI-FERRARIO@ec.europa.eu](mailto:Fabio.MONFORTI-FERRARIO@ec.europa.eu)

**Abstract.** The Paris Agreement has underlined the role of cities in combating climate change. The Global  
Covenant of Mayors for Climate and Energy (GCoM) is the largest ~~dedicated~~ international initiative  
15 ~~dedicated~~ to promote climate action at city level, covering globally over 10,000 cities and almost half the  
population of the European Union (EU) by end of March 2020. The ~~latest-fifth~~ Intergovernmental Panel  
on Climate Change (IPCC) report ~~denotes~~ that there is a lack of comprehensive, consistent datasets of  
cities' Greenhouse Gas (GHG) emissions inventories. ~~In order to partly address part of this gap,~~  
we present ~~here~~ a harmonised, complete and verified dataset of GHG inventories for 6,200 cities in  
20 European and Southern Mediterranean countries, signatories of the GCoM initiative. To complement the  
~~reported~~ emission data ~~reported~~, a set of ancillary data that have a direct or indirect potential impact on  
cities' ~~climate-action~~ Climate Action plans were collected from other datasets, supporting further  
research on local ~~climate-action~~ Climate Action and monitoring the EU-27's progress on Sustainable  
Development Goal (SDG) 13 on ~~Climate Action~~ Climate Action. The dataset is archived and publicly  
25 ~~available with the DOI number~~ <https://doi.org/10.2905/57A615EB-CFBC-435A-A8C5-553BD40F76C9>.

## 1. Background & Summary

Cities consume over two-thirds of the world's energy and generate about 70% of global GHG emissions  
(IPCC, 2014), ~~and are at while being at~~ the same time particularly vulnerable to the impacts of climate  
30 change (Reckien et al., 2018). An increasing number of cities have voluntarily adhered to transnational  
networks active in ~~climate-action~~ Climate Action (Busch et al., 2018; Heidrich et al., 2016; van der Ven  
et al., 2017). As these networks and initiatives have evolved, cities' ambition and climate targets have  
increased to match or even go beyond the ambition of countries (Bertoldi et al., 2018d).

However, the scientific community ~~denotes~~ the current lack of systemic knowledge of cities' quantified  
35 contribution to combating climate change (Acuto et al., 2018; IPCC, 2015). This knowledge gap  
originates from many issues, including dissimilarities in the methodologies used for developing local  
emission accountings and reference scenarios and for setting ambition targets, as well as the absence of

a global, open and harmonised dataset of cities' emissions inventories (Kona et al., 2018). Only as recently as in 2019, the first datasets were published in academic literature, aiming to fill regional gaps (Adami et al., 2020; Kilkis, 2019; Palermo et al., 2020b).

The dataset presented in this paper aims to fill these gaps in Europe and Southern Mediterranean countries (Table 1). It consists of a harmonised, comprehensive and verified dataset of GHG emissions based on data produced by 6,200 cities in the EU-27, European Free Trade Association (EFTA) countries and UK, Western Balkans, Eastern Europe and Southern EU neighbourhoods (Figure 1).

The dataset is extremely valuable for communities engaged in climate action, including policy makers at all levels pursuing informed decisions, as demonstrated by the several and diverse studies based on GCoM data that have appeared in literature in latest years. To illustrate, for example: (Palermo et al., 2020a) have discussed the mitigation policies at the local level. Other authors have extracted and analysed specific subsets of CoM data: Pablo-Romero et al. (Pablo-Romero et al., 2018) (2017) have analysed the so-called Benchmarks of Excellence actions, while (Crocì et al., 2016) Crocì et al. (2017) has analysed the major cities present at that time in the database and (Famoso et al., 2015) Famoso et al. (2015) has focused the study on the participation of signatories from Sicily.

Thanks to the improved accessibility of the dataset and the addition of a number of useful ancillary variables, we expect the number of studies to grow in the near future. In order to further support further the scientific community, Along with the dataset, we provide the method of producing the data, their corresponding metadata, as well as the technical validation performed.

The European Commission launched the Covenant of Mayors (CoM) in 2008 to endorse and support the effort of EU local authorities in mitigating climate change. In 2015, the Covenant expanded to also include climate adaptation. In 2011, the initiative was launched in the EU's Eastern and Southern neighbourhoods, and in 2016 the initiative became global, through the launch of the Global Covenant of Mayors for Climate and Energy (GCoM). The initiative registered a very rapid growth from 241 signatories in 2008 in the EU-27 to more than 10,000 signatories covering more than 869 million inhabitants worldwide as of March 2020.

The Joint Research Centre (JRC), the science and knowledge service of the European Commission, provides scientific and technical support to GCoM cities in the development and implementation of their climate action plans. The scientific support is given through guidance on methodologies for emission accounting and climate adaptation, as well as through the development of urban policy tools for climate action (Bertoldi et al., 2018a, 2018b; Kovac et al., 2020; Monforti-Ferrario et al., 2018; Peduzzi et al., 2020). While the technical support consists of checking and validating the data reported by cities in the MyCovenant platform (www.covenantofmayors.eu).

The published dataset (Kona et al., 2020) contains verified reported GHG emissions for 6,200 European and South Mediterranean cities for a set of reference years. Given the voluntary nature of the GCoM and the difficulty of local authorities to report using a harmonised framework, a statistical method for checking the reliability, cleaning and validating the reported data was developed and applied. The method allows building a coherent dataset and consists of four steps:

Field Code Changed

Formatted: Font: 10 pt, Not Bold

Formatted: Font: Calibri, 11 pt

Formatted: Font: Times New Roman, 10 pt

- Data+Reporting principles, data extraction and clustering of GHG reporting framework, data extraction and clustering of signatories into two groups (large /small areas) based on population size and degree of urbanization (threshold 50,000 inhabitants) (see section 2.1);
- Outlier identification and treatment in large urban areas: Data coherence and completeness checks on the: data reported in the platform are checked in terms of coherence and completeness with the official Sustainable Energy and Climate Action Climate Action Plan (SECAP) document in large urban areas(see section 2.2);(2.2);(2.2);(2.2);(2.2);(2.2);(2.2);(2.2);
- Outlier identification and treatment in small and medium towns: Detection of outliers: statistical method applied for the identification and correction of outliers in small urban areas in small medium towns (see section 2.3);(2.3);(2.3);(2.3);(2.3);(2.3);(2.3);(2.3);
- Matching emission data with ancillary data: signatories from the EU are matched with their respective administrative units in the EU official statistics for cities. Harmonised statistical information on signatories allows building a referenced structure for collecting, processing, storing, analysing and aggregating data to support the monitoring of the EU progress on the Sustainable Development Goal (SDG) 13 on Climate Action signatories from the EU are matched with their respective administrative units in the EU official statistics for cities. Harmonized statistical information on signatories allows building a referenced structure for collecting, processing, storing, analysing and aggregating data to support the monitoring of the EU progresses on the Sustainable Development Goal 13 on Climate Action(Eurostat, 2020). (Eurostat, 2020).
- (see section 2.4).

The dataset ([Figure 1 Table 1 online dataset](#)) thus contains adjusted self-reported data from cities (i.e. [GCoM dataset 2019: Emission Inventories](#)) coupled with ancillary data ([GCoM dataset 2019: Ancillary data](#)) related to geographic attributes (area, latitude and longitude, local administrative codes, heating degree-days), socio-economic aspects (GDP per capita) and demographic characteristics at city level (degree of urbanisation, population time series). A detailed technical evaluation at the city level was also performed against the independent estimates provided by the Emissions Database for Global Atmospheric Research [EDGAR v5.0](#) ([EDGAR-v5](#) provides time series of global anthropogenic emissions of greenhouse gases and air pollutants by country on a spatial grid (Crippa et al., 2020)).

In compliance with the EU data policy, we are now in a position to share with the community a ten years' dataset validated and harmonised with the EU statistical information system of local authorities. The validation process assesses the completeness of the data (i.e. to minimum reporting requirement), the coherence of the data (i.e. data reported in the platform are coherent with the Climate Action plan document) and includes a data cleaning step (i.e. detection of outliers and their treatment). In compliance with the EU data policy, we are now in a position to share with the community a ten years' dataset complete, cleaned, validated and harmonised with the EU statistical database on local authorities. In spite of Given the overall good quality of the dataset, some limitations and uncertainties remain and are described in under the “Limitation Conclusion, 4 limitations and future work” section.

The resulting dataset is of great value and interest and targets the needs expressed clearly by the scientific and academic community and governmental institutions. This is demonstrated through several data release requests that have been received from different groups. These include the IPCC working group on Climate mitigation chapter, the United Nations Framework Convention on Climate Change non-state-actors zone platform for Climate Action (UNFCCC-NAZCA), governmental and research institutions of EU-27 Member States interested in the local contribution to the national reduction targets, and other sub national levels interested in understanding the active territorial participation to Climate Action movement.

~~The resulting dataset is of great value and interest and responds to the clear needs expressed by the scientific and academic community and governmental institutions. Last~~ A distinctive characteristic ~~but not least, the peculiarity of the European GCoM initiative is that it includes in its members small towns interested and engaged in Climate Action, often absent from other initiatives~~ is the participation of small towns interested and engaged in climate action, often absent from other initiatives. Therefore, this dataset offers cities of all sizes the means to formulate a comparative analysis of the magnitude, efficiency and intensity of energy use and GHG emissions. Users are warned to read in detail the description of the dataset provided in this paper and to be aware of the overall scope of the GCoM initiative in designing their investigations. In particular, they should be aware that the initiative has never meant to be a method to create exhaustive inventories of all emission sources in the territory or to deal with emissions already included in national-scale control initiatives, such as the EU Emission Trading System (ETS) mechanisms. GHG emissions mainly reported are CO<sub>2</sub> emissions, and in rare cases CH<sub>4</sub> or N<sub>2</sub>O. Further to the information presented in this paper, a more comprehensive amount of information and the corresponding guidelines are available through the website. ~~Besides the forcedly limited amount of information presented in this paper, a relevant amount of information and guidelines is available throughout the CoM website.~~

## 2. Methods

Hereafter we describe the methods used to produce and consolidate the final dataset. Due to local authorities' difficulties in harnessing and reporting data within a harmonised framework, which may differ from the national emission reporting, not all the self-reported data could be considered reliable. Therefore, a method was developed to construct a robust dataset of emission inventories, organised into four steps:

- Step 1: Data reporting principles, extraction and clustering: accounting principles of GHG reporting framework, data extraction and clustering of signatories into two groups (large/small areas) based on degree of urbanization and/or population size (threshold 50,000 inhabitants);
- Step 2: ~~Detection of outliers from large urban areas~~ ~~Data coherence and completeness~~: Digital curation of data reported in the platform were performed in terms of completeness and coherence with the official ~~climate action~~ Climate Action plan document, the so-called SECAP in large urban areas;
- Step 3: Detection of outliers from small medium towns: statistical method for the identification and detection of outliers in the GHG emission dataset in small medium towns;

- Step 4: Matching emission data with ancillary data: signatories from the EU are matched with their respective local administrative units of the Geographic Information System of the European Commission.

## 2.1 Data principles, extraction and clustering

To streamline measurement and reporting procedures under the GCoM, a Common Reporting Framework (GCoM CRF) was developed during 2018 in consultation with partners and signatories: The GCoM has two officially recognised platforms for data reporting, the CDP ICLEI Unified Reporting System and the My Covenant platform. To streamline measurement and reporting procedures, a Common Reporting Framework (CRF) was developed during 2018 in consultation with partners and signatories of the GCoM. While the platforms differ in terms of the data collection approach, they are aligned with the GCoM CRF. The dataset provided in the current study is based on the information reported by signatories through the MyCovenant platform, following the JRC methodology (Bertoldi et al., 2018a), one of the officially recognized reporting platforms of the initiative, and the one used by majority of the signatories. Hereafter we report a brief description of the data collected on MyCovenant in alignment with the GCoM CRF.

The reporting framework is built upon the Emission Inventory Guidance, used by the European Covenant of Mayors and the Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC), used by the Compact of Mayors. Both refer to the 2006 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories.

The protocols for accounting the cities' emissions differ mainly in the principles and minimum reporting requirements on sources, the type of gases and the boundary of the inventory to be reported.

The protocol for accounting the emissions used by Covenant signatories is closely aligned with the IPCC 2006 guidelines regarding the source category of the in-boundary emissions (i.e. the administrative boundaries). It includes "sources" and "activities" rather than the scope framework used in other city protocols (for example the GHG protocol of WRI). Nevertheless, the emission inventory is not meant to be an exhaustive inventory of all emission sources in the territory. It focuses mainly on GHG emissions related to sectors (stationary energy, transport and waste/wastewater) upon which the local authority could intervene through sectoral measures and urban policies. They Signatories also can report as well GHG emissions from Industrial Processes and Product Use (IPPU) and Agriculture, Forestry and Other Land Use (AFOLU) sectors where these are significant (Table 2Table 2Table 4).

Moreover, one of the main differences on GHG emission reporting between non-state (e.g. cities) and state actors is the level of flexibility in choosing the inventory year with the most reliable data. The recommended baseline year for reporting is 1990, or the closest subsequent year for which the most comprehensive and reliable data can be provided (for example 2005).

The geographical boundaries of the "local territory" are the administrative boundaries of the entity (municipality, region) governed by the local authority which is a signatory to the GCoM. Regarding the type of gases, GCoM signatories shall report emissions of carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) converted into CO<sub>2</sub>-equivalents (CO<sub>2</sub>-eq.), according to their global warming potential

Formatted: Font: (Default) Times New Roman, 10 pt, Not Italic

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font color: Auto

([Local governments should disclose also which GWP factors they are using](#)). The three main GHG emission categories included in the inventories are:

- Direct emissions due to final energy consumption, excluding those from industrial plants involved in the ~~Emissions Trading Systems~~ [ETS](#).
- Indirect emissions related to grid supplied energy (electricity, heat, or cold) consumed in the local territory (Kona et al., 2019).
- Non-energy related direct emissions (such as from waste, wastewater) that occur in the local territory, if the ~~climate action~~ [Climate Action](#) plan contains measures to reduce such GHG emissions.

The GHG emissions are automatically derived in the platform as the product of activity data (detailing the energy consumption/waste per carrier/type) and emission factors, as reported by the signatories ([Table 2](#) ~~Table 1~~). The emission factors are coefficients, which quantify the emissions per unit of activity, and one out of three approaches can be used:

- IPCC ([2006](#)) – emission factors for fuel combustion – default values mostly based on the carbon content of each fuel;
- National or subnational emission factors for fuel combustion when these are different from the IPCC's.
- Life Cycle Assessment (LCA) – emission factors for the overall life cycle of each energy carrier, i.e. including not only the GHG emissions due to fuel combustion but also emissions of the entire energy supply chain – exploitation, transport and processing.

The procedure to verify and improve the coherence of the dataset starts with the extraction of complete emission inventories stored in a ~~relational SQL PostgreSQL~~ database. At the closing date of this study, (September 2019) 6,239 ~~climate-Climate Aca~~ [Climate Action](#) plans with complete inventories have been submitted by cities in the EU-27, EFTA countries and UK, Western Balkans, Eastern and Southern EU neighbourhoods.

Inventories and other data are self-reported to the online platform and must accurately reflect the content of the official ~~climate action~~ [Climate Action](#) plan (called ~~Sustainable Energy and E~~ [Climate Action Plan \(SECAP\)](#)) document. The SECAP document is a separate file, usually in PDF format and publicly available, that represents the official action plan endorsed and signed by the local council. ~~As a first step to address the quality of the data reported, yearly GHG emission per capita are plotted for each signatory.~~ The first step to understand the degree to which this is true and the quality of the ~~data-reported~~ [data](#), yearly GHG emission per capita are plotted for each signatory. ([Figure 2](#) ~~Figure 2~~). ~~shows the frequency distribution of GHG emissions per capita in emission inventories dataset in 2019, with observations that range from 0 to 80 tCO<sub>2</sub>-eq/cap, with a mean of 5.18 tCO<sub>2</sub>-eq/cap. In the vertical axis the density values are reported, i.e. the share of signatories with the same range of GHG emission per capita (i.e. emission from all the CoM sectors, excluding manufacturing and construction industries), by the width of the class (0.01 in this case).~~

The occurrence of outliers ([eg. large amounts of per capita values](#)) is a clear indication of errors in the data, ~~likely due to inputting error~~, therefore not all the data collected in the platform are consistent with the SECAP document. As the calculations of performance indicators for the dataset, such as the mean

Field Code Changed

Formatted: Font: Not Bold, English (United States)

Field Code Changed

and standard deviation, can be distorted by a single grossly inaccurate data point, checking and treating outliers is a routine part of data analysis.

Due to the high volume of information, it is not feasible to check individually the consistency of all the data objects with the SECAP document. The collection of the attributes (i.e. the variables: 15 energy carriers and 16 subsectors) describes the data objects (it is also known as record, point, case, sample, entity, or instance), which visually corresponds to the rows in the Excel files.

The original dataset comprises 6,239 signatories with a baseline inventory, out of which 1,845 with an additional monitoring inventory. In each inventory the cities report at maximum data for 15 energy carriers grouped into 16 subsectors, resulting therefore into 1.94 million data objects. The 16 subsectors have been grouped into 6 sectors (i.e. municipal, residential, tertiary, manufacturing and construction industries, transportation and waste sector, see Table 2Table 2Table) and null objects were deleted, leading in total to 61,207 data objects.

We therefore adopted a rule to treat the outliers, based on the benefits expected when scrutinizing the dataset for the overall assessment of the initiative. Data users willing of The overall assessment consists of producing performance indicators on the impact and the contribution of climate actionClimate Actions planned and implemented by CoM signatories must benefit of a robust dataset in order to avoid artefacts and unreliable results. In this context, it is evident that, the bigger a city is, the more impact any errors will have on the overall dataset. In order to have an accurate representation, it is then of utmost importance that large cities have highly accurate data.

Hence, we decided to adopt a customized method to treat outliers based on the signatories' degree of urbanization and population size (source <https://ec.europa.eu/eurostat/web/nuts/local-administrative-units>). The 6,239 signatories and their data were clustered into two groups:

- Large urban areas (densely populated area with a population density of at least 1,500 inhabitants per km<sup>2</sup> and a minimum population of 50,000): for this group manual curation of imputed errors in inventories was implemented, which significantly increased the performance indicators of the database by increasing their robustness (described in step 2).
- Small towns and rural areas (intermediate and thinly populated areas): for this group an automatic routine to identify and remove the outliers is applied. The rules governing the automatic detection and treatments of the outliers are detailed in Step 3.

## 2.2 Data coherence and completenessData cleaning – large urban areas

In this section, we describe the steps followed to detect and treat the outliers in inventories from large urban areas (i.e. cities and greater cities, with a population density of at least 1,500 inhabitants per km<sup>2</sup> and a minimum population of 50,000) along with correctness and completeness checks in the overall dataset. The identification and treatment of outliers in this group of cities has been performed qualitatively.

Because of the harmonization process of GCoM administrative data and local administrative units Eurostat database 2018 (Eurostat, 2018), 430 signatories covering 116.2 million inhabitants, are classified as cities and greater cities. In addition, in the other regions out of the EU-27 and EFTA regions (i.e. Eastern Europe; Western Balkans and Southern Mediterranean) where the classification was not available, we adopted as criteria only the population size as the threshold (i.e. a minimum population of

**Formatted:** Font: (Default) Times New Roman, 10 pt, Not Italic

50,000). Hence, within the GCoM 2019 dataset there are 701 baseline inventories presented by large urban areas, covering a total population of 165.26 million inhabitants.

As part of the evaluation process carried out by JRC on individual SECAPs, activity data were compared against the national/ EU averages (available at national/EU statistical systems [such as Eurostat, European Environment Agency](#)). In case of reported data that ranged out of one or more units higher than the average of the sectors national average, we double-checked the accuracy of ~~the platform's data~~ reported ~~data on the platform~~ with the SECAP document. As a result of the digital curation of outliers, identified through the comparison of self – reported data in the MyCovenant platform against the same data declared in the SECAP, twenty inventories (i.e., about 3 %) have been ~~manually~~ manually corrected. The SECAP document represents the official action plan endorsed and signed by the local council; therefore, ~~we assume as valid the data reported here the assumption is that the data reported within their the SECAP are the valid ones~~. The errors were often due to the misinterpretation of the unit measure to be reported in the online template (e.g. kWh/year instead of MWh/year, etc.).

At this point in the procedure, with the help of the statistical routine and the digital curation, we have consolidated the dataset related to activity data. The next step consists of comparing the emission factors used in GCoM inventories against the reference values from IPCC [AR4 \(4<sup>th</sup> Assessment Report](#) (IPCC, 2007) and the JRC databases (Lo Vullo et al., 2020) and their completeness (i.e. missing data on emissions were derived from reported activity data and vice versa). In case of reported emission factors that ranged out of  $\pm 50$  % of the reference value, we corrected them with the corresponding reference value. As a result of this procedure, there were 153 inventories from large urban areas, where 9.7 % (i.e. 526 out of 5,433 objects) of the data objects were corrected.

### 2.3 ~~Detection of outliers~~ Data cleaning – small and medium towns

In this section we describe the automatic routine implemented to detect and treat the outliers in inventories from small medium towns (number of inventories = 5,538 covering a total population of 46.78 million inhabitants).

Urban GHG emissions per capita may deviate significantly from national averages, due to the tendency of emissions to concentrate around human activities. Therefore, setting exclusion ranges of outliers in the per capita GHG emissions based on the national averages may lead to the exclusion of a high number of valid emission inventories from the GCoM dataset. To avoid this bias, we apply a statistical method based on intrinsic properties of the distribution of the emissions in the GCoM database. This allows identifying more accurately potentially unreliable emission inventories and the outliers likely to be the results of incorrect data entry.

The procedure starts with dividing the data into two groups based on the normalization process: the activity data in the residential/municipal/institutional/tertiary buildings and transport sector were normalised with the population size, whereas the activity data in manufacturing and construction industries were normalised with the GDP values. The majority of these industries are already governed by the cap and trade system ([EU Emission Trading scheme ETS](#)), therefore they are not recommended to be reported in the GCoM platform, although exceptions exist. In addition, signatories that report manufacturing emissions are generally large urban areas (80 % of the activity data within this sector is

**Formatted:** Font: (Default) Times New Roman, 10 pt, Not Italic



reported by cities and greater cities), which we have been already examined individually to check for outlying data.

The outliers identification method is based on a generalised ~~ESD~~ (extreme studentized deviate (ESD)) procedure for the detection of abnormal energy consumptions. The ESD is commonly used in literature (Cerquitelli et al., 2019; Gant., 2013; Rosner, 1983; Seem., 2007), because of its excellent performance under a variety of conditions to detect one or more outliers in a dataset that follows an approximately normal distribution. The per capita activity data in the residential/municipal/institutional/tertiary buildings and transport sector follows approximately a normal distribution.

The procedure iteratively identifies the extreme values in the dataset and then selects to remove those observations which are higher than the extreme values with a confidence level of 95 %. A detailed description of the routine is available at [Supplementary File 1](#) and [Supplementary File 2](#).

Applying this approach, 39 inventories were removed from the initial dataset (i.e. from initial 5,538 inventories). These signatories [received a further feedback in addition to the routinely checks already performed at the time of data submission](#) and have been approached ~~through the feedback report~~ to check and correct the data in the online platform. The clean and robust dataset thus contains 5,499 inventories. As a result, the original inventory containing 6,239 entries was reduced to a clean dataset of 6,200 signatories (i.e. 99 % of the original data), referred to hereafter as the “GCoM dataset 2019: Emission Inventories”.

To conclude, also a non-parametric statistical procedure, i.e. the Median Absolute Deviation (MAD), has been applied to identify outliers in dataset that ~~do are not follow a n-normal l-distributioned~~. This method is more robust than the ESD, but less efficient, and its validity increases as data approach ~~a -normal~~ distribution. Similarly to the ESD, the choice of the critical value is motivated by the reasoning that if ~~the~~ observations other than outliers have an approximately normal distribution, it picks up as an outlier any observations more than about three standard deviations from the means. The results of the MAD procedure produce the same outliers as the ESD procedure; therefore, we argue that the assumption on the quasi normal distribution is correct.

The next step consists of verifying the emission factors used in the inventories, against the reference values from IPCC [2006](#) and the JRC databases (Lo Vullo et al., 2020), and their completeness (i.e. missing data on emission were derived from reported activity data and vice versa). In case of reported emission factors that ranged out of  $\pm 50\%$  of the reference value, we corrected them with the corresponding reference value. Because of this procedure, there were 3,019 inventories from small towns, where 15 % (i.e. 8,008/52,496 records) of the data records were corrected ([Table 3](#)). ~~Error! Not a valid bookmark self-reference.~~ [Table 2](#) compares the main descriptive parameters of the two datasets. The main difference can be noted in the skewness parameter. Both frequency distributions have a positive skewness, meaning that the right tail is longer and the mass of the distribution is concentrated on the left of the figure.

#### 2.4 Matching emission data with ancillary data

GCoM signatories, when submitting their data ~~in-to~~ the MyCovenant platform, report the local authority name, the country and their centroids' coordinates. Through these three attributes, we have been able to digitally match the signatories with their corresponding local administrative units in the Geographic

**Formatted:** Font: (Default) Times New Roman, 10 pt, Not Italic

Information System of the European Commission (GISCO) (Eurostat, 2018). Harmonized statistical information on signatories allows building a referenced structure for collecting, processing, storing, analyzing and aggregating data.

In this way, we can derive all ancillary data related to institutional, demographic and socio-economic dimensions:

- Institutional dimension: the GCoM signatories are associated with their correspondent NUTS codes; the Local administrative units' codes; their Functional urban area and cities codes; the geographical coordinates; the area and shape files of their local administrative units;
- Demographic dimension: the GCoM signatories are associated with the population data in 2018; the degree of urbanisation;
- Socio-economic and climate dimension: the GCoM signatories are associated with the GDP at NUTS 3 level; and heating degree-days at NUTS 3 level.

The aim of the ancillary data is also to support the monitoring of the SDG 13 on ~~climate action~~ Climate Action in an EU-27 context, which focuses on climate mitigation, climate impacts and on initiatives that provide support to ~~climate action~~ Climate Action, as the ~~Global Covenant of Mayors~~ GCoM. More broadly, the ancillary data could support further research on investigating drivers of ~~climate action~~ Climate Action at city level and the development of urban policy design. In addition, we extracted the national values of GHG emissions per capita from EDGAR v5.0 for the corresponding GCoM activity sectors (~~Table 4~~ ~~Table 3~~ ~~Error! Reference source not found.~~).

### 3. Data availability

The dataset is archived and publicly available with the DOI number <https://doi.org/10.2905/57A615EB-CFBC-435A-A8C5-553BD40F76C9> (Kona et al., 2020).

### 4. ~~Benchmarking Comparison with ancillary emission~~ ~~other emission inventories~~

~~Matching reported emissions with cities' ancillary data enables an additional robustness check of the accuracy of inventories. In the case of GCoM, the uncertainty of reported emissions is particularly difficult to estimate since non-formal uncertainty analysis is applied by cities on the activity data and the emission factors. Hence, given this limitation, we argue that the best practical way to assess the uncertainty of reported data is to perform a detailed benchmark of the overall dataset comparison against international emission dataset such as EDGAR v5.0. A similar approach has been applied to validate cities emission data in United States (Nangini et al., 2019).~~

~~Although such a procedure does not necessarily implies an absolute validation of our data, could clarify to what extent the dataset is comparable with an internationally reputed source as EDGAR v5.0 is. It is also worth noticing that the two benchmarked datasets are different by principle, as GCoM collects data from local authorities with a supposedly good knowledge of their territory, while EDGAR v5.0 provides past and present global anthropogenic emissions of greenhouse gases and air pollutants by country on a spatial grid (Crippa et al., 2020). The methods used in EDGAR v5.0 (Crippa et al., 2020) downscale the~~

Formatted: Font: Palatino Linotype, English (United States)

Field Code Changed

Formatted: Font: Palatino Linotype

Formatted: Font: Palatino Linotype

emissions from a national or subnational scale to finer scales using spatial proxies and present results in gridded maps. EDGAR v5.0 combines several proxies ranging from population density to specific point source location maps for estimating emissions of different economic sectors. Regardless the different approach, the potential use of EDGAR gridded data for the examination of emission in large sample of cities worldwide has been already noticed in literature (Marcotullio et al., 2014). EDGAR v5.0 provides past and present global anthropogenic emissions of greenhouse gases and air pollutants by country on a spatial grid (Crippa et al., 2020).

Using ArcGIS, we overlaid the signatories' LAU-urban spatial boundaries onto the EDGAR v5.0 emissions grids. We then used the built-in Spatial Zonal Statistics tool to estimate total emissions for each urban area for-and two source categories: energy in buildings (RCO) and road transportation. EDGAR includes v5.0 includes emissions from a variety of sources (Solazzo et al., 2021) at the aggregate level of at least 0.1° spatial resolution (representing about 10 x 10 km<sup>2</sup> at the equator). Here we use the EDGAR v5.0 global grids of estimated emissions in metric tons for the year 2005 for the most prevalent GHGs: carbon dioxide excluding short cycle organic carbon (i.e. CO<sub>2</sub>\_excl\_short-cycle\_org\_C). Emissions of CO<sub>2</sub>\_excl\_short-cycle\_org\_C include all fossil CO<sub>2</sub> (such as fossil fuel combustion) and exclude all sources and sinks from land-use, land-use change and forestry (LULUCF) (Crippa et al., 2020). Overall, we compared data from 1945 signatories from EU-27 + UK countries with EDGAR v5.0 corresponding data on direct emissions in eEnergy in Buildings-buildings sector and road transportation (Online-only Table 3).

Annex 1 provides a detailed discussion of the benchmark of the two datasets in both transport and residentialbuilding sectors.

The correlation coefficient (R<sup>2</sup>) between the GCoM direct emission in the residential/municipal/institutional energy in buildings sector and their corresponding data in EDGAR v5.0 was 0.92, whereas for road transportation data was 0.66. We also calculated the root mean square error (RSME) between the GCoM dataset and EDGAR v5.0 direct emissions per capita in the two sectors: in the residential/municipal/institutional energy in buildings sector RSME<sub>Buildings</sub> = equalled to 0.08 CO<sub>2</sub>-eq/cap; and in the road transport sector RSME<sub>Transp</sub> equalled to = 0.11 CO<sub>2</sub>-eq/cap.

Regarding the direct emissions in the “Eenergy in buildinbuildinge” sector, the low value of the RSME uncertainty (0.088%) the correlation coefficients shows a good agreement between the two datasets. This might be attributed to the fact that, in this sector EDGAR v5.0 assumes that each resident is responsible for the same amount of emissions. Whereas GCoM collects direct observations (data collected from utilities mainly), that are good proxies of national of energy usage pattern and fuels deployed.

About the direct emissions in the transport sector, the slightly higher level of uncertainty (11.%) might be attributed to the fact that emissions in the Covenant framework covers only the urban fraction of the sector (i.e. in the EU urban mobility accounts for 40% of all CO<sub>2</sub> emissions of road transport). Moreover, the CoM dataset reported here does not reflect only the on road fraction of the transport sector, but all the emission in this sector. This is due to the old version of the reporting platform that collected data without distinguishing the modal share. Further on limitation is discussed in the next section. G v5.0G Overall, considering the completely different origin of EDGAR v5.0 and GCoM primary data the

agreement has to be considered ~~fully~~-satisfactory, taking into consideration the ~~well known~~well-known difficulties in matching inventories based on top-down or bottom-up approaches and the uncertainties affecting inventories in general. (Ssee e.g. (Solazzo et al., 2021) Solazzo et al. (2021) for a deep analysis of EDGAR v5.0 uncertainties).

## 5. Main findings

5. Local authorities that adhere to transnational networks active on climate action, by making publicly available the plan, without any obligation to do so, render themselves accountable both globally and locally (Gordon, 2016). This paper presents a major attempt to provide the scientific community with a reliable, consistent and complete dataset, derived from the cities' plan submissions. The following provides an overview of the results extracted from the analysis of the dataset in terms of signatories' participation, the submission status of the Climate Action plans, as well as its implementation progress in terms of the emission trend.

6. Starting with adhesion, Table 3 in the dataset reports the full list of the 8136 signatories and associated ancillary data. The ancillary data comprises institutional (i.e. statistical administrative information), demographic (i.e. population, degree of urbanization) and socio-economic data (i.e. GDP, heating degree-days, national GHG emissions per capita). Harmonised statistical information on signatories (i.e. the ancillary data) allows building a referenced structure for collecting, processing, storing, analysing and aggregating data to support the monitoring of the EU-27 progress on the SDG 13 on Climate Action (Eurostat, 2020). (Eurostat, 2020). (Eurostat, 2020). (Eurostat, 2020).

7. Regarding action planning, three-quarters of these signatories (i.e. 6200 local authorities) submitted an action plan, comprising a baseline emission inventory and a set of actions to reach their climate mitigation goals.

8. About the trend, less than one third of these submissions (i.e. 1845 signatories), reported progress on the implementation of the action plan by presenting a second inventory, called monitoring report. Table 2 in the dataset reports the activity data and related emissions mapped in the baseline and monitoring inventories aggregated into Stationary energy, Transport and Waste subsectors.

9. Hence, the trend on emissions uses these data to assess the progress made by the signatories in the implementation of their climate action plans is assessed in terms of emissions based on these data. Since the inventories might have different reporting sectors, we analyzed the trend ~~only in~~ considering only those sectors reported reporting data within the sector in both inventories, the baseline and the monitoring one. Overall, the trend on emissions from signatories reporting progress shows We found that the absolute reductions achieved from baseline inventories to monitoring inventories correspond to 23 %. Assuming that if -cities had progressed linearly towards their target, these signatories would have achieved 17 % of emission reduction by the inventory years, which is lower than the 23 % observed reduction. Consequently, we can assume that monitoring signatories are on track to reach their commitment.

## 10.6. Conclusions Limitations and future work

Despite the data mining and verification process, a few limitations and uncertainties remain in relation to the data quality. To start with, it is important to highlight the fact that the overall quality of the data reported in the platform depends mostly on the city's capacity to gather and report into the harmonized framework of GCoM. The JRC does not correct or adjust the reported data itself, but it is the responsibility of the signatory, on receipt of the feedback analysis from experts, to check and ~~possible-possibly~~ revise its data according to the ~~climate action~~ Climate Action plan. Indeed, we have noted an increasing quality of data reported from cities since 2010, mainly thanks to this feedback-rechecking system.

Therefore, the aim of the approach adopted here is not to validate the data as such (they are collected and reported by the signatories), but to guarantee as far as possible the internal consistency and completeness of the data reported in the online platform with the ~~climate action~~ Climate Action plan documents (i.e. the SECAP).

Secondly, ~~there is~~ a limited knowledge on the methods used by cities in determining the emissions, especially within the transport sector. The aim of ~~this-the~~ technical validation reported in section 4 is to compare the GCoM dataset against international datasets such as EDGAR, being well aware of the fact that GCoM reports direct observations, whereas EDGAR calculates emissions following a consistent Tier 1 approach based on AD and EF country specific information.

~~represents modeled values.~~ In small-medium urban ~~areas~~ areas, we ~~suppose-assume~~ that local authorities use the territorial approach based on the collected activity data ~~collected~~. ~~Indeed~~ For these areas, there is a good match with EDGAR v5.0 data, whereas in large urban areas, we note a significant deviation from EDGAR proxies. ~~As already mentioned. We argue that most probably~~ this is probably due, among other factors, to the differences between locally collected data (e.g. on local fleet) fact that CoM reports real data supplied from the transportation department, which are not caught in the modelling exercise of EDGAR. More precisely, EDGAR uses the and average national information. fleet that could be quite different from a local one. Moreover, due to the uncertainty on the methodological differences for accounting the emissions, embedded in the nature of the sector, the emissions in this sector can differ widely between cities with similar patterns or sizes.

~~About~~ Regarding waste, the mapping of emissions in this sector has only been added in the last revision of the reporting framework, therefore we expect more data in this sector to become available as cities integrate it in their inventories. For this reason, data from this category were not included in the present study.

~~To conclude~~ Finally, it has to be reminded that a major source of uncertainty originates from the use of emission factors developed for the national or sometimes even international scale, especially for the electricity and waste sectors. According to our information, signatories often apply the default emission factors provided by JRC and based on very wide scale IPCC Tier 1 approach. Clearly, such a coarse granularity is not always able to catch local peculiarities and features.

On the contrary, deploying city level emission factors for instance for electricity supplied through the grid, taking also into account local renewable energy production, would greatly increase the accuracy of the data.

Future work envisage the possibility of undertaking a comparable analysis also of the data reported by GCoM signatories through the [CDP-ICLEI Unified Reporting System](#), and other recognized/affiliated national and regional reporting platforms –with a view to expanding the coverage of a harmonised, complete and verified dataset of GHG inventories at city level.

**Code Availability:** Most data handling in the methods and technical validation was done in MATLAB (available at [Supplementary File 1](#)) and Microsoft Excel ([Supplementary File 2](#)).

**Acknowledgements:** We are grateful to local authorities who make public their engagement in ~~climate action~~Climate Action planning, through their participation in the Global Covenant of Mayors initiative. The authors would like to thank European Commission Directorate General for Energy, the CoM Office and JRC colleagues working in the CoM initiative for their support in giving visibility and effectiveness to the effort of cities and local governments in the climate change action. A special acknowledgment to our colleagues Monica Lanzoni for her in-depth reviewing of the statistical analyses, and to Andreas Kontogeorgos for his contribution to English language review.

**Author contributions:** Kona A., ~~Bertoldi P. and Ferrario~~ Monforti-Ferrario F. and Bertoldi P., designed the research. Kona A. prepared the manuscript. Kona A., Baldi M.G., Kakoulaki G. and Ahlgren C. assembled and prepared the dataset. All the authors revised the methodological steps and the manuscript.

**Competing interests:** The authors declare no competing interests. The views expressed are purely those of the authors and may not in any circumstances be regarded as stating an official position of the European Commission.

## References

Acuto, M., Parnell, S. and Seto, K. C.: Building a global urban science, Nat. Sustain., 1(January), 4–6, doi:10.1038/s41893-017-0013-9, 2018.

Adami, L., Tubino, M., Ragazzi, M., Conti, F. and Rada, E. C.: Local actions for reducing global greenhouse gas footprint: 10 years of covenant of mayors initiative, Int. J. Sustain. Dev. Plan., 15(2), 247–252, doi:10.18280/ijstdp.150216, 2020.

Bertoldi, P., Iancu, A., Kona, A., Suvi, M., Muntean, M., Lah, O. and Rivas, S.: Guidebook “How to develop a Sustainable Energy and Climate Action Plan (SECAP)”: PART 2 – Baseline Emission Inventory (BEI) and Risk and Vulnerability Assessment (RVA), EUR 29412., Publications Office of the European Union, Luxembourg. [online] Available from: <https://doi.org/10.2760/118857>, 2018a.

Bertoldi, P., Kona, A., Palermo, V., Zangheri, P., Serrenho, T., Rivas, S., Labanca, N., Kilkis, S., Lah, O., Glancy, R., Follador, M., Barbosa, P. and Andreanidou, K.: Guidebook “How to develop a Sustainable Energy and Climate Action Plan (SECAP)” PART 3-Policies, key actions, good practices for mitigation and adaptation to climate change and Financing SECAP(s), Publication Office of the European Union, Luxembourg. [online] Available from: <https://doi.org/10.2760/58898>, 2018b.

Formatted: Italian (Italy)

Formatted: Italian (Italy)

Bertoldi, P., Kona, A., Rivas, S. and Dallemand, J. F.: Towards a global comprehensive and transparent framework for cities and local governments enabling an effective contribution to the Paris climate agreement, *Curr. Opin. Environ. Sustain.*, 30, 67–74, doi:<https://doi.org/10.1016/j.cosust.2018.03.009>, 2018c.

545 Bertoldi, P., Kona, A., Rivas, S. and Dallemand, J. F.: Towards a Global Comprehensive and Transparent Framework for Cities and Local Governments enabling an Effective Contribution to the Paris Climate Agreement, *Curr. Opin. Environ. Sustain.*, 2018d.

Busch, H., Bendlin, L. and Fenton, P.: Urban Climate Shaping local response – The influence of transnational municipal climate networks on urban climate governance, *Urban Clim.*, 24(December 550 2017), 221–230, doi:[10.1016/j.uclim.2018.03.004](https://doi.org/10.1016/j.uclim.2018.03.004), 2018.

Cerquitelli, T., Corso, E. Di, Proto, S., Capozzoli, A., Bellotti, F., Cassese, M. G., Baralis, E., Mellia, M., Casagrande, S. and Tamburini, M.: Exploring energy performance certificates through visualization, in *Proceedings of the Workshops of the EDBT/ICDT 2019 Joint Conference*, EDBT/ICDT 2019, Lisbon, Portugal, March 26, 2019., edited by CEUR-WS.org, p. 7. [online] Available from: [http://ceur-  
ws.org/Vol-2322/BigVis\\_2.pdf](http://ceur-555 ws.org/Vol-2322/BigVis_2.pdf), 2019.

Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R. and Janssens-maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, *Sci. Data*, 1–17 [online] Available from: <http://dx.doi.org/10.1038/s41597-020-0462-2>, 2020.

560 Croci, E., Lucchitta, B., Janssens-Maenhout, G., Martelli, S. and Molteni, T.: Urban CO<sub>2</sub> mitigation strategies under the Covenant of Mayors: An assessment of 124 European cities, *J. Clean. Prod.*, 169, 161–177, doi:[10.1016/j.jclepro.2017.05.165](https://doi.org/10.1016/j.jclepro.2017.05.165), 2016.

Eurostat: GISCO, [online] Available from: [https://ec.europa.eu/eurostat/web/gisco/geodata/reference-  
data/administrative-units-statistical-units/lau#lau19](https://ec.europa.eu/eurostat/web/gisco/geodata/reference-data/administrative-units-statistical-units/lau#lau19), 2018.

565 Eurostat: SDG 13 - Climate action, [online] Available from: [https://ec.europa.eu/eurostat/statistics-  
explained/index.php?title=SDG\\_13\\_-\\_Climate\\_action#Climate\\_mitigation](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=SDG_13_-_Climate_action#Climate_mitigation) (Accessed 24 May 2021), 2020.

Famoso, F., Lanzafame, R., Monforte, P. and Scandura, P. F.: Analysis of the covenant of mayors initiative in sicily, *Energy Procedia*, 81, 482–492, doi:[10.1016/j.egypro.2015.12.122](https://doi.org/10.1016/j.egypro.2015.12.122), 2015.

570 Gant, F.: *Understanding Statistics: Basic Theory and Practice*. Basic Theory and Practise., 2013.

Gordon, J. D.: The Politics of Accountability in Networked Urban Climate Governance, *Glob. Environ. Polit.*, 16(2), 82–100, 2016.

Heidrich, O., Reckien, D., Olazabal, M., Foley, A., Salvia, M., Hurtado, S. D. G., Orru, H., Flacke, J.,

- Geneletti, D., Pietrapertosa, F., Hamann, J. J. and Tiwary, A.: National climate policies across Europe and their impacts on cities strategies, *J. Environ. Manage.*, 168, 36–45, doi:10.1016/j.jenvman.2015.11.043, 2016.
- IPCC: Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by B. Metz, O. R. Davidson, P. R. Bosch, R. Dave, and L. A. Meyer, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2007.
- IPCC: Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by O. Edenhofer, R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel, and J. C. Minx, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. [online] Available from: [http://www.ipcc.ch/pdf/assessment-report/ar5/wg3/ipcc\\_wg3\\_ar5\\_full.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg3/ipcc_wg3_ar5_full.pdf), 2014.
- IPCC, 2015: Human Settlements, Infrastructure, and Spatial Planning. In *Climate Change 2014: Mitigation of Climate Change: Working Group III Contribution to the IPCC Fifth Assessment Report*, pp. 923–1000, Cambridge University Press, Cambridge. [online] Available from: <https://doi.org/10.1017/CBO9781107415416.018>, 2015.
- Kilkis, S.: Data on cities that are benchmarked with the sustainable development of energy , water and environment systems index and related cross- sectoral scenario, *Data Br.*, 24, doi:10.1016/j.dib.2019.103856, 2019.
- Kona, A., Bertoldi, P., Monforti-Ferrario, F., Rivas, S. and Dallemand, J. F.: Covenant of mayors signatories leading the way towards 1.5 degree global warming pathway, *Sustain. Cities Soc.*, 41, 568–575, doi:10.1016/j.scs.2018.05.017, 2018.
- Kona, A., Bertoldi, P. and Kılış, Ş.: Covenant of Mayors: Local Energy Generation, Methodology, Policies and Good Practice Examples, *Energies*, 12(985), doi:10.3390/en12060985, 2019.
- Kona, A., Bertoldi, P., Baldi, M., Monforti-Ferrario, F., Palermo, V., Lo Vullo, E., Rivas-Calvete, S., Kakoulaki, G., Vetter, N., Thiel, C., Sgobbi, A., Ahlgren, C. and Posnic, B.: A dataset of GHG emissions for 6,200 cities in Europe and the Southern Mediterranean countries, Publications Office European Union, Luxembourg. [online] Available from: <https://doi.org/10.2905/57A615EB-CFBC-435A-A8C5-553BD40F76C9>, 2020.
- Kovac, A., Medaniel, S., Kona, A., Bertoldi, P. and Chavara, C.: *Aggregating Cities’ GHG Mitigation Targets with Modeled Emissions Scenarios*, Washington, DC. [online] Available from: <https://files.wri.org/s3fs-public/aggregating-cities-ghg-mitigation-targets.pdf>, 2020.
- Marcotullio, P. J., Sarzynski, A., Albrecht, J. and Schulz, N.: A Top-Down Regional Assessment of



Urban Greenhouse Gas Emissions in Europe, *Ambio*, 43, 957–968, doi:10.1007/s13280-013-0467-6, 2014.

610 Monforti-ferrario, F., Kona, A., Peduzzi, E., Pernigotti, D. and Pisoni, E.: The impact on air quality of energy saving measures in the major cities signatories of the Covenant of Mayors initiative, *Environ. Int.*, 118(February 2018), 222–234, doi:10.1016/j.envint.2018.06.001, 2020.

Nangini, C., Peregon, A., Ciaia, P., Weddige, U., Vogel, F., Wang, J., Bréon, F.-M., Bachra, S., Wang, Y., Gurney, K., Yamagata, Y., Appleby, K., Telahoun, S., Canadell, J. G., Grüber, A., Dhakal, S. and Creutzig, F.: A global dataset of CO<sub>2</sub> emissions and ancillary data related to emissions for 343 cities, *Sci. Data*, 6(1), 180280, doi:10.1038/sdata.2018.280, 2019.

Pablo-Romero, M. del P., Pozo-Barajas, R. and Sánchez-Braza, A.: Analyzing the effects of the benchmark local initiatives of Covenant of Mayors signatories, *J. Clean. Prod.*, 176, 159–174, doi:10.1016/J.JCLEPRO.2017.12.124, 2018.

620 Palermo, V., Bertoldi, P., Apostolou, M., Kona, A. and Rivas, S.: Assessment of climate change mitigation policies in 315 cities in the Covenant of Mayors initiative, *Sustain. Cities Soc.*, 102258 [online] Available from: <https://doi.org/10.1016/j.scs.2020.102258>, 2020a.

Palermo, V., Bertoldi, P., Apostolou, M., Kona, A. and Rivas, S.: Data on mitigation policies at local level within the Covenant of Mayors ' monitoring emission inventories, *Data Br.*, 32, 106217, doi:10.1016/j.dib.2020.106217, 2020b.

Peduzzi, E., Baldi, M. G., Pisoni, E., Kona, A., Bertoldi, P. and Monforti-ferrario, F.: Impacts of a climate change initiative on air pollutant emissions: Insights from the Covenant of Mayors, *Environ. Int.*, 145(April), 106029, doi:10.1016/j.envint.2020.106029, 2020.

630 Reckien, D., Salvia, M., Heidrich, O., Church, J. M., Pietrapertosa, F., De Gregorio-Hurtado, S., D'Alonzo, V., Foley, A., Simoes, S. G., Lorencová, E. K., Orru, H., Orru, K., Wejs, A., Flacke, J., Olazabal, M., Geneletti, D., Feliu, E., Vasilie, S., Nador, C., Krook-Riekkola, A., Matosović, M., Fokaides, P. A., Ioannou, B. I., Flamos, A., Spyridaki, N., Balzan, M. V., Fülöp, O., Paspaldzhiev, I., Grafakos, S. and Dawson, R.: How are cities planning to respond to climate change? Assessment of local climate plans from 885 cities in the EU-28, *J. Clean. Prod.*, 191, 207–219, doi:https://doi.org/10.1016/j.jclepro.2018.03.220, 2018.

635 Rosner, B.: Percentage Points for a Generalized ESD Many-Outlier Procedure, *Technometrics*, 25(2), 165–172, 1983.

Seem.: Using intelligent data analysis to detect abnormal energy consumption in buildings., *Energy Build.*, 39, 52–58, 2007.

640 Solazzo, E., Crippa, M., Guizzardi, D., Muntean, M., Choulga, M. and Janssens-Maenhout, G.: Uncertainties in the Emissions Database for Global Atmospheric Research (EDGAR) emission inventory

of greenhouse gases, *Atmos. Chem. Phys.*, 21, 5655–5683, doi:10.5194/acp-21-5655-2021, 2021.

van der Ven, H., Bernstein, S. and Hoffmann, M.: Valuing the Contributions of Nonstate and Subnational Actors to Climate Governance, *Glob. Environ. Polit.*, 17(1), 1–20, 2017.

- 645 Lo Vullo, E., Muntean, M., Duerr, M., Kona, A. and Bertoldi, P.: National and European Emission Factors for Electricity - Covenant of Mayors, Luxembourg. [online] Available from: <https://data.jrc.ec.europa.eu/collection/id-00172>, 2020.

## Tables

**Table 1. Datasets and their attributes.**

<u>GCoM dataset 2019: emission inventories</u>	<u>GCoM dataset 2019: ancillary data</u>
- <u>GCoM ID: identification code of the signatory</u>	- <u>GCoM ID: identification code of the signatory</u>
- <u>emission inventory id: identification code of the inventory</u>	- <u>signatory name and country code and Covenant Regions</u>
- <u>emission inventory sector: stationary sources (municipal, residential, institutional/tertiary buildings and facilities, as well as manufacturing and construction industries); mobile sources for transportation purposes (i.e. on-road, rail, waterborne navigation and off-road ) and Waste</u>	- <u>year of adhesion and population in the adhesion year and in 2018; signatory adhesion type</u>
- <u>type of emissions: direct emissions from fuel combustion as above described and from waste/wastewater sector; and indirect emissions due to consumption of grid-supplied energy consumption;</u>	- <u>longitude; latitude; area</u>
- <u>type of emission inventory: baseline or monitoring inventory</u>	- <u>regional identification code - level 3 (NUTS3): Nomenclature of Territorial Units for Statistics is a geocode standard for referencing the subdivisions of Member States for statistical purposes</u>
- <u>inventory year and population in the inventory year</u>	- <u>local Administrative Units identification code: level 1 and 2 (LAU1/LAU2)</u>
- <u>activity data and reporting unit: all activity data (i.e. final energy consumption) occurring in stationary sources and mobile sources for transportation purposes within the local authority boundary are reported in the baseline/monitoring inventories.</u>	- <u>Functional Urban Area identification code: (FUA)</u>
- <u>emission factor type: IPCC factor the activity-based approach or Life Cycle Approach</u>	- <u>Cities and greater cities identification code: CITY ID/GREATER CITY ID and the degree of urbanisation; type of local authority</u>
- <u>GHG emissions and reporting unit: occurring in stationary sources (excluding "energy generation" industries for and the ones under the EU- Emission Trading Schemes); occurring in mobile sources for transportation and non-energy related emissions from disposal and treatment of waste and wastewater generated within the city boundary are reported under waste/wastewater sector.</u>	- <u>Heating Degree-Days (HDD) and reference year HDD</u>
	- <u>GDP per capita at NUTS3 (average 2010-2018)</u>
	- <u>GHG emissions per capita in GCoM sectors in EDGAR v5.0 and reference years in EDGAR</u>
	- <u>mitigation reduction target 2020 or 2030: reduction target to be achieved through the implementation of the climate action plan by 2020/2030</u>
	- <u>reduction type: absolute or per capita</u>

**Formatted:** Font: (Default) Arial, 16 pt

**Formatted:** Font: (Default) Times New Roman, 10 pt

**Formatted:** Page break before

**Formatted:** Font: Times New Roman

**Formatted Table**

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: 9 pt

**Formatted:** Indent: Left: 0", Hanging: 0.09"

**Formatted:** Font: Times New Roman, 9 pt

**Formatted:** Font: Times New Roman

**Formatted:** Indent: Left: 0", Hanging: 0.09"

**Formatted:** Font: Times New Roman, 9 pt, English (United States)

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Indent: Left: 0", Hanging: 0.1"

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman, 9 pt

The "GCoM dataset 2019: Emission Inventories" are self-reported data in the MyCovenant platform of the GCoM reporting framework. Table 2 of the online dataset reports these emission related data, while Table 1 reports their metadata description. The "GCoM dataset 2019: ancillary data" comprises geographical attributes, socio-economic aspects and demographic characteristics at city level derived from Eurostat (i.e. the statistical office of the European Union) and from the EDGAR v5.0 (i.e. Emissions Database for Global Atmospheric Research). Table 3 of the online dataset reports these ancillary data, while Table 1 reports their metadata description.

**Table 221. Mapping of emission source categories in GCoM reporting framework with-based on the IPCC 2006 guidance**

Sectors and subsectors in GCoM reporting framework		IPCC (ref no.)	Description
Stationary energy	Residential buildings	1A4b; 1A1	All activities and related GHG emissions (direct emission from fuel combustion and indirect emission due to consumption of grid-supplied energy) occurring in stationary sources within the local authority boundary are reported. GHG emissions from sources covered by a regional or national <del>emissions</del> <u>Emissions trading Trading scheme Scheme</u> (ETS), or similar (i.e. industries with thermal energy in input below or equal to 20 MW) when ETS does not exist, are not accounted in the inventory. In addition, “energy generation” industries/facilities are not reported under this sector to avoid double counting with indirect emissions.
	Commercial building and facilities	1A4a; 1A1	
	Institutional buildings and facilities	1A4a; 1A1	
	Manufacturing, construction industries	1A1, 1A2; 1A1	
	Agriculture	1A4c; 1A1	
	Fugitive emissions	1B1, 1B2	
Transportation	On-road	1A3b; 1A1	All activities and related GHG emissions (direct emission from fuel combustion and indirect emission due to consumption of grid-supplied energy) occurring for transportation purposes within the local authority boundary will be reported. NB. <del>t</del> The GCoM dataset reported in this dataset does not reflect only the on road fraction of the transport sector, but all the emission in this sector.
	Rail	1A3c; 1A1	
	Waterborne navigation	1A3d; 1A1	
	Aviation	1A3a;	
	Off-road	1A3e; 1A1	
Waste	Solid waste disposal	4A	Sources related to disposal and treatment of waste and wastewater generating emissions within the city boundary are reported under Waste sector. Where waste/wastewater is used for energy generation, emissions are not reported under this sector to avoid double counting of indirect emission.
	Biological treatment	4B	
	Incineration and open burning	4C	
	Wastewater	4D	

Formatted: Font: 9 pt

Formatted Table

Formatted: Font: 9 pt

Formatted Table

Formatted: Font: 9 pt

Formatted Table

Formatted: Font: 9 pt, Font color: Black

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted Table

665

**Table 332. Statistical parameters of the GHG emissions in the GCoM datasets 2019**

Parameters	All CoM dataset 2019	Clean CoM dataset 2019
Number of signatories with complete GHG inventories in the Baseline year	6,239	6,200
Total population in the baseline year [Million inhabitants]	216.61	216.25
Mean [tCO <sub>2</sub> -eq/cap]	5.18	4.69
Median [tCO <sub>2</sub> -eq/cap]	4.78	4.75

Formatted: Font: Palatino Linotype, Not Bold

Formatted: Font: Times New Roman, 11 pt, Bold, English (United Kingdom)

Formatted: Font: 9 pt

Formatted Table

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Standard deviation [tCO <sub>2</sub> -eq/cap]	82.35	3.68
Skewness [tCO <sub>2</sub> -eq/cap]	76.39	2.37

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: Times New Roman

Table 443. Mapping of emission source categories with IPCC categories

GCoM sector	Sector code	Sector name	Inventory years	Country ISO code
Stationary energy/indirect emissions	1.A.1.a	1.A.1.a - Public Electricity and Heat Production	1990-2018	52 ISO codes
Stationary energy/residential	1.A.4.b	1.A.4.b - Residential	1990-2018	52 ISO codes
Stationary energy/Commercial/Institutional	1.A.4.a	1.A.4.a - Commercial/Institutional	1990-2018	52 ISO codes
Transportation	1.A.3.b	1.A.3.b - Road Transportation	1990-2018	52 ISO codes

Formatted: Font: Calibri, 11 pt, Not Bold

Formatted: Font: 9 pt

Formatted Table

Formatted: Font: 9 pt


Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: Times New Roman

## Figures



**Figure 1-GHG emissions per capita of signatories in the “CoM dataset 2019”. (Emission from all the CoM sectors, excluding manufacturing and construction industries). Global Covenant of Mayors signatories in Europe and South Mediterranean Countries as of 2019.**


The map “GCoM signatories in Europe and South Mediterranean Countries as of 2019” reports the location of the local authorities reporting to MyCovenant platform of the GCoM common reporting framework. Table 3 of the online dataset reports these signatories (8136) and their ancillary data, while Table 1 reports their metadata description.

This dataset covers the following countries, grouped into the Covenant region: European Union: EU-27; European Free Trade Association (EFTA) and UK: Iceland, Liechtenstein, Norway and Switzerland and United Kingdom; Western Balkans: Albania, Bosnia Herzegovina, Montenegro, Republic of North Macedonia, Serbia and Turkey; Eastern Europe: Armenia, Azerbaijan, Belarus, Georgia, Kazakhstan, Ukraine; Southern Mediterranean countries: Algeria, Lebanon, Jordan, Libya, Morocco, Palestine, Israel, Tunisia.

Formatted: Space Before: 12 pt, After: 3 pt, Line spacing: single, Page break before

City emission inventory data: "CoM dataset 2019: Emission Inventories"	Ancillary data institutional, demographic and socio-economic data: "CoM dataset 2019: Ancillary data"
<ul style="list-style-type: none"> <li>•CoM_ID: identification code of the signatory</li> <li>•signatory name</li> <li>•country code</li> <li>•Covenant region</li> <li>•Year of adhesion</li> <li>•population in the adhesion year</li> <li>•signatory adhesion type</li> <li>•type of emission inventory</li> <li>•inventory year</li> <li>•population in the inventory year</li> <li>•GHG emissions in stationary energy</li> <li>•GHG emissions in the transport sector</li> <li>•GHG emissions in the waste sector</li> <li>•GHG emissions total/per capita</li> <li>•mitigation target level by 2020/2030</li> </ul>	<ul style="list-style-type: none"> <li>•CoM_ID: identification code of the signatory</li> <li>•NUTS 3 region code: Nomenclature of Territorial Units for Statistics</li> <li>•LAU1:Local administrative unit 1 code</li> <li>•LAU2:Local administrative unit 2 code</li> <li>•CITY_ID: cities identification code</li> <li>•FUA_ID: Functional urban area</li> <li>•latitude</li> <li>•longitude</li> <li>•area</li> <li>•degree of urbanisation</li> <li>•population in the year 2018</li> <li>•heating degree days</li> <li>•GDP at NUTS 3 region (average 2010-2018)</li> <li>•GHG emissions per capita in EDGAR</li> </ul>

**Figure 1. Datasets and their attributes. Source of the “CoM dataset 2019: Emission Inventories” MyCovenant of Mayors platform, source of “CoM dataset 2019: Ancillary data” Eurostat and JRC.**



**Figure 222. -GHG emissions per capita of signatories in the “CoM dataset 2019”. (Emission from all the CoM sectors, excluding manufacturing and construction industries)**

Figure 2 shows the frequency distribution of GHG emissions per capita from emission inventories dataset, with observations that range from 0 to 80 tCO<sub>2</sub>-eq/cap, with a mean of 5.18 tCO<sub>2</sub>-eq/cap. These comprehends the emissions from all the GCoM sectors (stationary energy, transportation and waste), excluding manufacturing and construction industries. In the vertical axis the density values are reported, i.e. the share of signatories with the same range of GHG emission per capita, by the width of the range (0.01 in this case).

705



## Annex 1

Tables A.1 and A.2 show the values of absolute emissions (A.1) and three selected benchmark indicators – namely correlation, bias and NRMSE- (A.2) for both EDGAR v5.0 and GCoM datasets in road transport and energy in buildings sectors. Values are reported for both the whole set of selected signatories and for three subsets based on population size.

Firstly, it has to be noticed how correlation and NRMSE are consistently better for energy in buildings than road traffic and, for both sectors, tend to improve with increasing the city size, in agreement with the already mentioned fact that the coarse resolution of EDGAR v5.0 limits the description of smaller areas.

Analysing sectors, it is evident as GCoM provides consistently higher values of EDGAR v5.0 for energy in buildings sector, by an average value of 35 % with an overall correlation coefficient of 0.925. This relatively small bias and the good correlation values observed might be attributed to the fact that, in this sector EDGAR v5.0 splits sectorial emissions on the basis of population density, without at the moment considering regional data. Similarly, GCoM signatories collect data from utilities mainly, data that are themselves good proxies of national of energy usage pattern and fuels deployed.

Regarding traffic sector, table A.2 shows also how the bias between EDGAR v5.0 and GCoM emission data for road transport increases in function of the city size, with GCoM transport emissions lower than EDGAR data in smallest cities and overcoming them by a factor of three in largest cities. This is most probably due to the well known fact that largest cities act as traffic attractors: there the number of vehicles is not just proportional to the city population but depends also on the incoming and crossing traffic caused by cities activities, an effect that is captured by GCoM ground based data, but more difficult to be addressed by EDGAR that uses population density as a spatial proxy for emissions allocation. Moreover, the GCoM dataset reported here does not reflect only the on road fraction of the transport sector, but all the emission in this sector, due to the old version of the reporting platform that collected data without distinguishing the modal share. It has also to be considered that GCoM reports real data supplied from the transportation department, which are not necessarily caught in EDGAR: for instance, EDGAR v5.0 uses the average national fleet that could be quite different from a local one and, finally, that the default GCoM emission factors often used by signatories do not apply the very fine categorization of vehicle fleet applied in EDGAR.

On summary, observing the overall values of the benchmark indicators reported in the last line of Table A.2, and considering once again the deep differences in data sources, with GCoM based on local authorities knowledge, while EDGAR v5.0 is based on the use of population density as a main proxy, the behaviour observed does not come as a surprise and confirms the consistency between the two datasets, given the different methodologies applied.

**Formatted:** Heading 1, Space Before: 0 pt, After: 0 pt,  
Line spacing: single, Page break before

**Formatted:** Font: Times New Roman, 10 pt

**Table A.1 – Absolute emissions (MtCO<sub>2eq</sub>) for road transport and energy in buildings sectors in EDGAR v5.0 and GCoM datasets in selected signatories**

City Size	Number of signatories	Population in 2005 [million inh.]	Road transport emissions (EDGAR)	Road transport Emissions (GCoM)	Energy in Buildings emissions (EDGAR)	Energy in Buildings emissions (GCoM)
Pop. ≥ 500000	8	10.3	3.366	10.691	12.598	16.022
50000 < Pop. < 500000	96	16.3	10.909	25.097	12.525	24.566
Pop. ≤ 50000	1841	9.8	19.749	12.877	12.483	13.828
All Sample	1945	36.3	34.024	48.665	37.606	54.417

**Table A.2 – Benchmark indicators for road transport and energy in buildings sectors in EDGAR v5.0 and GCoM datasets in selected signatories**

City Size	Correlation (Road transport emissions)	Correlation (Energy in Buildings emissions)	BIAS (Road transport emissions)	BIAS (Energy in Buildings emissions)	NRMSE (Road transport emissions)	NRMSE (Energy in Buildings emissions)
Pop. ≥ 500000	0.928	0.945	218%	32%	1.481	0.449
50000 < Pop. < 500000	0.365	0.641	130%	48%	1.685	1.374
Pop. ≤ 50000	0.490	0.605	-35%	10%	1.536	1.411
All Sample	0.660	0.925	43%	35%	5.168	3.336