Response to Referee Comments

Manuscript "Reactive nitrogen fluxes over peatland and forest ecosystems using micrometeorological measurement techniques"

essd-2021-85

The authors sincerely thank the two reviewers for their thorough review. This document lists our responses to their comments.

Comment on essd-2021-85
Anonymous Referee #1

Referee comment on "Reactive nitrogen fluxes over peatland and forest ecosystems using micrometeorological measurement techniques" by Christian Brümmer et al., Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2021-85-RC1, 2021

Reviewer comment 1.1:

The auhors present the almost annual high-resolution (half-hourly) total Nr and NH3 flux datasets at two representive ecosystems. The dataset may be useful for model validation for atmospheric chemistry and land surface processes. However, considering publication I have to ask the authors to answer the following questions and comments;

- Could you demonstrate the novelty of the datasets more strongly, via comparing with the prior studies in terms of temporal resolution, data length, data quality, site characteristics (vegetation type)? For example, what about the quality or uncertainty of other past eddy covariance study for Nr over the vegetation compared with your datasets? How long is the longest record of past datasets, and where and what vegetation type?

Authors' response 1.1:

We appreciate the comment and agree that highlighting the novelty of the here presented datasets should be mentioned more strongly. The main reasons why there are not many eddy-covariance (EC) campaigns of reactive nitrogen (N_r) compounds and why campaign lengths are generally limited were already given in lines 51-62. These are – amongst others – high costs for maintenance, operation, and instruments, gas phase reactions, gas-aerosol interactions, basically a need for individual measurements for several compounds simultaneously as well as damping issues due to high-frequency losses.

We will further add an overview of reported EC studies of N_r compounds from literature, see Table R1 at the end of this document.

It can be derived that

- Most of the other presented campaigns are very limited in lengths. The only exceptions are the Munger et al. (1996) and the Horii et al. (2006) papers that explicitly mention the determination of a flux budget as an aim of their study.
- Most of the other studies follow different aims, some intend to test the suitability of the presented analyzer for EC measurements, others focus on correction factors or error analysis, while some are investigating processes of tropospheric chemistry.
- Many studies do not explicitly give flux uncertainties or detection limits for their systems. In those studies where numbers are mentioned, a large range of values is reported.

- No other EC ΣN_r (i.e. total $N_r)$ studies were found except for those by authors from this study.

We suggest to put the table into the Supplement of the article as some readers may find the information useful.

Additional text will be added in line 82 to properly link the novelty of the here presented datasets with the literature overview:

"The reader is referred to Table S1 in the supplementary section where an overview of previous eddy-covariance campaigns of different N_r compounds is given. The table highlights the limitations of campaign lengths, a wide range of flux uncertainties, and mainly different aims of previous studies such as testing the suitability of a certain analyzer, investigating processes of tropospheric chemistry, or derivation of correction factors."

Reviewer comment 1.2:

Another example is green-house gas flux; other ESSD papers demonstrated typically for greenhouse gas flux for longer time period (> several ten years), which enables us to discuss the annual carbon budget. Actually you mentioned about the impact of nitrogen deposition on carbon storage in 1443-448 in p15, but your WET site dataset for instance is collected for less than 1 yr and could not used for annual deposition/emission. Although this is partially described in Introduction section, more explanation is still required.

Authors' response 1.2:

It is true that we stress the possibility of the presented datasets for investigating potential effects of N deposition on CO₂ exchange. We agree that even for the >2.5-yr period at the forest site, it is hardly possible to indicate trends in the respective annual balances, let alone some sort of robust analysis on this time scale would be meaningful. It was meant that continuous N_r flux datasets like the ones that are here presented, offer a useful tool to investigate the relationship between N deposition and CO₂ exchange when carried out over multiple years. To clarify this, we rephrase line 443f. accordingly:

"Finally, continuous N_r flux datasets like the ones presented in this study can be used to analyse interactive effects with other components of biogeochemical cycles when carried out over multiple years."

Regarding the length of our presented datasets in comparison to typical greenhouse gas flux times series, it becomes clear that with the above-mentioned reasons such as the fact that campaigns have largely remained experimental due to high costs for devices, maintenance, and operation or the gas phase reactions requiring individual measurements of several species simultaneously (authors' response 1.1, also lines 51-62), the community is just simply not there yet where the CO_2 flux community is. It is the aim of this paper to provide a first piece into a direction of more and longer N_r flux measurements and should motivate other groups to work on joint initiatives as well as consolidated longer-term monitoring programs of N_r.

Reviewer comment 1.3:

- Please add the following data to your repository or if available from other projects (e.g., LTER, fluxnet, ...) please add these link address to your repository website: net radiation, sensible and latent heat, CO2 fluxes over the canopy, soil temp and moisture (for model validation), atmospheric pressure, precipitation, downward long-wave radiation (for model input), plant functional type (PFT), LAI, canopy height, leaf trait, soil type and texture (for input parameters). Management (fertilizer practice) data, too. They are

generally required for land surface modeling in addition to your concentration and flux data.

Authors' response 1.3:

We agree that providing the requested data is useful for running land-surface models. We checked the availability and will add those datasets to the repository that we either measured ourselves or where we have access to, see Table R2 for details.

Table R2: Availability of additional datasets to be added to the repository website from the peatland (WET) and the forest (FOR) site, where "x" indicates available and "o" indicates that these data will not be added as they were either not recorded or have an insufficient quality due to bad instrument performance. Fertilization is of no relevance at both sites (unmanaged).

	WET	FOR	
Rn	x	0	
LE	x	0	
Н	x	x	
F _{CO2}	x	(x), only 3 months	
Ts	x	0	
SWC	o, but WTD	0	
P _{bar}	x	x	
Precipitation	x	x	
Rlong,down	x	0	
PFT	х	х	
LAI	0	(x), modeled	
Canopy height	х	х	
Soil type	x	X	
Soil texture	х	x	
Fertilization	n.a.	n.a.	

Reviewer comment 1.4:

- I think your low-resolution data can be used for validation of your high-resolution data over time series and/or scatter plots. Currently this seems to be used just for dry deposition estimation though.

Authors' response 1.4:

We thank the reviewer for the suggestion and support the idea of verifying the agreement between monthly integrated DELTA (or passive sampler, PS) concentrations and averaged QCL values for the corresponding exposition periods. However, we think that the high-resolution data should be the reference and not the other way around due to precision, accuracy and rigorous calibration of the QCL and no further error sources from post-field wet-chemical lab analyses involved in the DELTA and passive sampler procedures.

For fluxes, a validation of N deposition estimation with a model using monthly concentration input data is not recommended as the time resolution does not resolve micrometeorological variability such as diurnal courses of temperature, radiation, friction velocity or humidity. This is extensively shown and discussed in the publication of Schrader et al. (2018) and would not add any new insight to the here presented dataset publication.

We limit our analysis to the NH₃ campaigns at the WET and the FOR site, because for ΣN_r at the WET site, DELTA and PS exposition did not overlap with the campaign period of the TRANC and at the FOR site the same topic is discussed in detail in the publications by Wintjen et al. (2022a, 2022b) for ΣN_r .

Figures R1 to R5 show the respective comparison of QCL vs. low resolution data from DELTA or PS. Data from the latter two low-resolution methods were consistently underestimated compared to QCL values. It must be noted that at the WET site, the data gap in QCL measurements in April 2014 led to a larger difference between QCL and DELTA as the values were generally decreasing from the early April peak due to fertilization on adjacent agricultural sites, hence biasing the QCL average. A full time series would have probably led to a slope much closer to 1 in Figure R2 as was found in the comparison between QCL and DELTA at the FOR site (*cf.* Figure R4).

We suggest adding these figures to the supplementary section and will extend lines 328f. in Section 5 (Data description) as follows:

"In the following, we briefly highlight key characteristics of the high-resolution TRANC and QCL data. A comparison with low-resolution concentration data from DELTA denuders and passive samplers is given in the supplementary section."



Figure R1: Concentration time series of NH_3 at the peatland (WET) site. Horizontal red lines correspond to the exposition time of the DELTA denuders. For better comparability, averages of the QCL are shown in blue for the same periods.



Figure R2: Scatter plot of NH₃ concentration from QCL and DELTA denuders corresponding to identical periods at the peatland (WET) site.



Figure R3: Concentration time series of NH_3 at the forest (FOR) site. Horizontal blue and red lines correspond to the exposition time of the DELTA denuders and passive samplers (PS), respectively. For better comparability, averages of the QCL are shown in black for the same periods.



Figure R4: Scatter plot of NH₃ concentration from QCL and DELTA denuders corresponding to identical periods at the forest (FOR) site.



Figure R5: Scatter plot of NH₃ concentration from QCL and passive samplers (PS) corresponding to the same periods at the forest (FOR) site.

Reviewer comment 1.5:

- please show the accuracy of gap-filling method using dry deposition models (4.3.2 and 4.4) via comparing the model outputs with original flux data over time series and/or scatter plots. Then you can tune the parameters such as Massad et al. (2010) model to reproduce the original flux data (unnecessary use of default parameter sets, if the results are improved)

Authors' response 1.5:

We appreciate the comment and intend to add Figures R6 to R8 to the supplementary section. Although the variability in the scatter plots showing valid measured vs. modeled fluxes is relatively high (Figure R6), the impact of the total cumulative exchange appears to be acceptable (Figures R7 and R8). The analysis is limited to the WET NH₃ and the FOR TRANC campaigns as no model application was performed for WET TRANC due to missing data of N_r species other than NH₃ and due to no flux calculation at FOR NH₃ (for reasons see the second part of the response to Comment 1.7).

For the WET NH₃ campaign, the model underestimates the QCL measurements by 13.5% when taking all valid measured half-hours into account. At the FOR TRANC campaign, we only show a comparison for the summer months in 2016, where the model overestimates the TRANC measurements by 65%. This is the period of highest fluxes and consequently the highest potential absolute deviations. It is shown in Wintjen et al. (2022b) that over the 2.5-yr campaign the difference between measurements and model is overall in the range of ~25%.

Information where the reader is referred to the respective figures will be added after line 295 (end of Section 4.4):

"The accuracy of the gap-filling method from this section and from 4.3.2 by comparing the model outputs with original flux data is shown in the supplementary section."

We appreciate the idea of using the valid measured fluxes for fine-tuning parameters in the models. However, as indicated multiple times in Section 6 (Potential applications), the aim of this paper is to present the datasets from the measurements by stressing all required steps for robust data processing and thorough QA/QC. A detailed model parameter investigation would go beyond the scope of this study and comprises one of many potential logical steps for a follow-up paper. We therefore do not intend to include such an analysis in the here presented measurement dataset presentation.



Figure R6: Measured vs. modeled deposition data in half-hourly time resolution at the forest (FOR, left panel) and peatland site (WET, right panel). FOR data comprise the period mid-July to end of September 2016. For the WET site the entire campaign from February to May 2014 is shown.



Figure R7: Time series (upper panel) and cumulative curves (lower panel) of measured vs. modeled deposition data in half-hourly time resolution at the peatland site (WET) from February to May 2014.



Figure R8: Time series (upper panel) and cumulative curves (lower panel) of measured vs. modeled deposition data in half-hourly time resolution at the forest site (FOR) from mid-July to September 2016.

Reviewer comment 1.6:

- Table 3 lacks the info about NH3 concentration and flux measurements and low-resolution datasets (while all data are available in the website).

Authors' response 1.6:

We will add the NH_3 related entries (concentration, fluxes, gap-filled time series with units and short descriptions) as new lines in Table 3 regardless of its presence in the respective site data table. To clarify this, we add the following sentence to the table caption:

"Note that not every column header exists in every site data table due to the focus on the specific gas and gap-filling method."

We will also add the headers from the low-resolution DELTA datasets. An additional column will be inserted to give the time resolution of the respective dataset.

Reviewer comment 1.7:

- Fig. 3: pls specity if the flux data is gap-filled or not. Also could you add the lower detection limit for the NH3 flux of "n.a." over the forest site?

Authors' response 1.7:

We thank the reviewer for the catch. The information was missing, but is surely important. The idea for the flux data presentation in Figure 3 is to show the most complete time series (column 3) and based on that a robust mean diurnal curve (column 4). Hence, for campaign 1 and 2 at the WET sites, the dataset gap-filled by mean diurnal variation (MDV) and for campaign 3 at the FOR site, the dataset gap-filled by the model is used. We will add the following information to the figure caption:

"Flux data shown in column 3 are the gap-filled time series using the mean diurnal variation (MDV) method for the WET campaigns and the model-filled time series for the FOR campaign. See Section 4.3 for details. The mean diurnal flux courses in column 4 are based on the given time series in column 3."

Regarding the lower detection limit, we state in lines 345-346 that the observed low signal-to-noise ratio as a consequence of insufficient high-frequency variability in the low-concentration regime made a robust flux detection impossible and referred to the publication of Zöll et al. (2016). For completeness, we will add the value in the caption of Figure 3:

"Insufficient high-frequency variability in the low-concentration regime of ammonia measured by QCL at the FOR site made a robust covariance detection unreliable with fluxes largely under the lower detection limit of 7.75 ng N m⁻² s⁻¹. NH₃ fluxes at the FOR site are therefore not given."

Referee comment on "Reactive nitrogen fluxes over peatland and forest ecosystems using micrometeorological measurement techniques" by Christian Brümmer et al., Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2021-85-RC2, 2021

Reviewer comment 2:

This paper describes an important database of nitrogen fluxes over two ecosystems. The site description is good and methodology is presented in details. Essential steps were carried out correctly by the authors, especially regarding the high frequency corrections due to sensor separation and lags between the sonic anemometer and gas analyzers. Gapfilling and uncertainty analysis were also carried out according to current methods, providing estimates of annual depositions.

Authors' response 2:

We appreciate the positive evaluation by the reviewer. We are glad to see that both dataset and text are apparently presented in a clear and understandable way.

Table R1: Overview of literature presenting eddy-covariance measurements of reactive nitrogen compounds. Some additional flux campaigns are listed in the publication of Walker et al. (2020).

Paper	Compound	Main aim of study	Dataset length	Flux uncertainty / detection limit	Vegetation type
Ammann et al. (2012)	ΣNr	Suitability of converter for EC measurements	Few weeks are shown for cross-validation with other techniques	~5 ng N m ⁻² s ⁻¹ (upper flux detection limit)	Managed grassland
Brümmer et al. (2013)	ΣNr	Temporal dynamics, controlling factors, and seasonal N budget	11 months	~6.6 ng N m ⁻² s ⁻¹ (upper flux detection limit)	Cropland (winter wheat)
Eugster and Hesterberg (1996)	NO ₂	Deriving transfer resistances	Four different periods with a total of 68 days	Not explicitly given	Rural litter meadow
Famulari et al. (2004)	NH ₃	Suitability of TDLAS system for EC; cross-validation with AGM	2 months	Not explicitly given, only standard deviation of fluxes for entire campaign	Managed grassland
Farmer and Cohen (2008)	HNO ₃ , Σ AN, Σ PN and NO ₂	In-canopy chemical analysis	12 months	Not explicitly given	Ponderosa pine plantation
Farmer et al. (2006)	HNO ₃ , Σ AN, Σ PN and NO ₂	Suitability of TD-LIF system for EC	12 months; shorter periods are shown from different seasons	<1 ng N m ⁻² s ⁻¹ ; <20% relative errors at low wind speed (<1 m s ⁻¹)	Ponderosa pine plantation
Farmer et al. (2011)	Aerosols (NH4, SO4, NO3)	Suitability of HR-AMS system for EC	15 days	~0.4 to 6.4 ng m ⁻² s ⁻¹ depending on substance and mode; typical single flux measurement was below DL for NH ₄ fragments	Ponderosa pine plantation
Ferrara et al. (2012)	NH ₃	Comparison of high-frequency correction methodologies using QC-TILDAS	13 days	~75 ng N m ⁻² s ⁻¹ (flux detection limit)	Cropland (sorghum)
Ferrara et al. (2016)	NH ₃	Temporal dynamics of NH ₃ volatilization after slurry application using QC-TILDAS	~14 days	Only MAE (4700 ng NH ₃ m ⁻² s ⁻¹) and RMSE (12000 ng NH ₃ m ⁻² s ⁻¹) given	Maize stubbles and Italian ryegrass

Ferrara et al.	NH ₃	Evaluation of measurement	21 days	13.6 and 20.7 ng m ⁻² s ⁻¹ at 95 and	Cropland (faba bean)
(2021)		errors using QCL spectrometer		99% CI, respectively	
Horii et al.	NO, NO ₂ ,	Impacts of temporal dynamics	7 months, but no time series	Not explicitly given	Mixed deciduous
(2004)	O₃	on tropospheric chemistry and	shown		forest
		parameterizations			
Horii et al.	NO _x , NO _y	Concentration and flux budgets	5 months, but only time	Not explicitly given	Mixed deciduous
(2006)		of N _r , inferring HNO ₃ ,	series of ~2 weeks are shown		forest
		validation of deposition			
		velocities			
Marx et al.	ΣN_r	Suitability of converter for	1-week validation, 11 months	Not explicitly given as aim was on	Managed grassland
(2012)		capturing all N _r species at high	field campaign	concentrations and fast response	and cropland (winter
		frequency			wheat)
Min et al.	NO, NO ₂	Comparison of gradient and	6 weeks, no time series	<8% for NO flux; <6% for NO ₂ flux;	Ponderosa pine
(2014)		direct flux measurements;	shown	0.08 ppt m s ⁻¹ (NO); 0.14 ppt m s ⁻¹	plantation
		within-canopy chemistry of		(NO ₂)	
		NO _x			
Moravek et	NH ₃	Quantify impact of adsorption	5 months	Median flux detection limit of 2.15	Corn crop field
al. (2019)		on time response of the system		ng m ⁻² s ⁻¹	
Munger et al.	NO _γ , O ₃	Response of NO _y deposition to	5 years	Only given for concentrations (~50	Mixed deciduous
(1996)		environmental conditions		ppt at the mixed forest site and	forest and spruce
				<10 ppt at the spruce woodland)	woodland
Rummel et	NO	Flux pattern within the canopy	3 months	0.07 ng N m ⁻² s ⁻¹	Amazonian rain forest
al. (2002)					
Sintermann	NH ₃	Suitability of a CIMS (chemical	Few days	5 ng N m ⁻² s ⁻¹	Crop stubble field and
et al. (2011)		ionization mass spectrometry)			cut grassland
		instrument for EC			
		measurements			
Sun et al.	NH ₃	Suitability of the open-path	2 weeks	1.3 +/- 0.5 ng m ⁻² s ⁻¹	Cattle feedlot
(2015)		NH ₃ sensor for EC			
		measurements and			
1					•
		comparison to other			

Wang et al.	NH ₃	Suitability of the open-path	1 week	7.1 ug N m ⁻² h ⁻¹	Subtropical rice paddy
(2021)		NH₃ sensor for EC			
		measurements			
Whitehead et	NH ₃	Suitability and inter-	2 campaigns, only few days	Not explicitly given	Managed grassland
al. (2008)		comparison of different	are presented		
		analyzers			

References in Table R1:

Ammann, C., Wolff, V., Marx, O., Brümmer, C., and Neftel, A.: Measuring the biosphere-atmosphere exchange of total reactive nitrogen by eddy covariance, Biogeosciences, 9, 4247–4261, https://doi.org/10.5194/bg-9-4247-2012, 2012.

Brümmer, C., Marx, O., Kutsch, W., Ammann, C., Wolff, V., Flechard, C. R., and Freibauer, A.: Fluxes of total reactive atmospheric nitrogen (ΣNr) using eddy covariance above arable land, Tellus B, 65, 19770, doi:10.3402/tellusb.v65i0.19770, 2013.

Eugster, W. and Hesterberg, R.: Transfer resistances of NO2 derived from eddy correlation flux measurements over a litter meadow at a rural site on the Swiss Plateau. Atmos. Environ., 30, 1247–1254, 1996.

Famulari, D., Fowler, D., Hargreaves, K., Milford, C., Nemitz, E., Sutton, M. A., and Weston, K.: Measuring eddy covariance fluxes of ammonia using tunable diode laser absorption spectroscopy, Water, Air Soil Pollut. Focus, 4, 151–158, 2004.

Farmer, D. K. and Cohen, R. C.: Observations of HNO3, ΣAN, ΣPN and NO2 fluxes: evidence for rapid HOx chemistry within a pine forest canopy. Atmos. Chem. Phys., 8(14), 3899–3917, doi:10.5194/acp-8-3899-2008, 2008.

Farmer, D. K., Wooldridge, P. J., and Cohen, R. C.: Application of thermal-dissociation laser induced fluorescence (TD-LIF) to measurement of HNO3, Σalkyl nitrates, Σperoxy nitrates, and NO2 fluxes using eddy covariance, Atmos. Chem. Phys., 6, 3471–3486, https://doi.org/10.5194/acp-6-3471-2006, 2006.

Farmer, D. K., Kimmel, J. R., Phillips, G., Docherty, K. S., Worsnop, D. R., Sueper, D., Nemitz, E., and Jimenez, J. L.: Eddy covariance measurements with high-resolution time-of-flight aerosol mass spectrometry: a new approach to chemically resolved aerosol fluxes, Atmos. Meas. Tech., 4, 1275–1289, https://doi.org/10.5194/amt-4-1275-2011, 2011.

Ferrara, R. M., Loubet, B., Di Tommasi, P., Bertolini, T., Magliulo, V., Cellier, P., Eugster, W., and Rana, G.: Eddy covariance measurement of ammonia fluxes: Comparison of high frequency correction methodologies, Agr. Forest Meteorol., 158–159, 30–42, 2012.

Ferrara, R. M., Carozzi, M., Di Tommasi, P., Nelson, D. D., Fratini, G., Bertolini, T., Magliulo, V., Acutis, M., and Rana, G.: Dynamics of ammonia volatilisation measured by eddy covariance during slurry spreading in north Italy. Agric. Ecosys. Environ., 219, 1–13, <u>https://doi.org/10.1016/j.agee.2015.12.002</u>, 2016.

Ferrara, R.M., Di Tommasi, P., Famulari, D., and Rana, G.: Limitations of an Eddy-Covariance System in Measuring Low Ammonia Fluxes. Boundary-Layer Meteorol 180, 173–186, <u>https://doi.org/10.1007/s10546-021-00612-6</u>, 2021.

Horii, C. V., Munger, J. W., and Wofsy, S. C.: Fluxes of nitrogen oxides over a temperate deciduous forest, J. Geophys. Res., 109, D08305, doi:10.1029/2003JD004326, 2004.

Horii, C. V., Munger, J. W., Wofsy, S. C., Zahniser, M., Nelson, D., and McManus, J. B.: Atmospheric reactive nitrogen concentration and flux budgets at a Northwestern US forest, Agr. Forest Meteorol., 136, 159–174, 2006.

Marx, O., Brümmer, C., Ammann, C., Wolff, V., and Freibauer, A.: TRANC – a novel fast-response converter to measure total reactive atmospheric nitrogen, Atmos. Meas. Tech., 5, 1045–1057, https://doi.org/10.5194/amt-5-1045-2012, 2012.

Min, K.-E., Pusede, S. E., Browne, E. C., LaFranchi, B. W., and Cohen, R. C.: Eddy covariance fluxes and vertical concentration gradient measurements of NO and NO2 over a ponderosa pine ecosystem: observational evidence for within-canopy chemical removal of NOx, Atmos. Chem. Phys., 14, 5495–5512, https://doi.org/10.5194/acp-14-5495-2014, 2014.

Moravek, A., Singh, S., Pattey, E., Pelletier, L., and Murphy, J. G.: Measurements and quality control of ammonia eddy covariance fluxes: A new strategy for high frequency attenuation correction, Atmospheric Measurement Techniques, 12, 6059–6078, https://doi.org/10.5194/amt-12-6059-2019, 2019.

Munger, J. W., Wofsy, S. C., Bakwin, P. S., Fan, S.-M., Goulden, M. L., Daube, B. C., Goldstein, A. H., Moore, K. E., and Fitzjarrald, D. R.: Atmospheric deposition of reactive nitrogen oxides and ozone in a temperate deciduous forest and a subarctic woodland: 1. Measurements and mechanisms, J. Geophys. Res., 101(D7), 12639–12657, doi:10.1029/96JD00230, 1996.

Rummel, U., Ammann, C., Gut, A., Meixner, F. X., and Andreae, M. O.: Eddy covariance measurements of nitric oxide flux within an Amazonian rain forest, J. Geophys. Res., 107, 8050, doi:10.1029/2001JD000520, 2002.

Sintermann, J., Spirig, C., Jordan, A., Kuhn, U., Ammann, C., and Neftel, A.: Eddy covariance flux measurements of ammonia by high temperature chemical ionisation mass spectrometry, Atmos. Meas. Tech., 4, 599–616, doi:10.5194/amt-4-599-2011, 2011.

Sun, K., Tao, L., Miller, D. J., Zondlo, M. A., Shonkwiler, K. B., Nash, C., and Ham, J. M.: Open-path eddy covariance measurements of ammonia fluxes from a beef cattle feedlot, Agr. Forest Meteorol, 213, 193–202, <u>https://doi.org/10.1016/j.agrformet.2015.06.007</u>, 2015.

Walker, J. T., Beachley, G., Zhang, L., Benedict, K. B., Sive, B. C., and Schwede, D. B.: A review of measurements of air-surface exchange of reactive nitrogen in natural ecosystems across North America, Science of The Total Environment, 698, 133975, <u>https://doi.org/10.1016/j.scitotenv.2019.133975</u>, 2020.

Wang, K., Kang, P., LU, Y., Zheng, X., Liu, M., Lin, T.-J., Butterbach-Bahl, K., and Wang, Y.: An open-path ammonia analyzer for eddy covariance flux measurement, Agr. Forest Meteorol, 308–309, 108570, <u>https://doi.org/10.1016/j.agrformet.2021.108570</u>, 2021.

Whitehead, J. D., Twigg, M., Famulari, D., Nemitz., E., Sutton, M. A., Gallagher, M. W., and Fowler, D.: Evaluation of Laser Absorption Spectroscopic Techniques for Eddy Covariance Flux Measurements of Ammonia, Environ. Sci. Technol., 42, 2041–2046, 2008.

References from response document:

Schrader, F., Schaap, M., Zöll, U., Kranenburg, R., and Brümmer, C.: The hidden cost of using low-resolution concentration data in the estimation of NH3 dry deposition fluxes. Scientific Reports, 8(1), 969, <u>https://doi.org/10.1038/s4159 8-017-18021-6</u>, 2018.

Wintjen, P., Schrader, F., Schaap, M., Beudert, B., and Brümmer, C.: Forest-atmosphere exchange of reactive nitrogen in a remote region – Part I: Measuring temporal dynamics. Biogeosciences, *in press*, 2022a.

Wintjen, P., Schrader, F., Schaap, M., Beudert, B., and Brümmer, C.: Forest-atmosphere exchange of reactive nitrogen in a remote region – Part II: Modeling annual budgets. Biogeosciences, *under review*, 2022b.