

Reconstructing 6-hourly PM_{2.5} datasets from 1960 to 2020 in China

Junting Zhong¹, Xiaoye Zhang^{1, 4*}, Ke Gui¹, Jie Liao², Ye Fei², Lipeng Jiang³, Lifeng Guo¹, Liangke Liu⁵, Huizheng Che¹, Yaqiang Wang¹, Deying Wang¹, Zijiang Zhou²

¹State Key Laboratory of Severe Weather & Key Laboratory of Atmospheric Chemistry of CMA, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

² National Meteorological Information Center, Beijing, 100081, China

³ Earth System Numerical Prediction Center, Beijing, 100081, China

⁴Center for Excellence in Regional Atmospheric Environment, IUE, Chinese Academy of Sciences, Xiamen, 361021, China.

⁵Department of Earth System Science, Tsinghua University, Beijing 100084, China

Correspondence to: Xiaoye Zhang (xiaoye@cma.gov.cn)

16 **Abstract**

17 Fine particulate matter (PM_{2.5}) has altered radiation balance on earth and raised environmental and
18 health risks for decades but has only been monitored widely since 2013 in China. Historical long-
19 term PM_{2.5} records with high temporal resolution are essential but lacking for both research and
20 environmental management. Here, we reconstruct a site-based PM_{2.5} dataset at 6-hour intervals from
21 1960 to 2020 that combines long-term visibility, conventional meteorological observations,
22 emissions, and elevation. The PM_{2.5} concentration at each site is estimated based on an advanced
23 machine learning model, LightGBM, that takes advantage of spatial features from 20 surrounding
24 meteorological stations. Our model's performance is comparable to or even better than those of
25 previous studies in by-year cross validation (CV) ($R^2=0.7$) and spatial CV ($R^2=0.76$) and is more
26 advantageous in long-term records and high temporal resolution. This model also reconstructs a
27 $0.25^\circ \times 0.25^\circ$, 6-hourly, gridded PM_{2.5} dataset by incorporating spatial features. The results show
28 PM_{2.5} pollution worsens gradually or maintains before 2010 from an interdecadal scale but mitigates
29 in the following decade. Although the turning points vary in different regions, PM_{2.5} mass
30 concentrations in key regions decreased significantly after 2013 due to clean air actions. In particular,
31 the annual average value of PM_{2.5} in 2020 is nearly the lowest since 1960. These two PM_{2.5} datasets
32 (publicly available at <https://doi.org/10.5281/zenodo.6372847>) provide spatiotemporal variations at
33 high resolution, which lay the foundation for research studies associated with air pollution, climate
34 change, and atmospheric chemical reanalysis.

35

1 Introduction

In the past decades, anthropogenic emissions of reactive gases and aerosols have been emitted increasingly in the atmosphere and thus led to a substantial increase in fine particulate matter (PM_{2.5}). Increased PM_{2.5} has strongly interacted with solar radiation through absorption and scattering, thereby reducing visibility and influencing the earth's radiance balance. Inhalable PM_{2.5} has increased human morbidity and mortality through penetrating the respiratory system (Pope et al., 2002; Beelen et al., 2007; Chen et al., 2016b). To evaluate the impacts of PM_{2.5} pollution on environment, climate, and health, the primary concern is to understand the spatiotemporal variations of PM_{2.5} concentrations. Namely, extended PM_{2.5} records with high temporal resolution lay the foundation for research studies associated with air pollution, climate change, and environmental health. Nevertheless, it was not until 2013 that the Ministry of Ecology and Environment (MEE) established a nationwide PM_{2.5} monitoring network. Long-term, accurate historical PM_{2.5} datasets are lacking for both research and environmental management.

Chemical transport models (CTMs) are expected to simulate the spatial and temporal variations of PM_{2.5} with reasonable emission inventories inputted. However, significant uncertainties still exist in historical emission inventories and physicochemical mechanisms, which resulted in inevitable biases in the simulated absolute values of PM_{2.5}. Satellite-based aerosol optical depth (AOD), which measures the aerosol extinction of the solar beam, is an indicator of ground-level aerosols. AOD data products from Moderate Resolution Imaging Spectroradiometer (MODIS) have broad spatial coverage and relatively long observation periods (~ 20 years). Therefore, assimilating satellite-retrieved AOD to construct atmospheric chemical reanalysis is a practical approach to reducing PM_{2.5} biases. In recent years, several international aerosol reanalysis datasets have been developed preliminarily, including the reanalysis data produced by the Copernicus Atmosphere Monitoring Service (CAMS) from the European Centre for Medium-Range Weather Forecasts (ECMWF) (Inness et al., 2019), the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) from the National Aeronautics and Space Administration (NASA) (Gelaro et al., 2017; Randles et al., 2017), aerosol reanalysis from the Navy Aerosol Analysis and Prediction System (NAAPS) (Lynch et al., 2016) and the Japanese Reanalysis for Aerosol (JRAero) from the Japanese Meteorological Agency (Yumimoto et al., 2017). In particular, CAMS produced gridded PM₁, PM_{2.5}, and PM₁₀ data at 80 km resolution since 2003 by assimilating satellite retrievals of total AOD, total tropospheric NO₂ column, total O₃ column, CO column, and vertical profiles (Inness et al., 2019). MERRA-2 reanalysis includes PM_{2.5} and PM₁₀ at 50 km resolution since 1980 by assimilating ground-based and satellite-retrieval (Gelaro et al., 2017; Randles et al., 2017). NAAPS generates gridded AOD data at ~100 km resolution from 2003 to 2013 by assimilating satellite-based AOD products (Lynch et al., 2016). JRAero provides PM_{2.5} and PM₁₀ at ~100 km resolution from 2011 to 2015 by assimilating satellite AOD data (Yumimoto et al., 2017). These reanalysis data have contributed significantly to research in aerosol-related fields. However, there are still some weaknesses in accuracy, spatial resolution, time span, and types of assimilated data. In China, the highest horizontal resolution of the four reanalysis is only 50 km, and this coarse grid setting may not be sufficient to capture the spatial differences in atmospheric pollutants at regional scales. In terms of the type of aerosol data assimilation, these reanalysis data mainly assimilate satellite-based and ground-based AOD, and do not take into account ground PM_{2.5} observations.

To overcome the reanalysis's weaknesses in low spatial resolution and high biases, numerical

researchers focus on constructing relatively long-term $PM_{2.5}$ datasets based on machine learning techniques that fuse multisource data, including satellite-retrieved AOD, CTM simulations, and even atmospheric chemical reanalysis. For example, Ma et al. (2016) estimated daily $PM_{2.5}$ records at 0.1° resolution between 2004-2013 with MODIS AOD. Liang et al. (2020) rebuilt monthly $PM_{2.5}$ concentrations at 1 km resolution during 2000-2016 based on the multiangle implementation of atmospheric correction (MAIAC) from MODIS and reanalysis AOD and $PM_{2.5}$ data from MERRA-2. Geng et al. (2021) reconstructed daily, 10 km $PM_{2.5}$ data between 2000-2020 with MODIS AOD and CTM simulations. Wei et al. (2021a) regenerated monthly, 1 km $PM_{2.5}$ records between 2000-2018 based on MAIAC AOD. Huang et al. (2021) estimated $1\text{ km} \times 1\text{ km}$ $PM_{2.5}$ concentrations daily between 2013-2019 based on MAIAC AOD and CTM outputs. However, some inherent limitations in satellite-based AOD are challenging to overcome. Due to the low sampling frequency of satellite-retrieved AOD, AOD-based $PM_{2.5}$ datasets are limited to a maximum temporal resolution of one day. With AOD over land unavailable before 2000, these $PM_{2.5}$ datasets can only be back-calculated to 2000 at the earliest. Although recent studies focus on estimating hourly $PM_{2.5}$ during the daytime based on AOD from geostationary satellites like Himawari 8 (Chen et al., 2019; Yan et al., 2020; Wang et al., 2021; Wei et al., 2021b), obtained $PM_{2.5}$ datasets can only extend for several years, and the data is missing at night or with cloud cover.

Compared with satellite data, ground-based meteorological observations have the advantages of long sequence time, high temporal resolution, and good data integrity. In China, the national meteorological observation network of the China Meteorological Administration (CMA) was established in the 1950s and is capable of continuously observing 6-hourly meteorological data on visibility and conventional meteorological variables, including temperature, pressure, wind, and relative humidity (RH). The number of national stations exceeded 2,000 in 1960 and stabilized at around 2,450 afterward. Studies have shown that visibility and conventional meteorological variables are closely related to $PM_{2.5}$ (Zhang et al., 2013a; Zhang et al., 2013b; Zhang et al., 2015; Wang et al., 2018; Zhu et al., 2018; Zhong et al., 2018). For example, low wind speed is highly unfavorable to the horizontal diffusion of pollutants (Zhang et al., 2013b). The increase in RH favors the hygroscopic growth of $PM_{2.5}$ and also promotes the accelerated conversion of gaseous precursors to particulate matter, leading to a rapid increase in $PM_{2.5}$ concentrations (Pilinis et al., 1989; Ervens et al., 2011; Kuang et al., 2016). Atmospheric visibility is directly related to $PM_{2.5}$ mass concentrations under dry conditions and non-linearly related to $PM_{2.5}$ and RH under humid conditions (Wang et al., 2019). Therefore, better results may be achieved if these ground-based meteorological data can be used to estimate historical $PM_{2.5}$ data in China. Liu et al. (2017) first estimated monthly visibility-based $PM_{2.5}$ concentrations between 1957-1964 and 1973-2014 based on 674 publicly available meteorological stations. Gui et al. (2020) constructed a virtual daily $PM_{2.5}$ network at 1180 meteorological sites between 2017-2018. Our previous research also shows that the visibility-based machine learning model that takes advantage of spatial features has great potential in reconstructing historical $PM_{2.5}$ datasets with long-term records and high temporal resolution (Zhong et al., 2021). In this study, we reconstruct a site-based $PM_{2.5}$ dataset at 6-hour intervals from 1960 to 2020 based on long-term visibility and conventional meteorological observations from ~2450 national stations, together with emissions and elevation. The $PM_{2.5}$ concentration at each site is estimated based on a Light Gradient Boosting Machine (LightGBM) model that takes advantage of spatial features from 20 surrounding meteorological stations. By incorporating spatial features, this model also reconstructs a $0.25^\circ \times 0.25^\circ$, 6-hourly, gridded $PM_{2.5}$ dataset. These two $PM_{2.5}$

123 datasets provide spatiotemporal variations at high resolution, which constitute the basis for research
124 studies associated with air pollution, climate change, and atmospheric chemical reanalysis.

125

2 Data and Methods

2.1 Multisource input data

Observational PM_{2.5} data. The MEE began laying out a PM_{2.5} monitoring network in January 2013, expanding the scope from key regions including the North China Plain (NCP), the Yangtze River Delta (YRD), the Pearl River Delta (PRD), and the Sichuan Basin (SB) as well as municipalities directly under the Central Government and provincial capitals, to 113 key and model cities for environmental protection, and eventually to all cities above prefecture level, with the number of observation sites expanded from the initial 520 to over 1,600. Since then, PM_{2.5} mass concentrations have been recorded continuously using the β -absorption methods or a micro-oscillating balance following a standard protocol (Huang et al., 2021). Hourly PM_{2.5} data of all sites between 2013-2020 are collected from the China National Environmental Monitoring Center (CNEMC, <http://www.cnemc.cn>). To produce high-quality PM_{2.5} data, a series of quality controls were conducted, including integrity checking, duplicate rejection, and outlier handling. All sites with a proportion of valid PM_{2.5} records exceeding 60% were considered. For each site, identical data for 3 consecutive hours were excluded first, and PM_{2.5} values over three standard deviations from 24-hour and 3-day moving average were regarded as outliers and discarded then. Eventually, PM_{2.5} data from 1485 sites remained for model development and application. In addition, pre-2013 PM_{2.5} measurements in US embassies in Beijing and Shanghai are used for independent validation evaluations (<http://www.stateair.net/web/historical>).

Visibility and conventional meteorological data. The CMA established a national meteorological observation network in the 1950s, with the station number exceeding 2000 at the beginning and stabilizing at ~2,450 afterward. The observation network can continuously record meteorological data on visibility and conventional meteorological variables, including temperature, pressure, wind, and RH. In recent years, meteorological observations, including 6-hourly records between 1960-2020 and gradually increasing hourly records after 2013, have been collected from the National Meteorological Information Center (NMIC). Due to the inconsistency of visibility data in terms of observation methods, we conducted a series of data conversions to ensure continuous and consistent data. Visibility data recorded on a scale ranging from 0 to 9 between 1960-1979 were converted to numerical data based on probability density distributions. Specifically, the probability density distribution of visibility for each of the ten years before and after 1980 was calculated at first. The numerical visibility from 1980 to 1989 was graded into classes, with the median value of each class being the corresponding value for each station, and finally, the class observations were converted into numerical observations. From September 2013 to 2016, visibility measurements gradually shifted from 6-hourly manual observations to 1-hourly automatic observations site-by-site. In keeping with manual measurements, the automatic records, which are slighter lower than manual measurements, were calibrated by dividing 0.75 following the guideline from the CMA (Cma, 2014).

Emission inventories and elevation. Historical anthropogenic emissions from 1960-2012 are taken from Peking global emission inventories, developed using a bottom-up approach with spatial resolution at 0.1°×0.1° and temporal resolution at 1-month intervals (<http://inventory.pku.edu.cn>) (Chen et al., 2016a; Huang et al., 2014; Huang et al., 2015; Wang et al., 2014). Current

anthropogenic emissions during 2013-2020 are from the multiresolution Emission Inventory in China (MEIC, <http://meicmodel.org>) (Zhang et al., 2009; Zheng et al., 2018; Zheng et al., 2021). Six emission variables from these two inventories are used as inputs for model development, including PM_{2.5}, NO_x, SO₂, NH₃, BC, OC, and CO. Thirty-meter elevation data are collected from the Global Digital Elevation Model (GDEM) version 2 (<https://earthexplorer.usgs.gov>). Both emission and elevation data are interpolated from grids to sites to match existing PM_{2.5} sites.

Auxiliary data. Monthly Normalized Difference Vegetation Index (NDVI) products are downloaded from Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive Center (LADDS DAAC, <https://ladsweb.modaps.eosdis.nasa.gov>). Land cover classification data are taken from National Geographic Information Resources Catalogue Service System (<https://www.webmap.cn/mapDataAction.do?method=globalLandCover>). Population data are taken from the Gridded Population of the World version 4 (GPWv4, <https://sedac.ciesin.columbia.edu/data/collection/gpw-v4>) and are calibrated based on the total population in China City Yearbooks. NDVI, Land cover, and population data are also interpolated according to PM_{2.5} sites and trained as inputs for model development. However, during the model training process, we found that these data had little or no improvement in the hindcast capability of the model, and the time span of these data is insufficient for long-term historical retrieval. Hence, these auxiliary data are not used in model building.

2.2 Spatiotemporal feature extraction

For each PM_{2.5} site, we extract five variables as temporal inputs, including year, month, day, hour, and day of year. The longitude and latitude variables are taken out as location inputs (Fig. 1b). Visibility, RH, and temperature from the nearest meteorological station of each PM_{2.5} are used as basic meteorological inputs. The distance between these two sites was also added as a feature. In addition to the influence of the nearest meteorological station, PM_{2.5} concentrations at a site are also affected by surrounding conditions. For example, transport of pollution due to air movement is the main cause of heavy pollution episodes in the early stage (Zhong et al., 2017; Zhong et al., 2018). Hence, we need to consider spatial effects from surrounding meteorological stations. Our previous study developed a novel feature engineering approach, which incorporated surrounding impact by extracting spatial features (Zhong et al., 2021). Specifically, the remaining 19 nearest stations were matched for each PM_{2.5} site, except the nearest meteorological station. Five variables, including longitude, latitude, temperature, visibility, and RH, were selected from the 19 stations. Then, we calculated the maximum, the minimum, the average value, the skewness value, and the standard deviation for each of the five variables. These produced features, which take advantage of surrounding conditions, are also considered as inputs. After spatiotemporal feature extraction, a total of 71 features were used as inputs for model training. To reduce computation and training time with guaranteed accuracy, the top 40 features in order of importance during small-sample-testing processes are used for the following model training and hindcasting. These features included visibility, temporal features, spatial features, emission features, and elevation.

2.3 Gridded input construction

In the previous construction of input features for PM_{2.5} sites, we used location information,

time information, meteorological information from 20 surrounding meteorological stations, emission information, and elevation. If we assume that each cell in grid cells is a virtual PM_{2.5} site, then it is possible to generate input features for each grid point. After the model is trained based on input features and PM_{2.5} concentrations at real PM_{2.5} sites, we can feed the gridded input data into the model in turn and consequently construct a gridded PM_{2.5} network. Therefore, we define a grid area at 0.25°×0.25° with longitude from 70° E to 150° E and latitude from 10° N to 60° N and select the grid points covering mainland China. For each grid point, we performed spatiotemporal feature extraction and generated the same 71 input features as those of real PM_{2.5} sites.

2.4 Model description

LightGBM is one of the state-of-the-art gradient boosting frameworks with better accuracy, lower memory usage, faster training speed, and capability of handling large-scale data (Ke et al., 2017). Our previous research used this machine learning model to predict PM_{2.5} mass concentrations, which shows an unprecedented predictive capacity on hourly, daily, monthly, and annual timescales (Zhong et al., 2021). This study will continue to use this algorithm and previously tuned hyperparameters for model development (Zhong et al., 2021). For hindcasting historical PM_{2.5} datasets prior to 2013, a LightGBM model is trained and validated based on PM_{2.5} observations and feature inputs from 2013 to 2020. The hindcast capability is validated using cross-validation methods, which are standard methods for parameter tuning and model validation in machine learning. The training dataset is divided into several parts, one of them is used as test data, and the remaining parts are used as training data in turn. Each result yields a corresponding evaluation value, which is then averaged to provide an estimate of the model's accuracy. This estimation is quantified by two metrics: the coefficient of determination (R^2) and root-mean-square error (RMSE). The hindcast capability is also validated using PM_{2.5} observations from the US embassies in Beijing and Shanghai, which have been observing PM_{2.5} data since as early as 2008. After model training and validation, historical 6-hourly input data are inputted into this model to reconstruct a site-based PM_{2.5} dataset at 6-hour intervals from 1960 to 2020; and gridded input data are inputted into the model to reconstruct a 0.25°×0.25°, 6-hourly, gridded PM_{2.5} dataset. The daily, monthly, yearly, and decadal average PM_{2.5} concentrations for each site and each grid are also calculated based on the two datasets. Monthly-average values were obtained with daily values no less than 20 days; otherwise, they will be missing. Year-average values were calculated with 12 valid month values, and decadal-average values were calculated with 10 valid year-average values. The flowchart for reconstructing PM_{2.5} datasets is shown in Fig. 1.




Fig. 1 A conceptual scheme for constructing long-term historical site-based and gridded PM_{2.5} records based on long-term visibility, conventional meteorological observations, emissions, and elevation.

3 Results and Discussion

3.1 Evaluation of model hindcast performance

The hindcast performance of our model is evaluated using two CV methods, including 10-fold CV and by-year CV. The 10-fold CV partitions the original training datasets into 10 subsamples, one of which is retained as the validation data in turn for testing the model, and the remaining 9 subsamples are used as training data. This method is the most common CV that can be compared with results in other studies. However, 10-fold CV often overestimates the model's ability to hindcast continuous historical data. Therefore, we also use by-year CV, during which one year of data is selected sequentially for testing, and the remaining data are used for model training. This method is specifically designed to evaluate the hindcast capability of the model.

Tab. 1 Model performance in primary predictors, temporal resolution, and hindcast capability compared with other national PM_{2.5} datasets in China.

Related studies	Primary predictors	Temporal resolution	CV type	CV resolution	CV R ²	CV RMSE
Ma et al., 2016	AOD	daily (2004-2013)	10-fold CV	daily	0.79	27.40
			by-year CV		0.41	
Fang et al., 2016	AOD	daily (2013-2014)	10-fold CV	daily	0.80	22.80
Liu et al., 2017	Visibility	monthly (1957-1964, 1973-2014)	10-fold CV	monthly	0.71	25.62
Xiao et al., 2018	AOD	daily (2013-2017)	10-fold CV	daily	0.79	21.00
Xue et al., 2019	AOD, CTM outputs	daily (2000-2016)	by-year CV	daily	0.61	27.80
Liang et al., 2020	AOD	monthly (2000-2016)	10-fold CV	monthly	0.93	6.20
Huang et al., 2021	AOD, CTM outputs	daily (2013-2019)	10-fold CV	daily	0.87-0.88	11.90-21.90
			by-year CV		0.62	27.70
Wei et al., 2021	AOD	monthly (2000-2020)	10-fold CV	monthly	0.86-0.90	10.00-18.40
			by-year CV		0.80	11.26
van Donkelaar et al. 2021	AOD, CTM outputs	monthly (1998-2020)	Non-CV	yearly	0.69	11.90
Geng et al., 2021	AOD, CTM outputs	daily (2000-2020)	out-of-bag CV	daily	0.80-0.88	13.90-22.10
			by-year CV		0.58	27.50
Bai et al., 2022	AOD	daily (2000-2020)	10-fold CV	daily	0.79	20.04
Our study	Visibility	6-hourly (1960-2020)	10-fold CV	hourly/6-hourly	0.79	20.07
				6-hourly	0.78	21.14
				daily	0.85	16.11
				monthly	0.92	7.90
			by-year CV	hourly/6-hourly	0.70	26.36
				6-hourly	0.71	25.63
				daily	0.78	20.90
				monthly	0.83	13.37

Table 1 compares our dataset and the available datasets in primary predictors, temporal resolution, and CV results (Ma et al., 2016; Fang et al., 2016; Liu et al., 2017; Xiao et al., 2018; Xue et al., 2019; Liang et al., 2020; Huang et al., 2021; Wei et al., 2021a; Van Donkelaar et al., 2021; Geng et al., 2021; Bai et al., 2022). AOD-based datasets are only available from around 2000 at the earliest, with temporal resolutions ranging from daily scale to monthly scale. In contrast, our visibility-based dataset spans 61 years from 1960 to 2020 at 6-hourly intervals, showing a clear advantage in terms of time span and resolution. The R² and RMSE values of our 10-fold CV results are 0.78 and 21.14 $\mu\text{g m}^{-3}$ for 6-hourly estimations, respectively, which indicates our model is quite robust in estimating PM_{2.5}. Due to a reduction in data amount, the R² and RMSE values further improved to 0.85 and 16.11 $\mu\text{g m}^{-3}$ for daily estimations and 0.92 and 7.90 $\mu\text{g m}^{-3}$ for monthly estimations. This result is comparable to or even better than those of other available datasets whose 10-fold CV R² ranges from 0.61 to 0.80 on a daily scale and from 0.71 to 0.93 on a monthly scale. Our by-year CV's R² and RMSE values are 0.71 and 25.63 $\mu\text{g m}^{-3}$ for 6-hourly estimations, which indicates our model is still robust in hindcast performance. The by-year CV R² values for daily and monthly estimations (0.78 and 0.83) are higher than those in other available datasets (0.41-0.62 and

0.80), which might be partly attributed to spatial feature extraction and the large volume of our training dataset. Zhong et al. (2021) has shown that extracting spatial features can result in a better hindcast performance by fully representing dimensional heterogeneity. Compared to hundreds of thousands to millions of training samples in AOD-based models, the training samples for the visibility-based model are over 100 million. An increase in the order of magnitude for training datasets will yield better results in machine learning.




Fig. 2 Density scatterplots of observed $\text{PM}_{2.5}$ and estimated $\text{PM}_{2.5}$ across China for by-year CV from 2013 to 2020. The time resolution for CV results is hourly and 6-hourly between 2013-2017 and hourly between 2017-2020. (Colors are probability distribution densities).

The refined by-year CV results for each year between 2013-2020 are shown in Fig. 2. The by-year CV R^2 lies between 0.58 and 0.79, with better hindcast performance after 2014. The potential reasons why the R^2 value in 2013 is slightly lower than those in other years are as follows. First, the $\text{PM}_{2.5}$ observation network was just established in 2013, during which dehumidification systems, processing procedures, and data quality control methods are incomplete, and therefore the overall data quality cannot be guaranteed. With the improvement of the observation network after 2014, both the quality and quantity of observations increase significantly. This situation where data quality is relatively low initially but increases over time is also found in O_3 observations. Second, the CMA began to convert some of the manual visibility observations to automatic observations in 2013, during which there were also some irregular procedures in instrument equipment, observation steps, and data quality control. Lastly, although we have corrected the biases between manual and automated observations, some biases may still exist. However, the biases are further reduced as we integrate all manual visibility observations in 2013 into our training dataset.

The model's hindcast capability is further evaluated independently using pre-2013 $\text{PM}_{2.5}$ observations. For the $\text{PM}_{2.5}$ data currently available, only the US embassies in Beijing and Shanghai have at least one year's $\text{PM}_{2.5}$ observations. Therefore, $\text{PM}_{2.5}$ data from these two sites are applied as an independent evaluation dataset. Figure 3 shows our estimated $\text{PM}_{2.5}$ are in close agreement with in-situ measurements in Beijing and Shanghai, where the overall R^2 between observations and estimations is 0.74 and 0.79, respectively. For each year between 2008-2012 in Beijing, the R^2 values fluctuated between 0.70 and 0.81, reflecting a stable and accurate by-year hindcast capability. As shown in Fig. 3 (c-h), the low values, high values, and temporal variations in $\text{PM}_{2.5}$ measurements are all well estimated. In particular, $\text{PM}_{2.5}$ measurements are lacking at the US

Embassy in Beijing in early 2008 and around 2009, but our model can provide reasonable and continuous estimations to fill in the gaps. This ability can also be used to fill in missing PM_{2.5} observations of MEE from 2013 onwards, building a complete PM_{2.5} dataset. Overall, the independent validation results show that historical PM_{2.5} data can be well reconstructed by our model.




Fig. 3 (a) Density scatterplots of observed PM_{2.5} and estimated PM_{2.5} between 2008-2012 at the US Embassy in Beijing; (b) Density scatterplots of observed PM_{2.5} and estimated PM_{2.5} in 2012 at the US Embassy in Shanghai; (c-g) Timeseries of observed PM_{2.5} and estimated PM_{2.5} for each year between 2008-2012 at the US Embassy in Beijing; and (h) Timeseries of observed PM_{2.5} and estimated PM_{2.5} for each year in 2012 at the US Embassy in Shanghai.

The model's ability to make PM_{2.5} predictions at locations outside the scope of the training stations is evaluated by spatial CV. For spatial CV, all the monitoring stations are randomly divided into five subsets, and the model is trained using data from four subsets and tested on the data from the remaining subset each time. As shown in Fig. 4, the R^2 for spatial cross-validation in different groups is between 0.75 and 0.79, reflecting robust predictive power for PM_{2.5} concentrations at sites outside the training sites. Our previous study also examined this predictive ability using PM_{2.5} data from 23 untouched regional PM_{2.5} stations (Zhong et al., 2021).




Fig. 4 Density scatterplots of observed $\text{PM}_{2.5}$ and estimated $\text{PM}_{2.5}$ for each group of spatial CV results

3.2 Spatiotemporal variations in the site-based $\text{PM}_{2.5}$ dataset during 1960-2020

Figure 5 shows the spatiotemporal variations in annual average site-based $\text{PM}_{2.5}$ between 1960-2020. The trend of $\text{PM}_{2.5}$ in China experiences three major stages, corresponding to a slow increase under low concentrations between 1960-1978, a continuous accumulation with high concentrations reached between 1979-2013, and a rapid decrease between 2014-2020. During the first stage, though $\text{PM}_{2.5}$ pollution occurred in parts of the NCP and the Guanzhong Plain (GZP), $\text{PM}_{2.5}$ concentrations remain low in the vast majority of areas. This is mainly because anthropogenic emissions of $\text{PM}_{2.5}$ precursors and primary $\text{PM}_{2.5}$ grow slowly at a low base, resulting in relatively low total emissions in different regions. However, $\text{PM}_{2.5}$ pollution still occurring in the NCP and GZP, even with relatively low emissions, indicates the low environmental capacity of these two regions. During the second stage, $\text{PM}_{2.5}$ reached an unprecedentedly high concentration after a continuous increase in nearly all regions in China. The heaviest $\text{PM}_{2.5}$ pollution occurred in the NCP and the GZP. The SB and the Northeast China Plain (NeCP) are the polluted regions with the next highest $\text{PM}_{2.5}$ pollution. Even the YRD and the PRD also experienced $\text{PM}_{2.5}$ pollution during this stage. This worsening of $\text{PM}_{2.5}$ pollution is closely associated with massive anthropogenic emissions from rapidly increasing living and industrial activities after reform and opening-up policies. From 1979 to 2013, primary $\text{PM}_{2.5}$, NO_x , SO_2 , NH_3 , BC, OC, and CO from the Peking emission inventory increased by 98%, 457%, 159%, 117%, 45%, -22%, and 243%, respectively. Despite a slow reduction in SO_2 after 2006, the total anthropogenic emissions each year still increased and thereby caused high-level $\text{PM}_{2.5}$ pollution after 2006. The results indicate that air pollutants cannot be emitted without restraint, even in regions with high atmospheric capacity. Otherwise, $\text{PM}_{2.5}$ pollution will inevitably occur. In addition to anthropogenic emissions, sand and dust storms, resulting in high $\text{PM}_{2.5}$ concentrations in western Xinjiang, worsened $\text{PM}_{2.5}$ pollution by trans-regional transport from the desert regions. During the last stage, $\text{PM}_{2.5}$ decreased nationwide with the mass concentrations in nearly all stations approximately or below $35 \mu\text{g m}^{-3}$ in 2020, even in the NCP and the GZP with limited environmental

capacity. The substantial declines in PM_{2.5} illustrate the effectiveness of implementing the toughest-ever clean air policy in China. The spatiotemporal variations of PM_{2.5} between 1960-2020 clearly show the long-term impact of economic development and energy consumption on our air quality and the effectiveness of recent years' unprecedented emission control policies.




Fig. 5 Spatial distribution of annual average PM_{2.5} mass concentration at 1485 stations from 1960 to 2020

The specific turning points in annual PM_{2.5} concentrations for different regions were investigated additionally. Figure 6 shows the temporal variations in national-average monthly and yearly PM_{2.5} mass concentrations and regional average 6-hourly, monthly, and yearly PM_{2.5} mass concentrations in “2+26” cities of the NCP, the YRD, the PRD, and the SB. The national-average yearly PM_{2.5} reached a peak of 67 $\mu\text{g m}^{-3}$ in 2007, declined in 2008, and then remained steady until 2013. A sharp fall followed after 2014, with PM_{2.5} concentrations decreasing from 63 $\mu\text{g m}^{-3}$ in 2013 to 34 $\mu\text{g m}^{-3}$ in 2020. The annual PM_{2.5} concentrations in the “2+26” cities also experienced similar changes with a peak in 2007 and a reduction in 2008, which might be related to emission reduction for the Beijing Olympics in 2008. For the YRD, the maximum value of PM_{2.5} mass concentration occurred in 2013 without a striking peak in 2007. For the PRD, the annual PM_{2.5} concentrations increased steadily between 1960-1978, then rose more and more steeply in the following years with a steep increase in 2003 and 2004 and peaked in 2004. A steady decrease with slight fluctuation occurred from 2005 to 2013, and then a sharp fall followed after 2014. This trend is different from that in the “2+26” cities and the YRD. For the SB, the turning point occurred in 2013, before which

the annual $\text{PM}_{2.5}$ concentrations increased steadily and remained steady.




Fig. 6 (a) Spatial distribution of average $\text{PM}_{2.5}$ mass concentrations between 1960-2020; (b-f) Timeseries of average $\text{PM}_{2.5}$ mass concentrations for all sites in China (b), “2+26” cities (c), Yangtze River Delta (d), Pearl River Delta (e) and Sichuan Basin (f), respectively.

3.3 Detailed spatial distributions from gridded $\text{PM}_{2.5}$ datasets

Figure 7 shows the annual spatial variations in $0.25^\circ \times 0.25^\circ$ gridded $\text{PM}_{2.5}$ between 1960-2020. Compared to site-based distributions, gridded $\text{PM}_{2.5}$ can portray the spatiotemporal variations in a clearer and more detailed way. For example, the most widespread and heaviest $\text{PM}_{2.5}$ pollution in western Xinjiang occurred in 1979. This abnormal pollution corresponds to the historical construction of northern severe dust storms, which recorded the event with the largest affected areas in April 1979 (Zhou and Zhang, 2003). As exposed to nearly the most frequent air stagnation in winter due to terrain and meteorological conditions (Wang et al., 2018), the NCP is the region with $\text{PM}_{2.5}$ pollution first to appear and last to disappear except areas affected by dust storms (Fig. 7). For year-to-year comparisons, it can be clearly seen that $\text{PM}_{2.5}$ concentrations in the NCP decreased slightly from 2007 to 2008 and from 2012 to 2013, respectively, and decreased significantly in 2014 relative to 2013. The $\text{PM}_{2.5}$ reduction is insignificant from 2015 to 2016 but striking from 2016 to 2017. In 2020, the nationwide $\text{PM}_{2.5}$ concentrations are comparable to those in 1960s and close to the lowest level ever recorded in almost 61 years.




Fig. 7 Gridded distribution of annual average $PM_{2.5}$ mass concentration from 1960 to 2020

Figure 8 shows inter-decadal spatial variations in gridded $PM_{2.5}$ between 1961–2020. $PM_{2.5}$ concentrations maintained at low levels in most areas over the first decade and increased to a certain extent in the NCP and western Xinjiang over the second decade. In the following decades, $PM_{2.5}$ pollution has worsened significantly in several key regions, including the NCP, the GZP, and the SB. This worsening was maintained until the last decade, during which $PM_{2.5}$ pollution mitigates significantly in nearly all populous and polluted regions in eastern China.




Fig. 8 Gridded distribution of decadal average $PM_{2.5}$ mass concentration from 1960 to 2020

The multi-year trend of our gridded PM_{2.5} dataset is also compared with those of publicly available datasets, including the TAP data (Geng et al., 2021), the GEFPM data (Van Donkelaar et al., 2021), the LGHAP data (Bai et al., 2022), and the CHAP data (Wei et al., 2021a), which have been interpolated to the same grid resolution. Figure 9 shows the spatial distributions of PM_{2.5} from those datasets at 5-year intervals between 2000-2020. One consistent trend across all datasets was that nationwide PM_{2.5} mass concentrations experienced an increase following a decrease from 2000 to 2020. However, the turning points are different for different datasets. From 2010 to 2015, PM_{2.5} pollution alleviated for TAP, CHAP, and our data but worsened for GEFPM and LGHAP. For the time (2015 and 2020) with ground observations available, all PM_{2.5} data show similar spatial distributions with the most severe pollution in the NCP in 2015 and significant improvement in nationwide air pollution in 2020. For the years (2000, 2005, and 2010) when ground observations were unavailable, significant disparities in pollution levels and regional distribution emerged from different datasets. Specifically, the LGHAP data are significantly lower than other data, while the TAP data are higher than others in nearly all regions except western Xinjiang. In western Xinjiang, PM_{2.5} concentrations from the GEFPM data are the highest among all the datasets. Due to a lack of ground PM_{2.5} observations before 2000, it is challenging to determine which dataset has the least bias and more reasonable distributions. In the future, applying ensemble average to multi-datasets might be an effective way to eliminate systematic bias.




Fig. 9 Distribution of reconstructed PM_{2.5} by different PM_{2.5} datasets in 2000, 2005, 2010, 2015, and 2020. From top to down are TAP, GEFPM, LGHAP, CHAP, and our dataset.

4 Data availability

The 6-hourly PM_{2.5} datasets from 1960 to 2020, including site-based and gridded data, are publicly accessible. Daily, monthly, and yearly sited-based and gridded PM_{2.5} datasets are also provided. The sited-based PM_{2.5} dataset is in the CSV format, and the gridded dataset PM_{2.5} is in the NETCDF format. All of them are available at <https://doi.org/10.5281/zenodo.6372847> (Zhong et al., 2022).

5 Conclusion

This study is among the first to generate long-term site-based and gridded PM_{2.5} datasets between 1960-2020 with 6-hourly resolution, based on long-term visibility, conventional meteorological observations, emissions, and elevation. A new feature engineering method that takes advantage of spatial features from 20 surrounding meteorological stations is employed in our LightGBM model to incorporate spatial effects of meteorological conditions. For by-year CV, the R² values of our model are 0.71, 0.78, and 0.83 for 6-hourly, daily, and monthly estimations, respectively, which are higher than those in other available datasets (0.41-0.62). This hindcast capability is further evaluated independently using pre-2013 PM_{2.5} data of 6 years from US embassies in Beijing and Shanghai. The low values, high values, and temporal variations in US-embassy PM_{2.5} measurements are all well estimated with the overall R² being 0.74 and 0.79 in Beijing and Shanghai, respectively. Both by-year CV and independent validation show that our model has a stable by-year hindcast capability and can reconstruct historical PM_{2.5} data in a relatively accurate way. Our datasets show that PM_{2.5} variations in China experience a slow increase under low concentrations between 1960-1978, a continuous accumulation with high concentrations reached between 1979-2013, and a rapid decrease between 2014-2020. The worsening of PM_{2.5} pollution is closely associated with massive anthropogenic emissions after reform and opening-up policies, while the substantial declines in PM_{2.5} are mainly due to the implementation of the toughest-ever clean air policy in China. In 2020, the nationwide PM_{2.5} concentrations were close to the lowest recorded level in almost 61 years. These two reconstructed PM_{2.5} datasets provide spatiotemporal variations at high resolution, which lay the foundation for research studies associated with air pollution, climate change, and atmospheric chemical reanalysis. It is worth noting that our datasets still have some weaknesses, with the main weakness being a lack of detailed bias estimations for each value in our datasets due to limited historical observations. In the future, we will collect as many PM_{2.5} observations as possible to validate the accuracy of our datasets and provide evaluations of uncertainty for our datasets.

Acknowledgements

The authors are grateful to all the organization and groups that provided indispensable datasets that we used in this study. We would like to acknowledge to CMA for providing the long-term visibility and MEE for the observational PM_{2.5} data. Also, we acknowledge the MEIC team and Peking for providing the emission inventories and GDEM data provider.

Financial support

This research was supported by the Major Project from Natural Science Foundation of China (42090030) and the Distinguished Young Scholars Project from Natural Science Foundation of China (41825011).

Author Contributions

XZ designed the research and led the overall scientific questions. JZ, KG, and LG carried out data processing and analysis based on suggestions from ZZ, DW, YW, and HC. LL calibrated PM_{2.5} data and meteorological observations. YF performed visibility conversions from class values to numeric values before 1980. JL and LJ provided calibrated visibility from 2013 to 2016 and suggestions about manuscript structures. JZ wrote the first draft of the manuscript, and XZ revised the manuscript. All authors read and approved the final version.

Competing financial interests

The authors declare no competing financial interests.

References

- Bai, K., Li, K., Ma, M., Li, K., Li, Z., Guo, J., Chang, N. B., Tan, Z., and Han, D.: LGHAP: the Long-term Gap-free High-resolution Air Pollutant concentration dataset, derived via tensor-flow-based multimodal data fusion, *Earth Syst. Sci. Data*, 14, 907-927, 10.5194/essd-14-907-2022, 2022.
- Beelen, R., Hoek, G., Den Brandt, P. A. V., Goldbohm, R. A., Fischer, P., Schouten, L. J., Jerrett, M., Hughes, E., Armstrong, B., and Brunekreef, B.: Long-term effects of traffic-related air pollution on mortality in a Dutch cohort (NLCS-AIR study), *Environmental Health Perspectives*, 116, 196-202, 2007.
- Chen, H., Huang, Y., Shen, H., Chen, Y., Ru, M., Chen, Y., Lin, N., Su, S., Zhuo, S., Zhong, Q., Wang, X., Liu, J., Li, B., and Tao, S.: Modeling temporal variations in global residential energy consumption and pollutant emissions, *Applied Energy*, 184, 820-829, <https://doi.org/10.1016/j.apenergy.2015.10.185>, 2016a.
- Chen, J., Yin, J., Zang, L., Zhang, T., and Zhao, M.: Stacking machine learning model for estimating hourly PM_{2.5} in China based on Himawari 8 aerosol optical depth data, *Science of The Total Environment*, 697, 134021, <https://doi.org/10.1016/j.scitotenv.2019.134021>, 2019.
- Chen, X., Zhang, L. W., Huang, J. J., Song, F. J., Zhang, L. P., Qian, Z. M., Trevathan, E., Mao, H. J., Han, B., Vaughn, M., Chen, K. X., Liu, Y. M., Chen, J., Zhao, B. X., Jiang, G. H., Gu, Q., Bai, Z. P., Dong, G. H., and Tang, N. J.: Long-term exposure to urban air pollution and lung cancer mortality: A 12-year cohort study in Northern China, *Science of the Total Environment*, 571, 855-861, 10.1016/j.scitotenv.2016.07.064, 2016b.
- CMA: Forecasting and Networking Department of China Meteorological Administration released letter No.4: Notice on the adjustments of the haze weather phenomenon observation and on the revision of the fog and haze observation data, 2014.
- Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmospheric Chemistry and Physics*, 11, 11069-11102, 10.5194/acp-11-11069-2011, 2011.
- Fang, X., Zou, B., Liu, X., Sternberg, T., and Zhai, L.: Satellite-based ground PM_{2.5} estimation using timely structure adaptive modeling, *Remote Sensing of Environment*, 186, 152-163, <https://doi.org/10.1016/j.rse.2016.08.027>, 2016.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., and Reichle, R.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *Journal of Climate*, 30, 5419-5454, 2017.
- Geng, G., Xiao, Q., Liu, S., Liu, X., Cheng, J., Zheng, Y., Xue, T., Tong, D., Zheng, B., Peng, Y., Huang, X., He, K., and Zhang, Q.: Tracking Air Pollution in China: Near Real-Time PM_{2.5} Retrievals from Multisource Data Fusion, *Environmental Science & Technology*, 55, 12106-12115, 10.1021/acs.est.1c01863, 2021.
- Gui, K., Che, H., Zeng, Z., Wang, Y., Zhai, S., Wang, Z., Luo, M., Zhang, L., Liao, T., Zhao, H., Li, L., Zheng, Y., and Zhang, X.: Construction of a virtual PM_{2.5} observation network in China based on high-density surface meteorological observations using the Extreme Gradient Boosting model, *Environment International*, 141, 105801, <https://doi.org/10.1016/j.envint.2020.105801>, 2020.
- Huang, C., Hu, J., Xue, T., Xu, H., and Wang, M.: High-Resolution Spatiotemporal Modeling for Ambient PM_{2.5} Exposure Assessment in China from 2013 to 2019, *Environmental Science & Technology*, 55, 2152-2162, 10.1021/acs.est.0c05815, 2021.
- Huang, Y., Shen, H., Chen, Y., Zhong, Q., Chen, H., Wang, R., Shen, G., Liu, J., Li, B., and Tao, S.:

Global organic carbon emissions from primary sources from 1960 to 2009, *Atmospheric Environment*, 122, 505-512, <https://doi.org/10.1016/j.atmosenv.2015.10.017>, 2015.

Huang, Y., Shen, H., Chen, H., Wang, R., Zhang, Y., Su, S., Chen, Y., Lin, N., Zhuo, S., Zhong, Q., Wang, X., Liu, J., Li, B., Liu, W., and Tao, S.: Quantification of Global Primary Emissions of PM_{2.5}, PM₁₀, and TSP from Combustion and Industrial Process Sources, *Environmental Science & Technology*, 48, 13834-13843, 10.1021/es503696k, 2014.

Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A. M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V. H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19, 3515-3556, 10.5194/acp-19-3515-2019, 2019.

Ke, G., Meng, Q., Finley, T., Wang, T., Chen, W., Ma, W., Ye, Q., and Liu, T.-Y.: Lightgbm: A highly efficient gradient boosting decision tree, *Advances in neural information processing systems*, 3146-3154, 2018.

Kuang, Y., Zhao, C., Tao, J., Bian, Y., and Ma, N. J. A. E.: Impact of aerosol hygroscopic growth on the direct aerosol radiative effect in summer on North China Plain, 147, 224-233, 10.1016/j.atmosenv.2016.10.013, 2016.

Li, X., Gao, C. Y., Gao, Z., and Zhang, X.: Atmospheric boundary layer turbulence structure for severe foggy haze episodes in north China in December 2016, *Environmental Pollution*, 264, 114726, <https://doi.org/10.1016/j.envpol.2020.114726>, 2020.

Liang, F., Xiao, Q., Huang, K., Yang, X., Liu, F., Li, J., Lu, X., Liu, Y., and Gu, D.: The 17-y spatiotemporal trend of PM_{2.5} and its mortality burden in China, *Proceedings of the National Academy of Sciences*, 117, 25601-25608, 10.1073/pnas.1919641117, 2020.

Liu, M., Bi, J., and Ma, Z.: Visibility-Based PM_{2.5} Concentrations in China: 1957–1964 and 1973–2014, *Environmental Science & Technology*, 51, 13161-13169, 10.1021/acs.est.7b03468, 2017.

Lynch, P., Reid, J. S., Westphal, D. L., Zhang, J., Hogan, T. F., Hyer, E. J., Curtis, C. A., Hegg, D. A., Shi, Y., and Campbell, J. R.: An 11-year global gridded aerosol optical thickness reanalysis (v1. 0) for atmospheric and climate sciences, *Geoscientific Model Development*, 9, 1489, 2016.

Ma, Z., Hu, X., Sayer, A. M., Levy, R., Zhang, Q., Xue, Y., Tong, S., Bi, J., Huang, L., and Liu, Y.: Satellite-Based Spatiotemporal Trends in PM_{2.5} Concentrations: China, 2004-2013, *Environmental Health Perspectives*, 124, 184-192, doi:10.1289/ehp.1409481, 2016.

Pilinis, C., Seinfeld, J. H., and Grosjean, D.: Water content of atmospheric aerosols, *Atmospheric Environment*, 23, 1601-1606, 10.1016/0004-6981(89)90419-8, 1989.

Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *Journal of the American Medical Association*, 287, 1132-1141, 2002.

Randles, C., Da Silva, A., Buchard, V., Colarco, P., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., and Hair, J.: The MERRA-2 aerosol reanalysis, 1980 onward. Part I: System description and data assimilation evaluation, *Journal of Climate*, 30, 6823-6850, 2017.

van Donkelaar, A., Hammer, M. S., Bindle, L., Brauer, M., Brook, J. R., Garay, M. J., Hsu, N. C., Kalashnikova, O. V., Kahn, R. A., Lee, C., Levy, R. C., Lyapustin, A., Sayer, A. M., and Martin, R. V.: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environmental Science & Technology*, 10.1021/acs.est.1c05309, 2021.

Wang, B., Yuan, Q., Yang, Q., Zhu, L., Li, T., and Zhang, L.: Estimate hourly PM_{2.5} concentrations from Himawari-8 TOA reflectance directly using geo-intelligent long short-term memory network, *Environmental Pollution*, 271, 116327, <https://doi.org/10.1016/j.envpol.2020.116327>, 2021.

Wang, R., Tao, S., Shen, H., Huang, Y., Chen, H., Balkanski, Y., Boucher, O., Ciais, P., Shen, G., Li, W., Zhang, Y., Chen, Y., Lin, N., Su, S., Li, B., Liu, J., and Liu, W.: Trend in Global Black Carbon Emissions from 1960 to 2007, *Environmental Science & Technology*, 48, 6780-6787, 10.1021/es5021422, 2014.

Wang, X., Zhang, R., and Yu, W.: The Effects of PM_{2.5} Concentrations and Relative Humidity on Atmospheric Visibility in Beijing, *Journal of Geophysical Research: Atmospheres*, 124, 2235-2259, <https://doi.org/10.1029/2018JD029269>, 2019.

Wang, X., Dickinson, R. E., Su, L., Zhou, C., and Wang, K.: PM 2.5 Pollution in China and How It Has Been Exacerbated by Terrain and Meteorological Conditions, *Bulletin of the American Meteorological Society*, 99, 105-119, 10.1175/bams-d-16-0301.1, 2018.

Wei, J., Li, Z., Lyapustin, A., Sun, L., Peng, Y., Xue, W., Su, T., and Cribb, M.: Reconstructing 1-km-resolution high-quality PM_{2.5} data records from 2000 to 2018 in China: spatiotemporal variations and policy implications, *Remote Sensing of Environment*, 252, 112136, <https://doi.org/10.1016/j.rse.2020.112136>, 2021a.

Wei, J., Li, Z., Pinker, R. T., Wang, J., Sun, L., Xue, W., Li, R., and Cribb, M.: Himawari-8-derived diurnal variations in ground-level PM_{2.5} pollution across China using the fast space-time Light Gradient Boosting Machine (LightGBM), *Atmos. Chem. Phys.*, 21, 7863-7880, 10.5194/acp-21-7863-2021, 2021b.

Xiao, Q., Chang, H. H., Geng, G., and Liu, Y.: An Ensemble Machine-Learning Model To Predict Historical PM_{2.5} Concentrations in China from Satellite Data, *Environmental Science & Technology*, 52, 13260-13269, 10.1021/acs.est.8b02917, 2018.

Xue, T., Zheng, Y., Tong, D., Zheng, B., Li, X., Zhu, T., and Zhang, Q.: Spatiotemporal continuous estimates of PM_{2.5} concentrations in China, 2000–2016: A machine learning method with inputs from satellites, chemical transport model, and ground observations, *Environment International*, 123, 345-357, <https://doi.org/10.1016/j.envint.2018.11.075>, 2019.

Yan, X., Zang, Z., Luo, N., Jiang, Y., and Li, Z.: New interpretable deep learning model to monitor real-time PM_{2.5} concentrations from satellite data, *Environment International*, 144, 106060, <https://doi.org/10.1016/j.envint.2020.106060>, 2020.

Yuan, R., Zhang, X., Liu, H., Gui, Y., Shao, B., Tao, X., Wang, Y., Zhong, J., Li, Y., and Gao, Z.: Aerosol vertical mass flux measurements during heavy aerosol pollution episodes at a rural site and an urban site in the Beijing area of the North China Plain, *Atmos. Chem. Phys.*, 19, 12857-12874, 10.5194/acp-19-12857-2019, 2019.

Yumimoto, K., Tanaka, T. Y., Oshima, N., and Maki, T.: JRAero: the Japanese reanalysis for aerosol v1.0, *Geoscientific Model Development*, 10, 3225, 2017.

Zhang, H. L., Wang, Y. G., Hu, J. L., Ying, Q., and Hu, X. M.: Relationships between meteorological parameters and criteria air pollutants in three megacities in China, *Environmental Research*, 140, 242-254, 10.1016/j.envres.2015.04.004, 2015.

Zhang, Q., Streets, D. G., Carmichael, G. R., He, K., Huo, H., Kannari, A., Klimont, Z., Park, I., Reddy, S., and Fu, J.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmospheric Chemistry and Physics*, 9, 5131-5153, 2009.

Zhang, R., Li, Q., and Zhang, R.: Meteorological conditions for the persistent severe fog and haze event over eastern China in January 2013, *Science China Earth Sciences*, 57, 26-35, 10.1007/s11430-013-4774-3, 2013a.

Zhang, X., Sun, J., Wang, Y., Li, W., Zhang, Q., Wang, W., Quan, J., Cao, G., Wang, J., Yang, Y., and Zhang, Y.: Factors contributing to haze and fog in China, *Chinese Science Bulletin*, 58, 1178,

10.1360/972013-150, 2013b.

Zheng, B., Cheng, J., Geng, G., Wang, X., Li, M., Shi, Q., Qi, J., Lei, Y., Zhang, Q., and He, K.: Mapping anthropogenic emissions in China at 1 km spatial resolution and its application in air quality modeling, *Science Bulletin*, 66, 612-620, <https://doi.org/10.1016/j.scib.2020.12.008>, 2021.

Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, 10.5194/acp-18-14095-2018, 2018.

Zhong, J., Zhang, X., Dong, Y., Wang, Y., Liu, C., Wang, J., Zhang, Y., and Che, H.: Feedback effects of boundary-layer meteorological factors on cumulative explosive growth of PM_{2.5} during winter heavy pollution episodes in Beijing from 2013 to 2016, *Atmos. Chem. Phys.*, 18, 247-258, 10.5194/acp-18-247-2018, 2018.

Zhong, J., Zhang, X., Gui, K., Wang, Y., Che, H., Shen, X., Zhang, L., Zhang, Y., Sun, J., and Zhang, W.: Robust prediction of hourly PM_{2.5} from meteorological data using LightGBM, *National Science Review*, 8, 10.1093/nsr/nwaa307, 2021.

Zhong, J., Zhang, X., Wang, Y., Sun, J., Zhang, Y., Wang, J., Tan, K., Shen, X., Che, H., and Zhang, L.: Relative contributions of boundary-layer meteorological factors to the explosive growth of PM_{2.5} during the red-alert heavy pollution episodes in Beijing in December 2016, *Journal of Meteorological Research*, 31, 809-819, 10.1007/s13351-017-7088-0, 2017.

Zhong, J., Zhang, X., Gui, K., Liao, J., Fei, Y., Jiang, L., Guo, L., Liu, L., Che, H., Wang, Y., Wang, D., and Zhou, Z.: Reconstructing 6-hourly PM_{2.5} datasets from 1960 to 2020 in China [dataset], 10.5281/zenodo.6372847, 2022.

Zhou, Z. and Zhang, G.: Typical severe dust storms in northern China during 1954—2002, *Chinese Science Bulletin*, 48, 2366-2370, 2003.

Zhu, W., Xu, X., Zheng, J., Yan, P., Wang, Y., and Cai, W.: The characteristics of abnormal wintertime pollution events in the Jing-Jin-Ji region and its relationships with meteorological factors, *Science of the Total Environment*, 626, 887-898, 2018.