

HMRFS-TP: long-term daily gap-free snow cover products over the Tibetan Plateau from 2002 to 2021 based on Hidden Markov Random Field model

5 Yan Huang^{1,2+*}, Jiahui Xu^{1,2+}, Jingyi Xu^{1,2}, Yelei Zhao^{1,2}, Bailang Yu^{1,2}, Hongxing Liu³, Shujie Wang⁴, Wanjia Xu^{1,2},
Jianping Wu^{1,2}, Zhaojun Zheng⁵

¹Key Laboratory of Geographic Information Science, Ministry of Education, East China Normal University, Shanghai 200241, China

²School of Geographic Sciences, East China Normal University, Shanghai 200241, China

³Department of Geography, the University of Alabama, Tuscaloosa, AL 35487, USA

10 ⁴Department of Geography, Earth and Environmental Systems Institute, Pennsylvania State University, University Park, PA 16802, USA

⁵National Satellite Meteorological Center, Beijing 100081, China

⁺These authors contributed equally to this work.

Correspondence to: Yan Huang (yhuang@geo.ecnu.edu.cn)

15 **Abstract.** Snow cover plays an essential role in climate change and the hydrological cycle of the Tibetan Plateau. The widely used Moderate Resolution Imaging Spectroradiometer (MODIS) snow products have two major issues: massive data gaps due to frequent clouds and relatively low estimate accuracy of snow cover due to complex terrain in this region. Here we generate long-term daily gap-free snow cover products over the Tibetan Plateau at 500 m resolution by applying a Hidden Markov Random Field (HMRF) technique to the original MODIS snow products over the past two decades. The data

20 gaps of the original MODIS snow products were fully filled by optimally integrating spectral, spatiotemporal, and environmental information within HMRF framework. The snow cover estimate accuracy was greatly increased by incorporating the spatiotemporal variations of solar radiation due to surface topography and sun elevation angle as the environmental contextual information in HMRF-based snow cover estimation. We evaluated our snow products, and the accuracy is ~~98.29%~~~~98.31%~~ in comparison with *in situ* observations and ~~91.36%~~~~92.44%~~ in comparison with high-resolution

25 snow maps derived from ~~Landsat 8 imagery~~~~Landsat images~~. Our evaluation also suggests that the incorporation of spatiotemporal solar radiation as the environmental contextual information in HMRF modelling, instead of the simple use of surface elevation as the environmental contextual information, results in the accuracy of the snow products increase by ~~2.71%~~~~2.72%~~ and the omission error decrease by ~~-3.59%~~~~4.59%~~. The accuracy of our snow products is especially improved during snow transitional period and over complex terrains with high elevation as well as sunny slopes. The new products can

30 provide long-term and spatiotemporally continuous information of snow cover distribution, which is critical for understanding the processes of snow accumulation and melting, analysing its impact on climate change, and facilitating water resource management in Tibetan Plateau. This dataset can be freely accessed from the National Tibetan Plateau Data Center at <https://doi.org/10.11888/Cryos.tpdc.272204> (Huang and Xu, 2022).

1 Introduction

35 Snow cover has the characteristics of low thermal diffusivity, high reflectivity, and strong water storage capacity, which has a profound effect on climate change (Gao et al., 2012), radiation budget (Yang et al., 2014; Huang et al., 2019), hydrological cycle (Dong, 2018), and human activities (Cereceda-Balic et al., 2020). The Tibetan Plateau (TP) has abundant snow cover, with highest elevation and largest snow cover area in the middle latitudes of Northern Hemisphere (Qiu, 2008; Yao et al., 2019). Snow is highly sensitive to climate change (Chen et al., 2018a), and snowmelt water quantity is closely connected to
40 the supply of soil moisture on the plateau (Wang et al., 2018) and the runoff of numerous rivers such as the Yangtze and Yellow rivers (Immerzeel et al., 2010). Long-term and detailed snow cover information is fundamental to investigating climate change and hydrological cycle of TP.

Remote sensing has allowed extraction of historical and near-real-time snow cover extent over inaccessible areas (Yang et al., 2015; Li et al., 2018). Snow products could be acquired via remote sensing satellites, such as the Landsat and Sentinel-2
45 series. Although these satellites have a high spatial resolution, they have a relatively coarse (10 or 16 days) temporal resolution, thereby rendering them insufficient to monitor the temporal variations in snow cover (Huang et al., 2022). Moderate Resolution Imaging Spectroradiometer (MODIS) has commenced producing snow products at 500 m resolution from 2000, which have been widely utilized as the primary datasets for monitoring snow cover (Muhammad and Thapa, 2020; Kilpys et al., 2020). The accuracy of MODIS snow products is greater than 85% at the global scale under clear sky
50 (e.g., Parajka et al., 2012; Yang et al., 2015). However, these products have many data gaps due to frequent clouds, causing discontinuity in the time and space of the products (Liu et al., 2020b). In addition, the complex terrain of TP makes it more challenging for accurate snow cover detection. Previous studies (Dong and Menzel, 2016; Dariane et al., 2017) have suggested the low accuracy of MODIS snow products for mountainous areas.

Various data-gap-filling techniques have been proposed to produce seamless MODIS snow cover products including multi-
55 source combination, temporal, spatial, and spatiotemporal filters (Xiao et al., 2021; Hussainzada et al., 2021; Richiardi et al., 2021). The multi-source combination method combines MODIS with passive microwave sensor data (Li et al., 2019b; Li et al., 2020). Although this method can be used to fill most data gaps, the accuracy of the combined product depends more on the accuracy of the microwave sensor data, and the spatial resolution (>1 km) is not sufficient to meet the demands for accurately assessing the snow cover variability (Wang et al., 2009). Many attempts have been directed toward filling the
60 gaps in MODIS optical remote sensing data based on spatial and temporal information. Temporal methods are used to reclassify the data-gap pixels by inferring the land cover types of the current pixels under clear sky within a few days before or after (e.g., the previous or next day) (Li et al., 2019a; Tran et al., 2019). Spatial methods estimate data-gap pixels based on gap-free pixels in the spatial neighborhood (Hou et al., 2019). Relevant environmental information (e.g., snow lines and topography) has been introduced into spatial methods (Wang et al., 2019; Kilpys et al., 2020). Spatiotemporal methods have
65 been integrated to fill as many data gaps as possible (Parajka and Blöschl, 2008; Kilpys et al., 2020). These studies integrated all information (i.e., spatiotemporal, and environmental information) according to heuristic rules instead of

rigorous quantitative model. Huang et al. (2018) developed a Hidden Markov Random Field (HMRF) framework to optimally combine all information. This technique not only fills data gaps but also provides fine improvement in accuracy compared to the original MODIS snow products.

70 The long-term series and high-precision products are the basis for snow cover research on the TP, which enable the monitoring and analysis of snow cover phenology and more comprehensive understanding of the snow cover trend. However, the existing daily products for the TP have an earlier end date (the latest version at 500 m resolution product ended in 2015), and some products still have a small number of data gaps ([Yu et al., 2016](#); [Qiu et al., 2017](#); [Xu et al., 2017](#); [Zheng and Cao, 2019](#); [Tang et al., 2013](#); [Yu et al., 2016](#); [Qiu et al., 2017](#); [Zheng and Cao, 2019](#)). Thus, long-term and high-precision daily

75 snow cover products for the TP should be generated using reliable data-gap-filling methods.

Here we generate long-term daily snow cover products over the TP by applying a ~~Hidden Markov Random Field (HMRF)~~ technique (Huang et al., 2018) to the original MODIS snow products from 2002 to 2021. In the previous HMRF modelling technique (Huang et al., 2018), surface elevation was considered as a surrogate for environmental information in mountainous regions. However, according to *in situ* photos from field survey in the TP, we found that the distribution of

80 snow cover differed from the model results, even at the same elevation level (Figs. 1b and 1c). The snow on the TP is strongly infected by complex topography (e.g., slope, aspect, sunlight duration, and solar incidence angle). Thus, while generating our new snow cover product, we incorporated solar radiation as a comprehensive indicator of topographical effects to correct the snow identification errors caused by the complex terrain of the TP.

Our study is outlined as follows: first, we present our study area and datasets. Then, we present the HMRF framework and

85 data-processing flowchart. By employing this modelling technique to the TP, we produced daily gap-free snow cover products for the TP from 2002 to 2021. We also validated our new snow products against *in situ* observations, snow cover mapped from ~~Landsat 8 imagery~~[Landsat series data](#), and snow cover data estimated from the initial MODIS and original HMRF modelling technique.

2 Study area and data


90 2.1 Study area

The TP is situated in western China, spanning 26°00'-39°46'N and 73°18'-104°46'E (Fig. 1). It encompasses approximately ~~2.56 × 10⁶~~[2.56 × 10⁶](#) km², with an average altitude exceeding 4000 m, and is one of the most susceptible regions to climate change (Jing et al., 2019). The air temperature in the TP increased from the northwest to southeast, with more precipitation in the southeast and less in the northwest (Wang et al., 2018). Generally, the TP has comparatively low temperatures and has

95 largest distribution of glaciers and snow in China (Liang et al., 2017). [Seasonal snow cover on the TP has a great potential to influence the hydrological cycle and heat wave frequency in northern China \(Wu et al., 2012\). In addition, seasonal snow accumulation on the TP is an important part of surface water accumulation in southwestern China and surrounding countries.](#)

Several major rivers in China and surrounding Asian countries, such as the Yangtze, Yellow, Mekong, Salween, Brahmaputra, Ganges, and Indus Rivers, all originate from the TP.

100 The status of the snow in the TP changes rapidly as a result of multiple factors, such as temperature, precipitation, synoptic forcing, and large-scale ocean-atmosphere oscillations (You et al., 2020), which may lead to sublimation and snowfall (Li et al., 2020). Affected by the Indian Ocean monsoon and East Asian summer monsoon, the southeast TP has plenty water vapor, which results in a large amount of cloud, particularly in spring and summer (Yang et al., 2015).






Figure 1: Topography, meteorological stations, and survey photos of the TP. (a) Surface elevation and distribution of meteorological stations in the TP. Landsat 8 imageryLandsat images, utilized for validation and sample area are also shown. (b) and (c) *in situ* photos from field survey in the TP.

2.2 Datasets

110 2.2.1 Daily MODIS snow cover products

We used version 6 MODIS daily snow products on Terra (MOD10A1 v6) and Aqua (MYD10A1 v6) from May 15, 2002, to December 31, 2021 (Hall and Riggs, 2016a; Hall and Riggs, 2016b). Google Earth Engine (GEE) (<https://earthengine.google.com>) platform was used for pre-processing the MODIS snow product. We used MODIS/006/MOD10A1 and MODIS/006/MYD10A1 datasets and NDSI_Snow_Cover and NDSI_Snow_Cover_Class bands. NDSI_Snow_Cover represents the value of the Normalized Difference Snow Index (NDSI), ranging 0–100. The

115

values in the NDSI_Snow_Cover_Class band include 200, 201, 211, 237, 239, 250, 254, and 255, which represent "missing data", "no decision", "night time", "inland water", "ocean", "clouds", "detector saturated", and "filled data", respectively (Riggs et al., 2019).

The original NDSI data were reclassified as snow, non-snow, and data-gaps classes (Huang et al., 2018). The pixels with an NDSI value of 40–100 in the NDSI_Snow_Cover band were reclassified as snow (Riggs et al., 2017). The pixels with an NDSI value of 0–40 in the NDSI_Snow_Cover band and with corresponding values of 211, 237, and 239 in the NDSI_Snow_Cover_Class band were reclassified as non-snow. The remaining pixels were reclassified as data-gaps. Then, we combined the MOD10A1 and MYD10A1 reclassification results of the same day according to the following rules: when a pixel included gap-free data in both MOD10A1 and MYD10A1 products, value in the MOD10A1 product was used; when a pixel included gap-free only in the one product, the gap-free data value was used. Finally, the combined Terra and Aqua results were re-projected onto Universal Transverse Mercator (UTM zone 45) at 500 m resolution, which was used as the initial snow cover products for filling data gaps in this research.

We identified the data gaps in original composite MODIS products, and found the annual average data gap proportion of the TP was 33.63%–38.75%, with an average of 36.12% (Fig. 2). The monthly average data gaps are also shown in Fig. 2, and found that the data gaps of the TP were the largest in summer (June to August), with an average of 45.20%. The data gaps declined rapidly starting from September and were the smallest in November (with an average of 28.00%), and gradually increased in winter and spring.




Figure 2: Annual and monthly average proportion of data gaps of original composite MODIS snow products.

135 2.2.2 Digital Elevation Models (DEMs)

We used the Shuttle Radar Topography Mission (SRTM) DEMs to calculate the topographical parameters (e.g., elevation, slope, and aspect). The SRTM DEM is at 90 m resolution in GeoTIFF format was from the United States Geological Survey (USGS). We then pre-processed the DEM data, including mosaicking, reprojection, resampling, and clipping.

2.2.3 *In situ* observations

140 We used snow depths from 137 *in situ* stations in the TP (Fig. 1a) obtained from October 1, 2002, to March 31, 2021, which were provided by the National Meteorological Centre of China. Fig. 1a shows all stations over the TP, which record coordinates, observation time, snow depth, and snow pressure. The surface elevation of these stations ranges from 1187 m to 5310 m, with an average of 3240 m. Overall, 72% stations are situated in the east ($>95^{\circ}\text{E}$), and only 22% are situated in high-altitude areas (>4000 m). A 3 cm threshold was utilized to evaluate the snow classes using *in situ* snow depth (Huang et al., 2022). The snow depth was reclassified as no-snow if it was less than 3 cm; if not, it was considered as snow.

2.2.4 ~~Landsat-8 imagery~~Landsat images

The snow cover mapped from ~~Landsat-8 imagery~~Landsat images were used to validate and evaluate our snow cover product. Using the GEE cloud platform, we selected ~~62-81~~ Landsat-8 series images with less than ~~10%-2%~~ cloud coverage under clear sky for verification, including Landsat-5 TM, Landsat-7 ETM+, and Landsat-8 OLI images. The detailed ~~Landsat-8 imagery~~Landsat images information is shown in the supplementary material (Table A1). The selected images were distributed throughout the study area from ~~November 22~~January 4, 20162002, to December 1634, 20212018. We used the C topographic correction model to correct for the terrain effect on Landsat series data (Teilet et al., 1982). Then, uUsing the SNOWMAP method proposed by Hall et al. (1995), we generated a snow cover map using ~~Landsat-8 imagery~~Landsat images for the TP. The SNOWMAP algorithm classified the pixels with NDSI values > 0.4 , green band > 0.10 , and SWIR1 band > 0.11 as snow (Huang et al., 2011). To improve the accuracy of resampling 30 to 500 m resolution, we adopted the following steps for resampling: first, for each MODIS pixel, we calculated the amount of Landsat snow-covered pixels in the current MODIS pixel. Second, we divided the snow-covered Landsat pixels by sum of the Landsat pixels contained in each MODIS pixel (a 500 m MODIS pixel is close to 277 Landsat pixels) (Crawford, 2015; Liu et al., 2020a). Finally, we reclassified pixels whose results in the previous step were less than 0.5 as no-snow; otherwise, they were reclassified as snow. Hence, we obtained resampled 500 m resolution Landsat images for further verification.

3 HMRF modelling technique

The HMRF framework optimally combines MODIS spectral, spatiotemporal, and environmental information to fill data gaps, thereby increasing snow estimate accuracy (Huang et al., 2018). This framework is expressed as a linear energy function in

165 which the total energy is modelled as the combination of all information (Eq. (1)). Thus, the HMRF framework requires to specify energy function for each information and to determine the optimal parameters that minimize the total energy.

$$U_T(\beta_n, x_i, N_{sp}, N_{tp}, I_{ev}) = \lambda_{xi}U_{xi}(\beta_n, x_i) + \lambda_{st}U_{st}(\beta_n, N_{sp}, N_{tp}) + \lambda_{ev}U_{ev}(\beta_n, I_{ev}) \quad (1)$$

where λ_{xi} , λ_{st} , and λ_{ev} are the spectral, spatiotemporal, and environmental parameters, respectively; and U_{xi} , U_{st} , and U_{ev} are the spectral, spatiotemporal, and environmental energy functions, respectively.

170 In the original HMRF modelling technique, Huang et al. (2018) used surface elevation to represent the environmental association in mountainous areas (hereafter referred to as HMRF_{ele}). However, because of the spatial heterogeneity of the TP, using only relative elevation cannot reflect the influence of complex terrain on snow cover (Figs. 1b and 1c). Therefore, the influence of additional topographic factors (e.g., slope, aspect, sunlight duration, and solar incidence angle) on snow cover must be considered. Based on the above reasons, we incorporated solar radiation as a surrogate for environmental information in HMRF framework (hereafter referred to as HMRF_{solar}). Further details are provided in Section 3.3.

175 The overall flowchart of the HMRF_{solar} modeling technique is shown in Fig. 3. We used daily composite MODIS snow cover products as initial dataset. First, we calculated the optimal parameters of each information using the Ho-Kashyap and DFBETAS algorithms (Bormann et al., 2012), based on randomly divided training samples on the initial snow cover product. Second, we determined the snow and non-snow classes of each pixel in the initial snow cover products using the optimal parameters and HMRF_{solar} algorithm. In the first round of the HMRF_{solar} algorithm, we employed a $3 \times 3 \times 3$
180 spatiotemporal cubic neighborhood to model spatiotemporal energy. If data gaps remained, we further expanded the cubic neighborhood of these pixels. Finally, we validated and evaluated our HMRF_{solar} -based products against the *in situ* observations, snow mapped from ~~Landsat 8 imagery~~[Landsat images](#), and snow cover data from the initial MODIS and original HMRF_{ele} modelling technique.




Figure 3: Overall flowchart of the HMRF_{solar}-based framework. (SCP stands for snow cover products)

3.1 Spectral information

The spectral energy is considered as the possibility of the pixel pertaining to non-snow or snow according to its spectral information. Fractional Snow Cover (FSC) represents the proportion of snow in a pixel, which can be calculated using the NDSI value provided by the MODIS snow product (Salomonson and Appel, 2004). ~~Aqua Band 6 was reinstated using an image restoration algorithm in MODIS version 6 (Gladkova et al., 2012); thus, a general equation can be used to calculate the FSC. The general snow fraction estimation method was derived over other regions, and many studies have shown that the derived linear relationship has limited accuracy in the TP. Therefore, we re-fitted the empirical relationship between FSC and NDSI of Terra and Aqua satellites in the TP using Landsat data over 20 years with 972884, and 952221 sample points, respectively.~~ The spectral probability $P(x_i|\beta_1)$ of snow class ~~can be was re-fitted defined~~ as follows:

$$P(x_i|\beta_1) = (-0.01 + 1.45 \times NDSI)/100 \quad (2)$$

$$P(x_i|\beta_1)_{Terra} = (1.222 \times NDSI + 0.038)/100 \quad (2)$$

$$P(x_i|\beta_1)_{Aqua} = (1.164 \times NDSI + 0.058)/100 \quad (3)$$

~~The correlation coefficients of the re-fitted equation of Terra and Aqua satellites were 0.86 and 0.89, respectively.~~ The spectral energy $U_{xi}(\beta_n, x_i)$ can be calculated as follows:

$$U_{xi}(\beta_n, x_i) \begin{cases} -[P(x_i|\beta_1)], & \text{if } n = 1 \\ -[1 - [P(x_i|\beta_1)]], & \text{if } n = 2 \end{cases} \quad (43)$$

where if pixel i is classified as no-snow class, then $n = 2$; if pixel i is the snow class, $n = 1$.

3.2 Spatiotemporal information

To fully take advantage of spatiotemporal information, we combined the temporal and spatial information to form a spatiotemporal cubic neighborhood N_{st} (Huang et al., 2018). In modeling, initial snow products ~~are~~ updated iteratively using classification results from the previous iteration. Convergence ends when the proportion of pixels whose type changes between two subsequent iterations is less than 0.1%, in which case the data gaps need to be calculate. If the data gaps still exist, the cubic neighborhood is temporally and spatially expanded next round.

We primarily set a $3 \times 3 \times 3$ cubic neighborhood; that is, the current pixel and its neighboring pixels formed a cube on day t , the day before ($t - 1$), and the day after ($t + 1$). In this condition, the energy equation of the current pixel is expressed as the weight of the proportion of snow and non-snow in the spatiotemporal neighborhood of the current pixel. The weight is reversely proportional to the distance from the neighboring pixel to the center pixel:

$$U_{st}(\beta_n, N_{sp}, N_{tp}) = \begin{cases} -\sum_{x=-1}^1 \sum_{y=-1}^1 \sum_{t=-1}^1 \frac{\frac{S(x,y,t)}{Dist(x,y,t)}}{\frac{V(x,y,t)}{Dist(x,y,t)}}, & \text{if } n = 1 \\ -\sum_{x=-1}^1 \sum_{y=-1}^1 \sum_{t=-1}^1 \frac{\frac{NS(x,y,t)}{Dist(x,y,t)}}{\frac{V(x,y,t)}{Dist(x,y,t)}}, & \text{if } n = 2 \end{cases} \quad (54)$$

where (x, y, t) is the relative coordinates of the pixel in the spatiotemporal cube; $S(x, y, t)$, $NS(x, y, t)$, $V(x, y, t)$,
 215 and $Dist(x, y, t)$ are defined respectively as follows:

$$S(x, y, t) = \begin{cases} 1, & \text{if } (x, y, t) \text{ pertains to snow} \\ 0, & \text{if } (x, y, t) \text{ pertains to non - snow} \end{cases} \quad (65)$$

$$NS(x, y, t) = \begin{cases} 1, & \text{if } (x, y, t) \text{ pertains to non - snow} \\ 0, & \text{if } (x, y, t) \text{ pertains to snow} \end{cases} \quad (76)$$

$$V(x, y, t) = \begin{cases} 1, & \text{if } (x, y, t) \text{ pertains to valid class} \\ 0, & \text{if } (x, y, t) \text{ pertains to data gaps} \end{cases} \quad (87)$$

$$Dist(x, y, t) = \sqrt{x^2 + y^2 + wt^2} \quad (98)$$

220 where $Dist(x, y, t)$ is the distance from each neighborhood pixel to the center pixel, which determines the weight of each area pixel; w denotes the weight of the temporal distance related to the spatial distance. This value was calculated via semi-variogram analysis and was determined as 3 in this study.

If data gaps still remain in $3 \times 3 \times 3$ spatiotemporal cubic neighborhood, we further expanded spatiotemporal neighborhood to $5 \times 5 \times 5$. In this study, when we used a $5 \times 5 \times 5$ cubic neighborhood, the data gaps were reduced to ~~0.01420~~0.0173%,
 225 which had little effect on snow cover estimate accuracy. Hence, the temporal neighborhood remained at 5, whereas the spatial neighborhood continued to expand. Finally, all data gaps could all be filled by continuously expanding the spatial neighborhood, which remarkably improved the MODIS estimate accuracy.

3.3 Environmental contextual information

The geographic location and seasons can determine the total solar radiation received on the TP, and the complex topography
 230 of the TP determines the availability of solar radiation at specific locations in the region, which further affects accumulation and melting of snow cover and determines its distribution (You et al., 2020). Here we use solar radiation to include the influence of environmental factors on snow cover, such as slope, aspect, sunlight duration, and solar incidence angle.

Kumar et al. (1997) originally proposed the solar radiation model, SHORTWAVE, to calculate the direct, reflected, and
 235 diffuse solar energy received by the ground. Huang et al. (2015) used a Graphics Processing Unit (GPU) methodology to improve the speed and effectiveness for modelling. In our study, we used the GPU-accelerated SHORTWAVE model as the environmental parameter in the $HMRF_{solar}$ model to calculate the daily solar radiation of the TP.

A general-purpose desktop computer was used to test the parallel computational efficiency. The computer has an Intel Core™ i7-10700 CPU (16 cores and max clock rate is at 2.90 GHz), an NVIDIA GeForce RTX 2070 SUPER Card with 2560 cores and 16240 MB global memory, and Windows 11 Ultimate 64-bit Operation System. The GPU-accelerated
 240 SHORTWAVE model uses latitude, slope, aspect, date, and interval time as inputs. To calculate the overall solar radiation in a day, we subdivided the day into subsequent time intervals, and set 15 min as the suitable interval time in this paper (Kumar et al., 1997; Antonic, 1998). The overall solar radiation I_a at time t can be calculated using Eq. (9):

$$I_a = I_p + I_d + I_r \quad (109)$$

where I_p , I_d , and I_r are the direct, diffuse, and reflected solar radiation, respectively, and can be estimated using the following equations:

$$I_p = I_0 \cdot \tau_b \cdot \cos\theta \quad (119)$$

$$I_d = \frac{I_0 \cdot \tau_d \cdot \cos^2\beta}{2\sin\alpha} \quad (124)$$

$$I_r = \frac{r \cdot I_0 \cdot \tau_r \cdot \sin^2\beta}{2\sin\alpha} \quad (132)$$

where I_0 is the extra-atmospheric solar flux; τ_b denotes the atmospheric transmittance; τ_d denotes the diffuse skylight transmittance; τ_r is the reflectance transmissivity (Gul et al., 1998); r is set to 0.2 in the current paper (Kumar et al., 1997); α is the solar elevation angle, which varies in terms of the latitude, season and time; θ is the solar incidence angle, which is determined by solar elevation angle α , the solar azimuth angle, surface slope (β) and aspect (Huang et al., 2015). The effect of the geographic location and seasons on the total solar radiation are represented by solar elevation angle variable α , and the effect of the complex topography on the total solar radiation is represented by surface slope (β) and aspect variables.

We repeated the calculation for each time interval. Last, we accumulated the solar radiation from sunrise to sunset to acquire the daily values.

Areas that receive more solar radiation generally experience later snow accumulation and earlier snowmelt than areas that receive less solar radiation. Therefore, if pixels in a spatial neighborhood have higher solar radiation and the classification result is snow, the probability that the center pixel is also snow increases. The environmental energy equation is defined as follows:

$$U_{ev}(\beta_n, N_{tp}, N_{sp}) = \begin{cases} -\sum_{x=-1}^1 \sum_{y=-1}^1 \frac{SE(x,y)}{SE(x,y)+NSE(x,y)}, & \text{if } n = 1 \\ -\sum_{x=-1}^1 \sum_{y=-1}^1 \frac{NSE(x,y)}{SE(x,y)+NSE(x,y)}, & \text{if } n = 2 \end{cases} \quad (143)$$

where (x,y) are coordinates of the pixel in the spatial neighborhood N_{sp} ; $SE(x,y)$ and $NSE(x,y)$ are defined as follows:

$$SE(x,y) = \begin{cases} 1, & \text{if } (x,y) \text{ belongs to snow and } solar(x,y) \geq solar(i) \\ 0, & \text{others} \end{cases} \quad (154)$$

$$NSE(x,y) = \begin{cases} 1, & \text{if } (x,y) \text{ belongs to non - snow and } solar(x,y) \leq solar(i) \\ 0, & \text{others} \end{cases} \quad (165)$$

where $solar(x,y)$ and $solar(i)$ represent the solar radiation received by the neighboring pixels and central pixel, respectively.

4 Results

4.1 Accuracy assessment based on *in situ* observations

Using HMRF_{solar}-based method, we produced gap-free snow cover products over the TP with a daily 500 m resolution from May 15, 2002, to December 31, 2021. We applied the validation indices based on the confusion matrix (Table 1), which is widely recommended by previous studies (Li et al., 2020).

Table 1. Confusion matrix comparing snow cover products with observation value

| Observation value | Snow cover products | |
|-------------------|---------------------|----------|
| | Snow | Non-snow |
| Snow | a | b |
| Non-snow | c | d |

The three validation indices are defined as follows:

$$OA = \left(\frac{a+d}{a+b+c+d} \right) \times 100\% \quad (17)$$

$$OE = \left(\frac{b}{a+b} \right) \times 100\% \quad (18)$$

$$CE = \left(\frac{c}{c+d} \right) \times 100\% \quad (19)$$

Overall accuracy (OA) is the probability a pixel that is correctly classified; Omission error (OE) and Commission error (CE) represent the underestimated and overestimated snow pixels, respectively.

We first compared our daily HMRF_{solar}-based and original MODIS snow products with snow depth observed from 137 *in situ* stations from 2002–2021 (Table 2). The overall accuracy (OA) of all gap-free pixels in original MODIS products was 97.96%. After employing the HMRF_{solar}-based method, the OA of these gap-free pixels was increased to ~~98.29%~~ 98.31%. The ~~omission error~~ OE was reduced by 0.57%. Since the *in situ* stations in the TP are primarily distributed in low-altitude areas (Fig. 1a), only using *in situ* observations may not be representative and lead to biases in the accuracy assessment. Therefore, we also conducted evaluation in comparison with snow cover mapped from ~~Landsat 8 imagery~~ Landsat data obtained from ~~2016~~ 2002–2021. ~~We selected Landsat 8 imagery acquired during 2016–2018 for two reasons. First, Fig. 2 shows the proportion of data gaps in the original MODIS products in these three years covered low, medium, and high values, which could represent the range of data gaps in original products covering 2002–2021. Second, according to our accuracy evaluation of HMRF_{solar}-based snow products in comparison with *in situ* observations from 2016–2018 (Table 2), we found that the OA of these three years was 97.76%, which is nearly the same as the OA from 2002–2021 (Table 1). This suggests that the accuracy from 2016–2018 of the products can represent that of the long-term series products. Therefore, in the follow-up accuracy evaluation, we used the snow cover mapped from the Landsat 8 imagery from 2016–2018 and regarded it as the true verification value in the following accuracy evaluation.~~

Table 2. Confusion matrices between HMRF_{solar}-based, original MODIS snow products, and *in situ* observation for gap-free pixels in original MODIS snow products during 2002–2021.

| <i>In situ</i> observation | HMRF _{solar} -based snow products | | | Original MODIS snow products | | |
|----------------------------|---|--|--------|------------------------------|--------------------|--------|
| | Snow | Non-snow | Total | Snow | Non-snow | Total |
| Snow | 4766 <u>4782</u> (66.61 <u>66.61</u> %) | 2389 <u>2373</u> (33.39%) | 7155 | 4725 (66.04%) | 2430 (33.96%) | 7155 |
| Non-snow | 3350 <u>3420</u> (1.01%) | 3281 <u>3280</u> (98.99%) | 331460 | 4493 (1.36%) | 326927 (98.64%) | 331460 |
| Total | 8116 <u>8202</u> | 3304 <u>993304</u> <u>13</u> | 338615 | 9218 | 329397 | 338615 |
| Overall accuracy | 98.31% <u>98.29%</u> | | | 97.96% | | |

295 **Table 2.** Confusion matrices between ~~HMRF_{solar}-based snow products and *in situ* observation during 2016–2018.~~

| <i>In situ</i> observation | HMRF _{solar} -based snow products | | |
|----------------------------|--|-------------------------------|-------------------|
| | Snow | Non-snow | Total |
| Snow | 1325 (43.15%) | 1746 (56.85%) | 3071 |
| Non-snow | 1488 (1.06%) | 139501 (98.94%) | 140989 |
| Total | 2813 | 141247 | 144060 |
| Overall accuracy | 97.76% | | |

4.2 Accuracy assessment based on ~~Landsat-8 imagery~~Landsat data

A total of ~~62–81~~ Landsat-8 images (Fig. 1a, Table A1) with less than ~~102~~% cloud coverage were selected for this accuracy assessment. We constructed the confusion matrices for the HMRF_{solar}-based snow cover products, HMRF_{ele}-based snow products, and original MODIS snow cover products against snow cover mapped from ~~Landsat-8 imagery~~Landsat series data products ~~during 2016–2018~~ (Table 3). The OA for gap-free pixels in original MODIS products was ~~89.31~~89.45%. When we used surface elevation to represent the environmental association in HMRF modelling technique, the OA of these gap-free pixels was increased to ~~89.72~~90.47%. Since using surface elevation only cannot well represent the influence of complex topography on snow cover distribution over the TP, this improvement is still limited. After incorporating the solar radiation to model the comprehensive influence of multiple topographic factors on snow cover, the OA of our HMRF_{solar}-based snow products was increased by ~~2.99~~2.06% compared with the OA of original MODIS products. Liu et al. (2020a) indicated that the classification errors in original MODIS snow products were strongly affected by ~~omission error~~-(OE). Our snow cover product provided a considerable improvement in this respect. The OE of our HMRF_{solar}-based snow products was ~~4.70~~3.24% lower than that of original MODIS products (Table 3).

Particularly, our HMRF method is able to fill data gaps in original MODIS products in which spectral information is unavailable caused by cloud by integrating spatiotemporal and environmental information together. Table 4 shows that the OA of the HMRF_{solar}- and HMRF_{ele}- based snow cover products for those data-gap pixels in original MODIS products was ~~83.86%~~83.18% and ~~81.15%~~80.32%, which is ~~6.27%~~7.50% and ~~9.32%~~9.13% lower than that of gap-free pixels. It is also showed that the OA of HMRF_{solar}-based snow cover product is ~~2.71%~~2.86% higher than that of HMRF_{ele}-based snow cover

product. Both Table 3 and Table 4 confirmed that using solar radiation as the environmental information in the HMRF
315 framework can develop more reliable snow cover products.

4.3 Accuracy improvement in different times of a snow season

To explore the details of snow cover variation, we define the snow season from September of previous year to August of the following year, for example, the time range of the 2002 snow season is 2002.9.1-2003.8.31 (Chen et al., 2018b). Previous studies (Huang et al., 2018; Klein, 2003; Li et al., 2019a) have suggested that the snow cover estimation accuracy was
320 generally low during snow transitional period (i.e., the beginning and end of a snow season). We plotted the temporal variations in the OA, OE, and CE of the $\text{HMRF}_{\text{solar}}$ - and HMRF_{ele} -based snow cover products (Fig. 4). Except November, December, and February, the OAs of the new snow products were more than 90% in all months. The accuracy was relatively low during snow transitional period (November, December, and February to April), whereas the accuracy was higher in snow stable period. The accuracy was comparatively low during snow transitional period (November to December, and
325 February to April), whereas the accuracy was higher in snow stable period (January). The trend of the CE was almost the same as that of snow cover; that is, the months with more snow cover had a larger CE. However, the CE was generally low in all months. By comparison with the HMRF_{ele} -based products, the accuracy of our new products was higher in almost all months. In addition to the stable snow period, the improvement for transition period was also notable, which increased by an average of ~~2.69%~~2.18%. In transition period, with rapid temperature changes, the snow status changed rapidly, and the snow
330 mapped by MOD10A1 (crossing at 10:30) and MYD10A1 (crossing at 13:30) were different because of the different temperatures and solar radiation. The original MODIS snow cover products have relatively high error during snow transitional period (Li et al., 2019). After incorporating solar radiation as an environmental contextual information in HMRF, the accuracy of snow cover products has been improved effectively.

Table 3. Confusion matrices between $\text{HMRF}_{\text{solarr}}$ -based snow products, HMRF_{ele} -based snow products, original MODIS snow products, and snow cover mapped from Landsat-8 imagery products for gap-free pixels during 2016–2018.



| Landsat-8 imagery | $\text{HMRF}_{\text{solarr}}$ -based snow products | | HMRF_{ele} -based snow products | | Original MODIS snow products | |
|-------------------|--|---------------------|---|---------------------|------------------------------|---------|
| | Snow | Non-snow | Total | Snow | Non-snow | Total |
| Snow | 1391834 (89.46%) | 164049 (10.54%) | 1555883 | 1320466 (84.87%) | 235417 (15.13%) | 1555883 |
| Non-snow | 77459 (4.73%) | 1561762 (95.27%) | 1639221 | 92900 (5.67%) | 1546321 (94.33%) | 1639221 |
| Total | 1469293 | 1725811 | 3195104 | 1413366 | 1781738 | 3195104 |
| Overall accuracy | 92.44% | | 89.72% | | 89.45% | |

Table 3. Confusion matrices between $\text{HMRF}_{\text{solarr}}$ -based snow products, HMRF_{ele} -based snow products, original MODIS snow products, and snow cover mapped from Landsat series data products for gap-free pixels during 2002–2021.

| Landsat series data | $\text{HMRF}_{\text{solarr}}$ -based snow products | | HMRF_{ele} -based snow products | | Original MODIS snow products | |
|---------------------|--|---------------------|---|--------------------|------------------------------|---------|
| | Snow | Non-snow | Total | Snow | Non-snow | Total |
| Snow | 916593 (85.06%) | 160936 (14.94%) | 1077529 | 901202 (83.64%) | 176327 (16.36%) | 1077529 |
| Non-snow | 108214 (5.31%) | 1931065 (94.69%) | 2039279 | 120590 (5.91%) | 1918689 (94.09%) | 2039279 |
| Total | 1024807 | 2092001 | 3116808 | 1021792 | 2095016 | 3116808 |
| Overall accuracy | 91.36% | | 90.47% | | 89.31% | |

Table 4. Confusion matrices between HMRF_{solar} - based snow cover products, HMRF_{ele} -based snow cover products, and snow cover mapped from ~~Landsat 8 imagery~~Landsat series data products for data-gap pixels during ~~2016–2018~~2002–2021.

| <u>Landsat series data</u> | <u>HMRF_{solar}-based snow cover product</u> | | | <u>HMRF_{ele}-based snow cover product</u> | | |
|----------------------------|--|----------------------------------|---------------|--|----------------------------------|---------------|
| | <u>Snow</u> | <u>Non-snow</u> | <u>Total</u> | <u>Snow</u> | <u>Non-snow</u> | <u>Total</u> |
| <u>Snow</u> | <u>87921</u> <u>(77.80%)</u> | <u>25093</u> <u>(22.20%)</u> | <u>113014</u> | <u>83868</u> <u>(74.21%)</u> | <u>29146</u> <u>(25.79%)</u> | <u>113014</u> |
| <u>Non-snow</u> | <u>13795</u> <u>(10.78%)</u> | <u>114148</u> <u>(89.22%)</u> | <u>127943</u> | <u>16269</u> <u>(12.72%)</u> | <u>111674</u> <u>(87.28%)</u> | <u>127943</u> |
| <u>Total</u> | <u>101716</u> | <u>139241</u> | <u>240957</u> | <u>100137</u> | <u>140820</u> | <u>240957</u> |
| <u>Overall accuracy</u> | <u>83.86%</u> | | | <u>81.15%</u> | | |




340 Figure 4: Temporal variations in OA (a), OE (b), and CE (c) of HMRFele- and HMRFsolar-based snow products from 2016-2018/2002-2021.


4.4 Accuracy improvement for different surface topography

We divided the elevation of the study area into four categories: <3000, 3000–4000, 4001–5000, and >5000 m, to explore the effect of elevation on snow cover. We then calculated the snow cover products accuracy in each category. Fig. 5 indicates that the OA of the products decreased with increasing elevation. The accuracy of our products was highest in the <3000 m category (94.2697.79%). With increasing elevation, the OE first decreased and then increased; that is, the OE value was the lowest in the 4001–5000 m category, whereas the CE was the highest in the 4001–5000 m category. Li et al. (2020) established effect of elevation on snow cover and also demonstrated the accuracy of original MODIS snow products generally decreased with increasing elevation. Areas at higher elevation generally receive more solar radiation; hence, the snow status changes more rapidly at higher elevations, resulting in decreased snow cover estimation accuracy.

By comparison with the $HMRF_{ele}$ -based products, the accuracy of our new products was higher in almost all elevation categories, and as the elevation increased, the accuracy improvement was more remarkable (reaching an improvement of 3.16%~~3.39%~~ in the >5000 m category). The improvements in OE first decreased and then increased; that is, the accuracy improvement of the OE value was the smallest in the 4001–5000 m category, with an improvement of 3.474.28%. The CE exhibited the opposite trend; that is, the accuracy improvement in the CE value was the largest in the 4001–5000 m category, at 2.334.2%. The accuracy improvement provided by our new snow cover product was substantially increased with the increase in elevation, indicating its effectiveness in high-altitude areas.



| Elevation (m) | Number of samples | OA-HMRFele (%) | OA-HMRFsolar (%) | OE-HMRFele (%) | OE-HMRFsolar (%) | CE-HMRFele (%) | CE-HMRFsolar (%) |
|---------------|-------------------|----------------|------------------|----------------|------------------|----------------|------------------|
| <3000m | 131774 | 97.10 | 97.79 | 36.03 | 25.82 | 2.02 | 1.58 |
| 3000-4000m | 657347 | 90.43 | 93.02 | 15.36 | 10.03 | 6.44 | 5.33 |
| 4001-5000m | 1845363 | 88.75 | 91.96 | 12.92 | 8.64 | 8.10 | 6.90 |
| >5000m | 992521 | 87.50 | 90.89 | 19.43 | 13.57 | 5.44 | 4.56 |



| Elevation (m) | OA-HMRFele | OA-HMRFsolar | OE-HMRFele | OE-HMRFsolar | CE-HMRFele | CE-HMRFsolar |
|---------------|------------|-----------------|------------|-----------------|------------|----------------|
| <3000m | 92.96% | 94.26% (+1.30%) | 27.31% | 20.98% (-6.33%) | 2.16% | 1.98% (-0.18%) |
| 3000-4000m | 89.15% | 91.33% (+2.18%) | 19.84% | 14.69% (-5.15%) | 7.23% | 5.91% (-1.32%) |
| 4001-5000m | 87.12% | 89.60% (+2.48%) | 13.24% | 9.77% (-3.47%) | 9.12% | 6.79% (-2.33%) |
| >5000m | 85.09% | 88.25% (+3.16%) | 20.37% | 16.12% (-4.25%) | 6.36% | 5.16% (-1.20%) |


360 **Figure 5: Effect of elevation on OA (a), OE (b), and CE (c) of HMRFele- and HMRFsolar-based snow products from 2016-2018/2002-2021.**

To explore the effect of slope on snow cover, we divided the slope of the study area into five categories: <math> <10^\circ </math>, $10^\circ\text{--}20^\circ$, $21^\circ\text{--}30^\circ$, $31^\circ\text{--}40^\circ$, and $>40^\circ$. ~~Because no Landsat 8 imagery met the conditions in areas with a slope > 40°, we only calculated the accuracy of the first four categories (Fig. 6).~~ The OA of the snow cover products decreased with increasing slope, which is consistent with the trend of the changes in accuracy with increasing elevation (Fig. 6). The accuracy of our new products was highest in the 0 –10° category, with a value of 95.91/93.78%. The ~~UE~~OE was high for relatively low-slope areas (<math> <30^\circ </math>), and the ~~OE~~CE was high for high-slope areas ($\geq 30^\circ$).


365

By comparison with the HMRF_{ele} -based products, the accuracy of our new products was higher in almost all slope categories, and the accuracy improvement was remarkable in the 10° – 20° category (reaching an improvement of 2.472.90%). This means topography of 10° – 20° slopes had the strongest impact on snow cover on the TP. The accuracy improvements in OE decreased with increasing slope, whereas the accuracy improvements in CE increased with increasing slope. In the $<10^{\circ}$ – 20° category, the improvement in CE was the smallest, at 0.360.56%, whereas the improvement in OE was the largest, at 5.566.83%. In the 31° – 40° – $>40^{\circ}$ category, the improvement in CE was the largest, at 1.972.20%, whereas the improvement in

in OE was the smallest, at 2.040.81%.



| Slope ($^{\circ}$) | Number of samples | OA-HMRFele(%) | OA-HMRFsolar(%) | OE-HMRFele(%) | OE-HMRFsolar(%) | CE-HMRFele(%) | CE-HMRFsolar(%) |
|----------------------|-------------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|
| 0-10 $^{\circ}$ | 1998584 | 94.38 | 95.91 | 17.54 | 11.98 | 2.17 | 1.81 |
| 10-20 $^{\circ}$ | 1094227 | 90.17 | 92.64 | 17.07 | 11.78 | 6.56 | 5.37 |
| 21-30 $^{\circ}$ | 418397 | 89.53 | 91.82 | 12.10 | 8.25 | 9.86 | 8.15 |
| 31-40 $^{\circ}$ | 116797 | 87.38 | 89.38 | 6.61 | 4.57 | 14.8 | 12.83 |



| Slope ($^{\circ}$) | OA-HMRFele | OA-HMRFsolar | OE-HMRFele | OE-HMRFsolar | CE-HMRFele | CE-HMRFsolar |
|--------------------------|------------|-----------------|------------|-----------------|------------|-----------------|
| <math><10^{\circ}</math> | 91.84% | 93.78% (+1.94%) | 26.94% | 20.11% (-6.83%) | 3.31% | 2.75% (-0.56%) |
| 10-20 $^{\circ}$ | 88.52% | 91.42% (+2.90%) | 21.06% | 16.74% (-4.32%) | 7.54% | 6.31% (-1.23%) |
| 21-30 $^{\circ}$ | 87.32% | 90.03% (+2.71%) | 17.23% | 14.36% (-2.87%) | 11.42% | 9.73% (-1.69%) |
| 31-40 $^{\circ}$ | 86.07% | 87.76% (+1.69%) | 13.35% | 10.91% (-2.44%) | 13.72% | 11.97% (-1.75%) |
| >40 $^{\circ}$ | 84.39% | 85.62% (+1.23%) | 7.54% | 6.73% (-0.81%) | 17.97% | 15.77% (-2.20%) |

Figure 6: Effect of slope on OA (a), OE (b), and CE (c) of HMRF_{ele} - and HMRF_{solar} -based snow products from ~~2016–2018~~2002–2021.

380 Finally, we explored the effect of aspect on snow cover (Fig. 7). The OA of the snow cover products was higher on shaded slopes than on sunny slopes. The OAs of our new products on sunny and shaded slopes were 91.1488.37% and 92.5690.07%, respectively. By comparison with the $HMRFele$ -based products, the accuracy improvement provided by our new products was remarkable on sunny slopes (improvement of 3.63.16%) than on shaded slopes (improvement of 2.541.75%), corresponding with the actual situation. The improvement in the OE on sunny slopes was also remarkable (improvement of 5.883.63%).





Figure 7: Effect of aspect on OA (a), OE (b), and CE (c) of $HMRFele$ - and $HMRFsolar$ -based snow products from 2016–20182002–2021.

5 Discussion

390 5.1 Effect of the $\text{HMRF}_{\text{solar}}$ -based snow cover products over complex terrain


Long-term and high-precision snow cover products are key to the snow hydrology research in the TP. Compared to the heterogeneous land cover, complex terrain has more severe effects on MODIS snow products in the TP (Liu et al., 2020a; Azizi and Akhtar, 2021). To enhance snow detection accuracy in mountainous areas, we used solar radiation as the environmental contextual information rather than surface elevation used in the original HMRF_{ele} modelling technique

395 (Huang et al., 2018). The calculated optimal parameters used in Eq. (1) could reflect the importance of each energy to a certain extent. The optimal parameters of the $\text{HMRF}_{\text{solar}}$ -based products for spectral, spatial-temporal, and environmental information were 0.117, 1.294, and 0.532, respectively, while that of the original HMRF_{ele} -based products were 0.338, 1.419, and 0.576, respectively. The proportion of environmental information of $\text{HMRF}_{\text{solar}}$ -based products (27.37%) were higher than that of HMRF_{ele} -based products (24.70%), which shows the improvement of $\text{HMRF}_{\text{solar}}$ -based products over complex


400 terrain. Moreover, Fig. 8a shows a true-color Sentinel-2B image. Figs. 8b, 8c, and 8d show the original MODIS, HMRF_{ele} - and $\text{HMRF}_{\text{solar}}$ -based snow cover products, respectively, on October 31, 2018. The examples in Fig. 8 (where the topography has a dominant effect on snow cover) show that the original HMRF_{ele} -based snow cover products mapped all data gaps in the shaded and sunny slopes at the same elevation as snow-covered pixels; however, after incorporating solar radiation, our new products accurately identified them as non-snow-covered pixels. In general, since sunny slopes receive

405 relatively more solar radiation, the area of snow cover on sunny slopes is smaller than that on shaded slopes. The new $\text{HMRF}_{\text{solar}}$ -based snow products more effectively filled the data gaps on sunny slopes, producing more realistic results. Solar radiation was particularly effective as an environmental contextual factor to correct the snow detection errors in mountainous areas.


(a) Sentinel-2B




(b) Original MODIS SCP



(c) HMRF_{Fele}-based SCP



(d) HMRF_{solar}-based SCP




Legend

□ Snow

■ Non-snow

■ Data gaps






410 **Figure 8: Comparison between true-color Sentinel-2B imagery and (a), original MODIS snow products (b), HMRFele-based snow products (c) HMRFsolar-based snow products.**

5.2 Potential applications of the HMRFsolar-based snow cover products

415 Compared with *in situ* observations, the overall accuracy of snow cover products on the TP reported by other studies is in the range of 90.74%-96.6%, and the omission error is greater than the commission error (Yu et al., 2016; Qiu et al., 2017; Xu et al., 2017; Zheng and Cao, 2019). The overall accuracy of our new snow products is 98.29% in comparison with *in situ* observations, and the new product has a considerable improvement in omission error. The new snow cover products provide spatiotemporally continuous snow cover distribution at 500 m resolution over the past two decades, thus having a great application potential for analyzing the processes of snow accumulation and melting. Figure 9 shows the snow cover trends over a sample area (location shown in Fig.1) from our daily gap-free snow products and 8-day composite MODIS snow

420

products on Terra (MOD10A2 v6.1) (Hall et al., 2021) during the 2017 snow season. We chose this sample region because it covers more mountainous areas including Hengduan Mountain and Bayankala Mountain, and has seasonal snow for multiple years. The MOD10A2 v6.1 products composite 8-days MODIS data to reduce the data gaps in the original daily MODIS snow products, and represent the maximum snow cover extent in 8 days, which means once snow is identified on any day in the 8-day time window, the pixel is mapped as snow. Although the overall variation trends of snow cover are similar between these two products, the snow cover percentage obtained from the MOD10A2 v6.1 data was 19.60~~19.70~~% higher than that of daily products during 2017 snow season. The status of the snow in the TP changes rapidly. Compared with the 8-day composite MODIS snow products (MOD10A2 v6.1), the HMRF_{solar}-based snow cover products can get more detailed and accurate information on snow accumulation and melting processes, particularly during the snow transitional periods (February to April, and November to December) when short-term snowfall events occur frequently (Fig.9). In addition, owing to its high temporal resolution (daily) and long-term spatiotemporal continuity, the generated snow cover products also provide important baseline information for monitoring climate change, calibrating hydrological models, and simulating snowmelt runoff in the TP.






Figure 9: Snow cover percentage for daily HMRF_{solar}-based snow cover products and 8-day composite MODIS snow products in a sample area during the 2017 snow season of the TP. (SCP stands for snow cover products.)

5.3 Limitations of the HMRF_{solar}-based snow cover products

The accuracy evaluation for the new snow cover products had some limitations. The majority of *in situ* observations in the TP are distributed in the low-altitude areas in the east, with only 22% located in high-altitude areas (Fig. 1a). The number of *in situ* observations in the snow category was considerably smaller than that in the non-snow category, which created challenges in verifying snow category classification (Zhang et al., 2021). Liu et al. (2020a) demonstrated that compared to *in situ* observations, snow cover mapped from Landsat series images is more related to MODIS snow cover products owing to the closer spatial match. Therefore, the conclusions about the new snow products were based on the accuracy evaluation against Landsat 8 imagery~~Landsat images~~, which may still contain some bias. Additionally, the solar radiation indicated used only imitates the solar radiation under clear-sky conditions. In a follow-up study, we will simulate the solar radiation throughout the year according to the weather conditions (e.g., the effect of cloud cover) and use parallel calculations to enhance calculation efficiency and further improve the snow cover product accuracy.

6 Data availability

The long-term daily snow cover products produced here are gap-free at 500 m resolution under the Universal Transverse Mercator (UTM zone 45) projection, and can be freely accessed from the National Tibetan Plateau Data Center at <https://doi.org/10.11888/Cryos.tpd.c.272204> (Huang and Xu, 2022), which is stored as a zip file (~ 1.56 GB) for each year. By uncompressing the zip file, the daily snow cover data is provided in GeoTIFF format, and the values in the snow cover

products are classified as snow (1) and non-snow (2). The name of each file is "HMRFSTP_yyyyddd.tif", in which
455 HMRFSTP is the abbreviation of "Hidden Markov Radom Field -based snow cover products for Tibetan Plateau", yyyy
stands for year and DDD stands for Julian day, such as 2002135.tif describes the snow cover on Tibet Plateau on the 135th
day of 2002.

7 Conclusions

In this study, we generated long-term daily gap-free snow cover products at 500 m resolution from original MODIS snow
460 products in the TP over the past two decades. The snow cover estimate accuracy was greatly improved by incorporating solar
radiation as a surrogate for environmental contextual information in HMRF framework in mountainous areas. We validated
and evaluated our snow cover products through comparison with *in situ* observations and high-resolution snow cover
mapped from ~~Landsat-8 imagery~~Landsat images, with accuracy estimate of ~~98.29% and 91.36%~~98.31% and 92.44%,
465 respectively. Our evaluation also suggests that incorporating solar radiation, instead of the simple use of surface elevation as
the environmental contextual information in HMRF framework, results in the accuracy of the snow products increase by
470 ~~2.722.71%~~ and the omission error decrease by ~~3.59%~~4.59%. Specifically, the accuracy of the new snow products is
particularly improved during snow transitional period and over complex terrains with high elevation as well as sunny slopes.
We believe that the long-term and spatiotemporally continuous snow cover products generated in this study have great
potential to analyse the processes of snow accumulation and melting, to monitor the climate change, and to understand the
hydrological cycling in the TP.

Author contributions.

HY and LHX developed the HMRF methodology and designed the study. XJH and ZYL applied the methodology and
performed the validation. XJY and XWJ performed the visualization. YBL, WSJ, and WJP analyzed the results. ZZJ worked
on data curation. Both the authors devoted to the writing of the paper and data quality control.

475 Competing interests.

The authors declare that they have no conflict of interest.

Disclaimer.

The views and interpretations in this publication are those of the authors and are not necessarily attributable to the ICI-MOD.

Special issue statement.

480 This article is part of the special issue “Extreme environment datasets for the three poles”. It is not associated with a conference.

Acknowledgments.

We would like to thank the National Meteorological Centre of China for providing the *in situ* observations over the Tibetan Plateau.

485 Financial support.

This work was supported by the National Natural Science Foundation of China (no. 42125604, 42071306 and 41701502).

Review statement.

This paper was edited by XX and reviewed by two anonymous referees.

Appendix

490 Table A1. Landsat ~~series-8-OLI~~ images used for assessment of the HMRF-based snow cover products in this study.

| <u>Image pair No.</u> | <u>Sensor</u> | <u>Tile path/row</u> | <u>Date of acquisition</u> | <u>Cloud cover (%)</u> |
|-----------------------|---------------|----------------------|----------------------------|------------------------|
| <u>1</u> | <u>ETM+</u> | <u>131/38</u> | <u>2002-11-22</u> | <u>1%</u> |
| <u>2</u> | <u>ETM+</u> | <u>136/38</u> | <u>2003-1-28</u> | <u>0%</u> |
| <u>3</u> | <u>ETM+</u> | <u>132/41</u> | <u>2003-2-17</u> | <u>1%</u> |
| <u>4</u> | <u>TM</u> | <u>136/33</u> | <u>2003-8-16</u> | <u>0%</u> |
| <u>5</u> | <u>TM</u> | <u>141/35</u> | <u>2003-9-20</u> | <u>0%</u> |
| <u>6</u> | <u>TM</u> | <u>135/33</u> | <u>2004-8-27</u> | <u>0%</u> |
| <u>7</u> | <u>TM</u> | <u>137/39</u> | <u>2004-12-15</u> | <u>1%</u> |
| <u>8</u> | <u>TM</u> | <u>132/38</u> | <u>2005-1-13</u> | <u>1%</u> |
| <u>9</u> | <u>TM</u> | <u>136/39</u> | <u>2005-3-14</u> | <u>1%</u> |
| <u>10</u> | <u>TM</u> | <u>132/34</u> | <u>2005-4-3</u> | <u>1%</u> |
| <u>11</u> | <u>TM</u> | <u>142/34</u> | <u>2005-6-28</u> | <u>0%</u> |
| <u>12</u> | <u>TM</u> | <u>136/36</u> | <u>2005-10-24</u> | <u>1%</u> |
| <u>13</u> | <u>TM</u> | <u>133/38</u> | <u>2005-11-4</u> | <u>1%</u> |
| <u>14</u> | <u>TM</u> | <u>135/39</u> | <u>2006-2-6</u> | <u>1%</u> |
| <u>15</u> | <u>TM</u> | <u>135/33</u> | <u>2006-8-1</u> | <u>0%</u> |

| | | | | |
|-----------|------------|---------------|-------------------|-----------|
| <u>16</u> | <u>TM</u> | <u>141/39</u> | <u>2006-9-28</u> | <u>1%</u> |
| <u>17</u> | <u>TM</u> | <u>132/34</u> | <u>2006-10-31</u> | <u>2%</u> |
| <u>18</u> | <u>TM</u> | <u>134/37</u> | <u>2006-11-30</u> | <u>1%</u> |
| <u>19</u> | <u>TM</u> | <u>136/39</u> | <u>2006-12-14</u> | <u>1%</u> |
| <u>20</u> | <u>TM</u> | <u>134/33</u> | <u>2007-3-6</u> | <u>1%</u> |
| <u>21</u> | <u>TM</u> | <u>141/34</u> | <u>2007-7-13</u> | <u>1%</u> |
| <u>22</u> | <u>TM</u> | <u>139/35</u> | <u>2007-9-17</u> | <u>1%</u> |
| <u>23</u> | <u>TM</u> | <u>132/37</u> | <u>2008-2-23</u> | <u>1%</u> |
| <u>24</u> | <u>TM</u> | <u>132/42</u> | <u>2008-3-10</u> | <u>1%</u> |
| <u>25</u> | <u>TM</u> | <u>134/36</u> | <u>2008-5-11</u> | <u>1%</u> |
| <u>26</u> | <u>TM</u> | <u>145/35</u> | <u>2008-6-25</u> | <u>1%</u> |
| <u>27</u> | <u>TM</u> | <u>142/34</u> | <u>2008-8-7</u> | <u>1%</u> |
| <u>28</u> | <u>TM</u> | <u>150/33</u> | <u>2008-10-2</u> | <u>1%</u> |
| <u>29</u> | <u>TM</u> | <u>130/37</u> | <u>2008-11-7</u> | <u>1%</u> |
| <u>30</u> | <u>TM</u> | <u>133/37</u> | <u>2008-12-14</u> | <u>1%</u> |
| <u>31</u> | <u>TM</u> | <u>132/38</u> | <u>2009-3-13</u> | <u>0%</u> |
| <u>32</u> | <u>TM</u> | <u>132/37</u> | <u>2009-4-14</u> | <u>1%</u> |
| <u>33</u> | <u>TM</u> | <u>147/35</u> | <u>2009-8-13</u> | <u>1%</u> |
| <u>34</u> | <u>TM</u> | <u>151/33</u> | <u>2009-9-10</u> | <u>1%</u> |
| <u>35</u> | <u>TM</u> | <u>138/37</u> | <u>2009-10-17</u> | <u>1%</u> |
| <u>36</u> | <u>TM</u> | <u>134/36</u> | <u>2009-11-22</u> | <u>1%</u> |
| <u>37</u> | <u>TM</u> | <u>133/38</u> | <u>2010-2-19</u> | <u>0%</u> |
| <u>38</u> | <u>TM</u> | <u>135/39</u> | <u>2010-3-21</u> | <u>1%</u> |
| <u>39</u> | <u>TM</u> | <u>150/32</u> | <u>2010-11-9</u> | <u>1%</u> |
| <u>40</u> | <u>TM</u> | <u>134/39</u> | <u>2011-3-1</u> | <u>1%</u> |
| <u>41</u> | <u>TM</u> | <u>141/35</u> | <u>2011-8-25</u> | <u>0%</u> |
| <u>42</u> | <u>TM</u> | <u>132/37</u> | <u>2011-10-29</u> | <u>0%</u> |
| <u>43</u> | <u>OLI</u> | <u>147/37</u> | <u>2013-4-18</u> | <u>2%</u> |
| <u>44</u> | <u>OLI</u> | <u>149/34</u> | <u>2013-5-18</u> | <u>2%</u> |
| <u>45</u> | <u>OLI</u> | <u>146/36</u> | <u>2013-8-1</u> | <u>1%</u> |
| <u>46</u> | <u>OLI</u> | <u>145/36</u> | <u>2013-9-27</u> | <u>2%</u> |
| <u>47</u> | <u>OLI</u> | <u>141/35</u> | <u>2013-11-18</u> | <u>1%</u> |
| <u>48</u> | <u>OLI</u> | <u>133/40</u> | <u>2014-1-13</u> | <u>1%</u> |
| <u>49</u> | <u>OLI</u> | <u>136/38</u> | <u>2014-2-19</u> | <u>1%</u> |
| <u>50</u> | <u>OLI</u> | <u>136/33</u> | <u>2014-7-13</u> | <u>0%</u> |
| <u>51</u> | <u>OLI</u> | <u>144/35</u> | <u>2014-8-22</u> | <u>1%</u> |
| <u>52</u> | <u>OLI</u> | <u>139/38</u> | <u>2015-1-10</u> | <u>1%</u> |
| <u>53</u> | <u>OLI</u> | <u>143/39</u> | <u>2015-3-11</u> | <u>1%</u> |
| <u>54</u> | <u>OLI</u> | <u>151/33</u> | <u>2015-10-13</u> | <u>1%</u> |

| | | | | |
|--------------------|---------------------|------------------------|----------------------------|--------------------|
| 55 | OLI | 136/38 | 2015-12-23 | 1% |
| 56 | OLI | 132/34 | 2016-5-3 | 2% |
| 57 | OLI | 151/33 | 2016-6-25 | 2% |
| 58 | OLI | 143/34 | 2016-9-21 | 1% |
| 59 | OLI | 133/37 | 2016-11-18 | 2% |
| 60 | OLI | 146/38 | 2017-2-1 | 1% |
| 61 | OLI | 151/33 | 2017-4-9 | 0% |
| 62 | OLI | 144/35 | 2017-7-29 | 1% |
| 63 | OLI | 133/37 | 2017-11-5 | 1% |
| 64 | OLI | 131/36 | 2018-4-16 | 1% |
| 65 | OLI | 133/38 | 2018-11-8 | 0% |
| 66 | OLI | 137/40 | 2018-12-22 | 0% |
| 67 | OLI | 131/38 | 2019-3-18 | 1% |
| 68 | OLI | 136/33 | 2019-8-28 | 0% |
| 69 | OLI | 151/33 | 2019-9-22 | 0% |
| 70 | OLI | 134/36 | 2019-11-2 | 1% |
| 71 | OLI | 132/38 | 2019-12-6 | 1% |
| 72 | OLI | 135/39 | 2020-1-12 | 1% |
| 73 | OLI | 136/38 | 2020-2-4 | 1% |
| 74 | OLI | 151/33 | 2020-8-23 | 1% |
| 75 | OLI | 149/35 | 2020-12-31 | 1% |
| 76 | OLI | 132/34 | 2021-1-25 | 1% |
| 77 | OLI | 143/34 | 2021-7-17 | 1% |
| 78 | OLI | 151/33 | 2021-9-27 | 0% |
| 79 | OLI | 135/36 | 2021-10-29 | 1% |
| 80 | OLI | 143/36 | 2021-11-22 | 0% |
| 81 | OLI | 151/33 | 2021-12-16 | 1% |

References

- Antonic, O.: Modelling daily topographic solar radiation without site-specific hourly radiation data, *Ecological Modelling*, 113, 31-40, doi: 10.1016/S0304-3800(98)00132-X, 1998.
- Azizi, A. H. and Akhtar, F.: Analysis of spatiotemporal variation in the snow cover in Western Hindukush-Himalaya region, *Geocarto International*, 2021. 1-23, doi: 10.1080/10106049.2021.1939442, 2021.
- Bormann, K. J., McCabe, M. F., and Evans, J. P.: Satellite based observations for seasonal snow cover detection and characterisation in Australia, *Remote Sensing of Environment*, 123, 57-71, doi: 10.1016/j.rse.2012.03.003, 2012.
- Cereceda-Balic, F., Vidal, V., Ruggeri, M. F., and Gonzalez, H. E.: Black carbon pollution in snow and its impact on albedo

- near the Chilean stations on the Antarctic peninsula: First results, *Science of the Total Environment*, 743, 140801, doi:
500 10.1016/j.scitotenv.2020.140801, 2020.
- Chen, X., Long, D., Liang, S., He, L., Zeng, C., Hao, X., and Hong, Y.: Developing a composite daily snow cover extent record over the Tibetan Plateau from 1981 to 2016 using multisource data, *Remote Sensing of Environment*, 215, 284-299, doi: 10.1016/j.rse.2018.06.021, 2018a.
- Chen, X., Long, D., Hong, Y., Hao, X., and Hou, A.: Climatology of snow phenology over the Tibetan plateau for the
505 period 2001-2014 using multisource data, *International Journal of Climatology*, 38, 2718-2729, doi: 10.1002/joc.5455, 2018b.
- Crawford, C. J.: MODIS Terra Collection 6 fractional snow cover validation in mountainous terrain during spring snowmelt using Landsat TM and ETM+, *Hydrological Processes*, 29, 128-138, doi: 10.1002/hyp.10134, 2015.
- Dariane, A. B., Khoramian, A., and Santi, E.: Investigating spatiotemporal snow cover variability via cloud-free MODIS
510 snow cover product in Central Alborz Region, *Remote Sensing of Environment*, 202, 152-165, doi: 10.1016/j.rse.2017.05.042, 2017.
- Dong, C.: Remote sensing, hydrological modeling and in situ observations in snow cover research: A review, *Journal of Hydrology*, 561, 573-583, doi: 10.1016/j.jhydrol.2018.04.027, 2018.
- Dong, C. and Menzel, L.: Producing cloud-free MODIS snow cover products with conditional probability interpolation and
515 meteorological data, *Remote Sensing of Environment*, 186, 439-451, doi: 10.1016/j.rse.2016.09.019, 2016.
- Gao, J., Williams, M. W., Fu, X., Wang, G., and Gong, T.: Spatiotemporal distribution of snow in eastern Tibet and the response to climate change, *Remote Sensing of Environment*, 121, 1-9, doi: 10.1016/j.rse.2012.01.006, 2012.
- ~~Gladkova, I., Grossberg, M., Bonev, G., Romanov, P., and Shahriar, F.: Increasing the accuracy of MODIS/Aqua snow product using quantitative image restoration technique, *IEEE Geoscience and Remote Sensing Letters*, 9, 740-743, doi: 10.1109/lgrs.2011.2180505, 2012.~~
520
- Gul, M. S., Muneer, T., and Kambezidis, H. D.: Models for obtaining solar radiation from other meteorological data, *Solar Energy*, 64, 99-108, doi: 10.1016/S0038-092x(98)00048-6, 1998.
- Hall, D. K., Riggs, G., and Salomonson, V. V.: Development of methods for mapping global snow cover using Moderate Resolution Imaging Spectroradiometer (MODIS) data, *Remote Sensing of Environment*, 54, 127-140, doi: 10.1016/0034-
525 4257(95)00137-P, 1995.
- Hall, D.K., Riggs, G.A.: MODIS/Aqua Snow Cover Daily L3 Global 500m SIN Grid, Version 6, available at: doi: 10.5067/MODIS/MOD10A1.006 (last access: 1 March 2022), 2016a.
- Hall, D.K., Riggs, G.A.: MODIS/Aqua Snow Cover Daily L3 Global 500m SIN Grid, Version 6, available at: doi: 10.5067/MODIS/MYD10A1.006 (last access: 1 March 2022), 2016b.
- 530 Hall, D. K. and G. A. Riggs.: MODIS/Terra Snow Cover 8-Day L3 Global 500m SIN Grid, Version 61. available at: doi: 10.5067/MODIS/MOD10A2.061, 2021.
- Hou, J., Huang, C., Zhang, Y., Guo, J., and Gu, J.: Gap-filling of MODIS fractional snow cover products via non-local

- spatio-temporal filtering based on machine learning techniques, *Remote Sensing*, 11, doi: 10.3390/rs11010090, 2019.
- Huang, G., Li, Z., Li, X., Liang, S., Yang, K., Wang, D., and Zhang, Y.: Estimating surface solar irradiance from satellites: 535 Past, present, and future perspectives, *Remote Sensing of Environment*, 233, doi: 10.1016/j.rse.2019.111371, 2019.
- Huang, K., Zhang, Y., Tagesson, T., Brandt, M., Wang, L., Chen, N., Zu, J., Jin, H., Cai, Z., Tong, X., Cong, N., and Fensholt, R.: The confounding effect of snow cover on assessing spring phenology from space: A new look at trends on the Tibetan Plateau, *Science of The Total Environment*, 756, doi: 10.1016/j.scitotenv.2020.144011, 2021.
- Huang, X., Liang, T., Zhang, X., and Guo, Z.: Validation of MODIS snow cover products using Landsat and ground 540 measurements during the 2001–2005 snow seasons over northern Xinjiang, China, *International Journal of Remote Sensing*, 32, 133-152, doi: 10.1080/01431160903439924, 2011.
- Huang, Y., Chen, Z., Wu, B., Chen, L., Mao, W., Zhao, F., Wu, J., Wu, J., and Yu, B.: Estimating roof solar energy potential in the downtown area using a GPU-accelerated solar radiation model and airborne LiDAR data, *Remote Sensing*, 7, 17212-17233, doi: 10.3390/rs71215877, 2015.
- 545 Huang, Y., Liu, H., Yu, B., Wu, J., Kang, E. L., Xu, M., Wang, S., Klein, A., and Chen, Y.: Improving MODIS snow products with a HMRF-based spatio-temporal modeling technique in the Upper Rio Grande Basin, *Remote Sensing of Environment*, 204, 568-582, doi: 10.1016/j.rse.2017.10.001, 2018.
- Huang, Y., Song, Z. C., Yang, H. X., Yu, B. L., Liu, H. X., Che, T., Chen, J., Wu, J. P., Shu, S., Peng, X. B., Zheng, Z. J., and Xu, J. H.: Snow cover detection in mid-latitude mountainous and polar regions using nighttime light data, *Remote Sensing of 550 Environment*, 268, doi: 10.1016/j.rse.2021.112766, 2022.
- Huang, Y., and Xu, J. H.: Daily cloud-free snow cover products for Tibetan Plateau from 2002 to 2021, National Tibetan Plateau Data Center, doi: 10.11888/Cryos.tpdc.272204, 2022.
- Hussainzada, W., Lee, H. S., Vinayak, B., and Khpalwak, G. F.: Sensitivity of snowmelt runoff modelling to the level of cloud coverage for snow cover extent from daily MODIS product collection 6, *Journal of Hydrology: Regional Studies*, 36, 555 doi: 10.1016/j.ejrh.2021.100835, 2021.
- Immerzeel, W. W., van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the asian water towers, *Science*, 328, 1382-1385, doi: 10.1126/science.1183188, 2010.
- Jing, Y.H., Shen, H.F., Li, X.H., and Guan, X.B.: A two-stage fusion framework to generate a spatio-temporally continuous MODIS NDSI product over the Tibetan Plateau, *Remote Sensing*, 11, doi: 10.3390/rs11192261, 2019.
- 560 Kilpys, J., Pipiraitė-Januškienė, S., and Rimkus, E.: Snow climatology in Lithuania based on the cloud-free moderate resolution imaging spectroradiometer snow cover product, *International Journal of Climatology*, 40, 4690-4706, doi: 10.1002/joc.6483, 2020.
- Klein, A.: Validation of daily MODIS snow cover maps of the Upper Rio Grande River Basin for the 2000–2001 snow year, *Remote Sensing of Environment*, 86, 162-176, doi: 10.1016/s0034-4257(03)00097-x, 2003.
- 565 Kumar, L., Skidmore, A. K., and Knowles, E.: Modelling topographic variation in solar radiation in a GIS environment, *International Journal of Geographical Information Science*, 11, 475-497, doi: 10.1080/136588197242266, 1997.

- Li, C., Su, F., Yang, D., Tong, K., Meng, F., and Kan, B.: Spatiotemporal variation of snow cover over the Tibetan Plateau based on MODIS snow product, 2001-2014, *International Journal of Climatology*, 38, 708-728, doi: 10.1002/joc.5204, 2018.
- Li, M., Zhu, X., Li, N., and Pan, Y.: Gap-filling of a MODIS Normalized Difference Snow Index product based on the similar pixel selecting algorithm: a case study on the Qinghai-Tibetan Plateau, *Remote Sensing*, 12, doi: 570 10.3390/rs12071077, 2020.
- Li, X., Jing, Y., Shen, H., and Zhang, L.: The recent developments in cloud removal approaches of MODIS snow cover product, *Hydrology and Earth System Sciences*, 23, 2401-2416, doi: 10.5194/hess-23-2401-2019, 2019a.
- Li, Y., Chen, Y., and Li, Z.: Developing daily cloud-free snow composite products from MODIS and IMS for the Tianshan 575 mountains, *Earth and Space Science*, 6, 266-275, doi: 10.1029/2018ea000460, 2019b.
- Liang, H., Huang, X., Sun, Y., Wang, Y., and Liang, T.: Fractional snow-cover mapping based on MODIS and UAV data over the Tibetan Plateau, *Remote Sensing*, 9, doi: 10.3390/rs9121332, 2017.
- Liang, T., Huang, X., Wu, C., Liu, X., Li, W., Guo, Z., and Ren, J.: An application of MODIS data to snow cover monitoring in a pastoral area: A case study in Northern Xinjiang, China, *Remote Sensing of Environment*, 112, 1514-1526, doi: 580 10.1016/j.rse.2007.06.001, 2008.
- Liu, C., Li, Z., Zhang, P., Zeng, J., Gao, S., and Zheng, Z.: An assessment and error analysis of MOD10A1 snow product using Landsat and ground observations over China during 2000–2016, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 13, 1467-1478, doi: 10.1109/jstars.2020.2983550, 2020a.
- Liu, Y., Chen, X., Hao, J.-S., and Li, L.-h.: Snow cover estimation from MODIS and Sentinel-1 SAR data using machine 585 learning algorithms in the western part of the Tianshan Mountains, *Journal of Mountain Science*, 17, 884-897, doi: 10.1007/s11629-019-5723-1, 2020b.
- Muhammad, S. and Thapa, A.: An improved Terra-Aqua MODIS snow cover and Randolph glacier inventory 6.0 combined product (MOYDGL06*) for high-mountain Asia between 2002 and 2018, *Earth System Science Data*, 12, 345-356, doi: 10.5194/essd-12-345-2020, 2020.
- 590 Parajka, J. and Blöschl, G.: Spatio-temporal combination of MODIS images - potential for snow cover mapping, *Water Resources Research*, 44, doi: 10.1029/2007wr006204, 2008.
- Parajka, J., Holko, L., Kostka, Z., and Blöschl, G.: MODIS snow cover mapping accuracy in a small mountain catchment – comparison between open and forest sites, *Hydrology and Earth System Sciences*, 16, 2365-2377, doi: 10.5194/hess-16-2365-2012, 2012.
- 595 Qiu, J.: China: The third pole, *Nature*, 454, 393-396, doi: 10.1038/454393a, 2008.
- Qiu, Y., Zhang, H., Chu, d., and Xuecheng, Z.: Cloud removing algorithm for the daily cloud free MODIS-based snow cover product over the Tibetan Plateau, *Journal of Glaciology and Geocryology*, 39(3), 515-526, doi: 10.7522/j.issn.1000-0240.2017.0058, 2017.
- 600 Richiardi, C., Blonda, P., Rana, F. M., Santoro, M., Tarantino, C., Vicario, S., and Adamo, M.: A revised snow cover algorithm to improve discrimination between snow and clouds: a case study in Gran Paradiso National Park, *Remote*

- Sensing, 13, doi: 10.3390/rs13101957, 2021.
- Riggs, G. A., Hall, D. K., and Román, M. O.: Overview of NASA's MODIS and Visible Infrared Imaging Radiometer Suite (VIIRS) snow-cover Earth System Data Records, *Earth System Science Data*, 9, 765-777, doi: 10.5194/essd-9-765-2017, 2017.
- 605 Riggs, G. A., Hall, D. K., and Román, M. O.: MODIS Snow Products Collection 6.1 User Guide, available at: https://modis-snow-ice.gsfc.nasa.gov/uploads/snow_user_guide_C6.1_final_revised_april.pdf (last access: 1 March 2022), 2019.
- Salomonson, V. V. and Appel, I.: Estimating fractional snow cover from MODIS using the normalized difference snow index, *Remote Sensing of Environment*, 89, 351-360, doi: 10.1016/j.rse.2003.10.016, 2004.
- Tang, Z., Wang, J., Li, H., and Yan, L.: Accuracy validation and cloud obscuration removal of MODIS fractional snow cover products over Tibetan Plateau, *Remote Sensing Technology and Application*, 28(3), 423-430, doi: 10.16089/j.cnki.1008-2786.000237, 2013.
- [Teilet, P. M., Guindon, B., Goodenough, D. G.: On the slope-aspect correction of multispectral scanner data, *Canadian Journal of Remote Sensing*, 8\(2\), 1537-1540, doi: 10.1080/07038992.1982.10855028, 1982.](#)
- Tran, H., Nguyen, P., Ombadi, M., Hsu, K. L., Sorooshian, S., and Qing, X.: A cloud-free MODIS snow cover dataset for the contiguous United States from 2000 to 2017, *Scientific Data*, 6, 180300, doi: 10.1038/sdata.2018.300, 2019.
- 615 Wang, G., Jiang, L., Shi, J., Liu, X., Yang, J., and Cui, H.: Snow-covered area retrieval from Himawari-8 AHI imagery of the Tibetan Plateau, *Remote Sensing*, 11, doi: 10.3390/rs11202391, 2019.
- Wang, X., Wu, C., Peng, D., Gonsamo, A., and Liu, Z.: Snow cover phenology affects alpine vegetation growth dynamics on the Tibetan Plateau: Satellite observed evidence, impacts of different biomes, and climate drivers, *Agricultural and Forest Meteorology*, 256-257, 61-74, doi: 10.1016/j.agrformet.2018.03.004, 2018.
- 620 [Wu, Z., Jiang, Z., Li, J., Zhong, S. and Wang, L.: Possible association of the western Tibetan plateau snow cover with the decadal to interdecadal variations of northern China heatwave frequency, *Climate Dynamics*, 39\(9-10\), 2393-2402, 2012.](#)
- Wang, X., Xie, H., Liang, T., and Huang, X.: Comparison and validation of MODIS standard and new combination of Terra and Aqua snow cover products in northern Xinjiang, China, *Hydrological Processes*, 23, 419-429, doi: 10.1002/hyp.7151, 2009.
- 625 Xiao, X., Liang, S., He, T., Wu, D., Pei, C., and Gong, J.: Estimating fractional snow cover from passive microwave brightness temperature data using MODIS snow cover product over North America, *The Cryosphere*, 15, 835-861, doi: 10.5194/tc-15-835-2021, 2021.
- [Xu, W. F., Ma, H. Q., Wu, D. H., and Yuan, W. P.: Assessment of the daily cloud-free MODIS snow-cover product for monitoring the snow-cover phenology over the Qinghai-Tibetan Plateau, *Remote Sensing*, 9, ARTN 585, doi: 10.3390/rs9060585, 2017.](#)
- 630 Yang, J., Jiang, L., Ménard, C. B., Luo, J., Lemmetyinen, J., and Pulliainen, J.: Evaluation of snow products over the Tibetan Plateau, *Hydrological Processes*, 29, 3247-3260, doi: 10.1002/hyp.10427, 2015.
- Yang, K., Wu, H., Qin, J., Lin, C., Tang, W., and Chen, Y.: Recent climate changes over the Tibetan Plateau and their impacts

- 635 on energy and water cycle: A review, *Global and Planetary Change*, 112, 79-91, doi: 10.1016/j.gloplacha.2013.12.001, 2014.
- Yao, T., Xue, Y., Chen, D., Chen, F., Thompson, L., Cui, P., Koike, T., Lau, W. K. M., Lettenmaier, D., Mosbrugger, V., Zhang, R., Xu, B., Dozier, J., Gillespie, T., Gu, Y., Kang, S., Piao, S., Sugimoto, S., Ueno, K., Wang, L., Wang, W., Zhang, F., Sheng, Y., Guo, W., Ailikun, Yang, X., Ma, Y., Shen, S. S. P., Su, Z., Chen, F., Liang, S., Liu, Y., Singh, V. P., Yang, K., Yang, D., Zhao, X., Qian, Y., Zhang, Y., and Li, Q.: Recent third pole's rapid warming accompanies cryospheric melt and
- 640 water cycle intensification and interactions between monsoon and environment: multidisciplinary approach with observations, Modeling, and Analysis, *Bulletin of the American Meteorological Society*, 100, 423-444, doi: 10.1175/bams-d-17-0057.1, 2019.
- You, Q., Wu, T., Shen, L., Pepin, N., Zhang, L., Jiang, Z., Wu, Z., Kang, S., and AghaKouchak, A.: Review of snow cover variation over the Tibetan Plateau and its influence on the broad climate system, *Earth-Science Reviews*, 201, doi: 10.1016/j.earscirev.2019.103043, 2020.
- 645 Yu, J., Zhang, G., Yao, T., Xie, H., Zhang, H., Ke, C., and Yao, R.: Developing daily cloud-free snow composite products from MODIS Terra–Aqua and IMS for the Tibetan Plateau, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 2171-2180, doi: 10.1109/tgrs.2015.2496950, 2016.
- Zhang, H., Zhang, F., Zhang, G., Yan, W., and Li, S.: Enhanced scaling effects significantly lower the ability of MODIS
- 650 normalized difference snow index to estimate fractional and binary snow cover on the Tibetan Plateau, *Journal of Hydrology*, 592, doi: 10.1016/j.jhydrol.2020.125795, 2021.
- Zheng, Z. and Cao, G.: Snow cover dataset based on multi-source remote sensing products blended with 1km spatial resolution on the Qinghai-Tibet Plateau (1995-2018), National Tibetan Plateau Data Center, doi: 10.11888/Snow.tpd.270102, 2019.