



How to learn more about hydrological conditions and phytoplankton dynamics and diversity in the eastern English Channel and the southern bight of the North Sea? the SRN data set (1992-2021).

Lefebvre Alain¹, Devreker David¹

5 ¹ Ifremer, Unité Littoral, Laboratoire Environnement et Ressources, 150 quai Gambetta, BP 699, 62321 Boulogne-sur-mer, France.

Correspondence to: Alain Lefebvre (alain.lefebvre@ifremer.fr)

Abstract. This article describes the historical data series produced by the SRN network managed by Ifremer. Since 1992, the SRN ('Suivi Régional des Nutriments' in French; Regional Nutrients Monitoring Programme) network has been analysing phytoplankton species and measuring twelve physicochemical parameters at ten different stations distributed along three different transects located in the Eastern English Channel, and the Southern Bight of the North Sea. The SRN collects a maximum of 184 samples per year and detects up to 281 taxa, including harmful algal bloom species (HABs), with a bi-weekly to monthly sampling frequency (depending on the location and the season). The objectives of this monitoring program are to assess the influence of continental inputs on the marine environment, and their implications on possible eutrophication processes. It also aims to estimate the effectiveness of development and management policies in the marine coastal zone. The regular acquisition of data allows the establishment of a long-term monitoring of the evolution of coastal water quality, as well as the observation of the consequences of large-scale alterations and modifications that are more related to regional anthropogenic activities. This paper provides an overview of the main characteristics of SRN data (descriptive statistics and data series main patterns) as well as an analysis of temporal trends using specific numerical tools available as an R package to help make use of such data. Main results of several research projects are also highlighted providing the readers with examples of what is doable with such a data set.

1 Introduction

25 Phytoplankton contribute to the biological pump that regulates the flow of carbon dioxide. This compartment is critically important as it forms the basis of marine food webs. Understanding the structure of this community is essential for any assessment of marine diversity (Garmendia et al., 2013). Hence, maintaining ecosystem goods and services is partly linked to



phytoplankton dynamics. There are several thousand phytoplankton species on the planet: the vast majority are completely harmless, but a few hundred can proliferate significantly, forming red, brown, or green waters, a few dozen are toxic to marine
30 fauna or humans through shellfish consumption (toxin bioaccumulation process), and others cause excessive organic matter inputs, affecting water quality. While some taxa contribute naturally to energy transfers to higher trophic levels, others are responsible for the development of toxic algal blooms that limit grazing (Nejstgaard et al., 2007). Indeed, Harmful Algal Blooms (HAB) can reduce benthic and pelagic biodiversity, and disrupt marine ecosystems (Rousseau et al., 1990). They also may lead to the degradation of seafood quality, rendering it unsuitable for human consumption (Berdalet et al., 2016). As a
35 result, any imbalance within the community, such as favouring dinoflagellates over diatoms, will have major effects on the biodiversity of higher trophic levels, as well as on the quality of the ecosystem in general. Consequently, such modifications of ecosystem goods and services will lead to several socio-economic consequences.

The observation and monitoring of ecosystems quality is often accomplished by setting up networks to monitor hydrological and biological parameters, which constitute the essential foundation for the overall characterization of the aquatic ecosystem.
40 As a result, the environmental descriptors are numerous, and are linked by cause and effect relationships; others are directly influenced by anthropogenic activities. Their ability to rapidly respond to changes in their environment allows the evaluation of a response to different sources of pressure. Phytoplankton is regularly used as a water quality indicator for directives and conventions (e.g. the Water Framework Directive (EU WFD, 2000), the Marine Strategy Framework Directive (MSFD - 2008/56/EC) (European Commission MSFD, 2008) and the Oslo and Paris Convention (OSPAR Commission, 2013)). The
45 metrics created for their purposes frequently employ phytoplankton biomass, abundance, and composition, as well as the frequency and intensity of blooms.

For a long time, the English Channel and the southern Bight of the North Sea have been subjected to intense anthropogenic pressure (fishing, tourism and leisure activities, marine aggregate extraction, maritime traffic, major port areas, degraded estuarine areas, off-shore wind turbine projects, etc.), with considerable economic stakes and subject to a diverse range of
50 users with frequently antagonistic interests (Carpentier et al., 2009). The English Channel ecoregion is an epicontinental sea that serves as a transition zone between the Atlantic and North Sea water bodies. The region has a temperate oceanic climate, influenced by wet and cold atmospheric currents coming from the Atlantic or more sporadically from the North Sea. It is characterized by its megatidal hydrodynamic regime, induced by tidal currents. This intense hydrodynamics influences the nature, distribution and dynamics of sediments, as well as the structure, distribution, dynamics and functioning of biological
55 compartments. In addition, a maritime zone under the impact of freshwater (ROFI: Region Of Freshwater Influence) arises along the French shores of the eastern Channel, mostly from the Seine river, and whose structure is maintained by contributions from other northern rivers (Somme, Canche, Authie). This structure will play an essential role in controlling the exchange of inert or living particles, organisms between the coast and the open sea (Brylinski et al., 1991).

Phytoplankton data allow us to address harmful algal blooms (HABs), particularly *Phaeocystis globosa* and *Pseudo-nitzschia*
60 ones. Since the 1970s, abnormal increase in intensity and duration of *Phaeocystis globosa* blooms, a naturally occurring taxon in the North Sea, have inspired scientific teams to launch research projects dedicated to this taxon in the context of excessive



nutrient enrichment of marine waters (Admiraal and Venekamp, 1986; Cadée and Hegeman, 1986; Eberlein et al., 1985; Lancelot and Mathot, 1985; Lancelot et al., 1997; Lefebvre et al., 2011; Lefebvre and Dezécache, 2020; Rahmel et al., 1995). *Phaeocystis globosa* was then identified as a potentially harmful taxon (in the HAB sense), not just due to its toxicity, but rather to the huge biomass created during its blooms and its consequences. *P. globosa* is also likely to produce precursors of dimethyl sulphide (DMS), a gas with a sulphurous and unpleasant odour. It was reported that DMS could cause respiratory, skin and ocular irritation in humans. It can also be found in the atmosphere and can be favourable to the formation of acid rain. At the end of its very complex, polymorphic development cycle, *P. globosa* appears in colonial form. These colonies are loaded with mucco-polysaccharides. They will break up in response to internal factors (ageing, lysis) and/or external factors (turbulence) and provoke by emulsion the accumulation of a thick, odorous foam on the coast. The *Pseudo-nitzschia* complex contains species that can produce an amnesic phycotoxin based on domoic acid. During *P. globosa* blooms, needles formed by *Pseudo-nitzschia* complex cells can stick into *P. globosa* colonies and form structures that irritate filter feeders. The lesions caused by these structures could promote viral and bacterial infections in fish.

Prior to 1992, french monitoring of phytoplankton populations and associated environmental factors in the English Channel and the southern bight of the North Sea was done episodically, via the RNO (Réseau National d'Observation) or the RNC (Réseau National de Contrôle). The Artois-Picardy Water Agency and Ifremer established the SRN (Suivi Régional des Nutriments) in 1992 in response to the need for precise monitoring of nutrient concentration variations over a longer period, as well as the response of the environment to this pressure, particularly in terms of phytoplankton development. The objective of this monitoring is to evaluate the influence of continental inputs on the marine environment (nitrogen, phosphate, silicate) and their consequences on possible eutrophication processes. It also aims to estimate the efficiency of wastewater treatment plants and policies for the development and management of the coastal zone and more generally the possible elimination of such discharges. The regular acquisition of data allows the establishment of a long-term monitoring of the coastal waters along three transects located off Dunkerque, Boulogne-sur-Mer and in the Bay of Somme, making it possible to pretend to be able to deconvolute the effects of large-scale changes from those linked to more local anthropic activities (Lefebvre et al., 2011). Since 1992, the SRN dataset has included long-term time series on marine phytoplankton and physico-chemical measurements along the east coast of the English Channel and the French section of the southern bight of the North Sea (Dunkerque). SRN data are complementary to REPHY and REPHYTOX datasets (PHYTOBS, 2021; REPHY, 2021; REPHYTOX, 2021). Phytoplankton data cover microscopic taxonomic identifications and counts, up to the species level, and pigments measures (Chlorophyll-*a* and pheopigment). Physico-chemical measurements include temperature, salinity, turbidity, suspended matters (organic, mineral), dissolved oxygen, and dissolved inorganic nutrients (ammonium, nitrite+nitrate, phosphate, silicate).

2 Objectives

The objective of this paper is to present the SRN dataset, from the sampling strategy to data collection (including associated Quality Assurance/ Quality Controls Steps) and data investigation and storage. The characteristics of the different datasets as



well as a general interpretation of their variability will be presented. Based on existing valorisations of the SRN dataset, we
95 will demonstrate that these data are of relevance not just for furthering understanding in marine phytoplankton ecology, but
also for public policy needs such as assessment of environmental or ecological status as requested by EU directives or Regional
Sea Conventions. We also propose some numerical tools based on an R package available for the scientific community and
developed specifically to rapidly process such data and therefore to valorise the findings.


3 Materials and Methods

100 3.1 Sampling area and sampling stations

The English Channel has a macro-tidal regime that ranges from 3m to 9m in the Dover Strait during neap tide and spring tide,
respectively. This regime produces high tidal currents that are basically parallel to the shore, as well as a north-eastward tidal
residual current from the English Channel to the North Sea. Along the French coast, fluvial supplies dispersed from the Bay
of Seine to the Cape Gris-Nez form a coastal water mass that floats nearshore, protected from the open sea by a frontal region
105 (Brylinski et al., 1991). Exchanges between inshore and offshore water masses, as well as particle and nutrient transport, are
tide-dependent and are larger during the neap than during the spring tide. This may seem counter-intuitive, but during the neap
tide, the frontal structure between inshore and offshore waters is more sloped from the vertical resulting in a greater surface of
exchange between coastal and offshore waters. This leads to enhanced exchange possibilities between the two water masses
(Brylinski et al., 1991).

110 Three sampling areas along an inshore offshore transect were studied by Ifremer from 1992 to 2021 (still ongoing) within the
frame of the SRN (“Suivi Régional des Nutriments”, i.e. Regional Nutrient Monitoring Programme), two of which are located
in the Eastern English Channel and one in the southern bight of the North Sea (Figure 1):

- 1) Boulogne-sur-Mer harbour (by the Dover Strait), a coastal zone separated from the open sea by a frontal area (Station name:
BL1, BL2, BL3);
- 115 2) Bay of Somme, the second ranked estuarine system after the Seine estuary on the French coasts of the English Channel
(Station name: Bif, Mimer, Atso, Mer2);
- 3) Dunkerque harbour, a shallow well-mixed coastal zone near the frontier between France and Belgium (Station name: DK1,
DK3, DK4).



120 **Figure 1. Map representing the different transects and sampling stations (Dunkerque, Boulogne and Bay of Somme) of the SRN network. The boundaries of the coastal and transitional water bodies of the European Water Framework Directive and the French Economic Exclusive Zone are also indicated.**

The main environmental characteristics of the areas are summarized in table 1.

125 **Table 1. Main environmental characteristics of the three ecosystems/transects: Dunkerque, Boulogne-sur-Mer and the Bay of Somme.**

	Dunkerque	Boulogne-sur-Mer	Bay of Somme
Tidal regime	macrotidal	macrotidal	macrotidal
Main characteristic	Shallow water (< 30 m)	Under the influence of a frontal structure	Estuary with medium turbidity
Main River	Aa	Liane	Somme



Main pressures	Industrial activities	Agricultural activities	Agricultural activities, shellfish aquaculture
Length of the main river (km)	89	37	263
Watershed (km ²)	1215	244	6550
Mean river Flow (m ³ s ⁻¹)	10	3	35
Sampling stations	Point 1 Dunkerque (DK1) Point 3 SRN Dunkerque (DK3) Point 4 SRN Dunkerque (DK4)	Point 1 Boulogne (BL1) Point 2 SRN Boulogne (BL2) Point 3 SRN Boulogne (BL3)	Atso Mimer Bif SRN Somme mer 2 (Mer2)

3.2. Hydrology

From March to June, water samples were taken twice a month from subsurface waters using a 5 l Niskin Bottle, and once a month the rest of the year. The approach used for chlorophyll-*a*, ammonia, nitrite, nitrate, phosphate and silicate analyses is provided by Aminot & K  rouel (2004). Chlorophyll-*a* concentrations were estimated by spectrophotometry after filtration through glass fiber filters and extraction with 90 % acetone.

An accurate test of nutrient limitation requires detailed measurements of algal growth under experimental nutrient addition (D'Elia et al., 1986). Nevertheless, in order to determine the potential limitation of primary production by nutrient availability when such experiment results are not available, the standard molar ratio for dissolved inorganic nitrogen (DIN = ammonium + nitrite + nitrate), phosphate and silicate were calculated and compared, according to the references of Redfield et al. (1963) and Brzezinski (1985) for the composition of the biogenic matter (Si: N: P = 16: 16: 1).

3.3. Phytoplankton

Phytoplankton samples have been collected along transects and were preserved with an acid lugol solution (0.25%). Sub-samples of 10 ml were settled for 24 hours in a counting chamber according to the Uterm  hl (1958). Cell enumerations were performed by inverted microscopy using a microscope within a month after the sample collection to prevent any significant changes in phytoplankton size and abundance. Except for *Phaeocystis globosa* enumeration, over 400 phytoplankton cells in each sample were counted with a 20X Plan Ph1 0.5NA objective, resulting in an error of 10%. For assessment of *P. globosa* counts, only the total number of cells are computed. A minimum of 50 solitary cells were enumerated from several randomly chosen fields (10 to 30) with a 40X Plan Ph2 0.75NA. Abundance of cells in colony was determined using a relationship between colony biovolume and cell number defined by Rousseau et al. (1990).

The total number of sample collected for complete determination of phytoplankton community since the creation of the SRN is shown in table 2.



Table 2. Number of samples of phytoplankton and number of species for each SRN stations on the investigated time span 1992-2021.

	Nb of sample	Nb of species
DK1	370	212
DK3	310	196
DK4	296	186
BL1	463	209
BL2	387	198
BL3	381	184
Bif	399	202
Mimer	257	188
Atso	449	214
Mer2	375	188
Total	3687	280

150 4. Database

To manage coastal monitoring data, Ifremer has developed the Quadrige² information system (<https://wwz.ifremer.fr/envlit/Quadrige-la-base-de-donnees>; last access on 28 April 2022), which combines a database with a variety of interpretation and information product development tools. As an information system, Quadrige² plays a crucial role in: (1) storing basic monitoring data, such as the results of analyses from all monitoring networks, in a safe, optimal, supervised, and scalable manner, and (2) interpreting and valuing the data. Once the data has been stored and a quality level assigned to it, it is ready for use in a wide range of applications. As a result, this system is the required link for monitoring data between data collection in the field and its availability in various formats. Quadrige² has been approved as the national reference information system for coastal waters by the French Ministry in charge of the Environment.

5. Quality Control

160 5.1. Data Validation

The data are collected in the field and/or laboratory, then enter into the Quadrige² database using an application with the same name. The control entails modifying the data entered (results and metadata) to ensure that it is consistent with the bench book (or field sheets). After this check is completed and any necessary corrections are made, the data are validated:

- Confirmation of the technical validity of the data which corresponds to the result of the analysis,
- 165 - Data locking, so that it can no longer be edited, even by the person who entered it,



- Data distribution: verified data may be taken and disseminated by all Quadriga² users who have access to the database.

5.2. Data Qualification

This initial round of data verification is followed by the qualifying procedure, which corresponds to:

- the search for questionable or even scientifically aberrant data,
- 170 - the correction of data when possible,
- the attribution of a qualification level to the data, which is:
 - good: data analysis are scientifically relevant,
 - doubtful: the data may be false: taking it into account may bias the analysis that will be made,
 - False: the data should not be included in the analysis because they are aberrant or present a problem (e.g., bad
- 175 analytical series and impossible to repeat).

The level of qualification corresponds to the level of confidence in the data. It determines the way in which the data is distributed (only data qualified as "good" and "doubtful" are distributed), and how it is used in specific data processing. The qualification is broken down into two main steps:

- 1) An "automatic" qualification that consists of looking for "gross" and easily identifiable errors,
- 180 2) An "expert" qualification, which consists in highlighting statistically aberrant data via adapted methods (time series analysis, statistical tests...). Only data qualified as "good" or "doubtful" from the previous step are used for the expert qualification.

6. Data Summary

Table 3 represents the descriptive statistics obtained for each physicochemical and biological parameter (excluding phytoplankton) and for each SRN station. For each of these series, the monotonic trend is estimated using a non-parametric
185 method, Mann-Kendall Seasonally adjusted autocorrelated series test (Devreker and Lefebvre, 2014).

Table 3. Statistical summary (minimum, first and third quantiles, mean, median, maximum, length of the data series) for the physico-chemical and biological variables collected within the SRN monitoring programme on the time span 1992-2021 and for coastal stations only (DK1, BL1, Atso). Increasing or decreasing monotonous¹ / non-monotonous² trends are indicated in the Sign. Trend column (1. One orange (green) arrow for an increasing (decreasing) monotonous trend - 2. Two arrows for a non-monotonous trend (shift in the time series) – Grey arrow indicates no significant trend).
190



Stations	Parameters	Min	1st Qu	Median	Mean	3rd Qu	Max	N	Sign. Trend
DK1	Temperature (°C)	1.00	8.50	12.10	12.36	16.59	21.70	378	→
	Chlorophyll- <i>a</i> (µg.l ⁻¹)	0.240	2.498	4.495	6.737	7.925	53.180	396	→
	Salinity	31.10	33.70	34.24	34.08	34.60	35.50	398	→
	SM (mg.l ⁻¹)	1.60	5.70	9.60	13.33	17.00	95.20	373	→
	NO ₃ +NO ₂ (µmol.l ⁻¹)	0.110	0.600	2.320	8.167	14.055	54.770	384	↓
	NH ₄ (µmol.l ⁻¹)	0.110	0.450	1.685	2.456	3.547	29.400	386	↓
	Oxygen (mg.l ⁻¹)	6.910	8.020	8.805	9.190	9.875	19.200	202	↓
	PO ₄ (µmol.l ⁻¹)	0.0100	0.1600	0.3900	0.5144	0.6900	9.8000	385	↓
	SIOH (µmol.l ⁻¹)	0.100	1.055	3.000	4.963	6.553	35.200	386	↗
BL1	Temperature (°C)	2.10	8.85	12.50	12.66	16.70	22.10	463	→
	Chlorophyll- <i>a</i> (µg.l ⁻¹)	0.010	1.708	3.375	5.117	7.060	29.600	472	↗
	Salinity	29.14	33.60	34.20	33.93	34.50	35.30	469	↗
	SM (mg.l ⁻¹)	0.050	3.400	5.650	8.237	9.575	46.400	446	→
	NO ₃ +NO ₂ (µmol.l ⁻¹)	0.1500	0.5775	2.0600	6.6585	11.2825	43.8100	450	→
	NH ₄ (µmol.l ⁻¹)	0.060	0.460	1.040	1.440	1.942	10.200	450	↓
	Oxygen (mg.l ⁻¹)	6.260	7.925	8.615	8.919	9.815	12.500	230	→
	PO ₄ (µmol.l ⁻¹)	0.0500	0.1225	0.2700	0.3987	0.5650	3.1000	454	↓
	SIOH (µmol.l ⁻¹)	0.100	0.465	1.640	3.173	4.303	19.010	452	↗
At so	Temperature (°C)	2.00	9.00	13.00	13.03	17.40	22.10	432	→
	Chlorophyll- <i>a</i> (µg.l ⁻¹)	0.210	2.990	5.780	8.269	10.680	58.530	434	→
	Salinity	26.00	32.20	33.20	32.74	33.70	35.10	438	↗
	SM (mg.l ⁻¹)	0.90	7.00	13.65	19.66	26.23	167.00	428	↓
	NO ₃ +NO ₂ (µmol.l ⁻¹)	0.100	1.645	6.820	11.972	19.725	56.860	423	→
	NH ₄ (µmol.l ⁻¹)	0.020	0.370	0.900	1.998	2.490	30.700	421	↓
	Oxygen (mg.l ⁻¹)	4.060	8.162	9.055	9.256	10.012	13.980	216	→
	PO ₄ (µmol.l ⁻¹)	0.0300	0.1300	0.2700	0.4167	0.6025	3.0300	424	↓
	SIOH (µmol.l ⁻¹)	0.060	1.030	3.570	6.099	9.910	41.000	423	→

The seasonal variability of phytoplankton populations corresponds to maximum abundance in spring and then a decrease in winter. This trend can be variable depending on the sites (different hydrodynamical conditions) and the environmental characteristics encountered (luminosity, nutrient inputs, etc.). The water masses sampled appear to be poorly structured vertically, while significant coastal to offshore gradients are evident. Thus, the general dynamics of the Eastern Channel-



Southern Bight of the North Sea ecosystem represents a classical functioning of a temperate system (Wafar et al., 1983; Gentilhomme and Lizon, 1998). The seasonal cycles of nutrients and phytoplankton biomass are well-defined. Inter-annual variability is high. The homogeneity of the sampling conditions makes it possible to avoid normalizing the results of nutrient concentration by salinity for the purpose of inter-site comparison. The analysis of the results shows a monotonic decreasing trend of phosphate concentration for all the studied sites, while silicate concentrations are relatively stable (except when considering inter-annual variability). The dynamics of nitrogen and phytoplankton biomass are far more complicated to handle, and such monotonic trends are not identifiable. These results may alter values of the stoichiometric ratios N: P: Si: N and Si: P (N: total nitrogen; P: Phosphate; Si: silicate) (Redfield et al., 1963; Brzezinski, 1985). Phytoplankton growth appears to be primarily limited by P and Si (for diatoms only). This notion of limiting phytoplankton growth and its consequences on phytoplankton communities deserves particular attention in a system bordered by coastal marine regions where eutrophication problems are of great importance (Bay of Seine and the North Sea) (Lefebvre and Devreker, 2020). Indeed, considerable changes in phytoplanktonic taxonomic productivity, abundance, dominance, and succession have been observed in recent decades as a result of increased human stresses, particularly, nutrient inputs (Billen et al., 2005; Gypens et al., 2013).

For a particular site, phytoplankton counts shows a high interannual and spatial variability. Maximum abundances are measured between March and June, due to the presence of *Phaeocystis globosa*, which dominates the phytoplankton population. Diatoms and dinoflagellates make up the majority of the community over the rest of the year. Between March and June, the prymnesiophyceae *Phaeocystis globosa* is sampled on a regular basis at all sites, and its concentration can exceed one million cells per liter (Figure 2). During the rest of the year, some isolated cells may be detected.

The Bay of Somme area had the highest concentration from 1992 to 2007, with more than $48 \cdot 10^6$ cells per litre at its peak. Likewise, maximum concentrations on the Dunkerque and Boulogne-sur-Mer transects are high, but to a lesser extent, reaching over $29 \cdot 10^6$ and $28 \cdot 10^6$ cells per litre, respectively.

The genera *Alexandrium*, *Dinophysis* and *Pseudo-nitzschia*, which are potentially responsible for the production of PSP (Paralytic Shellfish Poison), DSP (Diarrheic Shellfish Poison) and ASP (Amnesic Shellfish Poison) toxins, respectively, are regularly observed at the monitoring sites. It is worth noting that, even when the cell densities of these toxic genera exceed the alert thresholds, toxin analysis of shellfish collected from blooming regions can be surprisingly negative.




Figure 2. Inter-annual variability of phytoplankton major groups (Prymnesiophyceae, Bacillariophyceae and others phytoplankton) at the three coastal SRN stations: (a) Dunkerque (DK1), (b) Boulogne-sur-Mer (BL1) and (c) Atso in the bay of Somme. Vertical bars represents the relative abundances of these groups (%) and the black line the mean annual total abundance (10^5 cell.L^{-1}).



Figure 3 depicts the seasonal variability of the data. Despite the fact that the series shows drastically varied values depending on the station, the inter-seasonal variability for each of the characteristics depicted remains constant. In fact, nutrient concentrations are at their highest in winter. Phytoplankton that consume these nutrients increase during the spring, which explains why chlorophyll-*a* concentrations (a proxy for phytoplankton biomass) peak reaching its annual maximum, while oxygen concentrations decline reaching their annual minimum. At the end of summer, chlorophyll-*a* concentrations begin to diminish, while nutrients begin to replenish, eventually reaching high quantities in the fall. On the other hand, the temperature shows a classical variability of temperate marine waters with however some extreme values close to 0°C and above 20°C. Figure 4 shows the calculated monthly-scale anomalies for temperature, chlorophyll-*a* concentration, and nutrients at each SRN coastal station.

235





Figure 3. Monthly box and whisker plots of main physico-chemical parameters for the three coastal stations (DK1, BL1 and Atso) of the SRN network for the period 1992-2021.

240



245 **Figure 4. Monthly anomalies calculated for Chlorophyll-*a* and nitrite+nitrate concentrations, temperature and phosphate concentration measured for the three coastal stations (DK1, BL1 and Atso) of the SRN network for the period 1992-2021.**

7. Discussion and conclusion

The SRN data, which have been collected since 1992, represent one of the longest long-term datasets in the English Channel and the southern bight of the North Sea. It enables the study of phytoplankton dynamics and diversity, as well as changes in particular composition in response to anthropogenic and/or climatic change. Indeed, this data have been used by scientists from several backgrounds, and for a variety of research purposes. In the following paragraphs, main outputs from these research activities will be highlighted. This review can help the scientific community interested in this SRN dataset to identify topics that can be addressed or those that have not been addressed. It also allows for new topics to be discussed in more detail since the general patterns have been identified.

250 Lefebvre et al. (2011) proposed a break and trend analysis for nutrient concentrations using a simple and intuitive analytical method called the cumulative sum method (Ibanez et al., 1993). The authors defined the characteristics and patterns of seasonal variations in chlorophyll-*a*, nutrients and phytoplankton. They proposed a classification of years based on whether *Phaeocystis globosa* is dominant or not.



According to Lefebvre et al. (2011) and Hernández-Fariñas et al. (2014), the phytoplankton community in the Eastern English Channel and Southern Bight of the North Sea is dominated by Bacillariophyceae, Dinoflagellates, and Prymnesiophyceae, accounting for 81% of the total abundance. Hernández-Fariñas et al. (2014) estimated that the median contribution of *Phaeocystis globosa* during the period between March and May (data 1992-2014) ranges from 74% to 90% with the highest concentrations found at the coast. They highlighted two main periods with different environmental characteristics regardless of the SRN transect considered: 1992-2001 and 2002-2011. The latter period can be divided into two sub-periods: 2002-2007 and 2008-2011. These results highlighted the existence of a strong temporal structuring of the community under the influence of global and local factors. Globally, the abundance of *Pseudo-nitzschia* increases during the studied periods, while the abundance of other diatoms such as *Guinardia*, *Coscinodiscus* - *Stellarima* decreases. The dinoflagellates *Amphidinium*, *Alexandrium*, and *Polykrikos* mark the second great period. During the second sub-period, however, *Heterocapsa*, *Torodinium*, and *Eutreptiella* (Euglenoid) are widespread. *Melosira* and *Stephanopyxis* were frequent diatoms in the early years of monitoring, but became rare after 2002. There are changes in the abundance of some taxonomic units. The abundance of *Phaeocystis globosa* has not changed significantly in the Bay of Somme, although it has increased slightly in Dunkerque and Boulogne-sur-Mer. Between 2002 and 2007, the abundance of the *Gymnodinium-Gyrodinium* group of dinoflagellates increased significantly, corresponding to a log scale abundance increase of twofold.

More recently, Lefebvre & Dezécache (2020) highlighted a significant break in the evolution of *Phaeocystis globosa* and *Pseudo-nitzschia* complex abundance in the 2000s and different trajectories of abundances in response to changes in nutrient concentration observed over the period 1994-2018. The three contrasting SRN sites appear to respond differently depending on the intensity of the initial nutrient input pressure. While a recovery to a good ecological status is doubtful in the near future, these ecosystems appear to be in an unstable intermediate state that necessitates continuous efforts to reduce nutrient inputs, particularly nitrogen.

These considerations on the relationship between phytoplankton succession and environmental conditions inevitably lead us back to Margalef's (1978) mandala, who paved the way for phytoplankton ecology by proposing functional groups to represent the adaptation of different life forms to specific habitats. However, it seems very difficult to propose, based on these concepts, a typical pattern of phytoplankton succession in the eastern English Channel and southern Bight of the North Sea from the SRN data. Similarly, assigning phytoplankton succession to the classical path or the route leading to the harmful blooms of the Margalef's mandala is hard, as is proposing a logical transition scheme between the various strategies described. In fact, alternative routes, overlaps, and mixtures of taxa with distinct strategies do exist, and the same taxon can even display several strategies according to its morphotype, as in the case of *Phaeocystis globosa*.

The concept of niche is crucial in phytoplankton research because it helps in understanding the succession of taxa, their coexistence, exclusion, and the environmental conditions that control them, as well as their tolerance to environmental changes. Thus, Karasiewicz et al. (2018) used an improved version of the OMI (Outlying Mean Index) approach to evaluate the niches of diatoms and *Phaeocystis globosa*. Two different situations of *P. globosa* bloom amplitude were defined by two different environmental trajectories and two different diatom communities, whose key features are given in table 4. Karasiewicz &



Lefebvre (2022) also developed a new method for bloom detection (based on twenty-two phenological variables) within a time-series. A pairwise quantification of asymmetric dependencies between the phenological variables revealed the implication of different mechanism, common and distinct between the studied taxa. A Permanova assisted in revealing the significance of seasonal variation in environmental and community factors. They were able to locate the harmful taxonomic niches among the rest of the community and quantify how their respective phenology influences the dynamic of their subniches by using methodologies such as the Outlying Mean Index and the Within Outlying Mean Index.

These results are comparable to those of Hernández Fariñas et al. (2015) who, based on a similar approach but extended to the Dunkerque and Bay of Somme SRN sites, concluded that light, temperature, species richness and nutrient concentrations are the main factors controlling phytoplankton dynamics and community structure.

Table 4. Main biotic and abiotic characteristics during two contrasting situations of *Phaeocystis globosa* bloom intensity from SRN data in the coastal zone off Boulogne-sur-Mer (-: low value for the parameter under consideration; +: high value).

Bloom intensity for <i>P. globosa</i>	Low	High
Initiation of the <i>P. globosa</i> bloom	Late	Early
Temperature	-	+
Salinity	-	+
Turbidity	+	-
Winter nitrate stock	+	-
Winter phosphate stock	+	-
Competition with Diatoms	+	-

We also confirmed from SRN data that periods of *Phaeocystis globosa* dominance are generally associated with high concentrations of *Pseudo-nitzschia* complex. The simultaneous presence of *P. globosa* and *Pseudo-nitzschia* complex will cause the creation of structures resembling mini-bearings (*Pseudo-nitzschia* needles planted in *Phaeocystis* colonies) that will be passively swallowed by the fish. These structures can cause mechanical aggression of the gill and/or digestive tissues opening the way to viral and bacterial infections. This mechanical aggressiveness will become even more crucial as new needle-shaped species or those with pointed spicules have been found in our investigations, contributing to the creation of these irritating assemblages. These are *Chaetoceros* sp., some *Thalassiosira* sp., *Rhizosolenia imbricata*, *R. styliformis*. Furthermore, the impact of phytoplankton blooms on the pelagic compartment, and in particular on fish, needs further investigation, especially in light of the findings of Lefebvre et al. (2011), Hernández-Fariñas et al. (2014), who found an increase in the abundance of *Pseudo-nitzschia* sp. since the early 2000s, and Delegrange et al. (2018), who found a correlation between mortalities of farmed sea bass (*Dicentrarchus labrax*) and the spring phytoplankton bloom. Major blooms of *P. globosa* will lead to changes in viscosity (Seuront et al., 2006) that may cause behavioural changes in fish (e.g. inhibition of



larval swimming activity) or metabolic changes (e.g. inhibition of gill functions). Changes in viscosity will also inevitably affect prey/predator relationships within planktons (Seuront and Vincent, 2008).

320 In order to facilitate the analysis and valorisation of SRN data, Devreker & Lefebvre (2014) proposed the development of a
user interface developed in R (R Core Team, 2020). This interface is available on the Comprehensive R Archive Network
(CRAN) as the TTAinterfaceTrendAnalysis package. It allows quickly defining the main statistical characteristics of SRN
series and proposing classical time series analyses (data regularization and aggregation, detection of anomalies, breaks and
seasonal or global trends). The results are presented in the form of summary tables or graphs that are automatically saved in
325 the user's working directory.

SRN data are used for validation of coupled hydrodynamic-biogeochemical models such as ECO-MARS 3D, in addition to
improving scientific knowledge about phytoplankton dynamics, biodiversity, and water quality (Ménèsquen et al., 2019).
Including retrieved-chlorophyll-*a* and suspended matters, SRN data are an essential source for the development and
330 improvement of satellite water color algorithms (Gohin et al., 2019, 2020a). In terms of the latter parameter, they are the only
data with such geographical and temporal coverage available in the coastal area of the Eastern Channel and the southern bight
of North Sea. Other data from the Coastal Observation Service (Somlit; <https://www.somlit.fr/>; last access on 28 April 2022)
can supplement the SRN data for the Boulogne sur Mer coastal zone (Lheureux et al., 2021).

In the context of the implementation of the Water Framework Directive (WFD 2000/60/EC) since 2007 (COM, 2005, a, b, c),
335 some coastal points of SRN integrate the so-called Monitoring and Operational Control system. The new Marine Strategy
Framework Directive (MSFD) extends the WFD approach limited to the first nautical mile from the baseline (for biological
parameters) to the offshore waters (Exclusive Economic Zone). Thus, the offshore stations of SRN network also meet the
diagnostic and monitoring expectations advocated by this European directive (Lefebvre and Devreker, 2020).

As part of the Oslo and Paris Convention's strategy to combat eutrophication (OSPAR <http://www.ospar.org/>), SRN results
340 are used to define the eutrophication status of water bodies. The SRN data are also transmitted to the ICES working group
"Phytoplankton and Microbiol Ecology" (WG PME) in order to contribute to the drafting of the devoted annual report
(<http://www.ices.dk/community/groups/Pages/WGPME.aspx>; last access on 28 April 2022).

Data availability

SRN - Regional Observation and Monitoring program for Phytoplankton and Hydrology in the eastern English Channel (2022).
345 SRN dataset - Regional Observation and Monitoring Program for Phytoplankton and Hydrology in the eastern English
Channel. SEANOE. <https://doi.org/10.17882/50832>



Related datasets:

PHYTOBS (2021). PHYTOBS dataset - French National Service of Observation for Phytoplankton in coastal waters. SEANOE. <https://doi.org/10.17882/85178>

350 REPHY – French Observation and Monitoring program for Phytoplankton and Hydrology in coastal waters (2021). **REPHY dataset - French Observation and Monitoring program for Phytoplankton and Hydrology in coastal waters. Metropolitan data.** SEANOE. <https://doi.org/10.17882/47248>

REPHYTOX - French Monitoring program for Phycotoxins in marine organisms (2021). **REPHYTOX dataset. French Monitoring program for Phycotoxins in marine organisms. Data since 1987.** SEANOE. <https://doi.org/10.17882/47251>

355 Code availability

R package *TTAinterfaceTrendAnalysis* sur le site du CRAN (Comprehensive R Archive Network - <https://cran.r-project.org/web/packages/TTAinterfaceTrendAnalysis/index.html> ; last access on 28 April 2022).

Team list

360 Lefebvre Alain, Devreker David, Blondel Camille, Duquesne Vincent, Hebert Pascale, Fabien Lebon, Cordier Remy, Belin Catherine, Maud Lemoine, Nadine Neaud-Masson, Hugué Antoine, Durand Gaétane, Soudant Dominique.

Author contribution

AL led the SRN monitoring programme and also led the writing of the paper. DD collaborated to the writing of the paper. Colleagues from the team list either led cruises or complementary monitoring networks (linked to related datasets), either analysed the samples, or contributed to QA and QC procedures described in the paper.

365 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

370 The acquisition of this dataset was performed in the framework of the SRN network, created and implemented by Ifremer and the Artois Picardie Water Agency in 1992. The people cited in the team list and contributors include regional coordinators of the SRN, key analysts and experts in phytoplankton observations, key analysts in physico-chemical measurements, those who contributed to structuring SRN data in the Quadrige² database, and individuals who contributed to the data through their



personal expertise. Other contributors such as the REPHY and REPHYTOX teams, crewmembers, students and PhD who have contributed to valorisation of the SRN data are also gratefully acknowledged.

References

- 375 Admiraal, W. and Venekamp, L. A. H.: SIGNIFICANCE OF TINTINNID GRAZING DURING BLOOMS OF PHAEOCYSTIS POUCHETII (HAPTOPHYCEAE) IN DUTCH COASTAL WATERS*, Netherlands Journal of Sea Research, 61–66 pp., 1986.
- Aminot, A. and K erouel, R.: Hydrologie des  cosyst mes marins : param tres et analyses, 2004.
- Billen, G., Garnier, J., and Rousseau, V.: Nutrient fluxes and water quality in the drainage network of the Scheldt basin over the last 50 years, Hydrobiologia, 540, 47–67, <https://doi.org/10.1007/s10750-004-7103-1>, 2005.
- 380 Brzezinski, M. A.: THE Si:C:N RATIO OF MARINE DIATOMS: INTERSPECIFIC VARIABILITY AND THE EFFECT OF SOME ENVIRONMENTAL VARIABLES, J. Phycol., 21, <https://doi.org/10.1111/j.0022-3646.1985.00347.x>, 1985.
- Cad e, G. C. and Hegeman, J.: SEASONAL AND ANNUAL VARIATION IN PHAEOCYSTIS POUCHETII (HAPTOPHYCEAE) IN THE WESTERNMOST INLET OF THE WADDEN SEA DURING THE 1973 TO 1985 PERIOD,
- 385 Netherlands Journal of Sea Research, 29–36 pp., 1986.
- Carpentier, A., Martin, C., and Vaz, S.: Channel Habitat Atlas for marine Resource Management, final report, CHARM phase 2, 2009.
- Delegrange, A., Lefebvre, A., Gohin, F., Courcot, L., and Vincent, D.: Pseudo-nitzschia sp. diversity and seasonality in the southern North Sea, domoic acid levels and associated phytoplankton communities, Estuar. Coast. Shelf Sci., 214, 194–206, <https://doi.org/10.1016/j.ecss.2018.09.030>, 2018.
- 390 Devreker, D. and Lefebvre, A.: TTAinterfaceTrendAnalysis: An R GUI for routine Temporal Trend Analysis and diagnostics, J. Oceanogr. Res. Data, 6, 1–18, 2014.
- Eberlein, K., Leal, M. T., Hammer, K. D., and Hiekel, W.: Dissolved organic substances during a Phaeocystis pouchetii bloom in the German Bight (North Sea), Marine Biology, 311–316 pp., 1985.
- 395 Gentilhomme, V. and Lizon, F.: Seasonal cycle of nitrogen and phytoplankton biomass in a well-mixed coastal system (Eastern English Channel), Hydrobiologia, 361, 191–199, <https://doi.org/10.1023/A:1003134617808>, 1998.
- Gohin, F., Saulquin, B., Oger-Jeanneret, H., Lozac’h, L., Lampert, L., Lefebvre, A., Riou, P., and Bruchon, F.: Towards a better assessment of the ecological status of coastal waters using satellite-derived chlorophyll-a concentrations, Remote Sens. Environ., 112, 3329–3340, <https://doi.org/10.1016/j.rse.2008.02.014>, 2008.
- 400 Gohin, F., Van der Zande, D., Tilstone, G., Eleveld, M. A., Lefebvre, A., Andrieux-Loyer, F., Blauw, A. N., Bry re, P., Devreker, D., Garnesson, P., Hern andez Fari nas, T., Lamaury, Y., Lampert, L., Lavigne, H., Menet-Nedelec, F., Pardo, S., and Saulquin, B.: Twenty years of satellite and in situ observations of surface chlorophyll-a from the northern Bay of Biscay to the eastern English Channel. Is the water quality improving?, Remote Sens. Environ., 233, 111343,



- <https://doi.org/10.1016/j.rse.2019.111343>, 2019.
- 405 Gohin, F., Bryère, P., Lefebvre, A., Sauriau, P.-G., Savoye, N., Vantrepotte, V., Bozec, Y., Cariou, T., Conan, P., Coudray, S., Courtay, G., Françoise, S., Goffart, A., Hernández Fariñas, T., Lemoine, M., Piraud, A., Raimbault, P., and Rétho, M.: Satellite and In Situ Monitoring of Chl-a, Turbidity, and Total Suspended Matter in Coastal Waters: Experience of the Year 2017 along the French Coasts, *J. Mar. Sci. Eng.*, 8, 665, <https://doi.org/10.3390/jmse8090665>, 2020a.
- Gohin, F., Bryère, P., Lefebvre, A., Sauriau, P., Savoye, N., Vantrepotte, V., Bozec, Y., Cariou, T., Conan, P., Coudray, S.,
410 Courtay, G., Françoise, S., Goffart, A., Hernández Fariñas, T., Lemoine, M., Piraud, A., Raimbault, P., and Rétho, M.: Satellite and In Situ Monitoring of Chl-a, Turbidity, and Total Suspended Matter in Coastal Waters: Experience of the Year 2017 along the French Coasts, *J. Mar. Sci. Eng.*, 8, 665, <https://doi.org/10.3390/jmse8090665>, 2020b.
- Gypens, N., Delhez, E., Vanhoutte-Brunier, A., Burton, S., Thieu, V., Passy, P., Liu, Y., Callens, J., Rousseau, V., and Lancelot, C.: Modelling phytoplankton succession and nutrient transfer along the Scheldt estuary (Belgium, The Netherlands),
415 *J. Mar. Syst.*, 128, <https://doi.org/10.1016/j.jmarsys.2012.10.006>, 2013.
- Hernández-Fariñas, T., Soudant, D., Barillé, L., Belin, C., Lefebvre, A., and Bacher, C.: Temporal changes in the phytoplankton community along the French coast of the eastern English Channel and the southern Bight of the North Sea, *ICES J. Mar. Sci.*, 71, 821–833, <https://doi.org/10.1093/icesjms/fst192>, 2014.
- Hernández Fariñas, T., Bacher, C., Soudant, D., Belin, C., and Barillé, L.: Assessing phytoplankton realized niches using a
420 French National Phytoplankton Monitoring Network, *Estuar. Coast. Shelf Sci.*, 159, 1–13, <https://doi.org/10.1016/j.ecss.2015.03.010>, 2015.
- Ibanez F., Fromentin J.M., Castel J.: Application de la méthode des sommes cumulées à l'analyse des séries chronologiques en océanographie. *Compte Rendu de l'Académie des Sciences Paris*, 316, 745-748, 1993.
- Karasiewicz, S. and Lefebvre, A.: Environmental Impact on Harmful Species *Pseudo-nitzschia* spp. and *Phaeocystis globosa*
425 Phenology and Niche, *J. Mar. Sci. Eng.*, 10, 174, <https://doi.org/10.3390/jmse10020174>, 2022.
- Karasiewicz, S., Breton, E., Lefebvre, A., Hernández Fariñas, T., and Lefebvre, S.: Realized niche analysis of phytoplankton communities involving HAB: *Phaeocystis* spp. as a case study, *Harmful Algae*, 72, 1–13, <https://doi.org/10.1016/j.hal.2017.12.005>, 2018.
- Lancelot, C. and Mathot, S.: Biochemical fractionation of primary production by phytoplankton in Belgian coastal waters
430 during short- and long-term incubations with ¹⁴C-bicarbonate - II. *Phaeocystis poucheti* colonial population, *Mar. Biol.*, 86, <https://doi.org/10.1007/BF00397508>, 1985.
- Lancelot, C., Rousseau, V., Billen, G., and Van Eeckhout, D.: 4. Coastal eutrophication of the Southern Bight of the North Sea: Assessment and modelling, *Bull. la Soc. R. des Sci. Liege*, 66, https://doi.org/10.1007/978-94-011-5758-2_33, 1997.
- Lefebvre, A. and Devreker, D.: First Comprehensive Quantitative Multi-Parameter Assessment of the Eutrophication Status
435 from Coastal to Marine French Waters in the English Channel, the Celtic Sea, the Bay of Biscay, and the Mediterranean Sea, *J. Mar. Sci. Eng.*, 8, 561, <https://doi.org/10.3390/jmse8080561>, 2020.
- Lefebvre, A. and Dezécache, C.: Trajectories of Changes in Phytoplankton Biomass, *Phaeocystis globosa* and Diatom (incl.



- Pseudo-nitzschia sp.) Abundances Related to Nutrient Pressures in the Eastern English Channel, Southern North Sea, *J. Mar. Sci. Eng.*, 8, 401, <https://doi.org/10.3390/jmse8060401>, 2020.
- 440 Lefebvre, A., Guiselin, N., Barbet, F., and Artigas, F. L.: Long-term hydrological and phytoplankton monitoring (19922007) of three potentially eutrophic systems in the eastern English Channel and the Southern Bight of the North Sea, *ICES J. Mar. Sci.*, 68, 2029–2043, <https://doi.org/10.1093/icesjms/fsr149>, 2011a.
- Lefebvre, A., Guiselin, N., Barbet, F., and Artigas, F. L.: Long-term hydrological and phytoplankton monitoring (19922007) of three potentially eutrophic systems in the eastern English Channel and the Southern Bight of the North Sea, *ICES J. Mar. Sci.*, 68, 2029–2043, <https://doi.org/10.1093/icesjms/fsr149>, 2011b.
- 445 Lheureux, A., Savoye, N., Del Amo, Y., Goberville, E., Bozec, Y., Breton, E., Conan, P., L’Helguen, S., Mousseau, L., Raimbault, P., Rimelin-Maury, P., Seuront, L., Vuillemin, R., Caparros, J., Cariou, T., Cordier, M., Corre, A., Costes, L., Crispi, O., Crouvoisier, M., Crouvoisier, M., Derriennic, H., Devesa, J., Durozier, M., Ferreira, S., Garcia, N., Grossteffan, E., Gueux, A., Lafont, M., Lagadec, V., Lecuyer, E., Leroux, C., Macé, E., Maria, E., Mornet, L., Nowaczyk, A., Parra, M., Petit, F., and David, V.: Bi-decadal variability in physico-biogeochemical characteristics of temperate coastal ecosystems: from large-scale to local drivers, *Mar. Ecol. Prog. Ser.*, 660, 19–35, <https://doi.org/10.3354/meps13577>, 2021.
- Margalef R.: Life forms of phytoplankton as survival alternatives in an unstable environment. *Oceanologica Acta*, 493-509, 1978.
- Ménesguen, A., Dussauze, M., Dumas, F., Thouvenin, B., Garnier, V., Lecornu, F., and Répécaud, M.: Ecological model of the Bay of Biscay and English Channel shelf for environmental status assessment part 1: Nutrients, phytoplankton and oxygen, *Ocean Model.*, 133, 56–78, <https://doi.org/10.1016/j.ocemod.2018.11.002>, 2019.
- R Core Team. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, 2020.
- 460 Rahmel, J., Bätje, M., Michaelis, H., and Noack, U.: *Phaeocystis globosa* and the phytoplankton succession in the East Frisian coastal waters, 49, 399–408, <https://doi.org/10.1007/BF02368365>, 1995.
- Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of sea-water. In: *The Sea. Ideas and observations on progress in the study of the seas. Volume 2. The composition of sea-water. Comparative and descriptive oceanography.*, edited by: Hill, M. N., Interscience publishers, a division of Wiley J. & Sons (Eds), New York, London, 26–77 pp., 1963.
- 465 Rousseau, V., Mathot, S., and Lancelot, C.: Calculating carbon biomass of *Phaeocystis* sp. from microscopic observations, *Mar. Biol.*, 107, 305–314, <https://doi.org/10.1002/9781119072218.ch11>, 1990.
- Seuront, L. and Vincent, D.: Increased seawater viscosity, *Phaeocystis globosa* spring bloom and *Temora longicornis* feeding and swimming behaviours, *Mar. Ecol. Prog. Ser.*, 363, 131–145, <https://doi.org/10.3354/meps07373>, 2008.
- Seuront, L., Vincent, D., and Mitchell, J. G.: Biologically induced modification of seawater viscosity in the Eastern English Channel during a *Phaeocystis globosa* spring bloom, *J. Mar. Syst.*, 61, 118–133, <https://doi.org/10.1016/j.jmarsys.2005.04.010>, 470 2006.



Utermöhl, H.: Zur vervollkommnung der quantitativen phytoplankton-methodik, Mitt Int Ver Theor Angew Limnol, 9, 1–38, 1958.

475 Wafar, M. V. M., Le Corre, P., and Birrien, J. L.: Nutrients and primary production in permanently well-mixed temperate coastal waters, Estuar. Coast. Shelf Sci., 17, 431–446, [https://doi.org/10.1016/0272-7714\(83\)90128-2](https://doi.org/10.1016/0272-7714(83)90128-2), 1983.