



1 **Multi-year emission of carbonaceous aerosols from cooking, fireworks burning,**
2 **sacrificial incenses, joss paper burning, and barbecue and the key driving forces in**
3 **China**

4 **Yi Cheng**^{1,2}, **Shaofei Kong**^{1,2*}, **Liquan Yao**^{1,2}, **Huang Zheng**^{1,2}, **Jian Wu**^{1,2}, **Qin Yan**^{1,2}, **Shurui Zheng**
5 ^{1,2}, **Yao Hu**^{1,2}, **Zhenzhen Niu**^{1,2}, **Yingying Yan**¹, **Zhenxing Shen**³, **Guofeng Shen**⁴, **Dantong Liu**⁵,
6 **Shuxiao Wang**⁶, and **Shihua Qi**²

7 ¹ Department of Environment Science and Engineering, School of Environmental Studies, China University
8 of Geoscience, Wuhan 430074

9 ² Department of Atmospheric Science, School of Environmental Studies, China University of Geoscience,
10 Wuhan 430074

11 ³ Department of Environmental Science and Engineering, School of Energy and Power Engineering, Xi'an
12 Jiaotong University, Xi'an 710049

13 ⁴ Laboratory for Earth Surface Process, College of Urban and Environmental Sciences, Peking University,
14 Beijing 100871

15 ⁵ Department of Atmospheric Science, School of Earth Science, Zhejiang University, Hangzhou 310027

16 ⁶ State Key Joint Laboratory of Environmental Simulation and Pollution Control, School of Environment,
17 Tsinghua University, Beijing 100084

18 Corresponding author: Shaofei Kong (kongshaofei@cug.edu.cn)

19



20 **Abstract**

21 There has been controversy about the air pollutants emitted from sources closely related to people's
22 daily life (such as cooking, fireworks burning, sacrificial incenses and joss paper burning, and barbecue,
23 named as five missing sources, FMS) impacting the outdoor air quality to what extent. Till now, there is no
24 emission estimation of air pollutants from FMS, as the missing of both activity dataset and emission factors.
25 We attempted to combine the questionnaire data, various statistical data, and data of points of interest to
26 obtain a relatively complete set of activity data. The emission factors (EFs) of carbonaceous aerosols were
27 tested in our lab. Then, the emission inventories of carbonaceous aerosol with the high spatial-temporal
28 resolution for FMS were established firstly, and the spatial-variation trend and driving forces were discussed.
29 From 2000 to 2018, organic carbon (OC) increased from 4268 t to 4919 t. The OC emission from FMS was
30 1.48–2.18 % of its total emission in China. The emissions of black carbon, element carbon (EC), and brown
31 carbon absorption cross-section (ACS_{BrC}) emissions from FMS were in the ranges of 22.6–43.9 t, 213–324
32 t, and 14.7–35.6 Gm^2 , respectively. Their emissions tended to concentrate in special periods and areas. The
33 OC emission intensities in central urban areas were 3.85–50.5 times that of rural areas due to the high density
34 of human activities. While the ACS_{BrC} emissions in rural regions accounted for 63.0–79.5% of the total
35 emission result from uncontrolled fireworks burning. A mass of fireworks burning led to extremely higher
36 ACS_{BrC} and EC emissions on Chinese New Year's eve, as 1444 and 262 times their corresponding yearly
37 average values. Interestingly, significant ($p < 0.01$) correlations between human incomes and pollutant
38 emissions were found, but they were positive ($r = 0.94$) and negative ($r = -0.94$) for urban and rural regions,
39 indicating the necessity of regulating human lifestyle and increasing income for urban and rural peoples,
40 respectively.

41 This study provided the first-hand data for identifying the emissions, variation trends and impacting
42 factors of FMS, which is helpful for modeling works on air quality, climate effect, and human health risks at
43 specific periods or regions and for modifying their emission control policies. The data in this work could be
44 found at <https://doi.org/10.6084/m9.figshare.19999991.v1> (Cheng et al., 2022).

45 **Keywords:** Carbonaceous aerosols; Sources related to human activities; Emission inventory; Spatial-
46 temporal variation; Driving force

47



48 **1 Introduction**

49 China has experienced a period of serious air pollution, which produces a great health impact on
50 residents (Zheng et al., 2018; Zhang et al., 2019, 2020b; Tong et al., 2020). Carbonaceous aerosols (CA),
51 emitted from incomplete burning, include organic carbon (OC) and black carbon (BC, or element carbon,
52 EC), and they have attracted wide attention due to their adverse impacts on air quality, human health and
53 climate (Venkataraman et al., 2005; Ramanathan & Carmichael, 2008; Bond et al., 2013). The optical
54 properties of CA (especially brown carbon, BrC) are complex and mutative, which is also one of the
55 important factors affecting the global radiation balance (Feng et al., 2013; Laskin et al., 2015).

56 Several sources closely related to traditional human activities were potential emission sources of CA,
57 such as the burning of sacrificial incense and joss paper, traditional Chinese barbecue, Chinese style cooking,
58 and fireworks burning. The estimation of air pollutants from these sources was missing in the previous
59 emission inventory, and they were defined as five missing sources (FMS) in this study. FMS can lead to a
60 dramatic impact on ambient quality and human health in a short period or a specific region (Chiang & Liao,
61 2006; Wu et al., 2015; Kong et al., 2015; Ho et al., 2016; Wang et al., 2017; Lai & Brimblecombe, 2020).
62 For example, fireworks burning could contribute 60.1 % of PM_{2.5} during the Chinese Lunar New year's eve
63 in Nanjing, and sacrificial sources like sacrificial incense and joss paper burning could contribute 17.5% of
64 atmospheric PAHs in Chu-Shan, and 9.6% of PCDD/F in Taipei (Lin et al., 2008; Kong et al., 2015; Ho et
65 al., 2016). Recently, with the strengthened control of combustion-related sources, the important role of
66 cooking emissions in affecting air quality was gradually visible in densely populated downtown areas.
67 Cooking organic aerosols contributed to 15–34% of total OC and 6–9% of total PM_{2.5} in a downtown site in
68 Shanghai (Huang et al., 2022) and 31% of total organic aerosols in Beijing in winter (Hu et al., 2022). The
69 existing studies on FMS were mainly based on the ambient air monitoring datasets at certain sites or certain
70 periods (See & Balasubramanian, 2011; Wu et al., 2015; Shen et al., 2017; Lao et al., 2018; Tanda et al.,
71 2019; Yao et al., 2019; Hu et al., 2021; Huang et al., 2021). Till, no studies can provide their contributions
72 quantitatively on a large scale as the scare of emission inventory, which limited the identification of their
73 contributions in various regions or periods.

74 There also existed extensive queries that are these conclusions tenable as they were identified through
75 models or ambient monitoring data, but not from their real emission estimation. In China, the differences in



76 population and economy between urban and rural areas are increasing (Meng et al., 2019), and the
77 efficiencies and necessity of air quality control policies for FMS in urban and rural areas need to be assessed.
78 For instance, fireworks burning was generally banned in the central urban region, while suburbs and rural
79 regions were affected less by the same policies. The cooking smokes needed to be purified in city centers,
80 while in suburban and rural areas, the policy may be not strictly executed or it is even not necessary. Such
81 deviations in policy establishment and implementation could ultimately drive the differences in the
82 distribution of air pollutant emissions, which have not been addressed yet.

83 The emission inventory is the base for the quantitative description of anthropogenic pollutant emissions
84 (Li et al., 2017). The combination of the chemical transport models and high-resolution emission inventory
85 was paramount for understanding anthropogenic perturbations' impacts on the atmosphere, and for assessing
86 corresponding air pollution control strategies (Janssens-Maenhout et al., 2019; McDuffie et al., 2020). The
87 lack of emission inventories limits the large-scale model simulation, the optimization of corresponding
88 control measures, and the settlement of related disputes. In our previous work, Wu et al. (2021) have
89 established an emission inventory of levoglucosan which has included the emissions from FMS (Wu et al.,
90 2021). There were no other emission inventories of FMS reported, to the best of our knowledge.

91 To sum up, this study aimed to develop a methodological framework for establishing an emission
92 inventory of FMS, including the methods of various activity data acquisition, emission factors monitoring,
93 uncertainty assessment, and spatial-temporal allocation. The activity data were obtained based on household
94 investigation, statistic data, points of interest (POI), etc. The emission factors were monitored through a
95 unique emission monitoring test platform, especially for fireworks burning in our lab. Then a high spatial (1
96 km × 1 km) and temporal resolution (1 day for special festivals, and 1 month for the rest) emission inventory
97 was first established. The multi-year spatiotemporal variation of CA emissions from these sources was
98 analyzed and compared with other types of sources. Optimization pollution control measures were proposed
99 for these types of sources. The study provides a methodology for establishing an emission inventory of air
100 pollutants from the sources closely related to human activities. Other air pollutants emissions could also be
101 estimated in the future. The emission inventory obtained here could also provide the basic inputs for
102 corresponding modeling works.

103 **2 Methodology**



104 2.1 Combustion tests for emission factors

105 The combustion tests for FMS were performed with two custom-made combustion chambers. One of
106 them had an explosion-proof function and it was used in fireworks burning experiments. Another one was
107 used in sacrificial incense, joss paper, barbecue, and cooking emission experiments. A dilution sampling
108 system (Dekati FPS-4000, Finland) was employed to dilute the smoke. The smokes were diluted about 16–
109 30 times and aged for about 30 s in a residence chamber. The sampling system has been utilized in residential
110 fuel combustion experiments (Cheng et al., 2019; Yan et al., 2020; Zhang et al., 2021b; Wu et al., 2021).
111 Thirty-eight events were tested in this experiment, including 6 trials of sacrificial incense combustion
112 (abbreviations of materials: red incense: RI; environmental incense: EI; high incense: HI), 6 trials of joss
113 paper burning (red-printed paper: RP; small sacrificial paper: SP; large sacrificial paper: LP), 10 trials of
114 fireworks burning (firecrackers: FC; fountain fireworks: FF; handheld fireworks: HF; handheld fountain:
115 HT; spin fireworks: SF), 8 trials of barbecue (chicken: CK; beef: BF, lamb: LB; pork: PK), and 8 trials of
116 the cooking (cooking of meat: MT1; cooking of meat and pepper: MT2; cooking of meat and garlic: MT3;
117 cooking of meat, pepper, and garlic: MT4). The experiment materials were shown in **Figure S1**.

118 After diluted, the OC and EC in the smokes were drawn into an online carbonaceous aerosol analyzer.
119 This analyzer was developed by the Key Laboratory of Environmental Optics & Technology (Anhui Institute
120 of Optics and Fine Mechanics, CAS) based on the thermal-optical method (Ding et al., 2014). The analyzer
121 showed reliable stability and repeatability. A dual-spot Aethalometer (Model AE33, Magee Scientific, USA)
122 was employed to measure BC concentration and particulate optical properties (Drinovec et al., 2015). The
123 experiment system was shown in **Figure S2**.

124 2.2 Calculation of emission factors and optical properties

125 The emission factors of OC, EC, and BC were calculated by equation (1):

$$126 \quad EF_{ij} = \frac{v \times m_{ij} \times r}{v_0 \times M_j} \quad (1)$$

127 where i and j denoted pollutant and fuel; EF was the emission factor (mg kg^{-1}); v was the flue
128 gas flow (L min^{-1}); v_0 was the sampling flow (L min^{-1}); m was the mass of the pollutant detected by the
129 instruments (mg); r was the dilution ratio; M was the fuel consumption quality (kg) (Cheng et al., 2019;
130 Yan et al., 2020).

131 All filter-based optical measurements will be confronted with the underestimation caused by the loading



132 effect (Drinovec et al., 2015). The loading compensation could be calculated based on dual-spot
133 measurements. The detailed calculation process was referred to Drinovec et al. (2015). There was inferior
134 dependence of BC particles on light absorption in different light wavelengths. The absorption Ångström
135 exponent (AAE) was an exceptional parameter to describe this dependence, as shown in equations (2) and
136 (3):

$$137 \quad b_{abs} \sim \lambda^{-AAE} \quad (2)$$

$$138 \quad AAE = - \frac{\ln\left(\frac{b_{abs}(\lambda_1)}{b_{abs}(\lambda_2)}\right)}{\ln\left(\frac{\lambda_1}{\lambda_2}\right)} \quad (3)$$

139 where λ was the wavelength; b_{abs} denoted the total light absorption coefficient (Tian et al., 2019),
140 which could be calculated by equation (4) (Zotter et al., 2017):

$$141 \quad BC = b_{abs}(\lambda) / MAC(\lambda) \quad (4)$$

142 where MAC was the mass absorption cross-section, referring to the Aethalometer manufacturer.

143 As shown in equation (5), the b_{abs} of CA was aroused by BC and BrC. To calculate the $b_{abs}(\lambda, BC)$
144 at each wavelength, equation (6) was introduced. The AAE_{BC} was determined as a value of 1.0 according
145 to previous studies (Tian et al., 2019; Liakakou et al., 2020). $f_{BrC}(\lambda)$ (equation (7)) was utilized to estimate
146 the fraction of BrC light absorption ($b_{abs}(\lambda, BrC)$) in total light absorption ($b_{abs}(\lambda)$).

$$147 \quad b_{abs}(\lambda) = b_{abs}(\lambda, BC) + b_{abs}(\lambda, BrC) \quad (5)$$

$$148 \quad b_{abs}(\lambda, BC) = b_{abs}(880) \times \left(\frac{\lambda}{880}\right)^{-AAE_{BC}} \quad (6)$$

$$149 \quad f_{BrC}(\lambda) = 100\% \times b_{abs}(\lambda, BrC) / b_{abs}(\lambda) \quad (7)$$

150 Due to the complicated chemical properties of BrC, it was difficult to measure the accurate
151 concentration of BrC in the flue gas. Thus, previous studies have developed a peculiar EF called absorption
152 emission factor (AEF), as shown in equation (8) (Martinsson et al., 2015; Tian et al., 2019; Zhang et al.,
153 2020c). Most studies modeled the direct radiative forcing of BrC with its mass concentration and mass
154 absorption efficiency (MAE) as the input parameters. While the mass concentration and the total mass of
155 BrC in the atmosphere were still unclear, and the MAE values in the range of 0.08–3.8 m² g⁻¹ were also
156 variable (Park et al., 2010; Feng et al., 2013; Wang et al., 2014b; Zhang et al., 2020a). Thus, an inventory
157 established by AEF as the following equation could avoid the deviation raised by the mass concentration and



158 MAE of BrC (Tian et al., 2019).

$$159 \quad AEF_{ij} = \frac{\sum_{t_0}^{t_{sample}} (b_{absij} \times r \times v)}{M_j} \quad (8)$$

160 2.3 Acquisition of activity data

161 Data on the activity of sources directly affected the uncertainties of emission inventory, and an accurate
162 estimate of FMS consumption was a crucial prerequisite. Statistics on direct consumption of FMS were
163 scarce. Thus, multiple activity data and proxy variables were adopted, including the statistical yearbooks of
164 each province in China, datasets of POI (**Text S1, Figure S3**), and rural household investigation data in our
165 group (**Text S2**).

166 The original consumptions of sacrificial incenses, joss paper, and fireworks, were from a household
167 investigation. We got the per capita consumption of sacrificial incenses, joss paper, and fireworks in each
168 province. The data were adjusted to overcome the problem of insufficient sample size. In China, sacrificial
169 activities mean honoring ancestors, and they mainly take place in temples or graveyards. Most traditional
170 graveyards would be placed in hills that might be covered with vegetation. The data on the consumption of
171 sacrificial incenses and joss paper will be revised based on the number of temples (data from POI) and
172 frequency of forest fires caused by sacrifices (data from China Forestry Statistical Yearbook), as shown in
173 equation (9):

$$174 \quad C_{S_{adj,province}} = C_{S_{inv,province}} \times \left(\frac{N_{t_{province}}/POP_{province}}{2 \times N_{t_{nation}}/POP_{nation}} + \frac{F_{s_{province}}/F_{t_{province}}}{2 \times F_{s_{nation}}/F_{t_{nation}}} \right) \quad (9)$$

175 where C_s was the consumption of sacrificial incenses or joss paper per capita; N_t was the number of
176 temples; POP was the population, which came from the statistical yearbook of each province; F_s was the
177 forest fires raised by sacrificial activities; F_t was the forest fires raised by all anthropogenic activities; adj
178 represented the data after adjusted; inv represented the data from the household investigation; $province$
179 represented the data of each province; $nation$ represented the data of the entire nation.

180 Consumption of fireworks will be revised based on the number of retail shops of fireworks (data from
181 POI), and provincial fireworks export volume (statistical data), as shown in equation (10):

$$182 \quad C_{f_{adj,province}} = C_{f_{inv,province}} \times \left(\frac{N_{s_{province}}/POP_{province}}{2 \times N_{s_{nation}}/POP_{nation}} + \frac{V_{e_{province}}/POP_{province}}{2 \times V_{e_{nation}}/POP_{nation}} \right) \quad (10)$$

183 where C_f was the consumption of fireworks per capita; N_s was the number of retail shops of
184 fireworks; V_e was the export volume of fireworks (data from China Light Industry Yearbook). In addition,



185 the consumptions (C) of sacrificial incenses, joss paper, and fireworks at the municipal level were calculated
186 by combining the POI data and the consumptions at the provincial level, as shown in equation (11).

$$187 \quad C_{adj,city} = C_{inv,province} \times \frac{N_{city}/POP_{city}}{N_{province}/POP_{province}} \quad (11)$$

188 where N represented the number of temples or firework shops.

189 The original meats consumption per capita came from the statistical yearbook of each province. But,
190 the methods and radii of different provincial statistical yearbooks showed differences. Part of municipal level
191 statistics was missing. To complement the missing data, municipal per capita consumption expenditure was
192 introduced. Using the logarithmic relationship between per capita consumption expenditure and provincial
193 per capita meat consumption to complement municipal per capita meat consumption. As shown in equation
194 (12):

$$195 \quad y = a \times \ln x + b \quad (12)$$

196 where x represented the provincial per capita consumption expenditures in 2000–2018; y
197 represented provincial per capita meat consumption in 2000–2018. Parameters a and b were fit-out for
198 each province ($r = 0.60$, $p < 0.01$). Then, the parameters a , b of the province where each city is located,
199 and the per capita consumption expenditure of the city were substituted into equation (12) to calculate the
200 municipal per capita meat consumption.

201 **2.4 Calculation of the emissions**

202 Since some cities have established policies to forbid sacrificial incense, joss paper, and fireworks
203 burning in the main urban area, such policies were inoperative in non-urban regions. According to our survey,
204 the policies on forbidden sacrificial incense and joss paper were relatively vague. Thus, we assumed that if
205 one city forbade fireworks burning, then the burning of sacrificial incense and joss paper was also banned.
206 The total emissions E from sacrificial incense, joss paper, and fireworks were calculated by equation (13):

$$207 \quad E = \sum (POP_{urban,k} \times C_{urban,k} \times FB_k + POP_{non-urban,k} \times C_{non-urban,k}) \times EF \quad (13)$$

208 where k denoted the different cities; *urban* and *non-urban* represented urban regions and non-
209 urban regions (rural regions); POP was the population; C was the per capita consumption; EF was the
210 emission factor. $FB = 0$ or 1 depended on whether the burning of sacrificial incense, joss paper, and
211 firework was forbidden in urban regions. Unfortunately, there was no such detailed consumption data that
212 involved the fuel types of sacrificial incense, joss paper, and fireworks. Thus, for the calculation of emission



213 from sacrificial incense, joss paper, and fireworks, mean EFs were utilized here.

214 Emissions from barbecue were calculated by formula (14):

$$215 \quad E = \sum (POP_{urban,k} \times MC_{urban,k} + POP_{non-urban,k} \times MC_{non-urban,k}) \times \frac{T_{BBQ,k}}{T_{total,k}} \times OP \times EF \quad (14)$$

216 where MC was the meat consumption mass per capita; T_{BBQ} was the number of restaurants
217 specializing in barbecue; T_{total} was the total number of restaurants. The numbers of restaurants were
218 calculated by using POI data. OP was the percentage of meals eaten out (data from the National Institute
219 for Nutrition and Health, Chinese Center for Disease Control and Prevention). In this study, we assumed that
220 barbecue was a kind of eating out.

221 Emission from cooking was the sum of the emission from residential cooking and the emission from
222 the catering industry. And they were calculated by formulas (15) and (16):

$$223 \quad E_{RC} = \sum (POP_{urban,k} \times MC_{urban,k} + POP_{non-urban,k} \times MC_{non-urban,k}) \times \left(1 - \frac{T_{BBQ,k}}{T_{total,k}}\right) \times (1 - OP) \times EF \times RE_{RC,k} \quad (15)$$

$$224 \quad E_{CI} = \sum (POP_{urban,k} \times MC_{urban,k} + POP_{non-urban,k} \times MC_{non-urban,k}) \times \left(1 - \frac{T_{BBQ,k}}{T_{total,k}}\right) \times OP \times EF \times RE_{CI,k} \quad (16)$$

225 where E_{RC} was the emission from the residential cooking and E_{CI} was the emission from the catering
226 industry; RE was the removal efficiency. The removal efficiency of the catering industry (RE_{CI}) was from
227 the national standard (GB 18483-2001). The removal efficiency of residential cooking activity (RE_{RC}) was
228 calculated using the popularizing rate of the range hood (data from China Statistical Yearbook) and removal
229 efficiency of the range hoods (GB/T 17713-2011).

230 A Monte Carlo simulation was employed to analyze the uncertainties of the emission inventory (Wu et
231 al., 2018). The simulation was executed 10000 times. The uncertainties of activity data were set as 0.2 or 0.5
232 (Table S1), and the uncertainties of EFs came from the actual measurements. The result of uncertainty was
233 shown in Table S2.

234 2.5 Spatial-temporal distribution of emissions

235 As above-mentioned, the emissions from rural and urban FMS activity might differ greatly. During the
236 process of spatial allocation, this difference must be emphasized. Thus, we were tempted to use GIS data for
237 the classification of land use to divide urban and non-urban regions (Gong et al., 2019, 2020). Based on this
238 method, we got the data on the distribution of population (data from www.worldpop.org,
239 10.5258/SOTON/WP00674) in urban and rural regions and constructed the emission map from FMS (Text




240 S3).

241 The temporal allocation methods for FMS were also specific. For calculating the annual trends of
242 emissions, statistical data including annual fireworks sales (data from the statistic of the Ministry of
243 Emergency Management of the PRC) and annual restaurant sales (data from <https://data.stats.gov.cn/>) were
244 used. The monthly trends of sacrificial incenses, joss paper, and fireworks burning were calculated with data
245 from household investigations. We believed that the activities of these sources are mainly concentrated on
246 five traditional Chinese festivals (Chinese New Year's Eve (CNE), Chinese Spring Festival (CSF), Spring
247 Lantern Festival (LF), Qingming Festival (QF), and Zhongyuan Festival (ZF)). We calculated the percentage
248 of incense, joss paper, and fireworks that burned during these festivals, and spread the excess to other days.
249 The monthly trends of barbecue and cooking emissions were calculated by using the monthly restaurant sales
250 in each province (data from <https://data.stats.gov.cn/>).

251 It should be noted that the above methods were alternatives due to the lack of direct statistical data, and
252 the methods can be improved in the future.

253



254

255 **Figure 1** Methodological framework for establishing high-resolution emission inventory for FMS.



256

257 3 Results and discussion

258 3.1 Emission characterization and light absorption properties

259 The EFs obtained from the 38 tests were shown in **Table 1**. The mean EF_{OC} of sacrificial incense,
 260 barbecue, joss paper, fireworks burning, and cooking were $32.6 \pm 12.6 \text{ mg kg}^{-1}$, $33.2 \pm 13.5 \text{ mg kg}^{-1}$, $41.9 \pm$
 261 27.8 mg kg^{-1} , $51.9 \pm 45.5 \text{ mg kg}^{-1}$, and $159 \pm 34.0 \text{ mg kg}^{-1}$, respectively. While the EF_{EC} and EF_{BC} showed
 262 different tendencies. Barbecue exhibited higher EF_{EC} ($5.13 \pm 5.23 \text{ mg kg}^{-1}$) and EF_{BC} ($69.6 \pm 79.5 \text{ mg kg}^{-1}$)
 263 than those of sacrificial incense (EF_{EC}: $0.17 \pm 0.07 \text{ mg kg}^{-1}$, EF_{BC}: $1.80 \pm 0.92 \text{ mg kg}^{-1}$), joss paper (2.25
 264 $\pm 2.47 \text{ mg kg}^{-1}$, $3.79 \pm 2.23 \text{ mg kg}^{-1}$), cooking ($0.005 \pm 0.001 \text{ mg kg}^{-1}$, $1.54 \pm 0.17 \text{ mg kg}^{-1}$), and fireworks
 265 burning ($2.57 \pm 5.37 \text{ mg kg}^{-1}$, $14.8 \pm 17.3 \text{ mg kg}^{-1}$). The diversities can result from the fuel properties and
 266 burning conditions.

267

268

Table 1 BC, OC, and EC emission factors for five missing sources (mg kg^{-1}).

Sources	Materials*	BC	OC	EC
Sacrificial incense	RI	3.09 ± 0.05	49.2 ± 6.39	0.23 ± 0.07
	EI	1.24 ± 0.17	27.3 ± 2.63	0.17 ± 0.02
	HI	1.07 ± 0.04	21.4 ± 2.06	0.10 ± 0.02
Joss paper	RP	6.27 ± 1.59	35.5 ± 5.59	0.97 ± 0.10
	SP	1.65 ± 0.41	14.6 ± 1.64	0.64 ± 0.50
	LP	3.45 ± 1.18	75.5 ± 19.2	5.12 ± 2.28
Fireworks	FC	3.56 ± 0.32	8.72 ± 0.08	0.14 ± 0.03
	FF	2.89 ± 0.88	5.86 ± 1.28	0.06 ± 0.03
	HF	23.0 ± 8.63	124 ± 29.2	9.79 ± 8.49
	HT	7.49 ± 0.20	65.7 ± 10.5	2.39 ± 1.76
	SF	37.3 ± 22.8	55.1 ± 0.66	0.48 ± 0.29
Barbecue	CK	1.66 ± 0.30	21.5 ± 1.11	0.15 ± 0.02
	BF	37.2 ± 24.4	28.6 ± 8.85	3.78 ± 2.28
	LB	48.5 ± 17.7	32.2 ± 6.35	4.21 ± 0.58
	PK	191 ± 59.5	50.5 ± 12.2	12.4 ± 4.93
Cooking	MT1	1.79 ± 0.04	127	0.003
	MT2	1.54 ± 0.01	124	0.004
	MT3	1.34 ± 0.05	181	0.007
	MT4	1.48 ± 0.07	203	0.007


269

*: Abbreviations of fuels: RI: red incense; EI: environmental incense; HI: high incense; RP: red-printed paper; SP:



270 small sacrificial paper; LP: large sacrificial paper; FC: firecrackers; FF: fountain fireworks; HF: handheld fireworks; HT:
271 handheld fountain; SF: spin fireworks; CK: chicken; BF: beef, LB: lamb; PK: pork; MT1: cooking of meat; MT2: cooking
272 of meat and pepper; MT3: cooking of meat and garlic; MT4: cooking of meat, pepper, and garlic. FMS: five missing sources.
273

274 To quantify the light absorption properties of emissions from FMS, AAEs (370–880 nm) were calculated
275 (**Figure 2**). The average AAEs of FMS were in the range of 1.26–3.15. The mean AAE of sacrificial incenses
276 (2.69 ± 0.36) was slightly higher compared to joss paper (2.22 ± 0.65), fireworks burning (2.10 ± 0.50),
277 barbecue (1.40 ± 0.14), and cooking (1.39 ± 0.08). The AAEs in 370–880 nm wavelength of woody fuel
278 burning (1.0–2.7) (Martinsson et al., 2015; Zhang et al., 2020a, 2021c), crop residues burning (1.5–3.25)
279 (Tian et al., 2019; Zhang et al., 2020c, 2021a), coal combustion (1.1–2.5) (Tian et al., 2019; Zhang et al.,
280 2021a), and engines (1.1–2.4) (Corbin et al., 2018) were comparable to our results. $AAE > 1$ indicates that
281 there was BrC in aerosols (Saleh et al., 2013; Sun et al., 2017). Thus, it is necessary to investigate BrC
282 emission characteristics and the contribution of BrC to the total light absorption from various sources.
283



284

285 **Figure 2** The absorption Ångström exponents (370–880 nm) of FMS.

286




287 We have calculated f_{BrC} to estimate the light absorption ability of BrC in aerosols (**Figure S4**). f_{BrC}
288 showed a decreasing tendency toward the long wavelengths, which proved the understanding that the light
289 absorption ability of BrC had stronger spectral dependence than that of BC (Sun et al., 2017). At 370 nm
290 wavelength, f_{BrC} of sacrificial incense, joss paper, fireworks, barbecue, and the cooking were $71.5 \pm 5.32\%$,
291 $58.4 \pm 24.0\%$, $57.6 \pm 9.36\%$, $28.7 \pm 6.19\%$, and $29.2 \pm 4.30\%$, respectively. f_{BrC} of cooking sources (22.9–
292 37.4% with an average of 29.0%) like barbecue and cooking seemed to be much lower than other combustion
293 sources (34.8–82.7% with an average of 61.6%). f_{BrC} were 47% for coal combustion at 355 nm wavelength
294 (Sun et al., 2017), and 68–85% for biomass burning at 370 nm (Tian et al., 2019). As some emission sources
295 were neglected, the particulate absorption and warming effect contributed by BrC may be underestimated in
296 former modeling works (Laskin et al., 2015).

297 Furthermore, the AEF of BrC and BC have been calculated, as shown in **Figure S5**. As the wavelength
298 increased, the AEFs showed a decreasing trend. When $\lambda=370$ nm, AEF_{BrC} from fireworks burning was the
299 highest, as $2.65 \pm 3.23 \text{ m}^2 \text{ kg}^{-1}$, followed by barbecue ($0.45 \pm 0.49 \text{ m}^2 \text{ kg}^{-1}$), joss paper ($0.19 \pm 0.21 \text{ m}^2 \text{ kg}^{-1}$),
300 sacrificial incenses ($0.15 \pm 0.10 \text{ m}^2 \text{ kg}^{-1}$), and cooking ($0.012 \pm 0.004 \text{ m}^2 \text{ kg}^{-1}$). At 370 nm, the AEF_{BrC} of
301 coal combustion and biomass burning have been reported as $14.3\text{--}46.6 \text{ m}^2 \text{ kg}^{-1}$ and $2.01\text{--}24 \text{ m}^2 \text{ kg}^{-1}$
302 (Martinsson et al., 2015; Tian et al., 2019), which were 1–3 order of magnitude higher than those of FMS.

303 3.2 Characterization of activity

304 The total consumption of FMS has been shown in **Figure 3**. In 2018, 16.5 kt, 919 kt, and 5139 kt of
305 sacrificial incenses, joss paper, and fireworks were consumed in China. 30996 kt, 2872 kt, 1920 kt, and
306 12057 kt of pork, beef, lamb, and chicken were consumed. The total consumption amount of FMS was about
307 26.4% of the residential coal consumption in China (Peng et al., 2019). The consumptions of sacrificial
308 incense, joss paper, and fireworks were highest in Shandong (394 kt) and Sichuan (470 kt). Apart from lamb,
309 Guangdong province has the largest consumption of three kinds of meats (6197 kt), and the province with
310 the largest consumption of lamb was Xinjiang (361 kt). The consumption of FMS can be a reflection of the
311 local customs. For example, lamb consumption in Xinjiang was the highest in China. The reason was that
312 Xinjiang is the main producing area of lamb, one of the five pastoral areas in China, and Xinjiang's ethnic
313 structure makes mutton a dominant part of the daily diet (Xu et al., 2018).



314
315 **Figure 3** The total consumption of FMS during 2000–2018 in China (Abbreviation: SI: sacrificial incense; JP: joss paper;
316 FW: fireworks; PK: pork; BF: beef; LB: lamb; CK: chicken).

317
318 The FMS consumptions in most cities were at low levels. The top 30 cities (about 8% of the total
319 number of cities) with the largest fireworks consumption contributed 41.8% of the national consumption
320 (**Figure S6**). These cities have higher population densities, and the control measures for fireworks were not
321 yet in place. Per capita fireworks consumption in 52% of the cities was less than 5 kg yr⁻¹, and the mass of
322 a common firecracker can exceeds 5 kg. As shown in **Figure S7**, the distribution pattern of per capita
323 consumption of sacrificial incenses and joss paper was similar to that of fireworks. The differences in meat
324 consumption between cities were relatively smaller. The top 30 cities with the largest pork consumption only
325 contributed 30.8% of the national consumption. The lowest per capita pork consumption was only 2.31 kg
326 yr⁻¹ and the highest was 45.6 kg yr⁻¹ (**Figure S7**). While 71.9% of the cities had per capita pork consumption
327 of 10–30 kg yr⁻¹. From 2000 to 2018, the consumption of four types of meat increased by 49.3%, and the
328 trends of increased meat consumption were similar to a previous study (Batis et al., 2014). While the
329 consumption of sacrificial incense, joss paper, and fireworks showed a trend of increasing first and then
330 decreasing.




331 3.3 CA emission from FMS in China

332 3.3.1 Multi-year variation

333 In 2000–2018, OC, EC, BC, and BrC absorption cross-section (ACS_{BrC} , in 370 nm wavelength)
334 emissions from FMS were 4268–4919 t, 22.6–43.9 t, 213–324 t, and 14.7–35.6 Gm², respectively (**Figure**
335 **4**). Severe air pollution over the past decade has led China to enact a series of policies to limit emissions
336 from various sources. Thus, the total CA emission in China presented approximately monotonically
337 decreased tendencies. During 2010–2017, the total OC and BC emissions in China decreased from 3.2 Tg
338 and 1.7 Tg to 2.1 Tg and 1.3 Tg, mostly contributed by residential sources (76.9–80.3% of OC, 41.8–51.5%
339 of BC) (<http://meicmodel.org>, Li et al., 2017; Zheng et al., 2018). The emission tendencies of FMS showed
340 different variation tendencies compared with these above sources. 82.7–92.3% of OC emissions came from
341 cooking (**Figure S8**), and the OC emissions from FMS increased first and then decreased. From 2000 to
342 2018, meat consumption per capita increased from 22.7 kg to 33.9 kg with the rapid development of the
343 economy and living standards in China. However, the popularizing rates of range hoods in China have
344 increased by 44.9% in the urban regions and 23.3% in rural regions, and it led to a decrease in OC emissions.
345 The combined effect of these two factors made the emission of FMS OC raised by 14.4% from 2000 to 2013
346 and then decreased by 10.6% from 2013 to 2018. The EC, BC, and ACS_{BrC} emissions showed similar
347 tendencies. From 2000 to 2006, EC, BC, and ACS_{BrC} emissions from FMS increased by 52.3%, 45.4%, and
348 51.2%, respectively. Then they decreased by 48.7%, 27.8%, and 58.4% in 2006–2018. Fireworks burning
349 was one of the main contributors to CA emissions from FMS, which contributed 58.6–76.0%, 33.7–61.9%,
350 and 88.5–96.6% of the EC, BC, and ACS_{BrC} emissions. The consumption of fireworks showed a trend of
351 inverted U shape. It dominated the tendencies of EC, BC, and ACS_{BrC} emissions from FMS. Moreover, there
352 was a surge in emissions due to the high consumption of fireworks in 2014.

353



354

355 **Figure 4** Total CA emission from FMS in China from 2000 to 2018 (Abbreviations: SI: sacrificial incense; JP: joss paper;
356 FW: fireworks; BBQ: barbecue; RC: Residential cooking; CI: Catering industry).

357


358 3.3.2 Spatial variation

359 There are seven geographical regions in China (**Figure S9**), of which East China was the largest CA
360 emission region, and East China contributed 24.2–27.7%, 23.9–29.6%, 24.2–29.0%, and 23.5–29.9% of the
361 OC, EC, BC, and ACS_{BrC} total emissions from FMS, respectively (**Figure 5**). The dense distribution of the
362 population (28.2–30.3% of the population) was responsible for the high emissions in East China. The OC,
363 EC, and BC emission from FMS in Southernwest China was second to that of East China. OC, EC, and BC
364 emissions in Central China accounted for 21.5–27.2%, 18.0–21.3%, and 19.2–22.5% of their national total
365 emissions. Eating habits in Southernwest China led to its high emission. Southernwest China has a higher
366 density of barbecue restaurants (11.9% higher than the national average) and per capita meat consumption
367 (33.4% higher than the national average), as well as a large population (14.9% of the national total). The
368 ACS_{BrC} emission from Central China was second to that of East China, accounting for 14.3–21.6% of the
369 national total. 90.9–96.4% of ACS_{BrC} emission was from fireworks burning. The Hunan province in Central
370 China is one of the main fireworks-producing regions. The density of fireworks stores in Hunan province
371 was 2.29 times that of the national average. What's more, Central China was also one of the densely



372 populated regions, accounting for 16.0–17.6% of the national population. Due to the heating needs in winter,
373 the OC and BC emissions from other sources in North China contributed 14.8–17.2% and 17.6–21.1% of
374 the national total (Li et al., 2017). However, the contributions of OC and BC emissions from FMS in North
375 China were only 7.75–8.52% and 9.53–10.5%. The lower contribution was due to the lower per capita meat
376 consumption (25.4% lower than the national average) and fewer restaurants (5.5% lower than the national
377 average).

378 The emission distributions from different sources showed great differences, which came from the
379 regional culture and economic diversity (**Figure S10** and **Figure S11**). High emission regions of sacrificial
380 incense and joss paper overlapped with the areas with large numbers of temples and cradles of Chinese
381 Buddhism (**Figure S3**), where people in those areas may be more devout about sacrifice. The distributions
382 of cooking emissions (both residential cooking emissions and catering industry emissions) and barbecue
383 emissions were highly similar to the population distribution, especially in urban regions. This is consistent
384 with previous studies. For example, the cooking-related organic aerosol (COA) concentrations at urban sites
385 ($6.46\text{--}6.97\ \mu\text{g m}^{-3}$ in Beijing and $14.2\ \mu\text{g m}^{-3}$ in Shijiazhuang) were much higher than that at the rural site
386 ($2.96\text{--}3.74\ \mu\text{g m}^{-3}$ in Gucheng, a rural site near Beijing), and COA concentration was 0 at the background
387 site (Sun et al., 2013, 2020; Wang et al., 2015b, 2020; Huang et al., 2019). Areas with higher economic
388 consumption have more demands for repast styles and varieties, leading to more emissions. Emissions of
389 fireworks showed an obvious difference in urban and rural regions. Emissions from urban regions were near
390 zero, while emissions from suburbs and rural regions were much higher (**Section 3.3.3**).



391

392 **Figure 5** Spatial distribution of CA emission from FMS in China in 2018. The colorbar showed the emission in each grid.


393

394 3.3.3 Intense short-term and regionally concentrated emissions

395 As shown in **Figure S12** and **Figure S13**, CA emissions from residential sources in winter were




396 extremely higher than in summer resulting from the heating demand (Wang et al., 2012; Huang et al., 2015;
397 Li et al., 2017), while emissions from FMS showed a similar seasonal trend due to the fireworks burning.
398 During the Chinese Spring Festival, the fireworks burning results in massive pollutant emissions and severe
399 air pollution (Kong et al., 2015; Yao et al., 2019; Ding et al., 2019; Lai & Brimblecombe, 2020). We have
400 investigated the CA emissions from FMS in the month and at several related Chinese festivals (Chinese New
401 Year's Eve (CNE), Chinese Spring Festival (CSF), Spring Lantern Festival (LF), Qingming Festival (QF),
402 and Zhongyuan Festival (ZF) (Text S4)). As shown in Figure 6. The emissions were mostly concentrated in
403 January and February. 75.8% of fireworks were set off on CNE and CSF, and 20.4% were set off on LF
404 (Figure 7). Thus, the ACS_{Brc} emission on CNE was 1444 times of the yearly average, and the OC, EC, and
405 BC emissions were also 10.9, 262, and 74.6 times the average, respectively. The highly short-term emission
406 of fireworks led to a sharp increase in the concentration of air pollutants (Vecchi et al., 2008; Shi et al., 2011;
407 Cao et al., 2018; Lai & Brimblecombe, 2020).
408



409
410 **Figure 6** Averaged monthly CA emissions from FMS in China for the year of 2000–2018.

411




412
413 **Figure 7** Average CA emissions on Chinese New Year's Eve (CNE), Chinese Spring Festival (CSF), Spring Lantern Festival
414 (LF), Qingming Festival (QF), and Zhongyuan Festival (ZF) for the year of 2000–2018. The AVE showed the average daily
415 emissions except for the festivals mentioned above.

416
417

418 In addition to the feature of short-term, emission from FMS also shows obvious spatial distributions.
419 83.2–93.1% of the OC emission came from barbecue and cooking. The higher population density and living
420 quality led to higher OC emissions in urban regions. As shown in **Figure 8**, the OC emission intensities
421 (emission per square kilometer) in the urban regions of Chengdu, Xi'an, Beijing, and Tianjin were 62.6, 63.1,
422 27.0, and 14.6 times of those for rural regions in 2018. This situation was common in China. China set up
423 13 prevention and control regions (3 key regions and 10 city clusters, 3-10R) in 2013 to improve air pollution,
424 and they are relatively developed regions (https://www.mee.gov.cn/gkml/hbb/bgg/201303/t20130305_248787.htm). The OC emission intensities in the urban regions of 3-10R were 3.85–50.5 times of those for
425 the surrounding rural regions. Thus, OC emissions from FMS were concentrated in urban regions overall.
426 Since fireworks burning was concentrated in CNE or CSF, due to the emission from fireworks burning in
427 rural regions, the feature that OC emission concentrated in urban regions would be attenuated at CNE. In
428



429 contrast, ACS_{B_rC} emissions tended to concentrate in rural regions, especially during the periods of CNE and
430 CSF (**Figure 8**). Fireworks burning was the main contributor (>88.5%) to ACS_{B_rC} emissions, and the
431 fireworks burning was concentrated in CNE or CSF. The ACS_{B_rC} emission intensities in rural regions of
432 Chengdu, Xi'an, Beijing, and Tianjin were 18.8, 20.0, 107, and 150 times of those for urban regions at the
433 2018 CNE. Many cities have introduced policies to control firework burning, and civilized sacrifice is
434 encouraged. But these policies tend to be implemented effectively only in central urban regions. Urban
435 suburbs and surrounding rural regions, that are densely populated, are areas that policies do not consider or
436 be executed efficiently. The contribution of rural ACS_{B_rC} emissions in 3-10R was ~79.0% and even as high
437 as 96.7% at the 2018 CNE. However, the rural population in these regions only accounted for 14.1–41.9%.
438 In fact, 63.0–79.5% of ACS_{B_rC} emissions at CNE came from the rural regions in China. During the period
439 of CNE and CSF, pollutants emitted from rural residents' activities were likely to be transmitted to urban
440 areas, leading to serious air pollution in urban regions (Yao et al., 2019; Pang et al., 2021).




441

442 **Figure 8** Differences in OC (three figures above) and ACS_{BrC} (three figures below) emissions distributions in urban and
443 rural regions in Chengdu (core city of Sichuan Basin), Xi'an (core city of Guanzhong Plain), and Beijing & Tianjin (Core
444 city of North China Plain). The first and second lines showed the OC emissions in the total year of 2018, and on the day of
445 CNE, respectively. The third and fourth lines showed the ACS_{BrC} emissions in the total year of 2018, and on the day of CNE.
446 The arrows pointed to the city centers. The colorbars showed the emission in each grid for OC and ACS_{BrC}.



447 **3.3.4 Emissions impacted by economical development**

448 Barbecue and cooking contributed a significant portion of OC emissions from FMS, which led to a
449 distinctive feature of emissions from FMS. There was a certain correlation between OC emissions and local
450 economic development. We have gathered disposable income per capita data from 2000 to 2018 for each
451 city. The relationship between the disposable income and OC emission per capita has been assessed. As
452 shown in **Figure 9**, like other emission sources, OC emissions from FMS and disposable income showed an
453 inverted U-shape relationship (Environmental Kuznets Curves) (Wu et al., 2020; Zhong et al., 2020).
454 However, if we separated the emission-economical relationship in urban regions from rural regions, the
455 results would be different. The relationships were linear in both urban and rural regions (**Figure 9**). However,
456 the correlation was significantly negative ($r = -0.97, p < 0.01$) in urban regions compared to the positive one
457 ($r = 0.94, p < 0.01$) in rural regions. As discussed above, cooking sources dominated the OC emissions from
458 FMS in China. From 2000 to 2018, meat consumption per capita increased by 83.4%, and the OC emissions
459 per capita have increased by 36.9% in rural regions. In urban regions, meat consumption increased by 22.0%,
460 while OC emissions per capita decreased by 39.1%. The reason for this phenomenon was the higher
461 popularizing rate of range hoods in urban regions, which was 5.21 times that of rural regions. What's more,
462 the popularizing rate of range hoods in urban regions has also increased by 132% in urban regions. As a
463 result, OC emissions that would have been raised were eliminated by a large number of range hoods in urban
464 regions.
465



466

467 **Figure 9** The relationship between per capita disposable income and OC emissions from FMS for the years 2000–2018 in
468 China. The left figure showed the invert-U shape relationship between the income and the national average emissions. The
469 right figure showed the correlation between the income and emissions in the urban regions and the rural regions. The shaded
470 areas represent 95% confidence intervals.

471


472 In contrast to the relatively developed 3-10R, there are some contiguous poor regions (CPR) in China,
473 where located in the borderland or mountains (http://www.gov.cn/gzdt/2012-06/14/content_2161045.htm).
474 The other regions (OR) excluding 3-10R and CPR, were at a moderate level of development in China. The
475 OC emissions per capita in 3-10R, OR, and CPR, were 3.04–3.77 g, 3.49–4.00 g, and 3.54–4.11 g in 2000–
476 2018. OC emissions per capita in 3-10R, OR, and CPR have all crossed the inflection point of the emission-
477 economical correlation. Thus the relatively developed 3-10R have lower per capita emissions. It also verified
478 the economic impact of OC emissions. However, 3-10R has 76.2–77.4% of the population, thus the emission
479 intensities in 3-10R were still 3.11–3.39 times that of the national average.

480 3.3.5 The implication for modifying related air pollution control policies

481 To combat air pollution, China introduced its toughest air pollution control plan (Air Pollution
482 Prevention and Control Action Plan, APPCP) ever in 2013 (Zhang et al., 2019a). The implementation of the



483 APPCP had led to significant improvements in China's air quality. The control measures of FMS have also
484 begun to be widely promoted (**Figure S14**). There were 76.3% and 66.5% of cities have introduced policies
485 to restrict the emissions from the catering industry and fireworks burning before 2018. The removal
486 efficiencies of pollutants for small, medium, and large catering industries were higher than 60%, 75%, and
487 85% (GB 18483-2001). The local governments have the right to designate the areas where fireworks were
488 forbidden, usually urban areas, along with hospitals, factories, power plants, schools, and transportation hubs
489 (http://www.gov.cn/zwggk/2006-01/25/content_170906.htm). In addition, the government had proposed
490 residents install range hoods to control the emissions from cooking (APPCP), and the national popularizing
491 rate of range hoods increased by 43.6% from 2000 to 2018. The timing of implementation varies in local
492 governments, and the implementation of these policies has produced positive significance. OC, EC, BC, and
493 ACS_{BrC} emissions from have declined by 14.3–47.1%, 9.76–45.4%, 9.23–42.2%, and 10.4–48.2% in 2000–
494 2018 (**Figure 10**).
495



496
497 **Figure 10** The impact of policies on the reductions of CA emissions. The solid lines (left y-axis) represented the actual CA
498 emissions compared to the emissions without the policy impact (100% on the left y-axis). The shaded part of the solid line
499 represents uncertainties. The dotted line (right y-axis) represented the number of cities that issued policies to control FMS
500 emissions.



501 If we assume that there was also a quadratic fitting relationship between rural per capita OC emissions
502 and income, then rural per capita OC emissions would start to decline when rural per capita income reaches
503 16.8 k Yuan. The control of CA emissions from FMS like cooking should start from the perspective of
504 increasing the income of rural residents. With enough income, residents will tend to a more environmentally
505 friendly and green lifestyle. The green lifestyle is embodied by the installation of a range hood in this work.
506 In 2017, the impervious surfaces of urban regions only accounted for 1.52% of the national area (Gong et
507 al., 2019), and rural regions are vast by contrast. Thus, the cost of controlling the activities of fireworks
508 burning, sacrificial incenses, and joss paper burning in rural regions will be much higher than in urban
509 regions. For these sources, policies and standards should be set to limit their emissions from the burning
510 processes. In addition, it is questionable whether the environmentally friendly fireworks currently on the
511 market have a lower impact on the environment (Fan et al., 2021). Thus, manufacturers should be guided to
512 develop environmentally friendly fireworks, joss paper, and sacrificial incense to reduce emissions.

513 **3.4 Comparison with other studies**

514 As an emission source with less attention, most of the relevant studies focused on the EFs of PM (Jetter
515 et al., 2002; Lee & Wang, 2004; Wang et al., 2015a; Kuo et al., 2016; Jilla & Kura, 2017; Amouei
516 Torkmahalleh et al., 2018; Wang et al., 2018b; Zhao et al., 2018; Lin et al., 2019), PAHs (Yang et al., 2005,
517 2013; Zhao et al., 2019), and VOCs (Cheng et al., 2016; Wang et al., 2018b). Several metallic elements
518 (Croteau et al., 2010; Shen et al., 2017) and organic matters (Xiang et al., 2017; Que et al., 2019) have also
519 been tested. Few studies have tested OC and EC EFs of FMS (See & Balasubramanian, 2011; Zhang et al.,
520 2019b; Lin et al., 2021). Fireworks burning was the least studied emission source, while fireworks can emit
521 large amounts of particles. EFs of PM₁₀ from fireworks burning as 54–429 g kg⁻¹ have been reported
522 (Camilleri & Vella, 2016; Keller & Schragen, 2021), which were much higher than the CA EFs in this study.
523 The EFs in the literatures have been shown in **Table S3**.

524 Several studies have calculated the emission amount of the catering industry (**Table S4**). For example,
525 Wang et al. (2018a) have calculated the VOCs emission (66245 t) from restaurants in China based on
526 samplings of 9 types of restaurants. Jin et al. (2021) have calculated the OC from the catering industry in
527 China by investigations in two cities in Shandong and Shanxi provinces. The results showed that OC
528 emissions from the catering industry were 26.8 Gg, which was 66.0 times that of our results. The EFs used



529 in Jin et al. (2021) were the generation rates of pollutants, which were 0.48 mg m^{-3} for OC in oil fumes.
530 Different EFs and calculation methods may be the main reason for the discrepancy. Emissions from cooking
531 have been reported as the main driver of OC in urban regions, as it contributed large portions of organic
532 aerosols in Shanghai (20–35%) and Beijing (10–19%) (Liu et al., 2021; Zhu et al., 2021b). The effects of
533 cooking emissions on the urban atmosphere should not be neglected when other sources like residential or
534 industry sources were efficiently controlled (Zhang et al., 2021c; Zhu et al., 2021a).

535 Previous research has calculated the total OC and BC emissions in China, such as the widely used MEIC
536 (OC: 2080–3190 Gg, BC: 1253–1728 Gg) and PKU inventory (OC: 2345–3587 Gg, BC: 1455–1624 Gg)
537 (Wang et al., 2014a; Huang et al., 2015; Li et al., 2017). Residential sources or residential & commercial
538 source contributed most of OC (80.3% in MEIC, and 71.4% in PKU) and BC (51.5% in MEIC, and 51.0%
539 in PKU) emissions (Peng et al., 2019). The OC and BC emissions from FMS accounted for only 1.48–2.18 %
540 and 0.16–0.20 % of their national total emission. Thus, the OC and BC emissions from FMS were generally
541 meager. During the key period like the CNE, the contributions of FMS to the total OC and BC emissions can
542 rise to 2.25–3.46% and 1.05–1.58%. In key areas, the contribution rates would be relatively higher. For
543 instance, in 2014 CNE, FMS contributed 6.31% of OC emission in the Sichuan Basin and 2.91% of BC
544 emission in the Jiangxi-Hunan area. However, it should be noted that, the fireworks were always set off from
545 about 20:00 to 00:00 in CNE, so the intensive emission amounts could be considered at these times.
546 Therefore, the contribution of fireworks burning to CA in the atmosphere during CNE and CSF is still open
547 to debate.

548 It has caused widespread controversy that why the governments do not control the emissions from
549 industries and vehicles in CNE but emphasize the control of emissions from fireworks burning. The public
550 can not accept or believe the emissions from fireworks can lead to serious air pollution, which is the key
551 reason why they can not be completely eradicated in cities. From this study, the CA emissions are limited
552 compared with those from residential sources. An interesting question that the atmospheric scientists needed
553 to be solved in the future is that if the fireworks burning were not controlled, how many air pollutants from
554 other main sources of cities should be controlled alternatively.

555 **Summary and conclusions**

556 The absence of anthropogenic sources in the existing inventory prevents people from drawing accurate



557 conclusions about the control of short-term pollution. To calculate the emissions from these sources which
558 are difficult to estimate, we construct an emission inventory establishment method. We use the multiple
559 proxy data, such as the questionnaire, various statistics, and points of interest, to build a dataset of the activity
560 of five missing sources (FMS, including cooking, fireworks burning, sacrificial incenses, and joss paper
561 burning, and barbecue). The carbonaceous aerosols (CA) emission factors were tested in our lab using a self-
562 designed sampling platform. The OC, EC, and BC EFs varied in 5.86–203 mg kg⁻¹, 0.003–12.4 mg kg⁻¹,
563 and 1.07–191 mg kg⁻¹, respectively. BrC absorption EFs were in the range of 0.01–6.05 m² kg⁻¹ (370 nm).
564 From 2000 to 2018, the activity of FMS emitted 4268–4919 t, 22.6–43.9 t, 213–324 t, and 14.7–35.6 Gm²
565 of OC, EC, BC, and BrC absorption cross-section (ACS_{BrC}, in 370 nm wavelength). Emissions from FMS
566 would concentrate on special festivals. For example, CA emission in Chinese New Year’ eve was more than
567 10.8 times of the yearly average value. The distribution of pollutants also showed great differences between
568 urban and rural regions due to the demographic, economic, and policy implications. There was a negative
569 correlation ($r = -0.97$, $p < 0.01$) between individual emissions and disposable income in rural areas and a
570 positive correlation ($r = 0.94$, $p < 0.01$) in urban areas. The policy implications led to a reduction of over
571 42.2% of CA emissions from FMS. This study complements the lack of inventory research of such missing
572 sources and provides the prerequisite for modeling studies. Meanwhile, we suggest that raising residents’
573 income can be a feasible solution when reducing FMS emissions sources that are difficult to control, such as
574 fireworks, can be controlled from the manufacturer's side by guiding them to develop more environmentally
575 friendly products. We also suppose that whether it is possible to control other emission sources for providing
576 the environmental capacity for the emissions of fireworks burning.

577 **Data availability**

578 The dataset generated in this work is available at <https://doi.org/10.6084/m9.figshare.19999991.v1>
579 (Cheng et al., 2022).

580 The POI data (points of barbecue restaurants, temples, common restaurants, firework shops) was from
581 the Open Platform of Amap (<https://lbs.amap.com/>). The China Forestry Statistical Yearbook (forest fires),
582 China Light Industry Yearbook (fireworks export volume), and statistical yearbook of each province (urban
583 and rural population, meat consumption, consumption expenditure, and disposable income) came from
584 <https://data.cnki.net>. The percentage of meals eaten out came from <https://www.chinanutri.cn/>. The China



585 Statistical Yearbook (popularizing rate of the range hood) came from <https://data.cnki.net>. The distribution
586 of population was from www.worldpop.org. The land use data was from <http://data.ess.tsinghua.edu.cn/>. The
587 annual fireworks sales came from <https://www.mem.gov.cn/gk/tjsj/>. The annual and monthly restaurant sales
588 came from <https://data.stats.gov.cn/>.

589 **Author contribution**

590 **Yi Cheng**: Experiments, Visualization, Writing - original draft. **Shaofei Kong**: Conceptualization,
591 Methodology, Supervision, Writing - review & editing. **Liquan Yao**: Experiments, Formal analysis. **Huang**
592 **Zheng**: Experiments, Formal analysis. **Jian Wu**: Formal analysis. **Qin Yan**: Experiments. **Shurui Zheng**:
593 Experiments. **Yao Hu**: Validation, **Zhenzhen Niu**: Experiments. **Yingying Yan**: Supervision. **Zhenxing**
594 **Shen**: Supervision. **Guofeng Shen**: Supervision. **Dantong Liu**: Supervision. **Shuxiao Wang**: Supervision.
595 **Shihua Qi**: Supervision.

596 **Competing interests**

597 The authors declare no competing financial interest.

598 **Acknowledgments**

599 This study was financially supported by the National Natural Science Foundation of China (41830965;
600 42077202), and the Key Program of the Ministry of Science and Technology of the People's Republic of
601 China (2016YFA0602002; 2017YFC0212602).
602



603

604 **References**

605 Amouei Torkmahalleh, M., Ospanova, S., Baibatyrova, A., Nurbay, S., Zhanakhmet, G., & Shah, D. (2018).

606 Contributions of burner, pan, meat and salt to PM emission during grilling. *Environmental Research*,

607 *164*, 11–17. <https://doi.org/10.1016/j.envres.2018.01.044>

608 Batis, C., Sotres-Alvarez, D., Gordon-Larsen, P., Mendez, M. A., Adair, L., & Popkin, B. (2014).

609 Longitudinal analysis of dietary patterns in Chinese adults from 1991 to 2009. *British Journal of*

610 *Nutrition*, *111*(8), 1441–1451. <https://doi.org/10.1017/S0007114513003917>

611 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., et al. (2013). Bounding

612 the role of black carbon in the climate system: A scientific assessment. *Journal of Geophysical Research:*

613 *Atmospheres*, *118*(11), 5380–5552. <https://doi.org/10.1002/jgrd.50171>

614 Camilleri, R., & Vella, A. J. (2016). Emission factors for aerial pyrotechnics and use in assessing

615 environmental impact of firework displays: Case study from Malta. *Propellants, Explosives,*

616 *Pyrotechnics*, *41*(2), 273–280. <https://doi.org/10.1002/prop.201500205>

617 Cao, X., Zhang, X., Tong, D. Q., Chen, W., Zhang, S., Zhao, H., & Xiu, A. (2018). Review on

618 physicochemical properties of pollutants released from fireworks: Environmental and health effects and

619 prevention. *Environmental Reviews*, *26*(2), 133–155. <https://doi.org/10.1139/er-2017-0063>

620 Cheng, S., Wang, G., Lang, J., Wen, W., Wang, X., & Yao, S. (2016). Characterization of volatile organic

621 compounds from different cooking emissions. *Atmospheric Environment*, *145*, 299–307.

622 <https://doi.org/10.1016/j.atmosenv.2016.09.037>

623 Cheng, Y., Kong, S., Yan, Q., Liu, H., Wang, W., Chen, K., et al. (2019). Size-segregated emission factors

624 and health risks of PAHs from residential coal flaming/smoldering combustion. *Environmental Science*



- 625 *and Pollution Research*, 26, 31793–31803. <https://doi.org/10.1007/s11356-019-06340-2>
- 626 Cheng, Y., Kong, S., Yao, L., Zheng, H., Wu, J., Yan, Q., et al. (2022). Multi-year emission of carbonaceous
627 aerosols from cooking, fireworks burning, sacrificial incenses, joss paper burning, and barbecue and
628 the key driving forces in China. *figshare. Dataset*. <https://doi.org/10.6084/m9.figshare.19999991.v1>
- 629 Chiang, K.-C., & Liao, C.-M. (2006). Heavy incense burning in temples promotes exposure risk from
630 airborne PMs and carcinogenic PAHs. *Science of the Total Environment*, 372(1), 64–75.
631 <https://doi.org/10.1016/j.scitotenv.2006.08.012>
- 632 Corbin, J. C., Pieber, S. M., Czech, H., Zanatta, M., Jakobi, G., Massabò, D., et al. (2018). Brown and black
633 carbon emitted by a marine engine operated on heavy fuel oil and distillate fuels: Optical properties,
634 size distributions, and emission factors. *Journal of Geophysical Research: Atmospheres*, 123(11), 6175–
635 6195. <https://doi.org/10.1029/2017JD027818>
- 636 Croteau, G., Dills, R., Beaudreau, M., & Davis, M. (2010). Emission factors and exposures from ground-
637 level pyrotechnics. *Atmospheric Environment*, 44(27), 3295–3303.
638 <https://doi.org/10.1016/j.atmosenv.2010.05.048>
- 639 Ding, J., Guo, J., Wang, L., Chen, Y., Hu, B., Li, Y., et al. (2019). Cellular responses to exposure to outdoor
640 air from the Chinese Spring Festival at the air–liquid interface. *Environmental Science & Technology*,
641 53(15), 9128–9138. <https://doi.org/10.1021/acs.est.9b00399>
- 642 Ding, Q., Liu, J., Lu, Y., Wang, Y., Lu, F., & Shi, J. (2014). Research and development of an on-line
643 carbonaceous aerosol analyzer (In Chinese). *Chinese Journal of Scientific Instrument*, 35(06), 1246–
644 1253. <https://doi.org/10.19650/j.cnki.cjsi.2014.06.007>
- 645 Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., et al. (2015). The “dual-spot”
646 Aethalometer: An improved measurement of aerosol black carbon with real-time loading compensation.



- 647 *Atmospheric Measurement Techniques*, 8(5), 1965–1979. <https://doi.org/10.5194/amt-8-1965-2015>
- 648 Fan, S., Li, Y., & Liu, C. (2021). Are environmentally friendly fireworks really “green” for air quality? A
649 study from the 2019 National Day fireworks display in Shenzhen. *Environmental Science & Technology*,
650 55(6), 3520–3529. <https://doi.org/10.1021/acs.est.0c03521>
- 651 Feng, Y., Ramanathan, V., & Kotamarthi, V. R. (2013). Brown carbon: A significant atmospheric absorber of
652 solar radiation? *Atmospheric Chemistry and Physics*, 13(17), 8607–8621. [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-13-8607-2013)
653 13-8607-2013
- 654 Gong, P., Li, X., & Zhang, W. (2019). 40-Year (1978–2017) human settlement changes in China reflected by
655 impervious surfaces from satellite remote sensing. *Science Bulletin*, 64(11), 756–763.
656 <https://doi.org/10.1016/j.scib.2019.04.024>
- 657 Gong, P., Chen, B., Li, X., Liu, H., Wang, J., Bai, Y., et al. (2020). Mapping essential urban land use
658 categories in China (EULUC-China): Preliminary results for 2018. *Science Bulletin*, 65(3), 182–187.
659 <https://doi.org/10.1016/j.scib.2019.12.007>
- 660 Ho, C.-C., Chan, C.-C., Chio, C.-P., Lai, Y.-C., Chang-Chien, G.-P., Chow, J. C., et al. (2016). Source
661 apportionment of mass concentration and inhalation risk with long-term ambient PCDD/Fs
662 measurements in an urban area. *Journal of Hazardous Materials*, 317, 180–187.
663 <https://doi.org/10.1016/j.jhazmat.2016.05.059>
- 664 Hu, R., Wang, S., Zheng, H., Zhao, B., Liang, C., Chang, X., et al. (2021). Variations and sources of organic
665 aerosol in winter Beijing under markedly reduced anthropogenic activities during COVID-2019.
666 *Environmental Science & Technology*, acs.est.1c05125. <https://doi.org/10.1021/acs.est.1c05125>
- 667 Huang, D. D., Zhu, S., An, J., Wang, Q., Qiao, L., Zhou, M., et al. (2021). Comparative assessment of



- 668 cooking emission contributions to urban organic aerosol using online molecular tracers and Aerosol
669 Mass Spectrometry measurements. *Environmental Science & Technology*, 55(21), 14526–14535.
670 <https://doi.org/10.1021/acs.est.1c03280>
- 671 Huang, R.-J., Wang, Y., Cao, J., Lin, C., Duan, J., Chen, Q., et al. (2019). Primary emissions versus secondary
672 formation of fine particulate matter in the most polluted city (Shijiazhuang) in North China.
673 *Atmospheric Chemistry and Physics*, 19(4), 2283–2298. <https://doi.org/10.5194/acp-19-2283-2019>
- 674 Huang, Y., Shen, H., Chen, Y., Zhong, Q., Chen, H., Wang, R., et al. (2015). Global organic carbon emissions
675 from primary sources from 1960 to 2009. *Atmospheric Environment*, 122, 505–512.
676 <https://doi.org/10.1016/j.atmosenv.2015.10.017>
- 677 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., et al. (2019).
678 EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012.
679 *Earth System Science Data*, 11(3), 959–1002. <https://doi.org/10.5194/essd-11-959-2019>
- 680 Jetter, J. J., Guo, Z., McBrian, J. A., & Flynn, M. R. (2002). Characterization of emissions from burning
681 incense. *Science of the Total Environment*, 295(1–3), 51–67. [https://doi.org/10.1016/S0048-](https://doi.org/10.1016/S0048-9697(02)00043-8)
682 [9697\(02\)00043-8](https://doi.org/10.1016/S0048-9697(02)00043-8)
- 683 Jilla, A., & Kura, B. (2017). Particulate matter and carbon monoxide emission factors from incense burning.
684 *Environment Pollution and Climate Change*, 01(04). <https://doi.org/10.4172/2573-458X.1000140>
- 685 Jin, W., Zhi, G., Zhang, Y., Wang, L., Guo, S., Zhang, Y., et al. (2021). Toward a national emission inventory
686 for the catering industry in China. *Science of the Total Environment*, 754, 142184.
687 <https://doi.org/10.1016/j.scitotenv.2020.142184>
- 688 Keller, F., & Schragen, C. (2021). Determination of particulate matter emission factors of common



- 689 pyrotechnic articles. *Propellants, Explosives, Pyrotechnics*, 46(5), 825–842.
690 <https://doi.org/10.1002/prop.202000292>
- 691 Kong, S. F., Li, L., Li, X. X., Yin, Y., Chen, K., Liu, D. T., et al. (2015). The impacts of firework burning at
692 the Chinese Spring Festival on air quality: Insights of tracers, source evolution and aging processes.
693 *Atmospheric Chemistry and Physics*, 15(4), 2167–2184. <https://doi.org/10.5194/acp-15-2167-2015>
- 694 Kuo, S.-C., Tsai, Y. I., & Sopajaree, K. (2016). Emission characteristics of carboxylates in PM_{2.5} from
695 incense burning with the effect of light on acetate. *Atmospheric Environment*, 138, 125–134.
696 <https://doi.org/10.1016/j.atmosenv.2016.05.004>
- 697 Lai, Y., & Brimblecombe, P. (2020). Changes in air pollution and attitude to fireworks in Beijing.
698 *Atmospheric Environment*, 231, 117549. <https://doi.org/10.1016/j.atmosenv.2020.117549>
- 699 Lao, J.-Y., Xie, S.-Y., Wu, C.-C., Bao, L.-J., Tao, S., & Zeng, E. Y. (2018). Importance of dermal absorption
700 of polycyclic aromatic hydrocarbons derived from barbecue fumes. *Environmental Science &*
701 *Technology*, 52(15), 8330–8338. <https://doi.org/10.1021/acs.est.8b01689>
- 702 Laskin, A., Laskin, J., & Nizkorodov, S. A. (2015). Chemistry of atmospheric brown carbon. *Chemical*
703 *Reviews*, 115(10), 4335–4382. <https://doi.org/10.1021/cr5006167>
- 704 Lee, S.-C., & Wang, B. (2004). Characteristics of emissions of air pollutants from burning of incense in a
705 large environmental chamber. *Atmospheric Environment*, 38(7), 941–951.
706 <https://doi.org/10.1016/j.atmosenv.2003.11.002>
- 707 Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., et al. (2017). Anthropogenic emission inventories in
708 China: a review. *National Science Review*, 4(6), 834–866. <https://doi.org/10.1093/nsr/nwx150>
- 709 Liakakou, E., Kaskaoutis, D. G., Grivas, G., Stavroulas, I., Tsagkaraki, D., Paraskevopoulou, D., et al. (2020).



- 710 Long-term brown carbon spectral characteristics in a Mediterranean city (Athens). *Science of the Total*
711 *Environment*, 708, 135019. <https://doi.org/10.1016/j.scitotenv.2019.135019>
- 712 Lin, M.-D., Rau, J.-Y., Tseng, H.-H., Wey, M.-Y., Chu, C.-W., Lin, Y.-H., et al. (2008). Characterizing PAH
713 emission concentrations in ambient air during a large-scale joss paper open-burning event. *Journal of*
714 *Hazardous Materials*, 156(1–3), 223–229. <https://doi.org/10.1016/j.jhazmat.2007.12.015>
- 715 Lin, P., He, W., Nie, L., Schauer, J. J., Wang, Y., Yang, S., & Zhang, Y. (2019). Comparison of PM_{2.5} emission
716 rates and source profiles for traditional Chinese cooking styles. *Environmental Science and Pollution*
717 *Research*, 26, 21239–21252. <https://doi.org/10.1007/s11356-019-05193-z>
- 718 Lin, P., Gao, J., He, W., Nie, L., Schauer, J. J., Yang, S., et al. (2021). Estimation of commercial cooking
719 emissions in real-world operation: Particulate and gaseous emission factors, activity influencing and
720 modelling. *Environmental Pollution*, 289, 117847. <https://doi.org/10.1016/j.envpol.2021.117847>
- 721 Liu, J., Zhang, F., Xu, W., Chen, L., Ren, J., Jiang, S., et al. (2021). A large impact of cooking organic aerosol
722 (COA) on particle hygroscopicity and CCN activity in urban atmosphere. *Journal of Geophysical*
723 *Research: Atmospheres*, 126(8). <https://doi.org/10.1029/2020JD033628>
- 724 Martinsson, J., Eriksson, A. C., Nielsen, I. E., Malmberg, V. B., Ahlberg, E., Andersen, C., & Lindgren, R.
725 (2015). Impacts of combustion conditions and photochemical processing on the light absorption of
726 biomass combustion aerosol. *Environmental Science & Technology*, 49, 14663–14671.
727 <https://doi.org/10.1021/acs.est.5b03205>
- 728 McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., et al. (2020). A
729 global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific
730 sources (1970-2017): An application of the Community Emissions Data System (CEDs). *Earth System*



- 731 *Science Data*, 12(4), 3413–3442. <https://doi.org/10.5194/essd-12-3413-2020>
- 732 Meng, W., Zhong, Q., Chen, Y., Shen, H., Yun, X., Smith, K. R., et al. (2019). Energy and air pollution
733 benefits of household fuel policies in northern China. *Proceedings of the National Academy of Sciences*,
734 116(34), 16773–16780. <https://doi.org/10.1073/pnas.1904182116>
- 735 Pang, N., Gao, J., Zhao, P., Wang, Y., Xu, Z., & Chai, F. (2021). The impact of fireworks control on air
736 quality in four Northern Chinese cities during the Spring Festival. *Atmospheric Environment*, 244,
737 117958. <https://doi.org/10.1016/j.atmosenv.2020.117958>
- 738 Park, R. J., Kim, M. J., Jeong, J. I., Youn, D., & Kim, S. (2010). A contribution of brown carbon aerosol to
739 the aerosol light absorption and its radiative forcing in East Asia. *Atmospheric Environment*, 44(11),
740 1414–1421. <https://doi.org/10.1016/j.atmosenv.2010.01.042>
- 741 Peng, L., Zhang, Q., Yao, Z., Mauzerall, D. L., Kang, S., Du, Z., et al. (2019). Underreported coal in statistics:
742 A survey-based solid fuel consumption and emission inventory for the rural residential sector in China.
743 *Applied Energy*, 235, 1169–1182. <https://doi.org/10.1016/j.apenergy.2018.11.043>
- 744 Que, D. E., Hou, W.-C., Lin, S.-L., Tsai, Y.-I., Lu, I.-C., Wang, L.-C., et al. (2019). Emission of carbonyl
745 compounds from cooking oil fumes in the night market areas. *Aerosol and Air Quality Research*, 19(7),
746 1566–1578. <https://doi.org/10.4209/aaqr.2019.06.0289>
- 747 Ramanathan, V., & Carmichael, G. (2008). Global and regional climate changes due to black carbon. *Nature*
748 *Geoscience*, 1(4), 221–227. <https://doi.org/10.1038/ngeo156>
- 749 Saleh, R., Hennigan, C. J., McMeeking, G. R., Chuang, W. K., Robinson, E. S., Coe, H., et al. (2013).
750 Absorptivity of brown carbon in fresh and photo-chemically aged biomass-burning emissions.
751 *Atmospheric Chemistry and Physics*, 13(15), 7683–7693. <https://doi.org/10.5194/acp-13-7683-2013>



- 752 See, S. W., & Balasubramanian, R. (2011). Characterization of fine particle emissions from incense burning.
753 *Building and Environment*, 46, 1074–1080. <https://doi.org/10.1016/j.buildenv.2010.11.006>
- 754 Shen, H., Tsai, C.-M., Yuan, C.-S., Jen, Y.-H., & Ie, I.-R. (2017). How incense and joss paper burning during
755 the worship activities influences ambient mercury concentrations in indoor and outdoor environments
756 of an Asian temple? *Chemosphere*, 167, 530–540. <https://doi.org/10.1016/j.chemosphere.2016.09.159>
- 757 Shi, Y., Zhang, N., Gao, J., Li, X., & Cai, Y. (2011). Effect of fireworks display on perchlorate in air aerosols
758 during the Spring Festival. *Atmospheric Environment*, 45(6), 1323–1327.
759 <https://doi.org/10.1016/j.atmosenv.2010.11.056>
- 760 Sun, J., Zhi, G., Hitzenberger, R., Chen, Y., Tian, C., Zhang, Y., et al. (2017). Emission factors and light
761 absorption properties of brown carbon from household coal combustion in China. *Atmospheric*
762 *Chemistry and Physics*, 17(7), 4769–4780. <https://doi.org/10.5194/acp-17-4769-2017>
- 763 Sun, Y., He, Y., Kuang, Y., Xu, W., Song, S., Ma, N., et al. (2020). Chemical differences between PM₁ and
764 PM_{2.5} in highly polluted environment and implications in air pollution studies. *Geophysical Research*
765 *Letters*, 47(5). <https://doi.org/10.1029/2019GL086288>
- 766 Sun, Y. L., Wang, Z. F., Fu, P. Q., Yang, T., Jiang, Q., Dong, H. B., et al. (2013). Aerosol composition, sources
767 and processes during wintertime in Beijing, China. *Atmospheric Chemistry and Physics*, 13(9), 4577–
768 4592. <https://doi.org/10.5194/acp-13-4577-2013>
- 769 Tanda, S., Licbinsky, R., Hegrova, J., & Goessler, W. (2019). Impact of New Year’s Eve fireworks on the
770 size resolved element distributions in airborne particles. *Environment International*, 128, 371–378.
771 <https://doi.org/10.1016/j.envint.2019.04.071>
- 772 Tian, J., Wang, Q., Ni, H., Wang, M., Zhou, Y., Han, Y., et al. (2019). Emission characteristics of primary



- 773 brown carbon absorption from biomass and coal burning: Development of an optical emission inventory
774 for China. *Journal of Geophysical Research: Atmospheres*, 124, 1879–1893.
775 <https://doi.org/10.1029/2018JD029352>
- 776 Tong, D., Cheng, J., Liu, Y., Yu, S., Yan, L., Hong, C., et al. (2020). Dynamic projection of anthropogenic
777 emissions in China: methodology and 2015–2050 emission pathways under a range of socio-economic,
778 climate policy, and pollution control scenarios. *Atmospheric Chemistry and Physics*, 20, 5729–5757.
779 <https://doi.org/10.5194/acp-20-5729-2020>
- 780 Vecchi, R., Bernardoni, V., Cricchio, D., D’Alessandro, A., Fermo, P., Lucarelli, F., et al. (2008). The impact
781 of fireworks on airborne particles. *Atmospheric Environment*, 42(6), 1121–1132.
- 782 Venkataraman, C., Habib, G., Eiguren-Fernandez, A., Miguel, A. H., & Friedlander, S. K. (2005). Residential
783 biofuels in South Asia: Carbonaceous aerosol emissions and climate impacts. *Science*, 307(5714),
784 1454–1456. <https://doi.org/10.1126/science.1104359>
- 785 Wang, G., Cheng, S., Wei, W., Wen, W., Wang, X., & Yao, S. (2015a). Chemical characteristics of fine
786 particles emitted from different Chinese cooking styles. *Aerosol and Air Quality Research*, 15(6), 2357–
787 2366. <https://doi.org/10.4209/aaqr.2015.02.0079>
- 788 Wang, H., Xiang, Z., Wang, L., Jing, S., Lou, S., Tao, S., et al. (2018a). Emissions of volatile organic
789 compounds (VOCs) from cooking and their speciation: A case study for Shanghai with implications for
790 China. *Science of the Total Environment*, 621, 1300–1309.
791 <https://doi.org/10.1016/j.scitotenv.2017.10.098>
- 792 Wang, L., Xiang, Z., Stevanovic, S., Ristovski, Z., Salimi, F., Gao, J., et al. (2017). Role of Chinese cooking
793 emissions on ambient air quality and human health. *Science of the Total Environment*, 589, 173–181.



- 794 <https://doi.org/10.1016/j.scitotenv.2017.02.124>
- 795 Wang, L., Zheng, X., Stevanovic, S., Wu, X., Xiang, Z., Yu, M., & Liu, J. (2018b). Characterization
796 particulate matter from several Chinese cooking dishes and implications in health effects. *Journal of*
797 *Environmental Sciences*, 72, 98–106. <https://doi.org/10.1016/j.jes.2017.12.015>
- 798 Wang, Q., Sun, Y., Jiang, Q., Du, W., Sun, C., Fu, P., & Wang, Z. (2015b). Chemical composition of aerosol
799 particles and light extinction apportionment before and during the heating season in Beijing, China.
800 *Journal of Geophysical Research: Atmospheres*, 120(24), 12708–12722.
801 <https://doi.org/10.1002/2015JD023871>
- 802 Wang, R., Tao, S., Wang, W., Liu, J., Shen, H., Shen, G., et al. (2012). Black carbon emissions in China from
803 1949 to 2050. *Environmental Science & Technology*, 46(14), 7595–7603.
804 <https://doi.org/10.1021/es3003684>
- 805 Wang, R., Tao, S., Shen, H., Huang, Y., Chen, H., Balkanski, Y., et al. (2014a). Trend in global black carbon
806 emissions from 1960 to 2007. *Environmental Science & Technology*, 48(12), 6780–6787.
807 <https://doi.org/10.1021/es5021422>
- 808 Wang, X., Heald, C. L., Ridley, D. A., Schwarz, J. P., Spackman, J. R., Perring, A. E., et al. (2014b).
809 Exploiting simultaneous observational constraints on mass and absorption to estimate the global direct
810 radiative forcing of black carbon and brown carbon. *Atmospheric Chemistry and Physics*, 14(20),
811 10989–11010. <https://doi.org/10.5194/acp-14-10989-2014>
- 812 Wang, Y., Wang, Q., Ye, J., Li, L., Zhou, J., Ran, W., et al. (2020). Chemical composition and sources of
813 submicron aerosols in winter at a regional site in Beijing-Tianjin-Hebei region: Implications for the
814 Joint Action Plan. *Science of The Total Environment*, 719, 137547.



- 815 <https://doi.org/10.1016/j.scitotenv.2020.137547>
- 816 Wu, C., Bao, L., Guo, Y., Li, S., & Zeng, E. Y. (2015). Barbecue fumes: An overlooked source of health
817 hazards in outdoor settings. *Environmental Science & Technology*, 49(17), 10607–10615.
818 <https://doi.org/10.1021/acs.est.5b01494>
- 819 Wu, J., Kong, S., Wu, F., Cheng, Y., Zheng, S., Yan, Q., et al. (2018). Estimating the open biomass burning
820 emissions in central and eastern China from 2003 to 2015 based on satellite observation. *Atmospheric
821 Chemistry and Physics*, 18, 11623–11646. <https://doi.org/10.5194/acp-18-11623-2018>
- 822 Wu, J., Kong, S., Zeng, X., Cheng, Y., Yan, Q., Zheng, H., et al. (2021). First high-resolution emission
823 inventory of levoglucosan for biomass burning and non-biomass burning sources in China.
824 *Environmental Science & Technology*, 55(3), 1497–1507. <https://doi.org/10.1021/acs.est.0c06675>
- 825 Xiang, Z., Wang, H., Stevanovic, S., Jing, S., Lou, S., Tao, S., et al. (2017). Assessing impacts of factors on
826 carbonyl compounds emissions produced from several typical Chinese cooking. *Building and
827 Environment*, 125, 348–355. <https://doi.org/10.1016/j.buildenv.2017.08.045>
- 828 Xu M., Chao X., Zhang B., Liu S., Yin J., & Gan S. (2018). Study on the investigation, analysis and
829 countermeasures for the current status of mutton sheep industry in Xinjiang (In Chinese). *Acta
830 Ecologiae Animalis Domastici*, 39(3), 85–89.
- 831 Yan, Q., Kong, S., Yan, Y., Liu, H., Wang, W., Chen, K., et al. (2020). Emission and simulation of primary
832 fine and submicron particles and water-soluble ions from domestic coal combustion in China.
833 *Atmospheric Environment*, 224, 117308. <https://doi.org/10.1016/j.atmosenv.2020.117308>
- 834 Yang, C.-R., Ko, T.-H., Lin, Y.-C., Lee, S.-Z., Chang, Y.-F., & Hsueh, H.-T. (2013). Oyster shell reduces
835 PAHs and particulate matter from incense burning. *Environmental Chemistry Letters*, 11(1), 33–40.



- 836 <https://doi.org/10.1007/s10311-012-0374-2>
- 837 Yang, H.-H., Jung, R.-C., Wang, Y.-F., & Hsieh, L.-T. (2005). Polycyclic aromatic hydrocarbon emissions
838 from joss paper furnaces. *Atmospheric Environment*, 39(18), 3305–3312.
839 <https://doi.org/10.1016/j.atmosenv.2005.01.052>
- 840 Yao, L., Wang, D., Fu, Q., Qiao, L., Wang, H., Li, L., et al. (2019). The effects of firework regulation on air
841 quality and public health during the Chinese Spring Festival from 2013 to 2017 in a Chinese megacity.
842 *Environment International*, 126, 96–106. <https://doi.org/10.1016/j.envint.2019.01.037>
- 843 Zhang, A., Wang, Y., Zhang, Y., Weber, R. J., Song, Y., Ke, Z., & Zou, Y. (2020a). Modeling the global
844 radiative effect of brown carbon: A potentially larger heating source in the tropical free troposphere
845 than black carbon. *Atmospheric Chemistry and Physics*, 20(4), 1901–1920. [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-20-1901-2020)
846 [20-1901-2020](https://doi.org/10.5194/acp-20-1901-2020)
- 847 Zhang, L., Gao, Y., Wu, S., Zhang, S., Smith, K. R., Yao, X., & Gao, H. (2020b). Global impact of
848 atmospheric arsenic on health risk: 2005 to 2015. *Proceedings of the National Academy of Sciences*,
849 117(25), 13975–13982. <https://doi.org/10.1073/pnas.2002580117>
- 850 Zhang, L., Luo, Z., Du, W., Li, G., Shen, G., Cheng, H., & Tao, S. (2020c). Light absorption properties and
851 absorption emission factors for indoor biomass burning. *Environmental Pollution*, 267, 115652.
852 <https://doi.org/10.1016/j.envpol.2020.115652>
- 853 Zhang, L., Luo, Z., Li, Y., Chen, Y., Du, W., Li, G., et al. (2021a). Optically measured black and particulate
854 brown carbon emission factors from real-world residential combustion predominantly affected by fuel
855 differences. *Environmental Science & Technology*, 55(1), 169–178.
856 <https://doi.org/10.1021/acs.est.0c04784>



- 857 Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., et al. (2019a). Drivers of improved PM_{2.5} air
858 quality in China from 2013 to 2017. *Proceedings of the National Academy of Sciences*, 116(49), 24463–
859 24469. <https://doi.org/10.1073/pnas.1907956116>
- 860 Zhang, S., Zhong, L., Chen, X., Liu, Y., Zhai, X., Xue, Y., et al. (2019b). Emissions characteristics of
861 hazardous air pollutants from the incineration of sacrificial offerings. *Atmosphere*, 10(6), 332.
862 <https://doi.org/10.3390/atmos10060332>
- 863 Zhang, Y., Kong, S., Sheng, J., Zhao, D., Ding, D., Yao, L., et al. (2021b). Real-time emission and stage-
864 dependent emission factors/ratios of specific volatile organic compounds from residential biomass
865 combustion in China. *Atmospheric Research*, 248, 105189.
866 <https://doi.org/10.1016/j.atmosres.2020.105189>
- 867 Zhang, Z., Zhu, W., Hu, M., Liu, K., Wang, H., Tang, R., et al. (2021c). Formation and evolution of secondary
868 organic aerosols derived from urban-lifestyle sources: Vehicle exhaust and cooking emissions.
869 *Atmospheric Chemistry and Physics*, 21(19), 15221–15237. [https://doi.org/10.5194/acp-21-15221-](https://doi.org/10.5194/acp-21-15221-2021)
870 2021
- 871 Zhao, Y., Chen, C., & Zhao, B. (2018). Is oil temperature a key factor influencing air pollutant emissions
872 from Chinese cooking? *Atmospheric Environment*, 193, 190–197.
873 <https://doi.org/10.1016/j.atmosenv.2018.09.012>
- 874 Zhao, Y., Chen, C., & Zhao, B. (2019). Emission characteristics of PM_{2.5}-bound chemicals from residential
875 Chinese cooking. *Building and Environment*, 149, 623–629.
876 <https://doi.org/10.1016/j.buildenv.2018.12.060>
- 877 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., et al. (2018). Trends in China's anthropogenic



878 emissions since 2010 as the consequence of clean air actions. *Atmospheric Chemistry and Physics*, 18,
879 14095–14111. <https://doi.org/10.5194/acp-18-14095-2018>

880 Zhu, W., Guo, S., Zhang, Z., Wang, H., Yu, Y., Chen, Z., et al. (2021a). Mass spectral characterization of
881 secondary organic aerosol from urban cooking and vehicular sources. *Atmospheric Chemistry and*
882 *Physics*, 21(19), 15065–15079. <https://doi.org/10.5194/acp-21-15065-2021>

883 Zhu, W., Zhou, M., Cheng, Z., Yan, N., Huang, C., Qiao, L., et al. (2021b). Seasonal variation of aerosol
884 compositions in Shanghai, China: Insights from particle aerosol mass spectrometer observations.
885 *Science of The Total Environment*, 771, 144948. <https://doi.org/10.1016/j.scitotenv.2021.144948>

886 Zotter, P., Herich, H., Gysel, M., El-Haddad, I., Zhang, Y., Močnik, G., et al. (2017). Evaluation of the
887 absorption Ångström exponents for traffic and wood burning in the Aethalometer-based source
888 apportionment using radiocarbon measurements of ambient aerosol. *Atmospheric Chemistry and*
889 *Physics*, 17(6), 4229–4249. <https://doi.org/10.5194/acp-17-4229-2017>

890