

Global dataset on seagrass meadow structure, biomass ~~and~~, production ~~and~~ reproduction

Simone Strydom^{1,2}, [Roisin McCallum¹](#), [Anna Lafratta¹](#), Chanelle L. Webster¹, Caitlyn M. O’Dea¹, Nicole E. Said¹, ~~[Roisin McCallum¹](#)~~, ~~[Natasha Dunham¹](#)~~, Karina Inostroza³, Cristian Salinas¹, Samuel Billinghurst¹, ~~[Anna Lafratta¹](#)~~, Charlie M. Phelps¹, Connor Campbell¹, Connor Gorham¹, ~~[Natasha Dunham¹](#)~~, Rachele Bernasconi¹, Anna M. Frouws¹, Axel Werner¹, Federico Vitelli¹, Viena Puigcorbé^{1,4}, Alexandra D’Cruz¹, Kathryn M. McMahon¹, Jack Robinson¹, Megan J. Huggett⁵, Sian McNamara¹, Glenn A. Hyndes¹ and Oscar Serrano^{1,6*}

10 ¹Centre for Marine Ecosystems Research and School of Science, Edith Cowan University, Joondalup, 270 Joondalup Dr, WA 6027, Australia.

²Marine Science Program, Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, 17 Dick Perry Ave, Kensington, WA 6151, Australia.

³Biosfera, Associació d’Educació Ambiental, Catalonia, Spain

15 ⁴Institut de Ciències del Mar, Consejo Superior de Investigaciones Científicas, 08003 Barcelona, Spain.

⁵School of Environmental and Life Sciences, The University of Newcastle, 10 Chittaway Rd, Ourimbah, NSW 2258, Australia.

⁶Centro de Estudios Avanzados de Blanes, Consejo Superior de Investigaciones Científicas, Blanes, Spain.

20 *Correspondence to: Oscar Serrano (oserrano@ceab.csic.es)

Abstract. Seagrass meadows provide valuable socio-ecological ecosystem services, including a key role in climate change mitigation and adaptation. Understanding the natural history of seagrass meadows across environmental gradients is crucial to decipher the role of seagrasses in the global ocean. In this data collation, spatial and temporal patterns in seagrass meadow structure, biomass, ~~and~~ production ~~and~~ reproduction data are presented as a function of biotic and abiotic habitat characteristics. The biological traits compiled include measures of meadow structure (e.g., percent cover and shoot density), biomass (e.g., above-ground biomass), ~~and~~ production (e.g., shoot production), ~~and reproduction effort~~ (e.g.,

~~flowering intensity and seed bank density~~). Categorical factors include bioregion, geotype (coastal or estuarine), genera and year of sampling. This dataset contains data extracted from peer-reviewed publications published between 1975 and 2020 based on a Web of Science search, and includes 115 data variables across 12 seagrass genera. The dataset excludes data from mesocosm and field experiments, contains 141,773-271 data points extracted from 39064 publications, and it is publicly available on the PANGAEA® data repository (<https://doi.org/10.1594/PANGAEA.929968>). The top five most studied genera are *Zostera*, *Thalassia*, ~~*Halophila* and *Cymodocea*~~, *Halodule* and *Halophila* (8480% of data), and the least studied genera are *Phyllospadix*, *Amphibolis* and *Thalassodendron* (2.3% of data). The data hotspot bioregion is the Tropical Indo Pacific (25% of data) followed by the Tropical Atlantic (21%), whereas data for the other four bioregions are evenly spread (ranging between 13 and 156% of total data within each bioregion). From the data compiled, 5739% related to seagrass biomass and 33% to seagrass structure, while the least number of data were related to seagrass production (110% of data). This data collation can inform several research fields beyond seagrass ecology, such as the development of nature-based solutions for climate change mitigation, which include readership interested in blue carbon, engineering, fisheries, global change, conservation and policy.

45 **1 Introduction**

Approximately 65 million years ago, a group of marine angiosperms called seagrasses adapted to life within the coastal zone, and now, they rank among the most valuable ecosystems globally. Seagrasses encompass ~72 species within 12 genera spread across all continents except Antarctica (Short et al., 2011). Seagrasses are recognised as highly productive habitats that provide multiple ecosystem services relevant to human wellbeing, such as biodiversity, fisheries, sediment stabilisation and nutrient cycling across the coastal zone (Ascioti et al., 2022; Lamb et al., 2017; McMahan et al., 2013; Nordlund et al., 2016; Unsworth et al., 2018). (McMahan et al., 2013; Nordlund et al., 2016; Unsworth et al., 2018). Furthermore, the high primary production rates and capacity of seagrasses to sequester carbon is

relevant to mitigating climate change, while their role in stabilising the substrate, ameliorating
55 hydrodynamic energy and nourishing beaches with biogenic sands contributes to climate change
adaptation against storms and sea-level rise (Duarte et al., 2013).

Seagrass research initially focussed on understanding biology, distribution, ecology, taxonomy, and
phenology. More recently, the socio-ecological value of seagrass ecosystem services has received
recognition, in part owing to the extensive losses of seagrasses globally. Since the beginning of the 20th
60 century, widespread loss of seagrass meadows has been estimated at 0.9% yr⁻¹, linked to a variety of
factors including impacts associated with alterations to key drivers of growth (e.g., irradiance and
temperature) resulting from sediment loading, eutrophication, extreme climate events and flooding (Hall
et al., 1999; Short et al., 2011; Strydom et al., 2020; Waycott et al., 2009). Recent conservation and
management actions have resulted in the deceleration and reversal of declining trends in some locations
65 (de los Santos et al., 2019).

Duarte and Chiscano (1999) conducted a review on seagrass biomass and production, which has greatly
contributed to the advancement of seagrass research. Information on seagrass meadows structure,
production, biomass and reproduction is essential to understand the role of seagrasses in the global
ocean, while providing insights for developing restoration initiatives, informing management and
70 ultimately contributing to their conservation (Unsworth et al., 2018). Therefore, understanding global
patterns in the functioning of threatened natural ecosystems such as seagrass meadows, is crucial to
inform management strategies to protect natural assets (Cullen-Unsworth et al., 2014; Hoegh-Guldberg
and Bruno, 2010). Since Duarte and Chiscano (1999), new information across hundreds of peer-
reviewed manuscripts (past 24 years, 1996-2020) has not been synthesised and made available online,
75 which precludes gathering new knowledge around seagrass natural history based on data synthesis
studies. Indeed, data on seagrass reproduction has never been compiled.

In this review, data on key variables on seagrass meadow structure, biomass ~~and~~ production, ~~and~~
~~reproduction~~ published between 1975 and 2020 (data collected between 1972~~64~~ and 2020~~19~~) are
presented as a function of biotic and abiotic habitat characteristics. The main goals of this review are to

80 synthesize current literature on seagrass ecology to facilitate further multidisciplinary research, and to
identify research gaps and provide recommendations for future research. The dataset provides baseline
data that can inform science, management and policy. In particular, it provides critical and basic
knowledge to inform traditional seagrass biology and ecology fields, but also can contribute to advance
knowledge in other disciplines including fisheries, biodiversity, conservation, coastal biogeochemistry,
85 and emerging fields such as the Blue Economy.

2 Data compilation

2.1 Literature search

In order to create a global seagrass database containing relevant data on seagrass meadow structure,
biomass and production, ~~and reproduction~~, a Web of Science (www.webofknowledge.com) search was
90 conducted in June 2020 using these search terms for the growth, production and biomass variables:
((TS=((Seagrass* OR eelgrass OR SAV OR Amphibolis OR Cymodocea OR Enhalus OR Halophila
OR Halodule OR Posidonia OR Phyllospadix OR Ruppia OR Thalassia OR Thalassodendron OR
Zostera) AND (product* OR biomass OR growth OR exten))))

95 ~~Then another search for reproduction variables using these terms: ((TS=((Seagrass* OR eelgrass OR
SAV OR Amphibolis OR Cymodocea OR Enhalus OR Halophila OR Halodule OR Posidonia OR
Phyllospadix OR Ruppia OR Thalassia OR Thalassodendron OR Zostera) AND (germinat* OR reprod*
OR seed* OR flower* OR fruit* OR dispersal * OR gra\$ing*))).~~

Only data from peer-reviewed manuscripts was included and thereby, the dataset compiled excludes
100 data from non-peer reviewed manuscripts and reports. We acknowledge that our search has likely
missed a small portion of the peer-reviewed data published to date, owing to the use of different terms
across research fields, and the use of a single search engine WOS to conduct the review. Data for 115
variables (mean values) were extracted (Table 1, see 2.2) and compiled in a database
(<https://doi.org/10.1594/PANGAEA.929968>). These variables were selected based on their widespread
105 study among seagrass habitats, and to their usefulness for quantifying seagrass condition across papers

Formatted: Line spacing: 1.5 lines

with different aims (~~i.e., monitoring condition vs reproductive effort~~) (Short & Coles, 2001). Standardised units (spatially i.e., m² and temporally i.e., day⁻¹) are reported. Note that data from mesocosm experiments, field experiments with the exception of control sites, or meadows altered by direct anthropogenic disturbances (i.e., aquaculture, anchoring or dredging) were dismissed as these were considered as impacted meadows and were unlikely to reflect meadows in a 'natural' condition. Note that seagrass species were recorded following accepted convention as per Short et al. 2011 (e.g., *Zostera capricorni*, *Z. mucronata*, *Z. novaezealandica* were named as *Z. muelleri*).

2.2. Seagrass structure, biomass, production and reproduction variables

The ~~115~~ variables extracted from the peer-reviewed literature were classified within ~~threefour~~ categories: seagrass meadow structure (~~three3~~ variables), biomass (~~three3~~ variables) and, production (~~five5~~ variables) ~~and reproduction (4 variables)~~.

- Seagrass meadow structure: percent cover (%), shoot density (no. shoots m⁻²), leaf density (no. leaves m⁻²).
- Seagrass biomass: above-ground biomass dry weight (DW) (g DW m⁻²), below-ground biomass (g DW m⁻²) and total biomass (g DW m⁻²).
- Seagrass production: shoot production (g DW m⁻² day⁻¹), leaf production (g DW m⁻² day⁻¹), and above-ground production (g DW m⁻² day⁻¹), below-ground production (g DW m⁻² day⁻¹) and total production (g DW m⁻² day⁻¹).
- ~~Seagrass reproduction: flowering intensity or flowering shoots (no. flowers m⁻²), fruit density (no. fruits m⁻²), seed bank density (no. seeds m⁻²) and seedling density (no. seedlings m⁻²).~~

For all these ~~115~~ variables, relevant data points were extracted from results text, tables and when appropriate from figures using a web based tool that allow the extraction of data from plots, images and maps ([WebPlotDigitizer: https://automeris.io/WebPlotDigitizer/](https://automeris.io/WebPlotDigitizer/)). Datapoint in an individual row can be a mean of multiple replicates or a single unique measurement point for that variable and location. Other relevant spatial and site information was also extracted including the latitude and longitude (decimal

degrees), seagrass bioregion according to Short et al. 2007 (Fig. 1), geotype (coastal or estuarine
135 geomorphology), seagrass genera, the year of sampling when reported, and the doi of the publication
containing the data. When sampling site coordinates were not reported in the publication, study location
maps were consulted if applicable and corresponding coordinates estimated using *Google Earth*.
Similarly, geotype was classified as estuarine if the study site was on close proximity to riverine input or
within a coastal lagoon, conversely if there were no rivers nearby or the study site was located within an
140 embayment then it was considered coastal. ~~For the flowering intensity variable, reproductive shoots were
included in this dataset variable (i.e., studies on *Ruppia* counted reproductive shoots and as these had
flowers on them, they were considered an analogous term). Furthermore, if flowers were identified as
male or female in studies, they were included in the dataset as total number of flowers per m² regardless
of gender. Indeed, details on density of flowers, spathes, inflorescence shoots and reproductive shoots~~
145 ~~where combined into a single variable (i.e., flowering intensity).~~ If publications included data on above-
ground biomass and below-ground biomass for the same study site, these two values were summed to
estimate a value of total seagrass biomass. Publications that reported growth or production expressed as
grams of carbon were excluded. When sampling was conducted over multiple years, the year of sampling
was left blank and not reported in the dataset.

150 The seagrass natural history information reported and the way it was reported has evolved during the 45
years of research compiled. Overall, early publications provided comprehensive details regarding ~~the
description of flowers, seeds and fruits~~ meadow structure and production, while sampling procedures
were not clearly described. Later on, the sampling strategies and data reporting became more
155 standardized and comprehensive.

2.3 Statistical analyses

Descriptive parameters (e.g., count of data and publications, minimum, maximum and median values)
for all 15 variables were compiled. Median values are reported instead of mean values because the data
for most of the variables studied is not normally distributed. Boxplots for four key variables sorted by
160 bioregion and genera were produced in R using the ggplot2 package (Wickham, 2016) (version 4.0.1, R

Core Team 2020). In order to spatially illustrate the dataset, maps were also created in R, using the leaflet package (Graul, 2016).

3 Results and discussion

The highest number of data points were collected in year 2018, while the lowest occurred in 1975 (Fig. 2). Overall, all four data categories were represented well over time (1972–2019), with biomass data present in the majority of papers consistently over time, meadow structure data encompassing a larger proportion of data over the last decade, and reproduction and production data being the least studied categories. Data was extracted from a total 390 peer-reviewed publications, with approximately 66% of the studies were conducted in coastal marine areas (n = 263), with the remaining 32% of studies conducted in estuarine areas (n = 120), and 7 studies conducted in both coastal and estuarine areas (2%).

3.2 Spatial distribution of seagrass data

The seagrass database includes information collected across 115 variables on seagrass structure, biomass and production and reproduction from all 12 seagrass genera described to date, spanning all continents except Antarctica (Fig. 3). Based on the count of data, the top five most studied genera making up to 84% of the database were *Zostera* (n = 5,573), *Thalassia* (n = 2,081), *Cymodocea* (n = 1,456), *Halodule* (n = 1,416) and *Halophila* (n = 1,343), and *Cymodocea* (n = 1,241). The least studied genera were *Amphibolis* (n = 58), *Thalassodendron* (n = 115), and *Phyllospadix* (n = 126). The predominance of *Zostera* data could be related to their broad global distribution, including European countries which were the pioneers of seagrass science, while the least studied genera are more geographically restricted (Fig. 3). The bioregion with highest number of data was the Tropical Indo Pacific (n = 2,950), which also included 10 of the 12 genera, illustrating the seagrass biodiversity of this bioregion. The number of data across the Temperate North Pacific (n = 1,912; four genera), Mediterranean (n = 1,960; five genera), Temperate North Atlantic (n = 1,893; five genera), Temperate Southern (n = 1,634; seven genera), and Tropical Atlantic (n = 2,946; six genera) bioregions was similar. There was up to 679-fold difference between the most

and least studied seagrass genera, but only a 2-fold difference between bioregions. Overall, data for the production category was the least reported (n = 1,536; 11%) followed by structure (n = 4,643; 32%), whereas biomass variable types were the most reported (n = 8,092; 57%). The most prevalent data type was seagrass biomass (n = 6,087; 52%), followed by structure (n = 3,256; 28%), reproduction (n = 1,181; 10%) and production (n = 1,249; 11%) (Fig. 4).

3.3 Variability in seagrass data among variables

The dataset compiled includes data on shoot density (n = 32,212,366), percent cover (n = 1,285,734), leaf density (n = 159,146), above-ground biomass (n = 3,389,519), below-ground biomass (n = 24,008,488), total biomass (n = 2,695,080), shoot production (n = 15,140), leaf production (n = 737,670), above-ground production (n = 492,652), below-ground production (n = 125,890) and total production (n = 258,188), flowering intensity (n = 706), fruit density (n = 55), seed bank density (n = 312) and seedling density (n = 108). Overall, production was the least reported variable type (n = 1,249), followed by reproduction (n = 1,181). Seagrass structure and biomass variable types were the most reported (n = 3,256 and 6,807, respectively). Across all dataset, shoot density ranged from 0.08 to 28,682 shoots m⁻² (median = 675.4), percent cover from 0.03 to 100% (median = 335.2), leaf density from 5.1 to 48,978 leaves m⁻² (median = 3,287), above-ground biomass from 0.0010 to 1,509.9 g DW m⁻² (median = 523.2), below-ground biomass from 0.0340 to 53,635,076 g DW m⁻² (median = 694.0), total biomass from 0.0010 to 3,393 g DW m⁻² (median = 13448), shoot production from 0.00106 to 23.45 g DW m⁻² day⁻¹ (median = 02.344), leaf production from 0.0012 to 277.165 g DW m⁻² day⁻¹ (median = 1.454), above-ground production from 0.00103 to 23.5 g DW m⁻² day⁻¹ (median = 1.55), below-ground production from 0.0159 to 34 g DW m⁻² day⁻¹ (median = 21.3.20) and total production from 0.00218 to 398.5 g DW m⁻² day⁻¹ (median = 23.50), flowering intensity from 0.10 to 6,000 flower m⁻² (median = 16.1), fruit density from 0.5 to 3,229 fruits m⁻² (median = 142), seed bank density from 2.7 to 10,028 seeds m⁻² (median = 138), and seedling density from 0.001 to 7,560 seedlings m⁻² (median = 20.9).

There was high variability in most variables using pooled data across bioregions and genera, and in the amount of data for each variable across bioregion, geotype and genera (Table 1). The values of some variables varied substantially across the six bioregions (Fig. 5). Median total biomass was highest in the

215 Mediterranean bioregion (28369 g DW m⁻²), while the lowest was in the ~~Temperate North~~
~~Atlantic~~~~Tropical Indo Pacific~~ bioregion (40961 g DW m⁻²). The highest median shoot density values
were recorded in the ~~Temperate North Atlantic~~~~Temperate Southern~~ bioregion (1606967 shoots m⁻²) and
the lowest in the Temperate North Pacific bioregion (279466 shoots m⁻²), ~~whereas~~. The highest median
total production values ~~were~~ recorded in the Temperate Southern bioregion (9366 g DW m⁻² day⁻¹);
220 ~~while the highest median flowering intensity values were recorded in the Mediterranean bioregion (90~~
~~flowers m⁻²)~~. Of all genera, ~~median~~ total biomass was ~~generally~~ highest for seagrasses with persistent
life history stages, ~~such as~~ *Posidonia* and *Enhalus* (Kilminster et al., 2015). ~~Phyllospadix~~*Posidonia* had
the highest median total biomass (20131056 g DW m⁻²), followed by ~~Phyllospadix~~*Amphibolis*
(1055845 g DW m⁻²) (Fig. 6). Median shoot density values were highest for *Phyllospadix* (6,593 shoots
225 m⁻²) followed by ~~Ruppia~~*Halodule* (64,34344 shoots m⁻²). Total production was highest for
Phyllospadix (median 223 g DW m⁻² day⁻¹), followed by *Syringodium* (median 9.3 g DW m⁻² day⁻¹).
~~The highest median flowering intensity was recorded for~~ *Syringodium* (1,983 flowers m⁻²), followed by
~~Ruppia~~ (765 flowers m⁻²) and *Halophila* (600 flowers m⁻²).

230 3.4 Significant gaps

This global collation of seagrass data has illustrated some gaps in our collective peer-reviewed
knowledge. Across seagrass' worldwide distribution, limited peer-reviewed data were found for the
eastern Mediterranean, and the coastlines of South America and Africa. Data for some seagrass
variables were spatially depauperate, such as seagrass production at high latitudes (<50°N and S),
235 including the Temperate North Atlantic. Overall, ~~seagrass production variables~~ ~~were~~ the least reported
~~variable type followed by reproduction~~. When considering data among seagrass genera, the least studied
were *Amphibolis* (n = 586), *Thalassodendron* (n = 11587), and *Phyllospadix* (n = 126), with gaps in
most variables. ~~There was also a lack of reproductive information for~~ *Amphibolis*, *Phyllospadix* and
Thalassodendron. Lastly, there was no peer-reviewed published data found for production of *Ruppia*.

240 4 Conclusions

This database encompassing peer-reviewed data collected over the last ~~4858~~ years provides an overview of seagrass distribution, biomass, production ~~and~~, structure ~~and reproduction~~ on a global scale. The top ~~five~~ most prevalent studied genera encompassing 84% of data were *Zostera* (mostly from the Temperate North Pacific), *Thalassia* ~~and~~ *Halodule* (Tropical Atlantic), ~~*Halophila* and *Cymodocea* and~~ *Halophila* (Tropical Indo Pacific and Mediterranean), and the least studied genera *Amphibolis*, *Thalassodendron* ~~and~~ *Phyllospadix* (2.3% of data). Data hotspots include the Tropical Indo Pacific bioregion (25% of dataset; ~~from 89 unique publications~~), whereas the ~~Tropical-Temperate Southern Atlantic~~ bioregion had the least amount of data (13% of data; ~~79 publications~~). The strengths on seagrass natural history knowledge focus on seagrass biomass (57.4% of data), ~~while the least number of data was related to seagrass reproduction (9% of data)~~. Our review can inform several research fields beyond seagrass ecology, such as the development of Nature-Based Solutions for climate change mitigation and adaptation and Blue Economy, which include readership interested in blue carbon, engineering, fisheries, global change, conservation and policy.

255 5 Data availability

Data archived in the data repository PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.929968>) (Strydom et al., 2022)

6 Code availability

R scripts used to generate figures and maps can be found in the [supplementary information](#).

7 Team list

260 Simone Strydom, [Roisin McCallum](#), [Anna Lafratta](#), Chanelle L. Webster, Caitlyn M. O’Dea, Nicole E. Said, [Natasha Dunham](#) ~~Roisin McCallum~~, Karina Inostroza, Cristian Salinas, Samuel Billingham, ~~Anna~~

~~Lafatta~~, Charlie M. Phelps, Connor Campbell, Connor Gorham, ~~Natasha Dunham~~, Rachele Bernasconi, Anna M. Frouws, Axel Werner, Federico Vitelli, Viena Puigcorbé, Alexandra D’Cruz, Kathryn M. McMahon, Jack Robinson, Megan J. Huggett, Sian McNamara, Glenn A. Hyndes and Oscar Serrano.

265 **8 Author contribution**

OS conceived the idea. SS and OS lead the project, curated data and wrote the paper. SS, JR, SM, RB, MJH and OS conducted literature searches. SS, RM, KI, CW, NS, CS, SB, AL, CO, CC, CG, CMP, ND, AW, AF, RB, SM, FV, VP, AD, KM, JR, MJH, GH and OS contributed to the manuscript and/or extracted data from papers. SS, CW, RM, CMP and OS wrote scripts and created figures and tables. All

270 authors reviewed the manuscript.

9 Competing interests

The authors declare that they have no conflicting interest.

10 Disclaimer

275 **11 Acknowledgements**

A.L. was supported by Environmental Protection Authority (SA)/Water SA and Edith Cowan University (ECU) Industry Engagement Scholarship. C.S. was funded by ECU Higher Degree by Research Scholarship. OS was supported by I+D+i projects RYC2019-027073-I and PIE HOLOCENO 20213AT014 funded by MCIN/AEI/10.13039/501100011033 and FEDER. We would also like to
280 acknowledge the help of Madison Williams-Hoffman, Casper Avenant and Paul Lavery for their respective input into earlier versions of this manuscript, and the Centre for Marine Ecosystems
~~R~~research.

12 Reference List

- 285 [Ascioti, F. A., Mangano, M. C., Marcianò, C., & Sarà, G. \(2022\). The sanitation service of seagrasses—Dependencies and implications for the estimation of avoided costs. *Ecosystem Services*, 54, 10.1016/j.ecoser.2022.101418.](#)
- Cullen-Unsworth, L. C., Nordlund, L. M., Paddock, J., Baker, S., McKenzie, L. J. and Unsworth, R. K. F.: Seagrass meadows globally as a coupled social-ecological system: Implications for human wellbeing, *Mar. Pollut. Bull.*, 83(2), 387–397, doi:10.1016/j.marpolbul.2013.06.001, 2014.
- 290 Duarte, C. M. and Chiscano, C. L.: Seagrass biomass and production: A reassessment, *Aquat. Bot.*, 65(1–4), 159–174, doi:10.1016/S0304-3770(99)00038-8, 1999.
- Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I. and Marbà, N.: The role of coastal plant communities for climate change mitigation and adaptation, *Nat. Clim. Chang.*, 3(11), 961–968, doi:10.1038/nclimate1970, 2013.
- 295 Graul, C.: leafletR: Interactive Web-Maps Based on the Leaflet JavaScript Library, [online] Available from: <http://cran.r-project.org/package=leafletR>., 2016.
- Hall, M. O., Durako, M. J., Fourqurean, J. W. and Zieman, J. C.: Decadal changes in seagrass distribution and abundance in Florida Bay, *Estuaries*, 22(2), 445–459, doi:10.2307/1353210, 1999.
- Hoegh-Guldberg, O. and Bruno, J. F.: The impact of climate change on the world’s marine ecosystems, *Science* (80-.), 328(5985), 1523–1528, doi:10.1126/science.1189930, 2010.
- 300 Kilminster, K., McMahon, K., Waycott, M., Kendrick, G. A., Scanes, P., McKenzie, L., O’Brien, K. R.,

Lyons, M., Ferguson, A., Maxwell, P., Glasby, T. and Udy, J.: Unravelling complexity in seagrass systems for management: Australia as a microcosm, *Sci. Total Environ.*, 534, 97–109, doi:10.1016/J.SCITOTENV.2015.04.061, 2015.

305 de los Santos, C. B., Krause-Jensen, D., Alcoverro, T., Marbà, N., Duarte, C. M., van Katwijk, M. M., Pérez, M., Romero, J., Sánchez-Lizaso, J. L., Roca, G., Jankowska, E., Pérez-Lloréns, J. L., Fournier, J., Montefalcone, M., Pergent, G., Ruiz, J. M., Cabaço, S., Cook, K., Wilkes, R. J., Moy, F. E., Trayter, G. M. R., Arañó, X. S., de Jong, D. J., Fernández-Torquemada, Y., Auby, I., Vergara, J. J. and Santos, R.: Recent trend reversal for declining European seagrass meadows, *Nat. Commun.*, 10(1), 1–8, 310 doi:10.1038/s41467-019-11340-4, 2019.

[Lamb, J. B., Van De Water, J. A., Bourne, D. G., Altier, C., Hein, M. Y., Fiorenza, E. A., ... & Harvell, C. D. \(2017\). Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. *Science*, 355\(6326\), 731-733, 10.1126/science.aal1956.](#)

McMahon, K., Collier, C. and Lavery, P. S.: Identifying robust bioindicators of light stress in seagrasses: A meta-analysis, *Ecol. Indic.*, 30, 7–15, doi:10.1016/j.ecolind.2013.01.030, 2013.

Nordlund, L. M., Koch, E. W., Barbier, E. B. and Creed, J. C.: Seagrass ecosystem services and their variability across genera and geographical regions, edited by K. O. Reinhart, *PLoS One*, 11(10), e0163091, doi:10.1371/journal.pone.0163091, 2016.

Short, F., Carruthers, T., Dennison, W. and Waycott, M.: Global seagrass distribution and diversity: A bioregional model, *J. Exp. Mar. Bio. Ecol.*, 350(1–2), 3–20, doi:10.1016/j.jembe.2007.06.012, 2007.

Short, F. T. and Coles, R. G.: *Global seagrass research methods*, Elsevier., 2001.

Short, F. T., Polidoro, B., Livingstone, S. R., Carpenter, K. E., Bandeira, S., Bujang, J. S., Calumpang, H. P., Carruthers, T. J. B., Coles, R. G., Dennison, W. C., Erftemeijer, P. L. A., Fortes, M. D., Freeman, A. S., Jagtap, T. G., Kamal, A. H. M., Kendrick, G. A., Judson Kenworthy, W., La Nafie, Y. A.,
325 Nasution, I. M., Orth, R. J., Prathep, A., Sanciangco, J. C., Tussenbroek, B. van, Vergara, S. G.,
Waycott, M. and Zieman, J. C.: Extinction risk assessment of the world's seagrass species, *Biol. Conserv.*, 144(7), 1961–1971, doi:10.1016/J.BIOCON.2011.04.010, 2011.

Strydom, S., Murray, K., Wilson, S., Huntley, B., Rule, M., Heithaus, M., Bessey, C., Kendrick, G. A.,
Burkholder, D., Fraser, M. W. and Zdunic, K.: Too hot to handle: Unprecedented seagrass death driven
330 by marine heatwave in a World Heritage Area, *Glob. Chang. Biol.*, 26(6), 3525–3538,
doi:10.1111/gcb.15065, 2020.


Strydom, Simone; Webster, Chanelle L; O'Dea, Caitlyn M; Said, Nicole E; McCallum, Roisin;
Inostroza, Karina; Salinas, Cristian; Billinghamurst, Samuel; Lafratta, Anna; Phelps, Charlie M; Campbell,
Connor; Gorham, Connor; Dunham, Natasha; Bernasconi, Rachele; Frouws, Ankje; Werner, Axel;
335 Vitelli, Federico; Puigcorbé, Viena; D'Cruz, Alexandra; McMahon, Kathryn M; Robinson, Jack;
Huggett, Megan J; McNamara, Sian; Hyndes, Glenn A; Serrano, Óscar (2021): Global database of key
seagrass structure, biomass, production and reproduction variables. PANGAEA,
<https://doi.org/10.1594/PANGAEA.929968>

UNEP-WCMC and FT, S.: Global Distribution of Seagrasses (version 6), [online] Available from:
340 <http://data.unep-wcmc.org/datasets/7>, 2018.

Unsworth, R. K. F., McKenzie, L. J., Collier, C. J., Cullen-Unsworth, L. C., Duarte, C. M., Eklöf, J. S.,
Jarvis, J. C., Jones, B. L. and Nordlund, L. M.: Global challenges for seagrass conservation, *Ambio*,
doi:10.1007/s13280-018-1115-y, 2018.

Waycott, M., Duarte, C. M., Carruthers, T. J. B., Orth, R. J., Dennison, W. C., Olyarnik, S., Calladine,

345 A., Fourqurean, J. W., Heck, K. L., Hughes, A. R., Kendrick, G. A., Kenworthy, W. J., Short, F. T. and Williams, S. L.: Accelerating loss of seagrasses across the globe threatens coastal ecosystems., Proc. Natl. Acad. Sci. U. S. A., 106(30), 12377–81, doi:10.1073/pnas.0905620106, 2009. Wickham, H.: ggplot2: Elegant Graphics for Data Analysis, 2016.



350

Figure 1. Global distribution of seagrass meadows (green) overlaid within six seagrass bioregions. Seagrass distribution data sourced from UNEP-WCMC & Short, (2018). Seagrass bioregions adapted from Short, Carruthers, Dennison, & Waycott (2007).




Figure 2. Number of publications that included seagrass data (coloured by type: biomass, structure ~~and~~ production ~~and reproduction~~) based on the year of data collection. Data from peer-reviewed publications that did not report the year of sampling, were not included in this figure.

355

360




Figure 3. Global distribution map of data on seagrass structure, biomass and, production and reproduction coloured by genera. The coloured points indicate the genera of seagrass studied and where many studies overlap, the colour appears darker than key.

365





Figure 4. Global distribution map of seagrass study sites labelled as dots. The colours indicate the data type (biomass, ~~reproduction~~, production and structure), while the size of each dot illustrates the number of data points for each site.

370



375 **Figure 5.** Box whisker plots depicting seagrass total biomass (including above-ground and below-
 ground biomass; g DW m⁻²), shoot density (number of shoots m⁻²), total production (g DW m⁻² day⁻¹)
 and percent cover (%) and flowering intensity (number of flowers or inflorescence shoots m⁻²) values
 within each bioregion. The boxplots show the median value (black line within box), 75% and 25%
 percentiles create the top and bottom of the box and the tails are the maximum and minimum
 380 contributions within 1.5 interquartile range. Count of data (N) per bioregion is shown at the top of each
 whisker.




Figure 6. Box whisker plots depicting total biomass (g DW m⁻²), shoot density (number of shoots m⁻²), total net primary production (g DW m⁻² day⁻¹) and percent cover (%) flowering intensity (number of flowers or inflorescence shoots m⁻²)-values per genera. The boxplots show the median value (black line within box), 75% and 25% percentiles create the top and bottom of the box and the tails are the maximum and minimum contributions within 1.5 interquartile range. Count of data (N) per bioregion is shown at the top of each whisker.

395 **Table 1.** Summary table outlining the count of data for each of the 15 seagrass variables based on bioregion, geotype and genera categorical variables. Above-ground biomass (AG), below-ground biomass (BG), total biomass (TB), shoot density (ShD), leaf density (LD), percent cover (Cov), above-ground production (AGP), below-ground production (BGP), total production (TP), shoot production (SP), and leaf production (LP), ~~flowering intensity (FI), fruit density (FD), sediment seed bank density (SB), seedling density (SD).~~

Commented [SS1]: All values updated

Bioregion	Geotype	Genera	AG	BG	TB	Sh D	LD	Co v	AG P	BG P	TP	SP	LP	
Mediterranean	Coastal	<i>Cymodocea</i>	137	97	113	166	4	12	12	4	15		99	
		<i>Halophila</i>	16	16	16	24	16	12						
		<i>Posidonia</i>	37	17	21	136	27	5	7	21	17	12	46	
		<i>Zostera</i>	87	20	33	145	9	24			3		27	
	Estuarine	<i>Cymodocea</i>	24	24	24	24				12				12
		<i>Posidonia</i>	6											
		<i>Ruppia</i>	10	1	3									
		<i>Zostera</i>	80	63	60	134	35							
Temperate North Atlantic	Coastal	<i>Halodule</i>						3						
		<i>Thalassia</i>	6	6	6	8								
		<i>Zostera</i>	91	51	63	83	42	6	24	23			29	
	Estuarine	<i>Cymodocea</i>	5	5	5	5								
		<i>Ruppia</i>	26	26	26		7							
		<i>Zostera</i>	315	222	216	284	125	12	14	6	6	58	33	
Temperate North Pacific	Coastal	<i>Halophila</i>	23	23	42	25			13	13	13			
		<i>Phyllospadix</i>	12	12	12	12	18		12	12	12	12	12	
	Estuarine	<i>Zostera</i>	216	173	236	299	17	27		1	1		0	
		<i>Ruppia</i>	6			5								
		<i>Zostera</i>	171	132	191	166		15	11				76	
Temperate Southern	Coastal	<i>Amphibolis</i>	14		1	18		1	1					

		<i>Cymodocea</i>	2	2	2	2					2	
		<i>Halodule</i>	2	2	2	2					2	
		<i>Halophila</i>				3						
		<i>Posidonia</i>	125			59	24	8	9			1 11
		<i>Zostera</i>	2	2	2	2	19				2	
	Estuarine	<i>Halophila</i>	127	127	127		1		7	7	19	1
		<i>Posidonia</i>				40						37
		<i>Ruppia</i>	8	8	19	5						
		<i>Zostera</i>	186	93	136	137	322		55			2 2
Tropical Atlantic	Coastal	<i>Halodule</i>	90	90	86	45	16		18			1
		<i>Halophila</i>	12	9	47	21	1				3	
		<i>Ruppia</i>	42	42	42		19					
		<i>Syringodium</i>	14	3	22	12	2					1
		<i>Thalassia</i>	171	100	169	346	22		56	20	82	9 30
		<i>Zostera</i>	3	3	3		6					24 24
	Estuarine	<i>Halodule</i>	136	137	131	108	26					
		<i>Halophila</i>			3		2					
		<i>Ruppia</i>	58	52	57	11		14				
		<i>Syringodium</i>	11	13	8	5	25					5
		<i>Thalassia</i>	60	35	47	33	30				1	33
		<i>Zostera</i>	43	13	40	78						25
Tropical Indo Pacific	Coastal	<i>Amphibolis</i>	16		5	10	4	5	5		6	
		<i>Cymodocea</i>	163	44	100	141	101		2	10	25	2 34
		<i>Enhalus</i>	119	61	66	78	28		1		2	18
		<i>Halodule</i>	95	46	129	65	55		1	1	18	1 23
		<i>Halophila</i>	139	74	88	96	100	1	3	1		
		<i>Posidonia</i>	6			13	4				6	
		<i>Syringodium</i>	52	17	43	55	29			6	6	7
		<i>Thalassia</i>	285	87	117	152	78		1		7	1 63
		<i>Thalassodendron</i>	29	13	14	23	6		1		4	11
		<i>Zostera</i>	68	10	13	6	20	4				1
	Estuarine	<i>Cymodocea</i>	5	2	2	14					2	
		<i>Enhalus</i>	3		11	3	2					

<i>Halodule</i>	17	17	27	11	11							2		
<i>Halophila</i>	6	6	21	21	18									
<i>Syringodium</i>	1			1										
<i>Thalassia</i>	2	2	4	10	2									
<i>Thalassodendron</i>	2	3	2	2									3	2
<i>Zostera</i>	7	7	42	68	9							2		
Total # data	3389	2008	2695	3212	1285	146	265	125	258	151	737			

400