

1 **Long-Term Monthly 0.05° Terrestrial Evapotranspiration Dataset (1982–2018)**  
2 **for the Tibetan Plateau**

3  
4 Ling Yuan <sup>1,2</sup>, Xuelong Chen<sup>1,4,6\*</sup>, Yaoming Ma<sup>1,2,3,4,5,6\*</sup>, Cunbo Han<sup>1,4,6</sup>, Binbin Wang<sup>1,4,5,6</sup>, Weiqiang Ma<sup>1,4,6</sup>

5  
6 <sup>1</sup> State Key Laboratory of Tibetan Plateau Earth System, Environment and Resources (TPESER), Institute of  
7 Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China.

8 <sup>2</sup> College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

9 <sup>3</sup> College of Atmospheric Science, Lanzhou University, Lanzhou 730000, China

10 <sup>4</sup> National Observation and Research Station for Qomolangma Special Atmospheric Processes and  
11 Environmental Changes, Dingri 858200, China

12 <sup>5</sup> Kathmandu Center of Research and Education, Chinese Academy of Sciences, Beijing 100101, China

13 <sup>6</sup> China-Pakistan Joint Research Center on Earth Sciences, Chinese Academy of Sciences, Islamabad 45320,  
14 Pakistan

15  
16 **Corresponding author and address:**

17 Xuelong Chen, Dr., Prof., x.chen@itpcas.ac.cn

18 Yaoming Ma, Dr., Prof., ymma@itpcas.ac.cn

19 Building 3, No.16 Lincui Road, Chaoyang District, Beijing 100101, China

30 **Abstract**

31 Evapotranspiration (ET) plays a crucial role in the water balance of the Tibetan Plateau (TP), often referred  
32 to as the "Asian water tower" region. However, accurately monitoring and comprehending the spatial and  
33 temporal variations of ET components (including soil evaporation  $E_s$ , canopy transpiration  $E_c$ , and intercepted  
34 water evaporation  $E_w$ ) in this remote area remains a significant challenge due to the limited availability of  
35 observational data. This study generates a 37-year dataset (1982–2018) of monthly ET components for the TP  
36 using the MOD16-STM (MOD16 soil texture model). This model utilizes up-to-date soil properties,  
37 meteorological data, and remote sensing datasets. The estimated ET results strongly correlate with  
38 measurements from nine flux towers, demonstrating a low root mean square error of 13.48 mm/month, a mean  
39 bias of 2.85 mm/month, a coefficient of determination of 0.83, and an index of agreement of 0.92. The annual  
40 average ET for the entire TP, defined as elevations higher than 2500 meters, is approximately  $0.93 \pm 0.037$   
41 Gt/year. The predominant contributor to ET on the TP is  $E_s$ , accounting for 84% of the total ET. Our findings  
42 reveal a noteworthy upward trend in ET in most central and eastern parts of the TP, with a rate of approximately  
43 1–4 mm/year ( $p < 0.05$ ) and a significant downward trend with rates between  $-3$  and 1 mm/year in the  
44 northwestern part of TP during the period from 1982 to 2018. The average annual increase in ET for the entire  
45 TP over the past 37 years is approximately 0.96 mm/year. This upward trend can be attributed to the TP's  
46 warming and wetting climate conditions. The MOD16-STM ET dataset demonstrates a reliable performance  
47 across the TP compared to previous research outcomes. This dataset is valuable for research on water resource  
48 management, drought monitoring, and ecological studies. The entire dataset is freely accessible through the  
49 Science Data Bank (<http://doi.org/10.11922/sciencedb.00020>, Y. Ma\*, X. Chen\*, L. Yuan, 2021) and the  
50 National Tibetan Plateau Data Center (TPDC) ([https://data.tpdc.ac.cn/en/disallow/e253621a-6334-4ad1-b2b9-](https://data.tpdc.ac.cn/en/disallow/e253621a-6334-4ad1-b2b9-e1ce2aa9688f/)  
51 [e1ce2aa9688f/](http://doi.org/10.11888/Terre.tpdc.271913), <http://doi.org/10.11888/Terre.tpdc.271913>, L. Yuan, X. Chen\*, Y. Ma\*, 2021).

52

53 **Keywords:** Evapotranspiration; MOD16-STM; Climate change; Asian water tower; Tibetan Plateau

54

55

56

57

58

59

## 60 1. Introduction

61 The Tibetan Plateau (TP) (24–40°N, 70–105°E) is often referred to as the "Asian water tower" owing to  
62 its distinctive geographical and ecological characteristics, as acknowledged in studies by Immerzeel et al. (2010,  
63 2020), Yao et al. (2012), and Xu et al. (2019). Within this region, evapotranspiration (ET) plays a vital role in  
64 the overall water balance. The TP predominantly features grassland (covering more than 47% of the area) and  
65 sparse vegetation or bare soil (surrounding over 33%), as indicated by the Moderate Resolution Imaging  
66 Spectroradiometer (MODIS) land cover dataset (MCD12C1) (Fig. 1c). Arid or semi-arid conditions mostly  
67 characterize this vast expanse. The TP is currently undergoing significant changes in its hydrological cycle,  
68 driven by global warming, as documented in studies by Yang et al. (2014), Kuang et al. (2016), and Zohaib et  
69 al. (2017). Nevertheless, accurately monitoring the spatial and temporal fluctuations in ET on the TP remains a  
70 formidable challenge due to the intricate environmental conditions of the TP. Moreover, understanding how ET  
71 patterns on the TP will evolve in the context of global warming is essential for assessing the impacts of these  
72 changes on the local population's livelihoods.

73 In recent years, various datasets for estimating ET on the TP have been developed, including the  
74 complementary relationship (CR) model (Ma et al., 2019; Wang et al., 2020), the surface energy balance system  
75 (SEBS) model (Chen et al., 2014, 2021; Zhong et al., 2019; Han et al., 2017, 2021), and the Penman–Monteith  
76 model with remote sensing (RS-PM) (Wang et al., 2018; Song et al., 2017; Chang et al., 2019; Ma et al., 2022),  
77 among others. However, a considerable variance exists among these TP ET products (Peng et al., 2016; Baik et  
78 al., 2018; Li et al., 2018; Khan et al., 2018). Studies have utilized eddy-covariance measurements (Shi et al.,  
79 2014; You et al., 2017; Yang et al., 2019; Ma et al., 2020) and reanalysis datasets (Shi et al., 2014; Dan et al.,  
80 2017; Yang et al., 2019; De Kok et al., 2020) to investigate ET on the TP. A recent Han et al. (2021) study  
81 produced the region's 18-year ET dataset (2001–2018). Enhancements to the canopy conduction algorithm in  
82 the Penman–Monteith model have led to improved ET estimates in previous research (Leuning et al., 2008;  
83 Zhang et al., 2010; Li et al., 2015; Zhang et al., 2016, 2019; Gan et al., 2018). However, these ET products tend  
84 to perform poorly in TP areas with sparse vegetation and arid to semi-arid climates (Zhang et al., 2010; Li et al.,  
85 2014b; Song et al., 2017; Baik et al., 2018; Li et al., 2018; Khan et al., 2018).

86 The limitations of the MOD16 Penman–Monteith model in arid to semi-arid TP regions are primarily due  
87 to its failure to consider the dominant role of topsoil texture and topsoil moisture in governing  $E_s$  processes  
88 (Yuan et al., 2021). Accurately separating and validating ET components on the TP remains challenging, even

89 though total ET estimates tend to align across different products (Lawrence et al., 2007; Blyth and Harding,  
90 2011; Miralles et al., 2016). The TP is primarily characterized by short and sparse vegetation, and soil moisture  
91 is crucial in ET estimation for this region. Several studies have used the Penman–Monteith algorithm to estimate  
92 ET on the TP (Wang et al., 2018; Ma et al., 2022). However, these studies have not accounted for the effects of  
93 soil moisture (*SM*) on evaporation resistance and stomatal conductance.

94 The enhanced Penman–Monteith model, MOD16-STM (MOD16 soil texture model), has been developed  
95 to address these limitations. MOD16-STM redefines the modules for  $E_s$  to consider the impacts of *SM* on soil  
96 evaporation resistance. This modification is based on eddy-covariance (EC) observations conducted on the TP  
97 (Yuan et al., 2021), offering a promising opportunity to estimate ET components in this region accurately.  $E_s$   
98 often dominates ET in sparsely vegetated areas, especially in arid and semi-arid regions with large bare soil  
99 areas (Wilcox et al., 2003; Kool et al., 2014; Wang et al., 2018; Ma et al., 2015; Ma and Zhang, 2022). Previous  
100 studies have highlighted that 20% to 40% of global ET is attributed to  $E_s$ . Bare soil surface evaporation is a  
101 rapid process influenced by shallow surface water (Koster and Suarez, 1996).  $E_s$  is primarily controlled by water  
102 diffusion in the soil (Good et al., 2015; Yuan et al., 2022). Accurately quantifying and separating  $E_s$  is crucial  
103 for enhancing our understanding of water and energy cycles on the TP. However, quantifying ET and its  
104 components remains challenging due to the influence of atmospheric demand, soil moisture conditions, and  
105 complex interactions between heterogeneous vegetation and soil properties (Merlin et al., 2016; Wu et al., 2017;  
106 Philips et al., 2017; Lehmann et al., 2018). MOD16-STM holds the potential to generate a remote sensing  $E_s$   
107 and ET component dataset covering the satellite era since 1980. In this study, the MOD16-STM model,  
108 acknowledging its limitations, was employed to estimate a long-term ET and ET components dataset spanning  
109 37 years (1982–2018) (Yuan et al., 2021).


110 A preferable approach involves directly estimating ET based on topsoil moisture, significantly impacting  
111 the TP's surface water exchange. Thus, leveraging the advantages of the MOD16-STM model for ET estimation  
112 on the TP, this study aimed to achieve two main objectives: (1) develop a 37-year (1982–2018) monthly ET  
113 dataset for the TP at a  $0.05^\circ \times 0.05^\circ$  spatial resolution; (2) quantify the spatial distribution and spatiotemporal  
114 variability of ET and its components across the TP.

## 115 **2. Materials and methods**

### 116 **2.1 Study area**

117 The Tibetan Plateau, located between  $25\text{--}40^\circ\text{N}$  and  $74\text{--}104^\circ\text{E}$ , spans approximately 2.5 million  $\text{km}^2$  and

118 consists of land above 2,500 m in altitude (Fig. 1a). This region, as indicated by the FAO drought index dataset,  
 119 represents the largest landform unit in Eurasia and encompasses hyper-arid, arid, semi-arid, and sub-humid  
 120 climate zones (Fig. 1b). The land cover types primarily include mixed forests, grasslands, bare soil, glaciers,  
 121 and snow-covered areas (see Fig. 1c). The topsoil predominantly consists of sandy loam, loam, and clay (Fig.  
 122 1d). The annual average temperature in the region ranges from approximately  $-3.1^{\circ}\text{C}$  to  $4.4^{\circ}\text{C}$ . Average annual  
 123 precipitation gradually increases from less than 50 mm in the northwest to over 1000 mm in the southeast, with  
 124 the most precipitation occurring during summer (Ding et al., 2017). Over time, the TP has undergone significant  
 125 environmental changes, including increased precipitation, decreased wind speed (*wind*), fewer snow days,  
 126 reduced radiation, thawing permafrost, glacier melting, and increased vegetation (Kang et al., 2010; Yao et al.,  
 127 2012; Yang et al., 2014; Kuang et al., 2016; Bibi et al., 2018; Chen et al., 2019).




128  
 129 Figure 1. Maps of the (a) topography (STRM), (b) climate zones (FAO aridity index), (c) land cover types (MCD12C1),  
 130 and (d) soil textures (HWSD) in the study area. The red dots indicate the flux site locations.

131 **2.2 Generation of a long-term series of monthly ET products**

132 This study introduces a novel dataset comprising a long-term series of monthly ET generated using the

133 MOD16-STM model. The process of calculating monthly ET with the MOD16-STM model and the associated  
 134 driving datasets is illustrated in Figure 2.



135

136

Figure 2. Workflow of the MOD16-STM evapotranspiration product.

137 **2.2.1 Description of MOD16-STM ET model**

138 The MOD16-STM model computes the ET components using the Penman–Monteith equation as follows:

$$E_c = \frac{(\Delta \times f_c \times (R_n - G_0) + \rho_a \times C_p \times \frac{VPD}{r_a} \times f_c) \times (1 - F_{wet})}{\lambda \times \left( \Delta + \gamma \times \left( 1 + \frac{r_c}{r_a} \right) \right)} \quad (1)$$

$$E_s = \frac{(\Delta \times (1 - f_c) \times (R_n - G_0) + \rho_a \times C_p \times \frac{VPD}{r_a}) \times (1 - F_{wet})}{\lambda \times \left( \Delta + \gamma \times \left( 1 + \frac{r_s}{r_a} \right) \right)} \times \left( \frac{RH}{100} \right)^{\frac{VPD}{\beta}} \quad (2)$$

$$E_w = E_{wet\_s} + E_{wet\_c} \quad (3)$$

139 The total ET combines three distinct components:  $E_c$ ,  $E_s$ , and  $E_w$  (wet surface evaporation). For a more detailed  
 140 explanation of the calculations for  $E_{wet\_s}$  (evaporation from wet soil) and  $E_{wet\_c}$  (evaporation from wet canopy),  
 141 you can refer to Yuan et al. 2021. Here,  $r_a$  (s/m) is the aerodynamic resistance,  $r_c$  (s/m) is the aerodynamic  
 142 resistance of water vapor of the canopy, and  $r_s$  (s/m) is the surface (or canopy) resistance. Yuan et al. (2021)

143 optimized MOD16  $r_a$  based on the Monin-Obukhov similarity theory (MOST) and calibrated the empirical  
 144 values of  $r_c$  for grassland underlying surfaces. They also pointed out that the topsoil moisture content directly  
 145 affects the value of  $r_s$ , indirectly influencing the  $E_s$  process. Therefore, this study extended this optimization  
 146 algorithm from the site scale to the regional scale. The variables used in the above equations are defined as  
 147 follows:

- 148 •  $R_n$  represents the net radiation flux (W/m<sup>2</sup>).
- 149 •  $G_0$  denotes the soil heat flux (W/m<sup>2</sup>).
- 150 •  $\rho_a$  is the density of the air (kg/m<sup>3</sup>).
- 151 •  $C_p$  stands for the specific heat capacity of the air (J/(kg·K)).
- 152 •  $VPD$  represents the vapor pressure deficit (hPa).
- 153 •  $\Delta$  represents the slope of the saturated vapor pressure curve (hPa/K).
- 154 •  $\gamma$  is the psychrometric constant (hPa/K), calculated as  $\gamma = C_p \cdot P_a \cdot M_a / (\lambda \cdot M_w)$ , where  $\lambda$  is the latent heat  
 155 of vaporization (J/kg), and  $M_a$  and  $M_w$  are the molecular masses of dry air and wet air, respectively.
- 156 •  $r_a$  signifies the aerodynamic resistance (s/m).
- 157 •  $r_s$  represents the surface (or canopy) resistance (s/m).
- 158 •  $F_{wet}$  is the relative surface wetness.
- 159 • The vegetation cover fraction ( $f_c$ ) is estimated using the NDVI (Normalized Difference Vegetation  
 160 Index).

$$f_c = \left( \frac{NDVI - NDVI_{\min}}{NDVI_{\max} + NDVI_{\min}} \right)^2 \quad (4)$$

161  $R_n$  and  $G_0$  are calculated as follows:

$$R_n = (1 - \alpha) \times SWD + LWD - \varepsilon \times \sigma \times LST^4 \quad (5)$$

$$G_0 = R_n \times (I_c + (1 - f_c) \times (I_s - I_c)) \quad (6)$$

162 Here,  $SWD$  is the downward shortwave radiation,  $\alpha$  is land surface albedo,  $LWD$  is the downward longwave  
 163 radiation,  $\sigma$  represents the Stefan-Boltzmann constant ( $5.67 \times 10^{-8}$  W/(m<sup>2</sup>·K<sup>4</sup>)),  $\varepsilon$  is emissivity, and  $LST$  means  
 164 land surface temperature.  $I_c$  (= 0.05) and  $I_s$  (= 0.315) are the ratios of ground heat flux and net radiation for  
 165 surfaces with full vegetation cover (Su et al., 2002) and bare soil (determined by NDVI < 0.25 in this study)  
 166 (Yuan et al., 2021), respectively. When the air temperature ( $T_a$ ) is below 5°C, photosynthesis and transpiration

167 processes are not active, and therefore,  $E_c$  is not considered in the calculations. When the land surface  
 168 temperature is below 0°C, the sublimation equation is derived by modifying the surface energy balance equation  
 169 using the Clausius–Clapeyron equation, accounting for the equilibrium of water vapor in both liquid and frozen  
 170 states. It's important to note that this study did not estimate the evaporation from water surfaces. Previous  
 171 research has extensively examined water surface evaporation from lakes on the Tibetan Plateau in detail (Wang  
 172 et al., 2020). Therefore, this study focuses on land ET estimation, excluding water surface evaporation.

173 Numerous prior studies have employed optimized conductance to estimate  $E_c$  (Jarvis et al., 1976; Irmak  
 174 and Mutiibwa, 2010; Zhang et al., 2010; Leuning et al., 2008; Li et al., 2013, 2015), as well as  $E_s$  (Sun et al.,  
 175 1982; Camillo and Gurney, 1986; Sellers et al., 1996; Sakaguchi and Zeng, 2009; Ortega-Farias et al., 2010;  
 176 Tang et al., 2013). This study computed the  $r_a$  using the MOST (Thom, 1975; Liu et al., 2007).

$$r_a = \frac{\ln\left(\frac{z_h - d_0}{z_{0h}} - \psi_h\right) \ln\left(\frac{z_m - d_0}{z_{0m}} - \psi_m\right)}{k^2 u} \quad (7)$$

177 Where  $k$  represents the von Karman's constant (0.41),  $z_h$  and  $z_m$  denote the measurement heights for  $T_a$  and *wind*,  
 178 and  $d_0$  represents the displacement height. The stability correction functions for momentum ( $\psi_m$ ) and heat  
 179 transfer ( $\psi_h$ ) can be computed using universal parts. These correction terms' mathematical expressions (Eq. 8–  
 180 12) are as follows (Högström, 1996; Paulson, 1970).

181 For stable conditions:

$$\psi_m = -5.3 \frac{(z_m - z_{0m})}{L} \quad (8)$$

$$\psi_h = -8.0 \frac{z_h - z_{0h}}{L} \quad (9)$$

182 For unstable conditions:

$$\psi_m = 2 \ln\left(\frac{1+x}{1+x_o}\right) + \ln\left(\frac{1+x^2}{1+x_o^2}\right) - 2 \tan^{-1} x + 2 \tan^{-1} x_o \quad (10)$$

$$\psi_h = 2 \ln\left(\frac{1+y}{1+y_o}\right) \quad (11)$$

183 For neutral conditions:

$$\psi_m = \psi_h = 0 \quad (12)$$

184 In Equations (8–11), the following variables and parameters are defined:  $x = (1 - z_m/L)^{0.25}$ ,  $x_o = (1 - z_{0m}/L)^{0.25}$ ,  $y =$   
 185  $(1 - 11.6 z_h/L)^{0.5}$ , and  $y_o = (1 - 11.6 z_{0h}/L)^{0.5}$  (Högström, 1996; Paulson, 1970). Here,  $L$  represents the Obukhov  
 186 length (m), calculated as  $L = T_a \cdot u_*^2 / (k \cdot g \cdot T_*)$ , where  $g = 9.8 \text{ m/s}^2$  and  $T_*$  is the fractional temperature (K) and



187  $u_*$  denotes the friction velocity (m/s).  $T_*$  is further defined as  $T_* = -(\theta_s - \theta_a) / ((\ln(z_h/z_{oh}) - \psi_h))$ , where  $\theta_s$  can be  
 188 approximated using the *LST*, and  $\theta_a = T_a + z_h \cdot g / C_p$ . The parameterization of  $u_*$  and  $L$  has been successfully applied  
 189 in previous studies on the TP (Chen et al., 2013).  $z_{oh}$  represents the roughness length for heat transfer (m). A  
 190 parameterization scheme for  $z_{oh}$  developed by Yang et al. (2008) has been widely utilized in remote sensing land  
 191 surface fluxes and land surface models (LSMs) across the TP (Biermann et al., 2014; Chen et al., 2013; Ma et  
 192 al., 2015). This scheme has also been employed in the current study for consistency.

$$z_{oh} = \frac{70\nu}{u_*} \exp(-7.2u_*^{0.5}|T_*|^{0.25}) \quad (13)$$

193 where  $\nu$  is the fluid kinematic viscosity,  $\nu = 1.328 \times 10^{-5} \cdot (P_0/P_a) \cdot (T_a/T_0)^{1.754}$ ,  $P_0 = 1013$  hPa and  $T_0 = 273.15$  K. The  
 194 roughness height for momentum transfer ( $z_{om}$ ) was determined based on canopy height ( $h_c$ ), following the  
 195 method outlined by Chen et al. (2013). The water saturation degree of surface soil ( $SM/\theta_{sat}$ ) is utilized to impose  
 196 soil classification and soil texture constraints on the  $r_s$  and  $E_s$  estimates (Yuan et al., 2021), as follows:

$$r_s = \exp\left(a + b \times \frac{SM}{\theta_{sat}}\right) \quad (14)$$

197 Here, the parameters  $a$  and  $b$  are empirical coefficients that vary based on different soil textures, as documented  
 198 in Table 1. The estimation of  $\theta_{sat}$ , which considers soil organic content (SOC) and gravel content, can be  
 199 obtained using the Soc-Vg scheme (Chen et al., 2012; Zhao et al., 2018), and its calculation is as follows:

$$\theta_{sat} = (1 - V_{SOC} - V_g) \times \theta_{sat,m} + V_{SOC} \times \theta_{sat,sc} \quad (15)$$

200 Where  $\theta_{sat,m}$  represents the porosity of the mineral soil and can be calculated as  $\theta_{sat,m} = 0.489 - 0.00126 \cdot \%sand$   
 201 (Cosby et al., 1984). Additionally,  $\theta_{sat,sc}$  is the porosity of the SOC. It is assumed to be  $0.9 \text{ m}^3 \cdot \text{m}^{-3}$  in this study,  
 202 as per the work of Farouki (1981) and Letts et al. (2000). The variables  $V_{soc}$  and  $V_g$  denote the volumetric  
 203 fractions of the SOC and gravel, respectively, and their calculation is as follows:

$$V_{SOC} = \frac{\rho_p \times (1 - \theta_{sat,m}) \times m_{SOC}}{\rho_{SOC} \times (1 - m_{SOC}) + \rho_p \times (1 - \theta_{sat,m}) \times m_{SOC} + (1 - \theta_{sat,m}) \times \frac{\rho_{SOC} \times m_g}{1 - m_g}} \quad (16)$$

$$V_g = \frac{\rho_{SOC} \times (1 - \theta_{sat,m}) \times m_g}{(1 - m_g) \times \left(\rho_{SOC} \times (1 - m_{SOC}) + \rho_p \times (1 - \theta_{sat,m}) \times m_{SOC} + (1 - \theta_{sat,m}) \times \frac{\rho_{SOC} \times m_g}{1 - m_g}\right)} \quad (17)$$

204 In these equations,  $\rho_p$  represents the mineral particle density and is set at  $2700 \text{ kg/m}^3$ , while  $\rho_{soc}$  is the bulk  
 205 density of organic matter, maintained at  $130 \text{ kg/m}^3$ . Also,  $m_{soc}$  and  $m_g$  denote the percentages of organic matter  
 206 and gravel within the topsoil layer.

207 There are many parameters in equations 7–17. Some parameters have already been assessed for their

208 importance in ET estimation by Yuan et al. (2021). There are too many studies on investigating the empirical  
 209 paramters in equation 7–13. We will not repeat these analysis again. The parameterization method of  $\theta_{sat}$  in the  
 210 estimation of  $r_s$  in this study is composed of various empirical parameters ( $\rho_p$ ,  $\rho_{soc}$ , and  $\theta_{sat, sc}$ ) for different soil  
 211 types. We have conducted a uncertainty analysis of the estimated  $\theta_{sat}$  and sensitivity of its uncertainty to the  
 212 changes of empirical parameters. The impact of the empirical parameters on the estimation of ET is illustrated  
 213 in Figure A1 (a–c). The results indicate that with a 20% uncertainty range in the estimated parameters  $\rho_p$ ,  $\rho_{soc}$ ,  
 214  $\theta_{sat, sc}$  for  $\theta_{sat}$ , the loss in estimating ET is only below 3%. Thus, the conclusion is drawn that the estimation of  
 215 ET is not sensitive to uncertainty in the three parameters. Figure A2 also shows the accuracy of the estimated  
 216  $\theta_{sat}$  by the method used in this paper. Additionally, a sensitivity analysis is conducted on the empirical  
 217 parameters  $a$  and  $b$  for calculating  $r_s$ . Keeping  $\theta_{sat}$  and SM constant,  $r_s$  exhibits exponential changes with  
 218 variations in  $a$  and  $b$ , leading to significant fluctuations in the estimation of ET. Within a 20% range of variation  
 219 in  $a$  and  $b$ , the maximum loss in ET exceeded 50% in Figure A1 (d–e). Therefore, it is essential to perform  
 220 significance tests on the fitting results of the empirical parameters  $a$  and  $b$ , as well as independent validation of  
 221 the final ET estimates. The performance of soil surface resistance  $r_s$  estimated by the MOD16-STM model at  
 222 the site scale is demonstrated in Fig.3. The observations at the ten stations show that the soil surface resistance  
 223 exponentially decreases with the increasing  $SM/\theta_{sat}$ . The MOD16-STM has caught this exponential law. It has  
 224 a coefficient of determination ( $R^2$ ) higher than 0.34, which may enable the model to estimate the TP ET  
 225 reasonably.


226 It demonstrates that the parameterized method of  $\theta_{sat}$  maintain a high level of consistency with the  
 227 observed values. The sensitivity test show that the factors that have a significant impact on  $r_s$  and ET are the  
 228 topsoil moisture and soil organic matter content. Figure A3 present the impact of soil organic matter content on  
 229  $\theta_{sat}$  and ET estimation at different soil types. Hereby, we have collected the most updated soil texture and soil  
 230 moisture data to estimate the soil evaporation resistance.

231 Table 1. Parameters  $a$  and  $b$  were based on different soil textures used to calculate surface soil resistances.

Texture	$r_s = \exp\left(a + b \times \frac{SM}{\theta_{sat}}\right)$	
	$a$	$b$
Sandy Loam	7.65	-7.3
Sand	5.89	-8.17
Loamy Sand	8.02	-17.37
Silt Loam	7.09	-3.79
Loam	6.82	-4.33

Clay Loam	10.17	-7.43
Sandy Clay Loam	9.46	-4.52
Clay	10.02	-6.68
Silty Clay	11.67	-7.25
Silty Clay Loam	8.93	-9.14

232



233

234 Figure 3. The scatter point relationship between soil surface resistance ( $r_s$ ) and  $SM/\theta_{sat}$  observed at QOMS (sandy loam),  
 235 NAMOR (sandy), NASDE (loamy sand), Arou (silt loam), Maqu (loam), IT-CAS (clay loam), US-IHO (sandy clay  
 236 loam), US-Arm (clay), CH-Oe2 (silty clay), and US-Ib2 (silty clay loam). The red curves show the equations used by the  
 237 MOD16-STM model. The site information is given in Table 3 and A1.

### 238 2.2.2 Input data for calculating the TP ET

239 The MOD16-STM model relies on various remote sensing datasets, reanalysis datasets, and meteorological  
 240 forcing datasets to estimate monthly ET across the TP. Specific datasets are carefully selected to minimize  
 241 spatial and temporal gaps in the final product (Table 2). Here's a breakdown of the critical datasets and their  
 242 sources:

- 243 • Monthly meteorological forcing data, including *wind*,  $T_a$ , air specific humidity ( $q$ ),  $P_a$ ,  $SWD$ , and  $LWD$ ,  
 244 were obtained from the China Meteorological Forcing Dataset (CMFD) with a  $0.1^\circ$  spatial resolution from  
 245 1982–2018. This data source was accessed from the National Tibetan Plateau Data Center (Yang et al.,  
 246 2010; He et al., 2020). CMFD can be downloaded from TPDC (<https://data.tpdc.ac.cn/>).
- 247 •  $LST$  and precipitation data were sourced from ERA5-Land, which provides data with a spatial  
 248 resolution of  $0.1^\circ$  and a monthly temporal resolution. These datasets were obtained from the European

249 Centre for Medium-Range Weather Forecasts (ECWMF).

250 • Albedo ( $\alpha$ ) data, with a spatial resolution of  $0.05^\circ$  and an 8-day temporal resolution, were derived from

251 the Global Land Surface Satellite (GLASS) dataset (Liang et al., 2021).

252 • A long-term NDVI dataset, with a spatial resolution of  $0.05^\circ$  and daily temporal resolution, was

253 acquired from the National Oceanic and Atmospheric Administration's National Centers for Environmental

254 Information (NOAA-NCEI) ([https://www.ncei.noaa.gov/products/climate-data-records/normalized-](https://www.ncei.noaa.gov/products/climate-data-records/normalized-difference-vegetation-index)

255 [difference-vegetation-index](https://www.ncei.noaa.gov/products/climate-data-records/normalized-difference-vegetation-index)). This dataset calculates the canopy height and Leaf Area Index (LAI) (Chen

256 et al., 2013).

257 • A topsoil moisture dataset for the 0–10 cm depth, with a spatial resolution of  $0.25^\circ$  and a monthly

258 temporal resolution, was obtained from the Global Land Evaporation Amsterdam Model (GLEAM)

259 (Miralles et al., 2011).

260 • Upward longwave radiation ( $LWU$ ) was derived from LST using the Stefan-Boltzmann Law. The

261 emissivity of mixed pixels was calculated based on the specific emissivity values for vegetated and bare

262 land surfaces, following Sobrino et al. (2004).

263 • Soil texture and soil property data were obtained from the Harmonized World Soil Database v1.2

264 (HWSD) (Wieder et al., 2014). These data were used to calculate soil evaporation resistance.

265 Daily and 8-day input data were averaged over the temporal scale to create monthly datasets to ensure

266 consistency. The average value was considered invalid if the ratio of valid data in any given month was below

267 90%. Additionally, the spatial resolutions of all input datasets were interpolated to a standard  $0.05^\circ$  spatial

268 resolution using a widely used bilinear interpolation method.

269 Table 2. Input datasets are used to calculate the ET on the Tibetan Plateau.

	Data source	Temporal resolution	Availability	Domain	Spatial resolution	Method
<i>SWD</i>	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
<i>LWD</i>	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
$T_a$	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
$q$	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
<i>wind</i>	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
$P_a$	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis
Precipitation	CMFD	3 h	1979–2018	China land	$0.1^\circ \times 0.1^\circ$	Reanalysis

<i>LST</i>	ERA5	Monthly	1981–2021	Global	0.1° × 0.1°	Reanalysis
$\alpha$	GLASS	8 days	1981–2019	Global	0.05° × 0.05°	Satellite
NDVI	AVHRR	Daily	1981–2019	Global	0.05° × 0.05°	Satellite
<i>SM</i>	GLEAM	Monthly	1979–2019	Global	0.25° × 0.25°	Reanalysis
Soil Properties	HWSD	/	/	China land	0.083°/1 km	/

## 270 2.3 Validation methods

### 271 2.3.1 Model validation at site scale

272 Limited stations on the TP make it impossible to collect ET observations at all kinds of soil textures. We  
273 have collected datasets from 17 flux sites outside the TP (Table A1 in Appendix A). Five sites are used to verify  
274 the relationship between soil surface resistance and  $SM/\theta_{sat}$ . This result is presented in Fig. 3. The other twelve  
275 verification sites include ten different soil textures (sandy loam, sand, loamy sand, silt loam, loam, clay loam,  
276 sandy clay loam, clay, silty clay, and silty clay loam) and three surface cover types (grassland, evergreen forest,  
277 and cropland) (Table A1). These twelve sites are used to do model validation at the site scale. When evaluating  
278 the MOD16-STM at the site scale, the meteorological forcing data comes from the station measurement. This  
279 helps us to minimize the simulation uncertainty due to the errors in the model forcing datasets. This methodology  
280 can allow us to diagnose the model's limitation in representing the evapotranspiration process. Figure A4 shows  
281 that MOD16-STM can capture the ET variations at the twelve sites. Table A2 also lists the statistical values of  
282 the daily ET estimation. Since these sites include all kinds of soil textures and different canopy covers, we  
283 believe the MOD16-STM model could be applied to the TP regional scale.

### 284 2.3.2 ET product evaluation

285 The remote sensing ET product is validated through comparison with flux tower observations on the TP.  
286 Table 3 lists details of nine flux stations on the TP used for the ET product evaluation. These stations belong to  
287 the China-Flux (Dang-Xiong site (DX), Hai-Bei site (HB), Yu et al., 2006; Zhang et al., 2019a), the Tibetan  
288 Observation and Research Platform (TORP) (BJ, NADORS, SETORS, QOMS, NAMORS, and Shuang-Hu  
289 (SH), Ma et al., 2020), and the Heihe Watershed Allied Telemetry Experimental Research (HiWATER) (Arou,  
290 Liu et al., 2011, 2018; Che et al., 2019) networks. The nine stations are in areas with three different land cover  
291 types: alpine meadow, alpine steppe, and Gobi. Half-hourly flux data measured by eddy-covariance from the  
292 nine stations are collected. It's important to note that the energy balance closure ratio (ECR) indicates whether

293 the sum of sensible heat ( $H$ ), latent heat ( $LE$ ), and soil heat flux ( $G_0$ ) matches the  $R_n$ . Half-hourly data are  
 294 screened and corrected accordingly to ensure the reliability of eddy-covariance measurements. Half-hourly  $LE$   
 295 data is corrected using the Bowen ratio energy balance correction method (Chen et al., 2014).

$$ECR = \frac{H + LE}{R_n - G_0} \quad (18)$$

$$LE_{cor} = \frac{1}{ECR} \times LE \quad (19)$$

296

297 Table 3. Details of the nine flux observation stations on the TP used for the ET product evaluation

Sites	Long., Lat.	Land cover type	Elevation (m)	Availability	Climate zone	Reference
Shuang-Hu (SH)	88.83°E, 33.21°N	Alpine meadow	4947	2013–2018	Semi-arid	Ma et al. (2015b)
BJ	91.90°E, 31.37°N	Alpine meadow	4509	2010–2016	Semi-arid	
NADORS	79.60°E, 33.38°N	Alpine steppe	4264	2010–2018	Arid	
SETORS	94.73°E, 29.77°N	Alpine meadow	3326	2007–2018	Sub-humid	Ma et al., 2020
QOMS	86.95°E, 28.35°N	Gobi	4276	2007–2018	Semi-arid	
NAMORS	90.99°E, 30.77°N	Alpine meadow	4730	2008–2018	Semi-arid	
Arou	100.46°E, 38.05°N	Alpine meadow	3033	2008–2017	Sub-humid	Liu et al., 2011, 2018; Che et al., 2019
Dang-Xiong (DX)	91.06°E, 30.49°N	Alpine meadow	2957	2004–2010	Semi-arid	Yu et al., 2006;
Hai-Bei (HB)	101.32°E, 37.61°N	Alpine meadow	3190	2002–2010	Sub-humid	Zhang et al., 2019a

298

299 In the validation process, the half-hourly  $LE_{cor}$  data obtained from all the flux sites are subjected to further  
 300 processing, including conversion to daily and monthly averages, while employing a stringent quality control  
 301 procedure. Daily values are null if derived from valid data points amounting to less than 80% in a single day.  
 302 Similarly, monthly average values are disregarded in the validation if they are derived from valid data points  
 303 accounting for less than 80% of observations for that month. This approach ensured the robustness of the  
 304 validation process.

## 305 2.4 Accuracy metrics

306 The accuracy of the modeled ET was assessed by comparing the pixel values ( $M_i$ ), corresponding to the  
 307 latitude and longitude of the flux site, with the flux tower measurements ( $G_i$ ). Several statistical metrics are

308 employed for validation, including the Coefficient of Determination ( $R^2$ ), a measure of the proportion of the  
 309 variance in the observed data ( $G_i$ ) that is explained by the modeled data ( $M_i$ ). A higher  $R^2$  value indicates a  
 310 stronger linear relationship between the two datasets. Mean Bias (MB) represents the average difference  
 311 between the modeled ET ( $M_i$ ) and the observed flux tower measurements ( $G_i$ ). Positive MB values suggest  
 312 overestimation by the model, while negative values indicate underestimation. Root Mean Square Error (RMSE)  
 313 measures the standard deviation of the differences between modeled and observed values ( $M_i - G_i$ ). A smaller  
 314 RMSE implies greater accuracy in the model's predictions. Index of Agreement (IOA) indicates the degree of  
 315 agreement between modeled and observed data, with a value of 1 indicating perfect agreement. Higher IOA  
 316 values indicate better agreement between the two datasets. The equations for these parameters are as follows:

$$R^2 = \frac{\left(\sum_{i=1}^n (M_i - \bar{M})(G_i - \bar{G})\right)^2}{\sum_{i=1}^n (M_i - \bar{M})^2 \sum_{i=1}^n (G_i - \bar{G})^2}, 0 \leq R^2 \leq 1 \quad (20)$$

$$MB = \frac{1}{N} \sum_{i=1}^n (M_i - G_i) \quad (21)$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (M_i - G_i)^2} \quad (22)$$

$$IOA = 1 - \frac{\sum_{i=1}^n (M_i - G_i)^2}{\sum_{i=1}^n (|M_i - \bar{G}| + |G_i - \bar{G}|)^2} \quad (23)$$

317 The subscript  $i$  denotes individual samples, and  $n$  is the total number of samples used in the assessment. The  
 318 significance of each parameter helps evaluate the model's performance in estimating ET.

319

### 320 **3. Results**


#### 321 **3.1 Evaluation of ET products against flux tower measurements**

322 The reliability of remote sensing-based ET estimates is often questioned without ground measurement  
 323 verification. This study compares the simulated monthly ET rates from the  $0.05^\circ$  grid where each EC site is  
 324 located with the flux tower observational data to validate the MOD16-STM ET results. The validation outcomes  
 325 for monthly MOD16-STM ET, using flux tower data, are illustrated in Fig. 4. The modeled ET exhibits excellent  
 326 performance and high consistency across the TP compared to ET observations.

327 Specifically, the grassland sites (SETORS, Arou, DX, and HB) display strong agreement, with  $R^2$  and IOA  
 328 values exceeding 0.82 and 0.95, respectively. The NAMORS site has the lowest performance, with the highest

329 RMSE (17.8 mm/month) and the lowest  $R^2$  and IOA (0.63 and 0.87, respectively). On average, the mean  $R^2$  and  
330 IOA values exceed 0.83 and 0.93, respectively. All  $R^2$  values pass the significance test at the  $p < 0.05$  level. The  
331 mean  $|MB|$  and RMSE values are less than 3 mm/month and 14 mm/month. It's important to note that positive  
332 MB values indicate an overestimation of ET, particularly during the dry season over barren land (QOMS, DX,  
333 SH, and NADORS) (Fig. 4). Conversely, underestimation occurs at higher ET rates in the summer, likely  
334 because the soil is close to saturation, leading to an overestimation of  $r_s$  and underestimation of  $E_s$  and ET.  
335 Generally, the time series of ET variations at the nine flux tower stations exhibit seasonal and annual periodic  
336 variations (Fig. 5). The site-scale validation results demonstrate that MOD16-STM ET provides accurate  
337 estimates in the TP region.






338

339 Figure 4. The validation of the MOD16-STM monthly ET at (a) SETORS, (b) Arou, (c) HB, (d) QOMS, (e) DX, (f) NAMORS, (g) BJ,

340

(h) SH, (i) NADORS, and (j) all sites.



341

342

343

344

Figure 5. Time series variations in the MOD16-STM simulated ET (blue solid line with "\*" marks) and flux-tower-observed ET (red circles) at (a) SETORS, (b) Arou, (c) HB, (d) QOMS, (e) DX, (f) NAMORS, (g) BJ, (h) SH, and (i) NADORS.

345

### 3.2 Spatial pattern of the multiyear averaged ET across TP

346

347

348


349

Figure 6 displays the spatial distribution of the average annual ET and its three components across the TP. ET exhibits a decreasing trend from southeast to northwest, with the highest values exceeding 1000 mm/year in the southeastern TP (the Heng-duan Mountains) and the lowest values of less than 100 mm/year in the Qaidam Basin and northwestern TP. This spatial pattern of annual ET closely mirrored that of the aridity index (Fig. 1b),

350 which is influenced by atmospheric demand and water supply. The sub-humid zone, covering approximately  
351 32.9% of the TP, contributes the highest proportion (43% of the TP's total ET) compared to other climate zones.  
352  $E_s$  dominates the central and western TP, with its spatial distribution closely resembling the overall ET. The  
353 spatial distributions of  $E_c$  and  $E_w$  are in line with the distribution of vegetation. High values of  $E_c$  (>200 mm/year)  
354 and  $E_w$  (>50 mm/year) are primarily concentrated in densely vegetated areas such as the Heng-duan Mountains  
355 in the southeastern TP.

356 The multiyear average ET for each season on the TP is depicted in Figure 6, covering spring (March, April,  
357 and May), summer (June, July, and August), autumn (September, October, and November), and winter  
358 (December, January, and February). The estimated ET reflects the general seasonal patterns quite accurately.  
359 During spring, the average ET is higher than in autumn, ranging from 20 to 250 mm/month in spring and from  
360 20 to 150 mm/month in autumn. This difference can be attributed to the increase in surface water generated by  
361 the thawing of permafrost and snow and ice melting as temperatures rise in spring, intensifying surface  
362 evaporation processes. Additionally, vegetation transpiration increases during the growing season. In summer,  
363 ET exceeds 200 mm/month over most TP, except for large areas in the northwestern TP where ET remains  
364 below 100 mm/month. Conversely, in winter, lower ET values are observed primarily in the densely vegetated  
365 southeastern region of the TP due to reduced water availability (precipitation) and lower  $T_a$  across the entire TP  
366 during this season.

367 Over the TP, the multiyear seasonal ET averages across the entire TP are as follows:  $90.79 \pm 3.16$  mm/year  
368 ( $0.23 \pm 0.0081$  Gt/year) in spring,  $152.05 \pm 8.44$  mm/year ( $0.38 \pm 0.021$  Gt/year) in summer,  $71.96 \pm 2.86$  mm/year  
369 ( $0.18 \pm 0.0074$  Gt/year) in autumn, and  $30.54 \pm 1.85$  mm/year ( $0.077 \pm 0.0047$  Gt/year) in winter. The multiyear  
370 average ET is  $346.5 \pm 13.2$  mm/year, representing both the mean and standard deviation, which characterizes  
371 interannual variability. This corresponds to approximately  $0.88 \pm 0.034$  Gt/year. Among its components,  $E_s$   
372 accounted for  $292.36 \pm 10.39$  mm/year ( $0.74 \pm 0.027$  Gt/year),  $E_c$  amounted to  $47.85 \pm 3.34$  mm/year ( $0.12 \pm 0.006$   
373 Gt/year), and  $E_w$  is  $7.07 \pm 2.89$  mm/year ( $0.02 \pm 0.001$  Gt/year). Notably,  $E_s$  constitutes the majority of ET on the  
374 TP, exceeding 84%. Wang et al. (2020) accurately estimated that the water evaporated from all plateau lakes is  
375  $0.0517$  Gt/year. Therefore, utilizing the area-weighted average method, the annual average water evaporation  
376 across the entire TP is approximately  $0.93 \pm 0.037$  Gt/year. Furthermore, the TP has an average annual rainfall  
377 of about  $1.8 \times 10^3$  Gt/year, estimated by Jiang et al. (2022). Approximately 53% of the TP's precipitation returns  
378 to the atmosphere through ET.



379


380

Figure 6. Spatial distributions of the multiyear (1982–2016) mean seasonal ET in (a) Spring, (b) Summer, (c) Autumn,

381

and (d) Winter across the TP.

382



383

384

Figure 7. Spatial pattern of the multiyear (1982–2018) mean annual (a) ET (evapotranspiration), (b)  $E_s$  (soil evaporation),

385


(c)  $E_c$  (canopy transpiration), and (d)  $E_w$  (intercepted water evaporation) across the TP.

386



387 **3.3 Temporal variations in ET across TP**


388 Quantifying variations in ET, both inter- and intra-annual, holds significant importance in understanding  
389 monsoon phenomena and studying climate change patterns on the TP. Figure 7 presents the spatial distribution  
390 of annual ET and its component trends from 1982 to 2018. These trends exhibit spatial heterogeneity across the  
391 TP. The annual ET has seen a significant increase, with rates ranging from 1 to 4 mm/year ( $p<0.05$ ), primarily  
392 in the central and eastern TP, encompassing more than 86% of the TP. Conversely, there has been a notable  
393 decrease in annual ET, with rates ranging from  $-3$  to  $-1$  mm/year in the northwestern TP. Similarly, the trends  
394 for  $E_s$  mirror those of ET, albeit with slightly lower magnitudes (1–3 mm/year,  $p<0.05$ ).  $E_c$  and  $E_w$  have shown  
395 slightly increasing trends of 0–2 mm/year ( $p<0.05$ ). When averaged across the entire TP, ET,  $E_s$ , and  $E_c$   
396 exhibited significant increases during the period from 1982 to 2018, with rates of 0.96 mm/year, 0.64 mm/year,  
397 and 0.44 mm/year, respectively ( $p<0.05$ ; see Fig. 8). Seasonally, positive, and significant trends are observed in  
398 all seasons for ET (Fig. 9), with the strongest trends occurring in summer (0.46 mm/year). Furthermore,  
399 multisource ET products indicate that most regions of the TP have exhibited consistent ET changes over the  
400 past 30 years (Yin et al., 2013; Peng et al., 2016; Wang et al., 2018; Ma et al., 2019; Wang et al., 2020; Li et al.,  
401 2021; Ma et al., 2022).



402  
403 Figure 8. Spatial patterns of the trends (1982–2018) of the annual (a) ET (evapotranspiration), (b)  $E_s$  (soil evaporation),  
404 (c)  $E_c$  (canopy transpiration), and (d)  $E_w$  (intercepted water evaporation) across the TP. The stippling on the maps

405

indicates the statistically significant trends ( $p < 0.05$ ).




406

407 Figure 9. Time series of the (a) anomalies in the annual ET and its three components, and (b) anomalies in seasonal mean  
408 ET. The least squares fitted linear trend are demonstrated by the dashed colored lines.

409 The rise in ET across the entire TP from 1982 to 2018 can be attributed to the concurrent warming and  
410 increased precipitation experienced in this region during the same period. Since the 1980s, the TP has undergone  
411 a general trend of greening, warming, and heightened precipitation, as illustrated in Figure 10. ET has  
412 consistently increased over the past four decades, but there was a notable shift in climate factors around 2000.  
413 From 1982 to 2000, ET showed a continuous increase, accompanied by a rapid decline in wind speed, while the  
414  $R_n$  remained relatively stable. However, between 2000 and 2018, there was a sharp decrease in  $R_n$  alongside an  
415 unchanged wind speed, but ET continued to rise during this period.

416 Consequently,  $R_n$  and wind speed are not the dominant factors driving annual variations in ET. The  
417 significant increases in  $T_a$ ,  $SM$ , and precipitation have coincided with the greening of the land surface over the  
418 last two decades. These factors collectively contribute to the observed increase in ET. In the most recent decade,  
419 the substantial growth in  $SM$  has emerged as the primary control factor for ET growth.



420

421 Figure 10. Time series of the annual anomalies in the (a) NDVI, (b)  $T_a$ , (c)  $R_n$ , (d)  $u$ , (e)  $SM$ , and (f) precipitation and  
 422 their least squares fitted linear trends during two periods of 1982-2000 and 2000-2018.

423 In summary, the increase in ET over the TP can be attributed to multiple factors. The rise in available  
 424 surface water plays a significant role throughout the study. Additionally, there is evidence of a general increase  
 425 in precipitation across the TP (Fig. 10). The combined impact of warming (shown by  $T_a$  in Fig. 10) and

426 vegetation greening (shown by *NDVI* in Fig. 10) further facilitate the opening of vegetation stomata, promoting  
427 increased vegetation transpiration. Warming the land surface and increased wind speeds enhance the efficiency  
428 of turbulent water exchange between the land and atmosphere. Furthermore, land surface warming accelerates  
429 the melting permafrost and glaciers on the TP. The surface wetting and the thickening of the active soil layer  
430 facilitate water transport from the lower soil layers to the upper layers. These environmental changes, such as  
431 water availability, precipitation patterns, vegetation dynamics, and temperature trends, all contribute to the  
432 increase in ET over the TP.

### 433 **3.4 Comparison of the MOD16-STM product to other ET products over the TP**

434 We have compared the accuracy of the MOD16-STM product and other available TP region datasets. It is  
435 shown in Fig. 11. The MOD16-STM ET model demonstrates high performance on the TP, with an average  $R^2$   
436 value of 0.87 and an average RMSE of 13.48 mm/month. Wang et al. (2018) evaluated a modified PML model  
437 for ET estimation on the TP, reporting  $R^2$  values exceeding 0.85 and RMSE values lower than 14 mm/month.  
438 The spatially averaged ET for 1982–2012 is 378.1 mm/year. Wang et al. (2020) assessed the performance of  
439 the generalized nonlinear complementary principle for ET estimation based on flux tower observations from the  
440 TP. Their results indicated an  $R^2$  of 0.93 and a RMSE of 0.40 mm/day. The spatially averaged ET during 1982–  
441 2014 is 398.3 mm/year. Han et al. (2021) used a combination of the effective aerodynamic roughness length  
442 and the surface energy balance model to estimate ET for the entire TP from 2001 to 2018 (Han-ET). They found  
443 good agreement between modeled and in-situ measured values ( $R^2 > 0.81$ ,  $RMSE < 14.5$  mm/month), and the  
444 average annual ET is approximately  $496 \pm 23$  mm, which is higher than the  $346.5 \pm 13.2$  mm obtained in this study  
445 (Fig. 12). The discrepancy can be attributed to differences in models and periods used in the two studies. Ma et  
446 al. (2022) also employed the PML\_V2 model to estimate ET on the TP (PML), yielding  $R^2$  and RMSE values  
447 ranging from 0.4 to 0.9 and 0.3 to 0.8 mm/day, respectively. The 35-year mean annual ET rates from PML-Ma  
448 resulted in an average value of  $353 \pm 24$  mm/year for the entire TP. Notably, the proportion of soil evaporation  
449 estimated by PML-Ma was approximately 64% of the total ET, which is lower than the estimated 84% in this  
450 study. The primary reason for this difference may be attributed to variations in land cover classification. The  
451 MODIS land cover classification largely categorizes the land surface in the northwestern TP as bare soil,  
452 increasing the proportion of soil evaporation.







Figure 11. Taylor diagram of the monthly-scale ET dataset validated with flux ET observations.

#### 4. Discussions

##### 4.1 Cross-Comparison of the Spatial Distribution of ET on the TP

A cross-comparison of the multi-year average values of various ET products is conducted to assess the differences and consistency in their spatial patterns. From the spatial distribution of annual average ET (Figure 12), all the ET products for the TP exhibit a decreasing trend from southeast to northwest, consistent with the transition in surface types from forests to grasslands and bare soil. In the Hengduan Mountains region, all products show high values (>600 mm/year) due to the dense vegetation and ample precipitation. However, significant absolute differences are observed among these 15 ET products. There are high differences among the products in the sparsely vegetated western and central TP regions. In the central TP region, where the Han-ET product exhibits the highest annual ET (>600 mm/year), while GLDAS-VIC has the lowest (approximately 35 to 50 mm/year). In the northwestern TP, EB-ET, GLDAS, MERRA-2, and GLEAM-v3.5a products display low values (<50 mm/year), while others range between 100 and 300 mm/year. In the extremely arid Qiangtang Plateau, all products show low values due to limited available surface water. ERA-Interim, ERA5-Land, PML-Zhang, CR-Ma, MOD16-STM, and GLEAM-v3.5b have relatively balanced distributions in the central and western TP regions (200–350 mm/year). There are high differences in the distribution of ET among the products in the downstream area of the Yarlung Tsangpo River. The spatial resolution of our product is 0.05°. This might be the reason for MOD16-STM has low ET for this topographic complex region. It is worth noting that MOD16 ET product has many missing values in the northwestern TP region, making it inadequate for a comprehensive



474

475 Figure 12. Spatial distribution of annual averaged ET on the TP during 2000 to 2014 derived from 15 products.

476 **4.2 ET components partitioning**

477 It is also necessary to have ET components comparative validation to enhance the practicality of the  
 478 data generated in this study. Unfortunately, there are no measured ET component data publicly available at the  
 479 moment. Comparative validation can be conducted based on existing research findings. Cui et al. (2020)

480 estimated the  $E_c/ET$  at the Nagqu Station (31.37°N, 91.90°E; 4509 m above sea level) in the central region of  
481 the TP using laser spectroscopy and chamber methods. During the observation period, the isotopic-based  $E_c/ET$   
482 ranged from 15% to 73%, with an average value of 43%. We calculated  $E_c/ET$  from our dataset at the same  
483 location and time period (June and July). The values of  $E_c/ET$  from MOD16-STM are in the range of 13.1% to  
484 62.6%. The average of  $E_c/ET$  is  $38.4\% \pm 4.7\%$ , which has a difference of 4.6% relative to isotopic estimation.  
485 Our  $E_c/ET$  estimation is close to the observation at Nagqu. Guo et al. (2017) also pointed that  $E_c$  constituted less  
486 than half of total ET (41% annually, 29% during monsoon) in Magazangbu catchment over the TP.

487 Moreover, we assess the similarities and differences between MOD16-STM and GLEAM-v3.5a ET  
488 components on the TP. Figure 13 shows that GLEAM's  $E_s$  values are generally smaller than our estimation  
489 throughout the TP region. The most recent results from Zheng et al. (2022) also suggest that the GLEAM product  
490 underestimates global  $E_s$  outputs. Conversely, GLEAM's  $E_c$  values are overestimated in the central and eastern  
491 TP. The differences in  $E_w$  are minimal because the values in that region are inherently small. Previous research  
492 has indicated that in the central TP region,  $E_s/ET$  accounts for over 60% (Cui et al., 2020), and the average  
493  $E_s/ET$  ratio across the entire region exceeds 65% (Wang et al., 2018). The reason for the relatively higher  $E_s$  in  
494 the central TP is that this region primarily consists of high-altitude grassland as the underlying surface. In the  
495 summer, the dominant processes are  $E_c$  and  $E_s$ , but in the winter  $E_s$  becomes the predominant process.  
496 Consequently, the proportion of  $E_s$  is higher over the entire year. GLEAM's results show that the  $E_c$  process  
497 predominates in the central TP, which differs somewhat from the findings of this study. Zheng et al. (2022) also  
498 indicate that the  $E_s$  process predominates in the central TP, exceeding 300 mm/year. Therefore, the ET  
499 components in this study, when compared with previous research, are more in line with the actual conditions in  
500 the TP.





Figure 13. Spatial comparison of ET components and their differences (MOD16-STM minus GLEAM-v3.5a).

### 4.3 Discrepancy in the estimation of annual ET over the TP

Figure 14 provides a comprehensive overview of the periods covered by various ET datasets and their annual ET estimations for the TP. Yao et al. (2013) estimated TP's ET (PT-Yao) in China using a satellite-driven modified Priestley–Taylor algorithm, constrained by NDVI and apparent thermal inertia derived from temperature changes. Their reported mean annual ET for the TP is approximately 320 mm/year. Song et al. (2017) estimated TP's ET (PM-Song) from 2000 to 2010 using the improved Penman–Monteith method and meteorological and satellite remote sensing data at a 1 km spatial resolution. They concluded that the average annual ET on the TP is 350.3 mm/year. Additionally, 18 mean annual ET values are estimated using existing ET products (PML-Zhang (Zhang et al., 2019b), EB-ET (Chen et al., 2019), CR-Ma (Ma et al., 2019), CMIP6-ssp126 (Eyring et al., 2016), GLDAS-Noah (Rodell et al., 2004), GLASS (Liang et al., 2021), GLEAM-v3.5b (Miralles et al., 2011, 2016), ERAR-Land (Muñoz-Sabater et al., 2021), MTE (Jung et al., 2010), PM-Li (Li et al., 2014a, 2014b), LPJ-Yin (Yin et al., 2013)) are included for comparison. Han-ET, ERA5-Land, and CMIP6 produce the highest values (>400 mm/year), while LPJ-Yin, GLASS, EB-ET, GLDAS, and GLEAM values are less than 300 mm/year. The results demonstrate substantial variability in the TP's estimated mean annual ET values. These differences are influenced by objective factors such as data accuracy, limitations of validation

518 method, and algorithm flaws. The ensemble mean of all datasets yields an annual ET of 348.6 mm/year, with  
 519 the MOD16-STM model's estimation (346.5 mm/year) being the closest to this ensemble mean. Overall, the  
 520 MOD16-STM ET model again demonstrates acceptable TP performance.



521  
 522 Figure 14. (a) The annual mean ET values of 18 datasets. The x-axis is the time coverage of the ET datasets, and the y-  
 523 axis is the multiyear mean value. (b) The bars denote the mean values and variations of the annual ET.

#### 524 4.4 Errors caused by objective factors

525 The MOD16-STM and other models rely on remote sensing and reanalysis data as primary input sources.  
 526 However, it's essential to acknowledge the inherent uncertainty in these datasets (Ramoelo et al., 2014). For  
 527 example, the topsoil water content from satellite data includes some errors (Liu et al. 2021). This indicates that  
 528 SM from GLEAM may introduce uncertainties to our ET estimation. Some studies have documented the  
 529 greening of the TP. Figure 10a demonstrated a significant decrease in NDVI after 2000, contrasting with the  
 530 NDVI changes reported by Wang et al. (2022). This inconsistency highlights the considerable uncertainty in the  
 531 NDVI data.

532 Additionally, *LST* plays a fundamental role in calculating the surface energy balance. Consequently, errors  
 533 in ERA5 *LST* can also bring uncertainty to the ET estimation. This study used a threshold value of NDVI (0.25)  
 534 to categorize land surfaces as bare soil or canopy-covered pixels. This threshold value may miss-classify bare

535 soil and grassland on the TP. The land cover mismatches between the reality and the land surface types in the  
536 MOD16-STM ET model can also introduce errors in the model simulation.

537 It is worth noting that flux towers used for validation typically cover areas ranging from a few hundred  
538 square meters to several square kilometers. These validation sites' representativeness depends on observation  
539 instrument height, turbulence intensity, topography, environment, and vegetation conditions. While site-scale  
540 evaluations of the MOD16-STM ET are conducted in this study, it's essential to recognize the uncertainties  
541 stemming from the limited number of validation sites. Future research should consider validation across various  
542 land cover types, climate zones, elevations, and seasons to provide a more comprehensive assessment of model  
543 performance.

544 While the MOD16 model directly estimates ET, bypassing the need for calculating sensible heat, it still  
545 relies on empirical coefficients, particularly those redefined for different soil textures. However, the remaining  
546 empirical parameters, such as  $C_L$  (the mean potential and stomatal conductance per unit leaf area), can introduce  
547 uncertainties into simulation results. Thus, future studies should prioritize parameterizing these empirical factors  
548 based on physical processes to reduce simulation uncertainties. It's crucial to consider the influence of physical  
549 processes related to deeper soil water and heat transfer on resistance. The MOD16-STM algorithm's accuracy  
550 is highly dependent on higher-precision soil moisture products. Since a substantial portion of the TP is covered  
551 by permafrost and seasonally frozen soil, assessing soil moisture conditions during freezing and thawing periods  
552 becomes challenging. Consequently, it is essential to employ observations during freeze-thaw periods to validate  
553 the model's applicability.

554 In summary, enhancing the model by incorporating physical parameterizations, especially for empirical  
555 coefficients, and accounting for the complexities of soil moisture variations in frozen regions will reduce  
556 uncertainties in *ET* simulation results in future research.

## 557 **5. Conclusion**

558 In this study, we have developed a 37-year (1982–2018) monthly ET dataset with a high spatial resolution  
559 (0.05°) for the TP using the newly developed MOD16-STM model. This dataset covers the entire study area  
560 with high spatial resolution and a long time series, making it a valuable resource for climate studies. Then, we  
561 investigated ET's spatial distribution and temporal trends across the TP. Key findings are summarized below:

- 562 • The ET product generated by the MOD16-STM model exhibits strong performance on the TP.  
563 Compared to flux tower observation data, the model achieves high  $R^2$  and IOA values of 0.83 and 0.93,



564 respectively, with an RMSE of 13.48 mm/month and a modest bias (MB) of 2.58 mm/month. This ET  
565 dataset holds potential applications in water resource management, drought monitoring, and ecological  
566 studies.

567 • The ET on the TP displays spatial heterogeneity and temporal variations driven by a combination of  
568 atmospheric demand and water supply. Generally, annual ET decreases from the southeastern to the  
569 northwestern regions of the TP.  $E_s$  accounts for over 84% of the annual ET, and the estimated multiyear  
570 mean annual ET on the TP for 1982–2018 is  $346.5 \pm 13.2$  mm. This corresponds to an annual water  
571 evaporation of about  $0.93 \pm 0.037$  Gt from the entire TP.

572 • Significant temporal trends are observed in the ET. Most parts of the central and eastern TP exhibit  
573 increasing trends of about 1 to 4 mm/year ( $p < 0.05$ ), whereas the northwestern TP shows a decreasing trend  
574 of  $-3$  to  $-1$  mm/year ( $p < 0.05$ ). Averaged across the entire TP, the ET increased significantly at a rate of  
575  $0.96$  mm/year ( $p < 0.05$ ) from 1982 to 2018. This increase in ET over the entire TP from 1982 to 2018 can  
576 be attributed to the warming and wetting of the climate during this period.

577 These findings contribute to a better understanding of the ET dynamics on the Tibetan Plateau and provide  
578 a valuable dataset for climate research and related applications.

#### 579 **Data availability**

580 The monthly ET dataset presented and analyzed in this article has been released. It is freely available at  
581 the Science Data Bank (<http://doi.org/10.11922/sciencedb.00020>, Y. Ma\*, X.Chen\*, L. Yuan, 2021) and the  
582 National Tibetan Plateau Data Center (TPDC) ([https://data.tpdc.ac.cn/en/disallow/e253621a-6334-4ad1-b2b9-  
583 e1ce2aa9688f/](https://data.tpdc.ac.cn/en/disallow/e253621a-6334-4ad1-b2b9-e1ce2aa9688f/), <http://doi.org/10.11888/Terre.tpdc.271913>, L. Yuan, X.Chen\*, Y. Ma\*, 2021). The dataset is  
584 published under the Creative Commons Attribution 4.0 International (CC BY 4.0) license.

#### 585 **Author contributions**

586 YMM, LY, and XLC led the writing of this paper and acknowledge responsibility for the experimental  
587 data and results. LY, XLC, and YMM drafted the document, and LY led the consolidation of the input and  
588 simulation dataset. XLC revised the manuscript. This paper is written in cooperation with all the co-authors.

#### 589 **Declaration of Competing Interest**

590 The authors declare that they have no known competing financial interests or personal relationships that  
591 could have appeared to influence the work reported in this paper.

592 **Acknowledgments**

593 We are grateful for the datasets provided by the China-Flux (<http://www.chinaflux.org/>), Ameri-Flux  
594 (<https://ameriflux.lbl.gov/>), GHG-Europe (<http://www.europe-fluxdata.eu/ghg-europe>), the National Tibetan  
595 Plateau Data Center (<https://data.tpdc.ac.cn/zh-hans/data>), the European Centre for Medium-Range Weather  
596 Forecasts (ECWMF) (<https://www.ecmwf.int/>), NOAA-NCEI ([https://www.ncei.noaa.gov/products/climate-  
598 data-records/normalized-difference-vegetation](https://www.ncei.noaa.gov/products/climate-<br/>597 data-records/normalized-difference-vegetation)), the Global Land Evaporation Amsterdam Model  
599 (<https://www.gleam.eu/>), and the National Earth System Science Data Sharing Infrastructure ([http://glass-  
product.bnu.edu.cn/](http://glass-<br/>600 product.bnu.edu.cn/)). The authors would like to thank all their colleagues at the observation stations on the TP  
600 for their maintenance of the instruments.

601 **Financial support**

602 This study is funded by the Second Tibetan Plateau Scientific Expedition and Research (STEP) Program  
603 (2019QZKK0103 and 2019QZKK0105) and the National Natural Science Foundation of China (42230610,  
604 91837208, 41975009).

605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620



## Appendix A: MOD16-STM model validation at flux site out of the Tibetan Plateau

Table A1. Basic information about the five test sites (which are used to test the relationship between soil surface resistance ( $r_s$ ) and  $SM/\theta_{sat}$  in the MOD16-STM model) and 12 verification sites (used for the MOD16-STM model evaluation at site scale). All the stations are located outside of the Tibetan Plateau.

	Site	Lat; lon	Land cover	$\theta$ (cm)	$f_{sand}$	$f_{clay}$	$m_{soc}$ (%)	$\theta_{sat}$	Soil Texture	Reference
Test Sites	IT-Cas	45.07; 8.71	CRO	5	0.28	0.29	2.6	/	Clay loam	<i>Denef et al. (2013)</i>
	US-IHO	36.47; 100.62	Bare	5	0.58	0.28	/	0.53	Sandy Clay Loam	<i>Lemone et al. (2007)</i>
	US-Arm	36.61; -97.49	CRO	5	0.28	0.43	1.5	/	Clay	<i>Fischer et al. (2007)</i>
	CH-Oe2	47.29; 7.73	CRO	5	0.095	0.43	2.8	/	Silty Clay	<i>Alaoui and Goetz (2008)</i>
	US-IB2	41.84; -88.24	GRA	0~15	0.106	0.29	2.4	/	Silty clay Loam	/
Verification sites	US-Dk1	35.97; -79.09	GRA	10	0.48	0.09	/	0.52	Loam	<i>Novick et al. (2004)</i>
	US-Fwf	35.45; -111.77	GRA	5	0.30	0.13	3.2	/	Silt Loam	<i>Dore et al. (2012)</i>
	US-Wkg	31.74; -109.94	GRA	5	0.67	0.17	1.0	/	Sandy Loam	<i>Ameri-Flux</i>
	CA-Obs	53.98; -105.11	ENF	5	0.72	0.05	4.3	/	Sandy Loam	<i>Ameri-Flux</i>
	CA-Ojp	53.91; -104.69	ENF	5	0.94	0.03	2.5	/	Sand	<i>Ameri-Flux</i>
	CA-Ca2	49.87; -125.29	ENF	5	0.74	0.03	3.0	/	Loamy Sand	<i>Ameri-Flux</i>
	CA-Ca3	49.53; -124.90	ENF	5	0.39	0.20	4.9	/	Loam	<i>Ameri-Flux</i>
	US-Dk3	35.97; -79.09	ENF	5	0.25	0.34	2.4	/	Silt Loam	<i>Ameri-Flux</i>
	US-Fuf	35.08; -111.76	ENF	5	0.31	0.35	3.9	/	Clay Loam	<i>Ameri-Flux</i>
	US-Ib1	41.86; -88.22	CRO	2.5	0.10	0.35	1.8	/	Silty clay Loam	<i>Denef et al. (2013)</i>
	ES-ES2	39.28; -0.32	CRO	5	0.11	0.47	3.7	/	Silty Clay	<i>Kutsch et al. (2010)</i>
	IT-Bci	40.52; 14.96	CRO	5	0.32	0.46	1.5	/	Clay	<i>Denef et al. (2013)</i>

636 Table A2. Assessment results of the daily ET (mm/day) simulated by the MOD16-STM model at the 12 verification sites.

637

The in-situ meteorological observation data drive this simulation.

	Sites	R <sup>2</sup> ( $p < 0.05$ )	IOA	MB	RMSE
Grassland	US-DK1	0.71	0.91	0.27	0.74
	US-Fwf	0.59	0.84	0.06	0.55
	US-Wkg	0.69	0.84	0.005	0.58
Evergreen Forest	CA-Obs	0.88	0.96	0.05	0.33
	CA-Ojp	0.79	0.93	0.11	0.38
	CA-Ca2	0.77	0.92	0.23	0.49
	CA-Ca3	0.79	0.94	0.02	0.44
	US-Dk3	0.79	0.92	0.51	0.87
	US-Fuf	0.58	0.81	0.33	0.66
	Cropland	US-Ib1	0.65	0.88	0.39
ES-ES2		0.87	0.91	0.04	0.94
IT-Bci		0.41	0.76	0.14	1.14
Mean	/	0.72	0.89	0.18	0.68

638

639

640

641


642

643

644

645

646



647

648 Figure A1. A sensitivity analysis on the impact of the uncertainty in the empirical parameters ( $\rho_p$ ,  $\rho_{soc}$ ,  $\theta_{sat, sc}$ ,  $a$  and  $b$ ) to

649

the estimation of ET (test in August, 2018).

650

651

652

653

654

655

656

657

658




Figure A2. Validation of the consistency between the estimated and the observed values for  $\theta_{sat}$  over the TP.

659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672  
 673




Figure A3. The sensitivity of LE and  $\theta_{sat}$  to the changes of  $m_{soc}$  content at different sites..

674

675

676

677

678


679

680

681

682

683



684

685

686

687

688

689

690

691

692

693

694

695

696

## References

697

698

699

700

- Alaoui, A., and Goetz, B.: Dye tracer and infiltration experiments to investigate macropore flow, *Geoderma*, 144(1–2), 279–286, <https://doi.org/10.1016/j.geoderma.2007.11.020>, 2008.
- Baik, J., Liaqat, U. W., and Choi, M.: Assessment of satellite- and reanalysis-based evapotranspiration products with two blending approaches over the complex landscapes and climates of Australia, *Agric. For. Meteorol.*,

701 263, 388–398. <https://doi.org/10.1016/j.agrformet.2018.09.007>, 2018.

702 Bibi, S., Wang, L., Li, X., Zhou, J., Chen, D., Yao, T.: Climatic and associated cryospheric, biospheric, and  
703 hydrological changes on the Tibetan Plateau: a review, *Int. J. Climatol.*, 38, 1–17,  
704 <https://doi.org/10.1002/joc.5411>, 2018.

705 Biermann, T., Babel, W., Ma, W., Chen, X., Thiem, E., Ma, Y., Foken, T.: Turbulent flux observations and  
706 modeling over a shallow lake and a wet grassland in the Nam Co basin, Tibetan Plateau, *Theor. Appl.*  
707 *Climatol.*, 116(1–2), 301–316, <https://doi.org/10.1007/s00704-013-0953-6>, 2014.

708 Blyth, E., and Harding, R. J.: Methods to separate observed global evapotranspiration into the interception,  
709 transpiration, and soil surface evaporation components, *Hydrol. Process.*, 25(26), 4063–4068,  
710 <https://doi.org/10.1002/hyp.8409>, 2011.

711 Camillo, P. J. and Gurney, R. J.: A resistance parameter for bare soil evaporation models, *Soil Sci.*, 141(2), 95–  
712 105, <https://doi.org/10.1097/00010694-198602000-00001>, 1986.

713 Chang, Y., Qin, D., Ding, Y., Zhao, Q., Zhang, S.: A modified MOD16 algorithm to estimate evapotranspiration  
714 over the alpine meadow on the Tibetan Plateau, China, *J. Hydrol.*, 561, 16–30,  
715 <https://doi.org/10.1016/j.jhydrol.2018.03.054>, 2018.

716 Chen, D., Xu, B., Yao, T., Guo, Z., Cui, P., Chen, F., Zhang, R., Zhang, X., Zhang, Y., Fan, J., Hou, Z., Zhang,  
717 T.: Assessment of past, present, and future environmental changes on the Tibetan Plateau, *Kexue*  
718 *Tongbao/Chinese Science Bulletin*, 60(32), 3025–3035, <https://doi.org/10.1360/N972014-01370>, 2015.

719 Chen, X., Massman, W. J., Su, Z.: A Column Canopy-Air Turbulent Diffusion Method for Different Canopy  
720 Structures, *J. Geophys. Res. Atmos.*, 124(2), 488–506, <https://doi.org/10.1029/2018JD028883>, 2019.

721 Chen, X., Su, Z., Ma, Y., Liu, S., Yu, Q., Xu, Z.: Development of a 10-year (2001–2010) 0.1° data set of land-  
722 surface energy balance for mainland China, *Atmos. Chem. Phys.*, 14(23), 13097–13117,  
723 <https://doi.org/10.5194/acp-14-13097-2014>, 2014.

724 Chen, X., Su, Z., Ma, Y., Yang, K., Wen, J., Zhang, Y.: An Improvement of Roughness Height Parameterization  
725 of the Surface Energy Balance System (SEBS) over the Tibetan Plateau, *J. Appl. Meteorol. Climatol.*,  
726 52(3), 607–622, <https://doi.org/10.1175/JAMC-D-12-056.1>, 2013.

727 Chen, X., Su, Z., Ma, Y., Trigo, I., Gentile, P.: Remote Sensing of Global Daily Evapotranspiration based on a  
728 Surface Energy Balance Method and Reanalysis Data, *J. Geophys. Res. Atmos.*, 126(16), e2020JD032873,  
729 <https://doi.org/10.1029/2020JD032873>, 2021.

730 Chen, Y., Xia, J., Liang, S., Feng, J., Fisher, J. B., Li, X., Li, X., Liu, S., Ma, Z., Miyata, A., Mu, Q., Sun, L.,

731 Tang, J., Wang, K., Wen, J., Xue, Y., Yu, G., Zha, T., Zhang, L., Zhang, Q., Zhao, T., Zhao, L., Yuan, W.:  
732 Comparison of satellite-based evapotranspiration models over terrestrial ecosystems in China, *Remote*  
733 *Sens. Environ.*, 140, 279–293, <https://doi.org/10.1016/j.rse.2013.08.045>, 2014.

734 Chen, Y., Yang, K., Tang, W., Qin, J., and Zhao, L.: Parameterizing soil organic carbon's impacts on soil  
735 porosity and thermal parameters for Eastern Tibet grasslands, *Sci. China Earth Sci.*, 55(6), 1001–1011,  
736 <https://doi.org/10.1007/s11430-012-4433-0>, 2012.

737 Che, T., Li, X., Liu, S., Li, H., Xu, Z., Tan, J., Zhang, Y., Ren, Z., Xiao, L., Deng, J., Jin, R., Ma, M., Wang, J.,  
738 and Yang, X.: Integrated hydrometeorological, snow and frozen-ground observations in the alpine region  
739 of the Heihe River Basin, China, *Earth Syst. Sci. Data.*, 11, 1483–1499, [https://doi.org/10.5194/essd-11-](https://doi.org/10.5194/essd-11-1483-2019)  
740 1483-2019, 2019.

741 Cleugh, H. A., Leuning, R., Mu, Q., Running, S. W.: Regional evaporation estimates from flux tower and  
742 MODIS satellite data, *Remote Sens. Environ.*, 106(3), 285–304, <https://doi.org/10.1016/j.rse.2006.07.007>,  
743 2007.

744 Cosby, B. J., Hornberger, G. M., Clapp, R. B., and Ginn, T. R.: A Statistical Exploration of the Relationships  
745 of Soil Moisture Characteristics to the Physical Properties of Soils, *Water Resour. Res.*, 20(6), 682–690,  
746 <https://doi.org/10.1029/WR020i006p00682>, 1984.

747 Cui, J., Tian, L., Wei, Z., Huntingford, C., Wang, P., Cai, Z., ... Wang, L.: Quantifying the Controls on  
748 Evapotranspiration Partitioning in the Highest Alpine Meadow Ecosystem. *Water Resour. Res.*, 56(4).  
749 <https://doi.org/10.1029/2019WR024815>, 2020.

750 Dan, J., Gao, Y., and Zhang, M.: testing and Attributing Evapotranspiration Deviations Using Dynamical  
751 Downscaling and Convection-Permitting Modeling over the Tibetan Plateau, *Water.*, 13(15), 2096:  
752 <https://doi.org/10.3390/w13152096>, 2017.

753 De Kok, R.J., Kraaijenbrink, P.D.A., Tuinenburg, O.A., Bonekamp, P.N.J., Immerzeel, W.W.: Towards  
754 understanding the pattern of glacier mass balances in High Mountain Asia using regional climatic modeling,  
755 *Cryosphere*, 14, 3215–3234, <https://doi.org/10.5194/tc-14-3215-2020>, 2020.

756 Deneff, K., Galdo, I. D., Venturi, A., and Cotrufo, M. F.: Assessment of Soil C and N Stocks and Fractions  
757 across 11 European Soils under Varying Land Uses, *Open J. Soil Sci.*, 03(07), 297–313,  
758 <https://doi.org/10.4236/ojss.2013.37035>, 2013.

759 Dore, S., Montes-Helu, M., Hart, S. C., Hungate, B. A., Koch, G. W., Moon, J. B., Finkral, A., Kolb, T. E.:  
760 Recovery of ponderosa pine ecosystem carbon and water fluxes from thinning and stand-replacing fire,



761 Glob. Chang. Biol., 18(10), 3171–3185, <https://doi.org/10.1111/j.1365-2486.2012.02775.x>, 2012.

762 Ding, J., Chen, L., Ji, C., Hugelius, G., Li, Y., Liu, L., Qin, S., Zhang, B., Yang, G., Li, F., Fang, K., Chen, Y.,  
763 Peng, Y., Zhao, X., He, H., Smith, P., Fang, J., Yang, Y.: Decadal soil carbon accumulation across Tibetan  
764 permafrost regions, *Nature Geoscience*, 10(6), 420–424, <https://doi.org/10.1038/ngeo2945>, 2017.

765 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., Taylor, K. E.: Overview of the  
766 Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci.*  
767 *Model Dev.*, 9, 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

768 Farouki, O.T.: The thermal properties of soils in cold regions, *Cold Reg. Sci., Technol.* 5, 67–75,  
769 [https://doi.org/10.1016/0165-232X\(81\)90041-0](https://doi.org/10.1016/0165-232X(81)90041-0), 1981.

770 Fischer, M. L., Billesbach, D. P., Berry, J. A., Riley, W. J., and Torn, M. S.: Spatiotemporal variations in  
771 growing season exchanges of CO<sub>2</sub>, H<sub>2</sub>O, and sensible heat in agricultural fields of the Southern Great  
772 Plains, *Earth Interact.*, 11(17), <https://doi.org/10.1175/EI231.1>, 2007.

773 Gan, R., Zhang, Y., Shi, H., Yang, Y., Eamus, D., Cheng, L., Chiew, F., Yu, Q.: Use of satellite leaf area index  
774 estimating evapotranspiration and gross assimilation for Australian ecosystems, *Ecohydrology*, 11(5),  
775 e1974, <https://doi.org/10.1002/eco.1974>, 2018.

776 Good, S. P., Noone, D., and Bowen, G.: Hydrologic connectivity constrains partitioning of global terrestrial  
777 water fluxes, *Science*, 349(6244), 175–177, <https://doi.org/10.1126/science.aaa5931>, 2015.

778 Guo, X., Tian, L., Wang, L., Yu, W., Qu, D.: River recharge sources and the partitioning of catchment  
779 evapotranspiration fluxes as revealed by stable isotope signals in a typical high-elevation arid catchment.  
780 *J. Hydrol.*, 549, 616–630, <https://doi.org/10.1016/j.jhydrol.2017.04.037>, 2017

781 Han, C., Ma, Y., Chen, X., Su, Z.: Trends of land surface heat fluxes on the Tibetan Plateau from 2001 to 2012,  
782 *Int. J. Climatol.*, 37, 4757–4767, <https://doi.org/10.1002/joc.5119>, 2017.

783 Han, C., Ma, Y., Wang, B., Zhong, L., Ma, W., Chen, X., and Su, Z.: Long-term variations of actual  
784 evapotranspiration over the Tibetan Plateau, *Earth Syst. Sci. Data.*, 13, 3513–3524,  
785 <https://doi.org/10.5194/essd-13-3513-2021>, 2021.

786 He, J., Yang, K., Tang, W., Lu, H., Qin, J., Chen, Y., Li, X.: The first high-resolution meteorological forcing  
787 dataset for land process studies over China, *Sci. Data.*, 7(1), 25, [https://doi.org/10.1038/s41597-020-0369-](https://doi.org/10.1038/s41597-020-0369-y)  
788 [y](https://doi.org/10.1038/s41597-020-0369-y), 2020.

789 Immerzeel, W. W., Van Beek, L. P. H., Bierkens, M. F. P.: Climate change will affect the Asian water towers.  
790 *Science*, 2010, 328(5984), 1382–1385. <https://doi.org/10.1126/science.1183188>

791 Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies,  
792 B., Elmore, A., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J., Koppes, M., Kraaijenbrink,  
793 P., Kulkarni, A., Mayewski, P., Nepal, S., Pacheco, P., Painter, T., Pellicciotti, F., Rajaram, H., Rupper, S.,  
794 Sinisalo, A., Shrestha, A., Viviroli, D., Wada, Y., Xiao, C., Yao, T., Baillie, J. E. M.: Importance and  
795 vulnerability of the world's water towers, *Nature*, 577(7790), 364–369, [https://doi.org/10.1038/s41586-](https://doi.org/10.1038/s41586-019-1822-y)  
796 019-1822-y, 2020.

797 Irmak, S., and Mutiibwa, D.: On the dynamics of canopy resistance: Generalized linear estimation and  
798 relationships with primary micrometeorological variables, *Water Resour. Res.*, 46(8), W08526,  
799 <https://doi.org/10.1029/2009WR008484>, 2010.

800 Jarvis, P. G.: The Interpretation of the Variations in Leaf Water Potential and Stomatal Conductance Found in  
801 Canopies in the Field. *Philos. Trans. R. Soc. London. B, Biol. Sci.*, 273(927), 593–610,  
802 <https://doi.org/10.1098/rstb.1976.0035>, 1976.

803 Jiang, Y., Yang, K., Qi, Y., Zhou, X., He, J., Lu, H., Li, X., Chen, Y., Li, X., Zhou, B., Mamtimin, A., Shao, C.,  
804 Ma, X., Tian, J., and Zhou, J.: TPHiPr: a long-term (1979–2020) high-accuracy precipitation dataset (1/  
805 30°, daily) for the Third Pole region based on high-resolution atmospheric modeling and dense  
806 observations, *Earth Syst. Sci. Data*, 15, 621–638, <https://doi.org/10.5194/essd-15-621-2023>, 2023.

807 Jung, M., Reichstein, M., Ciais, P., Seneviratne, S. I., Sheffield, J., Goulden, M. L., Bonan, G., Cescatti, A.,  
808 Chen, J., De Jeu, R., Dolman, A., Eugster, W., Gerten, D., Gianelle, D., Gobron, N., Heinke, J., Kimball,  
809 J., Law, B., Montagnani, L., Mu, Q., Mueller, B., Oleson, K., Papale, D., Richardson, A., Rouspard, O.,  
810 Running, S., Tomelleri, E., Viovy, N., Weber, U., Williams, C., Wood, E., Zaehle, S., Zhang, K.: Recent  
811 decline in the global land evapotranspiration trend due to limited moisture supply, *Nature*, 467(7318), 951–  
812 954, <https://doi.org/10.1038/nature09396>, 2010.

813 Khan, M. S., Liaqat, U. W., Baik, J., and Choi, M.: Stand-alone uncertainty characterization of GLEAM,  
814 GLDAS, and MOD16 evapotranspiration products using an extended triple collocation approach, *Agric.*  
815 *For. Meteorol.*, 252, 256–268, <https://doi.org/10.1016/j.agrformet.2018.01.022>, 2018.

816 Kuang, X., and Jiao, J. J.: Review on climate change on the Tibetan plateau during the last half century, *J.*  
817 *Geophys. Res. Atmos.*, 121, 3979–4007, <https://doi.org/10.1002/2015JD024728>, 2016.

818 Kutsch, W. L., Aubinet, M., Buchmann, N., Smith, P., Osborne, B., Eugster, W., Wattenbach, M., Schrumpf,  
819 M., Schulze, E., Tomelleri, E., Ceschia, E., Bernhofer, C., Béziat, P., Carrara, A., Di Tommasi, P.,

820 Grunwald, T., Jones, M., Magliulo, V., Moureaux, C., Olioso, A., Sanz, M., Saunders, M., Sogaard, H.,  
821 Ziegler, W.: The net biome production of full crop rotations in Europe, *Agric. Ecosyst. Environ.*, 139(3),  
822 336–345, <https://doi.org/10.1016/j.agee.2010.07.016>, 2010.

823 Kool, D., Agam, N., Lazarovitch, N., Heitman, J.L., Sauer, T.J., Ben-Gal, A.: A review of approaches for  
824 evapotranspiration partitioning, *Agric. For. Meteorol.*, 184, 56–70,  
825 <https://doi.org/10.1016/j.agrformet.2013.09.003>, 2014.

826 Koster, R. D., and Suarez, M. J.: The Influence of Land Surface Moisture Retention on Precipitation Statistics.  
827 *J. Clim.*, 9(10), 2551–2567, [https://doi.org/10.1175/1520-0442\(1996\)009](https://doi.org/10.1175/1520-0442(1996)009), 1996.

828 Lawrence, D. M., Thornton, P. E., Oleson, K. W., and Bonan, G. B.: The Partitioning of Evapotranspiration into  
829 Transpiration, Soil Evaporation, and Canopy Evaporation in a GCM: Impacts on Land–Atmosphere  
830 Interaction, *J. Hydrometeorol.*, 8, 862–880, <https://doi.org/10.1175/JHM596.1>, 2007.

831 Lehmann, P., Merlin, O., Gentine, P., and Or, D.: Soil texture effects on surface resistance to bare soil  
832 evaporation, *Geophys. Res. Lett.*, 45(19), 10, 398–10, 405, <https://doi.org/10.1029/2018GL078803>, 2018.

833 Lemone, M. A., Chen, F., Alfieri, J. G., Cuenca, R. H., Hagimoto, Y., Blanken, P., Niyogi, D., Kang, S., Davis,  
834 K., Grossman, R. L.: NCAR/CU surface, soil, and vegetation observations during the International H2O  
835 Project 2002 field campaign, *Bull. Am. Meteorol. Soc.*, 88(1), 65–81, [https://doi.org/10.1175/BAMS-88-](https://doi.org/10.1175/BAMS-88-1-65)  
836 [1-65](https://doi.org/10.1175/BAMS-88-1-65), 2007.

837 Letts, M.G., Comer, N.T., Roulet, N.T., Skarupa, M.R., Verseghy, D.L.: Parametrization of peatland hydraulic  
838 properties for the Canadian land surface scheme, *Atmos. Ocean.*, 38, 141–160,  
839 <https://doi.org/10.1080/07055900.2000.9649643>, 2000.

840 Leuning, R., Zhang, Y.Q., Rajaud, A., Cleugh, H., Tu, K.: A simple surface conductance model to estimate  
841 regional evaporation using MODIS leaf area index and the Penman–Monteith equation, *Water Resour.*,  
842 *Res.* 44 (10), W10419, <https://doi.org/10.1029/2007WR006562>, 2010.

843 Liang, S., Cheng, J., Jia, K., Jiang, B., Liu, Q., Xiao, Z., Yao, Y., Yuan, W., Zhang, X., Zhao, X., Zhou, J.: The  
844 global land surface satellite (GLASS) product suite, *Bull. Am. Meteorol. Soc.*, 102, E323–E337,  
845 <https://doi.org/10.1175/BAMS-D-18-0341.1>, 2021.

846 Liu, J., Chai, L., Dong, J., Zheng, D., Wigneron, J. P., Liu, S., Zhou, J., Xu, T., Yang, S., Song, Y., Qu, Y., Lu,  
847 Z.: Uncertainty analysis of eleven multisource soil moisture products in the third pole environment based  
848 on the three-corned hat method, *Remote Sens. Environ.*, 255, 112225,  
849 <https://doi.org/10.1016/j.rse.2020.112225>, 2021.

- 850 Liu, S., Lu, L., Mao, D., and Jia, L.: Evaluating parameterizations of aerodynamic resistance to heat transfer  
851 using field measurements, *Hydrol. Earth Syst. Sci.*, 11, 769–783, <https://doi.org/10.5194/hess-11-769-2007>,  
852 2007.
- 853 Liu, S.M., Li, X., Xu, Z.W., Che, T., Xiao, Q., Ma, M.G., Liu, Q.H., Jin, R., Guo, J.W., Wang, L.X., Wang,  
854 W.Z., Qi, Y., Li, H.Y., Xu, T.R., Ran, Y.H., Hu, X.L., Shi, S.J., Zhu, Z.L., Tan, J.L., Zhang, Y., Ren, Z.G.:  
855 The Heihe Integrated Observatory Network: A Basin-Scale Land Surface Processes Observatory in China,  
856 *Vadose Zo. J.*, 17(1), 180072, <https://doi.org/10.2136/vzj2018.04.0072>, 2018.
- 857 Liu, S. M., Xu, Z. W., Wang, W. Z., Jia, Z. Z., Zhu, M. J., Bai, J., and Wang, J. M.: A comparison of eddy-  
858 covariance and large aperture scintillometer measurements with respect to the energy balance closure  
859 problem, *Hydrol. Earth Syst. Sci.*, 15, 1291–1306, <https://doi.org/10.5194/hess-15-1291-2011>, 2011.
- 860 Li, S., Hao, X., Du, T., Tong, L., Zhang, J., Kang, S.: A coupled surface resistance model to estimate crop  
861 evapotranspiration in the arid region of northwest China, *Hydrol. Process.*, 28(4), 2312–2323,  
862 <https://doi.org/10.2136/vzj2018.04.0072>, 2013.
- 863 Li, S., Wang, G., Sun, S., Chen, H., Bai, P., Zhou, S., Huang, Y., Wang, J., Deng, P.: Assessment of Multisource  
864 Evapotranspiration Products over China Using Eddy Covariance Observations, *Remote Sens.*, 10(11),  
865 1692, <https://doi.org/10.3390/rs10111692>, 2018.
- 866 Li, X., Liang, S., Yuan, W., Yu, G., Cheng, X., Chen, Y., Zhao, T., Feng, J., Ma, Z., Ma, M., Liu, S., Chen, J.,  
867 Shao, C., Li, S., Zhang, X., Zhang, Z., Sun, G., Chen, S., Ohta, T., Varlagin, A., Miyata, A., Takagi, K.,  
868 Saiqusa, N., Kato, T.: Estimation of evapotranspiration over the terrestrial ecosystems in China,  
869 *Ecohydrology*, 7(1), 139–149, <https://doi.org/10.1002/eco.1341>, 2014a.
- 870 Li, X., Wang, L., Chen, D., Yang, K., and Wang, A.: Seasonal evapotranspiration changes (1983–2006) of four  
871 large basins on the Tibetan Plateau, *J. Geophys. Res. Atmos.*, 119(23), 13,079–13,095,  
872 <https://doi.org/10.1002/2014JD022380>, 2014b.
- 873 Li, S., Zhang, L., Kang, S., Tong, L., Du, T., Hao, X., and Zhao, P.: Comparison of several surface resistance  
874 models for estimating crop evapotranspiration over the entire growing season in arid regions, *Agric. For.*  
875 *Meteorol.*, 208, 1–15, <https://doi.org/10.1016/j.agrformet.2015.04.002>, 2015.
- 876 Ma, N., Szilagyi, J., Zhang, Y., Liu, W.: Complementary-Relationship-Based Modeling of Terrestrial  
877 Evapotranspiration Across China During 1982–2012: Validations and Spatiotemporal Analyses, *J.*  
878 *Geophys. Res. Atmos.*, 124(8), 4326–4351, <https://doi.org/10.1029/2018JD029850>, 2019.
- 879 Ma, N., Zhang, Y.: Increasing Tibetan Plateau terrestrial evapotranspiration primarily driven by precipitation,

880 Agric. For. Meteorol., 317, 108887, <https://doi.org/10.1016/j.agrformet.2022.108887>, 2022.

881 Ma, N., Zhang, Y., Guo, Y., Gao, H., Zhang, H., Wang, Y.: Environmental and biophysical controls on the  
882 evapotranspiration over the highest alpine steppe, *J. Hydrol.*, 529, 980–992,  
883 <https://doi.org/10.1016/j.jhydrol.2015.09.013>, 2015.

884 Ma, N., Y. Zhang, C.-Y. Xu, and J. Szilagyi: Modeling actual evapotranspiration with routine meteorological  
885 variables in the data-scarce region of the Tibetan Plateau: Comparisons and implications, *J. Geophys. Res.*  
886 *Biogeosci.*, 120, 1638–1657, <https://doi.org/10.1002/2015JG003006>, 2015.

887 Ma, Y., Hu, Z., Xie, Z., Ma, W., Wang, B., Chen, X., Li, M., Zhong, L., Sun, F., Gu, L., Han, C., Zhang, L.,  
888 Liu, X., Ding, Z., Sun, G., Wang, S., Wang, Y., and Wang, Z.: A long-term (2005–2016) dataset of hourly  
889 integrated land–atmosphere interaction observations on the Tibetan Plateau, *Earth Syst. Sci. Data.*, 12,  
890 2937–2957, <https://doi.org/10.5194/essd-12-2937-2020>, 2020.

891 Merlin, O., Stefan, V. G., Amazirh, A., Chanzy, A., Ceschia, E., Er-Raki, S., Khabba, S.: Modeling soil  
892 evaporation efficiency in a range of soil and atmospheric conditions using a meta-analysis approach, *Water*  
893 *Resour. Res.*, 52(5), 3663–3684, <https://doi.org/10.1002/2015WR018233>, 2016.

894 Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., Dolman, A. J.: Global  
895 land-surface evaporation estimated from satellite-based observations, *Hydrol. Earth Syst. Sci.*, 15(2), 453–  
896 469, <https://doi.org/10.5194/hess-15-453-2011>, 2011.

897 Miralles, D. G., Jiménez, C., Jung, M., Michel, D., Ershadi, A., McCabe, M. F., Hirschi, M., Martens, B.,  
898 Dolman, A. J., Fisher, J. B., Mu, Q., Seneviratne, S. I., Wood, E. F., and Fernández-Prieto, D.: The  
899 WACMOS-ET project – Part 2: Evaluation of global terrestrial evaporation data sets, *Hydrol. Earth Syst.*  
900 *Sci.*, 20, 823–842, <https://doi.org/10.5194/hess-20-823-2016>, 2016.

901 Monteith, J.L.: Evaporation and environment. *Symp. Soc. Exp. Biol.*, 19, 205–234, 1965.

902 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S.,  
903 Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D., Piles, M., Rodríguez-Fernández, N.,  
904 Zsoter, E., Thépaut, J. N.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications,  
905 *Earth Syst. Sci. Data*, 13(9), 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.

906 Mu, Q., Heinsch, F. A., Zhao, M., and Running, S. W.: Development of a global evapotranspiration algorithm  
907 based on MODIS and global meteorology data, *Remote Sens. Environ.*, 111(4), 519–536.  
908 <https://doi.org/10.1016/j.rse.2007.04.015>, 2007.

909 Mu, Q., Zhao, M., and Running, S. W.: Improvements to a MODIS global terrestrial evapotranspiration

910 algorithm, *Remote Sens. Environ.*, 115(8), 1781–1800, <https://doi.org/10.1016/j.rse.2011.02.019>, 2011.

911 Novick, K. A., Stoy, P. C., Katul, G. G., Ellsworth, D. S., Siqueira, M. B. S., Juang, J., Oren, R.: Carbon dioxide  
912 and water vapor exchange in a warm temperate grassland, *Oecologia.*, 138(2), 259–274,  
913 <https://doi.org/10.1007/s00442-003-1388-z>, 2004.

914 Ortega-Farias, S., Poblete-Echeverría, C., and Brisson, N.: Parameterization of a two-layer model for estimating  
915 vineyard evapotranspiration using meteorological measurements, *Agric. For. Meteorol.*, 150(2), 276–286,  
916 <https://doi.org/10.1016/j.agrformet.2009.11.012>, 2010.

917 Peng, J., Loew, A., Chen, X., Ma, Y., and Su, Z.: Comparison of satellite-based evapotranspiration estimates  
918 over the Tibetan Plateau. *Hydrol, Earth Syst. Sci.*, 20, 3167–3182, [https://doi.org/10.5194/hess-20-3167-](https://doi.org/10.5194/hess-20-3167-2016)  
919 2016, 2016.

920 Phillips, T. J., Klein, S. A., Ma, H. Y., Tang, Q., Xie, S., Williams, I. N., Joseph, A., David, R., Margaret, S.:  
921 Using ARM observations to evaluate climate model simulations of land-atmosphere coupling on the U.S.  
922 Southern Great Plains, *J. Geophys. Res. Atmos.*, 122(21), 11,524–11, 548,  
923 <https://doi.org/10.1002/2017JD027141>, 2017.

924 Ramoelo, A., Majozi, N., Mathieu, R., Jovanovic, N., Nickless, A., and Dzikiti, S.: Validation of Global  
925 Evapotranspiration Product (MOD16) using Flux Tower Data in the African Savanna, South Africa,  
926 *Remote Sens.-Basel*, 6, 7406–7423, <https://doi.org/10.3390/rs6087406>, 2014.

927 Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C. J., Arsenault, K., Cosgrove, B.,  
928 Radakovich, J., Bosilovich, M., Entin, J., Walker, J., Lohmann, D., Toll, D.: The Global Land Data  
929 Assimilation System, *Bull. Am. Meteorol. Soc.*, 85, 381–394, <https://doi.org/10.1175/BAMS-85-3-381>,  
930 2004.

931 Sakaguchi, K., Zeng, X.: Effects of soil wetness, plant litter, and under-canopy atmospheric stability on ground  
932 evaporation in the Community Land Model (CLM3.5), *J. Geophys. Res., Atmos.* 114(D1),  
933 <https://doi.org/10.1029/2008JD010834>, 2009.

934 Sellers, P. J., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D. A., and Bounoua, L.: A Revised  
935 Land Surface Parameterization (SiB2) for Atmospheric GCMS. Part I: Model Formulation, *J. Clim.* 9(4),  
936 676–705, [https://doi.org/10.1175/1520-0442\(1996\)009<0676:ARLSPF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<0676:ARLSPF>2.0.CO;2), 1996.

937 Schlesinger, W. H., and Jasechko, S.: Transpiration in the global water cycle, *Agric. For. Meteorol.*, 189-190,  
938 115–117, <https://doi.org/10.1016/j.agrformet.2014.01.011>, 2014.

939 Shi, Q., Liang, S.: Surface-sensible and latent heat fluxes over the Tibetan Plateau from ground measurements,

940 reanalysis, and satellite data, *Amos. Chem. Phys.*,14, 5659–5677, <https://doi.org/10.5194/acp-14-5659->  
941 2014, 2014.

942 Sobrino, J. A., Jiménez-Muñoz, J. C., and Paolini, L.: Land surface temperature retrieval from LANDSAT TM  
943 5, *Remote Sens. Environ.*, 90(4), 434–440, <https://doi.org/10.1016/j.rse.2004.02.003>, 2004.

944 Song, L., Zhuang, Q., Yin, Y., Zhu, X., and Wu, S.: Spatio-temporal dynamics of evapotranspiration on the  
945 Tibetan Plateau from 2000 to 2010, *Environ. Res. Lett.*, 12(1), 014011, <https://doi.org/10.1088/1748->  
946 9326/aa527d, 2017.

947 Su, Z.: The Surface Energy Balance System (SEBS) for estimating turbulent heat fluxes, *Hydrol. Earth Syst.*  
948 *Sci.*, 6, 85–100, <https://doi.org/10.5194/hess-6-85-2002>, 2002.

949 Sun, S.F.: Moisture and heat transport in a soil layer forced by atmospheric conditions, Master thesis, Dept. of  
950 Civil Engineering, University of Connecticut, 72, 1982.

951 Tang, J. Y. and Riley, W. J.: A new top boundary condition for modeling surface diffusive exchange of a generic  
952 volatile tracer: theoretical analysis and application to soil evaporation, *Hydrol. Earth Syst. Sci.*, 17, 873–  
953 893, <https://doi.org/10.5194/hess-17-873-2013>, 2013.

954 Wang, B., Ma, Y., Su, Z., Wang, Y., and Ma, W.: Quantifying the evaporation amounts of 75 high-elevation  
955 large dimictic lakes on the Tibetan Plateau, *Sci. Adv.*, 6(26), <https://doi.org/10.1126/sciadv.aay8558>, 2020.

956 Wang, G., Lin, S., Hu, Z., Lu, Y., Sun, X., and Huang, K.: Improving Actual Evapotranspiration Estimation  
957 Integrating Energy Consumption for Ice Phase Change Across the Tibetan Plateau, *J. Geophys. Res.*  
958 *Atmos.*, 125(3), e2019JD031799, <https://doi.org/10.1029/2019JD031799>, 2020.

959 Wang, W., Li, J., Yu, Z., Ding, Y., Xing, W., Lu, W.: Satellite retrieval of actual evapotranspiration in the  
960 Tibetan Plateau: components partitioning, multi-decadal trends and dominated factors identifying, *J.*  
961 *Hydrol.*, 559, 471–485, <https://doi.org/10.1016/j.jhydrol.2018.02.065>, 2018.

962 Wang, Y., Lv, W., Xue, K. et al. Grassland changes and adaptive management on the Qinghai–Tibetan  
963 Plateau. *Nat. Rev. Earth. Environ* 3, 668–683 (2022). <https://doi.org/10.1038/s43017-022-00330-8>

964 Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S., Lee, X.: Revisiting the contribution of  
965 transpiration to global terrestrial evapotranspiration. *Geophysical Research Letters*, 44, 2792–2801.  
966 <https://doi.org/10.1002/2016GL072235>

967 Wieder, W.R., J. Boehnert, G.B. Bonan, and M. Langseth.: RegridDED Harmonized World Soil Database v1.2.  
968 Data set, Available online [<http://daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active  
969 Archive Center, Oak Ridge, Tennessee, USA, <http://dx.doi.org/10.3334/ORNLDAAAC/1247>, 2014.

- 970 Wilcox, B.P., Breshears, D.D., Seyfried, M.S.: Water balance on rangelands, In: Stewart, B.A., Howell, T.A.  
971 (Eds.), *Encyclopedia of Water Science*, Marcel Dekker Inc, New York, 791–794,  
972 <http://www.cprl.ars.usda.gov/wmru/pdfs/DekkerEvetTDR.pdf>, 2003.
- 973 Wu, C., Hu, B. X., Huang, G., and Zhang, H.: Effects of climate and terrestrial storage on the temporal variability  
974 of actual evapotranspiration, *J. Hydrol.*, 549, 388–403, <https://doi.org/10.1016/j.jhydrol.2017.04.012>, 2017.
- 975 Yang, K., He, J., Tang, W., Qin, J., Cheng, C. C. K.: On downward shortwave and longwave radiations over  
976 high altitude regions: Observation and modeling in the Tibetan Plateau, *Agric. For. Meteorol.*, 150(1), 38–  
977 46, <https://doi.org/10.1016/j.agrformet.2009.08.004>, 2010.
- 978 Yang, K., Koike, T., Ishikawa, H., Kim, J., Li, X., Liu, H., Liu S., Ma Y., Wang, J.: Turbulent flux transfer over  
979 bare-soil surfaces: Characteristics and parameterization, *J. Appl. Meteorol. Climatol.*, 47(1), 276–290,  
980 <https://doi.org/10.1175/2007JAMC1547.1>, 2008.
- 981 Yang, K., Wu, H., Qin, J., Lin, C., Tang, W., and Chen, Y.: Recent climate changes over the Tibetan Plateau  
982 and their impacts on energy and water cycle: A review, *Glob. Planet. Change.*, 112, 79–91,  
983 <https://doi.org/10.1016/j.gloplacha.2013.12.001>, 2014.
- 984 Yang, Y., Liu, Y., Li, M., Hu, Z., Ding, Z.: Assessment of reanalysis flux products based on eddy covariance  
985 observations over the Tibetan Plateau, *Theor. Appl. Climatol.*, 138, 275–292,  
986 <https://doi.org/10.1007/s00704-019-02811-1>, 2019.
- 987 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu,  
988 A., Xiang, Y., Kattel, D., Joswiak, D.: Different glacier status with atmospheric circulations in Tibetan  
989 Plateau and surroundings, *Nature Clim.Change*, 2(9), 663–667, <https://doi.org/10.1038/nclimate1580>,  
990 2012.
- 991 Yao, Y., Liang, S., Cheng, J., Liu, S., Fisher, J. B., Zhang, X., Jia, K., Zhao, X., Qin, Q., Zhao, B., Han, S.,  
992 Zhou, G., Li, Y., Zhao, S.: MODIS-driven estimation of terrestrial latent heat flux in China based on a  
993 modified Priestley-Taylor algorithm, *Agric. For. Meteorol.*, 171–172, 187–202,  
994 <https://doi.org/10.1016/j.agrformet.2012.11.016>, 2013.
- 995 Yin, Y., Wu, S., Zhao, D., Zheng, D., Pan, T.: Modeled effects of climate change on actual evapotranspiration  
996 in different eco-geographical regions in the Tibetan Plateau, *J. Geogr. Sci.*, 23(2), 195–207,  
997 <https://doi.org/10.1002/eco.1341>, 2013.
- 998 You, Q., Xue, X., Peng, F., Dong, S., Gao, Y.: Surface water and heat exchange comparison between alpine  
999 meadow and bare land in a permafrost region of the Tibetan Plateau, *Agric. For. Meteorol.*, 232, 48–65,



1000 <https://doi.org/10.1016/j.agrformet.2016.08.004>, 2017.

1001 Yuan, L., Ma, Y., Chen, X., Wang, Y., Li, Z.: An enhanced MOD16 evapotranspiration model for the Tibetan  
1002 Plateau during the unfrozen season, *J. Geophys. Res. Atmos.*, 126, e2020JD032787,  
1003 <https://doi.org/10.1029/2020JD032787>, 2021.

1004 Yu, G. R., Wen, X. F., Sun, X. M., Tanner, B. D., Lee, X., Chen, J. Y.: Overview of ChinaFLUX and evaluation  
1005 of its eddy covariance measurement, *Agric. For. Meteorol.*, 137(3–4), 125–137,  
1006 <https://doi.org/10.1016/j.agrformet.2006.02.011>, 2006.

1007 Zhang, G., T. Yao, H. Xie, K. Yang, L. Zhu, C. K. Shum, T. Bolch, S. Yi, S. Allen, L. Jiang, W. Chen, and C.  
1008 Ke: Response of Tibetan Plateau lakes to climate change: Trends, patterns, and mechanisms, *Earth-Science*  
1009 *Reviews*, 103269. <https://doi.org/10.1016/j.earscirev.2020.103269>, 2020.

1010 Zhang, L. M., Luo, Y. W., Liu, M., Chen, Z., Su, W., He et al.: Carbon and water fluxes observed by the Chinese  
1011 Flux Observation and Research Network (2003–2005) (in Chinese), *Sci. Data.*, 4(1),  
1012 <https://doi.org/10.11922/csdata.2018.0028.zh>, 2019.

1013 Zhang, K., Kimball, J. S., Nemani, R. R., and Running, S. W.: A continuous satellite-derived global record of  
1014 land surface evapotranspiration from 1983 to 2006, *Water Resour. Res.*, 46(9), W09522,  
1015 <https://doi.org/10.1029/2009WR008800>, 2010.

1016 Zhang, Y., Kong, D., Gan, R., Chiew, F. H. S., McVicar, T. R., Zhang, Q., and Yang, Y.: Coupled estimation  
1017 of 500 m and 8-day resolution global evapotranspiration and gross primary production in 2002–2017,  
1018 *Remote Sens. Environ.*, 222, 165–182, <https://doi.org/10.1016/j.rse.2018.12.031>, 2019b.

1019 Zhang, Y., Peña-Arancibia, J. L., McVicar, T. R., Chiew, F. H. S., Vaze, J., Liu, C., Lu, X., Zheng, H., Wang,  
1020 Y., Liu, Y., Miralles, D., Pan, M.: Multi-decadal trends in global terrestrial evapotranspiration and its  
1021 components, *Sci. Rep.*, 6, 19124, <https://doi.org/10.1038/srep19124>, 2016.

1022 Zhao, H., Zeng, Y., Lv, S., and Su, Z.: Analysis of soil hydraulic and thermal properties for land surface  
1023 modeling over the Tibetan Plateau, *Earth Syst. Sci. Data.*, 10, 1031–1061, [https://doi.org/10.5194/essd-10-](https://doi.org/10.5194/essd-10-1031-2018)  
1024 [1031-2018](https://doi.org/10.5194/essd-10-1031-2018), 2018.

1025 Zheng, C., Jia, L., Hu, G.: Global Land Surface Evapotranspiration Monitoring by ETMonitor Model Driven  
1026 by Multi-source Satellite Earth Observations. *J. Hydrol.*, 128444,  
1027 <https://doi.org/10.1016/j.jhydrol.2022.128444>, 2022.

1028 Zheng, D., Zhang, Q., Wu, S.: Mountain genecology and sustainable development of the Tibetan Plateau,  
1029 Springer Science & Business Media, 57, 2000.

1030 Zhong, L., Ma, Y., Hu, Z., Fu, Y., Hu, Y., Wang, X., Cheng, M., and Ge, N.: Estimation of hourly land surface  
1031 heat fluxes over the Tibetan Plateau by the combined use of geostationary and polar-orbiting satellites,  
1032 *Atmos. Chem. Phys.*, 19, 5529–5541, <https://doi.org/10.5194/acp-19-5529-2019>, 2019.

1033 Zohaib, M., H. Kim, and M. Choi.: Evaluating the patterns of spatiotemporal trends of root zone soil moisture  
1034 in major climate regions in East Asia, *J. Geophys. Res. Atmos.*, 122, 7705–7722,  
1035 <https://doi.org/10.1002/2016JD026379>, 2017.