

TiP-Leaf: A dataset of leaf traits across vegetation types on the Tibetan Plateau

Yili Jin¹, Haoyan Wang¹, Jie Xia¹, Jian Ni¹, Kai Li¹, Ying Hou¹, Jing Hu¹, Linfeng Wei¹, Kai Wu¹, Haojun Xia¹, Borui Zhou¹

¹College of Chemistry and Life Sciences, Zhejiang Normal University, Jinhua 321004, China

Correspondence [to](mailto:nijian@zjnu.edu.cn): Jian Ni (nijian@zjnu.edu.cn)

Abstract. Functional trait databases are emerging as a crucial tool for a wide range of ecological studies, including the next-generation vegetation modelling across the world. However, few large-scale studies have been reported on plant traits in the Tibetan Plateau (TP), the cradle of East Asian flora and fauna with specific alpine ecosystems, no report on plant trait databases could be found. ~~Here~~ In this work, an extensive dataset of 11 leaf functional traits (TiP-Leaf) mainly for ~~mainly~~ herbs and shrubs and a few trees on the TP was compiled through field surveys. The TiP-Leaf dataset, which was compiled from 336 sites distributed mainly in the plateau surface and the northern margin of the TP across alpine and temperate vegetation regions and sampled from 2018 to 2021, ~~contains~~ contained 1692 morphological trait measurements of leaf thickness, leaf fresh weight, leaf dry weight, leaf dry-matter content, leaf water content, leaf area, specific leaf area and leaf mass per area and 1645 chemical element trait measurements of leaf carbon, nitrogen and phosphorus contents. Thus, 468 species that ~~belonging~~ to 184 genera and 51 families were obtained and measured. In addition to leaf trait measurements, the geographic coordinates, bioclimate variables, disturbance intensity and vegetation types of each site were also recorded. The dataset could provide solid data support ~~for~~ to effectively ~~quantifying~~ quantify the modern ecological features of alpine ecosystems, thereby further evaluating the response of alpine ecosystem to climate change and human disturbances and improving the next-generation vegetation model. ~~It could be a great contribution to the regional and global plant trait databases.~~ The dataset, which is available from the National Tibetan Plateau Data Center (TPDC; Jin et al., 2022a; <https://doi.org/10.11888/Terre.tpd.272516>), can make a great contribution to the regional and global plant trait databases.

1 Introduction

Plant traits of morphological, anatomical, physiological and phenological characteristics respond to

28 changes in the living environment, affect ecosystem functions (Díaz & Cabido, 2001) and drive species
29 coexistence under environmental constraints (Violle et al., 2007). Over the past three decades, a growing
30 body of trait analyses has quantified the global and regional distribution patterns of key functional traits,
31 such as leaf (Reich & Oleksyn, 2004; Wright et al., 2004), seed size (Moles et al., 2007), plant height
32 (Moles et al., 2009), wood (Chave et al., 2009), plant form and function (Díaz et al., 2016), root (Ma
33 et al., 2018) and flower (Roddy et al., 2021). Such studies have successfully linked plant traits with
34 environmental changes (Meng et al., 2009, 2015; Myers-Smith et al., 2019; Maes et al., 2020; Wang et
35 al., 2022), natural and anthropogenic disturbances (Díaz et al., 2007) and ecosystem functions
36 (Reichstein et al., 2014). Findings from plant trait–environment–ecosystem function interaction could be
37 further utilised to map the spatial pattern of plant traits (Butler et al., 2018), build the next-generation of
38 vegetation model (Berzaghi et al., 2020), predict vegetation distribution (van Bodegom et al., 2014) and
39 function (Wang et al., 2017) and be incorporated into the Earth system model (Wullschleger et al., 2014).
40 New insights into ecosystem traits (He et al., 2019) and trait network (He et al., 2020) are bridging
41 multiple dimensions of biology, macroecology and geoscience. All these works require global and
42 regional plant trait databases, such as the TRY (Kattge et al., 2011, 2020), Growth-Form (Taseski et al.,
43 2019), Global Inventory of Floras and Traits (Weigelt et al., 2019), Fine-Root (Iversen et al., 2017),
44 GRoot (Guerrero-Ramírez et al., 2021), and tundra traits (Bjorkman et al., 2018), Plant Trait for
45 Mediterranean Basin Species (BROT) (Tavşanoğlu & Pausas, 2018), China traits (Wang et al., 2018),
46 Aus-Traits (Falster et al., 2021) and LT-Brazil (Mariano et al., 2021).

47 Plant trait databases across various biomes at global, continental and regional scales have been
48 largely raised largely, even in some remote areas with logistical difficulties, including the tundra
49 (Bjorkman et al., 2018) and tropical regions (Mariano et al., 2021). However, the Earth still has under-
50 sampled regions. The Tibetan Plateau (TP), includes the two major regions of Qinghai Province and
51 Xizang Autonomous Region, and partial areas from northwestern Gansu Province, western Sichuan
52 Province and northwestern Yunnan Province in China, is also called ‘Qinghai-Tibetan Plateau’. ~~known~~
53 ~~a~~As the world’s “Third Pole” and “Asia Water Tower” and the cradle of the East Asian flora, TP is
54 the most under-representative region in global and regional plant trait databases. The first version of the
55 Chinese plant trait database (Wang et al., 2018) ~~has~~ does not contain data from the TP, and the global
56 plant trait database TRY (Kattge et al., 2020) has only a few collections from various sources with non-

57 systematic sampling. Field-based, ~~small regional~~local studies of plant functional traits on the TP had
58 made some interesting advances. For example, Luo et al. (2005) linked the plant traits with ecosystem
59 functions, He et al. (2006) explored the influencing factors on plant traits, Geng et al. (2004) quantified
60 the patterns of plant trait correlations between above- and below-ground components, Wang et al. (2020)
61 compared their work with global dataset, and Xu et al. (2021) analysed the mechanism of plant trait
62 variation along the altitude pattern. Yet such works, where the sampling sites have been mostly along the
63 main roads in East TP, were also limited (Luo et al., 2005; He et al., 2006; He et al., 2010; Geng et al.,
64 2014; Wang et al., 2020; Xu et al., 2021), where the sampling sites have been mostly along the main
65 roads in East TP. Plant trait records from Central to West TP are ~~very extremely~~ rare. However, the TP
66 has the richest temperate alpine flora (Ding et al., 2020) and ~~the~~ most abundant plant diversity in the
67 world (Wang & Hong, 2022). It was also an evolutionary cradle for ~~the~~ herbaceous genera of China (Lu
68 et al., 2018). The uplift of the TP and its unique alpine vegetation are important to the monsoon climate
69 system and vegetation of East Asia (Chang et al., 1983) and regional and global climate change studies
70 (Piao et al., 2019).


71 As the largest and highest plateau in the world, the TP has not only changed the regional and global
72 climate system, geological structure, topography and hydrology (Yao et al., 2012) but also strongly
73 influenced the evolution of the flora, fauna and biodiversity strongly (Ding et al., 2020). It has 8876
74 vascular species from 1371 genera and 211 families, including 6475 herbaceous and 2401 woody plants,
75 of which 1706 were endemic to the TP (Yan et al., 2013). Here has three biodiversity hotspots of the
76 world (Sloan et al., 2014; Wang & Hong, 2022). Vegetation changes from the southeast to northwest,
77 from lowland broad-leaved evergreen forests, including tropical rainforest and subtropical evergreen
78 forest, montane mixed evergreen and deciduous forest, subalpine conifer forest to alpine shrubland,
79 meadow, steppe and desert, along an annual precipitation gradient from ca. 3000 mm to 50 mm (Chang,
80 1983). In physiognomy, ~~t~~The unique alpine vegetation looks ~~like-similar to~~ the arctic tundra ~~in~~
81 physiognomy but has different species composition. The plateau has amplified changes in climates (Chen
82 et al., 2015), and rapid climate change has led to profound changes in alpine species and ecosystems
83 (Zhang et al., 2015; Piao et al., 2019). Plant traits, as the link amongst species, environment and
84 ecosystem functions, are a ~~the~~ best tools to study the impacts of climate change on vegetation. Therefore,
85 the establishment of TP plant trait database and further analysis of plant trait–environment–ecosystem

86 function relationships are of great significance to understanding the future change and sustainable
87 development of the unique alpine vegetation on the roof of the world.

88 [HereIn this work](#), a TP leaf trait dataset (TiP-Leaf) was established, and 11 leaf traits from 468
89 species of 1692 leaf samples were collected from 336 sites across five of [the](#) six vegetation types on the
90 TP. [The Climate-climate](#) data of the sites were also provided. This dataset is not only an update of the
91 Chinese plant trait database but also a great contribution to the global trait database.

92 2 Study areas

93 The leaf traits of dominant and common plant species of the TP distributed mostly across the plateau
94 surface were sampled and measured from July to August in the summer of 2018–2021. Vegetation
95 surveys were conducted in four regions (Fig. 1), [namely](#), the source area of the Three Rivers in Qinghai
96 Province in Northeast TP (2018); Southern Xizang Autonomous Region in Southeast and Middle TP
97 (2019); Ngari Prefecture in Northwest TP and the Qaidam Basin of Northeast TP (2020); and the Qilian
98 Mountains, the Altun Mountains and the Kunlun Mountains in the northern margin of the TP, passing
99 through the southern margin of the Tarim Basin in Xinjiang Autonomous Region (2021).




100

101 **Figure 1.** Location and administrative division of the TP. The red line indicates the boundary of the TP in China,
102 which involves six administrative divisions (light black lines): Xizang, Qinghai, Sichuan, Xinjiang, Gansu and
103 Yunnan. The bold black lines represent important mountains. Four blocks with different colours represent the
104 approximate areas of four investigations conducted in various years. The background map is from the Chinese
105 National Bureau of Surveying and Mapping.

106 **3 Materials and methods**

107 **3.1 Sampling sites**

108 ~~Taking~~ Considering the zonal vegetation types and the precipitation gradient ~~into account~~, 336 sites
109 (Fig. 2) were selected to investigate the vegetation with less grazing and other anthropogenic disturbances.
110 Shrubby and herbaceous vegetation were mainly selected (332 sites), along with the forest vegetation in
111 the four sites (but removed for further analysis). In each site, 1–3 plots were set up to survey the species
112 composition, abundance, coverage and plant height. The plot areas for herbaceous vegetation, shrubby
113 vegetation and forest vegetation were 1 m × 1 m, 2 m × 2 m or 5 m × 5 m and 10 m × 10 m, respectively.
114 Geographical locations, natural and human disturbances, and vegetation types were also recorded (Jin et
115 al., 2022b). The dominant and common plant species in each site were determined by visual inspection,
116 the Plant-leaf samples of these plants were picked up, and the leaf traits were measured. Root samples
117 were obtained using soil pit method. The root traits were also measured but not shown in this paper.



118

119

120

121

122

123

124

125

126

127

Figure 2. Site distribution of the TiP-Leaf dataset. Vegetation regions were extracted from the vegetation regionalisation of China (ECVMC, 2007b). III, Warm Temperate Deciduous Broad-leaf Forest Region; IV, Subtropical Broad-leaf Evergreen Forest Region; V, Tropical Monsoon Rain Forest and Rain Forest Region; VI, Temperate Steppe Region; VII, Temperate Desert Region; VIII, TP Alpine Vegetation Region. VIII A, East TP Alpine Scrub and Alpine Meadow Subregion; VIII B, Middle TP Alpine Steppe Subregion; VIII C, Northwest TP Alpine Desert Subregion. Vegetation types were classified on the basis of field records. Numbers indicate the vegetation types recorded in the field. 1, coniferous forest; 2, alpine shrubland; 3, alpine meadow; 4, alpine steppe; 5, alpine desert; 6, temperate desert; 7, temperate steppe; 8, temperate meadow. The background map is from the Chinese National Bureau of Surveying and Mapping.

128

129

130

131

132

133

134

135

136

The vegetation type of the TP was classified ~~on the basis of field records~~ into eight types: high-cold (alpine) shrubland, meadow, steppe and desert, temperate steppe, meadow and desert, and coniferous forest, ~~on the basis of field records~~ (Fig. 2). Alpine shrubland is dominated by evergreen broad-leaved shrubs (*Rhododendron*), deciduous broad-leaved shrubs (*Salix*, *Dasiphora*, and *Sibiraea*) and evergreen coniferous shrubs (*Juniperus*); ~~that are~~ distributed in the cold and semi-humid Southeast TP (ca. 600–1000 mm/year). Alpine meadow is widely developed in East TP, where cold and wet climates ~~dominate~~ are prevalent (ca. 600 mm/year), dominated by several *Kobresia* species and mixed in with perennial forbs and cushion plants. Alpine steppe is in the middle TP, with a large continuous distribution, adapted to the cold and semi-dry continental climate (ca. 200 mm/year) and mainly composed of *Stipa* and

137 *Artemisia*. Alpine desert is mainly distributed in Northwest TP, where the climate is extremely continental
138 (ca. 50 mm/year) ~~and~~, dominated by *Krascheninnikovia compacta* and *Ajanía tibética*. Temperate
139 meadow, steppe and desert are distributed in the northern margin of the TP and Qaidam Basin of
140 Northeast TP, where elevations are lower, and the climate is relatively dry, dominated by several
141 xerophytes, especially *Haloxylon ammodendron*, *Halogeton glomeratus*, *Phragmites australis*, *Ephedra*,
142 *Kalidium*, *Calligonum* and *Tamarix*. Subalpine coniferous (*Abies* and *Picea*) forests are found on the
143 southeast and east margins. Therefore, the vegetation of the plateau is distributed along a transitional
144 gradient from southeast to northwest, arraying from subalpine forests, alpine meadow and scrub, through
145 alpine steppe and temperate desert to alpine desert. The alpine vegetation in the vegetation classification
146 of China was usually called high-cold vegetation ~~in the vegetation classification of China~~ [Editorial
147 Committee of Vegetation Map of China (ECVMC), The Chinese Academy of Sciences, 2007a]. Lowland
148 tropical and montane subtropical evergreen forests do not exist in the sampling area, ~~hence~~ Hence, they
149 are not included in this study.—

150 Each site was also assigned to a vegetation region on the basis of the vegetation regionalisation of
151 China (ECVMC, 2007b). The TP has six vegetation regions: ~~—~~, namely, Alpine Vegetation Region,
152 Temperate Steppe Region, Temperate Desert Region, Warm Temperate Deciduous Broad-leaf Forest
153 Region, Subtropical Broad-leaf Evergreen Forest Region, and Tropical Monsoon Rain Forest and Rain
154 Forest Region. The sampling sites were mainly concentrated in the Alpine Vegetation Region. Therefore,
155 in accordance with the degree of drought, the Selianinov drought index used in the Vegetation
156 Regionalisation Map of China (ECVMC, 2007b), TP vegetation was further divided into three subregions
157 from southeast to northwest: namely, East TP Alpine Scrub and Alpine Meadow Subregion, Middle TP
158 Alpine Steppe Subregion and Northwest TP Alpine Desert Subregion.

159 The plant name was determined in accordance with *Flora of China* (Editorial Committee of Flora
160 of China, 1959—2004), *Flora of Qinghai* (Editorial Committee of Flora of Qinghai, 1996—1999), *Flora*
161 *of Xizang* (Comprehensive Scientific Investigation Team of Qinghai-Tibet Plateau, Chinese Academy of
162 Sciences, 1985—1987), *Flora of Gansu* (Editorial Committee of flora of Gansu, 2005), *Flora of Xinjiang*
163 (Editorial Committee of flora of Xinjiang, 1993—1996) and *Flora of Deserts in China* (Liu, 1985). The
164 final species correction was based on the iPlant website (<http://www.iplant.cn/>), which merged all of the
165 information from the Chinese and English versions of *Flora of China* on the basis of APG IV

166 classification (Angiosperm Phylogeny Group, 2016).—

167 3.2 Leaf trait measurements

168 At each site, 2–~~to~~3 mature and disease-free complete leaves from each individual of dominant and
169 common plant species were collected, and at least 30 individuals were selected to meet the needs of trait
170 measurement and element analysis. When the single leaf was small, micro or leptophyllous, 100–200
171 leaves were picked. In total, 11 leaf functional traits (e.g. leaf thickness, LT; fresh weight, FW; dry weight,
172 DW; leaf dry-matter content, LDMC; leaf water content, LWC; leaf area, LA; specific leaf area, SLA;
173 leaf mass per area, LMA; leaf carbon content, LCC; leaf nitrogen content, LNC; and leaf phosphorus
174 content, LPC) were measured and calculated on the basis of the handbook of standardised measurement
175 for plant functional traits worldwide (Cornelissen et al., 2003; Pérez-Harguindeguy et al., 2013).—

176 LT (mm) was measured on the sampling day by using Vernier callipers with an accuracy of 0.01
177 mm. The thickness in the middle of the vein and margin of each leaf was measured, and then the average
178 of the five leaves was taken as the LT of a species. In addition to LT, 20–30 leaves for normal-leaved
179 plants and 100–200 leaves for small- to leptophyll-leaved plants were generally selected for other trait
180 variable measurements. FW (g) was obtained by weighing with 1/100 electronic balance. Subsequently,
181 ~~The~~ the fresh leaves were ~~then~~ oven dried at 75 °C for 48–72 hours to obtain the DW (g). LDMC was
182 measured as follows: $LDMC (g \cdot g^{-1}) = DW / FW$. LA (cm²) was measured using a scanner (EPSON
183 Perfection V 700 Photo Scanner) and a software (WinFOLIA Pro, Canada). SLA and LMA were
184 measured as follows: $SLA (cm^2 \cdot g^{-1}) = LA / DW$ and $LMA (g \cdot m^{-2}) = DW / LA \times 10^4$. The dried leaves were
185 further used for chemical analysis. LCC ($mg \cdot g^{-1}$), LNC ($mg \cdot g^{-1}$) and LPC ($mg \cdot g^{-1}$) were determined by
186 outside-temperature hot potassium dichromate oxidation–volumetric method (Wu, 2007), distillation–
187 titration method and vanadium molybdate yellow colorimetric method (Fang et al., 2011), respectively.

188 3.3 Climate data

189 The climate data of each sampling site were extracted from the climate and bioclimate datasets of
190 China (Cheng et al., submitted2022; Wei et al., 2022). China’s climate dataset consists of three variables
191 (monthly temperature, precipitation and sunshine percentage) that were averaged from long-term records
192 from 1981 to 2010 at 2152 meteorological stations across China (China Meteorological Data Service
193 Centre, <http://data.cma.cn>). These three climate factors and the absolute maximum and minimum

194 temperatures during the 30-year periods of 1981–2010 were interpolated into 1 km grid cells by using a
195 surface fitting technique of thin-plate smoothing spline (ANUSPLIN version 4.4, Hutchinson & Xu, 2013;
196 Xu & Hutchinson, 2013) that took-considered the impact of elevation on climates into-account on the
197 basis of the digital elevation model of the Shuttle Radar Topography Mission (~~SRTM~~) (Farr et al., 2007).
198 The interpolated climate data were used to drive a bioclimate software (Gallego-Sala et al., 2010) to
199 calculate the mean annual temperature (MAT), mean temperature of the coldest month (MTCO), mean
200 temperature of the warmest month (MTWA), annual growing degree days above 0 °C (GDD₀) and 5 °C
201 (GDD₅), mean annual precipitation (MAP), growing season precipitation (GP), annual drought index (1-
202 AET/PET) and annual moisture index (MAP/PET), where AET and PET refer to their annual actual
203 evapotranspiration and ~~PET is~~ annual potential evapotranspiration, respectively.

204 **3.4 Data analysis**

205 Beside the data description of leaf trait characteristics, sSix key leaf functional traits (e.g. LT, LDMC,
206 SLA, LCC, LNC and LPC), that reflect the key ecological significances of plants that grew in high
207 altitude and extremely cold environment, were selected in this paper for further simple statistical analyses.
208 LT affects the water supply and storage of leaves and the exchange process of matter and energy in
209 photosynthesis; LDMC reflects the ability of plants to acquire surrounding environmental resources; SLA
210 is considered the first choice index for studying plant physiological and ecological strategies under
211 specific environmental conditions; LCC is the main structural material of plants; LNC characterises the
212 ability of plants to absorb and utilise nutrient elements; and LPC is the second largest element that affects
213 plant growth. The mean, minimum, maximum, standard deviation (~~SD~~) and coefficient variation of traits
214 at each site were calculated to generally show the pattern of leaf traits of the Tibetan ecosystems. The
215 linear relationships between leaf traits of site average were analysed and mapped using the Origin
216 software (The Origin Lab, 2022) to reveal the trade-off between different traits in the special alpine
217 ecosystem. The detailed analyses of all the leaf traits, their variations and spatial patterns, within and
218 amongst functional groups and at species and site levels, will be further analysed in another paper.—

219 **4 Data description of sampling sites**

220 **4.1 Spatial distribution of sites**

221 ~~Eleven~~ A total of 11 key plant leaf traits of 1692 individuals of 468 species from 336 sites were
222 measured (Fig. 1 and 2).

223 The sampling sites were located in the northeast, middle to southwest, and north margin of the TP,
224 along with 145 sites in Xizang, 121 sites in Qinghai, 43 sites in Xinjiang, 16 sites in Gansu and 11 sites
225 in Sichuan (Fig. 1). Southeast TP, where forest ecosystems are distributed, has few plant trait data.
226 However, field measurements are being conducted in the Hengduan Mountains to measure the leaf, twig
227 and root traits of dominant and common trees and shrubs. Other ecologists have worked on some parts
228 of this region to perform leaf and other trait studies (Luo et al., 2005; Shi et al., 2012; Vandvik et al.,
229 2020; Xu et al., 2021). The Hoh Xil dead zone in Central North to Northwest TP is logistically not
230 accessible during the plant growing season when the frozen ground is melting. Therefore, the plant trait
231 data have been not available ~~up to now~~ to date.

232 **4.2 Altitudinal range of sites**

233 The altitudinal range of the sampling sites was between 805–5343 m, in which 69.3% of the sites
234 were located in the high altitudes (> 3500 m), 18.5% of the sites were located in the Qaidam Basin and
235 the East Qinghai with lower altitudes (2500–3500 m) and 12.2% of the sites were located in the northern
236 margin of the TP with lowest altitudes (< 2500 m).

237 **4.3 Vegetation types of sites**

238 In accordance with the field records, the vegetation was divided into eight types, along with 108
239 sites in alpine meadow, 87 sites in alpine steppe, 61 sites in temperate desert, 38 sites in alpine shrubland,
240 16 sites in alpine desert, 15 sites in temperate meadow, ~~seven~~ 7 sites in temperate steppe and ~~four~~ 4 sites
241 in forest. In addition, the number of sites in the TP Alpine Vegetation Region was the most abundant
242 (63.1%), and its three subregions, namely, ~~the~~ the Middle TP Alpine Steppe Subregion (33.9%), the East
243 TP Alpine Scrub and Alpine Meadow Subregion (20.8%) and the Northwest TP Alpine Desert Subregion
244 (8.3%), followed by the Temperate Desert Region (29.4%) and other vegetation regions (7.5%), which
245 are Subtropical Evergreen Broad-leaved Forest Region (3.9%), Temperate Steppe Region (2.4%) and
246 Warm Temperate Deciduous Broad-leaved Forest Region (1.2%), as shown in Fig. 2.—

247 **5 Data description of species and traits**

248 5.1 plant species

249 A total of 1692 leaf samples were collected and measured in the TiP-Leaf dataset, including 468
250 species ~~that~~ belonging to 184 genera in 51 families (amongst them, 17 samples were identified as genera,
251 ~~six-6~~ samples were identified as families and ~~one-1~~ sample could not be identified). Some species were
252 ~~frequently~~ sampled frequently. For example, *Kobresia pygmaea* occurred 52 times, mainly in East and
253 South TP; *Stipa purpurea* occurred 47 times, mainly in Middle and West TP; and *Potentilla bifurca*
254 occurred 41 times, mainly in South, West and Northeast TP. However, in some sites, only one or two
255 species were sampled, especially in the Qaidam Basin and the northern margin of TP. The top five
256 families with the largest number of sampled species were as follows: Asteraceae (83 species and 24
257 genera), Poaceae (47 species and 18 genera), Fabaceae (46 species and 11 genera), Cyperaceae (29
258 species and four genera) and Rosaceae (28 species and 10 genera). Amongst the 468 species, 79 species,
259 including *Rhodiola smithii*, *Pomatosace filicula*, *Oxytropis sericopetala*, *Arenaria gerzensis*, *Onosma*
260 *waltonii*, *Delphinium qinghaiense*, *Metaeritrichium microuloides* and *Androsace cuttingii*, were unique
261 to the TP, ~~such as *Rhodiola smithii*, *Pomatosace filicula*, *Oxytropis sericopetala*, *Arenaria gerzensis*,~~
262 ~~*Onosma waltonii*, *Delphinium qinghaiense*, *Metaeritrichium microuloides* and *Androsace cuttingii*.~~
263 Furthermore, two (e.g. *Rosa rugosa* and *Rheum globulosum*) were endangered species (~~*Rosa rugosa* and~~
264 ~~*Rheum globulosum*~~), seven (e.g. *Arnebia guttata*, *Tamarix taklamakanensis*, *Rhodiola smithii*, *Juniperus*
265 *tibetica*, *Reaumuria kaschgarica*, *Rheum tanguticum* and *Metaeritrichium microuloides*) were vulnerable
266 species (~~*Arnebia guttata*, *Tamarix taklamakanensis*, *Rhodiola smithii*, *Juniperus tibetica*, *Reaumuria*~~
267 ~~*kaschgarica*, *Rheum tanguticum* and *Metaeritrichium microuloides*~~) and 10 (e.g. *Myricaria prostrata*,
268 *Euphorbia kozlovii*, *Hippophae tibetana*, *Phlomis pygmaea*, *Physochlaina praealta*, *Gentiana*
269 *siphonantha*, *Astragalus handelii*, *Androsace cuttingii*, *Carex nakaoana* and *Leiospora exscapa*) were
270 near-threatened species (~~*Myricaria prostrata*, *Euphorbia kozlovii*, *Hippophae tibetana*, *Phlomis*~~
271 ~~*pygmaea*, *Physochlaina praealta*, *Gentiana siphonantha*, *Astragalus handelii*, *Androsace cuttingii*,~~
272 ~~*Carex nakaoana* and *Leiospora exscapa*~~) in the TiP-Leaf dataset.

273 5.2 Leaf trait variations


274 The site-level leaf traits are shown in Table 1. The variation of each leaf trait was significant. In
275 particular, DW, FW and LA varied by more than 150%, followed by LT and LWC. The variations of
276 LMA, SLA, LDMC, LCC, LNC and LPC were slightly stable.-

Table 1 Summary of leaf functional traits in the TiP-Leaf dataset.

Traits	Mean \pm SD	Max	Min	CV (%)
LT (mm)	0.42 \pm 0.22	1.55	0.06	52.38
FW (g)	0.14 \pm 0.31	3.82	0.0001	221.43
DW (g)	0.04 \pm 0.07	0.62	0.00003	175.00
LDMC (g·g ⁻¹)	0.37 \pm 0.09	0.75	0.08	24.32
LWC (g·g ⁻¹)	2.41 \pm 1.25	11.11	0.33	51.87
LA (cm ²)	3.22 \pm 5.23	44.51	0.01	162.42
SLA (cm ² ·g ⁻¹)	142.10 \pm 46.67	333.85	33.16	32.84
LMA (g·m ⁻²)	91.49 \pm 34.28	308.11	17.22	37.36
LCC (mg·g ⁻¹)	386.54 \pm 43.46	487.42	212.57	11.24
LNC (mg·g ⁻¹)	23.08 \pm 4.75	45.83	7.18	20.58
LPC (mg·g ⁻¹)	1.62 \pm 0.51	3.76	0.43	31.48

5.3 Leaf trait relationships

279 The fitting of linear models of the site averages of the six leaf traits (Fig. 3) showed that LT was
280 significantly negatively correlated with LDMC ($r^2 = 0.1339$; $p < 0.05$; Fig. 3a), SLA ($r^2 = 0.0533$; $p <$
281 0.05 ; Fig. 3b) and LCC ($r^2 = 0.2755$; $p < 0.05$; Fig. 3c); with downward trends. No relationship was found
282 between LT and LNC ($r^2 = -0.0028$; $p > 0.05$; Fig. 3d) nor LPC ($r^2 = 0.0055$; $p > 0.05$; Fig. 3e). The
283 results also revealed that LDMC was significantly negatively correlated with SLA ($r^2 = 0.0630$; $p < 0.05$;
284 Fig. 3f), LNC ($r^2 = 0.0335$; $p < 0.05$; Fig. 3h) and LPC ($r^2 = 0.0649$; $p < 0.05$; Fig. 3i) and significantly
285 positively correlated with LCC ($r^2 = 0.0526$; $p < 0.05$; Fig. 3g). In addition, linearly inversed relationships
286 were observed between SLA and LCC ($r^2 = 0.0371$; $p < 0.05$; Fig. 3j); and LNC ($r^2 = 0.0566$; $p < 0.05$;
287 Fig. 3k) and LPC ($r^2 = 0.1776$; $p < 0.05$; Fig. 3l). The three leaf chemical traits were also related to one
288 another, and the relationship between LNC and LPC ($r^2 = 0.2481$; $p < 0.05$; Fig. 3o) was closer than that
289 between LNC and LCC ($r^2 = 0.0552$; $p < 0.05$; Fig. 3m) and between LPC and LCC ($r^2 = 0.0879$; $p <$
290 0.05 ; Fig. 3n).



291

292

293

294

Figure 3. Relationship between site-based average of key leaf traits in the TiP-Leaf dataset. The black dot indicates the mean leaf trait measurements of all species in the site, and the straight line represents the fitting of the linear model. R^2 - r^2 is the adjusted r^2 , and p represents the probability value of the regression model.

295

6 Data availability

296

The TiP-Leaf dataset includes-comprises three data sheets in Microsoft Excel format, namely: (a) a

297 data sheet named ‘variables’, ~~which describing describes the~~ header information of the geographical
 298 coordinates, climate and traits in the dataset (Table 2); (b) a data sheet (site information) ~~that reporting~~
 299 ~~reports~~ the site location and climate data; and (c) a data sheet (plant traits) of the complete trait data of
 300 each plant species in each sampling site. As studies based on the TiP-Leaf dataset are already underway,
 301 researchers interested in using such data previously are strongly recommended to contact the authors to
 302 avoid overlapping studies. The dataset will be available through the National Tibetan Plateau Data Center
 303 (TPDC; Jin et al., 2022a; <https://doi.org/10.11888/Terre.tpcd.272516>); and shall also be made available
 304 via the global TRY plant trait database (Kattge et al., 2011, 2020; www.try-db.org/).

305 **Table 2** Summary information found in TiP-Leaf dataset

Heading	Description	Type
Site	Site number based on sampling time	Code
Lat	Latitude (decimal degrees)	Numeric
Lon	Longitude (decimal degrees)	Numeric
Elev	Elevation (m)	Integer
Animal intensity	Animal activity intensity	Character
Human intensity	Human interference intensity	Character
Vegetation type	Vegetation type from field survey	Character
Vegetation region	Vegetation region from vegetation map	Character
MAT	Mean annual temperature (°C)	Numeric
MTCO	Mean temperature of the coldest month	Numeric
MTWA	Mean temperature of the warmest month	Numeric
GDD ₀	Annual growing degree days above 0 °C	Numeric
GDD ₅	Annual growing degree days above 5 °C	Numeric
MAP	Mean annual precipitation (mm)	Numeric
GP	Growing season precipitation (mm)	Numeric
MI	Moisture index	Numeric
DI	Drought index	Numeric
<u>Soil type</u>	<u>Soil type from Resource and Environment Science and Data Center</u> <u>(http://www.resdc.cn)</u>	<u>Numeric</u>

Species	Scientific name	Character
Family	Botanical family	Character
Growth form	Trees, shrubs, semi-shrubs and herbs	Character
Life form	Deciduous and evergreen; Annual and perennial	Character
LT	Leaf thickness (mm)	Numeric
FW	Fresh weight (g)	Numeric
DW	Dry weight (g)	Numeric
LDMC	Leaf dry-matter content ($\text{g}\cdot\text{g}^{-1}$)	Numeric
LWC	Leaf water content ($\text{g}\cdot\text{g}^{-1}$)	Numeric
LA	Leaf area (cm^2)	Numeric
SLA	Specific leaf area ($\text{cm}^2\cdot\text{g}^{-1}$)	Numeric
LMA	Leaf mass per area ($\text{g}\cdot\text{m}^{-2}$)	Numeric
LCC	Leaf carbon concentration ($\text{mg}\cdot\text{g}^{-1}$)	Numeric
LNC	Leaf nitrogen concentration ($\text{mg}\cdot\text{g}^{-1}$)	Numeric
LPC	Leaf phosphorus concentration ($\text{mg}\cdot\text{g}^{-1}$)	Numeric

306 7 Summary

307 The TiP-Leaf dataset was compiled from direct field measurements, covering a great proportion of
308 plant species and vegetation types on the highest plateau in the world. The dataset provides important
309 data foundation not only for quantitative analyses of modern alpine vegetation but also for the prediction
310 of future responses of alpine ecosystem to climate change and improvement of next-generation
311 vegetation models. It could also be used to promote the vegetation protection and restoration on the TP
312 and contribute to the global plant trait database. However, the dataset also presents some unavoidable
313 limitations. For example, the establishment of sampling sites and the judgment of dominant and common
314 species are mostly subjective. The leaves of some plants are extremely small, resulting in incomplete
315 recognition when scanning the LA. Due to harsh field conditions, measuring the plant traits in time
316 occasionally becomes impossible. Prevent some leaves from losing too much water to withering is still
317 inevitable, although we have taken protective measures for the leaves. Inadequate collection of some leaf
318 samples results in less data of plant chemical element content than that of morphological traits. In any

319 case, performing large-scale collection of plant traits on the TP, which requires a lot of manpower and
320 material resources, as well as overcoming the adverse environment of high altitude and extreme
321 variability, is not easy.

322 The dataset in this study provides more leaf trait measurements and covers more sampling sites,
323 which were located not only along the main roads but also the accessible ~~small~~-paths, than previous
324 studies (Luo et al., 2005; He et al., 2006; He et al., 2010; Geng et al., 2014; Wang et al., 2020; Xu et al.,
325 2021). This dataset is the first plant trait dataset that represents all of the alpine vegetations on the TP.
326 However, more collections of trait data are needed in remote areas with assessable difficulty, such as the
327 Hoh Xil dead zone in Northwest TP (alpine meadow, steppe and desert vegetation) and in the
328 mountainous areas of East and Southeast TP with less trait studies (subalpine and alpine forest and
329 shrubland vegetation). These works could enhance the representativeness of the whole TiP-Leaf trait
330 database in terms of geographical space and vegetation type. Given the complex topography of the
331 plateau, more sites are requested to be surveyed. ~~and~~ Given the flourishing of alpine flora, traits from
332 more plant species should be measured. At present, the TiP-Leaf dataset consists of leaf traits only. The
333 TiP-Root trait dataset is underway and the trait data of twig and branch of woody species ~~should will~~ be
334 ~~further~~ measured further.

335 **Author contributions.** JN conceived the study. KL, YJ and JN led the field works. YJ and other co-
336 authors collected leaf samples and measured plant traits. YJ, HW and JX processed the dataset, performed
337 the analyses and wrote the first draft. JN and YJ improved the manuscript. All authors approved the final
338 version of the submitted manuscript.

339 **Competing interests.** The (co-)authors declare that they have no conflict of interest.

340 **Disclaimer.** ~~Publisher's~~ Publisher's note: Copernicus Publications remains neutral with regard to
341 jurisdictional claims in published maps and institutional affiliations.

342 **Acknowledgements.** This work was funded by the Second Tibetan Plateau Scientific Expedition and
343 Research Program (STEP, 2019QZKK0402) and the Strategic Priority Research Program of the Chinese
344 Academy of Sciences (XDA2009000003). The authors sincerely thank Chenyu Li, Tudan Luosang, Yezi
345 Sheng, Pingyu Sun and Deyu Xu for their help in the field survey and Jun Li, Ang Liu, Rui Tang and
346 Xinxin Zhou for helping with specimen identification.

347 **Financial support.** This work was supported by the Second Tibetan Plateau Scientific Expedition and

348 Research Program (STEP, 2019QZKK0402) and the Strategic Priority Research Program of the Chinese
349 Academy of Sciences (XDA2009000003).

350 **References**

351 Angiosperm Phylogeny Group.: An update of the Angiosperm Phylogeny Group classification for the
352 orders and families of flowering plants: APG IV, *Bot. J. Linn. Soc.*, 181, 1-20,
353 <https://doi.org/10.1111/boj.12385>, 2016.

354 Berzaghi, F., Wright, I. J., Kramer, K., Oddou-Muratorio, S., Bohn, F. J., Reyer, C. P. O., Sabate, S.,
355 Sanders, T. G. M., and Hartig, F.: Towards a New Generation of Trait-Flexible Vegetation Models,
356 *Trends Ecol. Evol.*, 35, 191-205, <https://doi.org/10.1016/j.tree.2019.11.006>, 2020.

357 Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Thomas, H. J. D., Alatalo, J. M.,
358 Alexander, H., Anadon-Rosell, A., Angers-Blondin, S., Bai, Y., Baruah, G., te Beest, M., Berner, L.,
359 Bjork, R. G., Blok, D., Bruelheide, H., Buchwal, A., Buras, A., Carbognani, M., Christie, K., Collier,
360 L. S., Cooper, E. J., Cornelissen, J. H. C., Dickinson, K. J. M., Dullinger, S., Elberling, B., Eskelinen,
361 A., Forbes, B. C., Frei, E. R., Iturrate-Garcia, M., Good, M. K., Grau, O., Green, P., Greve, M.,
362 Grogan, P., Haider, S., Hajek, T., Hallinger, M., Happonen, K., Harper, K. A., Heijmans, M. M. P.
363 D., Henry, G. H. R., Hermanutz, L., Hewitt, R. E., Hollister, R. D., Hudson, J., Hulber, K., Iversen,
364 C. M., Jaroszynska, F., Jimenez-Alfaro, B., Johnstone, J., Jorgensen, R. H., Kaarlejarvi, E., Klady,
365 R., Klimesova, J., Korsten, A., Kuleza, S., Kulonen, A., Lamarque, L. J., Lantz, T., Lavallo, A.,
366 Lembrechts, J. J., Levesque, E., Little, C. J., Luoto, M., Macek, P., Mack, M. C., Mathakutha, R.,
367 Michelsen, A., Milbau, A., Molau, U., Morgan, J. W., Morsdorf, M. A., Nabe-Nielsen, J., Nielsen,
368 S. S., Ninot, J. M., Oberbauer, S. F., Olofsson, J., Onipchenko, V. G., Petraglia, A., Pickering, C.,
369 Prevey, J. S., Rixen, C., Rumpf, S. B., Schaepman-Strub, G., Semenchuk, P., Shetti, R.,
370 Soudzilovskaia, N. A., Spasojevic, M. J., Speed, J. D. M., Street, L. E., Suding, K., Tape, K. D.,
371 Tomaselli, M., Trant, A., Treier, U. A., Tremblay, J. P., Tremblay, M., Venn, S., Virkkala, A. M.,
372 Vowles, T., Weijers, S., Wilmking, M., Wipf, S., and Zamin, T.: Tundra Trait Team: A database of
373 plant traits spanning the tundra biome, *Global Ecol. Biogeogr.*, 27, 1402-1411,
374 <https://doi.org/10.1111/geb.12821>, 2018.

375 Butler, E. E., Datta, A., Flores-Moreno, H., Chen, M., Wythers, K. R., Fazayeli, F., Banerjee, A., Atkin,
376 O. K., Kattge, J., Amiaud, B., Blonder, B., Boenisch, G., Bond-Lamberty, B., Brown, K. A., Byun,

377 C., Campetella, G., Cerabolini, B. E. L., Cornelissen, J. H. C., Craine, J. M., Craven, D., de Vries,
378 F. T., Diaz, S., Domingues, T. F., Forey, E., Gonzalez-Melo*, A., Gross, N., Han, W. X., Hattingh,
379 W. N., Hickler, T., Jansen, S., Kramer, K., Kraft, N. J. B., Kurokawa, H., Laughlin, D. C., Meir, P.,
380 Minden, V., Niinemets, U., Onoda, Y., Penuelas, J., Read, Q., Sack, L., Schamp, B., Soudzilovskaia,
381 N. A., Spasojevic, M. J., Sosinski, E., Thornton, P. E., Valladares, F., van Bodegom, P. M., Williams,
382 M., Wirth, C., and Reich, P. B.: Mapping local and global variability in plant trait distributions,
383 PNAS, 114, E10937-E10946, <https://doi.org/10.1073/pnas.170898411>, 2018.

384 Chang, D. H. S.: The Tibetan Plateau in relation to the vegetation of China, *Ann. Mo. Bot. Gard.*, 70,
385 564-570, <https://doi.org/10.2307/2992087>, 1983.

386 Chave, J., Coomes, D., Jansen, S., Lewis, S. L., Swenson, N. G., and Zanne, A. E.: Towards a worldwide
387 wood economics spectrum, *Ecol. Lett.*, 12, 351-366, [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2009.01285.x)
388 [0248.2009.01285.x](https://doi.org/10.1111/j.1461-0248.2009.01285.x), 2009.

389 Chen, D. L., Xu, B. Q., Yao, T. D., Guo, Z. T., Cui, P., Chen, F. H., Zhang, R. H., Zhang, X. Z., Zhang,
390 Y. L., Fan, J., Hou, Z. Q., and Zhang, T. H.: Assessment of past, present and future environmental
391 changes on the Tibetan Plateau, *Chin. Sci. Bull.*, 60, 3025-3035, [https://doi.org/10.1360/N972014-](https://doi.org/10.1360/N972014-01370)
392 [01370](https://doi.org/10.1360/N972014-01370), 2015.

393 [Cheng, Q., Wu, X. Q., Wei, L. F., Hu, X. F., and Ni, J.: 30-year average monthly/1-km climate variables](https://doi.org/10.3974/geodb.2022.06.03.V1)
394 [dataset of China \(1951–1980, 1981–2010\), Digital J. Global Change Data Repository,](https://doi.org/10.3974/geodb.2022.06.03.V1)
395 <https://doi.org/10.3974/geodb.2022.06.03.V1>, 2022.

396 Comprehensive Scientific Investigation Team of Qinghai-Tibet Plateau, Chinese Academy of Sciences.:
397 Vegetation of Xizang, Science Press, Beijing, 1985-1987.

398 Cornelissen, J. H. C., Lavorel, S., Garnier, E., Díaz, S., Buchmann, N., Gurvich, D. E., Reich, P. B., ter
399 Steege, H., Morgan, H. D., van der Heijden, M. G. A., Pausas, J. G., and Poorter, H.: A handbook
400 of protocols for standardised and easy measurement of plant functional traits worldwide, *Aust. J.*
401 *Bot.*, 51, 335-380, <https://doi.org/10.1071/BT02124>, 2003.

402 Díaz, S., and Cabido, M.: Vive la différence: plant functional diversity matters to ecosystem processes,
403 *Trends Ecol. Evol.*, 16, 646-655, [https://doi.org/10.1016/S0169-5347\(01\)02283-2](https://doi.org/10.1016/S0169-5347(01)02283-2), 2001.

404 Díaz, S., Lavorel, S., McIntyre, S., Falczuk, V., Casanoves, F., Milchunas, D. G., Skarpe, C., Rusch, G.,
405 Sternberg, M., Noy-Meir, I., Landsberg, J., Zhang, W., Clark, H., and Campbell, B. D.: Plant trait

406 responses to grazing – a global synthesis, *Global Change Biol.*, 13, 313-341,
407 <https://doi.org/10.1111/j.1365-2486.2006.01288.x>, 2007.

408 Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth,
409 C., Prentice, I. C., Garnier, E., Bonisch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T.,
410 Dickie, J., Gillison, A. N., Zanne, A. E., Chave, J., Wright, S. J., ~~Sheremet'ev~~Sheremet'ev, S. N.,
411 Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J.
412 S., Gunther, A., Falczuk, V., Ruger, N., Mahecha, M. D., and Gorne, L. D.: The global spectrum of
413 plant form and function, *Nature*, 529, 167-171, <https://doi.org/10.1038/nature16489>, 2016.

414 Ding, W. N., Ree, R. H., Spicer, R. A., and Xing, Y. W.: Ancient orogenic and monsoon-driven assembly
415 of the ~~world's~~world's richest temperate alpine flora, *Science*, 369, 578-581,
416 <https://doi.org/10.1126/science.abb4484>, 2020.

417 Editorial Committee of the Flora of China.: *Flora of China*, Science Press, Beijing, 1959-2004.

418 Editorial Committee of the Flora of Gansu.: *Flora of Gansu*, Gansu Science and Technology Press,
419 Lanzhou, 2005.

420 Editorial Committee of the Flora of Qinghai.: *Flora of Qinghai*, Qinghai ~~People's~~People's Press, Xining,
421 1996-1999.

422 Editorial Committee of the Flora of Xinjiang.: *Flora of Xinjiang*, Xinjiang Science and Health Press,
423 Urumqi, 1993-1996.

424 Editorial Committee of Vegetation Map of China, Chinese Academy of Sciences.: *Vegetation of China*
425 *and its Geographical Pattern – Illustration of the Vegetation Map of the People's Republic of China*
426 (1:1 000 000), Geology Press, Beijing, 2007a.

427 Editorial Committee of Vegetation Map of China, Chinese Academy of Sciences.: *Vegetation Map of the*
428 *People's Republic of China (1:1 000 000)*, Geology Press, Beijing, 2007b.

429 Falster, D., Gallagher, R., Wenk, E. H., Wright, I. J., Indiarto, D., Andrew, S. C., Baxter, C., Lawson, J.,
430 Allen, S., Fuchs, A., Monro, A., Kar, F., Adams, M. A., Ahrens, C. W., Alfonzetti, M., Angevin, T.,
431 Apgaua, D. M. G., Arndt, S., Atkin, O. K., Atkinson, J., Auld, T., Baker, A., von Balthazar, M., Bean,
432 A., Blackman, C. J., Bloomfield, K., Bowman, D. M. J. S., Bragg, J., Brodribb, T. J., Buckton, G.,
433 Burrows, G., Caldwell, E., Camac, J., Carpenter, R., Catford, J. A, Cawthray, G. R., Cernusak, L.
434 A., Chandler, G., Chapman, A. R., Cheal, D., Cheesman, A. W., Chen, S. C., Choat, B., Clinton, B.,

435 Clode, P. L., Coleman, H., Cornwell, W. K., Cosgrove, M., Crisp, M., Cross, E., Crous, K. Y.,
436 Cunningham, S., Curran, T., Curtis, E., Daws, M. I., DeGabriel, J. L., Denton, M. D., Dong, N., Du,
437 P. Z., Duan, H. L., Duncan, D. H., Duncan, R. P., Duretto, M., Dwyer, J. M., Edwards, C., Esperon-
438 Rodriguez, M., Evans, J. R., Everingham, S. E., Farrell, C., Firn, J., Fonseca, C. R., French, B. J.,
439 Frood, D., Funk, J. L., Geange, S. R., Ghannoum, O., Gleason, S. M., Gosper, C. R., Gray, E.,
440 Groom, P. K., Grootemaat, S., Gross, C., Guerin, G., Guja, L., Hahs, A. K., Harrison, M. T., Hayes,
441 P. E., Henery, M., Hochuli, D., Howell, J., Huang, G., Hughes, L., Huisman, J., Ilic, J., Jagdish, A.,
442 Jin, D., Jordan, G., Jurado, E., Kanowski, J., Kasel, S., Kellermann, J., Kenny, B., Kohout, M.,
443 Kooyman, R. M., Kotowska, M. M., Lai, H. R., Laliberte, E., Lambers, H., Lamont, B. B., Lanfear,
444 R., van Langevelde, F., Laughlin, D. C., Laugier-kitchener, B. A., Laurance, S., Lehmann, C. E. R.,
445 Leigh, A., Leishman, M. R., Lenz, T., Lepschi, B., Lewis, J. D., Lim, F., Liu, U., Lord, J., Lusk, C.
446 H., Macinnis-Ng, C., McPherson, H., Magallon, S., Manea, A., Lopez-Martinez, A., Mayfield, M.,
447 McCarthy, J. K., Meers, T., van der Merwe, M., Metcalfe, D. J., Milberg, P., Mokany, K., Moles, A.
448 T., Moore, B. D., Moore, N., Morgan, J. W., Morris, W., Muir, A., Munroe, S., Nicholson, A., Nicolle,
449 D., Nicotra, A. B., Niinemets, U., North, T., O'Reilly-Nugent, A., ~~O'Sullivan~~O'Sullivan, O. S.,
450 Oberle, B., Onoda, Y., Ooi, M. K. J., Osborne, C. P., Paczkowska, G., Pekin, B., Pereira, C. G.,
451 Pickering, C., Pickup, M., Pollock, L. J., Poot, P., Powell, J. R., Power, S. A., Prentice, I. C., Prior,
452 L., Prober, S. M., Read, J., Reynolds, V., Richards, A. E., Richardson, B., Roderick, M. L., Rosell,
453 J. A., Rossetto, M., Rye, B., Rymer, P. D., Sams, M., Sanson, G., Sauquet, H., Schmidt, S.,
454 Schonenberger, J., Schulze, E. D., Sendall, K., Sinclair, S., Smith, B., Smith, R., Soper, F., Sparrow,
455 B., Standish, R. J., Staples, T. L., Stephens, R., Szota, C., Taseski, G., Tasker, E., Thomas, F., Tissue,
456 D. T., Tjoelker, M. G., Tng, D. Y. P., de Tombeur, F., Tomlinson, K., Turner, N. C., Veneklaas, E. J.,
457 Venn, S., Vesk, P., Vlasveld, C., Vorontsova, M. S., Warren, C. A., Warwick, N., Weerasinghe, L. K.,
458 Wells, J., Westoby, M., White, M., Williams, N. S. G., Wills, J., Wilson, P. G., Yates, C., Zanne, A.
459 E., Zemunik, G., and Zieminska, K.: AusTraits, a curated plant trait database for the Australian flora,
460 Sci. Data, 8, 254, <https://doi.org/10.1038/s41597-021-01006-6>, 2021.
461 [Fang, J. B., Pang, R. L., Guo, L. L., Xie, H. Z., Li, J., Luo, J., Yu, H., Liu, Y., and Wu, F. K.: Determination](#)
462 [of nitrogen, phosphorus and potassium in plants \(NY/T 2017–2011\), 2011.](#)
463 Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez,

464 E., Roth, L., Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner M., Oskin, M., Burbank, D., and
465 Alsdorf, D.: The shuttle radar topography mission, *Rev. Geophys.*, 45, RG2004,
466 <https://doi.org/10.1029/2005RG000183>, 2007.

467 Gallego-Sala, A. V., Clark, J. M., House, J. I., Orr, H. G., Prentice, I. C., Smith, P., Farewell, T., and
468 Chapman, S. J.: Bioclimatic envelope model of climate change impacts on blanket peatland
469 distribution in Great Britain, *Clim. Res.*, 45, 151-162, <https://doi.org/10.3354/cr00911>, 2010.

470 Geng, Y., Wang, L., Jin, D. M., Liu, H. Y., and He, J. S.: Alpine climate alters the relationships between
471 leaf and root morphological traits but not chemical traits, *Oecologia*, 175, 445-455,
472 <https://doi.org/10.1007/s00442-014-2919-5>, 2014.

473 Guerrero-Ramírez, N. R., Mommer, L., Freschet, G. T., Iversen, C. M., McCormack, M. L., Kattge, J.,
474 Poorter, H., van der Plas, F., Bergmann, J., Kuyper, T. W., York, L. M., Bruelheide, H., Laughlin, D.
475 C., Meier, I. C., Roumet, C., Semchenko, M., Sweeney, C. J., van Ruijven, J., Valverde-Barrantes,
476 O. J., Aubin, I., Catford, J. A., Manning, P., Martin, A., Milla, R., Minden, V., Pausas, J. G., Smith,
477 S. W., Soudzilovskaia, N. A., Ammer, C., Butterfield, B., Craine, J., Cornelissen, J. H. C., de Vries,
478 F. T., Isaac, M. E., Kramer, K., König, C., Lamb, E. G., Onipchenko, V. G., Penuelas, J., Reich, P.
479 B., Rillig, M. C., Sack, L., Shipley, B., Tedersoo, L., Valladares, F., van Bodegom, P., Weigelt, P.,
480 Wright, J. P., and Weigelt, A.: Global root traits (GRooT) database, *Global Ecol. Biogeogr.*, 30, 25-
481 37, <https://doi.org/10.1111/geb.13179>, 2021.

482 He, J. S., Wang, X. P., Schmid, B., Flynn, D. F. B., Li, X. F., Reich, P. B., and Fang, J. Y.: Taxonomic
483 identity, phylogeny, climate and soil fertility as driver of leaf traits across Chinese grassland biomes,
484 *J Plant Res*, 123, 551-561, <https://doi.org/10.1007/s10265-009-0294-9>, 2010.

485 He, J. S., Wang, Z. H., Wang, X. P., Schmid, B., Zuo, W. Y., Zhou, M., Zheng, C. Y., Wang, M. F., and
486 Fang, J. Y.: A test of the generality of leaf trait relationships on the Tibetan Plateau, *New Phytol.*,
487 170, 835-848, [https://doi.org/10.1111/j.1469-
488 8137.2006.01704.x](https://doi.org/10.1111/j.1469-8137.2006.01704.x)~~<https://doi.org/10.1111/j.14698137.2006.01704.x>~~, 2006.

489 He, N. P., Liu, C. C., Piao, S. L., Sack, L., Xu, L., Luo, Y. Q., He, J. S., Han, X. G., Zhou, G. S., Zhou,
490 X. H., Lin, Y., Yu, Q., Liu, S. R., Sun, W., Niu, S. L., Li, S. G., Zhang, J. H., and Yu, G. R.: Ecosystem
491 traits linking functional traits to macroecology, *Trends Ecol. Evol.*, 34, 200-210,
492 <https://doi.org/10.1016/j.tree.2018.11.004>, 2019.

493 He, N. P., Liu, Y., Liu, C. C., Xu, L., Li, M. X., Zhang, J. H., He, J. S., Tang, Z. Y., Han, X. G., Ye, Q.,
494 Xiao, C. W., Yu, Q., Liu, S. R., Sun, W., Niu, S. L., Li, S. G., Sack, L., and Yu, G. R.: Plant trait
495 networks: improved resolution of the dimensionality of adaptation, *Trends Ecol. Evol.*, 35, 908-918,
496 <https://doi.org/10.1016/j.tree.2020.06.003>, 2020.

497 Hutchinson, M. F., and Xu, T. B.: ANUSPLIN Version 4.4 User Guide, Canberra: Fenner School of
498 Environment and Society, the Australian National University, 2013.

499 Iversen, C. M., McCormack, M. L., Powell, A. S., Blackwood, C. B., Freschet, G. T., Kattge, J., Roumet,
500 C., Stover, D. B., Soudzilovskaia, N. A., Valverde-Barrantes, O. J., van Bodegom, P. M., and Violle,
501 C.: A global Fine-Root Ecology Database to address below-ground challenges in plant ecology, *New
502 Phytol.*, 215, 15-26, <https://doi.org/10.1111/nph.14486>, 2017.

503 Jin, Y. L., Wang, H. Y., Wei, L. F., Hu, J., Wu, K., Xia, H. J., Xia, J., Zhou, B. R., Li, K., and Ni, J.: A
504 plot dataset of plant community of Qingzang Plateau, *Chinese Journal of Plant EcologyEcol.*,
505 <https://www.plant-ecology.com/CN/10.17521/cjpe.2022.017446>, 846-854, <https://www.plant-ecology.com/CN/10.17521/cjpe.2022.0174>, 2022b.

507 Jin, Y., Wang, H., Xia, J., Ni, J., Li, K., Hou, Y., Hu, J., Wei, L., Xia, H., and Zhou, B.: A dataset of leaf
508 traits on the Tibetan Plateau (2018-2021), National Tibetan Plateau Data Center,
509 <https://doi.org/10.11888/Terre.tpd.272516>, 2022a.

510 Kattge, J., Diaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönsch, G., Garnier, E., Westoby, M., Reich,
511 P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C., Harrison, S. P., van Bodegom, P. M., Reichstein,
512 M., Enquist, B. J., Soudzilovskaia, N. A., Ackerly, D. D., Anand, M., Atkin, O., Bahn, M., Baker, T.
513 R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond, W. J., Bradstock, R., Bunker, D. E.,
514 Casanoves, F., Cavender-Bares, J., Chambers, J. Q., Chapin, F. S., Chave, J., Coomes, D., Cornwell,
515 W. K., Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan,
516 W. F., Fang, J., Fernandez-Mendez, F., Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D.,
517 Freschet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A., Gutierrez, A. G., Hickler, T., Higgins,
518 S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C. A., Kerkhoff, A. J., Kirkup, D., Kitajima, K.,
519 Kleyer, M., Klotz, S., Knops, J. M. H., Kramer, K., Kuhn, I., Kurokawa, H., Laughlin, D., Lee, T.
520 D., Leishman, M., Lens, F., Lenz, T., Lewis, S. L., Lloyd, J., Llusia, J., Louault, F., Ma, S., Mahecha,
521 M. D., Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, A. T., Muller, S. C., Nadrowski,

522 K., Naeem, S., Niinemets, U., Nollert, S., Nuske, A., Ogaya, R., Oleksyn, J., Onipchenko, V. G.,
523 Onoda, Y., Ordonez, J., Overbeck, G., Ozinga, W. A., Patino, S., Paula, S., Pausas, J. G., Penuelas,
524 J., Phillips, O. L., Pillar, V., Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R., Rammig,
525 A., Reinsch, S., Reu, B., Sack, L., Salgado-Negre, B., Sardans, J., Shiodera, S., Shipley, B., Siefert,
526 A., Sosinski, E., Soussana, J. F., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram,
527 M., Weiher, E., White, M., White, S., Wright, S. J., Yguel, B., Zaehle, S., Zanne, A. E., and Wirth,
528 C.: TRY – a global database of plant traits, *Global Change Biol.*, 17, 2905-2935,
529 <https://doi.org/10.1111/j.1365-2486.2011.02451.x>, 2011.

530 Kattge, J., Bonisch, G., Diaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner, G. D.
531 A., Aakala, T., Abedi, M., Acosta, A. T. R., Adamidis, G. C., Adamson, K., Aiba, M., Albert, C. H.,
532 Alcantara, J. M., Alcazar, C. C., Aleixo, I., Ali, H., Amiaud, B., Ammer, C., Amoroso, M. M., Anand,
533 M., Anderson, C., Anten, N., Antos, J., Apgaua, D. M. G., Ashman, T. L., Asmara, D. H., Asner, G.
534 P., Aspinwall, M., Atkin, O., Aubin, I., Baastrup-Spohr, L., Bahalkeh, K., Bahn, M., Baker, T., Baker,
535 W. J., Bakker, J. P., Baldocchi, D., Baltzer, J., Banerjee, A., Baranger, A., Barlow, J., Barneche, D.
536 R., Baruch, Z., Bastianelli, D., Battles, J., Bauerle, W., Bauters, M., Bazzato, E., Beckmann, M.,
537 Beckman, H., Beierkuhnlein, C., Bekker, R., Belfry, G., Belluau, M., Beloiu, M., Benavides, R.,
538 Benomar, L., Berdugo-Lattke, M. L., Berenguer, E., Bergamin, R., Bergmann, J., Carlucci, M. B.,
539 Berner, L., Bernhardt-Romermann, M., Bigler, C., Bjorkman, A. D., Blackman, C., Blanco, C.,
540 Blonder, B., Blumenthal, D., Bocanegra-Gonzalez, K. T., Boeckx, P., Bohlman, S., Bohning-Gaese,
541 K., Boisvert-Marsh, L., Bond, W., Bond-Lamberty, B., Boom, A., Boonman, C. C. F., Bordin, K.,
542 Boughton, E. H., Boukili, V., Bowman, D. M. J. S., Bravo, S., Brendel, M. R., Broadley, M. R.,
543 Brown, K. A., Bruelheide, H., Brunnich, F., Bruun, H. H., Bruy, D., Buchanan, S. W., Bucher, S.
544 F., Buchmann, N., Buitenwerf, R., Bunker, D. E., Burger, J., Burrascano, S., Burslem, D. F. R. P.,
545 Butterfield, B. J., Byun, C., Marques, M., Scalon, M. C., Caccianiga, M., Cadotte, M., Cailleret, M.,
546 Camac, J., Camarero, J. J., Company, C., Campetella, G., Campos, J. A., Cano-Arboleda, L., Canullo,
547 R., Carbognani, M., Carvalho, F., Casanoves, F., Castagneyrol, B., Catford, J. A., Cavender-Bares,
548 J., Cerabolini, B. E. L., Cervellini, M., Chacon-Madriral, E., Chapin, K., Chapin, F. S., Chelli, S.,
549 Chen, S. C., Chen, A. P., Cherubini, P., Chianucci, F., Choat, B., Chung, K. S., Chytry, M., Ciccarelli,
550 D., Coll, L., Collins, C. G., Conti, L., Coomes, D., Cornelissen, J. H. C., Cornwell, W. K., Corona,

551 P., Coyea, M., Craine, J., Craven, D., Cromsigt, J. P. G. M., Csecserits, A., Cufar, K., Cuntz, M., da
552 Silva, A. C., Dahlin, K. M., Dainese, M., Dalke, I., Dalle Fratte, M., Anh, T. D. L., Danihelka, J.,
553 Dannoura, M., Dawson, S., de Beer, A. J., De Frutos, A., De Long, J. R., Dechant, B., Delagrangé,
554 S., Delpierre, N., Derroire, G., Dias, A. S., Diaz-Toribio, M. H., Dimitrakopoulos, P. G.,
555 Dobrowolski, M., Doktor, D., Drevojan, P., Dong, N., Dransfield, J., Dressler, S., Duarte, L.,
556 Ducouret, E., Dullinger, S., Durka, W., Duursma, R., Dymova, O., E-Vojtko, A., Eckstein, R. L.,
557 Ejtehadi, H., Elser, J., Emilio, T., Engemann, K., Erfanian, M. B., Erfmeier, A., Esquivel-Muelbert,
558 A., Esser, G., Estiarte, M., Domingues, T. F., Fagan, W. F., Fagundez, J., Falster, D. S., Fan, Y., Fang,
559 J. Y., Farris, E., Fazlioglu, F., Feng, Y. H., Fernandez-Mendez, F., Ferrara, C., Ferreira, J., Fidelis,
560 A., Finegan, B., Firn, J., Flowers, T. J., Flynn, D. F. B., Fontana, V., Forey, E., Forgiarini, C.,
561 Francois, L., Frangipani, M., Frank, D., Frenette-Dussault, C., Freschet, G. T., Fry, E. L., Fyllas, N.
562 M., Mazzochini, G. G., Gachet, S., Gallagher, R., Ganade, G., Ganga, F., Garcia-Palacios, P.,
563 Gargaglione, V., Garnier, E., Garrido, J. L., de Gasper, A. L., Gea-Izquierdo, G., Gibson, D., Gillison,
564 A. N., Giroldo, A., Glasenhardt, M. C., Gleason, S., Gliesch, M., Goldberg, E., Goldel, B.,
565 Gonzalez-Akre, E., Gonzalez-Andujar, J. L., Gonzalez-Melo, A., Gonzalez-Robles, A., Graae, B. J.,
566 Granda, E., Graves, S., Green, W. A., Gregor, T., Gross, N., Guerin, G. R., Gunther, A., Gutierrez,
567 A. G., Haddock, L., Haines, A., Hall, J., Hambuckers, A., Han, W. X., Harrison, S. P., Hattingh, W.,
568 Hawes, J. E., He, T. H., He, P. C., Heberling, J. M., Helm, A., Hempel, S., Hentschel, J., Herault, B.,
569 Heres, A. M., Herz, K., Heuertz, M., Hickler, T., Hietz, P., Higuchi, P., Hipp, A. L., Hiron, A., Hock,
570 M., Hogan, J. A., Holl, K., Honnay, O., Hornstein, D., Hou, E. Q., Hough-Snee, N., Hovstad, K. A.,
571 Ichie, T., Igic, B., Illa, E., Isaac, M., Ishihara, M., Ivanov, L., Ivanova, L., Iversen, C. M., Izquierdo,
572 J., Jackson, R. B., Jackson, B., Jactel, H., Jagodzinski, A. M., Jandt, U., Jansen, S., Jenkins, T.,
573 Jentsch, A., Jespersen, J. R. P., Jiang, G. F., Johansen, J. L., Johnson, D., Jokela, E. J., Joly, C. A.,
574 Jordan, G. J., Joseph, G. S., Junaedi, D., Junker, R. R., Justes, E., Kabzems, R., Kane, J., Kaplan,
575 Z., Kattenborn, T., Kavelenova, L., Kearsley, E., Kempel, A., Kenzo, T., Kerkhoff, A., Khalil, M. I.,
576 Kinlock, N. L., Kissling, W. D., Kitajima, K., Kitzberger, T., Kjoller, R., Klein, T., Kleyer, M.,
577 Klimesova, J., Klipel, J., Kloeppel, B., Klotz, S., Knops, J. M. H., Kohyama, T., Koike, F., Kollmann,
578 J., Komac, B., Komatsu, K., Konig, C., Kraft, N. J. B., Kramer, K., Kreft, H., Kuhn, I.,
579 Kumarathunge, D., Kuppler, J., Kurokawa, H., Kurosawa, Y., Kuyah, S., Laclau, J. P., Lafleur, B.,

580 Lallai, E., Lamb, E., Lamprecht, A., Larkin, D. J., Laughlin, D., Le Bagousse-Pinguet, Y., le Maire,
581 G., le Roux, P. C., le Roux, E., Lee, T., Lens, F., Lewis, S. L., Lhotsky, B., Li, Y. Z., Li, X. E.,
582 Lichstein, J. W., Liebergesell, M., Lim, J. Y., Lin, Y. S., Linares, J. C., Liu, C. J., Liu, D. J., Liu, U.,
583 Livingstone, S., Llusia, J., Lohbeck, M., Lopez-Garcia, A., Lopez-Gonzalez, G., Lososova, Z.,
584 Louault, F., Lukacs, B. A., Lukes, P., Luo, Y. J., Lussu, M., Ma, S. Y., Pereira, C. M. R., Mack, M.,
585 Maire, V., Makela, A., Makinen, H., Malhado, A. C. M., Mallik, A., Manning, P., Manzoni, S.,
586 Marchetti, Z., Marchino, L., Marcilio-Silva, V., Marcon, E., Marignani, M., Markesteijn, L., Martin,
587 A., Martinez-Garza, C., Martinez-Vilalta, J., Maskova, T., Mason, K., Mason, N., Massad, T. J.,
588 Masse, J., Mayrose, I., McCarthy, J., McCormack, M. L., McCulloh, K., McFadden, I. R., McGill,
589 B. J., McPartland, M. Y., Medeiros, J. S., Medlyn, B., Meerts, P., Mehrabi, Z., Meir, P., Melo, F. P.
590 L., Mencuccini, M., Meredieu, C., Messier, J., Meszaros, I., Metsaranta, J., Michaletz, S. T.,
591 Michelaki, C., Migalina, S., Milla, R., Miller, J. E. D., Minden, V., Ming, R., Mokany, K., Moles,
592 A. T., Molnar, V. A., Molofsky, J., Molz, M., Montgomery, R. A., Monty, A., Moravcova, L.,
593 Moreno-Martinez, A., Moretti, M., Mori, A. S., Mori, S., Morris, D., Morrison, J., Mucina, L.,
594 Mueller, S., Muir, C. D., Muller, S. C., Munoz, F., Myers-Smith, I. H., Myster, R. W., Nagano, M.,
595 Naidu, S., Narayanan, A., Natesan, B., Negoita, L., Nelson, A. S., Neuschulz, E. L., Ni, J., Niedrist,
596 G., Nieto, J., Niinemets, U., Nolan, R., Nottebrock, H., Nouvellon, Y., Novakovskiy, A., Nystuen, K.
597 O., ~~O'Grady~~O'Grady, A., ~~O'Hara~~O'Hara, K., ~~O'Reilly~~O'Reilly-Nugent, A., Oakley, S., Oberhuber,
598 W., Ohtsuka, T., Oliveira, R., Ollerer, K., Olson, M. E., Onipchenko, V., Onoda, Y., Onstein, R. E.,
599 Ordonez, J. C., Osada, N., Ostonen, I., Ottaviani, G., Otto, S., Overbeck, G. E., Ozinga, W. A., Pahl,
600 A. T., Paine, C. E. T., Pakeman, R. J., Papageorgiou, A. C., Parfionova, E., Partel, M., Patacca, M.,
601 Paula, S., Paule, J., Pauli, H., Pausas, J. G., Peco, B., Penuelas, J., Perea, A., Peri, P. L., Petisco-
602 Souza, A. C., Petraglia, A., Petritan, A. M., Phillips, O. L., Pierce, S., Pillar, V. D., Pisek, J.,
603 Pomogaybin, A., Poorter, H., Portsmouth, A., Poschlod, P., Potvin, C., Pounds, D., Powell, A. S.,
604 Power, S. A., Prinzing, A., Puglielli, G., Pysek, P., Raavel, V., Rammig, A., Ransijn, J., Ray, C. A.,
605 Reich, P. B., Reichstein, M., Reid, D. E. B., Rejou-Mechain, M., de Dios, V. R., Ribeiro, S.,
606 Richardson, S., Riibak, K., Rillig, M. C., Riviera, F., Robert, E. M. R., Roberts, S., Robroek, B.,
607 Roddy, A., Rodrigues, A. V., Rogers, A., Rollinson, E., Rolo, V., Romermann, C., Ronzhina, D.,
608 Roscher, C., Rosell, J. A., Rosenfield, M. F., Rossi, C., Roy, D. B., Royer-Tardif, S., Ruger, N.,

609 Ruiz-Peinado, R., Rumpf, S. B., Rusch, G. M., Ryo, M., Sack, L., Saldana, A., Salgado-Negret, B.,
610 Salguero-Gomez, R., Santa-Regina, I., Santacruz-Garcia, A. C., Santos, J., Sardans, J., Schamp, B.,
611 Scherer-Lorenzen, M., Schleuning, M., Schmid, B., Schmidt, M., Schmitt, S., Schneider, J. V.,
612 Schowanek, S. D., Schrader, J., Schrodt, F., Schuldt, B., Schurr, F., Garvizu, G. S., Semchenko, M.,
613 Seymour, C., Sfair, J. C., Sharpe, J. M., Sheppard, C. S., Sheremetiev, S., Shiodera, S., Shipley, B.,
614 Shovon, T. A., Siebenkas, A., Sierra, C., Silva, V., Silva, M., Sitzia, T., Sjoman, H., Slot, M., Smith,
615 N. G., Sodhi, D., Soltis, P., Soltis, D., Somers, B., Sonnier, G., Sorensen, M. V., Sosinski, E. E.,
616 Soudzilovskaia, N. A., Souza, A. F., Spasojevic, M., Sperandii, M. G., Stan, A. B., Stegen, J.,
617 Steinbauer, K., Stephan, J. G., Sterck, F., Stojanovic, D. B., Strydom, T., Suarez, M. L., Svenning,
618 J. C., Svitkova, I., Svitok, M., Svoboda, M., Swaine, E., Swenson, N., Tabarelli, M., Takagi, K.,
619 Tappeiner, U., Tarifa, R., Tauougourdeau, S., Tavsanoglu, C., te Beest, M., Tedersoo, L., Thiffault,
620 N., Thom, D., Thomas, E., Thompson, K., Thornton, P. E., Thuiller, W., Tichy, L., Tissue, D.,
621 Tjoelker, M. G., Tng, D. Y. P., Tobias, J., Torok, P., Tarin, T., Torres-Ruiz, J. M., Tothmeresz, B.,
622 Treurnicht, M., Trivellone, V., Trolliet, F., Trotsiuk, V., Tsakalos, J. L., Tsiripidis, I., Tyskland, N.,
623 Umehara, T., Usoltsev, V., Vadeboncoeur, M., Vaezi, J., Valladares, F., Vamosi, J., van Bodegom, P.
624 M., van Breugel, M., Van Cleemput, E., van de Weg, M., van der Merwe, S., van der Plas, F., van
625 der Sande, M. T., van Kleunen, M., Van Meerbeek, K., Vanderwel, M., Vanselow, K. A., Varhammar,
626 A., Varone, L., Valderrama, M. Y., Vassilev, K., Vellend, M., Veneklaas, E. J., Verbeeck, H.,
627 Verheyen, K., Vibrans, A., Vieira, I., Villacis, J., Violle, C., Vivek, P., Wagner, K., Waldram, M.,
628 Waldron, A., Walker, A. P., Waller, M., Walther, G., Wang, H., Wang, F., Wang, W. Q., Watkins, H.,
629 Watkins, J., Weber, U., Weedon, J. T., Wei, L. P., Weigelt, P., Weiher, E., Wells, A. W., Wellstein, C.,
630 Wenk, E., Westoby, M., Westwood, A., White, P. J., Whitten, M., Williams, M., Winkler, D. E.,
631 Winter, K., Womack, C., Wright, I. J., Wright, S. J., Wright, J., Pinho, B. X., Ximenes, F., Yamada,
632 T., Yamaji, K., Yanai, R., Yankov, N., Yguel, B., Zanini, K. J., Zanne, A. E., Zeleny, D., Zhao, Y. P.,
633 Zheng, J. M., Zheng, J., Zieminska, K., Zirbel, C. R., Zizka, G., Zo-Bi, I. C., Zotz, G., and Wirth,
634 C.: TRY plant trait database – enhanced coverage and open access, *Global Change Biol.*, 26, 119-
635 188, <https://doi.org/10.1111/gcb.14904>, 2020.

636 Liu, Y. X.: *Flora of Desert Plants of China*, Science Press, Beijing, 1985.

637 Lu, L. M., Mao, L. F., Yang, T., Ye, J. F., Liu, B., Li, H. L., Sun, M., Miller, J. T., Mathews, S., Hu, H.

638 H., Niu, Y. T., Peng, D. X., Chen, Y. H., Smith, S. A., Chen, M., Xiang, K. L., Le, C. T., Dang, V.
639 C., Lu, A. M., Soltis, P. S., Soltis, D. E., Li, J. H., and Chen, Z. D.: Evolutionary history of the
640 angiosperm flora of China, *Nature*, 554, 234-238, <https://doi.org/10.1038/nature25485>, 2018.

641 Luo, T. X., Luo, J., and Pan, Y. D.: Leaf traits and associated ecosystem characteristics across subtropical
642 and timberline forests in the Gongga Mountains, Eastern Tibetan Plateau, *Oecologia*, 142, 261-273,
643 <https://doi.org/10.1007/s00442-004-1729-6>, 2005.

644 Ma, Z. Q., Guo, D. L., Xu, X. L., Lu, M. Z., Bardgett, R. D., Eissenstat, D. M., McCormack, M. L., and
645 Hedin, L. O.: Evolutionary history resolves global organization of root functional traits, *Nature*, 555,
646 94-97, <https://doi.org/10.1038/nature26163>, 2018.

647 Maes, S. L., Perring, M. P., Depauw, L., Bernhardt-Romermann, M., Blondeel, H., Brumelis, G., Brunet,
648 J., Decocq, G., den Ouden, J., Govaert, S., Hardtle, W., Hedl, R., Heinken, T., Heinrichs, S., Hertzog,
649 L., Jaroszewicz, B., Kirby, K., Kopecky, M., Landuyt, D., Malis, F., Vanneste, T., Wulf, M., and
650 Verheyen, K.: Plant functional trait response to environmental drivers across European temperate
651 forest understorey communities, *Plant Biol.*, 22, 410-424,
652 <https://doi.org/10.1111/plb.13082>~~<https://doi.org/10.1038/nature26163>~~, 2020.

653 Mariano, E., Gomes, T. F., Lins, S. R. M., Abdalla- Filho, A. L., Soltangheisi, A., Araújo, M. G. S.,
654 Almeida, R. F., Augusto, F. G., Canisares, L. P., Chaves, S. S. F., Costa, C. F. G., Diniz-Reis, T. R.,
655 Galera, L. A., Martinez, M. G., Morais, M. C., Perez, E. B., Reis, L. C., Simon, C. D., Mardegan, S.
656 F., Domingues, T. F., Miatto, R. C., Oliveira, R. S., Reis, C. R. G., Nardoto, G. B., Kattge, J., and
657 Martinelli, L. A.: LT-Brazil: A database of leaf traits across biomes and vegetation types in Brazil,
658 *Global Ecol. Biogeogr.*, 30, 2136-2146, <https://doi.org/10.1111/geb.13381>, 2021.

659 Meng, T. T., Ni, J., and Harrison, S. P.: Plant morphometric traits and climate gradients in northern China:
660 a meta-analysis using quadrat and flora data, *Ann. Bot.*, 104, 1217-1229,
661 <https://doi.org/10.1093/aob/mcp230>, 2009.

662 Meng, T. T., Wang, H., Harrison, S. P., Prentice, I. C., Ni, J., and Wang, G.: Responses of leaf traits to
663 climatic gradients: adaptive variation versus compositional shifts, *Biogeosciences*, 12, 5339-5352,
664 <https://doi.org/10.5194/bg-12-5339-2015>, 2015.

665 Moles, A. T., Ackerly, D. D., Tweddle, J. C., Dickie, J. B., Smith, R., Leishman, M. R., Mayfield, M. M.,
666 Pitman, A., Wood, J. T., and Westoby, M.: Global patterns in seed size, *Global Ecol. Biogeogr.*, 16,
667 109-116, <https://doi.org/10.1111/j.1466-8238.2006.00259.x>, 2007.

668 Moles, A. T., Warton, D. I., Warman, L., Swenson, N. G., Laffan, S. W., Zanne, A. E., Pitman, A.,
669 Hemmings, F. A., and Leishman, M. R.: Global patterns in plant height, *J. Ecol.*, 97, 923-932,
670 <https://doi.org/10.1111/j.1365-2745.2009.01526.x>, 2009.

671 Myers-Smith, I. H., Thomas, H. J. D., and Bjorkman, A. D.: Plant traits inform predictions of tundra
672 responses to global change, *New Phytol.*, 221, 1742-1748, <https://doi.org/10.1111/nph.15592>, 2019.

673 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M.
674 S., Cornwell, W. K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter,
675 L., Wright, I. J., Ray, P., Enrico, L., Pausas, J. G., de Vos, A. C., Buchmann, N., Funes, G.,
676 ~~Quétier~~Quétier, F., Hodgson, J. G., Thompson, K., Morgan, H. D., ter Steege, H., van der Heijden,
677 M. G. A., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A. C., Aquino, S.,
678 and Cornelissen, J. H. C.: New handbook for standardised measurement of plant functional traits
679 worldwide, *Aust. J. Bot.*, 61, 167-234, <https://doi.org/10.1071/BT12225>, 2013.

680 Piao, S. L., Zhang, X. Z., Wang, T., Liang, E. Y., Wang, S. P., Zhu, J. T., and Niu, Ben.: Responses and
681 feedback of the Tibetan Plateau's alpine ecosystem to climate change, *Chin. Sci. Bull.*, 64, 2842-
682 2855, <https://doi.org/10.1360/TB-2019-0074>, 2019.

683 Reich, P. B., and Oleksyn, J.: Global patterns of plant leaf N and P in relation to temperature and latitude,
684 *PNAS*, 101, 11001-11006, <https://doi.org/10.1073/pnas.040358810>, 2004.

685 Reichstein, M., Bahn, M., Mahecha, M. D., Kattge, J., and Baldocchi, D. D.: Linking plant and ecosystem
686 functional biogeography, *PNAS*, 111, 13697-13702, <https://doi.org/10.1073/pnas.121606511>, 2014.

687 Roddy, A. B., ~~Martínez~~Martínez-Perez, C., Teixido, A. L., Cornelissen, T. G., Olson, M. E., Oliveira, R.
688 S., and Silveira, F. A. O.: Towards the flower economics spectrum, *New Phytol.*, 229, 665-672,
689 <https://doi.org/10.1111/nph.16823>, 2021.

690 Shi, W. Q., Wang, G. A., and Han, W. X.: Altitudinal variation in leaf nitrogen concentration on the
691 eastern slope of Mount Gongga on the Tibetan Plateau, China, *Plos One*, 7, e44628,
692 <https://doi.org/10.1371/journal.pone.0044628>, 2012.

693 Sloan, S., Jenkins, C. N., Joppa, L. N., Gaveau, D. L. A., and Laurance, W. F.: Remaining natural

694 vegetation in the global biodiversity hotspots, *Biol. Conserv.*, 177, 12-24,
695 <https://doi.org/10.1016/j.biocon.2014.05.027>, 2014.

696 Taseski, G. ~~M~~, Beloe, C. J., Gallagher, R. V., Chan, J. Y., Dalrymple, R. L., and Cornwell, W. K.: A
697 global-growth form database for 143,616 vascular plant species, *Ecology*, 100, e02614,
698 <https://doi.org/10.1002/ecy.2614>, 2019.

699 Tavşanoğlu, Ç ~~ağatay~~, and Pausas, J. G.: A functional trait database for Mediterranean Basin plants, *Sci.*
700 *Data*, 5, 180135, <https://doi.org/10.1038/sdata.2018.135>, 2018.

701 van Bodegom, P. M., Douma, J. C., and Verheijen, L. M.: A fully traits-based approach to modeling
702 global vegetation distribution, *PNAS*, 111, 13733-13738, <https://doi.org/10.1073/pnas.1304551110>,
703 2014.

704 Vandvik, V., Halbritter, A. H., Yang, Y., He, H., Zhang, L., Brummer, A. B., Klanderud, K., Maitner, B.
705 S., Michaletz, S. T., Sun, X. Y., Telford, R. J., Wang, G. X., Althuizen, I. H. J., Henn, J. J., Garcia,
706 W. F. E., Gya, R., Jaroszynska, F., Joyce, B. L., Lehman, R., Moerland, M. S., Nesheim-Hauge, E.,
707 ~~Nordås~~ ~~Nordas~~, L. H., Peng, A., Ponsac, C., Seltzer, L., Steyn, C., Sullivan, M. K., Tjendra, J., Xiao,
708 Y., Zhao, X. X., and Enquist, B. J.: Plant traits and vegetation data from climate warming
709 experiments along an 1100 m elevation gradient in Gongga Mountains, China, *Sci. Data*, 7, 189,
710 <https://doi.org/10.1038/s41597-020-0529-0>, 2020.

711 Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier, E.: Let the
712 concept of trait be functional!, *Oikos*, 116, 882-892, <https://doi.org/10.1111/j.0030-1299.2007.15559.x>, 2007.

714 Wang, C. S., Lyu, W. W., Jiang, L. L., Wang, S. P., Wang, Q., Meng, F. D., and Zhang, L. R.: Changes in
715 leaf vein traits among vein types of alpine grassland plants on the Tibetan Plateau, *J. Mountain-Mt.*
716 *Sci.*, 17, 2161-2169, <https://doi.org/10.1007/s11629-020-6069-4>, 2020.

717 Wang, H., Harrison, S. P., Prentice, I. C., Yang, Y. Z., Bai, F., Togashi, H. F., Wang, M., Zhou, S. X., and
718 Ni, J.: The China Plant Trait Database: toward a comprehensive regional compilation of functional
719 traits for land plants, *Ecology*, 99, 500, <https://doi.org/10.1002/ecy.2091>, 2018.

720 Wang, H., Prentice, I. C., Keenan, T. F., Davis, T. W., Wright, I. J., Cornwell, W. K., Evans, B. J., and
721 Peng, C. H.: Towards a universal model for carbon dioxide uptake by plants, *Nat. Plants*, 3, 734-
722 741, <https://doi.org/10.1038/s41477-017-0006-8>, 2017.

723 Wang, H., Wang, R. X., Harrison, S. P., and Prentice, I. C.: Leaf morphological traits as adaptations to
724 multiple climate gradients, *J. Ecol.*, [110, 1344-1355](https://doi.org/10.1111/1365-2745.13873), <https://doi.org/10.1111/1365-2745.13873>,
725 2022.

726 Wang, Q., and Hong, D. Y.: Understanding the plant diversity on the roof of the world, *The Innovation*,
727 3, 100215, <https://doi.org/10.1016/j.xinn.2022.100215>, 2022.

728 Wei, L. F., Hu, X. F., Cheng, Q., Wu, X. Q., and Ni, J.: A dataset of spatial distribution of bioclimatic
729 variables in China at 1 km resolution, *China Scientific-Sc. Data*, [https://doi.org/10.11922/11-
730 6035.csd.2022.0003.zh](https://doi.org/10.11922/11-6035.csd.2022.0003.zh), 2022.

731 Weigelt, P., König, C., and Kreft, H.: GIFT – A Global Inventory of Floras and Traits for macroecology
732 and biogeography, *J. Biogeogr.*, 47, 16-43, <https://doi.org/10.1111/jbi.13623>, 2019.

733 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J.,
734 Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J.,
735 Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C., Midgley, J. J., Navas, ~~Marie~~ Laure.,
736 Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V. I., Roumet, C.,
737 Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The worldwide leaf economics
738 spectrum, *Nature*, 428, 821-827, <https://doi.org/10.1038/nature02403>, 2004.

739 [Wu, D. X.: Protocols for Standard Biological Observation and Measurement in Terrestrial Ecosystems,](#)
740 [China Environmental Science Press, Beijing, 2007.](#)

741 Wullschleger, S. D., Epstein, H. E., Box, E. O., Euskirchen, E. S., Goswami, S., Iversen, C. M., Kattge,
742 J., Norby, R. J., van Bodegom, P. M., and Xu, X. F.: Plant functional types in Earth system models:
743 past experiences and future directions for application of dynamic vegetation models in high-latitude
744 ecosystems, *Ann. Bot.*, 114, 1-16, <https://doi.org/10.1093/aob/mcu077>, 2014.

745 Xu, H. Y., Wang, H., Prentice, I. C., Harrison, S. P., Wang, G. X., and Sun, X. Y.: Predictability of leaf
746 traits with climate and elevation: a case study in Gongga Mountain, China, *Tree Physiol.*, 41, 1336-
747 1352, <https://doi.org/10.1093/treephys/tpab003>, 2021.

748 Xu, T. B., and Hutchinson, M. F.: New developments and applications in the ANUCLIM spatial climatic
749 and bioclimatic modelling package, *Environ. Modell. Software*, 40, 267-279,
750 <https://doi.org/10.1016/j.envsoft.2012.10.003>, 2013.

751 Yan, Y. J., Yang, X., and Tang, Z. Y.: Patterns of species diversity and phylogenetic structure of vascular

752 plants on the Qinghai-Tibetan Plateau, *Ecol. Evol.*, 3, 4584-4595, <https://doi.org/10.1002/ece3.847>,
753 2013.

754 Yao, T. D., Thompson, L., Yang, W., Yu, W. S., Gao, Y., Guo, X. J., Yang, X. X., Duan, K. Q., Zhao, H.
755 B., Xu, B. Q., Pu, J. C., Lu, A. X., Xiang, Y., Kattel, D. B., and Joswiak, D.: Different glacier status
756 with atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Change*, 2, 663-667,
757 <https://doi.org/10.1038/nclimate1580>, 2012.

758 Zhang, X. Z., Yang, Y. P., Piao, S. L., Bao, W. K., Wang, S. P., Wang, G. X., Sun, H., Luo, T. X., Zhang,
759 Y. J., Shi, P. L., Liang, E. Y., Shen, M. G., Wang, J. S., Gao, Q. Z., Zhang, Y. L., and Ouyang, H.:
760 Ecological change on the Tibetan Plateau, *Chin. Sci. Bull.*, 60, 3048-3056,
761 <https://doi.org/10.1360/N972014-01339>, 2015.