

A new 2010 permafrost distribution map over the Qinghai-Tibet Plateau based on subregion survey maps: a benchmark for regional permafrost modeling

Zetao Cao¹, Zhuotong Nan^{1,2,*}, Jianan Hu¹, Yuhong Chen¹, Yaonan Zhang³

5 ¹Key Laboratory of Ministry of Education on Virtual Geographic Environment, Nanjing Normal University, Nanjing 210023, China;

²Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing 210023, China;

³National Cryosphere Desert Data Center, Lanzhou 730000, China

10

Correspondence to: Zhuotong Nan (nanzt@njnu.edu.cn)

Abstract. Permafrost over the Qinghai-Tibet Plateau (QTP) has received increasing attention due to its high sensitivity to climate change. Numerous spatial modeling studies have been conducted on the QTP to assess the status of permafrost, project future changes in permafrost, and diagnose contributors to permafrost degradation. Due to the scarcity of ground stations on the QTP, these modeling studies are often hampered by the lack of validation references, calibration targets, and model constraints, while a high-quality permafrost distribution map can be a good option as a benchmark for spatial simulations. Existing permafrost distribution maps on the QTP can poorly serve this purpose. An ideal benchmark map for spatial modeling should be methodologically sound, of sufficient accuracy, and based on observations from mapping years rather than all historical data spanning several decades. Therefore, in this study, we created a new permafrost distribution map over the QTP in 2010 with a novel permafrost mapping approach with satellite-derived ground surface thawing/freezing indices as inputs and survey-based subregion permafrost maps as constraints. This approach accounted for the effects of local factors by incorporating into the model an empirical soil parameter whose values were optimally estimated through spatial clustering and parameter optimization constrained by survey-based subregion permafrost maps, and was improved in this study to reduce parametric equifinality. This new map showed a total permafrost area of about 1.086×10^6 km² (41.2% of the QTP area) and seasonally frozen ground of about 1.447×10^6 km² (54.9%) in 2010, excluding glaciers and lakes. Validations using survey-based subregion permafrost maps (Kappa = 0.74) and borehole records (Overall Accuracy = 0.85 and Kappa = 0.43) showed higher accuracy of this map than two recent maps. Inspection of regions with obvious distinctions between the maps affirms that the permafrost distribution on this map is more realistic than that on the Zou et al. (2017) map. Given the demonstrated excellent accuracy, this map can serve as a benchmark map for constraining/validating land surface simulations on the QTP and as a historical reference for projecting future permafrost changes on the QTP in the context of global warming.

15
20
25
30

1 Introduction

Permafrost, defined as ground that remains at or below 0 °C for at least two consecutive years (Dobinski, 2011), underlies more than 20% of the exposed land area in the Northern Hemisphere (Obu et al., 2019) and constitutes an essential component of the Earth system. The Qinghai-Tibet Plateau (QTP), also known as the Earth's third pole, contains the largest mid- to low-latitude permafrost area in the world. Due to the complex topography and unique plateau climate, permafrost over the QTP is generally of low thermal stability and strongly influenced by complex local factors, making it different from high-latitude permafrost around the Arctic and more sensitive to global climate change (Li et al., 2008; Yang et al., 2019; Zhao et al., 2020).

In the context of global warming, significant permafrost degradation is occurring on the QTP and has strongly affected hydrological processes (Li et al., 2020), carbon cycling (Mu et al., 2020), and heat exchange processes (Zhao et al., 2020). In addition, hazards related to permafrost degradation threaten constructions and infrastructures on the QTP (Wang et al., 2020). Many researches have studied the complex responses and feedbacks of permafrost to climate change (Yang et al., 2019), while land surface model-based spatial modeling of permafrost has become an important approach (Ji et al., 2022). Using land surface models, many spatial modeling studies have attempted to project future changes of permafrost (Chang et al., 2018; Debolskiy et al., 2020; Yin et al., 2021), assess permafrost status under climate change (Koven et al., 2013; Burke et al., 2020), diagnose the contributors to regional permafrost degradation (Zhang et al., 2021; Zhang et al., 2021; Mekonnen et al., 2021), and project potential feedbacks on the climate system due to permafrost degradation (Zhang et al., 2020; Andresen et al., 2020; Yokohata et al., 2020; Wang et al., 2021). But evaluating the spatial simulations has often been challenging because ground observations on the QTP are often too sparse to provide references compared to the large spatial domain of the spatial modeling. Hence, there is a need for an accurate permafrost distribution map that would serve as a reference to validate the results of spatial simulations. The map could be used as a target for the calibration of model parameters and to provide a constraint for future projections to minimize biases resulting from the modeling process. Moreover, an accurate map of permafrost distribution could serve as a fundamental dataset for hydrological, carbon, ecological and engineering studies in cold regions (Hu et al., 2019; Li et al., 2020; Song et al., 2020; Mu et al., 2020).

Though many permafrost distribution maps have been compiled over the QTP (Cheng et al., 2011; Shi and Mi, 2013; Wang, 2013; Guo and Wang, 2013; Zou et al., 2017; Niu and Yin, 2018; Shi et al., 2018; Wu et al., 2018; Wang et al., 2019), few of them can serve as benchmarks for calibrating and validating land surface models that require sufficient accuracy (Wang et al., 2016; Cao et al., 2019a). The accuracy of existing maps is constrained by the limited availability and quality of data used to create them, as well as the inadequacy of mapping approaches. Early permafrost maps on the QTP (Cheng et al., 2011; Shi and Mi, 2013) were compiled through visual interpretation based on limited data and expert judgment. Subsequently, satellite data and reanalysis data have become the main data sources of permafrost mapping (Wang, 2013; Zou et al., 2017; Shi et al., 2018; Wang et al., 2019). But large gaps in satellite data caused by clouds would highly affect the accuracy of permafrost maps in absence of effective interpolation methods (Chen et al., 2020). Although the reanalysis products do not

suffer from cloud contamination, their coarse spatial resolutions and associated large uncertainties on the QTP (Hu et al., 2019; Qin et al., 2020; Cao et al., 2020) would limit the accuracy of the derived maps.

65 Uncertainties associated with mapping approaches also negatively impact the accuracy of existing permafrost maps. Common statistical learning methods for permafrost mapping (Wang et al., 2019; Ni et al., 2021) rely heavily on in situ observations as a training dataset and are often compromised when ground observations are unevenly distributed and have different observation periods, as was the case with the QTP. This led to misrepresentation and overfitting in permafrost maps (Marcer et al., 2017). Meanwhile, the lack of accurate soil properties, fine-tuned parameterization schemes, and high-resolution
70 forcing data on the QTP severely challenged the applications of land surface models in mapping permafrost (Wu et al., 2018). These physically-explicit models were often calibrated and validated at a point scale, leading to unpredictable uncertainties when extended to a large region with more variability and thus more complex conditions (Qin et al., 2017; Wu et al., 2018). In addition, permafrost distribution maps generated by land surface models are usually not well suited as independent benchmark maps because the land surface models are more or less similar in terms of model structure and forcing data.
75 Therefore, empirical and semi-physical approaches remain the mainstay of permafrost mapping on the QTP as they require fewer in situ observations than statistical learning methods and have a simpler structure with fewer parameters than physical models (Zou et al., 2017; Zhao et al., 2017). Nevertheless, these maps have been criticized for limited consideration of local factors (Cao et al., 2019a; Hu et al., 2020) and the lack of constraints imposed to avoid divergence. All these issues call into question the ability of the existing permafrost distribution maps to serve as benchmark maps for land surface simulations on
80 the QTP.


An ideal benchmark map for spatial modeling of permafrost should be, 1) a map based on an adequate number of robust observations that have not already been used for the simulation; 2) a map based on the data from the mapping year, rather than all data spanning decades while they were still available; this is especially important if the benchmark map is used to calibrate a transient model; and 3) a map that accounts for the influences of local factors and is well constrained during the mapping
85 process. Based on these criteria, this study aims to produce a permafrost distribution map over the QTP in 2010 through an effective permafrost mapping approach that considers the effects of local factors and utilizes observational data, including remote sensing data and survey-based subregion permafrost maps, as optimization targets and constraints. Our goal is to provide a new reference map of 2010 for permafrost studies on the QTP and to provide a benchmark map for transient simulations of QTP permafrost under climate change.

90 **2 Study area and data**

2.1 Study area

The QTP (bounded within 73.5–104.5 °E and 26–40 °N) has an area of about 2.6×10^6 km² and is surrounded by high mountain ranges with relatively flat terrain in the interior (Fig. 1). Most of the QTP lies between 3000–5000 m above sea level (a.s.l.) with an average of about 4000 m a.s.l.. The mean annual air temperature ranges from -5 to 5 °C in most areas between

95 3000–5000 m.a.s.l., with the highest monthly temperature recorded at about 10 °C in July and the lowest at -10 °C in January. In the last five decades preceding 2010, air temperature increased by about 0.3–0.4 °C per decade, which is more than twice the global warming rate (Zhang et al., 2019). Mean annual precipitation decreases from more than 700 mm in the southeast towards about 50 mm in the northwest, and about 90% of precipitation falls during the growing season from May to September (Peng et al., 2019). The QTP is mainly covered by alpine desert in the northwest and transitions to alpine meadow and forest in the southeast (Wang et al., 2016). Snow cover on the QTP is thin and of short duration (Wu and Zhang, 2008). Alpine permafrost has formed extensively on the QTP, with ice-rich layers often found near the permafrost table on the plateau, which is generally 2–3 m deep (Zhao et al., 2020). Permafrost thickness on the QTP ranges from several meters to about 350 m while the depth of zero annual amplitude varies from 3.5 to 17 m (Zhao et al., 2020). The QTP permafrost is also characterized by a high mean annual ground temperature (MAGT), which is above -3 °C in most permafrost regions (Zhao et al., 2020).



105

Figure 1. Map showing the topography of the Qinghai-Tibet Plateau (QTP), the locations of meteorological stations, and the subregions with extensive field surveys. Inset maps show the local permafrost distributions in the five subregions based on the survey data of ca. 2010. WK: West Kunlun; GZ: Gaize; AJ: Aerjin; WQ: Wenquan.

2.2 Subregion permafrost maps

110 From 2009 to 2014, a research project sponsored by China Minister of Science and Technology was carried out to investigate permafrost and its surroundings. Intensive surveys were conducted in five areas (i.e. West Kunlun, Gaize, Aejin, National Highway G308, and Wenquan, Fig. 1), each characterized by distinct climatic and geographic conditions and representative of the different permafrost environments on the QTP (Zhao et al., 2017). Comprehensive information was acquired through field observations, mechanical excavations, geophysical reconnaissance techniques (ground penetrating radar
115 (GPR), time-domain electromagnetic), and borehole drilling, which allowed mapping permafrost distribution with high accuracy in all five subregions. Permafrost distribution in the Wenquan and West Kunlun subregions was mapped by a multivariate adaptive regression splines (MARS) model trained on large samples from field surveys: 130 GPR profiles and 21 boreholes in Wenquan; and 103 GPR profiles, 50 pits and 13 boreholes in West Kunlun. In the Gaize, Aejin and G308 subregions, the maps were based on aspect-stratified relationships between altitudinal limits of permafrost and topography
120 (Chen et al., 2016).

These subregion permafrost maps have been widely used as ground truth in many modeling studies (Zou et al., 2017; Zhao et al., 2017; Shi et al., 2018; Wu et al., 2018; Wang et al., 2019). The original maps have a spatial resolution of 250 m. In this study, these maps were aggregated and then resampled to 1 km resolution and used to calibrate an empirical soil parameter representing a synthesized soil thermal and moisture condition.

125 2.3 Satellite land surface temperature product

The land surface temperature (LST) data product from the Moderate Resolution Imaging Spectroradiometer (MODIS) onboard the Terra and Aqua satellites is one of the most widely used LST products due to its high spatial and temporal resolutions (Wan, 2008). It has a global coverage and has been applied in many permafrost mapping studies to provide temperature conditions (Gisnås et al., 2017; Zou et al., 2017; Obu et al., 2019; Wang et al., 2019). In this study, the daily
130 MODIS LST/emissivity products (MOD11A1 and MYD11A1 version 6) were used, providing up to two daytime and two nighttime LST observations at 1 km resolution. These observations were used to estimate annual ground surface thawing (DDT) and freezing indices (DDF) driving the mapping approach. DDT and DDF are defined as the cumulative number of degree-days in a year when ground surface temperatures (GST) are above and below zero degrees Celsius, respectively (Nelson and Outcalt, 1987). In practice, multi-year average DDT (DDF) are used instead of a single-year DDT (DDF) in order to avoid
135 single-year meteorological anomalies. In addition, the presence of permafrost is by definition determined by the thermal conditions of the last two years, indicating that not only the DDT and DDF of the current year, but also those of the previous years influence the presence of permafrost. We finally used a period of 2005-2010 for deriving the DDT and DDF for 2010, partly because automatic weather stations have been commissioned on the QTP since 2005, which we needed to accomplish the DDT estimation from the MODIS LST data.

140 **2.4 Environmental factors influencing permafrost distribution**

In our approach, we spatially divided the study area into different soil clusters to represent the heterogeneity of permafrost environments on the QTP on the basis of several environmental factors. The composite 16-day 1-km normalized difference vegetation index (NDVI) product (MOD13A2) provides information on vegetation greenness and has been shown to be well suited to differentiate the main vegetation classes on the QTP (Zhao et al., 2015). We calculated the average
145 aggregate from 2005 to 2010 and used it as an attribute for spatial clustering and as a predictor variable for estimating DDT. Topographical factors, including elevation and slope, were derived from the Shuttle Radar Topography Mission 90m digital elevation database (SRTM/DEM, version 4) (Reuter et al., 2007), and then aggregated to a working spatial resolution of 1 km. The STRM-derived topographic wetness index (TWI), together with mean annual precipitation from 2005 to 2010 aggregated
150 from the 1-km monthly precipitation dataset for China (Peng et al., 2019), represents wetness conditions affecting permafrost distribution. Likewise, average aggregate fraction snow cover (FSC) data were processed for the same period from the 500 m Daily Fractional Snow Cover Dataset Over High Asia (Qiu et al., 2017). Soil texture type data derived from the China Data Set of Soil Properties for Land Surface Modeling (Shangguan et al., 2013) were also included.

2.5 In situ observations

2.5.1 Ground surface temperature (GST) observations

155 There are 131 national meteorological stations of China on the QTP (Fig. 1), which are mostly concentrated in the eastern QTP. At these stations, standard meteorological variables including air pressure, air temperature, precipitation, evaporation, relative humidity, wind speed/direction, sunshine hours, and 0 cm ground surface temperature, are measured four times a day, at 2, 8, 14, and 20 o'clock. We extracted the daily GST observations during the period from 2005 to 2010 at these stations from the daily meteorological dataset of basic meteorological elements of the China national surface weather stations (version
160 3.0) (National Meteorological Information Center, 2019). The in situ GST observations were used to estimate DDTs from the satellite LSTs on the QTP.

2.5.2 Permafrost presence/absence observations

We used permafrost presence/absence information from boreholes to evaluate the new permafrost distribution map of this study. These data are independent of those used to produce subregion permafrost maps and come primarily from three
165 sources. A newly published synthesis dataset of permafrost thermal state on the QTP (Zhao et al., 2021) provides 65 boreholes where soil temperatures at 10 m and 20 m depths were monitored for 2005-2018. The presence of permafrost at borehole locations around 2010 was thus determined based on mean annual soil temperatures at the two depths, since previous evidence suggests that the depths of zero annual amplitude at these locations are within the two depths. These boreholes were further classified into three categories: boreholes with stable permafrost (mean annual soil temperature below -0.1 °C at both depths),

170 boreholes with unstable permafrost (between $-0.1\text{ }^{\circ}\text{C}$ and $0\text{ }^{\circ}\text{C}$ at both depths), and boreholes with seasonal frost (above $0\text{ }^{\circ}\text{C}$ at both depths).

In addition, seven boreholes were collected from existing literature (Li et al., 2016) providing information on permafrost presence in the Yellow River source area, a key region in the eastern QTP. Ground temperatures in these boreholes were measured in the summers of 2013 and 2014 and assumed to reflect the thermal regimes in 2010. The borehole locations were
175 classified as seasonal frost if soil temperatures at the 15 m depth were above $0\text{ }^{\circ}\text{C}$, otherwise permafrost. In the Yangtze River source area, an important, ecologically vulnerable permafrost region of the QTP, recent observations of the presence/absence of permafrost in 2020 in 32 boreholes (Li et al., 2022) were also used as a reference. Because permafrost on the QTP has warmed in recent decades (Cheng et al., 2019), some boreholes indicative of the presence of seasonally frozen ground (SFG) in 2020 may have appeared to be permafrost in 2010. Therefore, these 32 boreholes were not used to quantitatively validate
180 the results, but only as an aid to comparison.

2.6 Existing QTP permafrost maps for comparison

To better evaluate the new map of this study, two peer permafrost distribution maps with a resolution of 1 km were used. One was compiled by Zou et al. (2017) using the temperature at the top of permafrost model (TTOP) together with MODIS LST data from 2003 to 2012 (hereinafter, Zou map). The other map was developed via a data-driven approach by Wang et al.
185 (2019) (hereinafter, Wang map) with samples from two previous maps: a 2006 map (Wang, 2013) with the QTP portion mapped using a multilinear regression model (Nan et al., 2002), and the Zou map. MODIS LST data were also used as a predictor variable for the Wang map. Recently, the Zou map has been widely used to represent permafrost distribution around 2010 and has served as ground truth in many QTP studies (Hu et al., 2019; Song et al., 2020; Mu et al., 2020; Ni et al., 2021; Yin et al., 2021). Cao et al. (2019a) evaluated the Zou map as the best performing permafrost map on the QTP based on an
190 inventory of field evidence.

For simplicity, we excluded lakes and glaciers from our analysis. Glacier inventory data on the QTP were a subset from Guo et al. (2015), and lake data for the period 2008-2010 from Zhang et al. (2017) provided the lake boundaries in this study.

3 Mapping method and validation

3.1 The FROSTNUM/COP method and the improvements


195 We applied the FROSTNUM/COP mapping method developed by Hu et al. (2020) to map the distribution of permafrost on the QTP. The general process of this method and the improvements we made in this study are outlined in Fig. 2. It is based on the extended ground surface frost number (FROSTNUM) model fed by satellite temperature data (Fig. 2a,b), and requires permafrost distribution maps for subregions as optimization constraints. This method accounts for local factors through a model parameter E , whose values were optimally determined for all spatial units following a procedure of spatial clustering
200 (Fig. 2c), parametric optimization (Fig. 2d), and decision tree (Fig. 2e).

The extended FROSTNUM model determines the occurrence of permafrost by a frost number F :

$$F = \frac{\sqrt{DDF}}{\sqrt{DDF} + E \cdot \sqrt{DDT}} \quad (1)$$

where DDF and DDT represent annual ground surface freezing and thawing indices ($^{\circ}\text{C}\cdot\text{day}$), respectively. E is a parameter accounting for the combined effect of shifting soil properties from the unfrozen to the frozen state and is determined by the soil thermal properties and moisture conditions in both states. If F is greater than 0.5, the ground is determined as permafrost, otherwise, it is seasonal frost. Under ideal circumstances, if soil conditions remain constant during the phase change, E equals 1 and Eq. (1) becomes Nelson's original frost number model (Nelson and Outcalt, 1987). Although the parameter E is physically well defined, in practice it is impossible to compute its value directly due to the lack of accurate information on soil properties and moisture conditions. Therefore, the FROSTNUM/COP method resorts to an optimization procedure to solve for E , which is detailed in Section 3.3. Once the freezing and thawing indices and parameters are ready, the extended FROSTNUM model is applied to map the distribution of permafrost across the QTP.

The optimization procedure in the original FROSTNUM/COP method uses Cohen's Kappa coefficient (Cohen, 1960) as an objective function, which measures the agreement of frozen soil type classification between the simulated map and survey-based subregion maps. This may lead to an equifinality problem given binary categorical raters for Kappa. In this study, we specially reduced the equifinality problem by modifying the objective function to additionally include a metric that guarantees boundary consistency and by introducing an ensemble simulation of 1000 runs of parametric optimization. These processes are explained in Section 3.3.



220 **Figure 2. Workflow of the permafrost mapping method. (a) The data required in this study. (b) Preparation of annual ground surface freezing indices (DDF) and thawing indices (DDT) based on satellite land surface temperature (LST) data. (c) Process of spatial clustering of soils in subregions and prediction of clusters for the study area based on local factors. (d) Determination of optimal values for soil parameter (E) in the model using the particle swarm optimization (PSO) algorithm constrained by the subregion maps. (e) The process of mapping the permafrost distribution on the QTP based on 1000 runs using the extended ground surface frost number model (FROSTNUM). Dashed blue lines mark the improved processes over the original FROSTNUM/COP**

225 method (Hu et al., 2020), including refinement of the optimization objective and ensemble runs. The diagram was modified from Hu et al. (2020). GST: ground surface temperature; NDVI: normalized difference vegetation index.

3.2 Preparation of ground surface freezing and thawing indices

DDF and DDT were calculated based on the MOD11A1 and MYD11A1 level 3 products (version 6). Gaps in the MODIS LST data due to cloudiness resulted in systematic cold biases (Westermann et al., 2012) and consequently uncertainties in mapping permafrost based on these data. Despite many all-weather LST products (Zhang et al., 2021; Xu and Cheng, 2021), we chose a stepwise interpolation approach based on the solar-cloud-satellite geometry (SCSG) effect (Chen et al., 2022) to interpolate the data-gap regions in the MODIS LST data. Compared to existing approaches, the SCSG-based approach requires only MODIS family data and is effective for extensive missing data (e.g. on the QTP). A brief introduction to this interpolation method is provided in Appendix A.

235 Due to the buffering effect of seasonal snow cover and vegetation, thermal offsets often exist between satellite-derived LST values and GST values. In most areas of the QTP, snow cover is thin and short-lived (Wu and Zhang, 2008; Zhao et al., 2017), thus the buffer effect of snow cover is limited and LSTs are close to GSTs during snow-free periods (Hachem et al., 2012), as also shown later in this study. Therefore, for the DDF values, we simply calculated the sum of negative degree-days of mean daily LST from four instantaneous LST observations, ignoring the effects of snow cover.

On the contrary, vegetation cover complicates DDT calculations by providing a strong thermal buffer between GST and LST, especially on the eastern QTP during growing seasons. Therefore, thermal offsets should be removed from the raw LST data before DDT can be derived as a sum of positive degree-days of mean daily LST. To this end, we developed a multilinear regression model where GST is a function of independent variables including the raw LST, NDVI, and latitude at weather stations (Huang et al., 2020). The correction was repeated over 23 time intervals a year, and the annual DDT was the aggregate of all these corrected positive degree-days. More information regarding this process can be found in Appendix B.

245 3.3 Determination of optimal values of soil parameter E

We followed the method developed by Hu et al. (2020) to spatially group the soils of five subregions into soil clusters. Because the QTP is a much larger region with more complex climate and terrain conditions than the experimental area in the previous study (Hu et al., 2020), the environmental variables we chose to account for the influences of local factors on permafrost distribution were slightly different. Apart from the previous factors of elevation, slope, TWI, precipitation, and soil texture type, we added NDVI and FSC as a response to the relatively strong heterogeneity of surface conditions on the QTP. Compared to Hu et al. (2020), we excluded the relief degree due to its high correlation with slope. To allow mixed clustering of categorical (soil texture type) and numerical variables, the k-prototype approach (Huang, 1998) was employed. Lakes were excluded during the clustering analysis.

255 The particle swarm optimization (PSO) algorithm (Wang et al., 2018) was used to find the optimal value of E associated with each soil cluster. In this population-based heuristic method, the candidate solutions are guided toward the best-known

positions in the search space, thus enabling a very rapid convergence to an optimal value. In the previous study (Hu et al., 2020), the only objective function was Cohen’s Kappa coefficient (Cohen, 1960), which quantifies the agreement between the simulation map and the survey-based subregion permafrost distribution maps. Despite the good performance achieved in the experimental study area (Gaize in Fig. 1), this relatively simple objective function inevitably leads to equifinality in larger regions such as the QTP. Recognizing that the Kappa coefficient is a good representation of the overall consistency between simulation results and subregion maps, we retained the Kappa coefficient (κ) and made the objective function more rigorous by adding a specially defined boundary consistency. The objective function is then a weighted sum of overall consistency (κ) and boundary consistency (β):

$$F_{ob} = \omega_{\kappa} \cdot \kappa + \omega_{\beta} \cdot \beta \quad (2)$$

where F_{ob} is the objective function value, ω_{κ} and ω_{β} are the weights imposed on κ and β , respectively, $\omega_{\kappa} + \omega_{\beta} = 1$. Since there is no simple way to determine ω_{κ} and ω_{β} , to minimize the effects of the weights a random value between 0.2 and 0.5 was chosen for ω_{β} and correspondingly for ω_{κ} in each of a total of 1000 ensemble runs. β represents boundary consistency, which measures how well the boundaries between permafrost and SFG zones in the subregion maps are represented by the simulation.

β is defined as the number of ‘positive boundary cells’ (N_m) normalized by the total number of ‘boundary cells’ (N_b), with a range of 0 to 1:

$$\beta = \frac{N_m}{N_b} \quad (3)$$

A boundary cell is a cell on the survey-based subregion maps, whose neighboring cells of a size $n \times n$ satisfy two conditions. First, the neighboring cells must contain both types of frozen ground (permafrost and SFG). Second, the neighboring cells contain at least two soil clusters. According to Eq. (1), the permafrost cells must have larger F values than those of SFG grid cells. Therefore, in the neighboring cells of any ‘boundary cell’, the F value averaged over the permafrost zone (\bar{F}_p) must be greater than that of the SFG zone (\bar{F}_s). A ‘boundary cell’ is ‘positive’ when this condition ($\bar{F}_p > \bar{F}_s$) is met, otherwise it is ‘negative’.

The optimization procedure aims to maximize β as part of the objective function, i.e., as high as possible boundary consistency of the simulated map relevant to the subregion maps. Since in the simulated map the DDTs and DDFs were already predetermined before the optimization procedure, frost number F in each grid cell depends on the E value associated with the specific soil cluster of that cell. This means that by adjusting the E values of the soil clusters in the neighboring cells of a ‘negative boundary cell’, that cell has the potential to turn into a ‘positive’ cell. In the other words, the number of ‘positive boundary cells’, or β , is a function of E , thus permitting parametric optimization. To illustrate this concept, we present a simple instance of a ‘boundary cell’ in Appendix C.

The lower and upper limits of E values were specified at 0.5 and 1.5, respectively, and the optimal E values were determined for all soil clusters occurring in the subregions. Finally, a C5.0 decision tree (Kuhn and Johnson, 2013) was trained

on the information of soil clusters in the subregions and then applied to predict soil clusters for all regions outside the subregions on the QTP on a cell basis, based on the same environmental factors used in spatial clustering. After the distribution map of soil clusters on the QTP was obtained, the values of soil parameter E for the QTP were determined by simply looking up the optimal E value associated with each soil cluster in the soil cluster distribution map.

290 **3.4 Mapping permafrost distribution and evaluation**

Once E values are known for all QTP cells, the extended FROSTNUM model was run to determine the type of frozen ground of each cell using a threshold of $F=0.5$ and then the permafrost distribution on the QTP can be mapped. However, this map may still be affected by local optima. To reduce these issues, parameter optimization was performed 1000 times, a number that warranted minimal variability of individual E values in our experiment, and the permafrost distribution on the QTP was estimated 1000 times in response to 1000 different sets of E values. Finally, an ensemble permafrost map on the QTP was generated by majority voting of the 1000 estimates.

We validated the resulting map (hereafter, our map) from multiple aspects. Although these survey-based subregion permafrost maps have been used as constraints during the optimization process, the optimal E values were obtained from all subregion maps as a whole. Therefore, the survey-based permafrost map in each subregion is still of value for validation. We also validated the maps using in situ permafrost presence/absence observations around 2010. We compared our map with two existing permafrost maps, the Zou map and Wang map (Zou et al., 2017; Wang et al., 2019), using the same references. In particular, we analyzed the spatial inconsistency between our map and the Zou map (Zou et al., 2017) in some typical regions. In these regions, we further evaluated the two maps by utilizing additional information from boreholes, satellite imagery, the permafrost zonation index (PZI) map (Cao et al., 2019b), and elevation characteristics. Satellite imagery provides indicative landscape evidence on permafrost occurrence. While the PZI rarely equates to the actual presence of permafrost, it indicates a probability of permafrost presence with a value ranging from 0 to 1 (Gruber, 2012). In some regions of the QTP where permafrost is thermally controlled by elevation, the dependency of permafrost occurrence on elevations provides useful information for evaluating permafrost distribution maps.

4 Results and discussion


310 **4.1 Ground surface thawing /freezing indices**

Figure. 3 demonstrates the discrepancies between average annual in situ DDT (DDF) values, calculated as averages over 2005-2010 from daily mean GSTs at 131 weather stations on the QTP, and the average annual satellite DDT (DDF) values derived directly from the daily mean MODIS LSTs at the corresponding MODIS pixels. The raw LST-derived DDF values matched perfectly with the in situ DDF values, echoing the limited effects of thin and short-duration snow cover on the thermal states of underlying soils on the QTP (Wu and Zhang, 2008; Zhao et al., 2017). In contrast, LST-derived annual DDT tended to underestimate in situ DDT, with substantial discrepancies at some sites. The discrepancies are mainly connected to the


thermal offset between remotely sensed LST and GST, which has also been reported by previous studies (Luo et al., 2018; Obu et al., 2019).

320 Obvious negative biases were observed in the raw LST-derived DDT values (Fig. 4a). However, after applying the interval-based approach, the negative biases were well removed. Moreover, the corrected data points were concentrated along the 1:1 line as shown in Fig. 4b while the coefficient of determination (R-squared) increased from 0.74 to 0.89. More importantly, the mean absolute error (MAE) value (334 °C day) of the corrected LST-derived DDTs was about one-third of the value (889 °C day) before the correction, i.e., the relative error dropped from 23.3% to 8.8%, below the 10% level. In addition, raw DDT data points with large deviations have been effectively corrected, resulting in a reduction in root mean square error (RMSE) of 1072 °C day to 421 °C day after correction. Most corrected data points fall into the ± 400 °C·day band (about 10%), indicating a well-controlled level of error after the removal of thermal offsets between GST and LST.


330 The distributions of annual DDT and DDF (Fig. 5a,b) were in close agreement with the characteristics of elevation (Fig. 1), which is one of the main factors controlling ground temperature distribution over the QTP. In general, annual DDT decreased and annual DDF increased with rising elevation. Over the relatively flat high plain between 80-90 °E and 33-37 °N, the annual DDT showed moderate latitudinal zonality, declining with increasing latitude, whereas the annual DDF showed the opposite. This is an indication of the influence of solar radiation on GST. The vast area and complex topography of the QTP resulted in a wide spectrum of annual DDT (from 0 to 9000 °C day) and DDF (from 0 to 8000 °C day). Most regions of the QTP lie between 3000 and 5500 m a.s.l., where DDT and DDF values were mostly between 1000 °C day and 2500 °C day. High DDT values appeared in the low mountains in the southeast QTP, in the Qaidam Basin in the north, and in the south valleys, whereas high DDF values appeared on the Qiangtang Plateau in the northern QTP and in the high mountain areas, favoring the formation of permafrost. The ratio of DDT-over-DDF indicates climatic controls on permafrost preservation (Fig. 5c). Regions with a DDT-over-DDF ratio < 1 would have the potential to form permafrost in absence of local factors that inhibit permafrost formation.



340 **Figure 3. Thawing/freezing indices calculated from the interpolated MODIS LST data (raw LST-derived DDT/DDF) and in situ observations of ground surface temperature (in situ DDT/DDF) at each QTP weather station. The ordinate indicates the annual thawing/freezing indices averaged over the period 2005 to 2010; and the abscissa shows 131 weather stations available on the QTP.**



345 **Figure 4. Bias correction of MODIS LST-derived DDT with the interval-based approach. (a) Before bias correction; (b) after applying the interval-based method. Data points represent annual DDT values in 2005-2010 from the 131 stations. The red dashed lines outline a range of ~10% (± 400 °C·day) from the 1:1 line (solid black line).**



350 **Figure 5. Maps of spatial distribution of (a) annual DDT, (b) annual DDF, and (c) the ratio of DDT-over-DDF on the QTP, averaged over 2005 to 2010. Lakes were excluded and are shown blank, whereas glaciers were included. Regions with a DDT-over-DDF ratio of <1 are climatically favorable for permafrost formation.**

4.2 Soil clusters


A total of eight soil clusters were determined by the k-prototype approach in the five subregions (Fig. 6), where lakes
 355 were excluded. Soils in one cluster share more similar environmental characteristics to each other, as reflected by a single value of model parameter E , than soils in other clusters. The dominant soil clusters in each subregion differed from each other (Table 1). Clusters 3 (30%) and 1 (30%) were dominant in West Kunlun, clusters 2 (58%) and 7 (23%) in Gaize, clusters 7 (49%) and 1 (23%) in Aertjin, clusters 8 (55%) and 7 (18%) in G108, and cluster 8 (85%) in Wenquan. This implies the distinctions of climatic and geographic conditions in these subregions.

360 Among all clusters, clusters 1, 2, and 3 differ in slope and TWI, but are all characterized by relatively high elevation (about 5000 m), low vegetation cover ($NDVI < 0.2$), thin snow cover ($FSC < 10$), and aridity (precipitation $< 200\text{mm}$) (Fig. 7), which generally may represent high plateaus. Cluster 4, with the highest TWI, represents the valley with low elevation and moderate slope. Cluster 5 has the highest elevation, highest FSC, and lowest NDVI (even below zero), thus it may represent high mountains covered by thick snow cover or glaciers. Cluster 6 has very varied elevations and steep slopes, and occurs on
 365 the hillslopes of high mountains. Except for the much lower TWI, cluster 7 is similar to cluster 4, and often appears around cluster 4, which represents valleys (Fig. 6). Therefore, it is likely that cluster 7 represents gentle slopes near valleys. Cluster 8 is mainly distributed in the two subregions (G308 and Wenquan) on the east QTP and characterized by the highest NDVI, highest precipitation, and lowest elevation, and it represents the soils with better hydrological, thermal and vegetation conditions on the east QTP.

370 The distribution of soil clusters on the QTP (Fig. 6) was predicted by the decision tree method. Soil cluster 8 covered the largest area of about 37.76% of the QTP and was mainly distributed in the east QTP, which is related to the training samples that were mainly located in G308 and Wenquan on the eastern QTP (Table 1). Soil clusters 2, 3, 6, and 7 covered roughly the same proportion of area, about 10% of the whole QTP, followed by clusters 1 (6.94%) and 4 (6.20%). Soil cluster 5,

representing glaciers and regions with thick snow cover, occupied the least area (2.85%), which is consistent with a previous study that thick snow coverage only represented a relatively small portion of the QTP (Dai et al., 2018).

The optimal values of parameter E associated with the soil clusters (Table 2) were determined from the 1000 optimization runs. The ranges of the optimal values were relatively narrow for all soil clusters, suggesting that equifinality was well mitigated due to a well-constrained objective. The mean values as the optima for clusters 4 and 5 were greater than 1, with an implication of unfavorable local conditions for permafrost formation and preservation. For example, heat advection by water flows occurring near rivers in valley areas represented by cluster 4 and the insulation effect of snow cover in regions of cluster 5 are not beneficial for permafrost formation and preservation. Clusters 1, 3, and 8 had relatively lower E values, suggesting favorable local environments for permafrost formation in these regions. Some characteristics in local factors, such as high elevation for clusters 1 and 3, and high precipitation for cluster 8, are beneficial for permafrost preservation (Zhang et al., 2021), as also reflected by their lower E values.




385

Figure 6. Resulting soil clusters in the five subregions and the predicted distribution of clusters on the QTP. A total of eight clusters were determined. Each soil cluster represents unique traits as reflected by a distinct value of model parameter E .

Table 1. Area percentages occupied by individual soil clusters in each subregion, over all subregions, and over the entire QTP.

Region	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6	Cluster 7	Cluster 8
West Kunlun	29.53	2.02	30.65	4.21	16.21	14.13	1.89	1.37

Gaize	1.11	57.75	5.34	12.62	0.12	0.44	22.55	0.06
Aerjin	22.97	12.39	8.51	3.88	0.40	2.88	48.76	0.20
G308	6.24	9.56	4.40	7.29	0.00	0.12	17.24	55.15
Wenquan	7.27	6.37	0.72	1.62	0.00	0.04	9.01	74.97
All subregions	14.63	25.93	16.27	7.93	7.17	6.44	12.91	8.72
QTP	6.94	11.45	11.69	6.20	2.85	9.31	13.79	37.76



390

Figure 7. Environmental characteristics of the soil clusters in the subregions: (a) elevation; (b) slope; (c) normalized difference vegetation index (NDVI); (d) fractional snow cover (FSC); (e) topographic wetness index (TWI); (f) precipitation. All clusters are shown in different colors to match those in Fig. 6. The center line in the box shows the median, the box shows the lower and upper quartiles, and the whiskers extend to the minimum and maximum data values.

395

Table 2. Ranges and mean values as the most optimal values of the soil parameter E associated with the eight soil clusters. The results were obtained from 1000 optimization trials.

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6	Cluster 7	Cluster 8
Max	0.778	0.831	0.784	1.014	1.149	1.011	0.938	0.785
Min	0.757	0.820	0.745	1.000	1.061	0.931	0.932	0.778
Mean	0.768	0.822	0.748	1.007	1.073	0.946	0.938	0.779

400 4.3 The resulting 2010 permafrost distribution map on the QTP

The resulting permafrost distribution in 2010 on the QTP is shown in Fig. 8 with a spatial resolution of 1km. Permafrost covered about 1.086×10^6 km², or 41.17% of the QTP, while SFG occupied about 1.447×10^6 km², or 54.85% of the total QTP

area. The non-frozen ground was about $2.24 \times 10^4 \text{ km}^2$, and the rest consisted of glaciers (about $4.08 \times 10^4 \text{ km}^2$) and lakes (about $4.17 \times 10^4 \text{ km}^2$).

405 The map shows that permafrost was prevalent throughout the north-central QTP, especially on the Qiangtang Plateau. In the north, the Qaidam Basin was occupied by SFG due to its low altitude, interrupting the continuity of permafrost that extended north to the Qilian Mountains. From the central Qiangtang Plateau southward, the spatial continuity of permafrost tended to decline due to decreasing latitude and elevation. Near the permafrost zone in the Bayan Har Mountains and Tanggula Mountains in the eastern QTP, SFG occurred extensively in the river source areas, namely the Three-River Headwaters Region, 410 probably due to the low latitude and the effects of heat advection by water flows that prevent permafrost formation. As the DDT-over-DDF ratios in these regions were generally greater than 1 (Fig. 5c), permafrost in the river source areas (e.g. the Yangtze River headwaters) was thermally vulnerable and very sensitive to climate warming (Zhang et al., 2022). On the southern QTP, permafrost was sporadically distributed at high elevations, mainly in the high mountains of the Eastern Himalayan Syntaxis, Gangdise Mountains, and Himalaya Mountains. Only a small amount of non-frozen ground existed in 415 the south QTP.




Figure 8. Map of permafrost distribution at 1 km resolution over the QTP in 2010 (our map), produced in this study. Areas and percentages of frozen soil types are provided. The map shows hill-shading with elevation.

420

4.4 Assessment based on survey maps and borehole data

Our map showed substantial spatial agreement with the survey-based permafrost maps in all subregions (Fig. 9), with a Cohen's Kappa coefficient (κ) of about 0.74 (Table 3), which was notably higher than that of the Zou map (Zou et al., 2017) and the Wang map (Wang et al., 2019), whose Kappa values were 0.55 and 0.50, respectively. Overall, compared with the survey-based maps (Fig. 9a) and our map (Fig. 9b), the Zou map significantly overestimated permafrost extents in Gaize and Aerjin and underestimated permafrost extent in G308 (Fig. 9c), while the Wang map severely overestimated permafrost extents in all subregions (Fig. 9d). The differences can be better discerned in the difference maps in Fig. E1.

More specifically, in permafrost-dominated West Kunlun, our map and the Zou map slightly overestimated the extent of the SFG around the lake, while the Wang map slightly underestimated the extent of SFG. There were only small differences between the three maps in West Kunlun, with almost the same κ for all three maps (0.62 for our map, 0.63 for both the Zou map and Wang map). All three maps were based on satellite LST data, which can generally capture the patterns of surface ground temperature. Although the interpolation methods for processing LST gaps differed and resulted in data with different accuracy for mapping, the cold climate in West Kunlun, with LST values well below zero in most areas, made the impact of these differences in input data on permafrost distribution negligible.

In Gaize, our map performed better ($\kappa = 0.71$) than the Zou map ($\kappa = 0.48$) and Wang map ($\kappa = 0.43$). Both the Zou and Wang maps severely overestimated permafrost distribution, whereas our map agreed well with the survey-based map in this region (Fig. 9b). The same trends occurred in Aerjin, and the comparison with the survey-based map indicated a much lower κ for the Zou map (0.38) and Wang map (0.00) than our map (0.71). Gaize has a warmer climate than West Kunlun and contains the southern limit of continuous permafrost. Therefore, the influence of input data accuracy and local factors on permafrost preservation is more profound in this area. The overestimated permafrost extents in Gaize and Aerjin in both the Zou map and Wang map may most likely be related to the relatively lower quality of the interpolated LST data as model input and insufficient consideration of local factors in the mapping approaches. Compared with the harmonic analysis of time series (HANTS) algorithm (Xu et al., 2013) used for the Zou and Wang maps to reconstruct the missing LST data under clear sky assumptions, the SCSG-based interpolation method with full consideration of cloud effects on LST used in this study was found to be effective in handling large areas of missing data with sufficient accuracy (Chen et al., 2022). Moreover, the daily GST data required to produce the Zou map and Wang map were the weighted sum of four MODIS LST observations per day through an empirical linear formula based on sample observations from three automatic weather stations in the central QTP (Zou et al., 2014). This relatively simple treatment of GST data in the Zou map and Wang map can lead to considerable systematic biases in some regions, especially in warm permafrost regions that are highly susceptible to thermal perturbations (e.g. the Gaize subregion), and ultimately cause large uncertainties in the final permafrost distribution maps. In contrast, the thermal offsets between GST and LST have been well handled in this study when estimating thawing indices from satellite LST observations by considering the effect of vegetation cover as a buffer layer based on 131 weather stations over the QTP. For the Wang map was produced using statistical learning methods, uncertainties also resulted from training samples selected

455 from two previous QTP permafrost maps produced in different years more than a decade apart and subject to varying levels of
uncertainty (Ran et al., 2012; Zou et al., 2017). All of these factors combined to overestimate permafrost extent in Gaize in the
Wang map.

In G308, the Wang map ($\kappa = 0.68$) indicated more permafrost areas than the local survey map, while both the Zou map
($\kappa = 0.48$) and our map ($\kappa = 0.68$) showed fewer permafrost areas, with the Zou map even more evident. The soil thermal
regime in G308 is strongly influenced by rivers and vegetation cover, and the effects were well accounted for in our mapping
460 approach, while they were absent in the Zou map. In Wenquan, both our map ($\kappa = 0.70$) and the Zou map ($\kappa = 0.65$) performed
generally satisfactorily, with a slight overestimation of permafrost extent. In contrast, the overestimation was more pronounced
in the Wang map ($\kappa = 0.46$), which is probably also related to the misrepresented training samples used for this map.

The maps were also verified by 72 permafrost presence/absence observations obtained by boreholes drilled within a five-
year time frame around 2010 (Li et al., 2016; Zhao et al., 2021). Only our map showed good agreement ($\kappa = 0.43$) with the
465 borehole observations in terms of κ (Table 4), compared to Zou map ($\kappa = 0.30$) and Wang map ($\kappa = 0.14$). According to the
borehole observations, SFG was underrepresented in all three maps, with our map performing the best in predicting SFG with
54.5% accuracy, as opposed to the Wang map with the worst performance, correctly identifying only 1 out of 11 seasonal frost
boreholes. Unlike SFG, permafrost was overestimated at borehole locations in three maps, as evidenced by relatively high
false positive (permafrost) rates (45.5% for our map, 54.5% for the Zou map, and 90.9% for the Wang map). As most of the
470 borehole locations were underlain by permafrost (Fig. 9), the Wang map predicted almost all locations as permafrost with no
discretion, as indicated by a 100% true position rate (Table 4), and consequently led to an inflated accuracy of 86.1%, which
was the highest among the three maps.

In our map, two out of six false negatives (misidentified as SFG) were the boreholes with unstable permafrost located in
the SFG zone close to the permafrost boundary. However, in the Zou map, all eight false negatives were boreholes with stable
475 permafrost, and those with unstable permafrost were in the permafrost zone. If we excluded all four boreholes with unstable
permafrost from the evaluation, the false negative rate of our map would drop from 9.8% to 7% and κ would rise from 0.43 to
0.49, whereas the false negative rate and κ of the Zou map would remain almost unchanged, leaving an even higher false
negative rate (~13%) than that of our map (~7%). This borehole-based verification may be biased by the mismatch between a
site and a 1 km \times 1 km grid cell. Nevertheless, those evidences as a whole point to the decent performance of our map in
480 predicting frozen ground distribution.




Figure 9. Spatial distributions of frozen ground in the subregions from (a) survey-based maps, (b) our map, (c) the map from Zou et al. (2017), and (d) the map from Wang et al. (2019). Triangle symbols mark the locations of boreholes drilled in around 2010. Difference maps between survey-based maps and the three simulated are provided in Figure E1.

485

Table 3. Kappa values measured between the evaluated permafrost maps (our map, Zou map, and Wang map) and survey-based maps in the subregions.

	West Kunlun	Gaize	Aerjin	G308	Wenquan	All subregions
Our map	0.62	0.71	0.71	0.68	0.70	0.74
Zou map	0.63	0.48	0.38	0.46	0.65	0.55
Wang map	0.63	0.38	0.00	0.68	0.46	0.50

490

Table 4. Measures of confusion matrices describing the performance of the evaluated permafrost maps (our map, Zou map, and Wang map) at the borehole locations. To fit the binary classification, permafrost is regarded as positive and SFG is considered negative. n=72.


	Our map	Zou map	Wang map
True positives (rate)	55 (90.2%)	53 (86.9%)	61 (100.0%)

False positives (rate)	5 (45.5%)	6 (54.5%)	10 (90.9%)
True negatives (rate)	6 (54.5%)	5 (45.5%)	1 (9.1%)
False negatives (rate)	6 (9.8%)	8 (13.1%)	0 (0.0%)
Accuracy	84.7%	80.6%	86.1%
Cohen's Kappa	0.43	0.30	0.14

4.5 Cross-comparison with the Zou map

495 The permafrost distributions in our map and Zou map were generally comparable, although there were discrepancies in some regions (Fig. 10), mainly in the transition region between the continuous permafrost zone of the Qiangtang Plateau to the north and the SFG zone to the south. In addition, the headwaters of China's major rivers (regions c, d in Fig. 10) in the eastern QTP showed noticeable spatial inconsistency between the two maps. These headwater regions were reported to be the critical regions where permafrost is warm and very susceptible to degradation due to climate change (Jin et al., 2011; Zhang et al., 2021). Permafrost there is characterized by high temperature (MAGT > -2.0 °C) and low thermal stability (Qin et al., 2017). The warm permafrost is difficult to distinguish from SFG, which poses a challenge to the accuracy of soil temperature modeling. Moreover, permafrost in transition areas is often controlled by many local factors (terrain, vegetation, soil properties and so on), and a model without adequate consideration of local factors often fails to accurately describe the soil thermal regime.

500



505 **Figure 10. Spatial inconsistencies in the distribution of frozen ground between our map and the Zou map. “Both P” represents areas**
identified as underlying permafrost in both maps; “Both SFG” represents areas identified as seasonally frozen ground (SFG) in both
maps; “Zou-P and Our-SFG” represents areas identified as underlying permafrost in the Zou map but SFG in our map; “Zou-SFG
and Our-P” represents areas identified as SFG in the Zou map but permafrost in our map. The dashed boxes highlight areas of
 510 **significant inconsistency. (a) Gaize and its vicinity, (b) the areas between the Altun Mountains and the Kunlun Mountains, (c) the**
headwaters of the Yangtze River, and (d) the headwaters of the Yellow River.

In and around Gaize (region a in Fig. 10), a larger extent of permafrost was simulated in the Zou map than in our map. Comparisons of the two maps with the survey-based Gaize map (Fig. 9) have already confirmed a better performance of our map in this region than the Zou map, due to the use of the survey-based Gaize map as part of the constraints in modeling our map (Table 3). The vicinity of Gaize is very similar to the Gaize subregion, also characterized by a relatively flat plateau with
 515 an arid climate and low vegetation cover. It can be inferred that our map in and around the Gaize subregion could likely have better accuracy than the Zou map.

In the areas between Altun and Kunlun Mountains (region b in Fig.10, Fig. 11) containing the Aejin subregion, our map estimated much more SFG than the Zou map. Referring to the survey-based Aejin subregion map (Fig. 9), the Zou map underestimated the extent of SFG, and our map showed a better performance despite a slight overestimation of SFG extent.
 520 According to borehole records (Fig. 9) in the Aejin subregion, some locations had ground temperature at 10 m of about -0.1 to 0 °C, and one borehole location was even above 0 °C but fell within a permafrost zone in the survey-based map. This reflects that permafrost in this region was extremely thermally unstable. We especially inspected inconsistency zones identified as

permafrost in the Zou map but as SFG in our map (Fig. 11a), where the DDT-over-DDF ratios were around 1.3 (Fig. 11b) and clusters 4 and 7 predominated with E values of about 1.07 and 0.94, respectively, (Fig. 6). Those characteristics are very similar to the SFG zone in the survey-based Aerjin map. Following Eq. (1), surface frost numbers were less than 0.5 in these areas with a climatic implication of no permafrost presence. From the satellite image (Fig. 11c), it can be seen that rivers are well developed in the basins. The presence of these rivers could potentially lead to greater degradation of permafrost due to the thermal advection of water flows. Overall, our map showed more acceptable distribution characteristics in this region than the Zou map. However, further field studies are necessary to provide more direct evidence to strengthen our understanding toward permafrost distribution in this critical region.





Figure 11. Maps of the areas between Altun Mountains and Kunlun Mountains (region b in Fig. 10) showing (a) detailed spatial differences in permafrost distribution between our map and the Zou map, (b) the ratios of DDT-over-DDF, and (c) a satellite image covering this region from Google Earth. The box indicates the Aerjin survey area for which the survey-based permafrost map is available. P (survey-based) and SFG (survey-based) represent permafrost/seasonally frozen ground in the survey-based map, respectively. See Figure 10 for notations. Ovals mark areas that were mostly likely to be thermally affected by the presence of water bodies.

In the source areas of the Yangtze River (region c in Fig. 10, Fig. 12), the riparian zones were generally identified as SFG in both our map and the Zou map. However, the SFG zones in our map spread on both sides along the rivers, whereas they were distributed on only one side of the rivers in the Zou map. In this region, 35 observations on permafrost presence/absence were collected. Of these, 32 were drilled in 2020 during the Second Tibetan Plateau Scientific Expedition and Research campaign (Li et al., 2022), so they were not included in the quantitative validation above. Boreholes QT11, QT15 and TGLGT (Fig. 12) collected from Zhao et al. (2021) were drilled before 2010, and the frozen ground types at these

borehole locations were correctly identified in both maps. For the 32 boreholes drilled in 2020, 2 of 6 boreholes with seasonal
545 frost and 24 of 26 boreholes with permafrost were correctly identified in our map, whereas no borehole with seasonal frost and
24 boreholes with permafrost were correctly identified in the Zou map. The misidentified boreholes were located near the
boundary of the permafrost zone on our map, whereas they were mostly located within permafrost zones on the Zou map. We
also noted that two upstream boreholes at 4870 m (Li et al., 2022), located within a permafrost zone in both maps (red box in
Fig. 12a,b), were revealed as seasonal frost in 2020. In a borehole labelled QTB15 (Fig. 12 a,b, Zhao et al., 2021) in this region,
550 ground temperature experienced a significant increase from 2006-2018 (Table E1), indicating a warming trend. Considering
the potential impact of climate warming occurring in this region over the past decade, it is possible that permafrost degradation
has occurred at the two upstream borehole locations, resulting in the conversion of permafrost in 2010 to SFG in 2020. In these
areas, the occurrence of permafrost degradation usually recedes to upstream areas with higher elevations and cooler air
temperatures. In other words, by reasonable inference, in 2010 permafrost would remain in upstream areas, while SFG was
555 present in downstream areas along the rivers, as well depicted by our map (Fig. 12a).

We further examined the two maps in the Yangtze River source areas using a PZI approach (Cao et al., 2019b). By
definition, permafrost regions should have higher PZI values than SFG regions. The PZI map (Fig. 12d) used here was
compiled based on 1475 in situ observations (Cao et al., 2019b), many of which were obtained between 2005-2018 in the
vicinity of the G109 National Highway traversing the Yangtze River headwaters, making the PZI map a possible reference in
560 this region. The PZI statistics for permafrost in our map were close to those in the Zou map (Fig. 12e). However, for the PZI
statistics in SFG regions, the lower and upper quartiles in the Zou map were 0.36 and 0.66, respectively, whereas the values
in our map were 0.34 and 0.53, respectively. The SFG regions shown in our map had lower PZI values. The upper quartile for
SFG regions (0.66) in the Zou map surpassed the lower quartile for permafrost regions, which was 0.55. The overlap is
questionable because it suggests that some SFG regions have higher PZI values than permafrost regions in the same map,
565 which contradicts the PZI definition. In contrast, the PZI ranges for both frozen ground types were more clearly distinguishable
in our map.




570 **Figure 12.** Maps in the Yangtze River headwaters (region c in Fig. 10) showing permafrost distributions in (a) our map and (b) Zou map and (c) the spatial differences, along with (d-e) spatial and statistical permafrost zonation index (PZI) distributions in this region. The boreholes QTB11, QTB15 and TGLGT in (a) and (b) were drilled before 2010 and provided by Zhao et al. (2021), while the others were drilled in 2020 during the Second Tibetan Plateau Scientific Expedition (Li et al., 2022). The red box covers two boreholes of particular concern. Both boreholes at an elevation of 4870 m a.s.l. were within a permafrost zone in both our map and Zou map, but were revealed seasonal frost in 2020. For (c), the same notations apply as in Figure 10. (e) Box plot showing the statistical distributions of PZI values for permafrost and SFG regions in our map (Our-P and Our-SFG) and those in the Zou map (Zou-P and Zou-SFG). The center line in the box shows the median, the box shows the lower and upper quartiles, and the whiskers extend to the minimum and maximum data values.

575

Similar to the Yangtze River source areas, there were considerable discrepancies between our map and the Zou map in the Yellow River source areas (region d in Fig. 10, Fig. 13). In this region, observations in 2013 and 2014 of seven boreholes collected from Li et al. (2016) were used as independent references. Our map proved to be more accurate, as five of the seven borehole locations were correctly identified in our map, but only 3 in the Zou map. Considering that elevation is the main factor controlling the permafrost distribution in this region, we conducted an analysis of elevation-related characteristics in this region. According to Li et al. (2016), the lower limit of permafrost occurrence in this region was around 4300 m. Our map showed greater consistency of permafrost distribution conforming to elevational characteristics than the Zou map, and the boundaries of permafrost zones of our map extended along the 4300 m contour in this region (Fig. 13a). In the Zou map, the permafrost area near the two lakes lower than 4300 m a.s.l. was overrepresented (Fig. 13b).

580

585



590 **Figure 13. Maps in the Yellow River headwaters (region d in Fig. 10) showing permafrost distributions in (a) our map and (b) the Zou map as well as (c) the spatial differences, along with (d-e) spatial and statistical elevation distributions in this region. The contour of 4300 m as the lower limit of permafrost occurrence in this region a.s.l. is shown in (a), (b), and (d). (e) Box plot showing the statistical distribution of elevations in the permafrost and SFG zones in both maps. The same notations apply as in Figure 12.**

4.6 Simulation limitations

Despite the relatively better performance of our map, some limitations of the mapping approach and room for potential improvements should be clarified. We leveraged GST observations from weather stations to estimate DDT from LST-derived thawing degree-days. However, GST sites are actually concentrated mainly in the eastern QTP, with few in the west (Fig. 1). This has a detrimental effect on the quality of the DDT estimate. Therefore, we developed multilinear regression models incorporating NDVI as a predictor. This not only properly reflects the thermal offset due to vegetation cover in the eastern QTP, where weather stations are concentrated, but also helps avoid overfitting in areas of low NDVI (<0.1) in the western QTP, where thermal offset tends to be low. It should also be noted that although the resulting DDT/DDF values still have some degree of bias, the residual errors were further reduced during the optimization process of our mapping approach by adjusting the E values to best match the simulated results with the survey-based subregion permafrost maps. We also imposed the boundary consistency as a part of the more stringent objectives during the optimization process, but the problem of parametric equifinality could not be fully solved and deserves further research, especially when working with binary classification maps (permafrost or SFG).

605 Our mapping approach relies on subregion survey maps to set up constraints on the simulation and to properly account for the influence of local factors through calibrating a model parameter. The quality and representativeness of subregion survey maps have a strong influence on the accuracy of the resulting permafrost map. In our approach, the heterogeneity of local factors in space is also represented by soil clusters. While more soil clusters can in theory better represent spatial heterogeneity,

there is a contradiction between the number of soil clusters and the effectiveness of parameter optimization. The more soil
610 clusters, the smaller the area for each soil cluster, and a smaller area would lead to a weaker constraint in the search of optimal
parameter values thus causing a stronger equifinality. Therefore, our mapping approach can benefit from more high-quality
subregion permafrost maps, which could provide more soil clusters to better represent the heterogeneous influences of local
factors.

5 Data availability

615 The new 2010 permafrost distribution map and associated data (annual DDT and DDF data derived from MODIS LST
data and soil clusters over the Qinghai-Tibet Plateau) are available on the repository hosted in figshare (Cao et al., 2022):
<https://doi.org/10.6084/m9.figshare.19642362>. Data are provided as GeoTIFF files (.tif). The sources of the datasets used for
mapping and comparison are listed in Appendix D. The related codes and sample data are accessible at
<https://github.com/nanzt/frostnumcop>.

620 6 Conclusions

This study provides a map of permafrost distribution at a spatial resolution of 1-km over the QTP in 2010 using a
modified version of the FROSTNUM/COP mapping approach. This approach estimated permafrost distribution through an
ensemble run of semi-physical model based on satellite temperature data and properly accounted for the effects of local factors
by adjusting a model parameter constrained by survey-based subregion permafrost maps. Ground surface thawing and freezing
625 indices with a relative error < 10% were obtained from interpolated all-weather MODIS LST data. The problem of parametric
equifinality was well mitigated by including boundary consistency as part of the objective function.

According to the new 2010 map, excluding glaciers and lakes, permafrost underlay about 1.086×10^6 km² (41.2% of the
total QTP area), while seasonally frozen ground covered about 1.447×10^6 km² (54.9% of the total QTP area) on the QTP in
2010. Permafrost spread continuously across the Qiangtang Plateau in the north-central QTP. The seasonally frozen ground
630 was mainly distributed in the south and east QTP. Our map also revealed that SFG were widespread in the headwaters of rivers
in the eastern QTP.

This map showed good consistency with the survey-based subregion permafrost maps, with a Kappa coefficient of 0.74,
which were much higher than that of two recently published maps (Zou et al., 2017; Wang et al., 2019). Upon validated against
72 borehole records of permafrost presence collected around 2010, we concluded that our map performed better than the Zou
635 map and Wang map. In typical regions with distinct differences, our map proved more acceptable than the Zou map based on
evidence from various aspects, including satellite imagery, PZI statistics, elevation features, and more independent boreholes
records. Our new 2010 permafrost distribution map provides accurate and fundamental information about QTP permafrost and

can thus serve as a benchmark map to calibrate/validate spatial simulations of land surface models on the QTP, as well as a historical reference for projecting future changes of QTP permafrost.

640 **Appendix A Solar-cloud-satellite geometry (SCSG) based interpolation approach**

We applied a stepwise interpolation approach to estimate missing cloudy-sky land surface temperature (LST) values of MODIS from informative samples due to the SCSG effect, by which satellite imagery records the cloudy-sky LST values of a portion of pixels. The satellite and sun have specific illumination and observation angles with respect to the ground. Based on the SCSG effect (Wang et al., 2019), each MODIS LST image was processed into four SCSG regions with one SCSG region containing known cloudy-sky LST values. A clear-sky interpolation method with the advantage of effectively handling large data gaps (Chen et al., 2020) was used to estimate clear-sky LST equivalents for every pixel in cloud-affected regions. This method estimated multiple initial estimates for each interpolated pixel by an empirically orthogonal function method based on multiple temporally proximate reference images, and then merged the initial estimates using a Bayesian approach to obtain a best estimate of clear-sky LST equivalent. Then, for each missing cloudy-sky pixel, a Multivariate Adaptive Regression Splines model (Friedman, 1991) was trained with the pixels in the specific SCSG region with known cloudy-sky LST values that were similar to that missing pixel in terms of environmental characteristics, and thereafter applied to recover missing cloudy-sky LST values (Chen et al., 2022). The fraction of pixels with null values for each image after the interpolation was small and was further interpolated by an ordinary Kriging method. This resulted in four all-weather LST values per day for all 1 km MODIS pixels. A sinusoidal method (Van Doninck et al., 2011) was applied to calculate the daily mean LSTs based on four instantaneous LST observations and the corresponding acquisition times.

660 **Appendix B Annual thawing index estimation**

We tested two methods to estimate the annual ground surface thawing index (DDT) from the raw LST-derived thawing degree-days at a MODIS pixel. One is a ‘one-year estimation’ in which a single regression model was fitted for each year. The other is a form of ‘interval-based estimation’, in which a full year was divided into 23 time intervals in line with the 16-day composite NDVI intervals each year and multilinear regression was made for each interval. Most intervals consist of 16 days, except for the last interval. The thawing degree-days over the 23 intervals per year were summed for the annual DDT.

The multilinear regression model for each time interval has the following form trained on data at meteorological sites:

$$DDT_{i,GST}' = f(DDT_{i,LST}', N_i, L) \quad (B1)$$

where $DDT_{i,GST}'$ is the ground surface thawing index for the i th interval of the year. $DDT_{i,LST}'$ is the thawing degree-days derived from the positive daily mean LST values of the pixel for the i th interval. N_i refers to the i th composite NDVI value of the pixel. L is the latitude. The index i ranges from 1 to 23. The training was based on meteorological records aggregated from all sites. The fitted functions (f) for individual intervals were then applied to the entire QTP to obtain the corrected interval

thawing degree-days of a year, before summing them for the annual DDT for that year. To minimize the risk of single-year meteorological anomalies, annual freezing index (DDF) and DDT values were averaged over the period from 2005 to 2010 and then used to drive the extended FROSTNUM model. The one-year estimation is on a yearly basis, rather than on an interval basis, following an approach similar to Eq.(B1) but without the need to sum the interval-based values.

To compare the performance of the ‘interval-based estimation’ method and the ‘one-year estimation’ method, we randomly divided the 131 weather stations into a training set (70%) and a testing set (30%) 100 times. Each time we performed both interval-based estimation and one-year estimation based on the same training set, and then assessed their prediction results using the testing set. Pearson’s correlation coefficient (r), root mean squared error (RMSE), and mean absolute error (MAE) were used as performance metrics.


By evaluating using annual in situ DDT values at QTP sites, the annual DDT values obtained by the interval-based estimation had generally lower errors and better linear correlation than the DDT values by the one-year estimation (Table B1). The ranges of metric values for the interval-based estimation were all narrower than those for the one-year estimation, indicating consistent improvements in performance across sites. This clearly demonstrates the advantage of the interval-based estimation over the one-year estimation in correcting thermal offsets between GST and LST when estimating DDT values from raw MODIS LST-derived degree-days.

Table B1. Comparison of performance between two approaches (interval-based and one-year) of estimating annual DDT from raw LST-derived thawing degree-days based on 100 trials with a random split of the training and test data sets from meteorological sites on the QTP. The values indicate the metric means from the 100 random trials, and the values in parentheses represent ranges. r : Pearson’s correlation coefficient; RMSE: root mean square error; MAE: mean absolute error.

	r	RMSE ($^{\circ}\text{C}\cdot\text{day}$)	MAE ($^{\circ}\text{C}\cdot\text{day}$)
Interval-based estimation	0.94 (0.88 - 0.97)	437 (344 - 554)	349 (252 - 458)
One-year estimation	0.92 (0.85 - 0.97)	486 (309 - 671)	368 (240 - 509)

Appendix C An instance of boundary cell

To illustrate the concept of boundary consistency introduced into the objective function, we present a simple instance of ‘boundary cell’ located in the survey-based map. In the neighboring cells (e.g. a size of 3×3) of the ‘boundary cell’, both frozen ground types (permafrost and seasonally frozen ground) and two soil clusters are present (Fig. C1). The DDT-over-DDF ratios in those cells are known as already calculated from the satellite LST data (Fig. C1). The ratios in the permafrost cells in the neighboring cells are presumably higher than those on the SFG cells to resemble a scenario where permafrost persists due to favorable local factors in areas despite unfavorable climatic conditions. This cell in the center would be considered as a ‘negative boundary cell’ if the E values associated with the two soil clusters equal 1, resulting in \bar{F}_p being smaller than \bar{F}_s (Fig. C1b). By adjusting E values accordingly, this negative boundary cell can become positive (Fig. C1c), i.e., with a larger \bar{F}_p versus \bar{F}_s . Thus, by enforcing boundary consistency, more rigorous constraints are helpful to mitigate parametric equifinality in the search for optima of E .



700 Figure C1. Illustration explaining the concept of a boundary cell and the optimization process to improve boundary consistency. Column (1) shows a boundary cell in a survey-based map whose 3×3 neighboring cells contain permafrost and seasonally frozen ground. Column (2) shows two soil clusters present in the neighboring cells. The numbers on the cells indicate the values of parameter E associated with the soil clusters of the cells. Column (3) shows the DDT-over-DDF ratios predetermined on the grid cells. In this case, permafrost cells have DDT-over-DDF ratios greater than one, indicating an unfavorable climate condition for permafrost formation. Column (4) shows the resulting ground surface frost numbers (F) for the cells. \bar{F}_p is an average of F over permafrost cells in the neighboring cells and \bar{F}_s an average of F over seasonal frost cells in the neighboring cells. A boundary cell is positive when \bar{F}_p is greater than \bar{F}_s . Row (b) indicates a negative boundary cell when the E values assume 1; and row (c) shows that this boundary cell becomes positive by adjusting the E values. Boundary consistency improves when negative boundary cells are converted to positive cells as much as possible (Row b to row c). We added boundary consistency as part of the objective function in an effort to mitigate parametric equifinality.

705

Appendix D Sources of open datasets used


710 The sources of data used in our mapping work are listed below. Daily MODIS LST/emissivity products (MOD11A1 and MYD11A1 version 6) and the NDVI product (MOD13A2) are provided by NASA and available at <https://www.earthdata.nasa.gov/>. The Shuttle Radar Topography Mission 90m digital elevation database (SRTM/DEM, version 4) (Reuter et al., 2007) is available at <https://cgiarcsi.community/data/srtm-90m-digital-elevation-database-v4-1/>. The

1-km monthly precipitation dataset for China (Peng et al., 2019) is available at <https://doi.org/10.5281/zenodo.3114194>. The
 715 500 m Daily Fractional Snow Cover Dataset Over High Asia (Qiu et al., 2017) is available at
<https://doi.org/10.11888/GlaciolGeocryol.tpe.0000016.file>. The China Data Set of Soil Properties for Land Surface Modeling
 (Shangguan et al., 2013) is available at <http://globalchange.bnu.edu.cn/research/soil2>. The China national surface weather
 stations (version 3.0) is provided by China National Meteorological Information Center and available at
<https://data.tpdc.ac.cn/en/data/52c77e9c-df4a-4e27-8e97-d363fdfce10a/>.

720 The borehole ground temperature data provided by (Zhao et al., 2021) is available at
<https://doi.org/10.11888/Geocry.tpdc.271107>. The new permafrost distribution map on the Tibetan Plateau by Zou et al. (2017)
 is available at <https://doi.org/10.11888/Geocry.tpdc.270468>. The permafrost distribution map by Wang et al. (2019) is
 available at <https://data.mendeley.com/datasets/ddj8ygdjbd/1>. Our new 2010 permafrost distribution map and associated data
 are available at (Cao et al., 2022): <https://doi.org/10.6084/m9.figshare.19642362>.

725 The above links have been checked and found accessible on 13 October 2022.

Appendix E Supplementary figures and tables



730 **Figure E1. Differences in spatial distribution of frozen ground type in the subregions between the survey-based maps and three simulated maps from (a) our study, (b) Zou et al. (2017), and (c) Wang et al. (2019). “Both P” represents areas identified as underlying permafrost in both survey-based and simulated maps; “Both SFG” represents areas identified as seasonally frozen ground (SFG) in both survey-based and simulated maps; “Simulated-SFG and Survey-P” represents areas identified as SFG in the simulated map but permafrost in the survey-based map; “Simulated-P and Survey-SFG” represents areas identified as permafrost in simulated map but SFG in the survey-based map.**

735 **Table E1. Annual average soil temperatures at three depths (3m, 6m, and 10m) in the borehole QTB15 (33.10°N, 91.90°E) within the source area of the Yangtze River. Data source: Zhao et al. (2021). Symbol ‘/’ denotes missing value.**

Year	Soil temperature at 3m (°C)	Soil temperature at 6m (°C)	Soil temperature at 10m (°C)
2006	/	/	-1.1
2007	-1.1	-1.2	-1.1
2008	/	/	-1.2
2009	-1.1	-1.1	-1.1
2010	-0.8	-1	-1
2011	-0.8	-0.9	-0.9
2012	-0.7	-0.8	-0.8
2013	-0.8	-0.8	-0.8
2014	/	/	-0.8
2015	-0.6	-0.7	-0.7
2016	-0.5	-0.6	-0.7
2017	-0.5	-0.6	-0.7
2018	-0.7	-0.8	-0.6

Author contributions.

740 Conceptualization, Z.N.; methodology, Z.N., Z.C., J.H. and Y.C.; formal analysis, Z.C.; writing—original draft, Z.C. and Z.N.; writing—review and editing, Z.N., Z.C. and Y.C.; supervision, Z.N.; funding acquisition, Z.N. and Y.Z.. All authors have read and agreed to the published version of the manuscript.

Competing interests.

The contact author has declared that neither they nor their co-authors have any competing interests.

Acknowledgements.

745 We would like to thank NASA (<https://search.earthdata.nasa.gov>), National Tibetan Plateau Data Center (<https://data.tpc.ac.cn>), the National Cryosphere Desert Data Center (<https://www.ncdc.ac.cn>), and Science Data Bank (<https://www.scidb.cn>) for providing the data. We also thank the scientists who participated in the Second Tibetan Plateau Scientific Expedition and Research campaign for their great work in collecting valuable in situ data on the QTP. We are especially grateful to Jan Beutel and Samuel Gagnon for their valuable and profound comments on the manuscript, which
750 greatly improved the quality of this paper.

Financial support.

This research has been financially supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (grant no. 2019QZKK0905-08) and National Natural Science Foundation of China (grant nos. 41971074 and 42171125). Zetao Cao also received the support from the Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX22_1583).

References

- Andresen, C.G., Lawrence, D.M., Wilson, C.J., McGuire, A.D., Koven, C., Schaefer, K., Jafarov, E., Peng, S., Chen, X. and Gouttevin, I.: Soil moisture and hydrology projections of the permafrost region-a model intercomparison, *The Cryosphere*, 14, 445 – 459, <https://doi.org/10.5194/tc-14-445-2020>, 2020.
- 760 Burke, E.J., Zhang, Y. and Krinner, G.: Evaluating permafrost physics in the Coupled Model Intercomparison Project 6 (CMIP6) models and their sensitivity to climate change, *The cryosphere*, 14, 3155 – 3174, <https://doi.org/10.5194/tc-14-3155-2020>, 2020.
- Cao, B., Gruber, S., Zheng, D. and Li, X.: The ERA5-Land soil temperature bias in permafrost regions, *The Cryosphere*, 14, 2581 – 2595, <https://doi.org/10.5194/tc-14-2581-2020>, 2020.
- 765 Cao, B., Zhang, T., Wu, Q., Sheng, Y., Zhao, L. and Zou, D.: Brief communication: Evaluation and inter-comparisons of Qinghai-Tibet Plateau permafrost maps based on a new inventory of field evidence, *The cryosphere*, 13, 511 – 519, <https://doi.org/10.5194/tc-13-511-2019>, 2019a.
- Cao, B., Zhang, T., Wu, Q., Sheng, Y., Zhao, L. and Zou, D.: Permafrost zonation index map and statistics over the Qinghai-Tibet Plateau based on field evidence, *Permafrost and Periglacial Processes*, 30, 178 – 194, <https://doi.org/10.1002/ppp.2006>, 2019b.
- 770 Cao, Z., Nan, Z., Chen, Y. and Hu, J.: A new 2010 permafrost distribution map over the Qinghai-Tibet Plateau based on subregion survey maps: a benchmark for regional permafrost modelling, *figshare* [data set], <https://doi.org/10.6084/m9.figshare.19642362>, 2022.
- Chang, Y., Lyu, S., Luo, S., Li, Z., Fang, X., Chen, B., Li, R. and Chen, S.: Estimation of permafrost on the Tibetan Plateau under current and future climate conditions using the CMIP5 data, *Int. J. Climatol.*, 38, 5659 – 5676, <https://doi.org/10.1002/joc.5770>, 2018.
- 775 Chen, J., Zhao, L., Sheng, Y., Li, J., Wu, X., Du, E., Liu, G. and Pang, Q.: Some characteristics of permafrost and its distribution in the Gaize area on the Qinghai—Tibet Plateau, China, *Arctic, Antarctic, and Alpine Research*, 48, 395 – 409, <http://dx.doi.org/10.1657/AAAR0014-023>, 2016.
- 780 Chen, Y., Nan, Z., Cao, Z., Ou, M. and Feng, K.: A stepwise framework for interpolating land surface temperature under cloudy conditions based on the solar-cloud-satellite geometry, *ISPRS Journal of Photogrammetry and Remote Sensing* (under

- review), 2022.
- Chen, Y., Nan, Z., Zhao, S. and Xu, Y.: A Bayesian Approach for Interpolating Clear-Sky MODIS Land Surface Temperatures on Areas With Extensive Missing Data, *IEEE J. Sel. Top. Appl. Earth Observ. Remote Sens.*, 14, 515 - 528, 785 <https://doi.org/10.1109/JSTARS.2020.3038188>, 2020.
- Cheng, G., Li, S., Nan, Z. and Tong, B.: Map of permafrost on the Qinghai-Tibet Plateau (1:3,000,000) (1983-1996), National Tibetan Plateau Data Center [data set], <https://doi.org/10.11888/Geocry.tpd.c.270014>, 2011.
- Cheng, G., Zhao, L., Li, R., Wu, X., Sheng, Y., Hu, G., Zou, D., Jin, H., Li, X. and Wu, Q.: Characteristic, changes and impacts of permafrost on Qinghai-Tibet Plateau (in Chinese), *Chinese Science Bulletin*, 64, 2783 - 2795, <https://doi.org/10.1360/TB-2019-0191>, 2019. 790
- Cohen, J.: A coefficient of agreement for nominal scales, *Educ. Psychol. Meas.*, 20, 37 - 46, <https://doi.org/10.1177/001316446002000104>, 1960.
- Dai, L., Che, T., Xie, H. and Wu, X.: Estimation of snow depth over the Qinghai-Tibetan Plateau based on AMSR-E and MODIS data, *Remote Sens.*, 10, 1989, <https://doi.org/10.3390/rs10121989>, 2018.
- 795 Debolskiy, M.V., Nicolsky, D.J., Hock, R. and Romanovsky, V.E.: Modeling Present and Future Permafrost Distribution at the Seward Peninsula, Alaska, *Journal of geophysical research: Earth surface*, 125, <https://doi.org/10.1029/2019JF005355>, 2020.
- Dobinski, W.: Permafrost, *Earth-Sci. Rev.*, 108, 158 - 169, <https://doi.org/10.1016/j.earscirev.2011.06.007>, 2011.
- Friedman, J.H.: Multivariate adaptive regression splines, *The annals of statistics*, 19, 1 - 67, 800 <https://www.jstor.org/stable/2241837>, 1991.
- Gisnås, K., Etzelmüller, B., Lussana, C., Hjort, J., Sannel, A.B.K., Isaksen, K., Westermann, S., Kuhry, P., Christiansen, H.H. and Frampton, A.: Permafrost map for Norway, Sweden and Finland, *Permafrost and periglacial processes*, 28, 359 - 378, <https://doi.org/10.1002/ppp.1922>, 2017.
- Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation, *The Cryosphere*, 6, 221 - 233, 805 <https://doi.org/10.5194/tc-6-221-2012>, 2012.
- Guo, D. and Wang, H.: Simulation of permafrost and seasonally frozen ground conditions on the Tibetan Plateau, 1981-2010, *Journal of geophysical research: Atmospheres*, 118, 5216 - 5230, <https://doi.org/10.1002/jgrd.50457>, 2013.
- Guo, W., Liu, S., Xu, J., Wu, L., Shangguan, D., Yao, X., Wei, J., Bao, W., Yu, P. and Liu, Q.: The second Chinese glacier inventory: data, methods and results, *J. Glaciol.*, 61, 357 - 372, <https://doi.org/10.3189/2015JoG14J209>, 2015.
- 810 Hachem, S., Duguay, C.R. and Allard, M.: Comparison of MODIS-derived land surface temperatures with ground surface and air temperature measurements in continuous permafrost terrain, *The Cryosphere*, 6, 51 - 69, <https://doi.org/10.5194/tc-6-51-2012>, 2012.
- Hu, G., Zhao, L., Li, R., Wu, X., Wu, T., Zhu, X., Pang, Q., Yue Liu, G., Du, E. and Zou, D.: Simulation of land surface heat

- fluxes in permafrost regions on the Qinghai-Tibetan Plateau using CMIP5 models, *Atmos. Res.*, 220, 155 – 168, 815 <https://doi.org/10.1016/j.atmosres.2019.01.006>, 2019.
- Hu, G., Zhao, L., Wu, X., Li, R., Wu, T., Su, Y. and Hao, J.: Evaluation of reanalysis air temperature products in permafrost regions on the Qinghai-Tibetan Plateau, *Theor. Appl. Climatol.*, 138, 1457 – 1470, <https://doi.org/10.1007/s00704-019-02888-8>, 2019.
- Hu, J., Zhao, S., Nan, Z., Wu, X., Sun, X. and Cheng, G.: An effective approach for mapping permafrost in a large area using 820 subregion maps and satellite data, *Permafrost and Periglacial Processes*, 31, 548 – 560, <https://doi.org/10.1002/ppp.2068>, 2020.
- Huang, R., Huang, J., Zhang, C., Ma, H., Zhuo, W., Chen, Y., Zhu, D., Wu, Q. and Mansaray, L.R.: Soil temperature estimation at different depths, using remotely-sensed data, *J. Integr. Agric.*, 19, 277 – 290, [https://doi.org/10.1016/S2095-3119\(19\)62657-2](https://doi.org/10.1016/S2095-3119(19)62657-2), 2020.
- Huang, Z.: Extensions to the k-means algorithm for clustering large data sets with categorical values, *Data Min. Knowl. 825 Discov.*, 2, 283 – 304, <https://doi.org/10.1023/A:1009769707641>, 1998.
- Ji, H., Nan, Z., Hu, J., Zhao, Y. and Zhang, Y.: On the Spin-up Strategy for Spatial Modeling of Permafrost Dynamics: a Case Study on the Qinghai-Tibet Plateau, *J. Adv. Model. Earth Syst.*, e2021M – e2750M, <https://doi.org/10.1029/2021MS002750>, 2022.
- Jin, H., Luo, D., Wang, S., Lü L. and Wu, J.: Spatiotemporal variability of permafrost degradation on the Qinghai-Tibet 830 Plateau, *Sci. Cold Arid Reg.*, 3, 281 – 305, <https://doi.org/10.3724/SP.J.1226.2011.00281>, 2011.
- Koven, C.D., Riley, W.J. and Stern, A.: Analysis of permafrost thermal dynamics and response to climate change in the CMIP5 Earth System Models, *J. Clim.*, 26, 1877 – 1900, <https://doi.org/10.1175/JCLI-D-12-00228.1>, 2013.
- Kuhn, M. and Johnson, K.: *Applied predictive modeling*, Springer, 2013.
- Li, J., Sheng, Y., Wu, J., Feng, Z., Ning, Z., Hu, X. and Zhang, X.: Landform-related permafrost characteristics in the source 835 area of the Yellow River, eastern Qinghai-Tibet Plateau, *Geomorphology*, 269, 104 – 111, <https://doi.org/10.1016/j.geomorph.2016.06.024>, 2016.
- Li, X., Cheng, G., Jin, H., Kang, E., Che, T., Jin, R., Wu, L., Nan, Z., Wang, J. and Shen, Y.: Cryospheric change in China, *Glob. Planet. Change*, 62, 210 – 218, <https://doi.org/10.1016/j.gloplacha.2008.02.001>, 2008.
- Li, Z., Li, Z., Feng, Q., Zhang, B., Gui, J., Xue, J. and Gao, W.: Runoff dominated by supra-permafrost water in the source 840 region of the Yangtze River using environmental isotopes, *J. Hydrol.*, 582, 124506, <https://10.1016/j.jhydrol.2019.124506>, 2020.
- Li, Z., Zhao, L., Liu, G., Zou, D., Wang, L., Yang, B., Du, E., Hu, G., Zhou, H., Wang, C., Xin, Z., Zhao, J., Yin, X., Chi, H., Tan, C. and Chen, W.: Analysis of soil moisture content in the active layer in the permafrost region of the Yangtze River source in frozen season (in Chinese), *Journal of Glaciology and Geocryology*, 40, 1 – 13, [https://doi.org/10.7522/j.issn.1000-0240.2022.0001](https://doi.org/10.7522/j.issn.1000-845 0240.2022.0001), 2022.

- Luo, D., Jin, H., Marchenko, S.S. and Romanovsky, V.E.: Difference between near-surface air, land surface and ground surface temperatures and their influences on the frozen ground on the Qinghai-Tibet Plateau, *Geoderma*, 312, 74 - 85, <https://doi.org/10.1016/j.geoderma.2017.09.037>, 2018.
- 850 Marcer, M., Bodin, X., Brenning, A., Schoeneich, P., Charvet, R. and Gottardi, F.: Permafrost favorability index: spatial modeling in the French Alps using a rock glacier inventory, *Front. Earth Sci.*, 5, 105, <https://doi.org/10.3389/feart.2017.00105>, 2017.
- Mekonnen, Z.A., Riley, W.J., Grant, R.F. and Romanovsky, V.E.: Changes in precipitation and air temperature contribute comparably to permafrost degradation in a warmer climate, *Environ. Res. Lett.*, 16, 24008, <https://doi.org/10.1186/s40645-020-00366-8>, 2021.
- 855 Mu, C., Abbott, B.W., Norris, A.J., Mu, M., Fan, C., Chen, X., Jia, L., Yang, R., Zhang, T. and Wang, K.: The status and stability of permafrost carbon on the Tibetan Plateau, *Earth-Sci. Rev.*, 211, 103433, <https://doi.org/10.1016/j.earscirev.2020.103433>, 2020.
- Nan, Z., Li, S. and Liu, Y.: Mean annual ground temperature distribution on the Tibetan Plateau: permafrost distribution mapping and further application (in Chinese), *Journal of Glaciology and Geocryology*, 24, 142 - 148, 860 <http://www.bcdt.ac.cn/CN/Y2002/V24/I2/142>, 2002.
- National Meteorological Information Center: Daily meteorological dataset of basic meteorological elements of China National Surface Weather Station (V3.0) (1951-2010), National Tibetan Plateau Data Center [data set], <https://data.tpdac.ac.cn/en/data/52c77e9c-df4a-4e27-8e97-d363fdfce10a/>, 2019.
- Nelson, F.E. and Outcalt, S.I.: A Computational Method for Prediction and Regionalization of Permafrost, *Arctic and Alpine Research*, 19, 279 - 288, <https://doi.org/10.1080/00040851.1987.12002602>, 1987.
- 865 Ni, J., Wu, T., Zhu, X., Hu, G., Zou, D., Wu, X., Li, R., Xie, C., Qiao, Y. and Pang, Q.: Simulation of the present and future projection of permafrost on the Qinghai-Tibet Plateau with statistical and machine learning models, *Journal of Geophysical Research: Atmospheres*, 126, e2020J - e33402J, <https://doi.org/10.1002/essoar.10503593.1>, 2021.
- Niu, F. and Yin, G.: Map of the frozen soil in the Tibetan Plateau (2003), National Tibetan Plateau Data Center [data set], 870 <https://doi.org/10.11888/GlaciolGeocryo.tpe.00000048.file>, 2018.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B. and Kholodov, A.: Northern Hemisphere permafrost map based on TTOP modelling for 2000-2016 at 1 km² scale, *Earth-Sci. Rev.*, 193, 299 - 316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.
- Peng, S., Ding, Y., Liu, W. and Li, Z.: 1 km monthly temperature and precipitation dataset for China from 1901 to 2017, *Earth Syst. Sci. Data*, 11, 1931 - 1946, <https://doi.org/10.5194/essd-11-1931-2019>, 2019.
- 875 Qin, Y., Wu, T., Zhao, L., Wu, X., Li, R., Xie, C., Pang, Q., Hu, G., Qiao, Y. and Zhao, G.: Numerical modeling of the active layer thickness and permafrost thermal state across Qinghai - Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 122, 11, 604 - 611, 620, <https://doi.org/10.1002/2017JD026858>, 2017.

- Qin, Y., Zhang, P., Liu, W., Guo, Z. and Xue, S.: The application of elevation corrected MERRA2 reanalysis ground surface
880 temperature in a permafrost model on the Qinghai-Tibet Plateau, *Cold Reg. Sci. Tech.*, 175, 103067,
<https://doi.org/10.1016/j.coldregions.2020.103067>, 2020.
- Qiu, Y., Wang, X., Han, L., Chang, L. and Shi, L.: Daily fractional snow cover dataset over High Asia (in Chinese), *Science
Data Bank*, 2, 1 - 13, <https://doi.org/10.11922/csdata.170.2017.0146>, 2017.
- Ran, Y., Li, X., Cheng, G., Zhang, T., Wu, Q., Jin, H. and Jin, R.: Distribution of permafrost in China: an overview of existing
885 permafrost maps, *Permafrost and Periglacial Processes*, 23, 322 - 333, <https://doi.org/10.1002/ppp.1756>, 2012.
- Reuter, H.I., Nelson, A. and Jarvis, A.: An evaluation of void - filling interpolation methods for SRTM data, *Int. J. Geogr. Inf.
Sci.*, 21, 983 - 1008, <https://doi.org/10.1080/13658810601169899>, 2007.
- Shangguan, W., Dai, Y., Liu, B., Zhu, A., Duan, Q., Wu, L., Ji, D., Ye, A., Yuan, H. and Zhang, Q.: A China data set of soil
properties for land surface modeling, *J. Adv. Model. Earth Syst.*, 5, 212 - 224, <https://doi.org/10.1002/jame.20026>, 2013.
- 890 Shi, Y. and Mi, D.: Map of snow, ice, and frozen ground in China (1988), National Tibetan Plateau Data Center [data set],
<https://doi.org/10.3972/westdc.020.2013.db>, 2013.
- Shi, Y., Niu, F., Yang, C., Che, T., Lin, Z. and Luo, J.: Permafrost presence/absence mapping of the Qinghai-Tibet Plateau
based on multi-source remote sensing data, *Remote Sens.*, 10, 309, <https://doi.org/10.3390/rs10020309>, 2018.
- Song, C., Wang, G., Mao, T., Dai, J. and Yang, D.: Linkage between permafrost distribution and river runoff changes across
895 the Arctic and the Tibetan Plateau, *Science China Earth Sciences*, 63, 292 - 302, <https://doi.org/10.1007/s11430-018-9383-6>,
2020.
- Van Doninck, J., Peters, J., De Baets, B., De Clercq, E.M., Ducheyne, E. and Verhoest, N.E.C.: The potential of multitemporal
Aqua and Terra MODIS apparent thermal inertia as a soil moisture indicator, *Int. J. Appl. Earth Obs. Geoinf.*, 13, 934 - 941,
<https://doi.org/10.1016/j.jag.2011.07.003>, 2011.
- 900 Wan, Z.: New refinements and validation of the MODIS Land-Surface Temperature/Emissivity products, *Remote Sens.
Environ.*, 112, 59 - 74, <https://doi.org/10.1016/j.rse.2006.06.026>, 2008.
- Wang, D., Tan, D. and Liu, L.: Particle swarm optimization algorithm: an overview, *Soft Comput.*, 22, 387 - 408,
<https://doi.org/10.1007/s00500-016-2474-6>, 2018.
- Wang, L., Du, Z., Wei, Z., Xu, Q., Feng, Y., Lin, P., Lin, J., Chen, S., Qiao, Y. and Shi, J.: High methane emissions from
905 thermokarst lakes on the Tibetan Plateau are largely attributed to ebullition fluxes, *Sci. Total Environ.*, 801, 149692,
<https://doi.org/10.1016/j.scitotenv.2021.149692>, 2021.
- Wang, S., Niu, F., Chen, J. and Dong, Y.: Permafrost research in China related to express highway construction, *Permafrost
and Periglacial Processes*, 31, 406 - 416, <https://doi.org/10.1002/ppp.2053>, 2020.
- Wang, T.: 1:4 million map of the Glaciers, Frozen Ground and Deserts in China (2006), National Tibetan Plateau Data Center
910 [data set], <https://doi.org/10.3972/westdc.015.2013.db>, 2013.

- Wang, T., Shi, J., Ma, Y., Husi, L., Comyn Platt, E., Ji, D., Zhao, T. and Xiong, C.: Recovering land surface temperature under cloudy skies considering the solar - cloud - satellite geometry: Application to MODIS and Landsat - 8 data, *Journal of Geophysical Research: Atmospheres*, 124, 3401 - 3416, <https://doi.org/10.1029/2018JD028976>, 2019.
- Wang, T., Wu, T., Wang, P., Li, R., Xie, C. and Zou, D.: Spatial distribution and changes of permafrost on the Qinghai-Tibet Plateau revealed by statistical models during the period of 1980 to 2010, *Sci. Total Environ.*, 650, 661 - 670, <https://doi.org/10.1016/j.scitotenv.2018.08.398>, 2019.
- Wang, T., Yang, D., Fang, B., Yang, W., Qin, Y. and Wang, Y.: Data-driven mapping of the spatial distribution and potential changes of frozen ground over the Tibetan Plateau, *Sci. Total Environ.*, 649, 515 - 525, <https://doi.org/10.1016/j.scitotenv.2018.08.369>, 2019.
- 920 Wang, W., Rinke, A., Moore, J.C., Cui, X., Ji, D., Li, Q., Zhang, N., Wang, C., Zhang, S. and Lawrence, D.M.: Diagnostic and model dependent uncertainty of simulated Tibetan permafrost area, *The Cryosphere*, 10, 287 - 306, <https://doi.org/10.5194/tc-10-287-2016>, 2016.
- Wang, Z., Wang, Q., Zhao, L., Wu, X., Yue, G., Zou, D., Nan, Z., Liu, G., Pang, Q. and Fang, H.: Mapping the vegetation distribution of the permafrost zone on the Qinghai-Tibet Plateau, *J Mt. Sci.*, 13, 1035 - 1046, [https://doi.org/10.1007/s11629-](https://doi.org/10.1007/s11629-015-3485-y)
- 925 015-3485-y, 2016.
- Westermann, S., Langer, M. and Boike, J.: Systematic bias of average winter-time land surface temperatures inferred from MODIS at a site on Svalbard, Norway, *Remote Sens. Environ.*, 118, 162 - 167, <https://doi.org/10.1016/j.rse.2011.10.025>, 2012.
- Wu, Q. and Zhang, T.: Recent permafrost warming on the Qinghai - Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 113, 1 - 22, <https://doi.org/10.1029/2007JD009539>, 2008.
- 930 Wu, X., Nan, Z., Zhao, S., Zhao, L. and Cheng, G.: Spatial modeling of permafrost distribution and properties on the Qinghai-Tibet Plateau, *Permafrost and periglacial processes*, 29, 86 - 99, <https://doi.org/10.1002/ppp.1971>, 2018.
- Xu, S. and Cheng, J.: A new land surface temperature fusion strategy based on cumulative distribution function matching and multiresolution Kalman filtering, *Remote Sens. Environ.*, 254, 112256, <https://doi.org/10.1016/j.rse.2020.112256>, 2021.
- Xu, Y., Shen, Y. and Wu, Z.: Spatial and temporal variations of land surface temperature over the Tibetan Plateau based on harmonic analysis, *Mt. Res. Dev.*, 33, 85 - 94, <https://doi.org/10.1659/MRD-JOURNAL-D-12-00090.1>, 2013.
- 935 Yang, M., Wang, X., Pang, G., Wan, G. and Liu, Z.: The Tibetan Plateau cryosphere: Observations and model simulations for current status and recent changes, *Earth-Sci. Rev.*, 190, 353 - 369, <https://doi.org/10.1016/j.earscirev.2018.12.018>, 2019.
- Yin, G., Niu, F., Lin, Z., Luo, J. and Liu, M.: Data-driven spatiotemporal projections of shallow permafrost based on CMIP6 across the Qinghai-Tibet Plateau at 1 km² scale, *Adv. Clim. Chang. Res.*, 12, 814 - 827, <https://doi.org/10.1016/j.accre.2021.08.009>, 2021.
- 940 Yokohata, T., Saito, K., Ito, A., Ohno, H., Tanaka, K., Hajima, T. and Iwahana, G.: Future projection of greenhouse gas emissions due to permafrost degradation using a simple numerical scheme with a global land surface model, *Prog. Earth Planet.*

- Sci., 7, 1 – 14, <https://doi.org/10.1186/s40645-020-00366-8>, 2020.
- Zhang, G., Nan, Z., Hu, N., Yin, Z., Zhao, L., Cheng, G. and Mu, C.: Qinghai - Tibet Plateau permafrost at risk in the late 21st century, *Earth's Future*, <https://doi.org/10.1029/2022EF002652>, 2022.
- Zhang, G., Nan, Z., Wu, X., Ji, H. and Zhao, S.: The role of winter warming in permafrost change over the Qinghai - Tibet Plateau, *Geophys. Res. Lett.*, 46, 11261 – 11269, <https://doi.org/10.1029/2019GL084292>, 2019.
- Zhang, G., Nan, Z., Yin, Z. and Zhao, L.: Isolating the Contributions of Seasonal Climate Warming to Permafrost Thermal Responses Over the Qinghai-Tibet Plateau, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2021JD035218>, 2021.
- Zhang, G., Nan, Z., Zhao, L., Liang, Y. and Cheng, G.: Qinghai-Tibet Plateau wetting reduces permafrost thermal responses to climate warming, *Earth Planet. Sci. Lett.*, 562, 116858, <https://doi.org/10.1016/j.epsl.2021.116858>, 2021.
- Zhang, G., Yao, T., Piao, S., Bolch, T., Xie, H., Chen, D., Gao, Y., O'Reilly, C.M., Shum, C.K. and Yang, K.: Extensive and drastically different alpine lake changes on Asia's high plateaus during the past four decades, *Geophys. Res. Lett.*, 44, 252 – 260, <https://doi.org/10.1002/2016GL072033>, 2017.
- Zhang, L., Xia, X., Liu, S., Zhang, S., Li, S., Wang, J., Wang, G., Gao, H., Zhang, Z. and Wang, Q.: Significant methane ebullition from alpine permafrost rivers on the East Qinghai - Tibet Plateau, *Nat. Geosci.*, 13, 349 – 354, <https://doi.org/10.1038/s41561-020-0571-8>, 2020.
- Zhang, X., Zhou, J., Liang, S. and Wang, D.: A practical reanalysis data and thermal infrared remote sensing data merging (RTM) method for reconstruction of a 1-km all-weather land surface temperature, *Remote Sens. Environ.*, 260, 112437, <https://doi.org/10.1016/j.rse.2021.112437>, 2021.
- Zhao, H., Liu, S., Dong, S., Su, X., Wang, X., Wu, X., Wu, L. and Zhang, X.: Analysis of vegetation change associated with human disturbance using MODIS data on the rangelands of the Qinghai-Tibet Plateau, *The Rangeland Journal*, 37, 77 – 87, <https://doi.org/10.1071/RJ14061>, 2015.
- Zhao, L., Wu, T., Xie, C., Li, R., Wu, X., Yao, J., Yue, G. and Xiao, Y.: Support geoscience research, environmental management, and engineering construction with investigation and monitoring on permafrost in the Qinghai-Tibet Plateau, China (in Chinese), *Bull Chin Acad Sci*, 32, 1159 – 1168, <https://doi.org/10.16418/j.issn.1000-3045.2017.10.015>, 2017.
- Zhao, L., Zou, D., Hu, G., Du, E., Pang, Q., Xiao, Y., Li, R., Sheng, Y., Wu, X. and Sun, Z.: Changing climate and the permafrost environment on the Qinghai-Tibet (Xizang) plateau, *Permafrost and Periglacial Processes*, 31, 396 – 405, <https://doi.org/10.1002/ppp.2056>, 2020.
- Zhao, L., Zou, D., Hu, G., Wu, T., Du, E., Liu, G., Xiao, Y., Li, R., Pang, Q. and Qiao, Y.: A synthesis dataset of permafrost thermal state for the Qinghai - Tibet (Xizang) Plateau, China, *Earth Syst. Sci. Data*, 13, 4207 – 4218, <https://doi.org/10.5194/essd-13-4207-2021>, 2021.
- Zhao, S.P., Nan, Z.T., Huang, Y.B. and Zhao, L.: The application and evaluation of simple permafrost distribution models on

- 975 the Qinghai-Tibet Plateau, *Permafrost and Periglacial Processes*, 28, 391 – 404, <https://doi.org/10.1002/ppp.1939>, 2017.
- Zou, D., Zhao, L., Sheng, Y., Chen, J., Hu, G., Wu, T., Wu, J., Xie, C., Wu, X. and Pang, Q.: A new map of permafrost distribution on the Tibetan Plateau, *The Cryosphere*, 11, 2527 – 2542, <https://doi.org/10.5194/tc-11-2527-2017>, 2017.
- Zou, D., Zhao, L., Wu, T., Wu, X., Pang, Q. and Wang, Z.: Modeling ground surface temperature by means of remote sensing data in high-altitude areas: test in the central Tibetan Plateau with application of moderate-resolution imaging spectroradiometer Terra/Aqua land surface temperature and ground-based infrared radiometer, *Journal of Applied Remote Sensing*, 8, 83516, <https://doi.org/10.1117/1.JRS.8.083516>, 2014.
- 980