

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

Reanalyses of Maskelyne’s Tidal Data at St. Helena in 1761

Philip L. Woodworth¹ and John M. Vassie¹

¹ National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool L3 5DA, United Kingdom

Correspondence to: P.L. Woodworth (plw@noc.ac.uk)

Abstract. The construction of an electronic data set of the tidal measurements made at St. Helena in 1761 by Nevil Maskelyne is described. These data were first analysed by Cartwright (1971, 1972) in papers which have importance within studies of changing tides. However, Cartwright’s data files were never archived for the benefit of other researchers, demonstrating that ‘old data’ at risk can sometimes take the form of electronic rather than paper records. In the present paper, the newly digitised Maskelyne data have been reanalysed by several techniques in order to obtain an updated impression of whether the tide has changed at that location in over two and a half centuries. Our main conclusion, consistent with that of Cartwright, is that the major tidal constituent (M2) has changed little. However, the results of the various techniques demonstrate how difficult it is to obtain reliable conclusions for the smaller constituents.

Keywords

Data reanalysis; Tidal science; Changing ocean tides; St. Helena; Nevil Maskelyne

28 **1 Introduction**

29
30 Almost fifty years ago, David Cartwright investigated whether the ocean tide at St. Helena had changed since 1761,
31 with his findings reported in two papers (Cartwright, 1971,1972). This was an interesting piece of work at the time,
32 but has gained additional importance since then, given our present understanding that the ocean tide has been
33 changing in recent decades in many parts of the world (Woodworth, 2010; ~~Haigh et al., 2020~~). There are many
34 possible reasons for such changes in the tide, of which change in water depth due to climate change is the most
35 obvious (Haigh et al., 2020). Consequently, there is a lot to be learned on this subject, with a re-examination of
36 historical data being especially important.
37

38 The comparison of the modern and historical tides at St. Helena was made using a year of high and low water data
39 at Ascension Island in 1958-9 in order to provide a reference tide for the use of the response method in analyses of
40 the short tidal records available from St. Helena. Ascension lies 1300 km northwest of St. Helena but can be
41 considered ‘nearby’ in the context of the ocean tide response to astronomical forcing. Cartwright himself had been
42 one of the developers of the response method (Munk and Cartwright, 1966). The modern tide at St. Helena was
43 determined by means of Cartwright’s own tidal measurements there for 39 days in 1969, while the historical tide
44 was calculated from measurements made by Nevil Maskelyne for over a month in 1761, both data sets analysed
45 using the response method and the Ascension reference record.
46

47 A listing of Maskelyne’s tidal measurements at St. Helena is given at the end of Maskelyne (1762) as shown in
48 Figure 1. Although Cartwright must have spent a lot of time putting these measurements into electronic form, it is
49 impossible for anyone to readily repeat his work now because he did not lodge his files in a data centre. Back in
50 1971 there was no culture of depositing data sets in centres such as the British Oceanographic Data Centre (BODC)
51 or even of providing Supplementary Material for a paper.
52

53 ‘Data reanalysis’ comes into this discussion because we wanted to see if we would obtain the same findings as
54 Cartwright, should the Maskelyne data be made available electronically once again, especially given the present
55 interest in changing tides (Haigh et al., 2020). As a result, a summary of our own conclusions on the tides at St.
56 Helena, based on analyses of Maskelyne’s data set, compared to those of Cartwright is given below.
57
58

1761.

Day of observation.	Apparent time	The height in divisions and tenths.	No. of observations.	N. B. The height is set down according to the divisions on the post, each of which is 3 inches.
Nov. 24 12.	h ' A. M.			
	8 56 A. M.	1 $\frac{1}{8}$	L 17	
	9 32 A. M.	1, 8	34	
	9 52 A. M.	2 $\frac{1}{4}$	18	
	10 37 A. M.	4, 0	16	
	0 58 P. M.	11	16	
	2 27 P. M.	12, 4	H 55	
	3 29 P. M.	12	18	
	3 49 P. M.	11, 3	16	
	4 32 P. M.	9, 4	20	
Nov. 25 13. High surf.	8 26 P. M.	1	L	
	9 54 P. M.	2 $\frac{1}{2}$		
	6 45 A. M.	4, 8	38	
	7 24 A. M.	3, 3	36	
	7 57 A. M.	2, 5	12	
	9 21 A. M.	1, 5	L 12	
	9 32 A. M.	2, 0	20	
	1 26 P. M.	10, 9	28	
	3 21 P. M.	13, 3	H 22	
	3 54 P. M.	12, 8	26	
6 35 P. M.	6, 3	20		

59
60
61
62
63

Figure 1. The start of the table of measurements of sea level at St. Helena in 1761 to be found at the end of Maskelyne's original paper. Source: Maskelyne (1762) reproduced with permission of the Royal Society.

2 The Maskelyne Data Set

The first person to make tidal measurements at St. Helena was the Rev. Nevil Maskelyne, Astronomer Royal 1765-1811. Maskelyne's reason for visiting the island was to observe the time of the transit of Venus on 6 June 1761, an objective that was prevented by the cloudy weather. His exercise in tide recording, from 12 November to 22 December 1761, must have compensated somewhat for the failure of that main objective.

As explained by Cartwright et al. (2017), knowledge of the tide was still rudimentary in 1761. Newton had shown that the main characteristics of the ocean tide followed from his gravitational theory. However, there was a lack of observational data from which one could learn more about tidal dynamics, especially from remote island locations. St. Helena was certainly remote but had one major drawback, in that its exposure to swell waves made it difficult to observe the modest tidal rise and fall (the mean tidal range is approximately 1 metre).

Maskelyne managed to largely eliminate the effect of swell waves by reading from a graduated vertical staff many times over the course of a few minutes and recording the average reading: "I therefore generally made 40 or 50 observations, and sometimes more than 100, if the rise and fall of the water seemed very irregular". The resulting averages traced out a smooth tidal curve, and simultaneous readings by Maskelyne and his assistant Charles Mason agreed consistently to better than half an inch (12 mm). (Mason is better known as the leading surveyor of the Mason-Dixon Line and for the measurement of a degree of latitude in North America.) Their observations were made for all states of the tide between 12 November and 22 December 1761, except for a short interruption when the swell damaged the vertical staff.¹

The resulting measurements can be found in a table at the end of Maskelyne (1762), and the present exercise involved the typing of those numbers into a single ASCII computer file. That file, called 'maskelyne_data', has a format which is essentially the same as Figure 1. It consists of spot measurements of heights at particular times, all of which come from Maskelyne's table apart from a couple of errors which Cartwright pointed out in a footnote at the bottom of Cartwright (1971, p617). In these cases, we have used the Cartwright numbers instead. The times corresponding to each tide level are given as hours and minutes and the levels themselves are in 'divisions and tenths' where one division is 3 inches. Also shown is the number of separate instantaneous estimates of height made rapidly by Maskelyne over several minutes, averaged and recorded to the nearest minute. There are some days which are almost complete, with measurements of tide level around the clock. However, as pointed out in Cartwright (1971), in the latter part of the data set the measurements are increasingly in daylight hours (Figure 2). Cartwright (1971) should be consulted for further explanation of how Maskelyne came to make his measurements and for additional details about them, while the header of 'maskelyne_data' contains more detailed information about the file itself. Section 6 mentions the locations from which this new data file might be obtained.

Before comparing our findings with those of Cartwright in the next sections, we can point to several remarks in his papers to do with the data that we have concerns about:

- On page 617 of Cartwright (1971) he says "Each sea level [measured by Maskelyne] was the mean of up to 100 or more observations at different states of the swell over several minutes, and recorded against the mean time to the nearest 1/4 min."

One can see from Maskelyne's table that there were certainly some sea level measurements obtained from over 100 separate observations. However, that was the case for only 10 of the measurements out of 478 in total. Normally, there were only a few 10s of observations. Therefore, it seems that this sentence of Cartwright (1971) over-states the quality of Maskelyne's data somewhat.


In addition, it is not clear where the "mean time to the nearest 1/4-min" statement came from. Maskelyne (1762, p589-590) says "I always looked at my watch before I began to note the height of the water, and looked at it again when I had finished the experiment; the medium of the two times I set down as the true time of the observation. The times set down are exact to the minute."

¹[The It is certainly possible to make visual measurements of 'still water level' to that accuracy using a tide staff \(tide pole\) in relatively calm conditions, by averaging visually over the incidence of swell waves. Similar tide poles continue to be used today to check the performance of modern tide gauges \(IOC, 2016\). One has at this point to assume that Maskelyne's staff was vertical, with no scale errors in the conversion of its 'divisions and tenths' into modern units. Unfortunately, the zero of Maskelyne's staff was not related to a land benchmark, so the historical data are not useful to studies of long-term sea level change.](#)

- 117 • The first footnote on Cartwright (1971, p617) says “I was unable to detect any sensible change in datum
118 after the pole was re-set.”
119

120 This is a reference to the entry in Maskelyne’s table showing that the tide pole was damaged on 3 December and
121 that Maskelyne put it back on 5 December saying “The post was set up again as near the former height as could be
122 judged.” However, as discussed below, there is some evidence for a datum shift of about 2 or 3 cm either side of this
123 event, and this datum shift impacts significantly on determination of changes in the diurnal tides in particular.
124

- 125 • There are also some additional minor inconsistencies. For example, there were 40 days with measurements
126 as stated in Cartwright (1971, p617). They spanned 41 days (12 Nov – 22 Dec) with a day with no data on
127 4 December. However, in Cartwright (1972, p337) he implies that the span was 42 days. And in his table
128 on p338, he says he used only 39 days of data which ‘exclude certain lacunae [gaps] in the data’. But the
129 only gap was the 4 December one. If he used only 39 days, it is not obvious which day of the 40 he dropped;
130 there were many days shown as having high surf which he might have been rejected otherwise.
131
- 132 • Finally, p617 contains a long paragraph concerning what the longitude was of the clock used by Maskelyne
133 and Mason. While this paragraph is interesting, it does not really seem relevant to the present comparison
134 of historical and modern tides. Even if there was an uncertainty of 0.1° in longitude as he suggests, that
135 propagates into uncertainties of only $0.1/0.2^\circ$ in the calculation of the phase lags of diurnal/semidiurnal
136 tides which is well within any realistic uncertainty in a comparison.
137



138
139 **Figure 2.** The recorded (local) times of Maskelyne’s measurements. Some measurements were made around the
140 clock in the earlier part of the data set. However, they can be seen to be restricted to daylight hours in the latter part.
141

3 Data Reanalyses

3.1 Time Corrections to Maskelyne's Measurements

Before a comparison can be made between the modern and historical tides, it is necessary to convert Maskelyne's times, ~~which are local apparent (sun dial) times, to Greenwich Mean Time (GMT) in a similar way to Cartwright (1971)~~ to Greenwich Mean Time (GMT), in a similar way to Cartwright (1971). Maskelyne's times were local apparent (sun dial) times, derived from measurements of local meridian transits of the sun using a transit telescope, with the aid of a clock to interpolate times between measurements. As Cartwright (1971) explains, the precision of these measurements could not be bettered since the work was undertaken alongside frequent astronomical checks as part of a study by Maskelyne of the going of clocks in different latitudes.

First, we adjusted for longitude using the same value (5.718 °W) used for the study of Manuel Johnson's data at St. Helena in 1826-1827 (Cartwright et al., 2017).² All such historical measurements of the tide at St. Helena have been made at the landing steps in Jamestown Bay, as they continue to be made to this day. Then, we corrected for the differences between local apparent and mean times due to the sun not always being on the equator, the difference being called the equation of time (EOT), using EOT values from the 1795 Nautical Almanac.

We used EOT values from the 1795 edition of the Almanac because we happened to have a copy available. The Nautical Almanac was first published in 1767 and so it did not exist in 1761, so we had to use EOT values for not too different a later year. Fortunately, 1761 and 1795 were both non-leap years and their EOT values should be very similar as any changes in the EOT over only three decades will be negligible; as a confirmation, we checked that the small differences in the EOT over two centuries between 1795 and 1991 in the tables we used were the same as those shown in Hughes et al. (1989). The resulting total time corrections for each day (i.e. longitude and EOT) were very similar to those listed in Table 5 of Cartwright (1971).

3.2 Initial Comparison of Modern and Historical Tides

In order to make our own comparison of the modern and historical tides, we initially made use of a set of 62 tidal constants derived from a record of sub-surface pressure (SSP) at St. Helena spanning one year (October 1995 - October 1996). This set is called STHL4. It contains constants for 5 long-period, 18 diurnal, 20 semidiurnal and 19 higher-frequency constituents. Although the record is of SSP and not real sea level, there is not much difference between the constants that would be obtained from the two as the product of density and acceleration due to gravity (ρg) happens to be almost exactly 1.0 at St. Helena. This was confirmed by comparison to another set of constants for real sea level during 1993-2006 computed by Richard Ray for the Manuel Johnson study (Cartwright et al., 2017).

However, there was the expected difference for S2 (the main solar semidiurnal tide) which has an amplitude of 10.25 cm in STHL4 and 11.39 cm in Ray's set. That can be explained by the S2 air tide at St. Helena having an amplitude of 1.1 mbar and a phase lag which is almost opposite that of sea level (Ray, 1998). As a result, S2 in SSP has an amplitude about 1 cm smaller than in sea level.

Therefore, for present purposes we defined a new set of constants based on STHL4 but with those for S2 taken from Ray's set. For consistency, we also replaced those for S1 by Ray's although this has an amplitude of only 1.4 mm, consistent with Figure 3 of Ray and Egbert (2004). This set of 62 harmonic coefficients is called STHL4.X. The amplitudes (H) and Greenwich phase lags (G) of the five main constituents (TC) and their origins (see Pugh and Woodworth, 2014) are listed in Table 1. (A full list of the harmonic coefficients in STHL4 and STHL4.X can be found in the Supplement.)

One obvious thing to point out is how small the diurnal tides are at St. Helena and, therefore, how difficult it would be to decide reliably on any changes in them from one epoch to another, even if one had a longer historical data set than Maskelyne's measurements of over a month and with irregular timing. Our findings, and those of Cartwright (1971) to be discussed below, should be considered with this reservation in mind.

² It was not possible to include in the present study information on the tide at St. Helena based on the Manuel Johnson measurements in 1826-1827 as they were not of sufficient quality for examining small changes in the tide through the years; see Cartwright et al. (2017) for details.

196 The STHL4.X harmonics were used to make 1-minute predictions of the tide for 1761 from which we picked out
 197 values at exactly the same times as Maskelyne’s measurements.³ These are called ‘Predicted’ values although
 198 ‘Hindcasted’ would probably be a better description (Cartwright’s expression for them was a ‘tidal synthesis’). There
 199 are then 478 of them corresponding to the same number of Maskelyne values.

200
 201 Figure 3(a) shows the time series of 1-minute predicted heights from 12 November - 22 December together with the
 202 Maskelyne sea levels shown by red dots. The two sets of values have been adjusted to have zero mean. It can be
 203 seen that many of Maskelyne’s measurements took place around high or low tide as was his intention (Maskelyne,
 204 1762). However, there were also many measurements around mid-tide. Figure 3(b) focusses on a subset of data for
 205 15-20 November, demonstrating the general good correspondence of predicted heights and Maskelyne’s
 206 measurements.

207
 208 Figure 3(c) shows sea level differences (Maskelyne - Predicted) with the overall mean difference removed. One can
 209 see that there is an apparent datum shift at the time that Maskelyne’s tide pole was damaged on 3 December and
 210 replaced on 5 December. Determining the size of a datum shift is difficult when the shift is comparable to the
 211 variability in the record due to fluctuations in the ocean water properties (especially temperature) and to
 212 meteorological effects, and it is sometimes difficult even deciding if there is a shift at all. However, simple inspection
 213 suggests a shift of about 2.8 cm at that time, estimated from the difference between the average sea level differences
 214 either side of the gap. Figure 4 (a) shows that after adjustment for the shift the Predicted and Maskelyne sea levels
 215 values correspond satisfactorily (as in fact do the unadjusted vales given that 2.8 cm is a small amount compared to
 216 the tidal range). Figure 4(b) shows that the sea level differences have no major dependence on tidal level.⁴

217
 218 An important issue at this point is that Cartwright (1971) did not believe that there was any evidence for a datum
 219 shift. Therefore, his findings were based on an analysis of the complete Maskelyne data set without consideration of
 220 either a datum shift or the possible importance of long-period tides (which in this case amounts to much the same
 221 thing). We made a considerable number of tests using predictions based on STHL4.X of whether findings on the
 222 tidal composition of Maskelyne’s data could be affected by his irregular temporal sampling and/or by a datum shift
 223 and/or by long-period tides. There were too many tests to be described in detail in this short note but our general
 224 conclusion was that a datum shift of 2-3 cm or the presence of long-period tides (or not) would not impact
 225 significantly on the determination of the main semidiurnal tides (M2 and S2) but would be important for the diurnals,
 226 with uncertainties of about 10% in their amplitudes. These initial tests informed our choice of methods employed in
 227 the next sections.


228
 229
 230
 231 **Table 1:** Amplitudes (H, cm) and Greenwich Phase Lags (G, deg) of the main tidal constituents (TC) in the STHL4.X
 232 set and their origins.

TC	H (cm)	G (deg)	Origin
M2	32.49	80.04	Principal lunar semidiurnal
S2	11.39	101.96	Principal solar semidiurnal
K1	3.46	349.22	Principal lunar/solar diurnal
O1	2.08	190.65	Principal lunar
N2	6.69	70.88	Larger elliptical lunar semidiurnal


³ These predictions assume the same nodal variations for the lunar tides as in the equilibrium tide, which is a reasonable assumption for an ocean island location. The end of 1761 is anyway not at a time of nodal maximum or minimum, so any uncertainty arising from this assumption will be small.

⁴ The apparent larger scatter at high and low waters than at mid-tide in Figure 4(a), which is counter intuitive as measurements are normally more accurate at the turning points than at mid-tide when the water level is changing rapidly, is an artefact of there being more measurements at the high and low water levels. The standard deviation of Maskelyne minus Predicted levels in Figure 4(b) is 3.2, 3.0 and 3.0 cm for bands of predicted level -60 to -20, -20 to 20 and 20 to 60 cm respectively.



243




244
245



(a)




(b)



(c)


247
248
249
250
251

Figure 3. (a) 1-minute predicted heights for 12 November - 22 December 1761 together with the Maskelyne sea levels shown by red dots. The two sets of values have been adjusted to have zero mean.



252
253
254
255
256
257
258
259
260
261
262
263


Figure 3. (b). A subset of (a) focussing on 15-20 November 1761 which corresponds to Figure 4 of Cartwright (1971), although the predicted tide will be slightly different in the two cases, and a couple of outlying Maskelyne's measurements shown here appear not to have been used by Cartwright. Before and after the six days shown the observations of Maskelyne are mostly confined to daylight hours. (c). Sea level differences (Maskelyne - Predicted) with the overall mean difference removed.





264
265
266
267

Figure 3(c). Sea level differences (Maskelyne—Predicted) with the overall mean difference removed.

268




269
270



271
272
273

Figure 4 (a). Predicted and Maskelyne sea levels values after datum shift adjustment.



274
275 **Figure 4** (b). Sea level differences (Maskelyne-Predicted) after datum shift correction showing no major dependence
276 on tidal level.
277

278 3.3 New Harmonic Tidal Analysis (First Method)

279
280 The next question is how to analyse more rigorously the modern and historical (i.e. Predicted and Maskelyne)
281 measurements in order to see how the tide might have differed between the two epochs. To do that, a form of
282 harmonic analysis was devised involving harmonic fitting to the two sets of 478 heights. The obvious drawback to
283 such a fit is that the data are not regularly spaced in time, so an initial concern was that fits could be biased towards
284 the earlier days with many measurements. However, the residuals of the fits described below look much like those
285 of Figure 3(c), with similar earlier and later residual variances. Therefore, this seems not to be the case.

286
287 Several variants of harmonic fits were tried and what seems to be the best is described here. This considered just
288 five main constituents (M2, S2, K1, O1 and N2) as it was believed that the available data justified using no more
289 than that. N2 was included in its own right, rather than as a related constituent, because it is so large (amplitude of
290 almost 7 cm, and larger than either K1 and O1) and in theory it should be determined adequately with a month of
291 data.

292
293 Several other harmonics were included as related constituents. These included K2 and T2 related to S2, P1 related
294 to K1, Q1 related to O1, and NU2, 2N2 and MU2 related to N2. Each of the related constituents were required to
295 have amplitudes relative to the main constituents in proportion to those in STHL4.X, and differences in phase lags
296 to those of the main constituents also as in STHL4.X, but with related amplitudes and phase lags adjusted for the
297 nodal cycle as appropriate for the end of 1761.

298
299 Various tests were made to ensure: (i) that the method and program code perform as required, and (ii), as mentioned
300 above, that the irregular sampling of the 478 Maskelyne times does not in itself unduly bias the determined
301 amplitudes and phases for the main constituents, especially the semidiurnals. The fit thereby contains 10 parameters
302 for the five main constituents and an optional 11th parameter to represent the possible datum shift on 4 December.

303
304 Changes were made before making these fits to both the Predicted and Maskelyne data considered previously in
305 Section 3.2, in order to remove the complications of long-period tides from both of them. This involved a simple
306 change for the Predicted data, in that we used only 57 of the 62 harmonics in STHL4.X (i.e. without the 5 long-
307 period components Sa, Ssa, Mm, MSf and Mf).

308
309 For Maskelyne's data, the contributions of Mm, MSf and Mf to the measurements were removed using values from
310 the Finite Element Solution (FES) 2014 model (Lyard et al., 2021). Of these, Mf is the most important and MSf the
311 least important. The model has H (cm) and G (deg) values of 1.49 and 358.64 for Mf, 0.69 and 355.46 for Mm, and
312 0.020 and 209.15 for MSf respectively. Corrections for the seasonal constituents Sa and Ssa were not made as it was
313 decided that they were unlikely to be important. Eight years of monthly mean sea level (MSL) values for St. Helena
314 in the Permanent Service for Mean Sea Level (PSMSL) data set (Holgate et al., 2013) indicated that November
315 levels were on average only 11 mm above December levels but could vary between +36 and -15 mm. Therefore,
316 given that we also had no way to correct for daily and weekly changes in MSL, no monthly MSL adjustments were
317 made.

318
319 The results when including the 11th parameter in the two fits to Predicted and Maskelyne data were as in Table 2(a).
320 AR is the ratio of historical to modern amplitudes and PHLD is the historical phase lag minus modern phase lag.
321 The fit to Predicted values suggested a possible datum shift of -0.2 cm, consistent with zero, as expected as of course
322 the predicted time series has no shift in it. On the other hand, the fit to the historical Maskelyne values resulted in
323 1.4 cm for the possible datum shift. This is a smaller shift than obtained in the initial look at the data in Section 3.2;
324 accounting for the long-period tides in the Maskelyne data using the FES2014 model values reduced its estimated
325 amount.

326
327 The fits indicate that there has been no change to the main semidiurnal tides (M2 and S2) although N2 is suggested
328 to be 15% smaller in the historical tide. On the other hand, they suggest that K1 had only 84% of its modern value
329 in historical times, and there was a 12% larger historical O1. These departures of the amplitudes of the historical
330 diurnals from their modern values of approximately 10% are consistent with the anticipated biases arising from the
331 various tests with STHL4.X mentioned above.

332
333 However, if one believed there to be no datum shift, as in Cartwright (1971), then fits using 10 parameters only
334 result in the values shown in Table 2(b). In this case, the M2 and S2 semidiurnal tides appear to be much the same
335 in the two epochs, with S2 just a few percent smaller in the historical data. However, it results in historical K1 being
336 only 72% of its modern value which is less plausible, while historical O1 is suggested to have been 8% larger. These

337 two sets of results demonstrate how sensitive the findings for the diurnals are to whether or not there was a small
 338 datum shift, even if only centimetric.

339
 340
 341
 342 **Table 2:** (a) Values of ratio of historical to modern amplitudes (AR) and historical phase lag minus modern phase
 343 lag (PHLD, deg) when including an 11th parameter in the fits to Predicted and Maskelyne data by the first harmonic
 344 method (Section 3.3). (b) Values of AR and PHLD when not including an 11th parameter in the first harmonic
 345 method. (c) Values obtained using the second harmonic method (Section 3.4).

346
 347 (a)

M2		S2		K1		O1		N2	
AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD
0.994	-0.50	0.992	1.263	0.841	1.758	1.124	9.613	0.846	-0.463

351
 352 (b)

M2		S2		K1		O1		N2	
AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD
0.992	-0.690	0.961	1.817	0.719	-6.029	1.080	12.021	0.841	-0.812

356
 357 (c)

M2		S2		K1		O1		N2	
AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD
0.997	-0.48	1.040	2.35	0.938	8.23	0.952	-5.38	0.935	-1.4
		[0.951]							

364 3.4 New Harmonic Tidal Analysis (Second Method)

365
 366 As explained above, a problem with analysing the Maskelyne data set is the irregular temporal sampling of his
 367 measurements. Therefore, in order to provide a more conventional time series for use in a second form of harmonic
 368 analysis, use was made of the predicted tide in order to interpolate at 1-minute intervals between Maskelyne's
 369 measurements, applying linear adjustments to the predictions between consecutive Maskelyne measurements so as
 370 to correspond faithfully to the original data when available. There is obviously some danger in this approach,
 371 particularly for calculation of the diurnal tides, given that many of the later Maskelyne measurements were during
 372 daytime only. Therefore, there is a possibility of some information content from the predicted tide passing into the
 373 interpolated Maskelyne tide. However, in spite of these reservations it was decided that this approach offered an
 374 interesting alternative method. The predicted tide in this case was taken from a set of 62 harmonics called STHL2
 375 calculated from an SSP record at St. Helena spanning November 1993 to February 1995. Its five main constituents
 376 have amplitudes and phase lags as shown in Table 3. [\(A full list of the harmonic coefficients in STHL2 can be found](#)
 377 [in the Supplement.\)](#) These STHL2 values are almost identical to those of STHL4, but the amplitude of S2 is 0.914
 378 of that in STHL4.X for the air tide reasons explained in Section 3.2.⁵

379
 380
 381 **Table 3:** Amplitudes (H, cm) and Greenwich Phase Lags (G, deg) of the main tidal constituents (TC) in the STHL2
 382 set.

TC	H (cm)	G (deg)
M2	32.37	79.92
S2	10.41	99.57

⁵ [A reviewer asked why we chose to use a different set of constants \(STHL2\) in this section instead of the STHL4.X used earlier, or the set derived from data during 1993-2006 mentioned above. The fact is that the two authors of this paper started on their analyses using their individually-chosen sets of data that were happy with as to their tidal information content. In practice, it does no harm to use different sets as it provides a feel for the stability of present-day constants.](#)

389	K1	3.49	349.71
390	O1	2.03	187.89
391	N2	6.67	70.87

392

393

394 Using STHL2 results in a time series of 59040 predicted 1-minute values between 12 November and 22 December
395 1761, and a separate time series of Maskelyne values interpolated using STHL2. Each of these was analysed using
396 a conventional monthly tidal analysis containing 27 independent constituents including two long-period constituents
397 (Mm and MSf), and 8 related constituents with relationships to corresponding independent constituents taken from
398 those in STHL2. Findings were as shown in Table 2(c), again suggesting similar semidiurnal tides (M2 and S2) in
399 modern and historical times, although the AR value for S2 slightly larger than 1.0 converts to 0.951 once correction
400 for the different S2 amplitudes in SSP and real sea level are allowed for (shown by the square brackets). There were
401 smaller historical diurnal tides than modern ones and smaller N2.

402

403 This tidal analysis results in interesting findings for the two long-period tides. These are calculated to have
404 amplitudes (cm) for Mm of 3.143 (historical) and 0.577 (predicted), and for MSf of 2.597 (historical) and 2.016
405 (predicted). However, as noted above, the amplitude of the real Mm at St. Helena is only 0.69 cm, if one assumes
406 the FES2014 model to be correct. It turns out that the much larger amplitude of Mm obtained in the analysis of the
407 historical data than that of the predicted data, together with its phase lag, is consistent with simply parameterising
408 the possible datum shift discussed in Section 3.3 in a different way.

409

410 In summary, The results of this second harmonic analysis for the main semidiurnal tides are essentially identical to
411 those of the first harmonic method in Section 3.3. In addition, both suggest smaller historical diurnal tides than
412 modern ones, although this second harmonic method suggests more stable K1 and O1 than does the first method.

413

414 3.5 New Response Analysis

415

416 In a further type of tidal analysis, we attempted to reproduce the work of Cartwright (1971) by modifying the
417 response analysis software which Cartwright used for the analysis of short records.

418

419 Unfortunately, the version available to us did not work with randomly spaced data and had to be modified to do so
420 and rewritten in the Delphi language. It filters a reference time series with 6 bandpass filters: diurnal, semidiurnal
421 and ter-diurnal, each having a real and conjugate part. In addition, the original version of the software lags the
422 reference data by 2 days relative to the observations and applies the same filters. As a result, there are 12 band-
423 passed series in total. The software then calculates a co-variance matrix between a reference series and the data under
424 investigation, and response weights are calculated relative to these 12 series. For present purposes, we extended the
425 method to employ reference series that also lead the observations by 2 days, thereby making the analysis
426 symmetrical, resulting in 18 band-passed reference series in all. The data themselves were not filtered.

427

428 We applied the program to the Maskelyne data using St. Helena predictions as a reference, derived from STHL2 as
429 described above. As regards the dominant M2 constituent, findings indicate AR = 0.992 and PHLD = -0.11°.
430 Meanwhile, AR for S2 = 0.972 after allowance for the air tide and PHLD = 4.19°. However, findings for the small
431 diurnal tides were unsatisfactory, yielding historical amplitudes only about 60% of their modern ones, which is less
432 plausible.

433

434 These results are not perfect but provide confidence in the findings using the harmonic methods, at least for the M2
435 constituent. The sparse Maskelyne data, and its daylight bias, are likely to be the major reasons for what appears to
436 be the poor performance of the response method for the diurnals.

437

438 4 Comparisons of Cartwright's and Our Own Analyses

439

440 Cartwright (1971) used a complicated form of response method, allowing for measurements at arbitrary fractions of
441 an hour so as to accommodate the irregular times of Maskelyne's measurements, and using data from Ascension
442 Island as a reference record in the method. Unfortunately, those Ascension data are no longer available and there are
443 details of his method which are hard to follow. Therefore, it is impossible to reproduce Cartwright (1971) in all
444 respects. The best we can do at this stage is to make our own analyses and see if our main conclusions agree with
445 his.

446

447 Cartwright (1971) decided that the 1761 M2 tide amplitude was 0.98 of the modern value (or 0.984 by a different
448 method) and that the historical phase lag was 2.9 ° (or 2.39 ° by the different method) less than now. We present

449 these in Table 4 in the same form as Table 2.⁶ These are almost the same as the conclusions for M2 in Section 3.3
 450 when allowing for a datum shift or not. They are also consistent with the M2 findings in Section 3.4. Cartwright
 451 concluded that his PHLD values were indistinguishable from zero given the noise in the records.

452
 453
 454 **Table 4:** Values of ratio of historical to modern amplitudes (AR) and historical phase lag minus modern phase lag
 455 (PHLD, deg) obtained in the two methods of Cartwright (1971).

456

	M2		S2		K1		O1	
	AR	PHLD	AR	PHLD	AR	PHLD	AR	PHLD
457 Method 1:	0.98	-2.9	1.02	2.9	1.0	13.0	0.83	13.0
458 Method 2:	0.984	-2.39	1.016	2.39	0.95	9.1	0.95	9.1

459
 460
 461

462
 463 In the case of S2, he stated that ‘the trends are almost exactly reversed [compared to M2]’. Cartwright used a pressure
 464 sensor for his 1969 measurements, as we did later for STHL2 and STHL4, but his 1971 paper makes no mention of
 465 the complication of the S2 air tide. However, the amplitude given for S2 in his Table 2 is 11.2 cm which indicates
 466 that, for one reason or other, he believed that the amplitude of S2 in the ocean tide at this location was essentially
 467 the same as we have used in STHL4.X, as discussed in Section 3.2.

468
 469 In the present study, the first harmonic method of Section 3.3 showed that, if one allows for the datum shift or not,
 470 then the ‘reverse S2 trend’ is indeed the case for phase lag, although the historical S2 amplitude is a little smaller
 471 than today, as for M2, rather than a little larger as Cartwright obtained. That is the same conclusion as for the second
 472 harmonic method in Section 3.4 once the air tide is allowed for. It cannot be important to agonise about the very
 473 small changes in S2 amplitude. That would require unreasonable assumptions concerning the accuracy of
 474 Maskelyne’s measurements: for example, on the accuracy of the 3-inch graduations of the tide pole and an
 475 assumption that it was perfectly vertical. In addition, there was the inevitable complication of making accurate tidal
 476 measurements in the frequent presence of high surf. Similar to Cartwright, we do not believe there is any significance
 477 in the ‘reverse trend’ for S2 phase lag, given the variability in the records, consistent with PHLD near zero for both
 478 of the main semidiurnals.

479
 480 The diurnals are more problematical. Cartwright (1971) considered that the historical amplitudes of the diurnals
 481 were 1.0 and 0.85 times the present ones for K1 and O1 respectively with both historical phase lags about 13 ° larger.
 482 In a second method, he considered both historical amplitudes to be about 95% of the present ones with historical
 483 phase lags 9 ° larger.

484
 485 The phase lag findings of Cartwright (1971) are consistent with those of the first harmonic method in Section 3.3
 486 when allowing for a datum shift, with phase lags larger in historical times (although not by as much as 13°). On the
 487 other hand, the first method has smaller/larger historical amplitudes for K1/O1, compared to similar or slightly
 488 smaller historical amplitudes for both constituents in Cartwright (1971). If one does not allow for a datum shift, the
 489 first method again suggests smaller/larger historical amplitudes for K1/O1 but PHLD values moving in opposite
 490 directions unlike in Cartwright (1971).

491
 492 The conclusions on the diurnals from the second harmonic method of Section 3.4 are consistent with Cartwright
 493 (1971), in there being historical amplitudes a few percent smaller than today, although the phase lags for K1 and O1
 494 move in opposite directions in the second method unlike in Cartwright (1971). One notes the large Mm obtained in
 495 Section 3.4 from the interpolated Maskelyne data is consistent with the datum shift considered in Section 3.3. Our
 496 own attempt at response analysis in Section 3.5 supported the case for essentially unchanged semidiurnal tides.

497
 498 Some differences in findings between this report and Cartwright (1971) are to be expected for several reasons. One
 499 important aspect concerns the data sets that he had available, which were a year of high and low water levels at
 500 Ascension in 1958-9, 39 days of his own measurements at St. Helena in 1969, and of course Maskelyne’s
 501 measurements in 1761. On the other hand, we have about three decades of continuous sea level measurements from
 502 St. Helena, acquired through South Atlantic programme of the National Oceanography Centre (Spencer et al., 1993).
 503 Those records, from which STHL4.X and STHL2 were derived, are much higher quality than Cartwright’s.
 504

⁶ We believe we have the correct signs for PHLD in this table: Cartwright chose to work with phase leads rather than lags.

505 Second, the methods used to analyse the data are different. Cartwright used the Ascension record as a reference in a
506 response analysis involving his 1969 data and Maskelyne’s data. On the other hand, we have used two forms of
507 harmonic analysis and our own response method. Less important, it seems that Cartwright did not include in his
508 analysis the several measurements that Maskelyne himself flagged as suspect’doubtful’ or ‘very doubtful’ because
509 of swell conditions, whereas we have used all 478 of Maskelyne’s measurements in our analyses.

510
511 Third, there is the question of whether or not there really was a datum shift on 4 December 1761. Cartwright (1971)
512 did not believe there was any evidence for a ‘sensible change in datum’, but the present work has shown that there
513 probably was a small shift. It is possible that the filters Cartwright used in his response method resulted in his analysis
514 being less sensitive to a small shift. However, that seems not to be the case when using the harmonic method.
515 Although findings on changes in the main semidiurnal tides (M2 and S2) are largely unaffected, consideration of the
516 shift does have an impact on findings for the diurnal tides. Unfortunately, the shift happens in almost the middle of
517 the 40 days of measurements and splitting the data sets into two and analysing them separately, as one might do with
518 a longer record, is not a suitable option.

520 5 Conclusions

521
522 To sum up, the headline results of Cartwright (1971) were that the semidiurnal tides had not changed at St. Helena
523 since 1761, that the amplitudes of the diurnals were on balance slightly smaller than today, and that they had about
524 10° larger phase lag in 1761. Both of our new analyses agree qualitatively with those conclusions for the
525 semidiurnals: we believe historical and modern M2 to be essentially the same and that the S2 amplitude was a couple
526 of percent smaller in historical times. On the other hand, one notes big differences between methods in Tables 2 and
527 4 for the diurnals which result from the difficulties of analysing the short Maskelyne data set. As a result, we would
528 be more hesitant to claim any changes at all in either the semidiurnal or diurnal tides. In retrospect, one wonders
529 why Cartwright chose to focus on the apparent changes in phase of the diurnals that he obtained, given that his own
530 discussion on top of p619 shows that their approximately 10° phase lag difference was only a 2-sigma effect.

531
532 The findings of Cartwright (1971) were important ones that have been referenced in major reviews of “tides a-
533 changin” (e.g. Haigh et al., 2020). However, from the perspective of ‘old records for new knowledge’, his study also
534 serves as an important example that electronic data sets can be as much at risk as paper records. His own
535 measurements at St. Helena in 1969 can no longer be found, while his version of the 1958-9 high and low water
536 record at Ascension and, of course, the Maskelyne data at St. Helena are also missing.⁷ Therefore, it is good at least
537 that a data set of the historical St. Helena measurements made by Maskelyne in 1761 is once again available for any
538 researcher to investigate. It has been interesting to make our own analyses from that recovered data set, confirming
539 Cartwright’s main findings on the similarity of the predominant M2 constituent in the historical and modern data
540 from St. Helena.

542 6 Data availability

543
544 The small file ‘maskelyne_data’ referred to above can be accessed via a DOI from [https://doi.org/10.5285/e0f2b6ea-](https://doi.org/10.5285/e0f2b6ea-d11d-3102-e053-6e86abc073ab)
545 [d11d-3102-e053-6e86abc073ab](https://doi.org/10.5285/e66db85a-eaae-6665-e053-6c86abc0bfb9).—<https://doi.org/10.5285/e66db85a-eaae-6665-e053-6c86abc0bfb9>. Its citation is
546 shown here as Woodworth and Vassie (2022). The file has also been deposited with the British Oceanographic Data
547 Centre in which it has Accession Number POL200133. Any information on the tidal analyses made in this study
548 may be obtained from both authors.

549
550 Supplement. A supplement to this paper is available containing a full list of harmonic constants in STHL4,
551 STHL4.X and STHL2.

552
553 **Author contributions.** PLW undertook the data rescue aspect of this work. Both PLW and JMV performed the data
554 analyses and prepared the manuscript.

555
556 **Acknowledgements.** We thank Richard Ray and David Pugh for comments on an earlier version of this article.
557 Richenda Houseago-Stokes of the British Oceanographic Data Centre is thanked for setting up the DOI for the
558 Maskelyne data set.

⁷ Cartwright obtained the 1958-9 Ascension high and low water data from the US Coast and Geodetic Survey (USCGS) so it is probable that a version of this data set is archived by a US centre. Cartwright (1971) also mentions a month of hourly sea levels from Ascension in 1959 that was obtained from the USCGS. The latter does not appear to have been used in the Cartwright (1971) study; if required, a manuscript copy of that short record may be obtained from the present authors.

559
560 **Competing interests.** The contact author has declared that neither author has any competing interests.

561
562
563
564
565

566 **ORCID**

567
568 P.L. Woodworth <http://orcid.org/0000-0002-6681-239X>

569
570 J.M. Vassie <http://orcid.org/0000-0002-9906-1019>

571

572 **References**

- 573
- 574 Cartwright, D.E.: Proceedings IHB Symposium on Tides held in Monaco, April 1967. p195-200. UNESCO
575 publication, 1969.
- 576
- 577 Cartwright, D.E.: Tides and waves in the vicinity of Saint Helena, Philos. T. R. Soc. London, A, 270, 603-649,
578 <https://doi.org/10.1098/rsta.1971.0091>, 1971.
- 579
- 580 Cartwright, D.E.: Some ocean tide measurements of the eighteenth century, and their relevance today, P. R. Soc.
581 Edinb. B: Biological Sciences, 72, 331-339, <https://doi.org/10.1017/S0080455X00001892>, 1972.
- 582
- 583 Cartwright, D.E., Spencer, R., Vassie, J.M., and Woodworth, P.L.: The tides of the Atlantic Ocean, 60N to 30S,
584 Philos. T. R. Soc. London, A, 324, 513-563, <https://doi.org/10.1098/rsta.1988.0037>, 1988.
- 585
- 586 Cartwright, D.E., Woodworth, P.L., and Ray, R.D.: Manuel Johnson's tide record at St. Helena, Hist. Geo-Space
587 Sc., 8, 9-19, <https://doi.org/10.5194/hgss-8-1-2017>, 2017.
- 588
- 589 Haigh, I.D., Pickering, M.D., Green, J.A.M., Arbic, B.K., Arns, A., Dangendorf, S., Hill, D., Horsburgh, K., Howard,
590 T., Idier, D., Jay, D.A., Lee, S.B., Müller, M., Schindelegger, M., Talke, S.A., Wilmes, S-B., and Woodworth, P.L.:
591 The tides they are a-changin', Rev. Geophys., 57, e2018RG000636, <https://doi.org/10.1029/2018RG000636>, 2020.
- 592
- 593 Holgate, S.J., Matthews, A., Woodworth, P.L., Rickards, L.J., Tamisiea, M.E., Bradshaw, E., Foden, P.R., Gordon,
594 K.M., Jevrejeva, S., and Pugh, J.: New data systems and products at the Permanent Service for Mean Sea Level, J.
595 Coastal Res., 29, 493-504, <https://doi.org/10.2112/JCOASTRES-D-12-00175.1>, 2013.
- 596
- 597 Hughes, D.W., Yallop, B.D., and Hohenkerk, C.Y.: The equation of time, Mon. Not. R. Astron. Soc., 238, 1529-
598 1535, <https://doi.org/10.1093/mnras/238.4.1529>, 1989.
- 599
- 600 [IOC \(Intergovernmental Oceanographic Commission\): Manual on sea-level measurements and interpretation \(IOC
601 Manuals and Guides No.14\), Volume V: Radar Gauges. \(ed. P.L. Woodworth\). Paris, Intergovernmental
602 Oceanographic Commission of UNESCO, 104pp, 2016. Available from
603 <http://unesdoc.unesco.org/images/0024/002469/246981E.pdf>.](http://unesdoc.unesco.org/images/0024/002469/246981E.pdf)
- 604
- 605 Lyard, F.H., Allain, D.J., Cancet, M., Carrère, L., and Picot, N.: FES2014 global ocean tide atlas: design and
606 performance, Ocean Sci., 17, 615-649, <https://doi.org/10.5194/os-17-615-2021>, 2021.
- 607
- 608 Maskelyne, N.: Observations on the tides in the Island of St. Helena: in a letter from the Rev. Nevil Maskelyne,
609 A.M. F.R.S. to Thomas Birch, D.D. Secretary to the Royal Society, Philos. T. R. Soc., 52, 586-606,
610 <https://doi.org/10.1098/rstl.1761.0100>, 1762.
- 611
- 612 Munk W.H. and Cartwright, D.E.: Tidal spectroscopy and prediction, Philos. T. R. Soc. London, A, 259, 533-581,
613 <https://doi.org/10.1098/rsta.1966.0024>, 1966.
- 614
- 615 Ray, R.D.: Diurnal oscillations in atmospheric pressure at twenty-five small oceanic islands, Geophys. Res. Lett.,
616 25, 3851-3854, <https://doi.org/10.1029/1998GL900039>, 1998.
- 617
- 618 Ray, R.D. and Egbert, G.D.: The global S1 tide, J. Phys. Oceanogr., 34, 1922-1935, [https://doi.org/10.1175/1520-0485\(2004\)034<1922:TGST>2.0.CO;2](https://doi.org/10.1175/1520-0485(2004)034<1922:TGST>2.0.CO;2), 2004.
- 620
- 621 Spencer, R., Foden, P.R., McGarry, C., Harrison, A.J., Vassie, J.M., Baker, T.F., Smithson, M.J., Harangozo, S.A.,
622 and Woodworth, P.L.: The ACCLAIM programme in the South Atlantic and Southern Oceans, Int. Hydrogr. Rev.,
623 70, 7-21, 1993.
- 624
- 625 Woodworth, P.L.: A survey of recent changes in the main components of the ocean tide, Cont. Shelf Res., 30, 1680-
626 1691, <https://doi.org/10.1016/j.csr.2010.07.002>, 2010.
- 627
- 628 Woodworth P.L. and Vassie J.M.: Maskelyne sea level observations from the Island of St. Helena in 1761. NERC
629 EDS British Oceanographic Data Centre NOC. [https://doi.org/10.5285/e0f2b6ea-d11d-3102-e053-6c86abc073ab-
630 2022](https://doi.org/10.5285/e0f2b6ea-d11d-3102-e053-6c86abc073ab-2022).
- 631 <https://doi.org/10.5285/e66db85a-eaae-6665-e053-6c86abc0bfb9>, 2022.

632 **Figure Captions**

- 633
- 634 1. The start of the table of measurements of sea level at St. Helena in 1761 to be found at the end of Maskelyne
635 (1762).
- 636
- 637 2. The recorded (local) times of Maskelyne's measurements. Some measurements were made around the clock in
638 the earlier part of the data set. However, they can be seen to be restricted to daylight hours in the latter part.
- 639 3. (a) 1-minute predicted heights for 12 November - 22 December 1761 together with the Maskelyne sea levels
640 shown by red dots. The two sets of values have been adjusted to have zero mean. (b) A subset of (a) focussing on
641 15-20 November 1761 which corresponds to Figure 4 of Cartwright (1971), although the predicted tide will be
642 slightly different in the two cases, and a couple of outlying Maskelyne's measurements shown here appear not to
643 have been used by Cartwright. Before and after the six days shown the observations of Maskelyne are mostly
644 confined to daylight hours. (c) Sea level differences (Maskelyne - Predicted) with the overall mean difference
645 removed.
- 646
- 647 4. (a) Predicted and Maskelyne sea levels values after datum shift adjustment. (b) Sea level differences (Maskelyne-
648 Predicted) after datum shift correction showing no major dependence on tidal level.
- 649