

ChinaCropSM1 km: a fine 1 km daily Soil Moisture dataset for dryland wheat and maize across China during 1993–2018

Fei Cheng¹, Zhao Zhang^{1, 2}, Huimin Zhuang¹, Jichong Han¹, Yuchuan Luo¹, Juan Cao¹, Liangliang Zhang¹, Jing Zhang¹, Jialu Xu¹ and Fulu Tao^{2, 3, 4}

¹Academy of Disaster Reduction and Emergency Management, Ministry of Emergency Management & Ministry of Education, School of National Safety and Emergency Management, Beijing Normal University, Beijing 100875, China

²Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographical Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

³College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

⁴Natural Resources Institute Finland (Luke), FI-00790 Helsinki, Finland

Correspondence to: Zhao Zhang (sunny_zhang@bnu.edu.cn)

Abstract. Soil moisture (SM) is a key variable of the regional hydrological cycle and has important applications for water resource and agricultural drought management. Various global soil moisture products have been mostly retrieved from microwave remote sensing data. However, there is currently rare spatially explicit and time-continuous soil moisture information with a high resolution at the national scale. In this study, we generated a 1 km soil moisture dataset for dryland wheat and maize in China (ChinaCropSM1 km) over 1993–2018 through a random forest (RF) algorithm based on numerous in situ daily observations of soil moisture. We independently used in situ observations (181327 samples) from the Agricultural Meteorological Stations (AMS) across China for training (164202 samples) and others for testing (17125 samples). An irrigation module was first developed according to crop type (i.e., wheat, maize), soil depth (0–10 cm, 10–20 cm) and phenology. We produced four daily datasets separately by crop type and soil depth, and their accuracies were all satisfactory (wheat r 0.93, ubRMSE 0.033 m^3m^{-3} ; maize r 0.93, ubRMSE 0.035 m^3m^{-3}). The spatiotemporal resolutions and accuracy of ChinaCropSM1 km were significantly better than those of global soil moisture products (e.g., r increased by 116%, ubRMSE decreased by 64%), including the global remote-sensing-based surface soil moisture dataset (RSSSM) and the European Space Agency (ESA) Climate Change Initiative (CCI) SM. The approach developed in our study could be applied to other regions and crops in the world, and our improved datasets are very valuable for many studies and field management, such as agricultural drought monitoring and crop yield forecasting. The data are published in Zenodo at <https://zenodo.org/record/6834530> (wheat₀₋₁₀) (Cheng et al., 2022a), <https://zenodo.org/record/6822591> (wheat₁₀₋₂₀) (Cheng et al., 2022b), <https://zenodo.org/record/6822581> (maize₀₋₁₀) (Cheng et al., 2022c) and <https://zenodo.org/record/6820166> (maize₁₀₋₂₀) (Cheng et al., 2022d).

刪除的內容: 1km

刪除的內容: 1km

刪除的內容: Crop

刪除的內容: s

刪除的內容:

刪除的內容: Minsitry

刪除的內容: a nation

刪除的內容: Here

刪除的內容: 1km

刪除的內容: stable crop

刪除的內容: s

刪除的內容: 1km

刪除的內容: .

刪除的內容: independently

刪除的內容: firstly

刪除的內容: i.e.

刪除的內容: accuracy is

刪除的內容: spatio-temporal

刪除的內容: 1km

刪除的內容: are

刪除的內容: 6 %

刪除的內容: 4 %

刪除的內容: into

刪除的內容: managements

刪除的內容: agriculture

刪除的內容: The data are published in Zenodo at <https://zenodo.org/record/6834530> (wheat₀₋₁₀) (Cheng et al., 2022a), <https://zenodo.org/record/6822591> (wheat₁₀₋₂₀) (Cheng et al., 2022b), <https://zenodo.org/record/6822581> (maize₀₋₁₀) (Cheng et al., 2022c) and <https://zenodo.org/record/6820166> (maize₁₀₋₂₀) (Cheng et al., 2022d).

1 Introduction

Soil moisture (SM) is closely associated with droughts and floods, and consequently agricultural production (Tao et al., 2003). Therefore, SM information at a high resolution is critical to improve crop yield prediction (Prasad et al., 2006; Chakrabarti et al., 2014) and drought impact assessment (Sheffield, 2004). However, such higher resolutions at both temporal (e.g., daily and

65 more than decade) and spatial scales are still unavailable across China, especially for dry croplands.

SM can be obtained in several ways, including in situ observations (Walker et al., 2004; Bogena et al., 2007), remote sensing retrieval (Mohanty et al., 2017; Wei et al., 2019), and process-based model simulations (Vergopolan et al., 2020; Ahmed et al., 2021).

Field observations provide the most accurate SM but are expensive and time-consuming, and there are large uncertainties from extrapolating the limited observations into larger regions with high heterogeneity (Collow et al., 2012; Crow et al., 2012).

70 Microwave sensors have been applied to retrieve SM in recent years (Schmugge et al., 2002; Wigneron et al., 2003; Amazirh et al., 2018). The microwave sensors can only monitor near-surface SM (0–10 cm) (Eagleman and Lin, 1976; Jackson et al., 1982). Passive microwave sensors can monitor daily SM but with a coarse resolution (25–40 km), compared

75 with a high spatial resolution (10–30 m) and a coarser repetition interval (15–25 days) for active sensors (Eagleman and Lin, 1976; Jackson et al., 1982; Mallick et al., 2009). Such SM products have large uncertainties due to the limitations of satellite coverage and downscaling methods, although they can easily cover large regions compared with in situ observations (Loew et al., 2013; Su et al., 2016; Peng et al., 2017). Deriving the SM from model simulation is also challenging because of its high requirements in input data, computing ability and large uncertainties from model parameters (Wang and Qu, 2009; Yilmaz et al., 2012; Petropoulos et al., 2015). In addition, many studies have found that irrigation, as an additional water supply source other than precipitation, reduces soil albedo (Chen and Dirmeyer, 2019), increases heat capacity (Wang et al., 2019), alters

80 local SM (Lawston et al., 2017), and affects the water/energy budget (Shen et al., 2013). However, few studies have taken irrigation into account in developing SM data products at the national or global scales (Drewniak et al., 2013; Qiu et al., 2016a).

Therefore, it is critical yet challenging to improve SM accuracy at both spatial and temporal resolutions.

As one part of the Climate Change Initiative (CCI), the European Space Agency (ESA) published a long-term surface SM dataset, and the latest version (v06.1) covered the period of 1978–2020 (<https://www.esa-soilmoisture-cci.org/>, last access: 10

85 Apr. 2022) (Dorigo et al., 2017; Gruber et al., 2019; Preimesberger et al., 2021). The ESA CCI SM products are consistent with the observed values at some grassland and farmland sites in China (Liu et al., 2011; Albergel et al., 2013; Dorigo et al., 2015, 2017); however, they have a coarse spatial resolution (~27 km) and many coverage gaps (Llamas et al., 2020; Guevara et al., 2021). More recently, based on multiple neural networks, the global remote-sensing-based surface soil moisture (RSSM) dataset covering 2003–2018 at 0.1° resolution was developed by using Soil Moisture Active Passive (SMAP) SM as the

90 primary training target. The RSSM improved the coefficient of determination (R^2) by 0.46 and the root mean squared error (RMSE) by 0.083 $\text{m}^3 \text{m}^{-3}$, with a 10-day resolution (Chen et al., 2021). In 2020, another new SM dataset in China from 2002 to 2018 was provided from different passive microwave SM products and model-based downscaling techniques (Meng et al., 2021). With an improved correlation coefficient (r) of 0.84 and an unbiased root-mean-squared error (ubRMSE) of 0.056

刪除的內容: .

刪除的內容: productions

刪除的內容: Thus

刪除的內容: resolution in

刪除的內容: by

刪除的內容: The field

刪除的內容: being

刪除的內容: The microwave

刪除的內容: The passive

刪除的內容: comparing

刪除的內容: whereas

刪除的內容: ones

刪除的內容: from

刪除的內容: the

刪除的內容: Thus

刪除的內容: .

刪除的內容: lots of

刪除的內容: Coefficient

刪除的內容: of

刪除的內容: Root Mean Squared Error

刪除的內容: of

115 m³ m⁻³, the new dataset has a 0.05° spatial resolution and a monthly time resolution. These SM products **have contributed** largely to related **agricultural** studies and **management**; however, they are still too coarse to assess agricultural drought risk and predict crop yield accurately.

Although numerous efforts have been devoted to **developing** SM products, major concerns should be addressed: (1) agricultural management activities such as irrigation have not been fully considered by **previous** studies, especially in countries **such as** 120 China with extensive **irrigated** areas **A** (Zhu et al., 2013); (2) both **the** spatial and temporal (e.g. daily) resolutions of SM products need to be improved for regional agricultural **management**; and (3) the SM accuracy **needs** to be further improved. In recent years, **in situ** observations **have become** available (Li et al., 2005). Some new methods, such as machine learning, are increasingly applied to many fields and have been shown to be robust in incorporating multiple sources **of** data to develop spatiotemporal datasets (Ahmad et al., 2010; Srivastava et al., 2013; Im et al., 2016).

125 **Therefore**, our main objectives in the study **were** to develop a novel method to generate a daily **1 km** SM dataset for dry croplands across China based on numerous field observations; to evaluate their accuracy and compare them with current products; **and** to explore the **spatiotemporal** characteristics of soil moisture for **dryland wheat and maize**. We **anticipate that** our methods and datasets will be valuable for **agricultural** drought monitoring and crop yield forecasting.

2 Materials and Method

130 2.1 Study area

The study area is dominated by dryland crops such as wheat and maize in China, with complex cultivation methods (Wu and Li, 2012) and various irrigation activities (Huang et al., 2015). According to the annual crop harvesting areas of crops across mainland China from 2000 to 2015 (Luo et al., 2020a, b), maize and wheat are the two main crops in China, accounting for 35.4% of the total harvested area (FAOSTAT, 2019). The study areas and SM in situ field monitoring sites for the two crops 135 are shown in Figure 1.

2.2 Data

2.2.1 In situ SM observations

The in situ SM observation data (http://data.cma.gov.cn/data/detail/dataCode/AGME_AB2_CHN_TEN.html, last **accessed**: 18 April 2021) from 1993 to 2018 **were** obtained from **agricultural** meteorological sites (**AMS**) in China, which recorded the 140 location, crop type, phenology, soil depth and SM. SM was measured at **depths** of 10 cm and 20 cm at each **AMS** on the 8th, 18th and 28th of each month. For each sample, crop phenology was observed and recorded by well-trained agricultural technicians in experimental fields (the average field size was 0.15 ha) and then **checked** and **qualified** by the Chinese Agricultural Meteorological Monitoring System (CAMMS). **The location of AMS is generally selected in areas with relatively homogeneous soil properties. Also the fact that crops were quite well managed by irrigation according to weather variability**

删除的内容: did contribute

删除的内容: agriculture

删除的内容: managements,

删除的内容: Despite

删除的内容: develop

删除的内容: the

删除的内容: like

删除的内容: irrigated

删除的内容: managements;

删除的内容: need

删除的内容: the

删除的内容: are becoming

删除的内容: Thus

删除的内容: are:

删除的内容: 1km

删除的内容: spatio-temporal

删除的内容: crop

删除的内容: s

删除的内容: are sure

删除的内容: agriculture

删除的内容: 4 %

删除的内容: access

删除的内容: in this study

删除的内容: agriculture

删除的内容: the

删除的内容: agro-meteorological station

删除的内容: were

带格式的: 字体: 非加粗

and crop growth status makes the crop SM records largely representative the overall level of pixels (1 km×1 km) (Zhang et al., 2020; Li et al., 2021). The first layer (0–10 cm) has been widely used to investigate the spatial and temporal characteristics of SM and validate SM retrieved from microwaves across China (Lacava et al., 2012; Zeng et al., 2015; Liu et al., 2018; Fang et al., 2020).

We collected the in situ observations of maize (287 sites) and wheat (240 sites), with a total of 181327 samples (maize: 36226 samples for the 0–10 cm soil layer, 36245 samples for the 10–20 cm soil layer; wheat: 54396 samples for the 0–10 cm soil layer, 54460 samples for the 10–20 cm soil layer).

2.2.2 Environmental factors

The environmental factors were classified into site features and gridded features, which both include meteorological data (MD), day of year (DOY), classified irrigation (CIR), soil properties (SP), remote sensing data (RSD), and geographical information (GI) (Table 1).

MD includes daily total precipitation (pre) and ante-accumulated precipitation over ten days (pre10) from meteorological stations across China (CNMSs) (<http://data.cma.cn>, last access: 10 April 2021) (Figure S1).

CIR was calculated using Eq. (1).

$$CIR = \begin{cases} 1, & C_i P_j D_k SM \geq SMI_{ijk}; \\ 0, & C_i P_j D_k SM < SMI_{ijk}; \end{cases} \quad (1)$$

where C_i , P_j , D_k , and SMI_{ijk} are crop type, phenology, soil depth, and the evaluation index of relative soil moisture (SMI) corresponding to the crop type i , phenology j , and soil depth k , respectively. SMI is a threshold to determine when irrigation is applied (Table 2), which was released by the Ministry of Water Resources of China (CNMWR) (<http://www.mwr.gov.cn>, last access: 10 July 2022) in July 2012.

SP includes sand, silt, gravel, organic carbon, clay contents, soil pH and bulk density, obtained from Harmonized World Soil Database Version 1.2 (<http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>, last access: 18 Aug. 2021). The original 30 arc-second raster spatial resolution data were resampled to a 1 km resolution based on nearest neighbor interpolation, and the site-related SP were extracted from values to points using ArcGIS 10.5 software (ESRI).

RSD includes reference evapotranspiration (pet) and field capacity (fc). pet was obtained from TerraClimate (<https://doi.org/10.7923/G43J3B0R>, last access: 28 Aug. 2021), which included monthly climate and climatic water balance from 1958-present with a resolution of 1/24° or ~4 km (Abatzoglou et al., 2018). fc was obtained from OpenLandMap (<https://zenodo.org/record/2784001>, last access: 18 July 2021) (Hengl and Gupta, 2019), which included fc under 33 kPa at 0 cm (b0) and 10 cm (b10) depths.

GI includes latitude (lat), longitude (lon), moisture index (im) (Thorntwaite, 1948), and river vector data, provided by the Data Centre for Resources and Environmental Sciences, Chinese Academy of Sciences (<http://www.resdc.cn/Default.aspx>, accessed on 18 April 2021). The distance from each AMS to river networks at all levels (R4, R5, R12) in China was calculated using the Euclidean distance analysis method.

带格式的: 字体: 非加粗

带格式的: 字体: 非加粗

带格式的: 字体: 非加粗

带格式的: 字体: 非加粗

带格式的: 字体: 非加粗

删除的内容: microwave

删除的内容: are

删除的内容: Site

删除的内容: Gridded

删除的内容: Classified Irrigation

删除的内容: the

删除的内容: PH

删除的内容: the

删除的内容: Reference Evapotranspiration

删除的内容: 33kPa

删除的内容: depth

215 2.2.3 Public SM products for comparison

We used two existing SM products for comparison: 1) The ESA CCI SM data are a merged multisatellite surface SM product, which consists of active, passive, or combined products. The SM retrievals were from four microwave radiometers (SMMR: Scanning Multichannel Microwave Radiometer, SSM/I: Special Sensor Microwave/Imager, TMI: Tropical Rainfall Measuring Mission (TRMM)'s Microwave Imager, and AMSR-E: Advanced Microwave Scanning Radiometer for the Earth Observing System) and two scatterometers (AMI: Active Microwave Instrument, ASCAT: Advanced Scatterometer) in a 0.25° global daily dataset. The assimilated data rely on their respective sensitivity to vegetation density and use a Global Land Data Assimilation System (GLDAS) surface SM product (Rodell et al., 2004) as a climatological reference (Wagner et al., 2012). The active/passive products were, the integration of the scatterometer/radiometer-based SM retrievals, while the ESA CCI SM product is the fusion of both the active and passive products. We used the v05.2 product for comparison because of its advantages compared with active/passive products (Liu et al., 2012; Dorigo et al., 2017). 2) The RSSSM is an improved global long-term remote-sensing-based surface SM dataset covering 2003–2018 at a 0.1° resolution (<https://doi.org/10.1594/PANGAEA.912597>). Considering their compatibility, we chose 1995 to 2018 for comparison between ChinaCropSM 1 km, and ESA CCI SM, and the 2003–2018 period for that of ChinaCropSM 1 km, and RSSSM.

2.3 Method

230 2.3.1 Variable selection and data treatment

For the site-related variables, we used the Extract Values to Points tool to extract the 1 km resolution raster information of the environmental (i.e., SP, RSD and GI) data to AMS point data, output point data attributes and save it in CSV format to obtain a dataset of environmental factors through ArcGIS 10.5, and then we deleted those with high multicollinearity ($|r| > 0.5$) according to the factor stacks (Figure S3 and S4). Therefore, the 11 independent variables (pre, pre10, DOY, CIR, T_REF_BULK, R4, im, pet, lat, lon and fc) were selected because they well characterize the impacts of meteorological, temporal, irrigation, soil properties, and geographical information on regional SM. We used the “Euclidean distance” option of the Spatial Analyst Tools in ArcGIS10.5 to obtain the variables related to river networks in China (Danielsson, 1980). We also applied the kriging interpolation method to obtain precipitation-related variables (e.g., pre- and pre-10) from CNMSs. Thereafter, all gridded maps were processed in the WGS84 UTM zone 45N Geographic Coordinate System (EPSG: 32645) and resampled to the same spatial resolution (1 km).

2.3.2 Model development

Ensemble learning was used to aggregate a collection of algorithms to predict the potential impacts, which represents a better method than that using any algorithm alone (Brownlee, 2016). Random forest (RF) is a typical ensemble learning algorithm that can be used to build predictive models for both classification and regression purposes. RF fits an ensemble of models that first train a multitude of decision trees and then obtain predictions by an average or vote through all individual trees (Breiman,

刪除的內容: multi-satellite

刪除的內容: are

刪除的內容: assimilated relies

刪除的內容: uses

刪除的內容: climatological

刪除的內容: are

刪除的內容: scatterometer /radiometer

刪除的內容: respectively, while

刪除的內容: comparing

刪除的內容: 1km

刪除的內容: .

刪除的內容: 1km

刪除的內容: 1km

刪除的內容: i.e.

刪除的內容: data set

刪除的內容: Thus

刪除的內容: with

刪除的內容: network

刪除的內容: pre10

刪除的內容: 32645),

刪除的內容: uses

刪除的內容: Forest

刪除的內容: which

270 2001). The algorithm introduces extra randomness when growing trees and searches for the best trees among a random subset
of features. This technique results in greater tree diversity, generally yielding an overall better model (Hutengs and Vohland,
2016; Lagomarsino et al., 2017). In addition, the bagging method, which constructs multiple training subdatasets by resampling
with replacement of the original dataset, is employed to reduce the variance and overfitting (Díaz-Uriarte and Alvarez de
Andrés, 2006; Zhang et al., 2018). Its high accuracy and stability in agricultural fields have been substantiated in several
previous studies, especially for predicting grain yield, identifying crop planting areas, and mapping soil properties (Hengl et
al., 2015; Jeong et al., 2016; Sun et al., 2019).

275 Hyperparameters in an RF model are very important to optimize its performance. Such parameters are initially defaulted, and
we need to investigate their appropriateness or find potentially better values during the development of an RF regression (RFR).
The important hyperparameters include the following:
n_estimators: the number of trees that the algorithm builds before taking the maximum voting or average overpredictions. A
280 high number of trees increases the performance and makes the predictions more stable, but demands more computations.
max_features: the maximum number of features that the random forest considers on a per-split level. The condition is based
on variance for regression.
min_samples_leaf: the minimum number of leaves that are required to split an internal node.
max_samples: ratio of samples needed for training each tree.

285 We applied the 10-fold cross-validation method to tune the four hyperparameters to avoid overfitting of the RF models (Figure
S5). Additionally, we used this 10-fold cross-validation to evaluate model performance (Figure S7).

The detailed irrigation module is shown in Figure S2. Given that SM is highly sensitive to irrigation application for dryland
wheat and maize in China, we first used RF classification (RFC) to build an irrigation module. This module aimed to predict
whether irrigation application occurred there, and assigns response variable “1” for irrigation and “0” for without irrigation
290 according to the response variables and predictor variables (the same environmental indicators used in producing
ChinaCropSM1 km).

The response variable (classified irrigation CIR) was calculated by the irrigation threshold (Table 2) and in situ information,
including crop type, phenology and soil depth. Then, we used the forecasted CIR as an additional predictor, integrating with
other key predictor variables, to drive RFR for forecasting SM. Considering the regional differences in SM, we randomly
295 sampled in situ SM observations (90% for training and 10% for testing) in each agricultural zone to develop the RF model. In
total, 98576 (65626) and 10820 (6845) observations were used for training and testing the model for wheat (maize),
respectively. All these point samples were used to develop the pointed SM model, and then these pointed models are applied
to inversely calculate the gridded SM by inputting 1-km raster environmental variables (Figure 2).

The hyperparameters in the optimal model were determined as 50, 1, 1 and 4 for the respective n_estimators, max_samples,
300 min_samples_leaf and max_features according to the highest accuracy during training (Figure S5). We implemented these
processes in MATLAB 9.8.0 (R2020a). More information can be found in the MATLAB help center
(<https://www.mathworks.com/help/stats/regressionlearner-app.html>, last access: 26 May 2022).

刪除的內容: constructed
刪除的內容: sub-dataset
刪除的內容: Hyper parameters in a
刪除的內容: a
刪除的內容: developing a
刪除的內容: hyper parameters include follows
刪除的內容: over predict
刪除的內容: .
刪除的內容: demand
刪除的內容: leafs
刪除的內容: hyper parameters for avoiding the
刪除的內容: Meanwhile
刪除的內容: use
刪除的內容: crop
刪除的內容: s
刪除的內容: firstly
刪除的內容: the
刪除的內容: s
刪除的內容: .
刪除的內容: assign
刪除的內容: -
刪除的內容: ones
刪除的內容: 1km
刪除的內容: As for the
刪除的內容: Classified Irrigation CIR), it
刪除的內容: is
刪除的內容: Totally
刪除的內容: are
刪除的內容: -
刪除的內容: applied
刪除的內容: -
刪除的內容: developed
刪除的內容: -
刪除的內容: 1km
刪除的內容: hyper parameters
刪除的內容: this
刪除的內容: Matlab9
刪除的內容: could
刪除的內容: referred to

The **feature** importance was evaluated for the RF model with the greatest regression accuracy by ordering the out-of-bag predictor observations, using the **MATLAB** ‘*oobPermutedPredictorImportance*’ function (<https://www.mathworks.com/help/stats/regressionbaggedensemble.oobpermutedpredictorimportance.html>, last access: 26 May 2022). We also used the method to measure the importance of each predictor variable **when** predicting ChinaCropSM.

删除的内容: features

删除的内容: ,

删除的内容: Matlab

删除的内容: during

2.3.3 Evaluation metrics for validation and comparison

The in situ observations provide the most accurate SM, **and** all performance measures were calculated using the testing dataset for evaluation purposes. All SM products were evaluated against the in situ observations (testing dataset) according to three metrics: **root mean square error** (RMSE; m^3m^{-3}), bias (m^3m^{-3}), unbiased RMSE (ubRMSE; m^3m^{-3}), **explained** variance (R^2), and the correlation coefficient (r), which are defined as follows Eqs. (2)–(6):

$$r = \frac{1}{N-1} \sum_{i=1}^N \left(\frac{P_i - \bar{P}}{\sigma_P} \right) \left(\frac{O_i - \bar{O}}{\sigma_O} \right) \quad (2)$$

$$R^2 = 1 - \frac{\sum_{i=1}^N (P_i - O_i)^2}{N \sum_{i=1}^N (P_i - \bar{P})^2} \quad (3)$$

$$\text{bias} = \frac{1}{N} \sum_{i=1}^N (P_i - O_i) \quad (4)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2} \quad (5)$$

$$\text{ubRMSE} = \sqrt{\overline{\text{RMSE}^2} - \text{bias}^2} \quad (6)$$

where the overbar indicates the mean; P_i is the i^{th} prediction SM from products; O_i is the i^{th} in situ observation SM; N is the total number of observations; and σ_o and σ_p are the standard deviations of the in situ observed or predicted SM, respectively.

In addition, we compared our four **subsets of** data with RSSSM and ESA CCI SM separately **by** evaluating their spatial and temporal **accuracies** related to in situ surface SM observations (Tables S1 and S2).

We evaluated our irrigation factor forecasting model results using the receiver operating **characteristic** (ROC) curve and **the area under the curve** (AUC) (Table S4) (Fawcett, 2006). **Additionally**, we calculated UA (Eq. 7), PA (Eq. 8), and overall accuracy (Eq. 9) based on confusion matrices (Table S3) containing the percentages of the four possible outcomes of a model:

true positive (TP), **true negative** (TN), **false-positive** (FP) and **false negative** (FN) (Fawcett, 2006).

$$\text{PA} = \frac{TP}{TP+FP} \quad (7)$$

$$\text{UA} = \frac{TP}{TP+FN} \quad (8)$$

$$\text{Accuracy} = \frac{TP+TN}{TP+FP+TN+FN} \quad (9)$$

删除的内容: Root Mean Square Error

删除的内容: Explained

删除的内容: subset

删除的内容: through

删除的内容: accuracy

删除的内容: Table

删除的内容: characteristics

删除的内容: their Area Under the Curve

删除的内容: Also

删除的内容: True Positive

删除的内容: True Negative

删除的内容: False Positive

删除的内容: false positive

删除的内容: False Negative

3 Results and discussion

3.1 Validation of ChinaCropSM1 km products

The scatterplots between the predicted SM and those observations were displayed by soil layers and crops (Figure 3). We found that the SM predicted by the RF model agreed well with the in situ SM observations, with an ubRMSE of 0.028–0.037, bias of -0.0011–0.0009, and r from 0.925–0.944. Additionally, the mean bias in predicting SM for wheat was negative (Figure 3-a, b), while those for maize were positive (Figure 3-c, d). These findings suggest that maize SM was overestimated, while that for wheat was underestimated. The absolute values of mean bias and RMSE in predicting SM at topsoil depth (0–10 cm) for both crops were relatively larger (e.g., 0–10 cm). RMSE 0.036>0.028) than that for a soil depth of 10–20 cm. This result indicates that the RF model performed better in predicting the SM content in the 10–20 cm layer than in the 0–10 cm layer, which was consistent with previous studies (O and Orth, 2020).

3.2 The improvement of ChinaCropSM1 km products with an irrigation module

Interestingly, all prediction accuracies of SM were consistently improved both for crops and depths (Figure 4) compared with those without an irrigation module (Table S5). Specifically, R^2 values were increased by 6.8–9.7%, and RMSE were decreased by 16–23% (Table S5). Among these, R^2 values for maize SM were slightly improved compared with those of wheat, and RMSE for maize was decreased more than that of wheat. This finding further suggests that the irrigation water requirements of maize are higher than those of wheat, which is consistent with the fact that summer maize requires large amounts of water to produce high yields. (Mohammed Karrou, 2012).

3.3 The significant scores of different factors for simulating SM

It is critical to select which independent variables are involved in a model, neither too many nor too few, while simultaneously avoiding multicollinearity among them. We have deleted 7 variables due to their high correlations ($|r| > 0.5$) with the 11 variables selected (Figure S3 and S4). Surprisingly, the top first was scored to irrigation factor (CIR), followed by pre10 (ante-accumulated precipitation over ten days) and fc (field capacity) (Figure 5). Current daily precipitation shows significantly different importance on SM planted by wheat and maize, with a similarity for DOY. However, all other factors show less importance in SM simulations. Compared with the significant roles of precipitation-related variables (e.g., pre10, pre) on SM in most rainfall-fed areas, irrigation shows overwhelming impacts on dryland soil moisture across China (Qiu et al., 2016b). Such results highlight that more accurately monitoring management activities, including irrigation times, areas and quantities, will further improve irrigation modules, consequently improving SM simulations (Wu et al., 2020; Zhang et al., 2015, 2022).

3.4 The temporal and spatial patterns between ChinaCropSM1 km and the in situ SM observations

The SM values in ChinaCropSM1 km were significantly correlated with the in situ SM observations, with a mean r of 0.92, 0.94, 0.93 and 0.94 for wheat₀₋₁₀, wheat₁₀₋₂₀, maize₀₋₁₀ and maize₁₀₋₂₀ respectively, during the whole growing period (Figure 6).

刪除的內容: 1km

刪除的內容: validation

刪除的內容: Moreover

刪除的內容: were

刪除的內容: those

刪除的內容: were

刪除的內容: value

刪除的內容: top soil

刪除的內容: relative

刪除的內容: eg

刪除的內容: It

刪除的內容: at

刪除的內容: that at

刪除的內容: the

刪除的內容: 1km

刪除的內容: accuracy

刪除的內容: comparison of

刪除的內容: than

刪除的內容: Such

刪除的內容: for

刪除的內容: was

刪除的內容: reason

刪除的內容: into

刪除的內容: or

刪除的內容: less

刪除的內容: show

刪除的內容: the

刪除的內容: Nevertheless

刪除的內容: on

刪除的內容: simulation. Comparing

刪除的內容: however,

刪除的內容: result highlights

刪除的內容: irrigating

刪除的內容: quantity

刪除的內容: module

刪除的內容: improve SM simulation

刪除的內容: 1km

刪除的內容: 1km

刪除的內容: are

刪除的內容: , respectively.

The spatial coefficients for wheat at 10–20 cm ~~were~~ generally higher than the surface SM (0.94 vs. 0.92), and the two soil depths SM in April and September ~~were~~ significantly higher (Figure 6–a, b). We attributed the high spatial correlations of surface SM to irrigation impacts, because April and September are planting ~~times~~ for both spring and winter wheat. The better relationships further substantiated ~~that~~ the irrigation module developed in our SM model improves the simulation accuracy for surface SM. Consistently, the spatial coefficients for maize at ~~the 10–20 cm depth~~ ~~were~~ higher than those for ~~the 0–10 cm depth~~ (0.94 vs. 0.93) (Figure 7-c, d). At the sowing (Apr.), heading (Jul.), and milking (Aug.) stages, maize usually ~~demands~~ a large water supply. The spatial coefficient for maize SM at both soil ~~depths~~ from May to Aug. ~~was~~ lower than the mean value, potentially due to the lack of irrigation applications (Yin et al., 2016) (Figure 6).

We ~~further~~ analyzed the temporal pattern of SM accuracy in different regions (Figure 7). The median ~~r~~ values for ~~the~~ Huang-Huai-Hai Plain and ~~the northern arid and semiarid regions~~ were higher than ~~those~~ in other agricultural regions because of ~~the~~ larger training samples. Our findings further substantiated that a larger training sample size will cause a higher temporal accuracy, indicated by a higher r and a lower RMSE (Figure S6). However, the poor performance in ~~the~~ Yunnan-Guizhou Plateau might be caused by smaller training samples (Figure S6).

3.5 Comparisons between ChinaCropSM ~~1 km~~ and public global SM products

We further compared our ChinaCropSM ~~1 km~~ with the two popular products through evaluating their spatial-temporal accuracy related to in situ surface SM observations. We summarized their evaluation ~~indices~~ by individual product in Table 3, which ~~consistently~~ indicated ~~the~~ bolds for our ChinaCropSM ~~1 km~~ (all $r > 0.90$, RMSE < 0.04), while RSSSM and ESA CCI SM were shown by $r < 0.50$ and RMSE > 0.1 .

To match the different spatial resolutions of ~~the~~ three products, we calculated the averages of all in-site observations in the same pixel (e.g., 1 km, 27 km or 0.1°) to make their spatial-temporal ~~accuracies~~ comparable. Interestingly, all ~~indices~~ of our products were consistently indicated by the higher accuracy (e.g., r 0.94, bias 0.005, RMSE 0.034, ubRMSE 0.034) (Figure 9). ~~The~~ RSSSM dataset significantly underestimated SM with an averaged bias of -0.114 , ~~accompanied by~~ a higher RMSE of 0.150. ESA CCI SM performed better than RSSSM (e.g., RMSE 0.11 vs. 0.15) derived from Soil Moisture Active Passive (SMAP) (Entekhabi et al., 2010), and we ascribed such improvement partly into some corrections based on in situ observations for ESA CCI SM (Dorigo et al., 2017). Such results highlight ~~that~~ SM products derived solely from remote sensing satellites should be corrected with ground observations. ~~Additionally~~, neither RSSSM nor ESA CCI SM ~~considered~~ irrigation activities; thus, their spatial correlations with ground ~~observations~~ are incomparable to those of our products (r 0.944 vs. 0.381 0.256) (Figure 8). Our study ~~strongly~~ substantiates ~~that~~ an irrigation module should be taken into account when developing SM simulation models ~~for~~ producing SM products.

删除的内容: are

删除的内容: are

删除的内容: .

删除的内容: time

删除的内容: 10–20cm

删除的内容: are

删除的内容: 0–10cm

删除的内容: Figure7

删除的内容: demanded water supplying a lot

删除的内容: depth

删除的内容: were

删除的内容: still

删除的内容: of

删除的内容: Northern Arid and Semiarid region

删除的内容: that

删除的内容: 1km

删除的内容: 1km

删除的内容: indexes

删除的内容: consistently

删除的内容: 1km

删除的内容: accuracy

删除的内容: indexes

删除的内容: accompanying with

删除的内容: Moreover

删除的内容: had

删除的内容: the

删除的内容: .

删除的内容: observes

删除的内容: strongly

删除的内容: of

515 4 Data availability

The 1 km gridded daily soil moisture dataset for main crops (i.e., wheat and maize) dryland in China from 1993 to 2018 (ChinaCropSM1 km) are publicly available at <https://zenodo.org/record/6834530> (wheat₀₋₁₀) (Cheng et al., 2022a), <https://zenodo.org/record/6822591> (wheat₁₀₋₂₀) (Cheng et al., 2022b), <https://zenodo.org/record/6822581> (maize₀₋₁₀) (Cheng et al., 2022c) and <https://zenodo.org/record/6820166> (maize₁₀₋₂₀) (Cheng et al., 2022d).

520 5 Discussion and Conclusions

We developed a daily 1 km soil moisture dataset based on numerous field observations (181327 samples) from 1993–2018, which significantly enriches the current SM datasets available. ChinaCropSM1 km shows higher temporal-spatial resolution and accuracy than the popular global SM products. Additionally, to date, few studies have provided a daily SM product with such a higher resolution, combining different soil depths and an irrigation module. ChinaCropSM1 km is the first SM product with a higher spatial resolution (~1 km) at depths of 0–10 and 10–20 cm in croplands in China, by compiling ground observations and using the RF method.

Our ChinaCropSM1 km, predicted by the RF model agreed well with in situ SM observations (ubRMSE ranges from 0.028–0.037, bias ranges from –0.0011–0.0009, r ranges from 0.925–0.944, and R^2 ranges from 0.860–0.895). An irrigation module was first developed according to crop type (i.e., wheat, maize), soil depth (0–10 cm, 10–20 cm) and phenology. All prediction accuracies of SM were consistently improved (R^2 values were increased by 6.8–9.7%, RMSE were decreased by 16–23%) both for crops and depths. Additionally, ChinaCropSM1 km generally has advantages over other popular gridded SM products (RSSM and ESA CCI SM) through evaluating their spatial-temporal accuracy related to in situ SM as the benchmark. Our ChinaCropSM1 km has relatively higher accuracy (all $r > 0.90$, RMSE < 0.04), while RSSM and ESA CCI SM showed $r < 0.50$ and RMSE > 0.1 .

The ChinaCropSM dataset is credible and accurate according to the results compared with the public datasets; however, some limitations still exist in our study. First, the limited AMS irrigation records may lead to uncertainty in the irrigation factor predictions. More detailed irrigation information will help to improve irrigation module performance. Second, our method for generating cropland SM is applicable to other regions and crops, but more environmental variables will be increasingly required considering that SM variabilities are complex processes controlled by many factors (Famiglietti et al., 2008; Qin et al., 2013; Guevara and Vargas, 2019), especially for irrigation activities. For example, to more accurately characterize irrigation activities, many field samples are required at both spatial and temporal resolutions. Other auxiliary data on information on crop growth, classification, and management (e.g., irrigation frequency, amount and method) will benefit the development of our irrigation module and the accurate derivation of SM datasets. Third, to provide more extensive SM data as possible as we can, a constant layer integrated with all pixels planting wheat/maize during 2000–2015 (<http://dx.doi.org/10.17632/jbs44b2hrk.2>) was applied to generate our ChinaCropSM1 km. Such merged areas could lead to uncertainties in their spatial distributions because annual wheat/maize planting areas are dynamic over time. To avoid the

- 刪除的內容: The 1km1 km gridded daily Soil Moisture for Croplandssoil moisture for croplands in China (ChinaCropSM1 ...
- 刪除的內容: 1km
- 刪除的內容: enrich
- 刪除的內容: Our
- 刪除的內容: 1km
- 刪除的內容: Moreover
- 刪除的內容: 1km
- 刪除的內容: ~1km
- 刪除的內容:
- 刪除的內容: .
- 刪除的內容: depth
- 刪除的內容: croplands
- 刪除的內容: the
- 刪除的內容:
- 刪除的內容: 1km
- 刪除的內容: firstly
- 刪除的內容: i.e.
- 刪除的內容: accuracy
- 刪除的內容: Meanwhile
- 刪除的內容: 1km
- 刪除的內容: 1km
- 刪除的內容: were shown by
- 刪除的內容: are
- 刪除的內容: comparing
- 刪除的內容: .
- 刪除的內容: are
- 刪除的內容: existed
- 刪除的內容: the
- 刪除的內容: performances
- 刪除的內容: the
- 刪除的內容: characterize
- 刪除的內容: the
- 刪除的內容: highly
- 刪除的內容: in
- 刪除的內容: of
- 刪除的內容: managements
- 刪除的內容: to develop
- 刪除的內容: derive
- 刪除的內容: more accurately

595 uncertainties, potential users should mask our products with explicitly annual wheat/maize planting maps to obtain accurate
SM data including spatial dynamic information. Fourth, different splitting methods during training and testing affect model
performance. Selecting a splitting method to improve the generalization performance is dependent on the data. Generally, the
larger the size of the data, the smaller the effect of the splitting methods on the results (Birba, 2020). Additionally, advanced
algorithms will be potential alternatives for random forest due to their strong dependence on inputs (Breiman, 2001; Rasmussen,
2004). Improving irrigation modules should focus on details such as irrigation amount and frequency, which will significantly
600 help to verify and improve the accuracy of both irrigation and SM predictions. We anticipate that a more accurate SM dataset
will be produced by applying the approach to other crops and areas in the future with all the above improvements.

刪除的內容: Third

刪除的內容: do

刪除的內容: which

刪除的內容: Moreover

刪除的內容: its

刪除的內容: module

刪除的內容: be focused

刪除的內容: are sure

刪除的內容: into

Author contributions

605 FC, HZ, ZZ, JH, JC, YL, LZ, JZ and JX contributed to the design of this research; FC and ZZ collectively prepared the
manuscript with contributions from all coauthors; JH, JC, YL, LZ, JX and JZ revised the manuscript; FC and HZ developed
the model code.

刪除的內容: co-authors

Competing interests

The authors declare that they have no conflicts of interest.

刪除的內容: conflict

Financial support

610 This study was supported by the National Key Research and Development Program of China (grant no. 2020YFA0608201)
and the National Natural Science Foundation of China (grant no. 41977405).

References

615 Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., and Hegewisch, K. C.: TerraClimate, a high-resolution global dataset of
monthly climate and climatic water balance from 1958–2015, Sci Data, 5, 170191, <https://doi.org/10.1038/sdata.2017.191>,
2018.

Ahmad, S., Kalra, A., and Stephen, H.: Estimating soil moisture using remote sensing data: A machine learning approach,
Advances in Water Resources, 33, 69–80, <https://doi.org/10.1016/j.advwatres.2009.10.008>, 2010.

Ahmed, A. A. M., Deo, R. C., Raj, N., Ghahramani, A., Feng, Q., Yin, Z., and Yang, L.: Deep Learning Forecasts of Soil
Moisture: Convolutional Neural Network and Gated Recurrent Unit Models Coupled with Satellite-Derived MODIS,
Observations and Synoptic-Scale Climate Index Data, Remote Sensing, 13, 554, <https://doi.org/10.3390/rs13040554>, 2021.

- Albergel, C., Dorigo, W., Reichle, R. H., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., Isaksen, L., de Jeu, R., and Wagner, W.: Skill and Global Trend Analysis of Soil Moisture from Reanalyses and Microwave Remote Sensing, *Journal of Hydrometeorology*, 14, 1259–1277, <https://doi.org/10.1175/JHM-D-12-0161.1>, 2013.
- Amazirh, A., Merlin, O., Er-Raki, S., Gao, Q., Rivalland, V., Malbeteau, Y., Khabba, S., and Escorihuela, M. J.: Retrieving surface soil moisture at high spatio-temporal resolution from a synergy between Sentinel-1 radar and Landsat thermal data: A study case over bare soil, *Remote Sensing of Environment*, 211, 321–337, <https://doi.org/10.1016/j.rse.2018.04.013>, 2018.
- Birba, D. E.: A Comparative study of data splitting algorithms for machine learning model selection, 2020.
- Bogena, H. R., Huisman, J. A., Oberdörster, C., and Vereecken, H.: Evaluation of a low-cost soil water content sensor for wireless network applications, *Journal of Hydrology*, 344, 32–42, <https://doi.org/10.1016/j.jhydrol.2007.06.032>, 2007.
- Breiman, L.: Random forests, *Machine learning*, 45, 5–32, 2001.
- Brownlee, J.: *Machine learning mastery with python*, Machine Learning Mastery Pty Ltd, 527, 100–120, 2016.
- Chakrabarti, S., Bongiovanni, T., Judge, J., Zotarelli, L., and Bayer, C.: Assimilation of SMOS Soil Moisture for Quantifying Drought Impacts on Crop Yield in Agricultural Regions, *IEEE J. Sel. Top. Appl. Earth Observations Remote Sensing*, 7, 3867–3879, <https://doi.org/10.1109/JSTARS.2014.2315999>, 2014.
- Chen, L. and Dirmeyer, P. A.: Global observed and modelled impacts of irrigation on surface temperature, *Int J Climatol*, 39, 2587–2600, <https://doi.org/10.1002/joc.5973>, 2019.
- Chen, Y., Feng, X., and Fu, B.: An improved global remote-sensing-based surface soil moisture (RSSSM) dataset covering 2003–2018, *Earth Syst. Sci. Data*, 13, 1–31, <https://doi.org/10.5194/essd-13-1-2021>, 2021.
- Cheng, F., Zhang, Z., Huimin Zhuang, Jichong Han, Yuchuan Luo, Cao, J., Liangliang Zhang, Zhang, J., Fulu Tao, and Jialu Xu: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for dryland wheat and maize across China during 1993–2018 (wheat0-10cm), <https://doi.org/10.5281/ZENODO.6834530>, 2022a.
- Cheng, F., Zhang, Z., Huimin Zhuang, Jichong Han, Yuchuan Luo, Cao, J., Liangliang Zhang, Zhang, J., Fulu Tao, and Jialu Xu: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for dryland wheat and maize across China during 1993–2018 (wheat10-20cm), <https://doi.org/10.5281/ZENODO.6822591>, 2022b.
- Cheng, F., Zhang, Z., Huimin Zhuang, Jichong Han, Yuchuan Luo, Cao, J., Liangliang Zhang, Zhang, J., Fulu Tao, and Jialu Xu: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for dryland wheat and maize across China during 1993–2018 (maize0-10cm), <https://doi.org/10.5281/ZENODO.6822581>, 2022c.
- Cheng, F., Zhang, Z., Huimin Zhuang, Jichong Han, Yuchuan Luo, Cao, J., Liangliang Zhang, Zhang, J., Fulu Tao, and Jialu Xu: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for dryland wheat and maize across China during 1993–2018 (maize10-20cm), <https://doi.org/10.5281/ZENODO.6820166>, 2022d.
- Collow, T. W., Robock, A., Basara, J. B., and Illston, B. G.: Evaluation of SMOS retrievals of soil moisture over the central United States with currently available in situ observations: EVALUATION OF SMOS WITH IN SITU DATA, *J. Geophys. Res.*, 117, n/a-n/a, <https://doi.org/10.1029/2011JD017095>, 2012.

Crow, W. T., Berg, A. A., Cosh, M. H., Loew, A., Mohanty, B. P., Panciera, R., de Rosnay, P., Ryu, D., and Walker, J. P.:
665 Upscaling sparse ground-based soil moisture observations for the validation of coarse-resolution satellite soil moisture
products: UPSCALING SOIL MOISTURE, *Rev. Geophys.*, 50, <https://doi.org/10.1029/2011RG000372>, 2012.

Danielsson, P.-E.: Euclidean distance mapping, *Computer Graphics and Image Processing*, 14, 227–248,
[https://doi.org/10.1016/0146-664X\(80\)90054-4](https://doi.org/10.1016/0146-664X(80)90054-4), 1980.

Díaz-Uriarte, R. and Alvarez de Andrés, S.: Gene selection and classification of microarray data using random forest, *BMC*
670 *Bioinformatics*, 7, 3, <https://doi.org/10.1186/1471-2105-7-3>, 2006.

Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M., Forkel, M., Gruber, A.,
Haas, E., Hamer, P. D., Hirschi, M., Ikonen, J., de Jeu, R., Kidd, R., Lahoz, W., Liu, Y. Y., Miralles, D., Mistelbauer, T.,
Nicolai-Shaw, N., Parinussa, R., Pratola, C., Reimer, C., van der Schalie, R., Seneviratne, S. I., Smolander, T., and Lecomte,
P.: ESA CCI Soil Moisture for improved Earth system understanding: State-of-the art and future directions, *Remote Sensing*
675 *of Environment*, 203, 185–215, <https://doi.org/10.1016/j.rse.2017.07.001>, 2017.

Dorigo, W. A., Gruber, A., De Jeu, R. A. M., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca, L., Chung, D., Parinussa,
R. M., and Kidd, R.: Evaluation of the ESA CCI soil moisture product using ground-based observations, *Remote Sensing of*
Environment, 162, 380–395, <https://doi.org/10.1016/j.rse.2014.07.023>, 2015.

Drewniak, B., Song, J., Prell, J., Kotamarthi, V. R., and Jacob, R.: Modeling agriculture in the Community Land Model, *Geosci.*
680 *Model Dev.*, 6, 495–515, <https://doi.org/10.5194/gmd-6-495-2013>, 2013.

Eagleman, J. R. and Lin, W. C.: Remote sensing of soil moisture by a 21-cm passive radiometer, *J. Geophys. Res.*, 81, 3660–
3666, <https://doi.org/10.1029/JC081i021p03660>, 1976.

Entekhabi, D., Njoku, E. G., O’Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., Entin, J. K., Goodman, S. D.,
Jackson, T. J., Johnson, J., Kimball, J., Piepmeier, J. R., Koster, R. D., Martin, N., McDonald, K. C., Moggaddam, M., Moran,
685 S., Reichle, R., Shi, J. C., Spencer, M. W., Thurman, S. W., Tsang, L., and Van Zyl, J.: The Soil Moisture Active Passive
(SMAP) Mission, *Proc. IEEE*, 98, 704–716, <https://doi.org/10.1109/JPROC.2010.2043918>, 2010.

Famiglietti, J. S., Ryu, D., Berg, A. A., Rodell, M., and Jackson, T. J.: Field observations of soil moisture variability across
scales: SOIL MOISTURE VARIABILITY ACROSS SCALES, *Water Resour. Res.*, 44,
<https://doi.org/10.1029/2006WR005804>, 2008.

690 Fang, B., Lakshmi, V., Bindlish, R., Jackson, T. J., and Liu, P.-W.: Evaluation and validation of a high spatial resolution
satellite soil moisture product over the Continental United States, *Journal of Hydrology*, 588, 125043,
<https://doi.org/10.1016/j.jhydrol.2020.125043>, 2020.

FAOSTAT (Food and Agriculture Organization Corporate Statistical Database): FAO online database, available
at: <http://www.fao.org/faostat/en/#data/QCL> (last access: October 2021), 2019, Crops and livestock products.

695 Fawcett, T.: An introduction to ROC analysis, *Pattern Recognition Letters*, 27, 861–874,
<https://doi.org/10.1016/j.patrec.2005.10.010>, 2006.

Gruber, A., Scanlon, T., van der Schalie, R., Wagner, W., and Dorigo, W.: Evolution of the ESA CCI Soil Moisture climate data records and their underlying merging methodology, *Earth Syst. Sci. Data*, 11, 717–739, <https://doi.org/10.5194/essd-11-717-2019>, 2019.

700 Guevara, M. and Vargas, R.: Downscaling satellite soil moisture using geomorphometry and machine learning, *PLoS ONE*, 14, e0219639, <https://doi.org/10.1371/journal.pone.0219639>, 2019.

Guevara, M., Taufer, M., and Vargas, R.: Gap-free global annual soil moisture: 15 km grids for 1991–2018, *Earth Syst. Sci. Data*, 13, 1711–1735, <https://doi.org/10.5194/essd-13-1711-2021>, 2021.

705 Hengl, T. and Gupta, S.: Soil water content (volumetric %) for 33kPa and 1500kPa suctions predicted at 6 standard depths (0, 10, 30, 60, 100 and 200 cm) at 250 m resolution (v0.1), <https://doi.org/10.5281/ZENODO.2629589>, 2019.

Hengl, T., Heuvelink, G. B. M., Kempen, B., Leenaars, J. G. B., Walsh, M. G., Shepherd, K. D., Sila, A., MacMillan, R. A., Mendes de Jesus, J., Tamene, L., and Tondoh, J. E.: Mapping Soil Properties of Africa at 250 m Resolution: Random Forests Significantly Improve Current Predictions, *PLoS ONE*, 10, e0125814, <https://doi.org/10.1371/journal.pone.0125814>, 2015.

710 Huang, S., Krysanova, V., Zhai, J., and Su, B.: Impact of Intensive Irrigation Activities on River Discharge Under Agricultural Scenarios in the Semi-Arid Aksu River Basin, Northwest China, *Water Resour Manage.*, 29, 945–959, <https://doi.org/10.1007/s11269-014-0853-2>, 2015.

Hutengs, C. and Vohland, M.: Downscaling land surface temperatures at regional scales with random forest regression, *Remote Sensing of Environment*, 178, 127–141, <https://doi.org/10.1016/j.rse.2016.03.006>, 2016.

715 Im, J., Park, S., Rhee, J., Baik, J., and Choi, M.: Downscaling of AMSR-E soil moisture with MODIS products using machine learning approaches, *Environ Earth Sci*, 75, 1120, <https://doi.org/10.1007/s12665-016-5917-6>, 2016.

Jackson, T. J., Schmugge, T. J., and Wang, J. R.: Passive microwave sensing of soil moisture under vegetation canopies, *Water Resour. Res.*, 18, 1137–1142, <https://doi.org/10.1029/WR018i004p01137>, 1982.

720 Jeong, J. H., Resop, J. P., Mueller, N. D., Fleisher, D. H., Yun, K., Butler, E. E., Timlin, D. J., Shim, K.-M., Gerber, J. S., Reddy, V. R., and Kim, S.-H.: Random Forests for Global and Regional Crop Yield Predictions, *PLoS ONE*, 11, e0156571, <https://doi.org/10.1371/journal.pone.0156571>, 2016.

Lacava, T., Matgen, P., Brocca, L., Bittelli, M., Pergola, N., Moramarco, T., and Tramutoli, V.: A First Assessment of the SMOS Soil Moisture Product With In Situ and Modeled Data in Italy and Luxembourg, *IEEE Trans. Geosci. Remote Sensing*, 50, 1612–1622, <https://doi.org/10.1109/TGRS.2012.2186819>, 2012.

725 Lagomarsino, D., Tofani, V., Segoni, S., Catani, F., and Casagli, N.: A Tool for Classification and Regression Using Random Forest Methodology: Applications to Landslide Susceptibility Mapping and Soil Thickness Modeling, *Environ Model Assess.*, 22, 201–214, <https://doi.org/10.1007/s10666-016-9538-y>, 2017.

Lawston, P. M., Santanello, J. A., and Kumar, S. V.: Irrigation Signals Detected From SMAP Soil Moisture Retrievals: Irrigation Signals Detected From SMAP, *Geophys. Res. Lett.*, 44, 11,860–11,867, <https://doi.org/10.1002/2017GL075733>, 2017.

- 730 [Li, H., Robock, A., Liu, S., Mo, X., and Viterbo, P.: Evaluation of Reanalysis Soil Moisture Simulations Using Updated Chinese Soil Moisture Observations, *Journal of Hydrometeorology*, 6, 180–193, <https://doi.org/10.1175/JHM416.1>, 2005.](#)
[Li, Z., Zhang, Z., and Zhang, L.: Improving regional wheat drought risk assessment for insurance application by integrating scenario-driven crop model, machine learning, and satellite data, *Agricultural Systems*, 191, 103141, <https://doi.org/10.1016/j.agsy.2021.103141>, 2021.](#)
- 735 [Liu, Y., Yang, Y., and Yue, X.: Evaluation of Satellite-Based Soil Moisture Products over Four Different Continental In-Situ Measurements, *Remote Sensing*, 10, 1161, <https://doi.org/10.3390/rs10071161>, 2018.](#)
[Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., van Dijk, A. I. J. M., McCabe, M. F., and Evans, J. P.: Developing an improved soil moisture dataset by blending passive and active microwave satellite-based retrievals, *Hydrol. Earth Syst. Sci.*, 15, 425–436, <https://doi.org/10.5194/hess-15-425-2011>, 2011.](#)
- 740 [Liu, Y. Y., Dorigo, W. A., Parinussa, R. M., de Jeu, R. A. M., Wagner, W., McCabe, M. F., Evans, J. P., and van Dijk, A. I. J. M.: Trend-preserving blending of passive and active microwave soil moisture retrievals, *Remote Sensing of Environment*, 123, 280–297, <https://doi.org/10.1016/j.rse.2012.03.014>, 2012.](#)
[Llamas, R. M., Guevara, M., Rorabaugh, D., Taufer, M., and Vargas, R.: Spatial Gap-Filling of ESA CCI Satellite-Derived Soil Moisture Based on Geostatistical Techniques and Multiple Regression, *Remote Sensing*, 12, 665, <https://doi.org/10.3390/rs12040665>, 2020.](#)
- 745 [Loew, A., Stacke, T., Dorigo, W., de Jeu, R., and Hagemann, S.: Potential and limitations of multidecadal satellite soil moisture observations for selected climate model evaluation studies, *Hydrol. Earth Syst. Sci.*, 17, 3523–3542, <https://doi.org/10.5194/hess-17-3523-2013>, 2013.](#)
[Luo, Y., Zhang, Z., Chen, Y., Li, Z., and Tao, F.: ChinaCropPhen1km: a high-resolution crop phenological dataset for three staple crops in China during 2000–2015 based on leaf area index \(LAI\) products, *Earth Syst. Sci. Data*, 12, 197–214, <https://doi.org/10.5194/essd-12-197-2020>, 2020a.](#)
[Luo, Y., Zhang, Z., Li, Z., Chen, Y., Zhang, L., Cao, J., and Tao, F.: Identifying the spatiotemporal changes of annual harvesting areas for three staple crops in China by integrating multi-data sources, *Environ. Res. Lett.*, 15, 074003, <https://doi.org/10.1088/1748-9326/ab80f0>, 2020b.](#)
- 750 [Luo, Y., Zhang, Z., Li, Z., Chen, Y., Zhang, L., Cao, J., and Tao, F.: Identifying the spatiotemporal changes of annual harvesting areas for three staple crops in China by integrating multi-data sources, *Environ. Res. Lett.*, 15, 074003, <https://doi.org/10.1088/1748-9326/ab80f0>, 2020b.](#)
- 755 [Mallick, K., Bhattacharya, B. K., and Patel, N. K.: Estimating volumetric surface moisture content for cropped soils using a soil wetness index based on surface temperature and NDVI, *Agricultural and Forest Meteorology*, 149, 1327–1342, <https://doi.org/10.1016/j.agrformet.2009.03.004>, 2009.](#)
[Meng, X., Mao, K., Meng, F., Shi, J., Zeng, J., Shen, X., Cui, Y., Jiang, L., and Guo, Z.: A fine-resolution soil moisture dataset for China in 2002–2018, *Earth Syst. Sci. Data*, 13, 3239–3261, <https://doi.org/10.5194/essd-13-3239-2021>, 2021.](#)
- 760 [Mohammed Karrou: Yield and water productivity of maize and wheat under deficit and raised bed irrigation practices in Egypt, *Afr. J. Agric. Res.*, 7, <https://doi.org/10.5897/AJAR11.2109>, 2012.](#)
[Mohanty, B. P., Cosh, M. H., Lakshmi, V., and Montzka, C.: Soil Moisture Remote Sensing: State-of-the-Science, *Vadose Zone Journal*, 16, vzj2016.10.0105, <https://doi.org/10.2136/vzj2016.10.0105>, 2017.](#)

- 765 [O. S. and Orth, R.: Global soil moisture from in-situ measurements using machine learning -- SoMo.ml, http://arxiv.org/abs/2010.02374, 5 October 2020.](http://arxiv.org/abs/2010.02374)
- [Peng, J., Loew, A., Merlin, O., and Verhoest, N. E. C.: A review of spatial downscaling of satellite remotely sensed soil moisture: Downscale Satellite-Based Soil Moisture, *Rev. Geophys.*, 55, 341–366, https://doi.org/10.1002/2016RG000543, 2017.](https://doi.org/10.1002/2016RG000543)
- 770 [Petropoulos, G. P., Ireland, G., and Barrett, B.: Surface soil moisture retrievals from remote sensing: Current status, products & future trends, *Physics and Chemistry of the Earth, Parts A/B/C*, 83–84, 36–56, https://doi.org/10.1016/j.pce.2015.02.009, 2015.](https://doi.org/10.1016/j.pce.2015.02.009)
- [Prasad, A. K., Chai, L., Singh, R. P., and Kafatos, M.: Crop yield estimation model for Iowa using remote sensing and surface parameters, *International Journal of Applied Earth Observation and Geoinformation*, 8, 26–33, https://doi.org/10.1016/j.jag.2005.06.002, 2006.](https://doi.org/10.1016/j.jag.2005.06.002)
- 775 [Preimesberger, W., Scanlon, T., Su, C.-H., Gruber, A., and Dorigo, W.: Homogenization of Structural Breaks in the Global ESA CCI Soil Moisture Multisatellite Climate Data Record, *IEEE Trans. Geosci. Remote Sensing*, 59, 2845–2862, https://doi.org/10.1109/TGRS.2020.3012896, 2021.](https://doi.org/10.1109/TGRS.2020.3012896)
- [Qin, J., Yang, K., Lu, N., Chen, Y., Zhao, L., and Han, M.: Spatial upscaling of in-situ soil moisture measurements based on MODIS-derived apparent thermal inertia, *Remote Sensing of Environment*, 138, 1–9, https://doi.org/10.1016/j.rse.2013.07.003, 2013.](https://doi.org/10.1016/j.rse.2013.07.003)
- 780 [Qiu, J., Gao, Q., Wang, S., and Su, Z.: Comparison of temporal trends from multiple soil moisture data sets and precipitation: The implication of irrigation on regional soil moisture trend, *International Journal of Applied Earth Observation and Geoinformation*, 48, 17–27, https://doi.org/10.1016/j.jag.2015.11.012, 2016a.](https://doi.org/10.1016/j.jag.2015.11.012)
- 785 [Qiu, J., Gao, Q., Wang, S., and Su, Z.: Comparison of temporal trends from multiple soil moisture data sets and precipitation: The implication of irrigation on regional soil moisture trend, *International Journal of Applied Earth Observation and Geoinformation*, 48, 17–27, https://doi.org/10.1016/j.jag.2015.11.012, 2016b.](https://doi.org/10.1016/j.jag.2015.11.012)
- [Rasmussen, C. E.: Gaussian Processes in Machine Learning, in: *Advanced Lectures on Machine Learning*, vol. 3176, edited by: Bousquet, O., von Luxburg, U., and Rätsch, G., Springer Berlin Heidelberg, Berlin, Heidelberg, 63–71, https://doi.org/10.1007/978-3-540-28650-9_4, 2004.](https://doi.org/10.1007/978-3-540-28650-9_4)
- 790 [Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., and Toll, D.: The Global Land Data Assimilation System, *Bull. Amer. Meteor. Soc.*, 85, 381–394, https://doi.org/10.1175/BAMS-85-3-381, 2004.](https://doi.org/10.1175/BAMS-85-3-381)
- [Schmugge, T. J., Kustas, W. P., Ritchie, J. C., Jackson, T. J., and Rango, A.: Remote sensing in hydrology, *Advances in Water Resources*, 25, 1367–1385, https://doi.org/10.1016/S0309-1708\(02\)00065-9, 2002.](https://doi.org/10.1016/S0309-1708(02)00065-9)
- 795 [Sheffield, J.: A simulated soil moisture based drought analysis for the United States, *J. Geophys. Res.*, 109, D24108, https://doi.org/10.1029/2004JD005182, 2004.](https://doi.org/10.1029/2004JD005182)

Shen, Y., Zhang, Y., R. Scanlon, B., Lei, H., Yang, D., and Yang, F.: Energy/water budgets and productivity of the typical croplands irrigated with groundwater and surface water in the North China Plain, *Agricultural and Forest Meteorology*, 181, 133–142, <https://doi.org/10.1016/j.agrformet.2013.07.013>, 2013.

800 Srivastava, P. K., Han, D., Ramirez, M. R., and Islam, T.: Machine Learning Techniques for Downscaling SMOS Satellite Soil Moisture Using MODIS Land Surface Temperature for Hydrological Application, *Water Resour Manage*, 27, 3127–3144, <https://doi.org/10.1007/s11269-013-0337-9>, 2013.

Su, C.-H., Zhang, J., Gruber, A., Parinussa, R., Ryu, D., Crow, W. T., and Wagner, W.: Error decomposition of nine passive and active microwave satellite soil moisture data sets over Australia, *Remote Sensing of Environment*, 182, 128–140, <https://doi.org/10.1016/j.rse.2016.05.008>, 2016.

805 Sun, C., Bian, Y., Zhou, T., and Pan, J.: Using of Multi-Source and Multi-Temporal Remote Sensing Data Improves Crop-Type Mapping in the Subtropical Agriculture Region, *Sensors*, 19, 2401, <https://doi.org/10.3390/s19102401>, 2019.

Tao, F., Yokozawa, M., Hayashi, Y., and Lin, E.: Changes in agricultural water demands and soil moisture in China over the last half-century and their effects on agricultural production, *Agricultural and Forest Meteorology*, 118, 251–261, [https://doi.org/10.1016/S0168-1923\(03\)00107-2](https://doi.org/10.1016/S0168-1923(03)00107-2), 2003.

810 Thornthwaite, C. W.: An Approach toward a Rational Classification of Climate, *Geographical Review*, 38, 55, <https://doi.org/10.2307/210739>, 1948.

Vergopolan, N., Chaney, N. W., Beck, H. E., Pan, M., Sheffield, J., Chan, S., and Wood, E. F.: Combining hyper-resolution land surface modeling with SMAP brightness temperatures to obtain 30-m soil moisture estimates, *Remote Sensing of Environment*, 242, 111740, <https://doi.org/10.1016/j.rse.2020.111740>, 2020.

815 Wagner, W., Dorigo, W., de Jeu, R., Fernandez, D., Benveniste, J., Haas, E., and Ertl, M.: FUSION OF ACTIVE AND PASSIVE MICROWAVE OBSERVATIONS TO CREATE AN ESSENTIAL CLIMATE VARIABLE DATA RECORD ON SOIL MOISTURE, *ISPRS Ann. Photogramm. Remote Sens. Spatial Inf. Sci.*, 1–7, 315–321, <https://doi.org/10.5194/isprsannals-I-7-315-2012>, 2012.

820 Walker, J. P., Willgoose, G. R., and Kalma, J. D.: In situ measurement of soil moisture: a comparison of techniques, *Journal of Hydrology*, 293, 85–99, <https://doi.org/10.1016/j.jhydrol.2004.01.008>, 2004.

Wang, C., Wang, Z.-H., and Yang, J.: Urban water capacity: Irrigation for heat mitigation, *Computers, Environment and Urban Systems*, 78, 101397, <https://doi.org/10.1016/j.compenvurbsys.2019.101397>, 2019.

825 Wang, L. and Qu, J. J.: Satellite remote sensing applications for surface soil moisture monitoring: A review, *Front. Earth Sci. China*, 3, 237–247, <https://doi.org/10.1007/s11707-009-0023-7>, 2009.

Wei, Z., Meng, Y., Zhang, W., Peng, J., and Meng, L.: Downscaling SMAP soil moisture estimation with gradient boosting decision tree regression over the Tibetan Plateau, *Remote Sensing of Environment*, 225, 30–44, <https://doi.org/10.1016/j.rse.2019.02.022>, 2019.

Wigneron, J.-P., Calvet, J.-C., Pellarin, T., Van de Griend, A. A., Berger, M., and Ferrazzoli, P.: Retrieving near-surface soil moisture from microwave radiometric observations: current status and future plans, *Remote Sensing of Environment*, 85, 489–506, [https://doi.org/10.1016/S0034-4257\(03\)00051-8](https://doi.org/10.1016/S0034-4257(03)00051-8), 2003.

Wu, B. and Li, Q.: Crop planting and type proportion method for crop acreage estimation of complex agricultural landscapes, *International Journal of Applied Earth Observation and Geoinformation*, 16, 101–112, <https://doi.org/10.1016/j.jag.2011.12.006>, 2012.

Wu, B., Ma, Z., and Yan, N.: Agricultural drought mitigating indices derived from the changes in drought characteristics, *Remote Sensing of Environment*, 244, 111813, <https://doi.org/10.1016/j.rse.2020.111813>, 2020.

Yilmaz, M. T., Crow, W. T., Anderson, M. C., and Hain, C.: An objective methodology for merging satellite- and model-based soil moisture products: OBJECTIVELY MERGING SOIL MOISTURE PRODUCTS, *Water Resour. Res.*, 48, <https://doi.org/10.1029/2011WR011682>, 2012.

Yin, X. G., Jabloun, M., Olesen, J. E., Öztürk, I., Wang, M., and Chen, F.: Effects of climatic factors, drought risk and irrigation requirement on maize yield in the Northeast Farming Region of China, *J. Agric. Sci.*, 154, 1171–1189, <https://doi.org/10.1017/S0021859616000150>, 2016.

Zeng, J., Li, Z., Chen, Q., Bi, H., Qiu, J., and Zou, P.: Evaluation of remotely sensed and reanalysis soil moisture products over the Tibetan Plateau using in-situ observations, *Remote Sensing of Environment*, 163, 91–110, <https://doi.org/10.1016/j.rse.2015.03.008>, 2015.

Zhang, D., Qian, L., Mao, B., Huang, C., Huang, B., and Si, Y.: A Data-Driven Design for Fault Detection of Wind Turbines Using Random Forests and XGboost, *IEEE Access*, 6, 21020–21031, <https://doi.org/10.1109/ACCESS.2018.2818678>, 2018.

Zhang, Q., Sun, P., Li, J., Singh, V. P., and Liu, J.: Spatiotemporal properties of droughts and related impacts on agriculture in Xinjiang, China: Spatiotemporal properties of droughts and related impacts, *Int. J. Climatol.*, 35, 1254–1266, <https://doi.org/10.1002/joc.4052>, 2015.

Zhang, Q., Shi, R., Singh, V. P., Xu, C.-Y., Yu, H., Fan, K., and Wu, Z.: Droughts across China: Drought factors, prediction and impacts, *Science of The Total Environment*, 803, 150018, <https://doi.org/10.1016/j.scitotenv.2021.150018>, 2022.

Zhang, Z., Li, Z., Chen, Y., Zhang, L., and Tao, F.: Improving regional wheat yields estimations by multi-step-assimilating of a crop model with multi-source data, *Agricultural and Forest Meteorology*, 290, 107993, <https://doi.org/10.1016/j.agrformet.2020.107993>, 2020.

Zhu, X., Li, Y., Li, M., Pan, Y., and Shi, P.: Agricultural irrigation in China, *Journal of Soil and Water Conservation*, 68, 147A-154A, <https://doi.org/10.2489/jswc.68.6.147A>, 2013.

删除的内容: Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., and Hegewisch, K. C.: TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015, *Sci Data*, 5, 170191, <https://doi.org/10.1038/sdata.2017.191>, 2018.

Ahmad, S., Kalra, A., and Stephen, H.: Estimating soil moisture using remote sensing data: A machine learning approach, *Advances in Water Resources*, 33, 69–80, <https://doi.org/10.1016/j.advwatres.2009.10.008>, 2010.

Ahmed, A. A. M., Deo, R. C., Raj, N., Ghahramani, A., Feng, Q., Yin, Z., and Yang, L.: Deep Learning Forecasts of Soil Moisture: Convolutional Neural Network and Gated Recurrent Unit Models Coupled with Satellite-Derived MODIS, Observations and Synoptic-Scale Climate Index Data, *Remote Sensing*, 13, 554, <https://doi.org/10.3390/rs13040554>, 2021.

Albergel, C., Dorigo, W., Reichle, R. H., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., Isaksen, I., de Jeu, R., and Wagner, W.: Skill and Global Trend Analysis of Soil Moisture from Reanalyses and Microwave Remote Sensing, *Journal of Hydrometeorology*, 14, 1259–1277, <https://doi.org/10.1175/JHM-D-12-0161.1>, 2013.

Amazirh, A., Merlin, O., Er-Raki, S., Gao, Q., Rivalland, V., Malbetreau, Y., Khabba, S., and Escorihuela, M. J.: Retrieving surface soil moisture at high spatio-temporal resolution from a synergy between Sentinel-1 radar and Landsat thermal data: A study case over bare soil, *Remote Sensing of Environment*, 211, 321–337, <https://doi.org/10.1016/j.rse.2018.04.013>, 2018.

Birba, D. E.: A Comparative study of data splitting algorithms for machine learning model selection, 2020.

Bogena, H. R., Huisman, J. A., Oberdörster, C., and Vereecken, H.: Evaluation of a low-cost soil water content sensor for wireless network applications, *Journal of Hydrology*, 344, 32–42, <https://doi.org/10.1016/j.jhydrol.2007.06.032>, 2007.

Breiman, L.: Random forests, *Machine learning*, 45, 5–32, 2001.

Brownlee, J.: Machine learning mastery with python, Machine Learning Mastery Pty Ltd, 527, 100–120, 2016.

Chakrabarti, S., Bongiovanni, T., Judge, J., Zotarelli, L., and Bayer, C.: Assimilation of SMOS Soil Moisture for Quantifying Drought Impacts on Crop Yield in Agricultural Regions, *IEEE J. Sel. Top. Appl. Earth Observations Remote Sensing*, 7, 3867–3879, <https://doi.org/10.1109/JSTARS.2014.2315999>, 2014.

Chen, L. and Dirmeyer, P. A.: Global observed and modelled impacts of irrigation on surface temperature, *Int J Climatol*, 39, 2587–2600, <https://doi.org/10.1002/joc.5973>, 2019.


Chen, Y., Feng, X., and Fu, B.: An improved global remote-sensing-based surface soil moisture (RSSM) dataset covering 2003–2018, *Earth Syst. Sci. Data*, 13, 1–31, <https://doi.org/10.5194/essd-13-1-2021>, 2021.

Cheng, F., Zhang, Z., Zhuang, H., Han, J., Luo, Y., Cao, J., Zhang, L., Zhang, J., Tao, F., and Xu, J.: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for Crop drylands across China during 1993–2018, <https://doi.org/10.5281/ZENODO.6834530>, 2022a.

Cheng, F., Zhang, Z., Zhuang, H., Han, J., Luo, Y., Cao, J., Zhang, L., Zhang, J., Tao, F., and Xu, J.: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for Crop drylands across China during 1993–2018, <https://doi.org/10.5281/ZENODO.6822591>, 2022b.

Cheng, F., Zhang, Z., Zhuang, H., Han, J., Luo, Y., Cao, J., Zhang, L., Zhang, J., Tao, F., and Xu, J.: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for Crop drylands across China during 1993–2018, <https://doi.org/10.5281/ZENODO.6822581>, 2022c.

Cheng, F., Zhang, Z., Zhuang, H., Han, J., Luo, Y., Cao, J., Zhang, L., Zhang, J., Tao, F., and Xu, J.: ChinaCropSM1km: a fine 1km daily Soil Moisture dataset for Crop drylands across China during 1993–2018, <https://doi.org/10.5281/ZENODO.6820166>, 2022d.



995 Figure 1 Study areas and SM in situ field monitoring sites in China. NAS: northern arid and semiarid region; LP: Loess Plateau; HP: Huang-Huai-Hai Plain; SCB: Sichuan Basin; MYP: middle-lower Yangtze Plain; YGP: Yunnan-Guizhou Plateau and southern China; QT: Qinghai-Tibet region; ChinaCropland 1 km: the harvesting areas of crops across mainland China.

删除的内容: the

删除的内容: Northern Arid and Semiarid

删除的内容: Middle-lower

删除的内容: 1km

Table 1 Environmental factors used in the study, including meteorological data (MD), day of year (DOY), classified irrigation (CIR), soil properties (SP), remote sensing data (RSD), and geographical information (GI).

Data type	Variable	Data description	Temporal resolution	Spatial resolution
MD	pre	daily precipitation	daily	1 km
	pre10	ante-accumulated precipitation over ten days	daily	1 km
DOY	DOY	day of year	daily	1 km
CIR	CIR	classified irrigation	-	-
SP	T_REF_BULK	unit: %kg dm ⁻³ .	-	1 km
	T_SAND	unit: % wt.	-	1 km
	T_CLAY	unit: % wt.	-	1 km
	T_PH_H2O	unit: %-log (H ⁺).	-	1 km
	T_GRAVEL	unit: % vol.	-	1 km
	T_SILT	unit: % wt.	-	1 km
RSD	pet	potential evapotranspiration	monthly	4 km
	fc	field capacity	-	250 m
GI	R4	river network vector I	-	-
	R5	river network vector II	-	-
	R12	river network vector III	-	-
	lat	Latitude	-	-
	lon	Longitude	-	-
	im	moisture index	-	-

Note: REF_BULK: soil bulk density; PH_H2O: hydrogen ion concentration; GRAVEL: volume percentage of crushed stone; T_v:topsoil layer. The dashed line represents no default values.

删除的内容: the

删除的内容: dash

删除的内容:


Table 2 Evaluation index of relative soil moisture (SMI) in different growth periods of crops at 0–10, and 10–20 cm depths.

SMI in different growth periods of wheat (%)						
seeding	seedling	tillering	greening	jointing	booting	grouting mature
70~90	75~95			80~95		55~60

SMI in different growth periods of maize (%)						
seeding	seedling	jointing	booting	tasseling	grouting	mature
75~85	65~75	70~80	75~85		65~75	


删除的内容: ,

删除的内容: depth



Symbols	Definitions
Crop	Wheat and Maize
SM	Soil moisture
C_i	The i th crop type
P_j	The j th phenology
D_k	The k th soil depth
SMI_{ijk}	The soil moisture evaluation index corresponding to the k th soil depth and the j th phenology of i th crop
MD	Meteorological data
DOY	Day of year
CIR	Classified irrigation
SP	Soil properties
GI	Geographical information
RSSSM	The global remote-sensing-based surface soil moisture dataset (2003–2018)
ESA CCI SM	The European Space Agency Climate Change Initiative SM v05.2 (1978–2020)
ChinaCropSM 1km	1-km daily Soil Moisture dataset for dryland wheat and maize across China
maize ₀₋₁₀	ChinaCropSM1km for maize harvesting areas at 0–10 cm soil depth
maize ₁₀₋₂₀	ChinaCropSM1km for maize harvesting areas at 10–20 cm soil depth
wheat ₀₋₁₀	ChinaCropSM1km for wheat harvesting areas at 0–10 cm soil depth
wheat ₁₀₋₂₀	ChinaCropSM1km for wheat harvesting areas at 10–20 cm soil depth

Figure 2 Flow chart for producing ChinaCropSM1 km with an irrigation module.



删除的内容:
 删除的内容: 1km
 删除的内容: irrigating




Figure 3 Comparison between the predicted soil moisture (ChinaCropSM 1 km) and in situ samples by crops and depths (cm). (a) wheat₁₀₋₁₀, (b) wheat₁₀₋₂₀, (c) maize₁₀₋₁₀ and (d) maize₁₀₋₂₀. The red lines are the trend lines, the color bar indicates the point density, and the black lines represent the 1:1 lines.

删除的内容: 1km

删除的内容: colorbar means

删除的内容: for




Figure 4. Comparison of soil moisture accuracy with and without an irrigation module.

删除的内容: between

删除的内容: irrigation






Figure 5 The importance scores of 11 independent variables and the irrigation factor (CIR).



005 Figure 6 Comparison of the spatial accuracy (r) between ChinaCropSM 1 km and in situ SM observations in each month by crops and depths. (a) wheat0-10, (b) wheat10-20, (c) maize0-10 and (d) maize10-20. The dashed lines represent the mean values.

删除的内容: 1km

删除的内容: dash



010 Figure 7 Comparison of the temporal accuracy (r , RMSE, bias, ubRMSE) between ChinaCropSM 1 km and in situ soil moisture observations by crops and depths. (a1, a2) wheat₀₋₁₀, (b1, b2) wheat₁₀₋₂₀, (c1, c2) maize₀₋₁₀ and (d1, d2) maize₁₀₋₂₀. The dashed lines represent the mean values.

删除的内容: 1km
删除的内容: dash

Table 3 Summary of means of evaluation indices (r , bias, RMSE, and ubRMSE) of three products (ChinaCropSM1 km, RSSSM and ESA CCI SM). All products were compared with in situ surface observations (0–10 cm).

Product	ChinaCrop	RSSSM	ESA CCI SM	ChinaCrop	RSSSM	ESA CCI SM
	SM1 km _{maize}			SM1 km _{wheat}		
r	0.93	0.43	0.35	0.93	0.29	0.33
RMSE	0.033	0.167	0.126	0.035	0.187	0.121
bias	0.0006	-0.1361	-0.0846	-0.0008	-0.1552	-0.0705
ubRMSE	0.033	0.097	0.093	0.035	0.105	0.099

删除的内容: on


删除的内容: indexes

删除的内容: 1km

删除的内容: , all


删除的内容: 1km

删除的内容: 1km



1025

Figure 8 Time series of comparison between in situ SM observations and products.



030 Figure 9 Boxplot of the temporal (a, c) and spatial (b, c) accuracies for ChinaCropSM1 km, RSSSM and ESA CCI SM by r , bias, RMSE, and ubRMSE. These evaluation indices were calculated by comparing the three products with in situ SM observations; the comparison period for ChinaCropSM1 km and RSSSM was from 2003 to 2018; and for ChinaCropSM1 km and ESA CCI SM was 1995–2018.

删除的内容: accuracy

删除的内容: 1km

删除的内容: indexes

删除的内容: 1km

删除的内容: 1km