

Mass Balance of the Greenland and Antarctic Ice Sheets from 1992 to 2020

Inès N. Otosaka¹, Andrew Shepherd^{1,2}, Erik R. Ivins³, Nicole-Jeanne Schlegel³, Charles Amory⁴, Michiel R. van den Broeke⁵, Martin Horwath⁶, Ian Joughin⁷, Michalea D. King⁷, Gerhard Krinner⁴, Sophie Nowicki⁸, Antony J. Payne⁹, Eric Rignot¹⁰, Ted Scambos¹¹, Karen M. Simon¹², Benjamin E. Smith⁷, Louise S. Sørensen¹³, Isabella Velicogna^{3,10}, Pippa L. Whitehouse¹⁴, Geruo A¹⁰, Cécile Agosta¹⁵, Andreas P. Ahlstrøm¹⁶, Alejandro Blazquez¹⁷, William Colgan¹⁶, Marcus E. Engdhal¹⁸, Xavier Fettweis¹⁹, Rene Forsberg¹³, Hubert Gallée⁴, Alex Gardner³, Lin Gilbert²⁰, Noel Gourmelen²¹, Andreas Groh⁶, Brian C. Gunter²², Christopher Harig²³, Veit Helm²⁴, Shfaqat Abbas Khan¹³, Christoph Kittel⁴, Hannes Konrad²⁵, Peter L. Langen²⁶, Benoit S. Lecavalier²⁷, Chia-Chun Liang¹⁰, Bryant D. Loomis²⁸, Malcolm McMillan²⁹, Daniele Melini³⁰, Sebastian H. Mernild³¹, Ruth Mottram³², Jeremie Mouginot⁴, Johan Nilsson³, Brice Noël⁵, Mark E. Pattle³³, William R. Peltier³⁴, Nadege Pie³⁵, Mònica Roca³⁶, Ingo Sasgen²⁴, Himanshu V. Save³⁵, Ki-Weon Seo³⁷, Bernd Scheuchl¹⁰, Ernst J.O. Schrama³⁸, Ludwig Schröder⁶, Sebastian B. Simonsen¹³, Thomas Slater¹, Giorgio Spada³⁹, Tyler C. Sutterley⁴⁰, Bramha Dutt Vishwakarma⁴¹, Melchior van Wessem⁵, David Wiese³, Wouter van der Wal¹², Bert Wouters^{12,5}

¹Centre for Polar Observation and Modelling, University of Leeds, Leeds, United Kingdom

²Department of Geography and Environmental Sciences, Northumbria University, Newcastle-upon-Tyne, United Kingdom

³Jet Propulsion Laboratory, California Institute of Technology, Pasadena, United States

⁴Institute of Environmental Geosciences, Université Grenoble Alpes, Grenoble, France

⁵Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

⁶Institut für Planetare Geodäsie, Technische Universität Dresden, Dresden, Germany

⁷Polar Science Center, University of Washington, Seattle, United States

⁸Department of Geology, University at Buffalo, Buffalo, United States

⁹School of Geographical Sciences, University of Bristol, Bristol, United Kingdom

¹⁰Earth System Science, University of California Irvine, Irvine, United States

¹¹Earth Science and Observation Center, CIRES, University of Colorado Boulder, Boulder, United States

¹²Faculty of Civil Engineering and Geoscience, Delft University of Technology, Delft, The Netherlands

¹³National Space Institute, Technical University of Denmark, Lyngby, Denmark

¹⁴Department of Geography, Durham University, Durham, United Kingdom

¹⁵Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL, CEA-CNRS-UVSQ, Gif-sur-Yvette, France

¹⁶Glaciology and Climate, Geological Survey of Denmark and Greenland, Copenhagen, Denmark

¹⁷Spatial Geophysics and Oceanography Studies Laboratory, Toulouse, France

¹⁸ESA-ESRIN, Frascati, Italy

¹⁹Geography, University of Liège, Liège, Belgium

²⁰Mullard Space Science Laboratory, University College London, West Sussex, United Kingdom

²¹University of Edinburgh, Edinburgh, United Kingdom

²²Aerospace Engineering, Georgia Institute of Technology, Atlanta, United States

²³Department of Geosciences, University of Arizona, Tucson, United States

²⁴Glaciology, Alfred-Wegener-Institute Helmholtz-Center for Polar and Marine Research, Bremerhaven, Germany

²⁵Satellite-based Climate Monitoring, Deutscher Wetterdienst, Offenbach/Main, Germany

²⁶Department of Environmental Science, iClimate, Aarhus University, Roskilde, Denmark

²⁷Department of Physics and Physical Oceanography, Memorial University, St. John's, Canada

- ²⁸Geodesy and Geophysics Laboratory, NASA GSFC, Greenbelt, United States
²⁹Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom
³⁰Istituto Nazionale di Geofisica e Vulcanologia, Rome, Italy
³¹SDU Climate Cluster, University of Southern Denmark, Odense, Denmark
³²Research and Development Department, Danish Meteorological Institute, Copenhagen, Denmark
³³isardSAT, Guildford, United Kingdom
³⁴Physics, University of Toronto, Toronto, Canada
³⁵Center for Space Research, University of Texas at Austin, Austin, United States
³⁶isardSAT, Barcelona, Spain
³⁷Department of Earth Science Education, Seoul National University, Seoul, Korea
³⁸Department SpE, Faculty of Aerospace Engineering, TU Delft, Delft, The Netherlands
³⁹Dipartimento di Fisica e Astronomia, Alma Mater Studiorum Università di Bologna, Bologna, Italy
⁴⁰Applied Physics Laboratory, University of Washington, Seattle, United States
⁴¹Interdisciplinary Centre for Water Research, Indian Institute of Science, Bengaluru, India

Correspondence to: Inès N. Otosaka (i.n.otosaka@leeds.ac.uk)

5 **Abstract.** Ice losses from the Greenland and Antarctic Ice Sheets have accelerated since the 1990s, accounting
for a significant increase in global mean sea level. Here, we present a new 29-year record of ice sheet mass balance
from 1992 to 2020 from the Ice Sheet Mass Balance Inter-comparison Exercise (IMBIE). We compare and
combine 50 independent estimates of ice sheet mass balance derived from satellite observations of temporal
changes in ice sheet flow, in ice sheet volume and in Earth's gravity field. Between 1992 and 2020, the ice sheets
10 contributed 21.0 ± 1.9 mm to global mean sea-level, with the rate of mass loss rising from 105 Gt yr^{-1} between
1992 and 1996 to 372 Gt yr^{-1} between 2016 and 2020. In Greenland, the rate of mass loss is $169 \pm 9 \text{ Gt yr}^{-1}$
between 1992 and 2020 but there are large inter-annual variations in mass balance with mass loss ranging from
 86 Gt yr^{-1} in 2017 to 444 Gt yr^{-1} in 2019 due to large variability in surface mass balance. In Antarctica, ice losses
continue to be dominated by mass loss from West Antarctica ($82 \pm 9 \text{ Gt yr}^{-1}$) and to a lesser extent from the
15 Antarctic Peninsula ($13 \pm 5 \text{ Gt yr}^{-1}$). East Antarctica remains close to a state of balance with a small gain of $3 \pm$
 15 Gt yr^{-1} , but is the most uncertain component of Antarctica's mass balance. The dataset is publicly available at
<https://doi.org/10.5285/77B64C55-7166-4A06-9DEF-2E400398E452> (The IMBIE Team, 2021).

1 Introduction

20 The Antarctic and Greenland Ice Sheets store the vast majority (99%) of Earth's freshwater ice on land. The rate
of change in ice sheet mass - or ice sheet mass balance - is the net difference between mass loss through solid ice
discharge at the grounding line, melting at the bed and at the ice-ocean interface and the surface mass balance
(SMB; precipitation minus meltwater runoff, sublimation, evaporation, and erosion). Over the past three decades
(between the 1990s and 2010s), ice losses from Antarctica and Greenland increased six-fold (The IMBIE Team,
25 2018, 2020), raising the global sea level (WCRP Global Sea Level Budget Group, 2018) and with it the risk of
coastal flooding worldwide (Kulp and Strauss, 2019; Vitousek et al., 2017; Hanson et al., 2011). In Antarctica,
the losses have arisen primarily due to ocean-driven melting of ice shelves (Adusumilli et al., 2020; Paolo et al.,
2015) and their collapse (Cook and Vaughan, 2010), which have accelerated the ice flow (Hogg et al., 2017;
Selley et al., 2021; Rignot et al., 2004), retreat (Konrad et al., 2018; Milillo et al., 2022; Jenkins et al., 2018) and
30 drawdown (Konrad et al., 2017; Shepherd et al., 2019) of numerous marine-terminating ice streams. In Greenland,
increasing air temperatures (Hanna et al., 2021) and decreasing cloud cover (Hofer et al., 2017) have exacerbated
summertime surface melting (Leeson et al., 2015; Tedesco and Fettweis, 2020) and runoff (Trusel et al., 2018;
Slater et al., 2021), in tandem with the speedup (Rignot and Kanagaratnam, 2006) and retreat (King et al., 2020)
of outlet glaciers responding to a warming ocean (Straneo and Heimbach, 2013). While ice sheet response to
35 climate forcing remains the least constrained component of the twenty-first-century sea level budget (Pattyn and
Morlighem, 2020; Fox-Kemper et al., 2021), maintaining the long-term observational record of ice sheet mass
balance is critical to improving ice sheet model skill (Edwards et al., 2021; Ritz et al., 2015) and confidence in
projections of sea level rise (Aschwanden et al., 2021; Slater et al., 2020; Shepherd and Nowicki, 2017).

Thanks to the launch of new satellite missions and the development of improved geophysical corrections and
40 models of SMB and glacial isostatic adjustment (GIA), it is now possible to routinely monitor ice sheet mass
changes using observations of ice-flow derived from satellite radar and optical imagery (e.g. Gardner et al., 2018;
Moon et al., 2012; Mouginot et al., 2017), surface elevation changes (derived from satellite altimetry) (e.g.

Sandberg Sørensen et al., 2018; Smith et al., 2020), and fluctuations in Earth's gravity field (derived from satellite gravimetry from GRACE and its follow on) (e.g. Tapley et al., 2019; Velicogna et al., 2020; Sasgen et al., 2020).
45 The Ice Sheet Mass Balance Inter-comparison Exercise (IMBIE) has shown that there is good agreement between these satellite methods (Shepherd et al., 2012) and that combining independent satellite-based ice sheet mass balance estimates reduces uncertainty in estimates of Greenland and Antarctica's contribution to sea level rise. By adopting a common framework to support the comparison and aggregation of ice sheet mass balance estimates generated by different participants, it is possible to assess differences between techniques and the impact of using
50 different geophysical corrections, SMB models, or GIA models in ice sheet mass balance estimation to produce a reconciled time-series of ice sheet mass changes. SMB models are required for estimating the net mass balance in the input-output method while GIA models are necessary to correct ice sheet mass balance estimates derived from satellite gravimetry and to a lesser extent those derived from satellite altimetry. The GIA is the result of solid Earth mass redistribution caused by changes in ice mass since the last glaciation. Gravimetry fields record the combined effect of mass redistribution due to the GIA and recent changes in ice sheet mass balance. The GIA
55 contribution therefore needs to be modelled separately and removed from the gravimetry fields, especially since it is of the same order of magnitude as the ice sheet mass balance signal (Caron and Ivins, 2020; Sutterley et al., 2014a). Altimetry elevation change estimates also need to be corrected for the GIA. However, contrary to gravimetry estimates, altimetry estimates are less sensitive to GIA as it manifests as an uplift (or subsidence) rate
60 of the order of a few millimetres per year, much smaller than the elevation changes recorded. The most recent IMBIE assessments for the Antarctic Ice Sheet and the Greenland Ice Sheet covered the periods 1992 to 2017 and 1992 to 2018, respectively, and reported a combined contribution of 17.8 ± 1.8 mm to global mean sea level (GMSL) between 1992 and 2017 (The IMBIE Team, 2018, 2020). Here, we extend these records to cover the same extended period (1st January 1992 to 31st December 2020) for both ice sheets. In the rest of the paper, all of
65 time periods cited refer to the period extending from 1st January of the first year quoted to 31st December of the second year quoted.

In the years since our most recent assessment there have been notable changes in ice sheet mass in both hemispheres, and in the availability of satellite observations and ancillary datasets with which to detect these changes. In Greenland, for example, atmospheric blocking and reduced summertime snowfall (Tedesco and
70 Fettweis, 2020) led to near-record levels of meltwater runoff in 2019 (Slater et al., 2021) which, in combination with progressively increasing ice discharge (Mouginot et al., 2019), set a new record for annual ice losses during the satellite era (Sasgen et al., 2020). In Antarctica, pervasive mass losses have continued in the Amundsen Sea Sector (Groh and Horwath, 2021) as a consequence of further grounding line retreat (Milillo et al., 2022) and the associated glacier speedup (Joughin et al., 2021). A follow on to the GRACE satellite mission (GRACE-FO) was
75 launched in May 2018 (Tapley et al., 2019), the ICESat-2 satellite laser altimeter mission was launched in September 2018 (Smith et al., 2020), and updated products have been released for many others - including swath altimetry from CryoSat-2 (Gourmelen et al., 2018). To accompany these observations, there have been updated models of GIA (e.g. Caron and Ivins, 2020) to correct mass and elevation changes associated with solid earth movement, of firn densification (e.g. Stevens et al., 2020) to correct changes in elevation for surface processes,
80 and of SMB (e.g. Fettweis et al., 2020; Mottram et al., 2021) to aid mass budget and mass balance partitioning calculations.

Here, we make use of new satellite observations, new methods and models to provide an updated IMBIE assessment of Greenland and Antarctic ice sheet mass balance, extending our most recent records by 3 and 4 years, respectively. We provide a description of the datasets incorporated in this updated assessment and of the aggregation methods employed. We also discuss differences between the ice sheet mass balance estimates derived from altimetry, gravimetry and the input-output method, and we present extended reconciled time-series of ice sheet mass change. We discuss the limitations of our dataset and outline a roadmap for future improvements.

2 Data

2.1 Data Background

Fluctuations in ice sheet mass are a key indicator of ice sheet stability and can be inferred using a range of satellite techniques (Shepherd et al., 2012). Satellite altimetry measures ice sheet elevation change, computed at orbit crossing points by calculating the difference in ice sheet elevation at a crossover point between ascending and descending satellite passes (e.g. Wingham et al., 1998), using clusters of data points acquired along all ground tracks (e.g. Pritchard et al., 2009), or by differencing height models separated over time (e.g. Csatho et al., 2014). Mass balance is estimated by accounting for changes in bedrock elevation (e.g. Caron and Ivins, 2020) and then by either prescribing the density associated to the elevation fluctuation (e.g. Shepherd et al., 2019) or by making a model-based correction for changes in firn compaction (Sørensen et al., 2011). The technique is unique in charting patterns of mass imbalance with fine (monthly) temporal sampling and fine (10^2 km^2) spatial resolution, and there are continental-scale measurements dating back to the early 1990s. Satellite measurements of ice velocity computed from sequential radar and optical imagery (e.g. Rignot and Kanagaratnam, 2006) are the basis of ice sheet input-output assessments (e.g. Rignot et al., 2019; Mouginot et al. 2019). Ice velocities are combined with estimates of ice thickness (e.g. Morlighem et al., 2017) to compute changes in marine-terminating glacier discharge, and then with regional climate model estimates of surface mass balance sources (snowfall, rainfall) and sinks (runoff, sublimation, evaporation, and erosion) (e.g. Fettweis et al., 2020; Mottram et al., 2021) to measure temporal changes in net mass balance. The technique provides monthly to annual temporal sampling and drainage basin scale spatial resolution, and there are continental-scale measurements dating back to the late 1970s. During the last decade, new satellite missions with a more frequent revisit time (down to 6 days using image pairs from Sentinel-1a and Sentinel-1b available during the period 2016 to 2021 until the end of Sentinel-1b mission) have been used to improve the temporal resolution of ice velocity measurements, allowing to investigate seasonal fluctuations in ice velocity (King et al., 2018; Lemos et al., 2018) and produce monthly estimates of ice discharge at the continental scale. Mankoff et al. (2021) even produced daily estimates of ice sheet mass balance from the input-output method by resampling the velocity data, however the original temporal resolution of ice velocity measurements does not exceed 12 days. Satellite gravimetry measures fluctuations in Earth's gravitational field, computed using either global spherical harmonic solutions (e.g. Velicogna and Wahr, 2006) or using spatially discrete mass concentration units (e.g. Luthcke et al., 2006). Ice sheet mass changes are determined after making model-based corrections for GIA (e.g. Caron and Ivins, 2020) and for the leakage of mass trends occurring elsewhere in the climate system, especially those arising from ocean mass variability and changes in land hydrology. The technique provides fine (monthly) temporal sampling and moderate (10^5 km^2) spatial resolution,

120 dating back to 2002 with the launch of the GRACE mission and the more recent launch of its follow on GRACE-
FO in 2018.

2.2 Input Data

To compile our assessment of Greenland Ice Sheet mass balance we use 27 satellite-based estimates of ice sheet mass change, including 8 estimates based on satellite altimetry, 16 based on satellite gravimetry, and 3 based on the input-output method. Compared to the most recent IMBIE assessment, 12 of these estimates have been updated to include more recent data for Greenland. This set of updated estimates is made of 2 estimates from the input-output method, 1 altimetry estimate, and 9 gravimetry estimates including data from the new GRACE Follow-On space gravimetry mission (GRACE-FO). For our assessment of Antarctica’s mass balance, we use 23 satellite-based estimates altogether, with 6 derived from altimetry, 16 from gravimetry, and 1 from the input-output method. More than half of these estimates have been extended in time compared to IMBIE-2. These updated estimates for Antarctica include the input-output method estimate, 2 altimetry estimates, and 10 gravimetry estimates combining GRACE and GRACE-FO data. In total, this new IMBIE assessment includes data from 14 satellite missions, spanning the years 1992 to 2020 – with results from all three geodetic techniques available between 2003 and 2018 in Greenland and 2002 and 2018 in Antarctica – and, for the first time, includes data from the GRACE-FO mission launched in 2018. A wide range of GIA models have been used to correct gravimetric and volumetric mass balance estimates. The models use in this assessment are all forward models, which combine a rheology model of the solid Earth with a model of past ice mass change. In this assessment, only two SMB models have been used in the input-output method estimates included – the RACMO (Regional Atmospheric Climate Model) and MAR (Modèle Atmosphérique Régional) models (Table 1).

140 **Table 1. Synthesis of satellite datasets, GIA, and SMB models used to derive the individual estimates of ice sheet mass balance included in this study. Details and references of the GIA and SMB models are available in Table A1.**

		1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020			
Satellite Missions																																	
IOM	ERS-1																																
	ERS-2																																
	RADARSAT-1																																
	ENVISAT																																
	ALOS/PALSAR																																
	RADARSAT-2																																
	TerraSAR-X																																
	COSMO-SkyMed																																
	Landsat-8																																
	Sentinel-1																																
ALT	ERS-1																																
	ERS-2																																
	ENVISAT																																
	ICESat																																
	CryoSat-2																																
GMB	GRACE																																
	GRACE-FO																																

GIA models		
AIS	A13	
	A13 and W12a	
	ICE-5G and W12a	
	ICE-6G	
	ICE-6G and A13	
	ICE-6G and IJ05_R2	
	IJ05 and W12a	
	IJ05_R2	
	IJ05_R2 and A13	
	IJ05_R2 and Paulson07	
	IJ05_R2 and Simpson09	
	IJ05_R2 and W12a	
	Khan_2016 and W12a	
	Schrama14	
	W12a	
GrIS	A13	
	ICE-5G	
	ICE-6G	
	ICE-5G and ICE-6G	
	ICE-6G and A13	
	Paulson07	
	Schrama14	
	Simpson09	
SMB models		
AIS	RACMO2.3	
GrIS	MAR 3.5	
	MAR 3.5.2	
	RACMO 2.3	

To achieve a meaningful comparison of ice sheet mass balance estimates, we analyse mass trends using common definitions of the Antarctic, West Antarctic, East Antarctic, Antarctic Peninsula, and Greenland Ice Sheet boundaries (AIS, WAIS, EAIS, APIS, and GrIS, respectively). We use two ice sheet drainage basin sets, both previously used in the past IMBIE assessments (Shepherd et al., 2012; IMBIE Team, 2018; 2020) and available at <http://imbie.org/imbie-3/drainage-basins/>. The first drainage basin set was derived based on ICESat surface elevation data and includes 27 basins in Antarctica covering an area of 11,885,725 km² and 19 in Greenland over an area of 1,703,625 km² (Zwally et al., 2012) and is retained for consistency with the first IMBIE assessment (Shepherd et al., 2012). The second set defines 18 basins in Antarctica covering 11,892,700 km² and 6 in Greenland covering 1,723,300 km² (Rignot et al., 2011a; Rignot et al., 2011b). The two ice sheet delineation differ by 1.1 % and 0.1 % of total ice sheet extent for the Greenland and Antarctic Ice Sheets, respectively, and thus using either of these definitions leads to a negligible difference in mass balance (The IMBIE Team, 2018; 2020). IMBIE participants were free to use either of these two definitions, and we combine mass trends over the GrIS,

155 AIS, WAIS, EAIS, and APIS together regardless of what definition was chosen. The different estimates included in this assessment are presented on Figure 1.




Figure 1. Individual rates of ice sheet mass balance from the input-output, altimetry, and gravimetry groups over the GrIS, APIS, EAIS, and WAIS included in this study and standardised following the procedure described in Section 3 (i). The grey shading shows the estimated uncertainty of the aggregated time-series per group calculated following the procedure described in Section 3 (ii).

2.3 Output Data

The output data consists of a single reconciled estimate of ice sheet mass balance covering the period 1st January 1992 to 31st December 2020 for the GrIS, AIS, APIS, WAIS, EAIS, and the sum of the GrIS and AIS. Two CSV

160 files are provided for each ice sheet region, one with the data provided in Gigatons (Gt) and one with the data provided in equivalent sea level contribution in millimetres (mm). These files contain annual rates of mass balance and cumulative mass changes with their corresponding uncertainties.

3 Methods

165 IMBIE participants contributed time-series of either relative mass change, $\Delta M(t)$, or of rate of mass change, $dM(t)/dt$, with their associated uncertainty, integrated over at least one of the ice sheet regions defined in the standard drainage basin sets. To produce a reconciled estimate of ice sheet mass change from these individual estimates, we compare and aggregate $dM(t)/dt$ from each satellite technique. The IMBIE assessment software used to produce the dataset presented in this study is available at <https://doi.org/10.5281/zenodo.7342481>. We apply a consistent processing scheme to all submitted datasets and for all ice sheet regions which consists of: i) 170 computing $dM(t)/dt$ for all datasets that were submitted as $\Delta M(t)$, ii) aggregating time-series of mass trends within each class of satellite observations, iii) combining the altimetry, gravimetry, and input-output time-series to derive a single reconciled time-series of mass trends, and iv) integrating this reconciled time-series of mass trends to produce the final reconciled time-series of cumulative mass change. In what follows, we summarise each of these processing steps:

175

i) Computing time-series of mass trends

180 First, we derive time-series of monthly rates of ice sheet mass change, $dM(t)/dt$, for all datasets that were submitted as $\Delta M(t)$ to allow the aggregation of datasets within each satellite observations class as $dM(t)/dt$ computed using a standardised approach. At each epoch, we estimate $dM(t)/dt$ by fitting a linear trend to the $\Delta M(t)$ data falling within a sliding window of 36 months, centred around the given epoch, using a weighted least-squares approach, with each point weighted by its error. The error on the derived time-series is taken as the sum in quadrature of the linear model structural error computed as the standard error of the linear regression s_e and the mean of the errors of the n_w points in the original $\Delta M(t)$ time-series falling within the 36-month sliding window as:

$$\sigma_{\frac{dM}{dt}}(t) = \sqrt{s_e^2 + \left(\frac{1}{n_w} \sum_{i=0}^{n_w-1} \sigma_{\Delta M,i} \right)^2} \quad (1)$$

185 Finally, the derived time-series of mass trends are truncated by half the window width at the start and end of their period.

ii) Aggregating time-series of mass trends from similar satellite observations

We aggregate the standardised time-series of mass trends within the altimetry, gravimetry, and input-output 190 groups separately to produce three time-series over each ice sheet region $\left. \frac{dM_{aggr}(t)}{dt} \right|_{group}$, where *group* refers to one of these three independent satellite techniques (i.e. altimetry, gravimetry, or input-output method). We calculate each aggregated time-series by taking the error-weighted average of the $n_{estimates\ per\ group}$ individual monthly rates of ice sheet mass change available from the same satellite technique group at each month:

$$\left. \frac{dM_{aggr}(t)}{dt} \right|_{group} = \frac{\sum_{i=0}^{n_{estimates\ per\ group}-1} \left. \frac{dM(t)}{dt} \right|_{group,i} / \sigma_{\left. \frac{dM(t)}{dt} \right|_{group,i}}}{\sum_{i=0}^{n_{estimates\ per\ group}-1} 1 / \sigma_{\left. \frac{dM(t)}{dt} \right|_{group,i}}} \quad (2)$$

195 The associated error is calculated as the sum in quadrature of the contributing individual time-series errors divided by the square root of the number of estimates in the group:

$$\sigma_{aggr,group}(t) = \sqrt{\frac{1}{n_{estimates\ per\ group}} \sum_{i=0}^{n_{estimates\ per\ group}-1} \sigma_{\left. \frac{dM(t)}{dt} \right|_{group,i}}^2} \quad (3)$$

iii) *Combining the altimetry, gravimetry, and input-output time-series of mass trends*

200 We combine the altimetry, gravimetry, and input-output time-series to produce a single reconciled time-series of mass trends by taking the error-weighted mean of the n_{group} independent estimates for which a mass trend estimate is available at each epoch (comprised between 1 and 3):

$$\frac{dM_{reconciled}(t)}{dt} = \frac{\sum_{i=0}^{n_{group}-1} \frac{dM_{aggr,i}(t)}{dt} / \sigma_{aggr,i}(t)}{\sum_{i=0}^{n_{group}-1} 1 / \sigma_{aggr,i}(t)} \quad (4)$$

We estimate the error on the reconciled mass trend time-series at each epoch as the sum in quadrature of the aggregated time-series errors divided by the square root of the number of independent estimates available:

$$\sigma_{reconciled}(t) = \sqrt{\frac{1}{n_{group}} \sum_{i=0}^{n_{group}-1} \sigma_{aggr,i}^2(t)} \quad (5)$$

Finally, when summing mass trends of multiple ice sheets, the combined uncertainty is estimated as the root sum square of the uncertainties for each region:

$$\sigma_{total}(t) = \sqrt{\sum_{regions} \sigma_{reconciled,i}^2(t)} \quad (6)$$

210

iv) *Generating the final reconciled time-series of cumulative mass change*

We generate a time-series of cumulative ice sheet mass change by integrating our reconciled time-series of mass trends over time for each region. We estimate the cumulative errors as the root sum square of errors, divided by 12 as our estimates are posted at monthly epochs:

$$215 \quad \sigma_{cumul}(t) = \sqrt{\frac{1}{12} \sum_{i=0}^{t-1} \sigma_{reconciled}^2(i)} \quad (7)$$

Here, we discuss the potential systematic bias introduced by the inclusion of the peripheral glaciers and ice caps (GICs) in the gravimetry estimates included in our assessment as the spatial resolution of satellite gravimetry is

220 not sufficient to resolve separately the mass change signals of these neighbouring ice masses. To examine this
further, we use Hugonnet et al. (2021) dataset (<https://doi.org/10.6096/13>, last access: 23 February 2023), which
provides mass balance estimates of the glaciers located at the periphery of the ice sheets derived from high
resolution digital elevation models. During the overlap of Hugonnet et al. study and the gravimetry recorded
employed in this study (2002-2019), Greenland peripheral glaciers lost mass at a rate of $35.5 \pm 1.6 \text{ Gt yr}^{-1}$. In
225 Antarctica (excluding the Sub Antarctic glaciers located further than 1000 km from the ice sheet), peripheral
glaciers lost mass at a rate of $11.8 \pm 3.4 \text{ Gt yr}^{-1}$, $0.7 \pm 1.1 \text{ Gt yr}^{-1}$, and $5.7 \pm 2.5 \text{ Gt yr}^{-1}$ at the APIS, EAIS, and
WAIS, respectively. To test the impact of the inclusion of the peripheral glaciers in our gravimetry estimates on
our reconciled ice sheet mass balance assessment, we use the peripheral glaciers mass trends time-series from
Hugonnet et al. to remove the contribution of the GICs on our aggregated gravimetry time-series. We use
230 consecutive 5-year rates of mass change for this analysis and their corresponding uncertainties. For 2020, which
is not covered by Hugonnet et al., we use the rate of mass change estimated over the 5-year period 2015-2019
instead. We combine in quadrature the uncertainty on the peripheral GICs mass balance and the uncertainty of
our aggregated gravimetry mass balance calculated from Eq. 3. Next, we follow the procedure described in step
(iii) to re-combine this modified gravimetry aggregated time-series with the altimetry and input-output aggregated
235 time-series. We compare this modified reconciled estimate to our original estimate and find that removing the
contribution of the GICs from the gravimetry time-series results in a reduction in mass loss of 4.1 % and 3.3 % in
Greenland and Antarctica, respectively, smaller than the uncertainty bounds of our reconciled estimate (Table
A2). This simple analysis shows that the inclusion of the peripheral ice masses in the gravimetry estimates
included in this study has a negligible impact on our reconciled mass balance assessment of the WAIS and EAIS,
240 and only a small impact (less than 10 Gt yr^{-1}) on our assessment of the GrIS and APIS.

4 Results

First, we compare individual estimates of ice sheet mass balance within each of the three geodetic technique
experiment groups, separately, to assess the level of agreement among estimates derived using the same technique.
Within each group, we compare annual rates of mass change and their standard deviation for each ice sheet region.
245 The input-output group includes significantly fewer mass balance estimates than the other technique experiment
groups, but these estimates have the advantage of providing information on the partitioning of mass trends between
signals related to SMB and ice dynamics, and they also cover relatively long periods of time (Figure 2). Ice
discharge is measured from satellite observations of ice velocities combined with estimates of ice thickness at
glaciers' termini, and SMB is derived from regional climate model outputs. To estimate the SMB anomaly in
250 Greenland, two estimates used MAR (version 3.2 and version 3.5.2) and one used RACMO (version 2.3). In
Antarctica, the input-output estimate used RACMO (version 2.3). In addition to using different SMB models,
those estimates also define different reference periods to calculate the SMB anomalies. All of the mass balance
estimates derived in this group were originally posted at annual resolution and we resample them over monthly
epochs to aggregate them with estimates from the other groups. We include 3 input-output method estimates of
255 GrIS mass balance, all at annual resolution and that together span the period 1992 to 2020 and overlap during the
period 2007 to 2011. During their common period, annual rates of mass change determined from these three input-
output datasets have a median difference of 28.5 Gt yr^{-1} with a standard deviation of 35 Gt yr^{-1} . For Antarctica

and its ice sheet components, we include one input-output mass balance estimate which covers the entire 1992 to 2020 period at annual resolution.

260 The altimetry group includes 8 mass balance estimates for the GrIS that together span the years 2003 to 2018, with 4 of these solutions derived from radar altimetry, 2 from laser altimetry, and 2 from a combination of both. We include 6 altimetry mass balance estimates for the AIS which together cover the period 1992 to 2019. In total we include 6 solutions for the EAIS, 6 for the WAIS, and 5 for the APIS. Of these, 2 solutions are derived from radar altimetry, 1 from laser altimetry, and 3 from a combination of both. To derive rates of surface elevation
265 change, various methods were applied to the laser and radar altimetry data including repeat-track, plane fit, or overlapping footprints techniques. For Greenland, half of the participants corrected the altimetry time-series for the GIA effect while for Antarctica, all participants applied a GIA correction. Next, to derive mass trends from rates of surface elevation change, either a constant density or a spatially and time varying density field from a firm density model forced by a regional climate model, were applied. These solutions have varying temporal
270 resolutions ranging from 1 month to 7.1 yr for an average effective temporal resolution of 3.0 yr for Greenland and 2.6 yr for Antarctica. The temporal resolution of the altimetry group is thus lower than annual, mainly due to the fact that solutions derived from laser altimetry data were all provided as constant rates spanning the duration of ICESat-1 mission while the radar altimetry solutions have a higher temporal resolution of 0.35 yr for Greenland and 0.47 yr for Antarctica. As there is no overlap period during which all altimetry estimates are available, we
275 compare solutions derived solely from radar altimetry and solutions incorporating laser altimetry data separately. In Greenland, radar altimetry solutions have a median difference of 144 Gt yr⁻¹ and standard deviation of 67 Gt yr⁻¹ during their two-year overlap period (2013 to 2014) while the median difference between laser and combination solutions is 29 Gt yr⁻¹ with a standard deviation of 29 Gt yr⁻¹ during their 6-year overlap (2004 to 2009). In Antarctica, the spread between laser solutions is largest at the EAIS with a standard deviation in annual
280 rates of 38 Gt yr⁻¹ between 2004 and 2008, followed by the WAIS and APIS with standard deviations of 23 Gt yr⁻¹ and 10 Gt yr⁻¹, respectively. On the other hand, radar altimetry solutions show a larger spread at the WAIS (21 Gt yr⁻¹) than at the EAIS (14 Gt yr⁻¹) during their overlap period (2013 to 2018).

The gravimetry group has the largest number of estimates, with 16 for each ice sheet that together span the period 2002 to 2020. All gravimetry solutions were submitted as time-series of cumulative mass change at monthly
285 resolution resulting in a collective effective resolution of 0.08 yr. All participants submitted estimates for all ice sheet regions, with 10 participants analysing spherical harmonic gravity field solutions using a wide range of approaches and 6 participants using mass concentration units (usually referred to as mascons) directly estimated from the GRACE and GRACE-FO level-1 K-band ranging data. Various GIA, hydrology leakage, and ocean leakage models were used to correct the gravimetry data for external signals. Overall, there is good agreement
290 between rates of ice sheet mass balance derived from satellite gravimetry. In Greenland, we compare the different gravimetry solutions over the period 2012 to 2014 and find that annual rates of mass have a median difference of 36 Gt yr⁻¹, and standard deviation is 30 Gt yr⁻¹. In Antarctica, the different gravimetry solutions overlap over a decade from 2004 to 2014 during which their annual rates of mass balance have a median difference of 41 Gt yr⁻¹. When comparing over the different regions of the Antarctic continent, the difference is greatest at the EAIS
295 with a median difference of 31 Gt yr⁻¹ and standard deviation of 26 Gt yr⁻¹. In the other regions, gravimetry estimates are in better agreement at the APIS with a median difference of 8 Gt yr⁻¹ and standard deviation of 10

Gt yr⁻¹, followed by the WAIS where the median difference between estimates reaches 19 Gt yr⁻¹ and their standard deviation is 17 Gt yr⁻¹.

300 Comparing mass balance estimates derived from similar satellite observations reveals that in Greenland, the median difference between estimates is the largest for the altimetry group and the smallest for the input-output group. In Antarctica, the median difference between altimetry estimates is less than 38 Gt yr⁻¹ and less than 41 Gt yr⁻¹ for gravimetry estimates during their respective overlap periods. However this comparison is limited by the varying temporal resolutions of the different datasets – especially for the altimetry group for which constant rates of mass change over long periods of time dampen temporal variation in ice sheet mass changes – and by the small
305 number of input-output estimates – in particular in Antarctica where only one estimate is available. This limits our ability to link differences between estimates derived from the same geodetic technique to methodological differences, or to the use of different geophysical corrections or auxiliary datasets.




Figure 2. Annual rates of mass change of the (a) GrIS, (b) AIS, (c) APIS, (d) EAIS, and (e) WAIS from the altimetry, gravimetry and input-output estimates included in this study (shown by the coloured bars) and the reconciled estimate produced from combining those estimates (shown by the thick black bars). The grey shading shows the uncertainty of our final reconciled estimate, calculated following the procedure described in Section 3 (iii). The number of individual mass balance estimates collated at each epoch is shown below each bar.

Next, we assess differences between the aggregated time-series derived within each class of satellite observations during the periods when estimates from all three geodetic techniques are available – from 2003 to 2018 for Greenland and from 2002 to 2019 for Antarctica (Figure A1). We compare rates of mass change during these overlap periods, which are 5 and 10 years longer than in the previous IMBIE assessments, respectively (Figure 3). We compare the standard deviation in aggregated rates of mass change altimetry, gravimetry and input-output estimates rates of mass change and to the uncertainty of our reconciled mass balance estimate (computed from Eq. 5) to assess whether differences between techniques are significant compared to the uncertainty of our reconciled assessment. In Greenland, rates of mass balance determined from altimetry, gravimetry, and the input-output method are in close agreement between 2003 and 2018, with a standard deviation of 19 Gt yr⁻¹ and a reconciled rate of mass loss of 221 ± 22 Gt yr⁻¹ from all three techniques. In Antarctica, the reconciled rate of mass loss between 2003 and 2019 is 115 ± 24 Gt yr⁻¹ but the spread of the altimetry, gravimetry and input-output estimates is 4 times larger than in Greenland (79 Gt yr⁻¹). Over the different regions of Antarctica, the spread of estimates of ice sheet mass balance increases with the size of the region considered, with standard deviations of 54 Gt yr⁻¹, 18 Gt yr⁻¹, and 16 Gt yr⁻¹, at the EAIS, WAIS, and APIS, respectively. Across all ice sheets, the input-output estimate is the most negative and the altimetry the most positive except at the EAIS, where the gravimetry estimate is the most positive. The greatest departure occurs at the EAIS where the three geodetic techniques disagree on even the sign of the mass change, with a maximum difference of 105 ± 33 Gt yr⁻¹ between rates of mass change from the input-output method and gravimetry estimates. This indicates that the EAIS remains a challenging region for which to monitor mass changes, likely due to the large extent of this region, the poorly constrained GIA signal and paleo-ice reconstruction (Bentley et al., 2014; Martín-Español et al., 2016; Small et al., 2019), and the relatively small mass imbalance in comparison to natural fluctuations in SMB in East Antarctica (Mottram et al., 2021).




Figure 3. Inter-comparison of rates of ice sheet mass balance of (a) the AIS, WAIS, EAIS, and APIS over the overlap period 2002-2019 and of (b) the GrIS during the overlap period 2003-2018 derived from the altimetry, gravimetry, and input-output techniques. The coloured bars represent the rates of mass balance and uncertainties of the aggregated technique time-series as calculated in Section 3 step (ii). The grey box represents the rate of mass balance of our final reconciled assessment calculated following the procedure detailed in Section 3 step (iii). The horizontal line in the middle of the box shows the reconciled rate of mass balance and the height of the box represents its associated uncertainty.

When examining the aggregated time-series of rate of mass change at annual resolution, we find the highest temporal correlation between the three time-series at the GrIS ($0.66 < r^2 < 0.83$). In addition, the gravimetry and input-output annual rates are also well-correlated at the APIS and WAIS ($r^2 = 0.83$). However, the altimetry mass balance time-series is poorly correlated with both the aggregated gravimetry and input-output time-series at the APIS and EAIS ($r^2 < 0.18$). The better correlation between the gravimetry and input-output time-series can be explained by their higher temporal resolutions, sufficient to resolve annual fluctuations in ice sheet mass balance which are substantial in these regions. Overall, we find that the vast majority of individual estimates of annual rates of mass balance included in this study fall within the uncertainty bounds of our reconciled estimate given their respective individual errors, with 96 %, 83 %, 83 %, 76 %, and 81 % of those annual rates of mass change falling within the reconciled uncertainty range at the GrIS, AIS, APIS, EAIS, and WAIS, respectively.

We integrate the combined mass balance estimates from gravimetry, altimetry, and the input-output method (Figure 2) to determine the cumulative mass lost from Antarctica and Greenland since 1992 (Figure 4). Antarctic mass loss continues to be dominated by ice discharge from West Antarctica where the signal is strongest – rising from $37 \pm 19 \text{ Gt yr}^{-1}$ between 1992 and 1996 to a maximum of $131 \pm 21 \text{ Gt yr}^{-1}$ between 2012 and 2016 (Table 2), before slowing slightly to $94 \pm 25 \text{ Gt yr}^{-1}$ during the last 5 years of our survey between 2017 and 2020. At the Antarctic Peninsula the increase in losses since the early 2000s that is generally associated with ice-shelf collapse (Rignot et al., 2004; Cook and Vaughan, 2010; Adusumilli et al., 2018) was masked briefly between 2012 and 2016, when the average rate of mass loss was reduced by 15 Gt yr^{-1} to $6 \pm 13 \text{ Gt yr}^{-1}$ in part due to an extreme snowfall event in 2016 (Wang et al., 2021; Chuter et al., 2021), before returning to $21 \pm 12 \text{ Gt yr}^{-1}$ between 2017 and 2020. East Antarctica remains the least certain component of Antarctic Ice Sheet mass balance, where the average 30-year mass trend is $3 \pm 15 \text{ Gt yr}^{-1}$. In all, the Antarctic Ice Sheet lost $2671 \pm 530 \text{ Gt}$ of ice between 1992 and 2020, raising the global sea level by $7.4 \pm 1.5 \text{ mm}$; after doubling in the mid-2000s from $62 \pm 41 \text{ Gt yr}^{-1}$ to $130 \pm 45 \text{ Gt yr}^{-1}$, increased Antarctic ice losses – largely driven by an acceleration in ice discharge from the Amundsen Sea Sector (Mouginot et al., 2014) – have persisted to the present-day. The rate of Greenland ice loss has remained highly variable during the last 5-year period of our updated assessment, ranging from $86 \pm 75 \text{ Gt yr}^{-1}$ in 2017 to a new maximum of $444 \pm 93 \text{ Gt yr}^{-1}$ in 2019 driven by exceptional surface melting during the summer (Tedesco and Fettweis, 2020). The majority of ice sheet losses have arisen from Greenland during our 29-year survey: $4892 \pm 457 \text{ Gt}$ in total at an average rate of $169 \pm 16 \text{ Gt yr}^{-1}$. Combined, Antarctica and Greenland lost $7563 \pm 699 \text{ Gt}$ of ice between 1992 and 2020, raising the global sea level by $21 \pm 2 \text{ mm}$.




Figure 4. Cumulative ice sheet mass changes. The shadings represent the associated uncertainties and are calculated following the procedure described in Section 3 (iv). The dashed lines show the results from our previous assessments (IMBIE-2).

Table 2. Rates of ice sheet mass change (Gt yr^{-1}). Rates are calculated from the first day (1st January) of the first year quoted to the last day (31st December) of the final year quoted in the table. For context, the last column gives the GMSL trend (mm yr^{-1}), calculated from the AVISO product (<https://www.aviso.altimetry.fr/mssl/>, last access: 12th April 2022) over the same period (as the GMSL record starts in 1993, we do not compute the fraction of sea level rise from the ice sheets for the first and last time period of the table). The percentage in brackets is the fraction of sea level rise driven by the ice sheets.

	GrIS (Gt yr^{-1})	AIS (Gt yr^{-1})	WAIS (Gt yr^{-1})	EAIS (Gt yr^{-1})	APIS (Gt yr^{-1})	GMSL (mm yr^{-1})
1992-1996	-35 ± 29	-70 ± 40	-37 ± 19	-27 ± 33	-7 ± 11	-
1997-2001	-48 ± 36 [4.0 %]	-19 ± 39 [1.6 %]	-42 ± 19 [3.5 %]	21 ± 32 [-1.7 %]	2 ± 11 [-0.2 %]	3.37 ± 0.11
2002-2006	-180 ± 39 [15.5 %]	-62 ± 41 [5.4 %]	-64 ± 20 [5.5 %]	21 ± 34 [-1.8 %]	-20 ± 11 [1.7 %]	3.23 ± 0.06
2007-2011	-280 ± 38 [31.8 %]	-130 ± 45 [14.8 %]	-129 ± 23 [14.6 %]	19 ± 36 [-2.2 %]	-21 ± 12 [2.3 %]	2.44 ± 0.11
2012-2016	-213 ± 40 [11.9 %]	-150 ± 43 [8.4 %]	-131 ± 21 [7.3 %]	-13 ± 35 [0.7 %]	-6 ± 13 [0.3 %]	4.96 ± 0.13
2017-2020	-257 ± 42 [17.7 %]	-115 ± 55 [7.9 %]	-94 ± 25 [6.5 %]	0 ± 47 [0.0 %]	-21 ± 12 [1.5 %]	4.03 ± 0.09
1992-2020	-169 ± 16	-92 ± 18	-82 ± 9	3 ± 15	-13 ± 5	-

5 Discussion

5.1. Comparison to previous IMBIE assessment

Finally, we assess the consistency of our results with our most recent assessment of ice sheet mass balance (IMBIE-2) to evaluate the impact of incorporating updated datasets and using an updated processing scheme. During their overlapping periods – 1992 to 2017 for Antarctica and 1992 to 2018 for Greenland – the results of this study and IMBIE-2 are in agreement within their respective uncertainties with rates of mass change of $-150.0 \pm 16 \text{ Gt yr}^{-1}$ and $-150 \pm 12 \text{ Gt yr}^{-1}$ for GrIS, respectively and rates of $-86 \pm 19 \text{ Gt yr}^{-1}$ and $-103 \pm 22 \text{ Gt yr}^{-1}$ for AIS, respectively. Next, comparing rates of mass balance within calendar years shows that results from this study and our previous assessment are consistent across all years for all ice sheets, except for two years at the start of our record (1992 and 1995) at the GrIS for which the difference between our mass balance assessments exceeds their respective uncertainty bounds. On average, the magnitude of the differences in annual rates of mass balance is 36 Gt yr^{-1} at GrIS, 33 Gt yr^{-1} at AIS, 12 Gt yr^{-1} at APIS, 31 Gt yr^{-1} at EAIS, and 23 Gt yr^{-1} at WAIS. The relatively small differences between our previous and current mass balance assessments originate from a combination of our inclusion of updated datasets and the implementation of an updated processing scheme in this study. In all ice sheet regions, participant datasets have been updated compared to our previous assessment. In addition, in this study we apply a common processing scheme to the AIS and GrIS, while in our previous study the mass balance assessments were aggregated with and without inverse-error weighting in the respective regions.

5.2 Limitations of this study and roadmap for future improvements

In this section, we discuss the limitations of our dataset and a roadmap to improve ice sheet mass balance assessments. The inclusion of the peripheral glaciers and ice caps in the vicinity of the Greenland and Antarctic Ice Sheets is ambiguous in our assessment as not all individual estimates of ice sheet mass balance included here account for those. This relates to the varying ability of satellite techniques to resolve mass balance over those small glaciated areas. Space gravimetry has a coarse spatial resolution of a few hundred kilometres which is not sufficient to separate signals of mass change originating from the ice sheet and its peripheral glaciers. On the other hand, the altimetry estimates included in this assessment exclude the peripheral glaciers and ice caps due to the complex terrain of these glaciers and their relatively small size compared to the footprint size of traditional pulse-limited altimeters. Finally, the input-output estimates do include mass changes from these glaciers, mostly by estimating their changes in SMB. Despite covering a relatively small area (around one tenth of the area of the ice sheets) (Pfeffer et al., 2014), these glaciers contribute significantly to global mean sea level rise with ice losses originating from the glaciers peripheral to the Greenland and Antarctic Ice Sheets amounting to $36 \pm 6 \text{ Gt yr}^{-1}$ and $21 \pm 5 \text{ Gt yr}^{-1}$ during the period 2010-2019, respectively (Hugonnet et al., 2021). In addition, ice losses have accelerated in the periphery of the Greenland Ice Sheet, with glacier mass loss increasing by 64 % between 2003-2009 and 2018-2021 (Khan et al., 2022). These glaciers therefore need to be accounted for without ambiguity in future IMBIE assessments to remove systematic biases between the different satellite techniques linked to their (non-)inclusion in individual mass balance estimates. Here, we performed a simple analysis to assess the potential impact of the ambiguous inclusion of these peripheral ice masses in our reconciled mass balance assessment and showed that this impact is limited thanks to the fact that we are aggregating different satellite techniques together – including some able to resolve separately ice sheet mass changes – and a different weighting has been applied

400 to the different estimates included. However, future approaches to address this issue will require careful treatment
of the leakage of mass signals between the ice sheets and their peripheral GICs within the gravimetry community,
rather than being limited to a subsequent removal of the contribution of these glaciers as we have done here. This
will nonetheless require robust mass balance estimates for developing and evaluating new methods. The recent
inventory of Earth's glaciers from satellite photogrammetry (Hugonnet et al., 2021), recent progress in satellite
405 altimetry – with the development of CryoSat-2 swath radar altimetry for measuring mass changes of mountain
glaciers (Foresta et al., 2016; Jakob et al., 2021) and the launch of ICESat-2 – and new community initiatives,
such as GlambIE (the Glacier mass balance Inter-comparison Exercise), will further contribute to this effort.

Continuing efforts to understand the remaining differences between altimetry, gravimetry, and the input-output
method is critical to provide more robust observational estimates of the contribution of the ice sheets to GMSL.
410 Producing estimates with a better temporal resolution by using data from the newest satellite missions,
reprocessing the satellite record with the newest geophysical corrections, and using a better uncertainty
characterisation, will undoubtedly help further reconcile satellite assessments of ice sheet mass balance produced
from different techniques. To achieve this, it is also important to assess the impact of SMB and GIA models. SMB
processes are responsible for a large proportion of Greenland's ice losses (and to a lesser extent of Antarctica's
415 ice losses) (Enderlin et al., 2014; Shepherd et al., 2020), and thus pursuing the efforts of recent model inter-
comparisons (Fettweis et al., 2020; Mottram et al., 2021) is key to improve the agreement between input-output
estimates but also to partition mass trends into SMB and ice dynamics components as it provides critical
information on the dominant processes at play. A model-inter-comparison of GIA models would also be timely
as new approaches have been developed in recent years to determine the GIA signal (Whitehouse, 2018). New
420 data-driven solutions that rely on present day geodetic observations (e.g. Riva et al., 2009; Vishwakarma et al.,
2022), and solutions derived from coupling a GIA model to an ice sheet mode (de Boer et al., 2017) have become
available. Examining the variability of GIA solutions determined from forward models, data inversion, and
coupled models will help reducing uncertainties in space gravimetry estimates of ice sheet mass balance.

Finally, improving the spatial resolution of the IMBIE assessment by producing time-series of mass changes
425 within the individual basins of the Greenland and Antarctic Ice Sheets will also contribute to further identify areas
of similarities and disagreement between satellite techniques (Sutterley et al., 2014) and will support the
identification of spatial biases in satellite estimates of ice sheet mass balance. In addition, regional assessments of
ice sheet mass balance could support the evaluation and calibration of ice sheet models, contributing to reducing
uncertainties in future sea level rise projections (Edwards et al., 2021; Nias et al., 2019).

430 **6 Conclusions**

We combine 50 estimates of ice sheet mass balance, 27 for Greenland and 23 for Antarctica, to produce a new
reconciled estimate of ice sheet mass balance showing that the ice sheets lost $7,563 \pm 699$ Gt of ice between 1992
and 2020. Ice losses have accelerated at both ice sheets over this 29-year record and the rate of ice loss is now 5
times higher in Greenland and 25 % higher in Antarctica compared to the early 1990s. Our assessment shows that
435 the altimetry, gravimetry, and input-output method are in close agreement in Greenland with a spread of 19 Gt yr^{-1}
over their common time period, which represents only 10.9 % of the rate of imbalance. In Antarctica, the spread

between techniques is 4 times larger than in Greenland, mostly due to large differences between estimates for the East Antarctic Ice Sheet. To further explore and interpret differences between geodetic techniques, producing altimetry estimates with a higher temporal resolution (especially during the first half of the satellite altimetry record), better GIA constraints for the gravimetry estimates, and additional estimates of ice sheet mass balance via the input-output method would improve the comparison and aggregation of ice sheet mass balance estimates. Continuously monitoring the mass balance of the ice sheets and producing annual updates of Greenland and Antarctica mass balance is critical to track their contribution to global mean sea level and constrain projections of future sea-level rise.

445 **7 Data Availability**

The aggregated Greenland and Antarctic Ice Sheets mass balance data and associated errors generated in this study are freely available at the NERC Polar Data Centre, <https://doi.org/10.5285/77B64C55-7166-4A06-9DEF-2E400398E452> (The IMBIE Team, 2021).

8 Code Availability

450 The code used to compute and aggregate rates of ice sheet mass change and their errors is freely available at <https://github.com/IMBIE>.

Author Contribution

The executive committee of IMBIE (A.S., E.R.I., N.J.S., I.N.O., C.A., M.B., M.H., I.J., M.D.K., G.K., S.N., A.J.P., E.R., T.S., K.M.S., B.E.S., L.S.S., I.V., P.L.W.) designed the study. A.S., L.S.S., A.G., L.G., N.G., B.C.G., 455 V.H., S.A.K., H.K., M.M., J.N., N.P., L.S., and S.B.S. contributed altimetry estimates. M.H., I.V., A.B., R.F., A.G., A. Groh, C.H., C.L., B.D.L., J.N., I.S., H.V.S., K.S., E.J.O.S., T.C.S., B.D.V., D.W., and B.W. contributed gravimetry estimates. E.R., A.P.A., W.C., J.M., and B.S. contributed input-output estimates. M.B., C. Agosta, X.F., H.G., C.K., P.L.L., S.H.M., R.M., and B.N. contributed SMB estimates. G.A., B.S.L., D.M., W.R.P., G.S., M.W., and W.W. contributed GIA estimates. I.N.O. and M.E.P. performed the mass balance data collation. I.N.O. 460 prepared the datasets comparison. I.N.O. led the writing and prepared the other figures and tables. I.N.O., A.S. and T. Slater wrote the manuscript. All authors participated in the data interpretation and commented on the manuscript.

Competing interests

C. A. is member of the editorial board of journal Earth System Science Data.

465 **Acknowledgements**

This work is an outcome of the Ice Sheet Mass Balance Inter-comparison Exercise (IMBIE) supported by the ESA EOEP-5 ‘EO Science for Society’, the ESA ‘Climate Change Initiative’, and the NASA Cryosphere Program. A portion of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under

a contract with the National Aeronautics and Space Administration (NASA) (80NM0018D0004). Funding for E.I. and N-J.S. was provided by NASA ROSES solicitation NNH20ZDA001N-CRYO in response to Proposal 20-CRYO2020-0003. GEUS data provided from the Programme for Monitoring of the Greenland Ice Sheet (www.PROMICE.dk) was funded by the Danish Ministry of Climate, Energy and Utilities. M.M. acknowledges the support of the UK NERC Centre for Polar Observation and Modelling (CPOM), and the Lancaster University-UKCEH Centre of Excellence in Environmental Data Science. P. L. L. gratefully acknowledges the contributions of Aarhus University Interdisciplinary Centre for Climate Change (iClimate, Aarhus University). N.G. used CryoSat data obtained from ESA at cs2eo.org and via the CryoTEMPO-EOLIS project <https://cryotempo-eolis.org/>. G.S. is funded by a research grant of DIFA (Dipartimento di Fisica e Astronomia “Augusto Righi”) of the Alma Mater Studiorum Università di Bologna. J.N. and A.G. were supported by the ITS_LIVE project awarded through NASA MEaSUREs program, and the NASA Cryosphere program through participation in the ICESat-2 science team. I.S. acknowledges funding by the Helmholtz Climate Initiative REKLIM (Regional Climate Change), a joint research project of the Helmholtz Association of German Research Centres (HGF). Ice velocity data for Greenland and Antarctica provided by UC Irvine is funded by NASA MEaSUREs program. BedMachine Antarctica is funded by NASA MEaSUREs program. BedMachine Greenland is funded by research grants from NASA Operation IceBridge Mission. BW was funded by NWO VIDI grant 016.Vidi.171.063.

References

- Adusumilli, S., Fricker, H. A., Medley, B., Padman, L., and Siegfried, M. B.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nat. Geosci.*, 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- Adusumilli, S., Fricker, H. A., Siegfried, M. R., Padman, L., Paolo, F. S., and Ligtenberg, S. R. M.: Variable Basal Melt Rates of Antarctic Peninsula Ice Shelves, 1994–2016, *Geophysical Research Letters*, 45, 4086–4095, <https://doi.org/10.1002/2017GL076652>, 2018.
- Aschwanden, A., Bartholomaus, T. C., Brinkerhoff, D. J., and Truffer, M.: Brief communication: A roadmap towards credible projections of ice sheet contribution to sea level, *The Cryosphere*, 15, 5705–5715, <https://doi.org/10.5194/tc-15-5705-2021>, 2021.
- Bentley, M. J., Ó Cofaigh, C., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C., Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith, J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E. A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W., Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Gолledge, N. R., Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A. S., Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P., Kristoffersen, Y., Kuhn, G., Leventer,

- A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J., Massé, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W., Nitsche, F. O., O'Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M., Simms, A. R., Spiegel, C., Stollendorf, T. D., Sugden, D. E., van der Putten, N., van Ommen, T., Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witus, A. E., and Zwartz, D.: A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum, *Quaternary Science Reviews*, 100, 1-9, 2014.
- Caron, L. and Ivins, E. R.: A baseline Antarctic GIA correction for space gravimetry, *Earth and Planetary Science Letters*, 531, 115957, <https://doi.org/10.1016/j.epsl.2019.115957>, 2020.
- 515 Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D., and Unnikrishnan, A. S.: Sea Level Change, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1137–1216, <https://doi.org/10.1017/CBO9781107415324.026>, 2013.
- 520 Chuter, S. J., Zammit-Mangion, A., Rougier, J., Dawson, G., and Bamber, J. L.: Mass evolution of the Antarctic Peninsula over the last two decades from a joint Bayesian inversion, *The Cryosphere*, 1–32, <https://doi.org/10.5194/tc-2021-178>, 2021.
- 525 Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years, *The Cryosphere*, 4, 77–98, <https://doi.org/10.5194/tc-4-77-2010>, 2010.
- 530 Csatho, B. M., Schenk, A. F., van der Veen, C. J., Babonis, G., Duncan, K., Rezvanbehbahani, S., van den Broeke, M. R., Simonsen, S. B., Nagarajan, S., and van Angelen, J. H.: Laser altimetry reveals complex pattern of Greenland Ice Sheet dynamics, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 18478-18424, 2014.
- 535 de Boer, B., Stocchi, P., Whitehouse, P. L., and van de Wal, R. S. W.: Current state and future perspectives on coupled ice-sheet – sea-level modelling, *Quaternary Science Reviews*, 169, 13-28, <https://doi.org/10.1016/j.quascirev.2017.05.013>, 2017.
- 540 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd, A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R., Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-Fenzi, B. K., Gladstone, R., Gолledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts, P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le Clec'h, S., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Mallet, J.-H., Martin, D.
- 545 F., Maussion, F., Morlighem, M., O'Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radić, V.,

Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N.-J., Shannon, S., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R., Zekollari, H., Zhao, C., Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, <https://doi.org/10.1038/s41586-021-03302-y>, 2021.

550

Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M. J., Angelen, J. H., and Broeke, M. R.: An improved mass budget for the Greenland ice sheet, *Geophysical Research Letters*, 41, 866–872, <https://doi.org/10.1002/2013GL059010>, 2014.

555

Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhout, L., Wilton, D., Wouters, B., Ziemen, F., and Zolles, T.: GrSMBMIP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet, *The Cryosphere*, 14, 3935–3958, <https://doi.org/10.5194/tc-14-3935-2020>, 2020.

560

Foresta, L., Gourmelen, N., Pálsson, F., Nienow, P., Björnsson, H., and Shepherd, A.: Surface elevation change and mass balance of Icelandic ice caps derived from swath mode CryoSat-2 altimetry, *Geophysical Research Letters*, 43(12), 138–12, <https://doi.org/10.1002/2016GL071485>, 2016.

565

Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., Slangen, A. B. A., and Yu, Y.: *Ocean, Cryosphere and Sea Level Change*, Cambridge University Press, 2021.

570

Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., van den Broeke, M., and Nilsson, J.: Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *The Cryosphere*, 12, 521–547, <https://doi.org/10.5194/tc-12-521-2018>, 2018.

Groh, A. and Horwath, M.: Antarctic Ice Mass Change Products from GRACE/GRACE-FO Using Tailored Sensitivity Kernels, *Remote Sensing*, 13, 1736, <https://doi.org/10.3390/rs13091736>, 2021.

575

Hanna, E., Cappelen, J., Fettweis, X., Mernild, S. H., Mote, T. L., Mottram, R., Steffen, K., Ballinger, T. J., and Hall, R. J.: Greenland surface air temperature changes from 1981 to 2019 and implications for ice-sheet melt and mass-balance change, *International Journal of Climatology*, 41, E1336–E1352, <https://doi.org/10.1002/joc.6771>, 2021.

580

Hanson, S., Nicholls, R., Ranger, N., Hallegatte, S., Corfee-Morlot, J., Herweijer, C., and Chateau, J.: A global ranking of port cities with high exposure to climate extremes, *Climatic Change*, 104, 89–111, <https://doi.org/10.1007/s10584-010-9977-4>, 2011.

- 585 Hofer, S., Tedstone, A. J., Fettweis, X., and Bamber, J. L.: Decreasing cloud cover drives the recent mass loss on the Greenland Ice Sheet, *Science*, 3, e1700584–e1700584, <https://doi.org/10.1126/sciadv.1700584>, 2017.
- Hogg, A. E., Shepherd, A., Cornford, S. L., Briggs, K. H., Gourmelen, N., Graham, J. A., Joughin, I., Mouginot, J., Nagler, T., Payne, A. J., Rignot, E., and Wuite, J.: Increased ice flow in Western Palmer Land linked to ocean melting, *Geophysical Research Letters*, 44, 4159–4167, <https://doi.org/10.1002/2016GL072110>, 2017.
- 590 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- 595 IMBIE Team: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- IMBIE Team: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, <https://doi.org/10.1038/s41586-019-1855-2>, 2020.
- 600 IMBIE Team: Antarctic and Greenland Ice Sheet mass balance 1992–2020 for IPCC AR6 (Version 1.0) [Data set]. UK Polar Data Centre, Natural Environment Research Council, UK Research & Innovation, available at: <https://doi.org/10.5285/77B64C55-7166-4A06-9DEF-2E400398E452>, 2021.
- 605 Jakob, L., Gourmelen, N., Ewart, M., and Plummer, S.: Spatially and temporally resolved ice loss in High Mountain Asia and the Gulf of Alaska observed by CryoSat-2 swath altimetry between 2010 and 2019, *The Cryosphere*, 15, 1845–1862, <https://doi.org/10.5194/tc-15-1845-2021>, 2021.
- Jenkins, A., Shoosmith, D., Dutrieux, P., Jacobs, S., Kim, T. W., Lee, S. H., Ha, H. K., and Stammerjohn, S.: West Antarctic Ice Sheet retreat in the Amundsen Sea driven by decadal oceanic variability, *Nature Geosci*, 11, 733–738, <https://doi.org/10.1038/s41561-018-0207-4>, 2018.
- Joughin, I., Shapero, D., Smith, B., Dutrieux, P., and Barham, M.: Ice-shelf retreat drives recent Pine Island Glacier speedup, *Science*, 7, eabg3080, <https://doi.org/10.1126/sciadv.abg3080>, 2021.
- 615 Khan, S. A., Colgan, W., Neumann, T. A., van den Broeke, M. R., Brunt, K. M., Noël, B., Bamber, J. L., Hassan, J., and Björk, A. A.: Accelerating Ice Loss From Peripheral Glaciers in North Greenland, *Geophysical Research Letters*, 49, e2022GL098915, <https://doi.org/10.1029/2022GL098915>, 2022.
- King, M. D., Howat, I. M., Jeong, S., Noh, M. J., Wouters, B., Noël, B., and van den Broeke, M. R.: Seasonal to decadal variability in ice discharge from the Greenland Ice Sheet, *The Cryosphere*, 12, 3813, <https://doi.org/10.5194/tc-12-3813-2018>, 2018.
- 620 King, M. D., Howat, I. M., Candela, S. G., Noh, M. J., Jeong, S., Noël, B. P. Y., van den Broeke, M. R., Wouters,

- 625 B., and Negrete, A.: Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat, *Communications Earth & Environment*, 1, 1–7, <https://doi.org/10.1038/s43247-020-0001-2>, 2020.
- Konrad, H., Gilbert, L., Cornford, S. L., Payne, A., Hogg, A., Muir, A., and Shepherd, A.: Uneven onset and pace of ice-dynamical imbalance in the Amundsen Sea Embayment, West Antarctica, *Geophysical Research Letters*, 44, 910–918, <https://doi.org/10.1002/2016GL070733>, 2017.
- 630 Konrad, H., Shepherd, A., Gilbert, L., Hogg, A. E., McMillan, M., Muir, A., and Slater, T.: Net retreat of Antarctic glacier grounding lines, *Nature Geosci*, 11, 258–262, <https://doi.org/10.1038/s41561-018-0082-z>, 2018.
- Kulp, S. A. and Strauss, B. H.: New elevation data triple estimates of global vulnerability to sea-level rise and coastal flooding, *Nat Commun*, 10, 1–12, <https://doi.org/10.1038/s41467-019-12808-z>, 2019.
- 635 Leeson, A. A., Shepherd, A., Briggs, K., Howat, I., Fettweis, X., Morlighem, M., and Rignot, E.: Supraglacial lakes on the Greenland ice sheet advance inland under warming climate, *Nature Climate Change*, 5, 51–55, <https://doi.org/10.1038/nclimate2463>, 2015.
- 640 Lemos, A., Shepherd, A., McMillan, M., and Hogg, A. E.: Seasonal Variations in the Flow of Land-Terminating Glaciers in Central-West Greenland Using Sentinel-1 Imagery, *Remote Sensing*, 10, 1878, <https://doi.org/10.3390/rs10121878>, 2018.
- 645 Luthcke, S. B., Zwally, H. J., Abdalati, W., Rowlands, D. D., Ray, R. D., Nerem, R. S., Lemoine, F. G., McCarthy, J. J., and Chinn, D. S.: Recent Greenland Ice Mass Loss by Drainage System from Satellite Gravity Observations, *Science*, 314, 1286-1289, <https://doi.org/10.1126/science.1130776>, 2006.
- 650 Mankoff, K. D., Fettweis, X., Langen, P. L., Stendel, M., Kjeldsen, K. K., Karlsson, N. B., Noël, B., van den Broeke, M. R., Solgaard, A., Colgan, W., Box, J. E., Simonsen, S. B., King, M. D., Ahlstrøm, A. P., Andersen, S. B., and Fausto, R. S.: Greenland ice sheet mass balance from 1840 through next week, *Earth Syst. Sci. Data*, 13, 5001-5025, <https://doi.org/10.5194/essd-13-5001-2021>, 2021.
- 655 Martín-Español, A., King, M. A., Zammit-Mangion, A., Andrews, S. B., Moore, P., and Bamber, J. L.: An assessment of forward and inverse GIA solutions for Antarctica, *Journal of geophysical research. Solid earth*, 121, 6947-6965, <https://doi.org/10.1002/2016JB013154>, 2016.
- 660 Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J. L., Prats-Iraola, P., and Dini, L.: Rapid glacier retreat rates observed in West Antarctica, *Nat. Geosci.*, 1–6, <https://doi.org/10.1038/s41561-021-00877-z>, 2022.
- Moon, T., Joughin, I., Smith, B., and Howat, I.: 21st-Century Evolution of Greenland Outlet Glacier Velocities, *Science*, 336, 576-578, <https://doi.org/10.1126/science.1219985>, 2012.

- 665 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44, 11,051-011,061, <https://doi.org/10.1002/2017GL074954>, 2017.
- 670 Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., Boberg, F., van de Berg, W. J., Fettweis, X., Gossart, A., van Lipzig, N. P. M., van Meijgaard, E., Orr, A., Phillips, T., Webster, S., Simonsen, S. B., and Souverijns, N.: What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates, *The Cryosphere*, 15, 3751–3784, <https://doi.org/10.5194/tc-15-3751-2021>, 2021.
- 675 Mouginot, J., Rignot, E., and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophysical Research Letters*, 41, 1576–1584, <https://doi.org/10.1002/2013GL059069>, 2014.
- 680 Mouginot, J., Rignot, E., Scheuchl, B., and Millan, R.: Comprehensive Annual Ice Sheet Velocity Mapping Using Landsat-8, Sentinel-1, and RADARSAT-2 Data, *Remote Sensing*, 9, 364, <https://doi.org/10.3390/rs9040364>, 2017.
- 685 Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, 116, 9239-9244, <https://doi.org/10.1073/pnas.1904242116>, 2019.
- Nias, I. J., Cornford, S. L., Edwards, T. L., Gourmelen, N., and Payne, A. J.: Assessing Uncertainty in the Dynamical Ice Response to Ocean Warming in the Amundsen Sea Embayment, West Antarctica, *Geophysical Research Letters*, 46, 11253-11260, <https://doi.org/10.1029/2019GL084941>, 2019.
- 690 Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327–331, <https://doi.org/10.1126/science.aaa0940>, 2015.
- Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331-1335, <https://doi.org/10.1126/science.aaz5487>, 2020.
- 695 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., and Sharp, M. J.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of Glaciology*, 60, 537-552, <https://doi.org/10.3189/2014JoG13J176>, 2014.
- 700

- Pritchard, H. D., Arthern, R. J., Vaughan, D. G., and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461, 971-975, <https://doi.org/10.1038/nature08471>, 2009.
- 705 Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A., and Thomas, R.: Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020697>, 2004.
- Rignot, E. and Kanagaratnam, P.: Changes in the Velocity Structure of the Greenland Ice Sheet, *Science*, 311, 986–990, <https://doi.org/10.1126/science.1121381>, 2006.
- 710 Rignot, E., Mouginot, J., and Scheuchl, B.: Antarctic grounding line mapping from differential satellite radar interferometry, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL047109>, 2011a.
- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A., and Lenaerts, J. T. M.: Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL046583>, 2011.
- 715 Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979-2017, *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 116, 1095-1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- 720 Ritz, C., Edwards, T. L., Durand, G., Payne, A. J., Peyaud, V., and Hindmarsh, R. C. A.: Potential sea-level rise from Antarctic ice-sheet instability constrained by observations, 528, 115–118, <https://doi.org/10.1038/nature16147>, 2015.
- Riva, R. E. M., Gunter, B. C., Urban, T. J., Vermeersen, B. L. A., Lindenbergh, R. C., Helsen, M. M., Bamber, J. L., van de Wal, R. S. W., van den Broeke, M. R., and Schutz, B. E.: Glacial Isostatic Adjustment over Antarctica from combined ICESat and GRACE satellite data, *Earth and Planetary Science Letters*, 288, <https://doi.org/10.1016/j.epsl.2009.10.013>, 516-523, 2009.
- 725 Sandberg Sørensen, L., Simonsen, S. B., Forsberg, R., Khvorostovsky, K., Meister, R., and Engdahl, M. E.: 25 years of elevation changes of the Greenland Ice Sheet from ERS, Envisat, and CryoSat-2 radar altimetry, *Earth and Planetary Science Letters*, 495, 234-241, <https://doi.org/10.1016/j.epsl.2018.05.015>, 2018.
- 730 Sasgen, I., Wouters, B., Gardner, A. S., King, M. D., Tedesco, M., Landerer, F. W., Dahle, C., Save, H., and Fettweis, X.: Return to rapid ice loss in Greenland and record loss in 2019 detected by the GRACE-FO satellites, *Commun Earth Environ*, 1, 1–8, <https://doi.org/10.1038/s43247-020-0010-1>, 2020.
- 735 Selley, H. L., Hogg, A. E., Cornford, S., Dutrieux, P., Shepherd, A., Wuite, J., Floricioiu, D., Kusk, A., Nagler, T., Gilbert, L., Slater, T., and Kim, T.-W.: Widespread increase in dynamic imbalance in the Getz region of Antarctica from 1994 to 2018, 12, 1133, <https://doi.org/10.1038/s41467-021-21321-1>, 2021.

740

Shepherd, A., Ivins, E. R., A. G., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Jan, T. M. L., Li, J., Stefan, R. M. L., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Ernst, J. O. S.,
745 Smith, B., Sundal, A. V., Jan, H. v. A., Willem, J. v. d. B., Michiel, R. v. d. B., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338, 1183-1189, <https://doi.org/10.1126/science.1228102>, 2012.

750

Shepherd, A. and Nowicki, S.: Improvements in ice-sheet sea-level projections, *Nature Climate Change*, 7, 672–674, <https://doi.org/10.1038/nclimate3400>, 2017.

755

Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophysical Research Letters*, 46, 8174–8183, <https://doi.org/10.1029/2019GL082182>, 2019.

760

Slater, T., Hogg, A. E., and Mottram, R.: Ice-sheet losses track high-end sea-level rise projections, *Nature Climate Change*, 10, 879–881, <https://doi.org/10.1038/s41558-020-0893-y>, 2020.

765

Slater, T., Shepherd, A., McMillan, M., Leeson, A., Gilbert, L., Muir, A., Munneke, P. K., Noël, B., Fettweis, X., van den Broeke, M., and Briggs, K.: Increased variability in Greenland Ice Sheet runoff from satellite observations, *Nat Commun*, 12, 6069, <https://doi.org/10.1038/s41467-021-26229-4>, 2021.

770

Small, D., Bentley, M.J., Jones, R.S., Pittard, M.L., and Whitehouse, P.L.: Antarctic ice sheet palaeo-thinning rates from vertical transects of cosmogenic exposure ages, *Quat. Sci. Rev*, 206, 65–80, <https://doi.org/10.1016/j.quascirev.2018.12.024>, 2019.

775

Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, 368, 1239-1242, <https://doi.org/10.1126/science.aaz5845>, 2020.

775

Sørensen, L. S., Simonsen, S. B., Nielsen, K., Lucas-Picher, P., Spada, G., Adalgeirsdottir, G., Forsberg, R., and Hvidberg, C. S.: Mass balance of the Greenland ice sheet (2003–2008) from ICESat data – the impact of interpolation, sampling and firn density, *The Cryosphere*, 5, 173–186, <https://doi.org/10.5194/tc-5-173-2011>, 2011.

Stevens, C. M., Verjans, V., Lundin, J. M. D., Kahle, E. C., Horlings, A. N., Horlings, B. I., and Waddington, E. D.: The Community Firn Model (CFM) v1.0, *Geoscientific Model Development*, 13(9), 4355-4377, 2020.

Straneo, F. and Heimbach, P.: North Atlantic warming and the retreat of Greenland's outlet glaciers, *Nature*, 504,

36–43, <https://doi.org/10.1038/nature12854>, 2013.

780

Sutterley, T. C., Velicogna, I., Csatho, B., van den Broeke, M., Rezvan-Behbahani, S., and Babonis, G.: Evaluating Greenland glacial isostatic adjustment corrections using GRACE, altimetry and surface mass balance data, *Environmental Research Letters*, 9, 014004, <https://doi.org/10.1088/1748-9326/9/1/014004>, 2014a.

785

Sutterley, T. C., Velicogna, I., Rignot, E., Mouginot, J., Flament, T., Van Den Broeke, M. R., info:eu, r. d. n., Van Wessem, J. M., Reijmer, C. H., and info:eu, r. d. n.: Mass loss of the Amundsen Sea Embayment of West Antarctica from four independent techniques, *Geophysical Research Letters*, 41, 8421-8428, <https://doi.org/10.1002/2014GL061940>, 2014b.

790

Tapley, B. D., Watkins, M. M., Flechtner, F., Reigber, C., Bettadpur, S., Rodell, M., Sasgen, I., Famiglietti, J. S., Landerer, F. W., Chambers, D. P., Reager, J. T., Gardner, A. S., Save, H., Ivins, E. R., Swenson, S. C., Boening, C., Dahle, C., Wiese, D. N., Dobslaw, H., Tamisiea, M. E., and Velicogna, I.: Contributions of GRACE to understanding climate change, *Nature Climate Change*, 5, 358-369, <https://doi.org/10.1038/s41558-019-0456-2>, 2019.

795

Tedesco, M. and Fettweis, X.: Unprecedented atmospheric conditions (1948–2019) drive the 2019 exceptional melting season over the Greenland ice sheet, *The Cryosphere*, 14, 1209–1223, <https://doi.org/10.5194/tc-14-1209-2020>, 2020.

800

Trusel, L. D., Das, S. B., Osman, M. B., Evans, M. J., Smith, B. E., Fettweis, X., McConnell, J. R., Noël, B. P. Y., and Broeke, M. R. van den: Nonlinear rise in Greenland runoff in response to post-industrial Arctic warming, *Nature*, 564, 104–108, <https://doi.org/10.1038/s41586-018-0752-4>, 2018.

805

Velicogna, I., Mohajerani, Y., Geruo, A., Landerer, F., Mouginot, J., Noel, B., Rignot, E., Sutterley, T., van den Broeke, M., van Wessem, J. M., and Wiese, D.: Continuity of ice sheet mass loss in Greenland and Antarctica from the GRACE and GRACE Follow-Onmissions, *Geophysical Research Letters*, 47, e2020GL087291, <https://doi.org/10.1029/2020gl087291>, 2020.

Velicogna I and Wahr J: Measurements of time-variable gravity show mass loss in Antarctica. *Science*, 2006, 311, 1754-1756, <https://doi.org/10.1126/science.1123785>, 2006.

810

Vishwakarma, B. D., Ziegler, Y., Bamber, J. L., and Royston, S.: Separating GIA signal from surface mass change using GPS and GRACE data, *Geophysical Journal International*, 232, 537-547, <https://doi.org/10.1093/gji/ggac336>, 2022.

815

Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L., and Storlazzi, C. D.: Doubling of coastal flooding frequency within decades due to sea-level rise, *Sci Rep*, 7, 1–9, <https://doi.org/10.1038/s41598-017-01362-7>, 2017.

Wang, L., Davis, J. L., and Howat, I. M.: Complex Patterns of Antarctic Ice Sheet Mass Change Resolved by Time-Dependent Rate Modeling of GRACE and GRACE Follow-On Observations, *Geophysical Research Letters*, 48, e2020GL090961, <https://doi.org/10.1029/2020GL090961>, 2021.

820

Wcrp Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth Syst. Sci. Data*, 10, 1551-1590, <https://doi.org/10.5194/essd-10-1551-2018>, 2018.

Whitehouse, P. L.: Glacial isostatic adjustment modelling: historical perspectives, recent advances, and future directions, *Earth Surf. Dynam.*, 6, 401-429, <https://doi.org/10.5194/esurf-6-401-2018>, 2018.

825

Wingham, D. J., Ridout, A. J., Scharroo, R., Arthern, R. J., and Shum, C. K.: Antarctic Elevation Change from 1992 to 1996, *Science*, 282, 456-458, <https://doi.org/10.1126/science.282.5388.456>, 1998.

830

Zwally, H. J., Giovinetto, M. B., Beckley, M. A. and Saba, J. L: Antarctic and Greenland Drainage Systems, GSFC Cryospheric Sciences Laboratory, http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php, 2012.

835

840

Appendix A

Table A1. References of the datasets, methods, GIA and SMB models employed by participants of the input-output, altimetry, and gravimetry experiment groups.

IOM	Andersen	Andersen, M. L. <i>et al.</i> Basin-scale partitioning of Greenland ice sheet mass balance components (2007–2011). <i>Earth and Planetary Science Letters</i> 409 , 89–95 (2015).
	Colgan	Colgan, W. <i>et al.</i> Greenland ice sheet mass balance assessed by PROMICE (1995–2015). <i>Geological Survey of Denmark and Greenland Bulletin</i> 43 (2019).
	Mouginot	Mouginot, J. <i>et al.</i> Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018. <i>PNAS</i> 116 , 9239–9244 (2019).
	Rignot	Rignot E. <i>et al.</i> Four decades of Antarctic Ice Sheet mass balance from 1979-2017. <i>PNAS</i> 116 (4), 1095-1103 (2019).
ALT	Gourmelen	Gourmelen, N. <i>et al.</i> CryoSat-2 swath interferometric altimetry for mapping ice elevation and elevation change. <i>Advances in Space Research</i> 62 , 1226–1242 (2018).
	Gunter	Gunter, B. C. <i>et al.</i> Empirical estimation of present-day Antarctic glacial isostatic adjustment and ice mass change. <i>The Cryosphere</i> 8 , 743–760 (2014).
	Helm	Helm, V., Humbert, A. & Miller, H. Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2. <i>The Cryosphere</i> 8 , 1539–1559 (2014).
	Khan	Khan, S. A. <i>et al.</i> Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming. <i>Nature Climate Change</i> 4 , 292–299 (2014).
	McMillan	McMillan, M. <i>et al.</i> A high-resolution record of Greenland mass balance. <i>Geophysical Research Letters</i> 43 , 7002–7010 (2016).
	Nilsson Gardner	Nilsson, J., Gardner, A., Sandberg Sørensen, L. & Forsberg, R. Improved retrieval of land ice topography from CryoSat-2 data and its impact for volume-change estimation of the Greenland Ice Sheet. <i>The Cryosphere</i> 10 , 2953–2969 (2016).
	Pie	Felikson, D. <i>et al.</i> Comparison of Elevation Change Detection Methods From ICESat Altimetry Over the Greenland Ice Sheet. <i>IEEE Transactions on Geoscience and Remote Sensing</i> 55 , 5494–5505 (2017).
	Sandberg Sørensen	Sørensen, L. S. <i>et al.</i> Mass balance of the Greenland ice sheet (2003–2008) from ICESat data – the impact of interpolation, sampling and firn density. <i>The Cryosphere</i> 5 , 173–186 (2011).
	Schroder	Schröder, L. <i>et al.</i> Four decades of Antarctic surface elevation changes from multi-mission satellite altimetry." <i>The Cryosphere</i> 13 (2), 427-449, (2019).
	Shepherd	Shepherd, A. <i>et al.</i> Trends in Antarctic Ice Sheet Elevation and Mass. <i>Geophysical Research Letters</i> 46 (14), 8174-8183 (2019).
Zwally	Zwally, H. J. <i>et al.</i> Mass gains of the Antarctic ice sheet exceed losses. <i>J. Glaciol.</i> 61 , 1019-1036 (2015).	
G	Blazquez	Blazquez, A. <i>et al.</i> Exploring the uncertainty in GRACE estimates of the mass redistributions at the

		Earth surface: implications for the global water and sea level budgets. <i>Geophys J Int</i> 215 , 415–430 (2018).
	Bonin	Bonin, J. & Chambers, D. Uncertainty estimates of a GRACE inversion modelling technique over Greenland using a simulation. <i>Geophys J Int</i> 194 , 212–229 (2013).
	Forsberg	Forsberg, R., Sørensen, L. & Simonsen, S. Greenland and Antarctica Ice Sheet Mass Changes and Effects on Global Sea Level. <i>Surv Geophys</i> 38 , 89–104 (2017).
	Gardner Nilsson	Gardner, A. S. <i>et al.</i> Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years. <i>The Cryosphere</i> 12 (2), 521–547 (2018).
	Groh	Groh, A., & Horwath, M. Antarctic Ice Mass Change Products from GRACE/GRACE-FO Using Tailored Sensitivity Kernels. <i>Remote Sensing</i> , 13, 1736, https://doi.org/10.3390/rs13091736 (2021).
	Harig	Harig, C. & Simons, F. J. Mapping Greenland’s mass loss in space and time. <i>PNAS</i> 109 , 19934–19937 (2012).
	Horvath	Horvath, A. G. Retrieving Geophysical Signals from Current and Future Satellite Missions. PhD thesis, Tech. Univ. Munich (2017).
	Luthcke	Luthcke, S. B. <i>et al.</i> Antarctica, Greenland and Gulf of Alaska land-ice evolution from an iterated GRACE global mascon solution. <i>Journal of Glaciology</i> 59 , 613–631 (2013).
	Moore	Andrews, S. B., Moore, P. & King, M. A. Mass change from GRACE: a simulated comparison of Level-1B analysis techniques. <i>Geophys J Int</i> 200 , 503–518 (2015).
	Save	Save, H., Bettadpur, S. & Tapley, B. D. High-resolution CSR GRACE RL05 mascons. <i>Journal of Geophysical Research: Solid Earth</i> 121 , 7547–7569 (2016).
	Schrama	Schrama, E. J. O., Wouters, B. & Rietbroek, R. A mascon approach to assess ice sheet and glacier mass balances and their uncertainties from GRACE data. <i>Journal of Geophysical Research: Solid Earth</i> 119 , 6048–6066 (2014).
	Seo	Seo, K.-W. <i>et al.</i> Surface mass balance contributions to acceleration of Antarctic ice mass loss during 2003–2013. <i>Journal of Geophysical Research: Solid Earth</i> 120 , 3617–3627 (2015).
	Velicogna	Velicogna, I., Sutterley, T. C. & Broeke, M. R. van den. Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data. <i>Geophysical Research Letters</i> 41 , 8130–8137 (2014).
	Vishwakarma	Vishwakarma, B. D., Horwath, M., Devaraju, B., Groh, A. & Sneeuw, N. A Data-Driven Approach for Repairing the Hydrological Catchment Signal Damage Due to Filtering of GRACE Products. <i>Water Resources Research</i> 53 , 9824–9844 (2017).
	Wiese	Wiese, D. N., Landerer, F. W. & Watkins, M. M. Quantifying and reducing leakage errors in the JPL RL05M GRACE mascon solution. <i>Water Resources Research</i> 52 , 7490–7502 (2016).
	Wouters	Wouters, B., Bamber, J. L., van den Broeke, M. R., Lenaerts, J. T. M. & Sasgen, I. Limits in detecting acceleration of ice sheet mass loss due to climate variability. <i>Nature Geoscience</i> 6 , 613–616 (2013).
GI	A13	A, Geruo, John Wahr, Shijie Zhong, Computations of the viscoelastic response of a 3-D compressible Earth to surface loading: an application to Glacial Isostatic Adjustment in Antarctica and Canada,

		<i>Geophysical Journal International</i> , Volume 192, Issue 2, February 2013, Pages 557–572, https://doi.org/10.1093/gji/ggs030
	W12a	Whitehouse, Pippa L., Michael J. Bentley, Glenn A. Milne, Matt A. King, Ian D. Thomas, A new glacial isostatic adjustment model for Antarctica: calibrated and tested using observations of relative sea-level change and present-day uplift rates, <i>Geophysical Journal International</i> , Volume 190, Issue 3, September 2012, Pages 1464–1482, https://doi.org/10.1111/j.1365-246X.2012.05557.x
	ICE-5G	Peltier, W. R. (2004). "Global Glacial Isostasy And The Surface Of The Ice-Age Earth: The ICE-5G (VM2) Model and GRACE." <i>Annual Review of Earth and Planetary Sciences</i> 32 (1): 111-149.
	ICE-6G	Peltier, W. R., Argus, D. F., & Drummond, R. (2015). Space geodesy constrains ice age terminal deglaciation: The global ICE-6G_C (VM5a) model. <i>Journal of Geophysical Research: Solid Earth</i> , <i>120</i> (1), 450-487.
	IJ05	Ivins, E., & James, T. (2005). Antarctic glacial isostatic adjustment: A new assessment. <i>Antarctic Science</i> , <i>17</i> (4), 541-553. doi:10.1017/S095410200500296)
	IJ05_R2	Ivins, E. R., T. S. James, J. Wahr, E. J.O. Schrama, F.W. Landerer & K. M. Simon, Antarctic contribution to sea-level rise observed by GRACE with improved GIA correction, <i>J. Geophys. Res., B - Solid Earth</i> , 118, 3126-3141, doi:10.1002/jgrb.50208 (2013).
	Paulson07	Paulson, A., Zhong, S. and Wahr, J. (2007), Inference of mantle viscosity from GRACE and relative sea level data. <i>Geophysical Journal International</i> , 171: 497-508. https://doi.org/10.1111/j.1365-246X.2007.03556.x
	Simpson09	Simpson, M. J. R. and Milne, G. A. and Huybrechts, P. and Long, A. J. (2009) 'Calibrating a glaciological model of the Greenland ice sheet from the last glacial maximum to present-day using field observations of relative sea level and ice extent.', <i>Quaternary science reviews.</i> , 28 (17-18). pp. 1631-1657.
	Khan_2016	Khan, S. A., I. Sasgen, M. Bevis, T. v. Dam, J. L. Bamber, J. Wahr, M. Willis, K. H. Kjær, B. Wouters, V. Helm, B. Csatho, K. Fleming, A. A. Bjørk, A. Aschwanden, P. Knudsen and P. K. Munneke (2016). "Geodetic measurements reveal similarities between post-Last Glacial Maximum and present-day mass loss from the Greenland ice sheet." <i>Science Advances</i> 2 (9): e1600931.
	Schrama14	Schrama, E. J. O., Wouters, B., and Rietbroek, R. (2014), A mascon approach to assess ice sheet and glacier mass balances and their uncertainties from GRACE data, <i>J. Geophys. Res. Solid Earth</i> , 119, 6048– 6066, doi: 10.1002/2013JB010923 .
SMB	RACMO 2.3	Van Wessem, J. M., Reijmer, C. H., Morlighem, M., Mouginit, J., Rignot, E., Medley, B., Joughin, I., Wouters, B., Depoorter, M. A., Bamber, J. L., Lenaerts, J. T. M., Van De Berg, W. J., Van Den Broeke, M. R. and Van Meijgaard, E. (2014) "Improved representation of East Antarctic surface mass balance in a regional atmospheric climate model," <i>Journal of Glaciology</i> , Cambridge University Press, 60(222), pp. 761–770.
	MAR 3.5	Fettweis, X., B. Franco, M. Tedesco, J. H. van Angelen, J. T. M. Lenaerts, M. R. van den Broeke and H. Gallée (2013). "Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR." <i>The Cryosphere</i> 7 (2): 469-489.




Figure A1. Mass balance time-series from the aggregated altimetry, gravimetry and input-output method over the a) WAIS, b) EAIS, c) APIS, and d) GrIS. The vertical dashed lines mark the overlap period of the three time-series. The aggregated time-series and corresponding uncertainties are calculated following the methods described in Section 3 (ii).

Table A2. Rates of mass change (in Gt yr⁻¹) over the gravimetry record (2002 to 2020) from our reconciled estimate and from a modified version of our reconciled estimate in which the contribution of the peripheral glaciers has been removed from the gravimetry estimates following the method described in Section 3.

	Reconciled assessment	Modified reconciled assessment
GrIS	-235.6 ± 20.6	-226.0 ± 20.6
APIS	-18.3 ± 6.0	-15.7 ± 5.8
EAIS	6.1 ± 19.7	6.2 ± 19.6
WAIS	-104.8 ± 11.2	-103.6 ± 10.8
 AIS	-117.0 ± 23.5	-113.1 ± 23.2