

Interdecadal glacier inventories in the Karakoram since the 1990s

Fuming Xie^{1,2}, Shiyin Liu^{1,2,3*}, Yongpeng Gao^{1,2}, Yu Zhu^{1,2}, Tobias Bolch⁴, Andreas Kääb⁵, Shimei Duan^{1,2}, Wenfei Miao^{1,2}, Jianfang Kang^{6,7}, Yaonan Zhang^{6,7}, Xiran Pan^{1,2}, Caixia Qin^{1,2}, Kunpeng Wu^{1,2}, Miaomiao Qi^{1,2}, Xianhe Zhang^{1,2}, Ying Yi^{1,2}, Fengze Han^{1,2},
5 Xiaojun Yao⁸, Qiao Liu⁹, Xin Wang¹⁰, Zongli Jiang¹⁰, Donghui Shangguan³, Yong Zhang¹⁰, Richard Grünwald², Muhammad Adnan^{1,2}, Jyoti Karki^{1,2}, Muhammad Saifullah¹¹

¹ Yunnan Key Laboratory of International Rivers and Transboundary Eco-Security, Yunnan University, Kunming 650500, China

² Institute of International Rivers and Eco-security, Yunnan University, Kunming, Yunnan 650500, China

10 ³ State Key Laboratory of Cryospheric Sciences, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

⁴ School of Geography and Sustainable Development, University of St Andrews, St Andrews KY19 9AL, Scotland, United Kingdom

⁵ Department of Geosciences, University of Oslo, Oslo, 0316, Norway

⁶ Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

15 ⁷ National Cryosphere Desert Data Center, Lanzhou 730000, China

⁸ College of Geography and Environmental Sciences, Northwest Normal University, Lanzhou 730070, China

⁹ Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu 610041, China

¹⁰ School of Resource Environment and Safety Engineering, Hunan University of Science and Technology, Xiangtan, China

¹¹ Department of Agricultural Engineering, Muhammad Nawaz Shareef University of Agriculture, Multan, Pakistan

20 *Correspondence: Shiyin Liu (shiyin.liu@ynu.edu.cn)

Abstract:

Multi-temporal glacier inventories provide key information about the glaciers, their characteristics and changes and are inevitable for glacier modelling and investigating geodetic mass changes. However, to date, no consistent multi-tempo glacier inventory for the whole of the Karakoram exists, negatively affecting the monitoring of spatiotemporal variations of glaciers' geometric parameters and their related applications. We used a semi-automatic method combining automatic segmentation and manual correction and produced multi-temporal Karakoram glacier inventories (KGI) compiled from Landsat TM/ETM+/OLI images for the 1990s, 2000s, 2010s, and 2020s. Our assessments using independent multiple digitization of 37 glaciers show that the KGI is sufficiently accurate, with an overall uncertainty of $\pm 3.68\%$. We also performed uncertainty evaluation for the contiguous glacier polygons using a buffer of half a pixel, which resulted in an average mapping uncertainty of $\pm 5.21\%$. We calculated more than 20 attributes for each glacier, including coordinates, area, supraglacial debris area, date information, and topographic parameters derived from the ASTER GDEM. According to the KGI-2020, approximately 10500 alpine glaciers ($> 0.01 \text{ km}^2$ each) cover an area of $22510 \pm 828 \text{ km}^2$ of which $10.18 \pm 0.38\%$ ($2290 \pm 84 \text{ km}^2$) are covered by supraglacial debris. Over the past three decades, the glaciers experienced a loss of clean ice/snow area but a gain in supraglacial debris. Supraglacial debris cover has increased by $17.63 \pm 1.44\%$ ($343.30 \pm 27.95 \text{ km}^2$) while non-debris-covered glaciers decreased by $1.56 \pm 0.24\%$ ($319.85 \pm 49.92 \text{ km}^2$). The total glacier area was relatively stable and showed only a slight insignificant increase of $23.45 \pm 28.85 \text{ km}^2$ ($0.10 \pm 0.13\%$). The glacier area has declined by $3.27 \pm 0.24\%$ in the eastern Karakoram while the glacier area slightly increased in central ($0.65 \pm 0.10\%$) and western Karakoram ($1.26 \pm 0.11\%$). Supraglacial debris has increased over whole Karakoram, especially in areas above 4200 m a.s.l., showing an upward shift. The glacier area changes were characterized by strong spatial heterogeneity, influenced by surging and

40 advancing glaciers. However, due to global warming, the glaciers are on average retreating. This is in particular true for small and
debris-free glaciers. The multi-temporal KGI data are available at the National Cryosphere Desert Data Center of China:
<https://doi.org/10.12072/ncdc.glacier.db2386.2022> (Xie et al., 2022a).

1 Introduction

The Karakoram mountains are centred in western Tibetan Plateau (see Fig. 1a and 1b) and host more than 20000 km² of glaciers,
45 making this region one of the most glacierized areas outside of the polar regions. As the main source of the rivers' natural flow,
meltwater from the Karakoram glaciers has a significant influence on the socio-economic development in the upper Indus River
basin and the Tarim River basin areas to the north of the mountains (Winiger et al., 2005; Liu et al., 2006; Immerzeel et al., 2020;
Liu et al., 2020; Nie et al., 2021). Due to global warming in recent decades, changes in glacier-related resources and hazards present
50 threats to the safety of critical infrastructure and the sustainable livelihoods of local communities living within the basin (Immerzeel
et al., 2020; Bazai et al., 2021; Gao et al., 2021). In the past few decades, remote sensing-based mass balance, glacier flow, and
surge or advance characteristics indicate that the glaciers in Karakoram are exhibiting anomalous behaviour (Hewitt, 2005; Gardelle
et al., 2012; Kääb et al., 2012; Bolch et al., 2017; Dehecq et al., 2018; Farinotti et al., 2020; Wu et al., 2020; Wu et al., 2021).

Glacier inventories provide fundamental baseline information about the glaciers and are needed for many applications (e.g.
glacier mass balance, glacier modelling). Generally, a glacier inventory contains an identifier (ID), name (if available), location (i.e.,
55 coordinates), size, and other relevant information concerning each glacier (Paul et al., 2009; Paul, 2017). Since the 1970s, thanks to
the development of remote sensing and GIS technology, a series of international glacier inventories such as the World Glacier
Inventory (WGI), the Randolph Glacier Inventory (RGI) or glacier inventories submitted to GLIMS (Global Land Ice Measurements
from Space) have been developed through the efforts of glaciologists from different countries (Müller and Scherler, 1980; Shih et
al., 1980; Paul et al., 2002; Pfeffer et al., 2014). To date, several glacier inventories exist in the Karakoram area, some of which
60 include information about supraglacial debris cover. These inventories include the Glacier Area Mapping for Discharge from the
Asian Mountains (GAMDAM) glacier inventory (GGI15) (Nuimura et al., 2015) and its updated version (GGI18) (Sakai, 2019),
the second Chinese glacier inventory (SCGI) (Liu et al., 2015; Guo et al., 2015), the International Centre for Integrated Mountain
Development (ICIMOD) glacier inventory (Bajracharya and Shrestha, 2011), and the Glaciers_cci (CCI) glacier inventory (Mölg et
al., 2018). However, due to differences and deficiencies in the remotely-sensed imagery, variable data years, and glacier outlines
65 extraction methods used in previous glacier inventories, their results have poor comparability, thus failing to meet the requirements
of glacier change analysis. Moreover, the presented areas of glacier coverage differ partially substantially for the different available
inventories (Bolch, 2019; Bolch et al., 2019; Bhambri et al., 2022).

The two versions of the GAMDAM inventory (GGI15 and GGI18) were generated by manual delineation with the exclusion
of glacierised areas steeper than 40°. These inventories have a difference in an area of 9430 km². The differences can be mainly
70 attributed to the omission of glacier areas above the bergschrund (Sakai, 2019). Moreover, the first one used Landsat images acquired
around the year 2000 and the second one used also scenes spanning 1990–2010 in case the scenes used in the first version were
unsuitable (Sakai, 2019). Whereas the ICIMOD inventory was compiled from Landsat TM/ETM+ images acquired between 2005
and 2009, the SCGI inventory only includes part of the Karakoram mountains and uses the Landsat TM/ETM+ satellite images
between 2007 and 2011. The CCI inventory differs somewhat because the clean ice/snow parts are identified automatically using
75 the image ratio technique and is based upon the Landsat images from 1998 to 2002. The debris-covered areas were manually mapped
with the support of coherence images generated using ALOS-1 PALSAR-1 scenes from 2007 and 2009 (Mölg et al., 2018). In the
RGI 6.0 (Rgiconsortium, 2017), the glaciers for the Karakoram region were extracted from the CCI inventory. A further approach
to constraining changes in Karakoram glaciers was presented by Rankl et al. (2014), who focused only on surge-type glaciers and

large glaciers with lengths > 3 km and areas > 0.15 km². Few studies present glacier area changes for smaller parts of the Karakoram (e.g. Minora et al. (2016) for the Central Karakoram National Park for 2001-2010 or Bhambri et al. (2013) for the Shyok Valley for 1973 to 2011). Both found overall no significant net are changes but large variability due to the presence of surging glaciers. However, consistent multi-temporal homogeneous glacier inventories covering the whole Karakoram mountains do not exist, thereby negatively affecting monitoring of spatiotemporal variations in the glaciers' geometric parameters and hampering explanation of their anomalous behaviour discussed by both international scientists and policymakers.

In most parts of the Karakoram, debris-covered glaciers are dominating (~10% of glacier area covered by debris). Thus, if this debris cover and its evolution are neglected, hydrological models may underestimate the longevity of glacier-sourced water resources and eventually overestimate the effects of glacier melt on sea-level rise (Herreid and Pellicciotti, 2020; Zhang and Liu, 2021). Many studies have reported that the supraglacial debris cover is expanding in different parts of the world (Mölg et al., 2019; Tielidze et al., 2020; Xie et al., 2020b). Herreid et al. (2017) report no net change of the debris-covered area in Central Karakoram from 1977 to 2014. However, variations in the debris cover across the whole Karakoram region remains unknown. Mapping glaciers with debris cover is still challenging (Paul et al., 2004; Pfeffer et al., 2014; Farinotti et al., 2020), with various data and approaches having been proposed to address this issue, including using SAR (Synthetic Aperture Radar) coherence maps, optical bands, thermal infrared features, terrain parameters, machine learning and other methods (Bhambri et al., 2011; Mölg et al., 2018; Alifu et al., 2020; Xie et al., 2020a).

The present study aims to present precise and homogeneous multi-temporal glacier inventories for the whole Karakoram. We developed a high-performance cloud-removal and image-compositing algorithm to produce high-quality images for the glacierised regions of Karakoram for the extraction of glacier outlines. These will provide important baseline data for glacier research and related applications.




Fig. 1 Overview of the study region (Karakoram Mountains). a) Karakoram Mountains physiography. b) General climatology trends in Central Asia. The orange and blue arrow lines represent the westerly and monsoon directions, respectively (Yao et al., 2012). The Karakoram boundary is a reasonable revised boundary with reference to Bhambri et al. (2017) and can be accessed freely via “<https://github.com/1923xfmingynu/Subdivision-Of-High-mountain-Asia>”. c) Worldwide reference system (WRS) Path/Row values of Landsat images used in this study. The blue area represents the glaciated area. d) Number of Landsat observations (COUNT) in each pixel, the details in the lower-left corner show the filtered image collection information: Landsat sensor name (TM/ETM+/OLI) and image acquisition time range,

including year range, day of year (DOY) and cloud cover threshold. Due to the poor quality of satellite imagery in some areas (red rectangle) in 1990s, two scenes from 1993 and 1994 are used in the image composite.

2 Data and methods

2.1 Datasets

110 This subsection lists all data sets covering the Karakoram that are used to produce and assist in the analysis of the multi-temporal glacier inventory, including optical images from different satellite sensor, digital elevation model (DEM), four previous glacier inventories, three supraglacial debris extents, two surge-type glacier inventories, two modelled ice thickness data, hydrological basins and river networks. Table 1 summarizes their key characteristics, presenting their sources, date and access link.

115 At least 12 Landsat images are required to cover the glacierised area of the Karakoram (Fig. 1c). We selected Level 1 terrain-corrected products of the Landsat TM/ETM+/OLI scenes (L1TP) images with 30 m spatial resolution representing the years 1990~2020, including 65 TM, 39 ETM+ and 82 OLI images (Table S1). The satellite images were identified and processed using Google Earth Engine (GEE). An Asia-North-Albers-Equal-Area-Conic projection coordinate system with the central meridian 76° and the standard parallels 33° and 38° was used for all spatial data in this study. Firstly, the available images were initially filtered from the Landsat image collection by selecting the melting season (days of year: 200-270), region of interest (Karakoram boundary), and cloud cover (< 25~30%). The filter criteria depend on the data quality and observations recorded in different periods as shown in the lower-left corner of Fig. 1d. We set the image interval to two years to ensure representativeness and reliability of the composite image as much as possible, a value smaller than the interval of three years deemed accepted in previous report (Paul et al., 2013). For the period of the 1990s, suitable images within one or two years were not sufficient to cover the whole Karakoram area, thus, two images from 1993 and 1994 were also selected for areas with no observations or unsuitable cloud conditions (see red box in Fig. 1d); this resulted in a more than three years interval in these areas, thus may increasing the inter-annual variation error of the glacier outlines. Then, we applied a GEE's own cloud-removal algorithm (*ee.Algorithms.Landsat.simpleCloudScore*) (Gorelick et al., 2017) to all the images and composited an image needed for glacier inventory. In the image compositing, if more than 10 observations were reached at a given point, only the observations from the first 10 cloudiness images were accepted to generate an image composite in order to reduce the data source complexity of the composite image.

130 To assess the uncertainty of our new glacier inventory data, we evaluated the glacier outlines derived from coarser spatial resolution satellite imagery using the glacier outlines identified from high resolution images or mean value of multiple digitisations from lower-resolution data. In an ideal case the outlines should be digitized several times (multi-manual digitization (MMD)) to consider the variability of the analysts ("round robin") (Paul et al., 2013) (See Sect. 2.6). Therefore, several Sentinel-2 10m scenes and a Planet 3m image are used.

135 Nuimura et al. (2015) used the ICESat GLA 14 data to evaluate DEM accuracy data in glaciers of high mountain Asia and found that the ASTER GDEM version 2 (v2) is more suitable in comparison to the SRTM DEM. Compared to ASTER GDEM v2, the updated v3 product has less voids areas due to an increased number of ASTER stereo image data and further developed processing that improves the vertical and horizontal accuracy of the data (Abrams, 2016). Therefore, topographic information (eg., slope, aspect and median elevation) for all glaciers and generation of ridgelines (see Sect. 3.3) in this study was obtained from ASTER GDEM v3 data. All ASTER GDEM v3 tiles were downloaded from <https://e4ftl01.cr.usgs.gov/ASTT/ASTGTM.003/>, and mosaicked in local Python GADL environment.

145 In the preliminary extraction of glacier outlines (see Sect. 2.2), we relied on the previous glacier boundaries. And we qualitatively compared area differences between different glacier inventories, supraglacial debris extents, as well as illuminated the variations of surge-type glaciers. Therefore, four previous glacier inventories, i.e., GGI18, CCI, SCGI and ICIMOD, three supraglacial debris extents (Mölg et al., 2018; Scherler et al., 2018; Herreid and Pellicciotti, 2020) and two surge-type glacier

inventories (Bhambri et al., 2017; Guillet et al., 2022) were used in this study without any modifications.

Furthermore, in the results and discussion, we analysed glacier changes in different sub-basins and highlighted the value of multi-temporal glacier inventories in ice volume calculation. Hydrological basins provided by the HydroSHEDS project (Lehner and Grill, 2013) and two published simulated ice thickness data (Farinotti et al., 2019; Millan et al., 2022a) were also used.

Table 1 Lists of data sets covering the Karakoram mountain that are used in this study.

Data Name	Sources	Date	Access		
Satellite images	Landsat TM, ETM and OLI +30-m/ 15-m images	1990, 1991, 1993, 1994; 2000, 2001; 2009, 2010; 2018, 2019, 2020 (details see Table S1)	GEE asset or https://earthexplorer.usgs.gov/		
	Sentinel-2 10-m images	2020-08-25, 2020-08-23	GEE asset or https://scihub.copernicus.eu/		
	Planet 3-m images	2019-05-29	Ordered and download via Planet's APIs		
DEM	30-m ASTER GDEM V3	2000-2013	https://e4ftl01.cr.usgs.gov/ASTT/ASTGTM.003		
Previous glacier Inventory	Glacier Inventories: (1) GGI18 (Sakai, 2019); (2) ICIMOD (Bajracharya and Shrestha, 2011); (3) SCGI (Guo et al., 2015); (4) CCI (Mölg et al., 2018)	(1) 1990-2010; (2) 2005-2009; (3) 2007-2011; (4) 1998-2002, 2007-2009	(1) https://doi.org/10.1594/PANGAEA.891422 (2) https://www.icimod.org/ (3) (4) https://doi.org/10.12072/ncdc.Westdc.db0006.2020 (4) https://doi.org/10.1594/PANGAEA.894707		
		Supraglacial debris extents: (5) from CCI inventory (Mölg et al., 2018); from global supraglacial debris-cover extents produced by (6) Scherler et al. (2018) and (7) Herreid and Pellicciotti (2020)	(5) 2007-2009; (6) 2013-2017; (7) 1986-2016 (median 2013)	(5) same as (4) (6) http://doi.org/10.5880/GFZ.3.3.2018.005 (7) https://doi.org/10.5281/zenodo.3866466	
		Surge-type glaciers inventory: from the data published by (8) Bhambri et al. (2017) and (9) Guillet et al. (2022)	(8) 1840s-2015 (9) 2000-2018	(8) https://doi.org/10.1038/s41598-017-15473-8 . (9) https://doi.org/10.5281/zenodo.5524861	
		Ice thickness	Ice thickness modelled by (1) Farinotti et al. (2019) and (2) Millan et al. (2022a)	-	(1) https://doi.org/10.3929/ethz-b-000315707 (2) https://doi.org/10.6096/1007
		Catchment boundaries and river networks	HydroSHEDS database (Lehner and Grill, 2013) produced based on SRTM DEM	~ 2000	https://www.hydrosheds.org/

2.2 Extraction of glacier outlines

Accurate glacier outline identification can be challenging, because glaciers exhibit a wide range of surface characteristics, they can be interpreted differently by the cartographer, surveyor, or remote sensing specialist (Paul, 2017; Paul et al., 2009). Moreover, poor quality satellite imagery (i.e., scenes with clouds, large shadows, or seasonal snow cover), debris-covered glaciers, rock glaciers, and perennial snow-covered depressions (potential presence of some ice underneath) are all sources of uncertainty in glacier outline mapping that may negatively affect the accuracy of the resulting outlines. Therefore, we used a semi-automatic method (band ratio + manual correction) to compile glacier inventories. This approach is widely applied (Bolch et al., 2010a; Guo et al., 2015; Paul et al., 2017).

Firstly, we calculated the NDSI $(\rho_{green} - \rho_{swir1}) / (\rho_{green} + \rho_{swir1})$ and initially considered areas with $NDSI \geq 0.4$ as clean ice/snow (or debris-free areas) (Dozier, 1989; Xie et al., 2020b) while other areas were labelled as initial debris-covered ice, followed manual correction (see next section). As previous studies have found, optimized thresholds for different periods can lead to overestimation of glacier area (Xie et al., 2020b); additionally, the glacier area has insensitivity to different thresholds when using high-quality satellite images on gentler terrain (Guo et al., 2015). Accordingly, we adopted the empirical NDSI threshold uniformly across the whole Karakoram for mapping clean ice/snow. A similar threshold was also used for generating glacier inventories for large regions elsewhere (e.g. Ke et al. (2016)). Second, since many pixels outside of the glacier extent are considered to be

170 supraglacial debris in the initial debris-covered data, glacier outlines from previous glacier inventories were used as a mask to
eliminate them, as suggested in similar studies (Bolch et al., 2010a; Scherler et al., 2018; Baumann et al., 2020). We combined two
earlier glacier inventories (90% CCI + 10% GGI18, to fully cover Karakoram glaciers) as the mask layer for KGI-1990s, and the
subsequent KGI-2000s, KGI-2010s, and KGI-2020s rely on the earlier revision of the glacier inventory (KGI-1990s, KGI-2000s
and KGI-2010s in that order). This approach is suitable with retreating glaciers but cuts off advancing glaciers. These were corrected
by manual post-processing. Moreover, to remove the “salt-and-pepper” effect in the initial binary map, we applied a morphological
filter with a 5×5 square kernel as described in Xie et al. (2020b). Then, the binary map was converted into vector format, retaining
175 a ‘Type’ attribute describing whether each polygon is debris-covered ice or clean ice/snow, followed by the manual correction.

2.3 Manual correction

There are five typical error types in initial glacier outlines that require revision, including errors due to glacier omission (Fig.
2a), clouds or shadow (Fig. 2b), glacier commission (Fig. 2c), glacier advance or surge (Fig. 2g), and errors due to debris and ponds.
To eliminate these errors, we have established detailed criteria and applied manual correction of multi-temporal glacier outlines
180 using Landsat SWIR-NIR-RED or NIR-RED-GRE (false) colour composites of images (30 m resolution). These were combined
with other optional baseline data, such as the 15 m-resolution panchromatic Landsat band, the land surface temperature (LST) map
derived from Landsat thermal infrared band (Kraaijenbrink et al., 2017; Liao et al., 2020), contour lines derived from ASTER GDEM
and Google Earth images. Most of the debris-covered ice sections were remapped using the panchromatic band, LST, topography,
and Google Earth images, while the ice tongues of debris-covered glaciers were identified by the location of their meltwater outlets
185 (see Fig. 2 (e) and (f)). The emergence of ice cliffs and ponds on glacier surfaces was also used in our present glacier inventory to
identify the site of debris-covered glaciers and classify them as part of the debris-covered ice. Paul et al. (2013) observed that there
are large uncertainties in manually identified glacier outlines, due to the influence of seasonal snow cover, shadows and complex
underlying glacier surface types, such as supraglacial debris cover, rock glaciers, stagnant ice, resulting in large deviations of the
glacier extent even by different experienced interpreters. This issue is especially prevalent for glacier tongues with thick debris,
190 which completely depends on visual interpretation to map the glacier, thus leading to further uncertainties. Therefore, to reduce the
artificial interferences caused by subjectivity, all the correction work in this study was performed by the main author only.




Fig. 2 Types of manual correction. **a)** Glaciers missed in previous inventories. **b)** Areas contaminated by hill shadows or cloud shadows. **c)** Glaciers that have retreated and require updating. **d)** Comparison of the revised glacier outlines with those from the CCI and GGI 18 glacier inventories, with a base map of Landsat 7 ETM+ composited in the 2000s. **e)** and **f)** are auxiliary methods for mapping debris-covered glaciers. The location of the water outlet is used to determine the position of the glacier tongue, and the land surface temperature images are used to distinguish between debris-covered glaciers and exposed bedrock. **g)** Glacier terminus advance may be caused by glacier surges (red rectangle). Except for subfigure d, the yellow and white lines represent the glaciers identified by the automatic threshold segmentation method as debris-covered and debris-free parts, respectively.

2.4 Ice divides

Segmentation of complex glacier polygons, such as glaciers connected in their accumulation areas, can be achieved by using ridgelines derived from a digital elevation model (DEM) (Mölg et al., 2018). Thus, we adopted the method proposed by Bolch et al. (2010a), using the ASTER GDEM v3 dataset. First, we merged the four periods of glacier outlines into a new version with a 1 km buffer. This was then used to clip the sink-filled DEM. A flow direction grid was then calculated to generate a raw drainage basin using watershed analysis (Fig. 3). Then manual corrections were performed to delete small basins ($<0.01\text{km}^2$), based on a three-dimensional topographic map derived from the DEM datasets and a hill-shadow (see the base map of Fig.3). The ridgelines were then extracted to split glacier complexes into single glaciers. After identifying the ice divides, we performed some batch processing on the revised glacier outlines, including filling voids (eliminating contained parts only) with an area of less than 4000 m^2 (i.e., less than 5 pixels), deleting small supraglacial debris patches ($<4000\text{ m}^2$) and eliminating glaciers smaller than 0.01km^2 . This resulted in glacier area changes of only 1.63 to max. 6.97 km^2 in the different periods.




Fig. 3 Glacier drainage basins derived from ASTER GDEM v3 and ridgelines extracted after manual correction.

2.5 Attribute data

We included more than 20 attributes for each analysed glacier. In addition to the GLIMS ID linked from the RGI inventory, each glacier was given a unique KGI ID according to the GLIMS ID encoding style that is automatically calculated via Python programming using the centroid latitude and longitude of the glacier polygon. For terrain attributes, the minimum, median, and maximum elevation, plus median slope and aspect were calculated from the ASTER GDEM v3. Since in particular large glaciers (> 5 km²) may consist of different branches and tributary glaciers with different aspects or glaciers have large bends, the resulting representativeness of the average aspect may be poor; we thus also appended an attribute of a manually identified aspect according to the approximate orientation of each glacier. In addition, other information, such as longitude, latitude, glacier area, debris area, name, date, glacier surface type (clean ice or debris), terminus status (whether advancing), as well as the name of the operator were included. For the corresponding date of all inventoried glaciers, we provide not only the year range of the composited image dates but also calculate the probability that the edge pixels of the glaciers originate from a certain date. This is on the premise that the interannual difference of pixels inside the glacier does not affect the accuracy of glacier extent. However, this assumption may affect the supraglacial debris cover. As shown in Fig. S1, we generated a date image to track the source image for each pixel in the composited image; this indicated that the composite images had little interannual variation for most glaciers in the inventory as the pixels in most composite scenes were derived from data from the same year.

2.6 Estimation of the uncertainty

Potential error-sources of glacier inventory include errors from satellite data quality (i.e., due to seasonal snow cover), DEMs, methods of glacier delineation, complex underlying glacier surface types, and quality of ridgelines (for single glaciers). In general, we applied two different uncertainty assessment methods. In the “round robin” method, seven experienced participants repeatedly digitized 37 glaciers between 0.10 km² and 100 km² in size, located in the Hunza valley, western Karakoram based on high-resolution images (Sentinel-2 and Planet) and lower-resolution images (Landsat 8), the spatial locations are shown in Fig. S2. The standard deviation (STD) of the digitized glacier area (MMD results) values is used to evaluate the precision of the analysts’ digitisation and the difference between its mean area value (as the “truth” of glacier area) and the area calculated from the glacier inventory are used to determine which glacier extent has lower uncertainty, i.e., the KGI or the MMD outlines. Results of MMD are shown in Table S2. The digitization on the Landsat OLI scene had STD of 1.34-25.93% (mean 2.29%), with the largest value for the small glacier

and partly debris-covered glaciers (Fig. 4), while the STD of on Sentinel-2 scene was smaller with a mean of 1.28% (from 0.48% to 7.39%). Apart from errors at debris-covered tongues or margins, interpretation differences in the accumulation area are considered a key error source, including differences in interpretation of snow patches and glacier extents (Fig. 4). Additionally, scatter plots (Fig. S3) of the linear regression for both the MMD and KGI-2020s glacier area showed a strong positive correlation between them, with a correlation coefficient (r) value of 1 and a root mean square error ($rmse$) of 0.78 km², thus confirming the high accuracy of the KGI data. The mean area difference of KGI outlines and the MMD mean values based on the Landsat image is -6.92%; 5.58% based on Sentinel-2 images. The mean area difference between the KGI outlines and the high-resolution image with only one digitization is -6.39%(Sentinel-2) and 2.56%(Planet). The area difference for the different glaciers varied from to -124.73% to 35.24% (-1.90 ~ 2.61 km²), influenced by glacier size, presence of tributary glaciers, and whether glaciers were debris-covered or debris free. The resultant total mapping uncertainty is $\pm 3.68\%$, which meets the requirements of the Global Observing System for Climate (Gcos, 2016). The results presented above are consistent with previous accuracy assessments that reported the mean area differences between the automatically derived and manually digitized outlines on high-resolution images between 2% and 5% (Andreassen et al., 2008; Paul et al., 2013; Paul et al., 2017).




Fig.4 Glacier extent digitized by different participants based on Sentinel-2 image (a, b), Landsat image (c) and Planet image (d) with KGI-2020s derived from Landsat 30-m imagery.

As a second measure of uncertainty, we applied the buffer method (Bolch et al., 2010a; Granshaw and G. Fountain, 2006) on the contiguous glacier polygons. Accordingly, a buffer of $\pm 1/2$ pixel (i.e., 15 m) for the KGI outlines (Fig. 5a) were generated and the difference between the area of the buffer and the KGI was used as the uncertainty measure. The uncertainty for the four periods KGI data are $\pm 5.31\%$, $\pm 5.18\%$, $\pm 5.12\%$ and $\pm 5.21\%$, with an average of $\pm 5.21\%$. In terms of the debris-covered areas, generally, a buffer of ± 1 or 2 pixels (30 or 60 m) buffer was suggested in previous research (Mölg et al., 2018; Paul et al., 2020). The uncertainty of the debris portion in this study was evaluated through the ratio of the glacier area to the debris cover area multiplied by the uncertainty of ± 1 pixel (30m) buffer (Fig. 5b), resulting in uncertainty of $\pm 27.89\%$, $\pm 29.39\%$, $\pm 28.20\%$, and $\pm 29.76\%$ for

the four periods, with a mean value of $\pm 28.81\%$.

For the whole glacier, the mapping uncertainty based on the “round robin” experiment is within the estimation range based on the buffer method, indicating that the ‘round robin’ value can be used as a reasonable estimation of the uncertainty. Hence, we used this value ($\sigma = \pm 3.68\%$) as the uncertainty value for all KGI data in this study. The area change uncertainty (σ_{Δ}) was estimated according to the standard error propagation, as root sum square of the uncertainty for outlines mapped from different periods, but only consider the glacier parts which showed change in the 1990s and 2020s (ΔA_{1990s} and ΔA_{2020s}) (Bhambri et al., 2011; Zhang et al., 2018; Li et al., 2022), calculated as : $\sigma_{\Delta} = \sqrt{(\Delta A_{1990s} * \sigma)^2 + (\Delta A_{2020s} * \sigma)^2}$.




Fig. 5. Overlaying of a 15m buffer (a) from the KGI-2020s glacier extent and a 30m buffer (b) from the supraglacial debris extent with MMD outlines on the base map of Sentinel-2 and Planet image.

3 Results

3.1 Characteristics and status of the Karakoram glaciers

We mapped a total of 10,498 glaciers with an area of $22510.73 \pm 828.39 \text{ km}^2$ (Table 2). All glaciers are divided into 9 areal classes of $\leq 0.05 \text{ km}^2$ to $\geq 100 \text{ km}^2$. 7941 (75.60% of all glaciers) are smaller than 1 km^2 , but cover only a total area of $2098.45 \pm 77.22 \text{ km}^2$ (9.30% of the total). 26 glaciers exceed 100 km^2 which covers $7430.64 \pm 273.45 \text{ km}^2$ or 32.5% of the total glacier area in Karakoram. Siachen ($1118.09 \pm 41.14 \text{ km}^2$), Baltoro ($849.44 \pm 31.26 \text{ km}^2$), Biafo ($579.25 \pm 21.32 \text{ km}^2$) and Hispar ($542.07 \pm 19.95 \text{ km}^2$) glaciers are larger than 500 km^2 . 20% (2175) of the glaciers with an area within size classes $1\sim 5 \text{ km}^2$ and $10\sim 50 \text{ km}^2$ cover a total area of $8519.63 \pm 313.52 \text{ km}^2$. Glaciers with size classes $5\sim 10 \text{ km}^2$ and $50\sim 100 \text{ km}^2$ are roughly equal in area, each class covering about $2,200 \text{ km}^2$. 581 glaciers, ranging from 5 to 100 km^2 cover $8736.78 \pm 321.51 \text{ km}^2$, meaning that all these glaciers plus the 26 glaciers exceeding 100 km^2 occupy 72% of the total area of the Karakoram glaciers.

We divided the Karakoram into northern and southern slopes (NK and SK) by ridgelines, and into eastern, central, and western parts (EK, CK and WK) using the Indus and Tarim sub-basin outlines (Fig.6). 55.3% of the glaciers are concentrated in CK, covering an area of $12547.12 \pm 461.73 \text{ km}^2$, followed by the glaciers in the WK. Less than one-fifth, i.e., $3954.30 \pm 145.52 \text{ km}^2$ or 17.60 \pm 0.89% of the total glaciers are located in the EK. A high proportion of debris cover is a typical feature of Karakoram glaciers, with a debris-cover percentage of $2.63 \pm 0.10\%$ ($104.94 \pm 3.83 \text{ km}^2$) in the eastern and $11.11 \pm 0.41\%$ ($1382.68 \pm 50.88 \text{ km}^2$) and $13.16 \pm 0.48\%$ ($804.32 \pm 29.60 \text{ km}^2$) in the central and western regions, respectively. Additionally, the glacier areas in NK and SK differ by about 20%, $12321.74 \pm 453.44 \text{ km}^2$ and $10188.99 \pm 374.95 \text{ km}^2$. However, the supraglacial debris cover in SK ($1500.85 \pm 55.23 \text{ km}^2$, $65.51 \pm 2.41\%$) is around twice than recorded in NK ($790.09 \pm 29.08 \text{ km}^2$).

Moreover, we divide the Karakoram mountains into five sub-basins: Gilgit-Hunza, Shigar, Shyok, Wakhan and sub-Tarim (Fig. 6). Among these, the Shyok sub-basin housed the largest extent of glaciers ($7918.54 \pm 291.40 \text{ km}^2$), followed by the Tarim sub-

basin ($5574.06 \pm 205.13 \text{ km}^2$), then Gilgit-Hunza ($4974.34 \pm 183.06 \text{ km}^2$), and Shigar ($3481.02 \pm 128.10 \text{ km}^2$). The percentage of debris cover is greatest in the Shigar (16.64%) and Gilgit-Hunza (14.72%) basins, while sub-Tarim basin has the lowest debris coverage (5.87%).




Fig. 6. The Karakoram Mountains were divided into western, central and eastern Karakoram (WK, CK and EK) according to the Indus and Tarim sub-basins, and into Northern and Southern Karakoram based on the main central ridgeline. The major sub-basin divisions in the Karakoram mountains also are shown. Hollow circles represent the glacier areas, as aggregated on a 20 km × 20 km grid.

3.2 Supraglacial debris cover of Karakoram glaciers

Supraglacial debris cover is widespread in the Karakoram. 1848 debris-covered glaciers (17.63% of all glaciers) with a total area of $2290.95 \pm 84.31 \text{ km}^2$ of supraglacial debris area ($\sim 10.18 \pm 0.38\%$ of total glacier area) were mapped in the KGI-2020s. The total area of debris-covered glaciers is more than 16800 km^2 , of which 74.4% of the glaciers (1374) are larger than 1 km^2 . The supraglacial debris area of glaciers smaller than 1 km^2 is only $49.73 \pm 1.83 \text{ km}^2$, accounting for $2.17 \pm 0.08\%$ of the total debris coverage. Hence, there is no substantial debris cover on these small glaciers. The debris cover area of the 26 glaciers larger than 100 km^2 dominates, representing almost half (47.5%) of the total debris cover area with an area of $1088.83 \pm 40.07 \text{ km}^2$. This is followed by the $10\text{--}50 \text{ km}^2$ size class glaciers, which cover a total area of $413.90 \pm 15.23 \text{ km}^2$ ($18.10 \pm 0.67\%$). Also, despite the glacier number being on the same order of magnitude in size classes $5\text{--}10 \text{ km}^2$ and $50\text{--}100 \text{ km}^2$, the latter's debris coverage ($321.21 \pm 11.82 \text{ km}^2$) is twice than of the former ($170.76 \pm 6.28 \text{ km}^2$), indicating that the larger the glacier, the greater the likelihood and severity of its surface debris cover.

From 1990 to 2020, the debris cover has increased by $17.63 \pm 1.44\%$ ($343.30 \pm 27.95 \text{ km}^2$), compared to a $1.56 \pm 0.24\%$ decrease ($319.85 \pm 49.92 \text{ km}^2$) in the clean ice or snow part. All different size classes show debris cover increases on their surfaces (Table S3). Debris cover area on glaciers $> 100 \text{ km}^2$ and in size classes in $1\text{--}5 \text{ km}^2$ present the most significant changes, increasing by $113.18 \pm 10.87 \text{ km}^2$ ($11.60 \pm 1.11\%$) and $79.12 \pm 5.08 \text{ km}^2$ ($44.16 \pm 2.84\%$) respectively. A similar trend was also found in glaciers smaller than 1 km^2 , in which the debris cover area increased by $19.47 \pm 1.32 \text{ km}^2$ ($64.34 \pm 4.36\%$) in total; however, this accounted reached for only 5.67% of the total recorded debris increase. Except for glaciers smaller than 1 km^2 , the growth rates of debris coverage in size classes between $1\text{--}5 \text{ km}^2$ ($\sim 44.16 \pm 2.84\%$) and $5\text{--}10 \text{ km}^2$ ($32.14 \pm 2.12\%$) are the most significant. Our results

indicate that supraglacial debris cover is expanding on large debris-covered glaciers, while many small and medium-sized debris-free glaciers are transforming into debris-covered glaciers. Over the study period, approximately 500 debris-free glaciers developed into debris-covered glaciers with a decadal growth rate of 12.20% (i.e., about 166 transforms into debris-covered glaciers every decade), more than 97% of which are smaller than 10 km². According to KGI-2020s, among the glaciers larger than 5 km², there are 457 (14379.96 ± 529.18 km²) glaciers with supraglacial debris coverage greater than 0.1 km². The area of these debris-covered glaciers increased by 98.06 ± 13.43 km² (0.69 ± 0.09%) from 1990 to 2020, while that of debris-free glaciers decreased by 12.72 ± 2.64 km² (0.68 ± 0.14%).

Table 2. Number and area of glaciers according to different size classes or surface types during 1990~2020.

Glacier Size (km ²) / Surface type	Glacier Area (GA)								GN (%)	GA (%)	GN Change NC(%)	GA Change AC(%)
	Glacier Number (GN)				km ²							
	1990s	2000s	2010s	2020s	1990s	2000s	2010s	2020s	2020s	(1990 ~ 2020)		
< 0.05	1120	1007	843	1096	40.14 ± 1.48	33.84 ± 1.25	29.53 ± 1.08	35.93 ± 1.32	10.44	0.16	-2.14	-10.49
0.05 ~ 0.1	1723	1659	1622	1654	124.89 ± 4.60	119.98 ± 4.42	117.39 ± 4.32	119.9 ± 4.41	15.76	0.53	-4	-4
0.1 ~ 0.5	3798	3764	3732	3752	936.61 ± 34.47	933.35 ± 34.34	928.24 ± 34.16	924.21 ± 34.01	35.74	4.11	-1.21	-1.32
0.5 ~ 1	1440	1459	1466	1439	1022.48 ± 37.63	1038.84 ± 38.23	1042.74 ± 38.37	1018.41 ± 37.48	13.71	4.52	-0.07	-0.40
1 ~ 5	1951	1945	1971	1950	4213.42 ± 155.05	4223.14 ± 155.41	4279.34 ± 155.47	4214.86 ± 155.11	18.57	18.72	-0.05	0.03
5 ~ 10	319	331	327	326	2199.79 ± 80.95	2294.42 ± 84.43	2271.23 ± 83.58	2231.65 ± 82.12	3.11	9.91	2.19	1.45
10 ~ 50	238	231	234	225	4549.64 ± 167.43	4419.86 ± 162.54	4462.6 ± 164.22	4304.77 ± 158.42	2.14	19.12	-5.46	-5.38
50 ~ 100	30	29	30	30	2259.27 ± 83.14	2149.36 ± 79.10	2223.37 ± 81.82	2200.36 ± 80.97	0.29	9.77	0	-2.61
> 100	24	26	26	26	7141.04 ± 262.79	7364.39 ± 271.01	7450.43 ± 274.18	7430.64 ± 273.44	0.25	33.01	8.33	4.06
Debris-free	9291	8914	8773	8650	20539.63 ± 755.86	20414.01 ± 751.24	20789.49 ± 765.05	20219.78 ± 744.09	82.40	89.82	-6.90	-1.56
Debris-covered*	1074	1199	1229	1374	1947.65 ± 71.67	2160.26 ± 79.50	2015.37 ± 74.17	2290.95 ± 84.31	13.08	10.18	27.93	17.63
Total	10643	10451	10251	10498	22487.28 ± 827.53	22574.27 ± 830.73	22804.86 ± 839.22	22510.73 ± 828.39	100	100	-1.36	0.10

* Glacier Number: number of debris-covered glaciers greater than 1 km²; Glacier Area: total area of all debris-covered ice

3.3 Glacier elevation, slope and aspect

The Karakoram glaciers have a wide range of elevations with a mean altitude of ~ 5300 m a.s.l. and a median altitude of ~5500 m a.s.l. (5200 ~ 6200 m a.s.l.). The hypsometry of the glacier area (Fig. 7a and 7b) indicates that the distribution of glaciers is concentrated at the altitude 5300~5700 m a.s.l., whereas the debris-covered sections are primarily distributed at altitudes of 4000~5000 m a.s.l. and gradually disappear at altitudes above 6,000 m a.s.l. The altitudinal profiles of the glacier surface area (Fig. 7b) indicate that glaciers lose area at low altitude, while near the altitude zone at 5000 ~ 6000 m a.s.l, it shows an increasing trend during 1990-2010, followed a decreasing trend since the 2010s. In terms of glacier orientation (Fig. 7c and 7d), most glaciers have a south, east or southeast aspect (~ 52.40%), with only a few glaciers facing to the north. While there are equal numbers of east and southeast-facing glaciers, the area of the southeast-facing glaciers is twice as large as that of the east-facing glaciers. Similarly, the area of the north-facing glaciers is not as small as that of the west-facing glaciers, which occupy the smallest area. The southeast and north-facing glaciers in the Karakoram mountains are dominated by large glaciers. As depicted in Fig. 7e and 7f, the slope of all glaciers is mainly concentrated between 0 and 50°, with almost no glaciers with slopes greater than 70°, whereas supraglacial debris is widespread in areas with slopes less than 15°. This may be due to broad, flat valley areas providing favourable terrain conditions for the accumulation and enrichment of supraglacial debris. Previous studies have used a slope value of less than 24°

(Paul et al., 2004) or 25° (Xie et al., 2020b) to limit the extent of debris, even though there are steep slopes with debris, which may result from steep ice cliffs or crevasses. Overall, compared to debris-free glaciers, debris-covered glaciers have a broader altitude range (i.e., lower minimum elevation and higher maximum elevation) and lower slopes (Table S4).




Fig.7 Altitudinal profiles of the glacier surface area at 50 m intervals for debris-covered ice (a) and all glaciers (b), showing variations from 1990

to 2020. Plots showing glaciers' number and area by aspect (c and d), in addition to surface slope versus latitude for glaciers and debris-covered sections (e and f).

4 Discussion

4.1 Comparison with previous glacier inventories

In order to determine major differences between previous glacier inventories and our glacier inventory in the Karakoram, we compared it with the SCGI (5114.68 km², glacier area within the KGI range), ICIMOD (11612.82 km²), CCI (20444.40 km²), and GGI18 (20036.55 km²) inventories. However, due to the different approaches, data sources and methods among different glacier inventories, it cannot be compared without a high level of uncertainty, so this is only a qualitative comparison. The Karakoram boundary used by us is different from that in previous studies (Bolch et al., 2012; Bolch et al., 2019; Bhambri et al., 2022), so the qualitative comparison is also only for areas covered by both inventories. Among these, the SCGI and ICIMOD inventory were compared with the KGI-2010s. We then contrasted the GGI18 and CCI inventory with the KGI-2000s to minimize the inter-annual variation between the compared data. For comparison the glacier areas were aggregated onto a 5 km × 5 km grid (Fig. 8). Our results indicate that there is a substantial difference between the ICIMOD inventory and KGI data. In some areas, the glacier area in a single grid cell is underestimated by 15.38 km² (253.40%), with a total *rmse* of 3.78 km². The GGI18 and SCGI inventories exhibited the differences from the KGI data between 1.45~7.11 km² and 1.34~8.60 km², with *rmse* values of 1.41 km² and 1.53 km², respectively. The difference between the CCI inventory and KGI-2000s was found to be the smallest, with *rmse* and *r* values of 0.65 km² and ~1 (0.998) ($p < 0.001$), while the difference in glacier area per grid was between 2.42~3.32 km². The clear underestimation of glacier area in the ICIMOD inventory may relate to the fact that clean ice or snow cover in high altitudes, steep headwalls, and areas affected by hill shadows or clouds were not considered (Bajracharya and Shrestha, 2011) (see also Fig. S4). Similar factors still affect the updated GAMDAM inventory (GGI18), resulting in glacier area values smaller than ours in most regions; in contrast, the difference between the CCI and KGI inventories was relatively small because a similar approach was adopted, with only differences in data source and interpreter (see Fig. 2d). In addition to the above factors, different interpreters' identification of debris-covered glaciers and inter-annual variation between comparative data are potential error sources in all glacier inventories. To sum up, such glacier inventories produced through different methodological standards, data sources, and interpreters contain many uncertainties from different sources, and therefore homogeneous multi-temporal glacier inventories are required to calculate glacier changes. Multi-temporal homogeneous glacier inventories are an important data source for either basic glacier outlies or as a validation set when computing glacier mass changes and associated runoff from projection models. Based on published ice thickness data, we calculated the ice volume in Karakoram is $2.03 \pm 0.52 \times 10^3 \text{ km}^3$ (Farinotti et al., 2019) or $2.81 \pm 1.08 \times 10^3$ (Millan et al., 2022b), which has a potential contribution to sea-level rise of $4.88 \pm 1.27 \text{ mm}$ or $7.11 \pm 3.07 \text{ mm}$. Taking into account different glacier extents in different periods, these projections will produce a variable error of 0.36 ~ 0.49%.

We performed another comparison in regions where both previous data and our inventory mapped supraglacial debris. The results indicate that the debris cover area in the CCI inventory is higher (18.47 km²) than that of KGI-2000s, while the total debris cover mapped by Scherler et al. (2018) is much lower than KGI-2020s (in total, ~18 % less). The latter is a particularly large difference, which may be related to the inter-annual variation (five years), lack of checks or manual correction, and overlooking the terminal changes caused by glacier surges or advances in the results extracted by Scherler et al. (2018). Moreover, the supraglacial debris cover on glaciers larger than 2 km² mapped by Herreid and Pellicciotti (2020) using Landsat images from 1986 to 2016 (median 2013) was 2095.96 km², while that of area for glaciers in four periods of KGI were 1859.24 km², 2045.21 km², 1931.95 km², and 2163.05 km² (mean 1999.86 km², ~5% less). Hence, compared to previous glacier inventory or supraglacial debris data, our KGI data has obvious advantages in aspects including Landsat image compositing, glacier initial outline mapping, and manual correction. Thus, the quality of KGI data is considered to be reliable and sufficiently accurate.




Fig.8. Comparison between the KGI inventory and previous glacier inventories. All data are aggregated on a $5\text{ km} \times 5\text{ km}$ grid. The map in the left column represents the area difference between previous glacier inventories and the KGI inventory from the same or closest period (SCGI and ICIMOD inventories are compared with KGI-2010s, while the updated GAMDAM (GGI18) and CCI inventories are compared with KGI-2000s). The right column illustrates, from top to bottom, the scattergrams of glacier area in each 5 km grid cell of the KGI inventory, plotted against the GGI18, CCI, ICIMOD and SCGI inventories in the Karakoram mountains.

4.2 Spatial-temporal variations of the Karakoram glaciers

The total glacier area in the Karakoram was overall relatively stable from 1990 to 2020, with a slight, insignificant increase of $23.45 \pm 28.85\text{ km}^2$ ($0.10 \pm 0.13\%$) on average. However, the total glacier number decreased by 1.36% in this period. Small glaciers ($< 1\text{ km}^2$) experienced a strong decrease in area, particularly those smaller than 0.05 km^2 . In contrast, large glaciers with areas

exceeding 100 km² showed an increase in area, which compensated for the total area decrease for smaller glaciers to some extent. In terms of time series, the total glacier area showed an increasing trend from 1990 to 2010, while a decreasing trend after 2010s. This may indicate a weakening of the abnormal behaviour of glaciers in the Karakoram owing to the continuous warming.

Our results indicate that the glacier area has declined by $3.27 \pm 0.24\%$ ($133.50 \pm 9.84 \text{ km}^2$) in EK, in contrast to a slightly increased area in CK ($0.65 \pm 0.10\%$, $80.89 \pm 12.61 \text{ km}^2$) and WK ($1.26 \pm 0.11\%$, $76.06 \pm 6.49 \text{ km}^2$) (Fig. 9, Table S5). This finding is also consistent with the negative mass balance for EK glaciers in the past decade and balanced budgets in other parts of the Karakoram (Vijay and Braun, 2018; Berthier and Brun, 2019). However, contrary to the area changes, the number of glaciers in CK and WK has decreased by 2.82% and 2.24%, respectively. Nevertheless, due to the influence of seasonal or annual variations in snow patches resulting in uncertainties in glacier number changes, especially for small glaciers, these changes might not be significant. Furthermore, our results show that glaciers in the northern slope of the Karakoram mountains are potentially undergoing a loss in area ($-0.49 \pm 0.08\%$), while the area of southern slope glaciers has potentially increased by $0.84 \pm 0.18\%$. Hence, it is of particular importance to consider topographic conditions when explaining of spatial heterogeneity in glacier change as also shown elsewhere (Xie and Liu, 2009; Garg et al., 2017). We suggest that this may be due to the wetter conditions, as well as the increased precipitation the south side of the main divide, allowing glaciers in this area to continue to grow and store (Xie and Liu, 2009; Dimri, 2021).

The abundance of supraglacial debris may also partially explain the variability in glacier change observed in the Karakoram. The debris cover increased across the whole region, with EK experiencing the fastest growth (an increase of $105.05 \pm 5.16\%$ or $53.25 \pm 2.62 \text{ km}^2$) during the past three decades. This trend was followed by CK ($18.99 \pm 1.47\%$, $220.65 \pm 17.09 \text{ km}^2$) and WK ($9.44 \pm 1.13\%$, $69.40 \pm 8.27 \text{ km}^2$). The debris cover in both NK and SK areas has increased by roughly the same order of magnitude ($\sim 170 \text{ km}^2$). On the southern slope of the Karakoram, $13.81 \pm 0.51\%$ of the glacierized areas are covered by debris, and the glacier area increased by $0.84 \pm 0.19\%$, while on the northern slope, the glacier area shrank by $0.49 \pm 0.08\%$ under a relatively small debris coverage ($5.60 \pm 0.21\%$). These results show that there is a relationship between debris coverage and glacier area change, i.e., the area of glaciers with higher debris coverage is increasing or remaining stable and vice versa. From 1990 to 2020, the debris coverage has notably increased in areas above 4,200 m a.s.l., indicating that supraglacial debris in the Karakoram was experiencing the same upward evolution. This was also observed in other mountains such as the Greater Caucasus (Tielidze et al., 2020) and western Swiss Alps (Mölg et al., 2019), and is an inevitable evolution process of debris-covered glaciers under global warming (Herreid and Pellicciotti, 2020). Increased snow avalanche activity and rockfalls at high altitudes may have brought more debris to the glacier (Hewitt, 2005), thus leading to an increase in supraglacial debris, in addition to the effects of global warming, such as thinning of the glaciers and expose more englacial debris, and increased numbers of paraglacial slope failures (Zhang et al., 2011; Zhong et al., 2022; Zhang et al., 2015). By focusing on terminus changes, we observed that the advance exceeds the recession for most of the glaciers. Similar to Rankl et al. (2014), we identified 65 terminus-advancing glaciers with a total area of $1,647.19 \pm 60.62 \text{ km}^2$. In the KGI-2020s, we found 63 single glacier units, since two of the glaciers have already become branches of other existing glaciers over the past three decades, all of which greatly contributed to the increase of glacier area by up to $25.33 \pm 0.93 \text{ km}^2$. Fig. 9 also shows a clear spatial coincidence between glacier regions with increased area and distribution of the advancing glaciers, which presumably could also be confirmed by the amount of increased area and terminal change area of the advanced glaciers. In addition, our data shows that most advancing glaciers ($\sim 70\%$) are surge-type glaciers, which represents abnormal glacier behaviour in the Karakoram compared to other regions in high mountain Asia (Farinotti et al., 2020). The most recent inventory of surge-type glaciers (Guillet et al., 2022) has compiled 223 glaciers in the Karakoram. The number of which is consistent with the results of Bhambri et al. (2017), but the glaciers were not completely identical. In KGI-2020s, there are 607 glaciers in size classes greater than 5 km^2 , including 157 surge-type glaciers, which have expanded by $89.92 \pm 6.97 \text{ km}^2$ ($0.89 \pm 0.07\%$) in total area and $136.03 \pm 14.73 \text{ km}^2$ ($10.61 \pm 1.15\%$) in supraglacial debris coverage in the past three decades; while the 450 non-surge-type glaciers lost area of $4.58 \pm$

6.61 km² (0.08 ± 0.11%). Compared to normal glaciers, these anomalous glaciers (i.e., surging and advancing glaciers) are directly responsible for the area growth of the Karakoram glaciers.




Fig.9 Glacier change in the Karakoram during 1990–2020. Glacier area aggregated on a 5 km × 5 km grid. Hollow dots represent the glaciers advanced during the period 1990-2020. The Karakoram Mountains were divided into western, central and eastern Karakoram (WK, CK and EK) according to the Indus and Tarim sub-basins, and into Northern and Southern Karakoram based on the main central ridgeline.

Moreover, referring to the suggestions of Braithwaite and Raper (2009) and (Sakai et al., 2015), we assume that median glacier elevation could act as a proxy for long-term equilibrium line altitude, which is correlated with the glacier mass balance budget, that can be used to describe the state and fate of glaciers (Fujita and Nuimura, 2011). Among the five sub-basins in the Karakoram, as shown in Table S6, the median elevations of glaciers in the three basins with increasing glacier coverage decreased, while the altitudes increased in the basins with decreasing glacier area. Spatially, as pointed out by Bolch et al. (2012), the median elevations increase with the distance from the moisture source (Fig. S5). The glaciers in the northwest exposed to the westerlies and heavily debris-covered have a relatively low median elevation, while the glaciers north or northeast of the main ridge of the Karakoram have a clearly higher median elevation. On the whole, the median elevation of the Karakoram glaciers showed an increasing trend during 1990-2020, indicating that glacier melting likely is becoming more intense, with runoff moving towards peak water (Huss and Hock, 2018; Nie et al., 2021). Especially in the melt-dominated Tarim and Indus basins, accelerated glacier melt is the main contributor to rising 21st century streamflow, which increases before peak water, then declines (Huss and Hock, 2018; Rounce et al., 2020; Nie et al., 2021).

4.3 Challenges in multi-temporal glacier inventories

Algorithms for identifying debris-covered glaciers have been developed but the accuracy is too low if no manual corrections

are applied. Like the method in the present study, semi-automated approaches are the most prevalent, which will always rely on existing glacier boundaries or time-consuming manual corrections (Paul et al., 2004; Bolch et al., 2007; Bhambri et al., 2011; Racoviteanu and Williams, 2012; Smith et al., 2015; Mölg et al., 2018). Although pixel-based machine learning has been used to map debris-covered ice (Alifu et al., 2020; Xie et al., 2020b; Xie et al., 2022b), isolated pixels, irregular outlines, or voids in the results have forced it to be revised to meet the needs of glacier inventories. In addition, available remotely sensed imagery with high quality (cloud-free, low or no seasonal snow, small shadows, geometrically calibrated) is required for precisely compiling glacier inventories. Even if satellite images from the end of the ablation season are used, some mapping errors may complicate the results for investigating glacier changes, due to interannual variations in snow cover in the melting season or remaining seasonal snowfields (Yi et al., 2021). For example, during the 2010s when the overall glacier area was abnormal and the area of clean ice/snow part was high, the identified debris cover was significantly lower as a result of the high snow cover in 2009 (see Fig. S6).

To obtain inventories of the time before higher resolution multi-spectral earth-observation satellites were launched (e.g. Landsat 4, launched in 1982), topographic maps were often used to map glacier outlines. However, these have large uncertainties (Bhambri and Bolch, 2009; Bolch et al., 2010b; Guo et al., 2015; Paul, 2017). For the 1960s and 1970s, declassified Corona KH-4 (spatial resolution 8-2m)/Hexagon KH-9 (12-8m) data, and Landsat MSS (60 m) images were used to compile inventories in some regions (Bolch et al., 2008; Bolch et al., 2010b; Schmidt and Nüsser, 2012; Bhambri et al., 2013; Gardent et al., 2014; Pieczonka and Bolch, 2015; Weber et al., 2020; Bhattacharya et al., 2021). However, the available optical imagery used in current glacier inventories is often limited due to occlusion by frequent cloud layers during the melting season; this, in turn, increases the interannual variation in the inventory results and further prolongs the time interval between multitemporal glacier inventories. Fortunately, this situation improved with the launch of Sentinel-2A and B and Landsat 8 and 9, except in cloudy areas, like south-eastern Tibetan Plateau. Although SAR coherence maps have been used in previous glacier inventories to overcome these limitations in optical imagery, most focus on the identification of supraglacial debris (Frey et al., 2012; Robson et al., 2015; Ke et al., 2016; Mölg et al., 2018; Alifu et al., 2020) and only a few studies have been used to map glacier outlines (Atwood et al., 2014; Yang et al., 2016; Lippl et al., 2018). Therefore, in the future, greater availability of cloud computing platforms to develop SAR image processing algorithms and make extensive use of optical remote imagery will greatly help the development of multitemporal glacier inventories. Moreover, if glacier extents in past periods can be reconstructed based on the characteristic glacial landform like lateral and terminal moraines (Shi and Liu, 2000) using existing aerial imagery and satellite maps, the temporal resolution of multi-temporal glacier inventories can be extended, such as for glacier inventories from the Little Ice Age (e.g. Lucchesi et al. (2014) and Meier et al. (2018)).

5 Conclusions and outlook

Glacier inventories are the baseline information for many applications such as hydrological modelling climate change studies. In this study, first we generated inventories which allowed us to systematically detect glacier change patterns in the Karakoram range over the past three decades by using Landsat scenes and ancillary data through a semi-automatic method based on the GEE cloud-based platform. Validation using the “round robin” method showed an overall mapping uncertainty of $\pm 3.68\%$ which is within the accuracy range required for glacier inventories by GCOS. We also compared the KGI with previous glacier inventories and evaluated it by applying uncertainties of $\pm 1/2$ pixel (15 m) and 1 pixel (30 m) buffers to the glacier outlines. These assessments indicate that our results are reliable and sufficiently accurate. In the 2020s, there were approximately 10500 glaciers in the Karakoram mountains, covering an area of $22510.73 \pm 828.39 \text{ km}^2$ of which $10.18 \pm 0.38\%$ ($2290.95 \pm 84.31 \text{ km}^2$) is covered by supraglacial debris. During the past 30 years, the glaciers experienced a loss of clean ice area and a gain in supraglacial debris. The total glacier area in the Karakoram remained relatively stable from 1990 to 2020, with a slight, but insignificant increase of $23.45 \pm 28.85 \text{ km}^2$ ($0.10 \pm 0.13\%$), while the supraglacial debris cover increased clearly by $17.63 \pm 1.44\%$ ($343.30 \pm 27.95 \text{ km}^2$), in contrast to a $1.56 \pm 0.24\%$ ($319.85 \pm 49.92 \text{ km}^2$) decrease in clean ice/snow. These glacier areal changes were characterized by strong spatial

heterogeneity and dominated by surging and advancing glaciers. However, the glaciers are experiencing the negative effects of faster snow/ice melting caused by global warming, with small and clean glaciers being more sensitive. This poses a growing threat for regional water scarcity and glacier-related hazards in the downstream basin. Most importantly, the KGI data in this study will provide basic glacier outlines for GLIMS and the forthcoming planned release of the next versions of the RGI, as well as for other glacier-related research applications.

6 Data availability

The KGI datasets are available from National Cryosphere Desert Data Center of China at <https://doi.org/10.12072/ncdc.glacier.db2386.2022> (Xie et al., 2022a) for users' assessments and applications. The data files contain shapefiles for the complete glaciers and the debris-covered section. Additionally, a description file is included.

Author contributions. SL designed the framework of glacier inventory. FX programmed the glacier mapping algorithms, manually revised glacier outlines and ridgelines, and wrote the first manuscript. SL edited the first draft. TB reviewed and improved the manuscript. FX, YZ, YG, SD, WM, FH, XR and CQ digitized glaciers using Sentinel-2 imagery. All authors discussed and improved the manuscript.

Acknowledgments. We thank the Google Earth Engine developer for the freely available cloud-computing platform and USGS/NASA for Landsat imagery, ASTER GDEM version 3 product. We acknowledge previous glacier inventory data from Glaciers_cci, GAMDAM glacier inventory, ICIMOD glacier inventory as well as the Second Chinese Glacier Inventory for supporting this study results.

Financial support. This research has been supported by the International Science and Technology Innovation Cooperation Program of the State Key Research and Development Program (Grant No. 2021YFE0116800), the National Natural Science Foundation of China (Grant No. 42171129), the Open Foundation from National Cryosphere Desert Data Center (Grant No.2021KF01), the Second Tibetan Plateau Scientific Expedition and Research program (Grant No. 2019QZKK0208), the Pakistan Science Foundation (Grant No. PSF/CRP/Consrn-488), the Director Fund of the International Research Center of Big Data for Sustainable Development Goals (Grant No. CBAS2022DF020), the Postgraduate Research and Innovation Foundation of Yunnan University (Grant No. 2021Z018) and the Scientific Research Fund project of Yunnan Education Department (Grant No.2022Y059).

References

- Abrams, M.: Aster Global Dem Version 3, and New Aster Water Body Dataset, ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, XLI-B4, 107-110, 10.5194/isprsarchives-XLI-B4-107-2016, 2016.
- Alifu, H., Vuillaume, J.-F., Johnson, B. A., and Hirabayashi, Y.: Machine-learning classification of debris-covered glaciers using a combination of Sentinel-1/-2 (SAR/optical), Landsat 8 (thermal) and digital elevation data, *Geomorphology*, 369, 10.1016/j.geomorph.2020.107365, 2020.
- Andreassen, L. M., Paul, F., Kaab, A., and Hausberg, J. E.: Landsat-derived glacier inventory for Jotunheimen, Norway, and deduced glacier changes since the 1930s, *Cryosphere*, 2, 131-145, 10.5194/tc-2-131-2008, 2008.
- Atwood, D. K., Meyer, F., and Arendt, A.: Using L-band SAR coherence to delineate glacier extent, *Canadian Journal of Remote Sensing*, 36, S186-S195, 10.5589/m10-014, 2014.
- Bajracharya, S. R. and Shrestha, B.: The Status of Glaciers in the Hindu Kush-Himalayan Region, International Centre for Integrated Mountain Development, GPO Box 3226, Kathmandu, Nepal2011.
- Baumann, S., Anderson, B., Chinn, T., Mackintosh, A., Collier, C., Lorrey, A. M., Rack, W., Purdie, H., and Eaves, S.: Updated inventory of glacier ice in New Zealand based on 2016 satellite imagery, *Journal of Glaciology*, 1-14, 10.1017/jog.2020.78, 2020.
- Bazai, N. A., Cui, P., Carling, P. A., Wang, H., Hassan, J., Liu, D., Zhang, G., and Jin, W.: Increasing glacial lake outburst flood

- hazard in response to surge glaciers in the Karakoram, *Earth-Science Reviews*, 212, 10.1016/j.earscirev.2020.103432, 2021.
- 545 Berthier, E. and Brun, F.: Karakoram geodetic glacier mass balances between 2008 and 2016: persistence of the anomaly and influence of a large rock avalanche on Siachen Glacier, *Journal of Glaciology*, 65, 494-507, 10.1017/jog.2019.32, 2019.
- Bhambri, R. and Bolch, T.: Glacier mapping: a review with special reference to the Indian Himalayas, *Progress in Physical Geography*, 33, 672-704, 10.1177/0309133309348112
10.5167/uzh-137192, 2009.
- 550 Bhambri, R., Bolch, T., and Chaujar, R. K.: Mapping of debris-covered glaciers in the Garhwal Himalayas using ASTER DEMs and thermal data, *International Journal of Remote Sensing*, 32, 8095-8119, 10.1080/01431161.2010.532821, 2011.
- Bhambri, R., Hewitt, K., Kawishwar, P., and Pratap, B.: Surge-type and surge-modified glaciers in the Karakoram, *Sci Rep*, 7, 15391, 10.1038/s41598-017-15473-8, 2017.
- Bhambri, R., Bolch, T., Kawishwar, P., Dobhal, D. P., Srivastava, D., and Pratap, B.: Heterogeneity in glacier response in the upper Shyok valley, northeast Karakoram, *The Cryosphere*, 7, 1385-1398, 10.5194/tc-7-1385-2013, 2013.
- 555 Bhambri, R., Chand, P., Nüsser, M., Kawishwar, P., Kumar, A., Gupta, A. K., Verma, A., and Tiwari, S. K.: Reassessing the Karakoram Through Historical Archives, in: *Environmental Change in South Asia: Essays in Honor of Mohammed Taher*, edited by: Saikia, A., and Thapa, P., Springer International Publishing, Cham, 139-169, 10.1007/978-3-030-47660-1_8, 2022.
- Bhattacharya, A., Bolch, T., Mukherjee, K., King, O., Menounos, B., Kapitsa, V., Neckel, N., Yang, W., and Yao, T.: High Mountain Asian glacier response to climate revealed by multi-temporal satellite observations since the 1960s, *Nature communications*, 12, 4133, 10.1038/s41467-021-24180-y, 2021.
- 560 Bolch, T.: Past and Future Glacier Changes in the Indus River Basin, in: *Indus River Basin*, edited by: Bolch, T., Elsevier Inc., 85-97, 10.1016/B978-0-12-812782-7.00004-7, 2019.
- Bolch, T., Menounos, B., and Wheate, R.: Landsat-based inventory of glaciers in western Canada, 1985–2005, *Remote Sensing of Environment*, 114, 127-137, 10.1016/j.rse.2009.08.015, 2010a.
- 565 Bolch, T., Buchroithner, M., Pieczonka, T., and Kunert, A.: Planimetric and volumetric glacier changes in the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data, *Journal of Glaciology*, 54, 592-600, 10.3189/002214308786570782, 2008.
- Bolch, T., Buchroithner, M. F., Kunert, A., and Kamp, U.: Automated delineation of debris-covered glaciers based on ASTER data, in: *GeoInformation in Europe*, M.A. Gomasca (ed.), edited by: Bolch, T., Buchroithner, M. F., Kunert, A., and Kamp, U., Millpress, Netherlands, 403-410, 2007.
- 570 Bolch, T., Pieczonka, T., Mukherjee, K., and Shea, J.: Brief communication: Glaciers in the Hunza catchment (Karakoram) have been nearly in balance since the 1970s, *The Cryosphere*, 11, 531-539, 10.5194/tc-11-531-2017, 2017.
- Bolch, T., Yao, T., Kang, S., Buchroithner, M. F., Scherer, D., Maussion, F., Huintjes, E., and Schneider, C.: A glacier inventory for the western Nyainqentanglha Range and the Nam Co Basin, Tibet, and glacier changes 1976–2009, *The Cryosphere*, 4, 419-433, 10.5194/tc-4-419-2010, 2010b.
- 575 Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., Bajracharya, S., and Stoffel, M.: The State and Fate of Himalayan Glaciers, *Science*, 336, 310-315, 10.1126/science.1215828, 2012.
- Bolch, T., Shea, J. M., Liu, S., Azam, F. M., Gao, Y., Gruber, S., Immerzeel, W. W., Kulkarni, A., Li, H., Tahir, A. A., Zhang, G., and Zhang, Y.: Status and Change of the Cryosphere in the Extended Hindu Kush Himalaya Region, in: *The Hindu Kush Himalaya Assessment*, edited by: Bolch, T., Shea, J. M., Liu, S., Azam, F. M., Gao, Y., Gruber, S., Immerzeel, W. W., Kulkarni, A., Li, H., Tahir, A. A., Zhang, G., and Zhang, Y., 209-255, 10.1007/978-3-319-92288-1_7, 2019.
- 580 Braithwaite, R. J. and Raper, S. C. B.: Estimating equilibrium-line altitude (ELA) from glacier inventory data, *Annals of Glaciology*, 50, 10.3189/172756410790595930, 2009.
- Dehecq, A., Gourmelen, N., Gardner, A. S., Brun, F., Goldberg, D., Nienow, P. W., Berthier, E., Vincent, C., Wagnon, P., and Trouvé, E.: Twenty-first century glacier slowdown driven by mass loss in High Mountain Asia, *Nature Geoscience*, 12, 22-27, 10.1038/s41561-018-0271-9, 2018.
- Dimri, A. P.: Decoding the Karakoram Anomaly, *Science of The Total Environment*, 788, 10.1016/j.scitotenv.2021.147864, 2021.
- Dozier, J.: Spectral Signature of Alpine Snow Cover from the Landsat Thematic Mapper, *Remote Sensing of Environment*, 28, 9-22, 10.1016/0034-4257(89)90101-6, 1989.
- 590 Farinotti, D., Immerzeel, W. W., and Dehecq, A.: Manifestations and mechanisms of the Karakoram glacier Anomaly, *Nature Geoscience*, 13, 8-16, 10.1038/s41561-019-0513-5, 2020.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12, 168-173, 10.1038/s41561-019-0300-3, 2019.
- 595 Frey, H., Paul, F., and Strozzi, T.: Compilation of a glacier inventory for the western Himalayas from satellite data: methods, challenges, and results, *Remote Sensing of Environment*, 124, 832-843, 10.1016/j.rse.2012.06.020, 2012.
- Fujita, K. and Nuimura, T.: Spatially heterogeneous wastage of Himalayan glaciers, *Proc Natl Acad Sci U S A*, 108, 14011-14014, 10.1073/pnas.1106242108, 2011.
- Gao, Y., Liu, S., Qi, M., Xie, F., Wu, K., and Zhu, Y.: Glacier-Related Hazards Along the International Karakoram Highway: Status and Future Perspectives, *Frontiers in Earth Science*, 9, 10.3389/feart.2021.611501, 2021.
- 600 Gardelle, J., Berthier, E., and Arnaud, Y.: Slight mass gain of Karakoram glaciers in the early twenty-first century, *Nature Geoscience*, 5, 322-325, 10.1038/ngeo1450, 2012.
- Gardent, M., Rabatel, A., Dedieu, J.-P., and Deline, P.: Multitemporal glacier inventory of the French Alps from the late 1960s to the late 2000s, *Global and Planetary Change*, 120, 24-37, 10.1016/j.gloplacha.2014.05.004, 2014.

- Garg, P. K., Shukla, A., and Jasrotia, A. S.: Influence of topography on glacier changes in the central Himalaya, India, *Global and Planetary Change*, 155, 196-212, 10.1016/j.gloplacha.2017.07.007, 2017.
- GCOS: The Global Observing System for Climate: Implementation Needs (GCOS-200), WMO., 2016.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sensing of Environment*, 202, 18-27, 10.1016/j.rse.2017.06.031, 2017.
- Granshaw, F. D. and G. Fountain, A.: Glacier change (1958–1998) in the North Cascades National Park Complex, Washington, USA, *Journal of Glaciology*, 52, 251-256, 10.3189/172756506781828782, 2006.
- Guillet, G., King, O., Lv, M., Ghuffar, S., Benn, D., Quincey, D., and Bolch, T.: A regionally resolved inventory of High Mountain Asia surge-type glaciers, derived from a multi-factor remote sensing approach, *The Cryosphere*, 16, 603-623, 10.5194/tc-16-603-2022, 2022.
- Guo, W., Liu, S., Xu, J., Wu, L., Shangguan, D., Yao, X., Wei, J., Bao, W., Yu, P., Liu, Q., and Jiang, Z.: The second Chinese glacier inventory: data, methods and results, *Journal of Glaciology*, 61, 357-372, 10.3189/2015JoG14J209, 2015.
- Herreid, S. and Pellicciotti, F.: The state of rock debris covering Earth's glaciers, *Nature Geoscience*, 13, 621-627, 10.1038/s41561-020-0615-0, 2020.
- Herreid, S., Pellicciotti, F., Ayala, A., Chesnokova, A., Kienholz, C., Shea, J., and Shrestha, A.: Satellite observations show no net change in the percentage of supraglacial debris-covered area in northern Pakistan from 1977 to 2014, *Journal of Glaciology*, 61, 524-536, 10.3189/2015JoG14J227, 2017.
- Hewitt, K.: The Karakoram Anomaly? Glacier Expansion and the 'Elevation Effect,' *Karakoram Himalaya, Mountain Research and Development*, 25, 332-340, 10.1659/0276-4741(2005)025[0332:Tkagea]2.0.Co;2, 2005.
- Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 135-140, 10.1038/s41558-017-0049-x, 2018.
- Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernandez, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., and Baillie, J. E. M.: Importance and vulnerability of the world's water towers, *Nature*, 577, 364-369, 10.1038/s41586-019-1822-y, 2020.
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J., and Arnaud, Y.: Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas, *Nature*, 488, 495-498, 10.1038/nature11324, 2012.
- Ke, L., Ding, X., Zhang, L. E. I., Hu, J. U. N., Shum, C. K., and Lu, Z.: Compiling a new glacier inventory for southeastern Qinghai-Tibet Plateau from Landsat and PALSAR data, *Journal of Glaciology*, 62, 579-592, 10.1017/jog.2016.58, 2016.
- Kraaijenbrink, P. D. A., Bierkens, M. F. P., Lutz, A. F., and Immerzeel, W. W.: Impact of a global temperature rise of 1.5 degrees Celsius on Asia's glaciers, *Nature*, 549, 257-260, 10.1038/nature23878, 2017.
- Lehner, B. and Grill, G.: Global river hydrography and network routing: baseline data and new approaches to study the world's large river systems, *Hydrological Processes*, 27, 2171-2186, 10.1002/hyp.9740, 2013.
- Li, Z., Wang, N., Chen, A. a., Liang, Q., and Yang, D.: Slight change of glaciers in the Pamir over the period 2000–2017, *Arctic, Antarctic, and Alpine Research*, 54, 13-24, 10.1080/15230430.2022.2028475, 2022.
- Liao, H., Liu, Q., Zhong, Y., and Lu, X.: Landsat-Based Estimation of the Glacier Surface Temperature of Hailuoguo Glacier, Southeastern Tibetan Plateau, Between 1990 and 2018, *Remote Sensing*, 12, 10.3390/rs12132105, 2020.
- Lippl, S., Vijay, S., and Braun, M.: Automatic delineation of debris-covered glaciers using InSAR coherence derived from X-, C- and L-band radar data: a case study of Yazgyl Glacier, *Journal of Glaciology*, 64, 811-821, 10.1017/jog.2018.70, 2018.
- Liu, S., Ding, Y., Shangguan, D., Zhang, Y., Li, J., Han, H., Wang, J., and Xie, C.: Glacier retreat as a result of climate warming and increased precipitation in the Tarim river basin, northwest China, *Annals of Glaciology*, 43, 10.3189/172756406781812168, 2006.
- Liu, S., Yao, X., Shangguan, D., Guo, W., Wei, J., Xu, J., Bao, W., and Wu, L.: The contemporary glaciers in China based on the Second Chinese Glacier Inventory (in Chinese), *ACTA GEOGRAPHICA SINICA*, 70, 3-16, 10.11821/dlxb201501001, 2015.
- Liu, S., Wu, T., Wang, X., Wu, X., Yao, X., Liu, Q., Zhang, Y., Wei, J., and Zhu, X.: Changes in the global cryosphere and their impacts: A review and new perspective., *Sciences in Cold and Arid Regions*, 12(6): 343–354., 343-354, 10.3724/SP.J.1226.2020.00343, 2020.
- Lucchesi, S., Fioraso, G., Bertotto, S., and Chiarle, M.: Little Ice Age and contemporary glacier extent in the Western and South-Western Piedmont Alps (North-Western Italy), *Journal of Maps*, 10, 409-423, 10.1080/17445647.2014.880226, 2014.
- Meier, W. J. H., Griebinger, J., Hochreuther, P., and Braun, M. H.: An Updated Multi-Temporal Glacier Inventory for the Patagonian Andes With Changes Between the Little Ice Age and 2016, *Frontiers in Earth Science*, 6, 10.3389/feart.2018.00062, 2018.
- Millan, R., Mouginit, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nature Geoscience*, 15, 124-+, 10.1038/s41561-021-00885-z, 2022a.
- Millan, R., Mouginit, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nature Geoscience*, 15, 124-129, 10.1038/s41561-021-00885-z, 2022b.
- Minora, U., Bocchiola, D., D'Agata, C., Maragno, D., Mayer, C., Lambrecht, A., Vuillermoz, E., Senese, A., Compostella, C., Smiraglia, C., and Diolaiuti, G. A.: Glacier area stability in the Central Karakoram National Park (Pakistan) in 2001–2010: The "Karakoram Anomaly" in the spotlight, *Progress in Physical Geography: Earth and Environment*, 40, 629-660, 10.1177/0309133316643926, 2016.
- Mölg, N., Bolch, T., Walter, A., and Vieli, A.: Unravelling the evolution of Zmuttgletscher and its debris cover since the end of the

- 665 Little Ice Age, *The Cryosphere*, 13, 1889-1909, 10.5194/tc-13-1889-2019, 2019.
- Mölg, N., Bolch, T., Rastner, P., Strozzi, T., and Paul, F.: A consistent glacier inventory for Karakoram and Pamir derived from Landsat data: distribution of debris cover and mapping challenges, *Earth Syst. Sci. Data*, 10, 1807-1827, 10.5194/essd-10-1807-2018, 2018.
- Müller, F. and Scherler, K.: Introduction to the world glacier inventory, 1980.
- 670 Nie, Y., Pritchard, H. D., Liu, Q., Hennig, T., Wang, W., Wang, X., Liu, S., Nepal, S., Samyn, D., Hewitt, K., and Chen, X.: Glacial change and hydrological implications in the Himalaya and Karakoram, *Nature Reviews Earth & Environment*, 2, 91-106, 10.1038/s43017-020-00124-w, 2021.
- Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A., Hoshina, Y., Takenaka, S., Omiya, S., Tsunematsu, K., Tshering, P., and Fujita, K.: The GAMDAM glacier inventory: a quality-controlled inventory of Asian glaciers, *The Cryosphere*, 9, 849-864, 10.5194/tc-9-849-2015, 2015.
- 675 Paul, F.: Glacier Inventory, in: *International Encyclopedia of Geography: People, the Earth, Environment and Technology*, edited by: Paul, F., 1-12, 10.1002/9781118786352.wbieg0877, 2017.
- Paul, F., Huggel, C., and Kääb, A.: Combining satellite multispectral image data and a digital elevation model for mapping debris-covered glaciers, *Remote Sensing of Environment*, 89, 510-518, 10.1016/j.rse.2003.11.007, 2004.
- 680 Paul, F., Kääb, A., Maisch, M., Hoelzle, M., and Haerberli, W.: The new remote-sensing-derived Swiss glacier inventory: I. Methods, *Annals of Glaciology*, 34, 355-361, 10.3189/172756402781817941, 2002.
- Paul, F., Barry, R. G., Cogley, J. G., Frey, H., Haerberli, W., Ohmura, A., Ommanney, C. S. L., Raup, B., Rivera, A., and Zemp, M.: Recommendations for the compilation of glacier inventory data from digital sources, *Annals of Glaciology*, 50, 119-126, 10.3189/172756410790595778, 2009.
- 685 Paul, F., Bolch, T., Briggs, K., Kääb, A., McMillan, M., McNabb, R., Nagler, T., Nuth, C., Rastner, P., Strozzi, T., and Wuite, J.: Error sources and guidelines for quality assessment of glacier area, elevation change, and velocity products derived from satellite data in the *Glaciers_cci* project, *Remote Sensing of Environment*, 203, 10.1016/j.rse.2017.08.038, 2017.
- Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemeč, J., Rabatel, A., Ramusovic, M., Schwaizer, G., and Smiraglia, C.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2, *Earth System Science Data*, 12, 1805-1821, 10.5194/essd-12-1805-2020, 2020.
- 690 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Konovalov, V., Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K., Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, *Annals of Glaciology*, 54, 171-182, 10.3189/2013AoG63A296, 2013.
- 695 Pfeiffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., and Sharp, M. J.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of Glaciology*, 60, 537-552, 10.3189/2014JoG13J176, 2014.
- Pieczonka, T. and Bolch, T.: Region-wide glacier mass budgets and area changes for the Central Tien Shan between ~1975 and 1999 using Hexagon KH-9 imagery, *Global and Planetary Change*, 128, 1-13, 10.1016/j.gloplacha.2014.11.014, 2015.
- 700 Racoviteanu, A. and Williams, M. W.: Decision Tree and Texture Analysis for Mapping Debris-Covered Glaciers in the Kangchenjunga Area, Eastern Himalaya, *Remote Sensing*, 4, 3078-3109, 10.3390/rs4103078, 2012.
- Rankl, M., Kienholz, C., and Braun, M.: Glacier changes in the Karakoram region mapped by multimission satellite imagery, *The Cryosphere*, 8, 977-989, 10.5194/tc-8-977-2014, 2014.
- 705 RGI Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0: Technical Report, Global Land Ice Measurements from Space, Colorado, USA. Digital Media., [10.7265/N5-RGI-60](https://doi.org/10.7265/N5-RGI-60), 2017.
- Robson, B. A., Nuth, C., Dahl, S. O., Hölbling, D., Strozzi, T., and Nielsen, P. R.: Automated classification of debris-covered glaciers combining optical, SAR and topographic data in an object-based environment, *Remote Sensing of Environment*, 170, 372-387, 10.1016/j.rse.2015.10.001, 2015.
- 710 Rounce, D. R., Hock, R., and Shean, D. E.: Glacier Mass Change in High Mountain Asia Through 2100 Using the Open-Source Python Glacier Evolution Model (PyGEM), *Frontiers in Earth Science*, 7, 10.3389/feart.2019.00331, 2020.
- Sakai, A.: Brief communication: Updated GAMDAM glacier inventory over high-mountain Asia, *The Cryosphere*, 13, 2043-2049, 10.5194/tc-13-2043-2019, 2019.
- 715 Sakai, A., Nuimura, T., Fujita, K., Takenaka, S., Nagai, H., and Lamsal, D.: Climate regime of Asian glaciers revealed by GAMDAM glacier inventory, *The Cryosphere*, 9, 865-880, 10.5194/tc-9-865-2015, 2015.
- Scherler, D., Wulf, H., and Gorelick, N.: Global Assessment of Supraglacial Debris-Cover Extents, *Geophysical Research Letters*, 45, 11,798-711,805, 10.1029/2018gl080158, 2018.
- Schmidt, S. and Nüsser, M.: Changes of High Altitude Glaciers from 1969 to 2010 in the Trans-Himalayan Kang Yatze Massif, Ladakh, Northwest India, Arctic, Antarctic, and Alpine Research, 44, 107-121, 10.1657/1938-4246-44.1.107, 2012.
- 720 Shi, Y. and Liu, S.: Estimation on the response of glaciers in China to the global warming in the 21st century, *Chinese Science Bulletin*, 45, 668-672, 10.1007/BF02886048, 2000.
- Shih, Y. f., Hsieh, T. c., Cheng, P. h., and Li, C. c.: Distribution, features and variations of glaciers in China, *World Glacier Inventory - Inventaire mondial des Glaciers*,
- Smith, T., Bookhagen, B., and Cannon, F.: Improving semi-automated glacier mapping with a multi-method approach: applications in central Asia, *The Cryosphere*, 9, 1747-1759, 10.5194/tc-9-1747-2015, 2015.

- Tielidze, L. G., Bolch, T., Wheate, R. D., Kutuzov, S. S., Lavrentiev, I. I., and Zemp, M.: Supra-glacial debris cover changes in the Greater Caucasus from 1986 to 2014, *The Cryosphere*, 14, 585-598, 10.5194/tc-14-585-2020, 2020.
- Vijay, S. and Braun, M.: Early 21st century spatially detailed elevation changes of Jammu and Kashmir glaciers (Karakoram–Himalaya), *Global and Planetary Change*, 165, 137-146, 10.1016/j.gloplacha.2018.03.014, 2018.
- 730 Weber, P., Andreassen, L. M., Boston, C. M., Lovell, H., and Kvarteig, S.: An ~1899 glacier inventory for Nordland, northern Norway, produced from historical maps, *Journal of Glaciology*, 66, 259-277, 10.1017/jog.2020.3, 2020.
- Winiger, M., Gumpert, M., and Yamout, H.: Karakorum-Hindukush-western Himalaya: assessing high-altitude water resources, *Hydrological Processes*, 19, 2329-2338, 10.1002/hyp.5887, 2005.
- 735 Wu, K., Liu, S., Jiang, Z., Liu, Q., Zhu, Y., Yi, Y., Xie, F., Ahmad Tahir, A., and Saifullah, M.: Quantification of glacier mass budgets in the Karakoram region of Upper Indus Basin during the early twenty-first century, *Journal of Hydrology*, 603, 10.1016/j.jhydrol.2021.127095, 2021.
- Wu, K., Liu, S., Jiang, Z., Zhu, Y., Xie, F., Gao, Y., Yi, Y., Tahir, A. A., and Muhammad, S.: Surging Dynamics of Glaciers in the Hunza Valley under an Equilibrium Mass State since 1990, *Remote Sensing*, 12, 10.3390/rs12182922, 2020.
- 740 Xie, F., Liu, S., Duan, S., Miao, W., Pan, X., and Qin, C.: Interdecadal glacier inventories in the Karakoram since the 1990s [dataset], 10.12072/ncdc.glacier.db2386.2022, 2022a.
- Xie, F., Liu, S., Gao, Y., Zhu, Y., Wu, K., Qi, M., Duan, S., and Tahir, A. M.: Derivation of Supraglacial Debris Cover by Machine Learning Algorithms on the GEE Platform: A Case Study of Glaciers in the Hunza Valley, *ISPRS Annals of Photogrammetry, Remote Sensing and Spatial Information Sciences*, V-3-2020, 417-424, 10.5194/isprs-annals-V-3-2020-417-2020, 2020a.
- 745 Xie, F., Liu, S., Wu, K., Zhu, Y., Gao, Y., Qi, M., Duan, S., Saifullah, M., and Tahir, A. A.: Upward Expansion of Supra-Glacial Debris Cover in the Hunza Valley, Karakoram, During 1990 ~ 2019, *Frontiers in Earth Science*, 8, 10.3389/feart.2020.00308, 2020b.
- Xie, Z. and Liu, C.: *Introduction to glaciology*(In Chinese), Shanghai Universal Science Press 2009.
- 750 Xie, Z., Haritashya, U. K., Asari, V. K., Bishop, M. P., Kargel, J. S., and Aspiras, T. H.: GlacierNet2: A hybrid Multi-Model learning architecture for alpine glacier mapping, *International Journal of Applied Earth Observation and Geoinformation*, 112, 10.1016/j.jag.2022.102921, 2022b.
- Yang, Y., Li, Z., Huang, L., Tian, B., and Chen, Q.: Extraction of glacier outlines and water-eroded stripes using high-resolution SAR imagery, *International Journal of Remote Sensing*, 37, 1016-1034, 10.1080/01431161.2016.1145365, 2016.
- 755 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D. B., and Joswiak, D.: Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings, *Nature Climate Change*, 2, 663-667, 10.1038/nclimate1580, 2012.
- Yi, Y., Liu, S., Zhu, Y., Wu, K., Xie, F., and Saifullah, M.: Spatiotemporal heterogeneity of snow cover in the central and western Karakoram Mountains based on a refined MODIS product during 2002–2018, *Atmospheric Research*, 250, 10.1016/j.atmosres.2020.105402, 2021.
- 760 Zhang, Y. and Liu, S.: Modeling of the Mass Balance of Glaciers with Debris Cover, in: *Geo-intelligence for Sustainable Development*, edited by: Zhang, Y., and Liu, S., Springer Singapore, Singapore, 191-212, 10.1007/978-981-16-4768-0_12, 2021.
- Zhang, Y., Fujita, K., Shiyin, L., Liu, Q., and Nuimura, T.: Distribution of debris thickness and its effect on ice melt at Hailuoguo glacier, southeastern Tibetan Plateau, using in situ surveys and ASTER imagery, *Journal of Glaciology*, 57, <https://doi.org/10.3189/002214311798843331>, 2011.
- 765 Zhang, Y., Hirabayashi, Y., Fujita, K., Liu, S., and Liu, Q.: Heterogeneity in supraglacial debris thickness and its role in glacier mass changes of the Mount Gongga, *Science China Earth Sciences*, 59, 170-184, 10.1007/s11430-015-5118-2, 2015.
- Zhang, Z., Liu, S., Zhang, Y., Wei, J., Jiang, Z., and Wu, K.: Glacier variations at Aru Co in western Tibet from 1971 to 2016 derived from remote-sensing data, *Journal of Glaciology*, 64, 397-406, 10.1017/jog.2018.34, 2018.
- 770 Zhong, Y., Liu, Q., Westoby, M., Nie, Y., Pellicciotti, F., Zhang, B., Cai, J., Liu, G., Liao, H., and Lu, X.: Intensified paraglacial slope failures due to accelerating downwasting of a temperate glacier in Mt. Gongga, southeastern Tibetan Plateau, *Earth Surface Dynamics*, 10, 23-42, 10.5194/esurf-10-23-2022, 2022.