

Reviewer comments ESSD-2022-288

RC1: 'Comment on essd-2022-288', Anonymous Referee #1, 02 Nov 2022

review of "Hydro-PE: gridded datasets of historical and future Penman-Monteith potential evaporation for the United Kingdom" by Robinson, Brown, Kay, Lane, Chapman, Bell and Blyth

The authors describe two new datasets for Potential EvapoTranspiration (PET) for the UK. These datasets are based on (processed) observations (provide an estimate of the PET values for the current climate) and model climate scenario from UKCP (and provide an estimate of future conditions of PET). The datasets come as an estimate of PET and include a parameterization of inception on days with rain. The approach leading to both datasets and the assumptions are discussed in the documentation in sufficient detail and the embedding in the literature is well done. The manuscript is a very good read and very clear.

However, there are a few concerns that need to be addressed before acceptance of the paper can be considered. These relate to 1) the uncertainty introduced by the need to use monthly mean values for sunshine duration and wind which are interpolated to the daily level and 2) inconsistencies between the input meteorological data for the historic period and the future period - and how these propagate into the PET estimates.

In addition, it would make the documentation of this dataset even more interesting if the manuscript would add a focus on climatology of more extreme situations and their projected changes. The monthly mean perspective is clearly relevant, but many users will be interested in the shortish warm spells with exceptionally sunny weather as these affect hydrology quite profoundly.

These concerns are detailed below.

My advice to the editor is to accept the paper with major revisions.

Main concerns:

1) It is a pity that daily maps for wind speed and sunshine duration (and air pressure) are not available in the HadUK-Grid dataset. The approach taken by the authors is then probably the best way forward. However, there are potentially large effects on the PET estimates because of the strong underestimation of the day-to-day variation in PET values. As the authors state in the manuscript, PET is non-linearly related to the input which makes the underestimation of day-to-day variations a potential problem. While section 5.3 comes some way in demonstrating that the approach of the manuscript works, but the answers are not quite satisfying. In my view, an additional analysis needs to be done. The UK Met Office's National Climate Information Centre has long records available (starting in the early 1960s) with daily values of sunshine duration. Long series with wind speed are available as well I guess, but can be obtained through NOAA's Global Summary of the Day (or through the ECA&D). These records allow for a day-to-day comparison of selected UK stations of PET using actual daily values and interpolated values.

The benefit of this approach - easy to do as you already have all the software in place - is that an assessment of the neglect of day-to-day variability on PET and PETI values can be made directly, also for the summer season (which is left-out in section 5.3). In this assessment, the monthly mean perspective is interesting (and in line with the rest of the manuscript), but a view on the under- or overestimation on the more extreme days (like total overcast days or warm and long sunny spells like those in the recent heat waves) is of particular value to the user.

2) Figs. 10 and 11 compare outcomes of PET values based on the (processed) observations and the UKCP climate scenario's. The manuscript discusses some issues with the model results, like the presence of a bias in the variables etc. What is missing a bit is the identification (and assessment of the severity) of a possible mis-match between model and observations. Apart from a possible bias, differences between observations and model meteorological data include the use of sunshine duration (obs) vs. radiation (model) (at least: this is what I expect that is used when processing the UKCP data). The difference between daily mean temperature (model) vs. Tmax and Tmin (obs) is another. Probably these differences are smaller than any model bias, but it would be good to establish this.

3) With the recent heat waves Europe has seen, many people will be interested in how the dataset performs for these more extreme situations. The current manuscript lacks any detail in this respect and that is a pity. An assessment of the quality of the dataset during dry spells/heat waves is required. The comparison of climatology of extreme PET values - like the climatology of 90th or 95th percentiles in daily PET - between the historical period and the projections is required for the user to gain trust that the dataset is reliable in this respect as well.

Other aspects that authors may want to look into

-) E5: (ground heat flux)

In this section, it appears that ground heat flux and ground heat storage are mixed-up. It was my understanding that the ground heat flux is often neglected when focusing on daily resolution estimates of PE.

-) line 325: the period 1985-1992 is remarkably short for any comparison. I think observational data are available for a more expansive period.

-) line 330: be clear if the standard deviation and correlation analysis involves the daily values or monthly mean values. The reader will see later in this paragraph (line 338) that daily values are used. Note that for the Pearson correlation, the seasonal cycle needs to be filtered-away (otherwise the correlation is likely to partly reflect that both datasets capture the seasonal cycle).

Citation: <https://doi.org/10.5194/essd-2022-288-RC1>

RC2: 'Comment on essd-2022-288', Anonymous Referee #2, 21 Mar 2023

This paper about two potential evapotranspiration datasets is well organized and documented. The datasets appear relevant and useful for hydrological modelling.

I agree with the thoughtful comments of the first reviewer (<https://doi.org/10.5194/essd-2022-288-RC1>). Some additional critical discussion of methodological constraints on the results could further increase the usefulness of the data for different applications. Besides the issues from the first review, I do not have other concerns with the manuscript.

Finally, I encourage the authors to consider making the code for their analysis available, which could allow testing other input data, application at different regions, etc.

Citation: <https://doi.org/10.5194/essd-2022-288-RC2>

Author response ESSD-2022-288

We thank both reviewers for their considered comments about the manuscript. Overall we agree that a further assessment of the methodology and analysis of extremes will help potential users to understand the utility of the data set, and will add these to the manuscript.

In addition to the specific responses below, we will also update the text to add some clarifying details about the use of the UKCP18 ensemble, particularly in the context of other available climate projections. We will also publish the potential evaporation (PE) calculation code and add it as an asset to this paper.

Response to RC1:

- 1) We agree that an assessment of the impact of using interpolated variables rather than actual daily values would be of use to potential users. We will identify a data set (or combination of data sets) containing all of the required variables at daily timestep. This will enable us to add an evaluation of the impact on the variability of PE when using interpolated monthly variables compared to the available daily variables. We will also compare with the use of raw monthly means.
- 2) Methods such as the calculation of shortwave radiation from sunshine hours and the use of daily minimum and maximum temperature to estimate the daily mean temperature are well-established when working with station data. However, it is still useful to demonstrate the impact of them on the PE calculations, particularly as the available input variables differ between our historical and future data sets. We will ensure that the data set(s) that we use to address point 1 can also be used to demonstrate this impact, and we will include this in the additional analysis.
- 3) We agree that an assessment of extremes would be a useful addition to this manuscript. We will add the suggested comparison of high percentiles between the historical and future PE data sets, and will also consider this when addressing point 1. We think that an analysis for particular meteorological regimes (eg heatwaves, dry spells) is more detailed than is necessary for this data paper, but that an analysis of high percentiles on a seasonal or monthly basis will address the need to assess the performance at extremes for potential users.

Other concerns:

E5: It is true that ground heat flux is often neglected in calculations of daily PE. However, it is included in the MORECS calculations, and we have retained it for compatibility with the existing MORECS PE. The use of the average daily heat storage to calculate the daily ground heat flux follows the MORECS methodology.

L325: Section 5.2 is specifically a comparison with site-level MORECS data, in order to help disentangle the differences between HadUK-Grid and MORECS that are due to the resolution mismatch from other differences. These site-level MORECS data are only available for this relatively short period, but we do not think that a comparison with other site level PE products would add to this particular comparison as they would not be equivalent to MORECS. We note that Section 5.1 does provide a comparison over a longer time period with gridded MORECS and CHES-PE, albeit at a broader spatial scale. We have targeted these comparisons with MORECS and CHES-PE due to their widespread use in UK hydrological and ecological modelling.

L330: We will clarify the text to specify that we used daily values to calculate these statistics. We will also calculate the Pearson correlation after filtering for the seasonal cycle for comparison.

Response to RC2:

We intend that addressing the comments of RC1 will provide the added critical discussion of the methodology that RC2 suggests.

We will make our PE calculation code available through github. This will allow researchers to apply it to other input data sets, either observational or model outputs, which will help to expand the understanding of uncertainty in historical and future projections of PE.

Manuscript changes ESSD-2022-288

As well as some miscellaneous grammar and word edits, we made the following changes to the manuscript:

L124-140 Added more details about the UKCP18 RCM ensemble and its relationship to CMIP5 and CMIP6.

L261-271 and Fig 11 Added a figure showing the climatology of the 90th percentiles of the data sets for the historical and future periods, and text discussing the differences.

L289-292, L297-298, L326-331 and Fig 13 Added a figure showing the climatology of the 90th percentiles in the historical period alongside the percentiles of CHES-PE, and text discussing the differences. We found that the Hydro-PE HadUK-Grid 90th percentiles were relatively high, but this is understood.

L362-365 and Fig 15 Specified that the metrics are calculated with daily data. Added the Pearson correlation calculated with deseasonalised data

L413-450 and Fig 18 and 19 We carried out an analysis using the PLUMBER2 quality-controlled and gap filled flux site data. This allowed us to investigate the impact of the interpolation and calculation of variables used for HadUK-Grid. We found that the PET and PETI calculated with the equivalents of the procedure applied to HadUK-Grid were consistent with those calculated from daily input variables, and that the differences could help explain the biases seen in the Hydro-PE HadUK-Grid data.

L457-460 Added a recommendation for users who are concerned about the high extremes

L509-511 Added a note about other data sets that may be used

L516 Added a code availability statement

L587-589 Added equation for calculating daily mean temperature from daily minimum and maximum

L678 Added some explanation of the treatment of ground heat flux

We have also updated **Fig 3** and **Fig 6**, to include the Hydro-PE HadUK-Grid data for 2021.

Hydro-PE: gridded datasets of historical and future Penman-Monteith potential evaporation for the United Kingdom

Emma L. Robinson¹, Matthew J. Brown¹, Alison L. Kay¹, Rosanna A. Lane¹, Rhian Chapman¹, Victoria A. Bell¹, and Eleanor M. Blyth¹

¹UK Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

Correspondence: Emma L. Robinson (emrobi@ceh.ac.uk)

Abstract. We present two new potential evaporation datasets for the United Kingdom: a historical dataset, Hydro-PE HadUK-Grid, which is derived from the HadUK-Grid gridded observed meteorology (1969–2021); and a future dataset, Hydro-PE UKCP18 RCM, which is derived from UKCP18 regional climate projections (1980–2080). Both datasets are suitable for hydrological modelling, and provide Penman-Monteith potential evapotranspiration parameterised for short grass, with and without a correction for interception on days with rainfall. The potential evapotranspiration calculations have been formulated to closely follow the methodology of the existing Meteorological Office Rainfall and Evaporation Calculation System (MORECS) potential evapotranspiration, which has historically been widely used by hydrological modellers in the United Kingdom. The two datasets have been created using the same methodology, to allow seamless modelling from past to future. Hydro-PE HadUK-Grid shows good agreement with MORECS in much of the United Kingdom, although Hydro-PE HadUK-Grid is higher in the mountainous regions of Scotland and Wales. This is due to differences in the underlying meteorology, in particular the wind speed, which are themselves due to the different spatial scales of the data. Hydro-PE HadUK-Grid can be downloaded from <https://doi.org/10.5285/9275ab7e-6e93-42bc-8e72-59c98d409deb> (Brown et al., 2022) and Hydro-PE UKCP18 RCM can be downloaded from <https://doi.org/10.5285/eb5d9dc4-13bb-44c7-9bf8-c5980fcf52a4> (Robinson et al., 2021).

1 Introduction

Evaporation is an important part of the hydrological cycle. Globally it is estimated that around 60% of the precipitation that falls on the land is returned to the atmosphere by evaporation from the land (Abbott et al., 2019). However, evaporation is difficult to observe, particularly over large areas, so estimates of potential evaporation (PE) can instead be derived from observed meteorology. PE is an estimate of the evaporative demand of the atmosphere, given an assumed land cover, and is a measure of how much evaporation would occur under specific meteorological conditions, given an unlimited water supply in the soil. In hydrological and crop modelling the actual evaporation (AE) from the land is calculated using potential evaporation (PE) as an estimate of the unlimited evaporation flux, scaled by a function of the soil wetness (Federer et al., 1996). Therefore PE is an

essential driving input for hydrological models, and accurate estimation of PE is essential for model performance, particularly in regions where rainfall is not limiting (Kay et al., 2013).

25 Evaporation from the land can include evaporation from the water contained in the soil, either through transpiration (evaporation through plant stomata) or evaporation from the soil surface directly, and evaporation from open water surfaces (either from permanent water bodies such as lakes, or from transient water on the surface of vegetation or ponding). In this paper we refer to the potential evaporation from the soil as potential evapotranspiration (PET), which may include both transpiration and evaporation from the bare soil. We refer to PE products that also include evaporation from water intercepted by the vegetation canopy as potential evapotranspiration with interception (PETI). We use potential evaporation (PE) as a catch-all term for both of these. The units of PE can be given as a mass flux, $\text{kg m}^{-2} \text{d}^{-1}$, or the equivalent depth of water, mm d^{-1} (assuming water density of 1000 kg m^{-3}). These are numerically equivalent. In this paper, we use mm d^{-1} .

There are a variety of formulations of PE, which rely on various combinations of meteorological inputs. The most parsimonious use just one (eg Thornthwaite, 1948; Oudin et al., 2005) or two (eg Blaney and Criddle, 1950; Priestley and Taylor, 35 1972) meteorological variables, but rely on empirical factors which are calibrated to historical or present-day climate. Different formulations of PE can show different trends under changing climate, adding to the uncertainty of climate change impacts on hydrology (Lemaitre-Basset et al., 2022). The more complex Penman-Monteith formulation (Monteith, 1965), while requiring a larger number of observed inputs, is derived from fundamental physics, so is able to represent temporal changes in drivers. Additionally, PE is also influenced by physical and physiological properties of the land surface, for example leaf area, plant 40 height, albedo and emissivity. These properties are implicitly part of the empirical parameterisation of the simpler models of PE, but can be explicitly included in the calculation of Penman-Monteith PE. Since both the meteorology and the properties of the land surface can be expected to change in the future, Penman-Monteith is most suitable as an input for hydrological modelling in a changing climate (Lemaitre-Basset et al., 2022).

Under climate change, the warming of the atmosphere is expected to intensify the hydrological cycle, with a warmer atmosphere able to hold more water leading to increased evaporation and precipitation (Trenberth, 1999). Alongside this, the rise 45 in atmospheric CO_2 concentrations is expected to lead to stomatal closure in plants, which would lead to increased water use efficiency and decreased transpiration per unit leaf area (Cao et al., 2010). However, it may also lead to increased plant growth, which would mean more transpiring leaf surface per unit area of ground (Alo and Wang, 2008). These are several possibly compensating effects, which can all be captured by the Penman-Monteith equation (Donohue et al., 2010).

50 The estimation of PE is highly dependent on land use, so any PE must be quoted for a specific parameterisation of the land cover and vegetation. To standardise this, the concept of a “reference crop” was introduced, usually short grass as recommended by the United Nations Food and Agricultural Organisation (FAO) (Pereira et al., 1999). Although some PE products are provided for different land covers, for many products a short grass parameterisation is assumed.

In the temperate maritime climate of the United Kingdom (UK), land evaporation is estimated to be around 40% of land 55 precipitation, based on observed river flow (Hannaford, 2015) and modelled runoff and evaporation (Blyth et al., 2019). It can be as high as 60% in the water limited regions of England, and as low as 30% in the cooler, wetter regions of Scotland and Wales (Blyth et al., 2019). Widely used PE datasets for hydrological modelling in the UK include: MORECS PE, a 40 km

gridded product for various land covers (Hough et al., 1995); MOSES, a 5 km gridded product calculated by the MOSES land surface model (Cox et al., 1998; Smith et al., 2006); and CHESS-PE, a 1 km gridded product for time-invariant short grass (Robinson et al., 2017) which has been aggregated to catchment scale as a component of the CAMELS-GB dataset (Coxon et al., 2020b, a). The FAO reference evapotranspiration method provides a formulation for calculating PE using local observed data, including a variety of adaptations to accommodate differing levels of data availability (Allen et al., 1998). MORECS PE calculated for short grass is the most widely used in hydrological modelling in the UK, so is considered a reference dataset throughout this paper. A comparison of PE products can be seen in Table 1.

The MORECS PE dataset is provided by the UK Met Office, for near real time and historical hydrological and crop modelling in the UK (Hough et al., 1995; Hough and Jones, 1997). It is a gridded estimate of weekly and monthly mean PE at 40 km resolution, derived from the MORECS 40 km gridded meteorological dataset. The gridded meteorology is interpolated from a network of meteorological observation stations before being used to calculate PE. It has been widely used as an input for hydrological models, both for research and operationally (Kay et al., 2013). MORECS PE is calculated for a reference short-grass as well as several other land covers. Some hydrological models use the short grass MORECS PE and apply an adjustment for known land cover (eg CLASSIC, Crooks and Naden, 2007). MORECS is purely observation-based, and has no corollary for studies of hydrology under future climate. It is also only available at a much lower resolution (40 km) than the typical resolution (1 km) of hydrological models (Bell et al., 2009; Crooks and Naden, 2007) and available rainfall datasets (Met Office et al., 2021; Tanguy et al., 2021; Lewis et al., 2022) in the UK. Since potential evapotranspiration is a non-linear function of the meteorology, using a lower resolution PE potentially misses important effects of local topography.

MORECS PE uses the Penman-Monteith formulation (Hough et al., 1995), but with some modifications. It includes a correction for the assumption that the surface temperature is equal to the air temperature, uses monthly-varying physiological parameters (leaf area index and stomatal resistance), and implements a correction on rain days to account for interception. This latter interception correction is important for driving models which do not explicitly calculate interception. Since interception is more efficient than transpiration, this combined potential evapotranspiration with interception (PETI) is higher than PET alone. The difference can be of the order of 10% (Robinson et al., 2017).

For hydrological modelling under future climates, PE can be calculated using climate model output in place of meteorological observations (eg Rudd and Kay, 2016). The current state of the art of climate modelling for the UK is UKCP18 (Lowe et al., 2018), which provides several strands of climate projections from global to regional. A high-resolution regional climate model (RCM), nested in a lower resolution global climate model (GCM) has been run as a perturbed parameter ensemble, providing twelve realisations of future climate for the UK at 12 km resolution, for 1980–2080 (Murphy et al., 2018), hereafter referred to as UKCP18 RCM.

In this paper we present two new potential evaporation datasets calculated using historical gridded observed meteorology and future climate projections over the UK: Hydro-PE HadUK-Grid and Hydro-PE UKCP18 RCM. The datasets consist of both PET and PETI calculated using methodology derived from MORECS. The parameterisation used for both has been chosen to be as similar to MORECS as possible, for consistency with existing historical modelling. The historical dataset, Hydro-PE HadUK-Grid, was calculated from the HadUK-Grid observation-based 1 km gridded dataset (Met Office et al., 2021; Hollis

et al., 2019). The future dataset, Hydro-PE UKCP18 RCM, was calculated from the 12 km RCM output from UKCP18 (Murphy et al., 2018). The combination of the two datasets enables seamless modelling from past to future climate projections.

95 In Section 2 we describe the input datasets for the historical calculations (Section 2.1) and the future calculations (Section 2.2). In Section 3 we describe the calculation of the historical PE (Section 3.2) and the future PE (Section 3.3), and in Section 4 we present summaries of each. In Section 5 we evaluate the two products against historical PE datasets, and we discuss the results in Section 6.

2 Input data

100 2.1 HadUK-Grid

HadUK-Grid is a historical gridded meteorological dataset derived from station observations interpolated to a 1 km grid (Hollis et al., 2019), using multiple regression combined with inverse-distance-weighted interpolation, to account for local geographic and topographic factors (Perry and Hollis, 2005). It has been published as a companion to the UKCP18 climate projections. Variables are available at a monthly timestep, and selected variables are also available daily. The start date of each variable is
105 dependent on the availability of station data. The earliest available data are monthly rainfall values from 1862, and coverage of variables increases with time, so all of the variables are available from 1969 onwards. This study used the data at 1 km resolution, using v1.0.3.0 for 1969-2020 inclusive, and v1.1.0.0 for 2021. There are no differences between the two versions for 1969-2020 for the variables used. It is also available at 5 km, 12 km, 25 km and 60 km resolutions. HadUK-Grid is representative of the meteorology at the centre of each grid box at the grid box centre elevation, rather than providing grid box
110 mean meteorology.

The HadUK-Grid daily climate variables used were:

- Maximum air temperature measured between 0900 UTC on day D and 0900 UTC on day D+1 ($t_{\text{asmax}}, T_{\text{max}}, ^\circ\text{C}$),
- Minimum air temperature measured between 0900 UTC on day D-1 and 0900 UTC on day D ($t_{\text{asmin}}, T_{\text{min}}, ^\circ\text{C}$),
- Daily total precipitation amount measured between 0900 UTC on day D and 0900 on day D+1 ($\text{rainfall}, P, \text{mm}$
115 d^{-1}).

The monthly climate variables used were:

- Duration of bright sunshine during the month ($s_{\text{un}}, t_S, \text{hours}$),
- Average of hourly mean wind speed at a height of 10 m above ground level over the month ($s_{\text{fcWind}}, u_{10}, \text{m s}^{-1}$),
- Average of hourly mean sea level pressure over the month ($p_{\text{sl}}, p_{\text{sl}}, \text{hPa}$),
- 120 – Average of hourly vapour pressure over the month ($p_{\text{v}}, e, \text{hPa}$).

2.2 UKCP18 RCM

The UKCP18 Regional Projections on a 12km grid over the UK for 1980–2080, v20190731 (Met Office Hadley Centre, 2018), are a perturbed parameter ensemble of regional climate model (RCM) output for the years 1980–2080, provided at various time resolutions from daily to decadal. The RCM was run on a rotated pole grid but the outputs are also available
125 regridded onto a 12 km resolution grid aligned with the British National Grid (OSGB36); the latter data are used here. The domain covers the UK and surrounding waters and includes a small part of northern France. The first ensemble member (EM) 01 uses the default parameterisation of the Hadley Centre climate model GC3.05 ([HadGEM3-GC3.05](#)) for the GCM ([Sexton et al., 2021](#); [Yamazaki et al., 2021](#)), and a regional version of this for the nested RCM ~~–The~~ ([Murphy et al., 2018](#)). ~~The~~ [HadGEM3-GC3.05 configuration is very similar to HadGEM3-GC3.1 \(Williams et al., 2018\), which was used for Met Office](#)
130 [contributions to the sixth phase of the IPCC’s Coupled Model Intercomparison Project \(CMIP6; Eyring et al., 2016; O’Neill et al., 2016\)](#), ~~except for some differences in the atmosphere model and the sea-ice model (Yamazaki et al., 2021).~~ The other ensemble members have had perturbations applied to several of the parameters ([Sexton et al., 2021](#)), within reasonable ranges ~~–in order to approximate the ensemble spread seen in~~ [informed by the fifth phase of the IPCC’s Coupled Model Intercomparison Project \(CMIP5; Taylor et al., 2012\).](#) ~~The HadGEM3-GC3.1 model has a relatively high climate sensitivity (Andrews et al., 2019), so~~
135 ~~the UKCP18 ensemble range of climate sensitivity is high compared to CMIP5~~ ([Murphy et al., 2018](#)). ~~All~~, but is consistent with the move to higher climate sensitivities in CMIP6 ([Andrews et al., 2019](#)). All UKCP18 ensemble members used the same emissions scenario: Representative Concentration Pathway 8.5 (RCP8.5), which is a high emissions scenario with no target for climate change mitigation ([Riahi et al., 2011](#)). ~~However, in order to provide a wider range of possible scenarios, each of the ensemble members was run with a different CO₂ concentration trajectory. Each of these are consistent with RCP8.5, but represent uncertainty in the emissions scenarios (Murphy et al., 2018). Details of the trajectories are given in Section 2.3.~~

The calculations were carried out on daily mean variables, only for grid boxes that were modelled as land in the RCM. The variables used for both PET and PETI were

- Daily mean specific humidity at 1.5 m (h_{uss}, q_a , kg kg^{-1}),
- Daily mean sea level pressure (p_{sl}, p_{sl} , hPa),
- 145 – Daily mean net surface short wave flux (r_{ss}, S_n , W m^{-2}),
- Daily mean net surface long wave flux (r_{ls}, L_n , W m^{-2}),
- Daily mean wind speed at 10 m ($sfcWind, u_{10}$, m s^{-1}),
- Daily mean air temperature at 1.5 m (t_{as}, T_a , $^{\circ}\text{C}$).

The calculation of PETI additionally used:

- 150 – Daily precipitation rate (p_r, P , mm d^{-1}).

2.3 CO₂ concentration

Future climate projections also include the global atmospheric CO₂ concentration. Although derived from the same emissions scenario, the atmospheric CO₂ concentration pathways used by the Met Office as input to the UKCP18 RCM runs were different for each ensemble member, to reflect global carbon cycle uncertainties (Murphy et al., 2018). Ensemble member 01 used the concentrations prescribed in RCP8.5 for concentration driven runs. The other ensemble members used CO₂ concentrations that were calculated by selected CMIP5 emissions-driven ensemble members. These had different future trajectories, resulting in CO₂ concentrations in 2080 ranging from 708 to 920 ppm across the ensemble (Murphy et al., 2018). These trajectories are shown in Fig. 1. The values of CO₂ were provided as annual values (Met Office Hadley Centre, 2020a).

2.4 Elevation

The surface elevation data used in the calculation of HadUK-Grid was the EU-DEM v1.1 (European Environment Agency, 2021), aggregated from 25m spatial resolution to 100m, then linearly interpolated to the centre point of each 1km grid box, following the method used in the generation of the HadUK-Grid 1km dataset (Hollis et al., 2019).

For the UKCP18 RCM ensemble, the climate model was run using elevation derived from EU-DEM v1.1 (European Environment Agency, 2021), smoothed appropriately for use as the lower boundary condition of the atmospheric model. After the model was run, the elevation was then regridded to the 12 km grid aligned with OSGB for distribution with the regridded climate variables (Met Office Hadley Centre, 2019) - this is the elevation used in this study.

3 Methods

Potential evapotranspiration, E_p (mm d⁻¹), was calculated using the Penman-Monteith equation (Monteith, 1965), derived in terms of specific humidity (Stewart, 1989), see Appendix A for details. The Penman-Monteith equation estimates the evaporation from an extensive vegetation-covered surface with unlimited water supply. It has several parameters which characterise the surface, including the roughness and the stomatal resistance of transpiration, and which are dependent on the type of vegetation that is present.

Historically, the idea of a reference crop – a hypothetical well-watered short grass – has been used for estimating PE (Allen et al., 1998). Although MORECS can be distributed for several land cover types, the most widely used for hydrological modelling in the UK is the short grass PE (eg Bell et al., 2009; Rudd and Kay, 2016). This is appropriate since short grass and similar short vegetation is one of the most widespread land cover types in the UK (Morton et al., 2021). For consistency with existing hydrological modelling, we thus calculated PE for a short grass surface. The surface parameters were chosen to match the short grass parameterisation of MORECS v2.0, details are given in Appendix E. The grass has a canopy height of 0.12 m. The leaf area index (LAI) and stomatal resistance vary by month and the monthly values of the parameters are given in Table 2.

180 In order to have consistent modelling from past to future, we applied the same methods to both HadUK-Grid and UKCP18 RCM meteorological data as far as possible. However, differences between the variables and temporal resolution of the two datasets engender some differences in the calculation procedures, which are noted below.

The calculations were carried out with daily mean variables. This differs from MORECS v2.0, who carried out separate calculations for day-time and night-time (Hough et al., 1995). This was not possible with either the HadUK-Grid or the
185 UKCP18 RCM data, as they do not provide enough information about the diurnal cycle. However tests with other datasets showed that the difference between whole day calculations and separate day-time and night-time calculations is negligible.

3.1 Interception correction

For these datasets, we first calculated Penman-Monteith PET. We then calculated PETI, by applying an interception correction on rain days. We used the methodology of MORECS v2.0, again using the parameters of a short grass to estimate the amount
190 of rainfall which is intercepted by the canopy (Hough et al., 1995). This was done by calculating the potential interception (PEI, mm d^{-1}), which is the rate of evaporation from water intercepted by the canopy. This is subject to the same aerodynamic resistance as PET, but is not limited by stomatal resistance, so was calculated by setting the canopy resistance r_s to be zero in the Penman-Monteith equation. The PETI was calculated as a combination of PET and PEI, dependent on how much water was intercepted by the canopy each day. The interception was dependent on the amount of precipitation and the LAI. A monthly
195 enhancement factor was applied to account for the different characteristics of rainfall in different months (see Table 2). Details of the calculations are in Appendix B.

3.2 Calculation of historical potential evaporation: Hydro-PE HadUK-Grid

The HadUK-Grid dataset does not provide exactly the variables required as input for the Penman-Monteith equation, so these were derived from the existing variables. First, all variables that were only available monthly — the sunshine hours, wind speed,
200 vapour pressure and sea-level air pressure — were interpolated to a daily timestep. These were then used in combination with the existing daily variables to calculate the appropriate input variables. Details are in Appendix C, and an overview of the interpolation procedure is shown in Fig. 2.

Note that, since we do not have an exact estimate of net radiation from HadUK-Grid, we use a modified form of the Penman-Monteith equation, which includes a correction for the radiative transfer between the surface and the screen height (Eq. A2).

205 3.3 Calculation of future potential evaporation: Hydro-PE UKCP18 RCM

The Penman-Monteith calculations were carried out using the daily mean values of the output from the climate model (see Appendix D). In this case, UKCP18 provides net short- and longwave radiation, so the unmodified Penman-Monteith equation was used (Eq. A1). In order to account for rising levels of atmospheric CO_2 , a fertilisation effect was applied to the stomatal resistance following the method of Kruijt et al. (2008) (see Appendix E5 for details).

210 Since the UKCP18 output is provided for all land and sea points in the domain, but PET and PETI are only valid over land, we only carried out the calculations for land points. The land points were selected using the land-sea mask provided by the Met Office (Met Office Hadley Centre, 2020b), which defines grid boxes to be either 100% sea or 100% land. There are some grid boxes which, although they are classified as sea in the RCM, do actually contain a small fraction of land. This is particularly important for hydrological models which run by disaggregating the meteorological inputs to a higher spatial resolution for
215 calculating river flows etc. In order to allow the PET and PETI to be used as input to such hydrological models, these grid boxes were filled with valid data. To do this, a mapping was created between each grid box which needed to be filled and the nearest comparable land grid box. Then the PET and PETI were copied from the existing land grid boxes to the target grid boxes. This was done rather than calculating the PET and PETI with the existing meteorology in the RCM output, because the meteorology over the sea would be unrepresentative of land.

220 4 Results

Maps of Hydro-PE HadUK-Grid PET and PETI can be seen in Fig. 3. There is a strong north-west to south-east gradient, with low values in the west of Scotland and high values in south-east England, which follows the climate of the regions from colder and wetter in the north-west to warmer and dryer in the south-east. Maps of Hydro-PE UKCP18 RCM PET and PETI can be seen in Figs. 4 and 5 respectively. Each panel shows the overall mean PET and PETI for each ensemble member. While there
225 is a range across the ensemble, reflecting the range in ensemble meteorology, there is a consistent spatial pattern across the ensemble, which is also consistent with the HadUK-Grid PET and PETI maps (Fig. 3).

Time-series of annual mean PET and PETI are shown in Fig. 6 for Hydro-PE HadUK-Grid and the Hydro-PE UKCP18 RCM ensemble. The mean monthly climatology of PET and PETI are shown in Fig. 7 for the first twenty years of the UKCP18 RCM ensemble (1980–2000) and the last twenty years (2060–2080). The Hydro-PE UKCP18 RCM ensemble is consistent
230 with Hydro-PE HadUK-Grid in the historical period, although Hydro-PE HadUK-Grid has an increase in PET and PETI in May, followed by a levelling off in June, that is not seen in Hydro-PE UKCP18 RCM. This is due differences in the seasonality of the input meteorology (see Section 5). Overall both PET and PETI increase over the course of the future projections, with PET increasing by 16% – 29% and PETI increasing by 14% – 25%. The largest increases are in the summer (21% – 36% for PET and 18% – 30% for PETI). Increases are more moderate in the winter and some ensemble members individually show a
235 decrease in PET and PETI for the winter months.

Maps of the difference between PETI and PET (as a percentage of PET) are shown in Fig. 8 for Hydro-PE UKCP18 RCM and in the right hand panel of Fig. 3 for Hydro-PE HadUK-Grid. The effect of including interception increases the PE estimate by 5-10% in low lying areas of south-east England, but has a larger increase of up to 35% in highland regions of Scotland. This is because the highlands of Scotland have relatively low evaporative demand (because they are cooler and wetter than other
240 regions), but larger amounts of rainfall than the rest of the country.

The mean monthly climatology of the difference between PETI and PET can be seen in Fig. 9. The mean value of the interception correction is largest in the summer months (0.21–0.32 mm d⁻¹) and lowest in the winter months (0.10–0.14 mm

d^{-1}) in the historical period of Hydro-PE UKCP18 RCM, although as the overall PET is lower in the winter this leads to a larger relative difference in the winter than the summer. This is consistent with Hydro-PE HadUK-Grid, which has mean
245 interception correction of 0.23–0.27 $mm\ d^{-1}$ for the summer months and 0.11–0.12 $mm\ d^{-1}$ for the winter months.

In the future the interception correction decreases in the summer months by 14% (to 0.16–0.30 $mm\ d^{-1}$) and increases in the winter by 8% (to 0.08–0.18 $mm\ d^{-1}$), leading to little change at the annual scale. The decrease in summer interception correction is why the relative increase in PETI is smaller than that of PET. The peak in the absolute difference between PETI and PET is shifted to earlier in the year (from June–September in the historical period to March–June at the end of the projections).
250 This may contribute to changing seasonality of river flows under future climates.

The Hydro-PE UKCP18 RCM interception correction is consistent with the Hydro-PE HadUK-Grid [interception correction](#) throughout the year. However, the ensemble mean is higher than Hydro-PE HadUK-Grid for most of the year, but lower in autumn. This is consistent with the model biases in the precipitation, as the model simulations overall have a higher precipitation than HadUK-Grid for most of the year, but slightly lower precipitation from August to October (Fig. 10).

255 Both variables show an increasing trend [in the mean](#) through the historical period and climate projections. The future projections of PET and PETI both increase by 0.29 $mm\ d^{-1}$ overall between 1980–2000 and 2060–2080, which is 22% of PET or 19% of PETI. The increase varies across the ensemble between 0.23 $mm\ d^{-1}$ and 0.38 $mm\ d^{-1}$. The relative increase ranges between 16 – 29% of PET, and 15 – 25% of PETI. ~~There~~ ~~However,~~ ~~there~~ is little change in the winter values ($< 0.1\ mm\ d^{-1}$), but in the summer both PET and PETI increase by around 0.7 $mm\ d^{-1}$ overall (ranging between 0.5 – 1.0 $mm\ d^{-1}$ across the
260 ensemble). ~~The increases in PET~~

[The 90th percentile values for each month in Hydro-PE HadUK-Grid and PETI-Hydro-PE UKCP18 RCM are shown in Fig. 11. The Hydro-PE UKCP18 RCM ensemble is consistent with Hydro-PE HadUK-Grid between June-September, but for the rest of the year the Hydro-PE HadUK-Grid 90th percentile is higher than Hydro-PE UKCP18 RCM. This is an artefact of using monthly HadUK-Grid inputs with smooth interpolation to daily combined with the calculations applied to obtain the
265 input variables from the available HadUK-Grid variables \(Section C\) causing the 90th percentiles in Hydro-PE HadUK-Grid to be high, rather than Hydro-PE UKCP18 RCM being too low \(see Sections 5.1 and 5.4 for further discussion of this\).](#)

[Similarly to the mean, there is little change in the Hydro-PE UKCP18 RCM 90th percentile values in winter, with some ensemble members showing a small decrease \(up to \$-0.10\ mm\ d^{-1}\$ for both \$PET_{90}\$ and \$PETI_{90}\$ \), and some a small increase \(up to \$0.17\ mm\ d^{-1}\$ for both \$PET_{90}\$ and \$PETI_{90}\$ \). The increases in the 90th percentiles in the summer are much larger, and are
270 larger than the increase in the mean, with \$PET_{90}\$ increasing between \$0.50 - 1.54\ mm\ d^{-1}\$ and \$PETI_{90}\$ increasing between \$0.47 - 1.47\ mm\ d^{-1}\$.](#)

[The increases in mean PET and PETI and in the 90th percentiles](#) are driven by increasing temperature, decreasing relative humidity, and increasing solar radiation (due to decreasing cloud cover) in the climate projections (Murphy et al., 2018). Increases in PETI are mitigated by projected decreases in rainfall in the summer (Murphy et al., 2018), so that the relative
275 contribution of the interception correction falls from 10% of summer PET (14% of annual PET) to 7% of summer PET (12% of annual) by the end of the projections.

5 Evaluation

5.1 Comparison with spatial PE datasets across Great Britain

As the Hydro-PE method has been designed to calculate PETI to be comparable with MORECS PE for short grass, we have compared Hydro-PE HadUK-Grid and Hydro-PE UKCP18 RCM with MORECS. We have also compared it with the existing CHES-PE PET and PETI datasets. We have calculated mean-monthly climatologies for the common period 1981-2017, and for Great Britain (GB) only, as this is the land area that is common to all four datasets. These can all be seen in Fig. 12. Both Hydro-PE HadUK-Grid PET and the ensemble mean of Hydro-PE UKCP18 RCM PET are lower than MORECS PE, which is as expected (because of the inclusion of the interception correction in MORECS), but Hydro-PE UKCP18 RCM PETI is higher than MORECS PE in the summer, and Hydro-PE HadUK-Grid PETI is higher than MORECS throughout the year. CHES-PE PET and PETI are both lower than MORECS, due to differences in the PE calculations and parameterisations. (The related FAO reference crop evaporation (Pereira et al., 1999) is a PET calculation that does not consider interception, so it will give a low estimate of PE when used for hydrology, compared to PETI.) The 90th percentiles calculated over 1980-2000 for GB are in Fig. 13. The Hydro-PE UKCP18 RCM values are consistent with CHES-PE throughout the year, while the Hydro-PE HadUK-Grid 90th percentiles are higher than CHES-PE, particularly between October - April. This is due to the temporal interpolation from monthly to daily combined with the assumptions used to calculate the required inputs from the available HadUK-Grid variables (Section C). This is further discussed in Section 5.4.

The differences between the four datasets are due to either: a) differences in the input meteorology; or b) differences in methodology. Since CHES-met is derived from MORECS meteorology, we expect the differences between MORECS PE and CHES-PE PETI to be due to methodological differences. However, since we have developed the Hydro-PE methodology to be as similar to MORECS as possible, the differences between Hydro-PE and MORECS PE should be due to differences between the meteorology in MORECS, HadUK-Grid and the UKCP18 RCM, including differences in the calculation of the input variables from the available data.

Since HadUK-Grid and MORECS meteorology are derived from the same network of station observations (although not exactly the same stations), and CHES-met is derived directly from MORECS, we might expect that the meteorology averaged over a region to be very similar in all three datasets. Indeed, there is good agreement in GB mean air temperature and precipitation between HadUK-Grid, MORECS and CHES-met (Fig. 10), although MORECS air temperature is a little lower in the winter and MORECS precipitation is a little higher in the summer - this is likely due to the slightly different spatial coverage of the different resolutions. However, the different methods of downscaling of some variables from stations or MORECS to the 1 km grid can introduce differences between datasets. Most notably, the GB mean HadUK-Grid wind speeds are much higher than all of the other datasets. This is due to very high wind speeds at high elevations in HadUK-Grid, particularly in Scotland, due to the elevation adjustment used in the HadUK-Grid calculations (Hollis et al., 2019). Figure 14 shows that in Scotland the HadUK-Grid wind speeds are much higher than the other observational datasets, while in England they are all comparable. Since MORECS is representative of a hypothetical site at mean grid box elevation (rather than mean grid box meteorology), and is a lower resolution, it does not represent these high wind speeds. The CHES wind speed corrections were

applied to the MORECS wind speed assuming that it does represent mean grid box wind speed, so the CHESSE corrections are also not able to reproduce high wind speeds. The HadUK-Grid wind speeds were adjusted based on topographic relationships, without assuming a preservation of the mean (Hollis et al., 2019). The high values of Hydro-PE HadUK-Grid are likely more representative of high elevations.

315 Other variables that are used in the calculation of PE are not directly observed, but have been derived from the variables provided by the station observations. There are differences in the methodology between HadUK-Grid, MORECS and CHESSE, which lead to some differences in inputs to the PE calculations (Fig. 10). The specific humidity of HadUK-Grid is lower than MORECS in the summer, which leads to a higher humidity deficit (as the air temperature is nearly the same). The net shortwave radiation is very similar between HadUK-Grid and MORECS, as they are both calculated using the same methodology. CHESSE-
320 met uses the same method to calculate downward shortwave, but adjusts for slope and aspect, and uses a different albedo parameterisation, so the CHESSE-met net short wave is slightly higher. The approximate total available energy (calculated with R_{ne} , Eq. C8) calculated from HadUK-Grid is very similar to that calculated from MORECS meteorology. In CHESSE-met the approximate available energy is higher, because the net radiation is calculated differently and because CHESSE does not include a ground heat flux. Overall, it is the combination of the higher wind speed and lower specific humidity in HadUK-Grid that
325 leads to the higher PETI in Hydro-PE HadUK-Grid than MORECS.

In addition to the differences in the means, the differences in methodologies have also had an impact on the extremes of the datasets. In particular the Hydro-PE HadUK-Grid 90th percentiles are relatively high due to a combination of the radiation calculations (Section 5.4) and the temporal interpolation of the monthly input variables. The CHESSE-PE 90th percentiles are also relatively high compared to the mean values in winter, which is likely to also be due to the use of the same short-wave radiation calculation, but the effect is not as large as for Hydro-PE HadUK-Grid as the CHESSE input variables are all daily. It was not possible to investigate the MORECS percentiles, as MORECS was only available as monthly means.

330

The UKCP18 variables have been obtained from a free-running climate model, which has not had any bias-correction applied. This means that although the climate model has been calibrated and parameterised against historical meteorological data, it is not expected to match the observed climate exactly. This may be down to deficiencies in the model structure and/or
335 calibration, but is also due to large scale climate variability. The UKCP18 RCM reproduces the historical air temperature well, although the ensemble is slightly cooler than the observations in the spring (Fig. 10). But it significantly overestimates precipitation through most of the year compared to MORECS, and slightly underestimates precipitation from August to October. This is likely due to the global circulation in the coarser resolution GCM within which the RCM is nested bringing more moisture to western Europe (Murphy et al., 2018). This results in a larger interception correction than is seen in the historical Hydro-PE
340 HadUK-Grid for much of the year.

Again, the UKCP18 RCM has a lower specific humidity than MORECS (Fig. 10), which in the spring and summer leads to a higher humidity deficit. The wind speed is very similar between the climate model and the observations, but the net shortwave radiation is much higher in the summer and autumn in the climate model. This is likely to be due to a lower amount of cloud in the model, as well as partly due to different albedo values. In particular, the climate model is run with a realistic land
345 cover, while the net short-wave radiation is calculated for all the other PE datasets assuming short grass everywhere. Overall

the available energy (calculated with the actual net radiation R_n , Eq.D1) calculated with UKCEP18 RCM is higher than that calculated with either HadUK-Grid or CHES-met. It cannot be compared to MORECS, because we do not have actual net radiation for MORECS, but since we can see that MORECS and HadUK-Grid have very similar approximate available energy, R_{ne} , we can infer that the UKCP18 RCM actual available energy is also higher than MORECS. Therefore, the high values of Hydro-PE UKCP18 RCM PETI in the summer are likely to be due to a combination of lower humidity and much higher available energy than in MORECS. The fact that the high May values of PETI seen in Hydro-PE HadUK-Grid and MORECS are not seen in Hydro-PE UKCP18 RCM is due to the seasonal cycle of temperature, humidity and available energy being shifted later in the year in the climate model than in the observations.

So although there are differences between the Hydro-PE products and MORECS in the historical period, these are due to understood differences in the input meteorology.

5.2 Comparison with daily MORECS site data

The Hydro-PE HadUK-Grid PETI data was further compared to daily MORECS PE data for 16 sites across GB, available for the period 01/01/1985 - 31/12/1992. The MORECS PE data are derived from observed station data at each site. These are compared with the HadUK-Grid PETI from the 1 km grid box that contains each site (see Table 3). We expect some differences between the MORECS PE and the Hydro-PE HadUK-Grid PETI due to the difference in location, and therefore meteorology, between the MORECS sites and the HadUK-Grid 1 km grid box centres.

Three metrics were calculated to evaluate daily Hydro-PE HadUK-Grid PE relative to daily MORECS PE; a) bias calculated as the difference in mean daily PE values annually and seasonally, b) difference in the standard deviation in daily PE, c) Pearson correlation coefficient of daily PE values and of deseasonalised daily PE values. The results are presented in Figure 15, with boxplots summarising results over the 16 sites (Figure 15a-c) and maps showing results at the site locations (Figure 15d-f).

The HadUK-Grid PETI is higher throughout the year than MORECS PE, shown by the positive biases across all sites and seasons (Figure 15a and d). This echoes the results from the GB-wide comparison in Figure 12. HadUK-Grid PETI also tends to have slightly higher variation than MORECS PE, shown by a higher standard deviation in PE time-series for 9 out of the 16 sites. Correlation between HadUK-Grid PETI and MORECS PE is generally good across all the sites, with Pearson correlation coefficients in the range 0.72 - 0.81. The high Pearson correlation coefficients are partly due to the fact that we represent the seasonal cycle well. With deseasonalised data, the Pearson correlation coefficients are in the range 0.30-0.47, which still shows a reasonable correlation with the MORECS PE. Even though the need for temporal interpolation of some HadUK-Grid variables from monthly to daily is likely to suppress the daily variability to some extent, the daily variability of the calculated PETI is still comparable to observations.

5.3 Comparison with eddy-covariance measurements

While PET and PETI are not directly observable quantities, they are an estimate of unconstrained evapotranspiration. In GB in the winter, spring and autumn evaporation rates are low and precipitation is high, so it can be assumed that most evaporation will be occurring at the potential rate. Thus we can compare eddy-covariance (EC) flux measurements of actual evaporation

(AE, mm d^{-1}) with PETI calculated from the meteorological measurements at each EC site using the Hydro-PE methodology (EC site PETI). We also compare this with the Hydro-PE HadUK-Grid PETI from the 1 km grid box containing each site with the EC site PETI.

We used measurements of AE and meteorology from two EC sites in England:

Berkshire Organic Grassland (BD-OG) Located on an organically managed grassland in the Berkshire downs, at an elevation of 184 m above sea level (Evans et al., 2016b). EC measurements are available from 01/01/2017 to 16/11/2019 (Morrison et al., 2019).

East Anglia Fens Grassland (EF-GF) Located on a managed grassland in a lowland peatland environment in the East Anglian Fens, at an elevation of 1 m below sea level (Evans et al., 2016a). EC measurements are available from 27/04/2017 to 31/03/2019 (Morrison et al., 2020).

Details of the site locations and the corresponding HadUK-Grid 1 km grid boxes are shown in Table 3.

The top two panels of Fig. 16 show the time-series of EC site PETI and the measured AE. The PETI is similar to the AE, except in the summer (JJA) when the AE drops below the PETI due to soil moisture limitation. The top row of Fig. 17 shows the EC site PETI plotted against measured AE (excluding JJA values) at each site. There is a near-one-to-one linear fit between the PETI and the AE at both sites, although there is a small negative bias, which may be due to differences between the sites and the idealised short grass used for the PETI parameterisation. In particular, EG-GF is sited on peat soils, which are not accounted for in the parameterisation. The difference in the standard deviation is also small. The Pearson correlation is very high at 0.92 for BD-OG and 0.90 for EF-GF. Thus it can be concluded that, at the EC sites, the PETI calculated with observed meteorology is representative of the observed AE during times of no water stress.

The top two panels of Fig. 16 also show the time-series of Hydro-PE HadUK-Grid PETI from the grid box that contains each EC site. The HadUK-Grid PETI compares well with the EC site PETI (Fig. 17, bottom row), with Pearson correlations of 0.78 and 0.86, similar to the comparison with daily MORECS site PE. For BD-OG there is a small positive bias, while for EG-GF there is a small negative bias. The latter is likely due to the low wind speed in HadUK-Grid compared to the wind speed observed at the EC site (Fig. 16 bottom row). There may also be an effect of the peat soil, which is not represented by the MORECS parameterisation.

The standard deviation of the Hydro-PE HadUK-Grid PETI is lower than the standard deviation of the EC site PETI, but the difference is small. There is a lack of daily variability in the temporally interpolated variables, which may reduce the variability of the Hydro-PE HadUK-Grid PETI. Figure 16 (bottom row) shows this for the wind speed – the HadUK-Grid wind speed has a much lower variability than the observed, but it replicates the monthly variability well. The bias in wind speed for EF-GF is likely due to the spatial interpolation of the HadUK-Grid dataset (Hollis et al., 2019).

Despite the differences in input meteorology due to a) temporal and spatial interpolation of the HadUK-Grid data and b) the spatial offset between the 1 km grid box centres and the location of the EC sites, the Hydro-PE HadUK-Grid PETI is a good approximation for EC site PETI. This shows that the use of daily data interpolated from monthly inputs has not introduced significant biases to the PETI calculation.

5.4 Evaluation of calculated meteorology

415 In order to evaluate the impact of temporal interpolation and the calculations applied to the HadUK-Grid input data we used
the PLUMBER2 dataset (Ukkola, 2020), which consists of gap-filled and quality controlled observational data from 170 global
flux sites. We used the half-hourly meteorological variables from the 33 grassland sites in the northern hemisphere. We first
applied the Hydro-PE calculations directly to daily means of the observed variables: precipitation, air temperature, specific
humidity, wind speed, air pressure and net radiation. We then used the half-hourly variables to calculate the equivalents of the
HadUK-Grid variables: daily minimum and maximum air temperature were calculated from the half-hourly air temperature;
420 sunshine hours were calculated from the half-hourly downward shortwave radiation, using the World Meteorological Organisation
threshold of 120 W m^{-2} ; and vapour pressure was calculated from specific humidity and air pressure by inverting Eq.A5. The
sunshine hours, vapour pressure, wind speed and surface air pressure were then averaged to monthly. We used these monthly
variables, plus the daily precipitation and the derived daily minimum and maximum air temperatures in the same calculations
as for the HadUK-Grid meteorology and performed the same interpolation to daily (Appendix C). Finally, we applied the
425 Hydro-PE calculations to these interpolated and derived daily variables.

We compared the PETI calculated using derived and interpolated variables with the PETI calculated using the original daily
mean variables using three metrics: the bias, the error in the standard deviation of the daily values, and the Pearson correlation
coefficient of the raw daily values and of deseasonalised daily values (Fig. 18). The PETI calculated using the interpolated
derived meteorology is overall higher than that calculated from the daily mean, although for some sites it is lower, with the bias
430 being between $-0.12 - 1.01 \text{ mm/d}$ across the sites. The largest difference is in the summer which has bias between $-0.26 - 1.65$
 mm/d . The errors in the standard deviation are roughly evenly distributed, with 19 sites having higher standard deviation and
14 sites lower. The Pearson correlation coefficient is between $0.75-0.94$ calculated with the raw data. With the deseasonalised
data it has a larger range, being between $0.14-0.70$; it is greater than 0.5 for 19 sites. Results were very similar for PET.

We also compared the monthly median and 10th and 90th percentiles of PETI for each site calculated a) using the daily
435 mean meteorology, b) using the meteorology derived and interpolated following the methods applied to HadUK-Grid and c)
using the monthly mean meteorology (Fig. 19). The median is well represented, as are the summer 90th percentiles and the
winter 10th percentiles. However, the PETI calculated using derived and interpolated input variables tends to overestimate
the 90th percentile in the autumn and winter, and overestimate the 10th percentile in the spring and summer compared to
using the original daily means. However, despite this, it does perform better than simply using monthly mean inputs, which
440 is equivalent to the use of MORECS monthly PE and which overestimates the 10th percentile and underestimates the 90th
percentile throughout the year. Most of the variability in the PETI calculated using derived and interpolated variables is due to
the use of the daily air temperature and precipitation, compared to using monthly means of all variables. There is a negligible
difference between using the actual daily mean air temperature and the approximation using daily minimum and maximum
temperatures (Eq. C1).

445 One source of the overall positive bias is the derived and interpolated net radiation, which has a high bias compared to
the daily PLUMBER2 net radiation values, with mean bias error ranging between $3.39 - 51.9 \text{ W m}^{-2}$ across the sites. This

is largely due to a high bias introduced in spring and summer by the calculation of shortwave radiation from sunshine hours (Section C6), which also causes the 10th percentiles of net radiation in the summer and 90th percentiles of net radiation outside of the summer to be higher than those of the daily PLUMBER2 data. The temporal interpolation further enhances this impact on the extremes, due to the effect of the reduced variability on the complex interplay between the different physical variables.

6 Discussion

These Hydro-PE datasets have been calculated with daily data. The Hydro-PE UKCP18 RCM data are self-consistent at a daily timestep, having been calculated from climate model output. Care should be taken with Hydro-PE HadUK-Grid however, as many of the input variables have been temporally downscaled from monthly to daily using a simple smooth interpolation. The variables have been interpolated independently of each other. The ~~Hydro-PE HadUK-Grid can be used at a daily timestep, although the daily values may not be completely consistent with PE calculated from observed daily data, due to small inconsistencies introduced by the temporal downscaling.~~ However, the monthly means of Hydro-PE HadUK-Grid are consistent with the monthly meteorology, and with monthly MORECS. The daily Hydro-PE HadUK-Grid is also consistent with daily MORECS PE (Section 5.2), and with daily EC site PETI (Section 5.3). However, there are some minor inconsistencies introduced by the temporal downscaling and conversion from the provided HadUK-Grid variables to the required inputs. In particular, this has caused an overestimation of winter and spring 90th percentiles. For applications sensitive to the winter and spring high extremes, users may consider winsorizing the data when using at a daily timestep.

Past studies have seen that good results can be obtained by calculating PET at a monthly timescale (Allen et al., 1998; Oudin et al., 2010), and hydrological modelling in the UK is often carried out using monthly PE, for example monthly MORECS is used to drive Grid-to-Grid (Bell et al., 2009) or CLASSIC (Crooks and Naden, 2007). For models which require daily PE inputs, using the temporally interpolated variables is an improvement on using monthly PE data, particularly as Hydro-PE HadUK-Grid uses daily (not interpolated) mean air temperature and precipitation.

Hydro-PE UKCP18 RCM has been calculated from an ensemble of climate model output, which provides a range of possible future scenarios. The UKCP18 RCM ensemble only considers a single high emissions scenario, RCP8.5 (Riahi et al., 2011), but the realisation of that scenario is different for each ensemble member due to differences in the input CO₂ concentration pathways as well as the differences in the parameterisation of the climate model. This encapsulates not only the uncertainty in the global climate response to increased CO₂ concentrations, but the uncertainty of conversion of increased emissions to increased CO₂ concentrations (Murphy et al., 2018). RCP8.5 is the highest of the four Representative Concentration Pathways (RCPs), which were developed by the climate modelling community after the Fourth Assessment Report (AR4; IPCC, 2007) to provide input to climate models and explore a range of emissions scenarios. The range of CO₂ concentration pathways used in the UKCP18 RCM ensemble was designed to approximate the spread of outcomes in the UKCP18 probabilistic projections, which included all four RCPs (Murphy et al., 2018). Hydro-PE UKCP18 RCM shows a large increase in both PET and PETI from the start to the end of the ~~data set~~dataset, alongside a change in the seasonality of the interception correction. Overall these changes would be likely to be smaller under less extreme future emissions scenarios. Although the overall Hydro-PE

480 UKCP18 RCM ensemble has large future changes, the lower end of the range of these changes may be consistent with more moderate future scenarios.

For users of climate model outputs, eg CMIP6 or UKCP18, calculating PET or PETI at daily timestep can involve large volumes of data, or require variables that are unavailable at the daily timestep. For PET, it is reasonable in this case to perform the calculations with monthly data (Oudin et al., 2010). However PETI, which is corrected for interception based on precipitation, 485 cannot reliably be calculated at a monthly timestep. This is because at a daily timestep, the interception correction is only applied to rain days; dry days remain equal to PET. If a monthly mean precipitation is used to calculate the interception correction to monthly PET, then this is the equivalent of applying the interception correction to all days in the month, thus overestimating the PETI. One may calculate PET and PEI at the monthly timestep, but it is essential to apply the interception correction based on the number of wet and dry days in the month. This can either be done by using daily precipitation to combine monthly PET 490 and PEI appropriately, or using the number of wet and dry days in a month in combination with monthly precipitation and monthly PET and PEI, depending on the data available.

These datasets have been calculated only for one surface: short grass. This was chosen as it is a standard that is widely used. However, hydrology may be better represented by PE calculated for realistic land use. This could be done by parameterising the PE calculations appropriately for several different vegetation and other land use types, then combining appropriately for 495 the given surface. The Hydro-PE calculations can easily be reparameterised using vegetation characteristics. Equally, although these Hydro-PE datasets have been specifically parameterised and created for the UK, the method is flexible and can be parameterised and applied to meteorological data globally.

Apart from the effect of CO₂ on stomatal resistance, all of the other parameters have been kept constant throughout the period of future calculations, in particular the LAI, following Rudd and Kay (2016). While increased CO₂ can be expected 500 to lead to increased biomass production, it is likely that the resultant effect on LAI, and the effect of the LAI on evaporation is actually small (Bunce, 2004), either due to the associated increase in shading, or due to nutrient limitation (Ainsworth and Long, 2005). Additionally, the short grass cover that is being modelled here is similar to managed grasslands, which are likely to be artificially limited in LAI and height by human intervention. However, for other vegetation types, the effect of rising CO₂ may well have an impact on both the magnitude and the seasonality of physiological parameters including LAI and canopy 505 height. Future work should include this effect.

7 Conclusions

We have demonstrated two new potential evaporation datasets, for consistent historical and future hydrological modelling. They have been calculated using a method consistent with the widely used MORECS PE dataset and are parameterised for compatibility with models that have been calibrated to MORECS PE. Hydro-PE HadUK-Grid allows historical studies and calibration 510 of hydrological models, while Hydro-PE UKCP18 RCM enables hydrological modelling to be carried out consistently into the future accounting for climate change. [This methodology is available to be applied to further climate model datasets, such as](#)

[the convection-permitting model strand of UKCP18 \(Kendon et al., 2021\) and EuroCORDEX-UK \(Barnes, 2023\), which will enable consistent hydrological modelling across a range of climate projections.](#)

8 Code and data availability

515 The data can be downloaded from the Environmental Information Data Centre in netCDF format. Hydro-PE HadUK-Grid is available at <https://doi.org/10.5285/9275ab7e-6e93-42bc-8e72-59c98d409deb> (Brown et al., 2022) and Hydro-PE UKCP18 RCM is available at <https://doi.org/10.5285/eb5d9dc4-13bb-44c7-9bf8-c5980fcf52a4> (Robinson et al., 2021).

The python code used to perform the Hydro-PE calculations is available at <https://github.com/NERC-CEH/hydro-pe>.

Appendix A: Penman-Monteith equation

520 Potential evapotranspiration, E_p (mm d⁻¹), is calculated using the Penman-Monteith equation (Monteith, 1965), derived in terms of specific humidity (Stewart, 1989),

$$E_P = \frac{t_D}{\lambda} \frac{\Delta_q (R_n - G) + \frac{\rho_a c_p}{r_a} (q_s - q_a)}{\Delta_q + \gamma_q \left(1 + \frac{r_s}{r_a}\right)}, \quad (\text{A1})$$

where $t_D = 86400$ s d⁻¹ is the length of a day, $\lambda = 2.5 \times 10^{-6}$ J kg⁻¹ is the latent heat of vaporisation of water, q_s (kg kg⁻¹) is specific humidity at saturation, Δ_q (kg kg⁻¹ K⁻¹) is the gradient of specific humidity at saturation with respect to temperature, R_n (W m⁻²) is net radiation, G (W m⁻²) is ground heat flux, $c_p = 1010$ J kg⁻¹ K⁻¹ is the specific heat capacity of air, ρ_a (kg m⁻³) is the density of air, q_a (kg kg⁻¹) is the specific humidity of air, $\gamma_q = c_p/\lambda = 0.004$ K⁻¹ is the psychrometric constant for specific humidity, r_s (s m⁻¹) is canopy resistance and r_a (s m⁻¹) is aerodynamic resistance.

Calculations for specific humidity at saturation and its gradient are given in Sections A2 and A3. The ground heat flux, canopy resistance and aerodynamic resistance are calculated following MORECS, see Section E2 – E4.

530 A1 Modified Penman-Monteith calculation

Since the meteorological variables provided in HadUK-Grid do not include net radiation values or surface temperature, we calculate the net radiation assuming that surface temperature T_* can be approximated by air temperature T_a . We then correct for this by calculating PE with a modified form of the Penman-Monteith equation, as used in MORECS v2.0 (Hough et al., 1995),

$$535 E_P = \frac{t_D}{\lambda} \frac{\Delta_q (R_{ne} - G) + \frac{\rho_a c_p}{r_a} (q_s - q_a) \left(1 + \frac{b_R r_a}{\rho_a c_p}\right)}{\Delta_q + \gamma_q \left(1 + \frac{r_s}{r_a}\right) \left(1 + \frac{b_R r_a}{\rho_a c_p}\right)}, \quad (\text{A2})$$

where R_{ne} is the net radiation calculated using T_a as a proxy for surface temperature (see Section C8) and

$$b_R = 4\epsilon\sigma_B T_a \quad (\text{A3})$$

is a correction for the radiative transfer between the surface and the screen height, with $\epsilon = 0.95$ the assumed surface emissivity (as in MORECS (Hough et al., 1995)) and $\sigma_B = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ the Stefan-Boltzmann constant.

540 A2 Specific humidity at saturation

Saturated specific humidity, q_s , is derived as a function of temperature from the empirical fit of saturated vapour pressure, e_s , as a function of temperature (Richards, 1971),

$$e_s = p_{sp} \exp \left(\sum_{i=1}^4 a_i \left(1 - \frac{T_{sp}}{T_a} \right)^i \right), \quad (\text{A4})$$

where $p_{sp} = 101325 \text{ Pa}$ is the steam point pressure, $T_{sp} = 373.15 \text{ K}$ is the steam point temperature, and $a = (13.3185, -0.9760, -0.6445, -0.1299)$ are empirical coefficients.

In general, specific humidity, q_a , can be calculated from vapour pressure, e_a (hPa), using the function

$$q_a = \frac{\epsilon_a e_a}{p_* - (1 - \epsilon_a) e_a}, \quad (\text{A5})$$

where p_* is surface air pressure and $\epsilon_a = 0.622$ is the mass ratio of water to dry air (Gill, 1982).

Substituting the saturated vapour pressure from Eq A4 into the conversion between vapour pressure and specific humidity in Eq. A5, gives an empirical function for saturated specific humidity of

$$q_s = \frac{\epsilon_a p_{sp} \exp \left(\sum_{i=1}^4 a_i \left(1 - \frac{T_{sp}}{T_a} \right)^i \right)}{p_* - (1 - \epsilon_a) p_{sp} \exp \left(\sum_{i=1}^4 a_i \left(1 - \frac{T_{sp}}{T_a} \right)^i \right)}. \quad (\text{A6})$$

A3 Derivative of specific humidity at saturation

The derivative of q_s with respect to air temperature, Δ_q ($\text{kg kg}^{-1} \text{ K}^{-1}$), can be calculated analytically from Eq. A6 and is given by

$$\Delta_q = \frac{T_{sp}}{T_a^2} \frac{q_s}{\epsilon_a} \left(\epsilon_a + (1 - \epsilon_a) q_s \right) \sum_{i=1}^4 a_i i \left(1 - \frac{T_{sp}}{T_a} \right)^{i-1}. \quad (\text{A7})$$

or

$$\Delta_q = \frac{T_{sp}}{T_a} \frac{p_* q_s}{p_* - e_s (1 - \epsilon_a)} \sum_{i=1}^4 i a_i \left(1 - \frac{T_{sp}}{T_a} \right)^{i-1}. \quad (\text{A8})$$

Appendix B: Interception correction

The PETI, E_{PI} (mm d^{-1}) is first calculated in the same way as PET. Then, on days with non-zero precipitation, an interception correction is applied. This is required because water that is intercepted by the canopy will evaporate at a faster rate than water

that is transpired. Again, this is implemented following the MORECS v2.0 calculations (Hough et al., 1995). These calculations treat all precipitation as rainfall, and do not allow for lying snow.

The amount of rainfall intercepted by the canopy, C_I (mm d⁻¹), is

$$C_I = \begin{cases} P f_P e_P & \text{if } P f_P < C_{\max} \\ C_{\max} e_P & \text{otherwise,} \end{cases} \quad (\text{B1})$$

565 where $f_P = 1 - 0.5^\Lambda$ is the LAI-dependent fraction of rainfall that would be intercepted if there were only one rain event in the day, $C_{\max} = 0.2\Lambda$ (mm) is the maximum capacity of the canopy to hold water, and e_P is an enhancement factor to allow for the different character of rainfall throughout the year. In the winter, e_P is 1, but it rises to 2 in the summer when the rain is likely to fall in multiple shorter, more intense, events. The monthly values of e_P are given in Table 2.

In a rain day, the intercepted fraction of rainfall is assumed to evaporate as an open water surface. This potential interception, 570 E_I (mm d⁻¹) is calculated using Eq. A1 with $r_s = 0$ (all other parameters have the same value as for PET). The time taken to dry the canopy at this rate, t_{dry} (d), is calculated to be

$$t_{\text{dry}} = \frac{C_I}{E_I}, \quad (\text{B2})$$

If this potential interception is enough to dry out the canopy in less than a day, then for the rest of the day then it reverts to the rate of PET (E_P), such that

$$575 \quad E_{PI} = t_{\text{dry}} E_I + (1 - t_{\text{dry}}) E_P. \quad (\text{B3})$$

If it is not enough to dry out the canopy, then the PETI is equal to the potential interception, E_I all day. The water remaining on the canopy is not carried over to the following day. By combining all these cases, and combining Eq. B2 and Eq. B3, the PETI is given by

$$E_{PI} = \begin{cases} E_P & \text{if } P = 0 \\ E_P + C_I \left(1 - \frac{E_P}{E_I}\right) & \text{if } P > 0 \text{ and } C_I < E_I \\ E_I & \text{if } P > 0 \text{ and } C_I \geq E_I. \end{cases} \quad (\text{B4})$$

580 **Appendix C: Calculation of input variables for Hydro-PE HadUK-Grid**

C1 Temporal interpolation of meteorology

The sunshine hours, wind speed, vapour pressure and sea-level air pressure were interpolated from a monthly to daily timestep using quadratic interpolation from python's scipy package (Virtanen et al., 2020), assuming that the monthly values represent the 15th of each month. The sunshine hours were divided by the number of days in each month to get an average daily value, 585 before interpolation. We constrained sunshine hours and vapour pressure to always be positive, to avoid some unphysical

negative values that were a result of extrapolation of the quadratic interpolation at either end of the time series. Following the temporal interpolation, the required input variables were calculated from the interpolated and existing daily variables as follows.

C2 Daily mean air temperature

590 We approximate the daily mean air temperature from the daily minimum and maximum air temperature using

$$T_a = \frac{T_{\min} + T_{\max}}{2}. \quad (\text{C1})$$

C3 Surface air pressure

Both the specific humidity at saturation and its derivative are a function of both air temperature and surface air pressure at the grid box elevation. As HadUK-Grid only provides sea level air pressure, p_{SL} (hPa), we must calculate the air pressure at the
 595 grid box elevation. The sea level air pressure is adjusted to the grid box elevation by assuming a constant local environmental lapse rate for air temperature of $\Gamma_T = -0.006 \text{ K m}^{-1}$ (as used in Hough et al., 1995; Robinson et al., 2017). The surface air pressure at a given elevation, p_* (hPa), is calculated using the integral of the hypsometric equation (Shuttleworth, 2012) so that

$$p_* = p_{\text{SL}} \left(\frac{T_a - z\Gamma_T}{T_a} \right)^{\frac{g}{r\Gamma_T}}, \quad (\text{C2})$$

600 where T_a (K) is the air temperature at the grid box elevation, z (m) is the grid box elevation above sea level, $g = 9.81 \text{ m s}^{-2}$ is acceleration due to gravity, $r = 287.05 \text{ J kg}^{-1} \text{ K}^{-1}$ is the specific gas constant of dry air.

C4 Air density

The density of air, ρ_a was calculated from the surface air pressure and the air temperature, such that

$$\rho_a = \frac{p_*}{rT_a}, \quad (\text{C3})$$

605 where $r = 287.05 \text{ J kg}^{-1} \text{ K}^{-1}$ is the gas constant of dry air.

C5 Specific humidity

The specific humidity is calculated using the water vapour pressure (e_a) and surface pressure (p_*) using Equation A5.

C6 Net shortwave surface radiation

Downward shortwave radiation, S_d (W m^{-2}), is calculated from sunshine hours following the MORECS procedure, using
 610 Eqs. 4.14 to 4.19 of Hough et al. (1995). Equation 4.14 yields the total radiation integrated over the day (W hr m^{-2}), so to get the daily average, this was divided by 24, the number of hours in a day, such that:

$$S_d = \frac{1}{24} R_a \left(\eta \left(a_A + b_A \frac{t_S}{t_N} \right) + c_A (1 - \eta) \right) \quad (\text{C4})$$

where a_A , b_A and c_A are the Ångström coefficients, R_a is the top-of-atmosphere radiation integrated over a day (W hr m^{-2}), t_S is measured hours of bright sunshine in the day, t_N is the length of daylight hours (sunrise to sunset) (Hough et al., 1995),
615 and η is

$$\eta = \begin{cases} 0 & \text{if } t_S = 0 \\ 1 & \text{if } t_S > 0. \end{cases} \quad (\text{C5})$$

The Ångström Coefficients are empirical constants that are used to calculate solar radiation from sunshine hours. They were calculated on the MORECS 40km grid by Cowley (1978). For this study they were then interpolated on to the 1km HadUK grid using a smooth bivariate spline. They were only available for England, Scotland and Wales, and so the coefficients for Northern
620 Ireland were spatially extrapolated. They are close to the ‘default’ values suggested for use by the FAO where local calibration is not available (see Eq. 35 in Allen et al. (1998)). a_A and c_A are constant in time, whereas b_A varies by month. Here, the b_A coefficient is interpolated on to a daily timestep using a periodic cubic spline (periodic so that there is not a discontinuity between the values for 31st Dec and 1st Jan).

R_a and t_N are calculated as

$$625 \quad R_a = t_S \left((t_2 - t_1) \sin \delta \sin \phi + \frac{12 \cos \delta \cos \phi \left(\sin \frac{\pi t_1}{12} - \sin \frac{\pi t_2}{12} \right)}{\pi} \right) \quad (\text{C6})$$

and

$$t_N = t_2 - t_1, \quad (\text{C7})$$

where t_1 and t_2 are sunrise and sunset times, ϕ is the latitude and δ is the solar declination angle calculated from the day of the year d using using

$$630 \quad \delta = 0.41 \cos 2\pi \frac{d - 172}{365}. \quad (\text{C8})$$

Sunrise, t_1 (hours), is calculated using

$$t_1 = \frac{12}{\pi} \arccos \left(\tan \delta \tan \phi + \frac{0.0145}{\cos \delta \cos \phi} \right), \quad (\text{C9})$$

and sunset, t_2 (hours), is given by

$$t_2 = 24 - t_1. \quad (\text{C10})$$

635 Net shortwave radiation is calculated from downward shortwave radiation using albedo, α , as

$$S_n = (1 - \alpha) S_d. \quad (\text{C11})$$

The values of albedo used are described in Section E1.

C7 Estimate of net longwave surface radiation

The calculation of L_{ne} followed MORECS, documented in more detail in Hough et al. (1995), using air temperature, vapour
640 pressure and sunshine hours via

$$L_{ne} = \epsilon \sigma_B T_a^4 \left(1.28 \left(\frac{e}{T_a} \right)^{\frac{1}{7}} - 1 \right) \left(0.2 + 0.8 \frac{t_S}{t_N} \right). \quad (\text{C12})$$

Note that this is only an approximation of net longwave radiation as it has made use of an approximation when calculating the upward component of the radiation, L_u . Normally, upward longwave radiation is calculated from surface temperature

$$L_u = \epsilon \sigma_B T_*^4. \quad (\text{C13})$$

645 However, if surface temperature is not available, then an approximate upward longwave radiation, L_{ue} can be calculated by substituting $T_a \approx T_*$ in Eq. C13. This approximation has been carried through to the approximate net longwave radiation in Eq. C12.

The approximate net longwave radiation is related to the actual net longwave radiation, L_n , by the relationship

$$L_{ne} = L_n - b_R (T_a - T_*), \quad (\text{C14})$$

650 where b_R is given by Eq. A3.

C8 Approximate net radiation

After deriving the approximate net longwave radiation and net solar radiation the approximate net radiation, R_{ne} (W m^{-2}), is simply calculated by summing these two components:

$$R_{ne} = L_{ne} + S_n. \quad (\text{C15})$$

655 That this is an approximation means that we must use a modified form of the Penman-Monteith equation, described in Section A1.

Appendix D: Calculation of input variables for Hydro-PE UKCP18 RCM

D1 Net radiation

The UKCP18 dataset contains net longwave radiation, L_n (W m^{-2}), and net shortwave radiation, S_n (W m^{-2}). Therefore the
660 total net radiation is given by

$$R_n = L_n + S_n. \quad (\text{D1})$$

D2 Surface air pressure

The daily pressure at mean sea level is adjusted to the pressure at the grid box surface height following Eq. C2.

D3 Air density

665 The air density was calculated from surface air pressure and air temperature using Eq. C3.

Appendix E: Surface parameters

E1 Albedo

The albedo α is also calculated following MORECS as a combination of the albedo of the grass, α_g , and the albedo of the underlying soil, α_s

$$670 \quad \alpha = \begin{cases} \alpha_g & \text{if } \Lambda > 4 \\ \alpha_s + 0.25\Lambda(\alpha_g - \alpha_s) & \text{otherwise} \end{cases} \quad (\text{E1})$$

where Λ is the LAI (Hough et al., 1995).

The albedo of grass is taken to be the MORECS value for soils with median available water content (AWC) $\alpha_g = 0.25$. Soil albedo is dependent on whether it is wet or dry; for wet soils $\alpha_s = 0.1$, and for dry soils $\alpha_s = 0.2$. A soil is considered wet if there has been precipitation in the day, dry otherwise. MORECS also provides values for soils with high AWC and low AWC,
675 but these were not used for this calculation, as soil properties are not considered.

E2 Ground heat flux

For the ground heat flux, the average monthly value of ground heat storage as used in MORECS v2.0 was used (see Table 2). This is a monthly mean of daily heat storage estimated from measurements of soil temperature (Wales-Smith and Arnott, 1980)

~~The monthly value is used for~~ was converted to a daily flux, and applied to all days in that month.

680 E3 Aerodynamic resistance

Aerodynamic resistance r_a (sm^{-1}) is calculated as a function of the roughness length of the canopy z_0 (h), using the MORECS equation

$$r_a = \frac{6.25}{u_{10}} \ln \frac{10}{z_0} \ln \frac{6}{z_0}, \quad (\text{E2})$$

where u_{10} is the wind speed at 10 m above the canopy and

$$685 \quad z_0 = \frac{h}{10}, \quad (\text{E3})$$

where h is the canopy height of grass. Following MORECS v2.0, grass is assumed to be 0.15 m high.

E4 Canopy resistance

The canopy resistance is a combination of the stomatal resistance of the canopy, r_{sc} (m s^{-1}), and the surface resistance of a bare soil surface, r_{ss} (m s^{-1}), following MORECS v2.0 (Hough et al., 1995), such that

$$690 \quad \frac{1}{r_s} = \frac{1 - 0.7^\Lambda}{r_{sc}} + \frac{0.7^\Lambda}{r_{ss}}, \quad (\text{E4})$$

where Λ is the LAI of the grass surface. The LAI was assumed to vary throughout the year, and monthly values are taken from MORECS v2.0 (see Table 2). This seasonal cycle of LAI was kept the same through the full time coverage of the calculations.

E5 CO₂ fertilisation effect

The stomatal resistance was also assumed to vary monthly through the year. In addition, it was assumed that stomatal resistance will increase in the future, due to projected rising CO₂ levels (Rudd and Kay, 2016). The baseline historical stomatal resistance of grass, r_{scM} (m s^{-1}), was taken from MORECS v2.0 (Hough et al., 1995) (see Table 2). For years up to the baseline year of 1981, the stomatal resistance was assumed to stay constant. From 1981 onwards the stomatal resistance was then adjusted for changes in CO₂ concentration, following the method of (Kruijt et al., 2008), so that for each month, m , and year, y , the stomatal resistance is

$$700 \quad r_{sc}[y, m] = \begin{cases} r_{scM}[m] & \text{if } y \leq 1981 \\ \frac{r_{scM}[m]}{1 - 0.00093 (\text{CO}_2[y] - \text{CO}_2[1981])} & \text{if } y > 1981, \end{cases} \quad (\text{E5})$$

where $\text{CO}_2[y]$ is the CO₂ concentration in ppmv in the year y and 1981 is the reference year. As each ensemble member of the UKCP18 RCM ensemble was run with a different CO₂ concentration (Murphy et al., 2018), this adjustment of the stomatal resistance was calculated separately for each ensemble member, using the corresponding CO₂ trajectory (Met Office Hadley Centre, 2020a).

705 The surface resistance of bare soil was set to 100 m s^{-1} , again following MORECS v2.0 (Hough et al., 1995).

Author contributions. ELR, ALK, RC, VAB and EMB designed the Hydro-PE algorithm. ELR wrote the Hydro-PE code. MJB designed and wrote the code to temporally interpolate the HadUK-Grid monthly data. MJB created the Hydro-PE HadUK-Grid dataset. ELR created the Hydro-PE UKCP18 dataset. RAL performed evaluation. All authors contributed to the manuscript.

Competing interests. The authors declare no competing interests.

710 *Acknowledgements.* This work was supported by the Natural Environment Research Council award number NE/S017380/1 as part of the Hydro-JULES programme.

References

- Abbott, B. W., Bishop, K., Zarnetske, J. P., Minaudo, C., Chapin, F., Krause, S., Hannah, D. M., Conner, L., Ellison, D., Godsey, S. E., et al.: Human domination of the global water cycle absent from depictions and perceptions, *Nature Geoscience*, 12, 533–540, 2019.
- 715 Ainsworth, E. A. and Long, S. P.: What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂, *New Phytologist*, 165, 351–372, <https://doi.org/https://doi.org/10.1111/j.1469-8137.2004.01224.x>, 2005.
- Allen, R. G., Pereira, L. S., Raes, D., Smith, M., et al.: Crop evapotranspiration — Guidelines for computing crop water requirements — FAO Irrigation and drainage paper 56, FAO, Rome, 300, D05 109, <https://www.fao.org/3/X0490E/x0490e00.htm>, 1998.
- 720 Alo, C. A. and Wang, G.: Potential future changes of the terrestrial ecosystem based on climate projections by eight general circulation models, *Journal of Geophysical Research: Biogeosciences*, 113, 2008.
- Andrews, T., Andrews, M. B., Bodas-Salcedo, A., Jones, G. S., Kuhlbrodt, T., Manners, J., Menary, M. B., Ridley, J., Ringer, M. A., Sellar, A. A., Senior, C. A., and Tang, Y.: Forcings, Feedbacks, and Climate Sensitivity in HadGEM3-GC3.1 and UKESM1, *Journal of Advances in Modeling Earth Systems*, 11, 4377–4394, <https://doi.org/https://doi.org/10.1029/2019MS001866>, 2019.
- 725 Barnes, C.: EuroCORDEX-UK: Regional climate projections for the UK domain at 12 km Resolution for 1980-2080, <https://doi.org/10.5285/B109BD69E1AF425AA0F661B01C40DC51>, 2023.
- Bell, V., Kay, A., Jones, R., Moore, R., and Reynard, N.: Use of soil data in a grid-based hydrological model to estimate spatial variation in changing flood risk across the UK, *Journal of Hydrology*, 377, 335–350, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2009.08.031>, 2009.
- 730 Blaney, H. F. and Criddle, W. D.: Determining water requirements in irrigated areas from climatological and irrigation data, Technical Paper 96, US Department of Agriculture (Soil Conservation Service), 1950.
- Blyth, E. M., Martinez-de la Torre, A., and Robinson, E. L.: Trends in evapotranspiration and its drivers in Great Britain: 1961 to 2015, *Progress in Physical Geography: Earth and Environment*, 43, 666–693, 2019.
- Brown, M., Robinson, E., Kay, A., Chapman, R., Bell, V., and Blyth, E.: Potential evapotranspiration derived from HadUK-Grid gridded climate observations 1969-2021 (Hydro-PE HadUK-Grid), <https://doi.org/10.5285/9275ab7e-6e93-42bc-8e72-59c98d409deb>, 2022.
- 735 Bunce, J.: Carbon dioxide effects on stomatal responses to the environment and water use by crops under field conditions, *Oecologia*, 140, 1–10, <https://doi.org/10.1007/s00442-003-1401-6>, 2004.
- Cao, L., Bala, G., Caldeira, K., Nemani, R., and Ban-Weiss, G.: Importance of carbon dioxide physiological forcing to future climate change, *Proceedings of the National Academy of Sciences*, 107, 9513–9518, 2010.
- 740 Cowley, J. P.: Distribution over Great-Britain of global solar irradiation on a horizontal surface, *Meteorological Magazine*, 107, 357–373, 1978.
- Cox, P., Huntingford, C., and Harding, R.: A canopy conductance and photosynthesis model for use in a GCM land surface scheme, *Journal of Hydrology*, 212-213, 79–94, [https://doi.org/https://doi.org/10.1016/S0022-1694\(98\)00203-0](https://doi.org/https://doi.org/10.1016/S0022-1694(98)00203-0), 1998.
- Coxon, G., Addor, N., Bloomfield, J., Freer, J., Fry, M., Hannaford, J., Howden, N., Lane, R., Lewis, M., Robinson, E., Wagener, T., and Woods, R.: Catchment attributes and hydro-meteorological timeseries for 671 catchments across Great Britain (CAMELS-GB), <https://doi.org/10.5285/8344e4f3-d2ea-44f5-8afa-86d2987543a9>, 2020a.
- 745

- Coxon, G., Addor, N., Bloomfield, J. P., Freer, J., Fry, M., Hannaford, J., Howden, N. J. K., Lane, R., Lewis, M., Robinson, E. L., Wagener, T., and Woods, R.: CAMELS-GB: hydrometeorological time series and landscape attributes for 671 catchments in Great Britain, *Earth System Science Data*, 12, 2459–2483, <https://doi.org/10.5194/essd-12-2459-2020>, 2020b.
- 750 Crooks, S. M. and Naden, P. S.: CLASSIC: a semi-distributed rainfall–runoff modelling system, *Hydrology and Earth System Sciences*, 11, 516–531, <https://doi.org/10.5194/hess-11-516-2007>, 2007.
- Donohue, R. J., McVicar, T. R., and Roderick, M. L.: Assessing the ability of potential evaporation formulations to capture the dynamics in evaporative demand within a changing climate, *Journal of Hydrology*, 386, 186–197, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2010.03.020>, 2010.
- 755 European Environment Agency: European Digital Elevation Model version 1.1, <http://land.copernicus.eu/pan-european/satellite-derived-products/eu-dem/eu-dem-v1.1/view>, 2021.
- Evans, C., Morrison, R., Burden, A., Williamson, J., Baird, A., Brown, E., Callaghan, N., Chapman, P., Cumming, A., Dean, H., Dixon, S., Dooling, G., Evans, J., Gauci, V., Grayson, R., Haddaway, N., He, Y., Heppell, K., Holden, J., Hughes, S., Kaduk, J., Jones, D., Matthews, R., Menichino, N., Misselbrook, T., Page, S., Pan, G., Peacock, M., Rayment, M., Ridley, L., Robinson, I., Rylett, D., Scowen, M., Stanley, K., and Worrall, F.: Final report on project SP1210: Lowland peatland systems in England and Wales — evaluating greenhouse gas fluxes and carbon balances, http://oro.open.ac.uk/50635/1/14106_Report_FINALDefraLowlandPeatPublished.pdf, 2016a.
- 760 Evans, J. G., Ward, H. C., Blake, J. R., Hewitt, E. J., Morrison, R., Fry, M., Ball, L. A., Doughty, L. C., Libre, J. W., Hitt, O. E., Rylett, D., Ellis, R. J., Warwick, A. C., Brooks, M., Parkes, M. A., Wright, G. M. H., Singer, A. C., Boorman, D. B., and Jenkins, A.: Soil water content in southern England derived from a cosmic-ray soil moisture observing system — COSMOS-UK, *Hydrological Processes*, 30, 4987–4999, <https://doi.org/https://doi.org/10.1002/hyp.10929>, 2016b.
- 765 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Federer, C. A., Vörösmarty, C., and Fekete, B.: Intercomparison of Methods for Calculating Potential Evaporation in Regional and Global Water Balance Models, *Water Resources Research*, 32, 2315–2321, <https://doi.org/https://doi.org/10.1029/96WR00801>, 1996.
- 770 Gill, A. E.: *Atmosphere-ocean Dynamics*, Academic Press, San Diego, CA, USA, 1982.
- Hannaford, J.: Climate-driven changes in UK river flows: A review of the evidence, *Progress in Physical Geography*, 39, 29–48, 2015.
- Hollis, D., McCarthy, M., Kendon, M., Legg, T., and Simpson, I.: HadUK-Grid – A new UK dataset of gridded climate observations, *Geoscience Data Journal*, 6, 151–159, <https://doi.org/https://doi.org/10.1002/gdj3.78>, 2019.
- 775 Hough, M. and Jones, R.: The United Kingdom Meteorological Office rainfall and evaporation calculation system: MORECS version 2.0-an overview, *Hydrology and Earth System Sciences*, 1, 227–239, 1997.
- Hough, M., Palmer, S., Weir, A., Lee, M., and Barrie, I.: The Meteorological Office Rainfall and Evaporation Calculation System: MORECS version 2.0, *Hydrological Memorandum 45*, Met Office, <https://library.metoffice.gov.uk/Portal/Default/en-GB/RecordView/Index/635882>, 1995.
- 780 IPCC: *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, IPCC, Geneva, Switzerland, 2007.
- Kay, A. L., Bell, V. A., Blyth, E. M., Crooks, S. M., Davies, H. N., and Reynard, N. S.: A hydrological perspective on evaporation: historical trends and future projections in Britain, *Journal of Water and Climate Change*, 4, 193–208, <https://doi.org/10.2166/wcc.2013.014>, 2013.

- Kendon, E., Short, C., Pope, J., Chan, S., Wilkinson, J., Tucker, S., Bett, P., and Harris, G.: Update to UKCP Local (2.2km) projections, https://www.metoffice.gov.uk/binaries/content/assets/metofficegovuk/pdf/research/ukcp/ukcp18_factsheet_snow_jul-2021.pdf, 2021.
- 785 Kruijt, B., Witte, J.-P. M., Jacobs, C. M., and Kroon, T.: Effects of rising atmospheric CO₂ on evapotranspiration and soil moisture: A practical approach for the Netherlands, *Journal of Hydrology*, 349, 257 – 267, <https://doi.org/10.1016/j.jhydrol.2007.10.052>, 2008.
- Lemaitre-Basset, T., Oudin, L., Thirel, G., and Collet, L.: Unraveling the contribution of potential evaporation formulation to uncertainty under climate change, *Hydrology and Earth System Sciences*, 26, 2147–2159, <https://doi.org/10.5194/hess-26-2147-2022>, 2022.
- 790 Lewis, E., Quinn, N., Blenkinsop, S., Fowler, H., Freer, J., Tanguy, M., Hitt, O., Coxon, G., Bates, P., Woods, R., Fry, M., Chevuturi, A., Swain, O., and White, S.: Gridded estimates of hourly areal rainfall for Great Britain 1990-2016 [CEH-GEAR1hr] v2, <https://doi.org/10.5285/fc9423d6-3d54-467f-bb2b-fc7357a3941f>, 2022.
- Lowe, J. A., Bernie, D., Bett, P., Bricheno, L., Brown, S., Calvert, D., Clark, R., Eagle, K., Edwards, T., Fosser, G., Fung, F., Gohar, L., Good, P., Gregory, J., Harris, G., Howard, T., Kaye, N., Kendon, E., Krijnen, J., Maisey, P., McDonald, R., McInnes, R., McSweeney, C., 795 Mitchell, J. F. B., Murphy, J., Palmer, M., Roberts, C., Rostron, J., Sexton, D., Thornton, H., Tinker, J., Tucker, S., Yamazaki, K., and Belcher, S.: UKCP18 Science Overview Report, 2018.
- Met Office, Hollis, D., McCarthy, M., Kendon, M., Legg, T., and Simpson, I.: HadUK-Grid gridded and regional average climate observations for the UK, <http://catalogue.ceda.ac.uk/uuid/4dc8450d889a491ebb20e724debe2dfb>, 2021.
- Met Office Hadley Centre: UKCP18 Regional Projections on a 12 km grid over the UK for 1980-2080, <https://catalogue.ceda.ac.uk/uuid/589211abeb844070a95d061c8cc7f604>, Accessed 2021-05-27, 2018.
- 800 Met Office Hadley Centre: UKCP18 land projections - 12km regional climate model, surface altitude (m) over Europe, <https://data.ceda.ac.uk/badc/ukcp18/data/land-rcm/ancil/orog>, Accessed 2020-06-03, 2019.
- Met Office Hadley Centre: UKCP18 land projections - atmospheric CO₂ concentrations, 2020a.
- Met Office Hadley Centre: UKCP18 land projections - land-sea mask, 2020b.
- 805 Monteith, J. L.: Evaporation and environment, in: *Symposia of the society for experimental biology*, vol. 19, pp. 205–234, Cambridge University Press (CUP) Cambridge, 1965.
- Morrison, R., Cooper, H., Cumming, A., Scarlett, P., Thornton, J., and Winterbourn, J.: Eddy covariance measurements of carbon dioxide, energy and water fluxes at an organically managed grassland, Berkshire, UK, 2017 to 2019, <https://doi.org/10.5285/5a93161f-0124-4650-a2c9-7e8aaea7e6bb>, 2019.
- 810 Morrison, R., Cooper, H., Cumming, A., Evans, C., Thornton, J., Winterbourn, J., Rylett, D., and Jones, D.: Eddy covariance measurements of carbon dioxide, energy and water fluxes at a cropland and a grassland on lowland peat soils, East Anglia, UK, 2016-2019, <https://doi.org/10.5285/2fe84b80-117a-4b19-a1f5-71bbd1dba9c9>, 2020.
- Morton, R., Marston, C., O’Neil, A., and Rowland, C.: Land Cover Map 2020 (land parcels, GB), <https://doi.org/10.5285/0e99d57e-1757-451f-ac9d-92fd1256f02a>, 2021.
- 815 Murphy, J. M., Harris, G. R., Sexton, D. M. H., Kendon, E. J., Bett, P. E., Brown, S. J., Clark, R. T., Eagle, K., Fosser, G., Fung, F., Lowe, J. A., McDonald, R. E., McInnes, R. N., McSweeney, C. F., Mitchell, J. F. B., Rostron, J., Thornton, H. E., Tucker, S., and Yamazaki, K.: UKCP18 Land Projections: Science Report, <https://www.metoffice.gov.uk/pub/data/weather/uk/ukcp18/science-reports/UKCP18-Land-report.pdf>, 2018.
- O’Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Krieglner, E., Lamarque, J.-F., Lowe, 820 J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geoscientific Model Development*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.

- Oudin, L., Hervieu, F., Michel, C., Perrin, C., Andréassian, V., Anctil, F., and Loumagne, C.: Which potential evapotranspiration input for a lumped rainfall–runoff model?: Part 2—Towards a simple and efficient potential evapotranspiration model for rainfall–runoff modelling, *Journal of Hydrology*, 303, 290–306, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2004.08.026>, 2005.
- 825 Oudin, L., titia Moulin, L., Bendjoudi, H., and Ribstein, P.: Estimating potential evapotranspiration without continuous daily data: possible errors and impact on water balance simulations, *Hydrological Sciences Journal*, 55, 209–222, <https://doi.org/10.1080/02626660903546118>, 2010.
- Pereira, L. S., Perrier, A., Allen, R. G., and Alves, I.: Evapotranspiration: Concepts and Future Trends, *Journal of Irrigation and Drainage Engineering*, 125, 45–51, [https://doi.org/10.1061/\(ASCE\)0733-9437\(1999\)125:2\(45\)](https://doi.org/10.1061/(ASCE)0733-9437(1999)125:2(45)), 1999.
- 830 Perry, M. and Hollis, D.: The generation of monthly gridded datasets for a range of climatic variables over the UK, *International Journal of Climatology*, 25, 1041–1054, <https://doi.org/https://doi.org/10.1002/joc.1161>, 2005.
- Priestley, C. H. B. and Taylor, R. J.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Monthly weather review*, 100, 81–92, 1972.
- Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N., and Rafaj, P.: RCP 8.5 — A scenario of comparatively high greenhouse gas emissions, *Climatic Change*, 109, 33–57, <https://doi.org/10.1007/s10584-011-0149-y>, 2011.
- 835 Richards, J. M.: A simple expression for the saturation vapour pressure of water in the range -50 to 140°C, *Journal of Physics D: Applied Physics*, 4, L15–L18, <https://doi.org/10.1088/0022-3727/4/4/101>, 1971.
- Robinson, E. L., Blyth, E. M., Clark, D. B., Finch, J., and Rudd, A. C.: Trends in atmospheric evaporative demand in Great Britain using high-resolution meteorological data, *Hydrology and Earth System Sciences*, 21, 1189–1224, <https://doi.org/10.5194/hess-21-1189-2017>,
840 2017.
- Robinson, E. L., Kay, A. L., Brown, M., Chapman, R., Bell, V. A., and Blyth, E.: Potential evapotranspiration derived from the UK Climate Projections 2018 Regional Climate Model ensemble 1980-2080 (Hydro-PE UKCP18 RCM), <https://doi.org/10.5285/eb5d9dc4-13bb-44c7-9bf8-c5980fcf52a4>, 2021.
- Rudd, A. C. and Kay, A. L.: Use of very high resolution climate model data for hydrological modelling: estimation of potential evaporation, *Hydrology Research*, 47, 660–670, <https://doi.org/10.2166/nh.2015.028>, 2016.
- 845 Sexton, D. M. H., McSweeney, C. F., Rostron, J. W., Yamazaki, K., Booth, B. B. B., Murphy, J. M., Regayre, L., Johnson, J. S., and Karmalkar, A. V.: A perturbed parameter ensemble of HadGEM3-GC3.05 coupled model projections: part 1: selecting the parameter combinations, *Climate Dynamics*, 56, 3395–3436, <https://doi.org/10.1007/s00382-021-05709-9>, 2021.
- Shuttleworth, W. J.: *Terrestrial Hydrometeorology*, John Wiley & Sons Ltd, <https://doi.org/10.1002/9781119951933.ch3>, 2012.
- 850 Smith, R. N. B., Blyth, E. M., Finch, J. W., Goodchild, S., Hall, R. L., and Madry, S.: Soil state and surface hydrology diagnosis based on MOSES in the Met Office Nimrod nowcasting system, *Meteorological Applications*, 13, 89–109, <https://doi.org/https://doi.org/10.1017/S1350482705002069>, 2006.
- Stewart, J. B.: On the use of the Penman-Monteith equation for determining areal evapotranspiration, in: *Estimation of Areal Evapotranspiration (Proceedings of a workshop held at Vancouver, B.C., Canada, August 1987)*, IAHS, Wallingford, Oxfordshire, UK, 1989.
- 855 Tanguy, M., Dixon, H., Prodocimi, I., Morris, D., and Keller, V.: Gridded estimates of daily and monthly areal rainfall for the United Kingdom (1890-2019) [CEH-GEAR], <https://doi.org/10.5285/dbf13dd5-90cd-457a-a986-f2f9dd97e93c>, 2021.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485 – 498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Thornthwaite, C. W.: An approach toward a rational classification of climate, *Geographical review*, 38, 55–94, 1948.

- 860 Trenberth, K. E.: Conceptual framework for changes of extremes of the hydrological cycle with climate change, in: *Weather and climate extremes*, pp. 327–339, Springer, 1999.
- Ukkola, A.: PLUMBER2: forcing and evaluation datasets for a model intercomparison project for land surface models v1.0, <https://doi.org/10.25914/5FDB0902607E1>, 2020.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J.,
865 van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Wales-Smith, B. and Arnott, J.: The evaporation calculation system used in the United Kingdom, 1980.
- 870 Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380, <https://doi.org/https://doi.org/10.1002/2017MS001115>, 2018.
- 875 Yamazaki, K., Sexton, D. M. H., Rostron, J. W., McSweeney, C. F., Murphy, J. M., and Harris, G. R.: A perturbed parameter ensemble of HadGEM3-GC3.05 coupled model projections: part 2: global performance and future changes, *Climate Dynamics*, 56, 3437–3471, <https://doi.org/10.1007/s00382-020-05608-5>, 2021.




Figure 1. The global mean atmospheric CO₂ concentrations used for the UKCP18 RCM ensemble (Met Office Hadley Centre, 2020a). The heavy black line shows ensemble member 01, the thin gray lines show the other ensemble members.




Figure 2. The temporal interpolation and calculation procedure applied to the HadUK-Grid meteorology, to create daily inputs for the calculation of Hydro-PE HadUK-Grid.




Figure 3. Annual mean PET (left) and PETI (centre) from Hydro-PE HadUK-Grid (~~1969-2020~~1969-2021). The right hand panel shows the relative difference between PETI and PET as a percentage of PET.




Figure 4. Mean PET for each ensemble member of Hydro-PE UKCP18 RCM over the historical period 1980–2020.




Figure 5. Mean PETI for each ensemble member of Hydro-PE UKCP18 RCM over the historical period 1980–2020.




Figure 6. Time-series of annual mean PET (top) and PETI (bottom). The black line shows ensemble member 01 of Hydro-PE UKCP18 RCM, the grey lines show the other ensemble members, and the blue line shows the ensemble mean. The orange line shows Hydro-PE HadUK-Grid. Note that the regions averaged are slightly different — Hydro-PE UKCP18 RCM averages over the UK, the Republic of Ireland and a small amount of northern France, while Hydro-PE HadUK-Grid only includes the UK.




Figure 7. Mean monthly climatology of PET (top row) and PETI (bottom row). The left hand column shows the mean over the first twenty years of the ensemble, the right hand column shows the mean over the final twenty years. Line styles as in Fig. 6




Figure 8. The difference between annual mean PETI and annual mean PET as a percentage of PET for each ensemble member over the historical period of Hydro-PE UKCP18 RCM (1980–2020).




Figure 9. Mean monthly climatology of the difference between PETI and PET as an absolute value (top row) and as a percentage of PET (bottom row). The left hand column shows the mean over 1980–2000, the right hand column shows the mean over 2060–2080. Line styles as in Fig. 6.




Figure 10. Monthly mean air temperature, specific humidity, wind speed, precipitation, net shortwave radiation and available energy averaged over GB (1981-2017) for MORECS (black line), UKCP18 RCM (blue line shows the ensemble mean and light blue area shows the ensemble range), HadUK-Grid (orange line), and CHES-met (brown dashed line). The MORECS specific humidity, shortwave radiation and available energy have been calculated from the daily MORECS meteorology, so they are an approximation of the full day- and night-time calculations. In the available energy panel, the MORECS, HadUK-Grid and CHES-met lines show the available energy used in the PE calculations, which was calculated using the approximate longwave radiation R_{ne} , which assumes the surface temperature can be approximated with the air temperature (Eq. C12). The dotted orange and brown lines show the available energy with an estimate of the correction from A3 applied, so are comparable with the UKCP18 RCM values, which are calculated using the actual net radiation components output by the climate model.




Figure 11. Monthly 90th percentiles of PET (top row) and PETI (bottom row). The left hand column shows the percentiles calculated over the first twenty years of the ensemble, the right hand column shows them calculated over the final twenty years. Line styles as in Fig. 6.




Figure 12. Monthly mean PET (top) and PETI (bottom) averaged over GB for 1981-2017. The black line in both panels is MORECS PE (which includes the interception correction, so is [equivalent to](#) PETI). The blue line shows the ensemble mean of Hydro-PE UKCP18 RCM, and the light blue area shows the ensemble range. The orange line shows Hydro-PE HadUK-Grid. The brown dashed line shows CHES-PE.




Figure 13. Monthly 90th percentile of PET (top) and PETI (bottom) averaged over GB for 1981-2000. The blue area shows the ensemble range of Hydro-PE UKCP18 RCM, and the black line shows the values for ensemble member 01. The orange line shows Hydro-PE HadUK-Grid. The brown dashed line shows CHES-PE.




Figure 14. Monthly mean PETI (top row) and wind speed (bottom row) averaged over England (left) and Scotland (right) (1981-2017) from MORECS (black line), UKCP18 RCM (blue line shows the ensemble mean and light blue area shows the ensemble range), HadUK-Grid (orange line), and CHES (brown dashed line).




Figure 15. Evaluation of the Hydro-PE HadUK-Grid PETI against daily MORECS PE (which is also PETI) across 16 MORECS sites. Boxplots summarise results over all 16 sites including: a) bias in mean [daily](#) PETI values calculated annually (Ann), over winter (DJF), spring (MAM), summer (JJA) and autumn (SON), b) difference in the standard deviation [of daily vailes](#), c) Pearson correlation coefficient [for the raw daily values \(black\) and the deseasonalised daily values \(grey\)](#). Maps show annual bias, difference in the standard deviation and Pearson correlation coefficient values [for the raw daily data](#) at each of the MORECS sites.




Figure 16. Upper two panels show time series of measured daily AE (black), calculated daily PETI (blue) and HadUK-Grid PETI (orange) for two EC sites (BD-OG and EF-GF). The grey regions show the summer data that are excluded from calculating the metrics. Lower two panels show observed daily wind speed (grey), observed monthly mean wind speed (black) and HadUK-Grid interpolated wind speed (orange) for two EC sites.




Figure 17. The top panels show PETI calculated from site meteorology plotted against observed AE, for values in winter (DJF), spring (MAM) and autumn (SON). The bottom panels show the HadUK-Grid PETI against the EC site PETI. The left-hand column is the BD-OG site, the right-hand column is the EF-GF site. Dashed lines show the least squares linear fit to the data.




Figure 18. Comparison of Hydro-PE PETI calculated using interpolated and derived daily variables and Hydro-PE calculated using daily mean variables for 33 northern hemisphere grassland flux sites in the PLUMBER2 dataset. The left panels show box plots of the annual and seasonal bias (top), error in the standard deviation (middle) and Pearson correlation (bottom) of the raw data (black) and the deseasonalised data (grey). The bars show the whole range of the data. The right hand plots show maps of the sites, coloured by the bias (top), error in the standard deviation (middle) and Pearson correlation of the raw data (bottom).




Figure 19. Monthly boxplots of the 10th, 50th and 90th percentiles of Hydro-PE PETI calculated using the daily PLUMBER2 data, the interpolated and derived PLUMBER2 variables, and the derived PLUMBER2 variables at a monthly timestep.

Table 1. Existing Penman-Monteith PE datasets and methods available in the UK.

Dataset/method name	Assumed land cover	Spatial resolution	Temporal resolution	Interception	CO ₂ response	Ground heat flux
MORECS (Hough and Jones, 1997)	A range of land cover types, with monthly varying parameters. The short grass parameterisation is widely used for hydrological modelling.	40 km	Weekly, monthly (calculated daily)	Yes	No	Yes
CHESS (Robinson et al., 2017)	Short grass with time-invariant parameters (GB only).	1 km	Daily	Both	No	No
MOSES (Smith et al., 2006)	Tiled combination of: broadleaf tree, needleleaf tree, C3 grass, C4 grass, crop, shrub, urban, lake, bare soil.	5 km	Hourly	No	Yes	Yes
FAO Reference Evapotranspiration (Allen et al., 1998)	Short grass with time-invariant parameters.	N/A ^a	Hourly, daily, weekly, monthly	No	No	Yes (except daily)

^aFAO is a method rather than a specific dataset. It can be applied to site or spatial data.

Table 2. Monthly values of LAI, stomatal resistance, daily mean ground heat flux and interception enhancement factor. All values are taken from the MORECS v2.0 documentation and are valid for short grass (Hough et al., 1995).

Month	LAI Λ (-)	Stomatal resistance r_{scM} ($s\ m^{-1}$)	Ground heat flux G ($W\ m^{-2}$)	Interception enhancement factor e_P (-)
January	2.0	80	-5.7	1.0
February	2.0	80	-3.1	1.0
March	3.0	60	1.3	1.2
April	4.0	50	7.0	1.4
May	5.0	40	9.8	1.6
June	5.0	60	10.5	2.0
July	5.0	60	8.9	2.0
August	5.0	70	2.9	2.0
September	4.0	70	-3.5	1.8
October	3.0	70	-8.6	1.4
November	2.5	80	-10.7	1.2
December	2.0	80	-8.6	1.0

Table 3. Sites used for evaluation, and the corresponding HadUK-Grid grid boxes. The top section gives details for MORECS sites, and the bottom section gives details for the eddy-covariance sites. Locations are given in British National Grid, with HadUK-Grid locations and elevations referring to the centre of the selected grid box. Distance (m) and Elevation difference (m) refer to the difference in location/elevation between the sites and the centre point of the HadUK-Grid grid box.

Site name	Site ID	Site location	Site elevation (m)	HadUK-Grid location	HadUK-Grid elevation (m)	Distance (m)	Elevation difference (m)
Kinloss	1057	306700E 862700N	6	306500E 862500N	5	283	1
Galashiels	1939	347900E 636700N	200	347500E 636500N	265	447	-65
Leeming	2245	430600E 489000N	35	430500E 489500N	31	510	4
Marham	3023	573700E 309100N	19	573500E 309500N	21	447	-2
Stansted	3626	553100E 222600N	100	553500E 222500N	100	412	0
Wilsden	4036	408800E 434900N	264	408500E 434500N	276	500	-12
Nottingham	4206	450300E 345600N	116	450500E 345500N	114	224	2
Ludlow	4750	350900E 274600N	108	350500E 274500N	116	412	-8
Shawbury	4757	355300E 322000N	74	355500E 322500N	70	539	4
Preston	4886	356400E 247500N	86	356500E 247500N	80	100	6
Ulcombe	5345	584300E 147500N	45	584500E 147500N	47	200	-2
Lyneham	5848	400600E 178200N	145	400500E 178500N	145	316	0
Eskmeals	7004	308500E 493100N	6	308500E 493500N	11	400	-5
Swansea	8413	264200E 192300N	6	264500E 192500N	10	361	-4
Yeovilton	8673	355100E 123700N	19	355500E 123500N	20	447	1
Okehampton	8825	260500E 91300N	398	260500E 91500N	395	200	3
Berkshire Organic Grassland	BD-OG	435932E 181436N	184	435500E 181500N	190	437	-6
East Anglia Fens Grassland	EF-GF	522934E 287366N	-1	522500E 287500N	0	455	-1