

OceanSODA-UNEXE: A multi-year gridded Amazon and Congo River outflow surface ocean carbonate system dataset

Richard P. Sims¹, Thomas M. Holding², Peter E. Land³, Jean-Francois Piolle⁴, Hannah L. Green^{1,3}, Jamie D. Shutler¹

5 ¹Centre for Geography and Environmental Science, College of Life and Environmental Sciences, University of Exeter, Penryn campus, United Kingdom

²Department of Human Behaviour, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig, 04103, Germany

³Plymouth Marine Laboratory, Plymouth, PL13DH, United Kingdom

10 ⁴Laboratoire d'Océanographie Physique et Spatiale (LOPS), IFREMER, Université of Brest, CNRS, IRD, IUEM, Brest, France

Correspondence to: Richard P. Sims (r.sims2@exeter.ac.uk)

Abstract

15 Large rivers play an important role in transferring water and all of its constituents including carbon in its various forms from the land to the ocean, but the seasonal and inter-annual variations in these riverine flows remain unclear. Satellite Earth observation datasets and reanalysis products can now be used to observe synoptic-scale spatial and temporal variations in the carbonate system within large river outflows. Here we present the OceanSODA-UNEXE time series, a dataset of the full carbonate system in the surface water outflows of the Amazon (2010-2020) and Congo Rivers (2002-2016). Optimal
20 empirical approaches were used to generate gridded Total alkalinity (TA) and dissolved inorganic carbon (DIC) fields in the outflow regions. These combinations were determined by equitably evaluating all combinations of algorithms and inputs against a [reference](#) matchup database of *in situ* observations. Gridded TA and DIC along with gridded temperature and salinity data enable the calculation of the full carbonate system in the surface ocean ([which includes pH and the partial pressure of carbon dioxide, \(\$p\text{CO}_2\$ \)](#)). The algorithm evaluation constitutes a Type A uncertainty evaluation for TA and DIC
25 where model, input and sampling uncertainties are considered. Total combined uncertainties for TA and DIC were propagated through the carbonate system calculation allowing all variables to be provided with an associated uncertainty estimate. In the Amazon outflow, the total combined uncertainty for TA was ~~identified as~~ 36 $\mu\text{mol kg}^{-1}$ (weighted RMSD 35 $\mu\text{mol kg}^{-1}$ and weighted bias 8 $\mu\text{mol kg}^{-1}$ for $n=82$) and for DIC was 44 $\mu\text{mol kg}^{-1}$ (weighted RMSD 44 $\mu\text{mol kg}^{-1}$ and weighted bias -6 $\mu\text{mol kg}^{-1}$ for $n=70$). The spatially averaged propagated [combined](#) uncertainties for the ~~partial pressure of carbon dioxide ($p\text{CO}_2$)~~ and pH are 85 μatm and 0.08 respectively, where the pH uncertainty is relative to an average pH of 8.19. In the Congo outflow, the combined uncertainty for TA was identified as 29 $\mu\text{mol kg}^{-1}$ (weighted RMSD 28 $\mu\text{mol kg}^{-1}$ and weighted bias 6 $\mu\text{mol kg}^{-1}$ for $n=102$) and for DIC was 40 $\mu\text{mol kg}^{-1}$ (weighted RMSD 37 $\mu\text{mol kg}^{-1}$ and weighted bias -16 $\mu\text{mol kg}^{-1}$ for $n=77$). The spatially averaged propagated [combined](#) uncertainties for $p\text{CO}_2$ and pH are 74 μatm and 0.08

respectively, where the pH uncertainty is relative to an average pH of 8.21. The combined uncertainties in TA and DIC in the Amazon and Congo outflows are lower than the natural variability within their respective regions allowing the time varying regional variability to be evaluated. Potential uses of these data would be for assessing the spatial and temporal flow of carbon from the Amazon and Congo rivers into the Atlantic and for assessing the riverine driven carbonate system variations experienced by tropical reefs within the outflow regions.

1 Introduction

Rivers connect the land and ocean providing a major pathway for carbon transport to the ocean (Regnier et al., 2013). The inorganic carbon content of rivers is poorly constrained ~~due because to the difficulties of sampling it is difficult to sufficiently sample~~ these highly spatial and temporally variable river outflows. Global estimates of the riverine flow of carbon from the land to the ocean (Friedlingstein et al., 2022) are determined from *in situ* upscaling (Regnier et al., 2013), ocean inverse model estimates (Jacobson et al., 2007), partial pressure of carbon dioxide ($p\text{CO}_2$) ocean sink based estimates (Watson et al., 2020) and atmosphere inversion based estimates (Rödenbeck et al., 2018). Clearly new methods that improve the characterisation of the magnitude, variability and temporal variations of carbon transported by rivers will help constrain uncertainties within global carbon budgets (Hauck et al., 2020).

The vast majority of $p\text{CO}_2$ measurements in the Surface ocean CO_2 Atlas (SOCAT) were made on research ships and ships of opportunity that mainly survey the open ocean and continental shelves (Bakker et al., 2016). Many bottle samples have been collected for total alkalinity (TA) and dissolved organic carbon (DIC) in rivers, but fully surveying the entirety of large rivers across all seasons is logistically challenging requiring extensive and economically expensive field campaigns (Ward et al., 2017). For example, the majority of the largest 100 rivers by discharge are found in South America and Asia (Dai and Trenberth, 2002) and have been historically under-sampled for carbonate system variables (Lauerwald et al., 2015). Issues related to scarcity of measurements is compounded by insufficient knowledge of the hydrology and spatial area extents of these systems (Allen and Pavelsky, 2018). Additionally, the amount of carbon in the rivers is a function of runoff rates, rainfall and land use, all of which have been disrupted by climate change and land use change (Piao et al., 2007; Kaushal et al., 2014; Regnier et al., 2013). This lack of large spatial and temporal scale baseline carbonate system observations in rivers means that assessing changes in these systems is challenging.

The carbon dynamics of the world's rivers also have implications on local biogeochemistry. Ocean acidification is the long-term process by which the oceans absorb atmospheric CO_2 making them ~~becomes~~ less alkaline ~~(as due to an increase in the~~ hydrogen ion concentration, lowering their pH ~~increases)~~ ~~whilst at the same time~~ ~~and Ocean acidification also~~ ~~decreases~~ decreasing the carbonate ion availability (Doney et al., 2009). Ocean acidification poses a threat to marine

Formatted: Subscript

65 organisms which build calcium carbonate structures and many rivers are sensitive to this due to their low buffering capacity
(Hu and Cai, 2013;Cai et al., 2011). River plumes can negatively influence wild fisheries and the aquaculture industry
(Mathis et al., 2015;Cattano et al., 2018) as plumes can transport low pH waters ~~that can~~with the ability to impact the growth
and life stages of many marine organisms (Cai et al., 2021). Additionally, river plumes can interact with high biodiversity
70 regions that are sensitive to sudden changes in the carbonate system such as sensitive coral reef systems (Mongin et al.,
2016;Dong et al., 2017). ~~Episodic~~Intermittent changes in the carbonate system caused by river plumes can ~~potentially~~
~~jeopardise ecosystem services like fisheries, aquaculture, and shoreline protection, the resultant result in~~financial and
biodiversity losses and ~~so are of~~ ~~paramount great~~ interest to local communities, businesses and policy makers (Doney et al.,
2020).

75 Satellite Earth observations provide a means of accurately assessing the carbonate content of large rivers by ~~using~~utilising
~~satellite observed~~ oceanographic variables ~~that can now be routinely observed from space combined~~with published
empirical algorithms ~~from the literature that link carbonate system variables to the satellite derived variables~~(Land et al.,
2019). ~~The Satellite Oceanographic Datasets for Acidification (OceanSODA) project (<https://esa-oceansoda.org>) was~~
~~established to further develop these approaches~~maximise the usefulness of current ~~insitu carbonate observation by exploiting~~
80 ~~the superb temporal and spatial resolution of satellite observations~~. We present the University Of Exeter (UNEXE)
OceanSODA dataset (OceanSODA- UNEXE), decadal datasets of riverine carbonate system variables for the two largest
rivers in the world by discharge, the Amazon and the Congo Rivers (Dai and Trenberth, 2002). This paper details how the
optimal combination of empirical algorithms and Earth observation datasets were selected and used to construct
OceanSODA-UNEXE. This paper provides an assessment of the uncertainty associated with the key TA and DIC parameters
85 using a standardised uncertainty framework and a large *in situ* database. The remaining ocean carbon system variables were
calculated from TA and DIC with propagated uncertainties.

2 Methods

2.1 Statistical terms overview

90 Root mean squared difference (RMSD) is a measure of accuracy and is calculated as the square root of the average of
squared errors e.g. $RMSD = ((\sum(x_0 - x_i)^2)/n)^{1/2}$, where x_0 are the estimated values, x_i are the reference values and n is the
number of observations. The bias of a dataset is defined as the mean difference between the estimation and reference, $Bias =$
 $\sum(x_0 - x_i)/n$. The mean absolute difference (MAD) of a dataset is a measure of variability and is calculated as $MAD = (\sum|x -$
 $\bar{x}|)/n$, where \bar{x} is the mean. The correlation coefficient (r_{ij}) is a measure of linear correlation between the estimate and the
reference variables and is defined as $r_{ij} = (\sum (x_i - \bar{x}_i) (x_j - \bar{x}_j)) / (\sum (x_i - \bar{x}_i)^2 \sum (x_j - \bar{x}_j)^2)^{1/2}$. Uncertainty representation and the

95 terminology used throughout this paper are consistent with the International Bureau of Weights and Measures (BIPM) Guide to the expression of uncertainty in measurement (GUM) methodology (JCGM, 2008).

Weighted statistics allow uncertainties in the reference dataset (*in situ* measurement uncertainty in this case) to be accounted for within the performance analysis (i.e. the reference data are not considered ‘truth’ as they also contain uncertainties).

100 Weights are calculated as the sum of the individual weight of each algorithm (w), where $w = 1/((in\ situ\ measurement\ uncertainty)^2 + (literature\ algorithm\ uncertainty)^2)^{1/2}$ (Ford et al., 2021). For clarity and easy comparison to previous published work both weighted and unweighted statistics of all metrics are given, for example weighted RMSD (wRMSD) is calculated as $wRMSD = (\sum(weights * (x_0 - x_i)^2))^{1/2}$.

105 ~~When evaluating algorithms using a statistical measure, in this case a statistical measure like wRMSD. When evaluating algorithms by wRMSD,~~ a further issue can arise where the valid region over which each algorithm can be applied overlaps with different *in situ* data. For example, an algorithm evaluated using data from highly-variable coastal waters may have a higher wRMSD than another algorithm evaluated using solely data from a less variable open ocean region; in this scenario it may be falsely concluded that the coastal ocean algorithm performs worse. This is a clear weakness of comparing wRMSD values from different sources and across differing regions (Land et al., 2019). Following the methodology of Land, Findlay et al. (2019) ~~we calculate RMSDe using the wRMSD result we derive RMSDe from wRMSD,~~ RMSDe is a more representative metric to compare accuracies as it allows algorithms to be evaluated in a like for like manner allowing their performance to be ranked. It is important to note that for the best algorithm, wRMSD = RMSDe as the best algorithm will always have a score of 1. To ensure the robustness of any statistics generated it was considered prudent to specify a
110 minimum data threshold. Only algorithms that had at least 30 matchups ($n=30$) between the algorithm and reference outputs were used in the calculation of RMSDe, this was done to prevent the selection of algorithms with low RMSDe values caused by evaluating the algorithm with a small number of data points. wRMSD is used as the preferred measure of accuracy in this
115 paper but unweighted RMSD values are also given.

2.2 Selection of empirical algorithms

120 In order to generate the full ocean carbonate system, two of the four carbonate system variables are needed. As TA is closely linked to salinity it is selected as one variable, for the second variable DIC is selected. There are many algorithms in the published literature for both TA and DIC and the required measurements in the river outflows are available to evaluate those algorithms. Other pairings of carbonate system parameters can be used to derive the full carbonate system (Land et al., 2015b) but are not explored here, ~~for example (e.g. Gregor and Gruber (2021) who use TA and pCO_2).~~

125

130 An exhaustive literature search using 24 search terms (~~listed in the supplementary materials~~) identified prospective algorithms that could be applied to the Amazon and Congo River outflows. The full list of **search terms and identified algorithms** can be found in the supplementary materials. **The region bounds of the Amazon outflow were defined as being 2° S to 24° N and 70° W to 31° W. The bounds of the Congo outflow were defined as being between 10°S to 4°N and 2° W to 16° E. To be included in the algorithm evaluation, algorithms needed to be applicable to these regions and to take the form of a linear or quadratic relationship with input variables that were easy to obtain and available as spatially and temporally varying datasets. These input variables included sea surface temperature (SST), sea surface salinity (SSS), potential temperature (which is assumed to be approximately equal to SST at the surface), dissolved oxygen (DO), nitrate (NO₃⁻), phosphate (PO₄⁻³), silicate (SiO₄⁻⁴) and chlorophyll-a.**~~To be included in the algorithm evaluation, algorithms needed to be applicable within the outflow region of either river (the Amazon between 2° S and 24° N and between 70° W and 31° W or the Congo between 10°S and 4°N and between 2° W and 16° E) and to take the form of a linear or quadratic relationship with accessible input variables e.g. sea surface temperature (SST), sea surface salinity (SSS), potential temperature (which is assumed to be approximately equal to SST at the surface), dissolved oxygen (DO), nitrate (NO₃⁻), phosphate (PO₄⁻³), silicate (SiO₄⁻⁴) and chlorophyll-a.~~ 26 of the identified algorithms were not included in the algorithm evaluation as they could not generate TA or DIC using the accessible input variables listed above or were not based on empirical algorithms (See Table S2 for the full list). **A (e.g. they were full-*in situ* approaches involving biogeochemical models or neural networks approaches were not included in the algorithm evaluation)** (See Table S2 for the full list). 10 TA algorithms (5 of 10 report RMSD values in their original publication) and 6 DIC algorithms (5 of 6 report RMSD) were evaluated for the Amazon and 4 TA algorithms (3 of 4 report RMSD) and 9 DIC algorithms (4 of 7 report RMSD) were evaluated for the Congo (Table S1). wRMSD and RMSDe can only be calculated for algorithms which report RMSD, so it is important to distinguish these. Table S1 details each algorithm's input variables, the published algorithm RMSD, the stated environmental ranges for which the algorithm is valid and some brief descriptive notes of how the algorithm was developed. The target output variable, the input variables, the mathematical algorithm, the valid geographical region and the valid geophysical conditions for each algorithm were then gathered for use in the algorithm evaluation process.

2.3 Algorithm evaluation

A multipurpose global reference matchups database (MDB) matching *in situ* carbonate system parameters from the surface 10 m with satellite, model and interpolated *in situ* datasets was used to perform the algorithm evaluation (Land et al., 2022). The MDB is optimised to reduce biases arising from uneven data density; this is achieved by grouping *in situ* observations into 100 km diameter regions of interest (ROI) which span a 10 day time period. The MDB was constructed from global datasets that include all of the variables required for all algorithms (SST, SSS, θ , DO, NO₃⁻, PO₄⁻³, SiO₄⁻⁴ and chlorophyll-a see section 2.2). The MDB includes three global SST datasets: European Space Agency Climate Change Initiative SST (ESACCI SST) v2.1 (Merchant et al., 2019; Good et al., 2019), Optimum Interpolation SST (OISST) v2.1 (Huang et al., 2021; Banzon et al., 2016) and the Coriolis Ocean dataset for Reanalysis (CORA) v5.2 (Szekely et al., 2019)) and four global SSS datasets: European Space Agency Climate Change Initiative SSS (ESACCI SSS) v2.31 (Boutin et al., 2020; Boutin et al., 2021), CORA v5.2 (Szekely et al., 2019), the *in situ* Analysis System (ISAS 15) (Kolodziejczyk et al., 2021; Gaillard et al., 2016) and Remote Sensing Systems data from the Soil Moisture Active Passive satellite (RSS-SMAP) level 3 v4.0 (Meissner et al., 2019; Meissner et al., 2018)). The MDB also contains TA, DIC and pH values, most of which come from the Global Ocean Data Analysis Project (GLODAPv2.2020) database (Olsen et al., 2016) and *p*CO₂ values most of which come

from SOCAT v2020 (Bakker et al., 2016) along with additional data for Arctic waters. A full list of references for data variable sources used in the MDB can be found in (Land et al., 2022). Before beginning this analysis we limited our analysis to only used MDB data that fall within these bounds: SST > -10 °C or < 40 °C, SSS > 0 or < 50, DIC > 500 $\mu\text{mol kg}^{-1}$ or < 3000 $\mu\text{mol kg}^{-1}$, TA > 500 $\mu\text{mol kg}^{-1}$ or < 3000 $\mu\text{mol kg}^{-1}$, pH > 6 or < 8.5 and $p\text{CO}_2$ > 100 μatm or < 3000 μatm and these constraints are consistent with the conditions likely within the Amazon and Congo regions.

Each valid algorithm was implemented by closely following the specific literature recommendations; this was done in such a way that empirical algorithms were only assessed for the geographical and geophysical ranges for which they were originally developed. The algorithm evaluation involved running each algorithm with inputs from the MDB to estimate the TA and DIC outputs (termed the “algorithm output”). Each algorithm output was then evaluated using the TA and DIC values from the MDB matchup database (termed the “reference output”). The RMSD, bias, correlation coefficient, and MAD are calculated from the algorithm output and the reference output. Additionally, wRMSD was calculated for algorithms where an RMSD is reported in the literature. Where wRMSD was calculated and the matchups (n) were > 30, RMSDe was calculated for those algorithms following the methodology of Land et al. (2019). As the MDB includes three global SST and four global SSS datasets, each algorithm was run for each of the twelve combinations of SST and SSS input datasets, and separate statistics were generated for each configuration. The subset of input variables from the MDB used to implement the algorithm, the algorithm output and the reference output for all 12 SST and SSS combinations are included in the supplementary materials. Additionally, summary statistics between the algorithm output and the reference output are also provided in the supplementary materials, these statistics are the (mean, wmean, standard deviation, RMSD, wRMSD, RMSDe, Bias, wBias, r, wr, MAD and wMAD) ~~are also provided in the supplementary materials.~~

The best performing TA and DIC algorithms for both regions were determined by ranking the algorithms by RMSDe, the best performing algorithms were the algorithms with the lowest RMSDe values. In order to select algorithms suitable for generating time series, observation datasets for SSS and SST needed a temporal overlap of at least 8 years. The remainder of this paper exclusively discusses the time series generated with these “optimal algorithms”.

The algorithm evaluation process follows a standardised framework for a Type A uncertainty evaluation for TA and DIC whereby the following sources of uncertainties are considered:

1. The TA or DIC measurement uncertainty. The measurement uncertainty for both TA and DIC is typically stated as $\pm 4 \mu\text{mol kg}^{-1}$ but following (Bockmon and Dickson, 2015), we use their more conservative measurement uncertainty of $\pm 10 \mu\text{mol kg}^{-1}$ (approximately equal to 0.5% of the nominal TA and DIC values for the open ocean);
2. The algorithm uncertainty which is the RMSD stated in the literature for each algorithm;
3. The spatial uncertainty arising from spatial heterogeneity in the region;

4. The uncertainty due to differences in the measurement depths in the MDB.

200 The measurement uncertainty of $10 \mu\text{mol kg}^{-1}$ and algorithm uncertainty are explicitly considered in the calculated of
wRMSD and the remaining uncertainties should be minimised through using the MDB (~~as the MDB has been~~ which was
designed to minimise these components (Land et al., 2022)). By accounting for all known sources of uncertainty this
constitutes a complete Type A uncertainty evaluation. The combined standard uncertainty (δQ) for TA and DIC is calculated
as $\delta Q = ((\delta a)^2 + (\delta b)^2)^{1/2}$ where δa is the RMSDe from the evaluation algorithm and δb is the wBias. The uncertainty
205 estimates provided within the dataset is the combined standard uncertainty (δQ) and these combined standard uncertainties
are reported without confidence intervals.

To aid interpretation of the combined uncertainty budgets, a second Type A uncertainty evaluation of the TA and DIC
approaches was calculated based purely on the empirical algorithm (e.g. literature RMSD value) and the uncertainties in the
210 input data propagated using standard techniques (Taylor, 1997) and assuming that any uncertainties were uncorrelated. This
is a second independent Type A uncertainty evaluation that does not include uncertainties due to spatial variability and
depth. Assuming that the first Type A uncertainty evaluation has captured all the uncertainties, the difference between the
two uncertainty evaluations enables estimation of the contribution of the spatial and depth variability on the uncertainty
budget. A similar approach was recently performed by Gregor and Gruber (2021) where they refer to these two approaches
215 as ‘top down’ (our first Type A uncertainty evaluation) and ‘bottom up’ (our second Type A uncertainty evaluation)
assessments.

Evaluating each algorithm exactly as it appears in the literature is required to ensure equitably in the algorithm evaluation.
Theoretically, all of the algorithms could have been modified to account for long-term changes in environmental conditions
220 (e.g. oceanic CO₂ uptake and increased freshwater input to the oceans) in these two outflow regions that have likely occurred
since the algorithms were first developed. However, accounting for changing environmental conditions is not
straightforward. Whilst secular trends for variables like surface CO₂ are accurate for the open ocean, these regions are
heavily impacted by the river outflows and so applying open ocean secular trends to these in these riverine regions is likely
problematic and so considered beyond the scope of this study.
225 are highly uncertain. Correcting each algorithm for secular
trends would either involve adding offsets to data and refitting the data or adding offsets directly to the variable being
derived with the algorithm. Any applied offsets would need to be generated with some form of model and would therefore
have associated uncertainties that would also need to be accounted for. A further complication is that even if all algorithms
were modified to a reference year, there is still a temporal mismatch between the algorithm reference year and the data in the
230 MDB, addressing that would require all the data in the MDB to be standardised to a reference year which would then
introduce additional uncertainties. Secular trends are not factored into this algorithm evaluation but they could be
incorporated into the open access toolbox if the above issues could be appropriately addressed. The impact of any secular

~~trends in any parameters is somewhat mitigated as the analysis focuses on the use of using TA and DIC and as these variables as two chosen carbonate system variables as they are likely have been less impacted by secular trends than $p\text{CO}_2$ and pH.~~

235 2.4 Creation of gridded monthly time series product

The full spatiotemporal resolution SST and SSS datasets and gridded World Ocean Atlas (WOA) DO, NO_3^- , PO_4^{3-} and SiO_4^{4-} datasets were downloaded from their online repositories, the total size of these full resolution datasets occupies ~500 GB of hard drive space and take several days to fully download. The SST and SSS datasets were uncompressed and were re-gridded onto a monthly $1^\circ \times 1^\circ$ standard World Geodetic System grid for use creating the gridded output, the grid has 180° of Latitude and 360° of Longitude with 0° centred on the Greenwich Meridian. Reformatting the datasets considerably reduced the file size to a more manageable ~50 GB. The scripts used to download these SST and SSS monthly datasets are fully automated, meaning that when new data are added to these datasets it can be easily incorporated into future versions of OceanSODA-UNEXE.

245 For each target variables (TA or DIC), the selected optimal SST and SSS gridded datasets and gridded WOA datasets were used as inputs to the respective TA and DIC algorithms. The algorithms were applied to the datasets in the two outflow regions where the SST and SSS dataset overlapped in time for the published environmental limits of each algorithm. The monthly SST and SSS inputs as well as the calculated monthly output variable (TA or DIC) are saved to a netCDF4 file as separate variables on the grid described above with dimensions of 768 by 180 by 360. ~~(where The time -768 is the time step~~
250 ~~in dimension is the number of months starting insince January 1957).~~ It is important to note that TA and DIC datasets (even if for the same region) can span different temporal domains and so should be treated as separate datasets. These differences are due to differences in the temporal range of the input data used to calculate them.

~~The oceanic carbon system equations (Millero, 2000) can be solved computationally (Lewis et al., 1998), using numerical a number of software packages exist to perform these calculations (Orr et al., 2015) and . The oceanic carbon system equations can now be solved natively in python with the established CO2SYS software package is now available in Python as PyCO2SYS (Humphreys et al., 2022). Where there was temporal overlap between the TA and DIC datasets in each region, the remaining oceanic carbonate system variables ($p\text{CO}_2$ and pH) were calculated using with PyCO2SYS v1.7.1 (Humphreys et al., 2022). A number of other output variables were also calculated including $f\text{CO}_2$, the carbonate ion content (CO_3^{2-}), the bicarbonate ion content (HCO_3^-), the hydrogen ion content (H^+) and the calcite (Ω Calcite) and aragonite (Ω Aragonite) saturation states. the pyCO2SYS v1.7.1 software package (Humphreys et al., 2022) was used to calculate the complete carbonate system variables ($p\text{CO}_2$, $f\text{CO}_2$ and pH on the free, total and seawater scales), the carbonate ion content (CO_3^{2-}), the bicarbonate ion content (HCO_3^-), the hydrogen ion content (H^+) and the calcite (Ω Calcite) and aragonite (Ω~~

265 ~~Aragonite) saturation states.~~ PpyCO2SYS was run with the selected optimal SST-dataset (SST, SSS and WOA datasets
where applicable, with the same data used for input and output conditions) and SSS-gridded-datasets, gridded WOA-datasets
at the surface (0 m depth) with the carbonic acid dissociation constants of Mehrbach (1973) refitted by Dickson and Millero
(1987) and the hydrogen sulphate dissociation constant of Dickson (1990). The full carbonate system is calculated twice,
once using the optimal SST and SSS datasets that were used to calculate TA then again using those used to calculate DIC.
270 Consequently each netCDF file contains a complete set of carbonate system outputs that are unique to that specific input
dataset.

Whilst some of the SST and SSS products used to create the gridded datasets contain only satellite observed SST at a
specified depth, some of the re-analysis SST and SSS products also incorporate surface ocean measurements from ships and
buoys which were made at differing depths. The TA and DIC data used to generate the MDB comes from the top 10 m of the
275 ocean and has not been adjusted for potential concentrations gradients in the surface ocean (Land et al., 2022). For these
reasons, OceanSODA-UNEXE is deemed relevant in the surface 10 m of the ocean.

2.5 Uncertainties in gridded monthly time series product


As stated in section 2.3, the assessed combined standard uncertainties for TA and DIC are single fixed values provided as
regionally-static (non-spatially varying) fields in the netCDF files. Uncertainty information for SST and SSS are taken from
280 the literature references and are provided as spatially varying fields in the netCDF ~~files (as provided in the original datasets).~~
Uncertainties in TA, DIC, SST and SSS are provided as optional uncertainty input arguments in PpyCO2SYS to generate
uncertainties for pH, $p\text{CO}_2$, Ω Calcite and Ω Aragonite. PyCO2SYS uses a forward finite difference approach to calculate
the derivatives needed to propagate uncertainties for all these output variables. This produces a spatially varying combined
uncertainty budget for each remaining carbonate system parameter. The spatially varying uncertainties in SST, SSS, pH,
285 $p\text{CO}_2$, Ω Calcite and Ω Aragonite are all provided in the respective netCDF files.

3 Results

3.1 Algorithm evaluation

The algorithm evaluation RMSDe values in the Amazon and Congo for TA and DIC are shown in Figure 1. From these
RMSDe values we selected the “optimal algorithms” for both regions as the algorithm combination with the lowest RMSDe.
290 The optimal algorithms were then used to produce OceanSODA-UNEXE. In the Amazon the optimal algorithm to generate
the TA data was (Cai et al., 2010) with ESACCI SST and ESACCI SSS (Table 1). To generate DIC for the Amazon the best
algorithm was (Lefèvre et al., 2010) using OISST SST and ESACCI SSS. In the Congo, the optimal algorithm to generate

the TA data was Lee et al. (2006) with CORA SST and ISAS SSS (Table 2). For DIC in the Congo the optimal algorithm was (Lee et al., 2000) with ESACCI SST and CORA SSS.



300 **Figure 1: Results of the algorithm evaluation, DIC and TA algorithms for the Amazon and Congo regions are compared by RMSDe. Each of the valid algorithms is listed by literature reference on the y axis, all of which correspond to algorithms in Table**

S1. Each SST and SSS dataset input configurations for which algorithms were evaluated are shown for each algorithm, with each SSS dataset given by a different symbol and each SST dataset given as a different colour in the legend. Entries are blank where there was no RMSD value in the literature, there were < 30 matchups or where the assessed RMSDe value was >100 $\mu\text{mol kg}^{-1}$.

305 Table 1: Summary table of the best combination of algorithm and SST/SSS datasets (aka the “optimal algorithms”) in the Amazon River domain. These optimal algorithms were determined by selecting the algorithm with the lowest RMSDe where there was at least 8 years of data.

	Algorithm	SST dataset	SSS dataset	Years of overlap	Number of matchups (n) for weighted statistics	RMSD ($\mu\text{mol kg}^{-1}$)	wBias ($\mu\text{mol kg}^{-1}$)	wRMSD ($\mu\text{mol kg}^{-1}$)	RMSDe ($\mu\text{mol kg}^{-1}$)	Combined standard uncertainty ($\mu\text{mol kg}^{-1}$)
Optimal TA algorithm	(Cai et al., 2010)	ESACCI SST	ESACCI SSS	10 years (2010 - 2020)	82	50.92	7.83	34.97	34.97	35.84
Optimal DIC algorithm	(Lefèvre et al., 2010)	OISST	ESACCI SSS	10 years (2010 - 2020)	70	53.33	-5.98	43.83	43.83	44.23

310 Table 2: Summary table of the best combination of algorithm and SST/SSS datasets (aka the “optimal algorithms”) in the Congo River domain. These optimal algorithms were determined by selecting the algorithm with the lowest RMSDe where there was at least 8 years of data.


	Algorithm	SST dataset	SSS dataset	Years of overlap	Number of matchups (n) for weighted statistics	RMSD ($\mu\text{mol kg}^{-1}$)	wBias ($\mu\text{mol kg}^{-1}$)	wRMSD ($\mu\text{mol kg}^{-1}$)	RMSDe ($\mu\text{mol kg}^{-1}$)	Combined standard uncertainty ($\mu\text{mol kg}^{-1}$)
Optimal TA algorithm	(Lee et al., 2006)	CORA	ISAS	14 years (2002 - 2016)	102	27.33	6.00	27.91	27.91	28.54
Optimal	(Lee et al., 2006)	ESACCI	CORA	14 years	77	37.37	-16.64	36.79	36.79	40.37

DIC algorithm	al., 2000)	SST		(2002 - 2016)						
------------------	---------------	-----	--	------------------	--	--	--	--	--	--

3.2 Dataset output

3.2.1 Amazon dataset

315 The primary output datasets are the monthly gridded carbonate system variables (DIC and TA). The seasonal progression of
both these variables is shown in the Amazon outflow region in Figures 2 and 3. There is a large range of DIC and TA in this
region, with high values (DIC $\sim 2000 \mu\text{mol kg}^{-1}$ and TA $\sim 2350 \mu\text{mol kg}^{-1}$) in the open ocean. The lowest values (DIC and TA
 $\sim 1400 \mu\text{mol kg}^{-1}$) are located at the mouth of the river. Intermediate values occur where the plume and ocean water are
mixing. The plume extends to the Northwest from the Amazon River mouth (Coles et al., 2013); this is especially visible
320 from April to September (Figures 2 and 3 plots b and c), less so from October to March (plots a and d), demonstrating the
seasonal variability. The region of influence of the Amazon Plume extends into the Caribbean between April and June
(Figure 2b and 3b), a feature reported elsewhere (Chérubin and Richardson, 2007). From July to September (Figure 2c and
3c) another large outflow region occurs just off the coast of Venezuela at the mouth of the Orinoco River (Hu et al., 2004).
The Orinoco plume is much less prominent in the other seasons. The proximity of the Orinoco River to the Caribbean
325 suggests that the Orinoco plume often reaches the islands between July and September (López et al., 2013). There are some
differences between the structure of the average TA and DIC plumes which may indicate active processes in the plume, for
example gas exchange which only affects DIC or biological production which affects TA and DIC at different rates. Annual
and seasonal data averages over the region shown in Figures 2 and 3 are provided in Table S3 for DIC, TA, pH, $p\text{CO}_2$, Ω
Calcite and Ω Aragonite.



330

Figure 2: Seasonally averaged DIC for the Amazon plume region in (a) January to March (b) April to June (c) July to September (d) October to December. Land outlines are shown in beige. Ocean regions out of bounds or where there was no algorithm output are left white. Algorithm data below $1600 \mu\text{mol kg}^{-1}$ at the river outflows is shown in mint green. The mouths of the Amazon, Orinoco and Maroni Rivers are labelled. The 52°W meridional section used for the Hovmöller plot used in Figure 4 is indicated as a black dashed line.




Figure 3: Seasonally averaged TA for the Amazon plume region in (a) January to March (b) April to June (c) July to September (d) October to December. Land outlines are shown in beige. Ocean regions out of bounds or where there was no algorithm output are left white. Algorithm data below $2000 \mu\text{mol kg}^{-1}$ at the river outflows is shown in mint green. Orinoco and Maroni Rivers are labelled. The 52°W meridional section used for the Hovmöller plot used in Figure 4 is indicated as a black dashed line.


335

The temporal aspect of the dataset is presented in Figure 4 as a meridional Hovmöller plot at 52°W (marked on Figures 2 and 3) which broadly cuts through the central plume outflow at 4°N , the mixing of the plume up to 10°N and the open ocean Atlantic water $>10^\circ \text{N}$. The lowest DIC values ($\sim 1400 \mu\text{mol kg}^{-1}$) are found in the plume at 4°N , there is regular seasonality with lower values in the October to December and higher values in April to June (Figure 4a). There does not appear to be much interannual variability in the magnitude of the DIC plume in the June to August period, although 2010 and 2013 appear to be weaker (Figure 4a). The maximum northerly extent of the plume varies from year to year, with higher DIC found at 15°N in the May to August period of 2011. The opposite is true in other years, for example the plume does not extend as far northwards in 2013 (Figure 4a). TA shows much more interannual variability (Figure 4b). The minimum plume values do not always occur at the same latitude or at the same time. In most years the minimum plume values are as low as $1750 \mu\text{mol kg}^{-1}$ but this was not the case in 2016. The northward extent of the plume is variable, in some years the plume extended much further North, e.g. June to July 2011, and in other years the plume influence was not as detectable

345

further North, e.g. May to July 2016. The period of time where the plume dominates the region is also variable; in some years the peak plume intensity which begins in May lasts through to September while in other years it is already declining in
350 July.

As $p\text{CO}_2$ and pH are derived from DIC and TA, the behaviour of these variables mirrors that of TA and DIC (Figure 4c and 4d). The river plume always has above neutral pH and occasionally extends far to the north, e.g. June to July 2011. There are low $p\text{CO}_2$ values ($<200 \mu\text{atm}$) in the plume which are much lower than some of the values observed in the plume (Lefèvre et al., 2017). Calcite and aragonite saturation states have the expected magnitude in open ocean regions and are lower in the
355 plume, with Ω Calcite levels ~ 3 in the plume.



360 **Figure 4:** Hovmöller plots in the Amazon outflow region for (a) DIC, (b) TA, (c) pH on the free scale, (d) $p\text{CO}_2$, (e) Ω Aragonite and (f) Ω Calcite. The plots span the 52° W meridian from 4 to 24° N. The plots cover the temporal overlap period of the TA and DIC datasets 2010- 2020.

The gridded data can also be spatially averaged to show the evolution of each of the carbonate system variables in time within the plume (defined as $\text{SSS} < 35$ (Grotsky et al., 2014)), outside the plume ($\text{SSS} > 35$) and over the whole region (Figures 5). The mean DIC calculated across whole region shows a very consistent seasonal pattern, with higher DIC in

365

December to February ($\sim 2000 \mu\text{mol kg}^{-1}$) and lower DIC in June to August ($\sim 1970 \mu\text{mol kg}^{-1}$). The mean values within the plume are much lower; the lowest DIC values ($\sim 1700 \mu\text{mol kg}^{-1}$) are seen in December to February and the highest values ($\sim 1900 \mu\text{mol kg}^{-1}$) are during September and October. The mean non-plume values show much smaller seasonal variability.

In the plume, outside of the plume and the whole region, surface DIC is very similar year to year. The mean DIC within the

370 | plume reaches a minimum in 2013 which was much lower than all other years. Non-plume DIC values increased by $0.49 \mu\text{mol kg}^{-1}$ per year. TA across the whole region is consistent inter annually, with a seasonal maximum in December to

February ($\sim 2360 \mu\text{mol kg}^{-1}$) and minimum in July to August ($\sim 2330 \mu\text{mol kg}^{-1}$), while TA within the plume shows a seasonal maximum in October ($\sim 2250 \mu\text{mol kg}^{-1}$) and minimum in April to May ($\sim 2150 \mu\text{mol kg}^{-1}$). The annual minimum within the

375 | plume varies from December to March, while the maximum consistently occurs in October and November, with considerable interannual variability of the timing of the annual minimum, with much lower TA in the December to February periods of 2012/2013 and 2017/2018 than over the same period in other years. Non-plume TA values increased by $0.76 \mu\text{mol kg}^{-1}$ per year. It should be noted that the whole region is out of phase with the plume, this is because whilst the plume is

more dilute this effect is outweighed by the larger plume between May and August.

380 | For pH there is a weak seasonal trend in the overall region and outside the plume. There is a lot more variability in the plume, with the highest pH values (~ 8.225) occurring in December and the lowest values (~ 8.2) in September. The best SST and SSS dataset from the DIC algorithm predict that the pH will be >0.1 higher than the best SST and SSS dataset from the TA algorithm between January and March. The two estimates agree within <0.1 pH units for the other 9 months of the year.

385 | For $p\text{CO}_2$, also computed from TA and DIC, mean values in the non-plume region and over the whole region were very stable, which is consistent with the minimal season variability seen in oligotrophic oceans. Whereas there was considerable variability in plume $p\text{CO}_2$, with average values of $\sim 325 \mu\text{atm}$ seen in the plume in March and values of $\sim 350 \mu\text{atm}$ in August. The differences between the $p\text{CO}_2$ calculated with the different SST and SSS datasets was very small, with differences $<5 \mu\text{atm}$ most of the year. The average calcite and aragonite saturation states are in the typical range for seawater

390 | and are much greater than the critical thermodynamic threshold for calcification with a value of 1 (Waldbusser et al., 2016) ~~(the critical thermodynamic threshold for calcification, Waldbusser et al., 2016)~~. It should be noted that an aragonite saturation state threshold of 3 has also been recommended for warm water corals (Guinotte et al., 2003), other studies have since shown that warm water corals can still adapt and survive in these conditions (Enochs et al., 2020; Uthicke et al., 2014). There are periods in the data such as at the start of 2019 where the aragonite saturation state falls below 3. The mean saturation states within the plume are lower than those found in the wider region.

395

Formatted: Superscript

Formatted: Superscript





Figure 5: Timeseries of spatially averaged (a) DIC, (b) TA, (c) pH, (d) $p\text{CO}_2$, (e) Ω Aragonite and (f) Ω Calcite in the Amazon region. The plots spans the temporal overlap period of the TA and DIC datasets 2010- 2020. Data are averaged across the whole region (solid line) as well as in the plume defined as $S < 35$ (dashed line) and outside of the plume $S > 35$ (dotted line). The line colour corresponds to variables that were calculated with SST and SSS datasets selected during the DIC algorithm evaluation (red) and the TA algorithm evaluation (blue).

3.2.1 Congo dataset


In the Congo outflow region, there is a strong seasonal variation in DIC (Figure 6). From July to September (Figure 6c) there are two regions of low DIC, one directly at the outflow point of the Congo River and another at the outflow point of the Niger River Delta. The outflows of both of these river dominated regions extend directly west, aligned to the South Equatorial Current in the eastern Atlantic (Hopkins et al., 2013). In October to December (Figure 6d) the spatial influence of the Congo outflow is smaller possibly due water masses moving northwards along the coast. Into January to March (Figure 6a) there is a region of low DIC along the whole coast, representing the intensification of the Niger and Congo discharges at this time of year (Chao et al., 2015). In April to June (Figure 6b) the coastal water mass separates becoming two distinct outflows, the outflows reach their greatest westward extent covering almost the whole domain.

Similar trends are seen in TA (Figure 7) with two distinct outflow regions in July to September (Figure 7c). The outflows begin to intensify in October to December (Figure 7d). The lowest TA values are observed in January to March to the east of the Niger Delta outflow (Figure 7a). Unlike in Figure 6a, the outflows remain distinct in Figure 7a. Between April and June the outflow reaches its maximum spatial extend as it flows out to the west where it is has a detectable impact across the region.



415 **Figure 6:** Seasonally averaged DIC for the Congo outflow region in (a) January to March (b) April to June (c) July to September (d) October to December. Land outlines are shown in beige. Ocean regions out of bounds or where there was no algorithm output are left white. Algorithm data below $1750 \mu\text{mol kg}^{-1}$ at the river outflows is shown in mint green. The Niger River Delta and the mouths of the Congo, Ogooué and Sanaga Rivers are labelled. The 6° S meridional section used for the Hovmöller plot used in Figure 8 is indicated a bold dashed line.

420



425 **Figure 7: Seasonally averaged TA for the Congo outflow region in (a) January to March (b) April to June (c) July to September (d) October to December** Land outlines are shown in beige. Ocean regions out of bounds or where there was no algorithm output are left white. The Niger River Delta and the mouths of the Congo, Ogooué and Sanaga Rivers are labelled. The 6° S meridional section used for the Hovmöller plot used in Figure 8 is indicated a bold dashed line.

A zonal section at 6° S was used to construct a Hovmöller plot of variables in the Congo (Figure 8). This latitude is centred across the outflow of the Congo River. Unlike in the Amazon, much of the plot region is masked in between May and July as the published algorithms were not valid for full environmental conditions experienced by the region.

430 The DIC plot (Figure 8a) shows that the outflow is low in DIC ($\sim 1800 \mu\text{mol kg}^{-1}$) and the open ocean is higher ($\sim 2050 \mu\text{mol kg}^{-1}$). The highest values are found in the January to March period, consistent with Figure 6a. The outflow is detectable over the widest area in the March to June period in all years. Whilst there is one period of intense outflow each year there are also some indication that a weaker outflow is intermittently detectable in the data (shown as vertical streaks in Figure 8a). TA (Figure 8b) shows an almost identical pattern to DIC. The lowest TA values ($\sim 2100 \mu\text{mol kg}^{-1}$) were observed between 435 January and March and slightly higher values ($\sim 2200 \mu\text{mol kg}^{-1}$) were seen when the outflow extended further West between April and June.

Higher pH values (~8.3) were observed in the outflow compared to the open ocean (~8.2) (Figure 8c. Previous studies have measured low pH values in the main body of the river and its tributaries (Wang et al., 2013; Bouillon et al., 2014), the higher pH values at the mouth of the river may be due to complex carbonate speciation. The minimum $p\text{CO}_2$ values are very low
440 (~200 μatm) in the inner part of the outflow (Figure 8d), while values in the outer part of the outflow are closer to the expected $p\text{CO}_2$ values around 350 μatm . (da Cunha and Buitenhuis, 2013). Mirroring the trends in pH, the calcite and aragonite saturation states (Figure 8e and 8f) are higher in the outflow relative to the open ocean. Anomalous pH, $p\text{CO}_2$ and the calcite and aragonite saturation states values in December 2003 are likely associated with the heavy rainfall over south-central Africa (Kadomura, 2005).

445




Figure 8: Hovmöller plots of (a) DIC, (b) TA, (c) pH, (d) $p\text{CO}_2$, (e) Ω Aragonite and (f) Ω Calcite for the Congo outflow region. The plots are centred on the 6° S slice that spans 2° W to 12° E. $\text{pH} > 8.4$, $p\text{CO}_2$ data $< 200 \mu\text{atm}$, Ω Calcite > 5 and Ω Calcite > 7 in the river outflow in December 2003 are shown in mint green. The plots spans the temporal overlap period of the TA and DIC datasets 2002- 2016 and thus the period for which the rest of the carbonate system was generated.

450

The mean DIC and TA across the whole Congo region, in the outflow and out of the outflow are very consistent year to year (Figure 9a and 9b). The yearly TA and DIC minima occur at approximately the same time and are the same magnitude most years. DIC and TA values are always higher in the non-outflow region than in the outflow. This consistency is also seen in the propagated variables pH (Figure 9c), $p\text{CO}_2$ (Figure 9d), and the aragonite and calcite saturation states (Figure 9e and 9f) all of which show very minor differences between the outflow and non-outflow regions.

455




Figure 9: Timeseries of spatially averaged (a) DIC, (b) TA, (c) pH, (d) $p\text{CO}_2$, (e) Ω Aragonite and (f) Ω Calcite in the Congo region. The plots spans the temporal overlap period of the TA and DIC datasets 2002- 2016. Data are averaged across the whole region (solid line) as well as in the outflow defined as $S < 35$ (dashed line) and outside of the outflow $S > 35$ (dotted line). The line colour corresponds to variables that were calculated with SST and SSS datasets selected during the DIC algorithm evaluation (red) and the TA algorithm evaluation (blue).

3.4 Uncertainty assessment

The TA and DIC combined standard uncertainties from the algorithm evaluation are shown in Table 1 and 2. The combined standard uncertainties for the optimal algorithms used to generate the time series dataset are reproduced in Table 3. The combined standard uncertainties are the first Type A uncertainty evaluation, which can be thought of as a “top down” estimate. The spatially averaged uncertainties of the remaining carbonate system variables using this top down estimate are shown in both outflow regions (Table 3). The uncertainties of the remaining carbonate system variables appear to be weakly dependent on the choice of SST and SSS datasets used to calculate them in PyCO2SYS (Table 3). Uncertainties are provided as absolute values rather than as percentages because of the steep gradient in values across the plume, but if required the uncertainty values can be expressed as a percentage using the minimum, maximum and mean values from table S4.

The second Type A uncertainty evaluation (using the algorithm RMSD from the literature and the uncertainties in the inputs) can be thought of as a “bottom up” estimate. For this bottom up estimate, the average combined standard uncertainties in TA and DIC are smaller for TA (8.52) and DIC (16.59) in the Amazon and in the Congo for TA (17.23) and DIC (14.25). These values are much lower than from the top down uncertainty evaluation as this uncertainty evaluation does not fully account for spatial and depth variability (by assuming that the literature RMSD values perfectly capture the variability), these values also do not account for measurement variability. The bottom up evaluation does account for uncertainty in the input SST and SSS datasets suggesting spatial and depth variabilities and differences between the in situ data (used to evaluate the uncertainties) are likely dominating the uncertainty budget

As the top down uncertainty evaluation is more robust it is the preferred uncertainty estimate. By comparing the uncertainties with the natural variability in TA and DIC (Table 4) it is clear that the dataset uncertainty is less than the natural variability data in both riverine regions. Whereas, the propagated uncertainties for $p\text{CO}_2$ and pH are larger than the natural variability, this is only because the $p\text{CO}_2$ and pH data generated in OceanSODA-UNEXE has full spatio temporal coverage (including in the high variability plume) compared to the $p\text{CO}_2$ and pH *in situ* data in the MDB.

Table 3: Output variable uncertainties in the two regions of OceanSODA-UNEXE. TA and DIC uncertainties are the combined standard uncertainties from the algorithm evaluation. The uncertainty for the remaining carbonate system variables are the average of the spatially varying propagated uncertainties for each of those variables calculated with the SST and SSS pair from either the TA or DIC algorithm. Note that due to the logarithmic nature of pH, uncertainty is calculated as 1σ in pH units and also in pH units where variability was determined in H^+ as $-\log_{10}(\overline{H^+}) + \log_{10}(\overline{H^+} + 1\sigma \text{H}^+)$.

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Variable	Using SSS and SST from TA algorithm							Using SSS and SST from DIC algorithm				
	TA	DIC	pCO ₂	pH converted from H ⁺	pH	ΩCal	ΩArag	pCO ₂	pH converted from H ⁺	pH	ΩCal	ΩArag
Combined uncertainty in the Amazon	34.74	44.34	85.19	0.08 relative to a pH of 8.20	0.07 relative to a pH of 8.19	0.91	0.61	86.17	0.08 relative to a pH of 8.19	0.08 relative to a pH of 8.19	0.91	0.60
Combined uncertainty in the Congo	28.54	33.25	73.14	0.07 relative to a pH of 8.21	0.08 relative to a mean pH of 8.21	0.79	0.52	74.02	0.07 relative to a pH of 8.20	0.08 relative to a mean pH of 8.20	0.79	0.52

Table 4: *In situ* measurement variabilities for TA, DIC, pH and pCO₂ are calculated as the standard deviation of the reference output (MDB). For the Amazon ESACCI SST and ESACCI SSS were the combination used with the MDB and for the Congo this was CORA SST and ISAS SSS. Note that due to the logarithmic nature of pH, uncertainty is calculated as 1σ in pH units and also in pH units where variability was determined in H⁺ as $-\log_{10}(\overline{H^+}) + \log_{10}(\overline{H^+} + 1\sigma H^+)$.

Variable	TA	DIC	pCO ₂	pH converted from H ⁺	pH	ΩCal	ΩArag
<i>In situ</i> measurement variability (1σ) in the Amazon from the MDB	104.47	102.96	25.86	0.037 relative to a pH of 8.07	0.043 relative to a mean pH of 8.07	–	–
<i>In situ</i> measurement variability (1σ) in the Congo from the MDB	46.13	51.71	31.36	0.003 relative to a pH of 8.03	0.003 relative to a mean pH of 8.03	–	–

4 Discussion

The algorithm evaluation (Figure 1) demonstrated that the choice of input SSS dataset to the TA and DIC algorithms makes the biggest difference in reducing the RMSDe in TA and DIC ~~(whereas SSS tends to be a secondary term in the majority of the algorithms so has less of a controlling effect)~~. The choice of literature TA and DIC algorithm itself impacts RMSDe to a much smaller extent than the choice of input SSS datasets. The prominence of SSS terms in the majority of the algorithms explains why the RMSDe is much more sensitive to the choice of SSS dataset compared to SST dataset, as SST is a secondary term in the majority of the algorithms. Whilst some algorithms did perform better than others the differences were so slight that it may not be that helpful to declare which of the algorithms are the best outright especially as this could change in the future with more data.

The requirement of having $n=30$ matchups to calculate the weighted statistics has a large impact on the choice of optimal algorithms. By including this stipulation, some of the more recent salinity input datasets are effectively de-selected as there are not enough contemporary measurements over their temporal range to allow their evaluation ~~(particularly in the Congo region)~~. For example, the SMAP satellite only launched in 2015 and there were not enough *in situ* matchups in either region to be able to fully assess the RSS-SMAP dataset. So using ISAS for salinity in the optimal TA algorithm in the Congo where data are available from 2002 onwards) increases the number of matchups by a factor of twenty compared to using RSS-SMAP (Table 2). Given time, the number of matchups with the satellite only products will increase allowing them to be fully evaluated. Considering the better spatial coverage of the satellite only products, it is likely that in the near future the best algorithm combinations to generate updates to this product will all use satellite derived SSS and SST. The recent proliferation of certified reference materials when measuring TA (Dickson et al., 2003) and DIC (Dickson, 2001) should also mean that newer *in situ* data will have lower uncertainties. The relatively low numbers amount of insitu data limits the potential use also a key factor in why empirical algorithms were chosen over state of the se of art neural network based approaches to estimate generate the carbonate system in these outflows and significantly more in situ data would be needed (for training and testing) before these approaches could likely be applied. There is currently insufficient training data in these outflows for neural network based approaches to accurately generate the carbonate system. Empirical algorithms that were derived with data from these outflows outperformed a global machine learning based approach of Gregor and Gruber (2021) in both these outflows.

By comparing the two type A uncertainty estimates, the standard combined uncertainty estimate (accounting for measurement uncertainty, spatial uncertainty, depth uncertainty and algorithm uncertainty, aka “top down”) and the second type A uncertainty evaluation (propagated input uncertainties through the literature algorithm, aka “bottom up”); the contribution of different factors to the uncertainty can be dissected. The bottom up uncertainty estimates are much smaller

530 than the top down uncertainty estimates reflecting the fact that less of the uncertainty has been considered in the bottom up
uncertainty estimates. The difference between the top down and bottom up estimates can be mainly attributed to the spatial
and depth uncertainty which is accounted for by using the MDB in the algorithm evaluation. Land et al. (2019) note that
reducing the uncertainty of the *in situ* measurements is just as important as all the remaining uncertainties in their full
methodology. Further reducing the standard combined uncertainty is challenging and would require either improvements in
535 satellite SSS retrievals (Vinogradova et al., 2019) or improvements in how uncertainty is quantifying in the GLODAP data (a
huge challenge give the different systems and protocols used by different laboratories) (Bockmon and Dickson, 2015).

The timeseries data demonstrates that there is low TA and DIC in the river outflows of both regions. The timeseries data
clearly shows that the discharge of the outflows and their zone of influence changes seasonally. Our results are consistent
540 with previous observations that the Amazon plume extent is smallest between January and March (Fournier et al., 2015) and
largest in the between April and June where it expels low TA and DIC waters (Cooley and Yager, 2006). The Amazon
plume reaches the Caribbean, as previously shown by Hellweger and Gordon (2002). The timeseries data identify that in
parts of the Amazon plume saturation states can drop below 3 (Table S4) which has implications for OA research, especially
for researchers studying coral reefs. The non-plume DIC and TA trends of 0.49 and 0.76 $\mu\text{mol kg}^{-1}$ per year are consistent
545 with the decadal increases in salinity normalised DIC (4.54 $\mu\text{mol kg}^{-1}$) and salinity normalised TA (7.39 $\mu\text{mol kg}^{-1}$) in the
2010s from the Bermuda Atlantic Time-series Study site (Bates and Johnson, 2020). The results identify that the discharge of
the Congo outflow is greatest between January and March and veers towards West Africa, which is consistent with work by
Hopkins et al. (2013).

550 The timeseries data comprises over ten years of carbonate system data for two of the world's largest rivers by discharge.
One immediate use for the timeseries data would be to assess the inorganic carbon flow of both of these rivers over time.
The timeseries data could also be used to investigate the impact of changes in land use within the river basin, e.g.
deforestation impacts on river discharge and inorganic carbon flow into rivers (Bass et al., 2014). If the impact of land use
555 changes were identifiable within the data then these approaches may prove a useful tool in monitoring the effectiveness of
policy or management actions addressing climate change and biodiversity loss in the Amazon Basin. The timeseries data
could also be used for ocean acidification research (Land et al., 2015a), as these rivers discharge enough fresh water to
influence carbonate saturation states. The tropical reefs of the Caribbean are infrequently impacted by the Amazon plume
(Chérubin and Richardson, 2007); the impact of the low pH waters on reef health is of great concern (Hoegh-Guldberg et al.,
560 2007) and could be explored with the timeseries data. Ocean acidification has been shown to impact foraging behaviour in
fish (Jiahuan et al., 2018) and sharks (Rosa et al., 2017), the timeseries data could be used in conjunction with GPS tracks of
fish or marine mammals to see if the outflows alter foraging behaviour. The timeseries data could also be used to explore

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

CO₂ fluxes in the Amazon outflow building upon more recent estimates (Olivier et al., 2022; Ibánhez et al., 2016; Mu et al., 2021).

565

5 Conclusions

OceanSODA-UNEXE is a time series dataset of the carbonate system in the outflow regions of the Amazon and Congo rivers. Optimal TA and DIC data are generated with the optimal combination of published algorithms and input datasets determined by an exhaustive round robin inter-comparison evaluation. By using a specially designed matchup database for the algorithm evaluation, uncertainties due to spatial and depth variability in the *in situ* references have been minimised. TA, DIC, SST and SSS are used as inputs into PyCO2SYS to calculate the ~~complete-remaining~~ carbonate system variables (pH and pCO₂). TA and DIC are provided with standard combined uncertainties from a Type A uncertainty evaluation whereas pH and pCO₂ are provided with propagated uncertainties from PyCO2SYS. The assessed uncertainties are lower than the natural variability within these regions and the main features of both river outflows are evident in all of the carbonate system variable outputs. Potential uses for these data could include evaluating the riverine carbon flux from the land into the ocean resulting from the Amazon and Congo rivers, or evaluating the extent that river-driven episodic changes in the carbonate system may be having on sensitive coral reefs that interact with the outflows.

570

575

Data availability

The dataset described in this paper is freely available at <https://doi.pangaea.de/10.1594/PANGAEA.946888>. The matchup database of (Land et al., 2022) which is needed to reproduce the algorithm evaluation is available at <https://data-cersat.ifremer.fr/data/ocean-carbonate/oceansoda-mmdb/>. All the remote sensing datasets needed to create the gridded products are all freely available from their respective online repositories.

580

Code availability

The code used to run this analysis and download the remote sensing datasets is provided in the supplement and is freely available at <https://github.com/Richard-Sims/OceanSODA>, upon acceptance of this manuscript the final version of the code will be issued a version number and DOI and will be located at <https://github.com/JamieLab/OceanSODA>. The code can be run on a desktop computer and requires no specialist computing facilities. For example, a laptop with Intel i7-4800MQ 2.70GHz CPU and 8 GB of memory can complete the algorithm evaluation in around one hour. On the same machine the

585

590

remote sensing datasets take several days to download and reprocess, this is partially subject to local internet speeds and host server speeds. Creating the gridded datasets for each region takes two hours on the same machine.

Acknowledgements

This work was funded by the European Space Agency via the OceanSODA project (4000112091/14/I-LG). We would like to
595 thank our OceanSODA colleges, Helen Findlay, Luke Gregor and Nicolas Gruber for their helpful discussions and feedback
on this work. Hannah Green was supported by a Ph.D. studentship funded by an AXA XL Ocean Risk Scholarship which
was awarded to Helen Findlay, Jamie Shutler, and Peter Land.

Author contributions

~~This manuscript was written by~~ Richard Sims led the writing of the manuscript, with inputs from Thomas Holding and Jamie
600 Shutler, and all co-authors made contributions to the final paper. Thomas Holding wrote the majority of the code used for the
algorithm evaluation and the creation of the dataset, Richard Sims made additional changes to the code for generating the
final dataset. Hannah Green assisted with debugging and testing the code and outputs. Peter Land and Jean Francois
provided essential updates to the matchup database. Thomas Holding performed the literature search to identify relevant
algorithms from the literature algorithms. Richard Sims produced the final manuscript figures partially based on earlier
605 versions produced by Thomas Holding. Jamie Shutler secured project funding and oversaw completion of the work.

References

- Allen, G. H., and Pavelsky, T. M.: Global extent of rivers and streams, *Science*, 361, 585-588,
<https://doi.org/10.1126/science.aat0636>, 2018.
- 610 Bakker, D. C., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., and Jones, S. D.: A
multi-decade record of high-quality fCO₂ data in version 3 of the Surface Ocean CO₂ Atlas (SOCAT), *Earth System Science*
Data, 8, 383, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Banzon, V., Smith, T. M., Chin, T. M., Liu, C., and Hankins, W.: A long-term record of blended satellite and in situ sea-surface
temperature for climate monitoring, modeling and environmental studies, *Earth System Science Data*, 8, 165-176,
<https://doi.org/10.5194/essd-8-165-2016>, 2016.
- 615 Bass, A. M., Munksgaard, N. C., Leblanc, M., Tweed, S., and Bird, M.: Contrasting carbon export dynamics of human impacted
and pristine tropical catchments in response to a short-lived discharge event, *Hydrological Processes*, 28, 1835-1843,
<https://doi.org/10.1002/hyp.9716>, 2014.
- Bates, N. R., and Johnson, R. J.: Acceleration of ocean warming, salinification, deoxygenation and acidification in the surface
subtropical North Atlantic Ocean, *Communications Earth & Environment*, 1, 1-12, 2020.
- 620 Bockmon, E. E., and Dickson, A. G.: An inter-laboratory comparison assessing the quality of seawater carbon dioxide
measurements, *Marine Chemistry*, 171, 36-43, <https://doi.org/10.1016/j.marchem.2015.02.002>, 2015.
- Bouillon, S., Yambélé, A., Gillikin, D. P., Teodoru, C., Darchambeau, F., Lambert, T., and Borges, A. V.: Contrasting
biogeochemical characteristics of the Oubangui River and tributaries (Congo River basin), *Scientific Reports*, 4, 1-10,
<https://doi.org/10.1038/srep05402>, 2014.

- 625 Boutin, J., Reul, N., Koehler, J., Martin, A., Catany, R., Guimbard, S., Rouffi, F., Vergely, J. L., Arias, M., Chakroun, M., Corato, G., Estella-Perez, V., Hasson, A., Josey, S., Khvorostyanov, D., Kolodziejczyk, N., Mignot, J., Olivier, L., Reverdin, G., Stammer, D., Supply, A., Thouvenin-Masson, C., Turiel, A., Vialard, J., Cipollini, P., Donlon, C., Sabia, R., and Mecklenburg, S.: Satellite-Based Sea Surface Salinity Designed for Ocean and Climate Studies, *Journal of Geophysical Research: Oceans*, 126, e2021JC017676, <https://doi.org/10.1029/2021JC017676>, 2021.
- 630 Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., Chou, W.-C., Zhai, W., Hollibaugh, J. T., and Wang, Y.: Acidification of subsurface coastal waters enhanced by eutrophication, *Nature Geoscience*, 4, 766-770, <https://doi.org/10.1038/ngeo1297>, 2011.
- Cai, W.-J., Feely, R. A., Testa, J. M., Li, M., Evans, W., Alin, S. R., Xu, Y.-Y., Pelletier, G., Ahmed, A., and Greeley, D. J.: Natural and anthropogenic drivers of acidification in large estuaries, *Annual Review of Marine Science*, 13, 23-55, <https://doi.org/10.1146/annurev-marine-010419-011004>, 2021.
- 635 Cai, W. J., Hu, X., Huang, W. J., Jiang, L. Q., Wang, Y., Peng, T. H., and Zhang, X.: Alkalinity distribution in the western North Atlantic Ocean margins, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2009JC005482>, 2010.
- Cattano, C., Claudet, J., Domenici, P., and Milazzo, M.: Living in a high CO₂ world: A global meta analysis shows multiple trait mediated fish responses to ocean acidification, *Ecological Monographs*, 88, 320-335, <https://doi.org/10.1002/ecm.1297>, 2018.
- 640 Chao, Y., Farrara, J. D., Schumann, G., Andreadis, K. M., and Moller, D.: Sea surface salinity variability in response to the Congo river discharge, *Continental Shelf Research*, 99, 35-45, <https://doi.org/10.1016/j.csr.2015.03.005>, 2015.
- Chérubin, L., and Richardson, P. L.: Caribbean current variability and the influence of the Amazon and Orinoco freshwater plumes, *Deep Sea Research Part I: Oceanographic Research Papers*, 54, 1451-1473, <https://doi.org/10.1016/j.dsr.2007.04.021>, 2007.
- 645 Coles, V. J., Brooks, M. T., Hopkins, J., Stukel, M. R., Yager, P. L., and Hood, R. R.: The pathways and properties of the Amazon River Plume in the tropical North Atlantic Ocean, *Journal of Geophysical Research: Oceans*, 118, 6894-6913, <https://doi.org/10.1002/2013JC008981>, 2013.
- Cooley, S., and Yager, P.: Physical and biological contributions to the western tropical North Atlantic Ocean carbon sink for med by the Amazon River plume, *Journal of Geophysical Research: Oceans*, 111, <https://doi.org/10.1029/2005JC002954>, 2006.
- 650 da Cunha, L. C., and Buitenhuis, E.: Riverine influence on the tropical Atlantic Ocean biogeochemistry, *Biogeosciences*, 10, 6357-6373, <https://doi.org/10.5194/bg-10-6357-2013>, 2013.
- Dai, A., and Trenberth, K. E.: Estimates of freshwater discharge from continents: Latitudinal and seasonal variations, *Journal of hydrometeorology*, 3, 660-687, [https://doi.org/10.1175/1525-7541\(2002\)003<0660:EOFDFC>2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003<0660:EOFDFC>2.0.CO;2), 2002.
- Dickson, A. G., and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep Sea Research Part A. Oceanographic Research Papers*, 34, 1733-1743, [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5), 1987.
- 655 Dickson, A. G.: Standard potential of the reaction: $\text{AgCl (s)} + 12\text{H}_2\text{(g)} = \text{Ag (s)} + \text{HCl (aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K, *The Journal of Chemical Thermodynamics*, 22, 113-127, [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.
- Dickson, A. G.: Reference materials for oceanic CO₂ measurements, *Oceanography*, 14, 21-22, 2001.
- 660 Dickson, A. G., Afghan, J., and Anderson, G.: Reference materials for oceanic CO₂ analysis: a method for the certification of total alkalinity, *Marine Chemistry*, 80, 185-197, [https://doi.org/10.1016/S0304-4203\(02\)00133-0](https://doi.org/10.1016/S0304-4203(02)00133-0), 2003.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other CO₂ problem, *Marine Science*, 1, 169-192, <https://doi.org/10.1146/annurev.marine.010908.163834>, 2009.
- 665 Doney, S. C., Busch, D. S., Cooley, S. R., and Kroeker, K. J.: The impacts of ocean acidification on marine ecosystems and reliant human communities, *Annual Review of Environment and Resources*, 45, 83-112, <https://doi.org/10.1146/annurev-environ-012320-083019>, 2020.
- Dong, X., Huang, H., Zheng, N., Pan, A., Wang, S., Huo, C., Zhou, K., Lin, H., and Ji, W.: Acidification mediated by a river plume and coastal upwelling on a fringing reef at the east coast of Hainan Island, Northern South China Sea, *Journal of Geophysical Research: Oceans*, 122, 7521-7536, <https://doi.org/10.1002/2017JC013228>, 2017.
- 670 Enochs, I., Formel, N., Manzello, D., Morris, J., Mayfield, A., Boyd, A., Kolodziej, G., Adams, G., and Hendee, J.: Coral persistence despite extreme periodic pH fluctuations at a volcanically acidified Caribbean reef, *Coral reefs*, 39, 523-528, <https://doi.org/10.1007/s00338-020-01927-5>, 2020.
- Ford, D., Tilstone, G. H., Shutler, J. D., Kitidis, V., Lobanova, P., Schwarz, J., Poulton, A. J., Serret, P., Lamont, T., and Chuqui, M.: Wind speed and mesoscale features drive net autotrophy in the South Atlantic Ocean, *Remote Sensing of Environment*, 260, 112435, <https://doi.org/10.1016/j.rse.2021.112435>, 2021.
- 675 Fournier, S., Chapron, B., Salisbury, J., Vandemark, D., and Reul, N.: Comparison of spaceborne measurements of sea surface salinity and colored detrital matter in the Amazon plume, *Journal of Geophysical Research: Oceans*, 120, 3177-3192, <https://doi.org/10.1002/2014JC010109>, 2015.

- 680 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W.,
and Pongratz, J.: Global carbon budget 2021, *Earth System Science Data*, 14, 1917-2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.
- Gaillard, F., Reynaud, T., Thierry, V., Kolodziejczyk, N., and Von Schuckmann, K.: In situ-based reanalysis of the global ocean
temperature and salinity with ISAS: Variability of the heat content and steric height, *Journal of Climate*, 29, 1305-1323,
<https://doi.org/10.1175/JCLI-D-15-0028.1> 2016.
- 685 Gregor, L., and Gruber, N.: OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate system for seasonal to
decadal studies of ocean acidification, *Earth System Science Data*, 13, 777-808, <https://doi.org/10.5194/essd-13-777-2021>, 2021.
- Grodsky, S. A., Reverdin, G., Carton, J. A., and Coles, V. J.: Year-to-year salinity changes in the Amazon plume: Contrasting
2011 and 2012 Aquarius/SACD and SMOS satellite data, *Remote Sensing of Environment*, 140, 14-22,
<https://doi.org/10.1016/j.rse.2013.08.033>, 2014.
- 690 Guinotte, J., Buddemeier, R., and Kleypas, J.: Future coral reef habitat marginality: temporal and spatial effects of climate
change in the Pacific basin, *Coral reefs*, 22, 551-558, <https://doi.org/10.1007/s00338-003-0331-4>, 2003.
- Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C., Bopp, L., Chau, T. T. T., Gürses, Ö., Ilyina, T., and Landschützer,
P.: Consistency and challenges in the ocean carbon sink estimate for the global carbon budget, *Frontiers in Marine Science*, 7, 852,
<https://doi.org/10.3389/fmars.2020.571720>, 2020.
- 695 Hellweger, F. L., and Gordon, A. L.: Tracing Amazon river water into the Caribbean Sea, *Journal of Marine Research*, 60, 537-
549, <https://doi.org/10.1357/002224002762324202>, 2002.
- Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., Harvell, C. D., Sale, P. F., Edwards, A.
J., and Caldeira, K.: Coral reefs under rapid climate change and ocean acidification, *Science*, 318, 1737-1742,
<https://doi.org/10.1126/science.1152509>, 2007.
- 700 Hopkins, J., Lucas, M., Dufau, C., Sutton, M., Stum, J., Lauret, O., and Channelliere, C.: Detection and variability of the Congo
River plume from satellite derived sea surface temperature, salinity, ocean colour and sea level, *Remote Sensing of Environment*,
139, 365-385, <https://doi.org/10.1016/j.rse.2013.08.015>, 2013.
- Hu, C., Montgomery, E. T., Schmitt, R. W., and Muller-Karger, F. E.: The dispersal of the Amazon and Orinoco River water in
the tropical Atlantic and Caribbean Sea: Observation from space and S-PALACE floats, *Deep Sea Research Part II: Topical
Studies in Oceanography*, 51, 1151-1171, <https://doi.org/10.1016/j.dsr2.2004.04.001>, 2004.
- 705 Hu, X., and Cai, W. J.: Estuarine acidification and minimum buffer zone—a conceptual study, *Geophysical research letters*, 40,
5176-5181, <https://doi.org/10.1002/grl.51000>, 2013.
- Huang, B., Liu, C., Banzon, V., Freeman, E., Graham, G., Hankins, B., Smith, T., and Zhang, H.-M.: Improvements of the daily
optimum interpolation sea surface temperature (DOISST) version 2.1, *Journal of Climate*, 34, 2923-2939,
<https://doi.org/10.1175/JCLI-D-20-0166.1> 2021.
- 710 Humphreys, M. P., Lewis, E. R., Sharp, J. D., and Pierrot, D.: PyCO2SYS v1. 8: marine carbonate system calculations in Python,
Geoscientific Model Development, 15, 15-43, <https://doi.org/10.5194/gmd-15-15-2022>, 2022.
- Ibáñez, J. S. P., Araujo, M., and Lefèvre, N.: The overlooked tropical oceanic CO₂ sink, *Geophysical research letters*, 43, 3804-
3812, <https://doi.org/10.1002/2016GL068020>, 2016.
- 715 Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for
surface fluxes of carbon dioxide: 2. Regional results, *Global Biogeochemical Cycles*, 21, <https://doi.org/10.1029/2006GB002703>,
2007.
- JCGM: Evaluation of measurement data—Guide to the expression of uncertainty in measurement, 134, 2008.
- 720 Jiahuan, R., Wenhao, S., Xiaofan, G., Wei, S., Shanjie, Z., Maolong, H., Haifeng, W., and Guangxu, L.: Ocean acidification
impairs foraging behavior by interfering with olfactory neural signal transduction in black sea bream, *Acanthopagrus schlegelii*,
Frontiers in physiology, 9, 1592, <https://doi.org/10.3389/fphys.2018.01592>, 2018.
- Kadomura, H.: Climate anomalies and extreme events in Africa in 2003, including heavy rains and floods that occurred during
northern hemisphere summer, *African study monographs. Supplementary issue 2005.*, 30, 165-181, <https://doi.org/10.14989/68453>,
2005.
- 725 Kaushal, S. S., Mayer, P. M., Vidon, P. G., Smith, R. M., Pennino, M. J., Newcomer, T. A., Duan, S., Welty, C., and Belt, K. T.:
Land use and climate variability amplify carbon, nutrient, and contaminant pulses: a review with management implications,
JAWRA Journal of the American Water Resources Association, 50, 585-614, <https://doi.org/10.1111/jawr.12204>, 2014.
- Land, P., Findlay, H., Shutler, J., Piolle, J.-F., Sims, R., Holding, T., and Green, H.: OceanSODA-MDB: a standardised surface
ocean carbonate system dataset for model-data intercomparisons, *Earth Syst. Sci. Data Discuss.*, <https://doi.org/10.5194/essd-2022-129>, 2022.
- 730 Land, P. E., Shutler, J. D., Findlay, H. S., Girard-Arduin, F., Sabia, R., Reul, N., Piolle, J.-F., Chapron, B., Quilfen, Y., and
Salisbury, J.: Salinity from space unlocks satellite-based assessment of ocean acidification, 49, 1987-1994,
<https://doi.org/10.1021/es504849s>, 2015a.

- 735 Land, P. E., Findlay, H. S., Shuttler, J. D., Ashton, I. G., Holding, T., Grouazel, A., Girard-Arduin, F., Reul, N., Piolle, J.-F., and Chapron, B.: Optimum satellite remote sensing of the marine carbonate system using empirical algorithms in the global ocean, the Greater Caribbean, the Amazon Plume and the Bay of Bengal, *Remote Sensing of Environment*, 235, 111469, <https://doi.org/10.1016/j.rse.2019.111469>, 2019.
- Lauerwald, R., Laruelle, G. G., Hartmann, J., Ciais, P., and Regnier, P. A.: Spatial patterns in CO₂ evasion from the global river network, *Global Biogeochemical Cycles*, 29, 534-554, 2015.
- 740 Lee, K., Wanninkhof, R., Feely, R. A., Millero, F. J., and Peng, T. H.: Global relationships of total inorganic carbon with temperature and nitrate in surface seawater, *Global Biogeochemical Cycles*, 14, 979-994, <https://doi.org/10.1029/1998GB001087>, 2000.
- Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G. H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, *Geophysical research letters*, 33, <https://doi.org/10.1029/2006GL027207>, 2006.
- 745 Lefèvre, N., Diverrès, D., and Gallois, F.: Origin of CO₂ undersaturation in the western tropical Atlantic, *Tellus B: Chemical and Physical Meteorology*, 62, 595-607, <https://doi.org/10.1111/j.1600-0889.2010.00475.x>, 2010.
- Lefèvre, N., Flores Montes, M., Gaspar, F. L., Rocha, C., Jiang, S., De Araújo, M. C., and Ibáñez, J.: Net heterotrophy in the Amazon continental shelf changes rapidly to a sink of CO₂ in the outer Amazon plume, *Frontiers in Marine Science*, 4, 278, <https://doi.org/10.3389/fmars.2017.00278>, 2017.
- 750 Lewis, E., Wallace, D., and Allison, L. J.: Program developed for CO₂ system calculations, Carbon Dioxide Information Analysis Center, managed by Lockheed Martin Energy Research Corporation for the US Department of Energy Tennessee, 1998.
- López, R., López, J. M., Morell, J., Corredor, J. E., and Del Castillo, C. E.: Influence of the Orinoco River on the primary production of eastern Caribbean surface waters, *Journal of Geophysical Research: Oceans*, 118, 4617-4632, <https://doi.org/10.1002/jgrc.20342>.
- 755 Mathis, J., Cooley, S., Lucey, N., Colt, S., Ekstrom, J., Hurst, T., Hauri, C., Evans, W., Cross, J., and Feely, R.: Ocean acidification risk assessment for Alaska's fishery sector, *Progress in Oceanography*, 136, 71-91, <https://doi.org/10.1016/j.pocean.2014.07.001>, 2015.
- Mehrbach, C., Culberson, C. H., Hawley, J. E., and Pytkowicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18, 897-907, <https://doi.org/10.4319/lo.1973.18.6.0897>, 1973.
- 760 Meissner, T., Wentz, F. J., and Le Vine, D. M.: The Salinity Retrieval Algorithms for the NASA Aquarius Version 5 and SMAP Version 3 Releases, *Remote Sensing*, 10, 1121, <https://doi.org/10.3390/rs10071121>, 2018.
- Merchant, C. J., Embury, O., Bulgin, C. E., Block, T., Corlett, G. K., Fiedler, E., Good, S. A., Mittaz, J., Rayner, N. A., and Berry, D.: Satellite-based time-series of sea-surface temperature since 1981 for climate applications, *Scientific data*, 6, 223, <https://doi.org/10.1038/s41597-019-0236-x>, 2019.
- 765 Millero, F. J.: The carbonate system in marine environments, in: *Chemical Processes in Marine Environments*, edited by: Gianguzza, A., Pelizzetti, E., and Sammartano, S., Environmental Science Springer, Berlin, Heidelberg, 9-41, 2000.
- Mongin, M., Baird, M. E., Tilbrook, B., Matear, R. J., Lenton, A., Herzfeld, M., Wild-Allen, K., Skerratt, J., Margvelashvili, N., and Robson, B. J.: The exposure of the Great Barrier Reef to ocean acidification, *Nature communications*, 7, 10732, <https://doi.org/10.1038/ncomms10732>, 2016.
- 770 Mu, L., Gomes, H. d. R., Burns, S. M., Goes, J. I., Coles, V. J., Rezende, C. E., Thompson, F. L., Moura, R. L., Page, B., and Yager, P. L.: Temporal Variability of Air-Sea CO₂ flux in the Western Tropical North Atlantic Influenced by the Amazon River Plume, *Global Biogeochemical Cycles*, 35, e2020GB006798, <https://doi.org/10.1016/j.csr.2021.104348>, 2021.
- Olivier, L., Boutin, J., Reverdin, G., Lefèvre, N., Landschützer, P., Speich, S., Karstensen, J., Labaste, M., Noisel, C., and Ritschel, M.: Wintertime process study of the North Brazil Current rings reveals the region as a larger sink for CO₂ than expected, *Biogeosciences*, 19, 2969-2988, <https://doi.org/10.5194/bg-19-2969-2022>, 2022.
- 775 Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., and Hoppema, M.: The Global Ocean Data Analysis Project version 2 (GLODAPv2)—an internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297, <https://doi.org/10.5194/essd-8-297-2016>, 2016.
- 780 Orr, J., Epitalon, J.-M., and Gattuso, J.-P.: Comparison of ten packages that compute ocean carbonate chemistry, *Biogeosciences*, 12, 1483-1510, <https://doi.org/10.5194/bg-12-1483-2015>, 2015.
- Piao, S., Friedlingstein, P., Ciais, P., de Noblet-Ducoudré, N., Labat, D., and Zaehe, S.: Changes in climate and land use have a larger direct impact than rising CO₂ on global river runoff trends, *Proceedings of the National Academy of Sciences*, 104, 15242-15247, <https://doi.org/10.1073/pnas.0707213104>, 2007.
- 785 Regnier, P., Arndt, S., Goossens, N., Volta, C., Laruelle, G. G., Lauerwald, R., and Hartmann, J.: Modelling estuarine biogeochemical dynamics: from the local to the global scale, *Aquatic geochemistry*, 19, 591-626, <https://doi.org/10.1007/s10498-013-9218-3>, 2013.

- Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO₂ data, *Biogeosciences*, 15, 2481-2498, <https://doi.org/10.5194/bg-15-2481-2018>, 2018.
- 790 Rosa, R., Rummer, J. L., and Munday, P. L.: Biological responses of sharks to ocean acidification, *Biology letters*, 13, 20160796, <https://doi.org/10.1098/rsbl.2016.0796>, 2017.
- Taylor, J.: *Introduction to error analysis, the study of uncertainties in physical measurements*, University Science Books, Mill Valley, CA, 1997.
- 795 Uthicke, S., Furnas, M., and Lønborg, C.: Coral reefs on the edge? Carbon chemistry on inshore reefs of the Great Barrier Reef, *PloS one*, 9, e109092, <https://doi.org/10.1371/journal.pone.0109092>, 2014.
- Vinogradova, N., Lee, T., Boutin, J., Drushka, K., Fournier, S., Sabia, R., Stammer, D., Bayler, E., Reul, N., and Gordon, A.: Satellite salinity observing system: Recent discoveries and the way forward, *Frontiers in Marine Science*, 6, 243, <https://doi.org/10.3389/fmars.2019.00243>, 2019.
- 800 Waldbusser, G. G., Hales, B., and Haley, B. A.: Calcium carbonate saturation state: on myths and this or that stories, *ICES Journal of Marine Science*, 73, 563-568, <https://doi.org/10.1093/icesjms/fsv174>, 2016.
- Wang, Z. A., Biennu, D. J., Mann, P. J., Hoering, K. A., Poulsen, J. R., Spencer, R. G., and Holmes, R. M.: Inorganic carbon speciation and fluxes in the Congo River, *Geophysical research letters*, 40, 511-516, <https://doi.org/10.1002/grl.50160>, 2013.
- 805 Ward, N. D., Bianchi, T. S., Medeiros, P. M., Seidel, M., Richey, J. E., Keil, R. G., and Sawakuchi, H. O.: Where carbon goes when water flows: carbon cycling across the aquatic continuum, *Frontiers in Marine Science*, 4, 7, <https://doi.org/10.3389/fmars.2017.00007>, 2017.
- Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G., Landschützer, P., Woolf, D. K., and Goddijn-Murphy, L.: Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon inventory, *Nature communications*, 11, 4422, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.

810