

Compare Results

Old File:

DB_manuscript_20220530.pdf

36 pages (7.63 MB)

30/05/2022, 10:41:48

versus

New File:

DB_manuscript_20220127.pdf

31 pages (65.74 MB)

27/01/2022, 11:06:42

Total Changes

499

Text only comparison

Content

197 Replacements

123 Insertions

179 Deletions

Styling and Annotations

0 Styling

0 Annotations

[Go to First Change \(page 1\)](#)

Global inventory of the stable isotopic composition of methane surface emissions, augmented by new measurements in Europe

Malika Menoud¹, Carina van der Veen¹, Dave Lowry², Julianne M. Fernandez², Semra Bakkaloglu^{2,3}, James L. France², Rebecca E. Fisher², Hossein Maazallahi^{1,*}, Mila Stanisavljević^{4,*}, Jarosław Nęcki^{4,*}, Katarina Vinković^{5,*}, Patryk Łakomiec^{6,*}, Janne Rinne^{6,7,*}, Piotr Korbeń^{8,*}, Martina Schmidt^{8,*}, Sara Defratyka^{9,*}, Camille Yver-Kwok^{9,*}, Truls Andersen^{5,*}, Huilin Chen^{5,*}, and Thomas Röckmann¹

¹Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University, Utrecht, The Netherlands

²Greenhouse Gas Research Laboratory (GGRL), Royal Holloway, University of London, Egham TW20 0EX, United Kingdom

³Now at: Sustainable Gas Institute, Imperial College London, London SW7 1NA, United Kingdom

⁴Faculty of Physics and Applied Computer Science, AGH University of Science and Technology, Krakow, Poland

⁵Centre for Isotope Research, Energy and Sustainability Institute Groningen (ESRIG), University of Groningen, Groningen, Netherlands

⁶Department of Physical Geography and Ecosystem Science, Lund University, Sweden

⁷Natural Resources Institute Finland, Helsinki, Finland

⁸Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

⁹Laboratoire des Sciences du Climat et de l'Environnement (LSCE-IPSL) CEA-CNRS-UVSQ Université Paris Saclay, Gif-sur-Yvette, 91191, France

*These authors contributed equally to this work.

Correspondence: Malika Menoud (m.menoud@uu.nl)

Abstract. Recent climate change mitigation strategies rely on the reduction of methane (CH₄) emissions. $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$ measurements can be used to distinguish sources and thus to understand the CH₄ budget better. The CH₄ emission estimates by models are sensitive to the isotopic signatures assigned to each source category, so it is important to provide representative estimates of the different CH₄ source isotopic signatures worldwide.

- 5 We present new measurements of isotope signatures of various, mainly anthropogenic, CH₄ sources in Europe, which represent a substantial contribution to the global dataset of source isotopic measurements from the literature, especially for $\delta^2\text{H}_{\text{CH}_4}$. They improve the definition of $\delta^{13}\text{C}_{\text{CH}_4}$ from waste sources, and demonstrate the use of $\delta^2\text{H}_{\text{CH}_4}$ for fossil fuel source attribution. Results from previous studies were combined in a common database that we updated with our new measurements, as well as with additional literature. We found that microbial sources are generally well characterised. The large variability in fossil fuels isotopic compositions requires particular care in the choice of weighting criteria for the calculation of a representative global value. The global dataset could be further improved by measurements from African, South American and Asian countries, as well as more measurements from pyrogenic sources.

The final version of the European methane isotope database coupled with a global inventory of fossil and non-fossil $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$ source signature measurements, is available at: <https://doi.org/10.24416/UU01-4PO56T> (Menoud et al., 2021a).

1 Introduction

The current change of the earth's climate is mainly caused by the emissions of greenhouse gases from anthropogenic activities (IPCC, 2013; IPCC 2021, 2021a). Methane (CH_4) is a strong greenhouse gas, with a global warming potential 32 times that of CO_2 over 100 years (Etminan et al., 2016). The increase in CH_4 concentration has contributed to an average warming of 0.5°C in 2010-2019 compared to 1850-1900, which is slightly smaller than the contribution of CO_2 (IPCC 2021, 2021b). The global CH_4 mole fraction ($\chi(\text{CH}_4)$) in the atmosphere has drastically increased since 1984, when direct regular measurements started, changing from 1645 ppb to 1850 ppb in 2017 (Nisbet et al., 2019). Compared to pre-industrial times (before 1750), the global $\chi(\text{CH}_4)$ has increased by 160%, from 720 to 1850 ppb (IPCC 2021, 2021a).

In the past 30 years, we have not observed a steady growth of atmospheric CH_4 mole fraction. Instead the increase in $\chi(\text{CH}_4)$ levelled-off between 2000 and 2007, and has been increasing again since then, from 2014 at the highest rate since the 1980's (Nisbet et al., 2019). This renewed increase presents a significant threat to reaching the goals of the Paris agreement, and mitigation policies are now also targeting CH_4 emissions (Shindell et al., 2017; Mayfield et al., 2017; Nisbet et al., 2020). Efficient strategies require good knowledge of the different kinds of CH_4 sources, their location and relative contributions. While emission estimates are reported at a country-level using statistical indicators, atmospheric inversions, based on observations, can be used to verify the inventories (Houweling et al., 2000; Zavala-Araiza et al., 2015; Henne et al., 2016; Maasackers et al., 2019). But the results from two approaches, respectively called bottom-up and top-down, are not fully compatible, reflecting a lack in our understanding of the CH_4 cycle (Etiope and Schwietzke, 2019; Saunio et al., 2020; Stavert et al., 2021).

Measurements of CH_4 isotopologues provide additional constraints on the relative contribution of the various source categories, because CH_4 isotopic composition depends on the formation processes (Schoell, 1980; Whiticar, 1999; Quay et al., 1999). Time series of ambient CH_4 isotopic ratios are already used to derive emission scenarios in global models (Bousquet et al., 2006; Schaefer et al., 2016; Turner et al., 2017; Thompson et al., 2018; Fujita et al., 2020; Lan et al., 2021), and at the regional scale (Röckmann et al., 2016; Stieger et al., 2019; Menoud et al., 2020, 2021b; Varga et al., 2021). In addition, isotope measurements have proven to be very successful for source attribution in cities (Phillips et al., 2013; Zazzeri et al., 2017; Maazallahi et al., 2020; Xueref-Remy et al., 2020; Defratyka et al., 2021; Fernandez et al., in review), and larger regions (Tarasova et al., 2006; Fisher et al., 2011; Beck et al., 2012; Warwick et al., 2016; Fisher et al., 2017; Lu et al., 2021). The uncertainties in the resulting emission rates of the different source categories depend on our knowledge of the different isotopic source signatures, and understanding of their variability (Rigby et al., 2012; Schwietzke et al., 2016; Sznéási, 2020).

Direct measurements of the isotopic signature of CH_4 sources allow us to characterise them well, and a lot of data is available in the literature. Several review articles on CH_4 isotopic source signatures were previously published (Rice and Claypool, 1981; Cicerone and Oremland, 1988; Bréas et al., 2001). The most recent one presented by Sherwood et al. (2017), and recently updated in Sherwood et al. (2021), gathered values from 13 489 locations (10 778 fossil fuel, 2711 non-fossil) from 347 published references. The 2017 study focused on (fugitive) fossil fuel sources, and allowed to re-evaluate the global $\delta^{13}\text{C}_{\text{CH}_4}$ value assigned to this emission category towards more depleted values (Schwietzke et al., 2016). A disadvantage of this database

50 is that it is rather US-centered, and that the dataset is strongest for fossil fuel sources, but less robust^x for non-fossil sources. Therefore the database can be completed by more studies, especially concerning non-fossil sources.

The MEMO² project (MEthane goes MOBILE - MEasurements and MOdeling) was a H2020 MSCA European Training Network¹ with the goal to use innovative mobile measurement and **modelling** tools to improve the quantification of CH₄ 55 emissions in Europe (Walter et al., 2019). An important component of MEMO² was the isotopic characterisation of CH₄ sources. Two laboratories involved in MEMO², at Utrecht University, The Netherlands, and at the Royal Holloway University of London, UK, carried out a large number of high-precision measurements with isotope ratio mass spectrometry (IRMS). Another method, using cavity ring-down spectroscopy (CRDS) was developed for the mobile measurements of ambient CH₄ isotopic composition. Several research groups were involved in field work with mobile measurements that targeted specific 60 sources or environments in several European countries. Using this network,^x numerous CH₄ sources could be sampled for isotopic measurements. The resulting isotopic source signatures were gathered in a publicly available database.^x The European Methane Isotope Database.

This article presents the data collected within^x MEMO², and the implications for the global understanding of CH₄ source isotopic composition. To place the new data in context, we analyse it together with an updated version of the Sherwood et al. 65 (2017, 2021) global database of measured CH₄ source signatures.

Methods

2.1 Measurements within the MEMO² project

2.1.1 Sampling

The data was collected by the research teams of 8 universities and research institutes: Utrecht University (UU), the Royal 70 Holloway University of London (RHUL), the Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Heidelberg University (UHEI), AGH University of Science and Technology (AGH), Lund University (LU), the University of Groningen (UG), and the Netherlands Organisation for Applied Scientific Research (TNO). They participated in several campaigns in the Netherlands, the United Kingdom, France, Germany, Poland, Sweden, Romania and Turkey. Several other teams collaborated in two intensive campaigns: the CoMet² campaign in the Upper Silsian Coal Basin (USCB) in Poland (Fiehn et al., 2020; 75 Galkowski et al., 2021), and the ROMEO campaign in Romania³ (Röckmann, 2020).

Different sampling methods were used:

¹Marie Skłodowska-Curie actions, Horizon 2020 Innovative Training Networks founded under the grant agreement No 722479; <https://cordis.europa.eu/project/id/722479>

²Carbon dioxide and Methane mission, May-June 2018

³Romanian Methane Emissions from Oil & gas, October 2019

- 80 – Mobile sampling on a road vehicle, using a fast analyser (0.1 to 10 Hz) on-board to detect CH₄ enhancements (G2301, G2201-i, and G4302, Picarro Inc., USA; MGGA-918 and UGGA, Los Gatos Research, ABB, USA; LI-7810 Trace Gas analyser, LI-COR, USA; Dual Laser Trace Gas Monitor, Aerodyne Research, USA). The samples were taken using a small electric pump connected to an inlet outside of the vehicle. The sample receptacles were bags of 1 to 3 L (SupelTM-Inert Multi-Layer Foil bags, Sigma-Aldrich Co. LLC, USA; Tedlar or FlexFoil sample bags, SKC Inc., USA). Surveys were made around known sources of CH₄, where we sampled the elevated mole fractions as well as background CH₄ on the same day. If it was not practical to approach a source with the vehicle during mobile surveys, samples were taken on
- 85 foot.
- Mobile sampling onboard of an aircraft, during the ROMEO campaign. A CRDS instrument (G4302, Picarro Inc., USA) was installed in the aircraft, and samples were taken from the outflow of the instrument into bags of 2 L (SupelTM-Inert Multi-Layer Foil bags, Sigma-Aldrich Co. LLC, USA) when an increase in CH₄ mole fractions was observed. The method is described in detail in Menoud et al. (in review).
- 90 – Mobile sampling on foot, without analyser. The samples were taken at regularly spread locations around a known CH₄ source, to make sure we collected air with CH₄ from the emission plume and background. In this case, the sample receptacles were bags of 2 to 3 L (SupelTM-Inert Multi-Layer Foil bags, Sigma-Aldrich Co. LLC, USA; Tedlar sample bags, SKC Inc., USA), filled with a portable hand pump.
- 95 – Soil chambers on wetlands in north Sweden and coal waste disposal areas in Poland. In wetlands, we installed transparent Plexiglas chambers on top of stainless steel collars that were pushed 20 cm into the peat. Samples from the chambers were taken during closure times, when $\chi(\text{CH}_4)$ increased, generally after 10 to 25 min. The soil chambers in Poland were made of plastic buckets covered with aluminum foil that were pushed about 5 cm in the ground and left for 30 min. In both cases, air was pumped into 2L sample bags (SupelTM-Inert Multi-Layer Foil, Sigma-Aldrich Co. LLC, USA) for further analysis in the lab.
- 100 – From an unmanned aerial vehicle (UAV), carrying an AirCore (coiled tubing) system to collect air samples (Andersen et al., 2018). The air samples were continuously pulled into the AirCore while flying transects across the plume of a CH₄ emission source, and were transferred to a 0.5 or 1 L bag sample after landing (SupelTM-Inert Multi-Layer Foil, Sigma-Aldrich Co. LLC, USA) for further analysis in the laboratory.

Some photos taken in the field illustrate the different sampling procedures that were used, and are available in Fig. A1 of the supplementary material.

105

2.1.2 Measurements of isotopic composition

The mass spectrometry measurements were performed at two laboratories: the IMAU (Institute for Marine and Atmospheric research Utrecht) at UU, and at the Department of Earth Sciences at RHUL. Both laboratories use a CF-IRMS (continuous flow isotopic ratio mass spectrometry) system to measure $\delta^{13}\text{C}$, and also $\delta^2\text{H}$ at IMAU. The system at IMAU was described

110 by Röckmann et al. (2016) and the one at RHUL by Fisher et al. (2006). The reproducibility both groups can achieve is of 0.05 to 0.1 ‰ for $\delta^{13}\text{C}_{\text{CH}_4}$. At IMAU, $\delta^2\text{H}$ measurements have a reproducibility lower than 2 ‰. For consistency of the results, the two laboratories measured a set of 5 cylinders that contained air with CH_4 of different isotopic composition. The resulting differences in $\delta^{13}\text{C}_{\text{CH}_4}$ for each cylinder ranged between 0.02 and 0.04 ‰. They were within the analytical error reported by the two laboratories, so that the isotopic results obtained within the MEMO² project are consistent across the laboratories. The inter-comparison exercise is presented in detail in a MEMO² deliverable report publically available⁴.

The UHEI and LSCE groups performed isotopic measurements using CRDS instruments (G2201-i, Picarro inc., USA). Their methods were described in Hoheisel et al. (2019) and Defratyka et al. (2021).

In the database, the method of isotopic measurements is specified by the "Measurement type" parameter, as either 'IRMS' or 'CRDS'. The laboratory where the measurements were performed is specified in the column "Measurement lab".

120 2.1 Calculation of isotopic signatures

The measurement results of $\delta^{13}\text{C}$ and $\delta^2\text{H}$ of CH_4 are for ambient air, and not the sources themselves. There are different methods to derive the isotopic source signatures from the sampled CH_4 enhancement signatures; the Keeling plot and Miller-Tans methods are commonly used mass balance approaches. The Keeling plot method is based on the assumption that the background is stable during the sampling period (Keeling, 1961; Pataki et al., 2003). The Miller-Tans method is also applicable when the condition of a stable background is not fulfilled (Miller and Tans, 2003). Because background samples were taken on each survey day and in the same region, the condition of stable background was usually fulfilled. Defratyka (2021) showed that in this case, both methods lead to similar results within their uncertainty.

Both methods involve a linear regression model to fit the observed data. Different models were used: ordinary least squares (OLS) minimising the difference in the y-axis coordinate, bivariate correlated errors and intrinsic scatter (BCES) (Akritas and Bershady, 1996), and ordinary distance regression (ODR) (Boggs and Rogers, 1990). Zobitz et al. (2006) compared different regression methods when applied in Keeling plots. The ODR can induce a bias towards lower values in the case the data points cover a relatively small range on the x-axis, so the OLS and BCES methods were usually preferred.

All the mass balance and regression methods are statistically valid. Therefore we did not work towards a uniform procedure, but the different approaches are specified for each entry of the database by the parameters "Mass balance approach" and "Regression method".

⁴Lowry, D., Röckmann, T., Fisher, R., Menoud, M., Fernandez, J., 2018. Isotopic measurements linked to common scale (Deliverable No. 2.1), WP2. MEMO²: MEthane goes MOBILE – MEasurements and MOdelling, Available at: <https://h2020-memo2.eu/wp-content/uploads/sites/198/2018/12/MEMO2-D2.1-Isotopic-measurements-linked-to-common-scale-final.pdf>.

2.2 Update of the global database

2.2.1 Structure of the database

We used the same parameters as in the database of Sherwood et al. (2017, 2021) for non-fossil data. That is because our objectives concern only values for $\delta^{13}\text{C}$ and $\delta^2\text{H}$ of emitted CH_4 , and do not include measurements of other gases or isotope signatures that Sherwood et al. (2017) reported in the fossil fuel database. The variables of interest are listed in Table 1 and include the site description (country, region, group, category and sub-category) and the $\delta^{13}\text{C}$ and $\delta^2\text{H}$ of CH_4 . We grouped the sources reported in the European Methane Isotope Database by region and sub-category in order to integrate it in the literature database. We kept the categories and sub-categories as defined in Sherwood et al. (2017, 2021), but when the new entries from MEMO² measurements and published literature required it, we added additional source categories or sub-categories.

The categories are grouped into the three main CH_4 formation pathways: modern microbial, pyrogenic, and fossil fuels. The "modern microbial" CH_4 is formed by microorganisms in surface ecosystems or in animals through enteric fermentation, and are referred simply as "microbial" throughout the paper. Microbial CH_4 formations in the subsurface related to petroleum systems belongs to the "fossil fuels" category. Compared to Sherwood et al. (2017, 2021), we extended the biomass burning type to include emissions from all combustion sources, such as traffic or industry (Table 1).

Table 1 Number of data entries ($\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$) per source category in the updated CH_4 isotopic signature database.

			Sherwood et al. (2017)	Additional literature	MEMO	
microbial	agriculture	ruminants	C3/C4	313	57	41
		rice paddies	flooded, flooded seasonally	499	15	
		piggery			20	
	waste	landfill		186	115	76
		sewage	wastewater, manhole	4	33	147
		biogas	manure, C4/C3	30	23	47
		manure	cattle		9	22
		compost				4
		abattoir	cattle		27	
	wetlands	temperate	marsh, bog, swamp, lake, estuary, pond, delta, fen, lagoon, reeds, flooded forest, wet prairie, river, mangrove	385	158	12
		tropical	floodplain, lake, swamp, marsh, river, riverine	199	94	
		polar (incl. boreal)	reeds, mixed bog, marsh, swamp, tundra, lake, estuary, fen, wet tundra, (thawn) permafrost, mire	557	72	30
	other	forest	boreal forest		2	
		termites		30	7	
	fossil fuels	exploitation	conventional	gas leak, gas installation, oil field, mixed, natural gas, oil refinery	8669	112
coal			active coal mine, inactive coal mine, coal seam	2904	184	111
shale		gas	737			
seeps	oceans	marine seep			8	
	coal seam gas			70		
	volcanoes			8		
pyrogenic	biomass burning	grass, pasture, brush, woodland, wood, forest, crop	113	2		
	fossil fuel burning	conventional	car, traffic, residential heating		71	5

150 2.2.2 Reported variables

The analytical parameters reported in the database are $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$, which are defined as:

$$\delta X = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right)$$

with $R = \frac{^{13}\text{C}}{^{12}\text{C}}$ for $X = ^{13}\text{C}$ or $R = \frac{^2\text{H}}{^1\text{H}}$ for $X = ^2\text{H}$

The unit is in per mille (‰), and values are relative to the international standard materials Vienna Pee Dee Belemnite (VPDB) for $\delta^{13}\text{C}$, and Vienna Standard Mean Ocean Water (VSMOW) for $\delta^2\text{H}$.

2.2.3 Literature data

We found additional data in the literature to complete the referred data listed in Sherwood et al. (2021). Because we aim at reflecting the actual CH_4 surface emissions to the atmosphere, we excluded studies that reported results from laboratory experiments, and of CH_4 dissolved in water (i.e. in oceans, wetlands and inland waters). We note that the search for data was biased because of the use of English language. The references we added concern published peer-reviewed articles and to a lesser extent thesis and conference papers. The studies were performed from 1982 to 2021 in various laboratories in the world. We did not perform additional data quality assessment.

2.3 Results and discussion

The data on isotopic source signatures from the measurement campaigns carried out within the MEMO² project (2017-2020) were compiled into one database: The European Methane Isotope Database. The first version was made accessible on October 1st 2020, and described in a publicly available report⁵. The European data was used in several publications over the past 2 years by Menoud et al. (2020, 2021b, in review); Maazallahi et al. (2020); Defratyka et al. (2021); Bakkaloglu et al. (2021, in review); Fernandez et al. (in review). The final version of the global database, including the MEMO² measurements and the additional literature are available at: <https://doi.org/10.24416/UU01-4PO56T>.

170 2.4 The European Methane Isotope Database

The isotopic signatures obtained within the MEMO² project concern 734 locations over 8 countries, with $\delta^2\text{H}$ source signatures being measured at 54 % of the sites (Table 2). Figure 1 shows the geographical distribution of the sampled sites in the different countries, according to the type of source. The number of sources we sampled does not necessarily represent the emission magnitudes.

⁵Menoud, M., Röckmann, T., Fernandez, J., Bakkaloglu, S., Lowry, D., Korben, P., Schmidt, M., Stanisavljevic, M., Necki, J., Defratyka, S., Kwok, C.Y., 2020. mamenoud/MEMO2_isotopes: v8.1 complete. Zenodo.

⁶Menoud, M., Röckmann, T., Lowry, D., Fernandez, J., 2020. Improved isotopic source signatures of local and regional CH_4 emissions (Deliverable No. 2.2), WP2. MEMO²: MEthane goes MOBILE – MEasurements and MOdelling. Available at: <https://h2020-memo2.eu/wp-content/uploads/sites/198/2021/03/MEMO2-D2.2-v3-final.pdf>.

Table 2. Number of locations where CH₄ isotopic source signatures were derived from sample measurements.

	$\delta^{13}\text{C}_{\text{CH}_4}$	$\delta^2\text{H}_{\text{CH}_4}$
The Netherlands	50	27
United Kingdom	240	54
Poland	98	73
Germany	73	23
France	46	23
Sweden	21	21
Romania	184	174
Turkey	2	0




Figure 1. Geographical distribution of isotopic signature measurements ($\delta^{13}\text{C}$ and/or $\delta^2\text{H}$ of CH₄) carried out within the MEMO² project (2017 to 2020), depending on the type of source. (a) All locations. (b) Only in the UK, Netherlands and Germany.

175 During mobile surveys, we mostly targeted anthropogenic emissions from the exploitation and use of fossil fuels and waste processing facilities (Fig. 1). These are the most obvious anthropogenic CH₄ sources in densely populated regions, and we acknowledge a deliberate sampling bias towards urbanised areas. No biomass burning emissions were characterised during the MEMO² project. The European Methane Isotope Database partially address the geographical bias pointed out by Sherwood et al. (2017): it particularly includes a large number of measurements made in Romania, where almost no data was available
180 before.

185 We characterised 376 locations by both $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values, and we compared the results to ranges reported in the literature in Fig. 2. The fossil fuel sources partly overlap with the range of thermogenic CH₄, but also spread towards lower $\delta^{13}\text{C}$ or higher $\delta^2\text{H}$. This is due to the presence of natural gas of microbial origin in the coal reservoirs of Silesia, in Poland (Kotarba, 2001; Kotarba and Pluta, 2009; Menoud et al., 2021b), as well as in Romania (Baciu et al., 2018; Fernandez et al., in review; Menoud et al., in review). We concluded that this microbial CH₄ originates from the CO₂ reduction pathway, because of a relatively enriched $\delta^2\text{H}$ (>-250 ‰), and relatively depleted $\delta^{13}\text{C}$ (<-60 ‰) (Milkov and Etiope, 2018). The $\delta^2\text{H}$ measurements were




Figure 2. Dual isotope plot presenting the new European Methane Isotope Database (circles); the literature data for the same source categories, taken from Sherwood et al. (2017, 2021) and completed with additional publications (squares); and shaded areas represent the different methanogenesis pathways from Milkov and Etiope (2018): MF = microbial fermentation, MC = microbial CO₂ reduction, T = thermogenic, A = abiotic.

in these cases particularly useful to distinguish fossil fuels from microbial sources (Menoud et al., 2021b; Fernandez et al., in review; Menoud et al., in review).

190 The waste-related source signatures are generally more enriched in $\delta^{13}\text{C}$ than the typical microbial fermentation range specified in previous reviews. The most enriched values concern sewage treatment plants and biogas plants. Changes in waste management practices towards less disposal and more biogas production can likely explain the higher $\delta^{13}\text{C}$ values found in recent studies (Bakkaloglu et al., 2021). A new study also reported surprisingly enriched $\delta^{13}\text{C}_{\text{CH}_4}$ (and $\delta^2\text{H}$) around a wastewater treatment plant in Australia: $\delta^{13}\text{C} = -47.6 \pm 2 \text{‰}$ (Lu et al., 2021). The $\delta^{13}\text{C}$ of CH_4 emitted from sewage treatment

195 plants depends on process parameters: oxic conditions lead to more enriched signatures than anaerobic treatment (Toyoda et al., 2011). Regarding biogas facilities, Bakkaloglu et al. (in review) emphasised the link between the type of substrate and the emitted CH_4 isotopic signatures: facilities that operate with C4 plant substrates emit CH_4 with higher $\delta^{13}\text{C}$ values in comparison with C3 plant substrates. Another driver for more or less enriched $\delta^{13}\text{C}_{\text{CH}_4}$ emissions from waste sources is isotopic fractionation when CH_4 reacts or diffuses. Diffusion and oxidation in the soil layers when CH_4 migrates from the deeper layers

200 are secondary processes that cause isotopic fractionation (Bergamaschi et al., 1998; De Visscher, 2004; Conrad, 2005; Gebert and Streese-Kleeberg, 2017; Obersky et al., 2018; Bakkaloglu et al., 2021), which increases the range of possible isotopic signatures of the emitted CH_4 .

205 The maps in Fig. 3 emphasise the similarities between $\delta^{13}\text{C}$ source signatures from modern microbial and fossil fuel sources in Poland and Romania. The average $\delta^{13}\text{C}_{\text{CH}_4}$ of fugitive emissions from coal, oil and gas extraction sites in Poland and Romania was $-48.5 \pm 0.6 \text{‰}$ (n=235), compared to $-38.9 \pm 0.3 \text{‰}$ (n=154) from gas leaks in only the UK and the Netherlands (and $-40.4 \pm 0.3 \text{‰}$ (n=217) including France and Germany). This distinction is also visible in the histograms of the Europe




Figure 3. Measured $\delta^{13}\text{C}$ (top) and $\delta^2\text{H}$ (bottom) in CH_4 from European anthropogenic sources sampled within MEMO². **A.** CH_4 fugitive emissions from the exploitation of fossil fuels (gas leaks, oil and gas extraction and processing sites). **B.** CH_4 emissions from modern microbial fermentation sources (ruminants, landfills, sewage treatment plants and biogas plants).

Methane Isotope Database in Fig. 5.A. In western Europe, $\delta^{13}\text{C}$ allows for a good separation between microbial and fossil fuel sources, which is well-established in the literature (Levin et al., 1993; Lowry et al., 2001; Röckmann et al., 2016; Zazzeri et al., 2017; Lowry et al., 2020). Yet we show that we can't use only $\delta^{13}\text{C}$ data to distinguish microbial and fossil fuel CH_4 from all European regions. Fortunately, the $\delta^2\text{H}_{\text{CH}_4}$ source signatures allow for a clear distinction between fossil fuel and modern microbial emissions of anthropogenic origin (Fig. 3 and 5.A).

2.5 Global data overview and representativeness

The extended global database including all literature data and the aggregated MEMO² data consists of 13313 and 4337 measurements of $\delta^{13}\text{C}$ and $\delta^2\text{H}$, respectively, from 64 countries. The map in Fig. 4 shows the partitioning of the data per country, and Table 1 the number of records per CH_4 source. The maps in Fig. A2 of the supplementary material illustrate the increase in number of measurements of each isotopologue.

The number of measurements made in fossil fuel reservoirs and compiled in the database by Sherwood et al. (2021) is comparatively larger than from studies of other CH_4 emission sources (Table 1), and the amount of measurements is not evenly spread geographically: significantly more measurements were made in North American and European countries, Australia, Brazil and Japan. In Russia and China, there were relatively more measurements as well, but only for fossil fuel sources. Despite including the first few measurements reported from Africa and the middle-east (France et al., 2021; Al-Shalan et al.,




Figure 4. Number of isotopic signature measurements ($\delta^{13}\text{C}$ and $\delta^2\text{H}$ of CH_4) carried out in different countries worldwide and reported in the present database.

2022), the data distribution remains unbalanced. Nevertheless, specific isotope signature dependencies can be further analysed for the different source categories:

225 **Fossil fuels** Fugitive emissions from fossil fuel reservoirs are highly variable not only on a large scale, but also from one basin to another, or even within the same basin (Sherwood et al., 2017; Milkov and Etiope, 2018; Lan et al., 2021). Therefore, CH_4 isotopic composition from one basin can't be simply upscaled to a country scale. Any new isotopic measurement from a production basin with large fugitive CH_4 emissions brings relevant information.

230 ✖ Sherwood et al. (2017) pointed out the lack of data for a list of conventional oil and gas and coal production countries, in Africa, the middle-east, central and southern Asia, and South America. Previous estimates of global CH_4 isotopic signatures from the exploitation of fossil fuels weighted the source signatures from one basin by its fuel production (Schwietzke et al., 2016). Recent work suggest that fuel production is not a reliable proxy to estimate CH_4 fugitive emissions (Zavala-Araiza et al., 2015; Alvarez et al., 2018; Rutherford et al., 2021; Chen et al., 2021; Maazallahi et al., 2021). Thus, the most relevant sampling locations would be ideally related to estimated emission rates from top-down measurements, instead of production or bottom-up emission estimates. Unfortunately, these data are lacking in many cases. Recently, particularly large CH_4 emissions were detected in central Asia (Varon et al., 2019), or measured in Mexico (Zavala-Araiza et al., 2021). ✖

235

Modern microbial The isotopic signatures of CH_4 from modern microbial sources (mainly wetlands, ruminants, waste degradation, rice paddies, termites) are largely dependent on environmental parameters such as the type of substrate and other ecosystem conditions. Figures A3 and A4 show that our new data confirm the trends previously observed: the $\delta^{13}\text{C}$ sensitivity to C3 or C4 plants in ruminant diet (Rust, 1981; Levin et al., 1993; Klevenhusen et al., 2010; Brownlow et al., 2017), to wetland latitudes ($\delta^{13}\text{C}$ depletion in polar regions because of less oxidation and the absence of C4 plants) (Fisher et al., 2017; Brownlow et al., 2017; Ganesan et al., 2018), and the $\delta^2\text{H}$ dependency on $\delta^2\text{H}_{\text{H}_2\text{O}}$ of precipitation,

240

and ultimately on the latitude (established for freshwater emissions) (Waldron et al., 1999; Chanton et al., 2006; Douglas et al., 2021; Stell et al., 2021). Based on the correlation with the plant metabolism (C3 or C4), $\delta^{13}\text{C}_{\text{CH}_4}$ from wetlands could be mapped on a global scale (Ganesan et al., 2018). Douglas et al. (2021) also suggested a spatial extrapolation of wetland $\delta^2\text{H}_{\text{CH}_4}$ using $\delta^2\text{H}_{\text{H}_2\text{O}}$ data, which can be interesting for under-sampled locations, such as the southern hemisphere. However, a certain variability will always remain because of the influence of other parameters such as the dominant methanogenic pathway (acetate fermentation or CO_2 reduction) (Waldron et al., 1998; De Visscher, 2004; Conrad, 2005; McCalley et al., 2014; Inglett et al., 2015; Chan et al., 2019; Douglas et al., 2021), or the composition of the substrate in organic matter for $\delta^{13}\text{C}$ (Conrad et al. (2011); Ganesan et al. (2018)).

2.6 Global data distribution

The global distribution of CH_4 isotopic signatures in the complete extended database is shown in Fig. 5. The values were grouped in categories that correspond to the largest reported emissions (Saunois et al., 2020). The categories agriculture, waste, wetlands, and partly other natural are all of modern microbial origin, mainly following the fermentation pathway (Milkov and Etiope, 2018). They show a normal distribution, and an overlap between the categories, except for the waste sources that are more enriched in ^{13}C . This difference is particularly visible in the MEMO² data, and from a relatively large number of sites from waste related sources. As mentioned in section 2.4, additional parameters control the isotopic signature of the emitted CH_4 , such as the type of substrate, the presence of oxygen, or secondary (e.g. oxidation) processes. We recommend to separate the waste category from the other microbial sources to minimise the uncertainty in the assigned isotopic signature. The fractionation factors derived for CH_4 microbial oxidation are larger for $\delta^2\text{H}$ (Coleman et al., 1981; Bergamaschi et al., 1998; Chanton et al., 2006), but we don't clearly see an influence of these additional parameters on the $\delta^2\text{H}_{\text{CH}_4}$ signatures of our dataset. Indeed, the range of $\delta^2\text{H}$ signatures for waste is the same for as agriculture and wetlands (Fig. 5), but these are based on few measurements compared to $\delta^{13}\text{C}$ (42 % of all measured waste sources reported $\delta^2\text{H}$ signatures). The relation between $\delta^2\text{H}_{\text{CH}_4}$ from wetlands and the $\delta^2\text{H}_{\text{H}_2\text{O}}$ from precipitation has been established (Waldron et al., 1999; Chanton et al., 2006; Douglas et al., 2021). But, further $\delta^2\text{H}$ measurements are required to better define the isotopic dependencies to secondary processes.

In Sherwood et al. (2017, 2021), the pyrogenic category only contained biomass burning data, and the binary distribution clearly illustrates the difference between C3 and C4 plants in terms of $\delta^{13}\text{C}_{\text{CH}_4}$ signatures. The additional biomass burning data we added from published literature confirms the dependency of $\delta^{13}\text{C}_{\text{CH}_4}$ on the plant metabolism. We also added pyrogenic data of fossil fuel burning from both our measurements and the literature. The resulting distribution of the $\delta^{13}\text{C}$ data is therefore smoother than in Sherwood et al. (2017) (Fig. 5), because $^{13}\text{C}_{\text{CH}_4}$ from fossil fuel burning is more variable than from biomass burning, and does not show a clear distinction between C3/C4 plant metabolisms. $\delta^2\text{H}_{\text{CH}_4}$ isotopic signatures from pyrogenic sources cover a wide range of values, and overlap with the ones of fossil fuels. Data on $\delta^2\text{H}_{\text{H}_2\text{O}}$ could help to parametrise the biomass burning $\delta^2\text{H}_{\text{CH}_4}$ in more detail (Vigano et al., 2010), similar to the above mentioned relation between $\delta^2\text{H}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{H}_2\text{O}}$ (Waldron et al., 1999; Chanton et al., 2006; Röckmann et al., 2010; Douglas et al., 2021).

Table 3. CH₄ isotopic source signatures assigned to the fossil fuel related emissions in global scale models.

Reference	$\delta^{13}\text{C}$ VPDB [‰]	$\delta^2\text{H}$ VSMOW [‰]
Gupta et al. (1996); Tyler et al. (2007)	-38 / -37 ¹	-175
Rigby et al. (2012)	-40 ²	-175
Monteil et al. (2011)	-40 / -35 ¹	
Rice et al. (2016)	-41.7	-175
Schaefer et al. (2016)	-37	
Schwietzke et al. (2016)	-44	
Fujita et al. (2020)	-45.2	-209
This database, mean \pm semi	-44.5 \pm 0.5 / -50.7 \pm 1.3 ¹	-183 \pm 3 / -210 \pm 5 ¹

¹ for natural gas/coal; ² also in Lassey et al. (2000); Houweling et al. (2000); Bousquet et al. (2006); Thompson et al. (2018)

Fugitive CH₄ emissions from fossil fuels cover a wide range of isotopic signatures: $\delta^{13}\text{C}$ from -72.5 to -18.3 ‰ and $\delta^2\text{H}$ from -349 to 14.0 ‰. The average $\delta^{13}\text{C}$ of all fugitive CH₄ emissions from the exploitation of fossil fuels in the European Methane Isotope Database was -44.6 ± 0.4 ‰ (n=452), and the weighted average was -46.6 ± 1.8 ‰ according to the relative emission from conventional and coal fuels production worldwide⁷. Our averages are lower than $\delta^{13}\text{C}$ values used in global models, and than the mean of the global database (Table 3). But the value of -44 ± 0.7 ‰ suggested by Schwietzke et al. (2016), based on the database from Sherwood et al. (2017) scaled with the fossil fuel production in the different regions, is relatively close to our average. The mean values we calculated in Table 3 (bottom row) do not necessarily represent the global isotopic signature of fossil fuel emissions, because this should be weighted by the magnitude of emissions in the different basins. However, our averages are an indication of the general $\delta^{13}\text{C}$ signatures from all measurements until now. Because of the high heterogeneity of the $\delta^{13}\text{C}$ of CH₄ from fossil fuel related activities, and the temporal variations in the production from the different regions (Stavert et al., 2021; US Energy Information Administration, 2021; Lan et al., 2021), it is important to keep a relatively large uncertainty when estimating in the global signature of fossil fuel emissions.

In section 2.4, we have shown the use of $\delta^{13}\text{C}_{\text{CH}_4}$ to distinguish fossil fuel emissions in western Europe, and the need for $\delta^2\text{H}_{\text{CH}_4}$ measurements in central and eastern Europe. In the global database, most fossil fuels records (83.5%) have $\delta^2\text{H}_{\text{CH}_4}$ values > -250 ‰. The few values of $\delta^2\text{H} < -300$ ‰, indicating microbial fermentation as gas origin, were found in some coal formations in the United States and Canada. Figure 5 still allows us to generally conclude that $\delta^2\text{H}$ measurements are more reliable to distinguish fossil fuel vs. biogenic CH₄ sources at the global scale than $\delta^{13}\text{C}$ only, which further emphasises the need for more $\delta^2\text{H}_{\text{CH}_4}$ measurements.

⁷Relative weights of 0.66 for conventional fuels (oil and natural gas) and 0.34 for coal. Emission data from Saunio et al. (2020)




Figure 5. Distribution of $\delta^{13}\text{C}$ (top) and $\delta^2\text{H}$ (bottom) in CH_4 for different source categories. **(a)** The European Methane Isotope Database (absolute numbers). "fossil fuels -E" shows fossil fuels data from Poland and Romania, and "fossil fuels -W" from the UK, the Netherlands, Germany, and France. **(b)** the global literature database, including the new European data (normalised probability density).

The updated database is beneficial for deriving a representative concept of the isotopic composition of CH_4 sources, but it is important to note that applying appropriate weighting arithmetic is essential. Users need to define the dominant CH_4 sources impacting an area, as well as the relative source type emission rates. Emission inventories provide such estimates, but top-down approaches are essential to identify potential biases and evaluate the bottom-up approaches (Alvarez et al., 2018; Etiopie and Schwietzke, 2019; Rutherford et al., 2021; Stavert et al., 2021).

3 Conclusions

This study presents an updated dataset of isotopic source signatures of CH₄ from recent atmospheric measurements, while including additional data from published literature which were not previously included. The new data is a contribution from the European Methane Isotope Database, that results from the sampling activities performed within the MEMO² project. It represents a substantial contribution to the global dataset for fugitive fossil fuels and waste sources, mainly sampled in urban areas.

We have highlighted two main improvements in our understanding of the CH₄ isotopic composition: (i) A more robust range of values for modern microbial sources, and a better characterisation of the $\delta^{13}\text{C}$ enrichment in CH₄ from waste sources. (ii) Fossil fuel related sources could have more depleted values than previous estimates used in global models. In this respect, our data confirm the analysis made by Schwietzke et al. (2016).

Finally, the new European data contain comparatively more $\delta^2\text{H}$ measurements. In the case of fossil fuel emissions, the use of $\delta^2\text{H}_{\text{CH}_4}$ is of particular interest. In general, utilizing both $\delta^{13}\text{C}$ and $\delta^2\text{H}$ for CH₄ improves our ability to clearly separate fossil fuel and microbial sources, compared to $\delta^{13}\text{C}$ alone. The use of $\delta^2\text{H}$ as additional constraint could help to answer open questions regarding the CH₄ global budget. To better understand the drivers of $\delta^2\text{H}$ variability (except for $\delta^2\text{H}$ of precipitation), more measurements are required, especially of pyrogenic and waste sources.

The present dataset can be used for CH₄ source attribution, and also to derive global signatures for the different types of emissions. Appropriate use of the database requires the analysis of specific parameters in relation to source type and the region of interest. A future improvement of this database would be to include more measurements on the African, Asian and South American continents, where experimental studies are lacking. Because of its potential for source characterisation, new studies should also focus on $\delta^2\text{H}_{\text{CH}_4}$ measurements.

4 Data availability

The database is made freely available to the scientific community in the belief that it provides the most complete picture of the stable isotopic composition of CH₄ sources. The free availability of these data does not constitute permission for publication of the data. For research projects, if the data used are essential to the work to be published, or if the conclusion or results largely depend on the data, co-authorship should be considered. Full contact details and information on how to cite the data are given in the accompanying database. The database is currently stored in a publically available repository: <https://doi.org/10.24416/UU01-4PO56T> (Menoud et al., 2021a).



Figure A1. Photos of the laboratory setup used for isotopic measurements during the MEMO² project and of the sample collection during field surveys. (a) IRMS system setup used at GGRL RHUL, for $\delta^{13}\text{C}_{\text{CH}_4}$ measurements. (b) IRMS system setup used by the UU team, here deployed at Krakow, for $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$ measurements. (c) Mobile measurement setup used by the Heidelberg University (UHEL) team, using cavity ring-down spectrometry (CRDS) isotope analyser and an air-core, as described in Hoheisel et al. (2019). *Photo: Piotr Korbeń* (d) Sample collection directly from an emission source on foot: a manhole in the city of Bucharest, Romania. CH_4 mole fractions were measured using a portable CRDS analyser. *Photo: Octavian* (e) Mobile measurements and sampling on-board of a car. Air was pumped from the top of the car and continuously measured with a CRDS analyser, and bag samples were taken when CH_4 enhancement were detected by the instrument. (f) Soil chamber placed on a coal waste disposal area in Poland. Samples were collected from the chamber several hours after it was placed. CH_4 mole fractions in the chamber were measured using a portable integrated cavity output spectroscopy (ICOS) instrument. *Photo: Mila Stanisavljević* (g) Full sample bag before analysis, collected on a landfill in Devon, UK.




Figure A2. Number of isotopic signature measurements reported in the present database, including the previous database of Sherwood et al. (2017, 2021), additional literature we found, and the European Methane Isotope Database, for $\delta^{13}\text{C}$ (left) and $\delta^2\text{H}$ (right) of CH_4 separately.




Figure A3. Measured $\delta^{13}\text{C}_{\text{CH}_4}$ signatures from ruminants in the literature¹ and MEMO² database according to the feed: a majority of C3 plants (red) or C4 plants (blue).

¹ Al-Shalan et al. (2022); Brownlow et al. (2017); Klevenhusen et al. (2009, 2010); Levin et al. (1993); Lu et al. (2021); Rust (1981); Townsend-Small et al. (2012); Wahlen et al. (1989)






Figure A4. Measured $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^2\text{H}_{\text{CH}_4}$ signatures from wetlands sites as reported in the literature² (solid circled) and MEMO₂ (open circles) database, color coded by the latitude zones. 

² Beck et al. (2012); Burke and Sackett (1986); Day et al. (2015); Happell et al. (1994); Kuhlmann et al. (1998); Lansdown (1992); Levin et al. (1993); Martens et al. (1992); Nakagawa et al. (2002); Smith et al. (2000); Sugimoto and Fujita (2006); Umezawa et al. (2011); Wahlen et al. (1989); Wassmann et al. (1992); Woltemate et al. (1984)

 *Author contributions.* MM, CV, DL, JF, SB, JF and RF performed the isotopic measurements. MM, TR, DL, JF, SR, JF, RF, HM, MS, JN, KV, PL, PK, MS, SD and TA took part in the collection of samples. MM gathered and analysed the data and prepared the figures; TR and DL contributed to the interpretation of the data. MM prepared the manuscript with contributions from TR, DL, JR, MS, PL, SB, HM, JF and HC.

Competing interests. The authors declare that they have no conflict of interest.

335 *Acknowledgements.* We thank all the staff from different organisations involved in the MEMO², CoMet and ROMEO projects who participated in the sample collection. We acknowledge the technical staff at UU and RHUL for the maintenance of the IRMS systems. This work was supported by ITN project “Methane goes Mobile – Measurements and Modelling” (MEMO2; <https://h2020-memo2.eu/>, last access: November 3rd, 2021). This project has received funding from the European Union’s Horizon 2020 Research and Innovation programme under the Marie Skłodowska-Curie grant agreement no. 722479.

- Akritas, M. G. and Bershad, M. A.: Linear Regression for Astronomical Data with Measurement Errors and Intrinsic Scatter, *The Astrophysical Journal*, 470, 706, <https://doi.org/10.1086/177901>, 1996.
- Al-Shalan, A., Lowry, D., Fisher, R., Nisbet, E., Zazzeri, G., Al-Sarawi, M., and France, J.: Methane Emissions in Kuwait: Plume Identification, Isotopic Characterisation and Inventory Verification, *Atmospheric Environment*, p. 118763, <https://doi.org/10.1016/j.atmosenv.2021.118763>, 2022.
- Alvarez, R. A., Zavala-Araiza, D., Lyon, D. R., Allen, D. T., Barkley, Z. R., Brandt, A. R., Davis, K. J., Herndon, S. C., Jacob, D. J., Karion, A., Kort, E. A., Lamb, B. K., Lauvaux, T., Maasackers, J. D., Marchese, A. J., Omara, M., Pacala, S. W., Peischl, J., Robinson, A. L., Shepson, P. B., Sweeney, C., Townsend-Small, A., Wofsy, S. C., and Hamburg, S. P.: Assessment of Methane Emissions from the U.S. Oil and Gas Supply Chain, *Science*, p. eaar7204, <https://doi.org/10.1126/science.aar7204>, 2018.
- Andersen, T., Scheeren, B., Peters, W., and Chen, H.: A UAV-based Active AirCore System for Measurements of Greenhouse Gases, *Atmospheric Measurement Techniques*, 11, 2683–2699, <https://doi.org/10.5194/amt-11-2683-2018>, 2018.
- Baciu, C., Ionescu, A., and Etiope, G.: Hydrocarbon Seeps in Romania: Gas Origin and Release to the Atmosphere, *Marine and Petroleum Geology*, 89, 130–143, <https://doi.org/10.1016/j.marpetgeo.2017.06.015>, 2018.
- Bakkaloglu, S., Lowry, D., Fisher, R. E., France, J. L., and Nisbet, E. G.: Carbon Isotopic Characterisation and Oxidation of UK Landfill Methane Emissions by Atmospheric Measurements, *Waste Management*, 132, 162–175, <https://doi.org/10.1016/j.wasman.2021.07.012>, 2021.
- Bakkaloglu, S., Lowry, D., Fisher, R. E., Menoud, M., Lanoisellé, M., Chen, H., Röckmann, T., and Nisbet, E. G.: Stable Isotopic Signatures of Methane from Waste Sources through Atmospheric Measurements, *Atmospheric Environment*, *in review*.
- Beck, V., Chen, H., Gerbig, C., Bergamaschi, P., Bruhwiler, L., Houweling, S., Röckmann, T., Kolle, O., Steinbach, J., Koch, T., Sapart, C. J., van der Veen, C., Frankenberg, C., Andreae, M. O., Artaxo, P., Longo, K. M., and Wofsy, S. C.: Methane Airborne Measurements and Comparison to Global Models during BARCA, *Journal of Geophysical Research: Atmospheres*, 117, [15310], <https://doi.org/10.1029/2011JD017345>, 2012.
- Bergamaschi, P., Lubina, C., Königstedt, R., Fischer, H., Veltkamp, A. C., and Zwaagstra, O.: Stable Isotopic Signatures ($\delta^{13}\text{C}$, δD) of Methane from European Landfill Sites, *Journal of Geophysical Research: Atmospheres*, 103, 8251–8265, <https://doi.org/10.1029/98JD00105>, 1998.
- Boggs, P. T. and Rogers, J. E.: Orthogonal Distance Regression, in: *Statistical Analysis of Measurement Error Models and Applications: Proceedings of the AMS-IMS-SIAM Joint Summer Research Conference Held June 10-16, 1989*, vol. 112 of *Contemporary Mathematics*, p. 186, American Mathematical Society, 1990.
- Bousquet, P., Ciais, P., Miller, J. B., Dlugokencky, E. J., Hauglustaine, D. A., Prigent, C., Van der Werf, G. R., Peylin, P., Brunke, E.-G., Carouge, C., Langenfelds, R. L., Lathière, J., Papa, F., Ramonet, M., Schmidt, M., Steele, L. P., Tyler, S. C., and White, J.: Contribution of Anthropogenic and Natural Sources to Atmospheric Methane Variability, *Nature*, 443, 439–443, <https://doi.org/10.1038/nature05132>, 2006.
- Bréas, O., Guillou, C., Reniero, F., and Wada, E.: The Global Methane Cycle: Isotopes and Mixing Ratios, Sources and Sinks, *Isotopes in Environmental and Health Studies*, 37, 257–379, <https://doi.org/10.1080/10256010108033302>, 2001.

- 375 Brownlow, R., Lowry, D., Fisher, R. E., France, J. L., Lanoisellé, M., White, B., Wooster, M. J., Zhang, T., and Nisbet, E. G.: Isotopic Ratios of Tropical Methane Emissions by Atmospheric Measurement: Tropical Methane $\delta^{13}\text{C}$ Source Signatures, *Global Biogeochemical Cycles*, 31, 1408–1419, <https://doi.org/10.1002/2017GB005689>, 2017.
- Burke, R. A. and Sackett, W. M.: Stable Hydrogen and Carbon Isotopic Compositions of Biogenic Methanes from Several Shallow Aquatic Environments, in: *Organic Marine Geochemistry*, edited by Sohn, M. L., vol. 305, pp. 297–313, American Chemical Society, Washington, DC, <https://doi.org/10.1021/bk-1986-0305.ch017>, 1986.
- 380 Chan, E. W., Shiller, A. M., Joung, D. J., Arrington, E. C., Valentine, D. L., Redmond, M. C., Breier, J. A., Socolofsky, S. A., and Kessler, J. D.: Investigations of Aerobic Methane Oxidation in Two Marine Seep Environments: Part 2—Isotopic Kinetics, *Journal of Geophysical Research: Oceans*, 124, 8392–8399, <https://doi.org/10.1029/2019JC015603>, 2019.
- Chan, J. P., Fields, D., and Hines, M. E.: Controls on the Hydrogen Isotopic Composition of Biogenic Methane from High-Latitude Terrestrial Wetlands, *Journal of Geophysical Research: Biogeosciences*, 111, <https://doi.org/10.1029/2005JG000134>, 2006.
- 385 Chen, Y., Sherwin, E., Berman, E., Jones, B., Gordon, M., Wetherley, E., Kort, E., and Brandt, A.: Comprehensive Aerial Survey Quantifies High Methane Emissions from the New Mexico Permian Basin, Preprint, *Atmospheric Sciences*, <https://doi.org/10.31223/X56D0D>, 2021.
- Cicerone, R. J. and Oremland, R. S.: Biogeochemical Aspects of Atmospheric Methane, *Global Biogeochemical Cycles*, 2, 299–327, 1988.
- Coleman, D. D., Risatti, J., and Schoell, M.: Fractionation of Carbon and Hydrogen Isotopes by Methane-Oxidizing Bacteria, *Geochimica et Cosmochimica Acta*, 45, 1033–1037, [https://doi.org/10.1016/0016-7037\(81\)90129-0](https://doi.org/10.1016/0016-7037(81)90129-0), 1981.
- 390 Conrad, R.: Quantification of Methanogenic Pathways Using Stable Carbon Isotopic Signatures: A Review and a Proposal, *Organic Geochemistry*, 36, 739–752, <https://doi.org/10.1016/j.orggeochem.2004.09.006>, 2005.
- Conrad, R., Noll, M., Claus, P., Klose, M., Bastos, W. R., and Enrich-Prast, A.: Stable Carbon Isotope Discrimination and Microbiology of Methane Formation in Tropical Anoxic Lake Sediments, *Biogeosciences*, 8, 795–814, <https://doi.org/10.5194/bg-8-795-2011>, 2011.
- 395 Day, S., Ong, C., Rodger, A., Etheridge, D., Hibberd, M., van Gorsel, E., Spencer, D., Krummel, P., Fry, R., Dell'Amico, M., Sestak, S., Williams, D., Loh, Z., and Barrett, D.: Characterisation of Regional Fluxes of Methane in the Surat Basin, Queensland: Phase 2: A Pilot Study of Methodology to Detect and Quantify Methane Sources, Technical report, CSIRO, Australia, 2015.
- De Visscher, A.: Isotope Fractionation Effects by Diffusion and Methane Oxidation in Landfill Cover Soils, *Journal of Geophysical Research*, 109, D18 111, <https://doi.org/10.1029/2004JD004857>, 2004.
- 400 Defratyka, S.: Characterizing Methane (CH₄) Emissions in Urban Environments (Paris), Ph.D. thesis, Université Paris-Saclay, 2021.
- Defratyka, S. M., Paris, J.-D., Yver-Kwok, C., Fernandez, J. M., Korben, P., and Bousquet, P.: Mapping Urban Methane Sources in Paris, France, *Environmental Science & Technology*, 55, 8583–8591, <https://doi.org/10.1021/acs.est.1c00859>, 2021.
- Douglas, P. M. J., Stratigopoulos, E., Park, S., and Phan, D.: Geographic Variability in Freshwater Methane Hydrogen Isotope Ratios and Its Implications for Global Isotopic Source Signatures, *Biogeosciences*, 18, 3505–3527, <https://doi.org/10.5194/bg-18-3505-2021>, 2021.
- 405 Etiope, G. and Schwietzke, S.: Global Geological Methane Emissions: An Update of Top-down and Bottom-up Estimates, *Elementa: Science of the Anthropocene*, 7, 47, <https://doi.org/10.1525/elementa.383>, 2019.
- Etminan, M., Myhre, G., Highwood, E. J., and Shine, K. P.: Radiative Forcing of Carbon Dioxide, Methane, and Nitrous Oxide: A Significant Revision of the Methane Radiative Forcing: Greenhouse Gas Radiative Forcing, *Geophysical Research Letters*, 43, 12,614–12,623, <https://doi.org/10.1002/2016GL071930>, 2016.
- 410 Fernandez, J. M., Maazallahi, H., France, J. L., Menoud, M., Corbu, M., Ardelean, M., Calcan, A., Townsend-Small, A., van der Veen, C., Fisher, R. E., Lowry, D., Nisbet, E. G., and Röckmann, T.: Street-Level Methane Emissions of Bucharest, Romania and the Influence of Urban Wastewater, *Atmospheric Environment X*, in review.

- Fiehn, A., Kostinek, J., Eckl, M., Klausner, T., Gałkowski, M., Chen, J., Gerbig, C., Röckmann, T., Maazallahi, H., Schmidt, M., Korbeń, P., Nęcki, J., Jagoda, P., Wildmann, N., Mallaun, C., Bun, R., Nickl, A.-L., Jöckel, P., Fix, A., and Roiger, A.: Estimating CH₄, CO₂, and CO Emissions from Coal Mining and Industrial Activities in the Upper Silesian Coal Basin Using an Aircraft-Based Mass Balance Approach, *Atmospheric Chemistry and Physics Discussions*, <https://doi.org/10.5194/acp-2020-282>, 2020.
- 415 Fisher, R., Lowry, D., Wilkin, O., Sriskantharajah, S., and Nisbet, E. G.: High-Precision, Automated Stable Isotope Analysis of Atmospheric Methane and Carbon Dioxide Using Continuous-Flow Isotope-Ratio Mass Spectrometry, *Rapid Communications in Mass Spectrometry*, 20, 200–208, <https://doi.org/10.1002/rcm.2300>, 2006.
- 420 Fisher, R. E., Sriskantharajah, S., Lowry, D., Lanoisellé, M., Fowler, C. M. R., James, R. H., Hermansen, O., Lund Myhre, C., Stohl, A., Greinert, J., Nisbet-Jones, P. B. R., Mienert, J., and Nisbet, E. G.: Arctic Methane Sources: Isotopic Evidence for Atmospheric Inputs, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL049319>, 2011.
- Fisher, R. E., France, J. L., Lowry, D., Lanoisellé, M., Brownlow, R., Pyle, J. A., Cain, M., Warwick, N., Skiba, U. M., Drewer, J., Dinsmore, K. J., Leeson, S. R., Bauguitte, S. J.-B., Wellpott, A., O’Shea, S. J., Allen, G., Gallagher, M. W., Pitt, J., Percival, C. J., Bower, K., George, C., Hayman, G. D., Aalto, T., Lohila, A., Aurela, M., Laurila, T., Crill, P. M., McCalley, C. K., and Nisbet, E. G.: Measurement of the ¹³C Isotopic Signature of Methane Emissions from Northern European Wetlands: Northern Wetland CH₄ Isotopic Signature, *Global Biogeochemical Cycles*, 31, 605–623, <https://doi.org/10.1002/2016GB005504>, 2017.
- 425 France, J., Fisher, R. E., Lowry, D., Lamoiselle, M., Nisbet-Jones, P., Andrade, M., Moreno, I., Forster, G., Oram, D., Helfter, C., Skiba, U., Stephens, M., Broderick, T., Hoyt, A., Gondwe, M., Jones, A. E., and Nisbet, E. G.: $\delta^{13}\text{C}$ Methane Source Signatures from Tropical Wetlands and Rice Field Emissions, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, Submitted for publication, 2021.
- 430 Fujita, R., Morimoto, S., Maksyutov, S., Kim, H.-S., Arshinov, M., Brailsford, G., Aoki, S., and Nakazawa, T.: Global and Regional CH₄ Emissions for 1995–2013 Derived From Atmospheric CH₄, $\delta^{13}\text{C}\text{-CH}_4$, and $\delta\text{D}\text{-CH}_4$ Observations and a Chemical Transport Model, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2020JD032903>, 2020.
- 435 Gałkowski, M., Fiehn, A., Swolkien, J., Stanisavljevic, M., Korben, P., Menoud, M., Necki, J., Roiger, A., Röckmann, T., Gerbig, C., and Fix, A.: Emissions of CH₄ and CO₂ over the Upper Silesian Coal Basin (Poland) and Its Vicinity, <https://doi.org/10.18160/3K6Z-4H73>, 2021.
-  Ganesan, A. L., Stell, A. C., Gedney, N., Comyn-Platt, E., Hayman, G., Rigby, M., Poulter, B., and Hornibrook, E. R. C.: Spatially Resolved Isotopic Source Signatures of Wetland Methane Emissions, *Geophysical Research Letters*, 45, 3737–3745, <https://doi.org/10.1002/2018GL077536>, 2018.
- 440 Gebert, J. and Streese-Kleeberg, J.: Coupling Stable Isotope Analysis with Gas Push-Pull Tests to Derive in Situ Values for the Fractionation Factor A_{ox} Associated with the Microbial Oxidation of Methane in Soils, *Soil Science Society of America Journal*, 81, 1107–1114, <https://doi.org/10.2136/sssaj2016.11.0387>, 2017.
-   Gupta, M., Tyler, S., and Cicerone, R.: Modeling Atmospheric $\delta^{13}\text{C}\text{CH}_4$ and the Causes of Recent Changes in Atmospheric CH₄ Amounts, *Journal of Geophysical Research: Atmospheres*, 101, 22 923–22 932, <https://doi.org/10.1029/96JD02386>, 1996.
- 445 Happell, J. D., Chanton, J. P., and Showers, W. S.: The Influence of Methane Oxidation on the Stable Isotopic Composition of Methane Emitted from Florida Swamp Forests, *Geochimica et Cosmochimica Acta*, 58, 4377–4388, [https://doi.org/10.1016/0016-7037\(94\)90341-7](https://doi.org/10.1016/0016-7037(94)90341-7), 1994.

- Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M., and Emmenegger, L.: Validation of the Swiss Methane Emission Inventory by Atmospheric Observations and Inverse Modelling, *Atmospheric Chemistry and Physics*, 16, 3683–3710, <https://doi.org/10.5194/acp-16-3683-2016>, 2016.
- Hoheisel, A., Yeman, C., Dinger, F., Eckhardt, H., and Schmidt, M.: An Improved Method for Mobile Characterisation of $\delta^{13}\text{C}\text{H}_4$ Source Signatures and Its Application in Germany, *Atmospheric Measurement Techniques*, 12, 1123–1139, <https://doi.org/10.5194/amt-12-1123-2019>, 2019.
- Houweling, S., Dentener, F., and Lelieveld, J.: Simulation of Preindustrial Atmospheric Methane to Constrain the Global Source Strength of Natural Wetlands, *Journal of Geophysical Research: Atmospheres*, 105, 17 243–17 255, <https://doi.org/10.1029/2000JD900193>, 2000.
- Inglett, K. S., Chanton, J. P., and Inglett, P. W.: Methanogenesis and Methane Oxidation in Wetland Soils, in: *SSSA Book Series*, edited by DeLaune, R., Reddy, K., Richardson, C., and Megonigal, J., pp. 407–425, American Society of Agronomy and Soil Science Society of America, Madison, WI, USA, <https://doi.org/10.2136/sssabookser10.c21>, 2015.
- IPCC: *Climate Change 2013: The Physical Science Basis ; Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge Univ. Press, Cambridge, UK and New York, NY, USA, 2013.
- IPCC 2021: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press., 2021a.
- IPCC 2021: *Summary for Policymakers*, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yeleğçi, O., Yu, R., and Zhou, B., p. In Press., Cambridge University Press, 2021b.
- Keeling, C. D.: The Concentration and Isotopic Abundances of Carbon Dioxide in Rural and Marine Air, *Geochimica et Cosmochimica Acta*, 24, 277–298, 1961.
- Klevenhusen, F., Bernasconi, S. M., Hofstetter, T. B., Bolotin, J., Kunz, C., and Soliva, C. R.: Efficiency of Monolaurin in Mitigating Ruminant Methanogenesis and Modifying C-isotope Fractionation When Incubating Diets Composed of Either C_3 or C_4 Plants in a Rumen Simulation Technique (Rusitec) System, *British Journal of Nutrition*, 102, 1308–1317, <https://doi.org/10.1017/S0007114509990262>, 2009.
- Klevenhusen, F., Bernasconi, S. M., Kreuzer, M., and Soliva, C. R.: Experimental Validation of the Intergovernmental Panel on Climate Change Default Values for Ruminant-Derived Methane and Its Carbon-Isotope Signature, *Animal Production Science*, 50, 159, <https://doi.org/10.1071/AN09112>, 2010.
- Kotarba, M. J.: Composition and Origin of Coalbed Gases in the Upper Silesian and Lublin Basins, Poland, *Organic Geochemistry*, 32, 163–180, [https://doi.org/10.1016/S0146-6380\(00\)00134-0](https://doi.org/10.1016/S0146-6380(00)00134-0), 2001.
- Kotarba, M. J. and Pluta, I.: Origin of Natural Waters and Gases within the Upper Carboniferous Coal-Bearing and Autochthonous Miocene Strata in South-Western Part of the Upper Silesian Coal Basin, Poland, *Applied Geochemistry*, 24, 876–889, <https://doi.org/10.1016/j.apgeochem.2009.01.013>, 2009.
- Kuhlmann, A. J., Worthy, D. E. J., Trivett, N. B. A., and Levin, I.: Methane Emissions from a Wetland Region within the Hudson Bay Lowland: An Atmospheric Approach, *Journal of Geophysical Research*, 103, 16 009–16 016, <https://doi.org/10.1029/98JD01024509.00>, 1998.
- Lan, X., Basu, S., Schwietzke, S., Bruhwiler, L. M. P., Dlugokencky, E. J., Michel, S. E., Sherwood, O. A., Tans, P. P., Thoning, K., Etiope, G., Zhuang, Q., Liu, L., Oh, Y., Miller, J. B., Pétron, G., Vaughn, B. H., and Crippa, M.: Improved Constraints on Global Methane Emissions and Sinks Using $\delta^{13}\text{C}\text{-CH}_4$, *Global Biogeochemical Cycles*, 35, <https://doi.org/10.1029/2021GB007000>, 2021.





- Lansdown, J. M.: The Carbon and Hydrogen Stable Isotope Composition of Methane Released from Natural Wetlands and Ruminants, Abstract, University of Washington, 1992.
- 490 Lassey, K. R., Lowe, D. C., and Manning, M. R.: The Trend in Atmospheric Methane $\delta^{13}\text{C}$ and Implications for Isotopic Constraints on the Global Methane Budget, *Global Biogeochemical Cycles*, 14, 41–49, <https://doi.org/10.1029/1999GB900094>, 2000.
- Levin, I., Bergamaschi, P., Dörr, H., and Trapp, D.: Stable Isotopic Signature of Methane from Major Sources in Germany, *Chemosphere*, 26, 161–177, [https://doi.org/10.1016/0045-6535\(93\)90419-6](https://doi.org/10.1016/0045-6535(93)90419-6), 1993.
- 495 Lowry, D., Holmes, C. W., Rata, N. D., O'Brien, P., and Nisbet, E. G.: London Methane Emissions: Use of Diurnal Changes in Concentration and $\delta^{13}\text{C}$ to Identify Urban Sources and Verify Inventories, *Journal of Geophysical Research: Atmospheres*, 106, 7427–7448, <https://doi.org/10.1029/2000JD900601>, 2001.
- Lowry, D., Fisher, R. E., France, J. L., Coleman, M., Lanoisellé, M., Zazzeri, G., Nisbet, E. G., Shaw, J. T., Allen, G., Pitt, J., and Ward, R. S.: Environmental Baseline Monitoring for Shale Gas Development in the UK: Identification and Geochemical Characterisation of Local Source Emissions of Methane to Atmosphere, *Science of The Total Environment*, 708, 134600, <https://doi.org/10.1016/j.scitotenv.2019.134600>, 2020.
- 500 Lu, X., Harris, S. J., Fisher, R. E., France, J. L., Nisbet, E. G., Lowry, D., Röckmann, T., van der Veen, C., Menoud, M., Schwietzke, S., and Kelly, B. F. J.: Isotopic Signatures of Major Methane Sources in the Coal Seam Gas Fields and Adjacent Agricultural Districts, Queensland, Australia, *Atmospheric Chemistry and Physics*, 21, 1–36, <https://doi.org/10.5194/acp-2021-76>, 2021.
- 505 Maasackers, J. D., Jacob, D. J., Sulprizio, M. P., Scarpelli, T. R., Nesser, H., Sheng, J.-X., Zhang, Y., Hersher, M., Bloom, A. A., Bowman, K. W., Worden, J. R., Janssens-Maenhout, G., and Parker, R. J.: Global Distribution of Methane Emissions, Emission Trends, and OH Concentrations and Trends Inferred from an Inversion of GOSAT Satellite Data for 2010–2015, *Atmospheric Chemistry and Physics*, 19, 7859–7881, <https://doi.org/10.5194/acp-19-7859-2019>, 2019.
- Maazallahi, H., Fernandez, J. M., Menoud, M., Zavala-Araiza, D., Weller, Z. D., Schwietzke, S., von Fischer, J. C., Denier van der Gon, H., and Röckmann, T.: Methane Mapping, Emission Quantification, and Attribution in Two European Cities: Utrecht (NL) and Hamburg (DE), *Atmospheric Chemistry and Physics*, 20, 14717–14740, <https://doi.org/10.5194/acp-20-14717-2020>, 2020.
- 510 Maazallahi, H., Röckmann, T., and Schwietzke, S.: Estimation of CH₄ Emissions from the Romanian Sector Using Field Measurementnets, In preparation, 2021.
- Martens, C. S., Kelley, C. A., Chanton, J. P., and Showers, W. J.: Carbon and Hydrogen Isotopic Characterization of Methane from Wetlands and Lakes of the Yukon-Kuskokwim Delta, Western Alaska, *Journal of Geophysical Research*, 97, 16689, <https://doi.org/10.1029/91JD02885>, 1992.
- 515 Mayfield, E. N., Robinson, A. L., and Cohon, J. L.: System-Wide and Superemitter Policy Options for the Abatement of Methane Emissions from the U.S. Natural Gas System, *Environmental Science & Technology*, 51, 4772–4780, <https://doi.org/10.1021/acs.est.6b05052>, 2017.
- McCalley, C. K., Woodcroft, B. J., Hodgkins, S. B., Wehr, R. A., Kim, E.-H., Mondav, R., Crill, P. M., Chanton, J. P., Rich, V. I., Tyson, G. W., and Saleska, S. R.: Methane Dynamics Regulated by Microbial Community Response to Permafrost Thaw, *Nature*, 514, 478–481, <https://doi.org/10.1038/nature13798>, 2014.
- 520 Menoud, M., van der Veen, C., Scheeren, B., Chen, H., Szénási, B., Morales, R. P., Pison, I., Bousquet, P., Brunner, D., and Röckmann, T.: Characterisation of Methane Sources in Lutjewad, The Netherlands, Using Quasi-Continuous Isotopic Composition Measurements, *Tellus B: Chemical and Physical Meteorology*, 72, 1–19, <https://doi.org/10.1080/16000889.2020.1823733>, 2020.
- Menoud, M., van der Veen, C., Lowry, D., Fernandez, J. M., Bakkaloglu, S., France, J. L., Fisher, R. E., Maazallahi, H., Stanisavljević, M., Nećki, J., Vinkovic, K., Łakomic, P., Rinne, J., Korbeň, P., Schmidt, M., Defratyka, S., Yver-Kwok, C., Andersen, T., Chen, H., and

- 525 Röckmann, T.: European Methane Isotope Database Coupled with a Global Inventory of Fossil and Non-Fossil $\delta^{13}\text{C}$ - and $\delta^2\text{H}$ - CH_4 Source Signature Measurements: V1.0.0, [dataset], Utrecht University, 2021a.
- Menoud, M., van der Veen, C., Necki, J., Bartyzel, J., Szénási, B., Stanisavljević, M., Pison, I., Bousquet, P., and Röckmann, T.: Methane (CH_4) Sources in Krakow, Poland: Insights from Isotope Analysis, *Atmospheric Chemistry and Physics*, 21, 13 167–13 185, 2021b.
- Menoud, M., van der Veen, C., Maazallahi, H., Hensen, A., Velzeboer, I., van den Bulk, P., Delre, A., Korbeň, P., Schwietzke, S., Ardelean, 530 M., Calcan, A., Etiope, G., Baciu, C., Scheutz, C., Schmidt, M., and Röckmann, T.: CH_4 Sotopic Signatures of Emissions from Oil and Gas Extraction Sites in Romania, *Elementa: Science of the Anthropocene*, in review.
- Milkov, A. V. and Etiope, G.: Revised Genetic Diagrams for Natural Gases Based on a Global Dataset of >20,000 Samples, *Organic Geochemistry*, 125, 109–120, <https://doi.org/10.1016/j.orggeochem.2018.09.002>, 2018.
- 535 Miller, J. B. and Tans, P. P.: Calculating Isotopic Fractionation from Atmospheric Measurements at Various Scales, *Tellus B: Chemical and Physical Meteorology*, 55, 207–214, <https://doi.org/10.3402/tellusb.v55i2.16697>, 2003.
- Monteil, G., Houweling, S., Dlugokenky, E. J., Maenhout, G., Vaughn, B. H., White, J. W. C., and Rockmann, T.: Interpreting Methane Variations in the Past Two Decades Using Measurements of CH_4 Mixing Ratio and Isotopic Composition, *Atmospheric Chemistry and Physics*, 11, 9141–9153, <https://doi.org/10.5194/acp-11-9141-2011>, 2011.
- Nakagawa, F., Yoshida, N., Sugimoto, A., Wada, E., Yoshioka, T., Ueda, S., and Vijarnsorn, P.: Stable Isotope and Radiocarbon 540 Compositions of Methane Emitted from Tropical Rice Paddies and Swamps in Southern Thailand, *Biogeochemistry*, 61, 1–19, <https://doi.org/10.1023/A:1020270032512>, 2002.
- Nisbet, E. G., Manning, M. R., Dlugokenky, E. J., Fisher, R. E., Lowry, D., Michel, S. E., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., Brownlow, R., Cain, M., France, J. L., Hermansen, O., Hossaini, R., Jones, A. E., Levin, I., Manning, A. C., Myhre, G., Pyle, J., 545 Vaughn, B. H., Warwick, N. J., and White, J. W. C.: Very Strong Atmospheric Methane Growth in the 4 Years 2014–2017: Implications for the Paris Agreement, *Global Biogeochemical Cycles*, 33, 318–342, <https://doi.org/10.1029/2018GB006009>, 2019.
- Nisbet, E. G., Fisher, R. E., Lowry, D., France, J. L., Allen, G., Bakkaloglu, S., Broderick, T. J., Cain, M., Coleman, M., Fernandez, J., Forster, G., Griffiths, P. T., Iverach, C. P., Kelly, B. F. J., Manning, M. R., Nisbet-Jones, P. B. R., Pyle, J. A., Townsend-Small, A., al-Shalaan, A., Warwick, N., and Zazzeri, G.: Methane Mitigation: Methods to Reduce Emissions, on the Path to the Paris Agreement, *Reviews of Geophysics*, 58, <https://doi.org/10.1029/2019RG000675>, 2020.
- 550 Obersky, L., Rafiee, R., Cabral, A. R., Golding, S. D., and Clarke, W. P.: Methodology to Determine the Extent of Anaerobic Digestion, Composting and CH_4 Oxidation in a Landfill Environment, *Waste Management*, 76, 364–373, <https://doi.org/10.1016/j.wasman.2018.02.029>, 2018.
- Pataki, D. E., Ehleringer, J. R., Flanagan, L. B., Yakir, D., Bowling, D. R., Still, C. J., Buchmann, N., Kaplan, J. O., and Berry, J. A.: 555 The Application and Interpretation of Keeling Plots in Terrestrial Carbon Cycle Research, *Global Biogeochemical Cycles*, 17, 1022, <https://doi.org/10.1029/2001GB001850>, 2003.
- Phillips, N. G., Ackley, R., Crosson, E. R., Down, A., Hutyra, L. R., Brondfield, M., Karr, J. D., Zhao, K., and Jackson, R. B.: Mapping Urban Pipeline Leaks: Methane Leaks across Boston, *Environmental Pollution*, 173, 1–4, <https://doi.org/10.1016/j.envpol.2012.11.003>, 2013.
- Quay, P., Stutsman, J., Wilbur, D., Snover, A., Dlugokenky, E., and Brown, T.: The Isotopic Composition of Atmospheric Methane, *Global 560 Biogeochemical Cycles*, 13, 445–461, <https://doi.org/10.1029/1998GB900006>, 1999.

- Rice, A. L., Butenhoff, C. L., Teama, D. G., Röger, F. H., Khalil, M. A. K., and Rasmussen, R. A.: Atmospheric Methane Isotopic Record Favors Fossil Sources Flat in 1980s and 1990s with Recent Increase, *Proceedings of the National Academy of Sciences*, 113, 10791–10796, <https://doi.org/10.1073/pnas.1522923113>, 2016.
- Rice, D. D. and Claypool, G. E.: Generation, Accumulation, and Resource Potential of Biogenic Gas, *Am. Assoc. Pet. Geol. Bull.*, 65, 1981.
- 565 Rigby, M., Manning, A. J., and Prinn, R. G.: The Value of High-Frequency High-Precision Methane Isotopologue Measurements for Source and Sink Estimation: Methane Isotopologues in Inversions, *Journal of Geophysical Research: Atmospheres*, 117, 1–14, <https://doi.org/10.1029/2011JD017384>, 2012.
- Röckmann, T.: ROMEO - Romanian Methane Emissions from Oil & Gas, in: EGU General Assembly 2020, EGU2020-18801, Online, <https://doi.org/10.5194/egusphere-egu2020-18801>, 2020.
- 570 Röckmann, T., Gómez Álvarez, C. X., Walter, S., van der Veen, C., Wollny, A. G., Gunthe, S. S., Helas, G., Pöschl, U., Keppler, F., Greule, M., and Brand, W. A.: Isotopic Composition of H₂ from Wood Burning: Dependency on Combustion Efficiency, Moisture Content, and δD of Local Precipitation, *Journal of Geophysical Research*, 115, D17 308, <https://doi.org/10.1029/2009JD013188>, 2010.
- Röckmann, T., Eyer, S., van der Veen, C., Popa, M. E., Tuzson, B., Monteil, G., Houweling, S., Harris, E., Brunner, D., Fischer, H., Zazzeri, G., Lowry, D., Nisbet, E. G., Brand, W. A., Necki, J. M., Emmenegger, L., and Mohn, J.: In Situ Observations of the Isotopic Composition of Methane at the Cabauw Tall Tower Site, *Atmospheric Chemistry and Physics*, 16, 10 469–10 487, <https://doi.org/10.5194/acp-16-10469-2016>, 2016.
- 575 Rust, F.: Ruminant Methane $\delta(^{13}C/^{12}C)$ Values: Relation to Atmospheric Methane, *Science*, 211, 1044–1046, <https://doi.org/10.1126/science.7466376>, 1981.
- Rutherford, J. S., Sherwin, E. D., Ravikumar, A. P., Heath, G. A., Englander, J., Cooley, D., Lyon, D., Omara, M., Langfitt, Q., and Brandt, A. R.: Closing the Methane Gap in US Oil and Natural Gas Production Emissions Inventories, *Nature Communications*, 12, 4715, <https://doi.org/10.1038/s41467-021-25017-4>, 2021.
- 585 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- 590 Schaefer, H., Fletcher, S. E. M., Veidt, C., Lassey, K. R., Brailsford, G. W., Bromley, T. M., Dlugokencky, E. J., Michel, S. E., Miller, J. B., Levin, I., Lowe, D. C., Martin, R. J., Vaughn, B. H., and White, J. W. C.: A 21st-Century Shift from Fossil-Fuel to Biogenic Methane Emissions Indicated by ¹³CH₄, *Science*, 352, 80–84, <https://doi.org/10.1126/science.aad2705>, 2016.
- Schoell, M.: The Hydrogen and Carbon Isotopic Composition of Methane from Natural Gases of Various Origins, *Geochimica et Cosmochimica Acta*, 44, 649–661, [https://doi.org/10.1016/0016-7037\(80\)90155-6](https://doi.org/10.1016/0016-7037(80)90155-6), 1980.

- Schwietzke, S., Sherwood, O. A., Bruhwiler, L. M. P., Miller, J. B., Etiope, G., Dlugokencky, E. J., Michel, S. E., Arling, V. A., Vaughn, B. H., White, J. W. C., and Tans, P. P.: Upward Revision of Global Fossil Fuel Methane Emissions Based on Isotope Database, *Nature*, 600 538, 88–91, <https://doi.org/10.1038/nature19797>, 2016.
- Sherwood, O. A., Schwietzke, S., Arling, V. A., and Etiope, G.: Global Inventory of Gas Geochemistry Data from Fossil Fuel, Microbial and Burning Sources, Version 2017, *Earth System Science Data*, 9, 639–656, <https://doi.org/10.5194/essd-9-639-2017>, 2017.
- Sherwood, O. A., Schwietzke, S., and Lan, X.: Global $\delta^{13}\text{C}$ -CH₄ Source Signature Inventory 2020, 2021.
- Shindell, D. T., Fuglested, J. S., and Collins, W. J.: The Social Cost of Methane: Theory and Applications, *Faraday Discussions*, 200, 605 429–451, <https://doi.org/10.1039/C7FD00009J>, 2017.
- Smith, L. K., Lewis, Jr., W. M., Chanton, J. P., Cronin, G., and Hamilton, S. K.: Methane Emissions from the Orinoco River Floodplain, Venezuela, *Biogeochemistry*, 51, 113–140, <https://doi.org/10.1023/A:1006443429909>, 2000.
- Stavert, A. R., Saunio, M., Canadell, J. G., Poulter, B., Jackson, R. B., Regnier, P., Lauerwald, R., Raymond, P. A., Allen, G. H., Patra, P. K., Bergamaschi, P., Bousquet, P., Chandra, N., Ciais, P., Gustafson, A., Ishizawa, M., Ito, A., Kleinen, T., Maksyutov, S., McNorton, 610 J., Melton, J. R., Müller, J., Niwa, Y., Peng, S., Riley, W. J., Segers, A., Tian, H., Tsuruta, A., Yin, Y., Zhang, Z., Zheng, B., and Zhuang, Q.: Regional Trends and Drivers of the Global Methane Budget, *Global Change Biology*, p. gcb.15901, <https://doi.org/10.1111/gcb.15901>, 2021.
- Stell, A. C., Douglas, P. M. J., Rigby, M., and Ganesan, A. L.: The Impact of Spatially Varying Wetland Source Signatures on the Atmospheric Variability of δ D-CH₄, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 379, 615 20200442, <https://doi.org/10.1098/rsta.2020.0442>, 2021.
- Stieger, J., Bamberger, I., Siegwolf, R. T. W., Buchmann, N., and Eugster, W.: Source Partitioning of Atmospheric Methane Using Stable Carbon Isotope Measurements in the Reuss Valley, Switzerland, *Isotopes in Environmental and Health Studies*, 55, 1–24, <https://doi.org/10.1080/10256016.2018.1561448>, 2019.
- Sugimoto, A. and Fujita, N.: Hydrogen Concentration and Stable Isotopic Composition of Methane in Bubble Gas Observed in a Natural 620 Wetland, *Biogeochemistry*, 81, 33–44, <https://doi.org/10.1007/s10533-006-9028-4>, 2006.
- Szénási, B.: Atmospheric Monitoring of Methane Emissions at the European Scale, Ph.D. thesis, Université Paris-Saclay, 2020.
- Tarasova, O., Brenninkmeijer, C., Assonov, S., Elansky, N., Rockmann, T., and Brass, M.: Atmospheric CH₄ along the Trans-Siberian Railroad (TROICA) and River Ob: Source Identification Using Stable Isotope Analysis, *Atmospheric Environment*, 40, 5617–5628, <https://doi.org/10.1016/j.atmosenv.2006.04.065>, 2006.
- 625 Thompson, R. L., Nisbet, E. G., Pisso, I., Stohl, A., Blake, D., Dlugokencky, E. J., Helmig, D., and White, J. W. C.: Variability in Atmospheric Methane From Fossil Fuel and Microbial Sources Over the Last Three Decades, *Geophysical Research Letters*, 45, <https://doi.org/10.1029/2018GL078127>, 2018.
- Townsend-Small, A., Tyler, S. C., Pataki, D. E., Xu, X., and Christensen, L. E.: Isotopic Measurements of Atmospheric Methane in Los Angeles, California, USA: Influence of “Fugitive” Fossil Fuel Emissions: LOS ANGELES METHANE EMISSIONS, *Journal of Geophysical 630 Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD016826>, 2012.
- Toyoda, S., Suzuki, Y., Hattori, S., Yamada, K., Fujii, A., Yoshida, N., Kouno, R., Murayama, K., and Shiomi, H.: Isotopomer Analysis of Production and Consumption Mechanisms of N₂O and CH₄ in an Advanced Wastewater Treatment System, *Environmental Science & Technology*, 45, 917–922, <https://doi.org/10.1021/es102985u>, 2011.
- 635 Turner, A. J., Frankenberg, C., Wennberg, P. O., and Jacob, D. J.: Ambiguity in the Causes for Decadal Trends in Atmospheric Methane and Hydroxyl, *Proceedings of the National Academy of Sciences*, 114, 5367–5372, <https://doi.org/10.1073/pnas.1616020114>, 2017.

- Tyler, S. C., Rice, A. L., and Ajie, H. O.: Stable Isotope Ratios in Atmospheric CH₄: Implications for Seasonal Sources and Sinks, *Journal of Geophysical Research*, 112, D03 303, <https://doi.org/10.1029/2006JD007231>, 2007.
- Umezawa, T., Aoki, S., Kim, Y., Morimoto, S., and Nakazawa, T.: Carbon and Hydrogen Stable Isotopic Ratios of Methane Emitted from Wetlands and Wildfires in Alaska: Aircraft Observations and Bonfire Experiments, *Journal of Geophysical Research*, 116, D15 305, <https://doi.org/10.1029/2010JD015545>, 2011.
- 640 US Energy Information Administration: International Energy Statistics Database, 2021.
- Varga, T., Fisher, R. E., France, J. L., Haszpra, L., Jull, A. J. T., Lowry, D., Major, I., Molnár, M., Nisbet, E. G., and László, E.: Identification of Potential Methane Source Regions in Europe Using $\delta^{13}\text{C}_{\text{CH}_4}$ Measurements and Trajectory Modeling, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD033963>, 2021.
- 645 Varon, D. J., McKeever, J., Jervis, D., Maasackers, J. D., Pandey, S., Houweling, S., Aben, I., Scarpelli, T., and Jacob, D. J.: Satellite Discovery of Anomalously Large Methane Point Sources From Oil/Gas Production, *Geophysical Research Letters*, 46, 13 507–13 516, <https://doi.org/10.1029/2019GL083798>, 2019.
- Vigano, I., Holzinger, R., Keppler, F., Greule, M., Brand, W., Geilmann, H., van Weelden, H., and Röckmann, T.: Water Drives the Deuterium Content of the Methane Emitted from Plants, *Geochimica et Cosmochimica Acta*, 74, 3865–3873, <https://doi.org/10.1016/j.gca.2010.03.030>, 2010.
- 650 Wahlen, M., Tanaka, N., Henry, R., Deck, B., Zeglen, J., Vogel, J., Southon, J., Shemesh, A., Fairbanks, R., and Broecker, W.: Carbon-14 in Methane Sources and in Atmospheric Methane: The Contribution from Fossil Carbon, *Science*, 245, 286–90, <https://doi.org/10.1126/science.245.4915.286>, 1989.
- Waldron, S., Watson-Craik, I. A., Hall, A. J., and Fallick, A. E.: The Carbon and Hydrogen Stable Isotope **Composi-**
- 655 **tion** of Bacteriogenic Methane: A Laboratory Study Using a Landfill Inoculum, *Geomicrobiology Journal*, 15, 157–169, <https://doi.org/10.1080/01490459809378073>, 1998.
- Waldron, S., Lansdown, J., Scott, E., Fallick, A., and Hall, A.: The Global Influence of the Hydrogen Isotope Composition of **Wa-**
- 660 **ter** on That of Bacteriogenic Methane from Shallow Freshwater Environments, *Geochimica et Cosmochimica Acta*, 63, 2237–2245, [https://doi.org/10.1016/S0016-7037\(99\)00192-1](https://doi.org/10.1016/S0016-7037(99)00192-1), 1999.
- Walter, S., Röckmann, T., and The MEMO2 team: MEMO2: MEthane Goes MObile – MEasurements and MOdelling, in: International Symposium on Non-CO₂ Greenhouse Gases (NCGG), Unpublished, Amsterdam, The Netherlands, 2019.
- Warwick, N. J., Cain, M. L., Fisher, R., France, J. L., Lowry, D., Michel, S. E., Nisbet, E. G., Vaughn, B. H., White, J. W. C., and Pyle, J. A.: Using $\delta^{13}\text{C}\text{-CH}_4$ and $\delta\text{D}\text{-CH}_4$ to Constrain Arctic Methane Emissions, *Atmospheric Chemistry and Physics*, 16, 14 891–14 908, <https://doi.org/10.5194/acp-16-14891-2016>, 2016.
- 665 Wassmann, R., Thein, U. G., Whiticar, M. J., Rennenburg, H., Seiler, W., and Junk, W. J.: Methane Emissions from the Amazon Floodplain: Characterization of Production and Transport, *Global Biogeochemical Cycles*, 6, 3–13, <https://doi.org/10.1029/91GB01767>, 1992.
- Whiticar, M. J.: Carbon and Hydrogen Isotope Systematics of Bacterial Formation and Oxidation of Methane, *Chemical Geology*, 161, 291–314, [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3), 1999.
- Woltemate, I., Whiticar, M. J., and Schoell, M.: Carbon and Hydrogen Isotopic Composition of Bacterial Methane in a Shallow Freshwater
- 670 Lake, *Limnology and Oceanography*, 29, 985–992, 1984.
- Xueref-Remy, I., Zazzeri, G., Bréon, F., Vogel, F., Ciais, P., Lowry, D., and Nisbet, E.: Anthropogenic Methane Plume Detection from Point Sources in the Paris Megacity Area and Characterization of Their $\delta^{13}\text{C}$ Signature, *Atmospheric Environment*, 222, 117 055, <https://doi.org/10.1016/j.atmosenv.2019.117055>, 2020.

-  Zavala-Araiza, D., Lyon, D. R., Alvarez, R. A., Davis, K. J., Harriss, R., Herndon, S. C., Karion, A., Kort, E. A., Lamb, B. K., Lan, X.,
675 Marchese, A. J., Pacala, S. W., Robinson, A. L., Shepson, P. B., Sweeney, C., Talbot, R., Townsend-Small, A., Yacovitch, T. I., Zimmerle,
D. J., and Hamburg, S. P.: Reconciling Divergent Estimates of Oil and Gas Methane Emissions, *Proceedings of the National Academy of
Sciences*, 112, 15 597–15 602, <https://doi.org/10.1073/pnas.1522126112>, 2015.
-  Zavala-Araiza, D., Omara, M., Gautam, R., Smith, M. L., Pandey, S., Aben, I., Almanza-Veloz, V., Conley, S., Houweling, S., Kort, E. A.,
680 Maasackers, J. D., Molina, L. T., Pusuluri, A., Scarpelli, T., Schwietzke, S., Shen, L., Zavala, M., and Hamburg, S. P.: A Tale of Two
Regions: Methane Emissions from Oil and Gas Production in Offshore/Onshore Mexico, *Environmental Research Letters*, 16, 024 019,
<https://doi.org/10.1088/1748-9326/abceeb>, 2021.
-  Zazzeri, G., Lowry, D., Fisher, R. E., France, J. L., Lanoisellé, M., Grimmond, C. S. B., and Nisbet, E. G.: Evaluating Methane Inventories
by Isotopic Analysis in the London Region, *Scientific Reports*, 7, 4854, <https://doi.org/10.1038/s41598-017-04802-6>, 2017.
-  Zobitz, J., Keener, J., Schnyder, H., and Bowling, D.: Sensitivity Analysis and Quantification of Uncertainty for Isotopic Mixing Relationships
685 in Carbon Cycle Research, *Agricultural and Forest Meteorology*, 136, 56–75, <https://doi.org/10.1016/j.agrformet.2006.01.003>, 2006.