

1 Geomorphological landslide inventory map of the Daunia Apennines, 2 southern Italy.

3 Francesca Ardizzone¹, Francesco Bucci¹, Mauro Cardinali^{1*}, Federica Fiorucci¹, Luca Pisano², Michele
4 Santangelo¹, Veronica Zumpano²

5 ¹CNR - IRPI, Via della Madonna Alta 126, 06128, Perugia, Italy

6 ²CNR - IRPI, Via Amendola 122, 70126, Bari, Italy

7 *Correspondence to: Mauro Cardinali (mauro.cardinali@irpi.cnr.it)

8 **Keywords**

9 Landslide inventory map, geomorphology, landslides, Daunia, expert mapping, Southern Italy

10 **Abstract**

11 Detailed and accurate geomorphological historical landslide inventory maps are an invaluable source of information for many
12 research topics and applications. Their systematic preparation worldwide has been advised by many researchers as it may
13 foster our knowledge on landslides, their spatial and temporal distribution, their potential interaction with the built
14 environment, their contribution to landscape dynamics, their response to climate change in the past. Due to the extreme
15 variability of the morphological and radiometric elements that can reveal historical landslides, geomorphological historical
16 landslide inventory maps are produced by expert interpretation, which makes it a time consuming and expensive process,
17 which often discourages wide area mapping activities. In this paper we present a new geomorphological historical landslide
18 inventory map for a 1,460 km² area in the Daunia Apennines, the northern-western sector of the Apulia (Puglia) region, in
19 Southern Italy. The inventory contains 17,437 landslides classified according to relative age, type of movement and estimated
20 depth. Landslides were mapped according to rigorous and reproducible criteria applied by two teams of expert photo-
21 interpreters to two sets of stereoscopic aerial photographs taken in 1954/55 and 2003. The dataset consists of a digital archive
22 publicly available at <https://doi.pangaea.de/10.1594/PANGAEA.942427> (Cardinali et al., 2022).

23 **1 Introduction**

24 Landslides are widespread natural hazards that occur worldwide posing threat to population, structures and infrastructures,
25 causing relevant economic damage to society (Donnini et al., 2017; Froude and Petley, 2018; Petley, 2012). Availability of
26 information on landslides locations, types, and sizes is a key basic information for many activities, from research to emergency
27 support and land planning. Nevertheless, expert landslide mapping is not an investment priority in many countries and
28 institutions, hence homogeneous wide scale landslide maps are rare (Guzzetti et al., 2012; [Bucci et al., 2021](#)), although the

29 availability of a landslide inventory map is fundamental to the production of landslide hazard and risk maps (Thiery et al.,
30 2020). As a result, several applications spanning from land management to susceptibility modelling that would benefit from
31 more homogeneous and complete datasets rely on often poor or incomplete/inconsistent data.

32 Any landslide mapping activity assumes (i) that landslides leave discernible features on the territory, i.e., a “landslide
33 signature” made by radiometric and morphologic elements (Fiorucci et al., 2018; Guzzetti et al., 2012; Santangelo et al., 2022),
34 (ii) that, in case of use of remote sensing, the data used (e.g. images) portray them, (iii) that the technique adopted is adequate
35 to the data used, and (iv) that the operator (or the automatic or semi-automatic classification system) is able to exploit those
36 features to detect and map landslides. Therefore, any landslide mapping activity is affected by a degree of incompleteness and
37 inconsistency that stems from these basic assumptions. Landslides can remain unnoticed since (i) features were cancelled by
38 erosion, other landslides and/or human activity (Malamud et al., 2004), (ii) the type of image or its spectral and spatial
39 resolution may be inadequate for some types or sizes of landslides or for some locations (e.g. slope orientation), (iii) interpreters
40 or (semi-)automatic systems may be unable to map some landslides using certain images or techniques (e.g. lack of experience,
41 poor or incorrect system training).

42

43 Landslide inventory maps (LIMs) are the simplest tools to represent landslides distribution in a territory. They usually display
44 landslide locations preferably as polygons rather than points or lines which should be used only for scale representation issues.
45 LIMs usually store landslide attributes such as type of movement, estimated time of occurrence, estimated depth, activity at
46 the date of the observation if available, and other more detailed data depending on availability and on the scale of the work
47 (Bucci et al., 2021; Guzzetti et al., 2012; Santangelo et al., 2015).

48 Different types of inventories can be prepared. Event inventories (E-LIMs, Antonini et al., 2002; Ardizzone et al., 2012;
49 Donnini et al., 2017; Fiorucci et al., 2017; Harp and Jibson, 2017; Mondini et al., 2012; Santangelo et al., 2022) report
50 landslides that were triggered by specific events (i.e., rainfall, rapid snowmelt, volcanic eruption, earthquake).
51 Geomorphological inventories (G-LIMs) report landslides that can be recognised by geomorphologists usually from the expert
52 interpretation of stereoscopic aerial photographs (Bucci et al., 2021; Cardinali et al., 2001; Guzzetti et al., 2012; Santangelo et
53 al., 2015) but also LiDAR derived images are widely used (Niculiță et al., 2016; Razak et al., 2013; Schulz, 2004; Van Den
54 Eeckhaut et al., 2007). They can be seen as the result of many landslide events over thousands of years (Malamud et al., 2004).
55 Multi-temporal landslide inventories (M-LIMs) include the information of G-LIMs and also report landslides that occurred in
56 the last tens of years as visible on historical images available at (possibly) regular time steps of several years (Galli et al., 2008;
57 Guzzetti et al., 2012; Zumpano et al., 2020).

58 G-LIMs can be prepared for wide areas and provide a fundamental source of information about landslides that occurred in the
59 last tens of thousands years. This particular type of inventory is an invaluable source of data but is time consuming and requires
60 a high level of training and experience to get accurate and reliable results (Guzzetti et al., 2012), which usually hampers the
61 systematic preparation of G-LIMs at regional or even national/continent scale.

62

63 This paper presents a new G-LIM prepared for the Daunia mountains in Puglia Region, Southern Italy. For this area,
64 historically affected by diffuse slow moving large landslides, the Regional government required the preparation of a new
65 landslide inventory to overcome the inhomogeneity among the existing inventories in terms of spatial distribution of landslides
66 and of techniques adopted and working and publication scales. The landslide inventory map was produced through the
67 interpretation of two sets of stereoscopic aerial photographs, taken in 1954/55 and 2003, supplemented by targeted field checks.
68 The new landslide inventory is an entirely original dataset openly accessible at
69 <https://doi.pangaea.de/10.1594/PANGAEA.942427> (Cardinali et al., 2022).

70 **2 Study area**

71 The study area extends for 1,460 km², in the northern-western sector of the Apulia (Puglia in Italian) region (Southern Italy)
72 (**Figure 1**). The area corresponds to the Daunia Apennine, located in the external (i.e. eastern) part of the Southern Apennine
73 fold and thrust belt system.


74 The southern Apennine is the result of compressive tectonic dynamics, characterised by an oblique collision of the Calabrian
75 forearc with the Apulian margin (Filice and Seeber, 2019) evolved from an antecedent subductive phase of thrust-emplacement
76 (Vitale et al., 2011; Vitale and Ciarcia, 2013). Currently, the external (eastern) margin preserves the compressive tectonics
77 while the internal sector is characterised by an extensive dynamic with tension faults that dissect the pre-existent fold and
78 thrust structure (Brozzetti et al., 2009; Schiattarella et al., 2003).

79 The western sector of the Daunia Apennine is characterised by a wide variety of formations, mainly clay-rich flysch lithologies
80 having different mechanical properties, highly affected by folds and faults (Cotecchia et al., 2020; Losacco et al., 2021;
81 Pellicani et al., 2014b), resulting in medium-to-high relief with the elevations ranging between 55 and 1152 m a.s.l., where the
82 highest peak is represented by Monte Cornacchia. Morphologically, it is strongly influenced by both the prolonged tectonic
83 stresses and by the differences in composition and erodibility, where selective erosion has alternatively produced gentler and
84 rounded slopes on the more erodible clayey formations, and steep slopes cut on harder rocks. In this sector, the orography
85 associated with the widespread presence of clay-rich materials and the intense deformation is the main cause of landslides
86 (Ciarcia et al., 2003; Losacco et al., 2021; Spalluto et al., 2021; Wasowski et al., 2010; Zumpano et al., 2020). The eastern
87 sector is characterised by the Sub-Apennine clay formation and alluvial deposits, and by gentle slopes slightly dipping towards
88 NE. In this sector, several orders of terraced fluvial deposits overlay the sub-Apennine clay formation. Here, mass movements
89 are mainly concentrated along the scarps of the terraces.

90 The Daunia Apennine is influenced by a Mediterranean sub-humid climate with mild and often wet winters and usually hot
91 and dry summers. In general, inter-annual temperature variations are significant and the total yearly precipitation seldom

92 exceeds 1000 mm (Wasowski et al., 2012, 2010). In the area, torrential streams flow NE-SW, draining the sediments from the
93 Daunia Mountains, through the Tavoliere Plain towards the Adriatic Sea.

94 In response to the litho-structural and morpho-climatic setting of this area, landslides are widespread. They are mostly slow
95 moving slide-type and flow-type movements, involving soil and earth, and secondarily rock material. Rapid moving (i.e. debris
96 flows) and fast moving (i.e. rock falls) landslides are less abundant (Wasowski et al., 2010, 2012; di Lernia et al., 2022;
97 Zumpano et al., 2020) . The widespread presence of landslides in the area is a major cause of damage ~~Landslides are widespread~~
98 ~~in the area, especially in the western sector, where landslides are a major cause of damage~~ to urban settlements, to the road
99 network, and to the stability of buildings. These features determine a direct effect on the development and the economy of the
100 area (Pellicani et al., 2014b, a; Zumpano et al., 2020).



101
102 **Figure 1** - Location map of the study area. Base map: DTM at 8 m resolution made available by the Regione Puglia
103 administration.
104

105 3 Materials and methods

106 The procedure adopted to prepare the landslide inventory map is illustrated in **Figure 2**. The overall procedure consists in three
107 main phases: (i) data collection; (ii) photo-interpretation; and, (iii) editing. The entire work was carried out by a “mapping
108 team” made up by four interpreters and one supervisor, and an “editing team” made by three GIS editors.

109 The first stage of the work consisted in the collection of data available for the study area (i.e., ancillary data, **Figure 2**; complete
110 list in the Appendix A) and the pre-processing of the aerial-photographs to produce oriented stereo-models in absolute
111 coordinates to be used for the visual interpretation in a photogrammetric GIS environment. Such photogrammetric pre-
112 processing included the interior and exterior orientation of each pair of aerial photographs. For the interior orientation, a non-
113 metric camera model was adopted for which the focal length of the camera, the flight altitude and the pixel size were required,
114 while for the representation of the ground-to-image geometry (exterior orientation) Ground Control Points (GCPs) were used.
115 The planimetric coordinates (x, y) of GCPs were manually chosen by visual comparison of the aerial photographs and an
116 orthophoto available taken in 2006 (at 1 m resolution available for download at <http://www.sit.puglia.it/>) and 1988 (at 1 m
117 resolution, available as WMS service at <http://www.pcn.minambiente.it/mattm/servizio-wms/>). The altimetric coordinate (z)
118 of the GCPs were obtained from a Digital Elevation Model (8 m resolution available for download at <http://www.sit.puglia.it/>).
119 Stereo models were prepared for both aerial photographs epochs, 1954/55 and 2003, i.e. the oldest and the latest aerial
120 photographs acquisitions available for the entire study area.

121 The second phase of the work is the photo-interpretation (**Figure 2**) of the aerial photographs. At the very initial stage of the
122 work, the entire mapping team defined a legend through an expeditious photo-interpretation in representative
123 geomorphological settings and according to well-established landslide classifications schemas (Cruden and Varnes, 1996,
124 Hungr et al., 2014, WP/WLI, 1993). Then, divided in pairs, for each stereo-pair the mapping team prepared a preliminary
125 interpretation (1st level G-LIM in **Figure 2**) that was then reviewed by the supervisor and discussed with the mapping team.
126 After this step, an updated version of the map was produced (2nd level G-LIM in **Figure 2**). Each of the two mapping team
127 pairs performed their preliminary photo-interpretation in alternating strips along the flight plan. The side-lap between
128 subsequent strips was therefore common between the different couples of interpreters, and it was used to compare, correct and
129 homogenise the interpretation made by different sub-teams independently. Such a continuous interaction among
130 geomorphologists made it possible to best define the characteristics of the identified landslides, limiting interpretation
131 inconsistencies among operators. Finally, at significant advancement steps, extensive field checks were performed by the entire
132 mapping team to check, validate and correct the inventory based on field evidence. After the field check stage, the interpretation
133 phase was concluded and the landslide mapping considered quasi-definitive, i.e., occasional changes might be needed if
134 inconsistencies were observed in the third stage (e.g., consistency with contour maps and hydrography).

135 The third and last step of the procedure, (editing) consists in the preparation of the geographical database. The interpreted
136 features drawn as polylines by the mapping teams were verified on the digital topography, i.e., contour lines at scale 1: 5,000
137 (editing phase in **Figure 2**). The checked polylines were then converted into polygon features, and verified with topological

138 checks to avoid overlapping. Finally, the polygons were coded according to the adopted legend (legend description is available
139 in Section 4 and the database schema is described in the metadata of the digital archive available at
140 <https://doi.pangaea.de/10.1594/PANGAEA.942427>, Cardinali et al., 2022). Landslides recognised in the study area were
141 classified by type according to Cruden and Varnes, (1996) and Hungr et al., (2014). Additionally, according to WP/WLI,
142 (1993) landslides were classified based on the estimated depth (as shallow or deep-seated), and the inferred relative age (as
143 relict, very old, old or recent). Examples and descriptions of landslides of different type, relative age, and depth, as well as of
144 other geomorphological elements are detailed in Section 4. A more detailed explanation of the criteria used for the definition
145 of the legend was given by Bucci et al., (2021).
146

147 **3.1 Available data**

148 The Geomorphological landslide inventory map (G-LIM) of Daunia Sub-Apennines was prepared through systematic visual
149 interpretation of a set of 270 b/w stereoscopic aerial photographs taken in 1954/55 at 1: 33,000 scale and of 384 b/w
150 stereoscopic aerial photographs taken in 2003 at 1: 30,000 scale. Aerial photographs were provided as 800 dpi scanned copies.
151 All the ancillary data used for interpretation are listed and described in detail in Appendix A.

152 **3.2 Hardware and software**

153 For the interpretation phase we used the digital stereoscopic vision 3D PluraView System, composed of two monitors for the
154 stereoscopic vision of digital images and a 45° inclined mirror placed on the bisector of the two monitors equipped with passive
155 3D glasses, (<https://www.3d-pluraview.com/en/>). A dedicated computer, equipped with IMAGINE Photogrammetry software
156 (<https://www.hexagongeospatial.com/products/power-portfolio/imagine-photogrammetry>) was used for the pre-processing of
157 digital stereoscopic images, and ArcGIS software with the 3D Analyst extension tool, was used to digitise the 3D data in a GIS
158 environment (Ardizzone et al., 2013; Fiorucci et al., 2015). The spatial resolution of the images (~1m for 1954/55 epoch,
159 ~0.9m for 2003 images) and the zoom capability of the 3D PluraView System allow the identification of very small features.
160 The stereo pairs were investigated using a zoom level at scale 1: 2500 to obtain mapped features compatible with a publication
161 scale of 1: 5000. The capabilities of ERDAS IMAGINE™ Photogrammetry Suite and ArcGIS™ Stereo Analyst of preparing,
162 managing and viewing multiple sets of oriented images all having the same reference system allowed the fast comparison of
163 landscape features in different epochs, improving the interpretation.
164




Figure 2. Flowchart of the procedure to prepare the geomorphological landslide inventory map. Pink rectangles indicate contributions of different teams/operators.

4 Geomorphological Landslide Inventory Map.

The inventory covers an area of 1,460 km² and include 17,347 landslides (**Figure 3**), corresponding to an average density of about 15.6 landslides per square kilometre (**Table 1**) if lowlands plains are excluded. Locally, landslide density reaches 60 landslides per square kilometre. Landslide size (Landslide Area, A_L in m²) is in the range $1.9 \times 10^1 < A_L < 6.8 \times 10^6$ (**Table 1**). Overall, landslides cover an area of 442 km², which represents the 39% of the hilly and mountainous portion of the study area.

In this section the different groups of elements composing the inventory will be analytically described. Sections 4.1 to 4.3 describe landslides classified according to their relative age, type classification and estimated depth. Section 4.4 describes the widespread landslides (i.e. landslides represented as groups of failures due to their small size and high spatial frequency). Section 4.5 describes the descriptive statistics of the landslide inventory, and Section 4.6 presents the geomorphological elements (i.e. elements that are considered in relation with slope evolution and that can provide useful information for landslide identification and mapping).

181 **Table 1.** Descriptive statistics of landslides by relative age.

	Number of features	Total Area [m ²]	Minimum area [m ²]	Maximum area [m ²]
Very old relict landslides	37	8.85×10 ⁷	1.38×10 ⁵	6.78×10 ⁶
Very old landslides	120	8.75×10 ⁷	5.82×10 ⁴	2.86×10 ⁶
Pre-2003 landslides	14,793	3.39×10 ⁸	7.1×10 ¹	2.3×10 ⁶
Recent landslides (2003)	2,049	5.01×10 ⁶	1.89×10 ¹	1.63×10 ⁵
Widespread landslides	348	1.13×10 ⁷	3.78×10 ²	3.54×10 ⁵
Entire inventory	17,347	4.421.89×10 ⁸	1.89×10 ¹	6.77×10 ⁶

182

183 **4.1 Landslide by relative age**

184 Based on their appearance in aerial photographs, landslides were classified according to their relative age, namely: (i) very old
185 relict landslides, (ii) very old landslides, (iii) old landslides, and (iv) recent landslides (**Figure 3**). The four relative age levels
186 are the result of a major subdivision, referring to the general morphologic appearance of landslides (Keaton and DeGraff,
187 1996), and are based on the assumption that evidence of landslides become less obvious with the increasing age, due, for
188 example, to erosion processes, vegetation growth, and occurrence of other landslides. Hence, older landslides are more difficult
189 to detect than more recent ones.

190 Landslides modifications within the two epochs (1954 and 2003) were negligible for very old relict and very old landslides,
191 whereas they were more frequent for old landslides (i.e. pre-2003). Therefore, the final landslide delineation considers by
192 definition the latest evidence within the 49-year time interval.

193 In our G-LIM, the most represented landslides in the study area are old landslides (**Table 1, Figure 3C**), corresponding to
194 14,793 landslides covering a total landslide area of 339 km². Recent landslides (**Figure 3D**) are also very numerous (2,049)
195 They are generally small, and cover a total area of 5 km². Very old relict and very old landslides (**Figure 3A, B**) are represented
196 by 37 and 120 landslides respectively, and cover an area of 88 and 87 km². The feature of being fewer and larger than more
197 recent landslides is common to other geomorphological inventories (Bucci et al., 2021, 2016a; Santangelo et al., 2015). The
198 spatial relationships between landslides belonging to the relative age groups are displayed in **Figure 4**, where a representative
199 example of the landslide relative age pattern for the entire inventory is shown.

200




Figure 3. Spatial distribution of (A) very old relict landslides, (B) very old landslides, (C) old landslides and (D) recent landslides.

201

202 **Figure 4B-E** shows the inventory disaggregated according to the four relative age classes recognised. The sequence of the
 203 four panels represents a timeline, where landslides of each time step are shown with their own colour, the same as **Figure 4A**,
 204 whereas landslides belonging to antecedent time steps are represented in white.

205

206 In addition to the four mentioned relative age classes, a generation principle based on crosscut relationships, defines minor
 207 subdivisions within each age class: younger landslides cover the older ones. For the overall inventory, this criterion allowed
 208 us to detect up to: i) two landslides generations within the relict landslides, ii) two generations within the very old landslides,
 209 iii) four generations within the old landslides, iv) two generations within the recent landslides. Such minor subdivisions are
 210 applicable only to landslides that overlap, i.e. it is not applicable for landslides that do not overlap. For instance, in **Figure 4D**
 211 the age class “old” records up to three overlapping classes, indicated respectively in light orange (first failure “old”), orange
 212 (second failure “old”) and red (third failure “old”).

213




Figure 4 - Example of the relative age classes in the area of the Pietramontecorvino municipality (see **Figure 1** for reference). (A) Detail of the inventory in the selected location. Black outline polygon indicates the area in B to I. Very old relict landslides (B), very old landslides (C), old landslides (D), and recent landslides (E) are portrayed in detail. Black outlined polygons in B to E are represented singularly in a perspective view on © GoogleEarth™ in B' to F'. In the legend, Rec: Recent landslides; O1, O2, O3: old landslides of first, second and third generation; VO1: Very old landslides of first generation; R: relict landslides; OAF: old alluvial fan; Te: terrace.

214

215

216 Landslides classified as very old relict (**Figure 4B**) generally affect entire slopes, where the energy of the relief is greatest.
 217 They are usually controlled by the geological and lithological structure and the presence of main faults (Bucci et al., 2016a).
 218 Very old relict landslides are deeply dismantled by the erosive action of watercourses and often reshaped by recurring
 219 gravitational phenomena. These morphological modifications are often related to the regional morphological and tectonic
 220 evolution, which has determined considerable base level variations in the study area. As a consequence, these landslides are
 221 often suspended with respect to the present day base level, and totally or partially isolated from the recent evolution of the
 222 drainage network. Therefore, very old relict landslides are considered to have occurred under geomorphological and climatic
 223 conditions different from the present day (WP/WLI, 1993).

224 Landslides classified as very old (**Figure 4C**) show evidence of active and passive interaction with river dynamics, being often
 225 partly eroded from them and locally modifying their path. Also, morphological modifications of very old landslides can be
 226 induced by the occurrence of other mass movements over time. Very old landslides are generally large and in agreement with
 227 the present day river network, they are mainly distributed where the local relief is high and they often occur in or near very
 228 old relict landslides, as local reactivations.

229 Landslides classified as old (**Figure 4D**) present recognizable evidence on both 1954/55 and 2003 aerial photographs epochs.
 230 These landslides generally present morphological characteristics typical of landslides which are not modified by erosion (e.g.,
 231 concave-convex shape of the slope, presence of steps and back slopes in the deposit area). Old landslides tend to cluster
 232 spatially, as it is evident from the several generations of mass movements that were recognised. Spatial clustering of old
 233 landslides is even more evident in the vicinity of very old or very old relict landslides. Old landslides are hypothesised to have
 234 occurred in more recent time (yet undefined) compared to very old and very old relict landslides.

235 Landslides classified as recent (**Figure 4E**) are represented by landslides that show evidence of having occurred close to the
 236 date of the 2003 aerial image. The diagnostic elements for the identification of recent landslides refer not only to the
 237 morphological evidence but above all to the spectral elements (photographic tone and contrast) which can be perceived even
 238 with a bi-dimensional (2D) view of the images. Recent landslides are mainly shallow failures essentially involving the soil
 239 horizons or the alteration of the debris cover for a few metres in depth. Their spatial distribution is ubiquitous and only limited
 240 to the areas where the triggering event induced the landslides. Therefore, they were found both in previously unaffected slopes
 241 or within pre-existing landslides.

242 As opposed to the recent landslides, which evidence is stronger, some uncertainty characterises the delineation of older
 243 landslide borders, which boundaries can be affected or even dismantled by different degrees of erosion, or covered by more
 244 recent slope failures. General considerations on relative age criteria of classification (Keaton and DeGraff, 1996) support our
 245 own experimental observations on landslide morphological appearance that the uncertainty degrees in landslide mapping
 246 increase with the increasing landslide age (Bucci et al., 2021). However, we do not have information on absolute age of the
 247 landslides in our study area, and only hypothesise that the majority of the mapped landslides occurred in the last 10–20 Kyr,
 248 as suggested by findings of recent studies on landslide age in Southern Apennine (Gioia et al., 2011) and elsewhere (Niculiță
 249 et al., 2016; Pánek et al., 2014).

250 4.2 Landslide by type

251 The type of the landslides is assigned based on the morphological and radiometric signature of each landslide on the aerial
 252 photographs. Landslide types included in the legend of the G-LIM are (i) slide type landslides (which include deep seated rock
 253 slides or earth slides, shallow soil slides or earth slides), (ii) earth flow, (iii) slide-earth flow, (iv) debris flow, (v) rock fall and
 254 (vi) sackung (**Table 2**). In Table 2, widespread landslides refer to polygons representing groups of landslides whose size is
 255 smaller than the smallest feature that can be represented at the publication scale of the final map.
 256

257 **Table 2.** Descriptive statistics of landslides by type.

Landslide type	Number of features	Total Area [m ²]	Minimum area [m ²]	Maximum area [m ²]
Slide type	9,867	2.9×10 ⁸	1.9×10 ¹	6.8×10 ⁶
Earth flow	2,435	4.6×10 ⁷	3.6×10 ¹	1.3×10 ⁷

Slide-earth flow	4,657	1.6×10^8	8.8×10^1	2.9×10^6
Debris flow	38	6.5×10^5	1.4×10^2	1.4×10^5
Rock fall	1	6.0×10^4	6.0×10^4	6.0×10^4
Sackung	1	5.0×10^5	5.0×10^5	5.0×10^5
Total	16,999	4.3×10^8	1.9×10^1	1.3×10^7
Widespread landslide	Number of features	Total Area [m²]	Minimum area [m²]	Maximum area [m²]
Shallow earth/soil slides and earth flows	138	7.1×10^6	5.2×10^2	3.5×10^5
Debris flow	193	4.1×10^6	3.8×10^2	1.4×10^5
Rock fall	17	1.3×10^5	5.2×10^2	2.4×10^4
Total	348	1.1×10^7	3.8×10^2	3.5×10^5

258

259 According to the type of movement, the most represented landslides in the study area are slides (9,867 landslides, **Figure 5A**)
 260 covering a total landslide area of 287 km². The 4,657 slide-earth flows (**Figure 5B**) cover an area of 158 km², whereas earth
 261 flows are represented by 2435 landslides (**Figure 5C**) covering an area of 46 km². **Table 2** reveals that these three classes
 262 represent more than 95% of the total landslides, while debris flows (38), rock falls (1) and sackungs (1) are rare.

263

264 The main features of the three more represented landslide types are illustrated in **Figure 6**. Inspection of the figure highlights
 265 the main cartographic differences (**Figure 6A, C, E**) between different landslide types (**Figure 6B, D, F**).

266 **Figures 6A, B** illustrates the slide type landslides. Such failures present a well-defined scarp which can be semi-circular
 267 (rotational slides) or angular (translational slides). The slide deposit is convex, with a morphologically depressed head of the
 268 deposit, characterised by a centripetal drainage and local counter-slopes, and a toe characterised by a typical upward bulge
 269 (**Figure 6B**). It is reasonable to expect a volume balancing between escarpment and deposit areas since slides are usually
 270 characterised by low mobility of landslide material. This is clear in **Figure 6A, B** that shows a planimetric and altimetric shift
 271 of about 25 metres of the displaced material, with no evidence of chaotic rearrangement and/or volumetric changes.

272 Slide-earth flows (**Figure 6C, D**) initiate as slides, then evolving into flows. Therefore, they show the characteristics of slides
 273 (most commonly rotational) in the escarpment area and in the head of the deposit, where local endorheic conditions can
 274 promote seasonal swamps with diffuse organic soils development (**Figure 6C, D**). On the other hand, the transport zone and
 275 accumulation zone are more similar to earth flows, characterised by the typical lobe shape of the accumulation zone (**Figure**
 276 **6C, D**).




Figure 5. Spatial distribution of (A) slide type landslides, (B) slide-earth flows, (C) earth flows, (D) widespread landslides.

277

278 Earth flows (**Figure 6E, F**) are characterised by an overall elongated planar shape, with the median part narrower than
 279 detachment and accumulation zones. The mobility of earth flows is generally higher than slides, which is also evident in their
 280 more elongated shape. A typical earth flow is usually bounded by lateral streams draining the deposits into the main river
 281 valley, which typically experiences narrowing of the valley section and erosion on the opposite side of the landslide toe (**Figure**
 282 **6F**)

283 Debris flows and rock falls are not statistically represented in our inventory because the Daunia Apennine lacks the
 284 environmental conditions favouring their development, such as steep rocky slopes and sub-vertical cliffs. However, local
 285 geomorphological conditions (e.g. steep edges of suspended fluvial terraces) can promote the formation of coalescing
 286 phenomena of small debris flows and small rock falls, which cannot be mapped individually but were included within areas
 287 with widespread landslides (**Figure 5D**, Section 4.4). Finally, sackungs are poorly represented in our study area.

288




Figure 6. Examples in map (left column) and on © GoogleEarth™. images (right column) of slide type failures (A,B), slide-earthflow landslides (C,D), earthflows (E,F). For each row, black outlined polygons and camera symbols in the left image identify the landslide shown in the corresponding image on the right and the point of view, respectively.

289 **4.3 Landslide estimated depth**

290 The estimated depth of a landslide is assigned based on the main morphological characteristics of the landslide, such as: slope
291 height, and extent of curvatures along the slope (convexity and concavity). Based on these morphological characteristics,
292 landslides were classified as deep-seated or shallow.

293 Deep-seated landslides are predominantly represented by failures that cover considerable extents, even exceeding one square
294 kilometre. Many of the deep-seated landslides can be classified as very old relict, very old and old. They often involve
295 considerable volumes of material, and can alter the local morphology and geological structure. In the inventory map, deep-
296 seated landslides are represented with two polygons, distinguishing the deposit area from the scarp area.

297 Shallow landslides are generally small ($\sim 10^4$ m²) and mainly represented by slide type and flow type landslides. They are
298 generally characterised by a scarp not very pronounced, with an estimated height of less than 2 metres, and by a deposit without
299 evident concavities and convexities on its surface. In the inventory, shallow landslides are mapped as single polygons including
300 both the scarp and the deposit area.

301 **4.4 Widespread landslides**

302 The areas of widespread landslides (**Table 2, Figure 5D**) are represented by landslides which size is smaller than the smallest
303 feature that can be represented at the publication scale of the final map. In the map we distinguished between: i) shallow
304 earth/soil slides and earth flows, ii) debris flows, iii) rock falls. Their spatial distribution is mainly related to locally steep
305 slopes, for instance in the vicinity of badlands, in the crown areas of pre-existing large landslides, or along the edges of ancient
306 suspended fluvial terraces (**Figure 7**). Widespread shallow slides and flows typically involve weathered clayey soils, while
307 widespread debris flows and rock falls are commonly fed by loose debris or poorly cemented conglomerates and breccias.
308 Finally, it was not possible to assign a well-defined age to widespread landslides, although their morphological evidence
309 suggests seasonal reactivations.

310 **4.5 Descriptive statistics of the G-LIM**

311 Descriptive statistics of landslide number and size are shown in **Figure 7**. Box plots in **Figure 7A, B** show the distribution of
312 landslide areas for the landslide classified based on the expected relative age (**Figure 7A**) and the type of movement (**Figure**
313 **7B**).

314 Inspection of the plot in **Figure 7A** confirms that slides, slide-earth flows and flows are the most represented landslides of the
315 inventory. Among these three classes, the slide type landslides are characterised by the largest size variability, while the slide
316 earth flows are on average slightly larger than the others. In the plot of **Figure 7A**, landslides (on the left) are separated by
317 areas of widespread landslides (on the right) because the latter are not related to a single feature and cannot be directly
318 compared to single landslides. This is the reason why the areas of widespread landslides are on average larger than the single

319 landslides. In particular, the areas of widespread soil slides and earth flows are larger than the others because they typically
320 involve edges of fluvial terraces characterised by strong lateral continuity.

321 The plot in **Figure 7B** shows the size distribution of the four relative age classes of landslides. The clear separation of the age
322 classes according to their median values is a statistically robust indication of the reliability of this landslide classification
323 strategy, which is underused in the landslide-oriented international literature.

324 Comparing **Figure 7A** and **7B**, it is worth noting that the median area of all landslide types is around 10^5 m², which corresponds
325 to the median area of the old landslides group. This was expected since this is the most represented landslide age class within
326 the inventory.

327 **Figure 7C** shows landslides count (represented by a colour gradient and labels) and cumulated area (proportional to circle
328 sizes), grouped by relative age classes and landslide type. Visual inspection of the plot reveals that relict and very old landslides
329 are not represented by small landslides, and in particular by debris flows and rock falls, which tend to be small and easily
330 obliterated by erosion and subsequent failures. Further evidence is that a large portion of the total landslide area is occupied
331 by few relict and very old kilometre-scale slope failures, whereas a decreasing trend both in cumulated area and total number
332 of landslides is evident through time for all landslide types within the old landslides age class. Such evidence suggests a size
333 threshold effect of previous landslides on subsequent slope movements, probably controlled by slope-scale morphological and
334 hydrological perturbations induced by the occurrence of the first failure.




Figure 7. Plots summarising landslide statistics. (A) Landslides number (represented by a colour gradient and labels) and cumulated area (proportional to circle sizes), grouped by relative age classes and landslide type (SK, sackung; SEF, slide earth flow; S, slide; RF, rock fall; DF, debris flow; EF, earth flow). (B) Distribution of landslide areas within each relative age class. The relative age classes are indicated through a letter (R, relict; V, very old; and O, old), and a subscript number that indicates the generation within the age class (1–4). (C) Scatterplot showing cumulative size and count of landslides grouped by relative age in the abscissa and type in the ordinate. Landslide number is represented by a colour gradient and labels; total area is proportional to circle sizes.

336 4.6 Geomorphological elements

337 In the landslide inventory map, some geomorphological elements are also reported in addition to the mass movements (**Figure**
338 **8**). Such elements can be considered in relation with slope evolution and can provide useful information for landslide
339 identification and mapping. In particular, our map includes alluvial deposits, fluvial terraces, and alluvial fans. Alluvial
340 deposits are flat and always located in the lowest portions of the *Tavoliere delle Puglie* Plain and along the main water courses
341 draining the hilly and mountain areas of the Daunia Apennine. Older alluvial/fluvial deposits are nowadays suspended over
342 the present-day base level and are organised in fluvial terraces at different elevations, recording the ongoing deepening of the
343 drainage network during the Quaternary. Alluvial fans were recognized and mapped in two relative age classes according to
344 their appearance. They were classified as relict if dismantled and dissected due to the incision of the present-day river network.
345 They were instead classified as recent if well preserved.

346




Figure 8 – (A) Excerpt of the landslide inventory map. Landslides were coloured in white for enhancing the geomorphological elements. (B) Birds Eye view of the same area as (A) prepared using © GoogleEarth™. **Figure 8** – (A) Excerpt of the landslide inventory map. Landslides were coloured in white for enhancing the geomorphological elements. (B) Birds Eye view of the same area as (A) prepared using © GoogleEarth™.

347

348 5 Conclusion

349 The landslide inventory map presented in this paper is a new catalogue of landslide phenomena in the Daunia Apennine. It
350 provides new data uniformly distributed over the study area at an unprecedented cartographic detail.

351 The new landslide inventory map was produced upon request of the authorities of the Puglia Region, to solve the heterogeneity
352 and limitations in the currently available landslide inventories (Pellicani et al., 2014b; Pellicani and Spilotro, 2015), mainly

353 due to: i) partial coverage of the territory, ii) heterogeneity in the degree of cartographic detail, iii) lack of metadata and other
354 methodological information, iv) geospatial inconsistencies where the different products overlap.

355 The new inventory represents a significant improvement upon earlier mapping because: (i) it presents a new homogeneous
356 landslide coverage for the Daunia Apennine, (ii) it is acquired with a uniform method and based on rigorous and reproducible
357 criteria, (iii) it keeps the same cartographic detail throughout the entire area, and (iv) it presents a systematic classification of
358 landslides by type, relative age, and depth, hence providing basic information at 1: 5000 scale for landslide characterization
359 and related land management activities. In addition, the spatially distributed information on landslide coverage portrayed in
360 the map, and the detailed scale of mapping, provide the fundamental information for landslide susceptibility and hazard
361 assessment (Guzzetti et al., 2005), and for supporting and integrating the study of seismic microzonation and the assessing of
362 seismic hazard both in urban and rural areas (Vignaroli et al., 2019). In addition to the technical and land management benefits,
363 our inventory provides new data to further investigation of the Quaternary evolution of the landscape of this part of the
364 Southern Apennines.

365 Analysis of the data revealed that the great majority of landslide volume was mobilised by relict and very old landslides,
366 whereas the landslide maximum size decreased over time and tended to cluster around pre-existing failures. The results have
367 relevance to determine the statistics of landslide size (Malamud et al., 2004), and to trigger detailed studies on the types,
368 patterns, and distribution of landslides in relation to geology (Bucci et al., 2016b) structure (Bucci et al., 2013), tectonic (Bucci
369 et al., 2014) and climatic forcing (Schiattarella et al., 2017). The presented dataset also documents the relationships between
370 landslides and other geomorphological elements, in particular with ancient and recent alluvial deposits and alluvial fans. These
371 fluvial related landforms can be used as input data to study quaternary depositional (Mirabella et al., 2018) and erosional events
372 (Mancini et al., 2020), and in combination with landslide information, to study the interplay of gravitational and fluvial
373 processes (Santangelo et al., 2013).

374 In the G-LIM presented, the vast majority of landslides are slow-moving, whereas rapid and fast landslides are rare (debris
375 flows and rock falls). Such evidence should be further confirmed by in-depth studies (e.g. multi-temporal inventories or
376 susceptibility models) as it may indicate a relatively low exposure of human life compared to other mountain areas of Southern
377 Apennines. On the other hand, it must be taken into account that the increasing frequency of extreme rainfall events may
378 trigger rapid and fast moving landslides also in this area, differently from what was observed in the past. This aspect further
379 shows that systematic landslide mapping to prepare new inventories and update the existing ones is crucial, particularly in the
380 ongoing climate change scenario (Donnini et al., under review).

381 We acknowledge that scarcity of data represents a limitation and a challenge for the landslide community. Most regional-scale
382 studies are less effective (or analysed) than expected due to the lack of easy sharing of landslide information in digital format.

383 Our work is also intended as a contribution to the broader geomorphological community, promoting the sharing of landslide
384 dataset at regional scale from various environments.

385 Finally, it is worth reminding that a landslide map, even if accurate, does not answer all questions regarding hazards or risk at
386 any scale. A Landslide map is informative about the area covered by landslides, but nothing can be said about the remaining
387 territory. The map dataset should be used as an indication to optimise resources and to plan investigations aimed at determining
388 landslide hazard and risk at a larger scale. In our case, the map dataset is published at 1: 5000 scale, and should be consulted
389 at most at the same detail level.

390

391 **6 Authors contributions**

392 We declare that all authors' contributions in preparing both the dataset and the manuscript are equivalent to that of a first
393 author. This is why all authors appear in alphabetical order.

394 **7 Acknowledgments**

395 This work was supported by the Civil Protection of the Puglia region, in the framework of the project “*Integrated assessment*
396 *of geo-hydrological instability phenomena in the Apulia region, interpretative models and definition of rainfall thresholds for*
397 *landslide triggering*” funded by the P.O.R. Puglia 2014-2020, Asse V - Azione 5.1. (Project identification number:
398 B82F16003840006).

399 **8 Data availability**

400 Dataset is available at: <https://doi.pangaea.de/10.1594/PANGAEA.942427> (Cardinali et al., 2022).

401

402 **Competing interests**

403 The authors declare that they have no conflict of interest.

404 **References**

405

406 Antonini, G., Ardizzone, F., Cardinali, M., Galli, M., Guzzetti, F., and Reichenbach, P.: Surface deposits and landslide
407 inventory map of the area affected by the 1997 Umbria-Marche earthquakes, *Bollettino della Società Geologica Italiana*,
408 Volume spe, 843–853, 2002.

409 Ardizzone, F., Basile, G., Cardinali, M., Casagli, N., Del Conte, S., Del Ventisette, C., Fiorucci, F., Garfagnoli, F., Gigli, G.,

410 Guzzetti, F., Iovine, G., Mondini, A. C., Moretti, S., Panebianco, M., Raspini, F., Reichenbach, P., Rossi, M., Tanteri,
411 L., and Terranova, O.: Landslide inventory map for the Briga and the Giampilieri catchments, NE Sicily, Italy, *Journal*
412 *of Maps*, 8, 176–180, <https://doi.org/10.1080/17445647.2012.694271>, 2012.

413 Ardizzone, F., Fiorucci, F., Santangelo, M., Cardinali, M., Mondini, A. C. A. C., Rossi, M., Reichenbach, P., Guzzetti, F.,
414 and Ardizzone F.: Very-High Resolution Stereoscopic Satellite Images for Landslide Mapping, in: *Landslide Inventory*
415 *and Susceptibility and Hazard Zoning*, Berlin, Heidelberg, Citation Key: CNRPRODOTTI89033, 95–101, 2013.

416 Brozzetti, F., Boncio, P., Lavecchia, G., and Pace, B.: Present activity and seismogenic potential of a low-angle normal fault
417 system (Città di Castello, Italy): Constraints from surface geology, seismic reflection data and seismicity,
418 *Tectonophysics*, 463, 31–46, <https://doi.org/10.1016/j.tecto.2008.09.023>, 2009.

419 Bucci, F., Cardinali, M., and Guzzetti, F.: Structural geomorphology, active faulting and slope deformations in the epicentre
420 area of the MW 7.0, 1857, Southern Italy earthquake, *Physics and Chemistry of the Earth, Parts A/B/C*, 63, 12–24,
421 <https://doi.org/10.1016/j.pce.2013.04.005>, 2013.

422 Bucci, F., Novellino, R., Tavarnelli, E., Prosser, G., Guzzetti, F., Cardinali, M., Gueguen, E., Guglielmi, P., and Adurno, I.:
423 Frontal collapse during thrust propagation in mountain belts: a case study in the Lucania Apennines, Southern Italy,
424 *Journal of the Geological Society*, 171, 571–581, <https://doi.org/10.1144/jgs2013-103>, 2014.

425 Bucci, F., Santangelo, M., Cardinali, M., Fiorucci, F., and Guzzetti, F.: Landslide distribution and size in response to
426 Quaternary fault activity: The Peloritani Range, NE Sicily, Italy, *Earth Surface Processes and Landforms*, 41,
427 <https://doi.org/10.1002/esp.3898>, 2016a.

428 Bucci, F., Mirabella, F., Santangelo, M., Cardinali, M., and Guzzetti, F.: Photo-geology of the Montefalco Quaternary Basin,
429 Umbria, Central Italy, *Journal of Maps*, 12, <https://doi.org/10.1080/17445647.2016.1210042>, 2016b.

430 Bucci, F., Santangelo, M., Fiorucci, F., Ardizzone, F., Giordan, D., Cignetti, M., Notti, D., Allasia, P., Godone, D.,
431 Lagomarsino, D., Pozzoli, A., Norelli, E., and Cardinali, M.: Geomorphologic landslide inventory by air photo
432 interpretation of the High Agri Valley (Southern Italy), *Journal of Maps*, 17, 376–388,
433 <https://doi.org/10.1080/17445647.2021.1943552>, 2021.

434 Cardinali, M., Antonini, G., Reichenbach, P., and Guzzetti, F.: Photo geological and landslide inventory map for the Upper
435 Tiber River basin, 2001.

436 Cardinali, M., Ardizzone, F., Bucci, F., Fiorucci, F., Pisano, L., Santangelo, M., and Zumpano, V.: Geomorphological
437 landslide inventory map of the Daunia Mountains, Southern Italian Apennines, PANGAEA [data set],
438 <https://doi.org/10.1594/PANGAEA.942427>, 2022.

439 Ciarcia, S., Di Nocera, S., Matano, F., and Torre, Ma. L.: Evoluzione tettono-sedimentaria e paleogeografica dei depocentri
440 «wedge-top» nell’ambito del «foreland basin system» pliocenico dell’Appennino meridionale (settore Irpino-Dauno),
441 *Boll. Soc. Geol. It.*, 122, 117–137, 2003.

442 Cotecchia, F., Santaloia, F., and Tagarelli, V.: Towards A Geo-Hydro-Mechanical Characterization of Landslide Classes:
443 Preliminary Results, *Applied Sciences*, 10, 7960, <https://doi.org/10.3390/app10227960>, 2020.

444 Cruden, D. M. and Varnes, D. J.: *Landslide Types and Processes*, Special Report National Research Council Transportation
445 Research Board, 247, 36–75, 1996.

446 [di Lernia, A., Cotecchia, F., Elia, G., Tagarelli, V., Santaloia, F., and Palladino, G.: Assessing the influence of the hydraulic
447 boundary conditions on clay slope stability: The Fontana Monte case study. *Engineering Geology*, 297, 106509,
448 <https://doi.org/10.1016/j.enggeo.2021.106509>, 2022.](https://doi.org/10.1016/j.enggeo.2021.106509)

449 Donnini, M., Napolitano, E., Salvati, P., Ardizzone, F., Bucci, F., Fiorucci, F., Santangelo, M., Cardinali, M., and Guzzetti,
450 F.: Impact of event landslides on road networks: a statistical analysis of two Italian case studies, *Landslides*, 14, 1521–
451 1535, <https://doi.org/10.1007/s10346-017-0829-4>, 2017.

452 [Donnini, M., Santangelo, M., Gariano, S.L., Bucci, F., Peruccacci, S., Alvioli M., Althuwaynee, O., Ardizzone, F., Bianchi,
453 C., Bornaetxea, T., Brunetti, M.T., Cardinali, M., Esposito, G., Grita, S., Marchesini, I., Melillo, M., Salvati, P.,
454 Yazdani, M., Fiorucci, F.: Landslides triggered by an extraordinary rainfall event in Central Italy on September 15,
455 2022, under review.](https://doi.org/10.1016/j.enggeo.2021.106509)

456 Filice, F. and Seeber, L.: The Culmination of an Oblique Time-Transgressive Arc Continent Collision: The Pollino Massif
457 Between Calabria and the Southern Apennines, Italy, *Tectonics*, 38, 3261–3280,
458 <https://doi.org/10.1029/2017TC004932>, 2019.

459 Fiorucci, F., Ardizzone, F., Rossi, M., and Torri, D.: The Use of Stereoscopic Satellite Images to Map Rills and Ephemeral

460 Gullies, *Remote Sensing*, 7, 14151–14178, <https://doi.org/10.3390/rs71014151>, 2015.

461 Fiorucci, F., Giordan, D., Santangelo, M., Dutto, F., Rossi, M., and Guzzetti, F.: Criteria for the optimal selection of remote
 462 sensing images to map event landslides, *Risk Assessment, Mitigation and Adaptation Strategies, Socioeconomic and*
 463 *Management Aspects*, <https://doi.org/10.5194/nhess-2017-111>, 2017.

464 Fiorucci, F., Giordan, D., Santangelo, M., Dutto, F., Rossi, M., and Guzzetti, F.: Criteria for the optimal selection of remote
 465 sensing optical images to map event landslides, 405–417, <https://doi.org/10.5194/nhess-18-405-2018>, 2018.

466 Froude, M. J. and Petley, D. N.: Global fatal landslide occurrence from 2004 to 2016, *Natural Hazards and Earth System*
 467 *Sciences*, 18, 2161–2181, <https://doi.org/10.5194/nhess-18-2161-2018>, 2018.

468 Galli, M., Ardizzone, F., Cardinali, M., Guzzetti, F., and Reichenbach, P.: Comparing landslide inventory maps,
 469 *Geomorphology*, 94, 268–289, <https://doi.org/10.1016/j.geomorph.2006.09.023>, 2008.

470 Gioia, D., Di Leo, P., Giano, S. I., and Schiattarella, M.: Chronological constraints on a Holocene landslide in an
 471 intermontane basin of the southern Apennines, Italy: Morphological evolution and palaeoclimate implications, *The*
 472 *Holocene*, 21, 263–273, <https://doi.org/10.1177/0959683610378879>, 2011.

473 Guzzetti, F., Reichenbach, P., Cardinali, M., Galli, M., and Ardizzone, F.: Probabilistic landslide hazard assessment at the
 474 basin scale, *Geomorphology*, 72, 272–299, <https://doi.org/10.1016/j.geomorph.2005.06.002>, 2005.

475 Guzzetti, F., Mondini, A. C., Cardinali, M., Fiorucci, F., Santangelo, M., and Chang, K.-T.: Landslide inventory maps: New
 476 tools for an old problem, *Earth-Science Reviews*, 112, 42–66, <https://doi.org/10.1016/j.earscirev.2012.02.001>, 2012.

477 Harp, E. L. and Jibson, R. W.: Inventory of landslides triggered by the 1994 Northridge, California earthquake,
 478 <https://doi.org/10.5066/F7Z60MKF>, 2017.

479 Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types, an update, *Landslides*, 11, 167–194,
 480 <https://doi.org/10.1007/s10346-013-0436-y>, 2014.

481 Keaton, J. R. and DeGraff, J. V.: Surface observation and geologic mapping, in: *Landslides: Investigation and mitigation.*,
 482 1996.

483 Losacco, N., Bottiglieri, O., Santaloia, F., Vitone, C., and Cotecchia, F.: The Geo-Hydro-Mechanical Properties of a
 484 Turbiditic Formation as Internal Factors of Slope Failure Processes, *Geosciences*, 11, 429,
 485 <https://doi.org/10.3390/geosciences11100429>, 2021.

486 Malamud, B. D., Turcotte, D. L., Guzzetti, F., and Reichenbach, P.: Landslide inventories and their statistical properties,
 487 *Earth Surf. Process. Landforms*, 29, 687–711, <https://doi.org/10.1002/esp.1064>, 2004.

488 Mancini, M., Vignaroli, G., Buccì, F., Cardinali, M., Cavinato, G. P., Di salvo, C., Giallini, S., Moscatelli, M., Polpetta, F.,
 489 Putignano, M. L., Santangelo, M., and Sirianni, P.: New stratigraphic constraints for the Quaternary source-to-sink
 490 history of the Amatrice Basin (central Apennines, Italy), *Geological Journal*, 55, 4226–4251,
 491 <https://doi.org/10.1002/gj.3672>, 2020.

492 Mirabella, F., Buccì, F., Santangelo, M., Cardinali, M., Caielli, G., De Franco, R., Guzzetti, F., and Barchi, M. R.: Alluvial
 493 fan shifts and stream captures driven by extensional tectonics in central Italy, *Journal of the Geological Society*,
 494 [jgs2017-138](https://doi.org/10.1144/jgs2017-138), <https://doi.org/10.1144/jgs2017-138>, 2018.

495 Mondini, A. C., Chang, K., Rossi, M., Marchesini, I., and Guzzetti, F.: Semi-automatic recognition and mapping of event-
 496 induced landslides by exploiting multispectral satellite images and DEM in a Bayesian framework, 852415,
 497 <https://doi.org/10.1117/12.977432>, 2012.

498 Niculiță, M., Mărgărint, M. C. M. C., and Santangelo, M.: Archaeological evidence for Holocene landslide activity in the
 499 Eastern Carpathian lowland, *Quaternary International*, 415, <http://dx.doi.org/10.1016/j.quaint.2015.12.048>, 2016.

500 Pánek, T., Hartvich, F., Jankovská, V., Klimeš, J., Tábořík, P., Bubík, M., Smolková, V., and Hradecký, J.: Large Late
 501 Pleistocene landslides from the marginal slope of the Flysch Carpathians, *Landslides*, 11, 981–992,
 502 <https://doi.org/10.1007/s10346-013-0463-8>, 2014.

503 Pellicani, R. and Spilotro, G.: Evaluating the quality of landslide inventory maps: comparison between archive and surveyed
 504 inventories for the Daunia region (Apulia, Southern Italy), *Bull Eng Geol Environ*, 74, 357–367,
 505 <https://doi.org/10.1007/s10064-014-0639-z>, 2015.

506 Pellicani, R., Van Westen, C. J., and Spilotro, G.: Assessing landslide exposure in areas with limited landslide information,
 507 *Landslides*, 11, 463–480, <https://doi.org/10.1007/s10346-013-0386-4>, 2014a.

508 Pellicani, R., Frattini, P., and Spilotro, G.: Landslide susceptibility assessment in Apulian Southern Apennine: heuristic vs.
 509 statistical methods, *Environ Earth Sci*, 72, 1097–1108, <https://doi.org/10.1007/s12665-013-3026-3>, 2014b.

- 510 Petley, D.: Global patterns of loss of life from landslides, *Geology*, 40, 927–930, <https://doi.org/10.1130/G33217.1>, 2012.
- 511 Razak, K. A., Santangelo, M., Westen, C. J. V., Straatsma, M. W., and Jong, S. M. D.: Generating an optimal DTM from
512 airborne laser scanning data for landslide mapping in a tropical forest environment, *Geomorphology (Amst.)*, 190, 112–
513 125, <https://doi.org/10.1016/j.geomorph.2013.02.021>, 2013.
- 514 Santangelo, M., Gioia, D., Cardinali, M., Guzzetti, F., and Schiattarella, M.: Interplay between mass movement and fluvial
515 network organization: An example from southern Apennines, Italy, *Geomorphology (Amst.)*, 188, 54–67,
516 <https://doi.org/10.1016/j.geomorph.2012.12.008>, 2013.
- 517 Santangelo, M., Gioia, D., Cardinali, M., Guzzetti, F., and Schiattarella, M.: Landslide inventory map of the upper Sinni
518 River valley, Southern Italy, *Journal of Maps*, 11, 444–453, <https://doi.org/10.1080/17445647.2014.949313>, 2015.
- 519 Santangelo, M., Cardinali, M., Bucci, F., Fiorucci, F., and Mondini, A. C.: Exploring event landslide mapping using Sentinel-
520 1 SAR backscatter products, *Geomorphology*, 397, 108021, <https://doi.org/10.1016/j.geomorph.2021.108021>, 2022.
- 521 Schiattarella, M., Di Leo, P., Beneduce, P., and Ivo Giano, S.: Quaternary uplift vs tectonic loading: a case study from the
522 Lucanian Apennine, southern Italy, *Quaternary International*, 101–102, 239–251, [https://doi.org/10.1016/S1040-6182\(02\)00126-X](https://doi.org/10.1016/S1040-6182(02)00126-X), 2003.
- 523 Schiattarella, M., Giano, S. I., and Gioia, D.: Long-term geomorphological evolution of the axial zone of the Campania-
524 Lucania Apennine, southern Italy: a review, *Geologica Carpathica*, 68, 57–67, <https://doi.org/10.1515/geoca-2017-0005>, 2017.
- 525
526 Schulz, W. H.: *Landslides mapped using LIDAR imagery*, Seattle, Washington, 2004.
- 527 Spalluto, L., Fiore, A., Miccoli, M. N., and Parise, M.: Activity maps of multi-source mudslides from the Daunia Apennines
528 (Apulia, southern Italy), *Nat Hazards*, 106, 277–301, <https://doi.org/10.1007/s11069-020-04461-3>, 2021.
- 529 Thiery, Y., Terrier, M., Colas, B., Fressard, M., Maquaire, O., Grandjean, G., and Gourdier, S.: Improvement of landslide
530 hazard assessments for regulatory zoning in France: STATE-OF-THE-ART perspectives and considerations,
531 *International Journal of Disaster Risk Reduction*, 47, 101562, <https://doi.org/10.1016/j.ijdrr.2020.101562>, 2020.
- 532 Van Den Eeckhaut, M., Poesen, J., Verstraeten, G., Vanacker, V., Nyssen, J., Moeyersons, J., Beek, L. P. H. van, and
533 Vandekerckhove, L.: Use of LIDAR-derived images for mapping old landslides under forest, *Earth Surface Processes
534 and Landforms*, 32, 754–769, <https://doi.org/10.1002/esp.1417>, 2007.
- 535 Vignaroli, G., Mancini, M., Bucci, F., Cardinali, M., Cavinato, G. P., Moscatelli, M., Putignano, M. L., Sirianni, P.,
536 Santangelo, M., Ardizzone, F., Cosentino, G., Di Salvo, C., Fiorucci, F., Gaudiosi, I., Giallini, S., Messina, P., Peronace,
537 E., Polpetta, F., Reichenbach, P., Scionti, V., Simionato, M., and Stigliano, F.: Geology of the central part of the
538 Amatrice Basin (Central Apennines, Italy), *Journal of Maps*, 15, 193–202,
539 <https://doi.org/10.1080/17445647.2019.1570877>, 2019.
- 540 Vitale, S. and Ciarcia, S.: Tectono-stratigraphic and kinematic evolution of the southern Apennines/Calabria–Peloritani
541 Terrane system (Italy), *Tectonophysics*, 583, 164–182, <https://doi.org/10.1016/j.tecto.2012.11.004>, 2013.
- 542 Vitale, S., Ciarcia, S., Mazzoli, S., and Zaghoul, M. N.: Tectonic evolution of the ‘Liguride’ accretionary wedge in the
543 Cilento area, southern Italy: A record of early Apennine geodynamics, *Journal of Geodynamics*, 51, 25–36,
544 <https://doi.org/10.1016/j.jog.2010.06.002>, 2011.
- 545 Wasowski, J., Lamanna, C., and Casarano, D.: Influence of land-use change and precipitation patterns on landslide activity
546 in the Daunia Apennines, Italy, *Quarterly Journal of Engineering Geology and Hydrogeology*, 43, 387–401,
547 <https://doi.org/10.1144/1470-9236/08-101>, 2010.
- 548 Wasowski, J., Lamanna, C., Gigante, G., and Casarano, D.: High resolution satellite imagery analysis for inferring surface–
549 subsurface water relationships in unstable slopes, *Remote Sensing of Environment*, 124, 135–148,
550 <https://doi.org/10.1016/j.rse.2012.05.007>, 2012.
- 551 WP/WLI: International Geotechnical societies UNESCO Working Party on World Landslide Inventory. Multilingual
552 landslide glossary, Richmond: BiTech Publishers Ltd, 1993.
- 553 Zumpano, V., Ardizzone, F., Bucci, F., Cardinali, M., Fiorucci, F., Parise, M., Pisano, L., Reichenbach, P., Santaloia, F.,
554 Santangelo, M., Wasowski, J., and Lollino, P.: The relation of spatio-temporal distribution of landslides to urban
555 development (a case study from the Apulia region, Southern Italy), *Journal of Maps*, 0, 1–8,
556 <https://doi.org/10.1080/17445647.2020.1746417>, 2020.
- 557
558
559

560 Appendix A

561 In the first phase of work, information and cartographic/thematic products were collected to support the preparation of the
562 geomorphological landslide inventory map of the Daunia Apennines. Some of the listed products have been provided by the
563 Regione Puglia, others can be consulted on the web, through Cartographic Portals or Web servers:

- 564 ● Regional Technical Maps in digital format at a scale of 1:5,000, supplied by Regione Puglia (www.sit.puglia.it);
- 565 ● Geological maps at scale 1:50.000 (CARG), available on the ISPRA cartographic portal;
- 566 ● Geological maps at a scale of 1:100,000 (Geological Map of Italy), available on the ISPRA cartographic portal;
- 567 ● Hydrogeomorphological map in scale 1:25.000 available on the SIT site of the Regione Puglia
568 (<http://webapps.sit.puglia.it/freewebapps/Idrogeomorfologia/index.html>);
- 569 ● Digital Elevation Model (DEM, cell 8×8m) provided by Regione Puglia (www.sit.puglia.it);
- 570 ● Digital Elevation Model (DEM) made by LiDAR (1×1m cell), carried out within the framework of the Extraordinary
571 Plan for Environmental Remote Sensing (PST-A) and provided by the Ministry of Ecological Transition (MiTE).
- 572 ● Urbanised and land use maps, provided by the Regione Puglia (www.sit.puglia.it);
- 573 ● Orthophoto maps provided by the Regione Puglia (www.sit.puglia.it);
- 574 ● Landslide inventory maps of Apulia from:
 - 575 ■ **IFFI Inventory**: Inventory of Phenomena Franosi in Italy, compiled by ISPRA;
 - 576 ■ **Official Archives**: *Piano Stralcio per l'Assetto Idrogeologico* (PAI) of the Basin Authorities of Puglia,
577 Basilicata, of the Rivers Trigno, Biferno and minor Saccione and Fortore, and of the Rivers Liri-Garigliano
578 and Volturno; *Project Inventory of Phenomena Franosi in Italy (IFFI - update 2006)*, compiled by ISPRA;
579 *Census Project of Italian Areas Historically Vulnerated by Geological and Hydraulic Disasters (AVI)*, drawn
580 up by the National Group for Defence against geo-hydrological Disasters of the National Research Council
581 (GNDCI-CNR);
 - 582 ■ **AdBP photo interpretation**: landslides from the photo interpretation study conducted by AdBP (Basin
583 Authority of Puglia Region);
 - 584 ■ **Province of Foggia**: it deals with the landslides surveyed by the Province of Foggia within the framework
585 of the agreements signed with the AdBP concerning the "Activity of collection and classification of data on
586 landslide phenomena in the Province of Foggia" and "Conduct of studies for the deepening of the aspects
587 related to the classified landslide phenomena".